

## **Response to review comments**

We first give a general response, before responding to the reviewers' comments.

### **General response**

*We thank the reviewer for helpful comments, which we feel have strengthened the article.*

*Based on the reviewers' comments, we made a number of small changes to increase clarity, corrected spelling mistakes, and expanded on examples in the results. We have also references other relevant work. We have clearly mentioned the importance of correlations in this analysis, both in the abstract, methods and conclusion. We have written two large paragraphs discussing the issue in the discussion section. We are happy with the changes and feel they have improved the article.*

### **Response to reviewer**

Reviewer's comments are in *italic*, while our responses are in standard font.

#### **Referee #1**

##### **General comments**

*The authors use a GTAP-based I/O model to simulate pollutant emissions from a consumption perspective. They estimate uncertainties in economic data and trace these uncertainties through the I/O model to emissions estimates by perturbing the GTAP tables used to calibrate the I/O model. The paper is well written and clear for the most part, and is a nearly-comprehensive look at the topic (with one major caveat described below).*

*Given the audience of ESD, I would suggest that some less abstract examples be included regarding the data and parameters that are being perturbed. For instance, the (rather complex) procedure for estimating variance in each economic flow value is described in depth in 2.3, but then examples of what this means for particular values are not given until 3.1 (for the examples of trade flows from China and etc.). I would suggest moving these examples to 2.3 and expanding on them a bit so the reader understands why this flow ends up with such a low uncertainty. This will have the added benefit of streamlining the results section to focus on the results in aggregate rather than the estimates of particular flow values, which is really more methods than results.*

We thank the reviewer for pointing this out. While we feel it is inappropriate to move one of the key results from the results section and in to the methods section, we see the need for better explanations. In the methods section, we have added text to indicate what other studies have found using the same general relationship: "Implementing this general methodology has given individual regions relatively small uncertainties in other studies (Lenzen et al., 2010; Wiedmann et al., 2008). Structural uncertainties have also recently been found to be relatively small for major economies (Moran and Wood, 2014)."

Furthermore, we have expanded on the examples mentioned by the reviewer in section 3.1 to help explain why the uncertainties end up relatively low: "These fluxes are mainly dominated by the largest sectors, which have been given the smallest uncertainties in the respective regions. The

export from China to USA is mainly coming from the manufacturing sectors, which combined is one of the largest Chinese sectors, hence with lower uncertainties. Annex B countries are given lower uncertainties than non-annex B countries, which explains the low uncertainty from USA to Western Europe.”.

*Important to connect this work to emissions uncertainty work using dynamic GTAP-based models. I'm not as familiar with what is available on the pure I/O accounting type models, so I assume the existing discussion of previous work is sufficient, but there is certainly some work using dynamic GE models that is relevant. Richard Plevin's dissertation work for instance is highly relevant and a good complete take on the subject from a CGE/life-cycle perspective (Plevin 2010). My group has also done work on this topic in the CGE context, which might be interesting (see for instance Elliott et al 2011 and Elliott et al 2010).*

We have mentioned the work done on dynamic CGE models in the introduction, and referenced Elliott et al 2012: “Other economic models, such as computable general equilibrium models, have also received attention recently with regards to uncertainties (Elliott et al., 2012).”

*One step that is ignored in the causal chain defined here (consumption->production->emissions->climate) but also introduces uncertainty is the mapping from emissions to atmospheric concentration. This is especially serious for the carbon cycle which has complex positive and negative feedbacks with both the ocean and land operating a different important time-scales (Glotter et al 2014).*

The carbon cycle is included in the metric equations, and reflected by CO<sub>2</sub>'s IRF. The uncertainty on the parameters is based on the model comparison (Joos et al., 2013). We perturb the parameters that go into explaining the feedbacks of the relatively quick response of the land surfaces, shallow oceans and atmosphere, and the relative slow response of the deep oceans. Additionally, the positive and negative feedback effects in the climate system are included in the climate sensitivity parameter, which is also given an estimated uncertainty. The uncertainties in the link from emissions to atmospheric concentrations is thus included, and mentioned in section 2.5 Emission metrics. To clarify this in the paper, we added text in the results section 2.5 (inserted text in *italic*): “Metric (temperature) values have an uncertainty range for the different pollutants and different time horizons, due to the perturbed metric parameters (RF, lifetime, and climate sensitivity). *This includes uncertainties from mapping emissions to atmospheric concentrations through the global carbon cycle, which is represented by the relatively uncertain climate sensitivity.*”

*Probably the biggest concern I have is correlation of data error. Certainly its understandable (as stated on page 1021 line 23-25) that little information exists about correlations, but in order to make the rather strong comparative assertions made in the paper (for example in the abstract pg1014 line 13-18) it is absolutely crucial to address some of these possible correlations, at least qualitatively and hopefully quantitatively. If you assume fully uncorrelated errors in data or parameters, then of course it's not surprising that uncertainty in high-resolution data/parameters such as sectors within countries would have a smaller effect on global variables like cumulative emissions or temperature than large-scale global parameters like climate sensitivity. It certainly may be that the estimated emissions factor for cement production (for example) has highly correlated error (if, for instance, sector emissions tend to be under-reported everywhere). A scenario using this sort of perturbation on the underlying data would likely find a much stronger impact on emissions and climate than one*

*assuming that the emissions factor for cement industry in each country had completely uncorrelated errors. At the very least this puts a strong caveat on the key assertion of the paper (that economic uncertainty is of less importance to global metrics) which seems fundamental to the point of the paper and really must be addressed.*

*Given that the paper is otherwise nicely comprehensive in most aspects, it's disappointing (and I think a real missed opportunity) to not include something on the topic of correlated error. At the least I would hope that some synthetic examples of correlated error could be evaluated in order to test whether the strong assertions still hold. For instance, a scenario assuming correlated errors in all sectors within a country but uncorrelated between countries and another considering correlated errors in a given sector between countries but no correlation between sectors. The actual amount of correlation in these scenarios is probably not even all that important, although one could estimate values of correlation that lead to certain critical thresholds based on the relative uncertainty of the factors described (what level of correlation would be required for the uncertainty from econ data in global temp to rival the uncertainty from climate sensitivity). I suspect you would find that this critical correlation is not actually very large at all.*

We agree with the reviewer on the importance of correlations, and it was amiss of us not to emphasize this in the article. We certainly acknowledge the need for addressing correlations in uncertainties of the datasets. In cases of high correlation, this may change the aggregated results substantially. We (as authors) have had many discussions on the issue of different types of correlations, particularly in the economic data, and how to feasibly and realistically include this. Through the review, we have revisited this issue by trying to implement correlations on a smaller scale. After trying to implement this, we have come to realize that this is a larger problem than we initially thought for a variety of reasons, which we discuss further down. However, we have now put more emphasis on the "caveat" by noting it clearly in the abstract, methods, discussion and conclusion section. In the discussion, we have added an extended section which discusses the issue of the correlation in economic data. Overall though, to do a thorough analysis of correlations in economic data is a significant undertaking which we see as beyond the scope of this paper. It is nevertheless an issue we continue to work on and would like to get a better understanding of.

The example mentioned by the reviewer about emission factors for cement production may be correlated across countries, is very plausible, as a similar chemical process is occurring everywhere where production is happening. However, the datasets we use consists of emissions, where the emission factors have been multiplied with energy use (and the emissions that follows from energy use). The emission factor used may be correlated across countries because calculations may have used a global or regional number for this, but the energy use is more likely to be uncorrelated across countries as production technology and production recipes varies across countries. Thus it is difficult to estimate the extent of correlation in the emissions data we use.

The review comment focused on correlation in errors, while we are of the opinion that correlation in data points is more relevant (particularly given balancing issues). As an example, we might have decent constraints on the total oil consumed in the country, and if the oil consumption in one sector is overestimated, then this implies an underestimate in another (anti-correlation). We included this correlation in the emissions data (using the quadratic programming routine), but because of the

extreme computational need we decided not to include in the IO data. Balancing the data will also introduce correlations, but to balance each MC run would be computationally expensive.

On the computation aspects, even if we had a correlation structure to apply, we at best could only do this in small scale problems. Given a dataset has  $N$  points, the correlation matrix has  $N^2$  data points. Our economic system has about 7500x7500 data points, which means the correlation matrix is prohibitively large for most computers. Given we have no real data on correlations, populating this matrix is guess work. But, given the size, even if we could populate it, the computational issue prohibits using it.

Generating log-normal numbers with correlation also poses a difficulty. Log-normal numbers are important in our context as they are never negative, which is important in terms of both economic and emissions data where many numbers are small in magnitude but with high uncertainty. If correlations were linear on a log scale, they would not be showing linear relationships between the parameters in a non-log domain, and that would make it difficult to implement what we want and to interpret the results.

As a final note, we stress that we agree that correlations are important and that they may take many forms in our datasets. The inclusion of them may also change our results and conclusions, mainly with respect to the economic errors in comparison to other errors. It is important to note that we have discussed this issue in detail, but it was a mistake not to emphasize this issue in the original submissions. However, with large difficulties in implementing them, and no actual data to base them on, we feel we have no other choice than to just flag the issue in the text and let the reader make up his own opinion about the value of the results. As mentioned, this is an issue we continue to work on.

To clarify this, we added text in the abstract: “Based on our assumptions, which does not account for correlations, the economic data appear to have a relatively small impact on uncertainty at the global and national level...”

We also mentioned this in the methods section: “Implementing correlations in such an analysis is a major difficulty, but may also have significant effects on the results. See the discussion in section 4.”

Furthermore, we discussed this in the discussion section: “A major difficulty not included in the economic and emissions data is the issue of correlations. There is a large need for addressing correlations in datasets and uncertainties, as this may have significant impacts and the results of such an analysis. We have explored correlations for metric uncertainties (temperature and CO<sub>2</sub> IRF), and in one sense introduced correlations in that we make all non-Annex B countries have double uncertainties of Annex B countries. However, correlations may further be an issue in several places in the datasets and methods which we have not included, and we see at least three places where they may be important: (1) in the way the MC analysis is build up where uncertainties given to certain region/sector combinations may be correlated in each run in order to simulate corresponding behavior in the model (if Norway’s emissions from cement production in one run is low, then Sweden’s emissions from the same sector may also be low in order to build a plausible world in each MC iteration), (2) overall (median) uncertainties in certain region/sector combinations could be similar (spatial correlations; if Norway’s emissions from agriculture is low, then Sweden’s emissions from the same sector could also be low, due to similar technology, statistical offices using similar

methods, etc.), and (3) between datasets (a perturbation in e.g. fossil fuel use in the economic dataset should be reflected by a similar correlating perturbation in the emissions dataset).

Implementing these correlations, which can be argued to be important for some sector/region combinations, may clearly change the uncertainty outcome. However, we have not included correlations in the economic and emissions data due to computational and conceptual issues: there is little or no data indicating correlations in uncertainties in sectoral economic data or emissions data although correlations might be plausible (thus populating a correlation matrix is guess work), the generation of log-normal numbers with correlations is not as straight forward as with a normal distribution as they would not be showing linear relationships between the parameters in a non-log domain, and due to the large datasets used in this analysis, the correlation matrix would be prohibitively large, posing serious computational issues. Thus, this is an issue for future work to investigate.”

We concluded in the conclusion section: “We did not account for correlations in the economic or emissions data, which may play an important role and have a significant impact on the results.”

### **Smaller questions**

*1018/12-13: Does the I/O model here use the full GTAP region/sector resolution without any aggregation?*

Yes. This has been mentioned explicitly in section 2.1: “We use these data to construct an MRIO model without any aggregation, which connects all regions at the sector level...”. We furthermore state in section 3.4: “To facilitate our discussion we aggregate the 58 economic sectors (post analysis) to 9 sectors.”

*1023: The explanation for how to construct relative uncertainty in each GTAP value needs clarification. My impression is that the MRIO model is a traditional accounting type I/O model and thus the only “parameters” in the model are relative consumption/production/trade shares. This means there is a very simply mapping from the GTAP values to the parameters, but this isn’t made clear. I understand that the model construction is explained elsewhere, but at least this one piece of information is fundamental enough that it should be described here.*

We thank the reviewer for pointing out this possible confusion. All GTAP values are given uncertainties and distributions, not only the trade shares. This has been mentioned in section 2.3: “In other words, we estimate the uncertainty of the MRIO data based on the uncertainty in the data used to construct it, which consists of all data points in the GTAP database used to construct the MRIO model.”

*1024/18: This should be restated for clarification: “To retain balance, we therefore choose not to rebalance. . .”*

We thank the reviewer for point out this typo. The sentence has been modified to: “We therefore choose not to rebalance, which effectively causes the “unbalanced” component to be shifted to the value added.”

1025/18: typo? “. . .emissions with roughly 12%...” should this be “. . .by roughly. . .”?

This has been changed to: “...by roughly...”.

1027/25: *You might take a look at Pierrehumbert (2014) for a useful critique of GTP. I’m convinced that you are using it appropriately in this case (though I’m no expert) but it may be useful for context in the discussion.*

We thank the reviewer for this reference. The reference has been cited in the methods section (inserted text in *italic*): “Although it has been shown that the GTP may have larger relative uncertainties than the alternative metric global warming potential (GWP) (Aamaas et al., 2013; Reisinger et al., 2010) *and it has been critiqued for some of its characteristics (Pierrehumbert, 2014)*, the GTP directly links to global temperature change and is thus arguably more policy relevant (Shine et al., 2005). In addition, the physical interpretation of the GWP is less clear and the metric has been criticized by many authors (Peters et al., 2011a; Shine, 2009; *Pierrehumbert, 2014*).”

1032/10: *these uncertainties seem very small to me intuitively, and I suspect the casual reader will agree. So this should be justified in much greater detail I think. I suppose the explanation is again due to cancellation of uncorrelated errors, and I again wonder how different the answer would be if you introduced even a small amount of correlation.*

This section has been moved to the Methods section according to a previous comment, and the discussion on the examples has been extended. In the results section, we added: “See the examples and the discussion on why the regional uncertainties are so low in the Methods section.” On the issue of correlations, see previous discussion.

1032/21-22: *a vast number? That’s pretty subjective. Can you say how many operations are required?*

Building a procedure to count all operations in the model is a large undertaking, but we have estimated that just the matrix inversion (which is done 10000 times in the MC model) requires more than  $10^{12}$  operations. This has now been mentioned in the text (inserted text in *italic*): “Since we start from the raw GTAP data to construct the MRIO table, and normalize and invert the MRIO table, a vast number of summations and multiplications are done with the initial perturbed data (*only inversion requires more than  $10^{12}$  operations*).”

1033/9-12: *I’m not convinced by the argument for discounting the GTAP uncertainty accounting in favor of the highly complex and somewhat ad hoc approach described in section 2. Surely if different methods for estimating uncertainty can give vastly different results then this must be accounted for, given that this is precisely the point of “uncertainty”. Its fine to choose one for the paper but you must at least test your conclusions against the alternatives, which it doesn’t seem that you do (indeed, given the very large uncertainty implied by the GTAP estimates, I’m assuming your conclusions would no longer be valid in this context). Explaining away the GTAP estimates because the model structure can’t handle it or because you don’t have the computational power to rebalance the matrices is not a convincing argument for saying that the uncertainty is not that large. . .*

According to GTAP (McDougall, 2006), the Table 19.6 in the GTAP documentation consists of “large sectors in large regions with large relative changes”. This is to show and explain some of the largest changes: “Of the more than eleven thousand (region, category) pairs, we select those that make the

largest contribution to the entropy distance measure". This is thus not a good representation of the overall uncertainties in the dataset, as it only deals with a small number of outliers. Further, these values represent those data points that move most in the GTAP balancing procedure, which may not reflect uncertainty. We use these numbers explicitly as a sensitivity analysis to see what happens if we assume that all data points are having the same data size/uncertainty relationship, but we cannot assume that this is representative for the whole dataset in the main analysis. Our approach may underestimate uncertainties, but we feel this is based on more sound assumptions than making Table 19.6 representative for the whole dataset. It is important to note that we do not exclude the GTAP data. We use the statistical relationship, but apply the parameters from the UK economic data.

*1033/21: I'm worried about the many uses of the phrase "we find small uncertainties". It seems to me that you are assuming small uncertainties by the structure of your methodology, rather than "finding" them.*

The structure of the methodology for estimating uncertainty in the economic data and emissions data is very similar, and thus they were expected to behave similarly. We found this not to be the case, and thus use the phrase "we find small uncertainties" several times to underline the sometimes counterintuitive results.

We should emphasize that others have found similarly small uncertainties (Moran and Wood, 2014). It is correct that this may be a consequence of "garbage in, garbage out", but we have attempted to make our error estimates follow any available literature. We additionally include a sensitivity analysis with larger uncertainties. To us, the issue of correlations is the main place that could introduce larger errors given the input we use.

*1035/15-18: how have you handled the natural gas sector? My experience is that GTAP has 2 sectors for gas, gas extraction and gas delivery, and these must actually be combined to get a consistent treatment of the gas sector. I recall this is quite tricky for tracking natural gas carbon.*

It is correct that GTAP has two natural gas sectors. Bear in mind that we use EDGAR emissions data, not emissions data from GTAP. Fugitive emissions from gas extraction is allocated to the gas extraction sector (which appear in the mining sector in our aggregated results), while a part of the emissions from public electricity and heat production are allocated to the gas distribution sector (which is part of the aggregated service sector). The gas distribution sector is the major consumer of the gas extraction sector, although a relatively small amount of emissions are allocated to this trade link as most emissions from gas (in a production view) are allocated to the gas distribution sector, which consuming regions and sectors purchase from.

*1035/15-18: how are you handling refined petroleum products? Refineries produce a huge diversity of products with different emissions profiles. Tracking this downstream to the consumers of refined petroleum is not easy. I think Elliott et al. 2010 (in the "carbon accounting" section) describes some of this using a simple example.*

The global supply-chain (using MRIO analysis) explains the link between extractions of petroleum (sectors including coal, oil, gas), production of petroleum based products and demand by consumers (sector refined petroleum). Thus, oil/gas extracted is sent to the refined petroleum sector. A consistent treatment is used in the emissions datasets (emissions from extraction in the coal, oil, gas

sectors, and emissions from refining in the refinery sector). Refined petroleum products is a single sector in the GTAP database, and we do not disaggregate this sector, which means that this sector is directly linked from production to consumption.

*1040/6: should be "individual MC ensembles" I believe, not "runs".*

This has been changed to "individual MC ensembles".

*1044/24: typo ". . .but is this. . ."*

This has been changed to "...this is...".

*1045/7-10: I'm not sure why you would expect consumption uncertainty to be higher if you don't account for factors such as the uncertain distribution of carbon in multiproduct outputs from different sectors, with the most important example being refined petroleum and coal products. Coke sold to steel manufactures has very different emissions than does gasoline sold to consumers or jet fuel sold to airline services. If you acknowledge that there is additional uncertainty introduced at every step of the carbon accounting flow from production to consumption, then I suspect you would find the consumption perspective much more uncertain.*

Intuitively, we expected the addition of uncertain data in the analysis (economic data) to significantly contribute to uncertainty in the end results. However, due to cancellations of errors, we found that there were only small errors from the economic dataset on an aggregated level. We do not track individual products, but agree that the uncertainties would be higher if we did (due to uncertainties in disaggregation).

*1046/26: typo "emissions uncertainties often dominate over emission uncertainties".*

This has been changed to: "emissions uncertainties often dominate over metric uncertainties".

*1047/5-6: I have a hard time with this. It seems like GTAP has tried, however imperfectly, to estimate uncertainty in their data. However the authors have chosen to ignore these estimates seemingly because they are "too uncertain". Instead they have specified a much small "uncertainty" which is really not uncertainty because obviously the true distribution of possible values is much larger if even the very group that synthesizes and releases this data is not comfortable putting anything smaller than huge error bars on it. The authors could described the ensembles they create as using "perturbations" specified using xyz methods and assumptions, but describing them as "uncertainty" is not right.*

The GTAP community has never published uncertainty data with their datasets, as far as we are aware. The table we refer to is only comparing numbers that is used as input and by how much they are changed after harmonization and balancing procedures are done (see answer to comment 1033/9-12 above). We choose to use this relationship as a function explaining the relationship between sector sizes and uncertainties. The table itself consists only of large sectors in large countries, having large changes. Thus it is only valid for these data points, and cannot be directly used to explain uncertainties on all sectors in all regions. We used this data as a test to see how this would affect the results (simple sensitivity analysis), but did no further analysis on this as we have no information on how this is representative for other data points. The word "uncertainty" referred to



here, points not only to economic uncertainties, but also uncertainties in metric parameters and emissions data, and thus the studies we refer to.

In the results section we refer to the GTAP table as “uncertainties”, which was unfortunate, and we have now changed the wording: “The “unfitted” and “fitted” data from Table 19.6 in the GTAP documentation (Fig. 2), however, act as a simple sensitivity analysis to our applied uncertainties.”

## Conclusion

*Overall I think this work has the potential to be a comprehensive take on carbon accounting and uncertainty, but it falls short in essential ways that must be addressed. The paper is detailed, comprehensive and well written, but it makes strong (and probably inaccurate or at least incomplete) conclusions that depend fundamentally on the assumptions made in setting up the problem (small uncorrelated errors in individual economic flow values). It does consider some limited alternative scenarios, but then discounts them without considering how they affect the conclusions.*

## References

- Plevin, Richard Jay. "Life cycle regulation of transportation fuels: Uncertainty and its policy implications." (2010). <http://plevin.berkeley.edu/docs/Plevin-Dissertation-2010.pdf>
- Elliott, J., M. Franklin, I. Foster, M. Loudermilk and T. Munson (2011). Propagation of Data Error and Parametric Sensitivity in Computable General Equilibrium Models, Computational Economics: 1-23. DOI: 10.1007/s10614-010-9248-5 <http://link.springer.com/article/10.1007/s10614-010-9248-5>
- Elliott, J., I. Foster, K. Judd, E. Moyer, and T. Munson (2010). CIM-EARTH: Framework and Case Study, The B.E. Journal of Economic Analysis & Policy: Vol. 10 : Iss. 2, Article 11. <https://ideas.repec.org/a/bpj/bejeap/v10y2010i2n11.html>
- M. Glotter, R. Pierrehumbert, J. Elliott, and E. Moyer. A simple carbon cycle representation for economic and policy analyses. Climatic Change 1-17, 2014.
- R.T. Pierrehumbert. Short-Lived Climate Pollution. Annual Review of Earth and Planetary Sciences, Vol. 42: 341-379 (Volume publication date May 2014).

## References

- Joos, F., Roth, R., Fuglestedt, J., Peters, G., Enting, I., Bloh, W. v., Brovkin, V., Burke, E., Eby, M., and Edwards, N.: Carbon dioxide and climate impulse response functions for the computation of greenhouse gas metrics: a multi-model analysis, Atmospheric Chemistry and Physics, 13, 2793-2825, 2013.
- McDougall, R. A.: Chapter 19: Updating and Adjusting the Regional Input–Output Tables, in: Global Trade, Assistance, and Production: The GTAP 6 Data Base, edited by: Dimaranan, B. V., Center for Global Trade Analysis, Purdue University, 2006.
- Moran, D., and Wood, R.: CONVERGENCE BETWEEN THE EORA, WIOD, EXIOBASE, AND OPENEU'S CONSUMPTION-BASED CARBON ACCOUNTS, Economic Systems Research, 26, 245-261, 10.1080/09535314.2014.935298, 2014.



## Response to review comments

We first give a general response, before responding to the reviewers' comments.

### General response

*We thank the reviewer for helpful comments, which have strengthened the article.*

*Based on the reviewer's comments, we made a number of small changes to increase clarity and changed a reference mistake. We also extended Table 7 to include uncertainty results on GWP, and discussed this in the results. Furthermore, we added references to better justify our assumptions. We are happy with the changes and feel they have improved the article.*

### Response to reviewer

Reviewer's comments are in *italic*, while our responses are in standard font.

#### Anonymous Referee #2

*This article addresses a very interesting and current topic, which is the estimation of consumption- vs production-based GHG emissions as well as related uncertainties. It is very well written. My scientific background is in the field of climate metrics and carbon accounting. I thus had a hard time understanding some parts of the method dealing with uncertainties associated to economic data. I know basically what GTAP and input/output models are, but I am not comfortable with all the details. The introduction and the first paragraphs of the method section (from the beginning to page 1019 line 14) present very clearly the scope of the paper and the work that has been done.*

*I do not understand where uncertainties for economic data are coming from. The authors are referring to McDougall 2001, but this reference is not listed, so I could not take a look at it. I understand that uncertainties are estimated using previous studies since no uncertainties are available for the GTAP datasets (Equation 1). This equation is parameterized using only two points (min and max) and Equations 2 and 3. However, I do not understand how  $v_{min}$ ,  $v_{max}$ ,  $r_{min}$  and  $r_{max}$  are determined. I do not understand neither how Figure 2 was obtained. If uncertainties are determined using Equation 1, we should see only a trend line. Where are the points coming from? At page 1023 line 14 and following, the authors are talking about the calibration of the uncertainty relationship given by Equation 1. I still do not understand what they mean. "For the smallest sectors we set  $v_{min}$  equal to 1 USD and assume  $r_{max} = 100\%$ , due to the lack of more precise regional uncertainty data." What is the basis of this assumption? My misunderstanding may be caused by my lack of scientific background in the subject (economic modeling), but I assume that other readers of this journal are in the same situation. The same approach is used to estimate uncertainties for emissions. Is Equation 1 also valid for this application?*

We thank the reviewer for pointing out this careless reference mistake. McDougall (2006) has now been updated and listed in the reference list. The data points in Table 19.6 in McDougall (2006) is plotted as red and blue circles in Figure 2, which differ depending on how you estimate the uncertainty (see discussion on page 1019:27). In order to clarify this, we added text in the figure caption (inserted text in *italic*): "Error distribution of selected GTAP input-output data (*taken from*

*Table 19.6 in McDougall (2006) and shown as colored circles), and trendlines showing the fit of the general functional relationship explained by Eq. (1). Red and blue circles differ due to different methods of estimating the uncertainty. See the discussion in the text.”*

Using first order power regression fit, we also show trend lines in Figure 2 using Equation 1, which fits well with the observations ( $R^2 > 0.9$ ). This indicates that Equation 1 can be used to represent the relationship between economic sectors' size and uncertainty. This is also what previous work on uncertainties in GTAP data has found (Lenzen et al., 2010; Wiedmann et al., 2008). We do not use the parameters from the regression directly since Table 19.6 is not a representative selection of the dataset, but rather show outliers. As stated in the manuscript, very little information is available in order to populate the parameters in Equation 1. Because of this, we use Equation 2 and 3 to parameterize the relationship with two extreme data points, both in terms of relative errors ( $r_{min}$  and  $r_{max}$ ) and at what sector sizes ( $v_{min}$  and  $v_{max}$ ) these errors should be given.

The uncertainties are difficult to estimate, but we use the trend lines of Figure 2 to explain the uncertainty on the largest sectors ( $r_{min}$ ), which the table is representing. The largest sector in each country varies, thus this is a function of the region's economy ( $v_{max} = 4\% \times GDP$ ). The lower threshold  $v_{min}$  has been set to 1 USD with  $r_{max} = 100\%$  uncertainty, which has been discussed in a paragraph, starting at page 1023:28, and has also been used in a recent study (Wiedmann et al., 2008). To clarify, we added text to section 2.3 (inserted text in *italic*): “For the smallest sectors we set  $v_{min}$  equal to 1 USD and assume  $r_{max}=100\%$  (following Wiedmann et al., 2008), due to the lack of more precise regional uncertainty data.”

Using these parameters for Equation 1, we show how this works for developed and developing countries in Figure 3. Each region has its own line, and every sector in each region will be on this line, depending on the sectors' size. To clarify this, we added text to the figure caption (inserted text in *italic*): “Functional relationship between sector sizes on horizontal axis (in kt CO2 emissions and million US dollars, respectively) and relative uncertainty on vertical axis. The red lines outline the range of developing regions, while the blue lines show the range of developed countries. *Each region has been estimated a single unique line, and all sectors, depending on their size, will fall on this line.* The form of this relationship is established independently for each pollutant.”

Equation 1 is also used to explain the relationship between emissions and uncertainties, due to lack of regional data, which has also been used in previous similar studies (Jackson et al., 2009; Lenzen et al., 2010). To clarify this, we added text in Section 2.2 (inserted text in *italic*): “Furthermore, we assume a similar relationship with the emissions data, based on a previous study of the UK Greenhouse Gas Inventory, where uncertainties were found using an error propagation model (Jackson et al., 2009). *This assumption is also shared by a recent study (Lenzen et al., 2010).*”

*Section 2.5: Everything looks as the state-of-the-art regarding climate metrics.*

*Page 1033 lines 5-20: I cannot understand this paragraph because I did not understand how uncertainties for economic data have been estimated. The authors seem to say that uncertainties are*

*provided in the GTAP documentation. Why did they not use them? I do not understand this explanation.*

See explanation above for how we estimated uncertainties. We thank the reviewer for pointing out the possible confusion in this paragraph. To clarify, we changed the text so that we are not referring to the Table 19.6 in McDougall (2006) as “uncertainties”, but rather as “unfitted” input data and “fitted” datasets (inserted text in *italic*): “The *“unfitted” and “fitted” data* from Table 19.6 in the GTAP documentation (Fig. 2), however, act as a simple sensitivity analysis to our applied uncertainties.” Furthermore: “*Thus, the results are sensitive to the input uncertainties, as this exercise has shown.*”

*The presentation of results is very clear.*

*Discussion:*

*Would there be a way to look at the sensitivity of the results to the different limitations identified regarding uncertainties for economic data which seem to be much less reliable than the two other types of uncertainty (crude assumptions, only parametric uncertainties, etc.)?*

Given the challenges with the Monte Carlo analysis with regards to the economic data, model comparisons (structural uncertainties) may be a better way of estimating economic uncertainties. This has just been done by Moran and Wood (2014), and we have now mentioned this in the manuscript (inserted text in *italic*): “It is often assumed that consumption-based emissions are more uncertain (Peters, 2008), but parametric uncertainty analysis shows that the uncertainties are small; structural uncertainties may be larger (Peters et al., 2012; Moran and Wood, 2014). In a recent study by Moran and Wood (2014), they find that most major economies’ carbon footprint results disagree by less than 10%.”

*What would be the impact on the results if GWP was used instead of GTP? Since the parameters used to calculate GWP are also used for GTP, this sensitivity analysis would not require more data. I understand the arguments in favor of GTP compared to GWP. However, GWP is still used everywhere and to see the difference using both indicators would be interesting.*

We thank the reviewer for pointing out this, and agree that it would add valuable information to the paper if results using GWP were also available. To make the comparison possible, we have extended Table 7 to also include results using GWP20, GWP50 and GWP100. We also added a paragraph discussion these results: “Table 7 illustrates the difference between uncertainties in AGTP, and GTP and GWP values.” “GWP calculations use the same parameters as with GTP, and although we do not use GWP in our results, we include the uncertainties in the table for comparison. Overall, we find less uncertainty using GWP than the other metrics, except for NOX. The GWP calculations are not dependent on the highly uncertain climate sensitivity, since it does not relate to global temperature change. Thus it is expected to have lower uncertainties. NOX has overlapping indirect effects, with highly uncertain RF values, which suggests that the GWP20 values can be both negative and positive, with a median close to zero. Thus it has a very high uncertainty.”

*Figures are not readable when printed in black and white. They are correct on the website with colors. However, if the authors wish that people could read the article when printed in black and white, it is currently not possible to understand most of the figures.*

We acknowledge the problem when printing in black and white, but feel that the colors add substantial information which would otherwise be difficult to show. E.g. Figure 6, 10 and 11 uses at least 14 different colors, which would be difficult to interpret if we could only use dotted and dashed lines and fills.

*Page 1019 line22: I cannot find the McDougall 2001 reference in the reference list.*

We thank the reviewer for pointing out this careless mistake, and have added and updated the reference to the list in the manuscript.

*Page 1020 line 23 and page 1026 line 13: What is the basis for this assumption (relative uncertainties for developing countries are twice the relative uncertainties for developed countries)? Why would it not be three, four or five times?*

We agree that multiplying uncertainties in developing regions with 2 may seem arbitrary. Many argue that uncertainties in developing regions would be higher, but no datasets exists that allow us to derive this relationship on a regional level as far as we know. However, other studies points in this direction: Andres et al. 2012 shows that global independent CO2 emission statistics generally agree within about 5% in developed regions and 10% for developing regions, thus twice the uncertainty for developing regions. To clarify this, we added references to the manuscript where we mention this (inserted text in *italic*): “The terms  $r_{min}$  and  $r_{max}$  define the smallest and largest relative errors, respectively, and are functions of developed and developing regions where the latter is given twice the uncertainties of the first group using the Kyoto Protocol groupings of Annex B and non-Annex B countries (e.g. CO2 emissions are found to have twice the uncertainty in developing regions than developed regions (Andres et al., 2012), see discussion in the next section).”.

## References

- Jackson, J., Choudrie, S., Thistlethwaite, G., Passant, N., Murrells, T., Watterson, J., Mobbs, D., Cardenas, L., Thomson, A., Leech, A., Li, Y., Manning, A., Walker, C., Brophy, N., Sneddon, S., Pierce, M., Thomas, J., and Brown, K.: UK Greenhouse Gas Inventory 1990 to 2007 - ANNEX 8: Uncertainties0955482380, 2009.
- Lenzen, M., Wood, R., and Wiedmann, T.: Uncertainty analysis for multi-region input–output models—a case study of the UK's carbon footprint, *Economic Systems Research*, 22, 43-63, 2010.
- McDougall, R. A.: Chapter 19: Updating and Adjusting the Regional Input–Output Tables, in: *Global Trade, Assistance, and Production: The GTAP 6 Data Base*, edited by: Dimaranan, B. V., Center for Global Trade Analysis, Purdue University, 2006.
- Moran, D., and Wood, R.: CONVERGENCE BETWEEN THE EORA, WIOD, EXIOBASE, AND OPENEU'S CONSUMPTION-BASED CARBON ACCOUNTS, *Economic Systems Research*, 26, 245-261, 10.1080/09535314.2014.935298, 2014.
- Wiedmann, T., Lenzen, M., and Wood, R.: Uncertainty Analysis of the UK-MRIO Model—Results from a Monte-Carlo Analysis of the UK Multi-Region Input-Output Model (Embedded Carbon Dioxide Emissions Indicator), Report to the UK Department for Environment, Food and Rural

Affairs by Stockholm Environment Institute at the University of York and Centre for Integrated Sustainability Analysis at the University of Sydney, Defra, London, Project Ref.: EV02033, 2008.

1 **Uncertainty in temperature response of current consumption-based emissions estimates**

2 J. Karstensen<sup>1</sup>, G. P. Peters<sup>1</sup> and R. M. Andrew<sup>1</sup>

3 | <sup>1</sup>Center for international climate and environmental research – Oslo (CICERO), P.O. Box. 1129  
4 Blindern, N-0318 Oslo, NORWAY

5 Correspondence to: [jonas.karstensen@cicero.oslo.no](mailto:jonas.karstensen@cicero.oslo.no)

6

7



## 8 Abstract

9 Several studies have connected emissions of greenhouse gases to economic and trade data to quantify  
10 the causal chain from consumption to emissions and climate change. These studies usually combine  
11 data and models originating from different sources, making it difficult to estimate uncertainties  
12 ~~in~~ along the ~~end results~~ entire casual chain. We estimate uncertainties in economic data, multi-pollutant  
13 emission statistics and metric parameters, and use Monte Carlo analysis to quantify contributions to  
14 uncertainty and to determine how uncertainty propagates to estimates of global temperature change  
15 from regional and sectoral territorial- and consumption-based emissions for the year 2007. We find  
16 that the uncertainties are sensitive to the emission allocations, mix of pollutants included, the metric  
17 and its time horizon, and the level of aggregation of the results. Uncertainties in the final results are  
18 largely dominated by the climate sensitivity and the parameters associated with the warming effects of  
19 CO<sub>2</sub>. ~~The~~ Based on our assumptions, which exclude correlations in the economic data, ~~the uncertainty~~  
20 ~~in the economic data appear to~~ have a relatively small impact on uncertainty at the ~~global and~~ national  
21 level, ~~while much in comparison to emission and metric uncertainty. Much~~ higher uncertainties are  
22 found at the sectoral level. Our results suggest that consumption-based national emissions are not  
23 significantly more uncertain than the corresponding production based emissions, since the largest  
24 uncertainties are due to metric and emissions which affect both perspectives equally. The two  
25 perspectives exhibit different sectoral uncertainties, due to changes of pollutant compositions. We find  
26 global sectoral consumption uncertainties in the range of  $\pm 9$ – $\pm 27\%$  using the Global Temperature  
27 Potential with a 50 year time horizon, with metric uncertainties dominating. National level  
28 uncertainties are similar in both perspectives due to the dominance of CO<sub>2</sub> over other pollutants. The  
29 consumption emissions of the top 10 emitting regions have a broad uncertainty range of  $\pm 9$ – $\pm 25\%$ ,  
30 with metric and emissions uncertainties contributing similarly. The Absolute Global Temperature  
31 Potential with a 50 year time horizon has much higher uncertainties, with considerable uncertainty  
32 overlap for regions and sectors, indicating that the ranking of countries is uncertain.

### 33 Introduction

34 Many studies have shown that national greenhouse gas (GHG) emission accounts can be viewed from  
35 either a production (territorial) or consumption perspective (Davis and Caldeira, 2010; Hertwich and  
36 Peters, 2009; Wiedmann, 2009; Peters and Hertwich, 2008). While the production view only looks at  
37 territorial emissions, the consumption view includes emissions from the production of imported  
38 products and excludes emissions from the production of exports. It has been shown that territorial  
39 emissions have decreased in most developed countries since 1990, but consumption-based emissions  
40 have increased (Peters et al., 2011c). This indicates that growth in consumption and international trade  
41 may undermine the effectiveness of climate policies that only limit emissions in a subset of countries,  
42 such as in the Kyoto Protocol (Wiebe et al., 2012; Kanemoto et al., 2013).

43 The concept of consumption-based emissions estimates can therefore be used to extend the cause-  
44 effect chain from consumption, to production, to emissions, and ultimately to global warming (Figure  
45 1). This is an important complement to the established territorial (Kyoto Protocol) viewpoint,  
46 particularly to link more directly to consumption as a key driver of emissions. More recent studies  
47 have broadened this concept to look at further consequences of increased global demand for traded  
48 products, such as deforestation (Karstensen et al., 2013), biodiversity loss (Lenzen et al., 2012),  
49 dependency on traded fossil fuels (Andrew et al., 2013), land-use change (Weinzettel et al., 2013), and  
50 water footprints (Hoekstra and Mekonnen, 2012).

51 In the estimation of consumption-based emissions accounts, various datasets and models are combined  
52 in the calculations, thus uncertainties and errors may arise in a number of datasets and models:  
53 emission data, metric data, economic data, etc. There are also uncertainties in assumptions and study  
54 design that can be more difficult to explicitly quantify, including which metric and time horizon to use  
55 for comparing pollutants, and how economic data for one specific year can be relevant to other years.

56 The uncertainty of many aspects of the cause-effect chain have been investigated previously (Höhne et  
57 al., 2008; Prather et al., 2012), but the link to consumption has not been made. There is a growing  
58 literature on the uncertainty in input-output (IO; economic) models used to estimate consumption-  
59 based emissions (Wilting, 2012; Lenzen et al., 2010; Peters et al., 2012; Moran and Wood, 2014;  
60 Inomata and Owen, 2014). Uncertainty in economic models, such as computable general equilibrium  
61 models, has also received attention recently (Elliott et al., 2012), but this literature is still not  
62 sufficiently robust. However, the literature on uncertainty in economic data and models is still  
63 relatively small, and large knowledge gaps remains (IPCC, 2014).

64 A number of studies have investigated uncertainty in emissions (European Commission, 2011; UNEP,  
65 2012; Marland et al., 2009; Macknick, 2011), both regional and global, but surprisingly there still does  
66 not exist an emission dataset with specified uncertainties at the country level across all climate-  
67 relevant species. In addition, there exist almost no estimates of uncertainty at the sector level. Many  
68 aspects of uncertainty have been investigated in the climate system (Skeie et al., 2013; Prather et al.,  
69 2012; Myhre et al., 2013a), but there is little literature on the uncertainties in emissions metrics (Olivie  
70 and Peters, 2013; Shine et al., 2007; Reisinger et al., 2010). We are not aware of any studies that have  
71 estimated the uncertainty introduced by each model and dataset (e.g. metric and IO uncertainties), or  
72 how uncertainty propagates when estimating climate change from consumption as a socio-economic  
73 driver.

74 We extend the uncertainty analyses done by Prather et al. (2009), Höhne et al. (2008) and den Elzen et  
75 al. (2005) by including consumption-based emissions for a single year and using a temperature-based  
76 emission metric, which is arguably a more policy-relevant method of weighting emissions. We use

77 Monte-Carlo analysis and draw on previous studies of uncertainties to perturb and highlight the  
78 different contributors: economic data, emission and metric parameters, and then compare our results  
79 with the previous studies.

## 80 **Methods**

81 We consider the propagation of uncertainty from the point of consumption of goods and services  
82 (products), to the production of products where emissions to air occur, to the climate impacts caused  
83 by those emissions (Figure 1). This can be thought of as a causal chain where consumption is assumed  
84 to be the primary driver, in turn driving production, which in turn leads to emissions, and then  
85 emissions lead to temperature change. These components of the cause-effect chain are linked by  
86 calculation methodologies, each requiring parameterization, and we break the analysis into those three  
87 components: economic data, emission statistics, and emission metrics. We estimate the uncertainty for  
88 each of the components individually, and finally connect the components to determine how  
89 uncertainty propagates through the cause-effect chain.

90 To determine the temperature response to a given level of consumption, we first map emission  
91 statistics for most important pollutants to producing regions and sectors (European Commission, 2011).  
92 Emissions are then converted to global temperature change using an emission metric (Aamaas et al.,  
93 2013). This means that we allocate a future global temperature change due to current production and  
94 consumption emissions. The allocations from producers to consumers (in sectors and regions) require  
95 the global supply chain to be enumerated using economic production and trade data (Peters, 2008).  
96 Production often goes through several steps from extraction and refining to manufacturing and  
97 packaging, and finally to consuming markets. These linkages are represented in the global supply  
98 chain through monetary transactions. We normalize emissions by monetary output in each sector in  
99 each region, and allocate emissions according to purchases made by consumers. The result connects  
100 production and consumption, which are potentially geographically separated, and estimates the  
101 consumption that is driving current production emissions and hence future global temperature  
102 response.

103 | All datasets and models introduce uncertainties in the analysis, thus we estimate uncertainties ~~on~~<sup>in</sup> the  
104 economic data, the emissions data and metric parameters in order to estimate uncertainties in the final  
105 results. We undertake the uncertainty analysis using Monte Carlo (MC) analysis, in which datasets and  
106 parameters are randomly perturbed according to predetermined distributions, and then sub-models are  
107 run sequentially to obtain distributions on the results (Granger Morgan et al., 1990). We isolate the  
108 individual contributions to uncertainty on the final results by perturbing individual components  
109 independently, before running everything together to estimate total uncertainty. The analysis considers  
110 parametric uncertainties on the components, as opposed to structural uncertainties, which would  
111 include the comparisons of different models and datasets (Peters et al., 2012). The next section lists  
112 the background data, and shows how uncertainties are estimated, before running the models and  
113 discussing the results.

### 114 ***Datasets and models***

115 We use multi-regional input-output (MRIO) analysis to link economic activities from production to  
116 consumption, capturing global supply chains at the sectoral level (Davis and Caldeira, 2010;  
117 Wiedmann, 2009). We source our economic input-output data from the Global Trade Analysis Project  
118 (GTAP) database version 8, which comprises domestic and trade data for the entire world economy in  
119 2007 divided into 129 regions and 58 sectors (Narayanan et al., 2012). We use these data to construct  
120 | an MRIO model, ~~which connects with the same regional and sectoral resolution, connecting~~ all regions  
121 at the sector level (Andrew and Peters, 2013; Peters et al., 2011b). While GTAP does not provide  
122 uncertainty estimates on the economic datasets, it is possible to generate realistic uncertainty estimates  
123 for the GTAP database from proxy data. Since an MRIO database is an aggregation of multiple

124 datasets, it inherits uncertainties from a number of sources, including: source data, base year  
125 extrapolations, balancing and harmonization procedures, allocations and aggregations (Wiedmann,  
126 2009; Weber, 2008).

127 We use emissions data for the year 2007 from the Emissions Database for Global Atmospheric  
128 Research (EDGAR), for a number of pollutants (see Table 1), mapping these data to the regions and  
129 sectors of the GTAP database. Uncertainties in emission statistics for each pollutant derive from  
130 multiple sources, e.g. for CO<sub>2</sub>: how much fuel is actually consumed, its carbon content, and how much  
131 of it is combusted. Additionally, to be consistent with top-down estimates, statistics are subject to  
132 adjustments and harmonization, and aggregated and grouped to economic sectors. Although national  
133 uncertainty may in some cases be large, global emissions are dominated by a small number of  
134 countries, thus the global uncertainty is mostly a reflection of these countries' data quality (Andres et  
135 al., 2012).

136 The estimated global temperature impact of emissions are calculated using the global temperature  
137 change potential (GTP) metric (Aamaas et al., 2013; Shine et al., 2005), which is essentially a  
138 parameterization of more complex climate models. The metric uses pollutant characteristics  
139 (atmospheric lifetime, radiative forcing) as input, and unlike the more commonly used Global  
140 Warming Potential (GWP) which only relates to radiative forcing, the GTP also includes estimates of  
141 climate temperature response (sensitivity) to changed radiative forcing in the atmosphere, which adds  
142 additional layers of uncertainties (Reisinger et al., 2010). We base our pollutant parameters on the  
143 ATTICA assessment (Fuglestvedt et al., 2010) and IPCC (2007) p. 212-213, and climate sensitivity  
144 and CO<sub>2</sub> uncertainties on the latest CMIP5 data (Olivié and Peters, 2013). The uncertainties on the  
145 other pollutants are drawn from several sources, but mostly following the IPCC Fifth Assessment  
146 Report (Myhre et al., ~~2013a~~2013b).

147

#### 148 ***General uncertainty relationships***

149 It has previously been shown that economic and emissions data show a general pattern where relative  
150 uncertainty is inversely related to magnitude (Lenzen et al., 2010; Wiedmann, 2009; Wiedmann et al.,  
151 2008; Lenzen, 2000). The GTAP data used in our analysis follows the same trends, based on selected  
152 input-output (IO) data where uncertainty is derived from differences between the reported input data  
153 and the final data in the database after harmonization is done and balancing constraints are met (Table  
154 19.6 in McDougall (2006)). These differences in data resulting from the harmonization process are  
155 available only for “*large sectors in large regions with large relative changes*”, which implies that this  
156 relationship indicate the high-end of uncertainties estimates (McDougall, 2006). Figure 2 shows the  
157 relationship for this subset of economic data and uncertainties, with first-order power regression fits to  
158 the observations ( $R^2 > 0.9$ ). The uncertainties are created from the difference between input and output  
159 values, relative to the input and output values, respectively. However, deriving uncertainties from  
160 these differences is not straightforward, as there are many different methods based on different  
161 assumptions which will add additional uncertainties (e.g. comparisons of the difference of input and  
162 output values to the input, output or mean values gives different results). Because of this, we only use  
163 the general relationship between sector size and uncertainty, and not the parameters from Table 19.6,  
164 when estimating sectoral uncertainties. Furthermore, we assume a similar relationship with the  
165 emissions data, based on a previous study of the UK Greenhouse Gas Inventory, where uncertainties  
166 were found using an error propagation model (Jackson et al., 2009). This assumption is also shared by  
167 other recent studies (Moran and Wood, 2014; Lenzen et al., 2010).

Field Code Changed

Field Code Changed

168 The dataset allows the parameterization of a function mapping relative uncertainties to the magnitude  
 169 of the data points. Following previous studies (Lenzen et al., 2010; Wiedmann et al., 2008), we  
 170 assume the data follows a power function

$$r_x = a x^b \quad (1)$$

171  
 172 where  $a$  and  $b$  are coefficients. As there is very little data available to parameterize Equation (1), we  
 173 parameterize the relationship using two extreme data points (generally the uncertainty on the  
 174 minimum and maximum values)

$$a = \frac{r_{min}}{v_{max}^b} \quad (2)$$

$$b = \frac{r_{max} - r_{min}}{v_{min} - v_{max}} \quad (3)$$

175  
 176 It is generally argued that developed countries have lower uncertainty than developing countries due to  
 177 the strength of institutions (Narayanan et al., 2012; Andres et al., 2012). The terms  $r_{min}$  and  $r_{max}$   
 178 define the smallest and largest relative errors, respectively, and are functions of developed and  
 179 developing regions ~~where the latter is given twice the uncertainties of the first group~~ (using the Kyoto  
 180 Protocol groupings of Annex B and non-Annex B countries). ~~We assume that developing countries~~  
 181 ~~have double the uncertainties of developed countries, based on estimates for CO<sub>2</sub> emissions (Andres et~~  
 182 ~~al., 2012; see further discussion in section 2.4).~~ This range is also sector- and region-dependent for the  
 183 economic and emissions data, which we define below. The terms  $v_{min}$  and  $v_{max}$  refer to fixed  
 184 minimum and maximum data values for sectors in a specific region, which is given the uncertainty of  
 185  $r_{max}$  and  $r_{min}$ , respectively. Figure 3 shows the functional relationship between sector sizes and  
 186 uncertainties for economic and emissions data, respectively.

187 The lower threshold  $v_{min}$  is fixed for all regions in the economic and emissions datasets, giving  
 188 sectors of the same size the same uncertainty, as the smallest sectors do not contribute much to the  
 189 national totals. The upper threshold  $v_{max}$  can also be fixed to a certain sector size. However,  
 190 uncertainties are likely to be regionally variable, as while a sector of e.g. 1 billion USD might be very  
 191 large for some countries, it might not be large in other regions. To account for this, we argue that the  
 192 sectors' importance should vary with their contribution to the nations' totals, e.g. gross domestic  
 193 product (GDP) or total emissions. We therefore scale  $v_{max}$  according to the regions' GDP and total  
 194 emissions, for the respective datasets, so that the sectors' importance in different regions is reflected  
 195 by their uncertainties. Sectoral values larger than  $v_{max}$  are given the same uncertainty as values equal  
 196 to  $v_{max}$ , to ensure that single large sectors do not affect the uncertainty on other large sectors (see  
 197 details below).

198 The estimated uncertainties are used to create distributions of perturbations. We impose log-normal  
 199 distributions so that distributions with small relative spreads closely resemble normal distributions,  
 200 while distributions with large relative spreads are skew but avoid negative values (Figure 4). The  
 201 distributions are characterized using reported data as medians, and the spreads are (in order of  
 202 decreasing preference) taken directly from the literature, derived from published analyses, or estimated.  
 203 We define uncertainties as the 5-95% confidence interval (90% CI; equivalent to 1.64 standard  
 204 deviations of a normal distribution).

205 By randomly perturbing each data point, we assume no correlations in the uncertainties of economic  
 206 and emissions data, which might not be accurate for some sector combinations (Peters et al., 2012).  
 207 ~~However, since little data exist, attempts to take this into account will further introduce other uncertain~~  
 208 ~~assumptions. Thus we do not adjust for correlations in these datasets. Implementing correlations in~~  
 209 ~~such an analysis is a major difficulty due to the size of the system under investigation and the lack of~~  
 210 ~~uncertainty data, but may also have significant effects on the results. We discuss this further in section~~  
 211 4. We do, however, undertake a simple sensitivity analysis on the parameter choices, by comparing the  
 212 final results on MRIO uncertainty with uncertainty from the GTAP table showing extreme  
 213 observations.

214 Aggregations of the results (from sectors to regions and from regions to global) usually decrease the  
 215 relative uncertainty, so that the national uncertainty is lower than individual sectors, and global  
 216 uncertainty is in some cases lower than national uncertainty. This is a result of the summation effect,  
 217 and the relationship between sector sizes and uncertainties. The largest sectors are given lowest  
 218 uncertainties, so that the national uncertainty is largely a reflection of the uncertainty of the largest  
 219 sectors. As an example of the summation effect, the relative uncertainty  $r$  of adding  $M \pm S$ ,  $n$  times, is

$$r = \frac{S/M}{\sqrt{n}} \quad (4)$$

220 assuming no correlations. To illustrate this effect, we show the uncertainty results at multiple levels.

#### 221 *Economic data (Multi-regional input–output model)*

222 The total sectoral output  $x$  of a region's economy (a vector) is the sum of intermediate consumption  $Ax$   
 223 and final consumption,  $y$  (Miller and Blair, 1985):

$$x = Ax + y \quad (5)$$

224 where  $A$  is the inter-industry requirements matrix, which is equivalent to the technology used in each  
 225 sector's production. We solve for the total output

$$x = (I - A)^{-1}y \quad (6)$$

226 where  $(I - A)^{-1}$  is the Leontief inverse  $L$ . Emissions are estimated for a given  $y$  by first estimating the  
 227 output, and then linking to sectoral emission intensities,  $F$ . This gives the direct and indirect emissions  
 228 (supply chain) emissions

$$f = F L y \quad (7)$$

229 The economic data from GTAP is represented in a multi-regional input–output (MRIO) model, which  
 230 is constructed from a number of smaller datasets. The GTAP dataset itself is based on a large number  
 231 of smaller datasets (such as national IO tables and trade data from UN's COMTRADE database),  
 232 which are harmonized to remove inconsistencies (Andrew and Peters, 2013; Peters et al., 2011b;  
 233 Narayanan et al., 2012). The construction of an MRIO table from the GTAP data is explained in detail  
 234 elsewhere (Peters et al., 2011b). In the MC analysis, we perturb the components of the GTAP database  
 235 (e.g., domestic IO data and international trade data) and not the resulting MRIO. In other words, we  
 236 estimate the uncertainty of the MRIO data based on the uncertainty in the data used to construct it  
 237 (Peters et al., 2011b). ~~which consists of all data points in the GTAP database used to construct the~~  
 238 MRIO model. This ensures that the uncertainties of the final model reflect the underlying uncertainties  
 239 of the various input data. We construct the perturbed  $L$  and  $y$ , before allocating the direct emissions  $F$   
 240 (which are also perturbed) to consuming regions and sectors.

Field Code Changed

241 We calibrate the uncertainty relationship (Equation 1) for the GTAP data using several datasets. From  
242 the trend lines created from the GTAP table (Figure 2), we find the smallest uncertainty on the largest  
243 sectors to be at approximately 5%. We therefore let 90% of perturbed values fall within 5% of the  
244 median, and set  $r_{min} = 5\%$  for the largest sectors (where  $v_{max}$  apply).

245 The upper threshold  $v_{max}$  is defined by the regions' GDP so that a sector of a specific size will have a  
246 larger importance (and hence a lower uncertainty) in a small region than in a large region. We use the  
247 UK data provided by Lenzen et al. (2010) to explain the range of uncertainties in a single economy. In  
248 this dataset the largest sectors have the smallest error, and following the trend line we find that the  
249 largest value is about 4% of UK GDP. We use this to define the upper threshold  $v_{max} = 4\% \times GDP_r$ ,  
250 which means that sectors at or above this value will be given the lowest national uncertainty ( $r_{min}$ ).  
251 Figure 3 shows the result of the implementations, where the lines indicate the range of developing and  
252 developed regions' sector sizes and uncertainties.

253 For the smallest sectors we set  $v_{min}$  equal to 1 USD and assume  $r_{max} = 100\%$  (following Wiedmann  
254 et al., 2008), due to the lack of more precise regional uncertainty data. The 1 USD relates to a small  
255 value often used in the GTAP database (Peters, 2006). These parameters may seem somewhat  
256 arbitrary, but these choices are not overly important. A value of 1USD in an IOT is exceedingly small  
257 (it represents the economic relationship between two sectors over one year). Indeed, analysis shows  
258 that removing small values has negligible effect on the estimates consumption based emissions (Peters  
259 and Andrew, 2012). Thus, 1 USD is effectively zero in our dataset. It could also be argued that the  
260 value of 1USD is highly uncertain and should have large uncertainty. Giving values smaller than this  
261 higher relative uncertainty causes highly skewed log-normal distributions for the perturbations (see  
262 Figure 4). The GTAP dataset has values as low as  $7e7 \times 10^{-35}$  causing  $r$  to be  $6e6 \times 10^6\%$ . Such highly  
263 skewed distributions for data points with small medians ( $\ll 1$  USD) can lead to large imbalances in the  
264 table.

265 An IO model is balanced so that gross input equals gross output, a fundamental characteristic of input-  
266 output models (Leontief, 1970). The same applies for a multiregional model (MRIO). When  
267 perturbing the coefficients in an IO table, it ultimately upsets the balance. In principal, the IO table can  
268 be rebalanced, but given the size of the systems (about  $7500 \times 7500$  matrices), rebalancing is  
269 prohibitively computationally expensive, and may reduce uncertainties as the perturbed values are  
270 changed. ~~To retain balance, we~~ therefore choose not to rebalance, which effectively causes the  
271 "unbalanced" component to be shifted to the value added. A concern is that the value added may  
272 become unrealistic (e.g., negative) as a consequence. The MC algorithm specifically outputs value  
273 added components to allow cross check imbalances with the raw data, and we find the distributions of  
274 the value added at the sector level to be within expected uncertainty bounds given the size of the value  
275 added. This is partially because of the parameterization of uncertainty we have used, and partially  
276 because the perturbations tend to cancel (the sum of random numbers). Thus, we can justify not  
277 rebalancing our perturbed IOTs and assume the imbalances are allocated to the value added (without  
278 having a large effect on the value added). Implementing this general methodology has also lead to  
279 relatively small regional uncertainties in other studies (Lenzen et al., 2010; Wiedmann et al., 2008).  
280 Structural uncertainties have also been found to be relatively small for major economies (Moran and  
281 Wood, 2014) ~~This approach is followed by others.~~

282 ~~For~~ As a simple sensitivity analysis of the input uncertainties, we also run the MC model with  
283 uncertainties according to the fit of the GTAP table uncertainties (trend line relative to final values,  
284 due to better fit; Figure 2). This vastly increases the uncertainties of all sectors, and we do not  
285 constrain the upper or lower uncertainties, meaning that very small sectors will be given unrealistically



286 | large uncertainties (1USD gives  $r = 1e^{-2}10^9\%$ ). This exercise is only valid for the data it represents;  
287 | large sectors in large countries, but is useful to facilitate the discussion about uncertainties in  
288 | economic data. We discuss these results when exploring MRIO uncertainties, but do not include this  
289 | when combining uncertainties.

## 290 *Emission statistics*

291 | The pollutants considered are listed in Table 1, which cover anthropogenic emissions for the year 2007  
292 | which have an effect on climate. We do not include emissions from short cycle biomass burning, as  
293 | this is considered to have a short lifetime in the atmosphere due to regrowth. The dataset originally  
294 | includes CO<sub>2</sub> emissions from forest fires and decay, which is a mix of natural and anthropogenic  
295 | emission. Extracting the anthropogenic emissions and mapping them to agricultural sectors would  
296 | require crude assumptions. We therefore do not include emissions related to forest loss, but  
297 | acknowledge that it would increase global CO<sub>2</sub> emissions ~~with~~by roughly 12% (van der Werf et al.,  
298 | 2009). The EDGAR dataset only provides crude information on uncertainty at the global level for  
299 | some species (European Commission, 2011). Therefore, global and regional uncertainties in emissions  
300 | are taken from a variety of sources (Table 1). Global fossil-fuel CO<sub>2</sub> emissions statistics are  
301 | independently produced by several organizations, but they generally agree with each other within  
302 | about 5% for developed countries and 10% for developing countries (Andres et al., 2012). The CO<sub>2</sub>  
303 | emission estimates are all based on energy data, and globally the emissions are thought to have an  
304 | uncertainty of  $\pm 10\%$  using a 95% CI (UNEP, 2012). Global SO<sub>2</sub> emissions have an estimated  
305 | uncertainty of between  $\pm 8\%$  and  $\pm 14\%$ , while regional uncertainties may be as large as  $\pm 30\%$  (Smith  
306 | et al., 2010). For CH<sub>4</sub>, N<sub>2</sub>O and F-gases, the uncertainty of global emissions have been estimated as  
307 |  $\pm 21\%$ ,  $\pm 25\%$  and  $\pm 17\%$ , respectively (UNEP, 2012).

308 | Table 1 shows parameters and uncertainties for each pollutant used as median values in the  
309 | perturbations. Very little data exist on uncertainty of emissions by sector, especially on a pollutant and  
310 | regional level. Lenzen et al. (2010) used a table of selected sectors of UK CO<sub>2</sub> emissions to find  
311 | uncertainties, originating from Jackson et al. (2009). According to the regression of the data points,  
312 | within the limits of the data points, there is a spread of uncertainties of roughly 10 times (Figure 2 in  
313 | Lenzen et al. (2010)). We therefore estimate sectoral uncertainty using the same general relationship  
314 | as with the economic data (Equation 1), where the uncertainty of global emissions is used as a proxy  
315 | for the lowest uncertainty estimate of the largest sectors ( $r_{min}$ ) and the smallest sectors' uncertainty is  
316 | scaled by 10 times ( $r_{max} = 10 r_{min}$ ).

317 | We assign developing countries an  $r_{min}$  and  $r_{max}$  which are double those of developed countries. We  
318 | define  $v_{min} = 1kt$  and  $v_{max} = 5\%$  of regional emissions. This dependence on total regional  
319 | emissions shifts the function so that a sector of a specific size will have a larger importance (and hence  
320 | a lower uncertainty) in a smaller region than in a larger region (Figure 3). We do not distinguish  
321 | between different sources of the same pollutant, due to lack of information at the sector level. This is,  
322 | in some cases, a crude simplification (e.g. when comparing uncertainties in emissions of certain  
323 | pollutants from agricultural sectors and power generation). Similarly, for the emissions data, we set  
324 |  $v_{min}$  equal to 1 kt emission. Values below this (as with economic data) have little impact on the  
325 | footprint of regions and sectors, and are therefore given zero uncertainty. ~~Estimates of uncertainty for  
326 | some pollutants for some of the nations do exist, which is included in the calculations. Where regional  
327 | information is available (e.g. for CO<sub>2</sub> emissions from China), we use that to set the minimum  
328 | uncertainty, which will also define the steepness of the uncertainty sector size relationship.~~

329 With every sector data point having an uncertainty, we create perturbations which we can sum to get a  
330 bottom-up estimate of the national uncertainty. Table 2 shows several perturbations of sectors ( $x_{in}$ ) for  
331 region  $r$ . Each perturbation  $i$  leads to a new national total ( $X_i$ ). However, independent uncertainty  
332 estimates of national totals (e.g. national emissions) that may be available for some regions may  
333 conflict with our bottom-up distributions on the national totals ( $X_N$ ). When summing the perturbed  
334 sectors  $x_{in}$  for a region, it is unlikely that the distribution of  $X_N$  will be the same as the known  
335 uncertainty in  $X$ .

336 Additionally, the uncertainty in  $X_N$  will depend on the number of elements contributing to the sum,  
337 according to standard propagation of uncertainty rules (RSS, root sum square; see earlier discussion on  
338 the summation effect). In practice, the uncertainty of  $X$  may be based on several lines of evidence,  
339 which may even exclude sector-based data. To ensure that we can reproduce the top-down uncertainty  
340 estimates of  $X$ , we use constrained optimization (using a quadratic programming (QP) methodology)  
341 to minimally adjust the perturbations of  $x_{in}$  to a given distribution of the  $X_N$  (Table 2).

342 Given that we can adjust one iteration so that it sums to a fixed  $X$ , we then give  $X$  a distribution based  
343 on known national uncertainties, and thus, each iteration of  $X$  is used to balance the same iteration of  
344 the disaggregated sector data ( $x_{in}$ ). This ensures that the sum of sectors ( $X_i$ ) always gives a  $X_N$  with a  
345 known uncertainty. The cost of this adjustment is that the spread of the large values in each region (e.g.  
346 a large sector) are adjusted to fit the constraints. To meet the criteria of e.g. a narrower distribution on  
347 the aggregated values, the large values have to be given a narrower distribution as well. This  
348 methodology allows us to give realistic uncertainties on each  $x_{in}$  leading to an  $X_N$  with a known  
349 uncertainty. We do not perform such balancing on the MRIO input data (previous section) as it is too  
350 computationally expensive, and there is little top-down data on uncertainties in economic data.

351

### 352 *Emission metrics*

353 To link emissions to temperature change, we use the global temperature change potential (GTP) as a  
354 metric to compare and aggregate pollutants (Shine et al., 2007). This gives an estimate of the global  
355 mean surface temperature change due to a pulse of emissions from a specific pollutant, and is a simple  
356 way of modeling the much more complex climate system, and its response. Uncertainties in metric  
357 values can arise from a range of factors: pollutant parameters (radiative forcing and lifetime) and the  
358 response of the climate system. Although it has been shown that the GTP may have larger relative  
359 uncertainties than the alternative metric global warming potential (GWP) (Aamaas et al., 2013;  
360 Reisinger et al., 2010) **and it has been criticized for some of its characteristics** (Pierrehumbert, 2014),  
361 the GTP directly links to global temperature change and is thus arguably more policy relevant (Shine  
362 et al., 2005). In addition, the physical interpretation of the GWP is less clear and the metric has been  
363 criticized by many authors (Peters et al., 2011a; Shine, 2009; Pierrehumbert, 2014). The GTP metric is  
364 calculated using impulse response functions, which explain the interaction of pollutant  $i$  in the  
365 atmosphere ( $IRF_i$ ) and the climate system (temperature) response to a pulse emission ( $IRF_T$ ) with  
366 specific radiative forcing (RF) and atmospheric lifetime.

367 We briefly describe the metric equations here, and refer to existing literature for more details (Aamaas  
368 et al., 2013; Fuglestedt et al., 2010; Olivie and Peters, 2013; Myhre et al., 2013a). The absolute GTP  
369 (AGTP) for each pollutant  $i$  is defined as

$$AGTP_i(H) = \int_0^H RF_i(t) IRF_T(H-t) dt \quad (8)$$

370 where the Radiative Forcing (RF) for a pulse emission is

$$RF_i(t) = RE \times IRF_i = A_i \exp\left(-\frac{t}{\tau_i}\right) \quad (9)$$

371 where  $t$  is time [years],  $H$  is the time horizon [years],  $A_i$  is the radiative efficiency for pollutant  $i$   
 372 [W/(m<sup>2</sup>kg)], and  $\tau_i$  is the decay time for pollutant  $i$  [years]. The AGTP metric is dependent on the IRF  
 373 of temperature, which incorporates the climate system response in global mean surface temperature to  
 374 a given radiative forcing. The climate response is modelled using two decaying exponential functions  
 375 representing: (1) the relative fast response of the atmosphere, the land surface and the ocean mixed  
 376 layer, and (2) the relative slow response of the deep ocean (Peters et al., 2011a),

$$IRF_T = \sum_{j=1}^J \frac{c_j}{d_j} \exp\left(-\frac{t}{d_j}\right) \quad (10)$$

377 where  $J$  is the number of decay terms (usually two),  $c_j$  is a component of the climate sensitivity  
 378 [K/(Wm<sup>2</sup>)], where the total climate sensitivity  $\lambda = \sum c_j$ , and  $d_j$  is the decay time [years] of component  
 379  $c_j$ . These two functions are explained by lifetimes and climate sensitivity for the individual  
 380 components (Table 3). The  $\lambda$  explains the change in equilibrium global-mean temperature due to  
 381 forcing by a pollutant in the atmosphere. We parameterize the IRF according to the results from  
 382 CMIP5 covering 15 different climate models (Olivié and Peters, 2013). This dataset is parameterized  
 383 by relatively short climate runs (140–150 years), and thus it is more representative of the short-term  
 384 climate response (less than 100 years) compared to the equilibrium response (see Olivié and Peters  
 385 (2013) for details). Nevertheless, the dataset leads to a median  $\lambda = 0.75$  K/Wm<sup>2</sup> (equivalent to 2.8°C  
 386 global-mean temperature increase), which is consistent with the climate response (sensitivity) of a  
 387 doubling of CO<sub>2</sub> concentration in the atmosphere within the range of 1.5 to 4.5°C (IPCC, 2013).

388 As CO<sub>2</sub> has a more complex interaction in the atmosphere and can not be sufficiently modelled with a  
 389 single exponential decay, we define the RF for CO<sub>2</sub> as a sum of exponentials (Aamaas et al., 2013):

$$RF_{CO_2}(t) = A_{CO_2} \left\{ a_0 + \sum_{i=1}^I a_i \left( 1 - \exp\left(-\frac{t}{\tau_i}\right) \right) \right\} \quad (11)$$

390 where  $a_i$  is the weight of each exponential, which by definition have to sum to one ( $\sum a_i = 1$ ), and  $I$  is  
 391 the number of exponentials. We follow Joos et al. (2013) and use four exponentials and weights, and  
 392 randomize the multiple lifetimes and coefficients so that the coefficients always sum to 1, following  
 393 Olivié and Peters (2013). The use of four different time scales was found to be sufficient to model  
 394 CO<sub>2</sub>'s behavior in the atmosphere compared to advanced climate models (Olivié and Peters, 2013).  
 395 Correlations between the parameters were implemented for CO<sub>2</sub> and IRF<sub>T</sub>, also based on Olivié and  
 396 Peters (2013), but the effect of the correlations on temperature results was found to be small (less than  
 397 1% of AGTP50 value for CO<sub>2</sub>).

398 Estimates from the literature are used as the median (Fuglestedt et al., 2010) and estimates of  
 399 uncertainty as spread of the distributions (Table 4 and 5). For the non-reactive pollutants, we  
 400 randomized the single RF and lifetime values, as these are represented by only a single decay function.  
 401 The RF used in the calculations includes the indirect effects of chemical reactions from the ozone  
 402 precursors (CO, NO<sub>x</sub> and NMVOC), which were perturbed similarly as the other pollutants. This

403 accounts for three indirect forcing effects: formation of O<sub>3</sub> (causing positive RF by CO, NO<sub>x</sub> and  
404 NMVOC), changing CH<sub>4</sub> levels (causing positive RF by CO and NMVOC, and negative RF by NO<sub>x</sub>),  
405 and CH<sub>4</sub> induced O<sub>3</sub>-effect (causing positive RF by CO and NMVOC, and negative RF by NO<sub>x</sub>)  
406 (Aamaas et al., 2013). The indirect effect of SO<sub>2</sub> is included by scaling the metric value, where the  
407 indirect effect of SO<sub>2</sub> is estimated to be about 175% of the direct effect (Aamaas et al., 2013). This is a  
408 crude estimate, and while the indirect effect may be more uncertain than the direct effect, we use the  
409 same uncertainty for the direct and indirect effects due to lack of pollutant specific data (Boucher et al.,  
410 2013).

411 Our analysis of uncertainty contributions from emissions and metric parameters uses Absolute GTP  
412 (AGTP) values with units of temperature change (in Kelvin or °C). When later allocating temperature  
413 data in the economic model, we also use GTP values in units of CO<sub>2</sub>-equivalent emissions for  
414 comparison. The GTP values are calculated by normalizing the AGTP values with reference to the  
415 AGTP values for CO<sub>2</sub>. When we connect the components for a full MC analysis, we choose a single  
416 time horizon for computational reasons. As discussed elsewhere (Fuglestedt et al., 2010), choosing a  
417 time horizon includes value judgment, and is not based solely on a scientific judgment. We choose to  
418 focus on the impact at 50 years (AGTP50 and GTP50), as this is both consistent with current literature  
419 (Myhre et al., 2013a), and within reasonable time for when to expect global warming to exceed 2  
420 degrees (Joshi et al., 2011; Peters et al., 2013).

421

## 422 **Results**

423 Estimated uncertainties are used to create distributions on all data points. To analyze how various  
424 stages of the cause-effect chain contribute to overall uncertainty, we introduce uncertainty separately  
425 in each part of the chain before combining them all together (Figure 1). We first show uncertainties  
426 resulting from (1) the economic data only, (2) the emissions data only, and (3) the metric calculations  
427 only. The final section (4) connects these three parts together to follow uncertainty through the entire  
428 cause-effect chain. The results show uncertainty propagation from consumption to global temperature  
429 change. The analysis is based on 10,000 MC runs.

430

### 431 ***MRIO uncertainty***

432 In this section, we assume there are no uncertainties on the territorial emissions data or emission  
433 metrics, thus the MRIO model uses unperturbed median estimates of GTP50 values for all pollutants  
434 when allocating emissions to consumers, and uncertainties are purely dependent on parametric  
435 uncertainty in the input data into the MRIO. In our analysis each of the 129 countries has 57 producing  
436 sectors (not including households as they are considered final demand in the model, and therefore not  
437 included in the processing), and thus the MRIO table has 7353 rows and columns. We emphasize here,  
438 but discuss later, that we consider parametric uncertainties and not structural uncertainties.

439 Table 6 shows uncertainties in emissions embodied in imports and exports, as well as consumption,  
440 due to perturbations only on the economic dataset. The exports indicate goods that are produced  
441 domestically but consumed abroad, while the imports indicate goods produced abroad but consumed  
442 domestically. The uncertainties in exported emissions are solely due to uncertainties in domestic  
443 economic data, thus reflecting the pattern of developed countries having higher uncertainties.  
444 Uncertainties in imported emission are generally higher than exported emissions, as the imports come

445 from a number of different regions of which many may have high uncertainties (e.g. emerging and  
446 developing economies).

447 For the largest consumption paths, the consumption perspective is not substantially more uncertain  
448 than the corresponding territorial view due to economic uncertainties. Following the largest  
449 international fluxes embodied in trade from Davis and Caldeira (2010) aggregated over all sectors, we  
450 find 2% uncertainty in emissions embodied in products exported from China to USA, 2% uncertainty  
451 from China to Western Europe, 3% from China to Japan and 1% from USA to Western Europe from  
452 economic uncertainties only. These fluxes are mainly dominated by the largest sectors, to which our  
453 method has assigned the smallest uncertainties. The export from China to USA mainly originates in  
454 the manufacturing sectors, which combined is one of the largest Chinese sectors, therefore with  
455 relatively low uncertainties. Annex B countries are assigned lower uncertainties than non-Annex B  
456 countries, which explains the relatively low uncertainty from USA to Western Europe.

457 For smaller paths, there are much higher economic uncertainties. More than 20% of the international  
458 trade routes have a higher uncertainty than 10% (total number of trade routes is 128 regions × 128  
459 regions), while the median of all is 6% uncertainty. The uncertainties in consumption emissions for the  
460 top emitters are very low for two reasons: (1) the effect of summations and aggregations reduce the  
461 uncertainties on the national level (Equation 4; much higher values are seen on a sectoral level), and (2)  
462 the distributions we give the perturbed data in the larger sectors are relatively small.

463 Since we start from the raw GTAP data to construct the MRIO table, and normalize and invert the  
464 MRIO table, a vast number of summations and multiplications are done with the initial perturbed data  
465 (inversion in a single MC ensemble requires more than  $10^{12}$  operations, which was estimated using the  
466 Lightspeed Matlab toolbox; (Minka, 2014)-). Following RSS uncertainty propagation, the relative  
467 uncertainty will decrease when adding equally sized numbers with equally sized uncertainty (not an  
468 unrealistic assumption for IOA). Thus, the relative uncertainty on the sum of a row in the MRIO (the  
469 output) will depend on the number,  $n$ , of large data points (Equation 4). This problem can be avoided  
470 by using a quadratic programming approach to rebalance the sum to a given uncertainty (as we do for  
471 the emissions data), but we do not do this as a) it is too computationally expensive, and b) it would  
472 require balancing the entire MRIO table to get consistent sums. This problem is difficult to negotiate  
473 given the size of the database we are using, and consequently this exerts a downward pressure on  
474 MRIO uncertainties. Because of this, and because uncertainty ranges of input values are small for the  
475 largest and most important sectors, the final results have small uncertainties. A valid question is then  
476 how reliable the uncertainties are.

477 The raw uncertainties “unfitted” and “fitted” data from Table 19.6 in the GTAP documentation  
478 (Figure Fig. 2), however,) act as a simple sensitivity analysis to our applied uncertainties, although  
479 since this table only samples the very largest deviations it is not representative of the uncertainties in  
480 the entire database. When we use these we find that the uncertainties are much larger for the largest  
481 emitters (between 160% and 400% uncertainty for consumption-based emissions), and for small and  
482 medium sized countries the uncertainties becomes unrealistically large. Thus, the results are clearly  
483 sensitive to the input uncertainties. This is expected as the input uncertainties are outliers in the GTAP  
484 database, thus the uncertainties are known to be large. As a consequence the vastly perturbed values  
485 lead to ill-defined MRIO tables (outside of machine precision), which will compromise accuracy in  
486 the final results (see Method discussion on skew distributions and small data points). However, as  
487 discussed earlier, using the difference between input and output values as a proxy of uncertainty is not  
488 straightforward. E.g. the first data point in Table 19.6 indicate an input values of 2 billion USD and an  
489 output value of 132 billion USD, where the difference (relative to the initial value) can be interpreted

490 as a change of 6500%. This *uncertainty* is vast, and many data points have much larger differences.  
491 Because of these difficulties, and since the results are only valid for specific sectors, we don't show  
492 regional results from this analysis, but only use it for illustrative purposes.

493 Overall, we find small uncertainties on the MRIO results, however, the uncertainties on the end results  
494 are a function of the uncertainties on the input values, as shown by the sensitivity analysis.  
495 Furthermore, the input uncertainties are estimated from small amounts of data and many assumptions,  
496 making the uncertainty estimates on the end results less robust. Although our results are supported by  
497 other studies that have performed parametric uncertainty analysis (Lenzen et al., 2010; Bullard and  
498 Sebald, 1988b; Peters, 2007), structural uncertainties in MRIO analysis is found to be larger (Peters et  
499 al., 2012). Thus we suggest that MRIO uncertainty may be best evaluated using a combination of  
500 structural uncertainties (model comparisons) and parametric Monte-Carlo uncertainties.

501

## 502 *Emissions*

503 At the global level, uncertainties in emissions are known from previous studies (Table 1), which are  
504 used to estimate uncertainties of emissions occurring from production at the sectoral and regional level.  
505 Figure 5 shows the uncertainty of all data points (7482 sectors, 129 regions and global aggregations)  
506 for all pollutants. Each data point's uncertainty is dependent on the sector size, the region's GDP and  
507 whether the region is a developed or developing country. Different activities are associated with  
508 different emissions, thus not all sectors in all regions include emissions from all pollutants.  
509 Additionally, the PFCs and HFCs groups are aggregates of several pollutants, thus the spreads are  
510 based on different amounts of data.

511 The red boxplots in Figure 5 shows the sectoral distributions of the relative uncertainties, not including  
512 data points with zero uncertainties. Aggregations of sectors to individual countries (blue boxplots)  
513 lower the uncertainty ranges, depending on the sectors' impact on national totals (NF<sub>3</sub> is a special case,  
514 where only one sector in each region has emissions, thus sectoral and regional uncertainties are the  
515 same). The median values for the boxplots indicate the skewness of the distributions. The distributions  
516 often have two distinct peaks (not visible in the boxplots), which are developed and developing  
517 countries, where the latter group has higher uncertainty. ~~For CO<sub>2</sub>, NH<sub>3</sub>, NO<sub>x</sub> and SO<sub>2</sub>, regional  
518 information has been used instead of global uncertainty as a proxy for the lowest uncertainty in the  
519 largest sectors in some countries.~~ The global aggregations are results of national totals, which are  
520 dominated by large regions (e.g. China and USA). The bottom-up global uncertainties are not  
521 constrained by top-down estimates, as we are not using aggregated global emissions in the end results.  
522 They are, however, all (except NF<sub>3</sub> due to few data points) lower than the input estimates from Table 1  
523 due to the aggregation effect. Small regions with low emission and high uncertainties thus have little  
524 effect on the global uncertainties.

525 The well-mixed GHGs (WMGHG; CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O, HFCs, PFCs, SF<sub>6</sub>, NF<sub>3</sub>, ~~CH<sub>4</sub>~~) generally have lower  
526 emissions uncertainties (~~59~~9% uncertainty for the aggregated sum) than the short lived pollutants (BC,  
527 OC, SO<sub>2</sub>, NH<sub>3</sub>; ~~12~~14% uncertainty) and precursors (CO, NMVOC, NO<sub>x</sub>; ~~20~~19% uncertainty). The  
528 WMGHGs accounted for  $39.4 \pm 0.915$  Gt CO<sub>2</sub>-eq. emissions (using GTP50), while the short-lived  
529 pollutants accounted for  $-4.6 \pm 0.56$  Gt CO<sub>2</sub>-eq. and the precursors accounted for  $0.4 \pm 0.1$  Gt CO<sub>2</sub>-eq.  
530 (where the two last groups have a mix of warming and cooling effects). Uncertainties in pollutant  
531 aggregates for emissions (tonnes) and GTP50 (CO<sub>2</sub>-eq.) values only include emission uncertainties,  
532 but are different due to different weighting of pollutants and due to mixing of cooling and warming  
533 effects. Uncertainties of territorial emissions from developing countries (54% of global emissions

534 using GTP50) have a median value of 32%, while developed regions have a median uncertainty of  
535 16%. These numbers are dominated by the uncertainty of CO<sub>2</sub>, and usually only small variations are  
536 seen due to other pollutants.

537 Globally, most emissions occur in the electricity generation sector (28% of global emissions using  
538 GTP50) and manufacturing sectors (25%) (see SI for sector aggregations). Uncertainties in emissions  
539 (tonnes) from electricity range from ~~+019%~~ for CO<sub>2</sub>, ~~+827%~~ for SO<sub>2</sub> and ~~5860%~~ for NO<sub>x</sub>, which are  
540 the most important pollutants (which has the largest contributions to the sectoral GTP50 value). For  
541 energy-intensive manufacturing, CO<sub>2</sub> (~~37%~~ uncertainty), SO<sub>2</sub> (~~58%~~), and CH<sub>4</sub> (~~5352%~~) are the most  
542 important pollutants. In the non energy-intensive manufacturing sectors, CO<sub>2</sub> (~~38%~~ uncertainty), SO<sub>2</sub>  
543 (~~+016%~~), and HFCs (~~+221%~~) dominate.

544 For agriculture, CH<sub>4</sub> (21% uncertainty) and N<sub>2</sub>O (26%) are equally important to the GTP50 value,  
545 while CO (37%) comes third. CH<sub>4</sub> has less uncertainty coming from agriculture than energy-intensive  
546 manufacturing, since for CH<sub>4</sub> the agriculture sector is much larger, which is consistent with top-down  
547 estimates (Kirschke et al., 2013). The household sector emits mainly CO<sub>2</sub> (~~78%~~ uncertainty), BC  
548 (~~+54156%~~) and OC (~~+39140%~~), due to household fuels and private transportation. The transport  
549 sectors consists mainly of CO<sub>2</sub> (5%), SO<sub>2</sub> (~~69%~~) and NO<sub>x</sub> (~~+617%~~). Mining, services, and food sectors  
550 are small in a production view, and consist mainly of CO<sub>2</sub> (~~24%~~), CH<sub>4</sub> (16%) and SO<sub>2</sub> (~~69%~~). These  
551 estimates are aggregates of sectors and regions (and gases for HFCs and PFCs), thus disaggregated  
552 data have larger uncertainties.

553

#### 554 ***Emission metrics***

555 Metric (temperature) values have an uncertainty range for the different pollutants and different time  
556 horizons, due to the perturbed metric parameters (RF, lifetime, and climate sensitivity). This includes  
557 uncertainties from mapping emissions to atmospheric concentrations through the global carbon cycle,  
558 which is represented by the relatively uncertain climate sensitivity. Figure 6 shows all pollutants on  
559 the same scale using AGTP for 2007 global emissions, with both relative and absolute uncertainties.  
560 The net temperature response (black dotted line) goes from negative to positive over the first few years,  
561 before the short-lived species decay and the net effect becomes dominated by CO<sub>2</sub> in the long run. The  
562 relative and absolute uncertainty of the net effect is largest in the first few years, and becomes roughly  
563 stable from 50 to 100 years. The strong temperature effects of SLCFs and thus the high absolute  
564 uncertainties of the mix of pollutants increase the net uncertainty in the first few years, but CO<sub>2</sub>  
565 dominates the uncertainty after 20 years.

566 The top contributors to absolute uncertainties in the first year are SO<sub>2</sub>, BC and NH<sub>3</sub>. BC and SO<sub>2</sub> have  
567 similar relative uncertainties, but since the emissions of SO<sub>2</sub> are much larger, it has five times the  
568 absolute uncertainty. OC, BC and SO<sub>2</sub> have the largest uncertainties after approximately 10 years  
569 (except for NH<sub>3</sub> due to its significantly larger RF uncertainty), as the uncertainties are dominated by  
570 RF and climate sensitivity uncertainties. NO<sub>x</sub> has a very high relative uncertainty after 7 years because  
571 its temperature effect goes from positive to negative around this time.

572 Figure 7 shows a breakdown of the parameters contributing to relative uncertainty of the AGTP values  
573 by pollutant (see SI Figure for absolute uncertainties). MC runs with separate metric components  
574 individually perturbed were done to isolate the individual contributions to uncertainties. For  
575 comparison, uncertainties on global emissions are also included in the graph, although not included  
576 when perturbing all components. Uncertainties on emissions and RF do not depend on time horizon,



577 thus they are straight lines. However, as the precursors have combined effects (see methods) the  
578 uncertainty on RF on CO, NMVOC and NO<sub>x</sub> actually change with time due to the different effects  
579 having different lifetimes.

580 For the first three years the total uncertainty for most pollutants (except the SLCFs: BC, OC, SO<sub>2</sub> and  
581 NH<sub>3</sub>) is completely dominated by the first decay parameter of the climate sensitivity, which has a  
582 median value of  $2.6 \pm 1.2$  years (Olivié and Peters, 2013). For the WMGHGs, the parameter continues  
583 to dominate to approximately 6-8 years where the uncertainty of the climate sensitivity component  
584 takes over and continues to dominate to at least 100 years. Between them they explain the largest  
585 contributions of uncertainties to the metric values for all time horizons. While the decay parameter  
586 explains the large uncertainties in the first years, the climate sensitivity parameter explains the  
587 increasing relative uncertainties towards 50 and 100 years. The climate sensitivity parameters are  
588 highly sensitive to time horizon since they have different effects at different times. For SO<sub>2</sub> and NH<sub>3</sub>,  
589 the first years are also effected by high uncertainties from RF. Other short lived pollutants (BC and  
590 OC) have large contributions from both emissions and RF values.

591 At 50 years, CO<sub>2</sub> and CH<sub>4</sub> have additional significant contributions to uncertainties from lifetimes.  
592 Since they both have lifetimes within the ranges of the graph, they show variability with time horizon.  
593 The shorter and longer lived pollutants show little variations in lifetime uncertainties over time  
594 horizons, as lifetimes are either too short or too long to have any effect within 100 years at this scale.  
595 The uncertainty on lifetime for several gases are assumed (Table 5), however, the small impact from  
596 lifetime uncertainties on the metric values indicate that small changes of the median lifetimes will for  
597 most pollutants have very little effect. At 50 years the short-lived pollutants have uncertainties in the  
598 range between  $\pm 95\%$  and  $\pm 165\%$ , while the WMGHGs have uncertainties in the range between  $\pm 35\%$   
599 and  $\pm 70\%$ . The precursors have uncertainties around  $\pm 65\%$ .

600 After 100 years, only the WMGHGs still have a significant temperature effect, which means that the  
601 SLCFs do not contribute with absolute uncertainties. In relative terms, shorter lived pollutants have a  
602 rise in uncertainties from 50 to 100 years, while the opposite is true for the longer lived pollutants. The  
603 last group is then completely dominated by climate sensitivity uncertainties. Most pollutants have  
604 relatively low uncertainty contributions from emissions as the global estimates are low, except for BC  
605 and OC. On a regional and sectoral level, the uncertainties from emissions are usually much more  
606 dominant, which shifts the total uncertainties at all time horizons.

607 The literature consists of both studies which allocate emissions using the absolute metric (AGTP) and  
608 the normalized metric (GTP). The GTP metric values are scaled with the AGTP values for CO<sub>2</sub>. When  
609 running the MC analysis we create AGTP values for every iteration, which implies that CO<sub>2</sub> always  
610 will be normalized by itself (by definition,  $GTP_{CO_2}=1$ ). Therefore, the uncertainties of total emissions  
611 using GTP values are quite different to AGTP uncertainties since the dominant species (CO<sub>2</sub>) has no  
612 metric uncertainty, and the uncertainties on other species are potentially amplified due to the  
613 uncertainty of AGTP<sub>CO<sub>2</sub></sub> values.

614 A second effect of using the GTP values is that the normalization of AGTP values include the climate  
615 sensitivity in both the numerator and denominator, which means that GTP values are less sensitive to  
616 climate sensitivity uncertainties than AGTP values (i.e. uncertainties are correlated). Table 7 illustrates  
617 the difference between uncertainties in AGTP, GTP and GTPGWP values. GTP uncertainties are  
618 typically  $\pm 10$ - $15$  percentage points below those of AGTP, and since the AGTP<sub>CO<sub>2</sub></sub> uncertainties are not  
619 strongly dependent on time horizons, they do not affect the uncertainties over different time horizons  
620 for other pollutants' GTP values much. GWP calculations use the same parameters as with GTP, and



621 although we do not use GWP in our results, we include the uncertainties in the table for comparison.  
622 Overall, we find less uncertainty using GWP than the other metrics (Reisinger et al., 2010), except for  
623 NO<sub>x</sub>. The GWP calculations are not dependent on the highly uncertain climate sensitivity, since it does  
624 not relate to global temperature change. Thus it is expected to have lower uncertainties. NO<sub>x</sub> has  
625 overlapping indirect effects, with highly uncertain RF values, which suggests that the GWP20 values  
626 can be both negative and positive, with a median close to zero. Thus it has a very high uncertainty.

627 A few other studies have investigated the uncertainties of AGTP and GTP values, but it is difficult to  
628 compare those which have as there are many different sources of uncertainties from many different  
629 models and datasets. Our GTP uncertainty results are generally higher than Olivie and Peters (2013)  
630 estimates, since we also include uncertainties on lifetimes and RF values of non-CO<sub>2</sub> species. Their  
631 GTP50 uncertainties for BC (-62–+67%), CH<sub>4</sub> (-38–+48%), N<sub>2</sub>O (-16–+25%) and SF<sub>6</sub> (-17–+25%) are  
632 higher than their GWP uncertainties, mainly due to the dependence on the uncertain climate response  
633 (Olivie and Peters, 2013). An other study (Fuglestedt et al., 2010) found similar uncertainties for  
634 GTP50 values for BC (around 200%) and smaller values for CH<sub>4</sub> (50%) compared to our results, and  
635 essentially zero for N<sub>2</sub>O, when only looking at sensitivity to the climate response. N<sub>2</sub>O is a special  
636 case as it has a similar average lifetime to CO<sub>2</sub>, thus it has similar climate sensitivity uncertainty as  
637 CO<sub>2</sub>, which can be seen in Figure 7 for AGTP values. The normalization of GTP therefore cancels the  
638 climate sensitivity effect. Based on an evaluation of several studies, (including Reisinger et al. (2010)),  
639 Myhre et al. (2013a) assessed the uncertainty of CH<sub>4</sub> for GTP100 to be ±75%, which is close to our  
640 estimate. Furthermore, Joos et al. (2013) found uncertainties for CO<sub>2</sub> AGTP values at 50 (±45%) and  
641 100 years (±90%), based on the spread of multiple climate models. Overall, we find the uncertainties  
642 to be consistent with other studies, but highly variable depending on datasets and choices.

#### 643 *Uncertainty on all components*

644 Total uncertainties in production- and consumption-based emission estimates reflect a combination of  
645 uncertainties from the economic data (IO data for regions and sectors), emissions data (tonnes of the  
646 pollutants occurring in regions and sectors), and metric parameters (RF and lifetime for the pollutants,  
647 and the resulting climate response). Additionally, the emissions of a region in a consumption  
648 perspective is a combination of domestic emissions as well as emissions occurring in other regions  
649 (due to emissions embodied in trade), which changes the mix of pollutants and inherits uncertainties  
650 from the regions and sectors they occur in. To facilitate our discussion we aggregate the 58 economic  
651 sectors (post analysis) to 9 sectors. The results are strongly dependent on different perspectives: (1)  
652 production and consumption, (2) relative or absolute metric values, (3) time horizon of metric, (4)  
653 global, regional or sectoral level, and (5) mix of pollutants included. To illustrate the largest  
654 differences, we focus on comparing points 1, 2 and 4, as 3 has been discussed extensively elsewhere  
655 (Myhre et al., ~~2013b~~2013a).

656 In the allocations of metric values in the MRIO model, we choose to use 50 year time horizon, as  
657 discussed earlier: it is consistent with other recent studies, and consistent with the 2 degree policy  
658 target. Because of the differences between absolute and relative metric uncertainties, we compare both  
659 when including perturbations on all components in the last section.

660 Figure 8 shows uncertainties from the components with aggregated sectors and the top emitting  
661 regions, using GTP50 production emissions. The three different bars represent individual MC  
662 ~~ensembles~~ with only the respective components perturbed. At the sector level, the uncertainties in  
663 emissions data is generally the smallest (from 46% to 2024% for sectors), except for households where  
664 large and highly uncertain emissions of BC and OC occur. Uncertainty in metrics has a range from 14%

Field Code Changed

Field Code Changed

665 | to 6463%, being especially large in sectors with non-CO<sub>2</sub> emissions (e.g. Agriculture and Mining).  
666 | Pollutants with higher relative uncertainty on emissions compared to uncertainty on metric values at  
667 | GTP50 (including BC, OC, and NF<sub>3</sub> at disaggregated levels), will tend to give higher uncertainty on  
668 | emissions, while the other pollutants will give higher uncertainty on metrics.

669 | The sector aggregation means that high and low uncertainties from different sector sizes are mixed,  
670 | and thus single sectors like construction have a higher uncertainty than the aggregated sector Services.  
671 | Disaggregation from the global sector perspective to national level and further to sector level reveals  
672 | that emissions uncertainties are a function of aggregations (sectoral uncertainties are adjusted to  
673 | specific national uncertainties), while the metric uncertainties are not directly dependent on sector  
674 | aggregation and will therefore not scale the same way. Consequently, disaggregated levels generally  
675 | find much higher emission uncertainties than metric uncertainties. For the top 10 emitters,  
676 | | disaggregated sectoral emission uncertainties have a median value between 13 and 6794 percentage  
677 | | points above the national aggregate, while the metric uncertainties have a median value between 4 and  
678 | | 16 percentage points above the national aggregated level.

679 | Furthermore, emission uncertainties are scaled according to sector sizes, whereas metric uncertainties  
680 | are not. This means that emission uncertainties are a combination of mix of pollutants and mix of  
681 | sector sizes, while metric uncertainties only reflect the mix of pollutants (where uncertainty is  
682 | dominated by temperature response). This makes the global sectoral and national level quite different,  
683 | since the national level represent various sector sizes with uncertainties according to the functional  
684 | relationship, while the global sectors might only represent large or small sectors. Because of this,  
685 | emission uncertainties usually dominate at the national level as the regions are less aggregated (each  
686 | region consists of 58 sectors) than the global sectors (each consisting of 129 regions). The difference  
687 | in regional uncertainties is attributed to different mix of territorial pollutants being emitted, the sector  
688 | sizes, size of economy and if the regions are developed or developing nations.

689 | Uncertainties from the different components do not linearly contribute to total uncertainty in the end  
690 | results, thus we calculate the total uncertainty in two different ways: an MC run with everything  
691 | perturbed, and a RSS approach combining the individual components. While the MC run is considered  
692 | the more robust method since it takes into account all data points, including the effect of error  
693 | cancelling, the RSS method is an approximation of error propagation which assumes no correlation  
694 | and normal distributions. The two methods agree in most cases, which imply that there are only small  
695 | correlations between the components and that the global-level data is close to normally distributed.  
696 | This further implies that a full computationally intensive MC run with all components perturbed might  
697 | not be necessary in ideal cases, as the RSS method can approximately derive the results.

698 | Figure 9 shows uncertainties from the consumption perspective, thus including MRIO uncertainties. In  
699 | general, the emissions embodied in imports and exports inherit uncertainties from the economic data  
700 | of the region where the emissions occur. Consumption emissions include territorial emissions and  
701 | emissions from imports, while they exclude emissions from exports. Since our MRIO uncertainties  
702 | only include parametric uncertainties they tend to be small due to the cancellation effect discussed  
703 | earlier, which is consistent with other similar studies (Lenzen et al., 2010; Wilting, 2012; Bullard and  
704 | Sebal, 1988a; Peters, 2007). Structural uncertainties, including differences in data sources, MRIO  
705 | models and definitions of consumption-based emissions, may be a larger source of uncertainty  
706 | (Andrew and Peters, 2013). The differences in the datasets and methods used to calculate  
707 | consumption-based CO<sub>2</sub> emissions have shown to be relatively small, with roughly 10% for USA for  
708 | 2007 (Peters et al., 2012). Although various studies use different input data and models, Peters et al.

709 (2012) found the results of major emitters to be robust across studies, even though 10% differences are  
710 not uncommon.

711 The top emitting regions are large economies, and therefore have mostly large economic sectors and  
712 therefore low aggregated uncertainties. The consumption perspective also mix pollutants in regions  
713 and sectors since the supply-chain is taken into account, leading to dilution of the sectoral and regional  
714 variability since multi-sectoral dependence for a single consuming sector is common (e.g. the  
715 production of a car needs input from other sectors, especially electricity). Households are considered  
716 final demand in the MRIO model, and therefore their emissions are not allocated through the  
717 economic model and thus do not inherit economic uncertainties.

718 Contrary to the production perspective, the national consumption-based emissions are more dominated  
719 by metric uncertainties, due to different mix of pollutants. Disaggregation of the consumption  
720 emissions reveals that metric uncertainties usually dominate the sectors for the top emitters, and that  
721 uncertainties in economic data also usually increase more than the emission uncertainties at the sector  
722 level. For these nations, disaggregated sectoral emission uncertainties have a median value between  
723 ~~0.32~~ and 11 percentage points above the national aggregate, while the metric uncertainties have a  
724 median value between ~~23~~ and ~~89~~ percentage points above the national aggregated level, and economic  
725 uncertainty have an increase between 4 and 10 percentage points.

726 Figure 10 show GTP values and uncertainties for the same sectors and regions, for both territorial and  
727 consumption perspectives. Comparing the allocation differences due to different perspectives help  
728 explain the change in uncertainties when going from production to consumption. Agriculture and  
729 mining see the largest sectoral decrease in uncertainties due mainly to different mix of pollutants  
730 (increased CO<sub>2</sub>), while transport and non-energy intensive manufacturing see an increase due to  
731 increased allocations of non-CO<sub>2</sub> emissions like SO<sub>2</sub>. Similar differences can be seen for regions: India  
732 and Brazil are uncertain due to SO<sub>2</sub> and CH<sub>4</sub> emissions, while the US consists mostly of CO<sub>2</sub>.

733 Most regions have quite similar uncertainty in both perspectives, indicating that the economic  
734 uncertainties do not play a major role for the large regions. The difference of uncertainties in the  
735 allocation perspectives can mainly be attributed to: (1) different mix of pollutants and (2) different  
736 allocations of emissions to sectors. The first effect gives net emission importers higher uncertainty in  
737 some sectors, due to highly uncertain pollutants (e.g. the share of non-CO<sub>2</sub> emissions in the UK is 30%  
738 higher using consumption-based emissions, assuming absolute values), while other sectors decrease  
739 uncertainties due to the increased allocation of CO<sub>2</sub>. The second effect is introduced when aggregating  
740 sectors to national level. The production emissions in a region are often dominated by a few large  
741 sectors, while the consumption-based emissions are distributed more evenly among the same sectors.  
742 This difference in distribution cause different relative errors on the aggregated result, even tough the  
743 sectoral uncertainties and the sum of emissions might be the same. Thus, on the national level, this  
744 effect creates smaller uncertainties. The combined results may give consumption-based emissions less  
745 uncertainty than production emissions on the national level (usually within 1-2% for the top emitters).

746 In the SI we demonstrate how to calculate consumption uncertainty analytically for a simple one-  
747 sector, two-region world economy. This reveals that the consumption uncertainty can be lower, under  
748 conditions that are not unusual. How this analytical solution generalizes to larger systems requires  
749 further research. A similar finding was also found by Peters et al. (2012).

750 The AGTP emissions include uncertainties on CO<sub>2</sub>, thus sectoral and regional uncertainties are larger  
751 and differences are reduced since it is the most common pollutant (Figure 11). In this view, e.g.  
752 Chinese and US emissions overlap greatly within the given uncertainties, suggesting that the ordering

753 is uncertain. The corresponding GTP values have less overlap. This may have large policy  
754 implications in terms of responsibility. Other choices may also change the relative importance and  
755 uncertainty of regions and sectors. Choosing 20 years as time horizon would give lower relative  
756 uncertainties for all pollutants because of lower uncertainties for lifetime and climate sensitivity,  
757 except for SO<sub>2</sub>, BC, OC and NH<sub>3</sub> due to their short-lived nature, thus regions and sectors with large  
758 emissions or consumption of SLCFs will be given larger uncertainties. Choosing 100 years will in  
759 most cases give higher relative uncertainties and give SLCFs less importance (see Figure 7). Overall,  
760 we find the uncertainties to be highly sensitive to methods and choices.

761

## 762 Discussion

763 This study investigates parametric uncertainties in the temperature response to territorial- and  
764 consumption-based emissions with uncertainty contributions from economic data, emissions data and  
765 metric parameters. Structural uncertainties (dataset and model differences) and other contributing  
766 factors such as emission metric, attribution methods and indicators of climate change may be equally  
767 important when assessing uncertainties, but we did not investigate those here (den Elzen et al., 2005;  
768 Höhne et al., 2008; Peters et al., 2012; Moran and Wood, 2014). Earlier studies have shown relatively  
769 low uncertainties when estimating countries' contributions to climate change. Prather et al. (2009)  
770 estimated an uncertainty range of -27% to +32% for the global warming caused by Annex I countries  
771 for the period 1990–2002 (0.11 ±0.03°C using 16–84 % confidence interval). Similar to them, we find  
772 that climate modeling generally has the largest contribution to total uncertainty on an aggregated level.

773 ~~Very few studies have looked at uncertainties in consumption-based emissions inventories. found~~  
774 ~~lower uncertainties for the UK carbon footprint (relative standard deviation of 5% in 2001) than our~~  
775 ~~results (±9%), but is this probably because we include other pollutants and metric uncertainties. Other~~  
776 ~~studies have indicated, similar to this, that the uncertainties in consumption-based emissions mostly~~  
777 ~~come from the emission datasets and not from the economic data.~~

778 Our analysis has shown that uncertainties change depending on the (1) allocation perspective, (2)  
779 pollutants included, (3) metric and (4) aggregation. These changes in uncertainties may have  
780 implications for future mitigation policies.

781 First, we found little difference in the uncertainties in production- and consumption-based emissions.  
782 It is often assumed that consumption-based emissions are more uncertain (Peters, 2008). Consistent  
783 with others, we find that parametric uncertainties are smaller, while structural uncertainties are  
784 generally larger (Peters et al., 2012; Moran and Wood, 2014). Lenzen et al. (2010) found lower  
785 uncertainties for the UK carbon footprint (relative standard deviation of 5% in 2001) than our results  
786 (±9%), but this is probably because we include other pollutants and metric uncertainties. In a recent  
787 study, Moran and Wood (2014), but parametric uncertainty analysis shows that the uncertainties are  
788 small; structural uncertainties may be larger. It found that parametric uncertainties in consumption-  
789 based emissions were generally lower than the uncertainty in territorial-based emissions and the  
790 structural uncertainties (model spread). They found that most major economies' carbon footprint  
791 results are within 10%, consistent with our results. However, it is difficult to gauge how robust the  
792 parametric consumption-based emission uncertainties are. On the one hand, our chosen input  
793 uncertainties may be underestimated but there exists scant data to verify this. Increasing the  
794 uncertainties requires the need to rebalance the MRIO tables used in the analysis, which may introduce  
795 correlations and additional uncertainties resulting from the balancing process. Due to the  
796 computationally expensive nature of this type of analysis, further work would be required to assess the

797 implications of rebalancing for each perturbation. On the other hand, the small uncertainties may  
798 reflect a realistic cancelling of numerous random errors (Lenzen et al., 2010). Settling these issues is a  
799 topic of future research.

800 ~~(2) Including~~Second, including SLCFs creates larger differences between regions' and sectors'  
801 uncertainties, where e.g. emissions from Brazil and India are much more uncertain than those of the  
802 other top 10 emitters due to large emissions in agriculture. Sectors such as agriculture, electricity and  
803 manufacturing have large non-CO<sub>2</sub> emissions, causing larger cooling and warming effects and  
804 additional uncertainties-on the net change. It is often ~~discussed~~argued that a shorter time horizon (e.g.  
805 20 years) places more emphasis on the short-lived pollutants relative to CO<sub>2</sub>, while with a longer time  
806 horizon (e.g. 100 years) the warming from CO<sub>2</sub> dominates. There is also a similar trade off with  
807 uncertainty: in the short term, the uncertainties are much larger due to the SLCFs, and thus the  
808 temperature effect of policies to reduce SLCFs ~~have~~has a more uncertain outcome; in the long-term,  
809 the more certain temperature effects of CO<sub>2</sub> dominate and the uncertainty due to the SLCFs becomes  
810 less relevant. Thus, uncertainty may tend to favor a more certain outcome on CO<sub>2</sub> mitigation compared  
811 to SLCFs. This hypothesis would require deeper analysis using economic and other models that  
812 incorporate uncertainty into decision making.

813 ~~(3) The~~Third, the GTP values have much smaller uncertainties than the AGTP metric, due to 1) the  
814 dominance of CO<sub>2</sub> which has  $GTP_{CO_2}=1$  and no uncertainty by definition and 2) the scaling by  
815  $AGTP_{CO_2}$  in the denominator which effectively reduces the impact of climate-sensitivity uncertainty in  
816 the GTP. This suggests that a normalized metric, GTP, may be better than an absolute metric, AGTP,  
817 in terms of reducing uncertainties. In perspective, the underlying uncertainties are ultimately the same,  
818 but they have just been shifted to different variables and scaled out. Thus, a GTP focus may give the  
819 impression of greater uncertainty in CO<sub>2</sub>, while the uncertainty is really translated to the GTP of other  
820 species. Other metrics, like the GWP, have lower uncertainties than the GTP as they do not include the  
821 response of the climate system (Olivié and Peters, 2013). Despite the metric uncertainties, it is unclear  
822 what role they should play in policy. From a scientific point of view the uncertainties are important,  
823 but in policy, once a metric and its parameters are chosen, their uncertainties are likely to be  
824 disregarded in subsequent analysis-policy applications. This is an area that needs further consideration.

825 ~~(4) Aggregation~~Fourth, aggregation changes the importance of the uncertainty contribution between  
826 the different components (economic data, emissions data and metric), as only the emissions data  
827 uncertainty have been estimated at both sector and regional level, while they all are affected by  
828 reduction in uncertainties by aggregation. On the global sectoral level, uncertainties are dominated by  
829 metrics. For the regions, emissions uncertainties often dominate over emission metric uncertainties. At  
830 the sector level, much larger variations are seen, with even economic uncertainties dominating in very  
831 small sectors. Thus, the role of uncertainties may differ depending on the level of aggregation.

832 These results presented are broadly in line with the existing literature on this topic (Wilting, 2012;  
833 Fuglestedt et al., 2010; Joos et al., 2013; Lenzen et al., 2010; Myhre et al., 2013a; Olivié and Peters,  
834 2013). However, our results are limited by the quality of the uncertainty information available as input  
835 into our analysis. Despite the widespread usage of the input data in a wide variety of studies, there still  
836 exists virtually no uncertainty information on economic data, and limited data on the uncertainties in  
837 emission statistics and metric parameters.

838 A major difficulty of uncertainty analysis is the issue of correlations. There is a large need for  
839 addressing correlations in datasets and uncertainties, as these may have significant impacts on the  
840 results. We see several places where correlations could be important: (1) correlations in the metric

841 parameters, (2) balancing constraints (e.g., if the production of electricity is low, then the consumption  
842 of electricity has to be low), (3) between datasets (e.g., a perturbation in fossil fuel use in the economic  
843 dataset should be reflected by a similar perturbation in the emissions dataset), and (4) in each MC  
844 ensemble the perturbation given to a particular region/sector combination may be correlated with other  
845 region/sectors (e.g. if Norway's emissions from cement production in one ensemble are low, then  
846 Sweden's emissions from the same sector may also be low due to correlations in emissions factors).

847 In our analysis we have explored correlations for metric parameters (temperature and CO2 IRF),  
848 which we found to have a small effect on the results, which is addressing point 1. The effect of  
849 correlations in the MRIO data, and linkages to emission data through energy consumption, has not  
850 previously been quantified, and this remains an important area of research. Although these correlations  
851 may change the uncertainty outcome, implementation of correlations in emissions and economic data  
852 faces considerable computational and conceptual hurdles. First, due to the large datasets used in this  
853 analysis, the correlation matrix would be prohibitively large (approximately 1015 elements), posing  
854 serious computational issues. Second, there are little or no data indicating correlations in uncertainties  
855 in sectoral economic data or emissions data, and populating a correlation matrix of the necessary size  
856 would therefore be largely guesswork. Given these constraints, we suggest that the best way forward is  
857 to generate small test cases to assess the importance of correlations in small datasets, but we leave this  
858 for future work.

859

## 860 **Conclusion**

861 We analyzed emissions from 129 countries and 58 sectors with 31 SLCFs and GHGs when estimating  
862 countries' territorial and consumption-based emissions for 2007. We use top-down uncertainty  
863 estimates to derive sector level uncertainties, and use these to perturb the economic data, emissions  
864 data and metric parameters in a Monte-Carlo model. We find the results are sensitive to some  
865 parameters (such as the uncertainty of the climate response and the datasets) and assumptions (such as  
866 developing countries are assigned twice the uncertainty for emissions and economic data), but  
867 especially to choices regarding allocation perspective, pollutants included, metric used and  
868 aggregation level of the results.

869 We find only minor uncertainty differences between allocation perspectives (production versus  
870 consumption) for the top regions, and uncertainties in the economic data are very small for the large  
871 countries. Since economic data generally does not have uncertainty information, it was necessary to  
872 estimate the uncertainties of the economic data and there is little data to verify our estimates. At the  
873 sectoral level, larger differences between production and consumption are found. The inclusion of  
874 SLCFs increases both the emissions and metric uncertainties, and gives larger variations between  
875 regions and sectors. A different choice of time horizon would change the prioritization of the gases  
876 and corresponding uncertainties. At the global level, the metric uncertainty (which is dominated by  
877 climate sensitivity) dominates over emission and economic uncertainty. At the regional level, the  
878 uncertainties from emissions are more important.

879 Our work points to key areas of future research required to reduce uncertainties. The climate  
880 sensitivity generally dominates uncertainties, and this is where the largest improvements can  
881 potentially be made. Most climate sensitivity literature focuses on the long-term sensitivity, whereas  
882 for metrics (and undoubtedly most mitigation analysis), the temporal path to the equilibrium response  
883 is most relevant (Impulse Response Function). Thus, we suggest much deeper analysis is needed on  
884 the time-evolution of the temperature response. Emission statistics are routinely collected, but



885 generally have poorly defined uncertainties. Our work indicates that large improvements in the  
886 reporting and analysis of emission uncertainties are needed. Additional metric uncertainties can be  
887 improved through a better characterization of metric parameters (radiative efficiencies and lifetimes).  
888 Reducing uncertainties in metrics and emission statistics will reduce both uncertainties in production-  
889 and consumption-based emissions. The uncertainty in the economic data was necessarily based on  
890 crude assumptions. While we found that the economic uncertainties were small, this result ~~needs to be~~  
891 ~~confirmed~~requires confirmation by more comprehensive analyses, critically including uncertainty  
892 correlations, which were excluded from our analysis. ~~This~~Reducing uncertainties in the economic data  
893 will have the effect of reducing uncertainties in consumption-based emissions only.

894

## 895 Acknowledgements

896 The authors acknowledge funding from the Norwegian Research Council project “Quantifying the  
897 global socio-economic and policy drivers for Brazil’s contribution to global warming” (project  
898 number: 196090).

## 899 References

- 900 Aamaas, B., Peters, G. P., and Fuglestedt, J. S.: Simple emission metrics for climate impacts, *Earth*  
901 *Syst. Dynam.*, 4, 145-170, 10.5194/esd-4-145-2013, 2013.
- 902 Andres, R. J., Boden, T. A., Bréon, F. M., Ciais, P., Davis, S., Erickson, D., Gregg, J. S., Jacobson, A.,  
903 Marland, G., Miller, J., Oda, T., Olivier, J. G. J., Raupach, M. R., Rayner, P., and Treanton, K.: A  
904 synthesis of carbon dioxide emissions from fossil-fuel combustion, *Biogeosciences*, 9, 1845-1871,  
905 10.5194/bg-9-1845-2012, 2012.
- 906 Andrew, R. M., Davis, S. J., and Peters, G. P.: Climate policy and dependence on traded carbon,  
907 *Environmental Research Letters*, 8, 034011, 2013.
- 908 Andrew, R. M., and Peters, G. P.: A multi-region input–output table based on the global trade analysis  
909 project database (GTAP-MRIO), *Economic Systems Research*, 25, 99-121, 2013.
- 910 Bond, T. C., Streets, D. G., Yarber, K. F., Nelson, S. M., Woo, J. H., and Klimont, Z.: A technology -  
911 based global inventory of black and organic carbon emissions from combustion, *Journal of*  
912 *Geophysical Research: Atmospheres* (1984–2012), 109, 2004.
- 913 Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V., Kondo,  
914 Y., Liao, H., Lohmann, U., P. Rasch, S.K. Satheesh, S. Sherwood, Stevens, B., and Zhang, X. Y.:  
915 Clouds and aerosols, in: *Climate Change 2013: The Physical Science Basis. Contribution of*  
916 *Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate*  
917 *Change*, edited by: Stocker, T. F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A.  
918 Nauels, Y. Xia, V. Bex, and Midgley, P. M., Cambridge University Press, Cambridge, United  
919 Kingdom and New York, NY, USA, 2013.
- 920 Bullard, C. W., and Sebald, A. V.: Monte Carlo Sensitivity Analysis of Input-Output Models, *The*  
921 *Review of Economics and Statistics*, 70, 708-712, 1988a.
- 922 Bullard, C. W., and Sebald, A. V.: Monte Carlo sensitivity analysis of input-output models, *The*  
923 *Review of Economics and Statistics*, 708-712, 1988b.
- 924 Clarisse, L., Clerbaux, C., Dentener, F., Hurtmans, D., and Coheur, P.-F.: Global ammonia distribution  
925 derived from infrared satellite observations, *Nature Geoscience*, 2, 479-483, 2009.
- 926 Collins, W., Derwent, R., Johnson, C., and Stevenson, D.: The oxidation of organic compounds in the  
927 troposphere and their global warming potentials, *Climatic Change*, 52, 453-479, 2002.
- 928 Davis, S. J., and Caldeira, K.: Consumption-based accounting of CO2 emissions, *Proceedings of the*  
929 *National Academy of Sciences*, 107, 5687-5692, 2010.
- 930 den Elzen, M., Fuglestedt, J. S., Höhne, N., Trudinger, C., Lowe, J., Matthews, B., Romstad, B.,  
931 Pires de Campos, C., and Andronova, N.: Analysing a countries' contribution to climate change:  
932 scientific and policy-related choices, *Environmental Science and Policy*, 8, 614-636, 2005.

933 Derwent, R., Collins, W., Johnson, C., and Stevenson, D.: Transient behaviour of tropospheric ozone  
934 precursors in a global 3-D CTM and their indirect greenhouse effects, *Climatic Change*, 49, 463-  
935 487, 2001.

936 Elliott, J., Franklin, M., Foster, I., Munson, T., and Loudermilk, M.: Propagation of Data Error and  
937 Parametric Sensitivity in Computable General Equilibrium Models, *Comput Econ*, 39, 219-241,  
938 10.1007/s10614-010-9248-5, 2012.

939 European Commission: Emission Database for Global Atmospheric Research (EDGAR), release  
940 version 4.2: <http://edgar.jrc.ec.europa.eu>, 2011.

941 Fuglestedt, J. S., Shine, K. P., Berntsen, T., Cook, J., Lee, D. S., Stenke, A., Skeie, R. B., Velders, G.  
942 J. M., and Waitz, I. A.: Transport impacts on atmosphere and climate: Metrics, *Atmospheric  
943 Environment*, 44, 4648-4677, 10.1016/j.atmosenv.2009.04.044, 2010.

944 Hertwich, E. G., and Peters, G. P.: Carbon footprint of nations: A global, trade-linked analysis,  
945 *Environmental science & technology*, 43, 6414-6420, 2009.

946 Hoekstra, A. Y., and Mekonnen, M. M.: The water footprint of humanity, *Proc Natl Acad Sci U S A*,  
947 109, 3232-3237, 10.1073/pnas.1109936109, 2012.

948 Höhne, N., Blum, H., Fuglestedt, J. S., Skeie, R. B., Kurosawa, A., Hu, G., Lowe, J., Gohar, L.,  
949 Matthews, B., de Salles, A. C. N., and Ellermann, C.: Contributions of individual countries'  
950 emissions to climate change and their uncertainty, Under Review, 2008.

951 Inomata, S., and Owen, A.: COMPARATIVE EVALUATION OF MRIO DATABASES, *Economic  
952 Systems Research*, 26, 239-244, 10.1080/09535314.2014.940856, 2014.

953 IPCC: Climate change 2007: Contribution of Working Group I to the Fourth Assessment Report of the  
954 Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, United  
955 Kingdom and New York, NY, USA., 2007.

956 IPCC: Summary for Policymakers, in: *Climate Change 2013: The Physical Science Basis.  
957 Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on  
958 Climate Change*, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K.,  
959 Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press,  
960 Cambridge, United Kingdom and New York, NY, USA., 2013.

961 IPCC: Chapter 5: Drivers, Trends and Mitigation, in: *Climate Change 2014: Mitigation of Climate  
962 Change. Contribution of Working Group III to the Fifth Assessment Report of the  
963 Intergovernmental Panel on Climate Change*, edited by: Edenhofer, O., R. Pichs-Madruga, Y.  
964 Sokona, E. Farahani, S. Kadner, K. Seyboth, A. Adler, I. Baum, S. Brunner, P. Eickemeier, B.  
965 Kriemann, J. Savolainen, S. Schlömer, C. von Stechow, Zwickel, T., and Minx, J. C., Cambridge  
966 University Press, Cambridge, United Kingdom and New York, NY, USA., p.17-18 and 61-62,  
967 2014.

968 Jackson, J., Choudrie, S., Thistlethwaite, G., Passant, N., Murrells, T., Watterson, J., Mobbs, D.,  
969 Cardenas, L., Thomson, A., Leech, A., Li, Y., Manning, A., Walker, C., Brophy, N., Sneddon, S.,  
970 Pierce, M., Thomas, J., and Brown, K.: UK Greenhouse Gas Inventory 1990 to 2007 - ANNEX 8:  
971 Uncertainties0955482380, 2009.

972 Joos, F., Roth, R., Fuglestedt, J., Peters, G., Enting, I., Bloh, W. v., Brovkin, V., Burke, E., Eby, M.,  
973 and Edwards, N.: Carbon dioxide and climate impulse response functions for the computation of  
974 greenhouse gas metrics: a multi-model analysis, *Atmospheric Chemistry and Physics*, 13, 2793-  
975 2825, 2013.

976 Joshi, M., Hawkins, E., Sutton, R., Lowe, J., and Frame, D.: Projections of when temperature change  
977 will exceed 2°C above pre-industrial levels, *Nature Clim. Change*, 1, 407-412, 2011.

978 Kanemoto, K., Moran, D., Lenzen, M., and Geschke, A.: International trade undermines national  
979 emission reduction targets: New evidence from air pollution, *Global Environmental Change*, 2013.

980 Karstensen, J., Peters, G. P., and Andrew, R. M.: Attribution of CO<sub>2</sub> emissions from Brazilian  
981 deforestation to consumers between 1990 and 2010, *Environmental Research Letters*, 8, 024005,  
982 2013.

983 Kirschke, S., Bousquet, P., Ciais, P., Saunoy, M., Canadell, J. G., Dlugokencky, E. J., Bergamaschi,  
984 P., Bergmann, D., Blake, D. R., and Bruhwiler, L.: Three decades of global methane sources and  
985 sinks, *Nature Geoscience*, 6, 813-823, 2013.

986 Lenzen, M.: Errors in Conventional and Input - Output—based Life—Cycle Inventories, *Journal of  
987 Industrial Ecology*, 4, 127-148, 2000.



988 Lenzen, M., Wood, R., and Wiedmann, T.: Uncertainty analysis for multi-region input–output  
989 models—a case study of the UK's carbon footprint, *Economic Systems Research*, 22, 43-63, 2010.

990 Lenzen, M., Moran, D., Kanemoto, K., Foran, B., Lobefaro, L., and Geschke, A.: International trade  
991 drives biodiversity threats in developing nations, *Nature*, 486, 109-112, 2012.

992 Leontief, W.: Environmental Repercussions and the Economic Structure: An Input-Output Approach,  
993 *The Review of Economics and Statistics*, 52, 262-271, 1970.

994 Levin, I., Naegler, T., Heinz, R., Osusko, D., Cuevas, E., Engel, A., Ilmberger, J., Langenfelds, R. L.,  
995 Neiningner, B., Rohden, C. v., Steele, L. P., Weller, R., Worthy, D. E., and Zimov, S. A.: The global  
996 SF<sub>6</sub> source inferred from long-term high precision atmospheric measurements and its comparison  
997 with emission inventories, *Atmos. Chem. Phys.*, 10, 2655-2662, 10.5194/acp-10-2655-2010, 2010.

998 Macknick, J.: Energy and CO<sub>2</sub> emission data uncertainties, *Carbon Management*, 2, 189-205,  
999 10.4155/cmt.11.10, 2011.

1000 Marland, G., Hamal, K., and Jonas, M.: How Uncertain Are Estimates of CO<sub>2</sub> Emissions?, *Journal of*  
1001 *Industrial Ecology*, 13, 4-7, 10.1111/j.1530-9290.2009.00108.x, 2009.

1002 McDougall, R. A.: Chapter 19: Updating and Adjusting the Regional Input–Output Tables, in: *Global*  
1003 *Trade, Assistance, and Production: The GTAP 6 Data Base*, edited by: Dimaranan, B. V., Center  
1004 for Global Trade Analysis, Purdue University, 2006.

1005 Miller, R., and Blair, P. D.: *Input-output analysis: Foundations and extensions*, Englewood Cliffs, NJ,  
1006 Prentice-Hall, 1985.

1007 Minka, T.: The Lightspeed Matlab toolbox: [http://research.microsoft.com/en-](http://research.microsoft.com/en-us/um/people/minka/software/lightspeed/)  
1008 [us/um/people/minka/software/lightspeed/](http://research.microsoft.com/en-us/um/people/minka/software/lightspeed/), access: 18.02.2015, 2014.

1009 Moran, D., and Wood, R.: CONVERGENCE BETWEEN THE EORA, WIOD, EXIOBASE, AND  
1010 OPENEU'S CONSUMPTION-BASED CARBON ACCOUNTS, *Economic Systems Research*, 26,  
1011 245-261, 10.1080/09535314.2014.935298, 2014.

1012 Myhre, G., D. Shindell, F.-M. Bréon, W. Collins, J. Fuglestedt, J. Huang, D. Koch, J.-F. Lamarque,  
1013 D. Lee, B. M., T. Nakajima, A. Robock, G. Stephens, Takemura, T., and Zhang, H.: Anthropogenic  
1014 and Natural Radiative Forcing, in: *Climate Change 2013: The Physical Science Basis. Contribution*  
1015 *of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate*  
1016 *Change*, Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y.  
1017 Xia, V. Bex and P.M. Midgley ed., Cambridge University Press, Cambridge, United Kingdom and  
1018 New York, NY, USA, 2013a.

1019 Myhre, G., D. Shindell, F.-M. Bréon, W. Collins, J. Fuglestedt, J. Huang, D. Koch, J.-F. Lamarque,  
1020 D. Lee, B. M., T. Nakajima, A. Robock, G. Stephens, Takemura, T., and Zhang, H.: Anthropogenic  
1021 and Natural Radiative Forcing Supplementary Material, in: *Climate Change 2013: The Physical*  
1022 *Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the*  
1023 *Intergovernmental Panel on Climate Change*, edited by: Stocker, T. F., D. Qin, G.-K. Plattner, M.  
1024 Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, Bex, V., and Midgley, P. M., Available from  
1025 [www.climatechange2013.org](http://www.climatechange2013.org) and [www.ipcc.ch](http://www.ipcc.ch), 2013b.

1026 Narayanan, B., Aguiar, A., and McDougall, R.: *Global Trade, Assistance, and Production: The GTAP*  
1027 *8 Data Base*: [https://www.gtap.agecon.purdue.edu/databases/v8/v8\\_doco.asp](https://www.gtap.agecon.purdue.edu/databases/v8/v8_doco.asp), 2012.

1028 Olivé, D., and Peters, G.: Variation in emission metrics due to variation in CO<sub>2</sub> and temperature  
1029 impulse response functions, *Earth Syst. Dynam.*, 4, 267-286, 2013.

1030 Peters, G.: From Production-Based to Consumption-Based National Emission Inventories, *Ecological*  
1031 *Economics*, 65, 13-23, 2008.

1032 Peters, G. P.: Efficient Algorithms for Life Cycle Assessment, Input-Output Analysis, and Monte-  
1033 Carlo Analysis, *International Journal of Life Cycle Assessment*, 12, 373-380, 2007.

1034 Peters, G. P., and Hertwich, E. G.: CO<sub>2</sub> Embodied in International Trade with Implications for Global  
1035 Climate Policy, *Environmental Science and Technology*, 42, 1401-1407, 2008.

1036 Peters, G. P., Aamaas, B., Berntsen, T., and Fuglestedt, J. S.: The integrated global temperature  
1037 change potential (iGTP) and relationships between emission metrics, *Environmental Research*  
1038 *Letters*, 6, 044021, 10.1088/1748-9326/6/4/044021, 2011a.

1039 Peters, G. P., Andrew, R., and Lennox, J.: Constructing an Environmentally-Extended Multi-Regional  
1040 Input–Output Table Using the Gtap Database, *Economic Systems Research*, 23, 131-152,  
1041 10.1080/09535314.2011.563234, 2011b.

1042 Peters, G. P., Minx, J. C., Weber, C. L., and Edenhofer, O.: Growth in emission transfers via  
1043 international trade from 1990 to 2008, *Proceedings of the National Academy of Sciences*, 108,  
1044 8903-8908, 2011c.

1045 Peters, G. P., and Andrew, R.: A multi-region input-output table based on the global trade analysis  
1046 project database, *Frontier of International Input-Output Analyses*, Tokyo, Japan, 2012,

1047 Peters, G. P., Davis, S. J., and Andrew, R.: A synthesis of carbon in international trade,  
1048 *Biogeosciences*, 9, 3247-3276, 10.5194/bg-9-3247-2012, 2012.

1049 Peters, G. P., Andrew, R. M., Boden, T., Canadell, J. G., Ciais, P., Le Quere, C., Marland, G.,  
1050 Raupach, M. R., and Wilson, C.: The challenge to keep global warming below 2°C, *Nature Clim.*  
1051 *Change*, 3, 4-6, 2013.

1052 Pierrehumbert, R. T.: Short-Lived Climate Pollution, *Annual Review of Earth and Planetary Sciences*,  
1053 42, 341-379, doi:10.1146/annurev-earth-060313-054843, 2014.

1054 Prather, M. J., Penner, J., Fuglestedt, J. S., Raper, S., de Campos, C. P., Jain, A., van Aardenne, J.,  
1055 Lal, M., Wagner, F., Kurosawa, A., Skeie, R. B., Lowe, J., Stott, P., and Höhne, N.: Tracking  
1056 uncertainties in the causal chain from human activities to climate, *Geophysical Research Letters*,  
1057 36, 2009.

1058 Prather, M. J., Holmes, C. D., and Hsu, J.: Reactive greenhouse gas scenarios: Systematic exploration  
1059 of uncertainties and the role of atmospheric chemistry, *Geophysical Research Letters*, 39, L09803,  
1060 10.1029/2012GL051440, 2012.

1061 Reisinger, A., Meinshausen, M., Manning, M., and Bodeker, G.: Uncertainties of global warming  
1062 metrics: CO2 and CH4, *Geophysical Research Letters*, 37, L14707, 10.1029/2010GL043803, 2010.

1063 Shindell, D., Kuylenstierna, J. C. I., Vignati, E., van Dingenen, R., Amann, M., Klimont, Z., Anenberg,  
1064 S. C., Muller, N., Janssens-Maenhout, G., Raes, F., Schwartz, J., Faluvegi, G., Pozzoli, L.,  
1065 Kupiainen, K., Höglund-Isaksson, L., Emberson, L., Streets, D., Ramanathan, V., Hicks, K., Oanh,  
1066 N. T. K., Milly, G., Williams, M., Demkine, V., and Fowler, D.: Simultaneously Mitigating Near-  
1067 Term Climate Change and Improving Human Health and Food Security, *Science*, 335, 183-189,  
1068 10.1126/science.1210026, 2012.

1069 Shindell, D. T., Faluvegi, G., Koch, D. M., Schmidt, G. A., Unger, N., and Bauer, S. E.: Improved  
1070 attribution of climate forcing to emissions, *Science*, 326, 716-718, 2009.

1071 Shine, K. P., Fuglestedt, J. S., Hailemariam, K., and Stuber, N.: Alternatives to the global warming  
1072 potential for comparing climate impacts of emissions of greenhouse gases, *Climatic Change*, 68,  
1073 281-302, 2005.

1074 Shine, K. P., Berntsen, T. K., Fuglestedt, J. S., Skeie, R. B., and Stuber, N.: Comparing the climate  
1075 effect of emissions of short- and long-lived climate agents, *Philosophical transactions. Series A,*  
1076 *Mathematical, physical, and engineering sciences*, 365, 1903-1914, 10.1098/rsta.2007.2050, 2007.

1077 Shine, K. P.: The global warming potential - the need for an interdisciplinary retrain, *Climatic Change*,  
1078 96, 467-472, 2009.

1079 Skeie, R., Berntsen, T., Aldrin, M., Holden, M., and Myhre, G.: A lower and more constrained  
1080 estimate of climate sensitivity using updated observations and detailed radiative forcing time series,  
1081 *Earth System Dynamics Discussions*, 4, 2013.

1082 Smith, S. J., van Aardenne, J., Klimont, Z., Andres, R., Volke, A., and Delgado Arias, S.:  
1083 Anthropogenic sulfur dioxide emissions: 1850–2005, *Atmos. Chem. Phys. Discuss.*, 10, 16111-  
1084 16151, 10.5194/acpd-10-16111-2010, 2010.

1085 SPARC: Report on the Lifetimes of Stratospheric Ozone-Depleting Substances, Their Replacements,  
1086 and Related Species, 2013.

1087 UNEP: The Emissions Gap Report 2012, Nairobi, 2012.

1088 van der Werf, G. R., Morton, D. C., DeFries, R. S., Olivier, J. G. J., Kasibhatla, P. S., Jackson, R. B.,  
1089 Collatz, G. J., and Randerson, J. T.: CO2 emissions from forest loss, *Nature Geosci.*, 2, 737-738,  
1090 [http://www.nature.com/ngeo/journal/v2/n11/supinfo/ngeo671\\_S1.html](http://www.nature.com/ngeo/journal/v2/n11/supinfo/ngeo671_S1.html), 2009.

1091 Weber, C. L.: Uncertainties in constructing environmental multiregional input-output models,  
1092 *International input-output meeting on managing the environment*, 2008, 1-31,

1093 Weinzettel, J., Hertwich, E. G., Peters, G. P., Steen-Olsen, K., and Galli, A.: Affluence drives the  
1094 global displacement of land use, *Global Environmental Change*, 23, 433-438,  
1095 <http://dx.doi.org/10.1016/j.gloenvcha.2012.12.010>, 2013.

1096 Weiss, R. F., Mühle, J., Salameh, P. K., and Harth, C. M.: Nitrogen trifluoride in the global  
1097 atmosphere, *Geophysical Research Letters*, 35, 2008.  
1098 Wiebe, K. S., Bruckner, M., Giljum, S., and Lutz, C.: Calculating Energy-Related Co2emissions  
1099 Embodied in International Trade Using a Global Input–Output Model, *Economic Systems Research*,  
1100 24, 113-139, 10.1080/09535314.2011.643293, 2012.  
1101 Wiedmann, T., Lenzen, M., and Wood, R.: Uncertainty Analysis of the UK-MRIO Model–Results  
1102 from a Monte-Carlo Analysis of the UK Multi-Region Input-Output Model (Embedded Carbon  
1103 Dioxide Emissions Indicator), Report to the UK Department for Environment, Food and Rural  
1104 Affairs by Stockholm Environment Institute at the University of York and Centre for Integrated  
1105 Sustainability Analysis at the University of Sydney, Defra, London, Project Ref.: EV02033, 2008.  
1106 Wiedmann, T.: A review of recent multi-region input–output models used for consumption-based  
1107 emission and resource accounting, *Ecological Economics*, 69, 211-222,  
1108 <http://dx.doi.org/10.1016/j.ecolecon.2009.08.026>, 2009.  
1109 Wild, O., Prather, M. J., and Akimoto, H.: Indirect long - term global radiative cooling from NOx  
1110 Emissions, *Geophysical Research Letters*, 28, 1719-1722, 2001.  
1111 Wilting, H. C.: Sensitivity and Uncertainty Analysis in Mrio Modelling: Some Empirical Results with  
1112 Regard to the Dutch Carbon Footprint, *Economic Systems Research*, 24, 141-171,  
1113 10.1080/09535314.2011.628302, 2012.  
1114  
1115

1116 **Table 1: Global emissions and uncertainties. The uncertainties indicate the 5%-95% (90%) percentile range. PFCs**  
 1117 **include: C2F6, C3F8, C4F10, C5F12, C6F14, C7F16, CF4, c-C4F8. HFCs include: HFC-125, HFC-134a, HFC-143a,**  
 1118 **HFC-152a, HFC-227ea, HFC-23, HFC-236fa, HFC-245fa, HFC-32, HFC-365mf, HFC-43-10-mee, following UNEP**  
 1119 **(2012).**

Pollutant	Global emissions (kt)	Uncertainty	Emissions references	Uncertainty references
PFCs	1.47E+01	±17%	European Commission (2011)	UNEP (2012)
CH <sub>4</sub>	3.25E+05	±21%	European Commission (2011)	UNEP (2012)
CO	9.47E+05	±25%	European Commission (2011)	European Commission (2011)
CO <sub>2</sub>	3.14E+07	±8%	European Commission (2011)	UNEP (2012)
HFCs	2.68E+02	±17%	European Commission (2011)	UNEP (2012)
N <sub>2</sub> O	1.02E+04	±25%	European Commission (2011)	UNEP (2012)
NF <sub>3</sub>	1.58E-01	±26%	European Commission (2011)	Weiss et al. (2008)
NH <sub>3</sub>	4.92E+04	±25%	European Commission (2011)	Clarisse et al. (2009)
NMVOC	1.60E+05	±50%	European Commission (2011)	European Commission (2011)
NO <sub>x</sub>	1.27E+05	±25%	European Commission (2011)	European Commission (2011)
SF <sub>6</sub>	6.17E+00	±10%	European Commission (2011)	Levin et al. (2010)
SO <sub>2</sub>	1.22E+05	±11%	European Commission (2011)	Smith et al. (2010)
BC	5.22E+03	±84%	Shindell et al. (2012)	Bond et al. (2004)
OC	1.34E+04	±84%	Shindell et al. (2012)	Bond et al. (2004)

1120

1121

1122 **Table 2: Example of perturbations of sectors for a single region  $r$ , and the resulting distribution on the national total.**  
 1123 **This bottom-up uncertainty estimate may not be consistent with top-down uncertainty estimates.**

Region $r$	Sector 1	Sector 2	Sector 3	Sector $n$	National total (sum of sectors)	Distribution on national totals
Perturbation 1	$x_{11}$	$x_{12}$	$x_{13}$	$x_{1n}$	$X_1$	$\rightarrow X_N$
Perturbation 2	$x_{21}$	$x_{22}$	$x_{23}$	$x_{2n}$	$X_2$	
Perturbation 3	$x_{31}$	$x_{32}$	$x_{33}$	$x_{3n}$	$X_3$	
Perturbation $i$	$x_{i1}$	$x_{i2}$	$x_{i3}$	$x_{in}$	$X_i$	

1124

1125

1126

1127

1128 **Table 3: Metric parameters with uncertainties. Note that the uncertainties are derived from CMIP5 data and Joos et**  
 1129 **al. (2013), but we use the corresponding distributions listed in Table 5 and 6 in the study by Olivié and Peters (2013)**  
 1130 **to account for correlations.**

Parameters	Values	Unit	Uncertainties
Climate sensitivity $f_1$	0.43	K/Wm <sup>2</sup>	±29%
Climate sensitivity $f_2$	0.32		±59%
Climate sensitivity decay $\tau_1$	2.57	year	±46%
Climate sensitivity decay $\tau_2$	82.24		±192%
CO <sub>2</sub> weight $a_0$	0.23		±20%
CO <sub>2</sub> weight $a_1$	0.28		±33%
CO <sub>2</sub> weight $a_2$	0.35		±28%
CO <sub>2</sub> weight $a_3$	0.14		±30%
CO <sub>2</sub> decay $\tau_0$	INF		—
CO <sub>2</sub> decay $\tau_1$	239.6	year	±58%
CO <sub>2</sub> decay $\tau_2$	18.42		±68%
CO <sub>2</sub> decay $\tau_3$	1.64		±63%

1131

1132 **Table 4: RF values and uncertainties. Note that CO, NMVOC and NO<sub>x</sub> are precursors, which have an effect on O<sub>3</sub>**  
 1133 **and CH<sub>4</sub> concentrations. Because of this, no single RF value can be given. The uncertainties indicate the 5%-95%**  
 1134 **(90%) percentile range. Parameters from IPCC (2007) are taken from Table 2.14, p. 212-213.**

Pollutant	RF (Wm <sup>-2</sup> kg <sup>-1</sup> )	Uncertainty	RF references	Uncertainty references
PFCs	6.40E-12 – 1.06E-11	±10%	IPCC (2007)	Myhre et al. (2013b)
CH <sub>4</sub>	1.82E-13	±17%	Fuglestedt et al. (2010)	Myhre et al. (2013b)
CO	-	±24%	Derwent et al. (2001)	Myhre et al. (2013b)
CO <sub>2</sub>	1.81E-15	±10%	Fuglestedt et al. (2010)	Myhre et al. (2013b)
HFCs	6.74E-12 – 1.53E-11	±10%	Fuglestedt et al. (2010), IPCC (2007)	Myhre et al. (2013b)
N <sub>2</sub> O	3.88E-13	±17%	Fuglestedt et al. (2010)	Myhre et al. (2013b)
NF <sub>3</sub>	1.66E-11	±10%	IPCC (2007)	Assumed
NH <sub>3</sub>	-1.03E-10	±123%	Shindell et al. (2009)	Myhre et al. (2013b)
NMVOC	-	±41%	Collins et al. (2002)	Myhre et al. (2013b)
NO <sub>x</sub>	-	±120%	Wild et al. (2001)	Myhre et al. (2013b)
SF <sub>6</sub>	2.00E-11	±10%	Fuglestedt et al. (2010)	Myhre et al. (2013b)
Sulphate	-3.20E-10	±50%	Fuglestedt et al. (2010)	Myhre et al. (2013b)
BC	1.96E-09	±66%	Fuglestedt et al. (2010)	Myhre et al. (2013b)
OC	-2.90E-10	±68%	Fuglestedt et al. (2010)	Myhre et al. (2013b)

1135

1136

1137 **Table 5: Lifetimes and uncertainties. The uncertainty on lifetime for several gases are assumed, but a sensitivity**  
 1138 **analysis revealed that a change of this uncertainty will not have a large impact on the results (see Metric results**  
 1139 **section below). Note that CO, NMVOC and NO<sub>x</sub> are precursors, which have an effect on O<sub>3</sub> and CH<sub>4</sub> concentrations.**  
 1140 **Because of this, no single RF value can be given. Values and uncertainties for CO<sub>2</sub> ~~is~~are given in Table 3. The**  
 1141 **uncertainties indicate the 5%-95% (90%) percentile range. Parameters from IPCC (2007) are taken from Table 2.14,**  
 1142 **p. 212-213.**

Pollutant	Lifetime (years)	Uncertainty	Lifetime references	Uncertainty references
PFCs	2600-50000	±20%	Fuglestvedt et al. (2010)	Assumed
CH <sub>4</sub>	12	±19%	Fuglestvedt et al. (2010)	Myhre et al. (2013b)
CO	-	±20%	Fuglestvedt et al. (2010)	Assumed
CO <sub>2</sub>	-	-	Fuglestvedt et al. (2010)	-
HFCs	1.4-270	[±12%-±29%]	Fuglestvedt et al. (2010), IPCC (2007)	Myhre et al. (2013b), SPARC (2013)
N <sub>2</sub> O	114	±13%	Fuglestvedt et al. (2010)	Myhre et al. (2013b)
NF <sub>3</sub>	740	±13%	Fuglestvedt et al. (2010)	SPARC (2013)
NH <sub>3</sub>	0.02	±20%	Fuglestvedt et al. (2010)	Assumed
NMVOC	-	±20%	Fuglestvedt et al. (2010)	Assumed
NO <sub>x</sub>	-	±20%	Fuglestvedt et al. (2010)	Assumed
SF <sub>6</sub>	3200	±20%	Fuglestvedt et al. (2010)	Assumed
Sulphate	0.01	±20%	Fuglestvedt et al. (2010)	Assumed
BC	0.02	±20%	Fuglestvedt et al. (2010)	Assumed
OC	0.02	±20%	Fuglestvedt et al. (2010)	Assumed

1143

1144



1145 **Table 6: Uncertainties in allocated emissions due to uncertainties in the economic dataset, by top 10 emitters. The**  
 1146 **territorial emissions are not perturbed, thus they have no uncertainty.**

	Region	Territorial	Exports	Uncertainty	Imports	Uncertainty	Consumption	Uncertainty
Top 10 emitters globally	1 China	7269	1966	1.7 %	400	2.1 %	5703	0.7 %
	2 United States of America	6380	744	1.1 %	1411	1.2 %	7047	0.3 %
	3 Russian Federation	2027	600	1.0 %	216	1.3 %	1642	0.5 %
	4 India	1812	232	2.0 %	186	2.6 %	1766	0.5 %
	5 Japan	1381	257	1.3 %	471	1.4 %	1595	0.5 %
	6 Germany	957	324	0.9 %	498	1.0 %	1130	0.6 %
	7 Brazil	750	127	2.1 %	116	3.1 %	739	0.7 %
	8 Canada	626	194	1.0 %	209	1.5 %	641	0.7 %
	9 United Kingdom	616	134	1.0 %	410	1.1 %	892	0.6 %
	10 Korea	547	158	1.9 %	214	2.4 %	602	1.2 %

1147

1148

1149 **Table 7: Metric values uncertainties for 20, 50 and 100 years time horizon. All metric parameters (excluding**  
 1150 **emissions) were perturbed. The uncertainties indicate the 5%-95% (90%) percentile range, where the plus-minus**  
 1151 **notation is half of the 90% CI. Numbers are rounded to nearest 5%, as multiple MC runs would give slightly different**  
 1152 **results (usually within 1-2%).**

Pollutants	AGTP20	AGTP50	AGTP100	GTP20	GTP50	GTP100	<u>GWP20</u>	<u>GWP50</u>	<u>GWP100</u>
PFCs	±30%	±35%	±35%	±20%	±20%	±20%	<u>±15%</u>	<u>±15%</u>	<u>±15%</u>
CH <sub>4</sub>	±45%	±70%	±75%	±35%	±55%	±70%	<u>±25%</u>	<u>±30%</u>	<u>±30%</u>
CO	±45%	±65%	±75%	±35%	±45%	±65%	<u>±20%</u>	<u>±20%</u>	<u>±25%</u>
CO <sub>2</sub>	±35%	±40%	±40%	±0%	±0%	±0%	<u>±0%</u>	<u>±0%</u>	<u>±0%</u>
HFCs	±30%	±40%	±40%	±20%	±20%	±20%	<u>±15%</u>	<u>±15%</u>	<u>±20%</u>
N <sub>2</sub> O	±35%	±40%	±40%	±25%	±25%	±30%	<u>±20%</u>	<u>±25%</u>	<u>±25%</u>
NF <sub>3</sub>	±35%	±35%	±35%	±20%	±25%	±25%	<u>±15%</u>	<u>±20%</u>	<u>±20%</u>
NH <sub>3</sub>	±180%	±165%	±170%	±165%	±150%	±165%	<u>±125%</u>	<u>±130%</u>	<u>±130%</u>
NMVOOC	±50%	±65%	±75%	±35%	±45%	±65%	<u>±20%</u>	<u>±20%</u>	<u>±25%</u>
NO <sub>x</sub>	±35%	±65%	±95%	±35%	±50%	±80%	<u>±295%</u>	<u>±150%</u>	<u>±125%</u>
SF <sub>6</sub>	±35%	±35%	±35%	±20%	±20%	±25%	<u>±15%</u>	<u>±20%</u>	<u>±20%</u>
SO <sub>2</sub>	±110%	±95%	±100%	±100%	±80%	±100%	<u>±55%</u>	<u>±55%</u>	<u>±55%</u>
BC	±125%	±110%	±110%	±110%	±95%	±110%	<u>±70%</u>	<u>±70%</u>	<u>±70%</u>
OC	±125%	±110%	±115%	±110%	±95%	±110%	<u>±70%</u>	<u>±75%</u>	<u>±75%</u>

Inserted Cells  
 Inserted Cells  
 Inserted Cells


1153

1154

1155

1156

1157




1158

1159 **Figure 1: Flow chart of activities (bold boxes) and the datasets that determine transitions between them (dashed boxes)**

1160

1161



1162

1163

1164

1165


1166

1167

1168

1169

Figure 2: Error distribution of selected GTAP input-output data, (taken from Table 19.6 in McDougall (2006) and trendlines shown as colored circles), and trend lines showing the fit of the general functional relationship explained by Equation Eq. (1). Red and blue circles differ due to different methods of estimating the uncertainty. See the discussion in the text.



1170

1171

1172

1173

1174


1175

1176

1177

1178

Figure 3: Functional relationship between sector sizes on horizontal axis (in kt CO<sub>2</sub> emissions and million US dollars, respectively) and relative uncertainty on vertical axis. The red lines outline the range of developing regions, while the blue lines show the range of developed countries. Each region has been estimated using a single unique curve, and all sectors, depending on their size, will fall on this curve. The form of this relationship is established independently for each pollutant.



1179

1180

1181

1182

1183

Figure 4: Distributions depending on median values and uncertainty. Both distributions have a median = 1, while the near-normal distribution (green) has a relative uncertainty of 100%, the skew distribution has a relative uncertainty of 500%. The green and red shaded areas indicate the 5-95% (90%) confidence intervals.






Figure 5: Relative uncertainties (90% CI) of all pollutants for all sectors (red boxplots), for national aggregates (blue boxplots) and global aggregates (green dots). The edges of the boxes indicate the 25<sup>th</sup> and 75<sup>th</sup> percentile, and the whiskers include extreme data points, but not outliers. The blue target symbol indicates the median value of the distributions. Pollutants are sorted according to global emissions in tonnes.




1192  
 1193 **Figure 6: a) The AGTP for a range of pollutants, with b) relative and c) absolute uncertainties due to metric**  
 1194 **parameters. Pollutants are sorted in the legend according to absolute temperature impact at 50 years. The box inside**  
 1195 **subplot a) shows the same figure on a different scale, and the shaded area around the net effect indicate the 90% CI**  
 1196 **uncertainty. Subplot b) has a log scale, showing relative uncertainties. Subplot c) (also using log scale) shows the**  
 1197 **absolute uncertainty for a 90% CI, of which half is the upper shaded area in a) and the other half is the lower shaded**  
 1198 **area.**

1199


1200



1201







1202



1203 **Figure 7: AGTP values (black lines) for all pollutants (sorted by absolute temperature impact at 50 years time horizon)**  
 1204 **and relative uncertainties (dashed lines) for metric parameters, on the right vertical axis. AGTP median values use**  
 1205 **parameters from the literature, while AGTP all show uncertainty with all parameters perturbed (excluding emissions).**  
 1206 **Uncertainties indicate the 90% CI range of the median values. Global emission uncertainties are derived from sector**  
 1207 **aggregations, and are the same as showed in Figure 5.**

1208







1209

1210

1211 **Figure 8: Territorial perspective of emissions and metric uncertainty using GTP50. Top graph shows global emissions**  
1212 **in sectors they occur in, while bottom graph shows regional emissions. Each of the components is represented by an**  
1213 **individual MC. The black circle indicates the aggregated RSS uncertainty. The uncertainty represents the 5-95% CI.**

1214

1215




1216

1217


1218 **Figure 9: Consumption perspective of emissions, metric and MRIO uncertainty using GTP50. Top graph shows global**  
1219 **emissions going to sectors, while bottom graph shows regional consumption.**

1220

1221



1222



1223


1224

1225

1226

Figure 10: GTP values and uncertainties for territorial (first bars) and consumption (second bars) perspectives. Percentages on top of the bars indicate total uncertainty (rounded to closest 5%).

1227



1228

1229

1230

1231

1232

1233

Figure 11: AGTP values and uncertainties for territorial (first bars) and consumption (second bars) perspectives. The uncertainty reflects a combination of all pollutants including CO<sub>2</sub>. Percentages on top of the bars indicate total uncertainty (rounded to closest 5%).