

***Final author's response to the Editor's
comments on "Global and regional
effects of land-use change on climate in
21st century simulations with interactive
carbon cycle"***

by L. R. Boysen et al.

Correspondence to: lboysen@pik-potsdam.de

Final response to the comments of Laurens Ganzeveld (Editor)

Thank you for your time-consuming effort of writing your own review on this manuscript. The points you have made are truly essential for the understanding of the paper. In the following you will find our reply to each of your comments and, again, you can also find the revised manuscript with the edited parts marked in red color.

We hope that our answers will satisfy your concerns and remarks.

With kind regards,
Lena R. Boysen

List of comments & Reply:

- 5
- Abstract: You modified the abstract in response to the comment raised by the referee; Since here the term storages refers according to me to CO₂ uptake whereas it is actually increases/decreases I would suggest to change this to: “Modifications of the land carbon pool by LULCC are disentangled in accordance with processes that can lead to increases and decreases in this carbon pool”

Reply: This sentence has been changed in accordance to the comment.

- Line 51: “...land–atmosphere energy and water exchange.”

10 **Reply:** Done.

- 15
- Line 175: It really took me some time again to carefully read the explanation of the different simulations you have done. I am convinced that you have made a maximum effort to find an optimal way to do so but think that any reader will have a hard time to easily understand what has been done going through all the selected acronyms. I suggest to change the main sentence that at the end summarizes and explains it well removing some more of these acronyms: “The difference between the ESM and L1A simulations therefore yields the biogeophysical effects of LULCC on climate (dT_{BGP}) whereas the difference between the L1A and L1B simulations yields the biogeochemical effects of LULCC on climate (dT_{BGC}). And is it not an idea to, instead of using the reference L1A and L1B to refer to the BGP and BGC simulations?”
- 20

Reply: We changed the sentence accordingly.

To the last point: Unfortunately, L1A and L1B alone do not represent BGP and BGC simulations and thus, this naming would be misleading. It is only the difference of simulations that gets to those, as explained in Table 2. Note that Table 2 will be placed within the text in the final version, which will help avoid confusion.

- Line 255; you added here the statement that “The modification of local BGC-induced temperature signals leads, for example, to a warming in all models in Australia where trees have been replaced by pastures (Fig. 1b).” I would suggest to add here that this is a clear example that despite replacing the trees by pasture, which results in a simulated increase in albedo, there is a simulated warming due to the BGC effect.

Reply: We agree that the previous phrasing caused the impression that the BGC effects in Australia were induced by the local LULCC and not by the global CO₂ signal caused by LULCC (the BGC signal was retrieved from simulations without LULCC but different CO₂ concentrations). Furthermore, we do want to state that there is a local BGC-induced warming signal but which does not necessarily have to be larger than the BGP-induced cooling (e.g. in MIR and IPSL). To clarify this sentence we have changed it as follows:

“On land the warming patterns differ among the models as the greenhouse effect of CO₂ is not homogeneously distributed. The modification of local BGC-induced temperature change leads, for example, to a small warming in all models in Australia (Fig. 1b).”

Furthermore, we added the following phrase on the net effect of LULCC on climate for Australia to the next part about BGP effects on climate:

5 *Line 280: "For two models, MIR and IPSL, this BGP effect in Australia (cooling caused by e.g. the simulated increase in albedo) dominates over the warming due to the above-mentioned BGC effect in this region."*

- Line 266: I had to read a couple of times through this explanation why your signals are less robust compared the Pitman result. You state that "In the RCP scenario, the area undergoing LULCC is relatively small and is mainly located outside regions with strong snowmasking effects, unlike in the past. BGP cooling..". I would rephrase this explaining more clearly that in the past LULCC BGP effects were more dominant in the mid- to high latitude regions, mainly associated with changes in albedo including changes in snow cover, the latter being less relevant in the analysis of future LULCC BGP effects where the LULCC is mainly found in tropical regions and where the albedo changes are counteracted by simulated decreases in evapotranspiration in the tropics.

Reply: We rephrased this and the following sentence:

20 *"In the underlying RCP8.5 scenario, the area undergoing LULCC is relatively small and is mainly located in the tropics. In the past, LULCC took mainly place in the mid- to high latitudes where a clear BGP signal was identified caused by albedo increase related to the seasonal snow-masking effect of trees. In the tropics however, changes in albedo are less dominant and may counteract in their effect on temperatures by changes in other BGP properties such as the latent heat fluxes."*

- Line 286: “While this holds for all models..”

Reply: This phrase has been changed.

- Line 299: “but also increases the snow-masking effect in periodically snow-covered regions.” I checked again what is exactly meant with an increase in the snow-masking effect and where this is introduced in the beginning of the paper referring to the work by Claussen et al. (2001). I infer that this effect is referring to the fact that with an increase in forests, the impact of snow cover on albedo is reduced due to the fact that the snow is mainly found under the forest canopy. It might be worthwhile to shortly mention this in the beginning of the ms to explain well what is meant with this effect.

Reply: To address this point we changed line 53 in the introduction from “For example, a local cooling may occur due to increased surface albedo and the seasonal snow-masking effect when forests are replaced by cropland in mid- to high latitudes (Claussen et al., 2001).” to

“... and the reduced seasonal snow-masking effect when forests are replaced by cropland in mid- to high latitudes, which leads to an increased albedo due to brighter snow cover (Bonan et al., 1992; Claussen et al., 2001).”

- Line 318: “is mainly attributed to enhanced changes in South America, Africa and Australia. These might be related to changes in latent heat fluxes or cloud cover.” Have to admit that this doesn’t make a strong impression in terms of really understanding the models signals. Is there a possibility that this explanation for having larger changes in these regions can be really corroborated?

Reply: This is true, the statement is a kind of speculative. However, it is very difficult to trace down explicit mechanisms for the temperature changes because this would require switching off mechanisms in simulations which is technically sophisticated and computationally very expensive. Figure 1 shows the changes in latent heat fluxes for 5 MIR which can partly explain some of the cooling patterns in Australia. I apologize for the positioning of this figure in the response text.
We added:

"... Australia which can partly be connected to changes in latent heat fluxes (not shown)."




Fig. 1. Latent heat flux in the EMS-L1A simulation for MIR.

- Line 342: Suggestion to change to “which emphasizes the importance of including pastures in the model simulations, and being another source of uncertainty of considering the LULCC effects in climate change, as IPSL, which does not consider the distinction between grassland and pastures, does not show significant changes”. I added here this suggestion but don’t know if this is correct. I anyhow suggest to mention here explicitly what is not considered by IPSL compared to the other models.

Reply: Yes, the IPSL model does not explicitly represent pastures but adjusts the grassland area to the required pasture area extent. However, pastures are harvested, managed and protected from fire which causes differences in the climate signals. CAN totally neglects pastures. Therefore we change the sentence to:

“This emphasizes the importance of including pastures in the model simulations and the currently large role of pasture implementation as source of uncertainty in simulating the LULCC effects on climate. Unlike those models that represent pastures as specific vegetation type, CAN, which neglects the representation of pastures, and IPSL, which does not consider the distinction between grasslands and pastures, do not show significant changes (see SI). “

- 5
- Line 407: “This suggests that the use of annual land-use transition maps rather than annual land cover states maps (gross instead of only net LULCC transitions; Hurtt et al., 2011) leads to substantial increases in land-use emissions”. This appears to be an important feature of your study pointing at some of the different approaches how to implement again the land use information in the different ESMs. When the reader would not be completely informed about some of these subtle differences it would be not that straightforward to appreciate this. So you are saying that MPI and MIR use the land-use transition maps whereas the other ESMs applied the land cover state maps? I thought that the models all implemented the same land use information in these simulations?
- 10

Reply: You are right; we do not mention these transition maps before and only refer to the SI material in the Method section. We therefore add to the Methods part in line 193:

15 *“... to different changes in vegetation cover (). MPI and MIR, for example, simulate LULCC patterns based on annual fractional changes given by a transition matrix (“gross LULCC transitions”) whereas CAN and IPSL only simulate annual LULCC state maps for each grid cell (“net transitions”). Details...”*

and changed line 405 to:

20 *“... is consistent with recent simulations showing that the net land-use change flux is substantially larger when gross transitions are accounted for in addition to net (Wilken-skjeld et al., 2014, in press; Stocker et al., 2014; see also Hurtt et al., 2011) like in MPI and MIR (see Supplement S1)”.*

Global and regional effects of land-use change on climate in 21st century simulations with interactive carbon cycle

L. R. Boysen^{1,*}, V. Brovkin¹, V. K. Arora², P. Cadule³, N. de Noblet-Ducoudré³, E. Kato⁴, J. Pongratz¹, and V. Gayler¹

¹Max Planck Institute for Meteorology, Hamburg, Germany

²Canadian Centre for Climate Modeling and Analysis, Meteorological Service of Canada, University of Victoria, Victoria, BC, V8W 2Y2, Canada

³Laboratoire des Sciences du Climat et de l'Environnement, Gif-sur-Yvette, France

⁴National Institute for Environmental Studies, Tsukuba, Japan

* now at: Potsdam Institute for Climate Impact Research, Research Domain 1: Earth System Analysis, Telegraphenberg A62, 14473 Potsdam, Germany

Correspondence to: L. R. Boysen (lboysen@pik-potsdam.de)

Abstract. Biogeophysical (BGP) and biogeochemical (BGC) effects of land-use and land cover change (LULCC) are separated at the global and regional scales in new interactive CO₂ simulations for the 21st century. Results from four Earth System models (ESMs) are analyzed for the future RCP8.5 scenario from simulations with and without land-use and land cover change (LULCC) contributing to the Land-Use and Climate, IDentification of robust impacts (LUCID) project. Over the period, 2006–2100, LULCC causes the atmospheric CO₂ concentration to increase by 12, 22, and 66 ppm in CanESM2, MIROC-ESM, and MPI-ESM-LR, respectively. Statistically significant changes in global near-surface temperature are found in three models with a BGC-induced global mean annual warming between 0.07 and 0.23 K. BGP-induced responses are simulated by three models in areas of intense LULCC of varying sign and magnitude (between –0.47 and 0.10 K). **Modifications of the land carbon pool by LULCC are disentangled in accordance with processes that can lead to increases and decreases in this carbon pool.** Global land carbon losses due to LULCC are simulated by all models: 218, 57, 35 and 34 Gt C by MPI-ESM-LR, MIROC-ESM, IPSL-CM5A-LR and CanESM2, respectively. On the contrary, the CO₂-fertilization effect caused by elevated atmospheric CO₂ concentrations due to LULCC leads to a land carbon gain of 39 Gt C in MPI-ESM-LR and is almost negligible in the other models. A substantial part of the spread in models' responses to LULCC is attributed to the differences in implementation of LULCC (e.g. whether pastures or crops are simulated explicitly) and the simulation of specific processes. Simple idealized experiments with clear

protocols for implementing LULCC in ESMs are needed to increase the understanding of model responses and the statistical significance of results, especially, when analyzing the regional-scale impacts of LULCC.

5 35

10

15

20

25

30

1 Introduction

About one-third of the global land surface has already been altered by land-use and land cover changes (LULCC) (Vitousek et al., 1997) primarily through deforestation and replacement of natural vegetation with cropland and pastures (Hurt et al., 2009; Ellis, 2011). The impacts of past, present and potential future LULCC on climate and the carbon cycle have been addressed in a number of recent studies (Matthews et al., 2004; Brovkin et al., 2004, 2013; Sitch et al., 2005; Shevliakova et al., 2009; Pongratz et al., 2010). The climatic consequences of LULCC can be expressed in terms of its biogeophysical (BGP) and biogeochemical (BGC) effects. BGP effects account for alterations of physical land surface characteristics such as changes in albedo and roughness length which in turn affect regional boundary layer dynamics and **land–atmosphere energy and water exchange**. For example, a local cooling may occur due to increased surface albedo and **the reduced seasonal snow-masking effect when forests are replaced by cropland in mid- to high latitudes, which leads to an increased albedo due to brighter snow cover** (Bonan et al., 1992; Claussen et al., 2001). However, a reduction in latent heat fluxes in tropical regions associated with a similar change in land cover may result in a warming (Davin and de Noblet Ducoudré, 2010; Brovkin et al., 2009) and decreases in cloud cover (Werth and Avissar, 2002). BGC effects alter the atmospheric greenhouse gas (GHG) composition which then affects the climate at the global scale. Over the historical period, LULCC-associated CO₂ emissions have increased atmospheric CO₂ concentration by 15–20 ppm (Matthews et al., 2004; Brovkin et al., 2004; Pongratz et al., 2010; Arora and Boer, 2010) and Shevliakova et al. (2013) even estimates a contribution of 43 ppm. The resulting global BGC warming effects may counteract regional BGP cooling effects of LULCC but may also intensify local temperature increases depending on the geographical location (Pongratz et al., 2011, 2009; Bathiany et al., 2010; Bala et al., 2007). Furthermore, LULCC affects land–atmosphere feedbacks which are triggered by changes in climate and atmospheric CO₂ concentration: the carbon-temperature feedback and the carbon-concentration feedback may act in opposite directions (Arora et al., 2013). The first one can either be a negative climate feedback due to increased plant productivity or a positive climate feedback as a result of enhanced heterotrophic respiration of soils in a warmer climate (Arnett et al., 2010; Bonan, 2008; Friedlingstein et al., 2006). The second one is a negative climate feedback due to the CO₂-fertilization effect of the vegetation. However, LULCC reduces the size of the land carbon sink and sources and thus may reduce these climate feedback effects.

The Land-Use and Climate, IDentification of robust impacts (LUCID) project is devoted to the detection of the impacts of LULCC on climate. Several studies have found robust climate signals associated with LULCC. Pitman et al. (2009), for example, showed that LULCC can affect latent

and sensible heat fluxes, albedo and near-surface temperatures in atmospheric general circulation models (AGCMs) with prescribed SSTs. Pitman et al. (2012) revealed changes in temperature extremes and Van der Molen et al. (2011) emphasized the latitudinal-dependent importance of cloud feedbacks in the context of climatic consequences of LULCC. Brovkin et al. (2013) found small regional impacts on albedo, available energy, near-surface temperature and land carbon storage by analyzing the output of six Earth System model simulations for the 21st century with prescribed CO₂ concentrations. However, large uncertainties remain, both in the sign and magnitude of BGP and BGC effects due to differences in model parameterizations and assumptions regarding the underlying processes. These mechanisms were investigated in detail, for example, by Boisier et al. (2012). Reducing the uncertainty associated with BGC and BGP effects of LULCC is one of the challenges for climate and Earth System modelers. Previous LUCID studies focused exclusively on BGP effects of LULCC with the exception of Brovkin et al. (2013), who compared BGP with BGC effects. However, their analysis, relying solely on simulations with prescribed CO₂, was restricted to changes in land carbon storage and first-order approximations of the consequences for global mean temperature. A consistent multi-model comparison of explicitly calculated BGP and BGC effects in terms of relevance for key climate variables is yet missing – a gap to be filled by the present study.

We use simulations for the 21st century following a specified emission-driven scenario called ESMRCP8.5 (Moss et al., 2010) which was carried out by four Earth System models participating in the fifth coupled model intercomparison project (CMIP5, Taylor et al., 2012). This scenario, provided by the integrated assessment model (IAM) MESSAGE (Riahi et al., 2011), includes spatially explicit LULCC patterns which reflect the expansion of crop and pasture land required to meet the increasing food demand of a growing world population. This scenario yields a total anthropogenic radiative forcing of about 8.5 W m⁻² in 2100. For the contribution to the LUCID project, the four climate modeling groups performed two additional ESMRCP8.5 simulations in which land cover was held constant at its year 2005 state, once with CO₂ concentrations calculated interactively and once with prescribed CO₂ concentrations from the ESMRCP8.5 simulation (see Table 1). This new approach uses the differences between the standard ESMRCP8.5 and the additional simulations to directly quantify the climatic consequences of regional BGP effects in comparison to the global BGC effects of LULCC on future climate. Thereby, we can also analyze the effect of interactively calculated CO₂ concentrations on land carbon pools and their contribution to temperature changes in contrast to estimated temperature changes from land carbon losses as it is usually done (Brovkin et al., 2013; Gillett et al., 2013). Finally, we identify major uncertainties arising in this multi-model approach.

2 Methods

Results from the ESMRCP8.5 simulations are used from four
 145 ESMS: MPI-ESM-LR (Giorgetta et al., 2013; Reick et al.,
 200 2013), MIROC-ESM (Watanabe et al., 2011), IPSL-CM5A-
 LR (Dufresne et al., 2013) and CanESM2 (Arora et al.,
 2011). Hereafter, the models are referred to as MPI, MIR,
 150 IPSL and CAN model, respectively. For the year 2006, MPI,
 MIR and CAN simulate 375, 387, and 386 ppm, respec-
 tively (no values for IPSL available), which compare well
 with the observed value of 382 ppm (Keeling et al., 2009)
 and close to the prescribed CO₂ concentration of RCP8.5
 155 with 377 ppm (for detailed benchmarking of these models,
 see Anav et al., 2013). The impacts of LULCC on climate
 and land-atmosphere fluxes of carbon are examined by dif-
 ferencing model simulations with and without LULCC. To
 distinguish BGP and BGC effects, three simulation set-ups
 160 between the years 2006 and 2100 are used (Table 1): ESM-
 RCP8.5 includes all RCP8.5 forcings with CO₂ freely ex-
 changed between the land, the ocean and the atmosphere
 components (i.e. CO₂ is simulated interactively; hereafter
 ESM simulation and $T_{LULCC}^{eCO_2}$ for resulting near-surface tem-
 165 peratures and $C_{LULCC}^{eCO_2}$ for simulated land carbon content
 in year 2100). The L1A simulation uses land cover corre-
 sponding to year 2005 and prescribes atmospheric CO₂ con-
 centration taken from the ESM simulation ($T_{no\ LULCC}^{eCO_2}$ and
 $C_{no\ LULCC}^{eCO_2}$). The L1B simulations also neglect LULCC but
 170 CO₂ is interactively simulated ($T_{no\ LULCC}^{eCO_2}$ and $C_{no\ LULCC}^{eCO_2}$). In
 general, the same terminology holds for the land carbon con-
 tent C ; however, changes in carbon pools due to BGP effects
 of LULCC are not separated by the ESM-L1A difference
 from the direct LULCC effects (deforestation, replacement
 175 of natural vegetation and regrowth), and are thus labeled
 $\Delta C_{\Delta LULCC}$. The difference between the ESM and L1A simu-
 lations therefore yields the biogeophysical effects of LULCC
 on climate (ΔT_{BGP}) whereas the difference between the L1A
 and L1B simulations yields the biogeochemical effects of
 180 LULCC on climate (ΔT_{BGC}). Finally, the difference between
 ESM and L1B simulations yields the net effect of LULCC on
 climate (ΔT_{net}) including all feedbacks (Table 2).

Additionally, BGP effects in our simulations with interac-
 tively simulated CO₂ are compared to BGP effects in simu-
 185 lations with prescribed CO₂ concentrations calculated from
 the difference of RCP8.5 and L2A simulations (hereafter,
 RCP simulation and ΔT_{BGP}^{RCP}) with prescribed CO₂ con-
 centrations (Brovkin et al., 2013).

The land-use change information was adapted from the
 190 land-use harmonization project by Hurtt et al. (2011). Al-
 though common land-use information were provided to all
 modeling groups, vegetation dynamics, land surface schemes
 and parameterizations differ substantially among the models
 leading to different changes in vegetation cover (Supplement
 195 Fig. S1). MPI and MIR, for example, simulate LULCC pat-
 terns based on annual fractional changes given by a transition

matrix (“gross LULCC transitions”) whereas CAN and IPSL
 only simulate annual LULCC state maps for each grid cell
 (“net transitions”). Details about participating models can be
 found in the Supplement Fig. S1 and Table S1 as well as
 in Brovkin et al. (2013). It needs to be noted that none of
 the participating models simulated plant growth with respect
 to nitrogen and phosphorus limitation and thus, land carbon
 uptakes by the biosphere and LULCC emissions might be
 overestimated (Goll et al., 2012).

Statistical methods were applied to test the significance
 of results. The modified Student’s t test was used which
 accounts for temporal autocorrelation (Zwiers et al., 1995;
 Findell et al., 2006) and removes linear trends for the averag-
 ing period of 2071–2100 caused by a strong CO₂ forcing. In
 the case of CAN, the average over three ensemble members
 is calculated. Since CAN did not perform L1A runs, BGP
 effects were estimated by the difference of RCP and L2A
 simulations for this model from Brovkin et al. (2013).

3 Results and discussion

3.1 Effects of LULCC on the atmospheric CO₂ concentration and on near-surface temperatures

3.1.1 Changes in atmospheric CO₂ concentrations

The exchange of carbon between the land and the atmosphere
 via plant and soil processes is modified by LULCC which
 thus affects atmospheric CO₂ concentrations. CO₂ concen-
 trations for interactive CO₂ simulations with and without
 LULCC are listed in Table 3 for MPI, CAN and MIR for
 the year 2100 (no data available for IPSL). All models show
 higher CO₂ concentrations in the ESM simulations at 2100
 (951 to 1134 ppm) than the MESSAGE model (926 ppm)
 upon which the RCP scenario is based. This is likely due to
 the underestimation of feedback mechanisms in IAMs rela-
 tive to Earth System models (Jones et al., 2013). The con-
 tribution of LULCC emissions is given by the difference be-
 tween simulations with and without LULCC (CO₂ $\Delta LULCC$)
 (Table 3; transient evolution of changes in Supplement
 Fig. S2). It is greatest for MPI and smallest for CAN which
 is also reflected and discussed in the changes of land carbon
 stocks in Sect. 3.3. Carbon emissions from LULCC enhance
 atmospheric CO₂ concentration above those due to fossil-
 fuel emissions by 7 % in MPI compared to only 1 and 2 % in
 CAN and MIR, respectively.

3.1.2 Biogeochemical effects on climate

Changes in the atmospheric GHG composition due to
 LULCC affect climate on the global scale. Global mean near-
 surface temperatures increase in all simulations until year
 2100 whereat MIR is the most sensitive model to rising GHG
 concentrations (see Supplement Fig. S3a). On a global av-

erage over the years 2071 to 2100, statistically significant increases in ΔT_{BGC} associated with LULCC are found in MPI (0.23 K), MIR (0.12 K) and CAN (0.07 K) (Table 4). LULCC emissions enhance the BGC warming associated with fossil-fuel emissions in a statistically significant manner by 8, 3 and 2 %, respectively (Table 4, first column). Maps of BGC effects for each model (Fig. 1b) show the wide-spread warming pattern of a well-mixed GHG, where the most pronounced temperature increases are found in polar regions due to the sea-ice-albedo feedback as well as temperature feedbacks (Pithan and Mauritsen, 2014) which contribute to the polar amplification. **On land the warming patterns differ among the models as the greenhouse effect of CO_2 is not homogeneously distributed. The modification of local BGC-induced temperature change leads, for example, to a small warming in all models in Australia Fig. 1b).**

3.1.3 Biogeophysical effects on climate

LULCC modifies the physical properties of the land surface which then affect near-surface climate, mainly on the local to regional scale. The model spread in ΔT_{BGP} signals is wide in the global mean and no statistical significance is detected (Table 4). This agrees with previous model intercomparisons of BGP effects of LULCC for historical times (e.g. Pitman et al., 2009); however, results must be expected to be less robust in our study due to the chosen scenario of LULCC. **In the underlying RCP8.5 scenario, the area undergoing LULCC is relatively small and is mainly located in the tropics. In the past, LULCC took mainly place in the mid- to high latitudes where a clear BGP signal was identified caused by albedo increase related to the seasonal snow-masking effect of trees. In the tropics however, changes in albedo are less dominant and may counteract in their effect on temperatures by changes in other BGP properties such as the latent heat fluxes.**

Here, the importance of LULCC implementation and its link to land–atmosphere processes in the models becomes visible when linking LULCC patterns (Supplement Fig. S1) with spatial ΔT_{BGP} responses in Fig. 1a. Conversions of forests (or shrubs as in Australia) to pasture areas (as dynamically implemented by MIR and MPI in Africa, South America and Australia) or grasslands (simulated in IPSL in Australia and South America) lead to BGP-induced cooling. **For two models, MIR and IPSL, this BGP effect in Australia (cooling caused by e.g. the simulated increase in albedo) dominates over the warming due to the above-mentioned BGC effect in this region.** CAN neglects pastures and thus only changes in cropland extent lead to a conversion of forested areas and natural grasslands. Latent heat fluxes are reduced over crop areas leading to a warming which overcompensates the cooling effect of increased albedo over these areas in tropical regions. While this holds true for all models in South America and Africa, IPSL simulates a cooling in those regions. This is rather untypical for IPSL as previous studies with this model (e.g. Davin and de Noblet Ducoudré,

2010) showed that the impact of LULCC on evapotranspiration dominates the total BGP response to LULCC in tropical regions. Note that the IPSL model also showed warming in the extratropics, due to particular assumptions in the seasonality of the leaf area index (LAI) for crops (Pitman et al., 2009). BGP warming is found over North America in MIR and IPSL where pastures (grassland in the latter model) and crops are abandoned for the regrowth of natural grassland and trees. This in turn not only decreases directly surface albedo but also increases the snow-masking effect in periodically snow-covered regions. This effect is also responsible for the observed warming in high northern latitudes of Eurasia, where the tree line shifts northward in a warmer climate in the dynamically simulated vegetation patterns of MPI and MIR.

However, there are more diverse temperature responses shown in Fig. 1 which cannot directly be linked to LULCC. Taking therefore only areas of intense LULCC (here defined as grid cells in which the area of LULCC equals or exceeds 10 % in 2100 compared to 2006) into account, results in statistically significant changes in three models (Table 4, see Supplement Fig. S3b): CAN, which neglects pastures, simulates a warming of 0.1 K (this value is based on results from Brovkin et al., 2013, as mentioned earlier in Sect. 2), whereas IPSL and MIR show a BGP cooling of 0.16 and 0.47 K, respectively. The prescribed CO_2 simulations analyzed by Brovkin et al. (2013) yield BGP cooling effect of 0.23 K for MIR. The stronger decrease in our analysis' near-surface temperature for MIR model is mainly attributed to enhanced changes in South America, Africa and **Australia which can partly be connected to changes in latent heat fluxes (not shown).** BGP cooling can therefore dampen or dominate the net effect on near-surface temperature in specific regions (and not coherently across the models, see Fig. 1c).

3.1.4 Role of LULCC in affecting regional climate

Here, we investigate whether BGP effects ($\Delta_t T_{\text{BGP}}$) can mitigate or rather enhance climate impacts caused by fossil and LULCC emissions alone (L1A simulation, $\Delta_t T_{\text{no LULCC}}^{\text{CO}_2}$) on the continental scale, where Δ_t means a difference between values averaged over the period 2071 to 2100 and the year 2006. Figure 2a illustrates the percentage impact of $\Delta_t T_{\text{BGP}} / \Delta_t T_{\text{no LULCC}}^{\text{CO}_2}$. Values are listed in the Supplement Table S2. Since CAN did not perform the $\Delta_t T_{\text{no LULCC}}^{\text{CO}_2}$ simulation it is not considered here. Overall, the models show inconsistent signs and magnitudes of how the BGP effects influence $\Delta_t T_{\text{no LULCC}}^{\text{CO}_2}$. However, the analysis shows that for the global land area the models coherently simulate a reduction of the fossil-fuel and LULCC emission-driven temperature increase ($\Delta_t T_{\text{no LULCC}}^{\text{CO}_2}$) by 2 % (0.1 K). Furthermore, MPI and MIR simulate the strongest (and statistically significant) potential of warming mitigation over Australia with -11 and -23 %. **This emphasizes the importance of including pastures in the model simulations and the currently**

large role of pasture implementation as source of uncertainty in simulating the LULCC effects on climate. Unlike those models that represent pastures as specific vegetation type, CAN, which neglects the representation of pastures, and IPSL, which does not consider the distinction between grasslands and pastures, do not show significant changes (for more detailed model descriptions see Supplement Table S1). Similarly, LULCC changes described in Sect. 3.1.3 are strong enough to counteract the warming caused by fossil and LULCC emissions in Africa in MIR and IPSL (-8 and -10% , respectively) but not in MPI with an insignificant warming signal of crops. Model responses are again uncertain and it is therefore difficult to link LULCC to adaptation or mitigation strategies, such as done by Pongratz et al. (2011) who analyzed the impact of reforestation.

3.2 Evaluation of the TRCE approach

Gillett et al. (2013) calculated the so-called transient response to cumulative emissions, TRCE, as the ratio of how global mean temperature changes in response to the cumulative increase of CO_2 in the atmosphere by 1 % per year until a doubling is reached. The TRCE for the participating models (in $^{\circ}\text{K Tg C}^{-1}$) is given in Table 5 (after Gillett et al., 2013). MPI and IPSL have a very similar low TRCE while CAN has the highest TRCE. By multiplying the TRCE with the loss of land carbon due to LULCC in 2100 found in each model, equivalent changes in near-surface temperature (ΔT_{TRCE}) can be estimated. Note, that the conversion factor from atmospheric CO_2 concentration to atmospheric carbon storages is $2.12 \text{ PgC ppm}^{-1}$. The availability of simulations that quantify ΔT_{BGC} interactively now allows us to evaluate the TRCE-approximation used by Brovkin et al. (2013) for prescribed CO_2 concentrations.

Results applying the TRCE-approximation for interactive and prescribed CO_2 simulations yield very similar results. For MIR, ΔT_{TRCE} agrees well with the interactively simulated temperature change ΔT_{BGC} (Table 4), and in CAN the TRCE estimate is only 0.01 K too high.

However, larger differences as found in MPI and IPSL hint to the relevance of effects other than the direct effects of LULCC emissions. The TRCE approach quantifies the climate response to cumulative carbon emissions before any BGP or BGC induced feedbacks occur but which are substantial for LULCC impacts (e.g. altered albedo). This linear approach therefore captures results only well in the absence of significant non-linearities in the models. Furthermore, we compared the instantaneous TRCE results to 30 year mean values which eliminate inter-annual variabilities. Overall, the TRCE approach serves as a good first estimate of the magnitude and direction of changes in near-surface temperatures due to LULCC emissions, but sensitivity analysis is needed for each model response.

3.3 Contribution of changes in land carbon storage

The modification of the land carbon sinks and sources via LULCC is responsible for the observed changes in the atmospheric CO_2 concentration (Table 3) and resulting climate effects. The effect of LULCC on the land carbon stocks is shown in Fig. 3. All models simulate land carbon losses due to LULCC (ΔC_{net} , dark solid lines) whereby the dominant carbon loss is mainly attributed to the deforestation ($\Delta C_{\Delta\text{LULCC}}$, light dashed lines) of carbon-rich tropical forest (see Supplement Fig. S1). In the extra-tropics, deforestation is less prevalent and the replacement of abandoned pastures by grasslands has almost no effect, because both are treated the same way in most models. The MPI model yields the strongest carbon loss of 218 Gt C in 2100 (Table 6, $\Delta C_{\Delta\text{LULCC}}$) which is partly attributed to its overestimation of initial carbon stocks in the tropics and dry-lands (Brovkin et al., 2013). The second largest decrease in land carbon in response to LULCC is found in MIR with 57 Gt C. This is consistent with recent simulations showing that the net land-use change flux is substantially larger when gross transitions are accounted for in addition to net (see also Hurtt et al., 2011; Wilkenskjaeld et al., 2014; Stocker et al., 2014) like in MPI and MIR (see Supplement S1). The reason is that cyclic conversions in fractional land cover might not be seen in the resulting vegetation distribution but lead to modified distributions of carbon among the reservoirs.

The increase in atmospheric CO_2 concentration and near-surface temperature following LULCC emissions affects land carbon storage differently across the models (ΔC_{BGC} , light solid lines). The carbon gain due to CO_2 -fertilization caused by LULCC emissions is strongest in MPI with 40 Gt C and is almost negligible in the other models with -3 to 4 Gt C. This probably explains the stronger difference in MPI to simulations with prescribed CO_2 concentration (Table 6, $\Delta C_{\Delta\text{LULCC}^{\text{rcp}}}$). Global mean annual atmosphere-to-land carbon fluxes reveal an increase until the mid-century in all models and all simulations (see Supplement Fig. S4). Around mid-century, the increasing respiration in a warmer climate reduces and more than overcompensates the enhanced carbon uptake associated with the CO_2 -fertilization effect, especially in MIR. The behavior of the MIR is consistent with the findings in Arora et al. (2013) who showed that the carbon-temperature feedback is strongest in the MIR.

The representation of modified land carbon sinks and sources by LULCC vary across the ESMs leading to the wide spread in carbon pool signals. The modeling groups used common land-use datasets and handled indirect effects coherently following the LUCID protocol so that only differences in simulated climate remain. However, intrinsic differences across the models remain such as the explicit simulation of some carbon cycle related processes (e.g. the representation of crops in CAN) and the neglect or parameterization of other processes (e.g. crops in MPI). One example is the simulation of fire emissions which was done by MPI

and IPSL (see Supplement Fig. S5). Interestingly, they both show that fire emissions are reduced by increased land management which would otherwise increase much stronger in a warmer climate. Following Houghton et al. (2012), these aspects cause uncertainties in modeling carbon emissions from LULCC in the order of $\pm 50\%$.

4 Conclusions

BGP and BGC impacts of LULCC on near-surface temperatures and land carbon pools are separated by using CMIP5-LUCID simulations with interactive CO_2 from four Earth Systems models. These results show that the BGP effect in the RCP scenario causes no statistically significant change in the globally-averaged near-surface temperature averaged over the period 2071–2100. This is the consequence of relatively small changes in land cover over the 2006–2100 period compared to that over the historical period. One further reason is the fact that over the 21st century LULCC primarily takes place in (sub)tropical regions where changes in latent heat fluxes have more impact than changes in albedo which are more effective in seasonally snow-covered regions. However, averaged over regions of intense LULCC (i.e. when LULCC impacts $\geq 10\%$ of a grid cell over the 2006–2100 period), three models simulate statistically significant changes of varying sign and magnitude (between 0.1 and -0.47 K). BGC effects of LULCC lead to statistically significant increases in global mean near-surface temperatures of 0.07, 0.12 and 0.23 K following increases in atmospheric CO_2 from LULCC emissions between 12, 22 and 66 ppm in CAN, MIR and MPI, respectively. The model spread is attributed to differences in modeling assumptions, parameterizations and included processes (e.g. fire) which lead to different manners in which the common LULCC pattern is implemented across models (e.g. with and without pastures) and induce a degree of uncertainty.

The BGP effects of LULCC may enhance or dampen its BGC effects. For example, in South America and Africa, MIR and IPSL both show that BGP effects dampen and, in the case of MPI, enhance BGC warming caused by land-use change and fossil-fuel emissions. A causal link between LULCC forcing and the climate impact is found for MIR where the presence of pastures in Europe and Australia tends to induce a local BGP cooling which offsets a BGC warming. Crops tend to warm climate in most areas and models. This is especially the case in CAN which is the only model that simulates an overall BGP warming in the absence of pasture representation. Conversion to pastures thus may have a climate change mitigation potential but more detailed and idealized experiments are required e.g. simulations with and without pasture cultivation in each model.

The approach of the transient response to cumulative emissions in 2100, TRCE (Gillett et al., 2013) captures the changes in temperature well for CAN and MIR but is less

precise for MPI and IPSL. Therefore, TRCE serves as a good first estimate but since it is a linear approach it is less reliable in case of non-linearities and strong variability in the models.

LULCC leads to carbon release from the land to the atmosphere. Accounting for gross LULCC transitions in both, MPI and MIR, results in stronger LULCC emissions than in the other two models. The global effect of CO_2 -fertilization due to LULCC is strong for MPI with 39 GtC in 2100 and almost negligible in the other models.

Land use change emissions are inherently uncertain. When implemented in ESMs, the diagnosed BGP and BGC effects of LULCC are even more uncertain because of the manner in which land-use change is interpreted and implemented across models. The BGC effects of LULCC are related to how the deforested biomass is treated, if or not transitions across land cover types are considered and how natural vegetation regrows after croplands/pastures are abandoned. All these factors determine the net LULCC emissions and thus the change in atmospheric CO_2 concentration. The BGP effects of LULCC are related to how changes in the physical appearance of the land surface affect the energy and water balance through changes in albedo, roughness length and other physical structural attributes of vegetation. Since models differ greatly in treating BGP and BGC effects of LULCC, the same LULCC pattern can yield differences in magnitude and even sign of the net effect. Simple idealized experiments with clear experimental protocols are needed to, for example, make actually simulated land-use patterns more comparable by coherently implementing or neglecting pastures. This would provide better understanding of why models respond differently to the same LULCC forcing and thus to help reducing uncertainty in the net effect of LULCC across models. Last but not least, some of the uncertainty could be eliminated by having several ensemble members which would make statistical significance testing more robust.

Supplementary material related to this article is available online at: <http://www.gmd.copernicus.org/@journalurl/@pvol/@fpage/@pyear/@journalnameshortlower-/@pvol-/@fpage-/@pyear-supplement.pdf>.

Acknowledgements. We acknowledge the World Climate Research Programme's Working Group on Coupled Modelling, which is responsible for CMIP, and we thank the climate modeling groups for producing and making available their model output. We thank Karl Taylor and Charles Doutriaux for help with setting up the CMOR tables for the LUCID-CMIP5 experiments. We appreciate the support by the staff of the German Climate Computing Center (DKRZ), in particular by Stephanie Legutke and Estanislao Gonzalez, in performing the LUCID-CMIP5 simulations and in making the model results available via DKRZ ESG gateway.

References

560 references

- Anav, A., Friedlingstein, P., Kidston, M., Bopp, L., Ciais, P., Cox, P., Jones, C., Jung, M., Myneni, R., and Zhu, Z.: Evaluating the land and ocean components of the global carbon cycle in the CMIP5 Earth System Models, *J. Climate*, 26, 6801–6843, doi:<http://dx.doi.org/10.1175/JCLI-D-12-00417.1>, 2013.
- Arneth, A., Harrison, S. P., Zaehle, S., Tsigaridis, K., Menon, S., Bartlein, P. J., Feichter, J., Korhola, A., Kulmala, M., O'Donnell, D., Schurgers, G., Sorvari, S., and Vesala, T.: Terrestrial biogeochemical feedbacks in the climate system, *Nat. Geosci.*, 3, 525–532, 2010.
- Arora, V. and Boer, G.: Uncertainties in the 20th century carbon budget associated with land use change, *Global Change Biol.*, 16, 3327–3348, 2010.
- Arora, V., Scinocca, J., Boer, G., Christian, J., Denman, K., Flato, G., Kharin, V., Lee, W., and Merryfield, W.: Carbon emission limits required to satisfy future representative concentration pathways of greenhouse gases, *Geophys. Res. Lett.*, 38, L05805, doi:<http://dx.doi.org/10.1029/2010GL046270>, 2011.
- Arora, V. K., Boer, G. J., Christian, J. R., Bopp, L., Brovkin, V., Eby, M., Friedlingstein, P., Ilyina, T., Jones, C. D., Hajima, T., Tjiputra, J. F., and Wu, T.: Carbon-concentration and carbon-temperature feedbacks in CMIP5 Earth system models, *J. Climate*, 26, 5289–5314, doi:<http://dx.doi.org/10.1175/JCLI-D-12-00494.1>, 2013.
- Bala, G., Caldeira, K., Wickett, M., Phillips, T., Lobell, D., Delire, C., and Mirin, A.: Combined climate and carbon-cycle effects of large-scale deforestation, *P. Natl. Acad. Sci. USA*, 104, 6550–6555, doi:<http://dx.doi.org/10.1073/pnas.0608998104>, 2007.
- Bathiany, S., Claussen, M., Brovkin, V., Raddatz, T., and Gayler, V.: Combined biogeophysical and biogeochemical effects of large-scale forest cover changes in the MPI earth system model, *Biogeosciences*, 7, 1383–1399, doi:<http://dx.doi.org/10.5194/bg-7-1383-2010>, 2010.
- Boisier, J., de Noblet Ducoudré, N., Pitman, A., Cruz, F., Delire, C., van den Hurk, B., van der Molen, M., Müller, C., and Voldoire, A.: Attributing the impacts of land-cover changes in temperate regions on surface temperature and heat fluxes to specific causes: results from the first LUCID set of simulations, *J. Geophys. Res.*, 117, D12116, doi:<http://dx.doi.org/10.1029/2011JD017106>, 2012.
- Bonan, G., Pollard, D., and Thompson, S.: Effects of boreal forest vegetation on global climate, *Nature*, 359, 716–718, 1992.
- Bonan, G.: Forests and climate change: forcings, feedbacks, and the climate benefits of forests, *Science*, 320, 1444–1449, 2008.
- Brovkin, V., Sitch, S., Von Bloh, W., Claussen, M., Bauer, E., and Cramer, W.: Role of land cover changes for atmospheric CO₂ increase and climate change during the last 150 years, *Global Change Biol.*, 10, 1253–1266, 2004.
- Brovkin, V., Raddatz, T., Reick, C., Claussen, M., and Gayler, V.: Global biogeophysical interactions between forest and climate, *Geophys. Res. Lett.*, 36, L07405, doi:<http://dx.doi.org/10.1029/2009GL037543>, 2009.
- Brovkin, V., Boysen, L., Arora, V., Boisier, J., Cadule, P., Chini, L., Claussen, M., Friedlingstein, P., Gayler, V., van den Hurk, B., Hurtt, G., Jones, C., Kato, E., de Noblet Ducoudré, N., Pacifico, F., Pongratz, J., and Weiss, M.: Effect of anthropogenic land-use and land cover changes on climate and land carbon storage in CMIP5 projections for the 21st century., *J. Climate*, doi:<http://dx.doi.org/10.1175/JCLI-D-12-00623.1>, in press, 2013.
- Claussen, M., Brovkin, V., and Ganopolski, A.: Biogeophysical vs. biogeochemical feedbacks of large-scale land cover change, *Geophys. Res. Lett.*, 28, 1011–1014, 2001.
- Davin, E. and de Noblet Ducoudré, N.: Climatic impact of global-scale deforestation: radiative vs. nonradiative processes, *J. Climate*, 23, 97–112, 2010.
- Dufresne, J.-L., Foujols, M.-A., Denvil, S., Caubel, A., Marti, O., Aumont, O., Balkanski, Y., Bekki, S., Bellenger, H., Benshila, R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Cheruy, F., Codron, F., Cozic, A., Cugnet, D., de Noblet, N., Duvel, J.-P., Ethe, C., Fairhead, L., Fichetef, T., Flavoni, S., Friedlingstein, P., Grandpeix, J.-Y., Guez, L., Guilyardi, E., Hauglustaine, D., Hourdin, F., Idelkadi, A., Ghattas, J., Jous-saume, S., Kageyama, M., Krinner, G., Labetoulle, S., Lahellec, A., Lefebvre, M.-P., Lefevre, F., Levy, C., Li, Z. X., Lloyd, J., Lott, F., Madec, G., Mancip, M., Marchand, M., Masson, S., Meurdesoif, Y., Mignot, J., Musat, I., Parouty, S., Polcher, J., Rio, C., Schulz, M., Swingedouw, D., Szopa, S., Talandier, C., Terray, P., Viovy, N., and Vuichard, N.: Climate change projections using the IPSL-CM5 Earth System Model: from CMIP3 to CMIP5, *Clim. Dynam.*, 40, 1–43, 2013.
- Ellis, E.: Anthropogenic transformation of the terrestrial biosphere, *Philos. T. Roy. Soc. A*, 369, 1010–1035, 2011.
- Findell, K. L., Knutson, T. R., and Milly, P.: Weak simulated extratropical responses to complete tropical deforestation, *J. Climate*, 19, 2835–2850, 2006.
- Friedlingstein, P., Cox, P., Betts, R., Bopp, L., Von Bloh, W., Brovkin, V., Cadule, P., Doney, S., Eby, M., Fung, I., Bala, G., John, J., Jones, C., Joos, F., Kato, T., Kawamiya, M., Knorr, W., Lindsay, K., Matthews, H. D., Raddatz, T., Rayner, P., Reick, C., Roeckner, E., Schnitzler, K.-G., Schnur, R., Strassmann, K., Weaver, A. J., Yoshikawa, C., and Zeng, N.: Climate-carbon cycle feedback analysis: results from the C4MIP model intercomparison, *J. Climate*, 19, 3337–3353, 2006.
- Gillett, N., Arora, V., Matthews, D., and Allen, M.: Constraining the ratio of global warming to cumulative CO₂ emissions using CMIP5 simulations, *J. Climate*, 26, 6844–6858, doi:<http://dx.doi.org/10.1175/JCLI-D-12-00476.1>, 2013.
- Giorgetta, M. A., Jungclaus, J., Reick, C. H., Legutke, S., Bader, J., Böttinger, M., Brovkin, V., Crueger, T., Esch, M., Fieg, K., Glushak, K., Gayler, V., Haak, H., Hollweg, H.-D., Ilyina, T., Kinne, S., Kornbluh, L., Matei, D., Mauritsen, T., Mikolajewicz, U., Mueller, W., Notz, D., Pithan, F., Raddatz, T., Rast, S., Redler, R., Roeckner, E., Schmidt, H., Schnur, R., Segschneider, J., Six, K. D., Stockhause, M., Timmreck, C., Wegner, J., Widmann, H., Wieners, K.-H., Claussen, M., Marotzke, J., and Stevens, B.: Climate and carbon cycle changes from 1850 to 2100 in MPI-ESM simulations for the Coupled Model Intercom-

- parison Project phase 5, *J. Adv. Model. Earth Syst.*, 5, 572–597, [doi:http://dx.doi.org/10.1002/jame.20038](http://dx.doi.org/10.1002/jame.20038), 2013.
- Goll, D. S., Brovkin, V., Parida, B. R., Reick, C. H., Kattge, J., Reich, P. B., van Bodegom, P. M., and Niinemets, Ü.: Nutrient limitation reduces land carbon uptake in simulations with a model of combined carbon, nitrogen and phosphorus cycling, *Biogeosciences*, 9, 3547–3569, [doi:http://dx.doi.org/10.5194/bg-9-3547-2012](http://dx.doi.org/10.5194/bg-9-3547-2012), 2012.
- Houghton, R. A., House, J. I., Pongratz, J., van der Werf, G. R., DeFries, R. S., Hansen, M. C., Le Quéré, C., and Ramankutty, N.: Carbon emissions from land use and land-cover change, *Biogeosciences*, 9, 5125–5142, [doi:http://dx.doi.org/10.5194/bg-9-5125-2012](http://dx.doi.org/10.5194/bg-9-5125-2012), 2012.
- Hurt, G. C., Chini, L. P., Frolking, S., Betts, R., Feddema, J., Fischer, G., Goldewijk, K. K., Hibbard, K., Janetos, A., Jones, C., Kindermann, G., Kinoshita, T., Riahi, K., Shevliakova, E., Smith, S., Stehfest, E., Thomson, A., Thornton, P., van Vuuren, D., and Wang, Y. P.: Harmonization of Global Land Use Scenarios for the Period 1500–2100 for IPCC AR5, Integrated Land Ecosystem Atmosphere Processes Study (iLEAPS) Newsletter, iLEAPS Newsletter, iLEAPS International Project Office, University of Helsinki, http://www.ileaps.org/sites/ileaps.org/files/newsletters/NL7_9MB.pdf, last access: 3 April 2014, 6–8, 2009.
- Hurt, G. C., Chini, L. P., Frolking, S., Betts, R. A., Feddema, J., Fischer, G., Fisk, J. P., Hibbard, K., Houghton, R. A., Janetos, A., Jones, C., Kindermann, G., Kinoshita, T., Klein Goldewijk, K., Riahi, K., Shevliakova, E., Smith, S., Stehfest, E., Thomson, A., Thornton, P., van Vuuren, D. P., and Wang, Y. P.: Harmonization of land-use scenarios for the period 1500–2100: 600 years of global gridded annual land-use transitions, wood harvest, and resulting secondary lands, *Climatic Change*, 109, 1–45, 2011.
- Jones, C., Robertson, E., Arora, V., Friedlingstein, P., Shevliakova, E., Bopp, L., Brovkin, V., Hajima, T., Kato, E., Kawamiya, M., Liddicoat, S., Lindsay, K., Reick, C., Roelandt, C., Segschneider, J., and Tjiputra, J.: 21st Century compatible CO₂ emissions and airborne fraction simulated by CMIP5 Earth System models under 4 Representative Concentration Pathways, *J. Climate*, 26, 4398–4413, [doi:http://dx.doi.org/10.1175/JCLI-D-12-00554.1](http://dx.doi.org/10.1175/JCLI-D-12-00554.1), 2013.
- Keeling, C. D., Piper, S. C., Bollenbacher, A. F., and Walker, J. S.: Atmospheric CO₂ Records from Sites in the SIO Air Sampling Network, Trends: a Compendium of Data on Global Change, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, Tenn., USA, 2009.
- Matthews, H., Weaver, A., Meissner, K., Gillett, N., and Eby, M.: Natural and anthropogenic climate change: incorporating historical land cover change, vegetation dynamics and the global carbon cycle, *Clim. Dynam.*, 22, 461–479, 2004.
- Moss, R. H., Edmonds, J. A., Hibbard, K. A., Manning, M. R., Rose, S. K., van Vuuren, D. P., Carter, T. R., Emori, S., Kainuma, M., Kram, T., Meehl, G. A., Mitchell, J. F. B., Nakicenovic, N., Riahi, K., Smith, S., Stouffer, R. J., Thomson, A. M., Weyant, J. P., and Wilbanks, T. J.: The next generation of scenarios for climate change research and assessment, *Nature*, 463, 747–756, 2010.
- Pithan, F. and Mauritsen, T.: Arctic amplification dominated by temperature feedbacks in contemporary climate models, *Nat. Geosci.*, 7, 181–184, [doi:http://dx.doi.org/10.1038/ngeo2071](http://dx.doi.org/10.1038/ngeo2071), 2014.
- Pitman, A. J., de Noblet-Ducoudré, N., Cruz, F. T., Davin, E. L., Bonan, G. B., Brovkin, V., Claussen, M., Delire, C., Ganzeveld, L., Gayler, V., van den Hurk, B. J. J. M., Lawrence, P. J., van der Molen, M. K., Müller, C., Reick, C., Seneviratne, S. I., Strengers, B. J., and Voldoire, A.: Uncertainties in climate responses to past land cover change: first results from the LUCID intercomparison study, *Geophys. Res. Lett.*, 36, L14814, [doi:http://dx.doi.org/10.1029/2009GL039076](http://dx.doi.org/10.1029/2009GL039076), 2009.
- Pitman, A. J., de Noblet-Ducoudré, N., Avila, F. B., Alexander, L. V., Boisier, J.-P., Brovkin, V., Delire, C., Cruz, F., Donat, M. G., Gayler, V., van den Hurk, B., Reick, C., and Voldoire, A.: Effects of land cover change on temperature and rainfall extremes in multi-model ensemble simulations, *Earth Syst. Dynam.*, 3, 213–231, [doi:http://dx.doi.org/10.5194/esd-3-213-2012](http://dx.doi.org/10.5194/esd-3-213-2012), 2012.
- Pongratz, J., Reick, C., Raddatz, T., and Claussen, M.: Effects of anthropogenic land cover change on the carbon cycle of the last millennium, *Global Biogeochem. Cy.*, 23, GB4001, [doi:http://dx.doi.org/10.1029/2009GB003488](http://dx.doi.org/10.1029/2009GB003488), 2009.
- Pongratz, J., Reick, C., Raddatz, T., and Claussen, M.: Biogeophysical vs. biogeochemical climate response to historical anthropogenic land cover change, *Geophys. Res. Lett.*, 37, L08702, [doi:http://dx.doi.org/10.1029/2010GL043010](http://dx.doi.org/10.1029/2010GL043010), 2010.
- Pongratz, J., Reick, C., Raddatz, T., Caldeira, K., and Claussen, M.: Past land use decisions have increased mitigation potential of reforestation, *Geophys. Res. Lett.*, 38, L15701, [doi:http://dx.doi.org/10.1029/2011GL047848](http://dx.doi.org/10.1029/2011GL047848), 2011.
- Reick, C., Raddatz, T., Brovkin, V., and Gayler, V.: The representation of natural and anthropogenic land cover change in MPI-ESM, *J. Adv. Model. Earth Syst.*, 5, 459–482, [doi:http://dx.doi.org/10.1002/jame.20022](http://dx.doi.org/10.1002/jame.20022), 2013.
- Riahi, K., Rao, S., Krey, V., Cho, C., Chirkov, V., Fischer, G., Kindermann, G., Nakicenovic, N., and Rafaj, P.: RCP 8.5 – A scenario of comparatively high greenhouse gas emissions, *Climatic Change*, 109, 1–25, 2011.
- Shevliakova, E., Pacala, S., Malyshev, S., Hurt, G., Milly, P., Caspersen, J., Sentman, L., Fisk, J., Wirth, C., and Crevoisier, C.: Carbon cycling under 300 years of land use change: importance of the secondary vegetation sink, *Global Biogeochem. Cy.*, 23, GB2022, [doi:http://dx.doi.org/10.1029/2007GB003176](http://dx.doi.org/10.1029/2007GB003176), 2009.
- Shevliakova, E., Stouffer, R. J., Malyshev, S., Krasting, J. P., Hurt, G. C., and Pacala, S. W.: Historical warming reduced due to enhanced land carbon uptake, *P. Natl. Acad. Sci. USA*, 110, 16730–16735, 2013.
- Sitch, S., Brovkin, V., von Bloh, W., van Vuuren, D., Eickhout, B., and Ganopolski, A.: Impacts of future land cover changes on atmospheric CO₂ and climate, *Global Biogeochem. Cy.*, 19, GB2013, [doi:http://dx.doi.org/10.1029/2004GB002311](http://dx.doi.org/10.1029/2004GB002311), 2005.

- 795 Stocker, B., Feissli, F., and Strassmann, K.: Past and
future carbon fluxes from land use change, shifting
cultivation and wood harvest, *Tellus B.*, 66, 23188,
doi:<http://dx.doi.org/http://dx.doi.org/10.3402/tellusb.v66.23188><http://dx.doi.org/10.3402/tellusb.v66.23188>,
2014.
- 800 Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An overview of
CMIP5 and the experiment design, *B. Am. Meteorol. Soc.*, 93,
485–498, 2012.
- Van der Molen, M., van den Hurk, B., and Hazeleger, W.: A damp-
ened land use change climate response towards the tropics, *Clim.
Dynam.*, 37, 2035–2043, 2011.
- 805 Watanabe, S., Hajima, T., Sudo, K., Nagashima, T., Take-
mura, T., Okajima, H., Nozawa, T., Kawase, H., Abe, M., Yoko-
hata, T., Ise, T., Sato, H., Kato, E., Takata, K., Emori, S.,
and Kawamiya, M.: MIROC-ESM 2010: model description
and basic results of CMIP5-20c3m experiments, *Geosci.
810 Model Dev.*, 4, 845–872, doi:<http://dx.doi.org/10.5194/gmd-4-845-2011>[10.5194/gmd-4-845-2011](http://dx.doi.org/10.5194/gmd-4-845-2011), 2011.
- Werth, D. and Avissar, R.: The local and global
effects of Amazon deforestation, *J. Geo-
phys. Res.-Atmos.*, 107, LBA55-1–LBA55-8.27,
815 doi:<http://dx.doi.org/10.1029/2001JD000717>[10.1029/2001JD000717](http://dx.doi.org/10.1029/2001JD000717),
2002.
- Wilkenskjeld, S., Kloster, S., Pongratz, J., Raddatz, T., and Re-
ick, C.: Comparing the influence of net and gross anthro-
pogenic land use and land cover changes on the carbon cycle
820 in the MPI-ESM, *Biogeosciences Discussions*, 11, 5443–5469,
doi:<http://dx.doi.org/10.5194/bgd-11-5443-2014>, 2014.
- Zwiers, F., von Storch, H., and Max-Planck-Institut für Me-
teorologie: Taking Serial Correlation into Account in Tests
of the Mean, Max-Planck-Institut für Meteorologie, *J.
825 Climate*, 8, 336–351, doi:[http://dx.doi.org/10.1175/1520-0442\(1995\)008%3C0336:TSCIAI%3E2.0.CO;2](http://dx.doi.org/10.1175/1520-0442(1995)008%3C0336:TSCIAI%3E2.0.CO;2)[10.1175/1520-0442\(1995\)008%3C0336:TSCIAI%3E2.0.CO;2](http://dx.doi.org/10.1175/1520-0442(1995)008%3C0336:TSCIAI%3E2.0.CO;2), 1995.

Table 1. Overview of CMIP5 and LUCID simulations based on CMIP5 standard simulations for RCP8.5 and the employed terminology exemplified with near-surface temperature T .
table

Simulation	terminology	CO ₂ concentration	LULCC
ESM	$T_{LULCC}^{eCO_2}$	Interactive (emission-driven)	As in RCP
L1B	$T_{no\ LULCC}^{eCO_2}$	Interactive	Fixed to year 2005
L1A	$T_{no\ LULCC}^{cCO_2}$	Prescribed (concentration-driven, output of the ESM run)	Fixed to year 2005
RCP	$T_{LULCC}^{cCO_2\ RCP}$	Prescribed from RCP8.5 (Moss et al., 2010)	Transient scenario (MESSAGE, Riahi et al., 2011) (Hurtt et al., 2011)
L2A	$T_{no\ LULCC}^{cCO_2\ RCP}$	As in RCP	Fixed to year 2005

Table 2. Overview of model setups and analysis strategies.

Difference	set-up differences	terminology/scientific interpretation
ESM-L1A	same CO ₂ concentration; with-without LULCC	BGP-effects: $\Delta T(\Delta LULCC, \Delta CO_2 = 0) = \Delta T_{BGP}$, $\Delta C(\Delta LULCC, \Delta CO_2 = 0) = \Delta C_{\Delta LULCC}$
L1A-L1B	different CO ₂ concentrations; both without LULCC	BGC-effects: $\Delta T(\Delta LULCC = 0, \Delta CO_2) = \Delta T_{BGC}$, $\Delta C(\Delta LULCC = 0, \Delta CO_2) = \Delta C_{BGC}$
ESM-L1B	different CO ₂ concentrations; with-without LULCC	net effects: $\Delta T(\Delta LULCC, \Delta CO_2) = \Delta T_{net}$, $\Delta C(\Delta LULCC, \Delta CO_2) = \Delta C_{net}$

Table 3. Atmospheric CO₂ (ppm) concentrations in 2100.

Model	CO ₂ LULCC	CO ₂ no LULCC	$\Delta CO_2 \Delta LULCC$
MPI	951	885	66
CAN	1037	1024	12
MIR	1134	1113	22
MESSAGE	926		

Table 4. ΔT_{BGP} and ΔT_{BGC} (K), averaged over the period 2071–2100: globally and over areas where LULCC $\geq 10\%$ of the grid cell. The asterisk (*) marks values with statistical significance ($\geq 95\%$) of a Student's t test accounting for autocorrelation. The temperature change over the 21st relative to 2006 century due to fossil fuel forcings only is given by $\Delta T_{no\ LULCC}^{eCO_2}$ (L1B simulation).

Model	$\Delta T_{no\ LULCC}^{eCO_2}$ Global	ΔT_{BGC} Global	ΔT_{BGP} LULCC $\geq 10\%$	ΔT_{BGP}
MPI	3.02	0.23*	0.02	0.03
CAN**	3.60	0.07*	0.02	0.10*
MIR	4.73	0.12*	−0.01	−0.47*
IPSL	3.70	−0.02	−0.03	−0.16*

** The BGP part in CAN is calculated as ΔT_{BGP}^{RCP} .

Table 5. Comparison of simulated ΔT_{BGC} (as in Table 4) to temperature changes derived from the TRCE approach (transient response of temperature to cumulative emissions; ΔT_{TRCE} Gillett et al., 2013). LULCC emissions are derived from the losses in land carbon storage ($\Delta C_{\Delta\text{LULCC}}$) multiplied by the TRCE values from Gillett et al. (2013) to approximate temperature changes. Results for RCP simulations ($\approx \Delta T_{\text{TRCE}}^{\text{RCP}}$) are taken from Brovkin et al. (2013). The asterisk * marks values of statistical significance ($p < 0.05$).

Model	ΔT_{BGC} (K)	TRCE (°K Tt C ⁻¹)	$\Delta C_{\Delta\text{LULCC}}^{\text{a}}$ (GtC)	$\approx \Delta T_{\text{TRCE}}$ (K)	$\Delta C_{\Delta\text{LULCC}}^{\text{RCP b}}$ (GtC)	$\approx \Delta T_{\text{TRCE}}^{\text{RCP b}}$ (K)
MPI	0.23*	1.604	218	0.35	205	0.33
CAN	0.07*	2.365	34	0.08	34	0.08
MIR	0.12*	2.151	57	0.12	62	0.13
IPSL	-0.02	1.585	31	0.06	37	0.06

^a Changes for CAN are calculated indirectly by $\Delta T_{\text{net}} - \Delta T_{\text{BGC}}^{\text{RCP}}$.

^b Brovkin et al. (2013).

Table 6. Global changes in cumulative land carbon fluxes ΔC (cumulative from 2006 until 2100 in GtC) in 2100 due to the various effects of LULCC: changes in vegetation distribution and climate ($\Delta C_{\Delta\text{LULCC}}$), net effect (ΔC_{net}), and BGC effects (ΔC_{BGC}).

Model	simulation-index	ΔC	$\Delta C^{\text{RCP b}}$
MPI	ΔLULCC	-218	-205
	net	-179	
	BGC	39	
CAN ^a	ΔLULCC	-34	-34
	net	-29	
	BGC	4	
MIR	ΔLULCC	-57	-62
	net	-56	
	BGC	2	
IPSL	ΔLULCC	-35	-37
	net	-38	
	BGC	-3	

^a Changes for CAN are calculated indirectly by $\Delta T_{\text{net}} - \Delta T_{\text{BGC}}^{\text{RCP}}$.

^b Brovkin et al. (2013).




Fig. 1. Maps displaying the change in near-surface temperature (K) averaged over 2071–2100 for each model. Only areas are shown where changes are statistically significant; (a) ΔT_{BGP} (for CAN ΔT_{BGP}^{RCP}); (b) ΔT_{BGC} (for CAN $\Delta T_{net} - \Delta T_{BGP}^{RCP}$); (c) ΔT_{net} . figure




Fig. 2. Relative changes in near-surface temperature: Comparison of ΔT_{BGP} relative to $\Delta T_{noLULCC}^{cCO_2}$ (L1A simulation), that is the BGP impacts of LULCC compared to the impacts of anthropogenic carbon emissions (both fossil-fuel and LULCC) on near-surface temperature (in %). Depicted are mean 2071–2100 values minus the 2006 state (indicated by “ Δ_t ”). Positive (negative) values indicate that BGP effects (ΔT_{BGP}) enhance (dampen) the change caused by LULCC and other anthropogenic emissions. Analysis is done for the following regions: Eurasia (EURA), North America (NOAM), South America (SOAM), Africa (AFRI), Australia (AUST), land (land area excluding ice sheets) and global (total area on Earth). A list of exact values can be found in the Supplement Table S2.




Fig. 3. 10 years-running global means of net changes due to LULCC in the terrestrial carbon content (in GtC). Dark solid lines represent ΔC_{net} , dashed lines $\Delta C_{\Delta LULCC}$ and light solid lines ΔC_{BGC} .