



**Global marine fish landings and atmospheric planetary wave forcing**

A. M. Powell Jr. and J. Xu

**Decadal regime shift linkage between global marine fish landings and atmospheric planetary wave forcing**

A. M. Powell Jr.<sup>1</sup> and J. Xu<sup>2</sup>

<sup>1</sup>Center for Satellite Applications and Research (STAR), NOAA/NESDIS, 5830 University Research Ct, College Park, MD 20740, USA

<sup>2</sup>Environmental Science and Technological Center(ESTC), College of Science, George Mason University, USA

Received: 26 June 2014 – Accepted: 15 July 2014 – Published: 7 August 2014

Correspondence to: A. M. Powell Jr. (al.powell@noaa.gov) and J. Xu (jxu14@gmu.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

This investigation focuses on a global forcing mechanism for decadal regime shifts and their subsequent impacts. The proposed global forcing mechanism is the global atmospheric planetary waves that can lead to changes in the global surface air–sea conditions and subsequently fishery changes. In this study, the five decadal regime shifts (1956–1957, 1964–1965, 1977–1978, 1988–1989, and 1998–1999) in the recent 59 years (1950–2008) have been identified based on student  $t$  tests and their association with global marine ecosystem change has been discussed. Changes in the three major oceanic (Pacific, Atlantic and Indian) ecosystems will be explored with the goal of demonstrating the linkage between stratospheric planetary waves and the ocean surface forcing that leads to fisheries impacts. Due to the multidisciplinary audience, the global forcing mechanism is described from a top-down approach to help the multidisciplinary audience follow the analysis. Following previous work, this analysis addresses how changes in the atmospheric planetary waves may influence the vertical wind structure, surface wind stress, and their connection with the global ocean ecosystems based on a coupling of the atmospheric regime shifts with the decadal regime shifts determined from marine life changes. The multiple decadal regime shifts related to changes in marine life are discussed using the United Nations Food and Agriculture Organization’s (FAO) global fish capture data (catch/stock). Analyses are performed to demonstrate the interactions between the atmosphere, ocean, and fisheries are a plausible approach to explaining decadal climate change in the global marine ecosystems and its impacts. The results show a consistent mechanism, ocean wind stress, responsible for marine shifts in the three major ocean basins. Changes in the planetary wave pattern affect the ocean wind stress patterns. A change in the ocean surface wind pattern from long wave (relatively smooth and less complex) to shorter wave (more convoluted and more complex) ocean surface wind stress creates changes in the ocean marine fisheries.

## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 1 Introduction

The global marine ecosystems in the world's major oceans exhibit long-term variations in time resembling "regime shifts". A regime shift is a transition from one state to another, with the transition period being much shorter than the length of the individual epochs (Overland et al., 2008). Regime shifts are associated with large, abrupt, persistent changes in both atmospheric and oceanic conditions that may be especially pronounced in physical and biological variables. Generally, regime shifts have been found in 1925, 1947, 1977, 1989, and 1998 (Mantua et al., 1997; Minobe, 1999; Beamish et al., 2004; King, 2005; Overland et al., 2008; Powell and Xu, 2011a). The 1976/77 regime shift is the most commonly identified shift in the literature currently and is often identified with high amplitude changes in numerous atmospheric, oceanic, and biological measures cited in the literature (Erbesmeyer et al., 1991; Minobe, 1997, 1999; Overland et al., 2008; Powell and Xu, 2012). A combination of atmospheric and oceanic forcing is thought to cause marine changes by affecting the physical ocean which impacts the ocean habitat or ecosystem. The purpose of this research is to determine whether a common global mechanism can be found consistent with the Chavez proposition (Chavez et al., 2003) that all the world's oceans and fisheries are affected in near synchrony by a simple and direct forcing that is similar in the different ocean basins. If a forcing can be identified, the regime shift strength and timing could be estimated and possibly forecast based on atmospheric, oceanic, and biological change. A regime shift forecast capability could provide decision makers with a significant tool for estimating ocean ecosystem impacts.

Regime shifts in fish populations are difficult to explain on the basis of biological relationships alone or fishing pressure (Chavez et al., 2003). Lehodey et al. (2006) stated that fish population variability is closely related to environmental variability. Based on the fishery landing data on the Canada's Pacific coast, Beamish et al. (2004) pointed out that the fish catches were related to trends in the climate and ocean environment that can be considered to be associated with regime shifts. In recent years numerous

# ESDD

5, 945–989, 2014

## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



long-term changes in physical forcing have been observed at global, regional and basin scales as a result of climate changes. In particular, the impacts of SST on ocean wind stress and vice versa have been investigated due to its key effects on atmosphere-ocean interaction (Chelton et al., 2001). Impacts of various forcings on biological processes supporting fish production in marine ecosystems have already been observed and may be used as proxies to estimate further global climate change impacts. As an example, El Nino's well known wind direction impacts on upwelling/downwelling off South America's west coast resulted in periods of extremely good and extremely poor fishing conditions that have had severe economic consequences. The effects of El Nino are felt in areas outside the west coast of South America. Physical factors that could impact fish production include atmospheric circulation and oceanic environmental (wind stress patterns, upwelling/downwelling, water temperature, ocean currents, spawning temperatures, etc.) variability patterns (Roy and Reason, 2001; Sugimoto et al., 2001). All of these factors are affected by the ocean wind stress. Chavez et al. (2003) remarked that the mechanism(s) responsible for the abrupt regime shifts should be relatively direct and simple, similar in the different regions, and likely linked with large-scale atmospheric and oceanic forcing. In this research, the key issue is to assess global atmospheric planetary wave structures to identify regime shifts in terms of the global atmospheric forcing (wind stress) that have ocean impacts and marine impacts (changes in normalized fish landings). The goal is to develop a global regime concept to support an improved understanding of the near synchronous marine changes in all the world's ocean basins. The variability of large marine populations and their associated ecosystems are likely impacted through global regime shift patterns or decadal climate change.

To detect regime shifts in the global marine environment, the United Nations Food and Agriculture Organization's (FAO) global fish capture (landings) data was used. The approach was to first detect the year(s) when regime shifts occurred in the global fish capture statistics using the student  $t$  test. A comparison with the planetary wave index

analysis was used to assess the likelihood the marine regime shifts were caused by global atmospheric forcing.

## 2 Data

The data used in this study includes the FAO global statistics on fish capture production. The atmospheric temperature and geopotential height fields used for the planetary wave amplitude index are from the NCEP/NCAR reanalysis (1948–2008).

### 2.1 FAO's fish capture data

The FAO fish capture data is the only available data set on the global fish catch (Froese et al., 2012) and it is also the most broadly used and accepted database (Liddel, personal communication, 2014). The annual series of capture production began in 1950. The data includes all quantities caught and landed for both food and feed purposes but excludes discards. Fish catches/landings are expressed in live weight which is the nominal weight of the aquatic organisms at the time of capture. According to the marine area where caught, capture production is also classified into 14 major marine fishing areas (Fig. 1) encompassing the waters of the Atlantic, Indian and Pacific oceans. The fish landing data for the Antarctic and Arctic Oceans with their adjacent seas (Fig. 1) are not included in this data and could be a source of bias. The FAO database contains the volume of aquatic species caught by country or area, by species, by FAO major fishing areas, and year, for all commercial, industrial, recreational and subsistence purposes. The nine families of FAO aquatic species are listed in Fig. 1 along with a map of the 14 sub-regional data collection areas. To minimize the effect of episodic or exceptional events, similar to the processing used in the previous studies (deYoung et al., 2004; Powell and Xu, 2013), this data undergoes normalization as described in Sect. 2.3.

## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The authors address the use of fish catch data in this analysis. There are difficulties with using this data. It is clear that reporting procedures have resulted in under reporting fish catch in some countries and over reporting in others, particularly China (Watson and Pauly, 2001). Policies and fishing rights changes in various regions of the world as well as changes in equipment can influence the catch statistics. In addition, catches can shift for many reasons including reclassifying the fish catch data categories. The adoption of marine protected areas and policies to improve protected/declining species seem to have positive impacts on the abundance of fish and potentially fish catch (Hilborn and Branch, 2013). The reliability and issues associated with the fish catch data comes from two articles Watson and Pauly (2001) and Hilborn and Branch (2013). Using the FAO fish catch data, Froese et al. (2012) showed explicitly that trends in catch data are not an artifact of the applied method and are consistent with trends in biomass data of fully assessed stocks.

The authors believe the information contained in previous articles discuss the basic issues with fish catch data. However, this is the only comprehensive global data base (Froese et al., 2012) from which to draw upon actual measurements. For the purposes of this analysis, the FAO fish catch data is the most prominent and comprehensive fishery data set available to assess linkages between global physical environmental forcings and the biological impacts on the world's fisheries. By normalizing the data, in aggregate, it should reflect real changes in fish catch whether the fish migrated to alternative regions, the ocean ecosystems changed, or other factors such as changes in fishing equipment or fishing policies played a role in the fish catch changes. Because this research seeks to identify the physical coupling between the atmospheric forcing and the resultant impact on fisheries, it is thought to be highly unlikely that changes in fishing strategies and policies are likely to significantly affect five major marine regime shifts. In addition, the coupling of the marine regime shifts to atmospheric regimes shifts should provide significant evidence that the wind driven ocean circulations play a role in the synchronous changes in global fish populations that have been identified in the literature. All the fish landing data were normalized based on the available record

## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



from 1950 through 2008. This normalization removes biases due to under and over reporting of fish landings and provides a reasonable comparison of the marine system. The purpose of using this data is to demonstrate that changes in environmental forcings impact the world's oceans in a similar manner, but not necessarily impacting the same fish species group(s) in each region or ocean basin. For the purposes of this analysis, the authors think the data is sufficiently robust to make a determination about environmental impacts on fish species groups. Analyses of this type help the community understand the data and the potential impacts of factors that influence the statistics. In this study, the landing data provides insights into environmental impacts on the biological processes related to the world's fisheries.

### 2.2 NCEP/NCAR reanalysis (NNR)

The monthly NCEP/NCAR reanalysis with a  $2.5^\circ \times 2.5^\circ$  grid resolution is used in the period of 1950–2008. It should be noted that the reanalysis period of 1958–1978 has no satellite data. The Television Infrared Observation Satellite (TIROS) Operational Vertical Sounder (TOVS) data includes the Microwave Sounding Unit (MSU), High Resolution Infrared Radiation Sounder (HIRS) and Stratospheric Sounding Units (SSU). The satellite information was not available for use in forecast models before the end of 1978; this is true for the atmospheric reanalysis used for this study. The Special Sensing Microwave/Imager (SSM/I) data was assimilated in the National Weather Service forecast system from 1993. The lack of satellite data in the early years could degrade the atmospheric analysis prior to the availability of the satellite data. The NNR temperature and geopotential height fields between 20–70 hPa (stratosphere) were used in this study to identify atmospheric regime shifts and their vertical connectivity to the ocean surface since previous analyses indicated the strongest planetary waves were in the stratosphere (Powell and Xu, 2012). Selected atmospheric pressure levels between the stratosphere and the surface (1000 hPa) were used to demonstrate the dynamic linkage between all levels of the atmosphere and the relationship to surface environmental conditions.

### 3 Methodology

#### 3.1 Normalization processing

To make a comparison of the interannual and decadal variability for each individual parameter over the time series, each parameter ( $x_i$ ,  $i = 1, 2, \dots, N$ ) must be normalized from the starting date to the end of the data set. The atmospheric data (20–70 hPa) and each fish species group were first normalized using the calculation expressed as follows:

$$\tilde{x}_i = \frac{x_i - \bar{x}}{\sigma}$$

Where  $\bar{x} = \frac{1}{N} \sum_{i=1}^N x_i$  and  $\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \bar{x})^2}$ .

See Powell and Xu (2011b) for more details. This provided the ability to determine the regime shift periods using a consistent approach between dissimilar data sets.

#### 3.2 Fourier spectrum analysis

The geopotential height, temperature and wind fields from the NCEP-NCAR reanalysis were decomposed into Fourier harmonics and the Fourier coefficients were used to recompose the temporal field for single zonal waves. Wavenumbers 1 thru 6 are Fourier decomposed from the 59 yr (1950–2008) data set. The geopotential height fields are consistent with the manner in which most meteorologists display the weather pattern for use in forecasting.

The height field anomalies were computed to highlight the changes in the global fields and to show the effect on the various levels of atmospheric low and high pressure systems. The height field anomalies across the multiple regime shifts will be discussed later in the paper.

## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





### 3.3 Identification of the regime shifts

To identify the regime shifts, the atmospheric planetary wave amplitude index [PWA] (Powell and Xu, 2011b) established the atmospheric planetary wave strength which was shown to influence ocean forcing through surface wind stress (Powell and Xu, 2012). The 55–75° N latitudes were used to identify mid-latitude forcing associated with global synoptic scale changes in the wind and vertical wave propagation which was previously analyzed as strong in the region. The 90% confidence level was established for the running student  $t$  test decadal period comparisons and provided an assessment of the most confident regime shift dates. For this analysis, the regime shift test was completed using 5 years on either side of the running target year (an 11 year interval in total, so it is called the 11 year (decadal) running  $t$  test in the following sections) to determine whether a significant regime shift on the decadal time scale had occurred. The most confident dates (positive or negative) are used as the dates of key atmospheric regime shifts. The planetary wave amplitude indices show regime shifts (Fig. 2) occurring at approximately 1956–1957, 1964–1965, 1977–1978, 1988–1989, and 1998–1999. The averaged values for each decadal period provide a way to assess the degree of the change across each shift and are shown in Fig. 2.

Since atmospheric forcing of the ocean is thought to be a critical factor, the atmospheric regime shifts are matched with the effects of the ocean surface stress and marine changes via fish landings in the research analysis.

### 3.4 Surface kinetic energy change with wavenumber

To measure the spatial characteristics of the surface atmospheric circulation and its forcing, the kinetic energy (KE) in the atmospheric surface waves over each major ocean domain is calculated. The KE is expressed by  $1/2(U^2(n) + V^2(n))$ , where  $n$  is the zonal wave number,  $U$  and  $V$  are the zonal and meridional wind components, respectively. The KE is assessed based on the wind anomalies associated with each ocean domain (Pacific, Atlantic and Indian oceans).

## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 4 Results

### 4.1 Regime shifts from the global FAO fish landing

The purpose of this section is to discuss the marine regime shifts identified in the global FAO fish landing data, indicating abrupt biological change, that can be associated with the atmospheric regime shifts. This begins the process of linking global biological change with physical changes in the ocean, ocean surface wind stress and atmospheric forcing. To detect the potential global regime shifts, similar to the approach conducted in the determination of global planetary wave regime shifts (Powell and Xu, 2012), the time series of marine landing data over each of the sub-regions were analyzed using the 11 year running student  $t$  test conducted on the normalized FAO fish landing data. The fish landing data included nine fish species groups in fourteen sub-regions (see Fig. 1 and Table A1 in the Appendix) over the Pacific, Indian and Atlantic oceans. The analysis period is 59 years from 1950 to 2008.

Figure 3 shows the detrended time series and 11 year running  $t$  test value for selected fish landings over the three ocean basins. The results clearly show that the year of regime shift occurrence depended on both the fish species group and sub-region. For example, the years of regime shift occurrence are identified (Fig. 3a) at 1966, 1977, 1987 and 2000 for the MDF fish species group (see Appendix A1 for definitions of the fish groups) over the West Central Pacific Ocean (PWC), are different from the years found (Fig. 3j) at 1963, 1972, 1977 and 1988 for the MCF over the East Central Atlantic Ocean (AEC). Regime shift transitions occur within a window of 2–3 years. The peak shift date within the transition window could come ahead or behind the central date during each decade. The individual fish species group graphics may appear to be inconsistent with the climate or ecosystem regime shifts found by previous studies from various data sets. In fact, each individual graph looks rather unique; although, each identifies distinct regime shifts with some similar and some not to other plots. This is likely due to the varying conditions in each region and the response of the various species to the changing conditions. These graphs were plotted for all 14 sub-regions

## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



with their 9 fish species categories for a total of 126 (14 times 9) regional fish species groups. However, when the statistical analysis for all nine fish species over the 14 sub-regions is analyzed, it shows a consistent result with the previous regime shift findings. By computing the frequency of occurrence of the various fish regime shifts for the Pacific, Atlantic and Indian Oceans, the temporal pattern of the most frequent regime shift occurrences matches closely with the regime shifts identified from the changing atmospheric patterns shown earlier.

Appendix Table A2 shows the year when a regime shift occurred in each decade. Based on the number of decadal regime shift occurrences collected in the running two-year interval (consistent with the short transition time for a regime shift), the percentage of fish landing regime shift occurrences (Fig. 4) indicates the regime shifts exceeding the statistical significance test at the 90 % confidence level are identified at 1955–1956, 1964–1965, 1977–1978, 1988–1989 and 1999–2000 in each decade over the Pacific Ocean (Fig. 4a). Similar shifts are observed within one or two years over the Atlantic Ocean (Fig. 4b) but two shifts at 1963–1964 and 1999–2000 satisfy the significance test at 85 % confidence levels. Both the Atlantic and Pacific regime shift patterns are consistent. The fish landing regime shifts over Indian Ocean show some different features from the other two ocean basins. Three regime shifts at 1957–1958, 1989–1990 and 1999–2000 passed the significance test at 90 % confidence level, and are consistent with those in the Atlantic and Pacific. The other three regime shifts are observed at 1969–1970, 1980–1981 and 1995–1996 using the 85 % confidence level. Overall, the results generally reproduced the five regime shifts from previous studies based on various data sets (Mantua et al., 1997; Minobe, 1999; Overland et al., 2008; Powell and Xu, 2011a). However, it is worth noting that the regime shifts in the Indian Ocean are different from their counterparts in the Pacific and Atlantic oceans.

The analysis shown in Fig. 4 identifies similar abrupt shift periods (1955–1958, 1963–1965, 1976–1978, 1987–1990, 1998–2000) in the fish catch data as the atmospheric data (Fig. 2) suggesting a potential link between the atmospheric forcing and the impacts on the fish species. The different regimes shifts in the Indian Ocean

## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



suggest a second forcing mechanism unrelated or indirectly related to direct atmospheric forcing. Since the atmospheric analysis was based on middle to high latitude wind forcing patterns indicative of non-tropical wind conditions, it is possible the Indian Ocean (which is also bounded to the north and west by land masses), may not react completely in concert with the other ocean basins since it is a “tropical ocean”. There is insufficient data to determine whether additional regime shifts exist from other forcings from the analysis technique applied. The remaining lesser shifts identified are substantially weaker and could be due to interannual variability or other factors and have been excluded from direct comparison.

The potential linkage between the five most frequent regime shifts identified in Fig. 4 from the fish landing data that correspond with the five abrupt shifts in the atmospheric planetary wave data (Fig. 2) will be described in the following paragraphs. It should be pointed out that the global fishery regime shifts produce a 15 to 25 % change (Fig. 4) in the sub-region fish species across the regime shifts with lesser percentage changes likely due to interannual variability. In addition, the sub-region species are affected differently even within the same ocean basin. This suggests the regime shift forcing mechanism may have sub-regional effects and will be addressed in Sects. 6 and 7.

### 4.2 Decadal shift in the stratospheric atmospheric planetary wave

The purpose of this section is to discuss the causality of atmospheric regime shifts, primarily in the Earth’s stratosphere, as they relate to the biological abrupt shifts of similar dates found in the FAO fish landing data. This begins the process of demonstrating the top–down linkage between abrupt atmospheric change leading to surface wind stress changes which have impacts on biological variability. This section discusses the global change in atmospheric state between the regime shifts at the strongest level identified in the atmospheric data (50 hPa in pressure located in the stratosphere).

In previous research, the changes in the atmospheric planetary wave pattern showed unique changes between the years with the strongest planetary waves and the weakest ones (Powell and Xu, 2011b). Also, the amplitudes of the planetary waves in the

## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



stratosphere seemed to be associated with ocean changes. Given the findings in the previous studies, an extension of this work investigated how the wave pattern changed across multiple regime shifts (Powell and Xu, 2011a). The analysis was performed at the 50 hPa level (in the stratosphere) where the effect appeared strongest (Powell and Xu, 2011b). The 50 hPa level regime shift periods of 1948–1956, 1957–1964, 1965–1977, 1978–1988, 1989–1998, 1999–2005 showed the global atmospheric planetary wave pattern essentially shifted (changed phase or location) in each of the periods (Fig. 5). However, the method of transferring wave energy from the various levels in the atmosphere to the surface was not specifically addressed in the earlier Powell and Xu (2011a) analysis, but will be addressed in the following section.

Using the five regime shift dates (six regime intervals) identified in both atmospheric and fish landing data sets, a top down analysis was undertaken from the atmosphere to the ocean surface. The atmospheric wave pattern was analyzed to determine whether a cause-effect mechanism could be identified to explain the changes between the regime shift periods. Fourier analysis of wavenumbers 1 through 6 derived from the height fields of the NCEP-NCAR atmospheric reanalysis shows the global wave pattern anomalies (Fig. 5). For the multidisciplinary audience, the patterns represent the significant features in the average northern hemispheric weather pattern for the identified periods. The colored (high pressure with clockwise wind circulation) and uncolored (low pressure with counter clockwise wind circulation) regions in Fig. 5 represent the changes in the strength and position of the dominant weather systems for the periods of interest. The closer the contours are together, the stronger the winds are blowing. The planetary wave anomalies at 50 hPa (in the stratosphere at approximately 20 km in height) are shown for each abrupt shift period: (a) 1948–1956, (b) 1957–1964, (c) 1965–1977, (d) 1978–1988, (e) 1989–1998, (f) 1999–2005. From Powell and Xu (2012), it was shown that the strongest wave amplitudes occurred in the NNR between 20–70 hPa with 50 hPa typically among the strongest levels. In addition, the wavenumber one global atmospheric anomaly pattern tended to change state by approximately reversing positions between the high and low pressure height anomalies (Powell and

## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Xu, 2011b). Since global atmospheric wavenumber one is the strongest (has the most energy) and has two primary “lobes” that represent the primary atmospheric wave (represented by the red-and-white region combination approximately 180° apart and outlined by the black boxes), its effects can be seen in each regime shift period’s height anomaly pattern. The combined impacts of wavenumbers 1 through 6 are shown to demonstrate how the additional, but weaker wavenumbers add complexity to the atmospheric forcing pattern while retaining the dominant wavenumber 1 pattern, although shifted in phase and amplitude.

Starting with the 1948–1956 period, the 1957–1965 pattern is shifted significantly when looking at the primary red (solid contours) and white shaded regions (dashed contours). Comparing the 1948–1956 pattern with each of the subsequent periods (1965–1977, 1978–1988, and 1989–1999) shows a significant shift or “reversal” of the global wave pattern anomalies from one period to the next. Only the last period representing 1999–2005, does not show a significant change in the pattern but the pattern is weaker in terms of the anomaly magnitudes observed in the previous panel. Also, the 1999–2005 period may be too long since another regime shift was possibly identified in 2003 within the period of interest (Peterson et al., 2006; Hatun et al., 2009). Note the wave anomaly positions suggest the continents or geography may play a key role in where the anomalies form and is consistent with meteorological dynamics. However, additional study will be required to confirm this potential geographic association with the pattern and the regime shifts.

There are clear distinguishing shifts and changes in the atmospheric pattern in the stratosphere in Fig. 5. Since the wind patterns change in accordance with the shifts in atmospheric wave pattern, it is reasonable to assume the surface winds and ocean will show corresponding signs of change. The global wave pattern anomalies are indicative of wind changes throughout the depth of the atmosphere and the surface wind stress patterns should reflect any changes.

### 4.3 Top-down connection from stratosphere through surface

The purpose of this section is to relate global stratospheric planetary wave changes to surface wind stress via top-down or vertical forcing processes. To demonstrate a mechanism exists that can transfer substantial energy from the stratosphere to the troposphere (the lower atmosphere in contact with the Earth's surface) through planetary waves, the global height anomalies in the vertical were analyzed. The height anomalies indicate the vertical change in intensity and placement of the global low and high pressure systems which are the lobes of the planetary wave pattern. The wind and temperature gradients associated with high and low pressure systems require a dynamic process(es) to maintain continuity vertically in the atmosphere. The tilting of the systems with height is a recognized consequence among meteorologists of the atmospheric dynamics required to support these systems. Consequently, the investigation focused on large scale planetary wave changes and their associated vertical adjustments in the height fields – used by meteorologists to assess the dynamic state of the atmosphere and make predictions.

The analysis in Fig. 6 was undertaken to demonstrate that dynamic changes in the vertical through the high and low pressure systems provide the surface wind forcing needed to modify the ocean's currents and subsequently impact marine ecosystems at the Earth's surface. The large scale anomaly features seen in the stratosphere at the 50 hPa level (and match the positions of the large pressure systems shown in polar plots in Fig. 5) can be followed to the surface via the generally red and blue colored regions (Fig. 6) which typically weaken in amplitude as they reach approach the surface (1000 hPa level). These height anomalies are consistent with the physical structure and processes for high and low pressure systems that generate the world's daily weather as well. The critical impact of this vertical structure is that it communicates to the Earth's surface a wind forcing pattern that affects the oceans.

Globally, there are strong wind and height anomalous regions over the Pacific, Atlantic and Indian oceans before and after each regime shift. However, a key difference

## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



is the anomaly areas are basically out of phase indicating the global wave pattern has shifted causing regional wind flow changes. For example, prior to 1977–1978 regime shift, the strongest two “height anomaly columns” representing high and low pressure circulations are over central Asia (negative anomaly [blue]) and the central Pacific (positive anomaly [green which turns red as one goes lower in height]). After 1978, the strongest two “height columns” of high and low pressure are over central Asia (positive anomaly, red) and over the central Pacific (negative anomaly, blue). This is essentially a change of phase between the two periods for both the positions and amplitudes of the core wave pattern features. In addition, it is clear the strongest anomaly patterns, both before and after the regime shift, are at the 50 hPa level and generally decrease in strength as they approach the surface. The height anomaly changes span the stratosphere to the surface. In these anomaly fields, a total of four high and low pressure system height anomalies (2 high, two low) can typically be seen – indicating a wavenumber 2 influence is highly probable. Wavenumber 2 has been implicated in vertical energy transport in several studies (Matsuno, 1970; Kodera, 2002; Matthes et al., 2006). The dynamics of weather systems have secondary circulations which may also contribute to the movement of energy in the central regions of the high and low pressure systems. The height anomalies make it clear that energy is being transferred vertically in the global or hemispheric scale waves associated with the high and low pressure systems that are the lobes of the planetary scale waves.

Figure 7 shows the global wind field (streamlines) and height anomalies (shaded areas) near the surface (1000 hPa) for the decadal periods (1948–1956, 1957–1964, 1965–1977, 1978–1988, 1989–1998, 1999–2005). The anomaly positions from the surface (1000 hPa) in Fig. 6 match the colored positions shown in Fig. 7 (Note the domain is from 60° N to 60° S to highlight the wind impact on the oceans compared to Fig. 6 where the domain was from pole to pole.). Comparing any two, time-sequenced sets of wind and height anomalies emphasize the differences from the mean surface atmospheric state in the periods before and after each regime shift. Positive height anomalies (yellow-red) are associated with high pressure systems and clockwise wind flows



## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

and negative height anomalies (blue-purple) are associated with low pressure systems and counter-clockwise wind flows *in the Northern Hemisphere* (note: opposite wind flows occur in the Southern Hemisphere for low and high pressure systems). Since these systems change position and amplitude, it indicates that regional changes are most likely the consequence of these global and hemispheric pattern changes. The key question is whether this can be demonstrated to impact the global marine ecosystems? For this, the analysis will target the three major ocean basins as a demonstration while using the global surface wind stress shown in Fig. 7 as the consistent comparison point for each ocean basin – except in more detail starting with Fig. 8.

### 4.4 Regime changes in surface atmospheric – oceanic conditions and corresponding changes in fish landings

The goal of this section is to demonstrate a connection between the surface atmospheric–oceanic condition and the marine ecosystems over the Pacific, Atlantic and Indian Oceans as represented by the change in fish landings. The change in fish landings is thought to be caused by the change in large to small scale surface wind structures which cause greater change of marine ecosystems and influences multiple key factors related to fish habitat such as upwelling/downwelling (typically related to feeding) and sea surface temperature advection (related to ecosystem stress and survivability). Each ocean basin will be shown with its wind streamline analysis, and its kinetic energy (KE) wind amplitude analysis compared with their associated normalized fish landings.

### 4.5 Pacific Basin

Figure 7 shows the atmospheric surface wind streamlines and the surface height field anomalies. These surface structures are associated with the stratospheric planetary wave that is dynamically coupled to the ocean surface wind stress through upper level atmospheric forcing. These initial figures are the basis for the ocean surface wind stress

discussion. The surface atmospheric height anomalies, representative of atmospheric low and high pressure circulations determine the surface wind stress (wind speed and direction) over the Pacific basin region and the Atlantic and Indian Ocean regions as well.

Looking specifically at the Pacific Ocean, Fig. 8 shows the detailed wind stress over the Pacific basin for each regime shift period. As one scans each regime shift periods, there is a noticeable difference in the complexity of the wind stress pattern in the streamline analysis. From the 1948–1956 period to the 1957–1964 and 1965–1977 periods the wind pattern complexity increases as indicated by more “wind stream swirls” and fewer large sweeping wind flow areas. The complexity of the wind stress pattern has a direct impact on ocean ecosystems.

To demonstrate the increase in complexity that is more rigorous than a simple visual review, a wavenumber analysis of the wind kinetic energy (KE) over each ocean domain shows the change in the wind stress wavenumber spectrum. Figure 9 shows the wind stress through the kinetic energy anomalies characterized by the change in the *local wave spectrum* over the Pacific Ocean basin (left panels). Figure 9 compares the kinetic energy surface stress wavenumber analysis (left panels) with the normalized fish landings by species (right panels) – where decreased (blue bars) and increased (red bars) fish landings can be compared for each sub-region/fish group for each abrupt shift for the Pacific Ocean.

The greater the amplitude of the strongest wavenumber (wavenumber 1), the larger and more “smooth” the wind field structures in each period even considering the greater structure is visible in Fig. 8 (compare with the streamline analysis in Fig. 7). The greater the amplitude of the higher wavenumbers (wavenumbers 2 thru 6) in the KE wind anomaly spectrum, the greater the number of small scale disturbances in the ocean wind stress. The wind flow in the upper left panel of Fig. 8a was dominated by wavenumber one (one tall red bar and lower bars for wavenumbers 2 thru 6) which shows a few large “swirls” with a number of relatively long connecting flows. Similar large scale relatively smooth wind flows can be found in the periods of 1950–1956,

**Global marine fish landings and atmospheric planetary wave forcing**

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1978–1988, and 1989–1998 (Fig. 8a, d, e). The strong local wavenumber one influence can be found in the kinetic energy analysis (left panel series) in Fig. 9a, d, e; the black circles rest on top of the red bar with the dominant wavenumber for each regime shift period. Similarly, the periods with the most fish species having positive normalized fish landings are the same periods with strong KE wavenumber one influence. In contrast, small scale vortex flow created with a reduction in wavenumber one amplitude and/or an increase of the smaller scale, high wavenumber KE features (wavenumbers 2 thru 6) in the Pacific ecosystem can be found in periods 1957–1964, 1965–1977 and 1999–2005 (Fig. 8b, c, f). These periods are associated with fewer fish species showing positive normalized fish landings (red) and greater numbers of fish species showing negative normalized fish landings (blue).

Based on the FAO sub-regions and groups of fish monitored in the Pacific ecosystem, the fish landings were normalized and presented on the right hand series of graphs in Fig. 9. The number on the axis represents the fish species and region identified per Table A1 in the Appendix. Colored bars are drawn to help highlight the impact of the abrupt shifts on the number of fish species affected in a particular ocean basin. Positive values (red bars) show the fish species with improved catch/landing data and those with blue bars have decreased fish landing/catch data. During relatively large scale stable surface wind field periods (with kinetic energy anomalies in the longest local ocean basin wave – i.e. strong wavenumber one influence), the number of fish species with increased normalized landings is greater than during relatively small scale wind structure periods (with a greater dependence on higher wavenumbers indicating increased small scale wind structures). In Fig. 9, the wavenumber with the highest amplitude kinetic energy is marked by a black circle on top of the wavenumber bar. The bars with the black circles on top tracks with the change in the number of fish species with the greatest fish landing improvements and inversely with the number of fish species with reduced landings.

Why should this be true? The ocean consists of multiple ecosystems, albeit fluid ones. When the wind pattern is influenced by large and relatively stable forcing, it

**Global marine fish landings and atmospheric planetary wave forcing**

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



creates the opportunity for large scale features (ecosystems) where the different fish species can thrive with less disruption than during periods where higher wavenumber wind forcing is dominant. While there are a number of factors that can affect the viability of an ecosystem, the impacts of sea surface temperature (SST) change, spawning temperature impacts, growth of algae for feeding and upwelling/downwelling are likely to be among the strongest, and all can be impacted by the wind pattern. This conclusion presumes the fish landing data accurately reflect changes in fish habitat and do not reflect changes in policy for fish catch or other artificial constraints that would change the conclusions of the analysis (Daw et al., 2009).

When the wind stress pattern is dominated by smaller scale features (higher KE wave numbers), it disrupts the ocean ecosystems creating more discontinuous and generally less favorable conditions across the ocean basin. This adds stress to each fish species seeking an environment in which it can thrive and is thought to affect their environmental stress and survivability. As a consequence, when the ocean wind stress is dominated by higher wavenumber kinetic energy anomalies, the general fish populations do less well over the entire ocean basin which is broken into multiple, distributed and more stressful living conditions. Which fish do better and worse is highly dependent on the needs of each fish species and is a convolution of the changes induced by the wind stress, the geographic ocean locations most affected, and the resulting environmental impacts (upwelling/downwelling, SST change, algal growth change, etc.).

In general, it appears the atmospheric circulation starting at stratospheric levels is vertically communicated to the earth's surface and impacts the ocean circulation. In this analysis, the mid to high latitude atmospheric wave and wind patterns are demonstrated as the drivers for complex ocean basin wind stress change that results in the fish landing patterns shown in this research. The global atmospheric wave pattern changes across the multiple regime shift boundaries appear responsible for the significant changes in the surface ocean corresponding to more and less complex surface wind stress patterns with the resulting change in the number of fish species doing better or worse. Whether this is due to actual fish populations changes, the migration of

fish to regions more conducive for survival (less stressful conditions), or other factors is not completely clear. However, the driving physical factors analyzed in this research are among those known to create changes in fish species abundance. More detailed analyses of individual physical changes and their impacts on specific fish species are an area for future research.

The net result of the Pacific Ocean analysis is that ocean wind stress forcing with large wind stress features (reflecting KE wavenumber one) produces greater numbers of fish species with improved fish catch/landings. Conversely, the regime shift periods with higher wavenumber wind stress impacts (more smaller scale wind stress features) leads to diminished numbers of fish species doing well (or more fish species with lower catch and landing numbers). The next step is to determine whether this holds true for the Atlantic and Indian Oceans.

#### 4.6 Atlantic Basin

Similar to the smooth (wavenumber one dominant) and complex (higher wavenumber dominant) wind stress patterns in the Pacific Ocean, the wind stress pattern in the Atlantic Ocean (Fig. 10) shows a similar close relationship with the normalized fish species landings (Fig. 11), even though the Atlantic Ocean basin has a number of characteristics dissimilar to the Pacific Ocean. The Atlantic Ocean is significantly narrower than the Pacific Ocean and the wind stress has fewer large fetch patterns like those seen in the Pacific Ocean. However, when the large scale wavenumber one wind stress anomaly pattern (left panels in Fig. 11) dominates in a regime period, the fish landings are positively impacted. The wavenumber one dominant wind stress pattern in the Atlantic ocean occurs in the periods of 1965–1977 and 1978–1988 (left panels in Fig. 11c and d) corresponding with more positive normalized fish landings in the same periods (right panels in Fig. 11c and d). The opposite patterns can be found in the other four periods (Fig. 11a, b, e, f) when higher KE wavenumber impacts (greater number of relatively small scale ocean surface wind stress features) are present.

## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Fish species change in the Atlantic Ocean is impacted by the changes in relatively smooth (large scale) and complex (small scale) wind stress change. When wavenumber one wind stress features dominate this region, more fish species do well. Conversely, when higher wavenumber wind features are present, fewer numbers of fish species do well (or conversely more fish species have lower fish catch).

## 4.7 Indian Basin

The Indian Ocean wind pattern is largely tropical, and the ocean is bounded on the north and west by continents. This creates circumstances different from the mid-latitude wind stress patterns in the Pacific and Atlantic oceans. However, the wind flow patterns over the Indian Ocean provide the same net tendency reinforcing the conclusion that the wind stress impacts the fish. Consequently, the wind stress pattern in the Indian Ocean (Fig. 12) shows a similar relationship with the fish landings as the Pacific and Atlantic Oceans which were related to wavenumber one wind stress patterns (Fig. 13). However, the Indian Ocean may be more dependent on the specific high wavenumber wind structures than the other two ocean basins. It appears the high wavenumber wind stress patterns coupled with a reduced interchange with the southern polar region may create larger impacts on the number of fish species with positive fish landings.

In the periods where KE wavenumber 2 and higher wavenumber wind stress patterns were dominant, more fish species showed fewer positive fish landings in the regime periods from 1957–1964 through 1989–1998 (panels in Fig. 13b–e). In looking at the wavenumber 2 dominant periods, the greatest impact occurred with wavenumber one at its lowest magnitude (1978–1988, note the change in scale for the figures in the left series of panels) and the north-south interchange with the southern polar region reduced (Fig. 12d). For the other wavenumber 2 wind stress dominant regime shifts, there is increased amplitude in wavenumbers 3 thru 6. During the four wavenumber 2 and higher dominant periods, the impact is addressed by both the reduction in wavenumber one features and generally increased wavenumber 3 thru 6 activity. The combination of wavenumber influence and cut-off polar flow reduces the number of

# Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



fish species doing well in the Indian Ocean with the 1957–1963, 1965–1977, 1978–1988 and 1989–1905 periods having the fewest fish species doing well as indicated by the normalized fish catch. This is contrasted with the periods of 1950–1956 and 1999–1905, where the wavenumber one dominant wind stress anomaly leads to substantially improved numbers of fish species doing well based on higher normalized fish landings. In these periods, large scale wind circulation patterns connecting with the mid-latitude and polar zones and are thought to create conditions more conducive for greater numbers of fish species thriving. Also, sustaining a higher wavenumber wind stress scenario appears to result in declining periods of positive fish landing anomalies. The periods with high wavenumber activity in the Indian Ocean tend to cut off the interchange of ocean flow with the south polar region and appears to worsen the impact of the high wavenumber wind stress conditions. Recovery of the number of normalized fish landings in the later three regime shift periods appears to also be a result of both the ocean wind stress and the increasing interchange with the south polar ocean region.

## 5 Discussion

Surface wind stress can create many changes in the ocean including zones of upwelling and downwelling which can impact feeding, the ocean temperatures and the availability of food. It can also affect the advection of warm or cold water, spawning temperatures, the creation of ocean fronts, etc. Many of these factors can affect an ocean ecosystem so the picture is not as simple as the analysis portrays since it does not identify individual factors affecting specific fish populations consistently, nor how the ecosystem improved to cause the improved fish landings. However, this analysis does satisfy Chavez' basic premise (2003) that the forcing mechanisms need to be global (hemispheric), simple, direct and couples with biological changes identified in the fish catch. In this analysis, the global atmospheric pattern is the significant driver of the ocean surface through wind stress. The atmospheric winds and circulation patterns

are the variables linked in this analysis to the normalized fish landings. The linkage is logical and explains how the synchrony of fish species changes in all the world's ocean basins could occur based on changes in the global atmospheric wave pattern that influences ocean systems.

The authors suggest that the global atmospheric wind stress forcing from the Earth's largest atmospheric waves is the driving mechanism. When the wind stress is more complex and more disruptive in the ocean basins, the number of fish species doing well decrease or conversely, the number of fish species doing poorly increase. This suggests that when an ocean basin has greater fine scale wind structure, it disrupts the ocean ecosystems and stresses the various fish species causing them to adapt, migrate or possibly suffer higher mortality due to stressful ocean habitat conditions. In terms of upwelling and downwelling alone, a case can be made that during low KE wavenumber dominant conditions, relatively large consistent areas with potential for easily increasing the ability of the fish to feed appear to develop. When high KE wavenumber winds develop, there are many smaller upwelling areas that could increase the stress in finding suitable ocean-assisted feeding areas, thereby reducing the size of the fish catch. The habitat stress will affect the ability of the fish to thrive. In response to the atmospheric wind stress forcing on the ocean surface, it is likely that multiple conditions change in an ocean ecosystem that contribute to the overall impacts on a complex food chain and other factors which may impact survival and overall stress. Spawning temperatures affect some species (Takasuka et al., 2008), while the wind impacts on upwelling and downwelling can affect food availability similar to the well known El Nino Southern Oscillation impacts off South America. Analogously, the ENSO changes in surface wind direction, the SST temperature and upwelling/downwelling impacts many fish species off South America and significantly reduces fish catch in the region. Fish species may be directly or indirectly impacted through the wind stress on ocean ecosystems through changes in the food chain, the ability to reproduce or may force the migration to more suitable ocean conditions. The fish catch changes according to how turbulent or disruptive the KE wind forcing flow

**Global marine fish landings and atmospheric planetary wave forcing**

A. M. Powell Jr. and J. Xu

Title Page

Abstract	Introduction
Conclusions	References
Tables	Figures

◀ ◻ ▶

◀ ◻ ▶

Back	Close
------	-------

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper



## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



is. There are likely multiple factors affecting the fish catch. However, the global wind forcing derived from the decadal global (hemispheric) weather pattern plays a dominant role in sub-regional/fish species groups that result in increased or decreased fish catch. The change in the wind stress conditions impact about 15–25 % of the FAO sub-region/fish species groups across the identified decadal regime shifts.

## 6 Conclusions

Based on the analysis of the global wave pattern forcing, the consequential changes in ocean surface wind stress and the subsequent impacts on fish catch were investigated. The global wave pattern structure (KE wavenumber influence) and its amplitude are likely the key forcing mechanism that causes near synchronous impacts in the world's oceans and subsequently impact ocean ecosystems resulting in changes in fish catch.

As extensions to previous research, the atmospheric planetary wave processes were investigated as a likely global forcing in the six regime shift periods (1948–1956, 1957–1964, 1965–1977, 1978–1988, 1989–1998 and 1999–2005) in both the atmospheric and fish catch data in the most recent 59 year period (1950–2008). The multiple regime shifts across 1956–1957, 1964–1965, 1977–1978, 1988–1989, and 1998–1999 appear to adjudicate the wave amplitude and wind stress coupling mechanism. The five regime shift and six period analysis has clearly identified a mechanism where stratospheric planetary wave amplitudes may cause changes in surface atmospheric wind conditions. The decadal wave amplitude changes modify the winds through large contiguous systems which transfer energy between the stratosphere, troposphere and the ocean surface via height field anomalies (indicative of vertical atmospheric processes) which extend throughout the atmosphere.

The surface atmospheric conditions show a significant consistent mechanism as responsible for the fishery regime shifts in the catch/landing numbers of the three major ocean basins. In all three of the Earth's major ocean basins (Pacific Ocean, Atlantic Ocean, and Indian Oceans), the large scale atmospheric circulation's kinetic energy

## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



wind stress anomaly is best associated with the local wavenumber one surface wind stress which results in increases in fish landings. Higher KE wavenumber wind stress is associated with decreases in fish landings. It is thought the larger scale ocean features lead to ocean ecosystem conditions conducive to most fish species thriving through reduced (improved) habitat stress. Conversely, when the wind stress anomalies have many smaller scale wind features (increased habitat stress), it is disruptive to the ocean ecosystems creating stress on the fish species to find more acceptable ocean conditions. The stress of adaptation, migration or general survivability are consequences of the many ocean ecosystem changes where smaller scale wind stresses can impact the ability of an ocean fish species to thrive as indicated by increased fish landings.

The decadal regime shift linkage between global marine fish landings and atmospheric planetary wave forcing was demonstrated. The link between atmospheric waves that propagate to the Earth's surface and the ocean wind stress was shown. Similarly, the linkage between the ocean surface wind stress and fish landings was portrayed in graphical form. The decadal global connection between the atmospheric wind forcing and changes in fish catch in the world's three major ocean basins verifies the Chavez hypothesis that the synchronous changes in fish populations are due to a consistent mechanism that is simple, direct and operates similarly in all the world's oceans. In addition, the analysis showed the sub-region/fish species changes associated with the decadal regime shifts are on the order of 15 to 25 %. This is a significant finding that helps quantify the impacts associated with decadal climate and regime shift change.

*Acknowledgements.* The NCEP/NCAR monthly reanalysis data were obtained from NOAA/CDC web site. The authors would like to thank these agencies for providing the data. Special thanks to FAO for the fisheries datasets that were provided.

This work was supported by the National Oceanic and Atmospheric Administration (NOAA), National Environmental Satellite, Data and Information Service (NESDIS), Center for Satellite Applications and Research (STAR).

The views, opinions, and findings contained in this publication are those of the authors and should not be considered an official NOAA or US Government position, policy, or decision.

## References

- Beamish, R. J., Benson, A. J., Sweeting, R. M., and Neville, C. M.: Regimes and the history of the major fisheries off Canada's west coast, *Prog. Oceanogr.*, 60, 355–385, 2004.
- Chavez, F. P., Ryan, J., Lluch-Cota, S. E., and Niquen, M. C.: From anchovies to sardines and back: multidecadal change in the Pacific Ocean, *Science*, 299, 217–221, 2003.
- Chelton, D. B., Esbensen, S. K., Schlax, M. G., Chelton, D. B., Esbensen, S. K., Schlax, M. G., Thum, N., Freilich, M. H., Wentz, F. J., Gentemann, C. L., McPhaden, M. J., and Schopf, P. S.: Observations of coupling+ between surface wind stress and sea surface temperature in the eastern tropical Pacific, *J. Climate*, 14, 1479–1497, 2001.
- Daw, T., Adger, W. N., Brown, K., and Badjeck, M. C.: Climate change and capture fisheries: potential impacts, adaptation and mitigation, in: *Climate Change Implications for Fisheries and Aquaculture: Overview of Current Scientific Knowledge*, edited by: Cochrane, K., De Young, C., Soto, D., and Bahri, T., *FAO Fisheries and Aquaculture Technical Paper*, No. 530, FAO, Rome, 107–150, 2009.
- DeYoung, B., Harris, R., Alheit, J., Beaugrand, G., Mantua, N., and Shannon, L.: Detecting regime shifts in the ocean: data considerations, *Prog. Oceanogr.*, 60, 143–164, 2004.
- Ebbesmeyer, C. C., Dr. Cayan, McLain, D. R., Nichols, F. H., Peterson, D. H., and Redmond, K. T.: 1976 step in the Pacific climate: forty environmental changes between 1968–1975 and 1977–1984, in: *Proceedings of the 7th Annual Pacific Climate (PACLIM) Workshop*, Betancourt, J. L. and Tharp, V. L., April 1990. California Department of Water Resources. Interagency Ecological Study Program Technical Report 26, 115–126, 1991.
- Froese, R., Zeller, D., Kleisner, K., and Pauly, D.: What catch data can tell us about the status of global fisheries, *Mar. Biol.*, 159, 1283–1292, doi:10.1007/s00227-012-1909-6, 2012.
- Hátún, H., Payne, M. R., Beaugrand, G., Reid, P. C., Sandø, A. B., Drange, H., Hansen, B., Jacobsen, J. A., and Bloch, D.: Large bio-geographical shifts in the north-eastern Atlantic Ocean: from the subpolar gyre, via plankton, to blue whiting and pilot whales, *Prog. Oceanogr.*, 80, 149–162, 2009.
- Hilborn, R. and Branch, T. A.: Does catch reflect abundance?, *Nature*, 494, 303–306, 2013.
- King, J. R.: Report of study group on fisheries and ecosystem responses to recent regime shifts, *PICES Scientific Report No. 28*, Sidney, BC, Canada, Marine Science Organization (PICES), 1–168, 2005.

## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Kodera, K.: Solar cycle modulation of the North Atlantic Oscillation: implications in the spatial structure of the NAO, *Geophys. Res. Lett.*, 29, 1281, doi:10.1029/2001GL014557, 2002.
- Lehodey, P., Alheit, J., Barange, M., Baumgartner, T., Beaugrand, G., Drinkwater, K., Fromentin, J.-M., Hare, S. R., Ottersen, G., Perry, R. I., Roy, C., van der Lingen, C. D., and Werner, F.: Climate variability, fish, and fisheries, *J. Climate*, 19, 1009–1029, 2006.
- Mantua, N. J., Hare, S. R., Zhang, Y., Wallace, J. M., and Francis, R. C.: A Pacific interdecadal climate oscillation with impacts on salmon production, *B. Am. Meteorol. Soc.*, 78, 1069–1079, 1997.
- Matsuno, T.: Vertical propagation of stationary planetary waves in the winter Northern Hemisphere, *J. Atmos. Sci.*, 27, 871–883, 1970.
- Matthes, K., Kuroda, Y., Kodera, K., and Langematz, U.: Transfer of the solar signal from the stratosphere to the troposphere: northern winter, *J. Geophys. Res.*, 111, D06108, doi:10.1029/2005JD006283, 2006.
- Minobe, S.: A 50–70 year climatic oscillation over the North Pacific and North America, *Geophys. Res. Lett.*, 24, 683–686, 1997.
- Minobe, S.: Resonance in bidecadal and pentadecadal climate oscillation over the North Pacific: role in climatic regime shifts, *Geophys. Res. Lett.*, 26, 855–858, 1999.
- Nitta, T. and Yamada, S.: Recent warming of tropical sea surface temperature and its relationship to the Northern Hemisphere circulation, *J. Meteorol. Soc. Jpn.*, 67, 375–383, 1989.
- Overland, J., Rodionov, S., Minobe, S., and Bond, N.: North Pacific Regime shifts: definitions, issues and recent transitions, *Prog. Oceanogr.*, 77, 92–102, 2008.
- Peterson, W. T., Hooff, R. C., Morgan, C. A., Hunter, K., Casillas, E., and Ferguson, J. W.: Ocean Conditions and Salmon Survival in the Northern California Current, Web Document, available at: <http://www.nwfsc.noaa.gov/research/divisions/fed/ecosysrep.pdf> (last access: 14 December 2013), 2006.
- Powell, A. M. and Xu, J.: Abrupt climate regime shifts, their potential forcing and fisheries impacts, *Atmos. Clim. Sci.*, 4, 33–47, 2011a.
- Powell, A. M. and Xu, J.: Possible solar forcing of interannual and decadal stratospheric planetary wave variability in the Northern Hemisphere: an observational study, *J. Atmos. Sol.-Terr. Phys.*, 73, 825–838, 2011b.
- Powell, A. M. and Xu, J.: The 1977 global regime shift: a discussion of its dynamics and impacts in the Eastern Pacific ecosystem, *Atmos.-Ocean*, 50, 421–436, doi:10.1080/07055900.2012.716023, 2012.

# ESDD

5, 945–989, 2014

## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

- Powell, A. M. and Xu, J.: Regime shifts of global marine ecosystem with FAO fishery landing and possible physical conditions, *Int. J. Ecosyst.*, 3, 95–105, 2013.
- Roy, C. and Reason, C.: ENSO related modulation of coastal upwelling in the eastern Atlantic, *Prog. Oceanogr.*, 49, 245–255, 2001.
- 5 Sugimoto, T., Kimura, S., and Tadokoro, K.: Impact of El Niño events and climate regime shift on living resources in the western North Pacific, *Prog. Oceanogr.*, 49, 113–127, 2001.
- Takasuka, A., Oozeki, Y., Kubota, H., and Lluch-Cota, S. E.: Contrasting spawning temperature optima: why are anchovy and sardine regime shifts synchronous across the North Pacific, *Prog. Oceanogr.*, 77, 225–232, 2008.
- 10 Watson, R. and Pauly, D.: Systematic distortions in world fisheries catch trends, *Nature*, 414, 534–536, 2001.
- Zhang, Y., Wallace, J. M., and Battisti, D. S.: ENSO-like interdecadal variability: 1900–1993, *J. Climate*, 10, 1004–1020, 1997.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

**Table A1.** Summary of FAO Fish Landing Codes by Nine FAO Species Groups and Fourteen locations.

Acronyms	Species	PNW(1)	PNE(2)	PWC(3)	PEC(4)	PSW(5)	PSE(6)	ANW(1)	ANE(2)	AWC(3)	AEC(4)	ASW(5)	ASE(6)	IOW(1)	IOE(2)
CHH(1)	Codes,hakes, haddocks	P11	P21	P31	P41	P51	P61	A11	A21	A31	A41	A51	A61	I11	I21
FHS(2)	Flounders,halibut, soles	P12	P22	P32	P42	P52	P62	A12	A22	A32	A42	A52	A62	I12	I22
HAS(3)	Herrings,anchovies, sardines	P13	P23	P33	P43	P53	P63	A13	A23	A33	A43	A53	A63	I13	I23
MFNI(4)	Marine fishes not identified	P14	P24	P34	P44	P54	P64	A14	A24	A34	A44	A54	A64	I14	I24
MCF(5)	Miscellaneous coastal fishes	P15	P25	P35	P45	P55	P65	A15	A25	A35	A45	A55	A65	I15	I25
MDF(6)	Miscellaneous demersal fishes	P16	P26	P36	P46	P56	P66	A16	A26	A36	A46	A56	A66	I16	I26
MPF(7)	Miscellaneous pelagic fishes	P17	P27	P37	P47	P57	P67	A17	A27	A37	A47	A57	A67	I17	I27
SRC(8)	Sharks, rays, chimaeras	P18	P28	P38	P48	P58	P68	A18	A28	A38	A48	A58	A68	I18	I28
TBB(9)	Tunas,bonitos, billfishes	P19	P29	P39	P49	P59	P69	A19	A29	A39	A49	A59	A69	I19	I29

Each region and species category (defined in Fig. 1) is identified by Pij, Aij, or Iij for Pacific (P), Atlantic (A) and Indian Oceans (I) with the first number being (i) the ocean region (i = 1, 2, ... 6) and the second number (j) the FAO species category (j = 1, 2, 3, ... 9). Definition example: A23 indicates the Atlantic Northeast (2) and Species Category of Herrings, Anchovies, and Sardines (3). Regions are defined geographically in Fig. 1. The codes shown in this appendix are keys to the horizontal axes in Figs. 9, 11, and 13.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



**Table A2.** Listed years of fish landing regime shift occurrence by Region and Species consolidated by decade from 1950–2008.

10 year	1951–1960	1961–1970	1971–1980	1981–1990	1991–2000	10 year	1951–1960	1961–1970	1971–1980	1981–1990	1991–2000
PNW-CHH	56		72 and 78	88	99	ANW-CHH	60	70	76	81	99
PNW-FHS	59		77	88	98	ANW-FHS		65	74	88	94
PNW-HAS	59		76		92	ANW-HAS		68	74	87	
PNW-MFNI	55	65	74	88		ANW-MFNI	55		76		
PNW-MCF		65				ANW-MCF		61	76	89	96
PNW-MDF	55		76	86	94	ANW-MDF	58		76	86	94
PNW-MPF	58				98	ANW-MPF	57		73 and 78	85	92
PNW-SRC		64	76	88	99	ANW-SRC	55		77	85	
PNW-TBB	57					ANW-TBB					94
PWC-CHH	60			86	91	ANE-CHH		63		89	
PWC-FHS				84	99	ANE-FHS		68		85	92
PWC-HAS			75	84	99	ANE-HAS		68	74	85	92
PWC-MFNI	55		73	84	99	ANE-MFNI	56	69	77	85	94
PWC-MCF			74	83		ANE-MCF		69	77		
PWC-MDF				87	97	ANE-MDF			77		
PWC-MPF				87	2000	ANE-MPF		66		89	
PWC-SRC	58	56		89	97	ANE-SRC	56	67	77	87	
PWC-TBB		66	79		94	ANE-TBB	57	63		84	97
PSW-CHH				81	91	AWC-CHH		61		86	2000
PSW-FHS		61	76	90		AWC-FHS		64	74	89	
PSW-HAS	57	70		89		AWC-HAS	59		78	88	
PSW-MFNI		63	74	86		AWC-MFNI	57			81	95
PSW-MCF	58	63	79	86	93	AWC-MCF	55	65		90	98
PSW-MDF	60	67	77	89		AWC-MDF	58	63	72	89	98
PSW-MPF	55	65	79		2000	AWC-MPF			78		
PSW-SRC	58	65		89	97	AWC-SRC	57	70		81	99
PSW-TBB	60			88		AWC-TBB	57	67		87	
PNE-CHH	55	61				AEC-CHH		65	78		
PNE-FHS		64	71 and 78			AEC-FHS	57		71	86	91
PNE-HAS	60		75	90		AEC-HAS	60	70	79	87	92
PNE-MFNI			74		99	AEC-MFNI	55	64			
PNE-MCF		67		90		AEC-MCF		63	72 and 77	88	
PNE-MDF	57	63	74		99	AEC-MDF	57	68			99
PNE-MPF		65	77		92	AEC-MPF	55	69			
PNE-SRC	59		78			AEC-SRC	57		71	81	97
PNE-TBB	55	70		85		AEC-TBB		68	74	90	
PEC-CHH		69		87		ASW-CHH	56		77		99
PEC-FHS	55		77		98	ASW-FHS		63		85	99
PEC-HAS		61	71		92	ASW-HAS	55	62	71	88	99
PEC-MFNI	56					ASW-MFNI	56	67		90	
PEC-MCF	60	70				ASW-MCF	56	68	76	88	
PEC-MDF	56	64	80			ASW-MDF	59		80	86	
PEC-MPF	56	62				ASW-MPF				87	
PEC-SRC	55	68	80			ASW-SRC	59	63			97
PEC-TBB		64		83		ASW-TBB		65	80		
PSE-CHH	58			87		ASE-CHH		66	79	90	
PSE-FHS		66			96	ASE-FHS			75	89	
PSE-HAS		61	72	85		ASE-HAS	56		80		2000
PSE-MFNI					92	ASE-MFNI					
PSE-MCF	57	64				ASE-MCF		66	80	89	98
PSE-MDF	55			86		ASE-MDF	59		77	90	
PSE-MPF	60		77	88	98	ASE-MPF		64		84	2000
PSE-SRC		66				ASE-SRC				84	99
PSE-TBB		64	73	88		ASE-TBB		62			

**Global marine fish landings and atmospheric planetary wave forcing**

A. M. Powell Jr. and J. Xu

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




**Table A2.** Continued.

10 year	1951–1960	1961–1970	1971–1980	1981–1990	1991–2000
IOW-CHH			75		2000
IOW-FHS				90	
IOW-HAS		64			2000
IOW-MFNI		64	78	84	97
IOW-MCF	58	67	80		91
IOW-MDF	56	61	74		2000
IOW-MPF		70	80	89	91
IOW-SRC		66		83	92
IOW-TBB	57		75	85	98
IOE-CHH	59		78	88	95
IOE-FHS	55	69	80	81	93 and 99
IOE-HAS	57	68	77	86	2000
IOE-MFNI		64		89	96
IOE-MCF	58		79	89	
IOE-MDF	60			86	
IOE-MPF	57	70		83	2000
IOE-SRC	55				93
IOE-TBB	58	70			96

The listed number is the year of the fish landing regime shift occurrence during each decade based on the 11 year running student  $t$  test for the normalized time series of one fish species from 1950–2008 at the 90 % significance confidence level. For example, the fish species group CHH over the Northwest Pacific (PNW-CHH) in Fig. 3b shows the  $t$  test value exceeds the statistical significance test at the 95 % confidence level in the years 1956, 1972, 1978, 1988 and 1999, which corresponds with 56, 72 and 78, 88, 99 in the table at the row for PNW-CHH.





**Figure 1.** Lists the Food and Agriculture Organization’s (FAO) 14 sub-ocean regions for marine fish species landings in the Pacific, Atlantic and Indian Oceans and the 9 groups of marine species tabulated.

**(a)** *Ocean Sub-regions on grey shade map:* Pacific, Northwest (PNW), Pacific, Western Central (PWC), Pacific, Southwest (PSW), Pacific, Northeast (PNE), Pacific, Eastern Central (PEC), Pacific, Southeast (PSE), Atlantic, Northwest (ANW), Atlantic, Western Central (AWC), Atlantic, Southwest (ASW); Atlantic, Northeast (ANE), Atlantic, Eastern Central (AEC); Atlantic, Southeast(ASE). Indian Ocean, Eastern (IOE); Indian Ocean, Western (IOW)

**(b)** *Nine FAO species groups* (1) Cod, Hakes, and Haddocks (CHH), (2) Flounders, Halibut and Soles (FHS), (3) Herrings, Anchovies and Sardines (HAS), (4) Marine Fishes Not Identified (MFNI), (5) Miscellaneous Coastal Fishes (MCF), (6) Miscellaneous Dermersal Fishes (MDF), (7) Miscellaneous Pelagic Fishes (MPF), (8) Sharks, Rays and Chimaeras (SRC), (9) Tunas, Bonitos and Billfishes (TRB).

The codes used in the graphics of this paper represent the combination of the FAO geographical ocean sub-regions (shown in this figure) and the fish species categories are listed above (for example PNW-CHH is the Pacific Northwest regions’ Cod, Hakes, and Haddocks fish species group). The shortened combined codes (ie. P21) used in the paper as the x-axis identifiers are described in Table A1 of the Appendix.

**Global marine fish landings and atmospheric planetary wave forcing**


A. M. Powell Jr. and J. Xu

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu



**Figure 2.** Normalized planetary wave amplitude index (PWAi) from 1948–2008 for 55–75° N, and 70–20 hPa with identified regime shift periods and approximate shift dates. Regime shifts are identified at 1956–1957, 1964–1965, 1977–1978, 1988–1989 and 1998–1999.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu



**Figure 3.** The detrended time series of the normalized fishery landing (NFL) with black curves over the three ocean basin (scaled by right y-axis), the 11 year running  $t$  test with red bar (scaled by left y-axis). The number indicates the year when the regime shift happened, based on passing the significance test at the 90 % confidence level. Each panel in this figure is the sample selected from the total 14 sub-regions and 9 species groups listed in Table 1A. Pacific Ocean: **(a)** PWC-MDF; **(b)** PNW-CHH; **(c)** PNW-SRC, **(d)** PSW-SRC ; **(e)** PNE-MDF; **(f)** PNE-FHS; Atlantic Ocean: **(g)** ANW-MPF; **(h)** ANE-MPF; **(i)** AWC-FHS; **(j)** AEC-MCF; Indian Ocean: **(k)** IOW-MFNI; **(l)** IOE-HAS. The distributions of the other fish species over the other sub-regions are not shown here, but all regime shift years are listed in Table A2.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu




**Figure 4.** The rate of fish landing regime shift occurrence (blue-red bar; left y-axis) changes with time based on the statistical results of Table A2 in Appendix and student *t* test value (black curves line, right y-axis) The red dashed line indicates the threshold of statistical significance *t* test at the 90 % confidence level (right y-axis). The red bar indicates the year when the regime shift occurred in each 10 year. **(a)** Pacific Ocean, **(b)** Atlantic Ocean, and **(c)** Indian Ocean.

The rate calculation is according to total counts of regime shift occurrence during two continuous years, the approach is made as follows. For example, during the 10 year from 1971–1980 over the Pacific Ocean in Table A2, there are 7 sub-region/fish species showing the regime shift at year 1977 and 3 sub-region/fish species at year 1978, so there are 10 sub-region/fish species showing the shift in 1977–1978. The total sub-region/fish species is 54 (6 sub-regions  $\times$  9 species = 54) over the Pacific Ocean, so that the rate of regime shift occurrence at 77–78 is  $10/54 = 18.5\%$ . The others were obtained in the same way.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu



**Figure 5.** Reconstructed planetary wave pattern using geopotential height for wavenumbers 1–6 at 50 hPa in the NCEP/NAR reanalysis. Wave anomalies in the following regime shift periods **(a)** 1948–1956; **(b)** 1957–1964; **(c)** 1965–1977; **(d)** 1978–1988, **(e)** 1989–1998, **(f)** 1999–2005. Units: gpm. Shaded areas indicate positive anomalies. Block boxes shown positions of net large scale features associated with the atmospheric regime shifts.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu



**Figure 6.** Reconstructed global wave amplitude anomalies from the height fields for planetary wavenumbers 1–6 with altitude (pressure level). The vertical panels arranged from left to right are the periods in 1948–1957; 1958–1964; 1965–1977; 1978–1988; 1989–1998; and 1999–2005, respectively. (Negative anomalies: blue-purple shading; positive anomalies: green-yellow-red). The vertical dashed line and “x” give an example during each decadal period representing the connection of the planetary wave active centers between stratosphere, troposphere and earth’s surface (1000 hPa).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu



**Figure 7.** Global regime shift period reconstructed height (shaded) and wind field anomalies (streamlines) at 1000 hPa model level for planetary wave numbers 1 thru 6. Shaded areas indicate height anomalies exceeding the significance tests at the 90 % confidence level. (Negative anomaly: blue-purple shading; positive anomaly: green-yellow-red). **(a)** 1948–1956; **(b)** 1957–1964; **(c)** 1965–1977; **(d)** 1978–1988, **(e)** 1989–1998, **(f)** 1999–2005. Note, the shown domain is from 60° S to 60° N.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu



**Figure 8.** *Pacific Ocean* reconstructed height (shaded) and wind field anomaly over the Pacific ocean at 1000 hPa model level for the planetary wave number 1–6. Shaded area indicate the height anomaly exceeding the significant tests at the 90 % confidence level. (Negative anomaly: blue-purple shading; positive anomaly: green-yellow-red). **(a)** 1948–1956; **(b)** 1957–1964; **(c)** 1965–1977; **(d)** 1978–1988, **(e)** 1989–1998, **(f)** 1999–2005. Coastline in orange. Note, the domain shown is from 60° S to 60° N to highlight regions of wind driven forcing over the Pacific Ocean. When the streamlines converge, the water is generally rising or sinking (upwelling and downwelling).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion





## Global marine fish landings and atmospheric planetary wave forcing


A. M. Powell Jr. and J. Xu



**Figure 9.** *Pacific Ocean.* Left panel indicates the wave kinetic energy anomaly  $1/2(u^2(n)+v^2(n))$  with wave number ( $x$ -axis). Right panel indicates the average normalized fish landings for the Pacific Ocean for each species in the periods coincident with abrupt climate regime shifts. **(a)** 1950–1956, **(b)** 1957–1964, **(c)** 1965–1977, **(d)** 1978–1988, **(e)** 1989–1998, **(f)** 1999–2005. The  $x$ -axis code number indicates both the geographical ocean sub-region (shown in Fig. 1) and the FAO fish species category. The codes are summarized in Table A1.

## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu



**Figure 10.** *Atlantic Ocean* reconstructed height (shaded) and wind field anomaly over the Atlantic ocean at 1000 hPa model level for the planetary wave number 1–6. Shaded area indicate the height anomaly exceeding the significant tests at the 90 % confidence level. (Negative anomaly: blue-purple shading; positive anomaly: green-yellow-red). **(a)** 1948–1956; **(b)** 1957–1964; **(c)** 1965–1977; **(d)** 1978–1988, **(e)** 1989–1998, **(f)** 1999–2005. Coastline in orange.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Global marine fish landings and atmospheric planetary wave forcing


A. M. Powell Jr. and J. Xu



**Figure 11.** *Atlantic Ocean.* Left panel indicates the wave kinetic energy anomaly  $1/2(u^2(n) + v^2(n))$  with wave number (x-axis). Right panel indicates the average normalized fish landings for the Pacific Ocean for each species in the periods coincident with abrupt climate regime shifts. **(a)** 1950–1956, **(b)** 1957–1964, **(c)** 1965–1977, **(d)** 1978–1988, **(e)** 1989–1998, **(f)** 1999–2005. The x-axis code number indicates both geographical ocean sub-region (shown in Fig. 1) and the FAO fish species category, the codes are summarized in Table A1.

## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu



**Figure 12.** *Indian Ocean* reconstructed height (shaded) and wind field anomaly over the Indian ocean at 1000 hPa model level for the planetary wave number 1–6. Shaded area indicate the height anomaly exceeding the significant tests at the 90 % confidence level. (Negative anomaly: blue-purple shading; positive anomaly: green-yellow-red) **(a)** 1948–1956; **(b)** 1957–1964; **(c)** 1965–1977; **(d)** 1978–1988, **(e)** 1989–1998, **(f)** 1999–2005. Coastline in orange. Note the latitude range for the Indian Ocean.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Global marine fish landings and atmospheric planetary wave forcing

A. M. Powell Jr. and J. Xu



**Figure 13.** Indian Ocean. Left panel indicates the wave kinetic energy anomaly  $1/2(u^2(n) + v^2(n))$  with wave number (x-axis). Right panel indicates the average normalized fish landings for the Pacific Ocean for each species in the periods coincident with abrupt climate regime shifts. **(a)** 1950–1956, **(b)** 1957–1964, **(c)** 1965–1977, **(d)** 1978–1988, **(e)** 1989–1998, **(f)** 1999–2005. The x-axis code number indicates both geographical ocean sub-region (shown in Fig. 1) and the FAO fish species category. The codes are summarized in Table A1.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion