

Changes to the manuscript (mainly additions) are marked in blue. The main additions are the six appendices.

Global warming projections derived from an observation-based minimal model

K. Rypdal¹

¹Department of Mathematics and Statistics, UiT The Arctic University of Norway, Norway

Correspondence to: Kristoffer Rypdal
(kristoffer.rypdal@uit.no)

Abstract. A simple conceptual model for the global mean surface temperature (GMST) response to CO₂ emissions is presented and analysed. It consists of linear long-memory models for the GMST anomaly response ΔT to radiative forcing and atmospheric CO₂-concentration response ΔC to emission rate. The responses are connected by the standard logarithmic relation between CO₂ concentration and its radiative forcing. The model depends on two sensitivity parameters, α_T and α_C , and two “inertia parameters,” the memory exponents β_T and β_C . Based on observation data, and constrained by results from the Climate Model Intercomparison Project Phase 5 (CMIP5), the likely values and range of these parameters are estimated, and projections of future warming for the parameters in this range are computed for various idealised, but instructive, emission scenarios. It is concluded that delays in the initiation of an effective global emission reduction regime is the single most important factor that influences the magnitude of global warming over the next two centuries. The **highlight** of this study is the simplicity and transparency of the conceptual model, which makes it a useful tool for communicating the issue to non-climate scientists, students, policy-makers, and the general public.

1 Introduction

In spite of five comprehensive reports from the Intergovernment Panel on Climate Change (IPCC) the perception of the threat of global warming to society remains highly diverse among the general public, decision makers, and the **scientific community at large**. This is in stark contrast to the general opinion among those who define themselves as climate scientists, where some studies suggest that as much as 97 percent recognise human activity as a main driver of **a main driver of global warming over the last century** (??). What distinguishes the climate science community from other scientists is the strong reliance among climate scientists on complex Earth System Models (ESMs), that is, on Atmospheric-Ocean General Circulation Models (AOGCMs) coupled to models that include biogeochemistry and cryosphere dynamics. The general skepticism against this “model science” is not hard to understand. Models are complex beyond comprehension, different models are not independent but consist of many common modules, and parametrisations are empirical to an extent that makes it

legitimate to question whether models are “massaged” to fit observations. The important point here is not whether this perception of climate modelling is correct or fair, but that the skepticism exists, and in many cases cannot be discarded as irrational.

30 The latest IPCC report from Work Group I on the climate system (?) contains a summary for policy makers that describes findings from observations and model studies which many physical scientists find unconvincing, and is not very easy read for the general public. The unconvincing part is the above mentioned [excessive](#) reliance on complex computer models. Most scientists want to understand and to be convinced by simple fundamental principles matched against clear-cut observations.
35 Decision makers and the informed layman want to see simple, clear alternatives for the future; not a myriad of incomprehensible scenarios labelled by [acronyms that carry no meaning to non-experts](#).

From the Co-Chair of Work Group I a very readable and important paper on the “The Closing Door of Climate Targets” (?) was published alongside the IPCC AR5 report, intended to demonstrate that as mitigation is delayed, climate targets formulated in international agreements become
40 unattainable. The results were based on the physical assumption of a linear relationship between the cumulated carbon emissions and peak global warming in scenarios where the cumulative emission is bounded. This relationship, and the constant of proportionality, were justified empirically from numerical experiments performed on a large number of ESMs which incorporate the global carbon cycle (??). Some readers, however, will find it unsatisfactory that they have to “believe” the models
45 in order to accept the conclusion of the paper. As a former plasma physicist, who only relatively recently has taken up research in Earth-system dynamics and climate science, I am often confronted with question from former colleagues of the type: “For half a century we have tried to model the transport properties of a magnetically confined plasma for controlled thermonuclear fusion, and we still haven’t succeeded very well, even though the physical system is infinitely simpler than the
50 climate. Why do you think these horrendously complex climate models perform any better?”

A major motivation for the present paper is to find ways to communicate with, and gain support from, the scientists who ask such questions. I do this by deriving results similar to those obtained in ? in a more transparent manner, and without resorting to complex ESMs as the primary justification. The underlying assumptions are justified from observations, although supporting evidence
55 from AOGCMs are also discussed. The conceptual models of the temperature and atmospheric carbon response are linear and simple enough to be understood by anyone with some background in elementary calculus and ordinary differential equations. The scenarios explored are idealised and the results presented in figures that should be comprehensible for readers without training in mathematics or physical sciences.

60 Sect. 2 describes and justifies the conceptual model. Sect. 3 presents projections for atmospheric CO₂ concentration and GMST for some idealised CO₂ emission scenarios, one which is very close to the “business as usual” Representative Concentration Pathway 8.5 (RCP8.5) scenario employed by the IPCC, and others which represent systematic emission reduction initiated at different times in

the future. This section also discusses policy implications that may follow from these projections,
65 and i Sect. 4, summarises and concludes. Six appendices elaborate on the physical interpretation and
justification of the minimal model, and on some mathematical aspects that may appear as paradoxes.
This material is referred to appendices in order to avoid interruption of the logical flow that leads
to the main results. The Supplementary Material contains data files and a well-documented *Mathe-*
matica notebook with routines that allow readers to replicate and extend all results presented in the
70 paper.

2 The conceptual model

A closed model for the evolution of the global mean surface temperature (GMST) could consist
of (i) a model for the GMST-anomaly response $\Delta T(t)$ to radiative forcing $F(t)$, (ii), a model for
the evolution of $\Delta C(t)$, given the CO₂ emission history $R(t)$, and (iii) a well established consti-
75 tutive relation between $F(t)$ and $\Delta C(t)$. This paper proposes extremely simple, linear models for
the GMST-response (i) and the CO₂-concentration response (ii). Each depends on two parameters
characterising the strength and the inertia (memory) of the response, respectively. In order to keep
the model sufficiently simple for a reader to be able to trace the connection between driver and re-
sponse, and the effect of variation of model parameters, major simplifying assumptions are made.
80 One is to neglect all other radiative forcing than CO₂. Although the main reason for this is to main-
tain simplicity, it is justified by forcing estimates that conclude that the non-CO₂ contributions tend
to cancel over the industrial period (?). Other important simplifications are linearity and stationarity:

Linearity. Global temperature has been found to respond quite linearly to forcing in general cir-
culation models (?), and as long as the climate system is far from a major tipping point, this linearity
85 may also pertain to the response of atmospheric CO₂ content to emissions. The effect of space-time
non-linearity is important primarily on variability on smaller than global scale. On global scale the
response function has an approximate power-law form that makes the system respond by a scale-
invariant stochastic process to a white-noise driver. This scale-invariance is characterised by a spec-
tral exponent β which gives rise to a power-law tail in the response function $G(s) \sim s^{\beta/2-1}$, where s
90 is the time following an impulse in the forcing. The physical interpretation of such a response is that
the climate system consists of a number of different interacting sub-systems with different response
times. There will be a maximum response time and hence there will be a cut-off of the power-law tail
in the response function for s larger than this maximal time constant. The justification, interpretation
and implication of this picture is further discussed in the appendices.

Stationarity. The response functions are assumed to be translation invariant, i.e., $G(t, t') = G(t -$
95 $t')$. This means that the GMST and the CO₂ concentration respond the same way in a future climate
as they do now. For the GMST this is a reasonable assumption as long as the global general circula-
tion pattern remains the same, i.e., as long as the climate system does not encounter a major tipping

point. Examples of such tipping points are the glacial-interglacial transitions, or the Dansgaard-
 100 Oeschger events during the last ice age (?). During the present interglacial period, the Holocene,
 there was a similar tipping event about 8.2 kyr ago. These events are believed to be associated
 with sudden influx of freshwater into the Northern Atlantic from the North-American Laurentide
 ice sheet and associated changes in the overturning ocean circulation. A number of potential tipping
 elements have been identified associated with global warming in the present Holocene climate (?).
 105 Among these are complete disappearance of the Arctic sea ice, runaway melting of the Greenland
 and West-Antarctic ice sheets, a radical change in the Atlantic thermohaline ocean circulation and
 the El Niño Southern Oscillation, shifts of the Indian and the West-African monsoons, and dieback
 of the Amazon and the boreal forests. Transitions associated with tipping elements of these types
 can change significantly the global temperature response as well as the carbon cycle response. Even
 110 in the absence of tipping points the stationarity assumption may be particularly wrong for the CO₂
 concentration, where e.g., saturation effects in the ocean mixed layer and the land biosphere may
 reduce fluxes in a future climate. It also neglects the coupling between sea surface temperature and
 the CO₂ flux, which will reduce the flux into the ocean in a warmer climate. However, experiments
 with Carbon cycle models subject to sudden CO₂ injections into the atmosphere indicate that the re-
 115 sponse in the CO₂ concentration can be described by a power-law response function. This response
 is not stationary in the sense that it will be the same for a new Carbon release in a future climate,
 but it may give an adequate description of the response to the present global warming event. Further
 details are given in Sects. ??, ?? and the appendices.

2.1 The temperature response

120 The simplest physics-based model of the GMST-response is the zero-dimensional, linearised energy
 balance model (EBM);

$$\frac{d}{dt}\Delta T = -\frac{1}{\tau_T}\Delta T + \frac{S}{\tau_T}F. \quad (1)$$

Here τ_T is the time constant for relaxation of the temperature anomaly, and S is the climate sen-
 sitivity. The model is often denoted the Budyko-Sellers model and first proposed by ? and ?. A
 125 simple derivation can be found in ?, where it is also pointed out that it is impossible to find a single
 time constant that describes adequately the response to forcing on all time scales. The reduction to
 a linear model from the nonlinear EBM with the full Stefan-Boltzmann radiation law is found in
 Appendix ??. This model is not only used for reproducing the global temperature to known (de-
 terministic) forcing, but can also be formulated as a stochastic differential equation by introducing
 130 a noise component to the forcing $F(t)$, representing the stochastic energy flux from atmospheric
 weather systems to the ocean and land surface (?). The solution to this equation can be written as a

convolution integral

$$\Delta T(t) = \int_0^t G_T(t-t')F(t') dt', \quad (2)$$

with an exponential response function

$$135 \quad G_T(t) = (S/\tau_T) \exp(-t/\tau_T). \quad (3)$$

The temperature response to a purely stochastic forcing, i. e., $F(t)$ is represented as a Gaussian white noise, is an Ornstein-Uhlenbeck stochastic process. In discrete time, this corresponds to a first-order autoregressive (AR(1)) process. If Eq. (??) provides an adequate description, with $F(t)$ separated into a deterministic and a white-noise component, then the residual obtained after subtracting the
 140 deterministic response from the observed annual GMST record should be a realisation of an AR(1) process. The time constant and the climate sensitivity can be determined by a maximum-likelihood estimation, and in ? they were estimated to $\tau \approx 4.3$ yr, and $S \approx 0.32 \text{ Km}^2\text{W}^{-1}$. However, the sensitivity obtained is lower than obtained from climate models, the fast response to volcanic eruptions is higher than in the observed record, and the residual does not conform well with an AR(1) process. ?
 145 demonstrated that the residual is better described by a model for persistent, fractional Gaussian noise (fGn). Such a noise can be produced by Eq. (??) if the exponential response function is replaced by a power-law function

$$G_T(t) = \alpha_T t^{\beta_T/2-1}, \quad (4)$$

where the memory exponent β_T is in the interval $0 < \beta_T < 1$. It can be shown that this process has
 150 a power spectral density on the form $\sim f^{-\beta_T}$, where f is the frequency (?). Hence, $\beta_T = 0$ corresponds to white noise, while increasing β_T signifies increasing degree of memory (or persistence) in the process. In this response model it replaces the time constant τ_T of the simple EBM. The parameter α_T replaces the climate sensitivity S . In ? the magnitude of the parameters α_T and β_T were estimated from the instrumental GMST record, revealing rather strong persistence, $\beta_T \approx 0.75$. Similar
 155 values were also found in multiproxy data for the Northern Hemisphere, and in ? they were found in data from a number of millennium-long AOGCM simulations. The long power-law tail in the response function may be interpreted as an effect of thermal exchange between the surface (e.g., the ocean mixed layer) and other components of the climate system with higher heat capacity (e.g., the deep ocean). A two-layer ocean energy balance model yields for instance a response function with
 160 two exponentials with different time constant. In ? such a two-layer model was compared to transient simulations of AOGCMs following an abrupt increase in CO_2 forcing, and the two time constants estimated from these data were typically 1-2 yr and 1-2 centuries, respectively. In ? it was shown that a power-law response provides an even better fit to the tail of the transient AOGCM-solutions, but the memory exponent is lower ($\beta_T \approx 0.35$) than found from the residuals in observations and AOGCM

165 simulations with historical forcing. One way of reconciling these conflicting results is to assume
that the forcing noise is not white, but rather a persistent noise which gives a contribution to the β_T
observed in the residuals. Details are shown in Appendix ???. On the other hand, it will be pointed
out in Sect. 3 that the Computer Model Intercomparison Project Phase 5 (CMIP5) in the RCP8.5
CO₂ concentration scenario yields results consistent with $\beta_T = 0.75$. Since this implies some uncer-
170 tainty with respect to the correct value of β_T for the temperature response I shall in Sect. 3 present
projections for the values $\beta_T = 0.35$ and $\beta_T = 0.75$, assuming that β_T is likely within this interval.

The significance of the inertia, or long-range memory (LRM), in the temperature response for
GMST projections is illustrated in Fig. ???. Panel (a) shows the estimated GMST response to forcing
scenario consisting of the anthropogenic forcing in the period 1880 – 2010 as presented in ?, linearly
175 projected to 2200 AD with the same mean growth rate as the the RCP8.5 scenario in the period 2010
– 2100 AD (?), and is shown as the blue curve in Fig. ??b. The blue and red curves in Fig. ??a are the
responses according to the power-law response models with $\beta_T = 0.35$, and $\beta_T = 0.75$, respectively.
The projection for an instant response ($\tau_T \rightarrow 0$, leading to $\Delta T(t) \rightarrow S F(t)$) is also shown as the
limit of zero inertia. Also shown as a light blue curve is the instrumental GMST record as given by ?.
180 These projections have been obtained by computing the integral $\int_0^t \alpha_T (t - t')^{(\beta_T/2 - 1)} F(t') dt'$ with
the specified β_T and then estimating α_T by regressing to the observed GMST record for the period
1880 – 2010 AD. The climate sensitivity S for the instantaneous response has also been found by
regressing $S F(t)$ to the instrumental data, and is found to be $S \approx 0.48 \text{ Km}^2 \text{ W}^{-1}$, which corresponds
to 1.8 K for a doubling of CO₂ concentration. The rising warming projected for increasing β_T is
185 a manifestation of the thermal inertia in parts of the climate system with high heat capacity that
exchange heat with the surface, and makes the surface temperature respond more slowly. The higher
surface warming in the distant future due to this inertia is a manifestation of “the warming in the
pipeline” (??).

The blue forcing path shown in Fig. ??b is an idealised “business as usual” (BAU) scenario.
190 Beyond 2100 AD there is every reason to believe that there will be a saturation of the rising trend,
even in the absence of active mitigation policies. In the RCP8.5 this takes place gradually during
the 22nd and first half of the 23rd century. This figure also shows some idealised scenarios where
the BAU is modified by mitigation action. One possible type of action is the sudden reduction of
emission that will stabilise the forcing at the level at the time of action. In the real world such an
195 action from one year to another is not possible, but it may be considered an approximation of certain
annual reduction over a period of a decade. For instance, 40% emission reduction can be achieved
by annual emission reduction of 5% over a decade. In Fig. ??b forcing scenarios for this type of
mitigation action are illustrated for three different years of onset of the action; 2030, 2070, and 2110
AD. The year 2030 gives the world fifteen years to prepare the action. Year 2070 leaves the problem
200 to those who are born today, i.e., to the next generation. Year 2110 leaves it to unborn generations.

The GMST projections for these scenarios are shown in Fig. ??c,d for the lower and higher memory exponents β_T . Under the low-inertia assumption in the temperature response ($\beta_T = 0.35$) the unmitigated forcing scenario in Fig. ??a yields approximately two degree of warming every 40 yr throughout the 21st century, and even higher rate of warming in the 22nd century. After stabilisation
 205 of the atmospheric CO₂ concentration, the temperature will continue to rise about one degree Celcius by the year 2200 AD, independent on when this stabilisation takes place. This one degree of additional warming is the “warming in the pipeline.” Under the high-inertia assumption ($\beta_T = 0.75$) the warming rate is approximately 30% higher, and the warming in the pipeline is about a 100% higher. The high-inertia projection with mitigation action in 2110 AD is very close to the multimodel
 210 mean RCP8.5 projection (?), suggesting some consistency between this simple global temperature response model and the models employed by the IPCC in the CMIP5 project.

Fig. ??c,d suggest that the two-degree Celcius target is unlikely to be attained by rapid stabilisation of atmospheric CO₂ concentration, if this action is started later than 2030 AD. If radical action is postponed until the GMST has passed the two-degree limit, it is likely that the global temperature
 215 will exceed three degrees by 2100 AD, and if action is postponed till the end of this century our descendants may experience a world that is 5 – 8 degrees warmer than before industrialisation.

2.2 The atmospheric CO₂ response

The dominant driver of climate change throughout the 20th century and beyond is anthropogenic radiative forcing, and in the 21st century, CO₂ forcing is expected to be the main anthropogenic
 220 driver. However, while AOGCMs traditionally have been driven by prescribing the atmospheric CO₂ concentration, the policy relevant quantity is the CO₂ emission rate. The main factor that determines future CO₂ forcing in a given emission scenario is the rate at which CO₂ is washed out of the atmosphere. This is where the carbon-cycle models incorporated in the ESMs become important. The model uncertainty is high, but they suggest the existence of a hierarchy of time scales, just as
 225 we have found in the temperature response (?). This hierarchy is not immediately apparent from the instrumental data records, but there is some indirect evidence, as will be demonstrated below. However, let us first consider a primitive model with only one response time scale, analogous to the simple EBM given by Eq. (??) for the surface temperature. In this model we assume that the Carbon flux out of the atmosphere is proportional to the anomaly ΔC of atmospheric Carbon content relative
 230 to the preindustrial concentration C_0 . This assumption follows from a Taylor expansion to first order of the Carbon flux $I(\Delta C) = (1/\tau_c)\Delta C + \dots$ around the preindustrial equilibrium $I(C_0) = 0$. The primitive equation for this perturbation is then

$$\frac{d}{dt}\Delta C = -\frac{1}{\tau_C}\Delta C + R, \quad (5)$$

where τ_C is the time constant for relaxation of CO₂ concentration to the preindustrial equilibrium.
 235 A first-order estimate of τ_C can be made from the estimates of the global carbon budget (?). The

annual carbon emission in the period 1960 – 2010 grew almost linearly from 4 GtC/yr to 11 GtC/yr. We can solve Eq. (??) for this period with $R = [4 + (7/50)]t$ GtC/yr in terms of τ_C and the initial atmospheric Carbon inventory anomaly, ΔC_{1960} . The conversion factor from concentration in ppm to GtC in total Carbon content is 2.12 (?), which yields $\Delta C_{1960} = (315 - 280) \times 2.12 \approx 74$ GtC if we assume a CO₂ concentration of 315 ppm in 1960 and preindustrial concentration 280 ppm. The preindustrial Carbon content, corresponding to 280 ppm, was $C_0 \approx 594$ GtC. This solution reproduces very well the observed evolution of the atmospheric CO₂ content in this period if one chooses $\tau_C = 33$ yr, as shown in Fig. ??a, and suggests that $\Delta C(t)$ is described by the response function,

$$\Delta G_C(t) = (r/\tau_C) \exp[-t/\tau_C]. \quad (6)$$

A calibration factor r has been introduced here because this response function is certainly too simplistic. For instance, Taylor expansion to first order does not take into account saturation of carbon flux into the ocean, which will invoke a much longer response time governed by biogeochemical processes of transport of Carbon from the mixed layer into the deep ocean. If we fix τ_c at value higher than 33 yr, r can be estimated by a simple, linear regression to the historic CO₂ concentration record. For $\tau_C = 33$ yr such regression yields of course $r \approx 1$, but for $\tau_C \geq 300$ yr it yields $r \approx 0.5$. This means that the “effective emission rate” in Eq. (??) is reduced to $rR(t)$. The natural interpretation is that approximately half of the emitted CO₂ is almost instantly removed from the atmosphere and the remainder has a lifetime of centuries, maybe millennia, i.e., that the response occurs on one fast and one slow time scale. Model studies, however, may suggest a hierarchy of time scales for the CO₂ concentration response. The large model comparison study of ? reveals a non-exponential tail in the response to a pulse of emitted CO₂. Fig. ??b shows that the multimodel mean is very well approximated by a power-law of the form

$$G_C(t) = \alpha_C t^{\beta_C/2-1}, \quad (7)$$

with $\beta_C \approx 1.6$. This power-law response suggests the simple, linear response model

$$\Delta C(t) = \int_0^t G_C(t-t')R(t') dt', \quad (8)$$

where the emission rate $R(t)$ may contain a stochastic contribution, giving rise to a stochastic component to ΔC . This stochastic component of ΔC is shown in Fig. ??c, as the residual obtained after subtracting a quadratic, polynomial fit to the Muana Lua record (the anthropogenic trend) and the seasonal variation. The power spectral density of this residual is shown in Fig. ??c, and indicates that the spectrum is consistent with a power law with spectral index $\beta_C \approx 1.6$ on time scales longer than a few years. The short duration of the record precludes *accurate* estimates of β_C from the spec-

trum, but it lends some support to the power-law response model with memory exponent in the range $1 < \beta_C < 2$.

270 2.3 The constitutive relation

A simple relation between CO₂ concentration anomaly and its radiative forcing is (?),

$$F = 5.35 \ln(1 + \Delta C/C_0) \text{ Wm}^{-2}. \quad (9)$$

Given an emission scenario $R(t)$, Eq. (??) can be used to compute $\Delta C(t)$ and from Eq. (??) one obtains $F(t)$. Finally this forcing is applied in Eq. (??) to compute $\Delta T(t)$.

275 3 Projections

3.1 Emission scenarios

Fig. ?? shows six different CO₂ emission scenarios. The baseline (unmitigated) scenario is the blue curve, which is an exponential fitted through the actual emission rates in 1960 and in 2010 AD. Interpreted as CO₂ equivalents of all well-mixed greenhouse gases it is close to the RCP8.5 emission scenario up till 2070, but is higher after this time, since the RCP8.5 emission rates saturate between 2070 and 2100. At 2030, 2070, and 2110 AD two types of mitigation action are considered. One where emissions are reduced by 1% per yr (50% reduction over 70 yr) and one with 5% per yr (50% reduction over 13.5 yr). The former is considered politically and economically feasible (?), the latter is at the limit of what is possible without total disruption of the world economy (?). The scenarios are similar to those considered by ?, although they are prescribed from 1880 AD, not from the present day. This is important for the response models employed here, since inertia (long memory) effects from the historical period of global emissions and warming influence the future projections.

3.2 Projections of CO₂ concentration

Atmospheric CO₂ concentrations $\Delta C(t)$ for the emission scenarios described in Fig. ?? are shown in Fig. ?. They are computed from Eq. (??), using the emission scenarios of Fig. ??, and subsequently estimating r by regressing to the historic $\Delta C(t)$ record. Fig. ??a shows the corresponding concentration scenarios estimated from the exponential response kernel with $\tau_c = 33$ yr. Few climate scientists believe that atmospheric, anthropogenic CO₂ are eliminated as fast as this, but it is interesting to examine, since this is still claimed by some “global warming skeptics” (?). In Fig. ??b and Fig. ??d the same scenarios are shown, assuming $\tau_c = 300$ yr and $\tau_c = \infty$, respectively. Here $r \approx 0.5$, i.e., 50% of the emitted CO₂ immediately removed from the atmosphere and the rest decaying exponentially with e-folding time τ_C . Fig. ??c employs the power-law response kernel with $\beta_C = 1.6$. Fig. ??b and Fig. ??c are almost identical, indicating that immediate removal of half of the emitted CO₂, followed by an exponential decay with $\tau_C = 300$ yr, has almost the same effect as a long-memory (power-law) response with $\beta_C = 1.6$.

The unmitigated concentration scenarios (blue curves) are almost the same in all models, and are very similar to the RCP8.5 scenario up to 2100AD. This is because the calibration factor r adjusts the scenario to fit the historic record. However, the evolution after mitigation action has started varies considerably between the models. The overly optimistic model in Fig. ??a, where $\tau_C = 33$ yr, predicts that the concentration starts declining a few decades after emission reduction has started, whereas in the other scenarios concentration continues to rise beyond 2200 AD in the 1% reduction scenarios. The scenario corresponding to the red full curves in Fig. ??b or Fig. ??c correspond closely to the full RCP8.5 scenario.

3.3 Projections of the GMST

The forcing $F(t)$ for the various concentration scenarios is computed from Eq. (??), and inserted into Eq. (??) to obtain the temperature evolution. Fig. ?? shows results for the concentration scenarios obtained from the exponential CO₂ concentration model with $\tau_C = 33$ yr and the power-law model with $\beta_C = 1.6$, considering these to represent low- and high-inertia ends of the CO₂ response. For each of these cases, low- and high-inertia ends ($\beta_T = 0.35$ and $\beta_T = 0.75$) of the GMST response are presented in the figure.

The projections for the high-inertia combination $\beta_C = 1.6$, $\beta_T = 0.75$ shown in Fig. ??d is the one that is most consistent with multi-model CMIP5 projections in the RCP8.5 scenario. As mentioned in Sect. 3.2, the red curve in Fig. ??c is close to the RCP8.5 CO₂-concentration pathway, and the corresponding GMST response shown by the red curve in Fig. ??d is close to the multimodel-mean GMST response given in Fig. 6 of ?. The high-end inertia ($\beta_T = 0.75$) for GMST response is also more consistent with analysis of instrumental records and multiproxy reconstructions of GMST (?) and millennium-long simulations of intermediate and high complexity (?). The high-end inertia for the CO₂-response is also more consistent with complex Carbon-cycle models, and the long-memory nature of the residual Mauna Lua record, as shown in Fig. ??d.

3.4 Policy implications

The range of the projections corresponding to given emission scenarios presented in Fig. ??a – d is much wider than the uncertainty of scientific knowledge reflected in the climate science literature. But it may give an indication of the doubts which are quite common outside the climate science community. Among these are the belief that CO₂ is removed from the atmosphere within decades (?), and that the GMST relaxes to a new radiative equilibrium within a few years after a sudden perturbation of radiative forcing (?). Fig. ??a presents projections which follow from these perceptions. Interestingly, the unmitigated projections up to 2110 AD (blue curves) are almost identical in all panels in Fig. ?. Hence, the inertia in the responses has little influence on the unmitigated response to the BAU emission scenario and uncertainty about the magnitude of the inertia parameters does not contribute much to uncertainty in the response to this scenario. Uncertainty in these pa-

rameters mainly plays a role for the projected effect of the emission reduction after action has been taken, as can be observed by comparing Fig. ??a and Fig. ??d. The effect of emission reduction is considerably greater under the optimistic low-inertia assumptions, but in all circumstances, delayed mitigation action increases the GMST in 2200 AD by 1-2 degrees for every 40 yr of delay.

340 One implication from this observation is that the global warming optimists have little reason for their optimism, since even the projections in Fig. ??a imply that the two-degree climate target will not be attained unless a radical and consistent emission reduction regime is initiated within a few decades from now. If this mitigation regime is delayed and initiated one generation later even the optimistic projections indicate that the temperature will peak close to 3 degrees during the next
345 century, and postponing yet another generation will let the temperature to rise beyond 4 degrees. If emission reductions are raised to the absolute pain threshold of 5% per yr, the peak temperature will not change much, but the temperature will come down faster after action has been initiated.

Under the more pessimistic, and presumably more realistic, circumstances presented in Fig. ??b and Fig. ??d the two-degree target is attainable only if extremely radical reductions (5% per yr) are
350 initiated within the coming two decades. Since such a strong emission reduction regime probably is politically infeasible, this target most likely is unattainable, and the globe will warm 3-7 degrees before the end of next century. Where the GMST will end within this range will essentially depend on the time it takes before radical global emission reductions is implemented. Hence, the slow socio-economic response may turn out to be the most detrimental of all inertia effects which threaten to
355 aggravate global warming.

4 Conclusions

It has been demonstrated that an extremely simple model for the global temperature response and the elimination of excess CO₂ from the atmosphere is all that is needed to make reasonable projections of global temperature under idealised emission scenarios. The model contains only four parameters;
360 characterising sensitivities and inertia in the temperature and CO₂ responses, respectively. All parameters can be estimated from observation data, although some constraining from high-complexity ESMS is useful. The model can be used as a pedagogical tool for students and scientists with some knowledge of elementary calculus, and projections can easily be produced under emissions scenarios different from those presented here.

365 The simplicity of the model may be perceived as an insult to “real” climate modellers, but as long as one deals only with global quantities, simplicity does not necessarily mean lack of accuracy. Global temperature has been found to respond quite linearly to forcing in general circulation models (?), and as long as the climate system is far from a major tipping point, this linearity may also pertain to the response of atmospheric CO₂ content to emissions. Under linearity and stationarity assump-

370 tions these two quantities are fully described in terms of their respective response functions, whose form can be postulated from basic physical principles and parameters estimated from observation.

For the policy makers of the world it is crucial to know to what extent an economically and politically painful mitigation scenario can be expected to be effective in constraining global warming. The analysis presented here confirms the main conclusion drawn by ?; the greatest threat against the stability of the global climate is the inability of humankind to respond in time.

Acknowledgements. This work was funded by project no. 229754 under the the Norwegian Research Council KLIMAFORSK programme.

Appendix A: Response to step forcing for one-box model

The linearised one-box model has the form

$$380 \quad C_1 \frac{dT_1}{dt} = -\frac{T_1}{S_{eq}} + F. \quad (A1)$$

Here T_1 is the perturbation of the mixed-layer temperature from an imagined equilibrium and F is the forcing relative to that equilibrium. C_1 is the heat capacity per square meter of the mixed layer, and the term T_1/S_{eq} is the linearised expression for the intensity of the outgoing long-wave radiation (OLR). It is determined by the (linearised) Stefan-Boltzmann (SB) law and the effective emissivity of the atmosphere, which also contains the effects of fast feedbacks. The nonlinear version and the linearisation procedure is described in Appendix ??. If a new equilibrium is attained with the forcing F we have

$$S_{eq} = \frac{T_1}{F},$$

which makes it natural to name S_{eq} the equilibrium climate sensitivity. It is determined from the SB constant and the effective atmospheric emissivity, i.e., it is totally determined by the atmosphere. The response function (Greensfunction: the response to $F = \delta(t)$) for the one-box model is

$$G(t) = \frac{1}{C_1} e^{-t/\tau_1} H(t), \text{ where } \tau_1 = C_1 S_{eq},$$

and $H(t)$ is the Heaviside unit step function. The response to a step-function forcing $F(t) = H(t)$ is,

$$T_1(t) = \int_{-\infty}^t G(t-t') dt' = S_{eq}(1 - e^{-t/\tau_1}). \quad (A2)$$

Appendix B: Response to step forcing for two-box model

385 The recent work by ? shows that a two-exponentials response can be fitted very well to a number of 150 yr AOGCM runs with step-function forcing. This raises the question whether the power-law LRM-response representation is really only an inaccurate expression of a response with two

exponential time scales, or vice versa. There is also an issue of whether the AOGCMs really capture the true scaling properties of the observed response. The two-box model couples the mixed layer to the deep ocean temperature T_2 through a simple heat conduction term

$$\begin{aligned} C_1 \frac{dT_1}{dt} &= -\frac{1}{S_{eq}} T_1 - \kappa(T_1 - T_2) + F \\ C_2 \frac{dT_2}{dt} &= \kappa(T_1 - T_2). \end{aligned} \quad (\text{B1})$$

where C_2 is the heat capacity of the deep ocean and κ is a heat conductivity. In the limit $C_2 \gg C_1$, the Greens-function for $T_1(t)$ correct to lowest order in the small parameter C_1/C_2 , is very simple and transparent;

$$G(t) = \left(\frac{S_{tr}}{\tau_{tr}} e^{-t/\tau_{tr}} + \frac{S_{eq} - S_{tr}}{\tau_{eq}} e^{-t/\tau_{eq}} \right) H(t), \quad (\text{B2})$$

The reponse to a step-function forcing; $F = H(t)$ then becomes

$$T_1(t) = S_{tr}(1 - e^{-t/\tau_{tr}}) + (S_{eq} - S_{tr})(1 - e^{-t/\tau_{eq}}), \quad (\text{B3})$$

where we have introduced some new parameters,

$$S_{tr} = \frac{S_{eq}}{1 + \kappa S_{eq}}, \quad \tau_{tr} = C_1 S_{tr}, \quad \tau_{eq} = \frac{C_2 S_{eq}}{1 - S_{tr}/S_{eq}}. \quad (\text{B4})$$

These parameters replace the heat capacities $C_{1,2}$ and the heat coupling constant κ , whose physical meaning is easy to grasp, but hard to measure directly. The meaning of the new parameters is apparent if we consider the response to a step-function forcing. Since $C_1/C_2 \ll 1$ we have $\tau_{tr} \ll \tau_{eq}$, and for $t \ll \tau_{eq}$ the response is completely dominated by the first term in equation (??), and hence relaxes exponentially with the transient time constant τ_{tr} to the new quasi-equilibrium S_{tr} , which is denoted the transient climate sensitivity. However, when t approaches τ_{eq} the second term comes into play, and there is a new delayed response with time constant τ_{eq} giving relaxation to the full radiative equilibrium S_{eq} .

From comparing the terms $-T_1/S_{eq}$ and $-\kappa(T_1 - T_2)$ in Eq. (??) we observe that κS_{eq} measures the ratio between the heat flux into the deep ocean and the OLR at the early stage of the response, i.e., when T_2 is still close to zero. From Eq. (??) we have that the part of the sensitivity caused by the slow response from interaction with the deep ocean is

$$S_{eq} - S_{tr} = (\kappa S_{eq}) S_{tr}.$$

Hence, it appears that κS_{eq} is an important parameter. If $\kappa S_{eq} \ll 1$ the inclusion of the deep ocean has little effect on the relaxation to equilibrium. If $\kappa S_{eq} \simeq 1$ or larger the slow response leads to a significant rise of the temperature after the transient equilibrium has been attained. The fast and the slow time constants are always well separated if $C_1 \ll C_2$ since

$$\frac{\tau_{tr}}{\tau_{eq}} = \frac{C_1}{C_2} \frac{\kappa S_{eq}}{(1 + \kappa S_{eq})^2} \leq \frac{C_1}{4C_2}.$$

Appendix C: Response to step forcing in LRM model and GCMs

The LRM-scaling response function $G_T(t) = \alpha_T t^{\beta_T/2-1}$ yields a response $T \sim t^{\beta_T/2}$ to a step in the forcing at time $t = 0$, while a linearly growing forcing yields a response $T \sim t^{\beta_T/2+1}$. Since the forcing is logarithmic in the CO₂ concentration the latter corresponds to exponentially growing concentration. Climate-model runs with linearly growing forcing are of course more realistic than
410 step-function runs, but both have been conducted as part of the CMIP5 project. Examples are 150 yr long simulations of the GISS-E2-H model with a sudden quadrupling of the CO₂-concentration (Fig. ??a) and a 1% per yr increase in the CO₂-concentration (Fig. ??b). A fit of the LRM-scaling response $T \sim t^{\beta_T/2}$ to the GISS-model result in Fig. ??a yields $\beta_T \approx 0.32$, and the solution is shown
415 as the red curve in the figure. The solution of the form $T \sim t^{\beta_T/2+1}$ is shown as the red curve in Fig. ??b. The fit to the tail of the step-function response looks good up to the 150 yr duration of the simulation, but the divergence of the solution as $t \rightarrow \infty$ indicates that the power-law tail with $\beta_T > 0$ is unrealistic for sufficiently large times. There exist few AOGCM simulations that investigate the response to such idealised forcing on millennium time scale. In ? some figures with results of such
420 runs are given. Fig. ??c is an adaptation of Fig. 3 in ?, which shows a 2000 yr long run of the GISS ModelE-R, and Fig. ??d shows a plot of the function $ct^{\beta_T/2+1}$ with $\beta = 0.32$. It demonstrates that at least this particular AOGCM exhibits the power-law tail in the temperature response on time scales up to two millennia.

Note that the $\beta_T \approx 0.32$ obtained for the LRM-model on long time scales is smaller than the
425 $\beta_T \approx 0.75$ estimated from the spectra of the residual of the instrumental data after the response to the deterministic forcing has been subtracted (?). If we produce such residuals by subtracting the red curves from the GISS-model curves in Fig. ??a,b the result looks like a fractional Gaussian noise (fGn) with spectral exponent $\beta \approx 0.65$. As mentioned in Sect. 2.1 an fGn $x_\beta(t)$ characterised by the spectral exponent β is produced by the convolution integral Eq. (2) in the main manuscript if
430 the response kernel is $G(t) \sim t^{\beta/2-1}$ and the forcing function $F(t)$ is a white Gaussian noise $x_0(t)$ (white noise is an fGn with $\beta = 0$). In other words we have,

$$x_\beta(t) = \int_{-\infty}^{\infty} t'^{\beta/2-1} H(t-t') x_0(t') dt', \quad (C1)$$

where $H(t)$ is the unit step function. By using the convolution theorem for the Fourier transform it is easily shown (?) that if $F(t)$ is an fGn with spectral exponent β_F , and the response function has
435 exponent β_T , then the convolution will produce an fGn with $\beta = \beta_T + \beta_F$;

$$x_\beta(t) = \int_{-\infty}^{\infty} t'^{\beta_T/2-1} H(t-t') x_{\beta_F}(t') dt'. \quad (C2)$$

In ? it was suggested that the discrepancy between the spectral exponent β of residuals in observed and simulated GMST records could be explained by assuming some long-range memory ($\beta_F > 0$)

in the stochastic forcing. It was pointed out there that this LRM could even be present in the CO₂-
 440 forcing, since some recent studies indicate strong spatiotemporal heterogeneity in the atmospheric
 CO₂ concentration which might give rise to a fluctuating global component of the global CO₂-
 forcing with long-memory properties.

Appendix D: Two-box vs. LRM fitting to GCM results

? have fitted the two-box model to 16 runs of 150 yr length to step-function forcing. There are four
 445 fitting parameters, and the fits are generally good. There is, however, a wide scatter in the fitting
 parameters between the different models, which may be an indication of overfitting. In Fig. ?? the
 surface temperature solution to the two-box model

$$T_1(t) = [S_{tr}(1 - \exp(-t/\tau_{tr})) + (S_{eq} - S_{tr})(1 - \exp(-t/\tau_{eq}))] F_{4 \times \text{CO}_2}, \quad (\text{D1})$$

and to the LRM model

$$450 \quad T_1(t) = ct^{\beta_T/2} F_{4 \times \text{CO}_2}, \quad (\text{D2})$$

have been fitted to simulation results for the GMST of climate models with step-forcing, $F(t) =$
 $F_{4 \times \text{CO}_2} H(t)$. Here $F_{4 \times \text{CO}_2} \approx 8.61 \text{ Wm}^{-2}$ is the forcing associated with a quadrupling of the atmo-
 spheric CO₂ concentration. The fitting parameters obtained are given in Table 1.

Model	τ_1 (months)	τ_2 (months)	S_{tr} (Km ² /W)	S_{eq} (Km ² /W)	c	β_T
GISS-E2-H	26	663	0.29	0.46	0.14	0.32
BNU-ESM	46	729	0.46	0.69	0.21	0.33
CCSM4	49	4.1×10^{10}	0.33	3.9×10^6	0.10	0.40
CNRM_CM5	38	390	0.37	0.58	0.20	0.31
MPI-ESM-LR	34	1061	0.46	0.75	0.20	0.33

Table 1. Parameters estimated by fitting Eqs. (??) and (??) to the climate model responses to an abrupt qua-
 drupling of atmospheric CO₂ shown in Fig. ?. The table shows the parameters obtained by the Mathematica
 routine FindFit.

The LRM-model in general gives a poorer fit on the short time scales. This is not surprising, since
 455 the LRM-response $ct^{\beta_T/2}$ has an infinite derivative at $t = 0$. However, a much better approximation
 is obtained if we fit the LRM model only in the interval (0, 100) months, but then β_T is raised to
 approximately 0.75. If we implement a four-parameter model with one power-law ($\beta_T \approx 0.75$) up
 to 100 months and another ($\beta_T \approx 0.35$) for $t > 100$ months, we obtain fits comparable to the two-
 exponential model. There is a wide scatter in the model parameters for the two-box model. Note
 460 particularly the huge values for τ_{eq} and S_{eq} for the CCSM4 model. The long time-scale tail is not
 captured by a reasonable exponential, but is well approximated by a reasonable power-law. On the

other hand, the scatter in the LRM-model parameters is small. All this indicates that the two-box model may suffer from overfitting in some cases.

When projections are limited to 2200 CE there is no practical difference between using a power-law response kernel (the LRM model) and the two-exponential kernel (the two-box model). This is illustrated in Fig. ??, where we compute the response for the exponential CO₂-concentration model with $\tau_C = 33$ yr and the two-box model parameters corresponding to the GISS-E2-H model and the CNRM_CM5 models, respectively. The parameters for the two models differ significantly, but the projections are almost identical. Moreover, they are very similar to the projections in Fig. ??a, where the temperature response is produced by the LRM-model with $\tau_C = 33$ yr and $\beta_T = 0.35$. This demonstrates that the mathematical divergence of the solution Eq. (??) for a step-function forcing has little impact on the projection up to 2200 CE for the forcing scenarios considered here. The advantage of the power-law kernel is that it provides a more parsimonious description (fewer fitting parameters) which provides a more precise parameter estimation.

475 Appendix E: Divergences, causality and initial conditions

If $G(t)$ is a power law the integral over prehistory $t \in (-\infty, 0)$ may lead to paradoxes, such as divergences of the integral. The solution to the paradox is to interpret the power-law as an approximation, for instance to a superposition of exponential response kernels. For a white-noise forcing this corresponds to an aggregation of Ornstein-Uhlenbeck (OU) processes, which are known to have the potential to produce a process that is a very good approximation to a fractional Gaussian noise (fGn) up to the time scale corresponding to the OU process with the greatest correlation time (?).

The scaling properties on scales of decades and longer arise from the heat transport within the oceans. This transport exhibits a maximum response time, which will provide an upper (exponential) cut-off of the power-law response function, but the characteristic time of this cut-off may be centuries or millennia. ? state in their abstract: “Scaling up to decades is demonstrated in observations and coupled atmosphere-ocean models with complex and mixed-layer oceans. Only with the complex ocean model the simulated power laws extend up to centuries.”

If we don’t treat the power-law as an approximation we have to deal with the divergences of the integral

$$490 \quad \Delta T(t) = \int_{-\infty}^t G(t-t') F(t') dt', \quad (\text{E1})$$

where $G(s) = s^{\beta_T/2-1}$. If we consider the unit step-function forcing $F(t) = H(t)$, and $\beta_T \neq 0$, the integral is

$$\Delta T(t) = \lim_{\epsilon \rightarrow 0^+} \int_{\epsilon}^t (t-t')^{\beta_T/2-1} dt' = \lim_{\epsilon \rightarrow 0^+} \int_{\epsilon}^t s^{\beta_T/2-1} ds = \lim_{\epsilon \rightarrow 0^+} \frac{2}{\beta_T} (t^{\beta_T/2} - \epsilon^{\beta_T/2}). \quad (\text{E2})$$

Clearly $\Delta T(t)$ diverges as $t \rightarrow \infty$ if $\beta_T > 0$, but it also diverges if $\beta_T < 0$ (as $\epsilon \rightarrow 0^+$). For $\beta_T = 0$
 495 there is a logarithmic divergence in both limits.

For physically meaningful results the $\beta_T > 0$ case requires some sort of cut-off (e.g., an exponential
 tail) for sufficiently large t , and the $\beta_T < 0$ case requires an elimination of the strong singularity
 of $G(s)$ at $s = 0$. As shown in Appendix ??, AOGCMs in the CMIP5 ensemble with step function
 forcing indicate a power-law response for large s at least up to 150 yr (and the GISS-E2-R model
 500 up to 2000 yr) with $\beta_T \approx 0.35$, so $\beta_T > 0$ is the case of interest for the global temperature response.
 The AOGCMs are also well approximated by an exponential response in the limit $s \rightarrow 0$ (for s up to
 a few years), so an exponential truncation in this high-frequency limit is also appropriate.

The truncation of the power-law kernels is a physical, and not a technical mathematical issue. It
 is an approximation to a hierarchy of exponential responses. With this interpretation the divergences
 505 evaporate. Below is a more detailed outline of this philosophy in an energy-balance context. Let
 us take as a starting point the simple zero-dimensional EBM before linearisation of the Stefan-
 Boltzmann law;

$$C \frac{dT}{dt} = -\epsilon \sigma_S T^4 + I(t), \quad (\text{E3})$$

where T is surface temperature in Kelvin, C is an effective heat capacity per area of the Earth's sur-
 face, σ_S is the Stefan-Boltzmann constant, ϵ is an effective emissivity of the atmosphere, and $I(t)$ is
 510 the incoming radiative flux density at the top of the atmosphere. Let $I_0 = I(0)$ be the initial incom-
 ing flux, $F(t) = I(t) - I_0$ is the radiative forcing, $T_{eq} = (I_0/\epsilon\sigma_S)^{1/4}$ is the equilibrium temperature
 at $t = 0$, $\Delta T(t) = T(t) - T_{eq}$ is the temperature anomaly measured relative to the initial equilibrium
 temperature, and $\Delta T_0 = \Delta T(0)$ is this anomaly at $t = 0$. Note that F here is the perturbation of the
 515 radiative flux with respect to the initial flux I_0 and not with respect to the flux $\epsilon\sigma_S T_0^4$ that would
 be in equilibrium with the initial temperature T_0 . The linearised EBM for the temperature change
 relative to the temperature T_0 (the one-box model) is

$$\frac{d\Delta T}{dt} = -\nu \Delta T + \mathcal{F}(t), \quad \Delta T(0) = \Delta T_0, \quad (\text{E4})$$

where $\nu = 4\epsilon\sigma_S T_0^3/C$, $\mathcal{F}(t) = F(t)/C$. By definition $\mathcal{F}(0) = [I(0) - I_0]/C = 0$. This is Eq. (??)
 520 and Eq. (??) with slightly different notation. The solution the initial value problem (i.v.p.) Eq. (??),
 with the initial condition $\Delta T(0) = \Delta T_0$, takes the form

$$\Delta T_{\text{i.v.p.}} = \int_0^t G(t-t') \mathcal{F}(t') dt' + \Delta T_0 e^{-\nu t}, \quad (\text{E5})$$

where $G(s) = \exp(-\nu s)$. The generalisation to a linear, causal response model, where $G(s)$ is
 not necessarily exponential, involves extending the integration domain in Eq. (??) to the interval
 525 $(-\infty, t)$;

$$\Delta T_{\text{r.m.}}(t) = \int_{-\infty}^t G(t-t') \mathcal{F}(t') dt'. \quad (\text{E6})$$

From the initial condition $\Delta T(0)_{\text{r.m.}} = \Delta T_0$ Eq. (??) yields,

$$\Delta T_0 = \int_{-\infty}^0 G(-t') \mathcal{F}(t') dt'. \quad (\text{E7})$$

For exponential response $G(s) = \exp(-\nu s)$ it is easy to verify that $\Delta T_{\text{i.v.p.}}(t) = \Delta T_{\text{r.m.}}(t)$, and
 530 Eq. (??) yields the following relation between the initial temperature anomaly and the forcing $\mathcal{F}(t)$
 for $t \in (t, 0)$;

$$\Delta T_0 = \int_{-\infty}^0 e^{\nu t'} \mathcal{F}(t') dt'. \quad (\text{E8})$$

For the exponential response there is no “divergence issue” in Eq. (??). Neither is there such an
 issue for the two-exponential solution to the two-box model (?). An “ N -box model” exhibits a
 535 response function for the temperature in each box which is a superposition of exponentials; $G(s) =$
 $\sum_{i=1}^N a_i \exp(-\nu_i s)$. For the surface (mixed layer) box the temperature anomaly takes the form

$$\Delta_{\text{r.m.}}(t) = \sum_{i=1}^N a_i e^{-\nu_i t} \int_{-\infty}^t e^{\nu_i t'} \mathcal{F}(t') dt'. \quad (\text{E9})$$

On the other hand, the N -box initial value problem has solution of the form

$$\Delta T_{\text{i.v.p.}}(t) = \sum_{i=1}^N a_i e^{-\nu_i t} \int_0^t e^{\nu_i t'} \mathcal{F}(t') dt' + \sum_{i=1}^N b_i e^{-\nu_i t}, \quad (\text{E10})$$

540 where the coefficients b_i are linearly related to the initial temperatures of each box; $b_i = \sum_{j=1}^N M_{ij} T_{0j}$.
 The condition $\tilde{T}_{\text{i.v.p.}}(t) = \tilde{T}_{\text{r.m.}}(t)$ now yields the relations between the initial temperatures and the
 prehistory of the forcing;

$$\sum_{j=1}^N M_{ij} \Delta T_{0j} = a_i \int_{-\infty}^0 e^{\nu_i t'} \mathcal{F}(t') dt' \quad \text{for } i = 1, \dots, N. \quad (\text{E11})$$

With a white-noise forcing $\mathcal{F}(t)$ the Eq. (??) is the Itô stochastic differential equation (in physics
 545 often called the Langevin equation). The solution is the Ornstein-Uhlenbeck (OU) stochastic process,
 which in discrete time corresponds to the first-order autoregressive (AR(1)) process. The power
 spectral density of this process is essentially a Lorentzian, which means that the high-frequency
 ($f \gg \nu$) part of the spectrum has the form $\sim f^{-2}$, and the low-frequency part $\sim f^0$. This means
 that if the climate response were well described by a one-box EBM we could use a power-law
 550 response model with $\beta_T \approx 2$ on time scales much shorter than the correlation time $\tau_c = \nu^{-1}$. On
 these time scales the stochastic process exhibits the characteristics of a Brownian motion (Wiener
 process), which is a self-similar process with spectral index $\beta = 2$. This process is non-stationary,
 and hence suffers from the divergences that we are worried about. But even though the Brownian
 motion diverges, the OU-process does not, because of the flattening of the spectrum for $f \ll \nu$.

555 Both observation data and AOGCMs indicate that the one-box EBM is inadequate, but the considerations above are equally valid for an N -box model, for which the white-noise forcing gives rise to an aggregation of OU-processes with different ν_i . Such an aggregation is known to be able to produce a process with approximate power-law spectrum with $0 < \beta < 2$ on time scales $\tau < \nu_{\min}^{-1}$ (?).

560 ? specifically argue that volcanic forcing may have a scaling exponent $\beta_F \approx 0.4$, and hence the convergence criterion $\beta = \beta_T + \beta_f < 1$ then requires $\beta_T < 0.6$. One remark to this is that the above discussion shows that the $\beta < 1$ criterion is not necessary on time scales shorter than $\tau < \nu_{\min}^{-1}$. However, observation indicates that $\beta < 1$, so this does not invalidate their argument. More important is that in recent papers the response to volcanic forcing has been subtracted from both instrumental
565 and multiproxy reconstruction data ? and from millennium-long AOGCM simulations (?), and the residuals have been analysed for β without finding a detectable influence of the volcanic forcing on β . The same is seen by comparing control runs of the AOGCMs with those driven by volcanic forcing (?).

Appendix F: Non-stationarity of the CO₂ response

570 In Sect. 2.2 we found (by comparing Figs. ??b and ??c that the LRM CO₂ response with $\beta_C = 1.6$ gives approximately the same evolution of CO₂ concentration up to 2200 CE as a response where 50% of the emitted CO₂ is absorbed by the surface almost immediately and the remainder decays exponentially with a time constant $\tau_C = 300$ yr. This is analogous to the situation with the temperature response, where where an LRM response gives very similar results to a two-exponential response
575 with appropriate fitting of model parameters (see Appendix ??). The most important difference is that the β_C -parameter is larger than unity. A step-function emission rate $R(t) = H(t)$ will give rise to a CO₂ concentration that grows like $(2\alpha_T/\beta_C)t^{\beta_C/2}$. This non-stationarity (divergence) of the response as $t \rightarrow \infty$ is reasonable, since the surface will not be able to absorb a sufficient fraction of the constantly emitted CO₂ to establish a new equilibrium. The exponential response kernel Eq. (??),
580 on the other hand yields the response $r[1 - \exp(-t/\tau_C)]$ to the step forcing. This implies establishment of a new equilibrium CO₂-concentration after $t \gg \tau_C$. This has little consequence as long as we consider projection only up to 2200 CE (and $\tau_C \approx 300$ yr). On millennium time scale we have the positive ice-age feedback, by which warming may lead to net release of CO₂ to the atmosphere, and hence lead to continuing growth of CO₂ concentration. It is assumed to be important in the triggering of glacial-interglacial transitions, although it is not very well understood. On time scales of
585 hundreds of kyr we have the negative Carbon weathering-cycle feedback that will eventually lead to a Carbon cycle equilibrium. The most interesting feature of this feedback in the present context is that it suggests that the anthropogenic global warming event may last for such a long time in absence of effective Carbon sequestration measures (?).

590 A more problematic non-stationarity of the Carbon-cycle response arises from stochastic forcing.
In this case the power-law response function will give rise to a fractional Brownian motion (fBm)
with power-spectral index $\beta_C \approx 1.6$. This is a non-stationary stochastic process in the sense that the
variance increases with time as t^{β_C-1} , which is not physically reasonable for sufficiently large t .
Here we may be saved by an exponential cut-off of the power-law tail, but this requires some sort
595 of negative Carbon-cycle feedback. It is difficult to assess the magnitude of the natural stochastic
component of the CO₂ emission rate. If it small the weathering-cycle feedback may be sufficient.

References

- Archer, D.: The Global Carbon Cycle, Princeton Primers in Climate, Princeton University Press, Princeton, 2010.
- 600 Allen, M. R., et al.: Warming caused by cumulative carbon emissions towards the trillionth tonne, *Nature*, 458, 1163-1166, doi:10.1038/nature08019, 2009.
- Anderegg, W.R.L.; Schneider, S.H., et al.: Expert credibility in climate change, *Proceedings of the National Academy of Sciences*, 107, 12107-12109. doi: 10.1073/pnas.1003187107, 2010.
- Bender, M. L.: Paleoclimate, Princeton Primers in Climate, Princeton University Press, Princeton, 2013.
- 605 Beran, J.: Statistics for Long-memory Processes, Monographs on statistics and applied probability, Chapman& Hall/CRC, Boca Raton, 1994.
- Brohan, P., Kennedy, J. J., Harris, I., Tett, S. F. B., and Jones, P. D.: Uncertainty estimates in regional and global observed temperature changes: A new data set from 1850, *J. Geophys. Res.*, 111, D12 106, 2006.
- Budyko, M. I.: The effect of solar radiation variations on the climate of the Earth, *Tellus*, 21, 611-619, 1969.
- 610 Uncertainty estimates in regional and global observed temperature changes: A new data set from 1850, *J. Geophys. Res.*, 111, D12 106, 2006.
- Cook J., et al.: Quantifying the consensus on anthropogenic global warming in the scientific literature, *Environ. Res. Lett.* 8, 024024, doi:10.1088/1748-9326/8/2/024024, 2013.
- den Elzen, M., Meinshausen, M., van Vuuren, D.: Multi-gas emission envelopes to meet greenhouse gas concentration targets: costs versus certainty of limiting temperature increase, *Glob. Environ. Change*, 17, 260-280, doi:10.1016/j.gloenvcha.2006.10.003, 2007.
- 615 Geoffroy, O., Saint-Martin, D., Olivié, D. J. L., Voldoire, A., Bellon, G., and Tytca, S.: Transient Climate Response in a Two-Layer Energy-Balance Model. Part I: Analytical Solution and Parameter Calibration Using CMIP5 AOGCM Experiments, *J. Climate*, 6, 1841-1857, doi:10.1175/JCLI-D-12-00195.1, 2013.
- 620 Fraedrich, K., and Blender, R., Scaling of Atmosphere and Ocean Temperature Correlations in Observations and Climate Models: *Phys. Rev. Lett.*, 90, 108501-4, doi:10.1103/PhysRevLett.90.108501, 2003.
- Granger, C. W. J.: Long Memory Relationships and the aggregation of dynamical models, *J. Econometrics*, 14, 227-238, 1980.
- Hansen, J., Sato, M., Kharecha, P., von Schuckmann, K.: Earth's energy imbalance and implications, *Atmos. Chem. Phys.*, 11, doi:10.5194/acp-11-13421-2011, 2011.
- 625 Lenton, T. M., Held, H., Kriegler, E., Hall J. W., Lucht, W., Rahmstorf, S., and Schellnhuber, H. J.: Tipping elements in the Earth's climate system, *Proc. Nat. Acad. Sci.*, 105, 1786-1793, doi: 10.1073/pnas.0705414105, 2008.
- Lovejoy, C., Schehrtzer, D., and Varon, D.: Do GCMs predict climate. . . or macroweather?, *Earth Syst. Dynam.*, 4, 439-454, doi: 10.5194/esd-4-439-2013, 2013.
- 630 Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P.M. (eds.): IPCC, 2013: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp., 2013.

- 635 Joos, F., Roth, R., Fugelstveidt, J. S., et al.: Carbon dioxide and climate impulse response functions for the computation of greenhouse gas metrics: a multi-model analysis. *Atmos. Chem. Phys.*, 13, doi:10.5194/acp-13-2793-2011, 2013, 2013.
- Le Quéré et al.: Global carbon budget 2014, *Earth Sys. Sci. Data Discuss* 7, 521-610, doi:10.5194/essdd-7-521-2014, 2014.
- 640 Matthews, H. D., et al.: The proportionality of global warming to cumulative carbon emissions, *Nature*, 459, 1129-132, doi:10.1038/nature08047, 2009.
- Meehl, G. A., Washington, W. M., Amman, C. M., Arblaster, J. M., Wigley, T. M., and Tebaldi, C.: Combinations of Natural and Anthropogenic Forcings in Twentieth-Century Climate. *J. Climate*, 17, 3721–3727, 2004.
- Meinshausen, M., Smith, S. J., Calvin, K., et al.: The RCP greenhouse gas concentrations and their extensions
645 from 1765 to 2300, *Climatic Change*, 11, 2013-241, doi: 10.1007/s10584-011-0156-z, 2011.
- Myhre, G., Highwood, J., Shine, K. P., and Stordahl, F.: New estimates of radiative forcing due to well-mixed greenhouse gases, *Geophys. Res. Lett.*, 25, 2715–2718, 1998.
- Rypdal, K. (2012) Global temperature response to radiative forcing: Solar cycle versus volcanic eruptions, *J. Geophys. Res.*, 117, D06115, doi:10.1029/2011JD017283, 2012.
- 650 Rypdal, K., Østvand, L., and Rypdal, M.: Long-range memory in Earth’s surface temperature on time scales from months to centuries, *J. Geophys. Res.*, 118, doi:10.1002/jgrd.50399, 2013.
- Rypdal, M., and Rypdal, K.: Long-memory effects in linear-response models of Earth’s temperature and implications for future global warming, *J. Climate*, 27, 5240-5258, doi:10.1175/JCLI-D-13-00296.1, 2014.
- Rypdal, K., Rypdal, M., and H.-B. Fredriksen: Spatiotemporal Long-Range Persistence in Earth’s Temperature
655 Field: Analysis of Stochastic-Diffusive Energy Balance Models, *J. Climate*, 28, 8379- 8395, 2015, DOI: 10.1175/JCLI-D-15-0183.1.
- Schwartz, S. E.: Heat capacity, time constant, and sensitivity of the Earth’s climate system, *J. Geophys. Res.*, 112, D24S05, doi:10.1029/2007JD008746, 2007.
- Sellers, W. D.: A global climate model based on the energy balance of the Earth-atmosphere system, *J. Appl. Meteorol.*, 8, 392-400, 1969.
660
- Solomon, L.: *The Deniers. The world-renowned scientists who stood up against global warming hysteria, political persecution and fraud*, Richard Viglante Books, 2008.
- Stocker, T.: The Closing Door of Climate Targets, *Science*, 339, 280-282, doi:10.1126/science.1232468, 2013.
- Stern, N.: *The Economics of Climate Change. The Stern Review*, Cambridge, 2007.
- 665 Østvand, L., Nilsen, T., Rypdal, K., Divine, D., and Rypdal, M.: Long-range memory in internal and forced dynamics of millennium-long climate model simulations, *Earth. Syst. Dynam.*, 5, 295-308, doi:10.5194/esd-5-295-2014, 2014.




Figure 1. (a): Light blue curve is the instrumental GMST for 1880 – 2010 AD. Black curve is the instantaneous response to the linearly extrapolated forcing scenario shown in panel (b). Blue curve is the response according to the model Eq. (??) with $\beta_T = 0.35$, and the red curve with $\beta_T = 0.75$. (b): The blue curve is a linearly projected forcing to 2200 AD with the same mean growth rate as the the RCP8.5 scenario in the period 2010 – 2100 AD. The brown curve is the stabilisation of this forcing in 2030 AD, the blue curve in 2070 AD, and the red curve in 2110 AD. (c): GMST responses to the forcing scenarios in (b) with $\beta_T = 0.35$. Colours correspond to those in (b). (d): Same as in (c), but with $\beta_T = 0.75$.




Figure 2. (a): Blue curve shows the atmospheric CO₂ concentration as measured by the Mauna Lua observatory. The red curve is the concentration computed from Eq. (??) with $\tau_C = 33$ yr, $\Delta C_{1960} = 74$ GtC (corresponding to an anomaly of $315-280=35$ ppm), and $C_0 = 594$ GtC (corresponding to 280 ppm). (b): Black curve is the multimodel mean CO₂ response to a pulse of emitted CO₂ as given in ?. The red, dashed curve is a least-square fit of a function of the form $\alpha_C t^{\beta_C/2-1}$ with the estimated $\beta_C \approx 1.6$. (c): The residual Mauna Lua signal after subtracting the quadratic polynomial and seasonal trends. (d): The power spectral density of the residual in (c) estimated by the periodogram presented in a log-log plot. The blue, dashed line has negative slope $\beta_C = 0.85$, and the red, dashed line $\beta_C = 1.6$.




Figure 3. Blue curve is carbon emission rate $R(t)$ scenario obtained by fitting the exponential $S_0 \exp(gt)$ to the emission rate 4 GtC/yr in 1960 and 11 GtC/yr in 2010 AD. The full, brown, orange, and red curves are the subsequent $R(t)$ after initiation of 1% reduction of emission rate per year. The dashed curves are the corresponding rates with 5% reduction per year.




Figure 4. Projections of CO₂ concentration under the emission scenarios in Fig. ?? using the modelling explained in Sect. 2. The colours correspond to those in Fig. ?. (a): $\tau_C = 33$ yr. (b): $\tau_C = 300$ yr. (c): $\beta_C = 1.6$. (d): $\tau_C = \infty$.




Figure 5. (a): The evolution of the GMST for the CO₂ concentration scenarios shown in Fig. ??a and Fig. ??c. (a): $\tau_C = 33$ yr and $\beta_T = 0.35$. (b): $\beta_C = 1.6$ and $\beta_T = 0.35$. (c): $\tau_C = 33$ yr and $\beta_T = 0.75$. (d): $\beta_C = 1.6$ and $\beta_T = 0.75$.




Figure 6. (a): LRM response model fit $c_1 t^{\beta_T/2}$ (red) to the GISS-E2-H model response to an abrupt quadrupling of atmospheric CO_2 (grey). The fit yields $\beta_T = 0.32$. (b): The LRM-reponse model solution $c_2 t^{\beta_T/2+1}$ with $\beta_T = 0.32$ (red) and the GISS-E2-H model response to a 1 % per yr increase in atmospheric CO_2 -concentration. (c): The 2000 yr response to a doubling of CO_2 in GISS ModelE-R as taken from Figure 3 in ?. (d) Response to the same forcing in the LRM model with $\beta_T = 0.32$.




Figure 7. Blue curves: Fit of the two-exponential response to the climate model responses to an abrupt quadrupling of atmospheric CO₂ concentration. Red curves: Fit of the LRM-scaling response. The expressions fitted are found in the caption of Table 1 and the coefficients estimated are shown in this table.




Figure 8. (a): The evolution of the GMST according to the two-box solution Eq. (??) for the CO₂ concentration scenarios shown in Fig. ??a and Fig. ??c. (a): $\tau_C = 33$ yr and the two-box parameters for the GISS-E2-H given in Table 1. (b): $\tau_C = 33$ yr and the two-box parameters for the CNRM_CM5 model given in Table 1.

Interactive
Comment

Interactive comment on “Global warming projections derived from an observation-based minimal model” by K. Rypdal

dr. Rypdal

kristoffer.rypdal@uit.no

Received and published: 13 November 2015

Reply to referee #1

Thanks for a positive and constructive review.

Referee: Abstract: Replace “Main value” by “highlights”.

Response: Good suggestion. Will be done.

Referee: Replace “the broad community of scientists” by “the scientific community.”

Response: “scientific community” could be interpreted as “the climate science community,” so that does not cover what I want to say. I can change it to “scientific community at large,” although I cannot really see what was wrong with the original phrase.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Interactive
Comment

Referee: Replace “. . . main driver of dangerous” by “main driver for.”

Response: There are two issues here. The use of the qualifier “dangerous” (global change). I can replace this by: “a main driver of global warming over the last century.” The preposition “of ” is correct here, “for” is not.

Referee: “The unconvincing part is the above mentioned reliance on complex computer models.” – is the author suggesting the reliance on ESMs is not necessary at all?

Response: First, I am not suggesting anything. I refer to views of “many physical scientists.” This sentence should not be read in isolation. In the full context it should be clear that it is EXCESSIVE reliance on ESMs that is perceived as unconvincing. I will add the qualifier “excessive.”

Referee: Remove “And” at the start of sentence.

Response: OK.

Referee: “meaningless acronyms”

Response: I replace this by “acronyms that carry no meaning to non-experts.”

Referee: The phrase: “. . . why do think these horrendously complex models perform any better.”

Response: Again, this is not my argument or opinion. I am just referring to comments I get from my former plasma physics colleagues, and these are formulated just like this. It is not my task to brush them up to make them look more reasonable.

Referee wants to remove the phrase where I claim that the present work is more “transparent” than the paper by Stocker.

Response: I don’t agree. A major motivation for this work is to use an approach that is more transparent in the sense that all assumptions are open to inspection to the reader. An approach that is based on empirical results from a large ensemble of ESMs

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

lacks this type of transparency.

Referee: Use of the first person singular “I.”

Response: In a paper I published in JGR-Atmospheres in 2012 I used the first person plural “we.” Here one referee instructed me to drop “the Royal we” and use “I.” The option which is left is to drop first person entirely, and use passive form everywhere. This inevitably leads to a very stiff style, which may be the desired norm of scientific writing in some languages, but not in English. I shall go through the paper to see if first person singular can be naturally be avoided in some places.

Referee: Elaborate the conclusions.

Response: That seems like a good suggestion, and I will do that.

Interactive comment on Earth Syst. Dynam. Discuss., 6, 1789, 2015.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Interactive
Comment

Interactive comment on “Global warming projections derived from an observation-based minimal model” by K. Rypdal

dr. Rypdal

kristoffer.rypdal@uit.no

Received and published: 7 January 2016

In the revised manuscript the issue of non-stationarity is discussed in an added paragraph in section 2 and in several of the added appendices (six in all).

Interactive comment on Earth Syst. Dynam. Discuss., 6, 1789, 2015.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Manuscript prepared for Earth Syst. Dynam. Discuss.
with version 2015/04/24 7.83 Copernicus papers of the \LaTeX class copernicus.cls.
Date: 7 January 2016

Response to Hébert and Lovejoy

K. Rypdal¹

¹Department of Mathematics and Statistics, UiT The Arctic University of Norway, Norway

Correspondence to: Kristoffer Rypdal (kristoffer.rypdal@uit.no)

The comment by Hébert and Lovejoy (H&L) raises a number of issues which would be interesting to discuss in depth, although some have little relevance to the present discussion paper. The main comment, which *is* relevant, is the following: Will the divergence of the convolution integral over a power-law kernel with $\beta > 0$ as $t \rightarrow \infty$ have an effect on the GMST-projection up to 2200 CE? The short answer is that it does not. In the attached supplement I demonstrate this. It is written as a group of appendices which will be added in the revised manuscript.

The comment of H&L is structured in sections. Below I will respond to each section in the given order. For more detail I refer to the Supplement attached to this reply.

1. Introduction

It is my impression that H&L base their physical interpretation of the power-law response kernel on a turbulent cascade paradigm for atmospheric dynamics (see e.g., Lovejoy and Schertzer, 2013, Chapter 1). The inertia effects (long-range memory) I am discussing in the paper do not arise from the atmosphere but from heat transport in the ocean, and possibly involves other high inertia components of the climate system, such as the cryosphere. This fact is well documented in AOGCMs, since the long-memory scaling only takes place if full ocean circulation is included. GCMs including only the mixed ocean layer does not exhibit this scaling (?).

My interpretation of the approximate scale-invariance of the temperature response is much simpler and based on energy-balance considerations. The climate system consists of a number of interacting subsystems with different response times. The simplest model is the so-called two-box model, where the zero-dimensional energy balance model is supplemented with an equation describing the heat exchange between the mixed ocean layer and the deep ocean. This gives rise to a response kernel consisting of a sum of two exponentials with one short and one long e-folding time. This kernel is already well approximated by a power law up to time scales comparable to the long time constant. In the supplement

30 I show examples from CMIP5 experiments with step-function forcing, where the response
is approximated by a power-law as well as two exponentials. Using the model param-
eters estimated from these experiments on the forcing scenarios used in my manuscript, I
demonstrate that the two kinds of response kernels give almost identical results. Since the
two-exponential kernel does not suffer from the divergence problem, this shows that this
35 divergence does not influence the projections on the time scales considered.

2. A Scaling Climate Response Function

In this section the paragraph below does not make sense to me:

40 “Notice that the initial conditions are important: the same linear operator with
different initial conditions will lead to a different Green’s function. Due to the
scaling symmetry respected by the dynamics, and boundary conditions, over a
wide range of scales, we take the basic Green’s function as a power law.”

The integral goes from $-\infty$ to t , so what does the “initial condition” mean? Let us define a
time origin ($t = 0$) at a specific date, and define T and its derivatives at this date as the initial
45 conditions. The mathematical fact is then that if \mathcal{L} is an ordinary n ’th order differential oper-
ator the evolution for $t > 0$ is uniquely determined by the $T(0)$ and the $n - 1$ first derivatives
at $t = 0$. On the other hand, I will show in Section 3 below that these initial conditions in turn
depend on the entire prehistory of the forcing in the time interval $t \in (-\infty, 0)$. The Green’s
function is determined by the dynamics, is not by the initial conditions. On the contrary, the
50 initial conditions are determined by the Green’s function and the past forcing.

However, if \mathcal{L} is a fractional derivative (i.e., if $G(t)$ is a power law), then the integral over
prehistory $t \in (-\infty, 0)$ may lead to paradoxes, such as divergences of the integral. The
solution to the paradox is to interpret the power-law as an approximation, for instance to a
55 superposition of exponential kernels. For a white-noise forcing this corresponds to an ag-
gregation of Ornstein-Uhlenbeck (OU) processes, which are known to have the potential to
produce a process that is a very good approximation to a fractional Gaussian noise (fGn)

up to the time scale corresponding to the OU process with the greatest correlation time. More on this in Section 3.

60

As discussed in the previous section, the scaling properties on scales of decades and longer do not arise from “the scaling symmetry respected by the (atmospheric) dynamics, and boundary conditions”, but from the heat transport within the oceans. This transport exhibits a maximum response time, which will provide an upper (exponential) cut-off of the power-law response function, but the characteristic time of this cut-off may be centuries or millennia. Let me cite from the abstract of ? : “Scaling up to decades is demonstrated in observations and coupled atmosphere-ocean models with complex and mixed-layer oceans. Only with the complex ocean model the simulated power laws extend up to centuries.”

65

70

3. Low-frequency divergence

In the present paper deterministic (anthropogenic) forcing is the only forcing considered. As discussed in ? the internal variability can be taken into account as a result of stochastic forcing whose variance can be estimated from the observation data. This allows to compute error bars due to internal variability (stochastic errors) on predictions based on the minimal model. I haven't included such error estimates in the paper because I consider model uncertainty to be more important.

75

The H&L comment discusses the conditions for divergence of the integral

80

$$T(t) = \int_{-\infty}^t G(t-t')F(t')dt', \quad (1)$$

where $G(s) = s^{\beta_T/2-1}$. If we consider the unit step-function forcing $F(t) = H(t)$ the integral is

$$T(t) = \lim_{\epsilon \rightarrow 0^+} \int_{\epsilon}^t (t-t')^{\beta_T/2-1} dt' = \lim_{\epsilon \rightarrow 0^+} \int_{\epsilon}^t s^{\beta_T/2-1} ds = \lim_{\epsilon \rightarrow 0^+} \frac{2}{\beta_T} (t^{\beta_T/2} - \epsilon^{\beta_T/2}), \quad (2)$$

85 for $\beta_T \neq 0$. Clearly $T(t)$ diverges as $t \rightarrow \infty$ if $\beta_T > 0$, but it also diverges if $\beta_T < 0$ (as $\epsilon \rightarrow 0^+$). For $\beta_T = 0$ there is a logarithmic divergence in both limits.

For physically meaningful results the $\beta_T > 0$ case requires some sort of cut-off (e.g., an exponential tail) for sufficiently large t , and the $\beta_T < 0$ case requires an elimination of the strong singularity of $G(s)$ at $s = 0$. AOGCMs in the CMIP5 ensemble with step function forcing indicate a power-law response for large s at least up to 150 yr (and the GISS-E2-R model up to 2000 yr) with $\beta_T \approx 0.35$, so $\beta_T > 0$ is the case of interest for the global temperature response. As discussed in Sect. 1 above (and in the Supplement) the inclusion of such a cut-off consistent with the AOGCM results will have very small effects on the predictions up to 2200 CE. The AOGCMs show an exponential response for $s \rightarrow 0$ (for s up to a few years), so an exponential truncation in this high-frequency limit is also appropriate.

I agree with H&L that the truncation of the power-law kernels is a “physical, and not a technical mathematical issue.” But their comment is nothing but a mathematical discussion of convergence, detached from the physical interpretation that I have given to the power-law response as an approximation to a hierarchy of exponential responses. With this interpretation the divergences evaporate. Below I outline the philosophy in some detail in an energy-balance context. Let us take as a starting point the simple zero-dimensional EBM before linearisation of the Stefan-Boltzmann law;

$$C \frac{dT}{dt} = -\epsilon \sigma_S T^4 + I(t), \quad (3)$$

105 where T is surface temperature in Kelvin, C is an effective heat capacity per area of the Earth's surface, σ_S is the Stefan-Boltzmann constant, ϵ is an effective emissivity of the atmosphere, and $I(t)$ is the incoming radiative flux density at the top of the atmosphere. Let $I_0 = I(0)$ be the initial incoming flux, $F(t) = I(t) - I_0$ is the radiative forcing, $T_{eq} = (I_0/\epsilon\sigma_S)^{1/4}$ is the equilibrium temperature at $t = 0$, $\tilde{T}(t) = T(t) - T_{eq}$ is the temperature anomaly measured relative to the initial equilibrium temperature, and $\tilde{T}_0 = \tilde{T}(0)$ is this
 110 anomaly at $t = 0$. Note that F here is the perturbation of the radiative flux with respect to the initial flux I_0 and not with respect to the flux $\epsilon\sigma_S T_0^4$ that would be in equilibrium with the initial temperature T_0 . The linearised EBM for the temperature change relative to the temperature T_0 (the one-box model) is

$$115 \quad \frac{d\tilde{T}}{dt} = -\nu\tilde{T} + \mathcal{F}(t), \quad \tilde{T}(0) = \tilde{T}_0. \quad (4)$$

where $\nu = 4\epsilon\sigma_S T_0^3/C$, $\mathcal{F}(t) = F(t)/C$. By definition $\mathcal{F}(0) = [I(0) - I_0]/C = 0$. The solution the initial value problem (i.v.p.) Eq. (??), with the initial condition $\tilde{T}(0) = T_0$, takes the form

$$\tilde{T}_{i.v.p.} = \int_0^t G(t-t')\mathcal{F}(t') dt' + T_0 e^{-\nu t}, \quad (5)$$

where $G(s) = \exp(-\nu s)$. The generalisation to a linear, causal response model, where
 120 $G(s)$ is not necessarily exponential, involves extending the integration domain in Eq. (??) to $-\infty$;

$$\tilde{T}_{r.m.}(t) = \int_{-\infty}^t G(t-t')\mathcal{F}(t') dt', \quad (6)$$

which corresponds to Eq. (5) in the H&L comment. From the initial condition $\tilde{T}(0)_{r.m.} = \tilde{T}_0$ Eq. (??) yields

$$125 \quad \tilde{T}_0 = \int_{-\infty}^0 G(-t') \mathcal{F}(t') dt'. \quad (7)$$

For exponential response $G(s) = \exp(-\nu s)$ it is easy to verify that $\tilde{T}_{i.v.p.}(t) = \tilde{T}_{r.m.}(t)$, and Eq. (??) yields the following relation between the initial temperature anomaly and the past forcing $\mathcal{F}(t)$ for $t < 0$;

$$T_0 = \int_{-\infty}^0 e^{\nu t'} \mathcal{F}(t') dt'. \quad (8)$$

130 For the exponential response there is no “divergence issue” in Eq. (??). Neither is there such an issue for the two-exponential solution to the two-box model (?). An “ N -box model” exhibits a response function for the temperature in each box which is a superposition of exponentials; $G(s) = \sum_{i=1}^N a_i \exp(-\nu_i s)$. For the surface (mixed layer) box the temperature anomaly takes the form

$$135 \quad \tilde{T}_{r.m.}(t) = \sum_{i=1}^N a_i e^{-\nu_i t} \int_{-\infty}^t e^{\nu_i t'} \mathcal{F}(t') dt'. \quad (9)$$

On the other hand, the N -box initial value problem has solution of the form

$$\tilde{T}_{i.v.p.}(t) = \sum_{i=1}^N a_i e^{-\nu_i t} \int_0^t e^{\nu_i t'} \mathcal{F}(t') dt' + \sum_{i=1}^N b_i e^{-\nu_i t}, \quad (10)$$

where the coefficients b_i are linearly related to the initial temperatures of each box; $b_i = \sum_{j=1}^N M_{ij} T_{0j}$. The condition $\tilde{T}_{i.v.p.}(t) = \tilde{T}_{r.m.}(t)$ now yields the relations between the initial temperatures and the prehistory of the forcing;

$$\sum_{j=1}^N M_{ij} T_{0j} = a_i \int_{-\infty}^0 e^{\nu_i t'} \mathcal{F}(t') dt' \quad \text{for } i = 1, \dots, N. \quad (11)$$

With a white-noise forcing $\mathcal{F}(t)$ the Eq. (??) is the Itô stochastic differential equation (in physics often called the Langevin equation). The solution is the Ornstein-Uhlenbeck (OU) stochastic process, which in discrete time corresponds to the first-order autoregressive (AR(1)) process. The power spectral density (PSD) of this process is essentially a Lorentzian, which means that the high-frequency ($f \gg \nu$) part of the spectrum has the form $\sim f^{-2}$, and the low-frequency part $\sim f^0$. This means that if the climate response were well described by a one-box EBM we could use a power-law response model with $\beta_T \approx 2$ on time scales much shorter than the correlation time $\tau_c = \nu^{-1}$. On these time scales the stochastic process exhibits the characteristics of a Brownian motion (Wiener process), which is a self-similar process with spectral index $\beta = 2$. This process is non-stationary, and hence suffers from the divergences that worries L&H. But even though the Brownian motion diverges, the OU-process does not, because of the flattening of the spectrum for $f \ll \nu$.

L&H argue that if the forcing is not white, but characterised by a spectral exponent β_f , then the criterion for convergence of the response requires $\beta = \beta_T + \beta_f < 1$. But this criterion is only necessary for the long time scales where the temperature process has a stationary character (in the one-box example on time scales greater than τ_c). If the time scales we study are shorter, we may accept $\beta > 1$.

Both observation data and AOGCMs indicate that the one-box EBM is inadequate, but the considerations above are equally valid for an N -box model, for which the white-noise forcing gives rise to an aggregation of OU-processes with different ν_i . Such an aggregation is

known to be able to produce a process with approximate power-law spectrum with $0 < \beta < 2$
165 on time scales $\tau < \nu_{\min}^{-1}$ (?).

L&H specifically argue that volcanic forcing may have a scaling exponent $\beta_f \approx 0.4$, and that
the convergence criterion $\beta = \beta_T + \beta_f < 1$ then requires $\beta_T < 0.6$. One remark to this is that
170 the above discussion shows that the $\beta < 1$ criterion is not necessary on time scales shorter
than $\tau < \nu_{\min}^{-1}$. However, observation indicates that $\beta < 1$, so this does not invalidate L&H's
argument. More important is that the response to volcanic forcing has been subtracted from
both instrumental and multiproxy reconstruction data ? and from millennium-long AOGCM
175 simulations (?), and the residuals have been analysed for β without finding a detectable
influence of the volcanic forcing on β . The same is seen by comparing control runs of the
AOGCMs with those driven by volcanic forcing (?).

There are a number of problems with the stochastic characterisation of volcanic forcing and
how such a shot-noise like process could be incorporated in a response model. This is an
180 interesting issue, but too far from the subject of the present paper to be discussed further
here.

References

- Fraedrich, K., and Blender, R., Scaling of Atmosphere and Ocean Temperature Correlations in Observations and Climate Models: *Phys. Rev. Lett.*, 90, 108501-4, doi:10.1103/PhysRevLett.90.108501, 2003.
- 185 Geoffroy, O., Saint-Martin, D., Olivié, D. J. L., Voldoire, A., Bellon, G., and Tytca, S.: Transient Climate Response in a Two-Layer Energy-Balance Model. Part I: Analytical Solution and Parameter Calibration Using CMIP5 AOGCM Experiments, *J. Climate*, 6, 1841-1857, doi:10.1175/JCLI-D-12-00195.1, 2013.
- Granger, C. W. J.: Long Memory Relationships and the aggregation of dynamical models, *J. Econometrics*, 14, 227-238, 1980.
- 190

Hansen, J., Sato, M., Kharecha, P., von Schuckmann, K.: Earth's energy imbalance and implications, *Atmos. Chem. Phys.*, 11, doi:10.5194/acp-11-13421-2011, 2011.

Lovejoy, S., and Schertzer, D.: *The Weather and Climate, Emergent laws and Multifractal Cascades*, Cambridge University Press, 2013.

195 Rypdal, M., and Rypdal, K.: Long-memory effects in linear-response models of Earth's temperature and implications for future global warming, *J. Climate*, 27, 5240-5258, doi:10.1175/JCLI-D-13-00296.1, 2014.

Rypdal, K., Rypdal, M., and H.-B. Fredriksen: Spatiotemporal Long-Range Persistence in Earth's Temperature Field: Analysis of Stochastic-Diffusive Energy Balance Models, *J. Climate*, 28, 8379-8395, 2015, DOI: 10.1175/JCLI-D-15-0183.1.

200 Østvand, L., Nilsen, T., Rypdal, K., Divine, D., and Rypdal, M.: Long-range memory in internal and forced dynamics of millennium-long climate model simulations, *Earth. Syst. Dynam.*, 5, 295-308, doi:10.5194/esd-5-295-2014, 2014.