

Differential climate impacts for policy-relevant limits to global warming: The case of 1.5° C and 2° C

Schleussner Carl-Friedrich^{1,2}, Lissner Tabea K.^{1,2}, Fischer Erich M.³, Wohland Jan², Perrette Mahé², Golly Antonius^{4,6}, Rogelj Joeri^{3,5}, Childers Katelin², Schewe Jacob², Frieler Katja², Mengel Matthias^{1,2}, Hare William^{1,2}, and Schaeffer Michiel^{1,7}

¹Climate Analytics, Friedrichstr 231 - Haus B, 10969 Berlin, Germany

²Potsdam Institute for Climate Impact Research, Potsdam, Germany

³Institute for Atmospheric and Climate Science, ETH Zurich, Zürich, Switzerland

⁴GFZ German Research Centre for Geosciences, Potsdam, Germany

⁵Energy Program, International Institute for Applied Systems Analysis, Laxenburg, Austria

⁶University of Potsdam, Institute of Earth and Environmental Science, Potsdam, Germany

⁷Wageningen University and Research Centre, Environmental Systems Analysis Group, Wageningen, The Netherlands

Correspondence to: Schleussner, Carl-Friedrich (carl.schleussner@climateanalytics.org)

Abstract.

Robust appraisals of climate impacts at different levels of global-mean temperature increase are vital to guide assessments of dangerous anthropogenic interference with the climate system. The 2015 Paris Agreement includes a two-headed temperature goal: "holding warming well below 2°C above pre-industrial levels, and pursuing efforts to limit the temperature increase to 1.5°C". Despite the prominence of these two temperature limits, a comprehensive overview of the differences in climate impacts at these levels is still missing. Here we provide an overview of key impacts of climate change at warming levels of 1.5°C and 2°C, including extreme weather events, water availability, agricultural yields, sea-level rise and risk of coral reef loss. Our results reveal substantial differences in impacts between a 1.5°C and 2°C warming that are highly relevant for the assessment of dangerous anthropogenic interference with the climate system. For heat-related extremes, the additional 0.5°C increase in global-mean temperature marks the difference between events at the upper limit of present-day natural variability and a new climate regime, particularly in tropical regions. Similarly, this warming difference is likely to be decisive for the future of tropical coral reefs. In a scenario with an end-of-century warming of 2°C, virtually all tropical coral reefs are projected to be at risk of severe degradation due to temperature induced bleaching from 2050 onwards. This fraction is reduced to about 90 % in 2050 and projected to decline to 70 % by 2100 for a 1.5°C scenario. Analyses of precipitation-related impacts reveal distinct regional differences and hot-spots of change emerge. Regional reduction in median water availability for the Mediterranean is found to nearly double from 9 % to 17 % between 1.5°C and 2°C, and the projected lengthening of re-

gional dry spells increases from 7 % to 11 %. Projections for agricultural yields differ between crop types as well as world regions. While some (in particular high-latitude) regions may benefit, tropical regions like West Africa, South-East Asia, as well as Central and Northern South America are projected to face substantial local yield reductions, particularly for wheat and maize. Best estimate sea-level rise projections based on two illustrative scenarios indicate a 50 cm rise by 2100 relative to year 2000-levels for a 2°C scenario, and about 10 cm lower levels for a 1.5°C scenario. A 1.5°C scenario would also reduce the rate of sea-level rise by about 30 % compared to a 2°C scenario. Our findings highlight the importance of regional differentiation to assess both future climate risks and different vulnerabilities to incremental increases in global-mean temperature. The article provides a consistent and comprehensive assessment of existing projections and a good basis for future work on refining our understanding of the difference between impacts at 1.5 °C and 2 °C warming.

1 Introduction

Recent decades have seen increasing climate impacts, many of which science is now able to attribute to anthropogenic carbon dioxide emissions and consequent global warming (IPCC, 2013; King et al., 2015). On-going temperature increase will escalate these impacts on ecological and human systems (IPCC, 2014a), which has made climate change a political issue of central importance. The response of the global community to that challenge laid out in the Paris Agreement under the United Nations Framework Convention on Climate Change (UNFCCC) provides a promising framework for global climate protection (UNFCCC, 2015). Specifically, the Agreement includes two long-term global goals (LTGGs): "holding warming well below 2°C above pre-industrial levels, and pursuing efforts to limit the temperature increase to 1.5°C, recognizing that this would significantly reduce the risks and impacts of climate change". LTGGs have been proven useful to guide climate action (SED, 2015) and their inclusion aims to operationalize the "ultimate objective" of the UNFCCC of a "stabilization of greenhouse gas concentrations in the atmosphere at a level that would prevent dangerous anthropogenic interference with the climate system" (UNFCCC, 1992). Although the assessment of levels of dangerous interference is primarily a political process that requires value judgements and depends on different world views (Knutti et al., 2015), it needs to be informed by the best available science outlining the impacts of climate change and mitigation efforts implied by different LTGGs. Based on the Fifth Assessment Report (AR5) of the Intergovernmental Panel on Climate Change (IPCC), a recent expert assessment focussing on the adequacy of the LTGGs in light of the ultimate objective of the convention concluded that "significant climate impacts are already occurring [...] and additional magnitudes of warming will only increase the risk of severe, pervasive and irreversible impacts" (SED, 2015). While the report emphasized that a warming of global mean surface air temperature (GMT) of 2°C above pre-industrial levels should not be seen as a 'safe' level, it also concluded that substantial research gaps exist regarding the differences in climate impacts between

a 1.5°C and 2°C temperature increase (SED, 2015). In particular, comprehensive, multi-sectoral assessments of differences in climate impacts between a 1.5°C and 2°C warming are lacking. The assessment of such differences would greatly profit from a regional and impact - centered approach that allows for a more differentiated picture than globally aggregated metrics (Seneviratne et al., 2016). For example, changes in the hydrological cycle as a result of temperature increase will be regionally dependent (Held and Soden, 2006).

The "Turn down the heat" - report series issued by the World Bank (Schellnhuber et al., 2012, 2013, 2014) assessed climate risks for a 2°C and a 4°C warming above pre-industrial levels for different world regions. The report of the Working Group 2 (WG2) of the IPCC AR5 includes both, impact and region specific chapters, and provides warming level dependent information on impacts where available. The range of emission scenarios which provide the basis for the climate impact projections in the IPCC AR5, the Representative Concentration Pathways (RCPs), however, do not allow for a straight-forward differentiation between impacts for warming levels of 1.5°C and 2°C. Only the lowest emission pathway RCP 2.6 is in line with keeping GMT increase above pre-industrial levels to below 2°C with a likely chance (66 % probability, IPCC, 2013) and no pathway in line with a 1.5°C limit is assessed in the AR5. Still, the IPCC AR5 WG2 report provides an expert assessment of key impacts at different levels of warming, summarized in five "Reasons-for-Concern" (RFCs, Oppenheimer et al., 2014). The risks for three out of five of these RFCs are assessed as at least moderate at 1.5°C GMT increase above pre-industrial levels, and as at least moderate-high at 2°C. In the RFC framework, moderate risks imply that associated impacts are both detectable and attributable to climate change with at least medium confidence, whereas high risks are associated with severe and widespread impacts (Oppenheimer et al., 2014). Among the three RFCs that show high risks at 2°C are *Risks to unique and threatened systems* (RFC1) that include coral reefs and other highly vulnerable human systems as well as ecosystems, *Risks associated with extreme weather events* (RFC2) and *Risks associated with the distribution of impacts* (RFC3).

Based on the Coupled Model Intercomparison Project 5 (CMIP5, Taylor et al., 2011) and the Inter-Sectoral Impact Model Intercomparison Project (ISI-MIP, Warszawski et al., 2013), this article provides an extensive assessment of regionally differentiated climate impacts at levels of 1.5°C and 2°C GMT increase above pre-industrial levels (henceforth 1.5°C and 2°C) for different climate impacts, including increases in extreme weather events (Section 3), changes in water availability (Section 4), crop yield projections (Section 5), sea-level rise (SLR, Section 6) and coral reef degradation (Section 7).

The following Section (2) outlines our methods for the assessment of changes in extreme weather indices, water availability and agricultural impacts. Analyses of sea-level rise and impacts on coral reefs contain additional details on sector-specific methods. Where impact-specific additional methodological specifications are needed, these are given in the respective section, followed by a presentation of the main results and a short discussion. A summarizing discussion and conclusions finalize

this contribution in Section 8. The Supplementary Material (SM) provides additional methodological information as well as further impact maps, regional overviews and summary tables.

95 2 Methods

This section provides an overview of the methods applied for the assessment of extreme weather indices, water availability and agricultural impacts. The individual subsections provide additional information on sector- and impact-specific methods as well as on the data analyzed. The meteorological extreme indices are derived from an ensemble of general circulation models (GCMs) from
100 CMIP5 (Taylor et al., 2011) while our assessment of water availability and agricultural impacts at 1.5°C and 2°C is based on the ISI-MIP Fast Track data (Warszawski et al., 2013; Frieler et al., 2015).

For both data archives, the impacts for a GMT increase of 1.5°C and 2°C above pre-industrial levels are derived for 20-year time slices with the respective mean warming for each model separately. To account for model deviations from observations over the historical period, the warming
105 levels are derived relative to the reference period 1986-2005, (this reference period is 0.6°C warmer than pre-industrial levels (1850 — 1900) , IPCC (2013)), which translates to a warming of 0.9°C and 1.4°C above reference period levels for the 1.5°C and 2°C limit, respectively. All time slices are derived from the RCP8.5 scenario (the time slices for the individual GCMs are given in the SM Tab. S1). 1986-2005 is also the common reference period to assess projected changes in extreme indices
110 and climate impacts.

All our results are calculated with respect to this common reference period. For consistency with the respective policy targets, however, we express the GMT differences of 0.9°C and 1.4°C by their implied pre-industrial warming of 1.5°C and 2 °C.

Analyzing time-slices centered around a specific level of warming relies on the assumption that
115 the changes in the climate and climate impact signals studied here are dominantly driven by changes in GMT and that the effect of changes in time-lagged systems such as large-scale ocean circulations (Schleussner et al., 2014a, b) on the quantities studied are of minor importance. In addition, this approach does not account for the effect of other anthropogenic climate forcings that may differ for the same level of total radiative forcing such as aerosols (Zopa et al., 2012). It comes, however, also
120 with several advantages. In particular, it eliminates the spread due to different transient climate responses across the model ensemble, which can deviate by up to a factor of two (Flato et al., 2013). Traditional approaches that analyze impacts over a given time period for all models in a model ensemble and relate this to a median GMT increase across the model ensemble do not account for this ensemble-intrinsic spread of global warming levels and will consequently overestimate the ensemble
125 uncertainty of the GMT-dependent indices studied. The time-slice approach has furthermore been shown to provide better accuracy than traditional pattern scaled approaches (Herger et al., 2015). Although also relying on the debatable assumption of scenario-independence of the projected signals,

which does not fully hold in climate stabilization scenarios (Tebaldi and Arblaster, 2014), time-slicing avoids known short-comings of classical pattern scaling analysis. In particular, it allows to
130 capture non-linearities in extreme indices and precipitation related signals that relate to non-linear local feedbacks (Lopez et al., 2013) or large-scale circulation changes (Chadwick and Good, 2013; Hawkins et al., 2014).

In addition to the anthropogenic forcing, natural variability is a dominant driver of the climate signal on multi-annual time scales for time-averaged quantities such as mean temperature and precipitation change (Knutti and Sedláček, 2012; Marotzke and Forster, 2014) and in particular for
135 extreme weather events (Kendon et al., 2008; Tebaldi et al., 2011). Thus, natural variability may mask an already present climate change signal and consequently lead to a delayed detection of the imprints of climate change (Tebaldi and Friedlingstein, 2013). To overcome this limitation, Fischer et al. (2013) proposed a spatial aggregation approach that allows for a robust detection of an anthropogenic footprint in climatic extremes despite natural variability – an approach that has also
140 been successfully applied to the observational record (Fischer and Knutti, 2014). Here, we adopt and extend this spatial-aggregation approach.

As in Fischer et al. (2013), we consider the distribution of changes in the selected impact indicator at each grid point over the global land-mass between 66°N and 66°S (henceforth referred
145 to as global land-mass) and additionally analyze changes for 26 world regions (as used in IPCC, 2012, see Tab. 1 for details). This yields distributions for 1.5°C and 2°C for each of the ensemble members and regions, where the sample size is given by the number of grid points included in the respective regions. In a next step, the statistical significance of differences between the 1.5°C and 2°C distributions is assessed for each region and ensemble member. This is done using a two-sample
150 Kolmogorov–Smirnov (KS) test with the null hypothesis that both distributions for 1.5°C and 2°C are drawn from the same probability distribution.

A rejection of the test’s null-hypothesis at a significance level of 95 % is taken as a robust difference in projections between these two warming levels. This pairwise test, based on the individual model ensemble members analyzed, allows for robust statements about differences between the two
155 warming levels, even if there is substantial overlap of uncertainty bands in the model ensemble. For GCMs that provide multiple realizations, the distributions are combined for each warming level leading to larger samples and higher discriminatory power of the KS test. Please note that this approach is only applied for the KS test and not for the ensemble projections. For the latter, the averaged signal over multiple realizations of a single GCM is included in the ensemble analysis ensuring equal
160 weight to all GCMs investigated (see SM Section 1 for further detail on the methods and the treatment of multiple realizations). A similar approach has been applied recently to investigate the timing of anthropogenic emergence in simulated climate extremes (King et al., 2015).

Based on the regionally specific distributions, cumulative density functions (CDF) of changes in the impact indices over the land area of the respective region are derived. As in Fischer et al.

165 (2013), we fit a probability density function to the empirical distribution of the climate signal using
a Gaussian kernel density estimator. Individual grid-cells are weighted according to their latitude-
dependent area. These CDFs are derived for each ensemble member (GCM or GCM-impact model
combination) and the ensemble median as well the likely range (66 % of the ensemble members are
within this range) are given. This land-area focused approach allows to directly assess not only the
170 median change over a region, but also changes for smaller fractions of the land area. At the same
time, the uncertainty estimates based on the model ensemble spread can be directly visualized.

Table 1. Overview of the world regions used as well as the respective acronyms based on IPCC (2012). Please
note that the Central American (CAM) region has been extended eastwards to also include the Caribbean.

ALA	Alaska, North-West Canada	NEB	North East Brazil
AMZ	Amazon	NEU	North Europe
CAM	Central America, Mexico, Caribbean	SAF	South Africa
CAS	Central Asia	SAH	Sahara
CEU	Central Europe	SAS	South Asia
CGI	East Canada, Greenland, Iceland	SAU	South Australia, New Zealand
CNA	Central North America	SEA	South-East Asia
EAF	East Africa	SSA	South-East South America
EAS	East Asia	TIB	Tibetan Plateau
ENA	East North America	WAF	West Africa
MED	Mediterranean	WAS	West Asia
NAS	North Asia	WNA	West North America
NAU	North Australia	WSA	West Coast South America

3 Extremes Weather Events

There is a growing body of evidence showing that the frequency and intensity of many extreme
weather events has increased significantly over the last decades as a result of anthropogenic climate
175 change, but confidence in the significance of the trend and attribution to anthropogenic origin differ
substantially between types of extreme weather events and regions (IPCC, 2013). With on-going
warming, these trends are projected to continue (IPCC, 2012). Impacts of extreme weather events
will particularly, but not exclusively, affect the most vulnerable with the lowest levels of adaptive
capacity and represent one of the biggest threats posed by climate change (IPCC, 2014b). In this
180 Section, the difference in impacts between a warming of 1.5°C and 2°C for four different types
of meteorological extreme event indices are assessed. Good agreement between the CMIP5 model
ensemble median estimates of extreme event indices and observational indices data sets has been
reported by Sillmann et al. (2013a). The indices used follow the recommendations of the Expert

Team on Climate Change Detection and Indices (Zhang et al., 2011) and are derived on an annual

185 basis:

- Intensity of hot extremes (**TXx**): Annual maximum value of daily maximum temperature.
- Warm spell duration indicator (**WSDI**): Annual count of the longest consecutive period in which the daily maximum temperature for each day exceeds the 90 % quantile for this day over the reference period. The minimum length is six consecutive days.
- 190 – Dry spell length or consecutive dry days (**CDD**): Annual maximum number of consecutive days for which the precipitation is below 1mm per day.
- Heavy precipitation intensity or maximum accumulated five-day precipitation (**RX5day**): Absolute annual maximum of consecutive 5-day precipitation.

3.1 Methods and Data

195 Projected changes in climate extreme indices are assessed using an ensemble of 11 CMIP5-models for TXx and WSDI and 14 for RX5day and CDD and follows the methods outlined in Section 2. The model selection was done based on data availability. All available GCMs are assessed on a uniform grid with a 2.5° x 1.9° resolution. Multiple realizations of scenario runs for individual models are included when available, and in that case allow to estimate CDFs for natural variability that are
200 derived based on pairwise realizations of model runs over the reference period (see SM Section 1.2 for further detail on the methodology applied).

We assess the changes in TXx and WSDI for a warming of 1.5°C and 2°C and derive changes of 20-year averages of extreme indices for the model dependent warming-level time-slices at each land grid point relative to the 1986-2005 reference period. Changes in precipitation-related indices are
205 described as relative changes while we consider absolute changes for the other indicators. For the CDF analysis for TXx, the absolute signal is normalized by the standard deviation over the reference period.

3.2 Results

3.3 Heat Extremes

210 Substantial increases of 3°C and more in TXx over large parts of the Northern Hemisphere, Central South America and South Africa as well as increases in warm-spell durations (WSDI) of 3 months and more are projected under a warming of 2°C. Fig. 1 depicts changes in TXx (left) and WSDI (right) for a 1.5°C (top) and 2°C (middle) GMT temperature increase, as well as the differences between the two warming levels (bottom) on a grid-cell basis. Particularly strong increases in WSDI
215 are found in some tropical coastal areas, which we attribute to a large share of ocean surface in the respective grid cells that lead to an amplification of the effect compared to pure land grid cells and

should not be over-interpreted. We excluded these grid-cells for the CDF analysis for the respective regions. The majority of GCMs agree on a robust increase in these heat-related indices and show significant differences between the two warming levels. The impacts are robustly smaller at 1.5°C warming in both cases (see results for the KS test listed in Tab. S2).




Figure 1. Median changes of TXx (left panels) and WSDI (right panels) for a warming of 2°C (upper panels), 1.5°C (middle panels) and the difference between the two warming levels (lower panels). Changes in TXx are given in °C, whereas changes in WSDI are given in days.

Globally and regionally resolved CDFs for TXx, normalized to the standard deviation (σ) over the reference period, are given in Fig. 2 and median values are listed in Tab. S2. 50 % of the global land-mass will experience a median TXx increase of more than 1.2 (1.8) standard deviations relative to the reference period for a warming of 1.5°C (2°C) above pre-industrial levels. The regional assessments indicate that the tropical regions in Africa, South America and South East Asia are projected to experience the strongest increase in land area covered by heat extremes relative to the regional natural variability, where 3- σ events become the new normal under a 2°C warming.

The pattern of a strong tropical signal is mainly due to the small natural variability of TXx in tropical regions. This is also apparent for the WSDI CDFs resolved in Fig. 3. For a warming of 1.5°C, a median increase in WSDI length by about one month is projected for 50 % of the global land area that increases by 50 % for a 2°C warming. Since this index is derived relative to natural variability over a reference period, the signal again is greatly amplified in tropical regions, where a




Figure 2. CDFs for projected regional aggregated changes in TXx (relative to the standard variation over the reference period) for the global land area between 66 °N and 66 °S (lower left corner) as well as resolved for 26 world regions separately (see Section 2 for further details). Changes are given relative to the standard deviation over the 1986-2005 reference period. Note that a change in 2 (3) standard deviations implies that events with a reference return time of several decades (centuries) become the new normal, whereas a new normal of 4- σ refers to an event that would be extremely unlikely to occur in a reference period climate. Region impact overviews are provided in the Supplementary Material.

median WSDI of up to three month is projected for Amazonia, East and West Africa and South-East Asia (see Tab. S2). Given that the WSDI only measures the longest consecutive interval, such an increase can be interpreted as entering a new climate regime for these tropical regions (Diffenbaugh and Scherer, 2011; Mora et al., 2013; King et al., 2015).

A meaningful assessment of impacts requires not only an assessment of absolute changes, but these also have to be interpreted in the light of regional climate conditions. It is the regional natural climate variability that arguably determines a "climate normal" to which human systems as well as ecosystems might be adapted to (Hansen et al., 2012; Coumou and Robinson, 2013). While this may hold as a general assumption for a range of impacts concerning human health as well as ecosystems, it is important to note that the severity of certain climate impacts may also depend on the exceedance of absolute thresholds, as has been shown for temperature effects on crop yields, for example (Deryng et al., 2014; Smith et al., 2014). The choice of an either relative or absolute representation of changes in climate impacts thus has to be made in light of the impact of interest. In addition, a normalization by the standard deviation similar to the one applied here has been shown to introduce statistical biases arising from a limited sample size of the reference period (Sippel et al., 2015) that we do not account for in the results presented here.




Figure 4. Same as Fig. 1, but for RX5day and CDD. Hatched areas indicate regions, where less than 66 % of the models in the ensemble agree with the sign of change of the median projections.

Amazon region and North-East Brazil, where precipitation extremes are likely related to the South American monsoon systems (Boers et al., 2014) and to a lesser extent for West Africa (see Fig 5 and Tab. S3).

A different picture emerges for CDD as an indicator for dry spell length. For the majority of the global land area, little to no differences in CDD are projected relative to the reference period (see Fig. 4). However, about 40 % of the global land area in the subtropical and tropical regions experience an increase in CDD length, including the Mediterranean, Central America, the Amazon as well as South Africa (compare Fig. 4 and Fig. 6). In these regions, the KS test also reveals robust indications for differences in impacts between 1.5°C and 2°C. This difference is particularly pronounced for the Mediterranean region, where the median CDD length increases from 7 % (likely range 4 to 10 %) to 11 % (likely range 6 to 15 %) between 1.5°C and 2°C.

It is important to highlight that CDD is only an indicator for dry spell length and does not account for changes in evapotranspiration and soil-moisture related effects. It should hence not be interpreted as a direct indicator for agricultural or hydrological (streamflow) drought (Mueller and Seneviratne, 2012; Orlowsky and Seneviratne, 2012). Furthermore, CDD is a metric for short dry spells, which represent only a snapshot of the overall changes in dryness (IPCC, 2012), while high-impact drought events like the Big Dry in Australia (Kiem and Verdon-Kidd, 2010) or the recent California drought




Figure 5. Same as Fig. 2 but for RX5day. Changes are given relative to the 1986-2005 reference period.




Figure 6. Same as Fig. 2 but for CDD. Changes are given relative to the 1986-2005 reference period.

stretch over month and potentially years (Ault et al., 2014). Nevertheless, CDD as well as RX5day can be seen as proxies for the precipitation-related component when assessing drought and flooding risks, respectively, and the results and impacted regions identified here are broadly consistent with projections based on more comprehensive indicators for droughts (Dai, 2012; Prudhomme et al., 2013) and flooding risk (Hirabayashi et al., 2013) alike.

4 Water Availability

Already today, water scarcity is among the biggest challenges for ecosystems and human societies in many regions globally. To assess changes in water availability (assessed here as the annual mean surface and subsurface runoff – QTOT) at 1.5°C and 2°C, we follow the approach outlined above in
295 Section 2. Projections are based on 11 global hydrological models (GHM) that participated in the ISI-MIP intercomparison (Schewe et al., 2013). These are forced with bias-corrected climate simulations from five CMIP5 GCMs (HadGEM2-ES, IPSL-CM5A-LR, MIROC-ESM-CHEM, GFDL-ESM2M, and NorESM1-M, see Hempel et al. (2013) for further details on the bias-correction methodology applied). Each of GCM-GHM combinations is treated as an individual ensemble member resulting in
300 a N=55 ensemble as a basis for the KS tests described above. Unlike for the CMIP5 ensemble, only one realization of each experiment is available and as a consequence the effect of natural variability cannot be assessed. ISI-MIP impacts are assessed at a 0.5° by 0.5° resolution.




Figure 7. Median projections for changes in annual mean runoff for a warming of 2°C (upper panel), 1.5°C (middle panel) and the difference between both levels (lower panel) relative to the 1986-2005 reference period. The projections are based on the ISI-MIP GCM-GHM model ensemble. Grid cells where less than 66 % of all GCM-GHM pairs agree with the median sign of change are hatched out. Grid cells with an annual mean runoff of less than 0.05 mm/day are masked white.

For a warming of 2°C, reductions in water availability of up to 30 % are projected in several – mainly subtropical – regions, in particular affecting the Mediterranean, South Africa, Central and Southern South America and South Australia (Fig. 7). A relative increase in runoff is projected in much of the high northern latitudes, as well as in parts of India, East Africa and parts of the Sahel. While many of these findings are consistent with earlier studies (James and Washington, 2013; Schewe et al., 2013), some may depend on the five GCMs chosen here and may be less robust in larger CMIP5 GCM ensembles (Knutti and Sedláček, 2012).




Figure 8. Same as Fig. 2 but for total annual runoff. Changes are given relative to the 1986-2005 reference period.

Fig. 7 (lower panel) and Fig. 8 illustrate the difference between a 1.5°C and 2°C warming. Differences are most prominent in the Mediterranean region where the median reduction in runoff almost doubles from about 9 % (likely range: 4.5 % – 15.5 %) at 1.5°C to 17 % (8 % – 28 %) at 2°C. For several other world regions such as Central America and Australia, there is an increasing risk of substantial runoff reductions exceeding 30 % for the upper limit of the 66 % quantile, although projections are highly uncertain (Tab. S4 and Fig. 8). The differences between 1.5°C and 2°C are smaller for many other regions, but the KS-test reveals that they are statistically significant for all world regions assessed (Tab. S4). These runoff results are also consistent with the findings on precipitation related extremes presented in Section 3.4.

In addition to changes in fresh water availability as a consequence of changes in the hydrological cycle, saltwater intrusion resulting from rising sea-levels or extreme coastal flooding has to be considered (Werner et al., 2013). Although strongly dependent on local circumstances including regional water management and coastal protection, saltwater intrusion might present a substantial

challenge in particular for low-lying coastal areas and small island states (Jiménez Cisneros et al., 2014).

325 5 Assessing agricultural risks

5.1 Methods and Data

We assess future agricultural crop yields in a 1.5°C and 2°C warmer world for the four major staple crops maize, wheat, rice and soy based on projections from the ISI-MIP Fast Track database (Warszawski et al., 2013; Rosenzweig et al., 2013). Projections for agricultural production depend
330 on a complex interplay of a range of factors, including physical responses to soils, climate and chemical processes, or nutrient and water availability, but are also strongly determined by human development and management. The representation of these processes differs strongly between different agricultural models. While studies suggest an increase in productivity for some crops as a result of higher CO₂-concentrations, large uncertainties remain with regard to temperature sensitivity, nutrient and water limitations, differences in regional responses and also the interactions between
335 these different factors (Rosenzweig et al., 2013). According to their metabolic pathways of carbon fixation in photosynthesis, main crops can be categorized as C3 and C4 plants. C4 plants such as maize, sorghum and sugar cane have a high CO₂ efficiency and as a consequence profit little from increased CO₂-concentrations, whereas for C3 plants including wheat, rice and soy a positive CO₂
340 - fertilization effect is to be expected. At the same time, increased CO₂-concentrations may lead to improved water use efficiency (Eamus, 1991). However, the effect of elevated CO₂ concentrations on plant growth is highly uncertain (McGrath and Lobell, 2013) and the representation of this effect greatly differs between different agricultural models. As a consequence, the ISI-MIP protocol has been conducted with and without accounting for CO₂-fertilization effects (further referred to
345 as the CO₂-ensemble and noCO₂-ensemble, respectively). Recent findings also underline the importance of elevated temperatures and heat extremes (Gourdji et al., 2013; Deryng et al., 2014), ozone concentrations (Tai et al., 2014) as well as the potential of increasing susceptibility to disease as a consequence of elevated CO₂ levels (Vaughan et al., 2014) for agricultural yields, which may counteract potential yield gains by CO₂-fertilization (Porter et al., 2014). Results for the CO₂ and
350 noCO₂-ensembles are presented separately, showing the range of potential manifestations and the additional risks of regional yield reductions, if effects of CO₂-fertilization turn out to be lower than estimated by the model ensemble.

The ISI-MIP ensemble contains simulations based on seven Global Gridded Crop Models (GGCM) for wheat, maize and soy and six GGCM for rice, run with input from five CMIP5 GCMs (for further information see Rosenzweig et al., 2013). For the CO₂-ensemble, all model combinations are
355 available (35, and 30 for rice), while for the noCO₂-ensemble runs have been provided for 23 (18 for rice) GGCM-GCM combinations. We restrict future crop growing areas to present day agricultural

areas (based on Monfreda et al., 2008) and assume no change in management type, meaning that "rainfed" and "irrigation" conditions are kept constant as well.

As in previous Sections, the results presented here are based on 20-year time slices from the RCP8.5 simulations and changes are given relative to the 1986-2005 reference period. The choice of displaying relative changes comes with several advantages, but will also lead to a disproportional visual amplification of minor absolute changes for regions with small present day yields, in particular in the high northern latitudes. An overview of the regionally resolved present day share in global production is given in Fig. S5.

Since agricultural impacts depend both on climatological changes and changes in the atmospheric CO₂-concentrations, the assumption of time-independent impacts underlying the time-slice-approach as discussed above does not fully hold for agricultural projections accounting for the effects of CO₂-fertilization (the CO₂-ensemble) and will lead to increased inner-ensemble spread as a consequence.

Please note that the regional aggregation for agricultural yields is not based on absolute yield change but on land area, as for the other indicators studied above. Since societal impacts of changes in agricultural production go beyond mere changes in yield, but also include for example local livelihood dependencies (Schellnhuber et al., 2013; Olsson et al., 2014), our assessment of local yield changes (on a grid-cell level) supplements and extends previous yield-centered analysis (Rosenzweig et al., 2013). Maps for the projected differences of yield changes on a grid-cell basis are provided in the supplementary information.

5.2 Results

5.2.1 Wheat

Our analysis reveals very small local median yield changes for 50 % of the global land area for a 1.5°C and 2°C warming. However, the uncertainties of these projections are substantial and reductions of about 6 % and 8 % for 1.5°C and 2°C, respectively, mark the lower end of the likely range (compare Tab. S5). For the noCO₂-ensemble, we find substantial median reductions in local wheat yields of 14 % at 1.5°C, with a statistically significant higher decrease of 19 % at 2 °C and potential reductions of up to 20 % (1.5°C) and 37 % (2°C) as lower limits for the likely range. The results of the KS-tests based on individual model combinations are given in Tab. S5 and for the global level as well as most regions, more than 83 % (90 %) of all ensemble members indicate a robust difference between projected impacts at 1.5°C and 2°C for the CO₂ (noCO₂)-ensemble.

Best estimate local yield reductions are projected for the tropical region of about 9 % (15 %) for 1.5°C (2°C) that are particularly pronounced in West African (median reduction of 13 % (19 %)). Under a 1.5 °C (2°C) warming, reductions of up to 25 % (42 %) are within the likely range of the CO₂ ensemble projections and for the noCO₂-ensembles, median reductions of 28 % (35 %) would be projected.




Figure 9. Same as Fig. 2 but for changes in wheat yields. Changes are given relative to the 1986-2005 reference period and ensemble projections excluding the effect of CO₂-fertilization are shown separately. The CDFs are derived only over the present day growing areas of the crop.

5.2.2 Maize

The effects of elevated CO₂ concentrations affect maize yields to a much lesser extent, as conditions are mostly saturated at present levels (see e.g. Leakey et al., 2006). Differences between runs are thus less pronounced for maize yields, where yield reductions are projected for both the CO₂ and the noCO₂-ensemble. As the number of runs differ between the two ensembles (see Methods), the small differences are likely due to the different ensemble size. Thus, we only discuss results for the CO₂-ensemble here. Differences between the warming levels are significant (all ensemble members indicate a significant difference for the global crop area, see Tab. S6), with median local yield reductions experienced by 50 % of the global crop area of around 1.5 % and 6 % for 1.5°C and 2°C warming, respectively. Risks of reductions of up to 26 % at 1.5°C and 38 % at 2°C are within the likely range globally (compare Fig. 9 and Tab. S6).

As apparent in Fig. 9, the likely range is deferred towards stronger reductions. Similar regional patterns compared to wheat projections are apparent. Again, the highest relative median changes occur in regions with a relatively low share of global production. For central North America, where at present about 10 % of global maize is produced, substantial differences between the two warming levels are projected, and risks for a strong negative effect in this region more than double between 1.5°C and 2°C warming from 15.5 % to 37 % (upper limit of the 66 % range). Tropical regions such as Central America, the Amazon and South-East Asia are projected to experience median local yield

reductions exceeding 5 % for 1.5°C and up to and more than 10 % for 2 °C, while projections for the full tropical region do not differ substantially from the global projections.




Figure 10. Same as Fig. 9, but for changes in maize yields.

5.2.3 Soy

Projections of changes in soy yields between the two assessed warming levels show robust differences (see Tab. S7). For the CO₂-ensemble, a median increase in global yields of 7 % is projected for 50 % of the global area under a warming of 1.5°C. This median increase vanishes for 2°C. Global differences between warming levels for the noCO₂-ensemble are smaller but nonetheless robust, with median reductions of 10 % and 12 %, respectively.

Regionally, the differences for the noCO₂-ensemble are more pronounced, especially in those regions with a large share in present-day global soy production. Median yields for the Amazon (AMZ) region, currently producing about 7 % of global soy (Monfreda et al., 2008, see also Fig. S5), are projected to reduce from 15 % under 1.5°C to 20 % under 2°C warming. Similar robust differences in yield reductions between 1.5°C and 2°C warming are also projected for the major soy producers in Central North America and South-East South America. For North Asia, where currently over 7 % of soy production takes place, median increases in yields of 28 % and 24 % are projected for a warming of 1.5°C for the noCO₂ and CO₂-ensembles, respectively. However, uncertainties for this region are high and a risk of substantial reductions of 25 % (1.5°C) and 20 % (2°C) in the CO₂-ensemble are within the likely range of the ensemble projections.




Figure 11. Same as Fig. 9, but for changes in soy yields.

5.2.4 Rice

430 Median changes in global rice yields for the CO₂-ensemble do not differ between the assessed warming levels, with projected increases of about 7 % although the respective local yield change distributions are significantly different (compare Tab. S8). The distribution of possible developments indicates risk of substantial reductions of up to 17 % and 14 % at 1.5°C and 2°C. For the noCO₂-ensemble, reductions of 8 % and 15 % are projected for the two warming levels.

435 The effects of CO₂-fertilization consistently indicates yield increases across regions for median projections. While differences between warming levels are apparent for some regions and the CO₂-ensemble, global estimates are very similar between both warming levels. For the noCO₂-ensemble, robust differences between 1.5°C and 2°C warming are apparent for all major rice producing regions, including all Asian regions where a total of 40 % of rice is produced today (EAS, SAS, SEA, TIB)
440 as well as the Amazon, and South American rice producers. Reductions are projected to double between the two warming levels, for example in South Asia, South-East South America and the Tibetan Plateau. For these regions, median projections are close to the lower end of the likely range (compare Fig. 12 and Tab. S8).

5.3 Discussion of agricultural impact projections

445 Our projections of local agricultural yields reveal substantial uncertainties in global median regional yield changes (Figs. 9 to 12) with a likely range (66 % - likelihood) comprising zero. For wheat, rice and soy, our projections indicate differences between the CO₂ and noCO₂ assessments, which are generally much larger than those between a 1.5°C and 2°C warming. While substantial uncertainty




Figure 12. Same as Fig. 9 but for changes in rice yields.

renders a differentiation between impacts at 1.5°C and 2°C warming difficult in most world regions, a clear signal emerges for the noCO₂-ensemble, that may serve as a high-risk illustration of potential climate impacts on agricultural production. In the noCO₂-ensemble, local yields are projected to decrease between 1.5°C and 2°C for all crop types.

Our results indicate that risks are region and crop specific and are in line with findings of previous model intercomparison studies (Asseng et al., 2014; Rosenzweig et al., 2013). While high-latitude regions may benefit, median projections for local yields in large parts of the tropical land area are found to be negatively affected already at 1.5°C. Risks increase substantially, if effects of CO₂-fertilization are less substantial or counter-acted by other factors such as extreme temperature response, land degradation or nitrogen limitation (Rosenzweig et al., 2013; Bodirsky and Müller, 2014; Bodirsky et al., 2014). In a statistical analysis of climate impacts on wheat and barley yields in Europe, Moore and Lobell (2015) report an overall negative contribution of climatic factors in line with findings of a meta-analysis by Asseng et al. (2014), which questions the positive effects projected in our CO₂-ensemble for this region and further support our approach of singling out noCO₂-ensemble projections. Given that a 1.5°C warming might be reached already around 2030, our findings underscore the risks of global crop yield reductions due to climate impacts outlined by Lobell and Tebaldi (2014), while giving further indications for the regional diversity of climate impacts with tropical regions being a hot-spot for climate impacts on local agricultural yields (Müller et al., 2014).

6 Sea-level rise

6.1 Methods

470 Projections for sea-level rise (SLR) cannot be based on a time-slice approach because of the importance of the time-lagged response of the ocean and cryosphere to the warming signal. Therefore, we selected two multi-gas scenarios illustrative of a 1.5°C and 2°C warming to assess SLR impacts over the the entire 21st century from a large emission scenario ensemble created by Rogelj et al. (2013). These scenarios were created with the integrated assessment modeling framework MESSAGE (the
475 Model for Energy Supply Strategy Alternatives and their General Environmental Impact, Riahi et al., 2007). For both scenarios, temperature projections are derived with the reduced complexity carbon-cycle and climate model MAGICC (Meinshausen et al., 2011) in a probabilistic setup (Meinshausen et al., 2009), which has been calibrated to be in line with the uncertainty assessment of equilibrium climate sensitivity of the IPCC AR5 (Rogelj et al., 2012, 2014). Each probabilistic setup ensemble
480 consists of 600 individual scenario runs. The first scenario keeps GMT to below 2°C relative to pre-industrial levels (1850-1875) during the 21st century with 50 % probability. The second scenario reduces emissions sooner and deeper, and keeps warming to below 1.5°C relative to pre-industrial levels during the 21st century with about 50 % probability and returns end-of-century warming to below 1.5°C with about 70 % probability. See Fig. 13 (upper panel) for median temperature projec-
485 tions for the 2°C and 1.5°C scenario and their associated uncertainty bands. Since the projections for coral reef degradation include a time dependent adaptation scenario, the same approach is taken for the coral reef projections (see Section 7).

SLR projections are based on Perrette et al. (2013), who developed a scaling approach for the various SLR contributions according to an appropriately chosen climate predictor – in this case
490 GMT increase and ocean heat uptake. Coupled with output from the MAGICC model, this allows to emulate the sea-level response of GCMs to any kind of emission scenario within the calibration range of the method that is spanned by the RCPs.

Consistent with the relationship found in CMIP3 and CMIP5 GCMs, ocean thermal expansion is assumed to be proportional to cumulative ocean heat uptake (Church et al., 2013). Mountain glacier
495 melt is computed following a widely-used semi-empirical relationship between rate of glacier melt, remaining surface glacier area, and temperature anomaly with respect to pre-industrial levels. This approach assumes constant scaling between area and volume (Wigley and Raper, 2005; Meehl et al., 2007), with parameters chosen to account for current melt rate and known glacier volume (Eq. 1 and Table 2 in Perrette et al., 2013). As already noticed by Gregory and Huybrechts (2006) (their
500 Fig. 5), the surface mass balance (SMB) anomaly from the Greenland ice-sheet can be approximated with reasonable accuracy as a quadratic fit to global mean temperature anomaly. Here we adopted the same functional form, but calibrated it to more recent projections by Fettweis et al. (2013). Following Hinkel et al. (2014), we scaled up these projections by $20\% \pm 20\%$ to account for missing

dynamic processes (elevation feedback $10\% \pm 5\%$, changes in ice dynamics $10\% \pm 5\%$, and $\pm 10\%$ arising from the skill of the SMB model to simulate the current SMB rate over Greenland). The climate-independent land-water contribution has been added for all scenarios following Wada et al. (2012).

Beyond the scaling approach, the main advancement of our approach compared to the IPCC AR5 (Church et al., 2013) stems from the inclusion of scenario-dependent Antarctic ice-sheet projections following Levermann et al. (2014). Linear response functions were derived from idealized step-forcing experiments from the SeaRISE project (Bindshadler et al., 2013) as a functional link between the rate of ice shelf melting and dynamical contribution to SLR over four Antarctic sectors and various ice-sheet models. Levermann et al. (2014) further assume linear scaling between global surface air warming, local ocean warming, and ice-shelf melting in each of the sectors. They adopted a Monte Carlo approach with 50,000 samples to combine the various parameter ranges, GCMs and ice-sheet models. To our knowledge, this is the most comprehensive attempt to date to link climate warming and Antarctic ice-sheet contributions to scenario-dependent sea-level rise over the 21st century.

6.2 Results

The results for the 2°C scenario are comparable with projections by Church et al. (2013) and Hinkel et al. (2014) for RCP2.6 (their Tables 13.5 and 4, respectively) that leads to a median GMT increase of about 1.6°C above pre-industrial levels. For this scenario, we project a median SLR of about 50 cm (36 – 65 cm, likely range) by 2100 and a rate of rise of 5.6 (4 – 7) mm/year over the 2081-2100 period. Under our illustrative 1.5°C scenario, projected SLR in 2100 is about 20 % (or 10cm) lower, compared to the 2°C scenario (See Tab. 2). The corresponding reduction in the expected rate of SLR over the 2081-2100 period is about 30 %. More importantly, and in contrast to the projections for the 2°C scenario, the rate for the 1.5°C scenario is projected to decline between mid-century and the 2081-2100 period by about 0.5 mm/year, which substantially reduces the multi-centennial SLR commitment (Schaeffer et al., 2012).

The projected difference in SLR between the 1.5°C and 2°C scenarios studied here is comparable to the difference between the RCP2.6 and RCP4.5 scenarios (Hinkel et al., 2014; Church et al., 2013), while the projected median GMT difference between the two RCP scenarios is about 0.8 °C for the 2081-2100 period. The relatively higher sensitivity of SLR in the 21st century to temperature increase at low climate warming is probably related to the earlier peaking of GMT under such scenarios and thus an already longer adjustment period for the time-lagged ocean and cryosphere. This leads to a larger share of committed multi-centennial SLR to occur in the 21st century. On multi-centennial timescales these scenario dependent differences are expected to vanish. A long-term difference, however, may arise from contributions by mountain glacier melt, which are particularly

vulnerable to GMT increase and thus differences in melted mountain glacier volume are higher for lower emission scenarios.

While SLR projections for the two illustrative 1.5°C and 2°C differ substantially, this effect is strongly scenario dependent. In particular, most emission pathways labeled as 1.5°C scenarios allow for a temporal overshoot in GMT and a decline below 1.5°C with a 50 % probability by 2100 (Rogelj et al., 2015), whereas the illustrative 1.5°C scenario used here does not allow for a GMT overshoot, but stays below 1.5°C over the course of the 21st century. For time-lagged climate impacts such as SLR that depend on the cumulative heat entry in the system, the difference between a scenario allowing for a GMT overshoot and one that does not will be significant.

Sea-level adjustment to climate warming has a time scale much larger than a century as a result of slow ice-sheet processes and ocean heat uptake. This means that in all emission scenarios considered, sea level will continue to rise beyond 2100. Levermann et al. (2013) have shown that on a 2000-year time-scale, sea-level sensitivity to global mean temperature increase is about 2.3m per °C. In addition to that, Levermann et al. (2013) report a steep increase in long-term SLR between 1.5°C and 2°C as a result of an increasing risk of crossing a destabilizing threshold for the Greenland ice-sheet (Robinson et al., 2012). The disintegration process that would lead to 5–7 m global SLR, however, is projected to happen on the time scale of several millennia.

Recent observational and modeling evidence indicates that a marine ice-sheet instability in the West Antarctic may have already been triggered, which could lead to an additional SLR commitment of about 1 m on a multi-centennial time scale. Spill-over effects of this destabilization on other drainage basins and their relation to GMT increase are as yet little understood (Rignot et al., 2014; Joughin et al., 2014; Favier et al., 2014) and there are indications that a destabilization of the full West Antarctic ice-sheet could eventually be triggered (Feldmann and Levermann, 2015). Similarly, Mengel and Levermann (2014) report a potential marine ice-sheet instability for the Wilkens Basin in West Antarctica containing 3–4 m of global SLR. The dynamics of these coupled cryosphere-oceanic systems remain a topic of intense research. Current fine-scale ocean models, suggest increased intrusion of warm deep water on the continental shelf as a result of anthropogenic climate change and thus indicate an increasing risk with increasing warming (Hellmer et al., 2012; Timmermann and Hellmer, 2013). Given the risk of potentially triggering multi-meter SLR on centennial to millennial time scales, this clearly calls for a precautionary approach that is further underscored by evidence from paleo-records, which reveals that past sea-levels might have about 6–9 m above present day for levels for a GMT increase not exceeding 2°C above pre-industrial levels (Dutton et al., 2015).




Figure 13. Upper panel: Probabilistic GMT projections for illustrative emission scenarios with a peak warming of 1.5°C (left panels) and 2°C (right panels) above pre-industrial levels during the 21st century. Lower panels: Probabilistic projections of global sea-level rise (SLR) for both scenarios relative to 1986-2005 levels. Uncertainty bands indicate the likely range (66 % probability within this range) and the very likely range (90 % probability), respectively.

Table 2. Projections for sea-level rise above the year 2000-levels for two illustrative 1.5°C and 2°C scenarios (see Fig. 13). Square brackets give the likely (66 %) range.

	1.5°C	2°C
SLR 2081-2100 [m]	0.37 [0.27,0.48]	0.44 [0.32,0.57]
SLR 2100 [m]	0.41 [0.29,0.53]	0.5 [0.36,0.65]
Rate of SLR 2041-2060 [mm/year]	4.6 [3.2,5.8]	5.6 [4.0,7.0]
Rate of SLR 2081-2100 [mm/year]	4.0 [2.7,5.5]	5.6 [3.8,7.7]

7 Coral reef systems

7.1 Methods

The projections of the degradation of coral reef sites uses the coral bleaching model developed in Frieler et al. (2012) based on the two illustrative 1.5°C and 2°C global emission pathways introduced
575 in Section 6.1. The framework applies a threshold-based bleaching algorithm by Donner (2009), which is based on degree heating months (DHMs), to sea surface temperature (SST) pathways of

2160 individual geospatial locations of coral reef sites (see www.reefbase.org) and generates as output the fraction of coral reef locations subject to long-term degradation. DHMs are a measure for the accumulated heat stress exerted on coral reefs due to elevated SST (see Fig. S6 for a graphical illustration of the methodology). Within a four months moving window the monthly SST above a reference value (here the mean of monthly maximal temperatures, MMM) are accumulated and compared to a threshold value (critical DHM threshold) that is associated with mass coral bleaching. The value of the critical DHM threshold depends on the scenario assumptions (see below). In order to translate coral bleaching events into long-term coral degradation, we refer to the assumption that reef recovery from mass coral bleaching is usually very limited within the first five years (Baker et al., 2008). Therefore, we assume a maximum tolerable probabilistic frequency of $0.2yr^{-1}$ (Donner, 2009) for bleaching events causing long-term degradation. The MMM is calculated from a 20-year climatological reference period (1980-2000) individually for every coral location and SST pathway. Thus, the MMM serves as an indicator of temperatures to which the corals of a certain reef location are generally adapted. In order to generate a scenario-independent description of coral reef response to different levels of global warming (e.g. any given global mean air temperature pathway) we apply the algorithm to a large number of SST pathways and reassign the fraction of 2160 mapped coral reef locations subject to long-term degradation back to global air temperature pathways. In total, we use the SST pathways of 19 Atmosphere-Ocean General Circulation Models (AOGCMs) from the multi-model CMIP3 project and seven different emission scenarios leading to 30,728 model years. We also used a wide range of critical DHMs (from 0° to 8°), which allows for the testing of risk scenarios with constant and variable critical DHM thresholds (e.g. thermal adaptation).

The condensed output of the global coral bleaching assessment allows for the implementation of different coral adaptation scenarios. In the standard scenario (*Constant*) a constant DHM threshold of 2°C is assumed. This means that corals can resist a cumulative heat stress of 2°C (accumulated over four month period) above the long-term maximum monthly mean (MMM) sea surface temperature for a given location. It has been demonstrated that this value serves as a good proxy for severe mass coral bleaching (Donner et al., 2005, 2007).

In addition to the constant scenario, an extremely optimistic scenario of strong thermal adaptation of the corals is assessed (*Adaptation*). Under this scenarios, the critical DHM threshold constantly increases from 2°C in the year 2000 up to 6°C in 2100. The assumption of a thermal adaptation of 0.4° per decade appears very ambitious given the long creation times of reef-building corals and the consequently slow rate at which evolutionary adaptation occurs. Furthermore, additional environmental stressors such as ocean acidification (Caldeira, 2005) and disease spreading (Maynard et al., 2015) have to be expected to slow-down coral growth and to reduce the adaptive capacity of tropical coral reefs. As a consequence, this scenario should be seen as an absolute lower boundary for degradation of coral reefs globally.

Finally, a third scenario takes the negative effect of the acidification of the oceans into account which reduces the calcification rates of the corals and thus promotes further degradation of coral reefs (*Saturation*). We derived a transfer function based on atmospheric CO_2 concentrations due to the fact that tropical surface aragonite saturation levels are in equilibrium with atmospheric CO_2 concentrations on a timescale of years to decades (Caldeira, 2005). With an assumption of the effect of the aragonite saturation on the critical DHM threshold (see supporting material of Frieler et al. (2012)) this translates into a measurable increased stress to corals.

7.2 Results

Coral reef systems are slow-growing, complex ecosystems that are particularly susceptible to the impacts of increased CO_2 concentrations, both through warming (and resulting coral bleaching) and ocean acidification (Pörtner et al., 2014). Our analysis re-iterates earlier findings that the risk of coral reefs to suffer from long-term degradation eventually leading to an ecosystem regime shift (Graham et al., 2015) will be substantial as early as 2030 (Meissner et al., 2012; Gattuso et al., 2015; Frieler et al., 2012). We find that this risk increases dramatically until the 2050s, where even under a $1.5^\circ C$ scenario, 90 % and more of all global reef grid cells will be at risk of long-term degradation under all but the most optimistic scenario assessed (the *Adaptation* case, see Sec. 7.1). However, long-term risks towards the end of the century are reduced to about 70 % of global coral reef cells under a $1.5^\circ C$ scenario but not under a $2^\circ C$ scenario (compare Fig. 14 and Tab. 3).

Our approach only includes the effects of increased CO_2 -concentrations, but does not account for other stressors for coral reef systems such as rising sea-levels, increased intensity of ENSO-events (Power et al., 2013), tropical cyclones (Knutson et al., 2010), invasive species and disease spreading (Maynard et al., 2015), and other local anthropogenic stressors, which ranks our projections of long-term coral reef degradation rather conservative. These projected losses will greatly affect societies, which depend on coral reefs as a primary source of ecosystem services e.g. in the fishery and tourism sector (Cinner et al., 2015). Teh et al. (2013) estimate that about 25 % of the world's small-scale fishers fish on coral reefs. Chen et al. (2015) report that a loss of less than 60 % of global coral reef coverage, that could very well be reached already in the 2030s, would inflict damages of more than US\$ 20 billion annually.

8 Discussion and Conclusions

The findings of our analysis support the IPCC AR5 Working Group 2 RFC assessment of differences in key impacts of climate change between warming of $1.5^\circ C$ and $2^\circ C$ above pre-industrial levels: we find that under a 1.5° scenario, the fractions of coral reef cells at risk of severe degradation are reduced significantly compared to a warming of $2^\circ C$ (RFC1), that the difference between $1.5^\circ C$ and $2^\circ C$ marks the transition between an upper limit of present-day natural variability and a new climate




Figure 14. Probabilistic projections of the fraction of global tropical coral reef cells suffering from long-term degradation under two illustrative 1.5°C (upper panel) and 2°C (lower panel) scenarios (see Fig. 13 upper panel) for two different assumptions about their adaptive capacity (see Section 7.1). Median projections and the 66 % range are shown. Note that uncertainties also include uncertainties in the GMT response (see Fig. 13). See Section 7.1 for further details on the methodology. Only the projections for the *Constant* and *Adaptation* scenario are shown, since the projections for the *Saturation* scenario differ only slightly from *Constant*. Tab. 3 gives results for all three scenarios assessed.

Table 3. Fraction of reef cells at risk of long-term degradation due to coral bleaching in 2050 and 2100 for three different assumptions about the adaptive capacity and susceptibility of corals to ocean acidification as described in Section 7.1 in percent. Median projections and the 66 % range (in square brackets) are given, accounting also for uncertainties in global mean temperature projections..

	1.5°C	2°C
2050		
Adaptation	9 [2,49]	39 [8,81]
Saturation	94 [60,100]	100 [95,100]
Constant	89 [48,99]	98 [86,100]
2100		
Adaptation	1 [0,2]	6 [1,50]
Saturation	69 [14,98]	100 [91,100]
Constant	69 [14,98]	99 [85,100]

regime in terms of heat extremes globally (RFC2), and that changes in water availability and local agricultural yields are already unevenly distributed between world regions at 1.5°C and even more

so at 2°C (RFC3). Central findings across the different indicators studied are summarized in Fig. 15

650 and regional summaries are given in the Supplementary Material (Fig. S.7–33)

Water availability reduction and dry spell length (CDD) increase are found to accelerate between 1.5°C and 2°C for several sub-tropical regions, in particular in the Mediterranean, Central America and the Caribbean, South Africa and Australia. Local agriculture production in tropical regions is projected to be strongly affected by ongoing warming, in particular, if effects of CO₂ -fertilization do
655 not play out as current models project them or are counter-balanced by other factors such as nitrogen and phosphor limitations or heat stress, which are not fully included in the models investigated here.

Our analysis of projected SLR reveals differences of about 10cm in global mean SLR between illustrative 1.5°C and 2°C scenarios by 2100. In addition, the end-of-century rate of sea-level rise for 1.5°C is about 30 % lower than for a 2°C pathway, indicating a substantially lower long-term sea-level rise commitment (Clark et al., 2016). Evidence from the paleo-record (Dutton et al., 2015) and
660 modeling studies (Levermann et al., 2013) further indicate that a multi-meter sea-level of potentially up to 9 m cannot be ruled out under a 2°C warming on multi-millennial time scales.

Our assessment based on this limited set of indicators implies that differences in climate impacts between 1.5°C and 2°C are most pronounced for particularly vulnerable regions and societal group-
665 ings with limited adaptive capacity (Olsson et al., 2014). Under a 2°C warming, coastal tropical regions and islands may face the combined effects of a near-complete loss of tropical coral reefs, which provide coastal protection and are a main source of ecosystem services, on-going sea-level rise above present day rates over the 21st century and increased threats by coastal flooding and inundation. The risks posed by extreme heat and potential crop yield reductions in tropical regions in
670 Africa and South East Asia under a 2°C warming are particularly critical given the projected trends in population growth and urbanization in these regions (O'Neill et al., 2013). In conjunction with other development challenges, the impacts of climate change represent a fundamental challenge for regional food security (Lobell and Tebaldi, 2014) and may trigger new poverty traps for several countries or populations within countries (Olsson et al., 2014).

Furthermore, the emergence of the Mediterranean region, including North Africa and the Levant, as a hot-spot for reductions in water availability and dry spell increases between 1.5°C and 2°C is of great relevance given the specific vulnerability of this region to water scarcity (Schellnhuber et al., 2014). The political instability in several countries in this region may further exacerbate the vulner-
675 ability of societies to climatic stresses, potentially increasing the risk of violent conflict outbreak (Kelley et al., 2015).

Taken together, we provide a consistent and comprehensive assessment of existing projections and a solid foundation for future work on refining our understanding of the difference between impacts at 1.5 °C and 2 °C warming. In particular, we illustrate how limiting warming to 1.5°C would "signif-
685 ically reduce the risks and impacts of climate change" as stated in the Paris Agreement. However, our analysis can only be a first step towards a more integrative post - Paris science agenda including

		1.5°C	2°C	
Heat wave (warm spell) duration [month]				
	Global	1.1 [1,1.3]	1.5 [1.4,1.8]	Tropical regions up to 2 (3) month at 1.5°C (2°C) warming
Reduction in annual water availability [%]				
	Mediterranean	9 [5,16]	17 [8,28]	Other dry subtropical regions like Central Amerika and South Africa also at risk
Increase in heavy precipitation intensity [%]				
	Global	5 [4,6]	7 [5,7]	Global increase in intensity due to warming, high latitudes (>45°N) and monsoon regions affected most.
	South Asia	7 [4,8]	10 [7,14]	
Global sea-level rise				
	in 2100 [cm]	40 [30,55]	50 [35,65]	1.5°C end-of-century rate about 30% lower than for 2°C reducing long-term SLR commitment.
	2081-2100 rate [mm/yr]	4 [3,5.5]	5.5 [4,8]	
Fraction of global coral reefs at risk of annual bleaching [Constant case, %]				
	2050	90 [50,99]	98 [86,100]	Only limiting warming to 1.5°C may leave window open for some ecosystem adaptation.
	2100	70 [14,98]	99 [85,100]	
Changes in local crop yields over global and tropical agricultural area [%]				
Wheat <small>local yield changes relative to 1981-2010</small>	Global	2 [-6,17]	0 [-8,21]	Projected reduction risks are largest for tropical regions, while high-latitude regions may benefit. Projections not including highly uncertain positive effects of CO ₂ -fertilization project reductions for all crop types of about 10% already at 1.5°C and further reductions at 2°C.
	Tropics	-9 [-25,12]	-16 [-42,14]	
Maize <small>local yield changes relative to 1981-2010</small>	Global	-1 [-26,8]	-6 [-38,2]	
	Tropics	-3 [-16,2]	-6 [-19,2]	
Soy <small>local yield changes relative to 1981-2010</small>	Global	7 [-3,28]	1 [-12,34]	
	Tropics	6 [-3,23]	7 [-5,27]	
Rice <small>local yield changes relative to 1981-2010</small>	Global	7 [-17,24]	7 [-14,27]	
	Tropics	6 [0,20]	6 [0,24]	

Figure 15. Summary of key differences in climate impacts between a warming of 1.5°C and 2°C above pre-industrial and stylized 1.5°C and 2°C scenarios over the 21st century. Square brackets give the likely (66 %) range.

the assessment of below 1.5°C impacts and requirements and costs of energy system transformation pathways in line with limiting warming to below 1.5°C (Rogelj et al., 2015).

Acknowledgements. We acknowledge the World Climate Research Programme's Working Group on Coupled Modelling, which is responsible for CMIP, and we thank the climate modeling groups for producing and making available their model output. For CMIP, the US Department of Energy's Program for Climate Model Diagnosis and Intercomparison provided coordinating support and led development of software infrastructure in partnership with the Global Organization for Earth System Science Portals. We would like to thank the modelling groups that participated in the Inter-Sectoral Impact Model Intercomparison Project (ISI- MIP).

The ISI-MIP Fast Track project underlying this paper was funded by the German Federal Ministry of Education and Research with project funding reference number 01LS1201A. The work was supported by the German Federal Ministry for the Environment, Nature Conservation and Nuclear Safety (11-II-093-Global-A SIDS and LDCs), within the framework of the Leibniz Competition (SAW-2013-PIK-5), from EU FP7 project HELIX (grant no. FP7-603864-2), and by the German Federal Ministry of Education and Research (BMBF, grant no. 01LS1201A1)

700 References

- Asseng, S., Ewert, F., Martre, P., Rötter, R., Lobell, D., Cammarano, D., Kimball, B. A., Ottman, M., Wall, G., White, J., Reynolds, M., Alderman, P., Prasad, P., Aggarwal, P., Anothai, J., Basso, B., Biernath, C., Challinor, A., De Sanctis, G., Doltra, J., Fereres, E., Garcia-Vila, M., Gayler, S., Hoogenboom, G., Hunt, L., Izaurralde, R., Jabloun, M., Jones, C., Kersebaum, K., Koehler, A.-K., Müller, C., Naresh Kumar, S., Nendel, C., O'Leary, G., Olesen, J., Palosuo, T., Priesack, E., Eyshi Rezaei, E., Ruane, A., Semenov, M., Shcherbak, I., Stöckle, C., Stratonovitch, P., Streck, T., Supit, I., Tao, F., Thorburn, P., Waha, K., Wang, E., Wallach, D., Wolf, J., Zhao, Z., and Zhu, Y.: Rising temperatures reduce global wheat production, *Nature Climate Change*, 5, 143–147, doi:10.1038/nclimate2470, 2014.
- 705 Ault, T. R., Cole, J. E., Overpeck, J. T., Pederson, G. T., and Meko, D. M.: Assessing the risk of persistent drought using climate model simulations and paleoclimate data, *Journal of Climate*, 27, 7529–7549, doi:10.1175/JCLI-D-12-00282.1, 2014.
- 710 Baker, A. C., Glynn, P. W. and Riegl, B.: Climate change and coral reef bleaching: An ecological assessment of long-term impacts, recovery trends and future outlook, *Estuarine, Coastal and Shelf Science*, 435–471, 2008. doi:10.1007/s00382-012-1408-y, 2012.
- 715 Bindshadler, R. A., Nowicki, S., Abe-OUCHI, A., Aschwanden, A., Choi, H., Fastook, J., Granzow, G., Greve, R., Gutowski, G., Herzfeld, U., Jackson, C., Johnson, J., Khroulev, C., Levermann, A., Lipscomb, W. H., Martin, M. A., Morlighem, M., Parizek, B. R., Pollard, D., Price, S. F., Ren, D., Saito, F., Sato, T., Seddik, H., Seroussi, H., Takahashi, K., Walker, R., and Wang, W. L.: Ice-sheet model sensitivities to environmental forcing and their use in projecting future sea level (the SeaRISE project), *Journal of Glaciology*, 59, 195–224, doi:10.3189/2013JoG12J125, 2013.
- 720 Bodirsky, B. L. and Müller, C.: Robust relationship between yields and nitrogen inputs indicates three ways to reduce nitrogen pollution, *Environmental Research Letters*, 9, doi:10.1088/1748-9326/9/11/111005, 2014.
- Bodirsky, B. L., Popp, A., Lotze-Campen, H., Dietrich, J. P., Rolinski, S., Weindl, I., Schmitz, C., Müller, C., Bonsch, M., Humpenöder, F., Biewald, A., and Stevanovic, M.: Reactive nitrogen requirements to feed the world in 2050 and potential to mitigate nitrogen pollution, *Nature Communications*, 5, doi:10.1038/ncomms4858, 2014.
- 725 Boers, N., Bookhagen, B., Barbosa, H., Marwan, N., and Kurths, J.: Prediction of extreme floods in the Central Andes by means of Complex Networks, *Nature Communications*, 16, 7173, doi:10.1038/ncomms6199, 2014.
- Caldeira, K.: Ocean model predictions of chemistry changes from carbon dioxide emissions to the atmosphere and ocean, *Journal of Geophysical Research*, 110, 1–12, doi:10.1029/2004JC002671, 2005.
- 730 Chadwick, R. and Good, P.: Understanding nonlinear tropical precipitation responses to CO₂ forcing, *Geophysical Research Letters*, 40, 4911–4915, doi:10.1002/grl.50932, 2013.
- Chen, P.-Y., Chen, C.-C., Chu, L., and Mccarl, B.: Evaluating the economic damage of climate change on global coral reefs, *Global Environmental Change*, 30, 12–20, doi:10.1016/j.gloenvcha.2014.10.011, 2015.
- 735 Church, J. A., Clark, P. U., Cazenave, A., Gregory, J. M., Jevrejeva, S., Levermann, A., Merrifield, M. A., Milne, G. A., Nerem, R. S., Nunn, P. D., Payne, A. J., Pfeffer, W. T., Stammer, D., and Unnikrishnan, A. S.: Sea Level Change, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate*. Chapter 13, edited by Stocker,

T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and P. M. Midgley, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.

Cinner, J. E., Pratchett, M. S., Graham, N. A. J., Messmer, V., Fuentes, M. M. P. B., Ainsworth, T., Ban, N., Bay, L. K., Blythe, J., Dissard, D., Dunn, S., Evans, L., Fabinyi, M., Fidelman, P., Figueiredo, J., Frisch, A. J., Fulton, C. J., Hicks, C. C., Lukoschek, V., Mallela, J., Moya, A., Penin, L., Rummer, J. L., Walker, S., and Williamson, D. H.: A framework for understanding climate change impacts on coral reef social–ecological systems, *Regional Environmental Change*, doi:10.1007/s10113-015-0832-z, 2015.

Clark, P. U., Shakun, J. D., Marcott, S. A., Mix, A. C., Eby, M., Kulp, S., Levermann, A., Milne, G. A., Pfister, P. L., Santer, B. D., Schrag, D. P., Solomon, S., Stocker, T. F., Strauss, B. H., Weaver, A. J., Winkelmann, R., Archer, D., Bard, E., Goldner, A., Lambeck, K., Pierrehumbert, R. T., and Plattner, G.-K.: Consequences of twenty-first-century policy for multi-millennial climate and sea-level change, *Nature Climate Change*, pp. 1–10, doi:10.1038/nclimate2923, 2016.

Coumou, D. and Robinson, A.: Historic and future increase in the global land area affected by monthly heat extremes, *Environmental Research Letters*, 8, 034 018, doi:10.1088/1748-9326/8/3/034018, 2013.

Dai, A.: Increasing drought under global warming in observations and models, *Nature Climate Change*, 3, 52–58, doi:10.1038/nclimate1633, 2012.

Deryng, D., Conway, D., Ramankutty, N., Price, J., and Warren, R.: Global crop yield response to extreme heat stress under multiple climate change futures, *Environmental Research Letters*, 9, 034 011, doi:10.1088/1748-9326/9/3/034011, 2014.

Diffenbaugh, N. S. and Scherer, M.: Observational and model evidence of global emergence of permanent, unprecedented heat in the 20th and 21st centuries, *Climatic Change*, 107, 615–624, doi:10.1007/s10584-011-0112-y, 2011.

Donner, S. D., Skirving, W. J., Little, C. M., Oppenheimer, M., and Hoegh-Gulberg, O.: Global assessment of coral bleaching and required rates of adaptation under climate change, *Global Change Biology*, 11, 2251–2265, doi:10.1111/j.1365-2486.2005.01073.x, 2005.

Donner, S. D., Knutson, T. R., and Oppenheimer, M.: Model-based assessment of the role of human-induced climate change in the 2005 Caribbean coral bleaching event., *Proceedings of the National Academy of Sciences*, 104, 5483–5488, doi:10.1073/pnas.0610122104, 2007.

Donner, S. D.: Coping with commitment: projected thermal stress on coral reefs under different future scenarios., *PloS one*, 4, e5712, doi:10.1371/journal.pone.0005712, 2009.

Dutton, A., Carlson, A. E., Long, A. J., Milne, G. A., Clark, P. U., DeConto, R., Horton, B. P., Rahmstorf, S., and Raymo, M. E.: Sea-level rise due to polar ice-sheet mass loss during past warm periods, *Science*, 349, doi:10.1126/science.aaa4019, 2015.

Eamus, D.: The interaction of rising CO₂ and temperatures with water use efficiency, *Plant, Cell & Environment*, 14, 843–852, doi:10.1111/j.1365-3040.1991.tb01447.x, 1991.

Favier, L., Durand, G., Cornford, S. L., Gudmundsson, G. H., Gagliardini, O., Gillet-Chaulet, F., Zwinger, T., Payne, A. J., and Le Brocq, A. M.: Retreat of Pine Island Glacier controlled by marine ice-sheet instability, *Nature Climate Change*, 4, 117–121, doi:10.1038/nclimate2094, 2014.

Feldmann, J., Levermann, A.: Collapse of the West Antarctic Ice Sheet after local destabilization of the Amundsen Basin, *Proceedings of the National Academy of Sciences*, 112, 112, 14191–14196, doi:10.1073/pnas.1512482112, 2015

780 Fettweis, X., Franco, B., Tedesco, M., van Angelen, J. H., Lenaerts, J. T. M., van den Broeke, M. R., and Gallée, H.: Estimating the Greenland ice sheet surface mass balance contribution to future sea level rise using the regional atmospheric climate model MAR, *The Cryosphere*, 7, 469–489, doi:10.5194/tc-7-469-2013, 2013.

Fischer, E. and Knutti, R.: Detection of spatially aggregated changes in temperature and precipitation extremes, *Geophysical Research Letters*, 41, doi:10.1002/2013GL058499, 2014.

785 Fischer, E. M., Beyerle, U., and Knutti, R.: Robust spatially aggregated projections of climate extremes, *Nature Climate Change*, 3, 1033–1038, doi:10.1038/nclimate2051, 2013.

Flato, G., Marotzke, J., Abiodun, B., Braconnot, P., Chou, S. C., Collins, W., Cox, P., Driouech, F., Emori, S., Eyring, V., Forest, C., Gleckler, P., Guilyardi, E., Jakob, C., Kattsov, V., Reason, C., and Rummukainen, M.: *Evaluation of Climate Models*, book section 9, pp. 741–866, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, doi:10.1017/CBO9781107415324.020, 2013.

790 Frieler, K., Meinshausen, M., Golly, A., Mengel, M., Lebek, K., Donner, S. D., and Hoegh-Guldberg, O.: Limiting global warming to 2C is unlikely to save most coral reefs, *Nature Climate Change*, 3, doi:10.1038/nclimate1674, 2012.

Frieler, K., Levermann, A., Elliott, J., Heinke, J., Arneth, A., Bierkens, M. F. P., Ciais, P., Clark, D. B., Deryng, D., Döll, P., Falloon, P., Fekete, B., Folberth, C., Friend, A. D., Gellhorn, C., Gosling, S. N., Haddeland, I., Khabarov, N., Lomas, M., Masaki, Y., Nishina, K., Neumann, K., Oki, T., Pavlick, R., Ruane, A. C., Schmid, E., Schmitz, C., Stacke, T., Stehfest, E., Tang, Q., Wissler, D., Huber, V., Piontek, F., Warszawski, L., Schewe, J., Lotze-Campen, H., and Schellnhuber, H. J.: A framework for the cross-sectoral integration of multi-model impact projections: land use decisions under climate impacts uncertainties, *Earth System Dynamics*, 6, 447–460, doi:10.5194/esd-6-447-2015, 2015.

800 Gattuso, J.-P., Magnan, A., Bille, R., Cheung, W. W. L., Howes, E. L., Joos, F., Allemand, D., Bopp, L., Cooley, S. R., Eakin, C. M., Hoegh-Guldberg, O., Kelly, R. P., Portner, H.-O., Rogers, A. D., Baxter, J. M., Laffoley, D., Osborn, D., Rankovic, A., Rochette, J., Sumaila, U. R., and Treyer, S.: Contrasting futures for ocean and society from different anthropogenic CO₂ emissions scenarios, *Science*, 349, doi:10.1126/science.aac4722, 2015.

Gourdji, S. M., Sibley, A. M., and Lobell, D. B.: Global crop exposure to critical high temperatures in the reproductive period: historical trends and future projections, *Environmental Research Letters*, 8, doi:10.1088/1748-9326/8/2/024041, 2013.

Graham, N. A. J., Jennings, S., Macneil, M. A., Mouillot, D., and Wilson, S. K.: Predicting climate-driven regime shifts versus rebound potential in coral reefs, *Nature*, 518, 94–97, doi:10.1038/nature14140, 2015.

810 Gregory, J. M. and Huybrechts, P.: Ice-sheet contributions to future sea-level change, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 364, 1709–1732, doi:10.1098/rsta.2006.1796, 2006.

Hansen, J., Sato, M., and Ruedy, R.: PNAS Plus: Perception of climate change, *Proceedings of the National Academy of Sciences*, 2012, doi:10.1073/pnas.1205276109, 2012.

815

- Hawkins, E., Joshi, M., and Frame, D.: Wetter then drier in some tropical areas, *Nature Climate Change*, 4, 646–647, doi:10.1038/nclimate2299, 2014.
- Held, I. M. and Soden, B. J.: Robust Responses of the Hydrological Cycle to Global Warming, *Journal of Climate*, 19, 5686–5699, 2006.
- 820 Hellmer, H. H., Kauker, F., Timmermann, R., Determann, J., and Rae, J.: Twenty-first-century warming of a large Antarctic ice-shelf cavity by a redirected coastal current, *Nature*, 485, 225–228, doi:10.1038/nature11064, 2012.
- Hempel, S., Frieler, K., Warszawski, L., Schewe, J., and Piontek, F.: A trend-preserving bias correction – the ISI-MIP approach, *Earth System Dynamics*, 4, 219–236, doi:10.5194/esd-4-219-2013, 2013.
- 825 Herger, N., Sanderson, B. M., and Knutti, R.: Improved pattern scaling approaches for the use in climate impact studies, *Geophysical Research Letters*, pp. 1–9, doi:10.1002/2015GL063569, 2015.
- Hinkel, J., Lincke, D., Vafeidis, A. T., Perrette, M., Nicholls, R. J., Tol, R. S. J., Marzeion, B., Fettweis, X., Ionescu, C., and Levermann, A.: Coastal flood damage and adaptation costs under 21st century sea-level rise, *Proceedings of the National Academy of Sciences*, 111, 3292–3297, doi:10.1073/pnas.1222469111, 2014.
- 830 Hirabayashi, Y., Mahendran, R., Koirala, S., Konoshima, L., Yamazaki, D., Watanabe, S., Kim, H., and Kanae, S.: Global flood risk under climate change, *Nature Climate Change*, 3, 1–6, doi:10.1038/nclimate1911, 2013.
- IPCC: Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation (SREX), Cambridge University Press, Cambridge, doi:10.1017/CBO9781139177245, 2012.
- IPCC: Summary for Policymakers, in: *Climate Change 2013: The Physical Science Basis. Contribution of*
- 835 *Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Stocker, T., Qin, D., Plattner, G.-K., Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P., pp. 1–100, IPCC AR5 WGI, Cambridge University Press, Cambridge, UK and New York, NY, USA, 2013.
- IPCC: *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, p. 151 pp, IPCC Geneva, Switzerland,
- 840 2014a.
- IPCC: Summary for Policy Makers, in: *Climate Change 2014: Impacts, Adaptation and Vulnerability - Contributions of the Working Group II to the Fifth Assessment Report*, edited by Field, C. B., Barros, V. R., Dokken, D., Mach, K. J., Mastrandrea, M. D., Bilir, T. E., Chatterjee, M., Ebi, K. L., Estrada, Y., Genova,
- 845 R., Girma, B., Kissel, E., Levy, A., MacCracken, S., Mastrandrea, P. R., and White, L., pp. 1–32, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, doi:10.1016/j.renene.2009.11.012, 2014b.
- James, R. and Washington, R.: Changes in African temperature and precipitation associated with degrees of global warming, *Climatic Change*, 117, 859–872, doi:10.1007/s10584-012-0581-7, 2013.
- 850 Jiménez Cisneros, B. E., Oki, T., Arnell, N. W., Benito, G., Cogley, J. G., Döll, P., Jiang, T., Mwakalila, S. S.: Freshwater resources, in: *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel of Climate Change*, edited by Field, C. B., Barros, V. R., Dokken, D. J., Mach, K. J., Mastrandrea, M. D., Bilir, T. E., Chatterjee, M., Ebi, K. L., Estrada, Y. O., Genova, R. C., Girma, B., Kissel, E. S., Levy,

- 855 A. N., MacCracken, S., Mastrandrea, P. R., and White, L. L., pp. 229–269, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2014.
- Joughin, I., Smith, B. E., and Medley, B.: Marine Ice Sheet Collapse Potentially Underway for the Thwaites Glacier Basin, West Antarctica, *Science*, 344, 735–388, doi:10.1126/science.1249055, 2014.
- Kelley, C. P., Mohtadi, S., Cane, M. A., Seager, R., and Kushnir, Y.: Climate change in the Fertile Crescent and
860 implications of the recent Syrian drought, *Proceedings of the National Academy of Sciences of the United States of America*, doi:10.1073/pnas.1421533112, 2015.
- Kendon, E. J., Rowell, D. P., Jones, R. G., and Buonomo, E.: Robustness of Future Changes in Local Precipitation Extremes, *Journal of Climate*, 21, 4280–4297, doi:10.1175/2008JCLI2082.1, 2008.
- Kharin, V. V., Zwiers, F. W., Zhang, X., and Wehner, M.: Changes in temperature and precipitation extremes in
865 the CMIP5 ensemble, *Climatic Change*, 119, 345–357, doi:10.1007/s10584-013-0705-8, 2013.
- Kiem, A. and Verdon-Kidd, D.: Towards understanding hydroclimatic change in Victoria, Australia—preliminary insights into the "Big Dry", *Hydrology and Earth System Sciences*, 14, 433–445, doi:10.5194/hess-14-433-2010, 2010.
- King, A. D., Donat, M. G., Fischer, E. M., Hawkins, E., Alexander, L. V., Karoly, D. J., Dittus, A. J., Lewis,
870 S. C., and Perkins, S. E.: The timing of anthropogenic emergence in simulated climate extremes, *Environmental Research Letters*, 10, 094 015, doi:10.1088/1748-9326/10/9/094015, 2015.
- Knutson, T. R., McBride, J. L., Chan, J., Emanuel, K., Holland, G., Landsea, C., Held, I., Kossin, J. P., Srivastava, A. K., and Sugi, M.: Tropical cyclones and climate change, *Nature Geoscience*, 3, 157–163, doi:10.1038/ngeo779, 2010.
- 875 Knutti, R. and Sedláček, J.: Robustness and uncertainties in the new CMIP5 climate model projections, *Nature Climate Change*, 3, 369–373, doi:10.1038/nclimate1716, 2012.
- Knutti, R., Rogelj, J., Sedláček, J., and Fischer, E. M.: A scientific critique of the two-degree climate change target, *Nature Geoscience*, 9, 13–18, doi:10.1038/ngeo2595, 2015.
- Leakey, A. D. B., Uribeharrea, M., Ainsworth, E. A., Naidu, S. L., Rogers, A., Ort, D. R., and Long, S. P.:
880 Photosynthesis, productivity, and yield of maize are not affected by open-air elevation of CO₂ concentration in the absence of drought., *Plant physiology*, 140, 779–790, doi:10.1104/pp.105.073957, 2006.
- Levermann, A., Clark, P. U., Marzeion, B., Milne, G. A., Pollard, D., Radic, V., and Robinson, A.: The multi-millennial sea-level commitment of global warming., *Proceedings of the National Academy of Sciences of the United States of America*, 110, 13 745–50, doi:10.1073/pnas.1219414110, 2013.
- 885 Levermann, A., Winkelmann, R., Nowicki, S., Fastook, J. L., Frieler, K., Greve, R., Hellmer, H. H., Martin, M. A., Meinshausen, M., Mengel, M., Payne, A. J., Pollard, D., Sato, T., Timmermann, R., Wang, W. L., and Bindshadler, R. A.: Projecting Antarctic ice discharge using response functions from SeaRISE ice-sheet models, *Earth System Dynamics*, 5, 271–293, doi:10.5194/esd-5-271-2014, 2014.
- Lobell, D. B. and Tebaldi, C.: Getting caught with our plants down: the risks of a global crop yield slowdown
890 from climate trends in the next two decades, *Environmental Research Letters*, 9, 074 003, doi:10.1088/1748-9326/9/7/074003, 2014.
- Lopez, A., Suckling, E. B., and Smith, L. A.: Robustness of pattern scaled climate change scenarios for adaptation decision support, *Climatic Change*, 122, 555–566, doi:10.1007/s10584-013-1022-y, 2013.

Marotzke, J. and Forster, P. M.: Forcing , feedback and internal variability in global temperature trends, *Nature*,
895 517, 565–570, doi:10.1038/nature14117, 2014.

Maynard, J., van Hooidonk, R., Eakin, C. M., Puotinen, M., Garren, M., Williams, G., Heron, S. F., Lamb,
J., Weil, E., Willis, B., and Harvell, C. D.: Projections of climate conditions that increase coral disease
susceptibility and pathogen abundance and virulence, *Nature Climate Change*, doi:10.1038/nclimate2625,
2015.

900 McGrath, J. M. and Lobell, D. B.: Regional disparities in the CO₂ fertilization effect and implications for crop
yields, *Environmental Research Letters*, 8, 014054, doi:10.1088/1748-9326/8/1/014054, 2013.

Meehl, G., Stocker, T., Collins, W., Friedlingstein, P., Gaye, A., Gregory, J., Kitoh, A., Knutti, R., Murphy,
J., Noda, A., Raper, S., Watterson, I., Weaver, A., and Zhao, Z.-C.: Global climate projections, in: *Climate
Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report
905 of the Intergovernmental Panel on Climate Change*, edited by Solomon, S., Qin, D., Manning, M., Chen, Z.,
Marquis, M., Averyt, K., Tignor, M., and Miller, H., chap. 10, pp. 747–845, Cambridge University Press,
Cambridge, United Kingdom and New York, NY, USA., 2007.

Meinshausen, M., Meinshausen, N., Hare, W., Raper, S. C. B., Frieler, K., Knutti, R., Frame, D. J., and Allen,
M. R.: Greenhouse-gas emission targets for limiting global warming to 2 °C, *Nature*, 458, 1158–1162,
910 doi:10.1038/nature08017, 2009.

Meinshausen, M., Raper, S. C. B., and Wigley, T. M. L.: Emulating coupled atmosphere-ocean and carbon
cycle models with a simpler model, *MAGICC6 - Part 1: Model description and calibration*, *Atmospheric
Chemistry and Physics*, 11, 1417–1456, doi:10.5194/acp-11-1417-2011, 2011.

Meissner, K. J., Lippmann, T., and Sen Gupta, A.: Large-scale stress factors affecting coral reefs: open ocean
915 sea surface temperature and surface seawater aragonite saturation over the next 400 years, *Coral Reefs*, 31,
309–319, doi:10.1007/s00338-011-0866-8, 2012.

Mengel, M. and Levermann, A.: Ice plug prevents irreversible discharge from East Antarctica, *Nature Climate
Change*, 27, 1–5, doi:10.1038/NCLIMATE2226, 2014.

Monfreda, C., Ramankutty, N., and Foley, J.: Farming the planet: 2. Geographic distribution of crop areas,
920 yields, physiological types, and net primary production in the year 2000., *Global Biogeochemical Cycles*,
22, 1–19, doi:10.1029/2007GB002947, 2008.

Moore, F. C. and Lobell, D. B.: The fingerprint of climate trends on European crop yields, *Proceedings of the
National Academy of Sciences*, p. 201409606, doi:10.1073/pnas.1409606112, 2015.

Mora, C., Frazier, A. G., Longman, R. J., Dacks, R. S., Walton, M. M., Tong, E. J., Sanchez, J. J., Kaiser,
925 L. R., Stender, Y. O., Anderson, J. M., Ambrosino, C. M., Fernandez-Silva, I., Giuseffi, L. M., and Gi-
ambelluca, T. W.: The projected timing of climate departure from recent variability, *Nature*, 502, 183–187,
doi:10.1038/nature12540, 2013.

Mueller, B. and Seneviratne, S. I.: Hot days induced by precipitation deficits at the global scale., *Pro-
ceedings of the National Academy of Sciences of the United States of America*, 109, 12 398–403,
930 doi:10.1073/pnas.1204330109, 2012.

Müller, C., Waha, K., Bondeau, A., and Heinke, J.: Hotspots of climate change impacts in sub-
Saharan Africa and implications for adaptation and development., *Global change biology*, 20, 2505–17,
doi:10.1111/gcb.12586, 2014.

- Olsson, L., Opondo, M., Tschakert, P., Agrawal, A., Eriksen, S. H., Ma, S., Perch, L. N., and Zakiideen, S. A.: Livelihoods and poverty, in: *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel of Climate Change*, edited by Field, C. B., Barros, V. R., Dokken, D. J., Mach, K. J., Mastrandrea, M. D., Bilir, T. E., Chatterjee, M., Ebi, K. L., Estrada, Y. O., Genova, R. C., Girma, B., Kissel, E. S., Levy, A. N., MacCracken, S., Mastrandrea, P. R., and White, L. L., pp. 793–832, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2014.
- Oppenheimer, M., Campos, M., Warren, R., Birkmann, J., Luber, G., O'Neill, B., and Kikkawa, T.: Emergent Risks and Key Vulnerabilities, in: *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Field, C. B., Barros, V. R., Dokken, D., Mach, K. J., Mastrandrea, M. D., Bilir, T. E., Chatterjee, M., Ebi, K. L., Estrada, Y., Genova, R., Girma, B., Kissel, E., Levy, A., MacCracken, S., Mastrandrea, P. R., and White, L., chap. 19, pp. 1039–1099, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2014.
- Orlowsky, B. and Seneviratne, S. I.: Global changes in extreme events: regional and seasonal dimension, *Climatic Change*, 110, 669–696, doi:10.1007/s10584-011-0122-9, 2012.
- O'Neill, B. C., Kriegler, E., Riahi, K., Ebi, K. L., Hallegatte, S., Carter, T. R., Mathur, R., and Vuuren, D. P.: A new scenario framework for climate change research: the concept of shared socioeconomic pathways, *Climatic Change*, 122, 387–400, doi:10.1007/s10584-013-0905-2, 2013.
- Perrette, M., Landerer, F., Riva, R., Frieler, K., and Meinshausen, M.: A scaling approach to project regional sea level rise and its uncertainties, *Earth System Dynamics*, 4, 11–29, doi:10.5194/esd-4-11-2013, 2013.
- Porter, J., Liyong, X., Challinor, A., Cochrane, K., Howden, M., Iqbal, M., Lobell, D., and Travasso, M.: Food Security and Food Production Systems, in: *IPCC 2014: Climate Change 2014: Impacts, Adaptation, and Vulnerability. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Chapter 7. Final Draft, October 2013*, pp. 1–82, IPCC AR5 WGII, Cambridge University Press, Cambridge and New York, 2014.
- Pörtner, H.-O., Karl, D. M., Boyd, P. W., Cheung, W. W. L., Lluich-Cota, S. E., Nojiri, Y., Schmidt, D. N., and Zavialov, P.: Ocean Systems, in: *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Field, C. B., Barros, V. R., Dokken, D., Mach, K. J., Mastrandrea, M. D., Bilir, T. E., Chatterjee, M., Ebi, K. L., Estrada, Y., Genova, R., Girma, B., Kissel, E., Levy, A., MacCracken, S., Mastrandrea, P. R., and White, L., chap. 06, pp. 411–484, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2014.
- Power, S., Delage, F., Chung, C., Kociuba, G., and Keay, K.: Robust twenty-first-century projections of El Niño and related precipitation variability., *Nature*, 502, pp. 541–545, doi:10.1038/nature12580, 2013.
- Prudhomme, C., Giuntoli, I., Robinson, E. L., Clark, D. B., Arnell, N. W., Dankers, R., Fekete, B. M., Franssen, W., Gerten, D., Gosling, S. N., Hagemann, S., Hannah, D. M., Kim, H., Masaki, Y., Satoh, Y., Stacke, T., Wada, Y., and Wisser, D.: Hydrological droughts in the 21st century, hotspots and uncertainties from a global multimodel ensemble experiment, *Proceedings of the National Academy of Sciences*, 111, 3262–3267, doi:10.1073/pnas.1222473110, 2013.

- Riahi, K., Grübler, A., and Nakicenovic, N.: Scenarios of long-term socio-economic and environmental development under climate stabilization, *Technological Forecasting and Social Change*, 74, 887–935, doi:<http://dx.doi.org/10.1016/j.techfore.2006.05.026>, 2007.
- Rignot, E., Mouginot, J., Morlighem, M., Seroussi, H., and Scheuchl, B.: Widespread, rapid grounding line retreat of Pine Island, Thwaites, Smith, and Kohler glaciers, West Antarctica, from 1992 to 2011, *Geophysical Research Letters*, 41, 3502–3509, doi:10.1002/2014GL060140, 2014.
- Robinson, A., Calov, R., and Ganopolski, A.: Multistability and critical thresholds of the Greenland ice sheet, *Nature Climate Change*, 2, 429–432, doi:10.1038/nclimate1449, 2012.
- Rogelj, J., Meinshausen, M., and Knutti, R.: Global warming under old and new scenarios using IPCC climate sensitivity range estimates, *Nature Climate Change*, 2, 248–253, doi:10.1038/nclimate1385, 2012.
- Rogelj, J., McCollum, D. L., Reisinger, A., Meinshausen, M., and Riahi, K.: Probabilistic cost estimates for climate change mitigation, *Nature*, 493, 79–83, doi:10.1038/nature11787, 2013.
- Rogelj, J., Meinshausen, M., Sedláček, J., and Knutti, R.: Implications of potentially lower climate sensitivity on climate projections and policy, *Environmental Research Letters*, 9, 031003, doi:10.1088/1748-9326/9/3/031003, 2014.
- Rogelj, J., Luderer, G., Pietzcker, R. C., Kriegler, E., Schaeffer, M., Krey, V., and Riahi, K.: Energy system transformations for limiting end-of-century warming to below 1.5 °C, *Nature Climate Change*, 5, 519–527, doi:10.1038/nclimate2572, 2015.
- Rosenzweig, C., Elliott, J., Deryng, D., Ruane, A. C., Müller, C., Arneth, A., Boote, K. J., Folberth, C., Glotter, M., Khabarov, N., Neumann, K., Piontek, F., Pugh, T. A. M., Schmid, E., Stehfest, E., Yang, H., and Jones, J. W.: Assessing agricultural risks of climate change in the 21st century in a global gridded crop model intercomparison., *Proceedings of the National Academy of Sciences of the United States of America*, pp. 1–6, doi:10.1073/pnas.1222463110, 2013.
- Schaeffer, M., Hare, W., Rahmstorf, S., and Vermeer, M.: Long-term sea-level rise implied by 1.5 °C and 2 °C warming levels, *Nature Climate Change*, pp. 20–23, doi:10.1038/nclimate1584, 2012.
- Schellnhuber, H.-J., Hare, W. L., Serdeczny, O., Adams, S., Coumou, D., Frieler, K., Marin, M., Otto, I. M., Perrette, M., Robinson, A., Rocha, M., Schaeffer, M., Schewe, J., Wang, X., and Warszawski, L.: Turn Down the Heat: Why a 4°C Warmer World Must be Avoided, Tech. rep., commissioned by the World Bank, Washington D.C., 2012.
- Schellnhuber, H. J., Hare, B., Serdeczny, O., Schaeffer, M., Adams, S., Baarsch, F., Schwan, S., Coumou, D., Robinson, A., Vieweg, M., and Others: Turn down the heat: climate extremes, regional impacts, and the case for resilience, Tech. rep., commissioned by the World Bank, Washington D.C., 2013.
- Schellnhuber, H. J., Reyer, C., Hare, B., Waha, K., Otto, I. M., Serdeczny, O., Schaeffer, M., Schleußner, C.-F., Reckien, D., Marcus, R., Kit, O., Eden, A., Adams, S., Aich, V., Albrecht, T., Baarsch, F., Boit, A., Canales Trujillo, N., Cartsburg, M., Coumou, D., Fader, M., Hoff, H., Jobbins, G., Jones, L., Krummenauer, L., Langerwisch, F., Le Masson, V., Ludi, E., Mengel, M., Möhring, J., Mosello, B., Norton, A., Perette, M., Pereznieto, P., Rammig, A., Reinhardt, J., Robinson, A., Rocha, M., Sakschewski, B., Schaphoff, S., Schewe, J., Stagl, J., and Thonicke, K.: Turn Down the Heat: Confronting the New Climate Normal, Tech. rep., commissioned by the World Bank, Washington D.C., 2014.

- Schewe, J., Heinke, J., Gerten, D., Haddeland, I., Arnell, N. W., Clark, D. B., Dankers, R., Eisner, S., Fekete, B. M., Colon-Gonzalez, F. J., Gosling, S. N., Kim, H., Liu, X., Masaki, Y., Portmann, F. T., Satoh, Y., Stacke, T., Tang, Q., Wada, Y., Wisser, D., Albrecht, T., Frieler, K., Piontek, F., Warszawski, L., and Kabat, P.: Multimodel assessment of water scarcity under climate change, *Proceedings of the National Academy of Sciences*, 2013, doi:10.1073/pnas.1222460110, 2013.
- Schleussner, C.-F., Levermann, A., and Meinshausen, M.: Probabilistic Projections of the Atlantic Overturning, under review, 2014a.
- Schleussner, C. F., Runge, J., Lehmann, J., and Levermann, A.: The role of the North Atlantic overturning and deep-ocean for multi-decadal global-mean-temperature variability, *Earth System Dynamics*, 4, 967–1013, doi:10.5194/esdd-4-967-2013, 2014b.
- SED: UNFCCC: Report on the Structured Expert Dialogue (SED) on the 2013–2015 review, FCCC/SB/2015/INF.1, 2015.
- Seneviratne, S. I., Donat, M. G., Pitman, A. J., Knutti, R., and Wilby, R. L.: Allowable CO₂ emissions based on regional and impact-related climate targets, *Nature*, 1870, 1–7, doi:10.1038/nature16542, 2016.
- Sillmann, J., Kharin, V. V., Zhang, X., Zwiers, F. W., and Bronaugh, D.: Climate extremes indices in the CMIP5 multimodel ensemble: Part 1. Model evaluation in the present climate, *Journal of Geophysical Research: Atmospheres*, 118, 1716–1733, doi:10.1002/jgrd.50203, 2013.
- Sillmann, J., Kharin, V. V., Zhang, X., Zwiers, F. W., and Bronaugh, D.: Climate extremes indices in the CMIP5 multimodel ensemble: Part 2. Future climate projections, *Journal of Geophysical Research: Atmospheres*, 118, 2473–2493, doi:10.1002/jgrd.50188, 2013.
- Sippel, S., Zscheischler, J., Heimann, M., Otto, F.E.L., Peters, J. and Mahecha M.D.: Quantifying changes in climate variability and extremes: Pitfalls and their overcoming, *Geophysical Research Letters*, 42, 9990–9998, doi:10.1002/2015GL066307, 2015.
- Smith, K. R., Woodward, A., Campbell-Lendrum, D., Chadee, D., Honda, Y., Liu, Q., Olwoch, J., Revich, B., and Sauerborn, R.: Human Health: Impacts, Adaptation, and Co-Benefits, in: *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Field, C. B., Barros, V. R., Dokken, D., Mach, K. J., Mastrandrea, M. D., Bilir, T. E., Chatterjee, M., Ebi, K. L., Estrada, Y., Genova, R., Girma, B., Kissel, E., Levy, A., MacCracken, S., Mastrandrea, P. R., and White, L., chap. 11, pp. 709–754, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2014.
- Tai, A. P. K., Martin, M. V., and Heald, C. L.: Threat to future global food security from climate change and ozone air pollution, *Nature Climate Change*, 4, 817–821, doi:10.1038/nclimate2317, 2014.
- Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An Overview of CMIP5 and the Experiment Design, *Bulletin of the American Meteorological Society*, 93, 485–498, doi:10.1175/BAMS-D-11-00094.1, 2011.
- Tebaldi, C. and Arblaster, J. M.: Pattern scaling: Its strengths and limitations, and an update on the latest model simulations, *Climatic Change*, 122, 459–471, doi:10.1007/s10584-013-1032-9, 2014.
- Tebaldi, C. and Friedlingstein, P.: Delayed detection of climate mitigation benefits due to climate inertia and variability, *Proceedings of the National Academy of Sciences of the United States of America*, 110, 17 229–34, doi:10.1073/pnas.1300005110, 2013.

- Tebaldi, C., Arblaster, J. M., and Knutti, R.: Mapping model agreement on future climate projections, *Geophysical Research Letters*, 38, L23701, doi:10.1029/2011GL049863, 2011.
- Teh, L. S. L., Teh, L. C. L., and Sumaila, U. R.: A Global Estimate of the Number of Coral Reef Fishers, *PLoS ONE*, 8, doi:10.1371/journal.pone.0065397, 2013.
- Timmermann, R. and Hellmer, H. H.: Southern Ocean warming and increased ice shelf basal melting in the twenty-first and twenty-second centuries based on coupled ice-ocean finite-element modelling, *Ocean Dynamics*, 63, 1011–1026, doi:10.1007/s10236-013-0642-0, 2013.
- UNFCCC: (United Nations Framework Convention on Climate Change) (1992) Framework Convention on Climate Change. United Nations (UN), New York., 1992.
- UNFCCC: Decision 1/CP.21. The Paris Agreement, 2015.
- Vaughan, M. M., Huffaker, A., Schmelz, E. A., Dafoe, N. J., Christensen, S., Sims, J., Martins, V. F., Swerbilow, J., Romero, M., Alborn, H. T., Allen, L. H., and Teal, P. E. A.: Effects of elevated [CO₂] on maize defence against mycotoxigenic *Fusarium verticillioides*, *Plant, Cell & Environment*, 37, 2691–2706, doi:10.1111/pce.12337, 2014.
- Wada, Y., van Beek, L. P. H., Sperna Weiland, F. C., Chao, B. F., Wu, Y.-H., and Bierkens, M. F. P.: Past and future contribution of global groundwater depletion to sea-level rise, *Geophysical Research Letters*, 39, 1–6, doi:10.1029/2012GL051230, 2012.
- Warszawski, L., Frieler, K., Huber, V., Piontek, F., Serdeczny, O., and Schewe, J.: The Inter-Sectoral Impact Model Intercomparison Project (ISI-MIP): Project framework., *Proceedings of the National Academy of Sciences of the United States of America*, pp. 1–5, doi:10.1073/pnas.1312330110, 2013.
- Werner, A. D., Bakker, M., Post, V.E.A., Vandenbohede, A., Lu, C., Ataie-Ashtiani, B., Simmons, C. T., Barry, D.A. : Seawater intrusion processes, investigation and management: Recent advances and future challenges, *Advances in Water Resources*, 51, 3-26, doi:10.1016/j.advwatres.2012.03.004, 2013.
- Wigley, T. M. L. and Raper, S. C. B.: Extended scenarios for glacier melt due to anthropogenic forcing, *Geophysical Research Letters*, 32, L05 704, doi:10.1029/2004GL021238, 2005.
- Winkelmann, R., Levermann, A., Ridgwell, A., and Caldeira, K.: Combustion of available fossil fuel resources sufficient to eliminate the Antarctic Ice Sheet, *Science Advances*, 1, doi:10.1126/sciadv.1500589, 2015.
- Zhang, X., Alexander, L., Hegerl, G. C., Jones, P., Tank, A. K., Peterson, T. C., Trewin, B., and Zwiers, F. W.: Indices for monitoring changes in extremes based on daily temperature and precipitation data, *Wiley Interdisciplinary Reviews: Climate Change*, 2, 851–870, doi:10.1002/wcc.147, 2011.
- Zopa, S., Schulz, Y. B. M., and Cugnet, S. B. D.: Aerosol and ozone changes as forcing for climate evolution between 1850 and 2100, *Climate Dynamics*, doi:10.1007/s00382-012-1408-y, 2012.