

This discussion paper is/has been under review for the journal Earth System Dynamics (ESD). Please refer to the corresponding final paper in ESD if available.

Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler^{1,2} and J. Tjiputra¹

¹Uni Research Climate, Bjerknes Centre for Climate Research, Bergen, Norway

²Pierre and Marie-Curie University, Paris, France

Received: 25 November 2015 – Accepted: 4 December 2015 – Published: 21 December 2015

Correspondence to: J. Tjiputra (jerry.tjiputra@uni.no)

Published by Copernicus Publications on behalf of the European Geosciences Union.

ESDD

6, 2645–2688, 2015

Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Earth System Model (ESM) simulations exhibit large biases compared to observations of the present ocean CO₂ sink. The inter-model spread in projections increases by nearly two-fold by the end of the 21st century, and therefore contributes significantly to the uncertainty of future climate projections. In this study, the Southern Ocean (SO) is shown to be one of the hot-spot regions for future uptake of anthropogenic CO₂, characterized by both the solubility pump and biological-mediated carbon drawdown in the Spring and Summer. We show, by analyzing a suite of fully-interactive ESMs simulations from the Coupled Model Intercomparison Project phase 5 (CMIP5) over the 21st century under the high CO₂ RCP8.5 scenario, that the SO is the only region where the atmospheric CO₂ uptake rate continues to increase toward the end of the 21st century. Furthermore, our study discovers a strong inter-model link between the contemporary CO₂ uptake in the Southern Ocean and the projected global cumulated uptake over the 21st century. The inter-model spread for the contemporary CO₂ uptake in the Southern Ocean is attributed to the variations in the simulated seasonal cycle of surface pCO₂. Two groups of model behaviour have been identified. The first one simulates anomalously strong SO carbon uptake, generally due to both too-strong net primary production and too-low surface pCO₂ in December–January. The second group simulates an opposite CO₂ flux seasonal phase, which is driven mainly by the bias in the sea surface temperature variability. We show that these biases are persistent throughout the 21st century, which highlight the urgent need for a sustained and comprehensive biogeochemical monitoring system in the Southern Ocean to better constrain key processes represented in current model systems.

1 Introduction

Since the industrial revolution, a steady increase in anthropogenic CO₂ emissions from fossil fuel burning, cement production and land-use change have led to an increase

ESDD

6, 2645–2688, 2015

Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



between the ocean, atmosphere and terrestrial biosphere reservoirs (Taylor et al., 2012).

Figure 1 presents the time-series of the global annual CO₂ uptake by the ocean computed from 9 different CMIP5 models (Sect. 2.2) from the year 2001 to 2099, including the observation-based estimate of carbon flux for the period 1998–2011 (Landschützer et al., 2014). In addition to the large present day inter-model spread, the figure also highlights the increase in the inter-model spread projected into the future. The magnitude of the standard deviation (i.e. of the inter-model variation) increases by a factor of two from 2001 ($\pm 0.3 \text{ PgCyr}^{-1}$) to 2099 ($\pm 0.6 \text{ PgCyr}^{-1}$). During this period, the projected cumulative oceanic carbon sink ranges from 340.4 to 488.5 Pg C. The 149 Pg C difference in the size of the ocean carbon sink translates into roughly a 70 ppm difference in atmospheric CO₂ concentration by the end of the 21st century. In order to improve the fidelity of future projections provided by the climate modeling community, it is necessary to identify and attribute the mechanisms responsible for the growth in the inter-model spread of ocean CO₂ uptake, and to determine methods to constrain this. Our study focuses on analysing relationships between regional and global uncertainties in ocean carbon uptakes, as simulated in CMIP5 models. The need to reduce this inter-model spread is imperative to reduce uncertainty in future climate projections and enable policy makers to make the most

ESDD

Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

The image shows a presentation navigation interface with a dark blue background and white text. At the top is a title bar with the text "Title Page". Below it is a grid of eight navigation buttons arranged in four rows and two columns. The first row contains "Abstract" and "Introduction". The second row contains "Conclusions" and "References". The third row contains "Tables" and "Figures". The fourth row contains navigation icons: a double left arrow and a double right arrow. Below the grid is a button with the text "Back" on the left and "Close" on the right. At the bottom is a large button with the text "Full Screen / Esc". Below that is a button with the text "Printer-friendly Version". At the very bottom is a button with the text "Interactive Discussion".

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀◀	▶▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



informed decisions. The simulated uncertainty in ocean carbon uptake could arise from different factors. Feedbacks from ocean and terrestrial biospheres on the CO₂ concentration are expected but highly uncertain and thus difficult to predict (Denman et al., 2007). Differences in (i) basin-scale ocean evolution of ocean carbon uptake rates, (ii) timing and amplitude of physical and biogeochemical processes driving the regional pCO₂ seasonal cycle, as well as (iii) responses to transient future climate change are among potential contributors to future uncertainty in ocean carbon uptake, and addressing these points will be the main focus of this study.

The paper is organized as follows: in the next section we describe the observations and models used in this study as well as the terms and metrics used to investigate the relationships between present-day and future carbon uptake and the regional boundaries. Section three discusses the results of the analyses. Additional discussions and comparison with previous studies are presented in section four. Finally, the study is summarized in section five.

2 Methods

2.1 Observations

We used the monthly data set documented by Landschützer et al. (2014) for the 2001–2010 period. It includes the surface ocean partial pressure of CO₂ (pCO₂) and the sea-air CO₂ flux (fgCO₂) gridded into a 1° × 1° horizontal resolution corresponding to 360 by 180 points in longitude and latitude, respectively. The pCO₂ data set is originally extrapolated in space and time from SOCAT (Surface Ocean CO₂ Atlas) version 2 (Bakker et al., 2014) by a two-step neural network approach as described in Landschützer et al. (2013). The sea-air CO₂ flux is computed based on this pCO₂ field applying a standard bulk formulation and high-resolution Cross-Calibrated Multi-Platform (CCMP) wind speeds (Atlas et al., 2011). The monthly averaged Sea Surface Salinity (SSS) was downloaded from the Simple Ocean Data Assimilation

Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(SODA, Carton and Giese, 2008) and has been regridded to the $p\text{CO}_2$ dataset. The Sea Surface Temperature (SST) is from the National Oceanic and Atmospheric Administration (NOAA) Optimum Interpolation (OI) sea surface v.2 (Reynolds et al., 2002).

The climatology of Net Primary Production (NPP) used for the seasonal model-data assessment in the Southern Ocean (Sect. 3.2) is documented by Nevison et al. (2015) and is computed over the 1997–2010 period using data derived from the Sea-viewing Wide Field-of-view Sensor (SeaWiFS). It uses an empirical Chlorophyll (Chl) algorithm for the Southern Hemisphere that was tuned to in situ Chl in the Southern Ocean and spatially blended with the standard SeaWiFS OC4 algorithm (Kahru and Mitchell, 2010).

2.2 $p\text{CO}_2$ decomposition

In order to allow decomposition of $p\text{CO}_2$ variability into its physical and biogeochemical components, we estimated the alkalinity (ALK) and dissolved inorganic carbon (DIC) at the same resolution as the $p\text{CO}_2$ data. The alkalinity was computed from the SST and SSS estimates depending on the region using the Lee et al. (2006) formulation. When temperatures are out of range in the selected region, the computation returns a missing value (NaN). The dissolved inorganic carbon (DIC) was computed using CO_2 inorganic carbon chemistry program CO2SYS developed in Matlab (<http://cdiac.ornl.gov/oceans/co2rprt.html>) using the gridded SST, SSS, $p\text{CO}_2$ and alkalinity as input parameters. The global average surface silicate and the phosphate concentrations were used: 10 and $0.75 \mu\text{mol kg SW}^{-1}$, respectively. We expect that this choice has a relatively small influence on our results since a shift of the concentration of silicate and phosphate by 4 and more than 6 times to their original mean values would generate only a 0.05 and 0.2 % change in the DIC computation. To complete the CO2SYS input, we applied the dissociation constants K1 and K2 introduced by Mehrbach et al. (1973) and refitted by Dickson and Millero (1987).

Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Decomposition of the total $p\text{CO}_2$ seasonal variability was based on the following equation (Tjiputra et al., 2014):

$$\frac{dp\text{CO}_2^{\text{tot}}}{dt} \approx \frac{dp\text{CO}_2^{\text{DIC}}}{dt} + \frac{dp\text{CO}_2^{\text{ALK}}}{dt} + \frac{dp\text{CO}_2^{\text{SST}}}{dt} + \frac{dp\text{CO}_2^{\text{SSS}}}{dt} \quad (1)$$

where the $p\text{CO}_2^{\text{tot}}$ variation in time (“ $dp\text{CO}_2^{\text{tot}}/dt$ ”) is approximately equal the sum of the four decomposed $p\text{CO}_2^x$ variation in time, where x is DIC, ALK, SST or SSS. The $p\text{CO}_2^x$ terms represent a set of thermodynamic equations that relates to the inorganic carbon species, taking into account variation in x while the other components are kept at their long-term local average values. In this way, $dp\text{CO}_2^{\text{DIC}}/dt$ is an estimate of the temporal variability of local $p\text{CO}_2$ field as a result of changing DIC only. The same estimates were applied for the other three parameters (i.e. ALK, SST, and SSS).

2.3 Model descriptions and post processing

The nine participating CMIP5 ESMs in alphabetical order, are the (1) Beijing Climate Center BCC-CSM1.1(m), (2) Canadian Centre for Climate Modeling and Analysis CanESM2, (3) Community Earth System Model CESM1-BGC, (4) Geophysical Fluid Dynamics Laboratory GFDL-ESM2G, (5) Hadley Global Environment Model 2 HadGEM2-ES, (6) Japan Agency for Marine-Earth Science and Technology MIROC-ESM, (7) Max Planck Institute for Meteorology MPI-ESM-LR, (8) Japanese Meteorology Research Institute MRI-ESM1 and (9) Norwegian Climate Centre NorESM1-ME. These models have also contributed to the last Intergovernmental Panel on Climate Change Assessment Report (IPCC-AR5). All outputs were downloaded directly from the Earth System Grid Federation (ESGF, <http://esgf.llnl.gov>) and we analyzed the fully interactive CO_2 emissions-based “esm” simulations. These “esm” simulations take into account carbon fluxes between the land-atmosphere and ocean-atmosphere interfaces to prognostically simulate the atmospheric CO_2 concentration. The selection of these models is based on the availability of all variables necessary

Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to discuss the impact of the $p\text{CO}_2$ seasonal cycle on the carbon uptake: fgCO_2 , $p\text{CO}_2$, SST, SSS, DIC, ALK and NPP. However, the BCC-CSM1.1 and the MIROC-ESM models do not provide some of these variables (ALK and DIC) and therefore were only analyzed for the contemporary and future uptake relationship (Sect. 3.1).

In order to compare the global and regional fgCO_2 between models presented in Sect. 3.1, fgCO_2 outputs were also interpolated to the observational grid of 360×180 points. Model outputs from the historical (esmHistorical) experiments were added to the RCP8.5 (esmrcp8.5) to complete the 2001–2009 period of study. We used the same “r1i1p1” realization from each model. Tables 1 and 2 summarize the physical and marine biogeochemical components and features of each model.

The marine primary productivity is one of the key components that governs the carbon cycle in the ocean, impacting the oceanic pump through alteration of the buffering capacity and the CO_2 remaining in the atmosphere. The ocean NPP is controlled by nutrient availability and other physical factors such as temperature and light. As presented in Table 2, the CMIP5 models use different representations of multiple nutrient limitations, varying from one to five explicit nutrients in CMOC-NPZD and BEC-TOPAZ2, respectively and from one to three phytoplankton species, NPZD-HAMOC-CMOC and BEC-TOPAZ2, respectively. This highlights the wide range of biogeochemistry complexity, which can also contribute to the inter-model spread in their respective outputs. We note that the MOM4-L40 uses the OCMIP2 biogeochemistry module, which doesn’t include an explicit marine ecosystem, therefore in this case the primary production is simulated only as a function of surface phosphate concentration.

2.4 Uptake efficiency

In Sect. 3.2 we compute for each model (at the original model resolution) the “uptake efficiency” ($\text{uptake}_{\text{eff}}^y$) of carbon in the ocean, where “y” represents the different basin regions as defined in Sect. 2.6 as well as the global (glb) and other ocean region excluding the SO (eSO). The $\text{uptake}_{\text{eff}}$ measures the efficiency of a specific water mass in taking up carbon for a given change in atmospheric $p\text{CO}_2$. A high $\text{uptake}_{\text{eff}}$

value represents a good capacity of the ocean to contain DIC for a certain change in atmospheric $p\text{CO}_2$ and vice versa. This term is computed as follow:

$$\text{uptake}_{\text{eff}}^y = \frac{\partial[\text{CT}^y]}{\partial[p\text{CO}_2^y]} = \frac{\text{DIC}^y}{p\text{CO}_2^y \text{RF}^y} \quad (2)$$

The “ DIC^y ” and “ $p\text{CO}_2^y$ ” in Eq. (2) represent the respective area-weighted mean surface concentration of DIC and $p\text{CO}_2^y$ within the domain y . “ RF^y ” is the regional mean Revelle Factor (Revelle and Suess, 1957), also known as the inverse of the ocean’s buffering capacity for atmospheric CO_2 uptake, i.e., to convert $\text{CO}_{2,\text{aq}}$ into different carbon species (carbonate and bicarbonate) within domain y . Water masses with a lower Revelle Factor are more efficient at taking up anthropogenic carbon (Sabine et al., 2004). The increase of atmospheric CO_2 has pushed the surface $\text{CO}_{2,\text{aq}}$ concentration to a higher level, resulting in an increase of the Revelle Factor (Zeebe and Wolf-Gladrow, 2001) and thus a decrease of ocean’s buffering capacity. This mechanism represents a positive climate feedback, which reduces the uptake rate of atmospheric CO_2 in the future (Wallace, 2001). The DIC and $p\text{CO}_2$ fields were taken directly from the model outputs whereas the RF were computed with CO2SYS.

2.5 Inter-model correlation

In order to assess qualitatively the existence of any patterns or consistencies between the simulated CO_2 uptakes among the different models, two metrics of inter-model correlation coefficients have been computed: R_{mean}^y and R_{cum}^y . A high correlation coefficient indicates a strong relationship between the contemporary and the projected inter-model spread. Moreover, a statistically significant positive correlation coefficient denotes that models that project weak uptake on the contemporary period tend to project weak future uptake, and vice versa. The two correlation coefficients were

Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



North Atlantic (NAt, 15–60° N), and Arctic Ocean (Arc, > 60° N). Computation for the analysis pertaining to the relationships between contemporary and future CO₂ uptake (Sect. 3.1) are made at the resolution of the observational data.

3 Results

3.1 Inter-model contemporary and future CO₂ uptake relationships

For each region defined in Sect. 2.6, we computed the inter-model correlation coefficient metrics following Eqs. (3) and (4). The SO region yields the highest correlation coefficient as depicted in Fig. 3 with $R_{\text{cum}}^{\text{SO}} = 0.66$ and $R_{\text{mean}}^{\text{SO}} = 0.77$, in blue- and red-bars, respectively. Other regions remain weakly correlated with a correlation coefficient close to zero or under 0.40 for both correlation fields (TPa, TAt, Ind, NPa, NAt and Arc regions). Only the mSO region reveals a quite strong negative correlation coefficient with $R_{\text{mean}}^{\text{mSO}} = -0.55$. Nevertheless, in all regions except the SO, the correlation coefficients are statistically insignificant at 90 % confidence level, while SO shows a statistically significance over 99 %. For the remainder of the analysis we therefore combined the 7 regions with insignificant correlation coefficient into one new region: global ocean excluding SO (eSO).

Globally, for the annual oceanic CO₂ uptake during the 2001–2010 period, all models are well within the range of the observations estimated by Landschützer et al. (2013) of about $1.99 \pm 0.50 \text{ Pg C yr}^{-1}$. The highest uptake estimates are simulated by the BCC-CSM1.1(m) and NorESM1-ME models (2.40 and 2.31 Pg C yr^{-1} , respectively) and the lowest uptake estimates come from the MPI-ESM-LR and MRI-ESM1 models (1.65 and 1.70 Pg C yr^{-1} , respectively).

Figure 4 depicts the inter-model relationships in the SO and eSO domains for $R_{\text{mean}}^{\text{SO}}$, $R_{\text{mean}}^{\text{eSO}}$, $R_{\text{cum}}^{\text{SO}}$ and $R_{\text{cum}}^{\text{eSO}}$. In SO, four models (BCC-CSM1.1, NorESM1-ME, MPI-ESM-LR and CESM1-BGC) overestimate the carbon uptake flux from the atmosphere to the ocean when compared with two independent observational-based estimates of about

0.15 ± 0.12 and 0.27 ± 0.13 PgCyr⁻¹ (Landschützer et al., 2014; Lenton et al., 2013). The highest estimates being simulated by the BCC-CSM1 and NorESM1-ME models at 1.02 ± 0.09 and 0.61 ± 0.10 PgCyr⁻¹, respectively. Two models (CanESM2 and MRI-ESM1) underestimate the flux, with the lowest estimate is simulated by the CanESM2 model, which is also the only model to simulate the Southern Ocean as a source of carbon to the atmosphere at about -0.66 ± 0.09 PgCyr⁻¹ (Fig. 4a). Three other models (GFDL-ESM2G, MIROC-ESM and HadGEM2-ES) projects CO₂ uptake within the two observational-based estimates.

Figure 4a and c illustrates the strong inter-model linear relationships between the contemporary CO₂ uptake rate in the SO and the projected future uptake rate (Fig. 4a) and cumulated carbon uptake over the 21st century (Fig. 4c). In the last decade of the 21st century, the CMIP5 models project ocean carbon uptake rate ranges from 4.2 to 6.0 PgCyr⁻¹. The cumulative oceanic CO₂ uptake during the 21st century is projected to be between 330 and 500 Pg C.

Figure 4 also demonstrates the peculiarity of the Southern Ocean as compared to the rest of the world's ocean (eSO), where the contemporary CO₂ uptake rate in the latter has a relative strong negative correlation with $R_{\text{mean}}^{\text{eSO}} = -0.56$ and $R_{\text{cum}}^{\text{eSO}} = -33$. To investigate the robustness of this correlation coefficient with time we computed R_{cum}^y for all of the original eight regional basins as well as for the eSO for all 10 year windows between 2001 and 2099 (shown in Fig. 4e). The SO region is shown to have a consistently strong positive correlation coefficient across time with $R_{\text{cum}}^{\text{SO}}$ values of 0.70 ± 0.03. Other regions have more pronounced temporal variations, particularly the North Atlantic (NAt) which goes from a negative correlation in early 21st century (-0.39) to a positive correlation after 2020 which increases to 0.55. However, it remains to be statistically insignificant at 90 % of confidence level.

Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.2 Carbon uptake evolution in the Southern Ocean

In this section, we examine why the SO has the highest $R_{\text{mean}}^{\text{SO}}$ and $R_{\text{cum}}^{\text{SO}}$ relative to the other regions. Only seven out of the nine models previously used to establish the correlations are used; the BCC-CSM1 and MIROC-ESM models are excluded because they do not provide the monthly ALK and DIC fields needed for the “uptake_{eff}” analysis. The remaining models are CanESM2, CESM1-BGC, GFDL-ESM2G, HadGEM2-ES, MPI-ESM-LR, MRI-ESM1 and NorESM1-ME. Figure 5 shows the time series anomalies (relative to year 2001) of CO₂ uptake, net primary production (NPP) and uptake efficiency (uptake_{eff}), in SO and eSO as simulated by the CMIP5 models.

There is a general increase in CO₂ uptake for both SO and eSO, as would be expected from the increasing atmospheric CO₂ concentrations under the RCP8.5 scenario. However, in SO (except for the CESM1-BGC model) the simulated uptake rates steadily increase towards the end of the 21st century, and the multi-model mean increases to $1.2 \pm 0.3 \text{ PgCyr}^{-1}$ higher than the present day (Fig. 5a). The CESM1-BGC model simulates stabilization of CO₂ uptake during the last two decades of the 21st century. In the other regions (Fig. 5b), the multi model mean reaches a saturation point of $1.9 \pm 0.4 \text{ PgCyr}^{-1}$, in the 2070s before the uptake strengths go down to $1.5 \pm 0.4 \text{ PgCyr}^{-1}$ in 2100. This “peak and decline” pattern is consistently shown in all models analyzed here.

The unique SO region benefits indeed from the strong link between deep and surface ocean through the southern upwelling (Sallée et al., 2013a). Earlier studies analyzing the previous generation of ESMs also demonstrated that this region will be an important sink of future atmospheric CO₂ although the efficiency of the sink may decrease (Roy et al., 2011). The increasing CO₂ sink in the SO was shown to be associated with a reduction in the fractional ice coverage which alleviates the light limitation on photosynthesis and increases of surface ocean temperature, both of which would increase the phytoplankton growing season.

ESDD

6, 2645–2688, 2015

Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A global mean decrease of NPP by about $-3.12 \pm 3.54 \text{ PgCyr}^{-1}$ is projected by the CMIP5 models, predominantly attributed to the increase in surface temperature leading to stronger stratification and hence reducing the nutrient supply to the surface ocean through vertical mixing (Bopp et al., 2013). The large differences between the structure of the ecosystem models of the CMIP5 models no doubt contributes to the large inter-model uncertainty. For example, the GFDL-ESM2G and MRI-ESM1 models simulate global annual NPP estimates which differ by more than a factor of two at 66.7 and 25.9 PgCyr^{-1} , respectively (not shown), and are outside of the multi-model standard deviation for the NPP estimates. The MRI-ESM1 model considers only one nutrient limitation and simulates only one type of phytoplankton while the GFDL-ESM2G model uses a more sophisticated ecosystem module with five types of nutrients and three classes of phytoplankton (Table 2). However, this alone is insufficient to determine the reason why the GFDL-ESM2G NPP is so strong. On the other hand, the MPI-ESM-LR NPP is in the low end of the inter-model range. Inter-model variations in the physical and biogeochemical interaction important for the surface primary productivity, such as irradiance and upwelling, should be analyzed further to seek to address this question. This, however, is beyond the scope of the present study.

Figure 5c–d show the NPP anomalies relative to 2001 in the SO and eSO regions through the 21st century, highlighting that surface primary production in SO is either stable or weakly increasing by roughly $0.5 \pm 0.3 \text{ PgCyr}^{-1}$ at the end of this century while it is clearly decreasing in the other regions (by $-2.6 \pm 0.1 \text{ PgCyr}^{-1}$). This is consistent with findings by Laufkötter et al. (2015), who show that the NPP increase in the SO is predominantly attributed to the weakening temperature limitation for phytoplankton growth projected in the future. They also indicate that, despite the inter-model agreement in long-term trend, the regional inter-model variation is substantial.

In SO, steady biological production may also be responsible for maintaining low $p\text{CO}_2$ in the Summer and keeping a higher buffer capacity than in the other regions (Hauck and Völker, 2015). The two last panels of Fig. 4 show the anomaly of uptake_{eff} (Sect 2.4 for definition) which is expected to decrease as the Revelle Factor increases

Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



under future high ambient atmospheric CO₂ concentrations (Heinze et al., 2015). Nevertheless, the decreasing trend is weaker in the SO than the eSO region, at -0.28 ± 0.01 vs. $-0.37 \pm 0.15 \mu\text{mol kg}^{-1} \text{ppm}^{-1}$ respectively by year 2099. Indeed, for the same change in $p\text{CO}_2$ and roughly the same Revelle Factor change, the SO experiences a smaller change in DIC (not shown), indicating a unique process occurring in the high latitude SO. Deep winter mixing at polar regions is very efficient in transporting the anthropogenic carbon from surface to depth, resulting in an increase of uptake efficiency (Marinov et al., 2007). Moreover, the increase in the meridional temperature gradient from the tropic to the high latitudes projected in the future could lead to an enhancement of the Antarctic Circumpolar Current via a stronger wind stress at the surface (Gillett and Fyfe, 2013). As a result, this could translate to enhancement in intermediate water formation and more efficient transport of anthropogenic carbon from the surface into depth.

The steady increase in carbon uptake in the SO region could be the reason for the strong correlation of this region with future global sinks (Sect. 3.1 and Fig. 4a). We therefore focus on analyzing the mechanism for ocean uptake in the SO region as simulated by the different models in the next subsections.

3.3 Inter-model division in the Southern Ocean carbon uptake

Figure 4a–b show that the CMIP5 models exhibit diverse contemporary carbon uptake sinks in SO, from an outgassing of $-0.66 \pm 0.09 \text{ Pg C yr}^{-1}$ (CanESM2) to an uptake of $1.02 \pm 0.09 \text{ Pg C yr}^{-1}$ (BCC-CSM1.1). To investigate the mechanisms driving the inter-model heterogeneity, we compute the carbon uptake at a seasonal time-scale, and compare to estimates from observations (Landschützer et al., 2014).

We divided the seven CMIP5 models into two groups. The first group (hereafter referred as “G1”) represents those that simulate anomalously stronger annual CO₂ uptake rate in the SO as compared to the observational-based estimates and consists of the CESM1-BGC, HadGEM2-ES, MPI-ESM-LR, and NorESM1-ME models. The second group (hereafter referred as “G2”) comprises models that simulate anomalously

weaker CO₂ uptake, consisting of the CanESM2, GFDL-ESM2G and MRI-ESM1 models. Figure 6 shows the inter-model mean and spread of these two groups in their projections of the seasonal cycle of carbon fluxes, NPP and anomalies of SST for both the contemporary (2001–2010) and future (2090–2099) periods.

Figure 6 illustrates that the G1 models have nearly the opposite seasonal cycle as the G2 models. The G1 models (Fig. 6a, red lines) simulate a strong mean ocean CO₂ uptake in December–January of about $-0.30 \text{ mol C m}^{-2} \text{ month}^{-1}$, which in the same direction as, but more than four times stronger in magnitude than, that estimated from the observations of $-0.07 \text{ mol C m}^{-2} \text{ month}^{-1}$. This overestimation corresponds to the period of the highest NPP, where the G1 models mean simulate an NPP maximum of more than three times stronger than the observations: 35 compared to $11 \text{ g C m}^{-2} \text{ month}^{-1}$, respectively (Fig. 6c, black and red lines). Moreover, the SST anomaly during this two-month periods has a negligible effect on the CO₂ flux; there is no significant change in carbon flux occurring when the SST anomaly increases from $+0.3$ to $+1.5^\circ\text{C}$ (Fig. 6e, red line). This highlights that the biological activity in G1 models is the primary driver for the CO₂ flux seasonal cycle in SO during high-productivity season, while the impact of the seasonal temperature on the surface $p\text{CO}_2$ appears to play only a secondary role (Takahashi et al., 2002).

In contrast to the G1, G2 models (Fig. 6, blue lines) simulate strong outgassing during the summertime with a positive CO₂ flux of nearly $0.10 \text{ mol C m}^{-2} \text{ month}^{-1}$. This is in disagreement to the observations and is predominantly driven by the SST changes. The magnitude of the SST anomaly from the G2 models is two times stronger than the observations, whereas the NPP cycle is similar in amplitude. The rapid warming and cooling of SST simulated in the G2 models during the Spring and Fall seasons lead to a higher and lower surface $p\text{CO}_2$, respectively. As a result, the G2 models simulate strong CO₂ uptake during the Fall season, which also implicates the solubility pump as a primary driver.

During the austral Winter, in August, observations show a maximum of outgassing, roughly $0.03 \text{ mol C m}^{-2} \text{ month}^{-1}$ (Fig. 6a, black line). G1 models simulate the same

Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



mechanism at this period but the maximum of outgassing is reached earlier in May–June instead of August. Concurrently, G1 models simulate a minimum of NPP at this time, pushing up the $p\text{CO}_2$ at the surface. Thus, the SO turns into a source of CO_2 for the atmosphere despite the SST anomaly of about -0.6°C which would tend to push in the opposite direction. The same shift appears for G2 models but for an opposite CO_2 flux seasonal cycle, with a maximum of uptake in May of nearly $-0.10 \text{ mol C m}^{-2} \text{ month}^{-1}$. The NPP is twice as strong in May–June for the G2 models than for the G1 models at 1.5 vs. $0.6 \text{ g C m}^{-2} \text{ month}^{-1}$, respectively. This, in addition to a stronger magnitude of SST anomaly, leads to a CO_2 uptake in April–June being projected by the G2 models as depicted in Fig. 6a (blue line).

Figure 7 illustrates the mean spatial variation in CO_2 uptake for each season from the observations and as simulated by the G1 and G2 models during the contemporary period. In the G1 models, the too-strong seasonal cycle as compared to the observations occurs throughout most parts of the SO, especially during October–December and January–March with considerably stronger carbon sinks found in south of the circumpolar current. Between 50 and 60°S , the outgassing in G1 models is noticeably stronger during April–June and July–September. In the G2 models, the largest source of bias during October–December and January–March when compared to the observations is found in the Atlantic and Indian sectors of SO where the models simulate a relatively uniformly strong outgassing.

Future period simulations (Fig. 6, right panels) show that the seasonal phase in the carbon flux will be relatively similar, but the amplitude will grow considerably as compared, to the current seasonality. The distinctions in NPP and SST seasonal cycle between G1 and G2 models are also maintained. Therefore, the bias in the present day seasonal phase of CO_2 fluxes is projected to persist toward the end of the 21st century. We note that there is a one month shift of simulated SST anomaly seasonal cycle where maximum of SST anomaly appears in March instead of February.

3.4 Drivers for the Southern Ocean carbon uptake

Following Eq. (1) in the methods section, we decomposed the $p\text{CO}_2$ seasonal cycle anomalies into four drivers: DIC, ALK, SST and SSS for both the contemporary and future periods (shown in Fig. 8). The SSS-induced variations are not shown because the magnitude is negligible relative to the other variables. As with the previous subsection, we focused our analysis for the two contrasting model groups (G1 and G2). The amplitude of $p\text{CO}_2$ from the G1 (Fig. 8a, red line) overestimates the Landschützer et al. (2014) estimations. Nevertheless, it fits closer with later measurements from Merlivat et al. (2015), though they estimated the sea-air carbon flux only from a section of the Southern Ocean where $p\text{CO}_2$ anomalies amplitude can reach roughly $120 \mu\text{atm}$. In both groups as well as the observations, the SST- and ALK-induced $p\text{CO}_2$ anomalies are generally in phase with each other and the opposite with the DIC-induced variability.

The DIC-induced $p\text{CO}_2$ anomaly from the G1 models simulates a minimum in January instead of March, a two-month shift as compared to the observations. The G1 models, which simulate anomalously strong SO carbon uptake, generally simulate too-low surface $p\text{CO}_2$ during December–January (an anomaly of $-40 \mu\text{atm}$; Fig. 8a, red line) due to the too-strong NPP (Fig. 6c, red line). The driving parameter seems indeed to be the DIC at the water surface (-38 to $-64 \mu\text{atm}$ of anomalies in December–January, respectively). Thus DIC consumption by the phytoplankton via photosynthesis confirms the importance of biological activity for carbon uptake in this group of models during this high-productivity period. The G2 models project nearly the same amplitude as the observations, but depicts an opposite phase for carbon uptake as compared to the observations. Figure 8 shows that G2 models simulate anomalously too-strong surface $p\text{CO}_2$ during December–January, of roughly 7 to $12 \mu\text{atm}$, respectively (Fig. 8c, blue line). There, the driving parameters are the SST and alkalinity with anomalies of about 38 and $19 \mu\text{atm}$ in January, respectively. Indeed, these two components tend to push up the $p\text{CO}_2$ at the surface in the Summer season (December–March) and

ESDD

6, 2645–2688, 2015

Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



present also a one-month shift as compared to the observations (Fig. 8c, black squares and circles).

The future simulations accentuate even more the $p\text{CO}_2$ seasonal cycles for the G1 models (Fig. 8b, red line). The amplitude of this seasonal cycle approximately doubles in 2090–2099 relative to 2001–2010 (i.e., the standard deviation increases from 24.5 to 50.3 μatm), mostly due to the DIC-induced variability (Fig. 8b, red triangles). The amplitudes of the SST-induced and ALK-induced variability are projected to double as well, however, their combined magnitude is still weaker than DIC-induced variability. The G2 models maintain roughly the same amplitude of $p\text{CO}_2$ total through the 21st century (Fig. 8d, blue line), though the $p\text{CO}_2$ -induced components increase also by a factor of two. Nevertheless, the DIC-induced seasonality cycle of the G2 models, conversely to the G1, is about the same order of magnitude as compared to the SST-induced seasonal cycle, thus balancing the change in the $p\text{CO}_2$ total seasonal cycle.

4 Discussions

The ocean plays an instrumental role in buffering the increasing atmospheric CO_2 concentration and the ongoing climate change. In this study, for the first time, we evaluate the relationships between present day regional ocean carbon sinks with future cumulative carbon sinks over the 21st century under the high CO_2 RCP8.5 scenario as simulated by a suite of fully-interactive CMIP5 ESMs. The SO is found to be a good predictor for future global carbon uptake. We therefore examined the representation of oceanic carbon uptake and its future evolution in the SO. Specifically, we assess the model capability to simulate the observed seasonal $p\text{CO}_2$ cycle for the present day period.

With respect to the annual mean CO_2 uptake in the Southern Ocean, Jiang et al. (2014) evaluate a set of CMIP5 models but from different simulations, i.e., with prescribed atmospheric CO_2 concentrations, and over a slightly smaller domain (56–62° S). Despite these differences, our present findings are very much comparable to the

Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



prior study, with the CanESM2 and GFDL-ESM2G models simulating net outgassing and close to neutral CO₂ fluxes. On the contrary, the HadGEM2-ES and MPI-ESM-LR were shown to have relatively stronger carbon sinks, especially during austral summer. For the 2001–2010 period, only one model (MRI-ESM1, ~ 362 μatm) simulates a mean surface pCO₂ in the SO that is lower than the observed ~ 364 μatm; the majority of models simulate a stronger mean carbon sink than the observational estimate. This indicates the need to consider the seasonal cycle when evaluating carbon uptake projections in the SO.

Jiang et al. (2014) also examine the CMIP5 models simulated pCO₂ seasonal cycle in the Drake Passage as compared to shipboard measurements. They show that the pCO₂ seasonal cycle in this confined domain is representative of the broader circumpolar region. Our analysis using Landschützer et al. (2014) data further indicates that pCO₂ seasonal cycle in the SO region is reasonably homogeneous in phase, with the maximum pCO₂ in austral Winter (August) and minimum in Summer (January), as illustrated in Figs. 6 and 8. We show that most models simulate larger seasonal pCO₂ amplitude than the observations. Despite overestimating the mean carbon sinks, both HadGEM2-ES and MPI-ESM-LR reproduce the seasonal phase of the data. The MPI-ESM-LR simulates too-strong biological-mediated carbon uptake in the early Spring period, consistent with the anomalously strong late Winter mixing (August–September) (Jiang et al., 2014; Sallée et al., 2013b), which upwells the required nutrients to fuel biological production. The GFDL-ESM2G, one of the two models that simulate carbon sinks closest to the observations, simulates the opposite pCO₂ seasonal phase to the observations, which largely attributed by the bias in absolute values and amplitude of SST seasonal cycle. None of the CMIP5 models analyzed here are able to reproduce the observed seasonal cycle and annual mean carbon sinks within the uncertainty range in the SO. This highlights the difficulty in simulating the correct variability in hydrography and biogeochemistry in this region.

In their model study applying an ad hoc parameterization of wind stirring, Rodgers et al. (2014) demonstrate that changes in wind stirring has large impact on the mean

carbon uptake and seasonal cycle phasing in the SO (south of 45° S). They show that resultant changes in seasonal onset of stratification influences both entrainment and the biological pump. Furthermore, Sallée et al. (2013b) identify the annual mean freshwater flux as the primary source of error for the SO mixed layer depth in CMIP5 models. This uncertainty arises from the lack of accurate estimates of buoyancy fluxes from observations in the region.

According to the analysis preformed in this study, improving the representation of amplitude and seasonal phase of contemporary surface $p\text{CO}_2$ in SO has the potential to reduce the uncertainty of the future ocean carbon uptake in CMIP5 models. Bias in amplitude is identified to be associated with the magnitude of primary production in the Spring–Summer seasons, whereas bias in the the seasonal phase is attributed by poor representation of SST seasonal cycle. Seasonally varying surface primary production data along with relevant biogeochemical and ecosystem state variables (e.g., nutrients and oxygen) would help constrain process parameterization in the model. In order to improve the SST simulation, improvements in representation of physical processes across the air–sea interface and between mixed layer and ocean interior supported with high quality observation would be needed.

Despite a steady increase in surface $p\text{CO}_2$ observations in the SO region over recent decades, it remains markedly undersampled, both spatially and temporally. Presently, there are only a very few locations where the full annual cycle of observations are available (Bakker et al., 2014). The SO region also has the largest differences in the net CO_2 fluxes as estimated from different methods involving observations and models (Landschützer et al., 2014). Recently, Resplandy et al. (2014) find that DIC-induced small spatial-scale $p\text{CO}_2$ structures existing in the SO is non-negligible. Such small-scale processes are generally missing in the coarse resolution CMIP5 models and sparse observations. In addition, the strong interannual variations in the air–sea CO_2 fluxes identified in this region (e.g., Landschützer et al., 2015; Lovenduski et al., 2015) further emphasizes the potentially large uncertainty of the seasonal $p\text{CO}_2$ cycle determined from the observations as presented in this study.

ESDD

6, 2645–2688, 2015

Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Beyond surface processes, uncertainties in the the subsurface circulation patterns could also contribute to the surface biases simulated in the SO. Here, regions critical for biological production and carbon uptake are associated with mode and intermediate water formation locations (Sarmiento et al., 2004; Sallée et al., 2012). Despite that the simulated net uptake rates of atmospheric CO₂ in the SO are mostly overestimated compared to the observations (as shown in this study), Frölicher et al. (2015) show that the CMIP5 models underestimate the anthropogenic carbon storage in the Southern Ocean. This indicates either a shortcoming in the simulated large scale overturning circulation or a too-strong sink of non-anthropogenic carbon simulated here. Therefore, a better constrain of the former mechanism should be prioritized in order to improve the projection of long-term evolution of air–sea CO₂ uptake.

New US-led initiatives that aim to enhance our understanding of the Southern Ocean processes are emerging, for instance, the Southern Ocean Carbon and Climate Observation and Modeling (SOCCOM, <http://soccom.princeton.edu>). The biogeochemical Argo floats planned to be deployed will provide novel measurements that will help tease out how the changing physical processes influence the biogeochemistry dynamics, and vice versa. The EXport Processes in the Ocean from RemoTe Sensing (EXPORT, <http://cce.nasa.gov/cce/ocean.htm>) campaign, which studies the export and fate of ocean NPP using remote sensing observations, will also provide better constrain for ecosystem processes parameterization in the model. In addition, multi-models intercomparison involving observational data such as this study is useful to elucidate the complex interplay among physical and biogeochemical processes, which ultimately would reduce uncertainties in climate projections.

5 Conclusions

The latest generation ESMs project ocean carbon uptake with considerable uncertainty, and this uncertainty is projected to grow twofold by the end of the 21st century under high future CO₂ emissions scenario. In this study, the evaluation of the CMIP5 ESMs

was focused on assessing the ability of the models to project the future CO₂ fluxes between ocean and atmosphere by looking at the correlation coefficient of each region with the global future ocean CO₂ uptake.

We found that the highest inter-model correlation is in the Southern Ocean (SO) region ($R_{\text{mean}}^{\text{SO}} = 0.77$, $R_{\text{cum}}^{\text{SO}} = 0.66$), meaning that most models agree with the evolution of their CO₂ uptake behaviour through the 21st century. The majority of models simulate a steady increase in CO₂ sink rate due to a weaker decrease in buffer capacity and to a relatively stable NPP throughout the 21st century. We show that models that take up anomalously low CO₂ in the SO today would project low cumulative CO₂ uptake throughout the 21st century and vice versa. This suggests that the carbon uptake in the SO can be used to constrain future global uptake uncertainty. We highlighted that in other regions, the models simulate a decrease of CO₂ uptake during the second half of this century but with large inter-model spread in the timing of the decreasing trend, thus affecting the multi-model correlation in these areas.

We have identified a strong bias in the amplitude of carbon uptake simulated by the CMIP5 models for the period 2001–2010 in the Southern Ocean, ranging from a source to the atmosphere to a sink almost three times more powerful than has been observed (1.02 ± 0.09 vs. 0.27 ± 0.13 Pg C yr⁻¹, Lenton et al., 2013). Inter-model spread in the SO carbon sink arises from variations in the surface *p*CO₂ seasonality, which is attributed by the bias in the simulated timing and amplitude of primary production and SST. By analyzing the differences in the simulated *p*CO₂ seasonalities, we classified two groups of models according to two different behaviours. Models that simulate anomalously strong 2001–2010 CO₂ uptake in SO (Fig. 6a, red color) reproduce the observed *p*CO₂ seasonal cycle but its amplitude is 2.5 times stronger than the observations. This is because of the strong surface DIC variations, which pushes down the *p*CO₂ by simulating too-strong biological production. The effect on the projected simulations is an increase of CO₂ uptake due to the weaker decrease in seawater CO₂ buffering capacity. Other models that simulate anomalously low 2001–2010 CO₂ uptake in SO (Fig. 6a, blue color) simulate comparable seasonal *p*CO₂ amplitude with

Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the observations but in the opposite direction. This is due to the bias in the SST seasonal cycle with stronger amplitude which tend to push up the $p\text{CO}_2$ in January–March, therefore simulating the SO as evolving towards a source of carbon for the atmosphere.

Seasonal timing and amplitude in $p\text{CO}_2$ are shown to be critical in order to accurately simulate the present and future CO_2 uptake and therefore accurate monitoring of these biogeochemical processes in the SO is critical in order to constrain the assessment of the contemporary and future ocean carbon sink, and subsequently the uncertainty in future climate change. However, the Southern Ocean remains one of the most poorly sampled ocean regions with respect to biogeochemistry. The observational data analysed here were generated through extrapolations of the limited direct measurements, which could add an extra uncertainty into the analysis. This emphasizes the urgent need for a sustained and comprehensive observational campaign in this region, which is emerging as the key region to better constrain the evolution of future ocean carbon sinks.

ESDD

6, 2645–2688, 2015

Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiptura

The navigation bar consists of a series of blue buttons with white text. The top button is 'Title Page'. Below it are two buttons: 'Abstract' and 'Introduction'. The next row contains 'Conclusions' and 'References'. The following row has 'Tables' and 'Figures'. Below these are four navigation buttons: a double left arrow, a double right arrow, a single left arrow, and a single right arrow. The next row contains 'Back' and 'Close'. Below these is a wide button labeled 'Full Screen / Esc'. The final row contains a wide button labeled 'Printer-friendly Version'. The bottom-most button is labeled 'Interactive Discussion'.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀◀	▶▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



ns1002k). We also acknowledge the EU FP7 Marie Curies International Research Staff Exchange Scheme (IRSES) Fellowship SOCCLI “The role of Southern Ocean carbon cycle under climate change (no. 317699)”. This is a contribution to the Bjerknes Centre for Climate Research.

References

- Arora Vivek, K., Boer, G. J., Friedlingstein, P., Eby, M., Jones, C., Christian, J., Bonan, G., Bopp, L., Brovkin, V., Cadule, P., Hajima, T., Ilyina, T., Lindsay, K., Tjiputra, J., and Wu, T.: Carbon concentration and carbon–climate feedbacks in CMIP5 Earth system models, *J. Climate*, 26, 5289–5314, 2013. 2647, 2676
- Assmann, K. M., Bentsen, M., Segschneider, J., and Heinze, C.: An isopycnic ocean carbon cycle model, *Geosci. Model Dev.*, 3, 143–167, doi:10.5194/gmd-3-143-2010, 2010. 2676
- Atlas, R., Hoffman, R. N., Ardizzone, J., Leidner, S. M., Jusem, J. C., Smith, D. K., and Gombos, D.: A cross-calibrated multiplatform ocean surface wind velocity product for meteorological and oceanographic applications, *B. Am. Meteorol. Soc.*, 92, 157–174, 2011. 2649
- Bakker, D. C. E., Pfeil, B., Smith, K., et al.: An update to the Surface Ocean CO₂ Atlas (SOCAT version 2), *Earth Syst. Sci. Data*, 6, 69–90, doi:10.5194/essd-6-69-2014, 2014. 2649, 2665
- Bopp, L., Resplandy, L., Orr, J. C., Doney, S. C., Dunne, J. P., Gehlen, M., Halloran, P., Heinze, C., Ilyina, T., Séférián, R., Tjiputra, J., and Vichi, M.: Multiple stressors of ocean ecosystems in the 21st century: projections with CMIP5 models, *Biogeosciences*, 10, 6225–6245, doi:10.5194/bg-10-6225-2013, 2013. 2658
- Carton, J. A. and Giese, B. S.: A reanalysis of ocean climate using Simple Ocean Data Assimilation (SODA), *Mon. Weather Rev.*, 36, 2999–3017, 2008. 2650
- Collins, W. J., Bellouin, N., Doutriaux-Boucher, M., Gedney, N., Halloran, P., Hinton, T., Hughes, J., Jones, C. D., Joshi, M., Liddicoat, S., Martin, G., O’Connor, F., Rae, J., Senior, C., Sitch, S., Totterdell, I., Wiltshire, A., and Woodward, S.: Development and evaluation of an Earth-System model – HadGEM2, *Geosci. Model Dev.*, 4, 1051–1075, doi:10.5194/gmd-4-1051-2011, 2011. 2676
- Denman, K. L., Brasseur, G., Chidthaisong, A., Ciais, P., Cox, P. M., Dickinson, R. E., Hauglustaine, D., Heinze, C., Holland, E., Jacob, D., Lohmann, U., Ramachandran, S.,

ESDD

6, 2645–2688, 2015

Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



da Silva Dias, P. L., Wofsy, S. C., and Zhang, X.: Couplings between changes in the climate system and biogeochemistry, in: Climate Change 2007: The Physical Science Basis, Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge University Press, Cambridge, UK and New York, NY, USA, 499–587, 2007. 2649

Dickson, A. G., and Millero, F. J.: A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media, Deep-Sea Res., 34, 1733–1743, 1987. 2650

Dunne, J. P., John, J. G., Adcroft, A. J., Griffies, S. M., Hallberg, R. W., Shevliakova, E., Stouffer, R. J., Cooke, W., Dunne, K. A., Harrison, M. J., Krasting, J. P., Malyshev, S. L., Milly, P. C. D., Philipps, P. J., Sentman, L. T., Samuels, B. L., Spelman, M. J., Winton, M., Wittenberg, A. T., and Zadeh, N.: GFDL's ESM2 global coupled climate–carbon Earth System Models Part I: Physical formulation and baseline simulation characteristics, J. Climate, 25, 6646–6665, doi:10.1175/JCLI-D-11-00560.1, 2012. 2676

Friedlingstein, P., Andrew, R. M., Rogelj, J., Peters, G. P., Canadell, J. G., Knutti, R., Luderer, G., Raupach, M. R., Schaeffer, M., Van Vuuren, D. P., and Le Quéré, C.: Persistent growth of CO₂ emissions and implications for reaching climate targets, Nat. Geosci., 7, 709–715, 2014. 2648

Frölicher, T. L., Sarmiento, J. L., Paynter, D. J., Dunne, J. P., Krasting, J. P., and Winton, M.: Dominance of the Southern Ocean in anthropogenic carbon and heat uptake in CMIP5 models, J. Climate, 28, 862–886, doi:10.1175/JCLI-D-14-00117.1, 2015. 2666

Fuss, S., Canadell, J. G., Peters, G. P., Tavoni, M., Andrew, R. M., Ciais, P., Jackson, R. B., Jones, C. D., Kraxner, F., Nakicenovic, N., Le Quéré, C., Raupach, M. R., Sharifi, A., Smith, P., and Yamagata, Y.: Betting on negative emissions, Nature Clim. Change, 4, 850–853, doi:10.1038/nclimate2392, 2014. 2648

Gent, P. R., Danabasoglu, G., Donner, L. J., Holland, M. M., Hunke, E. C., Jayne, S. R., Lawrence, D. M., Neale, R. B., Rasch, P. J., Vertenstein, M., Worley, P. H., Yang, Z.-L., and Zhang, M.: The community climate system model version 4., J. Climate, 24, 4973–4991, 2011. 2676

Gillett, N. P. and Fyfe, J. C.: Annular mode changes in the CMIP5 simulations, Geophys. Res. Lett., 40, 1189–1193, 2013. 2659

Giorgetta, M. A., Jungclaus, J. H., Reick, C. H., Legutke, S., Brovkin, V., Crueger, T., Esch, M., Fieg, K., Glushak, K., Gayler, V., Haak, H., Hollweg, H.-D., Ilyina, T., Kinne, S., Kornblueh, L.,

Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Matei, D., Mauritsen, T., Mikolajewicz, U., Mueller, W. A., Notz, D., Raddatz, T., Rast, S., Redler, R., Roeckner, E., Schmidt, H., Schnur, R., Segschneider, J., Six, K., Stockhause, M., Wegner, J., Widmann, H., Wieners, K.-H., Claussen, M., Marotzke, J., and Stevens, B.: Climate change from 1850 to 2100 in MPI–ESM simulations for the Coupled Model Intercomparison Project 5, *J. Adv. Model. Earth Syst.*, 5, 572–597, 2013. 2676

Hauck, J. and C. Völker: Rising atmospheric CO₂ leads to large impact of biology on Southern Ocean CO₂ uptake via changes of the Revelle factor, *Geophys. Res. Lett.*, 42, 1459–1464, 2015. 2658

Heinze, C., Meyer, S., Goris, N., Anderson, L., Steinfeldt, R., Chang, N., Le Quéré, C., and Bakker, D. C. E.: The ocean carbon sink – impacts, vulnerabilities and challenges, *Earth Syst. Dynam.*, 6, 327–358, doi:10.5194/esd-6-327-2015, 2015. 2647, 2659

Ilyina, T., Six, K. D., Segschneider, J., Maier-Reimer, E., Li, H., and Nunez-Riboni, I.: The global ocean biogeochemistry model HAMOCC: model architecture and performance as component of the MPI–Earth System Model in different CMIP5 experimental realizations, *J. Adv. Model. Earth Syst.*, 5, 1–29, 2013. 2676

Jiang, C. L., Gille, S. T., Sprintall, J., and Sweeney, C.: Drake passage oceanic pCO₂: evaluating CMIP5 coupled carbon–climate models using in situ observations, *J. Climate*, 27, 76–100, doi:10.1175/JCLI-D-12-00571.1, 2014. 2663, 2664

Kahru, M. and Mitchell, G.: Blending of ocean colour algorithms applied to the Southern Ocean, *Remote Sens. Lett.*, 1, 119–124, 2010. 2650

Landschützer, P., Gruber, N., Bakker, D. C. E., Schuster, U., Nakaoka, S., Payne, M. R., Sasse, T. P., and Zeng, J.: A neural network-based estimate of the seasonal to inter-annual variability of the Atlantic Ocean carbon sink, *Biogeosciences*, 10, 7793–7815, doi:10.5194/bg-10-7793-2013, 2013. 2649, 2655

Landschützer, P., Gruber, N., Bakker, D. C. E., and Schuster, U.: Recent variability of the global ocean carbon sink, *Global Biogeochem. Cy.*, 28, 927–949, doi:10.1002/2014GB004853, 2014. 2648, 2649, 2656, 2659, 2662, 2664, 2665, 2678, 2682, 2686, 2687, 2688

Landschützer, P., Gruber, N., Haumann, F. A., Rödenbeck, C., Bakker, D. C. E., van Heuven, S., Hoppema, M., Metzl, N., Sweeney, C., Takahashi, T., Tilbrook, B., and Wanninkhof, R.: The reinvigoration of the Southern Ocean carbon sink, *Science*, 349, 1221–1224, 2015. 2665

Laufkötter, C., Vogt, M., Gruber, N., Aita-Noguchi, M., Aumont, O., Bopp, L., Buitenhuis, E., Doney, S. C., Dunne, J., Hashioka, T., Hauck, J., Hirata, T., John, J., Le Quéré, C., Lima, I. D., Nakano, H., Seferian, R., Totterdell, I., Vichi, M., and Völker, C.: Drivers and uncertainties

Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of future global marine primary production in marine ecosystem models, Biogeosciences Discuss., 12, 3731–3824, doi:10.5194/bgd-12-3731-2015, 2015. 2658

Lee, K., Tong, L. T., Millero, F. J., Sabine, C. L., Dickson, A. G., Goyet, C., Park, G.-H., Wanninkhof, R., Feely, R. A., and Key, R. M.: Global relationships of total alkalinity with salinity and temperature in surface waters of the world's oceans, Geophys. Res. Lett., 33, L19605, doi:10.1029/2006GL027207, 2006. 2650

Lenton, A., Tilbrook, B., Law, R. M., Bakker, D., Doney, S. C., Gruber, N., Ishii, M., Hoppema, M., Lovenduski, N. S., Matear, R. J., McNeil, B. I., Metzl, N., Mikaloff Fletcher, S. E., Monteiro, P. M. S., Rödenbeck, C., Sweeney, C., and Takahashi, T.: Sea–air CO₂ fluxes in the Southern Ocean for the period 1990–2009, Biogeosciences, 10, 4037–4054, doi:10.5194/bg-10-4037-2013, 2013. 2656, 2667

Le Quéré, C., Raupach, M. R., Canadell, J. G., Marland, G., Bopp, L., Ciais, P., Conway, T. J., Doney, S. C., Feely, R., Foster, P., Friedlingstein, P., Gurney, K., Houghton, R. A., House, J. I., Huntingford, C., Levy, P. E., Lomas, M. R., Majkut, J., Metzl, N., Ometto, J. P., Peters, G. P., Prentice, I. C., Randerson, J. T., Running, S. W., Sarmiento, J. L., Schuster, U., Sitch, S., Viovy, T. T. N., van der Werf, G. R., and Woodward, F. I.: Trends in the sources and sinks of carbon dioxide, Nat. Geosci., 2, 831–836, 2009. 2647

Lindsay, K., Bonan, G. B., Doney, S. C., Hoffman, F. M., Lawrence, D. M., Long, M. C., Mahowald, N. M., Moore, J. K., Randerson, J. T., and Thornton, P. E.: Preindustrial control and twentieth-century carbon cycle experiments with the Earth system model CESM1(BGC), J. Climate, 27, 8981–9005, doi:10.1175/JCLI-D-12-00565.1, 2014. 2676

Lovenduski, N. S., Fay, A. R., and McKinley, G. A.: Observing multidecadal trends in Southern Ocean CO₂ uptake: What can we learn from an ocean model?, Global Biogeochem. Cy., 29, 416–426, 2015. 2665

Merlivat, L., Boutin, J., and Antoine, D.: Roles of biological and physical processes in driving seasonal air–sea CO₂ flux in the Southern Ocean: new insights from CARIOCA pCO₂, J. Marine Syst., 147, 9–20, 2015. 2662

Marinov, I., Gnanadesikan, A., Toggweiler, J. R., and Sarmiento, J. L.: The Southern Ocean biogeochemical divide, Nature, 441, 964–967, 2007. 2659

Mehrbach, C., Culbertson, C. H., Hawley, J. E., and Pytkwicz, R. M.: Measurement of the apparent dissociation constants of carbonic acid in seawater at atmospheric pressure, Limnol. Oceanogr., 18, 897–907, 1973. 2650

- Mikaloff Fletcher, S. E., Gruber, N., Jacobson, A. R., Gloor, M., Doney, S. C., Dutkiewicz, S., Gerber, M., Follows, M., Joos, F., Lindsay, K., Menemenlis, D., Mouchet, A., Müller, S. A., and Sarmiento, J. L.: Inverse estimates of the oceanic sources and sinks of natural CO₂ and the implied oceanic carbon transport, *Global Biogeochem. Cy.*, 21, GB1010, doi:10.1029/2006GB002751, 2007. 2654
- Nevison, C. D., Manizza, M., Keeling, R. F., Kahru, M., Bopp, L., Dunne, J., Tiputra, J., Ilyina, T., and Mitchell, B. G.: Evaluating the ocean biogeochemical components of Earth system models using atmospheric potential oxygen and ocean color data, *Biogeosciences*, 12, 193–208, doi:10.5194/bg-12-193-2015, 2015. 2650, 2686
- Oschlies, A.: NAO-induced long-term changes in nutrient supply to the surface waters of the North Atlantic, *Geophys. Res. Lett.*, 28, 1751–1754, 2001. 2676
- Palmer, J. R. and Totterdell, I. J.: Production and export in a global ecosystem model, *Deep-Sea Res. Pt. I*, 48, 1169–1198, 2001. 2676
- Peters, G. P., Andrew, R. M., Boden, T., Canadell, J. G., Ciais, P., Le Quéré, C., Marland, G., Raupach, M. R., and Wilson, C.: The challenge to keep global warming below 2 °C, *Nature Climate Change*, 3, 4–6, 2013. 2648
- Rahmstorf, S., Box, J. E., Feulner, G., Mann, M. E., Robinson, A., Rutherford, S., and Schaffernicht, E. J.: Exceptional twentieth-century slowdown in Atlantic Ocean overturning circulation, *Nature Climate Change*, 5, 475–480, 2015. 2647
- Resplandy, L., Boutin, J., and Merlivat, L.: Observed small spatial scale and seasonal variability of the CO₂ system in the Southern Ocean, *Biogeosciences*, 11, 75–90, doi:10.5194/bg-11-75-2014, 2014. 2665
- Revelle, R. and Suess, H. E.: Carbon dioxide exchange between atmosphere and ocean and the question of an increase of atmospheric CO₂ during the past decades, *Tellus*, 9, 18–27, 1957. 2653
- Reynolds, R. W., Rayner, N. A., Smith, T. M., Stokes, D. C., and Wang, W.: An improved in situ and satellite SST analysis for climate, *J. Climate*, 15, 1609–1625, 2002. 2650
- Rodgers, K. B., Aumont, O., Mikaloff Fletcher, S. E., Plancherel, Y., Bopp, L., de Boyer Montégut, C., Iudicone, D., Keeling, R. F., Madec, G., and Wanninkhof, R.: Strong sensitivity of Southern Ocean carbon uptake and nutrient cycling to wind stirring, *Biogeosciences*, 11, 4077–4098, doi:10.5194/bg-11-4077-2014, 2014. 2664
- Roy, T., Bopp, L., Gehlen, M., Schneider, B., Cadule, P., Frolicher, T. L., Segschneider, J., Tjiputra, J., Heinze, C., and Joos, F.: Regional impacts of climate change and atmospheric

Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- CO₂ on future ocean carbon uptake: a multimodel linear feedback analysis, *J. Climate*, 24, 5195–5195, 2011. 2657
- Sabine, C. L., Feely, R. A., Gruber, N., Key, R. M., Lee, K., Bullister, J. L., Wanninkhof, R., Wong, C. S., Wallace, D. W. R., Tilbrook, B., Millero, F. J., Peng, T., Kozyr, A., Ono, T., and Rios, A. F.: The oceanic sink for anthropogenic CO₂, *Science*, 305, 367–374, 2004. 2653
- Sabine, C. L., Hankin, S., Koyuk, H., et al.: Surface Ocean CO₂ Atlas (SOCAT) gridded data products, *Earth Syst. Sci. Data*, 5, 145–153, doi:10.5194/essd-5-145-2013, 2013.
- Sallée, J. B., Matear, R. J., Rintoul, S. R., and Lenton, A.: Localized subduction of anthropogenic carbon dioxide in the Southern Hemisphere oceans, *Nat. Geosci.*, 5, 579–584, 2012. 2666
- Sallée, J. B., Shuckburgh, E., Bruneau, N., Meijers, A. J. S., Bracegirdle, T. J., Wang, Z., and Roy, T.: Assessment of Southern Ocean water mass circulation and characteristics in CMIP5 models: historical bias and forcing response, *J. Geophys. Res.-Oceans*, 118, 1830–1844, doi:10.1002/jgrc.20135, 2013a. 2657
- Sallée, J. B., Shuckburgh, E., Bruneau, N., Meijers, A. J. S., Bracegirdle, T. J., and Wang, Z.: Assessment of Southern Ocean mixed layer depths in CMIP5 models: historical bias and forcing response, *J. Geophys. Res.-Oceans*, 118, 1845–1862, doi:10.1002/jgrc.20157, 2013b. 2664, 2665
- Sarmiento, J. L., Gruber, N., Brzezinski, M. A., and Dunne, J. P.: High-latitude controls of thermocline nutrients and low latitude biological productivity, *Nature*, 427, 56–60, 2004. 2666
- Takahashi, T., Sutherland, S. C., Sweeney, C., Poisson, A., Metzl, N., Tilbrook, B., Bates, N., Wanninkhof, R., Feely, R. A., Sabine, C., Iafsson, J., and Nojiri, Y.: Global sea-air CO₂ flux based on climatological surface ocean pCO₂ and seasonal biological and temperature effects, *Deep-Sea Res. Pt. II*, 49, 1601–1622, 2002. 2660
- Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An overview of CMIP5 and the experiment design, *B. Am. Meteorol. Soc.*, 93, 485–498, 2012. 2648
- Tjiputra, J. F., Roelandt, C., Bentsen, M., Lawrence, D. M., Lorentzen, T., Schwinger, J., Seland, Ø., and Heinze, C.: Evaluation of the carbon cycle components in the Norwegian Earth System Model (NorESM), *Geosci. Model Dev.*, 6, 301–325, doi:10.5194/gmd-6-301-2013, 2013. 2676
- Tjiputra, J. F., Olsen, A., Bopp, L., Lenton, A., Pfeil, B., Roy, T., Segschneider, J., Totterdell, I., and Heinze, C.: Long-term surface pCO₂ trends from observations and models, *Tellus B*, 66, 23083, doi:10.3402/tellusb.v66.23083, 2014. 2651

Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Wallace, D.: Storage and transport of excess CO₂ in the oceans: the JGOFS/WOCE global CO₂ survey, in: Ocean Circulation and Climate: Observing and Modelling the Global Ocean, edited by: Siedler, J., Church, J., and Gould, J., Academic, San Diego, CA, 489–521, 2001. 2653

5 Watanabe, S., Hajima, T., Sudo, K., Nagashima, T., Takemura, T., Okajima, H., Nozawa, T., Kawase, H., Abe, M., Yokohata, T., Ise, T., Sato, H., Kato, E., Takata, K., Emori, S., and Kawamiya, M.: MIROC-ESM 2010: model description and basic results of CMIP5-20c3m experiments, Geosci. Model Dev., 4, 845–872, doi:10.5194/gmd-4-845-2011, 2011. 2676

10 Wu, T., Li, W., Ji, J., Xin, X., Li, L., Wang, Z., Zhang, Y., Li, J., Zhang, F., Wei, M., Shi, X., Wu, F., Zhang, L., Chu, M., Jie, M., Liu, Y., Wang, F., Liu, X., Li, Q., Dong, M., Liang, X., Gao, Y., and Zhang, J.: Global carbon budgets simulated by the Beijing Climate Center Climate System Model for the last century, J. Geophys. Res.-Atmos., 118, 4326–4347, 2013. 2676

15 Yukimoto, S., Adachi, Y., Hosaka, M., Sakami, T., Yoshimura, H., Hirabara, M., Tanaka, T. Y., Shindo, E., Tsujino, H., Deushi, M., Mizuta, R., Yabu, S., Obata, A., Nakano, H., Koshiro, T., Ose, T., and Kitho, A.: A New global climate model of the Meteorological Research Institute: MRI-CGCM3, J. Meteorol. Soc. Jpn., 90A, 23–64, 2012. 2676

Zeebe, R. and Wolf-Gladrow, D.: CO₂ in Seawater: Equilibrium, Kinetics, Isotopes, Elsevier Oceanogr. Ser., vol. 65, Elsevier, Amsterdam, the Netherlands, 346 pp., 2001. 2653

Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Table 1. Description of the models analyzed in this study, indicating ocean model resolutions (levels on the vertical, horizontal resolution in degrees), the thickness of the surface layer and the marine biogeochemical components included in the ESM.

Model	Ocean (level, zonal, meridional)	1st layer thickness [m]	Marine biogeochemical components	References
BCC-CSM1.1(m)	40 lev, 0.3–1, 1°	10	MOM4–L40	Wu et al. (2013)
CanESM2	40 lev, 1.41, 0.94°	10	CMOC	Arora et al. (2013)
CESM1–BGC	60 lev, 1.25, 0.27–0.54°	10	BEC	Gent et al. (2011) Lindsay et al. (2014)
GFDL–ESM2G	63 lev, 0.3–1°	2	TOPAZ2	Dunne et al. (2013)
HadGEM2–ES	40 lev, 0.3–1°	10	Diat–HadOCC	Collins et al. (2011) Palmer and Totterdell (2001)
MIROC–ESM	44 lev, 1.4, 0.5–1.7°	2.5	NPZD	Watanabe et al. (2011)
MPI–ESM–LR	40 lev, 1.5°	10	HAMOCC5.2	Oschlies (2001) Giorgetta et al. (2013)
MRI–ESM1	51 lev, 0.5–1°	4	NPZD	Ilyina et al. (2013)
NorESM1–ME	53 lev, 1.125°	10	HAMOCC5.1	Yukimoto et al. (2012) Assmann et al. (2010) Tjiputra et al. (2013)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Table 2. Main characteristics of the marine biogeochemical components of the 10 ESMs used in this study: list of nutrients limiting the phytoplankton growth and the number of explicitly represented phytoplankton and zooplankton groups.

Model	Nutrients	Phytoplankton	Zooplankton
BEC	5(NO ₃ , NH ₄ , PO ₄ , SiO ₄ , Fe)	3(diatoms, nanophyto–, diazotrophs)	1
CMOC	NO ₃	1	1
Diat–HadOCC	4(NO ₃ , NH ₄ , SiO ₄ , Fe)	2(diatoms, non–diatoms)	1
HAMOCC5.1	4(NO ₃ , PO ₄ , SiO ₄ , Fe)	1	1
HAMOCC5.2	3(PO ₄ , NO ₃ , Fe)	1	1
MOM4–L40	PO ₄	–	–
NPZD	NO ₃	1	1
TOPAZ2	5(NO ₃ , NH ₄ , PO ₄ , SiO ₄ , Fe)	3(diatoms, other eukaryotes, diazotrophs)	1

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra




Figure 1. Time-series of annual ocean carbon uptake as simulated by nine ESMs for the 2001–2100 period. Thick grey lines indicates the multi-model mean, shaded areas represent $\pm$ one standard deviation of the inter-model variations. The observational estimates (black circles) are from Landschützer et al. (2014).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra




Figure 2. Global annual contemporary carbon uptake rate vs. **(a)** global future 2090–2099 carbon uptake and **(b)** cumulative global carbon uptake over 2001–2099. The straight grey lines show the best fit linear regression across all models.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra





Figure 3. (a) The eight regional ocean basins adopted in this study, defined according to different ocean basins and latitudinal lines of 70, 45, 15° S, 15, and 60° N, and (b) correlation coefficient between regional contemporary CO₂ uptake rate with (red-bars) future uptake rate in the 2090s and (blue-bars) cumulative carbon uptake in the 21st century.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra



Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ESDD

6, 2645–2688, 2015

Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra




Figure 6. Seasonal cycle of: **(a–b)** Carbon flux, **(c–d)** NPP anomalies, and **(e–f)** SST anomalies, for the Southern Ocean region simulated by CMIP5 models for the 2001–2010 (left panels) and 2090–2099 (right panels) decades. The red (blue) shadings represents the range of G1 (G2) models that simulate anomalously strong (weak) carbon uptake (see also text on Sect. 3.3). The coloured lines represent the mean of the respective model groups, black lines represent observational estimate adopted from Landschützer et al. (2014) for CO₂ uptake and SST anomaly, and the NPP anomaly were taken from SeaWiFS data as described in Nevison et al. (2015).

Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra




Figure 7. Contemporary (2001–2010) seasonal mean of sea-to-air CO₂ fluxes from (first-column) observations-based estimates of Landschützer et al. (2014), (second-column) G1 models, and (third-column) G2 models. Each row panels depict values for different seasons (JFM, AMJ, JAS, and OND).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Southern Ocean as a constrain to reduce uncertainty in future ocean carbon sinks

A. Kessler and J. Tjiputra




Figure 8. Anomalies of decomposed $p\text{CO}_2$ components in the Southern Ocean for the 2001–2010 (left panels) and future 2090–2099 (right panels) periods: **(a, b)** overestimating models; **(c, d)** underestimating models. Values shown are from multi-model mean. The black lines and markers are estimates from observations according to Landschützer et al. (2014).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

