

Climate and carbon cycle dynamics in a CESM simulation from 850-2100 CE

Flavio Lehner^{1,2,*}, Fortunat Joos^{1,2}, Christoph C. Raible^{1,2}, Juliette Mignot^{1,2,3},
Andreas Born^{1,2}, Kathrin M. Keller^{1,2}, and Thomas F. Stocker^{1,2}

¹Climate and Environmental Physics, University of Bern, Switzerland

²Oeschger Centre for Climate Change Research, University of Bern, Switzerland

* now at National Center for Atmospheric Research, Boulder, USA

³LOCEAN Laboratory, Sorbonne Universités, France

Correspondence to: Flavio Lehner (lehner@climate.unibe.ch)

Abstract. Under the protocols of the Paleoclimate and Coupled Modelling Intercomparison Projects a number of simulations were produced that provide a range of potential climate evolutions from the last millennium to the end of the current century. Here, we present the first simulation with the Community Earth System Model (CESM), which includes an interactive carbon cycle, that covers the
5 last millennium. The simulation is continued to the end of the twenty-first century. Besides state-of-the-art forcing reconstructions, we apply a modified reconstruction of total solar irradiance to shed light on the issue of forcing uncertainty in the context of the last millennium. Nevertheless, we find that structural uncertainties between different models can still dominate over forcing uncertainty for quantities such as hemispheric temperatures or the land and ocean carbon cycle response. Comparing
10 with other model simulations we find forced decadal-scale variability to occur mainly after volcanic eruptions, while during other periods internal variability masks potentially forced signals and calls for larger ensembles in paleoclimate modeling studies. At the same time, we fail to attribute millennial temperature trends to orbital forcing, as has been suggested recently. The climate-carbon cycle sensitivity in CESM during the last millennium is estimated to be between 1.0-2.1 ppm °C⁻¹. How-
15 ever, the dependence of this sensitivity on the exact time period and scale illustrates the prevailing challenge of deriving robust constraints on this quantity from paleoclimate proxies. In particular, the response of the land carbon cycle to volcanic forcing shows fundamental differences between different models. In CESM the tropical land dictates the response to volcanoes with a distinct behavior for large and moderate eruptions. Under anthropogenic emissions, global land and ocean carbon uptake
20 rates emerge from the envelope of interannual natural variability as simulated for the last millennium by about year 1947 and 1877, respectively.

1 Introduction

The last about 1,000 years constitute the best opportunity previous to the instrumental period to study the transient interaction of external forcing and internal variability in climate, atmospheric CO₂, and the carbon cycle on interannual to multi-decadal time scales. In fact, the instrumental record is often too short to draw strong conclusions on multi-decadal variability. The relatively stable climate together with the abundance of high-resolution climate proxy and ice core data makes the last millennium an interesting target and testbed for modeling studies. Yet, the large and sometimes controversial body of literature on the magnitude and impact of solar and volcanic forcing on interannual to multi-decadal climate variability illustrates the challenges inherent in extracting a robust understanding from a period that is characterized by a small signal-to-noise ratio in many quantities and for which uncertainties in the external forcing remain (e.g., Wanner et al., 2008; PAGES 2k network, 2013; Schurer et al., 2013). In addition, a process-based quantitative explanation of the reconstructed preindustrial variability in atmospheric CO₂ and carbon fluxes is largely missing.

Compared to glacial-interglacial climate change, the last millennium experienced little climate variability, yet there is evidence for distinct climate states during that period (e.g., Lehner et al., 2012b; Keller et al., 2015). Within the last millennium the Medieval Climate Anomaly (MCA, ~950-1250 AD) and the Little Ice Age (LIA, ~1400-1700 AD) are two key periods of documented regional or global temperature excursions suggested to be driven by a combination of stronger solar irradiance and reduced volcanic activity and vice versa, respectively (e.g., Crowley, 2000; Mann et al., 2009; PAGES 2k network, 2013). Despite large efforts in reconstructing (e.g., PAGES 2k network, 2013) and simulating (e.g., Fernandez-Donado et al., 2013; Masson-Delmotte et al., 2013) the transition from the MCA to the LIA, substantial uncertainties remain with respect to the mechanisms at play. Recent studies point towards solar insolation playing a minor role for climate over the last millennium (Ammann et al., 2007; Schurer et al., 2014), while in turn regional feedback processes in response to volcanic eruptions and solar variability need to be considered to explain decadal-scale climate variability (e.g., Lehner et al., 2013; Moffa-Sanchez et al., 2014). At high northern latitudes, the importance of millennial-scale orbital forcing is another debated issue (e.g., Kaufman et al., 2009).

Further, the last millennium offers the possibility to study the natural variability of the carbon cycle and its response to external forcing. Models with a carbon cycle module are extensively tested against present-day observations and widely used for emission-driven future projections (e.g., Hoffman et al., 2014). Yet, there are only few last millennium simulations including a carbon cycle (e.g., Gerber et al., 2003; Jungclaus et al., 2010; Brovkin et al., 2010; Friedrich et al., 2012). The sensitivity of the carbon cycle to climate has been shown to be mostly positive, i.e., with warming additional CO₂ is released to the atmosphere (Ciais et al., 2013). However, the magnitude of this feedback remains poorly constrained by observations (Frank et al., 2010) and models (e.g. Friedlingstein et al., 2006). In particular, determining the role of the land in past and future car-

bon cycle variability and trends is still challenging. In both idealized (Doney et al., 2006; Joos et al.,
60 2013) and scenario-guided multi-model studies (Friedlingstein et al., 2014) the land constitutes the
largest relative uncertainty in terms of intermediate- to long-term carbon uptake.

As for physical climate quantities, explosive volcanic eruptions constitute an important forcing
for the carbon cycle. The sensitivity of the carbon cycle to such eruptions have been investigated
by Jones and Cox (2001), Frölicher et al. (2011), Brovkin et al. (2010), or tji using different Earth
65 System Models. For this short-lived forcing, the land response appears to be the driver of most post-
eruption carbon cycle changes, with a range of magnitudes and time horizons associated with the
different models. Further, Frölicher et al. (2013) pointed out that the magnitude of the carbon cycle
response to volcanoes depends critically on the climate state during the eruption.

The third Paleoclimate Modelling Intercomparison Project (PMIP3; Schmidt et al., 2011) and
70 fifth Coupled Model Intercomparison Project (CMIP5; Taylor et al., 2012) represent joint efforts, in
which different modeling groups perform identical experiments, allowing for a systematic compari-
son of the models (e.g., Schmidt et al., 2014). Here we contribute to the existing set of simulations an
integration from 850-2100 CE with the Community Earth System Model, including a carbon cycle
module. The aims of this study are (i) to detect coherent large-scale features of forced variability in
75 temperature and carbon cycle quantities, in particular in response to volcanic eruptions, (ii) to inves-
tigate the relative role of forcing uncertainty and model structural uncertainty, and (iii) to provide
an estimate of preindustrial variability and the time of emergence from it under anthropogenic cli-
mate change. The setup chosen here is unique in a number of ways and tailored to address the aims
mentioned before: first, the carbon cycle is fully interactive with the other model components with
80 the exception of the radiation code, which is fed by reconstructed CO₂. This allows us to study the
isolated effect of climate on the carbon cycle, while guaranteeing an external forcing consistent with
existing reconstructions. Second, the orbital parameters are held constant to study their importance
relative to simulations with transient orbital parameters. Third, the solar forcing incorporated in the
simulation has a larger amplitude than the majority of PMIP3 simulations and hence enables us to
85 investigate whether the results are sensitive to this amplitude.

This paper is structured as follows: a description of the model and experimental setup is presented
in Section 2. In Sections 3 and 4 we address the general simulated climate and carbon cycle evolution
and investigate forced and unforced variability of the simulated climate by comparing models to
reconstructions and models to models. Sections 5 focuses on the response of the climate and carbon
90 cycle to volcanic forcing. Section 6 deals with estimating the climate-carbon cycle sensitivity in
CESM. A discussion and conclusions follow in Section 7.

2 Data and methods

2.1 Model description

The Community Earth System Model (CESM; Hurrell et al., 2013) is a fully-coupled state-of-the-art Earth System Model developed by the National Center for Atmospheric Research (NCAR) and was released in 2010. In terms of physics, CESM relies on the fourth version of the Community Climate System Model (CCSM4; Gent et al., 2011). Additionally, a carbon cycle module is included in CESM's atmosphere, land, and ocean components. The CESM version used here is release 1.0.1 in the so-called 1° version and includes components for the atmosphere, land, ocean, and sea ice, all coupled through a flux coupler.

The atmospheric component of CESM 1.0.1 is the Community Atmosphere Model version 4 (CAM4; Neale et al., 2010), which has a finite volume core with a uniform horizontal resolution of $1.25^\circ \times 0.9^\circ$ at 26 vertical levels. The land component is the Community Land Model version 4 (CLM4; Lawrence et al., 2011), which operates on the same horizontal grid as CAM4 and includes a prognostic carbon-nitrogen cycle that calculates vegetation, litter, soil carbon, vegetation phenology, and nitrogen states. Further, it includes a prognostic fire module, which is governed by near-surface soil moisture conditions and fuel availability.

The ocean component is the Parallel Ocean Program version 2 (POP2; Smith et al., 2010; Danabasoglu et al., 2012) with an nominal 1° horizontal resolution and 60 depth levels. The horizontal resolution varies and is higher around Greenland, to where the North Pole is displaced, as well as around the equator. Embedded in POP2 is the Biogeochemical Elemental Cycle model (BEC; Moore et al., 2004) that builds on a nutrient-phytoplankton-zooplankton-detritus food web model and distinguishes three phytoplankton functional types (Long et al., 2013). Carbon export and remineralization are parameterized according to Armstrong et al. (2002). Alkalinity, pH, partial pressure of CO₂, and concentrations of bicarbonate, and carbonate ions are diagnosed from prognostic dissolved inorganic carbon, alkalinity, and temperature- and salinity-dependent equilibrium coefficients. Biogenic calcification is implemented as being proportional to a fraction of small phytoplankton production, which is temperature-dependent. An exponential curve is prescribed to simulate the dissolution of sinking CaCO₃ (Moore et al., 2004). There exists no dependence of calcification-dissolution rates on saturation state. Organic material reaching the ocean floor is remineralized instantaneously, i.e., no sediment module is included. River discharge from CLM4 does not carry dissolved tracers but nitrogen deposition to the ocean surface has been prescribed. The sea ice component is the Community Ice Code version 4 (CICE4) from the Los Alamos National Laboratories (Hunke and Lipscomb, 2010), including elastic-viscous-plastic dynamics, energy-conserving thermodynamics, and a subgridscale ice thickness distribution. It operates on the same horizontal resolution as POP2.

2.2 Experimental setup

Table 1 provides an overview of the simulations conducted for this study. First, a 500-year control simulation with perpetual 850 CE forcing (hereafter CTRL) was branched from a 1850 CE control simulation with CCSM4 (Gent et al., 2011). However, restart files for the land component were
130 taken from a 850 CE control simulation, kindly provided by the NCAR, in which the land use maps by Pongratz et al. (2008) were applied. This procedure has the advantage that the slow-reacting soil and ecosystem carbon stocks are closer to 850 CE conditions than in the 1850 CE control simulation. A transient simulation covering the period 850-2099 CE was then branched from year 258 of CTRL. Despite the shortness of CTRL leading up to the start of the transient simulation, most
135 quantities of the surface climate, such as air temperature, sea ice, or upper ocean temperature, can be considered reasonably equilibrated at the start of the transient simulation, as the forcing levels due to TSI and most greenhouse gases are similar between 1850 and 850 CE (Landrum et al., 2013). However, weak trends in CTRL are still detectable in slow-reacting quantities such as deep ocean temperature (below 2,000 m; $\sim -0.04^\circ\text{C } 100\text{ yr}^{-1}$), Atlantic Meridional Overturning Circulation ($\sim -0.22\text{ Sv } 100\text{ yr}^{-1}$), Antarctic Circumpolar Current ($\sim 0.70\text{ Sv } 100\text{ yr}^{-1}$), dissolved in-
140 organic carbon in the ocean ($\sim -0.01\% 100\text{ yr}^{-1}$), or soil carbon storage ($\sim 0.2\% \text{ C } 100\text{ yr}^{-1}$). The Antarctic Bottom Water formation rate shows no drift.

The applied transient forcing largely follows the PMIP3 protocols (Schmidt et al., 2011) and the Coupled Model Intercomparison Project 5 (CMIP5; Taylor et al., 2012), consisting of total solar irradiance (TSI), greenhouse gases (GHGs), volcanic and anthropogenic aerosols, and land use changes
145 (Fig. 1). Here, the TSI reconstruction by Vieira and Solanki (2010, $TSI_{V_{S09}}$) is used, to which a synthetic 11-year solar cycle is added (Schmidt et al., 2011). In light of the recently enlarged envelope of reconstructed TSI amplitude (Schmidt et al., 2012), we scale TSI by a factor of 2.2635 to have an amplitude of 0.2% between present-day (1961-1990 CE) and the late Maunder Minimum
150 (1675-1704 CE), which is about twice as large as the 0.1 % used in most PMIP3 simulations:

$$TSI = 2.2635 \cdot (TSI_{V_{S09}} - \overline{TSI_{V_{S09}}}) + \overline{TSI_{V_{S09}}}. \quad (1)$$

Fig. 1a shows that the TSI used here lies in between the large-amplitude reconstruction by Shapiro et al. (2011) and the bulk of small-amplitude reconstructions of the original PMIP3 protocol (Schmidt et al.,
155 2011). Note, that a recent detection and attribution study indicates small amplitude TSI reconstructions to agree better with temperature reconstructions over the last millennium than large amplitude reconstructions (Schurer et al., 2014), in agreement with Ammann et al. (2007). For the twenty-first century the last three solar cycles of the data set are repeated continuously. The insolation due to Earth's orbital configuration is calculated according to Berger (1978) with the orbital parameters
160 held constant at 1990 CE values.

The volcanic forcing follows Gao et al. (2008) from 850-2001 CE. It provides estimates of the stratospheric sulfate aerosol loadings from volcanic eruptions as a function of latitude, altitude, and month and is implemented in CESM as a fixed single-size distribution in the three layers in the lower stratosphere (Neale et al., 2010). Post-2001 CE volcanic forcing remains zero.

165 Land use and land use changes (LULUC) are based on Pongratz et al. (2008) from 850 to 1500 CE, when this dataset is splined into Hurtt et al. (2011), a synthesis dataset that extends into the future. The two datasets do not join smoothly but exhibit a small step-wise change in the distribution of crop land and pasture at the year 1500 CE. Up until about 1850 CE global anthropogenic LULUC are small, however, can be significant regionally (Hurtt et al., 2011). Towards the industrial era LULUC
170 accelerate, dominated by the expansion of crop land and pasture. Here, only net changes in land use area are considered. The impact of shifting cultivation and wood harvest on carbon emissions from land use is neglected; these processes are estimated to have contributed order 30% to the total carbon emissions from land use (Shevliakova et al., 2009; Houghton, 2010; Stocker et al., 2014).

The temporal evolution of long-lived greenhouse gases (GHGs: CO₂, CH₄, and N₂O) is prescribed
175 based on estimates from high-resolution Antarctic ice cores that are joined with measurements at mid-twentieth century (Schmidt et al., 2011, and references therein). While the carbon cycle module of CESM interactively calculates the CO₂ concentration originating from land use changes, fossil fuel emissions (post-1750 CE, following Andres et al., 2012), and carbon cycle-climate feedbacks, it is radiatively inactive. Instead, the ice core and measured data are prescribed in the radiative code,
180 keeping the physical model as close to reality as possible. As a result, the impact of the interactive coupling of the carbon cycle module is minor for simulated climate, and limited to changes in surface conditions due to changing vegetation. For the extension of the simulation post-2005 CE the Representative Concentration Pathway 8.5 (RCP 8.5) is used, representing the unmitigated "business-as-usual" emission scenario, corresponding to a forcing of approximately 8.5 W m⁻² at the year 2100
185 (Moss et al., 2010).

Aerosols such as sulfate, black and organic carbon, dust, and sea salt are implemented as non-time-varying up to 1850 CE, perpetually inducing the spatial distributions of the 1850 CE control simulation during this time (Landrum et al., 2013). Post-1850 CE, the time-varying aerosol datasets provided by Lamarque et al. (2010, 2011) are used, whereby CESM only includes a representation
190 of direct aerosol effects. Similarly, nitrogen (NH_x and NO_y) input is held constant until it starts to be time-varying from 1850 CE onwards, also following Lamarque et al. (2010, 2011). Iron fluxes from sediments are held fixed (Moore and Braucher, 2008).

2.3 Other model simulations

Besides to output from current Model Intercomparison Projects, we compare CESM results to those
195 from a similar simulation with CCSM4 (Landrum et al., 2013) and IPSL-CM5A-LR (Sicre et al., 2013), two simulations without interactive carbon cycle. The solar and volcanic forcing reconstruc-

tions applied to CCSM4 and IPSL-CM5A-LR are identical to ours with the exception of the scaling of TSI that we applied to CESM. The goal here is to investigate the question whether different solar forcing amplitudes applied to the same physical model (CESM vs. CCSM4) have a larger effect than
200 applying the same solar forcing to two different physical model (IPSL-CM5A-LR vs. CCSM4).

Further, we compare CESM to MPI-ESM (ECHAM5/MPIOM; Jungclaus et al., 2010; Friedrich et al., 2012), a model that includes an interactive carbon cycle, to assess the robustness of the simulated climate and carbon cycle variations in response to external forcing. MPI-ESM uses Krivova et al. (2007) as TSI forcing and Crowley et al. (2008) as volcanic forcing. These differ from the CESM
205 forcing in amplitude much more than in timing and therefore allow for a comparison of the forced response. All simulations, except ours, apply transient orbital forcing and are summarized in Table 2. If not using the full ensemble of MPI-ESM, we focus on the member "mil0021". Another difference in terms of experimental setup between CESM and MPI-ESM is that MPI-ESM was run with a fully interactive carbon cycle, i.e., the prognostic CO₂ interacts with the radiation and through that again
210 influences climate, while in our setup this is a one-directional interaction only. Further, MPI-ESM is coarser resolved than CESM in both ocean and atmosphere and applies the A1B scenario for the twenty-first century (IPCC, 2000), which corresponds roughly to the current intermediate scenario as compared to the high scenario RCP8.5 used in CESM.

3 General climate and carbon cycle evolution

215 3.1 Temperature

The simulated annual mean Northern Hemisphere (NH) surface air temperature (SAT) follows the general evolution of proxy reconstructions: a warm Medieval Climate Anomaly (MCA, ~950-1250 CE), a transition into the colder Little Ice Age (LIA, ~1400-1700 CE), followed by the anthropogenically driven warming of the nineteenth and twentieth century (Fig. 3). The NH MCA-
220 to-LIA cooling amounts to $0.26 \pm 0.18^\circ\text{C}$ (taking the time periods defined above, which are as in Mann et al., 2009), placing it in the lower half of reconstructed amplitudes that range from about 0.1 to 0.7°C (IPCC, 2013). The subsequent warming from 1851-1880 CE to 1981-2010 CE amounts to $1.23 \pm 0.15^\circ\text{C}$, while observations report only $0.71 \pm 0.13^\circ\text{C}$ (Cowtan and Way, 2014). This overestimation by CESM takes place almost entirely after 1960 and arises largely from missing negative
225 forcing from the indirect aerosol effect, which is not implemented in CAM4 (Meehl et al., 2012). The late twentieth century being the warmest period in the NH in the past millennium is consistent with reconstructions (e.g., PAGES 2k network, 2013).

In CESM, the inception of the NH LIA occurs in concert with decreasing TSI and a sequence of strong volcanic eruptions during the thirteenth century. Reconstructions differ substantially in this
230 matter and start to cool as early as 1100 CE or as late as 1400 CE. Further, new regional multi-proxy reconstructions of temperature provide no support for a hemispherical or globally synchronous MCA

or LIA but show a clear tendency towards colder temperatures and exceptionally cold decades over most continents in the second half of the millennium (PAGES 2k network, 2013; Neukom et al., 2014).

235 The last millennium simulation with CCSM4 shows a largely coherent behavior with CESM in terms of amplitude and decadal variability of NH SAT (850-1850 CE correlation of 5-year filtered annual means $r = 0.88$, $p < 0.001$). The difference in NH SAT due to the different TSI amplitudes in CESM and CCSM4 scales roughly with the regression slope of NH SAT vs. TSI of both CESM and CCSM4 (~ 0.13 °C per W m^{-2}), although internal variability can easily mask this effect at times.
240 For example, the Maunder Minimum (1675-1704 CE), the 30-year period with the lowest TSI values and – by construction of the TSI scaling – with the largest difference between CESM and CCSM4 (1.5 W m^{-2}), is only 0.14 °C cooler than in CCSM4 and not 0.20 °C as expected from the regression.

The NH temperature evolution of additional PMIP3 and CMIP5 simulations shows that the multi-model range is within the one of the reconstructions and encompasses the instrumental-based observations (Fig. 3). Disagreement between models and reconstructions exists in particular on the
245 magnitude of response to the eruptions at 1258 CE and around 1350 CE. The 1258 CE eruption is the largest volcanic event recorded for the last millennium and its climatic impact was likely enhanced through the cumulative effect of three smaller eruptions following shortly after (Gao et al., 2008; Crowley et al., 2008; Lehner et al., 2013). However, the pronounced cooling that is simulated
250 by the models for this cluster of eruptions is largely absent in temperature reconstructions. Conversely, around 1350 CE temperature reconstructions show a decadal-scale cooling presumably due to volcanoes that is absent in the models, as the reconstructed volcanic forcing shows only two relatively small eruptions around that time. Part of this incoherent picture may arise from the unknown aerosol size distribution (Timmreck et al., 2010) and geographic location of past volcanic eruptions
255 (Schneider et al., 2009), and differences in reconstruction methods. As many proxy reconstructions of temperature rely heavily on tree ring data it is worth noting that the dendrochronology community currently debates whether the trees' response to volcanic eruptions resembles the true magnitude of the eruption (Mann et al., 2012; Anchukaitis et al., 2012; Tingley et al., 2014).

Disagreement among the models exists on the relative amplitude of the MCA, where most models
260 show colder conditions than CESM and CCSM4. Remarkably, the simulation by IPSL-CM5A-LR applied the same TSI and volcanic forcing as CCSM4, yet it comes to lie at the lower end of the PMIP3 model range during the MCA. In other words, the way how models respond to variations in TSI and other forcings can still make a larger difference in the simulated amplitude than the scaling of TSI by a factor of 2, which in turn complicates a proper detection and attribution of solar
265 forcing during the last millennium (Servonnat et al., 2010; Schurer et al., 2014). Further disagreement among the models exists on the response to volcanic eruptions, where CESM and CCSM4 are among the more sensitive models (an oversensitivity of CCSM4 to volcanoes based on twentieth century simulations was reported by Meehl et al., 2012). Turning to the century-scale change

over the industrial era, CESM and CCSM4 are on the upper end of the CMIP5 range and show an
270 overestimation of the observed warming.

The simulated mean SAT of the Southern Hemisphere (SH) generally shows a similar evolution
as for the NH with the signature of the MCA and LIA superimposed on a weak millennial cooling
trend. Models and reconstructions disagree to a larger extent in the SH than in the NH, in particular
regarding cold excursions due to large volcanic eruptions, which are largely absent in the reconstruc-
275 tions. Similar results have been reported in a recent study on interhemispheric temperature variations
that finds much less phasing of the two hemispheres in reconstructions than in models, potentially
related to underestimated internal variability on the SH in models (Neukom et al., 2014). A linger-
ing question of climate modeling in general is whether models are too global in their response to
external forcing. That is, they might show too little regional variability that is independent from the
280 global mean response during a forced period. However, the uncertainties in the early period of the
reconstructions prohibits to robustly answer this question. Similar to the NH, the industrial warm-
ing in the SH from 1851-1880 CE to 1981-2010 CE ($0.53 \pm 0.07^\circ\text{C}$) is overestimated by CESM
($0.71 \pm 0.13^\circ\text{C}$).

The differential warming between the hemispheres in CESM is among the smallest among CMIP5
285 models (not shown). This is mainly due to the underestimated deep water formation in the South-
ern Ocean, leading to a comparably strong warming of the SH and likely an underestimation of
the oceanic uptake of anthropogenic carbon (Long et al., 2013). With a transient climate response
of 1.73°C and an equilibrium climate sensitivity of 3.20°C (Meehl et al., 2012), CESM lies in the
middle of recent estimates of 1.0 to 2.5°C and 1.5 to 4.5°C , respectively (IPCC, 2013).

290 3.2 Orbital forcing

To detect and attribute the influence of orbital forcing on SAT trends during the last millennium, we
compare our simulation with fixed orbital parameters to the CCSM4 simulation with time-varying or-
bital parameters (Fig. 2). While both models experience a negative long term trend in global TSI until
about 1850 CE (Fig. 1), the difference arising from the different orbital setup can be seen best in Arc-
295 tic summer land insolation (Fig. 2). Hence, Arctic summer land SAT has been proposed as a quantity
to be affected by orbital forcing already on time scales of centuries to millennia (Kaufman et al.,
2009). However, we find no detectable difference between the two simulations in the trend of Arctic
summer land SAT (Fig. 2b). In fact, the Arctic multi-decadal to centennial summer land SAT anom-
alies in CESM span a very similar range as in CCSM4, despite CESM not accounting for time-varying
300 orbital parameters: Fig. 2c shows non-overlapping 100- and 200-year mean SAT anomalies plotted
against the corresponding mean solar insolation. The results from CCSM4 suggest a clear relation
of the two quantities, however, the results of CESM show that nearly identical SAT anomalies are
possible without orbital forcing. In other words, while we detect a long-term cooling trend in Arctic
summer SAT in both CESM and CCSM4, we fail to attribute this trend to orbital forcing alone, as

305 suggested by Kaufman et al. (2009). This is confirmed in new simulations with decomposed forcing, again comparing simulations with fixed and time-varying orbital parameters (B. Otto-Bliesner, personal communication).

3.3 Carbon cycle

The prognostic carbon cycle module in CESM allows us to study the response of the carbon cycle to transient external forcing. The land biosphere is a carbon sink during most of the first half of the last millennium, but becomes a source as anthropogenic land cover changes start to have a large-scale impact on the carbon cycle (Table 3). The ocean is a carbon source at the beginning and becomes a sink in the second half of the last millennium. The residual of these fluxes represents changes in the atmospheric reservoir of carbon, illustrated in Fig. 3c by the prognostic CO₂ concentration. 310
315 The amplitude of the simulated concentration does not resemble the one reconstructed from ice cores (i.e., imposed on the radiative code of CESM), in particular the prominent CO₂ drop in the seventeenth century is not captured by CESM. This raises the question whether the sensitivity of the carbon cycle to external forcing is too weak in CESM, whether the imposed land use changes are too modest (Kaplan et al., 2011; Pongratz et al., 2011), whether major changes in ocean circulation are not captured by models (Neukom et al., 2014), or whether the ice core records are affected by 320 uncertainties due to in-situ production of CO₂ (Tschumi and Stauffer, 2000). Ensemble simulations with MPI-ESM also do not reproduce the reconstructed amplitudes or the drop (Jungclaus et al., 2010). Further, Earth System Models of Intermediate Complexity or vegetation models driven by GCM output do not reproduce the uptake of carbon by either ocean or land needed to explain the 325 reconstructed amplitudes (Stocker et al., 2011; Gerber et al., 2003).

The rise in atmospheric CO₂ due to fossil fuel combustion is in good agreement with ice cores until about the 1940s. After that, a growing offset exists, leading to an overestimation of about 20 ppm by 2005 in CESM, qualitatively similar to the CMIP5 multi-model mean (Hoffman et al., 2014). From the observational estimates one can diagnose that the discrepancy arises primarily from 330 overestimated carbon release from land (Fig. 4a; see also Hoffman et al., 2014; Lindsay et al., 2014). From 1750 to 2011 CE the cumulative total land release (including LULUC) is 83 Pg C (compared to 30±45 Pg C from observational estimates Ciais et al., 2013), while the cumulative net land uptake is 95 Pg C (160±90 Pg C Ciais et al., 2013). The ocean cumulative uptake of 151 Pg C compares more favorably to current estimates of 155±30 Pg C (Ciais et al., 2013). Note, however, that given the 335 overestimation of atmospheric CO₂, one would expect a higher ocean uptake. This bias originates largely from an underestimation of the uptake in the Southern Ocean (Long et al., 2013). Along with this goes an underestimated seasonal cycle in CESM, originating from a too weak growing season net flux in CLM4 (Keppel-Aleks et al., 2013). MPI-ESM, on the other hand, underestimates atmospheric CO₂ due to weak emissions from LULUC (Pongratz et al., 2008).

340 The twenty-first century sees substantial emissions from fossil fuel burning under RCP 8.5 (Fig. 3c).
In addition, LULUC is associated with a positive flux into the atmosphere, particularly until around
2050 CE (Table 3). After accounting for LULUC (which constitutes a carbon loss for the land) the
net land sink increases to about 7 Pg C yr^{-1} at the end of the twenty-first century (Fig. 4a). The rate
of ocean uptake, on the other hand, peaks around 2070 at about 5 Pg C yr^{-1} , despite that atmospheric
345 CO_2 continues to rise (Fig. 3c). This decoupling of the trends in atmospheric CO_2 growth and ocean
uptake flux is linked to non-linearities in the carbon chemistry (Schwinger et al., 2014). The change
in dissolved inorganic carbon per unit change in the partial pressure of CO_2 decreases with increas-
ing CO_2 , and thus the uptake capacity of the ocean. Additionally, differences in the ventilation time
scales of the upper and the deep ocean likely play a role. While the surface ocean and the thermo-
350 cline exchanges carbon on annual-to-multi-decadal time scales with the atmosphere, it takes century
to ventilate the deep ocean as evidenced by chlorofluorocarbon and radiocarbon data (Key et al.,
2004; Wunsch and Heimbach, 2007, 2008). CESM has a documented low bias in Southern Ocean
ventilation due to too shallow mixed layer depths, contributing to the underestimated carbon uptake
of the ocean (Long et al., 2013).

355 The prognostic atmospheric CO_2 increases to 1,156 ppm by 2100 CE. This would imply a forcing
of 7.6 W m^{-2} from CO_2 relative to 850 CE, significantly more than the approximately 6.5 W m^{-2} that
are imposed by the radiative code (calculated according to IPCC, 2001, , see also Fig. 1c). This prop-
agation of the twentieth century bias is consistent with the CMIP5 multi-model mean (Friedlingstein et al.,
2014) and has motivated attempts to reduce such biases by using observational constraints for ocean
360 ventilation (Matsumoto et al., 2004), the tropical land carbon storage sensitivity to temperature vari-
ations (Cox et al., 2013; Wenzel et al., 2014; Wang et al., 2014) and for the oceanic and terrestrial
carbon fluxes (Steinacher et al., 2013). CESM with CLM4, however, shows very little sensitivity in
tropical land carbon, in part due to the inclusion of an interactive nitrogen cycle, which – through
enhanced photosynthetic uptake due to nitrogen fertilization – tends to counteract accelerated soil
365 decomposition from warming (Lawrence et al., 2012; Wenzel et al., 2014). Together with the under-
estimated oceanic uptake this leads to the roughly 20% larger airborne fraction in CESM as com-
pared to what is actually prescribed as atmospheric concentration in the radiative code according to
the RCP 8.5.

Fig. 4 puts the current and projected changes into perspective of preindustrial variability. Esti-
370 mated interannual variability prior to 1750 CE is $\pm 0.94 \text{ Pg C yr}^{-1}$ (1 standard deviation) for the
net atmosphere-land and $\pm 0.42 \text{ Pg C yr}^{-1}$ for the net atmosphere-ocean flux. The much larger in-
terannual variability in land than ocean flux is consistent with independent estimates and results
from other models (e.g., Ciais et al., 2013). Large volcanic eruptions, as they have occurred in
the last millennium, cause anomalously high uptake rates that for a short period of time are on
375 par with current uptake rates (Fig. 4a and b, full range). We estimate when the anthropogenically
forced, global-mean land and ocean uptake fluxes leave the bound of preindustrial natural variabil-

ity (Hawkins and Sutton, 2012; Keller et al., 2014). As a threshold criteria, it is required that the decadal-smoothed uptake fluxes are larger than the upper bound of 2 standard deviations of the annual fluxes prior to 1750 CE. Then, the simulated global-mean land and ocean uptake fluxes have emerged from natural interannual variability by 1947 CE and by 1877 CE, respectively. The prognostic atmospheric CO₂ concentration emerges already in 1755 CE, while the simulated global-mean temperature does not emerge until 1966 CE.

4 Model-model coherence

A classical approach to assess the robustness of model results is to rely on the multi-model mean response to a given forcing (IPCC, 2013). However, as there are only very few last millennium simulations with comprehensive Earth System Models to date, this approach is not feasible to investigate the decadal-scale climate-carbon cycle responses to external forcing in the period before 1850 CE. Instead, we estimate periods of forced variability with a 100-year running-window correlation of CESM and MPI-ESM, indicating phasing of the two models. The time series are anomalies to their 850-1849 mean and are smoothed with a 5-year local regression filter before calculating the correlation. Thereby, we focus on the preindustrial period, as the twentieth and twenty-first century are dominated by anthropogenic trends, which are non-trivial to remove for a proper correlation analysis. In addition, regression analysis is used.

4.1 Temperature

Fig. 5a and b show anomalies of zonal mean annual SAT from CESM and MPI-ESM. In both models the northern high latitudes show the strongest trend from positive anomalies during the MCA to negative anomalies during the LIA. This is consistent with the current understanding of polar amplification during either warm or cold phases (Holland and Bitz, 2003; Lehner et al., 2013). The twentieth and twenty-first century then see the strong anthropogenic warming, although this occurs earlier in CESM due to missing negative forcings from indirect aerosol effects (section 2). Superimposed on the preindustrial long-term negative trend are volcanic cooling events. In CESM many of these are global and are able to considerably cool the SH extra-tropics around 60° S, while in MPI-ESM the SH extra-tropics are only weakly affected. These differences are likely related to the Southern Ocean heat uptake rates in the two models (arising from under- and overestimation of Southern Ocean mixed layer depths in CESM and MPI-ESM, respectively; Danabasoglu et al., 2012; Marsland et al., 2003). This is evident also in the delayed warming at these latitudes in the twenty-first century in MPI-ESM as compared to CESM. The consistent SH high latitude positive anomalies before the thirteenth century, on the other hand, appear to be related to a positive phase of the Southern Annular Mode (SAM) in both models (not shown), a behavior common to most PMIP3 models. Note, however, that a recent reconstruction of the SAM finds the models to lack amplitude

in their simulated variability, challenging the models' capabilities to represent SAM (Abram et al., 2014).

The phasing on interannual to decadal scales between the two models is largely restricted to periods of volcanic activity and within those mainly to land-dominated latitudes (except Antarctica, 415 which shows no forced variability on these time scales; Fig. 5c). Despite the largest absolute temperature anomalies occurring in the Arctic, the correlations are highest in the subtropics, due to the smaller interannual variability there. Periods of centennial trends, such as the MCA or the Arctic cooling during the Maunder Minimum around 1700 CE, do not show up in the correlation analysis that focuses on 100-year windows, suggesting multi-decadal low-frequency forcing, such as centennial TSI trends, or internal feedback mechanisms to be responsible for the missing correlation. A 420 regression analysis between the 5-year filtered annual TSI and SAT at each gridpoint (different filter lengths of up to 50 years have been tested as well without changing the results) reveals a clear link of the two quantities at high latitudes. In CESM this seems to be driven primarily by a displacement of the sea ice edge (Arctic) and Southern Ocean heat uptake (Fig. 6a). As the sea ice response has not 425 been detected in an earlier model version (Ammann et al., 2007, their Fig. 4), it warrants the questions whether the regression of SAT on TSI might be biased by imprints of volcanoes (Lehner et al., 2013), even when the timeseries are filtered, especially in a model like CESM that has a very strong volcanic imprint.

Forthcoming simulations with solar-only forcing will be able to answer that question. MPI-ESM, 430 on the other hand, shows a similar polar amplification signal from solar forcing, but not as clearly linked to sea ice (Fig. 6b). MPI-ESM also displays a stronger land-ocean contrast than CESM (see also Geoffroy et al., 2015).

In addition to the comparison with MPI-ESM, Fig. 5d shows results from the correlation analysis between CESM and CCSM4, two simulations that in terms of physics differ only in their applied TSI 435 amplitude and orbital parameters. Not unexpected, there are generally more robust signals of forced variability as compared to CESM vs. MPI-ESM (Fig. 5c), very likely due to the identical physical model components in CESM and CCSM4. Similarly, global mean SAT shows generally stronger phasing between CESM and CCSM4 (Fig. 5e). However, the latitudinal and temporal pattern of the CESM vs. CCSM4 analysis agrees well with the one arising from CESM vs. MPI-ESM (Fig. 5c; 440 with exception of the much stronger phasing in CESM and CCSM4 during the volcanic eruptions in the 1450s) and suggest the physical mechanism behind periods of phasing to be robust across the two models.

Applied to ocean temperature, the above approach enables us to investigate the penetration depth of a forced signal seen at the surface (Fig. 7). Indeed, most of the surface signals also show up as 445 significant correlations down to depths of about 150-200 m, whereby their timing suggests again volcanic forcing as the origin. Reduced heat loss from the tropical equatorial Pacific together with reduced heat uptake at high latitudes are responsible for ocean cooling after volcanoes (not shown).

The Atlantic Meridional Overturning Circulations (AMOC) in the CESM and MPI-ESM shows no significant correlation, however, the highest correlation occurs during the thirteenth century and coincides with a phasing of the upper ocean temperatures due to strong volcanic forcing (Fig. 7d). The correlation between CESM and CCSM4 at that time is even higher and points to a significant imprint of the volcanic forcing on ocean circulation (Otterå et al., 2010; Swingedouw et al., 2013, 2015). However, during the remainder of the millennium, no phasing of the AMOC is found.

4.2 Carbon cycle

We apply the same correlation analysis to zonally integrated land and ocean carbon fluxes from the two models to detect forced variability in the carbon cycle. Compared to SAT hardly any phasing can be found between the models in atmosphere-to-land carbon fluxes (not shown), which is due to its large interannual variability and to distinctly different responses to external forcing in the two models, as will be illustrated in section 5. Similarly, there is little model phasing in net atmosphere-to-ocean carbon fluxes (not shown). Results become somewhat clearer when considering globally integrated upper-ocean dissolved inorganic carbon (DIC; Fig. 8). There appear to exist spurious trends in CESM, likely related to model drift. We repeated the analysis, but with the CESM output detrended in each grid cell by subtracting the CTRL over the corresponding period 850-1372 CE. Due to the shortness of CTRL, we cannot apply this to the whole simulation. However, these tests showed that the correlation between the two simulation is largely insensitive to the drift in CESM. In Fig. 8c there are periods of coherent carbon draw-down coinciding with volcanic eruptions around 1450 CE and 1815 CE in response to temperature-driven solubility changes. Interestingly, MPI-ESM shows a distinct behavior for the strong eruption of 1258 CE, with a prolonged ocean carbon loss after a weak initial uptake. CESM shows a stronger and more sustained carbon uptake, leading to no correlation between the two models for this eruption. The reasons for this discrepancy are discussed in section 5.

Generally, the largest changes in upper-ocean carbon storage occur in response to volcanoes and take place in the tropical Pacific (Chikamoto et al., prep), with other significant changes occurring in the North and South Pacific, the subtropical Atlantic and the Arctic (section 5). Within the tropical oceans, the models show different characteristics: CESM shows a larger variability in DIC than MPI-ESM and, when influenced by anthropogenic emissions in the twentieth and twenty-first century, takes up a larger portion of the total ocean carbon uptake than in MPI-ESM (not shown). In MPI-ESM, the Southern Ocean shows stronger variability and larger carbon uptake in the twenty-first century, illustrating the different behavior of the two models in terms of ocean carbon cycle variability and trend magnitude, closely related to the different mixed layer depth in the Southern Ocean region.

5 Volcanic forcing

To further isolate the response of the climate system and carbon cycle to volcanic eruptions, a Superposed Epoch Analysis is applied to both simulations. Thereby, composite time series for the strongest three (top3) and following strongest seven eruptions (top10), by measure of optical depth anomaly, over the period 850-1850 CE are calculated for the CESM and MPI-ESM (Fig. 9). The time series are calculated as deseasonalized monthly anomalies to the 5 years preceding an eruption.

The physical parameters global mean surface air temperature and global mean precipitation decrease in both models after volcanic eruptions, although the response of CESM is stronger by roughly a factor 2-2.5 (Fig. 9a, b, f, g). Consequently, CESM temperature and precipitation take longer (~15 years) to relax back to pre-eruption values than MPI-ESM (~9 years).

The atmospheric carbon inventory, on the other hand, shows a remarkably different response in the two models. In CESM the atmosphere initially loses about 2-3 Pg C, irrespectively of the eruption strength, with the minimum occurring after about 1-2 years. In the top10 case values return to normal after about 16 years, while in the top3 case they tend to return already after about six years, and overshoot. This overshoot is not straightforward to understand and did not seem to occur in earlier versions of the model (Frölicher et al., 2011; Rothenberg et al., 2012). In MPI-ESM the response is a priori more straightforward and slower: in the top10 case the atmosphere loses about 2.5 Pg C, reaches a minimum after 2-4 years, and returns to pre-eruption values after 10-16 years. The top3 case reaches its minimum (-6 Pg C) a bit faster, but then takes about 20 years to return to pre-eruption values (Brovkin et al., 2010).

Partitioning these atmospheric carbon changes into land and ocean changes indicates that the land is primarily responsible for the differing response behavior of the two models, confirming the findings in the previous section. While in both models the land drives the atmospheric change by taking up carbon initially, it is released back to the atmosphere within about 3 years in CESM, but kept in the land for at least 15 years in MPI-ESM (and up to 50 years for the 1258 CE eruption; Brovkin et al., 2010). In the top3 case of CESM the land starts to even lose carbon after about 5 years, causing the overshoot seen in the atmospheric carbon.

A closer look at CESM reveals a distinct response to the top3 and the top10 volcanoes. The response to top3 must be understood as an interplay of a number of processes: the initial global cooling triggers a La Niña-like response and a corresponding cloud and precipitation reduction that is particularly pronounced over tropical land, where also large changes in carbon storage occur (see Fig. 11a-c for the spatial pattern). Fig. 10 and the following analysis therefore focuses on tropical land. Direct solar radiation decreases, indirect radiation increases, with a net decrease (Fig. 10d). These unfavorable conditions cause a reduction in net primary productivity and a strong decrease of vegetation (-8 Pg C; Fig. 10a and e). At the same time, decomposition of dead biomass becomes less efficient due to reduced temperature (similar to, e.g., Frölicher et al., 2011). Despite the simultaneous decrease in net primary production this results in a build-up of dead biomass of about 5 Pg C

(Fig. 10b). Due to the dry conditions and availability of dead biomass there is increased fire activity, leading to increased carbon loss from land. However, the fire cannot get rid of the large amount of dead biomass immediately (Fig. 10f). While vegetation decrease and dead biomass buildup balance each other, the soil takes up about 2 Pg C (Fig. 10c), stores it for at least 16 years, and is therefore responsible for the initial net land uptake seen in Fig. 9e (see also Fig. 11c left). After about two years, tropical precipitation increases again and puts a halt to the decrease in vegetation (Fig. 10a and Fig. 11b right). The vegetation does not recover fully for another about 20 years. The dead biomass, on the other hand, gets decomposed entirely within about 15 years and therefore turns the land into a carbon source, causing the overshoot in CO₂. In the top10 case, the precipitation and radiation response is about half of the top3 case, and so is the vegetation decrease. Consequently, vegetation recovers faster. The decomposition of dead biomass, however, takes about the same amount of time as in the top3 case as the decomposition rates are similar for both cases. Hence, the land acts as a more sustainable carbon sink in the top10 case. In MPI-ESM it is the soil as well which acts as main land carbon storage pool, while the vegetation decrease is significantly less than in CESM (Brovkin et al., 2010), leading to the different response behavior of the two land models, particularly striking in the top3 case. Note that there are subtle regional differences between CESM and the earlier version of the carbon cycle-enabled NCAR model CSM1.4-carbon (Frölicher et al., 2011): tropical Africa sees a reduction of land carbon in CESM, related to a persistent increase in cloud cover and precipitation after volcanoes, while CSM1.4-carbon saw a decrease in precipitation and an increase in land carbon.

The ocean, on the other hand, shows a qualitatively similar response in CESM and MPI-ESM with an uptake of carbon and a gradual relaxation back to pre-eruption values over 20 or more years. In CESM the radiative cooling leads to increased uptake in the Western Pacific, while in the Eastern Pacific, cooling is less as this region is more controlled by upwelling rather than direct radiative forcing, as suggested by Maher et al. (2014) (Fig. 11d). Two or more years after the volcano a La Niña-like pattern settles in both surface temperature as well as carbon uptake. Some model differences exist, e.g., in the top3 case of MPI-ESM the ocean starts to release carbon, compensating the persistent positive anomaly in the land inventory (imposed on the ocean via atmospheric CO₂ concentration Brovkin et al., 2010), a feature not present in CESM, in which the land does not store the anomalous carbon as long. In CESM the tropical oceans appear to be more sensitive to volcanic forcing than to TSI variations. The equatorial Pacific shows the strongest response in DIC to volcanoes (Fig. 11d), while the response to TSI variations of comparable radiative forcing is up to an order of magnitude weaker and confined to higher latitudes (not shown). Overall it seems therefore that the response of the land vegetation governs the overall different responses in the two models.

In an attempt to validate the two models, one is restrained to the well-observed eruption of Pinatubo in 1991 CE, as the CO₂ records from ice cores do not adequately resolve short-term variations induced by volcanoes over the last millennium. Fig. 12 shows the global temperature and atmo-

spheric carbon response to Pinatubo as extracted from observations, CESM, and the 3-member ensemble of MPI-ESM. Note that the effects of El Niño-Southern Oscillation and anthropogenic emissions have been removed from the CO₂ observations to obtain a tentative estimate of the actual CO₂ response to the Pinatubo eruption (Frölicher et al., 2013). The initial cooling of about -0.5°C and the relaxation back to initial temperatures around 1998 CE is captured well by both models. The MPI-ESM ensemble, however, shows a large and robust variation around 1995 CE, seemingly related to a phasing of ENSO variability in response to the eruption (see also Zanchettin et al., 2012). Further, the magnitude of atmospheric carbon response matches better in CESM, although the overshoot of the observation-based estimate is not captured. CESM's response also falls within the range of the earlier model version (Frölicher et al., 2013). It remains unclear whether this mismatch reflects a model-deficiency or is due to uncertainties arising from removing the ENSO signal from the CO₂ observations. However, the mechanisms described above that lead to an atmospheric CO₂ overshoot for large eruptions in CESM offer an opportunity for reconciliation of this discrepancy. Further, the precipitation response (and therewith the cloud and surface short-wave response) to volcanic eruptions is not well constrained due to the small number of observed eruptions (Trenberth and Dai, 2007). Biases in the representation of these processes can influence a model's carbon cycle response.

6 Climate-carbon cycle sensitivity

Due to the absence of large anthropogenic disturbances of the carbon cycle, the last millennium represents a testbed to estimate the climate-carbon cycle sensitivity, expressed as $\text{ppm } ^{\circ}\text{C}^{-1}$, and can thus potentially help to constrain this quantity (e.g., Woodwell et al., 1998; Joos and Prentice, 2004; Scheffer et al., 2006; Cox and Jones, 2008; Frank et al., 2010). Note, however, that there exist important differences between studies in how this sensitivity is calculated and what it implies. Studies using observations and fully-coupled simulations (Frank et al., 2010; Jungclaus et al., 2010) estimate the sensitivity from the ratio of changes in CO₂ over changes in temperature. This quantity folds in feedbacks, as the initial sensitivity of the carbon cycle to climate change modifies itself via the climate change that arises from the changed carbon cycle. This is distinct from the climate-carbon cycle feedback sensitivity γ , which uses idealized simulations with atmospheric CO₂ held constant, while the climate varies naturally, to isolate the feedback parameter (Friedlingstein et al., 2006). The sensitivity that can be derived from our CESM transient simulation is subtly different again in that the carbon cycle will respond to changes in climate and this response will feed back on the carbon stocks through increased or decreased atmospheric CO₂ concentrations, yet these changes in CO₂ are not allowed to feed back on the climate. Such a sensitivity is expected to be lower than γ , which we can derive from CTRL.

Here, we estimate the climate-carbon cycle sensitivity for CESM as follows. We focus on the period before significant LULUC (850-1500 CE) and apply different low-pass filters of 20 to 120 years,

taking 5-year increments, to the time series of NH SAT and global CO₂. The filtering aims at minimizing the influence of short-lived forcings such as volcanic eruptions that have a relatively direct impact on temperature and CO₂ (as seen above) and thus may hinder the detection of a low-frequency influence of temperature on CO₂. For each filter length we determine the highest lag correlation of
595 the two time series, considering lags of up to 100 years. By design of our simulation we expect NH SAT to lead CO₂, which is confirmed by all lag correlations indicating positive lags for NH SAT (peak of lag correlation at 80.5 ± 3.4 years). We regress the lagged time series and find a median estimate of $1.3 \text{ ppm } ^\circ\text{C}^{-1}$ with a range from 1.0 to $1.8 \text{ ppm } ^\circ\text{C}^{-1}$, depending on the filter length. About $-1 \text{ ppm } ^\circ\text{C}^{-1}$ is explained by the land carbon cycle, while the ocean shows smaller sensitivities of
600 about $-0.4 \text{ ppm } ^\circ\text{C}^{-1}$. Note, that we use NH SAT in order to be comparable with existing studies (Frank et al., 2010; Jungclaus et al., 2010). Using global SAT instead of NH SAT can influence the sensitivity estimate, especially for the forced simulation: including the vast ocean area of the SH tends to dampen temperature variability induced by volcanoes and TSI variations. With temperature variability dampened, the sensitivity increases to $1.7 \text{ ppm } ^\circ\text{C}^{-1}$ (1.4-2.1).

605 This estimate is barely within the reconstruction-constrained range of 1.7 - $21.4 \text{ ppm } ^\circ\text{C}^{-1}$ (Frank et al., 2010) and suggests a comparably low sensitivity of the carbon cycle in CESM. This low sensitivity is in agreement with, e.g., Arora et al. (2013). Note that Frank et al. (2010) found different sensitivities for the early and late part of the last millennium with the mean for the period 1050-1549 CE being $4.3 \text{ ppm } ^\circ\text{C}^{-1}$. Indeed, a strong temporal dependence of the climate-carbon cycle sensitivity is also
610 found in CESM when looking at individual 200-year windows (Fig. 13). The period 1300-1500 CE even shows negative sensitivity, which seems to be related to the different time scales with which SAT and CO₂ relax back to the pre-eruption conditions after perturbations from large volcanic eruptions (Fig. 9a and c): atmospheric CO₂ decreases from having overshoot while SAT increases after the initial cooling, leading to a negative correlation of the two quantities.

615 This illustrates the time-variant character of the climate-carbon cycle sensitivity, which substantially complicates any attempt to constrain it by last millennium data and warrants caution when making inferences from past to future sensitivities. Besides Frank et al. (2010), Frölicher et al. (2011) found the sensitivity to vary greatly in a coupled model with the time scale and magnitude of volcanic forcing considered. This issue is further highlighted by the larger sensitivity derived for idealized $+1\% \text{-CO}_2 \text{ year}^{-1}$ simulations with CESM ($11.9 \text{ ppm } ^\circ\text{C}^{-1}$), for which a dependence on the
620 background state, the scenario, and even the method is reported (Plattner et al., 2008; Arora et al., 2013). Further, it is worth stressing that such sensitivity estimates cannot be extrapolated easily across time scales, as different processes might be at play (Ciais et al., 2013).

Applying the identical analysis to CTRL reveals other time scales of climate-carbon cycle feedback, suggesting maximum lags of less than 10 years and a sensitivity of 2.3 (1.4-2.9) $\text{ppm } ^\circ\text{C}^{-1}$.
625 Using global SAT instead of NH SAT has no discernible effect ($2.3 \text{ ppm } ^\circ\text{C}^{-1}$), as the CTRL does not see volcanoes or TSI variations. A later peak in the lag correlation of NH SAT and CO₂ clus-

ters at 73.3 ± 1.1 years in CTRL, i.e., close to where the forced simulation shows its highest lag correlation, but these lag correlations are much weaker ($r \sim 0.4$ compared to $r \sim 0.7$ in the forced simulation). This is generally consistent with the finding by Jungclaus et al. (2010) that a forced simulation exhibits increased power on lower frequencies compared to a control simulation.

7 Discussion and conclusions

This study presents a simulation from 850 to 2100 CE with the fully-coupled CESM, including carbon cycle, and provides an overview on the imprint of external forcing on different climate and carbon cycle diagnostics in the simulation. For comparison we draw on a number of PMIP3 simulations, particularly, comparable simulations with CCSM4 and MPI-ESM. The evolution of NH SAT during the preindustrial era in CESM is in reasonable agreement with both reconstructions and other models, albeit the uncertainties in reconstructions and forcing still being considerable. Comparing to more reliable data in the twentieth century, the anthropogenic warming in CESM is overestimated due to a lack of negative forcing from indirect aerosol effects. On the SH, CESM and most other models do not capture the evolution of the mean SAT as well. The discrepancies could be explained by (i) significant model biases in SH and also interhemispheric SAT variability (Neukom et al., 2014), (ii) spectral biases in proxies used in the reconstructions (Franke et al., 2013), (iii) uncertainties in the external forcing (Masson-Delmotte et al., 2013), or (iv) natural internal variability (Bothe et al., 2013). Unfortunately, these potential explanations are neither exclusive nor independent. Arguments for model bias come from the fact that reconstructed interhemispheric SAT variability lies outside the models' range over 40% of the time (Neukom et al., 2014); but these arguments are weakened by the uncertainty in external forcing. We show here that implementing the same TSI forcing in two different models results in a larger difference in simulated SAT than implementing two different TSI forcings in the same model. Hence, model structural uncertainty remains an issue in determining the role of external forcing over the last millennium.

Albeit beyond the scope of this study, detecting structural and spatial dependencies such as illustrated here offers an opportunity to reconcile the discrepancies (e.g., regarding SH volcanic signals) between reconstructions and simulations, which might originate from sampling bias, model deficiencies, a combination of these, or the fact that reality may be one realization by chance not encompassed by a multi-model ensemble (Deser et al., 2012; Lehner et al., 2012a; Bothe et al., 2013; Neukom et al., 2014).

Further, we compare simulations with and without orbital forcing and fail to attribute northern high latitude SAT trends over the last millennium to orbital forcing. This hampers, if not challenges, the validation of recent findings based on proxy archives that claim a distinct low-frequency orbital component in millennial trends (Kaufman et al., 2009; Esper et al., 2012). Instead, the decreasing

trend in annual TSI – as opposed to seasonal and regional insolation – together with local feedbacks are able to account for a similar magnitude of trend.

When forced with emissions from LULUC, TSI variations, and volcanic eruptions over the last millennium, both CESM and MPI-ESM do not reproduce atmospheric CO₂ variability as suggested by ice cores. Notably, the large drop of CO₂ in the seventeenth century is not reproduced, similar as in earlier studies (Gerber et al., 2003; Stocker et al., 2011; Jungclaus et al., 2010). Neukom et al. (2014) hypothesized that the unique, globally synchronous cooling during the LIA (which might be related to ocean dynamics) can serve as an explanation for this drop. While both CESM and MPI-ESM show a global cooling during the LIA, they develop no apparent phasing of ocean dynamics or carbon uptake and do not show any marked CO₂ reduction around that time, leaving this issue unresolved. The strong volcanic forcing during the thirteenth century, on the other hand, is able to synchronize the AMOC on decadal scales, confirming similar results from the Bergen Climate Model and IPSL-CM5A-LR (Otterå et al., 2010; Swingedouw et al., 2013). Under anthropogenic emissions, land and ocean carbon uptake rates emerge from the envelope of natural variability as simulated for the last millennium by about 1947 CE and 1877 CE, respectively. Atmospheric CO₂ and global temperature emerge by 1755 CE and 1966 CE, suggesting that changes in carbon-cycle related variables would be easier to detect than temperature given sufficient observational data (Keller et al., 2015).

We find forced decadal-scale variability in CESM and MPI-ESM in response to major volcanic eruptions in both SAT and upper-ocean temperature, while the response in carbon cycle quantities is less coherent among models (see also Resplandy et al., 2015). Outside volcanically active periods large parts of the decadal-scale variations cannot be attributed to external forcing, suggesting that internal variability masks external forcing influence. Note, however, that recent work suggest that small volcanic eruptions, which are typically not well-resolved in reconstructions of volcanic activity, exert a significant cumulative effect on global temperature and climate (Ridley et al., 2014).

Volcanoes trigger a coherent global response in SAT and precipitation that is qualitatively in line with earlier studies on the volcanic influence on climate and carbon cycle (e.g., Jones and Cox, 2001; Brovkin et al., 2010; Frölicher et al., 2011, 2013). However, the carbon cycle response, in particular on land, shows fundamental model differences in terms of perturbation amplitude and persistence after volcanic eruptions. These differences arise from a differing land vegetation responses in the two models. The extent to which such structural uncertainties matter is illustrated by the large spread in the airborne fraction of CO₂ between these two (and other) models in the twenty-first century (see also Friedlingstein et al., 2014). In particular, known biases in CESM’s carbon uptake in response to anthropogenic emissions in the twentieth and twenty-first century lead to a 20% overestimation of the atmospheric CO₂ concentration and the corresponding prognostic radiative forcing as compared to the prescribed RCP8.5 at year 2100 CE.

The climate-carbon cycle sensitivity of CESM as estimated from the anthropogenically unperturbed first part of the last millennium is between 1.0 and 2.1 ppm °C⁻¹, with a dependency on the filtering and the exact time period considered. Generally, the sensitivity of the carbon cycle to temperature variations in CESM is comparably small (Frank et al., 2010) and reveals a strong component of unforced natural variability. In a transient last millennium simulation with small temperature variations, the proper detection of a lead-lag relation between temperature and the carbon cycle is complicated by the superposition of perturbations and responses. In addition to the classic climate-carbon cycle sensitivity experiments (e.g., Arora et al., 2013) it is therefore desirable to conduct step function-like sensitivity experiments in order to isolate the response of the carbon cycle to a particular external forcing (Gerber et al., 2003).

Despite the challenges that paleoclimate modelling faces, a number of lessons regarding forcing and structural uncertainties can be learned from these experiments. In order to better understand the role of internal versus externally-forced variability – which remains particularly critical for a period of relatively weak external forcing, such as the last millennium – larger simulation ensembles as well as ensembles with decomposed forcing should become a standard procedure in paleoclimate modelling. Since these are computationally expensive simulations, this calls for an informed discussion on the optimal usage of computing resources, to which studies like the one here can contribute valuable information. At the same time, uncertainties in forcings and reconstructions need to be further reduced to be able to better validate models in the past with the goal of constraining their future response. Key targets for such constrains are the sensitivity of temperature to solar and volcanic forcing and the climate-carbon cycle sensitivity.

Appendix A

720 A1

Acknowledgements. We gratefully acknowledge Axel Timmermann, Bette Otto-Bliesner, Peter Lawrence, and Rosie Fisher for valuable discussions as well as four anonymous reviewers for very helpful comments. We are grateful to the NCAR in Boulder, USA, for providing the code of the CESM, to the World Climate Research Programme's Working Group on Coupled Modelling, which is responsible for CMIP, to the climate modeling groups for producing and making available their model output. This study is supported by the Swiss National Science Foundation (grant no. 200020 147174), the European Commission through Seventh Framework Program (FP7) projects CARBOCHANGE (grant no. 264879) and Past4Future (grant no. 243908). J.M. has benefited from the support of the French Agence Nationale de la Recherche (HAMOC : ANR 13-BLAN-06-0003). The simulations for this study were performed on a CRAY XT5 and XE6 at the Swiss National Supercomputing Centre (CSCS) in Lugano.

References

- Abram, N. J., Mulvaney, R., Vimeux, F., Phipps, S. J., Turner, J., and England, M. H.: Evolution of the Southern Annular Mode during the past millennium, *Nature Climate Change*, 4, 564–569, 2014.
- 735 Ammann, C. M., Joos, F., Schimel, D., Otto-Bliesner, B. L., and Tomas, R.: Solar influence on climate during the past millennium: Results from transient simulations with the NCAR Climate System Model, *Proc. Natl. Acad. Sci.*, 104, 3713–3718, 2007.
- Anchukaitis, K. J., Breitenmoser, P., Briffa, K. R., Buchwal, A., Buentgen, U., Cook, E. R., D'Arrigo, R. D., Esper, J., Evans, M. N., Frank, D., Grudd, H., Gunnarson, B. E., Hughes, M. K., Kirilyanov, A. V., Koerner, C., Krusic, P. J., Luckman, B., Melvin, T. M., Salzer, M. W., Shashkin, A. V., Timmreck, C., Vaganov, E. A., 740 and Wilson, R. J. S.: Tree rings and volcanic cooling, *Nature Geoscience*, 5, 836–837, doi:10.1038/ngeo1645, 2012.
- Andres, R. J., Boden, T. A., Breon, F. M., Ciais, P., Davis, S., Erickson, D., Gregg, J. S., Jacobson, A., Marland, G., Miller, J., Oda, T., Olivier, J. G. J., Raupach, M. R., Rayner, P., and Treanton, K.: A synthesis of carbon dioxide emissions from fossil-fuel combustion, *Biogeosciences*, 9, 1845–1871, 745 doi:10.5194/bg-9-1845-2012, 2012.
- Armstrong, R., Lee, C., Hedges, J., Honjo, S., and Wakeham, S.: A new, mechanistic model for organic carbon fluxes in the ocean based on the quantitative association of POC with ballast minerals, *Deep Sea Res. Part II*, 49, 219–236, 2002.
- Arora, V. K., Boer, G. J., Friedlingstein, P., Eby, M., Jones, C. D., Christian, J. R., Bonan, G., Bopp, 750 L., Brovkin, V., Cadule, P., Hajima, T., Ilyina, T., Lindsay, K., Tjiputra, J. F., and Wu, T.: Carbon-Concentration and Carbon-Climate Feedbacks in CMIP5 Earth System Models, *J. Clim.*, 26, 5289–5314, doi:10.1175/JCLI-D-12-00494.1, 2013.
- Berger, A. L.: Long-term variations of caloric insolation resulting from the Earth's orbital elements, *J. Atm. Sci.*, 35, 2362–2367, doi:10.1016/0033-5894(78)90064-9, 1978.
- 755 Bothe, O., Jungclauss, J. H., and Zanchettin, D.: Consistency of the multi-model CMIP5/PMIP3-past1000 ensemble, *Clim. Past*, 9, 2471–2487, doi:10.5194/cp-9-2471-2013, 2013.
- Bretagnon, P. and Francou, G.: Planetary theories in rectangular and spherical variables - VSOP-87 solutions, *Astronomy & Astrophysics*, 202, 309–315, 1988.
- Brovkin, V., Lorenz, S. J., Jungclauss, J., Raddatz, T., Timmreck, C., Reick, C. H., Segsneider, J., and Six, K.: 760 Sensitivity of a coupled climate-carbon cycle model to large volcanic eruptions during the last millennium, 62, 674–681, doi:10.1111/j.1600-0889.2010.00471.x, 2010.
- Chikamoto, M. O., Timmermann, A., Yoshimori, M., Lehner, F., Laurian, A., Abe-Ouchi, A., Mouchet, A., Joos, F., and Cobb, K.: Delayed tropical Pacific biological productivity response to strong volcanic forcing, *Geophys. Res. Lett.*, prep.
- 765 Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Jones, C., Le Quere, C., Myneni, R.B. Piao, S., and Thornton, P.: Carbon and Other Biogeochemical Cycles, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- Cowan, K. and Way, R. G.: Coverage bias in the HadCRUT4 temperature series and its impact on recent 770 temperature trends, *Quarterly J. Royal Meteor. Soc.*, 140, 1935–1944, doi:10.1002/qj.2297, 2014.

- Cox, P. and Jones, C.: Climate change - illuminating the modern dance of climate and CO₂, *Science*, 321, 1642–1644, doi:10.1126/science.1158907, 2008.
- Cox, P. M., Pearson, D., Booth, B. B., Friedlingstein, P., Huntingford, C., Jones, C. D., and Luke, C. M.: Sensitivity of tropical carbon to climate change constrained by carbon dioxide variability, *Nature*, 494, 341–344, doi:10.1038/nature11882, 2013.
- 775 Crowley, T. J.: Causes of climate change over the past 1000 years, *Science*, 289, 270–277, 2000.
- Crowley, T. J., Zielinski, G., Vinther, B., Udisti, R., Kreutz, K., Cole-Dai, J., and Castellano, E.: Volcanism and the Little Ice Age, *PAGES newsletter*, 16, 22–23, 2008.
- Danabasoglu, G., Bates, S. C., Briegleb, B. P., Jayne, S. R., Jochum, M., Large, W. G., Peacock, S., and Yeager, S. G.: The CCSM4 ocean component, *J. Clim.*, 25, 1361–1389, doi:10.1175/JCLI-D-11-00091.1, 2012.
- 780 Deser, C., Knutti, R., Solomon, S., and Phillips, A. S.: Communication of the role of natural variability in future North American climate, *Nature Climate Change*, 2, 775–779, doi:10.1038/NCLIMATE1562, 2012.
- Doney, S. C., Lindsay, K., Fung, I., and John, J.: Natural variability in a stable, 1000-yr global coupled climate-carbon cycle simulation, *J. Clim.*, 19, 3033–3054, doi:10.1175/JCLI3783.1, 2006.
- 785 Esper, J., Frank, D. C., Timonen, M., Zorita, E., Wilson, R. J. S., Luterbacher, J., Holzkaemper, S., Fischer, N., Wagner, S., Nievergelt, D., Verstege, A., and Buentgen, U.: Orbital forcing of tree-ring data, *Nature Climate Change*, 2, 862–866, doi:10.1038/NCLIMATE1589, 2012.
- Fernandez-Donado, L., Gonzalez-Rouco, J. F., Raible, C. C., Ammann, C. M., Barriopedro, D., Garcia-Bustamante, E., Jungclaus, J. H., Lorenz, S. J., Luterbacher, J., Phipps, S. J., Servonnat, J., Swingedouw, D., Tett, S. F. B., Wagner, S., Yiou, P., and Zorita, E.: Large-scale temperature response to external forcing in simulations and reconstructions of the last millennium, *Clim. Past*, 9, 393–421, doi:10.5194/cp-9-393-2013, 2013.
- 790 Frank, D. C., Esper, J., Raible, C. C., Buentgen, U., Trouet, V., Stocker, B., and Joos, F.: Ensemble reconstruction constraints on the global carbon cycle sensitivity to climate, *Nature*, 463, 527–532, doi:10.1038/nature08769, 2010.
- 795 Franke, J., Frank, D., Raible, C. C., Esper, J., and Broennimann, S.: Spectral biases in tree-ring climate proxies, *Nature Climate Change*, 3, 360–364, doi:10.1038/NCLIMATE1816, 2013.
- Friedlingstein, P., Meinshausen, M., Arora, V. K., Jones, C. D., Anav, A., Liddicoat, S. K., and Knutti, R.: Uncertainties in CMIP5 climate projections due to carbon cycle feedbacks, *J. Clim.*, 27, 511–526, doi:10.1175/JCLI-D-12-00579.1, 2014.
- 800 Friedlingstein, P., Cox, P., Betts, R., Bopp, L., Von Bloh, W., Brovkin, V., Cadule, P., Doney, S., Eby, M., Fung, I., Bala, G., John, J., Jones, C., Joos, F., Kato, T., Kawamiya, M., Knorr, W., Lindsay, K., Matthews, H. D., Raddatz, T., Rayner, P., Reick, C., Roeckner, E., Schnitzler, K. G., Schnur, R., Strassmann, K., Weaver, A. J., Yoshikawa, C., and Zeng, N.: Climate-carbon cycle feedback analysis: Results from the C⁴MIP model intercomparison, *J. Clim.*, 19, 3337–3353, doi:10.1175/JCLI3800.1, 2006.
- 805 Friedrich, T., Timmermann, A., Abe-Ouchi, A., Bates, N. R., Chikamoto, M. O., Church, M. J., Dore, J. E., Gledhill, D. K., Gonzalez-Davila, M., Heinemann, M., Ilyina, T., Jungclaus, J. H., McLeod, E., Mouchet, A., and Santana-Casiano, J. M.: Detecting regional anthropogenic trends in ocean acidification against natural variability, *Nature Climate Change*, 2, 167–171, doi:10.1038/NCLIMATE1372, 2012.

- 810 Frölicher, T. L., Joos, F., and Raible, C. C.: Sensitivity of atmospheric CO₂ and climate to explosive volcanic eruptions, *Biogeosciences*, 8, 2317–2339, doi:10.5194/bg-8-2317-2011, 2011.
- Frölicher, T. L., Joos, F., Raible, C. C., and Sarmiento, J. L.: Atmospheric CO₂ response to volcanic eruptions: The role of ENSO, season, and variability, *Global Biogeochem. Cyc.*, 27, 239–251, doi:10.1002/gbc.20028, 2013.
- 815 Gao, C., Robock, A., and Ammann, C.: Volcanic forcing of climate over the past 1500 years: An improved ice core-based index for climate models, *J. Geophys. Res.*, 113, doi:10.1029/2008JD010239, 2008.
- Gent, P. R., Danabasoglu, G., Donner, L. J., Holland, M. M., Hunke, E. C., Jayne, S. R., Lawrence, D. M., Neale, R. B., Rasch, P. J., Vertenstein, M., Worley, P. H., Yang, Z.-L., and Zhang, M.: The Community Climate System Model Version 4, *J. Clim.*, 24, 4973–4991, doi:10.1175/2011JCLI4083.1, 2011.
- 820 Geoffroy, O., Saint-Martin, D., and Voldoire, A.: Land-sea warming contrast: the role of the horizontal energy transport, *Clim. Dyn.*, pp. 1–19, doi:10.1007/s00382-015-2552-y, 2015.
- Gerber, S., Joos, F., Brugger, P., Stocker, T., Mann, M., Sitch, S., and Scholze, M.: Constraining temperature variations over the last millennium by comparing simulated and observed atmospheric CO₂, *Clim. Dyn.*, 20, 281–299, doi:10.1007/s00382-002-0270-8, 2003.
- 825 Hawkins, E. and Sutton, R.: Time of emergence of climate signals, *Geophys. Res. Lett.*, 39, doi:10.1029/2011GL050087, 2012.
- Hoffman, F. M., Randerson, J. T., Arora, V. K., Bao, Q., Cadule, P., Ji, D., Jones, C. D., Kawamiya, M., Khatiwala, S., Lindsay, K., Obata, A., Shevliakova, E., Six, K. D., Tjiputra, J. F., Volodin, E. M., and Wu, T.: Causes and implications of persistent atmospheric carbon dioxide biases in Earth System Models, *J. Geophys. Res.*, 119, 141–162, doi:10.1002/2013JG002381, 2014.
- 830 Holland, M. M. and Bitz, C. M.: Polar amplification of climate change in coupled models, *Clim. Dyn.*, 21, 221–232, 2003.
- Houghton, R. A.: How well do we know the flux of CO₂ from land-use change?, *Tellus*, 62, 337–351, doi:10.1111/j.1600-0889.2010.00473.x, 2010.
- 835 Hunke, E. C. and Lipscomb, W. H.: CICE: the Los Alamos sea ice model documentation and software user’s manual version 4.1, Tech. rep., Los Alamos National Laboratory (LANL), 76 pp., 2010.
- Hurrell, J. W., Holland, M. M., Gent, P. R., Ghan, S., Kay, J. E., Kushner, P. J., Lamarque, J. F., Large, W. G., Lawrence, D., Lindsay, K., Lipscomb, W. H., Long, M. C., Mahowald, N., Marsh, D. R., Neale, R. B., Rasch, P., Vavrus, S., Vertenstein, M., Bader, D., Collins, W. D., Hack, J. J., Kiehl, J., and Marshall, S.: The Community Earth System Model: A Framework for Collaborative Research, *Bull. Am. Meteorol. Soc.*, 94, 1339–1360, doi:10.1175/BAMS-D-12-00121.1, 2013.
- 840 Hurtt, G. C., Chini, L. P., Frolking, S., Betts, R. A., Feddema, J., Fischer, G., Fisk, J. P., Hibbard, K., Houghton, R. A., Janetos, A., Jones, C. D., Kindermann, G., Kinoshita, T., Goldewijk, K. K., Riahi, K., Shevliakova, E., Smith, S., Stehfest, E., Thomson, A., Thornton, P., van Vuuren, D. P., and Wang, Y. P.: Harmonization of land-use scenarios for the period 1500–2100: 600 years of global gridded annual land-use transitions, wood harvest, and resulting secondary lands, *Clim. Change*, 109, 117–161, doi:10.1007/s10584-011-0153-2, 2011.
- IPCC: Special Report on Emissions Scenarios (SRES), Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2000.

- IPCC: Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2001.
- IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- 855 Jones, C. and Cox, P.: Modeling the volcanic signal in the atmospheric CO₂ record, *Global Biogeochem. Cyc.*, 15, 453–465, doi:10.1029/2000GB001281, 2001.
- Joos, F. and Prentice, I. C.: A paleo-perspective on changes in atmospheric CO₂ and climate, Island Press, 2004.
- Joos, F., Roth, R., Fuglestedt, J. S., Peters, G. P., Enting, I. G., von Bloh, W., Brovkin, V., Burke, E. J., Eby, M., Edwards, N. R., Friedrich, T., Froelicher, T. L., Halloran, P. R., Holden, P. B., Jones, C., Kleinen, T., 860 Mackenzie, F. T., Matsumoto, K., Meinshausen, M., Plattner, G. K., Reisinger, A., Segschneider, J., Shaffer, G., Steinacher, M., Strassmann, K., Tanaka, K., Timmermann, A., and Weaver, A. J.: Carbon dioxide and climate impulse response functions for the computation of greenhouse gas metrics: a multi-model analysis, *Atmos. Chem. Phys.*, 13, 2793–2825, doi:10.5194/acp-13-2793-2013, 2013.
- Jungclaus, J. H., Lorenz, S. J., Timmreck, C., Reick, C. H., Brovkin, V., Six, K., Segschneider, J., Giorgetta, 865 M. A., Crowley, T. J., Pongratz, J., Krivova, N. A., Vieira, L. E., Solanki, S. K., Klocke, D., Botzet, M., Esch, M., Gayler, V., Haak, H., Raddatz, T. J., Roeckner, E., Schnur, R., Widmann, H., Claussen, M., Stevens, B., and Marotzke, J.: Climate and carbon-cycle variability over the last millennium, *Clim. Past*, 6, 723–737, doi:10.5194/cp-6-723-2010, 2010.
- Kaplan, J. O., Krumhardt, K. M., Ellis, E. C., Ruddiman, W. F., Lemmen, C., and Goldewijk, K. K.: 870 Holocene carbon emissions as a result of anthropogenic land cover change, *The Holocene*, 21, 775–791, doi:10.1177/0959683610386983, 2011.
- Kaufman, D. S., Schneider, D. P., McKay, N. P., Ammann, C. M., Bradley, R. S., Briffa, K. R., Miller, G. H., Otto-Bliesner, B. L., Overpeck, J. T., Vinther, B. M., and 2k Project Members, A. L.: Recent warming reverses long-term Arctic cooling, *Science*, 325, 1236–1239, doi:10.1126/science.1173983, 2009.
- 875 Keller, K. M., Joos, F., Lehner, F., and Raible, C. C.: Detecting changes in marine responses to ENSO 850-2100 CE: Insights from the ocean carbon cycle, *Geophys. Res. Lett.*, doi:10.1002/2014GL062398, 2015.
- Keller, K. M., Joos, F., and Raible, C. C.: Time of emergence of trends in ocean biogeochemistry, *Biogeosciences*, 11, 3647–3659, doi:10.5194/bg-11-3647-2014, 2014.
- Keppel-Aleks, G., Randerson, J. T., Lindsay, K., Stephens, B. B., Keith Moore, J., Doney, S. C., Thornton, P. E., 880 Mahowald, N. M., Hoffman, F. M., Sweeney, C., Tans, P. P., Wennberg, P. O., and Wofsy, S. C.: Atmospheric carbon dioxide variability in the Community Earth System Model: Evaluation and transient dynamics during the twentieth and twenty-first centuries, *J. Clim.*, 26, 4447–4475, doi:10.1175/JCLI-D-12-00589.1, 2013.
- Key, R., Kozyr, A., Sabine, C., Lee, K., Wanninkhof, R., Bullister, J., Feely, R., Millero, F., Mordy, C., and Peng, T.: A global ocean carbon climatology: Results from Global Data Analysis Project (GLODAP), *Global Biogeochem. Cyc.*, 18, doi:10.1029/2004GB002247, 2004.
- 885 Krivova, N. A., Balmaceda, L., and Solanki, S. K.: Reconstruction of solar total irradiance since 1700 from the surface magnetic flux, *Astronomy & Astrophysics*, 467, 335–346, doi:10.1051/0004-6361:20066725, 2007.

- Lamarque, J. F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liousse, C., Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K., and van Vuuren, D. P.: Historical (1850-2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: Methodology and application, *Atmospheric Chemistry and Physics*, 10, 7017–7039, doi:10.5194/acp-10-7017-2010, 2010.
- Lamarque, J.-F., Kyle, G. P., Meinshausen, M., Riahi, K., Smith, S. J., van Vuuren, D. P., Conley, A. J., and Vitt, F.: Global and regional evolution of short-lived radiatively-active gases and aerosols in the Representative Concentration Pathways, *Clim. Change*, 109, 191–212, doi:10.1007/s10584-011-0155-0, 2011.
- Landrum, L., Otto-Bliesner, B. L., Wahl, E. R., Conley, A., Lawrence, P. J., Rosenbloom, N., and Teng, H.: Last millennium climate and its variability in CCSM4, *J. Clim.*, 26, 1085–1111, doi:10.1175/JCLI-D-11-00326.1, 2013.
- Lawrence, D. M., Oleson, K. W., Flanner, M. G., Thornton, P. E., Swenson, S. C., Lawrence, P. J., Zeng, X., Yang, Z.-L., Levis, S., Sakaguchi, K., Bonan, G. B., and Slater, A. G.: Parameterization improvements and functional and structural advances in version 4 of the Community Land Model, *J. of Adv. in Modeling Earth Systems*, 3, doi:10.1029/2011MS000045, 2011.
- Lawrence, P. J., Feddema, J. J., Bonan, G. B., Meehl, G. A., O'Neill, B. C., Oleson, K. W., Levis, S., Lawrence, D. M., Kluzek, E., Lindsay, K., and Thornton, P. E.: Simulating the biogeochemical and biogeophysical impacts of transient land cover change and wood harvest in the Community Climate System Model (CCSM4) from 1850 to 2100, *J. Clim.*, 25, 3071–3095, doi:10.1175/JCLI-D-11-00256.1, 2012.
- Le Quéré, C., Andres, R. J., Boden, T., Conway, T., Houghton, R., House, J. I., Marland, G., Peters, G. P., van der Werf, G., Ahlström, A., et al.: The global carbon budget 1959–2011, *Earth System Science Data*, 5, 165–185, doi:10.5194/essd-5-165-2013, 2013.
- Lean, J., Rottman, G., Harder, J., and Kopp, G.: *SORCE* contributions to new understanding of global change and solar variability, *Solar Physics*, 230, 27–53, doi:10.1007/s11207-005-1527-2, 2005.
- Lehner, F., Raible, C. C., and Stocker, T. F.: Testing the robustness of a precipitation proxy-based North Atlantic Oscillation reconstruction, *Quat. Sci. Rev.*, 45, 85–94, doi:10.1016/j.quascirev.2012.04.025, 2012a.
- Lehner, F., Raible, C. C., Stocker, T. F., and Hofer, D.: The freshwater balance of polar regions in transient simulations from 1500 to 2100 AD using a comprehensive coupled climate model, *Clim. Dyn.*, 39, 347–363, doi:10.1007/s00382-011-1199-6, 2012b.
- Lehner, F., Born, A., Raible, C. C., and Stocker, T. F.: Amplified inception of European Little Ice Age by sea ice-ocean-atmosphere feedbacks, *J. Clim.*, 26, 7586–7602, doi:10.1175/JCLI-D-12-00690.1, 2013.
- Lindsay, K., Bonan, G. B., Doney, S. C., Hoffman, F. M., Lawrence, D. M., Long, M. C., Mahowald, N. M., Moore, J. K., Randerson, J. T., and Thornton, P. E.: Preindustrial-Control and Twentieth-Century Carbon Cycle Experiments with the Earth System Model CESM1(BGC), *J. Clim.*, 27, 8981–9005, doi:10.1175/JCLI-D-12-00565.1, 2014.
- Long, M. C., Lindsay, K., Peacock, S., Moore, J. K., and Doney, S. C.: Twentieth-century oceanic carbon uptake and storage in CESM1(BGC), *J. Clim.*, 26, 6775–6800, doi:10.1175/JCLI-D-12-00184.1, 2013.
- Maher, N., Sen Gupta, A., and England, M. H.: Drivers of decadal hiatus periods in the 20th and 21st centuries, *Geophys. Res. Lett.*, 41, 5978–5986, doi:10.1002/2014GL060527, 2014.

- Mann, E. M., Zhang, Z., Rutherford, S., Bradley, R. S., Hughes, M. K., Shindell, D., Ammann, C., Faluvegi, G., and Ni, F.: Global signatures and dynamical origins of the Little Ice Age and Medieval Climate Anomaly, *Science*, 326, 1256–1260, doi:10.1126/science.1177303, 2009.
- 930 Mann, M. E., Fuentes, J. D., and Rutherford, S.: Underestimation of volcanic cooling in tree-ring-based reconstructions of hemispheric temperatures, *Nature Geoscience*, 5, 202–205, doi:10.1038/NNGEO1394, 2012.
- Marsland, S., Haak, H., Jungclaus, J., Latif, M., and Roske, F.: The Max-Planck-Institute global ocean/sea ice model with orthogonal curvilinear coordinates, *Ocean Modelling*, 5, 91–127, doi:10.1016/S1463-5003(02)00015-X, 2003.
- 935 Masson-Delmotte, V., Schulz, M., Abe-Ouchi, A., Beer, J., Ganopolski, A., González Rouco, J., Jansen, E., Lambeck, K., Luterbacher, J., Naish, T., Osborn, T., Otto-Bliesner, B., Quinn, T., Ramesh, R., Rojas, M., Shao, X., and Timmermann, A.: *Information from Paleoclimate Archives*, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- Matsumoto, K., Sarmiento, J., Key, R., Aumont, O., Bullister, J., Caldeira, K., Campin, J., Doney, S., Drange, H., Dutay, J., Follows, M., Gao, Y., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., Lindsay, K., Maier-Reimer, E., Marshall, J., Matear, R., Monfray, P., Mouchet, A., Najjar, R., Plattner, G., Schlitzer, R., Slater, R., Swathi, P., Totterdell, I., Weirig, M., Yamanaka, Y., Yool, A., and Orr, J.: Evaluation of ocean carbon cycle models with data-based metrics, *Geophys. Res. Lett.*, 31, doi:10.1029/2003GL018970, 2004.
- 940 Meehl, G. A., Washington, W. M., Arblaster, J. M., Hu, A., Teng, H., Tebaldi, C., Sanderson, B. N., Lamarque, J.-F., Conley, A., Strand, W. G., and White, III, J. B.: Climate system response to external forcings and climate change projections in CCSM4, *J. Clim.*, 25, 3661–3683, doi:10.1175/JCLI-D-11-00240.1, 2012.
- Moffa-Sanchez, P., Born, A., Hall, I. R., Thornalley, D. J. R., and Barker, S.: Solar forcing of North Atlantic surface temperature and salinity over the past millennium, *Nature Geoscience*, 7, 275–278, doi:10.1038/ngeo2094, 2014.
- 950 Moore, J., Doney, S., and Lindsay, K.: Upper ocean ecosystem dynamics and iron cycling in a global three-dimensional model, *Global Biogeochem. Cyc.*, 18, doi:10.1029/2004GB002220, 2004.
- Moore, J. K. and Braucher, O.: Sedimentary and mineral dust sources of dissolved iron to the world ocean, *Biogeosciences*, 5, 631–656, 2008.
- Moss, R. H., Edmonds, J. A., Hibbard, K. A., Manning, M. R., Rose, S. K., van Vuuren, D. P., Carter, T. R., 955 Emori, S., Kainuma, M., Kram, T., Meehl, G. A., Mitchell, J. F. B., Nakicenovic, N., Riahi, K., Smith, S. J., Stouffer, R. J., Thomson, A. M., Weyant, J. P., and Wilbanks, T. J.: The next generation of scenarios for climate change research and assessment, *Nature*, 463, 747–756, doi:10.1038/nature08823, 2010.
- Neale, R. B., Richter, J. H., Conley, A. J., Park, S., Lauritzen, P. H., Gettelman, A., Williamson, D. L., Rasch, P. J., Vavrus, S. J., Taylor, M. A., Collins, W. D., Zhang, M., and Lin, S.-J.: Description of the NCAR 960 Community Atmosphere Model (CAM 4.0), Tech. rep., National Center for Atmospheric Research (NCAR), 212 pp., 2010.
- Neukom, R., Gergis, J., Karoly, D. J., Wanner, H., Curran, M., Elbert, J., Gonzalez-Rouco, F., Linsley, B. K., Moy, A. D., Mundo, I., Raible, C. C., Steig, E. J., van Ommen, T., Vance, T., Villalba, R., Zinke, J., and Frank, D.: Inter-hemispheric temperature variability over the past millennium, *Nature Climate Change*, 4, 965 362–367, doi:10.1038/NCLIMATE2174, 2014.

- Otterå, O. H., Bentsen, M., Drange, H., and Suo, L.: External forcing as a metronome for Atlantic multidecadal variability, *Nature Geoscience*, 3, 688–694, doi:10.1038/NGEO955, 2010.
- PAGES 2k network: Continental-scale temperature variability during the past two millennia, *Nature Geoscience*, 6, 339–346, 2013.
- 970 Plattner, G. K., Knutti, R., Joos, F., Stocker, T. F., von Bloh, W., Brovkin, V., Cameron, D., Driesschaert, E., Dutkiewicz, S., Eby, M., Edwards, N. R., Fichefet, T., Hargreaves, J. C., Jones, C. D., Loutre, M. F., Matthews, H. D., Mouchet, A., Mueller, S. A., Nawrath, S., Price, A., Sokolov, A., Strassmann, K. M., and Weaver, A. J.: Long-term climate commitments projected with climate-carbon cycle models, *J. Clim.*, 21, 2721–2751, doi:10.1175/2007JCLI1905.1, 2008.
- 975 Pongratz, J., Reick, C., Raddatz, T., and Claussen, M.: A reconstruction of global agricultural areas and land cover for the last millennium, *Global Biogeochem. Cyc.*, 22, doi:10.1029/2007GB003153, 2008.
- Pongratz, J., Caldeira, K., Reick, C. H., and Claussen, M.: Coupled climate-carbon simulations indicate minor global effects of wars and epidemics on atmospheric CO₂ between ad 800 and 1850, *The Holocene*, 21, 843–851, doi:10.1177/0959683610386981, 2011.
- 980 Resplandy, L., Séférian, R., and Bopp, L.: Natural variability of CO₂ and O₂ fluxes: What can we learn from centuries-long climate models simulations?, *J. Geophys. Res.*, doi:10.1002/2014JC010463, 2015.
- Ridley, D. A., Solomon, S., Barnes, J. E., Burlakov, V. D., Deshler, T., Dolgii, S. I., Herber, A. B., Nagai, T., Neely, III, R. R., Nevzorov, A. V., Ritter, C., Sakai, T., Santer, B. D., Sato, M., Schmidt, A., Uchino, O., and Vernier, J. P.: Total volcanic stratospheric aerosol optical depths and implications for global climate change, *Geophys. Res. Lett.*, 41, 7763–7769, doi:10.1002/2014GL061541, 2014.
- 985 Rothenberg, D., Mahowald, N., Lindsay, K., Doney, S. C., Moore, J. K., and Thornton, P.: Volcano impacts on climate and biogeochemistry in a coupled carbon-climate model, *Earth Sys. Dyn.*, 3, 121–136, doi:10.5194/esd-3-121-2012, 2012.
- Scheffer, M., Brovkin, V., and Cox, P.: Positive feedback between global warming and atmospheric CO₂ concentration inferred from past climate change, *Geophys. Res. Lett.*, 33, doi:10.1029/2005GL025044, 2006.
- 990 Schmidt, G. A., Jungclaus, J. H., Ammann, C. M., Bard, E., Braconnot, P., Crowley, T. J., Delaygue, G., Joos, F., Krivova, N. A., Muscheler, R., Otto-Bliesner, B. L., Pongratz, J., Shindell, D. T., Solanki, S. K., Steinhilber, F., and Vieira, L. E. A.: Climate forcing reconstructions for use in PMIP simulations of the last millennium (v1.0), *Geoscientific Model Development*, 4, 33–45, doi:10.5194/gmd-4-33-2011, 2011.
- 995 Schmidt, G. A., Jungclaus, J. H., Ammann, C. M., Bard, E., Braconnot, P., Crowley, T. J., Delaygue, G., Joos, F., Krivova, N. A., Muscheler, R., Otto-Bliesner, B. L., Pongratz, J., Shindell, D. T., Solanki, S. K., Steinhilber, F., and Vieira, L. E. A.: Climate forcing reconstructions for use in PMIP simulations of the Last Millennium (v1.1), *Geoscientific Model Development*, 5, 185–191, doi:10.5194/gmd-5-185-2012, 2012.
- Schmidt, G. A., Annan, J. D., Bartlein, P. J., Cook, B. I., Guilyardi, E., Hargreaves, J. C., Harrison, S. P., Kageyama, M., LeGrande, A. N., Konecky, B., Lovejoy, S., Mann, M. E., Masson-Delmotte, V., Risi, C., Thompson, D., Timmermann, A., Tremblay, L. B., and Yiou, P.: Using palaeo-climate comparisons to constrain future projections in CMIP5, *Clim. Past*, 10, 221–250, doi:10.5194/cp-10-221-2014, 2014.
- 1000 Schneider, D. P., Ammann, C. M., Otto-Bliesner, B. L., and Kaufman, D. S.: Climate response to large, high-latitude and low-latitude volcanic eruptions in the Community Climate System Model, *J. Geophys. Res.*, 114, doi:10.1029/2008JD011222, 2009.
- 1005

- Schurer, A. P., Tett, S. F., and Hegerl, G. C.: Small influence of solar variability on climate over the past millennium, *Nature Geoscience*, 7, 104–108, 2014.
- Schurer, A. P., Hegerl, G. C., Mann, M. E., Tett, S. F. B., and Phipps, S. J.: Separating forced from chaotic climate variability over the past millennium, *J. Clim.*, 26, 6954–6973, doi:10.1175/JCLI-D-12-00826.1, 2013.
- 1010 Schwinger, J., Tjiputra, J. F., Heinze, C., Bopp, L., Christian, J. R., Gehlen, M., Ilyina, T., Jones, C. D., Salas-Melia, D., Segschneider, J., Seferian, R., and Totterdell, I.: Nonlinearity of Ocean Carbon Cycle Feedbacks in CMIP5 Earth System Models, *jc*, 27, 3869–3888, doi:10.1175/JCLI-D-13-00452.1, 2014.
- Servonnat, J., Yiou, P., Khodri, M., Swingedouw, D., and Denvil, S.: Influence of solar variability, CO₂ and orbital forcing between 1000 and 1850 AD in the IPSLCM4 model, *Clim. Past*, 6, 445–460, doi:10.5194/cp-6-445-2010, 2010.
- 1015 Shapiro, A. I., Schmutz, W., Rozanov, E., Schoell, M., Haberreiter, M., Shapiro, A. V., and Nyeki, S.: A new approach to the long-term reconstruction of the solar irradiance leads to large historical solar forcing, *Astronomy & Astrophysics*, 529, doi:10.1051/0004-6361/201016173, 2011.
- Shevliakova, E., Pacala, S. W., Malyshev, S., Hurr, G. C., Milly, P. C. D., Caspersen, J. P., Sentman, L. T., Fisk, J. P., Wirth, C., and Crevoisier, C.: Carbon cycling under 300 years of land use change: Importance of the secondary vegetation sink, *Global Biogeochem. Cyc.*, 23, doi:10.1029/2007GB003176, 2009.
- 1020 Sicre, M. A., Khodri, M., Mignot, J., Eiriksson, J., Knudsen, K. L., Ezat, U., Closset, I., Nogues, P., and Masse, G.: Sea surface temperature and sea ice variability in the subpolar North Atlantic from explosive volcanism of the late thirteenth century, *Geophys. Res. Lett.*, 40, 5526–5530, doi:10.1002/2013GL057282, 2013.
- 1025 Smith, R., Jones, P., Briegleb, B., Bryan, F., Danabasoglu, G., Dennis, J., Dukowicz, J., Eden, C., Fox-Kemper, B., Gent, P., Hecht, M., Jayne, S., Large, M. J. W., Lindsay, K., Maltrud, M., Norton, N., Peacock, S., Vertenstein, M., and Yeager, S.: The Parallel Ocean Program (POP) Reference Manual, Tech. rep., Los Alamos National Laboratory (LANL), 140 pp., 2010.
- Steinacher, M., Joos, F., and Stocker, T. F.: Allowable carbon emissions lowered by multiple climate targets, *Nature*, 499, 197–203, doi:10.1038/nature12269, 2013.
- 1030 Stocker, B. D., Strassmann, K., and Joos, F.: Sensitivity of Holocene atmospheric CO₂ and the modern carbon budget to early human land use: Analyses with a process-based model, *Biogeosciences*, 8, 69–88, doi:10.5194/bg-8-69-2011, 2011.
- Stocker, B. D., Feissli, F., Strassmann, K. M., Spahni, R., and Joos, F.: Past and future carbon fluxes from land use change, shifting cultivation and wood harvest, *Tellus*, 66, doi:10.3402/tellusb.v66.23188, 2014.
- 1035 Swingedouw, D., Mignot, J., Labetoulle, S., Guilyardi, E., and Madec, G.: Initialisation and predictability of the AMOC over the last 50 years in a climate model, *Clim. Dyn.*, 40, 2381–2399, doi:10.1007/s00382-012-1516-8, 2013.
- Swingedouw, D., Ortega, P., Mignot, J., Guilyardi, E., Masson-Delmotte, V., Butler, P. G., Khodri, M., and Seferian, R.: Bidecadal North Atlantic ocean circulation variability controlled by timing of volcanic eruptions, *Nature Communications*, 6, doi:10.1038/ncomms7545, 2015.
- 1040 Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An overview of CMIP5 and the experiment design, *Bull. Am. Meteorol. Soc.*, 93, 485–498, doi:10.1175/BAMS-D-11-00094.1, 2012.

- Thompson, D. W. J., Wallace, J. M., Jones, P. D., and Kennedy, J. J.: Identifying signatures of natural climate variability in time series of global-mean surface temperature: Methodology and insights, *J. Clim.*, 22, 6120–6141, doi:10.1175/2009JCLI3089.1, 2009.
- Timmreck, C., Graf, H.-F., Lorenz, S. J., Niemeier, U., Zanchettin, D., Matei, D., Jungclaus, J. H., and Crowley, T. J.: Aerosol size confines climate response to volcanic super-eruptions, *Geophys. Res. Lett.*, 37, doi:10.1029/2010GL045464, 2010.
- Tingley, M. P., Stine, A. R., and Huybers, P.: Temperature reconstructions from tree-ring densities overestimate volcanic cooling, *Geophys. Res. Lett.*, 41, 7838–7845, doi:10.1002/2014GL061268, 2014.
- Tjiputra, J. F. and Otter, O. H.: Role of volcanic forcing on future global carbon cycle, *Earth Sys. Dyn.*, 2, 53–67, doi:10.5194/esd-2-53-2011, 2011.
- Trenberth, K. E. and Dai, A.: Effects of Mount Pinatubo volcanic eruption on the hydrological cycle as an analog of geoengineering, *Geophys. Res. Lett.*, 34, doi:10.1029/2007GL030524, 2007.
- Tschumi, J. and Stauffer, B.: Reconstructing past atmospheric CO₂ concentration based on ice-core analyses: open questions due to in situ production of CO₂ in the ice, *J. Glaciology*, 46, 45–53, doi:10.3189/172756500781833359, 2000.
- Vieira, L. E. A. and Solanki, S. K.: Evolution of the solar magnetic flux on time scales of years to millenia, *Astronomy & Astrophysics*, 509, doi:10.1051/0004-6361/200913276, 2010.
- Wang, J., Zeng, N., Liu, Y., and Bao, Q.: To what extent can interannual CO₂ variability constrain carbon cycle sensitivity to climate change in CMIP5 Earth System Models?, *Geophys. Res. Lett.*, 41, 3535–3544, doi:10.1002/2014GL060004, 2014.
- Wanner, H., Beer, J., Buetikofer, J., Crowley, T. J., Cubasch, U., Flueckiger, J., Goosse, H., Grosjean, M., Joos, F., Kaplan, J. O., Kuettel, M., Mueller, S. A., Prentice, I. C., Solomina, O., Stocker, T. F., Tarasov, P., Wagner, M., and Widmann, M.: Mid- to Late Holocene climate change: An overview, *Quat. Sci. Rev.*, 27, 1791–1828, doi:10.1016/j.quascirev.2008.06.013, 2008.
- Wenzel, S., Cox, P. M., Eyring, V., and Friedlingstein, P.: Emergent constraints on climate-carbon cycle feedbacks in the CMIP5 Earth System Models, *J. Geophys. Res.*, 119, 794–807, doi:10.1002/2013JG002591, 2014.
- Woodwell, G., Mackenzie, F., Houghton, R., Apps, M., Gorham, E., and Davidson, E.: Biotic feedbacks in the warming of the Earth, *Clim. Change*, 40, 495–518, doi:10.1023/A:1005345429236, 1998.
- Wunsch, C. and Heimbach, P.: Practical global oceanic state estimation, *Physica D – nonlinear phenomena*, 230, 197–208, doi:10.1016/j.physd.2006.09.040, Data Assimilation for Geophysical Systems Program, Stat & Appl Math Sci Inst, Research Triangle Pk, NC, 2005, 2007.
- Wunsch, C. and Heimbach, P.: How long to oceanic tracer and proxy equilibrium?, *Quat. Sci. Rev.*, 27, 637–651, doi:10.1016/j.quascirev.2008.01.006, 2008.
- Zanchettin, D., Timmreck, C., Graf, H.-F., Rubino, A., Lorenz, S., Lohmann, K., Krueger, K., and Jungclaus, J. H.: Bi-decadal variability excited in the coupled ocean-atmosphere system by strong tropical volcanic eruptions, *Clim. Dyn.*, 39, 419–444, doi:10.1007/s00382-011-1167-1, 2012.




Figure 1. Forcings used in the last millennium simulation with CESM. (a) TSI in comparison with the different TSI reconstructions proposed by PMIP3. (b) Volcanic forcing as total volcanic aerosol mass. (c) Radiative forcing (RF, calculated according to IPCC, 2001) from the greenhouse gases CO₂, CH₄, and N₂O. (d) Major changes in land cover (as fraction of global land area). See text for details.




Figure 2. (a) Mean June-August (JJA) Arctic (>60° N land) solar insolation in CCSM4 with time-varying orbital parameters and CESM with fixed orbital parameters. (b) Arctic JJA temperature difference between CCSM4 and CESM. The least-squares linear trend of this temperature difference is given in red. (c) Arctic JJA temperature anomalies (from their 850-1850 AD mean) versus solar insolation as 100-year and 200-year averages (10 and 5 circles, respectively) from CCSM4 and CESM (red and blue, respectively). The least-squares linear trend for each cloud of 100-year and 200-year averages is given in the respective color. The shading envelops the range of temperature versus solar insolation for each cloud of means.

Table 1. List of simulations conducted for this study. See text for details regarding the forcing. TSI=total solar irradiance, GHGs = greenhouse gases, E_{CO_2} = anthropogenic CO_2 emissions from fossil fuel burning and cement production. LULUC = land use and land use change.

	Control simulation (CTRL)	Transient simulation (CESM)
Forcing	850 CE (500 years)	850-2099 CE
TSI	$1360.228 \text{ W m}^{-2}$	adjusted Vieira and Solanki (2010) and Lean et al. (2005)
Volcanic	none	Gao et al. (2008)
GHGs	CO_2 (279.3 ppm) CH_4 (674.5 ppb) N_2O (266.9 ppb)	Schmidt et al. (2011)
E_{CO_2}	none	Andres et al. (2012) and Moss et al. (2010)
Aerosol	1850 CE from Lamarque et al. (2010)	Lamarque et al. (2010, 2011)
Orbital	1990 CE after Berger (1978)	1990 CE after Berger (1978)
LULUC	850 CE from Pongratz et al. (2008)	Pongratz et al. (2008) and Hurtt et al. (2011)




Figure 3. (a) Northern Hemisphere and (b) Southern Hemisphere temperature anomalies in model simulations and reconstructions. The anomalies are with reference to 1500-1899 CE (left panels) and 1850-1899 CE (right panels). Gray shading in (a) indicates the reconstruction overlap (IPCC, 2013), in (b) the reconstruction by Neukom et al. (2014). The 5-95% range of the simulations from the third Paleoclimate Modelling Intercomparison Project (PMIP3) and the fifth Coupled Model Intercomparison Project (CMIP5; applying the RCP 8.5) are given in green and red shading, respectively. Note that MPI-ESM applies the A1B scenario (IPCC, 2000), which has a weaker forcing than RCP 8.5. Hemispheric means from observations are shown as thick black line (Cowtan and Way, 2014). All time series have been smoothed by a local regression filter which suppresses variability higher than 30 years. The Medieval Climate Anomaly (MCA) and the Little Ice Age (LIA) are indicated as defined in Mann et al. (2009). (c) Evolution of atmospheric CO₂ in CESM (black), MPI-ESM (grey; ensemble range), from ice cores (red), from measurements (orange), and from RCP8.5 used to force the radiative code in CESM (magenta). The small inset in the middle panel shows the observed annual cycle at Mauna Loa, Hawaii, and a 2° × 2° average over Hawaii from CESM, both derived from the period 1958-2012.




Figure 4. Annual mean net carbon flux from the atmosphere to (a) land and (b) ocean. Green bars given the full and 10-90% range from the preindustrial part of the simulation. Observational estimates are from Le Quéré et al. (2013)

Table 2. Selected forcing details of simulations used in comparisons with CESM. TSI=total solar irradiance, LULUC=land use and land use change.

Forcing	CESM	CCSM4	IPSL-CM5A-LR	MPI-ESM
TSI	adjusted Vieira and Solanki (2010) and Lean et al. (2005)	Vieira and Solanki (2010) and Lean et al. (2005)	Vieira and Solanki (2010) and Lean et al. (2005)	Krivova et al. (2007)
Volcanic	Gao et al. (2008)	Gao et al. (2008)	Gao et al. (2008)	Crowley et al. (2008)
Orbital	1990 CE, Berger (1978)	transient, Berger (1978)	transient, Berger (1978)	transient, Bretagnon and Francou (1990)
LULUC	Pongratz et al. (2008) and Hurtt et al. (2011)	Pongratz et al. (2008) and Hurtt et al. (2011)	non-transient	Pongratz et al. (2008)




Figure 5. 5-year filtered zonal mean anomalies of surface air temperature (SAT), relative to 850-1849 CE from (a) CESM and (b) MPI-ESM. (c) 100-year running-window correlation of zonal mean SAT from CESM and MPI-ESM. 0.75 Tukey window has been applied to the data before correlation to weaken sharp transitions. Stippling indicates significance (5% level), taking into account autocorrelation estimated from the entire time period. (d) As (c) but for the correlation of CESM with CCSM4. (e) As (d) but for global mean SAT. Small inset on top shows volcanic and solar forcing of CESM and MPI-ESM. Volcanic forcing of CESM scaled to have the same radiative forcing as MPI-ESM for Pinatubo in 1991 CE. Solar forcing relative to 1850 CE.




Figure 6. Regression of total solar irradiance (TSI) on surface air temperature (SAT) for the period 850-1850 CE in (a) CESM and (b) MPI-ESM. Time series at each gridpoint have been 5-year filtered. Only significant regression coefficients at the 5% level are shown. The small panel shows zonal means.

Table 3. Cumulative carbon emissions by different components over different time periods in CESM, in Pg C. Positive (negative) values indicate emission to (uptake from) the atmosphere.

	850-1500 CE	1501-1750 CE	1751-2011 CE	2012-2100 CE
Ocean	26.0	-4.0	-151.3	-413.0
Land	-15.0	10.3	82.5	139.4
Land (without LULUC)	-24.4	-9.3	-94.7	-436.3
Fossil Fuels	0.0	0.0	358.0	1,851.5




Figure 7. 5-year filtered zonal mean anomalies of horizontally averaged ocean temperature, relative to 850-1849 CE from (a) CESM and (b) MPI-ESM. (c) 100-year running-window correlation of zonal mean SAT from CESM and MPI-ESM. A 0.75 Tukey window has been applied to the data before correlation to weaken sharp transitions. Stippling indicates significance at the 5% level, taking into account autocorrelation estimated from the entire time period. (d) 100-year running-window correlation of the Atlantic Meridional Overturning Circulation (AMOC) in CESM and MPI-ESM.




Figure 8. 5-year filtered zonal mean anomalies of horizontally integrated dissolved inorganic carbon (DIC), relative to 850-1849 CE from (a) CESM and (b) MPI-ESM. (c) 100-year running-window correlation of zonal mean SAT from CESM and MPI-ESM. A 0.75 Tukey window has been applied to the data before correlation to weaken sharp transitions. Stippling indicates significance at the 5% level, taking into account autocorrelation estimated from the entire time period.




Figure 9. Superposed Epoch Analysis of the strongest three (top3) and following strongest seven eruptions (top10) of the period 850-1850 CE in (a-e) CESM and (f-j) MPI-ESM for (a, f) global mean surface air temperature, (b, g) global mean precipitation, (c, h) atmospheric carbon given in Pg C on the left y-axis and in ppm CO₂ on the right y-axis, (d, i) ocean carbon, and (e, j) land carbon. Time series are deseasonalized and calculated as anomalies to the mean of the preceding five years. The shading shows the 10-90% range.




Figure 10. Superposed Epoch Analysis of the strongest three (top3) and following strongest seven eruptions (top10) for tropical land ($25^{\circ} S$ to $25^{\circ} N$) in CESM during the period 850-1850 CE. Land carbon inventory changes split up in (a) vegetation, (b) dead biomass (litter and wooden debris), and (c) soil. Further, changes in (d) solar radiation, (e) net primary production (NPP), and (e) loss of carbon through fire. Time series are deseasonalized and calculated as anomalies to the mean of the preceding five years. The shading shows the 10-90% range.




Figure 11. Composites of top 10 post-volcanic eruption years as anomalies to the preceding 5 years, averaged over (left) the first 2 years starting with the year of the eruption, and (right) the following three years. (a) Surface air temperature, (b) precipitation, (c) total land carbon, (d) dissolved inorganic carbon (DIC) integrated over the top 200 meters. Shading or stippling indicates significance at the 5% level. Note, that for land carbon at an individual grid cell hardly any significant changes are detected due to the large inter-annual variability.




Figure 12. Global mean changes in response to Pinatubo. (a) Global mean surface air temperature and (b) atmospheric carbon, both deseasonalized and linearly detrended over 30 years centered on June 1991; temperature observations were corrected for El Niño-Southern Oscillation and other dynamical components (Thompson et al., 2009), CO₂ observations were corrected for El Niño-Southern Oscillation and anthropogenic emissions (Frölicher et al., 2013).




Figure 13. Temporal dependence of the climate-carbon cycle sensitivity γ in CESM. Normalized probability density functions (PDF) of γ for 200-year windows overlapping by 50 years (color-filled), for the full period 850-1500 CE (black solid), and for the CTRL (black dashed). The spread of each PDF arises from the range of low-pass filters applied (20 to 120 years).