

# Quantifying differences in land use emission estimates implied by definition discrepancies

Benjamin D. Stocker<sup>1</sup> and Fortunat Joos<sup>2,3</sup>

<sup>1</sup>Department of Life Sciences, Imperial College London, Silwood Park, Ascot, SL5 7PY, UK

<sup>2</sup>Climate and Environmental Physics, Physics Institute, University of Bern, Switzerland

<sup>3</sup>Oeschger Centre for Climate Change Research, University of Bern, Switzerland

*Correspondence to:* Benjamin D. Stocker (b.stocker@imperial.ac.uk)

**Abstract.** The quantification of CO<sub>2</sub> emissions from anthropogenic land use and land use change (eLUC) is essential to understand the drivers of the atmospheric CO<sub>2</sub> increase and to inform climate change mitigation policy. Reported values in synthesis reports are commonly derived from different approaches (observation-driven bookkeeping and process-modelling) but recent work has emphasized that inconsistencies between methods may imply substantial differences in eLUC estimates. However, a consistent quantification is lacking and no concise modelling protocol for the separation of primary and secondary components of eLUC has been established. Here, we review differences of eLUC quantification methods and apply an Earth System Model (ESM) of Intermediate Complexity to quantify them. We find that the magnitude of effects due to merely conceptual differences between ESM and offline vegetation model-based quantifications is  $\sim 20\%$  for today. Under a future business-as-usual scenario, differences tend to increase further due to slowing land conversion rates and an increasing impact of altered environmental conditions on land-atmosphere fluxes. We establish how coupled Earth System Models may be applied to separate secondary component fluxes of eLUC arising from the replacement of potential C sinks/sources and the land use feedback and show that secondary fluxes derived from offline vegetation models are conceptually and quantitatively not identical to either, nor their sum. Therefore, we argue that synthesis studies should resort to the “least common denominator” of different methods, following the bookkeeping approach where only primary land use emissions are quantified under the assumption of constant environmental boundary conditions.

## 20 1 Introduction

Anthropogenic emissions of CO<sub>2</sub> are the main driver for observed climate change (Stocker et al., 2013b) and primarily result from the combustion of fossil fuels and anthropogenic land use and land use change (LUC) (Le Quéré et al., 2015). Conceptually, fossil fuel emissions can be regarded as an external forcing acting upon the C cycle-climate system. In contrast, LUC additionally modi-

25 fies the response of terrestrial ecosystems to elevated CO<sub>2</sub> and changes in climate (Gitz and Ciais, 2003; Strassmann et al., 2008) and thereby affects the C cycle-climate feedback (Joos et al., 2001; Friedlingstein et al., 2006; Stocker et al., 2013a). This leaves room for interpretations as to how exactly land use change emissions (*eLUC*) are to be defined and where the system boundaries are to be drawn.

30 The definition of *eLUC* is relevant for the accounting of the global C budget (Ciais et al., 2013). Top-down derived land-atmosphere C fluxes that are not explained by bottom-up estimates of *eLUC* are commonly ascribed to the *residual terrestrial C sink*. Differences in the definition of *eLUC* thus directly translate into differences in estimates for the residual terrestrial C sink. This budget term is a major source of uncertainty in climate projections (Jones et al., 2013) and its quantitative

35 understanding motivates a large part of current research in biogeochemistry and terrestrial ecology.

Common to almost all approaches to quantify “CO<sub>2</sub> emissions from land use change” using global process-based models, is that *eLUC* is calculated as the difference in the global total land-to-atmosphere flux (*F*) between a realistic world where land vegetation cover and C pools are affected by prescribed, time-varying LUC maps (subscript LUC) and a hypothetical world, where no LUC is

40 occurring (subscript 0):

$$eLUC = F_{LUC} - F_0 \quad (1)$$

However, the definition or model setup, under which  $F_{LUC}$  and  $F_0$  are calculated, is relevant as it implies the inclusion of secondary fluxes. Strassmann et al. (2008) (henceforth termed SM08) lay out a framework to distinguish between different component fluxes arising from land use, including pri-

45 mary emissions from converted land, and secondary emissions arising from the interactions between climate, CO<sub>2</sub> and LUC. Pongratz et al. (2014) (henceforth termed PG14) show that numerous different definitions of *eLUC* have been used in the published literature, implying a bewildering array of different combinations of component fluxes that are counted towards *eLUC* in the different studies. SM08 and PG14 demonstrate conceptually that due to this, typical *eLUC* estimates derived from

50 observation-driven bookkeeping models, offline Dynamic Global Vegetation Models, and coupled Earth System Models give systematically different results.

Substantial, setup-related differences in *eLUC* estimates have been found in earlier studies (Strassmann et al., 2008; Arora and Boer, 2010; Gasser and Ciais, 2013), and different component fluxes have been identified and quantitatively separated within their respective modelling framework (Gitz

55 and Ciais, 2003; Strassmann et al., 2008). SM08 distinguished between primary emissions that cap-

ture the direct effects of land conversion, and secondary effects arising from the interaction of land conversion and environmental change (CO<sub>2</sub> and climate). SM08 further separated the secondary fluxes into the *land use feedback flux* and the *replaced sinks/sources flux*. We term these *eLFB* and *eRSS*, respectively, and provide definitions in Sect. 3 and quantifications in Sect. 5. Recently, Gasser and Ciais (2013) (GC13) provided quantitative estimates of historical *eLUC* following different definitions. However, their analysis is limited to offline vegetation model quantifications and thus cannot address the aforementioned discrepancies between offline and ESM methods.

Here, we apply a single model, use a simple formalistic description of *eLUC* flux components inspired by GC13 and SM08, and follow the classification of PG14 to distinguish different methods of *eLUC* quantification. We quantify these differences for the historical period and a future business-as-usual scenario (RCP8.5). In contrast to earlier studies (Strassmann et al., 2008; Arora and Boer, 2010), we designed model setups to limit differences in *eLUC* to merely conceptual ones by using climate and CO<sub>2</sub> outputs from the coupled simulations to drive offline simulations, instead of using observational data for the latter. We will demonstrate that such definition differences imply inconsistencies of estimated land use emissions on the order of 20% on the global scale and may increase to 30% under a future business-as-usual scenario. This is directly relevant for territorial C balance accountings and national greenhouse gas balances under the Kyoto Protocol and thus inherently carries a political relevance.

We elucidate the implications of the choice of definition for the residual terrestrial C sink and global C budget accountings and discuss how *eLUC* quantifications may most appropriately be defined in studies that rely on multiple methodological approaches. In such cases, we propose, following Houghton (2013), to resort to the “least common denominator”, following the bookkeeping approach (method D1 in PG14), where LUC emissions are defined without accounting for any indirect effects on terrestrial C storage caused by transient changes in CO<sub>2</sub> or climate.

## 2 Brief overview of methods D1, D3, and E2

We start by revisiting the classification of PG14 for a subset of *eLUC* quantification methods identified in their study. We focus our analysis on the discrepancy between *eLUC* derived from bookkeeping and offline vegetation models (D1 and D3 methods) and coupled ESMs (E2 method). Results the D3 method feature prominently in model intercomparison studies (McGuire et al., 2001; Sitch et al., 2008), the Global Carbon Project (Le Quéré et al., 2015) and the IPCC (Ciais et al., 2013), and are often presented along with and compared against D1-type estimates. Yet, a consistent separation of commonly identified component fluxes can only be achieved by ESMs (see below).

## 2.1 Bookkeeping method (D1)

The first global quantifications of CO<sub>2</sub> emissions from LUC were based on bookkeeping models that track the fate of C after conversion from natural to cropland or pasture vegetation or vice versa (Houghton et al., 1983). Updated bookkeeping estimates of *e*LUC (Houghton, 1999; Houghton et al., 2012) still represent the benchmark against which process-based models with prognostic vegetation C density are often compared (Le Quéré et al., 2015). Bookkeeping models use observational information of C density in natural and agricultural vegetation and in different biomes to calculate *e*LUC (Houghton et al., 1983). Environmental boundary conditions thus implicitly represent fixed conditions under which the observations are taken, i.e. climate, CO<sub>2</sub>, and N-deposition levels of recent decades. Process-based vegetation models can be run in a conceptually corresponding setup (“bookkeeping method” in SM08 and thereafter) by holding environmental boundary conditions constant. While bookkeeping models are designed to derive LUC-related C emissions from a single simulation, process-based models commonly take the difference in net land-to-atmosphere carbon flux (*F*) between a simulation with and one without LUC:

$$eLUC_{D1} = F_{LUC}^0 - F_0^0 \quad (2)$$

In general, *F* refers to a global annual flux, but equations provided here are valid also for cumulative fluxes and smaller spatial domains. Constant environmental boundary conditions (CO<sub>2</sub>, climate, nitrogen deposition etc.) in both simulations are reflected by superscript ‘0’. *F*<sub>0</sub><sup>0</sup> is the land-atmosphere flux in the reference state, which may either be forced with the land use distribution at the beginning of the transient simulation (year 1700 here, see Section 4) or zero anthropogenic land use. This choice affects secondary fluxes but, after model spinup and equilibration of C pools, *F*<sub>0</sub><sup>0</sup> is zero in either case except for net land-atmosphere CO<sub>2</sub> fluxes occurring due to unforced climate variability.

Internal, unforced climate variability may affect the quantification of *e*LUC as climate variability affects the land-atmosphere carbon flux *F*. Ideally, the model setup should be such that internal, unforced variability evolves identically in both simulations. Then the land-atmosphere fluxes from land not affected by LUC and caused by internal variability would cancel when evaluating Eq. 2. In practice, this may be difficult to achieve for some state-of-the-art Earth System Models as LUC affects heat and water fluxes and thus climate. A potential solution is to run the land module offline in both simulations or to force the land module in the simulation with LUC by using climate output from the reference simulation without LUC.

*e*LUC<sub>D1</sub> is equivalent to primary emissions (see Section 3) and capture instantaneous CO<sub>2</sub> emissions occurring during deforestation and C uptake during regrowth, as well as delayed (legacy) emissions from wood product decay and the gradual re-adjustment of soil and litter C stocks to altered input levels and turnover times. Depending on the model, *e*LUC<sub>D1</sub> may also include effects of shifting cultivation (cycle of cutting forest for agriculture, then abandoning), and wood harvest. *e*LUC<sub>D1</sub>

is determined by the spatio-temporal information of land use change, C inventories in natural and agricultural land and the response time scales of C pools after conversion.

## 125 **2.2 Climate and CO<sub>2</sub>-driven offline models (D3 method)**

Prognostically simulating vegetation C density instead of prescribing it has the advantage that secondary effects under environmental change can be simulated. The first such study using a set of process-based vegetation models with prescribed, transiently varying climate and CO<sub>2</sub> from observed historical data was presented by McGuire et al. (2001). This method is termed D3 following  
130 the classification of PG14 and is also referred to as an 'offline' setup, commonly applied to stand-alone Dynamic Global Vegetation Models (DGVM) or Terrestrial Ecosystem Models (TEM).

$$eLUC_{D3} = F_{LUC}^{FF+LUC} - F_0^{FF+LUC} \quad (3)$$

Here, the superscripts indicate that actually observed, time-varying environmental conditions (climate, CO<sub>2</sub>, N-deposition, etc.) are the result of fossil fuel emissions and other non-LUC related  
135 forcings (FF), and land use change (LUC), and are prescribed in the LUC and in the non-LUC simulation. This also corresponds to the setup used in GC13 for quantifying "emissions from land use change". Their "CCN" perturbation is analogous to what the superscript 'FF+LUC' represents.

## **2.3 Emission driven coupled Earth System Models (E2)**

For a consistent separation of total CO<sub>2</sub> emissions related to LUC, emission-driven, coupled Earth  
140 System Models (ESM) may be applied. In such a setup, climate and atmospheric CO<sub>2</sub> interactively evolve in response to anthropogenic land use change, fossil fuel emissions, and other forcings. This method is termed E2 following the classification of PG14 and is typically computed with ESM or simpler atmosphere-ocean-land climate-carbon models:

$$eLUC_{E2} = F_{LUC}^{FF+LUC} - F_0^{FF} \quad (4)$$

Here, the superscript 'FF' corresponds to the environmental conditions simulated with prescribed fossil emissions and other non-LUC related anthropogenic or natural forcing, whereas superscript 'FF+LUC' refers to a simulation where environmental conditions evolve interactively in response to LUC-related emissions, as well as the 'FF' forcing. As noted also in earlier publications (Strassmann et al., 2008; Arora and Boer, 2010; Pongratz et al., 2014), here, in contrast to the D3 method,  
150 environmental conditions in the LUC and non-LUC simulation differ. In the non-LUC case, climate and CO<sub>2</sub> are consistent with absent LUC, and hence CO<sub>2</sub> is lower in the non-LUC simulation. This implies a systematic difference in flux quantifications following the D3 and E2 methods. This difference may be expressed as flux components that are either ascribed to total  $eLUC$  or not. Below, we will identify a set of commonly defined flux components and investigate the discrepancies between  
155 methods D1, D3, and E2 conceptually (Section 3) and quantitatively (Section 5).

Unforced climate variability will evolve differently in the two ESM simulations as the applied forcing is different. The component in  $F_{\text{LUC}}^{\text{FF}+\text{LUC}}$  and  $F_0^{\text{FF}}$  arising from differences in internal variability will be attributed to  $e\text{LUC}_{\text{E2}}$  according to Eq. 4. This misattribution could be significant in particular when considering small regions and short time scales. Ensemble simulations would be required to quantify the impact of internal climate variability on  $e\text{LUC}_{\text{E2}}$ . Alternatively, averaging over a large spatial domain and temporal smoothing tends to moderate the influence of unforced variability on  $e\text{LUC}_{\text{E2}}$ .

### 3 Defining flux components

SM08, PG14, and GC13 establish a formalism to describe and discuss the different definitions of total  $e\text{LUC}$  and its component fluxes. Here, we synthesize these previous frameworks to a minimal description that allows us to identify the different flux components contained in  $e\text{LUC}$  provided by the offline DGVM setups (D3 method), coupled ESM model setups (E2 method), and the bookkeeping approach (D1 method). We then show that  $e\text{LUC}_{\text{E2}} = e\text{LUC}_0 + e\text{RSS} + e\text{LFB}$  plus nonlinearities. We propose a definition for the delineation between component fluxes that follows a separation along underlying drivers of environmental changes, and that allows a consistent identification of component fluxes in coupled model setups with and without the FF forcing. The formalism presented below sets the basis for the analysis and discussion in subsequent sections.

Following GC13, the total flux in a simulation with complete forcings (FF and LUC) can be written as the sum of fluxes occurring on undisturbed land and disturbed land.

$$F_{\text{LUC}}^{\text{FF}+\text{LUC}} = \underbrace{(A_0 - \Delta A) \Delta f_{\text{nat}}^{\text{FF}+\text{LUC}}}_{\text{undisturbed land}} + \underbrace{\Delta A (f^0 + \Delta f_{\text{agr}}^{\text{FF}+\text{LUC}})}_{\text{disturbed land}} \quad (5)$$

$F$  denotes again a carbon flux (e.g. in  $\text{GtC yr}^{-1}$ ),  $f$  a carbon flux per unit area, and  $\Delta$  a change with respect to the reference period/start of the simulation. Superscripts '0', 'LUC', and 'FF' refer to the driver of environmental conditions: no forcing, emissions from LUC, and fossil fuel plus other non-LUC forcings. Subscript 'agr' refer to converted land and subscripts 'nat' to land that has not changed its status over the course of the simulation.  $\Delta A$  is the total area that has been converted, e.g., from natural to agricultural, up to the point in time of interest.  $A_0$  is the initial (reference) area.  $\Delta f_{\text{nat}}^{\text{FF}+\text{LUC}}$  is the change of the area-specific flux occurring on unconverted land due to environmental impacts caused by the combination of FF and LUC. GC13 treat fluxes on disturbed land as a vector representing land area cohorts that have transitioned from natural to agricultural land at a given time. Here, we drop the vector notation for individual age cohorts after conversion and lump these into a scalar representing non-natural (agricultural) land of varying age ( $\Delta A (f^0 + \Delta f_{\text{agr}}^{\text{FF}+\text{LUC}})$ ).  $f^0$  are direct emissions in response to land conversion under constant environmental conditions and comprise instantaneous and legacy fluxes as identified by  $I_u$  and  $L_u$  in PG14;  $\Delta f_{\text{agr}}^{\text{FF}+\text{LUC}}$  is its

modification due to environmental change ( $\delta I$  and  $\delta L$  in PG14). Note that on long time scales, the cumulative flux of ( $\Delta f_{\text{agr}}^{\text{FF}+\text{LUC}}$ ) is independent of the magnitude of  $f^0$ .

Using  $\Delta f^{\text{FF}+\text{LUC}} = \Delta f^{\text{FF}} + \Delta f^{\text{LUC}} + \delta$  and  $F_0^{\text{FF}} = A_0 \Delta f_{\text{nat}}^{\text{FF}}$  allows us to expand and rearrange terms in Eq. 5 and to write the total C flux induced by LUC as a sum of commonly separated component flux components, primary emissions ( $e\text{LUC}_0$ ), replaced sinks/sources ( $e\text{RSS}$ ), and the land use feedback flux ( $e\text{LFB}$ ):

$$e\text{LUC} = F_{\text{LUC}}^{\text{FF}+\text{LUC}} - F_0^{\text{FF}} \quad (6)$$

$$= \Delta A (\Delta f_{\text{agr}}^{\text{FF}} - \Delta f_{\text{nat}}^{\text{FF}}) \quad (e\text{RSS}) \quad (7)$$

$$+ (A_0 - \Delta A) \Delta f_{\text{nat}}^{\text{LUC}} + \Delta A \Delta f_{\text{agr}}^{\text{LUC}} \quad (e\text{LFB}) \quad (8)$$

$$+ \Delta A f^0 \quad (e\text{LUC}_0) \quad (9)$$

$$+ (A_0 - \Delta A) \delta_{\text{nat}} + \Delta A \delta_{\text{agr}} \quad (\text{non-linear}) \quad (10)$$

Figure 1 reveals that global fluxes due to FF and due to LUC forcing alone combine in an almost perfectly linear fashion to the flux induced by the combined effect of FF and LUC up to present and discernible deviations ( $\delta$ ) emerge only in a future scenario of continuously rising  $\text{CO}_2$  and changing climate and contribute  $\sim 10\text{--}20\%$  by 2100 in RCP8.5.  $A_0 \Delta f_{\text{nat}}^{\text{FF}}$  can be interpreted as the potential land C sink ( $e\text{PS}$ ) under environmental change caused by FF.

$$e\text{PS} = A_0 \Delta f_{\text{nat}}^{\text{FF}} \quad (11)$$

Note that the reference state may also include agricultural land. But under commonly used model setups, the extent of agricultural land in the reference state is small in comparison to the area under natural vegetation. Above definition (Eqs. 6-10) of the total C flux induced by LUC corresponds to the E2 method,  $e\text{LUC}_{\text{E2}}$  (Eq. 4).  $e\text{LUC}_0$  are primary emissions and equivalent to  $e\text{LUC}_{\text{D1}}$ , as quantified using a bookkeeping approach. Analogously, component fluxes of the land-atmosphere  $\text{CO}_2$  exchange in the different model setups  $F_i^k$  can now be identified (see Table 2).

In spite of the variety of terminologies presented in the published literature, studies generally agree that total C fluxes induced by LUC can be split into primary emissions,  $e\text{LUC}_0$ , that capture the direct effects of land conversion, and secondary effects arising from the interaction of land conversion and environmental change ( $\text{CO}_2$ , climate). However, the exact delineation between secondary emissions  $e\text{LFB}$  and  $e\text{RSS}$  differs (Strassmann et al., 2008; Pongratz et al., 2009, 2014). Here, we chose a definition so that  $e\text{RSS}$  arises due to environmental changes (e.g.  $\text{CO}_2$ , climate, N-deposition, ozone, air pollution, etc.) that are not caused by LUC, whereas  $e\text{LFB}$  is due to environmental changes driven by LUC. According to Eq. 8 and for a reference state without land under use,  $e\text{RSS}$  can be interpreted as the difference in sources/sinks between land under potential natural vegetation ( $\Delta f_{\text{nat}}^{\text{FF}}$ ) and agricultural land ( $\Delta f_{\text{agr}}^{\text{FF}}$ ) and scales with the area of land converted  $\Delta A$ . The LUC-feedback flux  $e\text{LFB}$  (Eq. 9) describes the flux arising as a consequence of LUC-induced environmental changes (e.g.,  $\text{CO}_2$ , climate change).  $e\text{LFB}$  occurs on non-converted (natural) and converted (agricultural)

land, with different sink strength ( $\Delta f_{\text{nat}}^{\text{LUC}}$  and  $\Delta f_{\text{agr}}^{\text{LUC}}$ ). To sum up,  $e\text{RSS}$  arises from secondary  
 225 effects of fossil fuel emissions (and N deposition, etc.), whereas  $e\text{LFB}$  is driven only by LUC. This  
 is reflected by the fact that only superscript 'LUC' occurs in the definition of  $e\text{LFB}$ , whereas only  
 'FF' occurs in the definition of  $e\text{RSS}$ . The definitions of  $e\text{RSS}$ , and hence of  $e\text{LFB}$  differ slightly  
 between publications (Strassmann et al., 2008; Pongratz et al., 2014). SM08 defined  $e\text{LFB}$  so that  
 this flux only occurs on remaining natural land. Specifically, the term ( $\Delta A \Delta f_{\text{agr}}^{\text{LUC}}$ ) appears in  $e\text{LFB}$   
 230 here, while it is ascribed to  $e\text{RSS}$  in SM08. However, this flux component is relatively small (see  
 Fig. 1). As indicated by PG14,  $e\text{RSS}$  may also be defined as  $e\text{RSS} = \Delta A (\Delta f_{\text{agr}}^{\text{FF+LUC}} - \Delta f_{\text{nat}}^{\text{FF+LUC}})$ ,  
 implying that  $e\text{LFB} = A_0 \Delta f_{\text{nat}}^{\text{LUC}}$ . Our choice of  $e\text{RSS}$  and  $e\text{LFB}$  has the advantage that it follows  
 an intuitive separation between underlying environmental drivers (FF vs. LUC) and that  $e\text{LFB}$  can  
 identically be separated in coupled ESM-type simulations where the FF forcings are excluded. This  
 235 corresponds to the E1 definition in PG14, with  $e\text{LUC}_{\text{E1}} = F_{\text{LUC}}^{\text{LUC}} - F_0^0 = e\text{LUC}_0 + e\text{LFB}$ , and was  
 applied by Pongratz et al. (2009) and Stocker et al. (2011).

For clarity, we have dropped the temporal and spatial dimensions of fluxes and areas and have  
 reduced the formalism to a distinction only between natural and agricultural land; the latter being  
 representative for croplands, pastures, built-up area etc. This is a simplification for a formal illustra-  
 240 tion and we note that the simulations presented in Sect. 5 account for the full complexity of fluxes  
 across space, different agricultural and natural vegetation types, and time.

As pointed out in earlier publications by SM08, PG14 and Arora and Boer (2010), as well as  
 in Section 1,  $e\text{LUC}_{\text{D3}}$  and  $e\text{LUC}_{\text{E2}}$  are not identical and hence  $e\text{LUC}_{\text{D3}}$  cannot be written as the  
 sum of component fluxes identified above. In other words, while primary emissions  $e\text{LUC}_0$  can be  
 245 consistently derived from offline DGVMs by simply holding environmental conditions constant, the  
 secondary fluxes derived from such studies are neither equal to  $e\text{RSS}$ , nor  $e\text{LFB}$ , nor the sum of the  
 two. In other words,  $e\text{RSS}$  and  $e\text{LFB}$  cannot be separated as shown here using offline vegetation  
 models.

$$e\text{LUC}_{\text{D3}} - e\text{LUC}_0 \neq e\text{RSS} + e\text{LFB} \quad (12)$$

250 By expanding terms analogously to above derivation, the difference between  $e\text{LUC}$  quantifications  
 from the E2 and the D3 methods turns out to

$$e\text{LUC}_{\text{E2}} - e\text{LUC}_{\text{D3}} = A_0 (\Delta f_{\text{nat}}^{\text{LUC}} + \delta_{\text{nat}}). \quad (13)$$

Ignoring the non-linearity term  $\delta_{\text{nat}}$ , the discrepancy can thus be interpreted as a flux, triggered  
 by environmental changes caused by LUC, but occurring on land not converted since the reference  
 255 period ( $f_{\text{nat}}^{\text{LUC}}$ ). Note that this is not identical to  $e\text{LFB}$  as defined here. The same theoretical result can  
 be found when applying the formalism of PG14 and their definition of flux components in  $e\text{LUC}_{\text{E2}}$   
 and  $e\text{LUC}_{\text{D3}}$ , with the difference turning out to  $(\delta_l + \sigma_{l,f})(E_n + E_p)$ .

In the literature,  $e\text{LUC}$  estimates from bookkeeping (corresponding to D1) and offline vegetation  
 models following the D3 method are often presented alongside (Ciais et al., 2013; Le Quéré et al.,



2015). Conceptually, they are not identical and estimates thus imply systematic differences. We can analogously decompose the fluxes in each simulation (see also Table 2) and write this difference as

$$eLUC_{D3} - eLUC_{D1} = eRSS + \Delta A (\Delta f_{agr}^{LUC} - \Delta f_{nat}^{LUC}) + \Delta A (\delta_{agr} - \delta_{nat}) \quad (14)$$

Note that the term  $\Delta A (\Delta f_{agr}^{LUC} - \Delta f_{nat}^{LUC})$  is sometimes included in  $eRSS$  implying that the difference between D3 and D1 is described simply by  $eRSS$ . However, our definition of  $eRSS$  differs.

## 4 Methods

In order to quantify the individual flux components and the discrepancy between the different quantifications of  $eLUC$  outlined in previous sections, we apply the emission-driven, coupled Bern3D-LPX Earth System Model of Intermediate Complexity as described in Stocker et al. (2013a) and the offline DGVM model setup where the LPX DGVM is driven in an offline mode as described in Stocker et al. (2014). Results from the offline vegetation model were also used in global C budget accountings (Le Quéré et al., 2013; Le Quéré et al., 2014, 2015), following the D3 method for estimating  $eLUC$  therein. The model is spun up at constant boundary conditions representing year 1700 ( $CO_2$  insolation, HYDE-based (Goldewijk, 2001) land use distribution from the LUH dataset (Hurt et al., 2006), and recycled 1901-1931 CRU TS 2.1 climate (Mitchell and Jones, 2005). Model drift is absent after the spin-up. During the transient simulation (1700-2100), climate is simulated by adding an anomaly pattern, scaled by global mean temperature change relative to 1700, to the continuously recycled CRU climatology (temperature, precipitation, cloud cover). This implies that unforced variability is identical in all simulations. We focus on results after 1800 but chose an early start of the transient simulation (1700) in order to minimise effects of the initial equilibrium assumption for LUC-related fluxes. For the historical period and the future “business-as-usual” scenario (RCP8.5), we apply CMIP5 standard inputs (Taylor et al., 2012). Land use change is simulated following the Generated Transitions Method, including shifting cultivation-type agriculture and wood harvesting, as described in Stocker et al. (2014). In contrast to the previous studies by Stocker et al. (2013a) and Stocker et al. (2014), we apply the model at a coarser spatial resolution ( $2.5^\circ \times 3.75^\circ$ , instead of  $1^\circ \times 1^\circ$ ). This has negligible effects (see Section 5). LUC-related  $CO_2$  emissions are calculated as the difference in the land-atmosphere  $CO_2$  exchange flux between the simulation with and without LUC using Eq. 2 for the bookkeeping, Eq. 4 for the coupled, and Eq. 3 for the offline setup. In the coupled ESM setup, atmospheric  $CO_2$  concentrations and climate evolve interactively in response to the respective forcings. In the offline model setup following the D3 method, we directly prescribe climate fields and  $CO_2$  concentrations to the vegetation component (LPX model). In this case, climate and  $CO_2$  are taken from the output of the coupled ESM simulation, driven by FF and LUC ( $F_{LUC}^{FF+LUC}$ ) and are prescribed to both offline simulations, with and without LUC. This corresponds conceptually to the common setup chosen for D3-type simulations, but instead of prescribing  $CO_2$  and climate from observations (which is the result of FF and LUC as well), we prescribe it from

295 the coupled model output here in order to exclude differences in forcings between the coupled (E2) and offline (D3) setups, and to focus on differences in computed emissions implied by the different definitions.

The model is run in a set of simulations that allows us to disentangle flux components  $eRSS$  and  $eLFB$ . Using the description of decomposed fluxes given in Table 2 and the definition of  $eRSS$  in  
300 Eq. 7, the replaced sinks/sources flux component can be derived from simulations described in Table 1 as

$$eRSS = F_{LUC}^{FF+LUC} - F_{LUC}^{LUC} - F_0^{FF} + (A_0 - \Delta A) \delta_{nat} + \Delta A \delta_{agr} \quad (15)$$

Again, we may ignore the non-linearity terms  $\delta$ . The expression of Eq. 15 also follows intuition. It represents the flux induced by environmental conditions caused by fossil fuel emissions in a world  
305 with LUC ( $F_{LUC}^{FF+LUC} - F_{LUC}^{LUC}$ ) and a world without LUC ( $F_0^{FF} - F_0^0$ ). The last term is zero, except for unforced variability, as neither LUC nor changing environmental conditions are acting. Alternatively,  $eRSS$  can also be derived as  $(F_{LUC}^{FF+LUC} - F_0^{FF+LUC}) - (F_{LUC}^{LUC} - F_0^{LUC})$ , which is formally identical to Eq. 15, assuming linearity. Analogously, the land use feedback flux can be derived as

$$eLFB = F_{LUC}^{LUC} - F_{LUC}^0. \quad (16)$$

310 Also this can be understood intuitively.  $eLFB$  represents the total land-atmosphere flux in a world with LUC (but without fossil fuel emissions),  $F_{LUC}^{LUC}$ , minus the direct effects of LUC,  $F_{LUC}^0$ . In other words, it represents the secondary flux caused by LUC alone. Again, alternatively  $eLFB$  can be derived as  $F_{LUC}^{FF+LUC} - F_{LUC}^{FF}$ , which is identical to Eq. 16, except for non-linearities.

## 5 Results

315 Figure 2 illustrates annual emissions from LUC as quantified from the different approaches. During the historical period, the offline quantification (D3) suggests  $\sim 23\%$  higher emissions than the coupled setup (E2). Cumulative emissions amount to 164 GtC with D3 and 133 GtC with E2 (1850 – 2005 AD, see Table 3). SM08 applied observational  $CO_2$  and climate in simulations used for D3. They found slightly higher differences of D3 vs. E2 (30% higher in their D3). Arora and Boer  
320 (2010) report a difference of  $\sim 100\%$  for a case where they only used  $CO_2$  concentrations from their interactive  $F_{LUC}^{FF+LUC}$  to force their  $F_0^{FF+LUC}$  simulation. A stronger effect in this case appears plausible as the replaced sinks/sources flux due to climate and  $CO_2$  effects are generally opposing (Strassmann et al., 2008). Stocker et al. (2014) applied the same model at a  $1^\circ \times 1^\circ$  resolution following the D3 and D1 methods to quantify “total” and “primary” LUC emissions. Results at the  
325 finer resolution (165 GtC for “total GNT” in their Table 3) are virtually identical to the present estimate. The bookkeeping method yields cumulative historical fluxes of 152 and 177 GtC under preindustrial and present-day environmental conditions. Primary emissions under preindustrial and present-day background exhibit largely identical temporal trends but differ in absolute magnitude.

16% higher emissions under present-day conditions are due to generally larger C density in natural  
 330 (non-cropland and non-pasture) vegetation and soils simulated under elevated CO<sub>2</sub> (364 ppm) and  
 the warmer climate (corresponding to years 1982 – 2012 AD in the CRU TS 3.21 dataset (Mitchell  
 and Jones, 2005). Differences in constant environmental conditions thus have qualitatively the same  
 effect as uncertainty in C stocks on natural and agricultural land. I.e.,  $eLUC_{D1}$  scales linearly with  
 simulated differences in natural and agricultural land and the trends in  $eLUC_{D1}$  derived under prein-  
 335 dustrial and present-day environmental conditions are identical, but markedly different from trends  
 in  $eLUC_{D3}$  and  $eLUC_{E2}$ .

Cumulative historical emissions following the D1 method under preindustrial (present-day) con-  
 ditions are 14% (33%) higher than suggested by the E2 method. These differences are substantial  
 and are on the order of the model range as presented in intercomparison studies (Sitch et al., 2008;  
 340 Le Quéré et al., 2015) or on the order of effects of accounting for wood harvest and shifting culti-  
 vation (Stocker et al., 2014). For the future period (2006-2099 AD) following RCP8.5, cumulative  
 emissions (2004-2099) for the D3 and E2 method are on the same order (192 and 188 GtC), but  
 considerably higher than for the D1 method (133 and 153 GtC under preindustrial and present-  
 day conditions). Differences with respect to the relative increase from present-day emission levels  
 345 (average over 1995-2004) to projected levels in the last decade of the 21st century are even larger.  
 Following the D1 method, the increase is 22% (34%) when holding conditions constant at prein-  
 dustrial (present-day) levels. Due to different inclusion of secondary fluxes, the projected increase  
 following the D3 method is 67% and 121% following E2.

Figure 3 illustrates the different flux components of total emissions from LUC following the E2  
 350 method and reveals the underpinnings of the discrepant levels and trends of emissions when quan-  
 tified with different methods. During the historical period (1850 – 2005 AD),  $eRSS$  cumulatively  
 adds 6% to primary emissions, similar as in SM08 (5%), while  $eLFB$  reduces them by 17%, similar  
 as in SM08 (18%) but less than in Pongratz et al. (2009) and Stocker et al. (2011) (30-40%). At  
 present-day,  $eRSS$  and  $eLFB$  are of similar magnitude, hence total ( $eLUC_{E2}$ ) and primary emissions  
 355 ( $eLUC_0$ ) are at approximately the same level. In RCP8.5, atmospheric CO<sub>2</sub> and temperatures con-  
 tinue to grow, while land conversion rates and primary emissions are stabilised. As a result  $eLFB$  is  
 stabilised, while  $eRSS$  continues to increase and contributes ~50% to total emissions in 2100. This  
 explains the different trends in “total” (based on E2 and D3) versus primary emissions.

The difference between  $eLUC_{E2}$  and  $eLUC_{D3}$  is of approximately the same magnitude as  $eLFB$ ,  
 360 although slightly smaller, and exhibits a trend that is closely matched by  $eLFB$  until roughly 2030  
 AD (see dashed line in Fig. 3). This is expected as the difference, derived in Eq. 13, is equal to  
 $A_0 (\Delta f_{nat}^{LUC} + \delta_{nat})$ , and thus resembles the definition of  $eLFB$  (see Eq. 8).

Secondary emissions are determined by the magnitude of C sinks and sources induced by envi-  
 ronmental change, occurring differently on disturbed (agricultural) and undisturbed (natural) land.  
 365 Fig. 4 reveals that the C sink capacity on natural land under rising CO<sub>2</sub> and a changing climate (year

2100, RCP8.5) is greatest in semi-arid regions of the Tropics and Subtropics and along the boreal treeline. In contrast, agricultural land at low latitudes acts as a net C source under environmental change and a net sink at high latitudes. The difference between the sink strength on natural and agricultural land is related to the  $e$ RSS component flux and reveals that the Tropics are the most efficient potential C sinks. Interestingly, at high latitudes, agricultural vegetation is an even more efficient C sink than natural vegetation. Fig. 4 also provides information about the spatial distribution of non-linearities from the combination of the FF and LUC forcings, corresponding to the differences between the red and the black curves in Fig. 1 in year 2100. The sum of individual effects is greater than their combination in almost all vegetated areas, but most pronounced along the transition zone between forest and open woodland. Opposite effects are simulated in individual gridcells and are likely related to the threshold-behavior of the dominant vegetation type.

## 6 Discussion

To quantify the differences in  $e$ LUC quantifications by coupled ESM (E2 method), offline DGVMs (D3 method), and the bookkeeping method (D1 method), we applied a model setup where differences stemming from driving data are removed. Then, discrepancies in total  $e$ LUC arise exclusively from the applied methods (D1, D3, E2). Our results suggest that such discrepancies in global  $e$ LUC estimates are substantial for the historical period and imply strikingly different trends in  $e$ LUC for a future business-as-usual scenario. These differences stem from the implicit inclusion of secondary flux components. As we have pointed out, secondary fluxes derived from offline vegetation model setups are conceptually not identical to what is commonly referred to as the replaced sinks/sources flux or the land use feedback, nor the sum of the two.

Land use change is a substantial driver of the observed CO<sub>2</sub> increase and has contributed about 25% to total anthropogenic CO<sub>2</sub> emissions for the period 1870-2014 (Le Quéré et al., 2015). Current (2004-2013) emission levels are  $0.9 \pm 0.5$  GtC yr<sup>-1</sup> (Le Quéré et al., 2015). Reducing emissions from deforestation and forest degradation is now an important part of international climate change mitigation efforts under the United Nation Framework Convention on Climate Change. Periodically issued synthesis reports by the IPCC (Ciais et al., 2013), annually updated CO<sub>2</sub> flux quantifications by the Global Carbon Project (Le Quéré et al., 2015), as well as multi-model intercomparison projects (CMIP5, 2009; CMIP6, 2014; TRENDY) provide valuable information on LUC CO<sub>2</sub> emissions. However, values derived from different approaches are commonly presented alongside and respective uncertainty ranges partly stem from implicit methodological differences. The lack of a standard methodological protocol for LUC emission estimates and the inclusion of secondary fluxes also obscures the scientific interpretation of model results and their comparison with observational data. Below, we outline two different perspectives on what “emissions from LUC” may represent.

## 400 6.1 Carbon budget accounting

On local to regional scales, the land C budget on natural (or weakly managed) land is derived from forest inventory data (Pan et al., 2011), net ecosystem exchange estimates from eddy flux towers (Valentini et al., 2000; Friend et al., 2007), growth assessments from tree ring data, satellite data (Baccini et al., 2012; Harris et al., 2012), and atmospheric inversions of the CO<sub>2</sub> distribution using  
 405 transport models (Gatti et al., 2014). As pointed out also by Houghton (2013) and PG14, it is in general not possible to disentangle to which extent such observation-based estimates of the local net air-land C flux are driven by environmental change induced by fossil fuel combustion or by remote LUC. Fossil fuel emission estimates do not, by definition, include any such secondary effects. *e*LUC estimates including the *e*LFB component are thus conceptually inconsistent with reported values  
 410 for fossil fuel emissions. Similarly, comparing *e*LUC quantifications that include *e*RSS with up-scaled local-to-regional scale observation-based information is confounded by this virtual, because not realised, flux component.

This is relevant for continental-to-global scale C budget accountings, where CO<sub>2</sub> exchange fluxes between the major reservoirs (ocean, atmosphere, land, fossil fuel reserves) and the airborne fraction  
 415 of anthropogenic CO<sub>2</sub> emissions are quantified (Canadell et al., 2007; Le Quere et al., 2009; Knorr, 2009; Ballantyne et al., 2012; Le Quéré et al., 2015). By definition, estimates for *e*LUC directly translate into the magnitude of the implied residual terrestrial C sink (see Fig. 5) and the airborne fraction. Inclusion of secondary LUC fluxes thus determines where the system boundaries between *e*LUC and the residual terrestrial sink are drawn. The D3 method ascribes replaced sinks/sources  
 420 (*e*RSS) to *e*LUC. This implies that the residual terrestrial sink represents a flux occurring in a hypothetical state before land conversion. This may be misleading in view of the actual reduction of land C sinks due to the reduction of natural vegetation. This reduction of the residual sink due to the replacement of natural by agricultural vegetation is only captured when basing its quantification on D1-type *e*LUC estimates.

Processes determining primary emissions are directly observable (i.e. C stocks in vegetation and  
 425 soils, C loss during deforestation, fate of product pools, soil C evolution after conversion). Such information may be used to benchmark simulated *e*LUC<sub>D1</sub>. As discussed by Houghton (2013), separating environmental effects from management effects (direct effects from LUC) also serves to lower uncertainty in *e*LUC estimates as it excludes effects of CO<sub>2</sub> fertilisation and climate impacts  
 430 on C stocks – processes less well understood and notoriously challenging to simulate. These uncertainties explain the relatively large differences in quantifications of *e*LFB as indicated in Section 5. Houghton (2013) argued that this type of uncertainty should be solely ascribed to the residual budget term to reflect which terms are subject to the largest uncertainties.

Our results also demonstrated the differences in *e*LUC<sub>D1</sub> implied by prescribing preindustrial  
 435 versus present-day environmental conditions (see Fig. 2). It may be argued that prescribing present-day conditions allows best comparability with bookkeeping estimates where observational data of

C density in natural and agricultural land is used, that inherently represents conditions of the recent past. However, we note that total terrestrial C storage is 1775, 1838, and 1982 GtC in our simulations for  $F_{LUC}^{0-PI}$ ,  $F_{LUC}^{FF+LUC}$ , and  $F_{LUC}^{0-PD}$  (mean over years 2000-2004; superscript '0-PI' ['0-PD'] refers to constant preindustrial [present-day] environmental conditions). I.e., the case where C stocks are responding to transient changes in CO<sub>2</sub> and climate ( $F_{LUC}^{FF+LUC}$  – the closest analogue to what observational data represent) is farther from its equilibrium to be attained under present-day conditions than its equilibrium under preindustrial conditions. In other words, quantifying  $eLUC_{D1}$  under preindustrial conditions is a viable and pragmatic solution.

Adopting the D1 method for benchmarking, model-intercomparison studies and syntheses based on multiple methods has the critical practical advantage of being the “least common denominator” that can be followed using empirically-based bookkeeping methods, offline vegetation models, as well as Earth System Models. Quantification of  $eLUC_{D1}$  simply requires a preindustrial control simulation (no forcings, constant environmental conditions) which is already part of the CMIP6 DECK simulations (CMIP6, 2014), and one additional run with transient LUC while environmental conditions are held constant at preindustrial levels (see Sect. 4). This could be achieved by Earth System Models without computationally demanding coupled model setups involving interactive atmosphere and ocean, but using prescribed preindustrial climate and CO<sub>2</sub> and their land models in a stand-alone mode instead. Serving as an “entry card” for future model intercomparisons, this would guarantee continuity and comparability between model development cycles and periodically repeated syntheses.

## 6.2 LUC in the Earth system

LUC effects on climate and the Earth system are not fully captured by their direct (primary) CO<sub>2</sub> emissions. Vegetation cover change also affects the local surface energy and water balances (biogeophysical effects) and emissions of other greenhouse gases. Deforestation by purposely set fires is associated with emissions of a range of radiatively active compounds (e.g., CH<sub>4</sub>, CO, NO<sub>x</sub>), wetland management may have strong effects on CH<sub>4</sub> emissions, and the application of mineral fertiliser and manure on agricultural land increases soil N<sub>2</sub>O emissions and sets in motion a cascade of detrimental environmental effects (Galloway et al., 2003), many of which directly or indirectly affect climate (Erisman et al., 2011).

Apart from these direct effects where LUC can be regarded as a forcing acting upon the Earth system, LUC also modifies the land response to external forcings. E.g., the replacement of woody vegetation with crops reduces the CO<sub>2</sub>-driven fertilisation sink. Thus, LUC affects the strength of the land-climate feedback (Stocker et al., 2013a). Furthermore, primary LUC emissions induce a secondary C uptake flux as a feedback to elevated CO<sub>2</sub> concentrations caused by primary emissions. These feedback effects are captured by the LUC flux components  $eRSS$  and  $eLFB$ . Coupled Earth System Models featuring an active C cycle require a preindustrial control simulation and a fossil

C emission-driven simulation over the industrial period where transient LUC and other climate and environmental forcings are activated to quantify the sum of primary and secondary land use C emissions (method E2). Such an emission-driven, land use-enabled simulation may become part of the CMIP6 protocol. Additional simulations are required to quantify individual components separately (see Tab. 2).

The results presented here demonstrate the importance of secondary fluxes under slowing land conversion rates and continuously increasing CO<sub>2</sub>. In RCP 8.5,  $eRSS$  is set to increase to  $\pm 1$  GtC yr<sup>-1</sup> and make up around half of  $eLUC_{E2}$  by the end of the 21st century. Hence, in order to capture the overall effect of LUC on the terrestrial C cycle feedback, these must be accounted for. However, we recommend to account for the effect of secondary LUC-related fluxes in global C budget assessments as an anthropogenic modification of the terrestrial C sink. We emphasize that offline vegetation model setups are not capable of separating  $eRSS$  and  $eLFB$  as defined here.

## 7 Conclusions

Estimates of CO<sub>2</sub> emissions from land use are essential to quantify the global C budget and inform climate change mitigation policy. However, inconsistent methodologies have been applied in syntheses based on multiple models and methods. In order to guarantee comparability and continuity, we recommend that modelling studies provide estimates derived under constant, preindustrial boundary conditions (D1 method). This method can be followed by offline vegetation models and Earth System Models, and is best comparable to observation-based estimates following the bookkeeping approach. This implies that the residual terrestrial sink derived from the global C budget includes the sink flux stimulated by environmental changes in response to LUC and reflects effects of replacement of potential C sinks due to land conversion. We have suggested how coupled, emission-driven Earth System Models may be applied to separate component fluxes defined here. Such analyses are essential to capture the full impact of LUC on climate and CO<sub>2</sub>.

*Acknowledgements.* This study received support by the Swiss National Science Foundation (SNF), the iTREE project (CR-SII3 136295) of the SNF, and by the European Commission through the FP7 projects CARBOCHANGE (grant no. 264879) and EMBRACE (grant no. 282672). We thank Raphael Roth for help with coupled simulations.

## References


- Arora, V. K. and Boer, G. J.: Uncertainties in the 20th century carbon budget associated with land use change, *Glob. Change Biol.*, 16, 3327–3348, doi:10.1111/j.1365-2486.2010.02202.x, <http://dx.doi.org/10.1111/j.1365-2486.2010.02202.x>, 2010.
- 505 Baccini, A., Goetz, S. J., Walker, W. S., Laporte, N. T., Sun, M., Sulla-Menashe, D., Hackler, J., Beck, P. S. A., Dubayah, R., Friedl, M. A., Samanta, S., and Houghton, R. A.: Estimated carbon dioxide emissions from tropical deforestation improved by carbon-density maps, *Nature Clim. Change*, 2, 182–185, doi:10.1038/nclimate1354, 2012.
- Ballantyne, A. P., Alden, C. B., Miller, J. B., Tans, P. P., and White, J. W. C.: Increase in observed net carbon  
510 dioxide uptake by land and oceans during the past 50 years, *Nature*, 488, 70–72, <http://dx.doi.org/10.1038/nature11299>, 2012.
- Canadell, J. G., Le Quere, C., Raupach, M. R., Field, C. B., Buitenhuis, E. T., Ciais, P., Conway, T. J., Gillett, N. P., Houghton, R. A., and Marland, G.: Contributions to accelerating atmospheric CO<sub>2</sub> growth from economic activity, carbon intensity, and efficiency of natural sinks, *PNAS*, 104, 18 866–18 870,  
515 doi:10.1073/pnas.0702737104, 2007.
- Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., A., C., DeFries, R., J., G., Heimann, M., Jones, C., Le Quéré, C., Myneni, R., S., P., and Thornton, P.: Carbon and Other Biogeochemical Cycles, in: *Climate Change 2013: The Physical Science Basis. Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Stocker, T., Qin, D., Plattner, G.-K.,  
520 Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P., pp. 571–658, Cambridge University Press, 2013.
- CMIP5: CMIP5 Coupled Model Intercomparison Project, Website, available from <http://cmip-pcmdi.llnl.gov/index.html> accessed 27/10/2011, 2009.
- CMIP6: CMIP Phase 6, Website, available from <http://www.wcrp-climate.org/wgcm-cmip/wgcm-cmip6> ac-  
525 cessed 19/02/2015, 2014.
- Erisman, J. W., Galloway, J., Seitzinger, S., Bleeker, A., and Butterbach-Bahl, K.: Reactive nitrogen in the environment and its effect on climate change, *CURRENT OPINION IN ENVIRONMENTAL SUSTAINABILITY*, 3, 281–290, doi:10.1016/j.cosust.2011.08.012, 2011.
- Friedlingstein, P., Cox, P., Betts, R., Bopp, L., von Bloh, W., Brovkin, V., Cadule, P., Doney, S., Eby, M., Fung, I., Bala, G., John, J., Jones, C., Joos, F., Kato, T., Kawamiya, M., Knorr, W., Lindsay, K., Matthews, H. D.,  
530 Raddatz, T., Rayner, P., Reick, C., Roeckner, E., Schnitzler, K. G., Schnur, R., Strassmann, K., Weaver, A. J., Yoshikawa, C., and Zeng, N.: Climate-Carbon Cycle Feedback Analysis: Results from the C4MIP Model Intercomparison, *J. Clim.*, 19, 3337–3353, doi:10.1175/JCLI3800.1, 2006.
- Friend, A. D., Arneeth, A., Kian, N. Y., Lomas, M., Ogee, J., Rodenbeck, C., Running, S. W., Santaren, J.-D.,  
535 Sitch, S., Viovy, N., Woodward, I. F., and Zaehle, S.: FLUXNET and modelling the global carbon cycle, *Global Change Biology*, 13, 610–633, doi:10.1111/j.1365-2486.2006.01223.x, <http://dx.doi.org/10.1111/j.1365-2486.2006.01223.x>, 2007.
- Galloway, J., Aber, J., Erisman, J., Seitzinger, S., Howarth, R., Cowling, E., and Cosby, B.: The nitrogen cascade, *Bioscience*, 53, 341–356, doi:10.1641/0006-3568(2003)053[0341:TNC]2.0.CO;2, 2003.




- 540 Gasser, T. and Ciais, P.: A theoretical framework for the net land-to-atmosphere CO<sub>2</sub> flux and its implications in the definition of "emissions from land-use change", *Earth System Dynamics*, 4, 171–186, doi:10.5194/esd-4-171-2013, <http://www.earth-syst-dynam.net/4/171/2013/>, 2013.
- Gatti, L. V., Gloor, M., Miller, J. B., Doughty, C. E., Malhi, Y., Domingues, L. G., Basso, L. S., Martinewski, A., Correia, C. S. C., Borges, V. F., Freitas, S., Braz, R., Anderson, L. O., Rocha, H., Grace, J., Phillips, O. L., and Lloyd, J.: Drought sensitivity of Amazonian carbon balance revealed by atmospheric measurements, *Nature*, 506, 76–80, <http://dx.doi.org/10.1038/nature12957>, 2014.
- 545 Gitz, V. and Ciais, P.: Amplification effect of changes in land use and concentration of atmospheric CO<sub>2</sub>., *Comptes Rendus Geoscience*, 335, 1179–1198, 2003.
- Gitz, V. and Ciais, P.: Amplifying effects of land-use change on future atmospheric CO<sub>2</sub> levels, *Global Biogeochem. Cycles*, 17, doi:10.1029/2002GB001963, 2003.
- 550 Goldewijk, K. K.: Estimating global land use change over the past 300 years: the HYDE database, *Global Biogeochem. Cycles*, 15, 417–434, 2001.
- Harris, N. L., Brown, S., Hagen, S. C., Saatchi, S. S., Petrova, S., Salas, W., Hansen, M. C., Potapov, P. V., and Lotsch, A.: Baseline Map of Carbon Emissions from Deforestation in Tropical Regions, *Science*, 336, 1573–1576, doi:10.1126/science.1217962, 2012.
- 555 Houghton, R. A.: The annual net flux of carbon to the atmosphere from changes in land use 1850-1990, *Tellus*, 51B, 298 – 313, 1999.
- Houghton, R. A.: Keeping management effects separate from environmental effects in terrestrial carbon accounting, *Glob. Change Biol.*, 19, 2609–2612, 2013.
- 560 Houghton, R. A., Hobbie, J. E., Melillo, J. M., Moore, B., Peterson, B. J., Shaver, G. R., and Woodwell, G. M.: Changes in the carbon content of terrestrial biota and soils between 1860 and 1980 - a net release of CO<sub>2</sub> to the atmosphere, *Ecological Monographs*, 53, 235–262, 1983.
- Houghton, R. A., House, J. I., Pongratz, J., van der Werf, G. R., DeFries, R. S., Hansen, M. C., Le Quéré, C., and Ramankutty, N.: Carbon emissions from land use and land-cover change, *Biogeosciences*, 9, 5125–5142, doi:10.5194/bg-9-5125-2012, <http://www.biogeosciences.net/9/5125/2012/>, 2012.
- 565 Hurtt, G. C., Frolking, S., Fearon, M. G., Moore, B., Shevliakova, E., Malyshev, S., Pacala, S. W., and Houghton, R. A.: The underpinnings of land-use history: three centuries of global gridded land-use transitions, wood-harvest activity, and resulting secondary lands, *Global Change Biol.*, 12, 1208–1229, doi:10.1111/j.1365-2486.2006.01150.x, 2006.
- 570 Jones, C., Robertson, E., Arora, V., Friedlingstein, P., Shevliakova, E., Bopp, L., Brovkin, V., Hajima, T., Kato, E., Kawamiya, M., Liddicoat, S., Lindsay, K., Reick, C. H., Roelandt, C., Segschneider, J., and Tjiputra, J.: Twenty-First-Century Compatible CO<sub>2</sub> Emissions and Airborne Fraction Simulated by CMIP5 Earth System Models under Four Representative Concentration Pathways, *JOURNAL OF CLIMATE*, 26, 4398–4413, doi:10.1175/JCLI-D-12-00554.1, 2013.
- 575 Joos, F., Prentice, I. C., Sitch, S., Meyer, R., Hooss, G., Plattner, G.-K., Gerber, S., and Hasselmann, K.: Global warming feedbacks on terrestrial carbon uptake under the Intergovernmental Panel on Climate Change (IPCC) emission scenarios, *Global Biogeochem. Cycles*, 15, 891–907, 2001.
- Knorr, W.: Is the airborne fraction of anthropogenic CO<sub>2</sub> emissions increasing?, *Geophysical Research Letters*, 36, n/a–n/a, doi:10.1029/2009GL040613, <http://dx.doi.org/10.1029/2009GL040613>, 2009.

- 580 Lamarque, J.-F., Kyle, G. P., Meinshausen, M., Riahi, K., Smith, S. J., van Vuuren, D. P., Conley, A. J., and Vitt, F.: Global and regional evolution of short-lived radiatively-active gases and aerosols in the Representative Concentration Pathways, *Clim. Change*, 109, 191–212, doi:10.1007/s10584-011-0155-0, 2011.
- Le Quere, C., Raupach, M. R., Canadell, J. G., and Marland et al., G.: Trends in the sources and sinks of carbon dioxide, *Nature Geosci*, 2, 831–836, <http://dx.doi.org/10.1038/ngeo689>, 2009.
- 585 Le Quéré, C., Andres, R. J., Boden, T., Conway, T., Houghton, R. A., House, J. I., Marland, G., Peters, G. P., van der Werf, G. R., Ahlström, A., Andrew, R. M., Bopp, L., Canadell, J. G., Ciais, P., Doney, S. C., Enright, C., Friedlingstein, P., Huntingford, C., Jain, A. K., Jourdain, C., Kato, E., Keeling, R. F., Klein Goldewijk, K., Levis, S., Levy, P., Lomas, M., Poulter, B., Raupach, M. R., Schwinger, J., Sitch, S., Stocker, B. D., Viovy, N., Zaehle, S., and Zeng, N.: The global carbon budget 1959–2011, *Earth System Science Data*, 5,
- 590 165–185, doi:10.5194/essd-5-165-2013, <http://www.earth-syst-sci-data.net/5/165/2013/>, 2013.
- Le Quéré, C., Peters, G. P., Andres, R. J., Andrew, R. M., Boden, T. A., Ciais, P., Friedlingstein, P., Houghton, R. A., Marland, G., Moriarty, R., Sitch, S., Tans, P., Arneeth, A., Arvanitis, A., Bakker, D. C. E., Bopp, L., Canadell, J. G., Chini, L. P., Doney, S. C., Harper, A., Harris, I., House, J. I., Jain, A. K., Jones, S. D., Kato, E., Keeling, R. F., Klein Goldewijk, K., Körtzinger, A., Koven, C., Lefèvre, N., Maignan, F., Omar, A., Ono,
- 595 T., Park, G.-H., Pfeil, B., Poulter, B., Raupach, M. R., Regnier, P., Rödenbeck, C., Saito, S., Schwinger, J., Segschneider, J., Stocker, B. D., Takahashi, T., Tilbrook, B., van Heuven, S., Viovy, N., Wanninkhof, R., Wiltshire, A., and Zaehle, S.: Global carbon budget 2013, *Earth System Science Data*, 6, 235–263, doi:10.5194/essd-6-235-2014, <http://www.earth-syst-sci-data.net/6/235/2014/>, 2014.
- Le Quéré, C., Moriarty, R., Andrew, R. M., Peters, G. P., Ciais, P., Friedlingstein, P., Jones, S. D., Sitch, S.,
- 600 Tans, P., Arneeth, A., Boden, T. A., Bopp, L., Bozec, Y., Canadell, J. G., Chini, L. P., Chevallier, F., Cosca, C. E., Harris, I., Hoppema, M., Houghton, R. A., House, J. I., Jain, A. K., Johannessen, T., Kato, E., Keeling, R. F., Kitidis, V., Klein Goldewijk, K., Koven, C., Landa, C. S., Landschützer, P., Lenton, A., Lima, I. D., Marland, G., Mathis, J. T., Metzl, N., Nojiri, Y., Olsen, A., Ono, T., Peng, S., Peters, W., Pfeil, B., Poulter, B., Raupach, M. R., Regnier, P., Rödenbeck, C., Saito, S., Salisbury, J. E., Schuster, U., Schwinger, J., Séférian,
- 605 R., Segschneider, J., Steinhoff, T., Stocker, B. D., Sutton, A. J., Takahashi, T., Tilbrook, B., van der Werf, G. R., Viovy, N., Wang, Y.-P., Wanninkhof, R., Wiltshire, A., and Zeng, N.: Global carbon budget 2014, *Earth System Science Data*, 7, 47–85, doi:10.5194/essd-7-47-2015, <http://www.earth-syst-sci-data.net/7/47/2015/>, 2015.
- McGuire, A. D., Sitch, S., Clein, J. S., Dargaville, R., Esser, G., Foley, J., Heimann, M., Joos, F., Kaplan, J.,
- 610 Kicklighter, D. W., Meier, R. A., Melillo, J. M., Moore III, B., Prentice, I. C., Ramankutty, N., Reichenau, T., Schloss, A., Tian, H., Williams, L. J., and Wittenberg, U.: Carbon balance of the terrestrial biosphere in the twentieth century: Analyses of CO<sub>2</sub>, climate and land use effects with four process-based ecosystem models, *Global Biogeochem. Cycles*, 15, 183–206, 2001.
- Mitchell, T. D. and Jones, P. D.: An improved method of constructing a database of monthly climate observations and associated high-resolution grids, *Int. J. Climatol.*, 25, 693–712, doi:10.1002/joc.1181, 2005.
- 615 Pan, Y., Birdsey, R. A., Fang, J., Houghton, R., Kauppi, P. E., Kurz, W. A., Phillips, O. L., Shvidenko, A., Lewis, S. L., Canadell, J. G., Ciais, P., Jackson, R. B., Pacala, S. W., McGuire, A. D., Piao, S., Rautiainen, A., Sitch, S., and Hayes, D.: A Large and Persistent Carbon Sink in the World's Forests, *Science*, 333, 988–993, doi:10.1126/science.1201609, <http://www.sciencemag.org/content/333/6045/988.abstract>, 2011.


- 620 Pongratz, J., Reick, C. H., Raddatz, T., and Claussen, M.: Effects of anthropogenic land cover change on the carbon cycle of the last millennium, *Global Biogeochemical Cycles*, 23, GB4001+, doi:10.1029/2009GB003488, 2009.
- Pongratz, J., Reick, C. H., Houghton, R. A., and House, J. I.: Terminology as a key uncertainty in net land use and land cover change carbon flux estimates, *Earth System Dynamics*, 5, 177–195, doi:10.5194/esd-5-177-2014, <http://www.earth-syst-dynam.net/5/177/2014/>, 2014.
- 625 Sitch, S., Huntingford, C., Gedney, N., Levy, P. E., Lomas, M., Piao, S. L., Betts, R., Ciais, P., Cox, P., Friedlingstein, P., Jones, C. D., Prentice, I. C., and Woodward, F. I.: Evaluation of the terrestrial carbon cycle, future plant geography and climate-carbon cycle feedbacks using five Dynamic Global Vegetation Models (DGVMs), *Global Change Biology*, 14, 2015–2039(25), 2008.
- 630 Stocker, B. D., Roth, R., Joos, F., Spahni, R., Steinacher, M., Zaehle, S., Bouwman, L., Xu-Ri, X.-R., and Prentice, I. C.: Multiple greenhouse-gas feedbacks from the land biosphere under future climate change scenarios, *Nature Clim. Change*, 3, 666–672, 2013a.
- Stocker, B. D., Feissli, F., Strassmann, K., Spahni, R., and Joos, F.: Past and future carbon fluxes from land use change, shifting cultivation and wood harvest, *Tellus B*, 66, doi:10.3402/tellusb.v66.23188, <http://www.tellusb.net/index.php/tellusb/article/view/23188>, 2014.
- 635 Stocker, B. D., Strassmann, K., and Joos, F.: Sensitivity of Holocene atmospheric CO<sub>2</sub> and the modern carbon budget to early human land use: analyses with a process-based model, *Biogeosciences*, 8, 69–88, doi:10.5194/bg-8-69-2011, 2011.
- Stocker, T., Qin, D., Plattner, G.-K., Alexander, L., Allen, S., Bindoff, N., Bréon, F.-M., Church, J., Cubasch, U., Emori, S., Forster, P., Friedlingstein, P., Gillett, N., Gregory, J., Hartmann, D., Jansen, E., Kirtman, B., Knutti, R., Krishna Kumar, K., Lemke, P., Marotzke, J., Masson-Delmotte, V., Meehl, G., Mokhov, I., Piao, S., Ramaswamy, V., Randall, D., Rhein, M., Rojas, M., Sabine, C., Shindell, D., Talley, L., Vaughan, D., and Xie, S.-P.: Technical Summary, book section TS, pp. 33–115, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, doi:10.1017/CBO9781107415324.005, [www.climatechange2013.org](http://www.climatechange2013.org), 2013b.
- 640 Strassmann, K. M., Joos, F., and Fischer, G.: Simulating effects of land use changes on carbon fluxes: past contributions to atmospheric CO<sub>2</sub> increases and future commitments due to losses of terrestrial sink capacity, *Tellus B*, 60, 583–603, doi:10.1111/j.1600-0889.2008.00340.x, 2008.
- Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An overview of CMIP5 and the experiment design, *Bulletin of the American Meteorological Society*, 93, 485–498, doi:10.1175/BAMS-D-11-00094.1, 2012.
- 650 TRENDY: TRENDY, Website, available from <http://dgvn.ceh.ac.uk/node/21> accessed 19/02/2015.
- Valentini, R., Matteucci, G., Dolman, A., Schulze, E., Rebmann, C., Moors, E., Granier, A., Gross, P., Jensen, N., Pilegaard, K., Lindroth, A., Grelle, A., Bernhofer, C., Grunwald, T., Aubinet, M., Ceulemans, R., Kowalski, A., Vesala, T., Rannik, U., Berbigier, P., Loustau, D., Guomundsson, J., Thorgeirsson, H., Ibrom, A., Morgenstern, K., Clement, R., Moncrieff, J., Montagnani, L., Minerbi, S., and Jarvis, P.: Respiration as the main determinant of carbon balance in European forests, *Nature*, 404, 861–865, doi:10.1038/35009084, 2000.




**Figure 1.** Global cumulative net land-to-atmosphere CO<sub>2</sub> fluxes induced by environmental change caused by FF ( $F_0^{\text{FF}}$ ), LUC ( $F_0^{\text{LUC}}$ ), their combined effect ( $F_0^{\text{FF}+\text{LUC}}$ ), and the sum of individual effects ( $F_0^{\text{FF}} + F_0^{\text{LUC}}$ ). Curves represent cumulative global fluxes induced by environmental change, weighted by their time-varying area of natural vegetation (dashed lines), agricultural land (dotted lines), and their sum (solid lines). Note that this excludes all direct effects of LUC. The difference between the combined and the sum of effects corresponds to the non-linearity  $(A_0 - \Delta A(t)) \delta_{\text{nat}}(t) + \Delta A(t) \delta_{\text{agr}}(t)$ . The model setups are described in Tables 1 and 2.




**Figure 2.** (a) Annual land use change emissions as quantified following different methods. (b) Difference of different eLUC definitions relative to  $e\text{LUC}_{\text{D1}}$ , quantified under preindustrial boundary conditions. Total emissions derived from an offline, concentration-driven DGVM setup (D3 method) are given by black solid lines. Total emissions derived from a coupled, emission-driven ESM setup (E2 method) are given by black dashed lines. Primary emissions are given by colored lines under constant pre-industrial (red) and constant present-day (green) environmental conditions (climate, CO<sub>2</sub>, N deposition). Time series are calculated following Eqs. 2-4, where  $F$  is the global total land-atmosphere CO<sub>2</sub> flux in the respective simulation. Bold lines are splines of annual emissions given by thin lines. Results are from simulations following CMIP5 model inputs (historical until 2005, RCP8.5 until 2099).



**Figure 3.** Flux components of land use change emissions. Total emissions as derived from an emission-driven, coupled ESM setup (E2 method), and calculated with Eq. 4, are given by the black lines. Primary emissions under preindustrial boundary conditions are given by red lines. These correspond to curves in Figure 2. The replaced sinks/sources flux ( $eRSS$ ) and the land use change feedback flux ( $eLFB$ ) are given by magenta and blue lines, respectively. The difference between total emissions quantified by D3 method (see black solid line in Fig. 2) and E2 method is given by the black dashed line. Time series are calculated following Eqs. 2, 4, 15, and 16. Bold lines are splines of annual emissions given by thin lines. Results are from simulations following CMIP5 model inputs (historical until 2005, RCP8.5 until 2099).



**Figure 4.** Top row: Cumulative atmosphere-land C flux ( $\text{kgC m}^{-2}$ ) induced by environmental change from 1700 to 2100 on natural (a) and agricultural land (b; mean of cropland and pasture, weighted by respective area shares). The period 2005-2100 follows the RCP8.5 scenario. Climate and  $\text{CO}_2$  are prescribed from the outputs of the coupled simulation (offline simulation  $F_0^{\text{FF}+\text{LUC}}$  uses outputs from  $F_{\text{LUC}}^{\text{FF}+\text{LUC}}$ ). (c) Difference of flux occurring on natural and agricultural land  $f_{\text{nat}}^{\text{FF}+\text{LUC}} - f_{\text{agr}}^{\text{FF}+\text{LUC}}$ . (d) Spatial distribution of non-linear effects, cumulative in year 2100. Its global total over time is expressed also in Figure 1 (difference between black and red curves).



**Figure 5.** Land use change emissions ( $eLUC$ , dark blue bars) calculated from different methodologies and their implied residual terrestrial C sink (annual flux in  $GtC\ yr^{-1}$ , mean over 1996-2005). The total terrestrial C balance is constrained by atmospheric measurements and is  $-0.8\ GtC\ yr^{-1}$  (mean over 1996-2005, Le Quéré et al. (2014), left vertical line). It is independent of  $eLUC$  estimates. The residual terrestrial C sink (green arrow) is defined as the difference of  $eLUC$  and the total terrestrial C balance. Depending on the definition of  $eLUC$ , the residual C sink is affected by inclusion of secondary fluxes (light blue bars,  $eRSS$  and  $eLFB$ ) into  $eLUC$ .

**Table 1.** Model setups.  $F$  is the simulated total net flux of C from the terrestrial biosphere to the atmosphere. Subscript 0 refers to a setup where the area under use is kept constant at 1700 conditions and subscript LUC to a setup where the area under use is transiently varying following the land cover data by Hurtt et al. (2006). Superscript LUC and FF refer to environmental changes ( $CO_2$ , climate, etc.) due to LUC forcing and non-LUC related forcing (FF) or their combination (FF+LUC). Simulations with superscript '0' are forced by constant environmental (climate and  $CO_2$ ) conditions (e.g, preindustrial or modern). In coupled simulations, climate and  $CO_2$  evolve interactively as simulated by the coupled Bern3D-LPX model. The offline model mode uses either outputs from the coupled simulations or constant climate and  $CO_2$  and  $F$  is computed with the stand-alone vegetation model LPX. N-deposition ('N-dep.') is prescribed from Lamarque et al. (2011).

Setup	model mode	Climate	$CO_2$	LUC	FF	N-dep.
$F_{LUC}^{FF+LUC}$	coupled	interactive	interactive	on	on	on
$F_{LUC}^{LUC}$	coupled	interactive	interactive	on	off	const.
$F_0^{FF}$	coupled	interactive	interactive	const.	on	on
$F_0^{LUC}$	offline	from $F_{LUC}^{LUC}$	from $F_{LUC}^{LUC}$	const.	–	const.
$F_0^{FF+LUC}$	offline	from $F_{LUC}^{FF+LUC}$	from $F_{LUC}^{FF+LUC}$	const.	–	on
$F_{LUC}^0$	offline	constant	constant	on	–	const.
$F_0^0$	offline	constant	constant	const.	–	const.

**Table 2.** Flux decomposition for model setups described in Table 1.  $A_0$  is land area at the reference state,  $\Delta A$  is the area of land converted relative to the reference state.  $\Delta f_{\text{nat}}$  and  $\Delta f_{\text{agr}}$  are the fluxes on unconverted and converted land induced by environmental change. The underlying driver of environmental change is given by the superscripts.  $f^0$  is the flux due to direct impacts of land conversion, not including effects of environmental change.  $F_0^0$  is zero except for the flux arising from unforced climate variability. The component flux  $A_0 \Delta f_{\text{nat}}^{\text{LUC}}$  has not been named explicitly. Non-linearity terms are ignored in this table. Note that fluxes  $F$  generally refer to global totals for a given point in time  $t$ . Thus, for example  $F_0^{\text{FF}}(t) = \int_{x,y} A_0(x,y) \Delta f_{\text{nat}}^{\text{FF}}(x,y,t) dx dy$ . For simplicity, we have dropped the time and space dimensions.

Setup	Decomposed flux	Component fluxes
$F_{\text{LUC}}^{\text{FF}+\text{LUC}}$	$(A_0 - \Delta A) \Delta f_{\text{nat}}^{\text{FF}+\text{LUC}} + \Delta A f^0 + \Delta A \Delta f_{\text{agr}}^{\text{FF}+\text{LUC}}$	$e\text{PS} + e\text{LUC}_0 + e\text{RSS} + e\text{LFB}$
$F_{\text{LUC}}^{\text{LUC}}$	$(A_0 - \Delta A) \Delta f_{\text{nat}}^{\text{LUC}} + \Delta A f^0 + \Delta A \Delta f_{\text{agr}}^{\text{LUC}}$	$e\text{LUC}_0 + e\text{LFB}$
$F_0^{\text{FF}}$	$A_0 \Delta f_{\text{nat}}^{\text{FF}}$	$e\text{PS}$
$F_0^{\text{LUC}}$	$A_0 \Delta f_{\text{nat}}^{\text{LUC}}$	$A_0 \Delta f_{\text{nat}}^{\text{LUC}}$
$F_0^{\text{FF}+\text{LUC}}$	$A_0 \Delta f_{\text{nat}}^{\text{FF}+\text{LUC}}$	$e\text{PS} + A_0 \Delta f_{\text{nat}}^{\text{LUC}}$
$F_{\text{LUC}}^0$	$\Delta A f^0$	$e\text{LUC}_0$
$F_0^0$	0	0

**Table 3.** Cumulative emissions (GtC) over historical and future period for different methods ( $e\text{LUC}_{\text{D1}}$ ,  $e\text{LUC}_{\text{D3}}$ ,  $e\text{LUC}_{\text{E2}}$ ) and component fluxes ( $e\text{RSS}$ ,  $e\text{LFB}$ ).  $e\text{LUC}_{\text{D1-PI}}$  and  $e\text{LUC}_{\text{D1-PD}}$  refer are quantified under preindustrial (PI) and present-day (PD) environmental conditions.

	1850-2004	2005-2099
$e\text{LUC}_{\text{D1-PI}}$	152	133
$e\text{LUC}_{\text{D1-PD}}$	177	153
$e\text{LUC}_{\text{D3}}$	164	192
$e\text{LUC}_{\text{E2}}$	133	188
$e\text{RSS}$	9	71
$e\text{LFB}$	-26	-17