We thank the reviewers for their careful reading of the manuscript. We have revised the manuscript to take into account the reviewers' comments as described below in red.

Reviewer #1

General comments

The authors must be congratulated for improving the general state of the paper. It reads easier, some of the confusing results have been removed and the new analyses are interesting. However, this reviewer still feels the manuscript lacks organization and clear objectives. The current version still feels like a collection of results, without a clear storyline and defined objectives. The authors justification that future projections in LAI can be constraint based on current model performance is incomplete (See Section 5 comments).

We thank the reviewer for acknowledging our work to accommodate the reviewer's previous comments and revise the manuscript to accommodate the reviewer's new comments as described below. We would like to note that most of the reviewer's strongest complaints come from a misunderstanding of what we are arguing, and thus we focus most of our modifications towards clarifying the text. For example we are not arguing that you should BELIEVE the future projections of the 'better' models more, we are just using model skill to CHARACTERIZE model future projections.

Abstract Generally the abstract has improved and it is easier to follow the main results of the paper. However, this reviewer feels it fails to summarize the main points of the paper clearly.

We modify the abstract to more clearly state the main points of the paper and summarize the results.

"Here our goal is to characterize the LAI projections from the latest generation of Earth system models (ESMs) for the Representative Concentration Pathway (RCP) 8.5 and RCP4.5 scenarios. ."

We also rewrite some parts of the abstract to be more clear.

Introduction Generally the introduction has improved and it reads better. However, this reviewer feels it lacks a clear definition of the objectives and the logic behind them.

We revise the introduction following the reviewer's comments (below) to make it more clear what our objectives are.

Page 4, line 15. This should not be included here, but in the methodology.

It is not clear what the reviewer is objecting to, perhaps the mention of the CMIP5? We think it is appropriate to introduce the CMIP5 in the introduction of the paper.

Page 4 lines 14 – page 4 line 7. This reviewer feels the introduction fails to clearly state the objectives of the paper and to justify them.

We modify the last paragraph in the introduction to be more clear about the goals of this paper:

"Here, our goal is to characterize the ESM projections of future LAI in order to better understand how LAI is projected to change."

The justification for our goal lies in the first paragraph, where we say:

The surface energy budget, as well as plant-based emissions and deposition of aerosols and chemically or radiatively important gases, are also sensitive to predicted LAI (e.g. Oleson et al., 2013). Therefore, small errors in simulated LAI can become large errors in many ESMs' biophysical and biogeochemical processes, and changes in LAI alone can change climate (e.g. Bounoua et al., 2000; Ganzefeld et al., 1998; Lawrence and Slingo, 2004; Oleson et al., 2013; Kala et al., 2014). Unlike many biophysical attributes, LAI can be observed from satellite (Zhu et al., 2013), and thus represents one of the few land carbon or vegetation variables that can be directly evaluated in coupled models (e.g. Randerson et al., 2009; Luo et al., 2012, Anav et al., 2013b). Finally changes in LAI, and the related normalized difference vegetation index (NDVI), can indicate ecosystem health and natural resource availability. As such, LAI is widely used within the famine prediction community (Funk and Brown, 2006; Groten, 1993) and represents a variable that is easy to use in climate impacts studies. Thus it is important to consider the 21st century projections for LAI in Earth System Models.

Methods Generally this section is still poorly written. The section could greatly benefit from better sub---headers for each section (datasets, data analysis, etc.). In the current version the authors mixed the explanation of the datasets they used with the analysis.

We try to accommodate the reviewer's style requests, but also need to make sure the text is clear, and not just a laundry list of analysis without explanation.

Page 5 lines 12---20 Poorly written.

We rewrite lines 12-20:

"We use evaluations of LAI, based on satellite retrieved variables (e.g. Zhu et al., 2013; Anav et al. 2013b; Sitch et al. 2015), to characterize the relationship between model skill and projections (e.g. Steinacher et al., 2010; Cox et al., 2013; Flato et al., 2013; Hoffman et al., 2014) (Section 4). "

Page 6 lines 16---20 should be in the introduction

We move some of the text in these lines to the introduction:

"Changes in LAI variability are also important for understanding the impact of climate change, since they can lead to an increase in the length and frequency of low LAI events, even as mean LAI increases. We consider, therefore, both changes in the mean and the frequency of low LAI events, and how this information compares to precipitation projections, which are commonly used for climate impact studies (e.g. Field et al., 2014)."

Page 7 line 20 I am failing to see why precipitation was not included.

As discussed in page 11, the pattern of the correlation precipitation versus LAI was not robust to which precipitation anomalies were included, as discussed in the next point. We add this back into the supplemental material.

Page 11 line 5. This should be shown at least as supplementary. If the analysis of future LAI is so dependent on precipitation I would expect to see similar patterns in the present---time.

Per the reviewers' request in the last iteration, we removed most of the historical analysis, but we add this back into the supplemental material. The precipitation-LAI relationship is sensitive to the meteorological analysis, which means that getting the right precipitation-LAI correlation in the current climate relies on the models getting the right mean and IAV in precipitation in the current climate. Unfortunately the models are unable to do this. We try to make this point more clear with the following text on page 11:

"The LAI-precipitation relationship across IAV was very sensitive to the meteorology used, and thus is not shown or used to evaluate the current climate simulations of LAI (supplemental material). This implies that errors in the simulations of the mean and variability in precipitation in the current climate, which are very difficult for ESMs to simulate well (e.g. Flato et al., 2014) are very important for the simulation of IAV in LAI."

Page 11 line 15: close to perfect is incorrect, should sate "equal to the observed data".

We modify as suggested.

Results Generally the section is clearer than before and has improved, however it still feels like a collection of different papers, lacking coherence across sections. Some of the results are not well summarized and the reader has to jump from one figure to another to understand them. Additionally the authors mix results with discussion, making it difficult for the reader to fully follow the story. There are several wrong generalization regarding the link between LAI, climate and food security that subtract value form the paper.

The reviewer suggests rewriting this section and the next, dividing the results from the discussion. This is a style issue, and the style that the reviewer suggests is typical for Global Change Biology, for example, but is not intuitive nor the simplest style for the lead author on this paper (she finds this style extremely difficult to read, preferring the journal styles that she usually publishes in, for example Global Biogeochemical Cycles), and thus we do not chose to rewrite the paper in this way. However, we modify the text following many of the reviewer's specific comments below.

Section 3.1 I fail to understand why the authors choose 3 time periods across the century and 1 RCP8.5; instead of focusing in the changes by the end of the 21st century across all RCPs (or at least the two used here). It is hardly surprising to see the increasing change in LAI across time for the same RCP. I also found hardy surprising to find a correlation across RCPs, in spite of the different land use forcing.

This paper represents the first time that LAI is analyzed, and a very important point is how LAI projections behave with time, and whether the LAI changes are gradual or there is a 'tipping' point type behavior (not seen here). It is normal for IPCC variables to be analyzed at various times across the future projections to better understand them and characterize their behavior (e.g. as in the IPCC SPM or papers like Tebaldi et al., 2011 looking at climate projections).

We add text into the methods this point more clear (Section 2.2: at the end of the first paragraph): "We focus on three time periods throughout the twenty-first century because the change in LAI can potentially switch between positive and negative in these different time periods (e.g. Lombardozzi et al. 2014), and we want to identify whether the changes through time are gradual, or if there is a tipping point."

The land use changes in the future projections are very small in all of the RCPs compared to the land use changes that occurred already in the historical period, so we add in this point (in Section 2.4 where we discuss the impact of land use on the simulations):

"Future simulations are unlikely to be more sensitive than the historical simulations to land use and land cover change, because the scenarios include less future land cover change than has occurred historically (Hurtt et al., 2011; Van Vuuren et al., 2011)."

Section 3.2. I do not agree with the usage of LAI as a proxy for food security, although this

approach has been used to analyse present---day changes I believe it is not solid to use it for future projections. It is easier to justify this section saying LAI is a general indicator of vegetation health.

We add in discussion of vegetation health, as suggested by the reviewer. Since two of the authors on this paper, as well as some of the funding for this paper comes from the food security field (specifically for pasture usage), we are more interested in this point than the reviewer apparently is, and thus leave in the references to the potential linkages to food security issues.

In section 3.2: "This is because LAI and the closely related variable, NDVI, are used for identification and forecasting of drought and famine (e.g Funk and Brown, 2006; Groten, 1993) as well as a general indicator of ecosystem health (e.g. Field et al., 1998)."

Spatial links between P and LAI could have been shown better, using simple linear regressions. Figure 6 is remarkably confusing and overly complicated.

We think Figure 6 is a very clear way to show the difference in projections between P and LAI, and thus prefer to explain this figure better rather than remove. We add in appropriate text, focusing on how this figure allows us to identify specifically WHERE the precipitation and LAI projections differ, which cannot be seen from a correlation coefficient. Note that elsewhere the cross model correlations with precipitation are shown (Section 3.3). Here we add:

"General correlations between precipitation and LAI will be discussed in the next few sections, but here we consider the spatial distribution of at-risk regions as defined by LAI or precipitation changes. "

Section 3.3. The analysis on C stocks appears out of the blue and is simply wrong. The analysis on the CO2 fertilization is too simplistic and the interpretation is likely to be wrong. The authors are not separating the effects of CO2, climate and temperature on LAI; which leads to the false assumption that CO2 is not the main driver of LAI. On the opposite, CO2 is probably the main reason why LAI increased, but also lead to an increase in T and changes in P, which in turn drive LAI trends to decrease in the tropics (dryness) and increase on the NH (temperature). The correct way to perform this analysis is using simulation where CO2 changes but not the radiative forcing and compare them with the full simulations.

We agree completely with the arguments of the reviewer, and these are actually the points we are trying to make in this section, but apparently did not communicate well to this reviewer. Our point is that LAI evolution is likely to be very complicated, very important and needs more analysis explicitly in order to understand what is likely to happen in the future. The other reviewer seemed to find this section clear, so we not completely rewrite, but rather try to modify the text to make these arguments more clear

in the text.

At the beginning of Section 3.3 we add:

"Here we use different model output attributes to characterize the future projections, and focus on the following variables: temperature, precipitation, and vegetation carbon. We also characterize the relationship of carbon dioxide fertilization as simulated in the models to the model projections. Note that there are many other potential drivers of the projected LAI changes that are likely to be important, and thus our study only seeks to consider the most obvious interactions, and highlights the uncertainties in the model-specific drivers of LAI projections. "

and at the end of Section 3.3 we add:

"Across these simulations, whether or not the model includes dynamic vegetation does not significantly correlate with changes in LAI in any of the regions.

Overall, the relationship of land model characteristics and LAI is not straightforward, which argues that more analysis of the complicated interactions between the details of the land biophysics and biogeochemistry, as well as biogeography changes is required in order to better understand and improve model projections "

Section 4.0 the authors still lack a proper discussion on weather current model performance is link to better future simulation. No discussion of the N---effect on limiting NPP (and therefore LAI) is included; also no discussion on how dynamic vs. prescribed vegetation will affect this.

Arguing that others have used a similar methodology is not correct; the cited papers (e.g. Cox et al.) are based on very robust scientific evidence of links between the vegetation NBP and climate (e.g. in Cox et al. is the C residence time in the soils over the tropics). The relationship between LAI and climate is not as straight forward (e.g. the growing season cannot be extended forever, the increase in T also leads to earlier snow---melt which can lead to drought latter on the growing season), hence one cannot assume that current model performance will be the same in the future.

Additionally, ranking model performance against 1 observation may be lead to errors due to the obs---dataset bias. See Forkel et al. 2014 biogeosciences and Forkel et al. 2015 Global Change Biology.

There seems to be a deep misunderstanding on what we are doing in this section. The reviewer seems to think we are arguing that one should BELIEVE that the better performing models have better projections, as the reviewer seems to believe that the Cox et al., 2014 represents. That is not a testable hypothesis, because we don't know what will happen in the future. In contrast what we are trying to do here is to test the

hypothesis that model skill in the current climate is correlated with model projection. This is one the standard ways that climate models are CHARACTERIZED in order to understand their behavior. It does not mean that you should believe that the model projections with the better current climate simulation are more accurate. We also think that the LAI projections themselves should get more attention from scientists, instead of only the carbon storage on land, which is why we are writing this paper.

We add in text to clarify this point and rewrite the introductory sentence for this section: "There are large differences between the different models' projections of future LAI (e.g. Figure 3; Figure S5; Figure 7b). Previous studies have hypothesized that they could reduce the uncertainty in future projections by looking for relationships between model metrics and future projections of climate, and then choosing the models which best match the observations in the current climate (e.g. Cox et al., 2013; Hoffman et al., 2014) or by subsampling models for different regions by their performance (e.g. Steinacher et al., 2010). In this section we explore both approaches. In essence, we are looking for a correlation between current model performance and future projections, this correlation has been used in some studies to argue for a more correct answer, and to reduce the uncertainty in the future projections. In many cases in climate modeling and projections, there is no correlation between model skill in current climate conditions and projections (e.g. Cook and Vizy, 2006), however in some limited cases there is a correlation between metric score and a projection, and one is able to constrain future projections (e.g. Cox et al., 2013; Steinacher et al., 2010). Here we consider whether such a case applies. In doing this type of analysis, we are making an assumption that model skill in the current climate translates into better model projections, which may be a product of real model differences or a statistical error. The advantages and disadvantages of using this type of approach are discussed in more detail in Flato et al. (2013). Here we do not advocate that such an approach leads to a better projection, but rather simply use this approach to characterize the future model projections."

We do discuss the N limitation in the models, as that is related to the CO2 fertilization (Section 3.3), but we can't really span the entire space on this issue, as there is only the CLM-CN included in this set of models as the only N-limited model. We try to make this more clear. Of course, all the models included here have prognostic vegetation, or we wouldn't be evaluating their LAI. But perhaps the reviewer means the biogeography or the inclusion of dynamic vegetation? We add in a statement that the inclusion of dynamic vegetation does not correlate with the LAI projections (in Section 3.3, as described in the point above).

Thank you for bringing to our attention the Forkel et al., papers. We add in a reference to the Forkel et al., 2015 paper into the section discussing the many issues with the LAI data (Section 4.1):

"However, the satellite derived LAIs have biases; for example, they underestimate high LAIs due to being unable to see all the leaf layers in closed canopies or

overestimate LAIs in more arid regions, and thus there may also be an error in the observational dataset (see discussion in Anav et al. 2013b; Pfeifer et al. 2014 or Forkel et al., 2015, for example)."

Section 5.0 again the author interpretation to some of the results is not correct (e.g. the LAI trends in RCP4.5 are smaller than in RCP8.5 due to CO2 not climate forcing).

We agree completely with the reviewer's point, and try to clarify this point. CO2 has both a biogeochemical (co2 fertlization) AND a climate impact, and we cannot deconvolve these in this paper. We modify in Section 5.0:

"The change in LAI appears to grow with carbon dioxide and temperature increases across regions over the 21st century (Figure 3). Changes in LAI projected in the RCP4.5 are largely consistent with changes in RCP8.5, but have a reduced amplitude due to the smaller carbon dioxide and climate forcing."

I am missing an analysis linking P to low---LAI over the historical period.

As discussed above, we add this into the supplemental material.

Tables Table 4 seems unnecessary.

We think that Table 4 shows important results suggesting that there is a relationship in the models between skill in the present time and future projections, and thus this is an important Table for characterizing the model projections of LAI.

Figures

Figure quality and control has improved and the new figures and clearer and better. However they are hardly self---explanatory, lacking information such as titles, legend names, etc.

We carefully review each figure and add in more information in the figures, although perhaps the problem is that in revision the figure captions are not attached to the figures, as they will be in the final paper.

Reviewer #2

The focus of manuscript by Mahowald et al. is on changes in LAI as simulated by the CMIP5 models. In revised manuscript, Mahowald et al. accounted for most of the reviewer suggestions. This improved the paper considerably and I support its publication. I have only minor comments listed below.

We appreciate the reviewer's careful reading of the paper and suggestions for revisions.

- In response to the reviewer comments, the authors included analysis of the 2nd scenario (RCP4.5). Since results appear to be almost linearly scalable with the forcing, this had not added much value to the manuscript, but rather led to more complicated figures such as Fig. 3 and Fig. 11. Not sure it was a useful suggestion, but it is up to the authors whether to keep it as it is.

We agree that the addition of the RCP4.5 is perhaps not as useful as it could be, and move the Figure 3 into the supplemental material, which allows the figure to add to the argument that the LAI seems to be scalable, without complicating the main text. Fig 11 doesn't reference RCP4.5, so we do not move that figure.

p.5, l. 13: The reference to Friedlingstein et al. 2006 is not appropriate here since it analyses CMIP3 models. You can cite instead the paper by Arora et al., 2013.

Good point: we revise to use Arora et al., 2013.

Figure 3c, what are the units: %? The scale is from 0 to 0.7, if these are %, numbers are negligible.

The units are fractions: we revise to make more clear.

Figure 4 shows changes in LAI in relation to changes in temperature. This looks informative, but I wonder how significant is an effect of CO2 on LAI, since CO2 modifies plant productivity and affect LAI directly, without changes in climate. There is some discussion of CO2 fertilization on page 19, however, it is unclear what are consequences for sensitivities shown on Fig. 4.

We agree completely with the reviewer, which we had thought we had discussed in the original text, that this is likely to be strongly a function of CO2, and try to make this point more clear in the text. We cannot de-convolve the impacts of CO2 fertilization from climate in this paper. We rewrite: "Recognize that the change in temperature scales with the change in CO_2 forcing from carbon dioxide fertilization as well as other physical variables such as precipitation (e.g. Mitchell, 2003; Moss et al., 2010)."

Figure 4: The color code of different models is difficult to distinguish in the printed copy, especially for color-blind readers. Could models be represented by unique symbols, eg as on the Fig. 11? Also, adding RCP4.5 results on this plot is confusing as they usually align with RCP8.5 results at the top of the other lines. Could results of RCP4.5 analysis be removed from these plots?

We revise this plot consistent with the reviewer's suggestions, and put the additional lines from RCP4.5 into a supplemental plot, so show that the RCP4.5 is scalable from RCP8.5.

1	Projections of Leaf Area Index in Earth System Models
2	
3	N. Mahowald*1, F. Lo ¹ , Y. Zheng ¹ , L. Harrison ² , C. Funk ² , D. Lombardozzi ³ , C. Goodale ⁴
4	
5	
6	¹ Department of Earth and Atmospheric Sciences, Cornell University, Ithaca, NY,
7	14853
8	² Department of Geography, University of California at Santa Barbara, Santa Barbara,
9	CA 93106
10	³ Climate and Global Dynamics Division, National Center for Atmospheric Research,
11	Boulder, CO, 80307
12	⁴ Department of Ecology and Evolutionary Biology, Cornell University, Ithaca, NY
13	14853
14	
15	*Corresponding author: mahowald@cornell.edu
16	
17	

Abstract

1

- $2\qquad \hbox{The area of leaves in the plant canopy, measured as leaf area index (LAI), modulates}$
- 3 key land-atmosphere interactions, including the exchange of energy, moisture,
- 4 carbon dioxide (CO₂), and other trace gases and aerosols, and is therefore an
- 5 essential variable in predicting terrestrial carbon, water, and energy fluxes. Here
- 6 <u>our goal is to characterize the</u> LAI projections from the latest generation of Earth
- 7 system models (ESMs) for the Representative Concentration Pathway (RCP) 8.5 and
- 8 RCP4.5 scenarios. On average, the models project increases in LAI in both RCP8.5
- 9 and RCP4.5 over most of the globe, but also show decreases in some parts of the
- tropics. Because of projected increases in variability, there are <u>also</u> more frequent
- 11 periods of low LAI <u>across broad regions of the tropics</u>. Projections of LAI changes
- varied greatly among models: some models project very modest changes, while
- 13 others project large changes, usually increases. Modeled LAI typically increases with
- modeled warming in the high latitudes, but often decreases with increasing local
- warming in the tropics. The models with the most skill in simulating current LAI in
- the tropics relative to satellite observations tend to project smaller increases in LAI
- 17 in the tropics in the future compared to the average of all the models. Using LAI
- 18 projections to identify regions that may be vulnerable to climate change presents a
- 19 slightly different picture than using precipitation projections, suggesting LAI may be
- 20 an additional useful tool for understanding climate change impacts. Going forward,
- 21 users of LAI projections from the CMIP5 ESMs evaluated here should be aware that
- 22 model outputs do not exhibit clear-cut relationships to vegetation carbon and
- 23 precipitation. Our findings underscore the need for more attention to LAI

Cornell University 11/30/2015 8:18 AM

Deleted: We

Chris Funk 12/27/2015 3:45 PM

Deleted:

Cornell University 11/30/2015 8:19 AM

Deleted: examine

Cornell University 1/7/2016 11:33 AM

Deleted: across broad regions of the tropics

Cornell University 1/7/2016 11:33 AM

Deleted: .

Cornell University 1/7/2016 11:33 AM

Deleted: Projections for both RCP8.5 and 4.5 produce similar LAI trends, with reduced magnitude for RCP4.5.

Cornell University 1/7/2016 11:34 AM

Deleted: Projected increases in LAI generally occur in the same regions that are projected to experience increases in precipitation.

Cornell University 1/7/2016 11:32 AM

Deleted: incorporating temperature influences,

Cornell University 1/7/2016 11:34 AM

Deleted: to

Cornell University 1/7/2016 11:34 AM

Deleted: the climate change impacts community

<u>projections</u>, in terms of understanding the <u>drivers of projected changes and</u>

Providing future projections of climate change feedbacks and impacts is one of the

improvements to model skill.

3

4

5

1

2

1.0 Introduction

6 goals motivating the development of Earth system models (ESMs). The latest 7 generation of ESMs includes land models that simulate the temporal evolution of 8 carbon and vegetation (Friedlingstein et al., 2006). To do so, these models predict 9 leaf area index (LAI) and other carbon cycle variables. LAI represents the amount of 10 leaf area per unit land area, and is an important land carbon attribute. Many ESMs 11 calculate leaf-level carbon and water fluxes, which are then scaled regionally and 12 globally based on LAI (e.g. Oleson et al., 2013). The surface energy budget, as well as 13 plant-based emissions and deposition of aerosols and chemically or radiatively 14 important gases, are also sensitive to predicted LAI (e.g. Oleson et al., 2013). 15 Therefore, small errors in simulated LAI can become large errors in many ESMs' 16 biophysical and biogeochemical processes, and changes in LAI alone can change 17 climate (e.g. Bounoua et al., 2000; Ganzefeld et al., 1998; Lawrence and Slingo, 2004; 18 Oleson et al., 2013; Kala et al., 2014). Unlike many biophysical attributes, LAI can be 19 observed from satellite (Zhu et al., 2013), and thus represents one of the few land 20 carbon or vegetation variables that can be directly evaluated in coupled models (e.g. 21 Randerson et al., 2009; Luo et al., 2012, Anav et al., 2013b). Finally changes in LAI, 22 and the related normalized difference vegetation index (NDVI), can indicate 23 ecosystem health and natural resource availability. As such, LAI is used within the

Cornell University 1/7/2016 11:32 AM

Deleted: edictions

Laura Harrison 12/23/2015 11:44 AM

Deleted: LAI projections are not simply related to vegetation carbon or precipitation, and yet are extremely important, suggesting that further analysis and improvements in predictions of LAI is required.

Laura Harrison 12/23/2015 11:57 AM

Deleted: widely

1 famine prediction community (Funk and Brown, 2006; Groten, 1993) and 2 represents a variable that is easy to use in climate impacts studies. Thus it is 3 important to consider the 21st century projections for LAI in Earth System Models. 4 The current generation of ESMs has prepared historical and future scenario 5 simulations within the Coupled Modeling Intercomparison Project (CMIP5) (Taylor et al., 2009). There have been extensive evaluations and comparisons of the future 6 7 projections of the land, ocean, and atmospheric carbon cycle in the ESMs in the Laura Harrison 12/23/2015 11:58 AM Deleted: at 8 CMIP5 (e.g. Arora et al., 2013a; Friedlingstein et al., 2013; Jones et al., 2013). There 9 has also been comparison of ESM-simulated seasonal variability in LAI against 10 satellite-based observations for the high latitudes (Anav et al., 2013a; Murray-11 Tortarolo et al., 2013), as well as comparisons of LAI and other variables in ESMs 12 across the globe (Anav et al., 2013b). Additionally, Shao et al. (2013), Mao et al. 13 (2013), and Sitch et al. (2015) evaluated the relationship between the carbon cycle 14 and other variables, such as temperature, or LAI, over decadal and longer time 15 scales. These ESM-based comparisons build on the long history of evaluation of 16 model simulations of vegetation properties and carbon balance (e.g. Cramer et al., 1999). 17 18 Here, our goal is to characterize the ESM projections of future LAI in order to Cornell University 11/30/2015 8:20 AM Deleted: we 19 better understand how LAI is projected to change. Most of our analysis emphasizes Cornell University 11/30/2015 8:20 AM Deleted: examine 20 the Representative Concentration Pathway (RCP) 8.5, the most extreme future 21 scenario, and we contrast it with RCP4.5, a less extreme scenario (van Vuuren et al., 22 2011) (Section 3). We characterize both the model mean LAI projected change, as Cornell University 11/30/2015 8:20 AM Deleted: evaluate 23 well as the model mean divided by the standard deviation (e.g. Meehl et al., 2007;

Tebaldi et al., 2011). In addition, we consider whether LAI projections can help the 1 Cornell University 1/7/2016 11:36 AM Deleted: also 2 climate impact community anticipate regions that may experience increased climate 3 exposure and increased risk of food insecurity in the future. Changes in LAI Chris Funk 12/27/2015 3:59 PM Deleted: increased variability are also important for understanding the impact of climate change, since 4 5 they can lead to an increase in the length and frequency of low LAI events, even as mean LAI increases. We consider, therefore, both changes in the mean and the 6 7 frequency of low LAI events, and how this information compares to precipitation 8 projections, which are commonly used for climate impact studies (e.g. Field et al., 9 2014). We also consider what model traits may be related to the spread in the future Cornell University 12/7/2015 12:53 PM Deleted: 10 model projections (Section 3). We use evaluations of LAI, based on satellite-based 11 observations (e.g. Zhu et al., 2013; Anav et al. 2013b; Sitch et al. 2015), to Chris Funk 12/27/2015 4:01 PM Deleted: variables 12 characterize the relationship between model skill and projections (e.g. Steinacher et Cornell University 12/7/2015 2:50 PM **Deleted:** see if there is 13 al., 2010; Cox et al., 2013; Flato et al., 2013; Hoffman et al., 2014) (Section 4). Cornell University 12/7/2015 2:50 PM Deleted: a 14 Section 5 presents our summary and conclusions. Cornell University 12/7/2015 12:54 PM Deleted: , which could be used to constrain 15 future model projections 2.0 Methods and datasets 16 17 18 2.1 Model datasets 19 Coupled carbon model experiments were included as part of the CMIP5 Cornell University 11/30/2015 8:26 AM Deleted: in 20 experiments (e.g. Arora et al., 2013: Taylor et al., 2009). The historical simulations Cornell University 12/7/2015 2:36 PM Deleted: Friedlingstein et al., 2006 21 and Representative Concentration Pathway for 8.5 (RCP8.5; van Vuuren et al., 2011; Riahi et al., 2011), using prescribed carbon dioxide concentrations, were analyzed 22 23 here (Table 1). We chose to focus on the RCP8.5 scenario as it has the largest

changes in carbon dioxide and climate. Analysis of the RCP4.5 scenario (Wise et al. 1 2 2009; van Vuuren et al., 2011) is also included for comparison for the models which Cornell University 11/30/2015 8:26 AM **Deleted:** using the models for which 3 included the RCP4.5 simulations at the CMIP5 archive (all models except BNU-ESM Chris Funk 12/27/2015 4:02 PM Deleted: data and CESM-BGC). 4 Cornell University 11/30/2015 8:27 AM Deleted: were available for download 5 Model variables analyzed included monthly-mean precipitation, near surface air temperature, vegetation carbon stock and LAI. Only models which had data for 6 7 all these variables for both historical and RCP8.5 scenarios were included in this 8 study. Some models submitted multiple versions, at different resolutions or with 9 slightly different physics (Table 1). Even though some of the models are closely 10 related (e.g. CESM1-BGC and NorESM-ME), we include different configurations of 11 the same model. 12 2.2 Model future projection analysis, Cornell University 11/30/2015 8:43 AM Formatted: Font:Bold 13 This analysis examines model mean changes between the current climate (1981-Cornell University 11/30/2015 8:39 AM Deleted: 14 2000) and future climate time periods (2011-2030, 2041-2060 and 2081-2100). To Cornell University 11/30/2015 8:41 AM Formatted: Indent: First line: 0" 15 identify the location where models project statistically significant changes, we Fiona A. Lo 12/11/2015 10:15 AM **Deleted:** these changes will be 16 analyze the ratio of the mean change to variability; this is accomplished by dividing 17 the mean changes over 20 year time periods by the standard deviation over the 18 current climate (1981-2000) and shown in terms of standard deviation units (e.g. 19 Mahlstein et al., 2012; Tebaldi et al., 2011). Previous studies have shown that the 20 spatial and temporal scale used to define these changes can <u>determine</u> whether Chris Funk 12/27/2015 4:05 PM **Deleted:** be important for 21 these signals are statistically significant (Lombardozzi et al. 2014). We focus on Cornell University 1/7/2016 11:39 AM Deleted: 22 three time periods throughout the twenty-first century because the change in LAI 23 can potentially switch between positive and negative in these different time periods

(e.g. Lombardozzi et al. 2014), and we want to identify whether the changes through time are gradual, or if there is a tipping point.

3 Changes in LAI variability are also important for understanding the impact of climate change. To estimate the periods of low LAI and low precipitation, we 4 5 calculate the fraction of the time during which the variable is one standard deviation (evaluated in the 1981-2000 time period) below the current mean (1981-2000). By 6 7 definition, if the variables have a Gaussian distribution, each gridbox would be 8 considered having a "Low LAI" for 1/6 (16%) of the time, and this is approximately 9 true at most grid points (not shown). We use this metric to estimate the fraction of 10 the time in the future that this condition exists, and specifically whether it increases 11 in the future.

2.3. Observational data

1

2

12

13

14

15

16

17

18

19

20

21

22

23

LAI, data derived from satellite over the 30-year period 1981-2010 is used to evaluate the CMIP5 model skill in the current climate. This observational dataset is derived using neural network algorithms using the Global Inventory Modeling and Mapping (GIMMS) Normalized Difference Vegetation Index (NDVI3g) and the Terra Moderate Resolution Spectroradiameter (MODIS) LAI (Zhu et al., 2013). The satellite data are only available over regions with green vegetation, and thus are lacking over desert and arid regions. A detailed description of the algorithm and comparison to ground-truth observations are shown in Zhu et al. (2013). Compared with field-measured LAI, Mean Squared Errors (RMSE) in the satellite LAI estimates are estimated to be approximately 0.68 LAI, for spanning LAI ranges from < 1 to

Cornell University 11/30/2015 8:30 AM

Deleted: For example, in some regions there is a predicted increase in the mean LAI as well as an increase in the variability. This can lead to an increase in the length and frequency of low LAI events, even as mean LAI increases. The length and frequency of these periods matter for understanding the potential for drought and ramifications for agriculture or ecosystems.

Cornell University 12/7/2015 3:04 PM

Deleted: percent

Cornell University 11/30/2015 8:41 AM

Deleted: 2

Chris Funk 12/27/2015 4:06 PM

Deleted: Leaf Area Index (

Chris Funk 12/27/2015 4:06 PM

Deleted:)

Laura Harrison 12/23/2015 12:16 PM

Deleted: s

almost 6 (Zhu et al., 2013). Comparisons with ground-based observations confirm 1 2 that the new LAI product also seems to capture observed interannual variability 3 patterns (Zhu et al., 2013). Gridded temperature data for the period 1981-2010 were derived from the 4 5 Global Historical Climatology Network and Climate Anomaly Monitoring System 6 (GHCN_CAMS) 2m temperature dataset (Fan and Dool, 2008). Estimates of the 7 uncertainty in temperature gridded datasets suggest that the uncertainty in 8 temperatures at a grid box level is estimated to be between 0.2 and 1°C (Jones et al., 9 1997; Fan and Dool, 2008). 10 11 2.4 Methodology for evaluation of current climate LAI simulation Cornell University 11/30/2015 8:41 AM Deleted: 3 12 Several recent studies have used the same new satellite-derived LAI dataset 13 (GIMMS LAI3g) in land model evaluation (e.g. Murray-Tortarolo et al 2013; Anav et 14 al. 2013a; 2013b, Mao et al. 2013, Sitch et al. 2015), including some of the same land 15 models used here. Thus we do not repeat a complete evaluation of model LAI 16 compared to satellite LAI. We use the satellite LAI dataset to consider whether there 17 is a relationship between the models' ability to simulate LAI in the current climate 18 and the models' climate projections. We use a few basic metrics in this study (Table 19 2), which are described briefly below.

Results for the model and observations are evaluated on a 2.5°x2.5° grid based on the observed temperature data grid (see Section 2.3). For the metric analysis here, the averages shown are grid-box means, not areal averages. This allows us to use similar weighting for both the averages and the rank correlation

20

21

22

23

Cornell University 11/30/2015 8:43 AM

Deleted: 2

1 coefficients, and tends to weight the global analysis towards high latitudes. However, 2 most of the analysis focuses on regional areas (tropical (<30°), mid-latitudes (>30° 3 and <60°) and high-latitudes (>60°), where the differences between weighting by 4 area and weighting by grid box are reduced. 5 We compare the satellite-based observed (LAI3g) and model-simulated mean 6 LAI for the current climate (similar to previous studies e.g. Randerson et al., 2009; 7 Luo et al., 2012; Anav et al., 2013b). The period 1981-2010 is used for this 8 comparison. To examine regional differences in LAI simulations, the annual mean 9 LAI in the models and observations are averaged and compared over different 10 areas: global, tropical (<30°), mid-latitudes (>30° and <60°) and high-latitudes 11 (>60°) (Table 2: mean LAI: model/obs.). A second metric evaluates the models' 12 ability to capture spatial variations in LAI, using the spatial correlation across the 13 grid-boxes of the annual mean LAI in the model compared to the observations (e.g. 14 Anav et al., 2013b; Table 2: Mean: Corr.). 15 Important for this study is the consideration of the temporal variability 16 simulated in the model. The magnitude of the seasonal cycle is calculated as the 17 standard deviation of the climatological monthly means at each grid box. This metric 18 is slightly different than how LAI has previously been evaluated in some studies (e.g. 19 Anav et al., 2013a; Murray-Tortarolo et al. 2013; Sitch et al. 2015), but is more 20 similar to analyses of other climate variables (Glecker et al., 2008), facilitating 21 inclusion of LAI within climate model evaluations. Metrics for the seasonal cycle 22 were computed using a spatial average over each region (Table 2: Std. Dev. 23 Seasonal: Model/obs.). For the seasonal cycle, the ability to capture the timing of

phenology can be important (e.g. Anav et al., 2013a, Zhu et al., 2013). To analyze this 1 2 ability, we computed the temporal correlation of observed and model-simulated 3 monthly means at every grid box, and then averaged over each region (Table 2: Seasonal Avg. Corr.). 4 5 To evaluate the models' ability to simulate LAI interannual variability (IAV), we consider the magnitude of the interannual variability, which is calculated as the 6 7 standard deviation of annual mean LAI across years at each grid box (e.g. Zhu et al., 8 2013). The IAV is then spatially averaged and compared between the model and 9 satellite observations (Table 2: Std. Dev. IAV: Model/obs.). We focus our study on 10 IAV, based on the inter-annual means, but there may be important changes in the 11 seasonal cycle or length of growing season on an interannual time basis, which our 12 simple approach does not consider (e.g. Murray-Tortarolo et al. 2013). 13 Previous studies have examined correlations between temperature and 14 satellite- derived LAI (e.g. Anav et al., 2013a; 2013b, Zhu et al., 2013) or the closely 15 related normalized difference vegetation index (NDVI; Zeng et al. 2013). Observed 16 variations of LAI at high latitudes tend to be dominated by changes in temperature, 17 while the tropics are more dominated by moisture (Anav et al., 2013a; 2013b, Zeng 18 et al., 2013), which is also seen in coupled-carbon climate models for carbon cycle 19 variables (e.g. Fung et al., 2005). In order to understand what may be driving the 20 IAV in the LAI, we calculate metrics to examine the rank correlation between 21 anomalies in LAI and anomalies in temperature and trends with time. Although 22 correlations do not identify causation, they can help identify the strength of

23

relationships among various driving factors.

This analysis focuses on the relationship between temperature and LAI for comparing interannual variability in the modeled and observed datasets. Sensitivity studies have indicated that the grid-box level relationship between temperature and LAI is a good indicator of features intrinsic to the model, rather than to the meteorology forcing the model (supplemental material Figure S1; as seen also in Anav et al., 2013a; Murray-Tortarolo et al., 2013). This was not the case for the relationship between precipitation and LAI. In sensitivity studies conducted as part of this study, we forced the Community Land Model (Lawrence et al., 2012; Lindsay et al., 2014), which is the land model used in the CESM (Table 1), with reanalysis derived data data (Qian et al., 2006; Harris et al., 2013) instead of model derived winds. The LAI-precipitation relationship across IAV was very sensitive to the meteorology used, and thus is not shown or used to evaluate the current climate simulations of LAI (supplemental material; Figure S1). This implies that errors in the simulations of the mean and variability in precipitation in the current climate, which are very difficult for ESMs to simulate well (e.g. Flato et al., 2014), are very important for the simulation of IAV in LAI.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

Land use, especially the conversion from natural vegetation to agricultural use, can heavily perturb the mean and evolution of the seasonal cycle and interannual variability in current climate LAI. To determine whether this changes our model evaluation, we exclude grid boxes with more than 50% of agricultural land use, based on Ramankutty et al. (2008). Results of the model evaluation with and without agricultural grid-box were quantitatively and qualitatively similar to those presented here, and thus we include all grid-boxes in this analysis. Future

Chris Funk 12/27/2015 4:13 PM Deleted:

Chris Funk 12/27/2015 4:14 PM

Deleted: e

Chris Funk 12/27/2015 4:14 PM

Deleted: ,

1 simulations are unlikely to be more sensitive than the historical simulations to land 2 use and land cover change, because the scenarios include less future land cover 3 change than has occurred historically (Hurtt et al., 2011; Van Vuurden et al., 2011). 4 For ease of interpretation, we present the metrics described above in Figure 5 9, in which higher numbers represent a better simulation. For correlations, this Cornell University 12/7/2015 1:24 PM Deleted: 10 representation is straightforward: 1 is a perfect correlation and lower values 6 7 represent a worse simulation. For the other metrics that are not correlations, we 8 convert the statistics to values similar ranges to facilitate ease of display. The mean 9 model bias metric (model/obs) is normalized to a value that varies between 0 and 1, 10 with 1 being close to the observed data. This approach penalizes models which have Cornell University 11/30/2015 8:45 AM Deleted: perfect 11 too high of a mean equally with model that have too low of a mean, using the 12 following formula (Figure 9): Cornell University 12/7/2015 1:24 PM Deleted: 10 13 $Model\ Evaluation\ Value = \frac{2}{\left\{\frac{Model\ Mean}{Observed\ Mean} + \frac{Observed\ Mean}{Model\ Mean}\right\}}$ 14 (1) 15 We use this method to convert mean biases and standard deviation biases to a 16 model evaluation value (MEV). This is a slightly different method than used in 17 previous studies (e.g. Gleckler et al., 2008), as the MEV does not square the standard 18 deviations. Since we use ranks and rank correlations, the difference between these 19 methods is unlikely to be important, and allows us to use a similar ranking method 20 for mean and standard deviation comparisons. 21 22 3.0 Results 23 3.1 Future projections

First we consider the model mean projections of change in LAI for RCP8.5, similar to
analyses for other standard model variables, and show their evolution through the

3 21st century (e.g. Meehl et al., 2007). Across most of the globe, LAI is projected to

4 increase through 2081-2100, with small decreases projected for parts of Central and

South America and Southern Africa (Figure 1). The increases in LAI are largest in

high latitudes, mountainous regions (e.g. Tibetan plateau) and some parts of the

7 mid-latitudes and tropics (Figure 1; for reference, mean satellite observed LAIs in

the current climate are presented in Figure S2). Notice that in this study we use

9 projections of human land use based on the RCP8.5 or RCP4.5, and thus an

important human role in future land cover change is driven by the assumptions of

11 the scenario chosen for these studies. Generally, for all the RCPs, there is less land

 $\,$ 12 $\,$ $\,$ use and land cover change projected in the future than occurred in the past (e.g. van

13 Vuuren et al., 2011; Ward et al., 2014).

5

6

8

10

14

15

16

17

18

19

20

21

22

In order to isolate the changes that are statistically significant, for each model we divided the change in LAI by the IAV standard deviation. Values over 1 are considered statistically significant (e.g. following Tebaldi et al. 2011; Mahlstein et al. 2012). Using this approach, statistically significant changes in LAI start over the high latitudes, and spread over much of the globe with time (Figure 2). By 2081-2100, the increases in LAI are 8 times as large as IAV over large parts of high latitude regions, as well as the Tibetan plateau and some desert regions, indicating large changes (Figure 2c). Part of the reason for these very large normalized LAI

values is that they have low IAV in the current climate. A few isolated tropical

Cornell University 12/7/2015 3:08 PM

Formatted: Superscript

Cornell University 12/7/2015 1:12 PM

Deleted: .

Cornell University 12/7/2015 1:00 PM

Deleted: 1

regions are projected to have statistically significant reductions in mean LAI, such as in Central America and the Amazon basin.

3 Examination of the RCP4.5 shows a similar pattern of an increase in LAI over most of the globe, although lower in magnitude, based on either the mean change in 4 5 LAI, or the normalized LAI change (Figure \$3a and \$3b). This result suggests that the pattern of change in LAI, as seen in the literature for temperature or even to a 6 7 lesser extent for precipitation, is similar across different climate change scenarios, with the magnitude dependent on the magnitude of the forcing (e.g. Mitchell, 2003; 8 9 Moss et al., 2010). There is a consistent relationship between changes in LAI and 10 temperature across the different time periods for each model; that is, most models 11 and regions show a constant slope between changes in LAI and temperature (Figure 12 3). Most models even show a similar slope between LAI and temperature for the 13 RCP4.5 as the RCP8.5 (Figure <u>S</u>4). Recognize that the change in temperature <u>scales</u> 14 with the change in CO₂ forcing from carbon dioxide fertilization as well as other physical variables such as precipitation (e.g. Mitchell, 2003; Moss et al., 2010). This 15 16 similarity in slope for each model across RCPs and time periods breaks down in the 17 tropics for a few of the models, as some show steeper increases in LAI at warmer 18 temperatures and others shift from LAI increases to declines as warming continues 19 (GFDL, IPSL, MIROC and MPI models) (Figure 3b). Across the tropics, LAI is 20 projected to increase in some regions and decrease in others, so small changes in 21 the relative area of these changes can lead to large shifts in the regional net mean 22 LAI change. The value of spatial correlations between the RCP4.5 and RCP8.5 mean 23 LAI change at each gridbox for the 2081-2100 time period is 0.81, 0.70, 0.79 and

Cornell University 1/7/2016 11:45 AM

Deleted: s

Cornell University 12/7/2015 1:22 PM

Deleted: 4

Cornell University 12/7/2015 3:05 PM

Deleted: is likely to scale with a change

Cornell University 12/7/2015 3:05 PM

Formatted: Subscript

Cornell University 12/7/2015 3:05 PM

Deleted: in

Fiona A. Lo 12/11/2015 10:16 AM

Deleted: as well

Cornell University 12/7/2015 1:23 PM

Deleted: 4

1 0.89, for the globe, tropics, mid-latitudes and high-latitudes, respectively (averaged 2 across the models), showing the spatial coherence in the LAI projections between 3 these two RCPs. Even the models with the lowest spatial correlations between the two RCPs (GFDL, IPSL, MIROC and MPI) have statistically significant correlation 4 5 coefficients of 0.45 or higher in the tropics, where correlations are the lowest. 6 The models project a wide range of future changes in LAI (Figure 3). One Cornell University 12/7/2015 1:25 PM Deleted: 4 7 model (BNU-ESM) projects a large global mean increase of over 1 m²/m² by 2081-8 2100. For the other models, projected global mean increases in LAI amounted to 0.5 9 m²/m² or less. Some models (inmcm4, IPSL, MIROC and MPI model versions) 10 projected small net decreases in LAI in the tropics (Figure 3). <u>Inter-model</u> Cornell University 12/7/2015 1:25 PM Deleted: 4 11 differences become even more apparent at the grid-box level, with very different Chris Funk 12/27/2015 4:31 PM Deleted: Between 12 changes in LAI projected by the different models (Figure S5). The spread in model Cornell University 12/7/2015 1:12 PM Deleted: 2 13 projections is discussed further below (section 4.0) in relation to whether there is a 14 relationship between model skill at predicting LAI in the current climate and future Cornell University 12/7/2015 1:26 PM Deleted: can be used to reduce model 15 model projections (e.g. Steinacher et al., 2010; Flato et al., 2013; Cox et al., 2013; spread in these 16 Hoffman et al., 2014). 17 18 3.2 Identifying regions at risk due to climate change 19 In addition to being important for land-atmosphere biophysical and biogeochemical 20 interactions, LAI is also one of the few ESM model variables that is directly usable by Chris Funk 12/27/2015 4:32 PM Deleted: potentially 21 the climate impacts community, along with temperature and precipitation. This is 22 because LAI and the closely related variable, NDVI, are used for identification and 23 forecasting of drought and famine (e.g Funk and Brown, 2006; Groten, 1993) as well

1 as a general indicator of ecosystem health (e.g. Field et al., 1998). Thus LAI 2 projections that identify the regions that are most at risk can help guide and 3 motivate climate adaptation by identifying emergent areas of vulnerability. The model mean view of the future projections of LAI is quite optimistic (Figure 1, 2 and 4 5 3), however, if variability also increases, some regions may experience years with lower LAI more frequently than in current climate, despite having a constant or 6 7 higher mean LAI. In fact, many regions, especially in the tropics, are at risk for 8 more Low LAI years (Figure 4). Here we define % Low LAI as the % of years when Cornell University 12/7/2015 1:26 PM Deleted: 5 9 the annual average is one standard deviation below the current mean (Section 2.2). Cornell University 11/30/2015 8:41 AM Deleted: 1 10 If the variability and mean stayed constant, the % Low LAI would remain at 16%. 11 More Low LAI years are projected for large areas of the tropics and subtropics 12 where projected increases to mean LAI are small in magnitude or negligible (Figure 13 1c vs 4c, for example). Model mean changes between the current climate (1981-Cornell University 12/7/2015 1:26 PM Deleted: 5 14 2000) and future climate time periods indicate substantial (>2x) increases in the 15 frequency of low LAI in important agricultural areas (South America, Australia, 16 Southeast Asia, and parts of Southern Africa) (Figure 4). Increased risk areas in Fig. Cornell University 12/7/2015 1:27 PM Deleted: 5 17 4 also coincide, in some cases, with some of the most food insecure regions of the Cornell University 12/7/2015 1:27 PM Deleted: 5 18 world (e.g. Brown and Funk, 2008; Field et al., 2014). Similar to mean changes in LAI, 19 the %Low LAI for the RCP4.5 at 2081-2100 is similar in pattern and magnitude to 20 that seen earlier in the century for the RCP8.5 scenarios (Figure \$3c vs. Figure 4). Cornell University 12/7/2015 1:27 PM Deleted: 3 21 Next we consider whether using LAI adds information compared to Cornell University 12/7/2015 1:27 PM Deleted: 5 22 precipitation, which is more traditionally used in climate change impacts 23 assessments (e.g. Stocker et al. 2013; Field et al. 2014). General correlations

1 between precipitation and LAI will be discussed in the next few sections, but here 2 we consider the spatial distribution of at-risk regions as defined by LAI or 3 precipitation changes. To do this, we first consider the mean change in normalized Cornell University 12/7/2015 2:56 PM Deleted: F precipitation (Figure 5a) and the % Low Precipitation (Figure 5b), both defined 4 Cornell University 12/7/2015 2:56 PM Deleted: we 5 equivalently to the LAI values (Section 2.2; Figure 2c and Figure 4c, respectively) for Cornell University 12/7/2015 1:27 PM Deleted: 6 the model simulations considered here. Broadly speaking, the changes in 6 Cornell University 12/7/2015 1:27 PM Deleted: 6 7 precipitation seem to occur in similar regions as the changes in LAI, with large Cornell University 11/30/2015 8:41 AM Deleted: 1 8 increases in precipitation over the high latitudes, and decreases over the subsidence Cornell University 12/7/2015 1:28 PM Deleted: 5 9 zones of the tropics, as seen previously (e.g. Meehl et al., 2007; Tebaldi et al., 2012). 10 Note that requiring the mean change to be statistically significant is a much stricter 11 criteria than just an increase in low LAI, and thus the area identified in the two 12 methods is quite different (Figure 5a vs. 5b). Overlaying the regions from LAI and Cornell University 12/7/2015 1:28 PM Deleted: 6 13 precipitation which are either one standard deviation below the mean on average in Cornell University 12/7/2015 1:28 PM Deleted: 6 14 the models (Figure 5c) or see an increase in % Low values (Figure 5d) suggests that Cornell University 12/7/2015 1:28 PM Deleted: 6 15 LAI and precipitation largely show similar areas being at risk due to climate change, Cornell University 12/7/2015 1:28 PM Deleted: 6 16 but there are significant regions which do not overlap. This suggests that there is 17 potentially additional information for climate impact studies using LAI projections 18 than using precipitation alone (Figure 5c and 5d). One of the most noticeable Cornell University 12/7/2015 1:28 PM Deleted: 6 19 differences between LAI and precipitation projections is in the Mediterranean Cornell University 12/7/2015 1:28 PM Deleted: 6 20 region where precipitation is projected to decrease, but LAI is not. Conversely, LAI Chris Funk 12/27/2015 4:39 PM Deleted: in 21 projections suggest that some parts of South America and southern Africa are likely 22 to experience more stress, which are not identified using precipitation. Future

1 studies should consider whether the results of the LAI projections are useful for

2 impact studies specifically in these regions.

3

4

13

14

15

16

17

18

19

20

21

22

23

3.3 Drivers of LAI projections

drivers of LAI projections.

5 Next we consider what drives the differences in model projections for LAI, using the 6 example of RCP8.5 at 2080-2100. Here we use different model output attributes to 7 characterize the future projections, and focus on the following variables: 8 temperature, precipitation, and vegetation carbon. We also characterize the 9 relationship of carbon dioxide fertilization as simulated in the models to the model 10 projections. Note that there are many other potential drivers of the projected LAI 11 changes that are likely to be important, and thus our study only seeks to consider 12 the most obvious interactions, and highlights the uncertainties in the model-specific

By correlating temperature and LAI projections at each grid box for each model we can look for potentially causal relationships between model projections of temperature and LAI (Figure 6). This is analogous to using a ranked correlation coefficient to summarize the scatter in RCP8.5 points in Figure 3, but at each grid box instead of the regional average. There are strong positive correlations between model simulated changes in temperature and LAI in some regions, especially in parts of the northern high latitudes (Figure 6a), suggesting that models with a projected larger warming in the high latitudes also simulate larger increases in LAI. Higher temperatures may drive higher LAI; higher LAIs may also be driving higher

temperatures because of the importance of LAI in changing surface energy fluxes

Cornell University 12/7/2015 1:52 PM

Formatted: Indent: First line: 0.5"

Chris Funk 12/27/2015 4:40 PM

Deleted: ,

Cornell University 12/7/2015 1:51 PM

Deleted: ,

Cornell University 12/7/2015 1:28 PM

Deleted: 7

Cornell University 12/7/2015 1:28 PM

Deleted: 4

Cornell University 12/7/2015 1:28 PM

Deleted: 7

(e.g. Lawrence and Slingo, 2004; Kala et al. 2014). By contrast, there are strong negative correlations across most of the tropics and subtropics (Figure 6a).

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

The projected changes in precipitation are strongly correlated with projected changes in LAI across different models in many locations (Figure 6b). This is consistent with the model mean analysis (Section 3.2) that showed that for most locations changes in LAI occur in the same locations as changes in precipitation (Figure 5). Again, because LAI changes the surface energy fluxes, there may be a feedback from LAI changes to precipitation (e.g. Lawrence and Slingo, 2004; Kala et al. 2014). The correlations seen in this analysis for RCP 8.5 are similar for the RCP4.5 (Figure S6).

Next, we examine the correlation across models between the modeled changes in vegetation carbon stocks and change in LAI between current conditions and 2081-2100 (Figure 6c). The relationship between LAI and vegetation carbon is not straightforward, and depends on the specific biophysics and biogeochemistry algorithms used in the models. Many ESMs calculate photosynthetic rates per unit leaf area; these rates are then extrapolated to canopy-level gross primary production using LAI and other variables (e.g., light, nitrogen and CO₂ availability and leaf physiological parameters) (e.g., See Bonan et al., 2011, Piao et al., 2013). The simulated increases in LAI are correlated across models with simulated increases in plant carbon stocks in many low-LAI regions, including many deserts, grasslands, and tundra ecosystems (Figure 6c). Leaves compose most or all of the aboveground plant biomass in these ecosystems (e.g., Friedlingstein et al. 1999), such that increases in LAI relate directly to increases in plant carbon stocks.

Cornell University 12/7/2015 1:29 PM

Deleted: 7

Cornell University 12/7/2015 1:29 PM

Deleted: 7

Cornell University 1/7/2016 11:50 AM

Deleted:), suggesting that changes in precipitation are correlated with the differences in LAI projections between models

Cornell University 11/30/2015 8:42 AM

Deleted: 1

Chris Funk 12/27/2015 4:43 PM

Deleted: ,

Cornell University 12/7/2015 1:29 PM

Deleted: 6

Cornell University 12/7/2015 1:12 PM

Deleted: 3

Cornell University 12/7/2015 1:53 PM

Deleted: Last

Cornell University 12/7/2015 1:32 PM

Deleted: 7

Cornell University 12/7/2015 1:33 PM

Deleted: 7

1 Changes in LAI correlate more poorly with simulated changes in plant carbon stocks

2 in other regions, with small or negative correlations in many boreal, temperate, and

3 tropical forested regions (Figure 6c). Leaves typically compose only 3-5% of

4 aboveground plant biomass in forests (Friedlingstein et al. 1999), and closed-

canopy forests can contain widely variable stocks of woody biomass that typically

depend more on successional status than LAI or growth rate. Differences in the

fractional composition and turnover of these leaf- and woody tissues should

decouple changes in LAI from changes in carbon stocks in woody biomass. As an

example, in the CLM, the land model for the CESM-BGC, CO₂ fertilization causes a

larger increase to wood allocation (62%) than to leaf allocation (21%) in the

11 Southeastern US (Lombardozzi, personal communication, 2015). Thus, the issue of

12 how LAI responds in different models is interesting and should be considered in

13 future studies.

5

6

7

8

9

10

14

15

16

17

18

19

20

21

22

23

Another important potential contributor to the future projections of LAI is the effectiveness of carbon fertilization in the models (e.g. Arora et al., 2013). Using the carbon dioxide fertilization factor (β -land) from the Arora et al. (2013) study we use a rank correlation to explore the importance of the carbon dioxide fertilization strength for predicting future vegetation carbon and LAI across the models. Naively, we might expect models that respond more strongly with increased carbon uptake under higher CO_2 conditions (i.e. larger β -land) to have greater vegetation carbon and LAI in the future. Globally the correlation with β -land is 0.46 for vegetation carbon and -0.21 for LAI, suggesting that while some of the differences in future

vegetation carbon projections across models are due to differences in the model

Cornell University 12/7/2015 1:33 PM

Deleted: 7

Laura Harrison 12/23/2015 2:59 PM

Deleted: ia

Cornell University 12/7/2015 2:10 PM

Deleted: We would

Chris Funk 12/27/2015 4:51 PM

Deleted: is

1 simulation of CO₂ fertilization, LAI changes are not necessarily related to CO₂ 2 fertilization. The β -land correlation for vegetation carbon is 0.29, 0.47 and 0.60 for 3 tropical, mid-latitude and high latitude, regions, respectively, while for LAI these values are -0.18, -0.09 and 0.21. Thus for high latitudes, especially, the projections 4 5 of LAI appear to be dependent on the way the models' simulate the carbon dioxide fertilization in the different models. This could also be, however, an artifact that the 6 7 two models with the lowest carbon dioxide effect (CESM-BGC and NOR-ESM) use 8 the same land carbon model (Thornton et al., 2009), which predicts low values of 9 LAI in high latitudes for present day and does not tend to increase LAI much in the 10 future. These models also have low carbon dioxide fertilization effects, because of 11 the inclusion of nitrogen limitation, which could be driving the weak positive 12 correlation between model projections of LAI and carbon dioxide fertilization in the 13 high latitudes.

It is interesting that in the tropics the carbon dioxide fertilization is negatively correlated to future LAI changes, and only slightly correlated with vegetation carbon. Again, this could be an artifact of having only two related low carbon fertilization models, as these models see a strong increase in nitrogen mineralization in the tropics in a warming climate, which allows an increase in productivity in the future tropics (Thornton et al., 2009). In other words, the negative correlation in the tropics between LAI projections and CO_2 fertilization could be due to the smaller temperature impact on carbon cycle (γ -land from Arora et al., 2013) in the N-limited models (i.e. the β -land and γ -land are negatively correlated in Table 2 of Arora et al., 2013).

14

15

16

17

18

19

20

21

22

23

Cornell University 12/7/2015 2:42 PM

Deleted: their

Cornell University 12/7/2015 2:43 PM

Formatted: No widow/orphan control, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers Across these simulations, whether or not the model includes dynamic

vegetation does not significantly correlate with changes in LAI in any of the regions.

Overall, the relationship of land model characteristics and LAI is not straightforward, which argues that more analysis of the complicated interactions between the details of the land biophysics and biogeochemistry, as well as biogeography changes is required in order to better understand and improve model projections

8

10

11

12

13

14

15

16

17

18

19

20

21

22

23

1

2

3

4

5

6

7

4.0 The relationship between model skill and future projections

There are large differences between the different models' projections of future LAI (e.g. Figure 3; Figure S5; Figure 7b). Previous studies have hypothesized that they could reduce the uncertainty in future projections by looking for relationships between model metrics and future projections of climate, and then choosing the models which best match the observations in the current climate (e.g. Cox et al., 2013; Hoffman et al., 2014) or by subsampling models for different regions by their performance (e.g. Steinacher et al., 2010). In this section we explore both approaches, In essence, we are looking for a correlation between current model performance and future projections, this correlation has been used in some studies to argue for a more correct answer, and to reduce the uncertainty in the future projections. In many cases in climate modeling and projections, there is no correlation between model skill in current climate conditions and projections (e.g. Cook and Vizy, 2006), however in some limited cases there is a correlation between metric score and a projection, and one is able to constrain future projections (e.g.

Chris Funk 12/27/2015 4:55 PM

Deleted: statistically

Cornell University 12/7/2015 2:42 PM

Formatted: Font:+Theme Body

Cornell University 12/7/2015 1:55 PM

Deleted:

Cornell University 12/7/2015 1:34 PM

Deleted: Reducing spread in the

Cornell University 12/7/2015 1:48 PM

Deleted: f

Cornell University 12/7/2015 1:33 PM

Deleted: 4

Cornell University 12/7/2015 1:14 PM

Deleted: 2

Cornell University 12/7/2015 1:33 PM

Deleted: 8

Cornell University 12/7/2015 1:48 PM

Deleted: tried to

Cornell University 12/7/2015 1:35 PM

Deleted: use

Cornell University 12/7/2015 1:35 PM

Deleted: to try to reduce the spread in LAI projections at the end of the 21st century (2081-2100)

Chris Funk 12/27/2015 4:57 PM

Deleted: the

Chris Funk 12/27/2015 4:58 PM

Deleted: which

Cornell University 12/7/2015 1:35 PM

Deleted: in order

Deleted: 9

1 Cox et al., 2013; Steinacher et al., 2010). Here we consider whether such a case 2 applies. In doing this type of analysis, we are making an assumption that model skill 3 in the current climate translates into better model projections, which may be a product of real model differences or a statistical error. The advantages and 4 5 disadvantages of using this type of approach are discussed in more detail in Flato et al. (2013). Here we do not advocate that such an approach leads to a better 6 7 projection, but rather simply use this approach to characterize the future model 8 projections. 9 4.1 Evaluation of model LAI 10 11 Several recent studies have evaluated the land models in ESMs using the LAI 12 satellite records (e.g. Anav et al. 2013a; 2013b; Mao et al. 2013; Sitch et al. 2015). 13 Thus we do not repeat those assessments, but rather briefly summarize the results 14 of the comparisons here. 15 Most models tend to overestimate the mean LAI compared to the 16 observations (Figure 8a), and this is true at all latitudes (Figure 8a, Table S2). Cornell University 12/7/2015 1:37 PM Deleted: 9 17 Several models have a large overestimates (>50% too high), including bcc-csm1, Cornell University 12/7/2015 1:37 PM Deleted: 9 18 bcc-csm1-1, BNU-ESM, GFDL-ESM2G, GFDL-ESM2M, MIROC-ESM. The over-19 prediction relative to the satellite data tend to be larger in tropical regions for most 20 models, but the GFDL model estimates are also larger in the high latitudes (Figure Chris Funk 12/27/2015 5:00 PM Deleted: for 21 &a, Table S2). However, the satellite derived LAIs have biases; for example, they Cornell University 1/7/2016 11:57 AM **Deleted:** versions 22 underestimate high LAIs due to being unable to see all the leaf layers in closed Laura Harrison 12/23/2015 3:06 PM **Deleted:** for the GFDL model versions 23 canopies or overestimate LAIs in more arid regions, and thus there may also be an Cornell University 12/7/2015 1:38 PM

Cornell University 12/7/2015 2:15 PM Deleted: or 2 2014 or Forkel et al., 2015, for example). 3 Some models also tend to over predict the strength of the seasonal cycle (e.g. bcc-csm1, BNU-ESM, MIROC-ESM) (Figure 9b; Table S1), where the strength of the 4 5 seasonal cycle is measured by the globally averaged standard deviations of the monthly mean climatology. But the region in which they over-predict the strength of 6 7 the seasonal cycle differs between models. Of course, there is not a strong seasonal 8 cycle in the tropics, where the lowest standard deviations tend to occur (Figure &e; Cornell University 12/7/2015 1:38 PM Deleted: 9 9 Table S2a). Again, because of the difficulties of retrieving accurate LAI from 10 satellites in closed canopies, the observations may underestimate the seasonal cycle 11 in tropical forests. 12 Interannual variability tends to be over-predicted in some of the models (e.g. 13 bcc-csm1, bcc-csm1_1, BNU-ESM, CESM1-BGC, GFDL-ESM2G, GFDL-ESM2M, MIROC-14 ESM, MIROC-ESM_CHEM) (Figure &c, Table S1). For this calculation, the interannual Cornell University 12/7/2015 1:38 PM Deleted: 9 15 variability (IAV) is calculated as the standard deviation of the annual average across 16 multiple years. Generally, the models do a decent job simulating the spatial 17 variability in the annual mean LAI (Figure &d; Table S1), with the correlations being Cornell University 12/7/2015 1:38 PM Deleted: 9 18 strongest in the tropics, and weakest in the high latitudes (Figure 8d; Table S2). This Cornell University 12/7/2015 1:38 PM Deleted: 9 19 is likely partly due to the strength of the LAI differences in tropics and the limitation Cornell University 1/7/2016 11:59 AM Deleted: its 20 of LAI primarily by moisture alone (with low LAI in arid regions and high LAI in Cornell University 1/7/2016 11:58 AM Deleted: the deserts 21 tropical forests). The timing of the seasonal cycle (Figure &e; Table S1) is less well Cornell University 12/7/2015 1:38 PM Deleted: 9 22 simulated in the models, with several models not having an average statistically Fiona A. Lo 12/11/2015 10:20 AM Deleted: o 23 significant correlation (\sim 0.5 for 95% significance for 12 month seasonal cycle) on Fiona A. Lo 12/11/2015 10:20 AM Deleted: a

error in the observational dataset (see discussion in Anav et al. 2013b; Pfeifer et al.

1

Cornell University 12/7/2015 1:15 PM

Deleted: 4

1 the global scale, or in the mid- and high latitudes (e.g. GFDL, MPI-ESM-MR on global 2 scale, GFDL, inmcm4 and MPI-ESM-MR for various regions). 3 Next we explore the observed and modeled relationship between LAI and temperature, and the observed and modeled trend in LAI (e.g. Anav et al., 2013a; 4 5 Anav et al., 2013b; Ichii et al., 2002; Zeng et al., 2013; Mao et al., 2013; Zhu et al., 2013). As previously shown, there are positive relationships between modeled and 6 7 measured LAI and temperature in high latitudes (Figure 6a; Figure S5; e.g. Anav et al. Cornell University 12/7/2015 1:38 PM Deleted: 7 8 2013a; Ichii et al., 2002; Zeng et al. 2013; Zhu et al. 2013). In the tropics (<30°), the Cornell University 12/7/2015 1:14 PM Deleted: 4 9 relationship can be positive or negative but some regions tend towards a negative 10 relationship (Figure S5; Figure 6a). This is consistent with our understanding that Cornell University 12/7/2015 1:14 PM Deleted: 4 11 many places in the tropics are close to the optimal growing temperature already, Cornell University 12/7/2015 1:39 PM Deleted: 7 12 and increases may lead to reduced productivity (Lobell et al., 2011), although this 13 also could be related to moisture stress (Fung et al., 2005). Compared to the 14 observed correlations, most models have too strong of a negative relationship 15 between LAI and temperature in the tropics, and too strong of a positive relationship in the high latitudes (Figure &f, Table S2a-c). In the tropics, the BNU-16 Cornell University 12/7/2015 1:39 PM Deleted: 9 17 ESM model has a weakly positive impact of temperature, while in the high latitudes, 18 especially the CanESM2, HadGEM2-CC, HadGEM2-ES, MPI-ESM-MR models have a 19 much stronger correlation than observed. The model and observations show 20 similarly weak correlations between the temperature and LAI in the mid-latitudes. 21 Some regions show substantial trends over time (1981-2010) in measured

LAI (Figure S7b), especially in high latitudes in the Northern Hemisphere (e.g. Zhu et

al., 2013; Mao et al. 2013). This could be associated with the longer growing season

22

23

due to warming (e.g. Lucht et al., 2002; Zeng et al. 2013). It is also possible that this trend is due to CO_2 fertilization effects (e.g. Friedlingstein and Prentice, 2010). For high latitudes, we find a rank correlation of 0.58 across the models between the CO_2 fertilization factor on land for the Earth system models (called the β -land in Arora et al., 2013, as discussed above) and the average correlation of observed LAI with time, suggesting that there may be a component of carbon dioxide fertilization in the models' temporal trends. These trends are stronger in the models than the observations, which may be related to an overestimate of the fertilization effect.

With regard to LAI interannual variability correlations with temperature or time, there are also strong correlations among temperature, precipitation and time themselves (e.g. IPCC, 2007). Here we do not attempt to differentiate these signals because of the statistical complexity and the shortness of the time record. The shortness of the record considered could also lead to aliasing of the real variability, especially in regions like the Sahel that have strong decadal scale variations (e.g. Loew, 2014). The observational datasets also contain measurement noise, while the model values do not. We expect the measurement noise to reduce the correlations of LAI with the environmental variables in the observations relative to the true values, as seen compared to many models (Figure 3f). Thus, our metrics for interannual variability are likely to be more impacted by uncertainty in the observations than for the annual mean or seasonal cycle, and thus they may be less useful for evaluation of the models, although potentially interesting. For this study, we consider the IAV in the annual mean, but there may be important changes in the seasonal cycle or length of growing season on an interannual time basis, which our

Fiona A. Lo 12/11/2015 10:21 AM

Formatted: Indent: First line: 0.5"

Fiona A. Lo 12/11/2015 10:21 AM

Deleted: that

Cornell University 12/7/2015 1:40 PM

Deleted: 9

1 simple approach does not consider (e.g. Murray-Tortarolo et al. 2013). In addition,

2 the regional or global average of some of these correlations may be difficult to

3 interpret, as not statistically significant (e.g. Figure af), thus making the LAI IAV

4 correlations less helpful.

5

8

9

10

11

13

14

15

18

Figure <u>9</u> summarizes our comparisons of the models <u>with</u> the observations

6 for LAI for the different metrics in Table 2 (Tables S1, S2). In order to show both

7 correlations and model mean biases in the same figure, we have converted the

model-data comparisons into Model Evaluation Values using equation (1) in Section

2.3, where 1 is a perfect model simulation and lower values represent worse model

simulations. Overall, none of the models does a perfect job, and improving

simulation of LAI for all models will be important. In addition, as discussed above,

12 some models perform better in some regions than others. In order to more easily

see how the models compare, we also show the ranking of the different models in

each region (Table 3). For this comparison, we exclude the magnitude and

correlations in the IAV, because the observational estimates for this are more likely

16 to be in error than for the annual mean and seasonal analysis, as discussed above.

17 Thus our overall evaluation of LAI in the models includes the following metrics:

annual mean LAI, spatial correlation of annual mean, standard deviation of seasonal

19 cycle and temporal correlation of the seasonal cycle. In the tropics the top three

20 models are the INMCM4, the IPSL-CM5A-LR and the IPSL-CM5B-LR. For the mid-

21 latitudes the top models are the CanESM2, IPSL-CM5A-MR and the HADGEM2-ES.

For high-latitudes the top models are the BNU-ESM, bcc-csm1 and the MIROC-

23 ESM_CHEM (Table 3; Figure 9).

Cornell University 12/7/2015 1:40 PM

Deleted: 9

Cornell University 12/7/2015 1:40 PM

Deleted: 10

Chris Funk 12/27/2015 5:38 PM

Deleted: to

Cornell University 12/7/2015 1:40 PM

4

5

6

7

8

9

10

12

14

17

18

23

4.2 Future projections constrained by current model performance

3 Across broad regions, we evaluate which metrics are the most useful for potentially

constraining future climate projections by considering how the metric is correlated

with the projections (Figures 8 and 9; Tables S1; S2). We consider 4 regions: the

globe, tropics (latitudes < 30°), mid-latitudes (latitudes between 30 and 60°), and

high latitudes (latitudes >60°). For the first approach, we look for the metrics that

have the highest correlation coefficient to constrain the future estimate of change in

LAI (similar to Cox et al., 2013) (Figure 10a and 10b). Using this approach, we look

for the model metrics (from Table 2) which have the highest correlations with

11 future projections across the models, for each of the regions. If we choose the

models which do the best job with the metrics, this reduces the number of models

included in the projections, and may reduce model spread in projections.

As an example, for the globe, there are two metrics that correlate the highest

with future projections: the average <u>correlation of IAV in LAI with date (i.e. the</u>

16 trend, and the global mean LAI ratio of model to observation. This analysis

suggests that models with the largest relative change in LAI over the last 30 years

(1980-2010) will have the largest change in LAI in the future (Figure 10a). It also

19 suggests that models with higher LAI in the current climate, will have a larger

20 change in the future (Figure 10b). In Fig 10a and 10b, the observation-based

21 estimates are indicated by the gray vertical bar. Notice that the projected change in

22 LAI given by models that match best with the observations differs for different

metrics, and thus it does not allow us to uniquely constrain the future projections

Cornell University 12/7/2015 1:40 PM

Deleted: 9

Cornell University 12/7/2015 1:40 PM

Deleted: 10

Cornell University 12/7/2015 1:41 PM

Deleted: 1

Cornell University 12/7/2015 1:41 PM

Deleted: 1

Cornell University 1/7/2016 12:00 PM

Deleted: IAV

Cornell University 1/7/2016 12:00 PM

Deleted: vs. date correlation

Cornell University 12/7/2015 1:41 PM

Deleted: 1

Cornell University 12/7/2015 1:41 PM

Deleted: 1

Cornell University 12/7/2015 1:41 PM

Deleted: 1

Cornell University 12/7/2015 1:41 PM

(although it does suggest that the highest values are the least likely). There is one
 model with a very large change in LAI in the future (BNU-ESM), We use Spearman
 rank correlations instead of Pearson, correlations, so that these results are largely
 insensitive to the removal of one model.

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

For both the tropical region and in the global analysis, the change with time (LAI IAV correlation with date) and the mean model/observation have the largest correlations (Figure 10c and 10d). Thus models that predict high LAIs in the current climate and/or currently have large trends with time, tend to project higher LAI changes in the future. Again, these two metrics would constrain our future projections to two different LAI values, as they grey lines intersect with the slope at different LAI changes (Figure 10c and 10d). For mid-latitudes, the highest correlation (and only statistically significant correlation) is between the model predicted change in precipitation and LAI (Figure 10e). Thus mid-latitude projections of LAI are difficult to constrain based on model metrics, but are sensitive to modeled changes in precipitation (as seen also in Figure 5). For high latitudes there are three metrics with similar correlation coefficients: the average temporal correlation in the seasonal cycle, the size of the interannual variability and the size of the seasonal cycle in LAI (Figure 10f., 10g, 10h). Unfortunately again, these three metrics suggest a different projected change in LAI when the observed value is used to identify the models that are most realistic (grey line in Figure 10f, 10g and 10h).

Overall, this analysis of multiple metrics suggests that there is no single

metric available that is the most important in all circumstances for improving our

estimates for the changes in LAI. Thus, deduction of a more probable future LAI

Cornell University 1/7/2016 12:01 PM

Deleted:), which can drive much of the correlation

Fiona A. Lo 12/11/2015 9:53 AM

Deleted: simple

Fiona A. Lo 12/11/2015 9:53 AM

Deleted: however,

Cornell University 12/7/2015 1:41 PM

Deleted: 1

Cornell University 12/7/2015 1:41 PM

Deleted: 6

Cornell University 12/7/2015 1:41 PM

Deleted: 1

Cornell University 12/7/2015 1:42 PM

projection is not available to us in this case (as opposed to Cox et al., 2013, where only one metric is presented).

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

The second approach for characterizing the relationship between model simulations in the current climate and future climate projections, and potentially for reducing spread in the future projections follows the ideas of Steinacher et al. (2010). Here for each region, we chose the models that performed the best for several metrics (i.e. using the rankings in Table 3), instead of just one metric at a time (as above). For this study, we chose to use the top half of the models, based on their performance for each region (Table 3), so we include 9 models out of the available 18 models for each region. Using this approach does change the mean future projections, especially for the tropics and high latitudes (Table 4; Figure 7a vs. Zb), and does reduce the spread in the model values in the tropical region, but does not reduce the mean spread in mid-latitudes or high latitudes (Table 4; Figure 7c vs. **7**d). In the tropics, the top models tend to have lower future projections of LAI than the average of all the models (0.07 m^2/m^2 instead of 0.16 m^2/m^2). This is actually consistent with the analysis in Figure 10, since the models with the higher skill (close to grey line) would tend to have lower or middle values of future LAI projections (Figure 10a,b). For the mid-latitudes, there is not as much difference between using all models or the top performing models (Table 6), while for high latitudes, the top models tend to project slightly higher LAI in the future, also consistent with Figure 10 (f,g,h), where the observations tend to suggest higher LAI projections are more consistent for the metrics with the highest correlation.

Fiona A. Lo 12/11/2015 10:25 AM

Deleted: instead of including 18 models,

Cornell University 12/7/2015 1:42 PM

Deleted: 8

Cornell University 12/7/2015 1:42 PM

Deleted: 1

Cornell University 12/7/2015 1:42 PM

Deleted: 1

Cornell University 12/7/2015 1:42 PM

1 The spatial distribution of the change in the future projections using the all 2 models in comparison to, the top models is consistent with the mean over the Fiona A. Lo 12/11/2015 10:26 AM Deleted: vs 3 regions, with the largest change being seen across the tropics, with a reduction in both the mean LAI projection (Figure 7a vs. 7b) as well as the standard deviation 4 Cornell University 12/7/2015 1:43 PM Deleted: 8 5 (Figure 7c vs. 7d). The changes in mid-latitudes and high latitudes from Cornell University 12/7/2015 1:43 PM Deleted: 8 subsampling only the top performing models are not very large in most locations 6 Cornell University 12/7/2015 1:43 PM Deleted: 8 7 (Figure 7a vs. 7b). Only in the tropics is the spread in the models reduced in the Cornell University 12/7/2015 1:43 PM Deleted: 8 future projections (Figure $\frac{7}{4}c$ vs. $\frac{7}{4}d$). The fraction of the time in drought in the 8 Cornell University 12/7/2015 1:43 PM Deleted: 8 9 future is increased in the tropics, if we only consider the top models (Figure Ze vs. Cornell University 12/7/2015 1:43 PM Deleted: 8 10 Cornell University 12/7/2015 1:43 PM Deleted: 8 11 Our results suggest that the better performing models tend to project lower Cornell University 12/7/2015 1:43 PM Deleted: 8 12 LAIs in the future in the tropics in contrast to Cox et al. (2013), which focused on Cornell University 12/7/2015 3:03 PM Deleted: percent 13 carbon-temperature relationships in the Amazon and which showed that Cornell University 12/7/2015 1:43 PM Deleted: 8 14 observational constraints on the models tend to suggest less loss in carbon under Cornell University 12/7/2015 1:43 PM Deleted: 8 15 higher temperatures. However these results may not be inconsistent as they 16 consider different metrics in different regions, and LAI is not necessarily linearly 17 related to vegetative carbon or carbon uptake in the models (see discussion in 18 Section 3.4), suggesting that more analysis of how allocation is parameterized in the Cornell University 11/30/2015 8:42 AM Deleted: 3 19 land carbon models is warranted. 20 Our analysis suggests that using multiple metrics does provide information 21 that allows us in some cases (especially the tropics) to change our mean future 22 projection, and potentially reduce the spread between models predictions. Overall, 23 including only the top models in the tropics project a more pessimistic future, with

1 small increases in mean LAI, and an expansion in the regions at risk for a low LAI, 2 while at high latitudes, it tends to increase the already large increase in mean in LAI. 3 4 **5.0 Summary and Conclusions** 5 LAI is an important term for scaling leaf-level biogeophysical and biogeochemical 6 processes to regional and global areas, and thus it is vital to consider its change in 7 future projections. Here for the first time we consider LAI projections across the 8 CMIP5 models and find that over much of the globe in the future, the models project 9 an increase in mean LAI in the RCP8.5 scenario over the 21st century. Decreases are 10 projected in the limited regions where there is also a projected decrease in mean 11 precipitation; these regions are constrained primarily to the tropics. The change in Fiona A. Lo 12/11/2015 10:27 AM Deleted:, 12 LAI appears to grow with <u>carbon dioxide and</u> temperature increases across regions 13 over the 21st century (Figure 3). Changes in LAI projected in the RCP4.5 are largely Cornell University 12/7/2015 1:43 PM Deleted: 4 14 consistent with changes in RCP8.5, but have a reduced amplitude due to the smaller 15 carbon dioxide and climate forcing. 16 For assessing climate change impacts, we propose that both mean LAI and 17 LAI variability are important in identifying vulnerable regions in future projections. 18 The models project an increased incidence of Low LAI conditions despite higher Chris Funk 12/27/2015 5:43 PM Deleted: 19 mean LAIs, especially in the tropics (Figure 4). While much of the variability in LAI Cornell University 12/7/2015 1:44 PM Deleted: 5 20 is driven by changes in precipitation, projections of lower mean LAI or Low-LAI 21 incidence can identify a slightly different set of vulnerable regions (Figure 5), and Cornell University 12/7/2015 1:44 PM Deleted: 6 22 add to the information that precipitation projections provide.

1 In order to characterize the model projections and evaluate whether we can Cornell University 12/7/2015 1:44 PM Deleted: explore 2 potentially use model skill in the current climate to reduce the spread in the future 3 projections (e.g. Flato et al., 2013), we conducted a brief comparison of the models to available satellite-derived LAI data (Zhu et al., 2013), similar to previous analyses 4 5 (e.g. Anav et al., 2013a; 2013b; Mao et al., 2013; Sitch et al., 2015). Our results support the previous conclusions that the modeled LAI could be improved in many 6 Fiona A. Lo 12/11/2015 10:28 AM Deleted: ly 7 aspects of the mean, seasonal and interannual variability, although difficulties in the 8 observational data may preclude definitive assessment (Figure 8). Cornell University 12/7/2015 1:45 PM Deleted: 9 9 We use two different methods for <u>relating current model skill to model</u> 10 projections, and find that combining multiple metrics to choose better models (e.g. Cornell University 12/7/2015 1:45 PM Deleted: reducing the large spread in 11 similar to Steinacher et al., 2010) seems to work more robustly than simply future projections 12 correlating one metric against future projections (e.g. Cox et al., 2013; Hoffman et al., 13 2014), because the different metrics suggest different future projections (Figure 14 10). Overall, the top-performing models (top half of the models from Table 4) Cornell University 12/7/2015 1:45 PM Deleted: 1 15 suggest smaller future increases in LAI in the tropics, and more regions with more 16 incidences of low-LAI conditions than assessments that include all the models. This 17 approach also reduces the spread among models in the tropics. However, using only 18 the top models did not make a large difference in projections in the mid- and high 19 latitudes (Figure 7). Realize, however, that it is not clear that the models that Cornell University 12/7/2015 1:45 PM Deleted: 8 20 perform best in the current climate have more accurate projections, as discussed in 21 more detail in Flato et al. (2013). Fiona A. Lo 12/11/2015 10:28 AM Deleted: ' 22 Finally, the spread among the models' projections of LAI was correlated with Cornell University 12/7/2015 1:46 PM Deleted: 7 23 the models' projections of precipitation (Figure 6b, and Figure 5). Thus our Cornell University 12/7/2015 1:46 PM Deleted: 6

1 projections of LAI ultimately rest on the ability of models to project future 2 precipitation. <u>Unfortunately</u> in many regions the projected changes in precipitation 3 are not large enough to be statistically significantly outside natural variability (e.g. 4 Tebaldi et al., 2011) and there are discrepancies between climate model and 5 statistical model predictions (e.g. Funk et al., 2014 vs. Tebaldi et al., 2011). In 6 addition to precipitation affecting the future projections of LAI, increasing 7 temperatures are likely to stress systems, even if there is additional rainfall (e.g. 8 Lobell et al., 2011), expanding the regions at risk to increased drought (Figure 5). 9 Because of the importance of LAI for biophysical and biogeochemical interactions, 10 as well as the potential for LAI to be useful to the impacts community, we encourage 11 more analysis of the drivers of LAI variability and changes in the future, as well as 12 improvements in the model mechanisms responsible for the simulation of LAI. 13 14 Acknowledgements 15 We acknowledge the World Climate Research Programme's Working Group on 16 Coupled Modelling, which is responsible for CMIP, and we thank the climate 17 modeling groups (listed in Table 1 of this paper) for producing and making available 18 their model output. For CMIP the U.S. Department of Energy's Program for Climate 19 Model Diagnosis and Intercomparison provides coordinating support and led 20 development of software infrastructure in partnership with the Global Organization 21 for Earth System Science Portals. We acknowledge NSF-0832782 and 1049033 and

assistance from C. Barrett and S. Schlunegger and the anonymous reviewers. We

acknowledge the assistance of the LAI development group for making the LAI 3g

22

23

Fiona A. Lo 12/11/2015 10:28 AM **Deleted:** Yet,

Cornell University 12/7/2015 1:47 PM Deleted: 6

- 1 product available, and the NOAA/OAR/ESRL PSD group for making the GPCP and
- 2 GHCN gridded products available online at http://www.esrl.noaa.gov/psd/.. This
- 3 work was made possible, in part, by support provided by the US Agency for
- 4 International Development (USAID) Agreement No. LAG---A---00---96---90016---00
- 5 through Broadening Access and Strengthening Input Market Systems Collaborative
- 6 Research Support Program (BASIS AMA CRSP). All views, interpretations,
- 7 recommendations, and conclusions expressed in this paper are those of the authors
- 8 and not necessarily those of the supporting or cooperating institutions.

- 1 **Table 1** Model simulations from the Climate Modeling Intercomparison Projection
- 2 (CMIP5) included in this study. All models listed here were available for the RCP8.5
- 3 analysis, while the all models except BNU-ESM and CESM-BGC were available for the
- 4 RCP4.5 analysis.

Model	Land Model	Land	N-	Dynamic	Citation
		Resolution	Cycle	Veg.	
BCC-CSM1	BCC-AVIM1.0	2.8°x2.8°	N	Y	(Wu et al., 2013)
BCC-CSM1-M	BCC-AVIM1.0	1.1°x1.1°	N	Y	(Wu et al., 2013)
	CoLM + BNU-DGVM	2.8°x2.8°	N	Y	(BNU-ESM,
					http://esg.bnu.edu.cn/BNU_ESM_webs/h
BNU-ESM					tmls/index.html)
CanESM2	CLASS2.7+CTEM1	2.8°x2.8°	N	N	(Arora et al., 2011)
CESM1-BGC	CLM4	0.9°x1.2°	Y	N	(Lindsay et al., 2014)
GFDL-ESM2G	LM3	2.5° x 2.5°	N	Y	(Dunne et al., 2013)
	LM3 (uses different	2.5° x 2.5°	N	Y	(Dunne et al., 2013)
GFDL-ESM2M	physical ocean model)				
HadGEM2-CC	JULES+TRIFFID	1.9° x 1.2°	N	Y	(Collins et al., 2011)
	JULES+TRIFFID	1.9° x 1.2°	N	Y	(Collins et al., 2011)
HadGEM2-ES	(includes chemistry)				
INM-CM4	Simple model	2° x 1.5°	N	N	(Volodin et al., 2010)
IPSL-CM5A-LR	ORCHIDEE	3.7° x1.9°	N	N	(Dufresne et al., 2013)
IPSL-CM5A-MR	ORCHIDEE	2.5° x 1.2°	N	N	(Dufresne et al., 2013)
	ORCHIDEE (improved	3.7° x 1.9°	N	N	(Dufresne et al., 2013)
IPSL-CM5B-LR	parameterization)				
MIROC-ESM_	MATSIRO+SEIB-DGVM	2.8° x 2.8°	N	Y	(Watanabe et al., 2011)
	MATSIRO+SEIB-DGVM	2.8° x 2.8°	N	Y	(Watanabe et al., 2011)
MIROC-ESM-CHEM	(adds chemistry)				
MPI-ESM-LR	JSBACH+BETHY	1.9° x 1.9°	N	Y	(Raddatz et al., 2007)
	JSBACH+BETHY	1.9° x 1.9°	N	Y	(Raddatz et al., 2007)
	(ocean model higher				
MPI-ESM-MR	resolution)				
NorESM1-ME	CLM4	2.5° x 1.9°	Y	N	(Bentsen et al., 2013)

 $1 \qquad \text{Table 2: Table of Metrics for LAI comparisons between model and observation used in the following} \\$

tables. More description of these metrics are provided in Section 2.4

2

3 4

Metrics		Description
Mean	Model	Ratio of mean LAI from the model and
	/obs	observations
	Corr.	Spatial correlation of Mean LAI
Std. Dev.	Model	Ratio of seasonal cycle strength: Ratio of standard
Seasonal	/obs	deviation of the climatological monthly mean LAI
		from the model and observations
	Avg.	Avg. Corr. of the temporal evolution of the
	Corr.	climatological seasonal cycle in the model vs.
		observations at each grid box
Std. Dev.	Model	Ratio of IAV strength: ratio of standard deviation of
IAV	/obs	the annual mean LAI from the model and
		observations
IAV LAI	Avg.	Avg. Corr. between LAI and temperature in IAV
vs. T	Corr.	
IAV LAI vs	Avg.	Avg. Corr. between LAI and date in IAV
date	Corr	

Cornell University 11/30/2015 8:42 AM

		T	
	Tropical	Midlatitude	High latitude
bcc-csm1	10	10	2
bcc-csm1-1	9	8	11
BNU-ESM	18	18	1
CanESM2	17	1	16
CESM1-BGC	6	11	17
GFDL-ESM2G	14	15	17
GFDL-ESM2M	16	17	6
HadGEM2-CC	10	5	7
HadGEM2-ES	14	3	11
inmcm4	1	8	13
IPSL-CM5A-LR	2	5	13
IPSL-CM5A-MR	4	1	9
IPSL-CM5B-LR	3	4	5
MIROC-ESM	12	15	4
MIROC- ESM_CHEM-	13	14	2
MPI-ESM-LR	5	7	9
MPI-ESM-MR	7	12	15
NorESM1-ME	8	13	7

Table 4: Mean and standard deviation across models for future projections (LAI change in m^2/m^2) (2081-2100) for all models and for the top half of the models

	Tropics	Mid-latitude	High-latitude
Mean Change (all			
models)	0.16	0.35	0.31
Mean Change (top			
models)	0.07	0.31	0.37
Standard Deviation			
across models (all			
models)	0.35	0.23	0.20
Standard Deviation			
across models (top			
models)	0.25	0.24	0.24

1 Figure captions 2 Figure 1: Mean of all models for the annual mean change in LAI (m²/m²) over time relative to current (1981-2000) for 2011-2030 (a), 2041-2060 (b) and 2081-2100 3 (c) for RCP8.5. 4 5 6 Figure 2: Mean of all models for the annual mean change in LAI over time relative to 7 current (1981-2000), normalized by each model's current (1981-2000) standard 8 deviation at each grid point, for 2011-2030 (a), 2041-2060 (b) and 2081-2100 (c) 9 for RCP8.5. 10 Cornell University 12/7/2015 1:19 PM Deleted: . 11 **Figure 3:** Scatter plot of the change in annual average surface temperature (Ts C) Cornell University 12/7/2015 1:19 PM Deleted: 4 12 (x-axis) against the change in annual average LAI (m2/m2) (y-axis) for the global (a), 13 tropics (b), mid-latitudes (c) and high-latitudes (d). Averages over four time periods 14 are shown; 1981-2000(with 0 changes), 2011-2030, 2041-2060 and 2081-2100, Cornell University 12/7/2015 1:21 PM Deleted: for each RCP 15 connected by a line. The final point (2081-2100) for RCP8.5 is a triangle, The Cornell University 12/7/2015 1:21 PM **Deleted:**, while RCP4.5 is a filled circle 16 temperatures increase in all simulations with time, so increases in the x-axis 17 indicate an increase in time. Note that there are 4 points along each line, and thus if 18 there is no inflection point, the slope of the line is constant across the 21st century. A 19 similar plot including RCP4.5 is included in Figure S4. 20 21 **Figure 4:** Mean of the models for the <u>fraction</u> of the time during which the annual Cornell University 12/7/2015 1:19 PM Deleted: 5 22 mean LAI is considered "Low" (model projected annual mean LAI is less than one Cornell University 12/7/2015 3:03 PM Deleted: percent 23 standard deviation of the current mean at each gridbox) is shown for 2011-2030 (a),

1 2041-2060 (b) and 2081-2100 (c) for RCP85, where the current mean and standard 2 deviation are defined for each grid box for 1981-2000. For the current climate, the 3 <u>fraction</u> of time below one standard deviation will be <u>0.16</u>, which is colored in grey, Cornell University 12/7/2015 3:03 PM Deleted: percentage 4 so all colors represent an increase in low LAI. Cornell University 12/7/2015 3:03 PM Deleted: 16% 5 6 **Figure 5**: Mean of all models for the change in annual mean precipitation for 2081-Cornell University 12/7/2015 1:19 PM Deleted: 6 7 2100 compared to current (1981-2000), normalized by the model standard 8 deviation for RCP8.5 (similar to Figure 2c, but for precipitation) (a). Mean of the 9 models % of the time during which the annual mean precipitation is one standard 10 deviation below current values (similar to figure 5c, but for precipitation) for 2081-11 2100 in RCP8.5 (b). Grid-boxes identified as statistically significantly decreasing in 12 LAI (green) or precipitation (blue) or both (red) (i.e. the blue regions in Figure 2a 13 and Figure 6a contrasted) (c). Grid-boxes identified as having an increase in the 14 amount of time with Low LAI (green) or precipitation (blue) or both (red) (i.e. the 15 blue regions in Figure 5c and Figure 6b contrasted) (c). 16 17 18 Figure 6: Rank correlation across models at every grid box of the mean model Cornell University 12/7/2015 1:19 PM Deleted: 7 19 change in LAI (2081-2100 minus 1981-2000) for RCP8.5 against the model change 20 over the same time period of temperature (a), precipitation (b) and vegetation 21 carbon stock (c). 22

1 Figure 7: Mean of all models for the annual mean change in LAI over time (2081-Cornell University 12/7/2015 1:19 PM Deleted: 8 2 2100) relative to current (1981-2000), normalized by each model's current (1981-3 2000) standard deviation at each grid point (a) for all models (same as Figure 1c) and (b) for the top models, defined as the models performing in the top half (Table 4 5 4) for each region, tropical, mid-latitude or high-latitude. Because different models 6 are included in different regions, there can be discontinuities at the boundaries in 7 Figure 8b (e.g. 30 and 60 degrees latitude). The standard deviation in the mean 8 future projection at 2081-2100 across the models at each grid point are shown for 9 (c) all models and (d) top models. Indication of "Low" LAI is the model mean 10 <u>fraction</u> of the time that LAI is more than one standard deviation below the current Cornell University 12/7/2015 3:03 PM Deleted: percent 11 mean LAI and is shown for (e) all models (same as figure 5c) and (f), top models for 12 the period 2081-2100, where the current mean and standard deviation are defined 13 for each grid box for 1981-2000. For the current climate, the <u>fraction</u> of <u>the</u> time Cornell University 12/7/2015 3:04 PM Deleted: percentage 14 below one standard deviation will be 0.16, which is colored in grey, so all colors Cornell University 12/7/2015 3:04 PM Deleted: 16% 15 represent an increase in drought. 16 17 **Figure 8:** Comparison of model metrics for the LAI comparisons from Table 2 Cornell University 12/7/2015 1:19 PM Deleted: 9 18 across the models, for each region (global, tropical, mid-latitude and high latitude) 19 for a) Mean model/observations, b) seasonal std deviation model/observations, c) 20 IAV standard deviation model/observations, d) spatial correlation of model to 21 observed LAI, e) average temporal correlation for seasonal variability, f) average 22 IAV LAI correlation with temperature (* indicates observed value), g) average IAV 23 LAI correlation with time (* indicates observed value).

3

5

7

8

2 | **Figure 9:** Comparison of model metrics for the annual mean and seasonal metrics

Cornell University 12/7/2015 1:19 PM

Deleted: 10

from Table 2 across the models for a. global, b. tropical, c. mid-latitude and d. high-

4 latitude regions. Similar information is shown in Table S1 and S2, but here

converted to the Model Evaluation Value (equation 1) so that 1 is a perfect model

6 simulation and lower values indicate worse simulations. Models are shown in Table

1, and listed in the figure. Metrics are mean annual (+), spatial correlation of mean

annual (*), seasonal cycle standard deviation(diamond), mean seasonal cycle

9 correlation (triangle) and interannual variability (IAV) standard deviation (square).

1011

12

13

14

15

16

17

18

19

Figure 10: Scatterplot of the metrics with the highest absolute value of the

correlation between the metric and future LAI changes across the globe (LAI IAV

correlated with date (a) and mean LAI model/obs (b)) tropics (<30°) (LAI<u>IAV</u>

correlated with date (c) and mean LAI model/obs (d)), mid-latitudes (between 30°

and 50°) projected change in precipitation (e)) and high-latitudes (>50°) (seasonal

cycle average correlation (f), strength of IAV model/obs (g), and seasonal cycle

strength model/obs (h). The symbols are in the shown colors for each model. The

grey represents the value an ideal model would have based on the observations.

The black line is the line that results from a linear regression of the x and y-axis.

20

21

Cornell University 12/7/2015 1:19 PM

Deleted: 1

Cornell University 12/7/2015 1:23 PM

Deleted: which

Bibliography

- Anav, A., Murray-Tortarolo, G., Friedlingstein, P., Stich, S., Piao, S., and Zhu, Z.:
 Evaluation of Land Surface Models in Reproducing Satellite Derived Leaf
 Area Index over the High Latitude-Northern Hemisphere. Part II: Earth
 System Models, Remote Sensing, 5, 3637-3661, 2013a.
- Anav, A., P. Friedlingstein, M. Kidston, L. Bopp, P. Ciais, P. Cox, C. Jones, M. Jung, R.
 Myneni, and Z. Zhu, Evaluating the land and ocean components of the global carbon cycle in the CMIP5 earth system models, *Journal of Climate*, 26, 6801-6843, 2013b
 - Arora, V. K., Boer, G. J., Friedlingstein, P., Eby, M., Jones, C., Christian, J., Bonan, G., Bopp, L., Brovkin, V., Cadule, P., Hajima, T., Ilyina, T., Lindsay, K., Tjiputra, J. F., and Wu, T.: Carbon-Concentration and carbon-climate feedbacks in CMIP5 earth system models, Journal of Climate, 26, 5289-5314, 2013.
 - Arora, V. K., Scinocca, J., Boer, G. J., Christian, J., Denman, K. L., Flato, G., Kharin, V., Lee, W., and Merryfield, W.: Carbon emission limits required to satisfy future representative concentration pathways of greenhouse gases, Geophysical Research Letters, 38, L05805, doi:10.1029/2010GL046270, 2011.
 - Bentsen, M., Bethke, I., Debernard, J., Iversen, T., Kirkevag, A., Seland, Ø., Drange, H., Roelandt, C., Seierstad, I., Hoose, C., and Kristjansson, J.: The Norwegian Earth System Model, NorESM1-M- Part 1: Description and basic evaluation of the physical climate, Geoscientific Model Development, 6, 687-720, 2013.
 - Bonan, G., P. Lawrence, K. Oleson, S. Levis, M. Jung, M. Reichstein, D. LAwrence, and S. Swenson, Improving canopy processes in the Community Land Model version 4 (CLM4) using global flux fields empirically inferred from FLUXNET data, *Journal of Geophysical Research*, 116(G02014), doi:10.1029/2010JG001593, 2011
 - Bounoua, L., Collatz, G., Los, S. O., Sellers, P., Dazlich, D., Tucker, C., and Randall, D.: Sensitivty of climate to changes in NDVI, Journal of Climate, 13, 2277-2292, 2000.
- Collins, W. J., Bellouin, N., Doutriaux-Boucher, M., Gedney, N., Halloran, P., Hinton, T.,
 Hughes, J., Jones, C., Joshi, M., Liddicoat, S., Martin, G., O'Connor, F., Rae, J. G.,
 Senior, C., Stitch, S., Totterdell, I., Wiltshire, A., and Woodward, S.:
 Development and evaluation of an Earth-system model--HadGEM2,
 Geoscientific Model Development, 4, 997-1062; doi:1010.5194/gmdd-1064-1997-2011, 2011.
 - Cook, K. and Vizy, E.: Coupled model simulations of the West African Monsoon System: Twentieth- and Twenty-First-Century Simulations, Journal of Climate, 19, 3681-3703, 2006.
 - Cox, P., Pearson, D., Booth, B., Friedlingstein, P., Huntingford, C., Jones, C., and Luke, C.: Sensitivity of tropical carbon to climate change constrained by carbon dioxide variability, Nature, 494, 341-344, 2013.
- Cramer, W., Kicklighter, D. W., Bondeau, A., Moore, B., Churkina, G., Nemry, B., Ruimy,
 A., and Schloss, A. e. a.: Comparing global models of terrresrial net primary
 production (NPP): overview and key results, Global Change Biology, 5, 1-15,
 1999.

```
Doherty, R., Stich, S., Smith, B., Lewis, S., and Thornton, P.: Implications of future climate and atmospheric CO2 content for regional biogeochemistry, biogeography and ecosystem services across East Africa, Global Change Biology, 16, 617-640; doi:610.1111/j.1365-2486.2009.01997.x, 2010.
```

- Dufresne, J.-L., Foujols, M.-A., Denvil, S., Caubel, A., Marti, O., Aumont, O., Balkanski, Y., Bekki, S., Bellenger, H., Benshila, R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Cheruy, F., Codron, F., Cozic, A., Cugnet, D., de Noblet, N., Duvel, J.-P., Ethe', C., Fairhead, L., Fichefet, T., Flavoni, S., Freidlingstein, P., Lefebvre, M., Lefrevre, F., Levy, C., Li, Z., Lloyd, J., Lott, F., Madec, G., Mancip, M., Marchand, M., Masson, S., Merurdesoif, Y., Mignot, J., Musat, I., Parouty, S., Polcher, J., Rio, C., Schulz, M., Swingedouw, D., Szopa, S., Talandier, C., Terray, P., Viovy, N., and Vuichard, N.: Climate change projections using the IPSL-CM5 Earth system modl: From CMIP3 to CMIP5, Climate Dynamics, 40, 2123-2165, 2013.
- Dunne, J., John, J., Sheviliakova, E., Stouffer, R. J., Krasting, J., Malyshev, S., Milly, P., Sentman, L., Adcroft, A., Cooke, W., Dunne, K., Harrison, M., Krasting, J., Malyshev, S., Milly, P., Phillips, P., Sentman, L., Samuels, B., Spelman, M., Winton, M., Wittenberg, A., and Zadeh, N.: GFDL's ESM2 global cupoled climate-carbon Earth system models. Part II: Carbon System formation and baseline simulation characteristics, Journal of Climate, 26, 2247-2267, 2013.
- Fan, Y. and Dool, v. d.: A global monthly land surface air temperature analysis for 1948-present, Journal of Geophysical Research, 113, D01103, doi:10.1029/2007JD008470, 2008.
- Field, C., Behrenfeld, M., Randerson, J., Falkowski, P., 1998. Primary Production of the Biosphere: Integrating Terrestrial and Oceanic Components. Science 281, 237; doi:210.1126/science.1181.5374.1237.
- Field, C. B., et al., Technical Summary, in *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by C. B. Field, et al., pp. 35-94, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2014.
- Flato, G., et al., Chapter 9: Evaluation of climate models, in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by T. F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P. M. Midgley, Cambridge University Press, Cambridge, UK and New York, NY, 2013.
- Forkel, M., Carvalhais, N., Verbesselt, J., Mahecha, M., Neigh, C., and Reichstein, M.: Trend Change Detection in NDVI Time Series: Effects of Inter-annual Variability and Methodology, Remote Sensing, 5, 2113-2144:doi: 2110.3390/rs5052113, 2013.
- Forkel, M., Migliavacca, M., Thonicke, K., Reichstein, M., Schaphoff, S., Weber, U., Carvalhais, N., 2015. Codominant water control on global interannual variability and trends in land surface phenology and greennes. Global Change Biology 21, 3414-3435; doi:3410.3111/gcb.12950.

```
Friedlingstein P, G Joel, CB Field, and IY Fung, Toward an allocation scheme for global terrestrial carbon models. Global Change Biology 5:755-770, 1999.
```

- Friedlingstein, P., Cox, P., Betts, R., Bopp, L., Bloh, W. v., Brovkin, V., P. Cadule, Doney, S., Eby, M., I. Fung, G. Bala, J. John, C. Jones, F. Joos, T. Kato, M. Kawamiya, Knorr, W., K. Lindsay, H. D. Mathews, T. Raddatz, P. Rayner, Reick, C., E. Roeckner, Schnitzler, K.-G., Schnurr, R., K. Strassmen, A. J. Weaver, C. Yoshikawa, and Zeng, N.: Climate-carbon cycle feedback analysis, results from the C4MIP Model intercomparison, J. Climate, 19, 3337-3353, 2006.
- Friedlingstein, P., Meinshausen, M., Arora, V. K., Jones, A., Anav, A., Liddicoat, S., and Knutti, R.: Uncertainties in CMIP5 climate projections due to carbon cycle feedbacks, Journal of Climate, 2013. *27*, 511–526, doi:510.1175/JCLI-D-1112-00579.00571.
- Friedlingstein, P. and Prentice, I. C.: Carbon-climate feedbacks: a review of model and observation based estimates, Current Opinion in Environmental Sustainability, 2, 251-257, 2010.
- Fung, I., Doney, S., Lindsay, K., and John, J.: Evolution of carbon sinks in a changing climate, Proceedings of National Academy of Science, 102, 11201-11206, 2005.
- Funk, C. and Brown, M.: Intra-seasonal NDVI change projections in semi-arid Africa, Remote Sensing of Environment, 101, 249-256, 2006.
- Funk, C., Hoell, A., Shukla, S., Bladé, I., Liebmann, B., Roberts, J. B., and Husak, G.: Predicting East African spring droughts using Pacific and Indian Ocean sea surface temperature indices, Hydrology and Earth System Sciences, 11, 3111-3136, 2014.
- Ganzeveld, L., Lelieveld, J., and Roelofs, G.-J.: A dry deposition parameterization for sulfur oxides in a chemistry and general circulation model, J. Geophys. Res., 103, 5679–5694, 1998.
- Gleckler, P., Taylor, K. E., and Doutriaux, C.: Performance metrics for climate models, Journal of Geophysical Research, 113, D06104, doi:10.1029/2007JD008972, 2008.
- Groten, S.: NDVI-crop monitoring and early yield assessment of Burkino Faso, International Journal of Remote Sensing, 14, 1495-1515, 1993.
- Harris, I., Jones, P., Osborn, T., and Lister, D.: Updated high-resolution grids of monthly climatic obsevations--the CRU TS3.10 dataset, International Journal of Climatology, 2013. 623-642, doi:10.1002/joc.3711.
- Hawkins, E. and Sutton, R.: The potential to narrow uncertainty in regional climate predictions, Bulletin of the American Meteorological Society, 2009. 1095-1107, DOI:10.1175/2009BAMS2607.1, 2009.
- Hoffman, F., Randerson, J., Arora, V. K., Bao, Q., Cadule, P., Ji, D., Jones, C., Kawamiya, M., Khatiwala, S., Lindsay, K., Obata, A., Sheviliakova, E., Six, K., Tjiputra, J. F., Volodin, E., and Wu, T.: Causes and implications of persistent atmospheric carbon dioxide biases in Earth System Models, Journal of Geophysical Research-Biogeosciences, 119, 141-162, doi:110.1002/2013JG002381, 2014.

Hurtt, G.C., Chini, L.P., Frolking, S., Betts, R.A., Feddema, J., Fischer, G., Fisk, J.P., Hibbard, K., Houghton, R.A., Janetos, A., Jones, C.D., Kindermann, G., Kinoshita, Cornell University 12/7/2015 12:58 PN Formatted: Indent: Left: 0", Hanging:

- T., Goldewijk, K.K., Riahi, K., Shevliakova, E., Smith, S., Stehfest, E., Thomson, A., Thornton, P., Vuuren, D.P.v., Wang, Y.P., 2011. Harmonization of land-use scenarios for the period 1500-2100: 600 years of global gridded annual land-use transitions, wood harvest, and resulting secondary lands. Climatic Change 109, 117-161, doi:110.1007/s10584-10011-10153-10582.
- Jchii, K., Kawabata, A., and Yamaguchi, Y.: Global correlation analysis for NDVI and climatic variables and NDVI trends: 1982-1990, Int. J. Remote Sens., 23, 3873-3878, 2002.
- IPCC: Summary for Policymakers. In: Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Avery, K. B., Tignor, M. Miller, H. (Eds.), Cambridge University Press, Cambridge, UK and New York, NY, USA, 2007.
- Jones, C., Robertson, E., Arora, V. K., Friedlingstein, P., Sheviliakova, E., Bopp, L., Brovkin, V., Hajima, T., Kato, E., Kawamiya, M., Liddicoat, S., Lindsay, K., REICK, C., Roelandt, C., Segschneider, J., and Tjiputra, J. F.: 21st Century compatible CO2 emissions and airborne fraction simulated by CMIP5 Earth System models under 4 Representative Concentration Pathways., Journal of Climate, 26, 4398-4413; doi:.4310.1175/JCLI-D-4312-00554.00551, 2013.
- Jones, P., Osborn, T., and Briffa, K.: Estimating sampling errors in large-scale temperature averages, Journal of Climate, 10, 2548-2568, 1997.
- Jong, R., Verbesselt, J., Zeileis, A., and Schaepman, M.: Shifts in Global Vegetation Activity Trends, Remote Sensing, 5, 1117-1133; doi: 1110.3390/rs5031117, 2013.
- Kala, J., M. Decker, J.-F. Exbrayat, A. Pitman, C. Carouge, J. Evans, G. Abramowitz, and D. Mocko, Influence of Leaf Area Index prescriptions on simulations of heat, moisture and carbon fluxes, *Journal of Hydrometeorology*, 15, 489-503, 2014.
- Lawrence, D. and Slingo, J.: An annual cycle of vegetatio in a GCM. Part I: implementation and impact on evaporation, Climate Dynamics, 22, 87-105, 2004
- Lawrence, D. M., K.W. Oleson, M.G. Flanner, C.G. Fletcher, P.J. Lawrence, S. Levis, S., Swenson, C., and G.B. Bonan: The CCSM4 land simulation, 1850-2005:

 Assessment of surface climate and new capabilities, J. Climate, 25, 2240-2260, 2012.
- Lin, S.-J., and R. B. Rood (1997), An explicit flux-form semi-Lagrangian shallow-water model on the sphere, *Quarterly Journal of the Royal Meteorological Society*, *123*, 2477-2498.
- Lindsay, K., Bonan, G., Doney, S., Hofffman, F., Lawrence, D., Long, M. C., Mahowald, N., Moore, J. K., Randerson, J. T., and Thornton, P.: Preindustrial and 20th century experiments with the Earth System Model CESM1-(BGC), Journal of Climate, 27, 8981-9005, 2014.
- Lobell, D., Schlenker, W., and Costa-Roberts, J.: Climate trends and global crop production since 1980, Science, 333, 616-620, 2011.
- Loew, A.: Terrestrial satellite records for climate studies: how long is long enough? A test case for the Sahel, Theoretical Applied Climatology, 115, 427-440; doi: 410.1007/s00704-00013-00880-00706, 2014.

Cornell University 12/7/2015 12:57 PM

Deleted:

Lombardozzi, D., Bonan, G., and Nychka, D.: The emerging anthropogenic signal in
 the land-atmosphere carbon cycle, Nature Climate Change, 4, 796-800, DOI:
 710.1038/NCLIMATE2323, 2014.

- Lucht, W., Prentice, I. C., Myneni, R., Stich, S., Friedlingstein, P., Cramer, W., Bousquet, P., Buermann, W., and Smith, B.: Climate control of the high-latitude vegetation greening trend and the Pinatubo effect, Science, 296, 1687-1689, 2002.
- Luo, Y. Q., , J. T. R., Abramowitz, G., Bacour, C., Blyth, E., Carvalhais, N., Ciais, P., Dalmonech, D., Fisher, J. B., Fisher, R., Friedlingstein, P., Hibbard, K., Hoffman, F., Huntzinger, D., Jones, C. D., , C. K., Lawrence, D., Li, D. J., Mahecha, M., Niu, S. L., Norby, R., Piao, S. L., Qi, X., Peylin, P., Prentice, I. C., Riley, W., Reichstein, M., Schwalm, C., Wang, Y. P., Xia, J. Y., Zaehle, S., and Zhou, X. H.: A framework for benchmarking land models, Biogeosciences, 9, 3857-3874, 2012.
- Mao, J., Shin, X., Thornton, P., Hoffman, F., Zhu, Z., and Myneni, R.: Global Latitudinal-Asymetric Vegetation Growth Trends and Their Driving Mechanisms: 1982-2009, Remote Sensing, 5, 1484-1497; doi: 1410.3390/rs5031484, 2013.
- Maxino, C., McAvaney, B., Pitman, A., and Perkins, S.: Ranking the AR4 climate models over the Murray-Darling Basin using simulated maximum temperature, minimum temperature and precipitation, International Journal of Climatology, 28, 1097-1112, 2008.
- Meehl, G., Stocker, T., Collins, W., Frielingstein, P., Gaye, A., Gregory, J. M., Kitoh, A., Knutti, R., Murphy, J., Noda, A., Raper, S., Watterson, I., Weaver, A., and Zhao, Z.-C.: Global Climate Projections. In: Climate Change 2007: The Physical Science Basis, Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate change, Solomon, S., Qin, D., Manning, M. Chen, Z., Marquis, M., Avert, K., Tignor, M., Miller, H. (Ed.), Cambridge University Press, Cambridge, UK, 2007.
- Mitchell, T., Pattern scaling: an examination of the accuracy of he technique for describing future climates, *Climatic Change*, *60*, 217-242, 2003.
- Moss, R., et al., The next generalton of scenarios for climate change research and assessment, *Nature*, 463, 747-756; doi:710.1038/nature08823, 2010.
- Murray-Tortarolo, G., Anav, A., Friedlingstein, P., Stich, S., Piao, S., Zhu, Z., Poulter, B., Zaehle, S., Alhstrom, A., Lomas, M., Levis, S., Viovy, N., and Zeng, N.: Evaluation of Land Surface Models in Reproducing Satellite-Derived LAI over the High-Latitude Northern Hemisphere. Part I: Uncoupled DGVMs., Remote Sensing, 5, 4819-4838; doi:4810.3390/rs5104819, 2013.
- Oleson, K., Lawrence, D., Bonan, G., Drewniak, B., Huang, M., Koven, C., Levis, S., Li, F., Riley, W., Subin, Z., Swensen, S., Thornton, P., Bozbiyik, A., Fisher, R., Kluzek, E., Lamarque, J. F., Lawrence, P., Leung, L. R., Lipscomb, W., Muszala, S., Ricciuto, D., Sacks, W., Sun, Y., Tang, J., and Yang, Z.-L.: Technical Description of version4.5 of the Community Land Model (CLM), NCAR, Boulder, CO, 2013.
- Pfeifer, M., Lefebvre, V., Gonsamo, A., Pellikka, P., Marchant, R., Denu, D., and Platts,
 P.: Validating and linking the GIMSS Leaf Area Index (LAI3g) with
 Environmental Controls in Tropical Africa, Remote Sensing, 6, 1973-1990;
 doi: 1910.3390/rs6031973, 2014.

Qian, T., Dai, A., Trenberth, K., and Oleson, K.: Simluation of global land surface
 conditions from 1948 to 2004. part I: forcing data and evaluations, American
 Meteorological Society, 2006. 953-975, 2006.

- Raddatz, T., Reick, C. H., Knorr, W., Kattge, J., Roeckner, E., Schnur, R., Schnitzler, K.-G., Wetzel, P., and Jungclaus, J.: Will the tropical land biosphere dominate the climate-carbon cycle feedback during the twenty-first century?, Climate Dynamics, 29, 565-574, 2007.
- Ramankutty, N., Evan, A., Monfreda, C., and Foley, J.: Farming the planet: the geographic distribution of global agricultural lands in the year 2000, Global Biogeochemical Cycles, 22, BG1003, 10.1029/2007GB002952, 2008.
- Randerson, J., Hoffman, F., Thornton, P., Mahowald, N., Lindsay, K., Lee, Y.-H., Nevison, C. D., Doney, S., Bonan, G., Stockli, R., Covey, C., Running, S., and Fung, I.: Systematic assessment of terrestrial biogeochemistry in coupled climate-carbon models, Global Change Biology, 15, 2462: doi:2410.1111/j.1365-2486.2009.01912x, 2009.
- Rasch, P., D. Coleman, N. Mahowald, D. Williamson, S.-J. Lin, B. Boville, and P. Hess (2006), Characteristics of atmospheric transport using three numerical formulations for atmospheric dynamics in a single GCM framework, *Journal of Climate*, 19, 2243-2266
- Riahi, K., S. Rao, V. Krey, C. Cho, V. Chikov, G. Fischer, G. Kindermann, N. Nakicenovic, and P. Rafaj, RCP 8.5--A scenario of comparatively high greenhouse gas emissions, *Climatic Change*, 109, 33-57; doi:10.1007/s10584-10011-10149-y, 2011.
- Shao, P., Zeng, X., Sakaguchi, K., Monson, R., and Zeng, X.: Terrestrial carbon cycle:
 climate relations in eight CMIP5 Earth System MOdels, Journal of Climate, 26,
 8744-8764, 2013.
- Sitch, S., Friedlingstein, P., Gruber, N., Jones, S., Murray-Tortarolo, G., Ahlstrom, A.,
 Doney, S., Graven, H., Heinze, C., Huntingford, C., Levis, S., Levy, P., Lomas, M.,
 Poulter, B., Viovy, N., Zaehle, S., Zeng, N., Piao, S., LeQuere, C., Smith, B., Zhu, Z.,
 and Myneni, R.: Recent trends and drivers of regional sources and sinks of
 carbon dioxide, Biogeosciences, 12, 653-679; doi: 610.5194/bg-5112-5653 2015, 2015.
 - Steinacher, M., Joos, F., Frolicher, T. L., Bopp, L., Cadule, P., Cocco, V., Doney, S. C., Lindsay, K., Moore, J. K., Schneider, B., and Segschneider, J.: Projected 21st century decrease in marine productivity: a multi-model analysis, Biogeosciences, 7, 979-1005, 2010.
 - Stocker, T., et al., Technical Summary, in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by T. F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P. M. Midgley, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
 - Taylor, K. E.: Summarizing multiple aspects of model performance in a single diagram, Journal of Geophysical Research, 106, 7183-7192, 2001.
- Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: A summary of the CMIP5 Experimental Design, available at: http://cmip-

pcmdi.llnl.gov/cmip5/docs/Taylor_CMIP5_design.pdf (last access: , April 8, 2015), 2009..

- Tebaldi, C., J. Arblaster, and R. Knutti (2011), Mapping model agreement on future climate projections, *Geophysical Research Letters*, *38*(L23701), doi:10.1029/2011/GL049863.
- Thornton, P., Doney, S., Lindsay, K., Moore, J. K., Mahowald, N., Randerson, J., Fung, I., Lamarque, J. F., Feddema, J., and Lee, Y.-H.: Carbon-nitrogen interactions regular climate-carbon cycle feedbacks: results from an atmosphere-ocean general circulation model, Biogeosciences-discussion, 6, 3303-3354, 2009.
- van Vuuren, D. P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G., Kram, T., Krey, V., Nakicenovic, N., Smith, S., and Rose, S.: The representative concentration pathways: an overview, Climatic Change, 109, 5-31, 2011.
- vanVuuren, D., Elzen, M. G. d., Lucas, P., Eickhout, B., Strengers, B., Ruijven, B. v., Wonink, S., and Houdt, R. v.: Stabilizing greenhouse gas concentrations at low levels: an assessment of reduction strategies and costs, Climatic Change, 81, 119-159; doi:110.1007/s10584-10006-19172-10589, 2007.
- Volodin, E., Dianskii, N., and Gusev, A.: Simulating present day climate with the INMCM4.0 coupled model of the atmospheric and oceanic general circulations, Izv. Ocean. Atmos. Phys., 46, 414-431, 2010.
- Vrieling, A., de Leeuw, J., and Said, M.: Length of Growing Period over Africa: Variability and Trends from 30 years of NDVI Time Series, Remote Sensing, 5, 982-1000: doi: 1010.3390/rs5020982, 2013.
- Wang, W., Ciais, P., Nemani, R., Canadell, J., Piao, S., Stich, S., White, M., Hashimoto, H., Milesi, C., and Myneni, R.: Variations in atmospheric CO2 growth rates coupled with tropical temperature, Proceedings of the National Academy of Science USA, 110, 13061-13066; doi: 13010.11073/pnas.1219683110, 2013.
- Ward, D. S., Mahowald, N., and Kloster, S.: Potential climate forcing of land use and land cover change, Atmospheric Chemistry and Physics, 14, 12701-12724, 2014.
- Watanabe, S., Hajima, T., Sudo, K., Nagashima, T., Takemura, T., Okajima, H., Nozawa, T., Kawase, H., Abe, M., Yokohata, T., Ise, T., Sato, H., Kato, E., Takata, K., Emori, S., and Kamamiya, M.: MIROC-ESM 2010: Model description and basics results of CMIP5-20c3m experiments, Geoscientific Model Development, 4, 845-872, 2011.
 - Wise, M., K. V. Calvin, A. Thomson, L. E. Clarke, B. Bond-Lamberty, R. Sands, S. J. Smith, A. C. Janetos, and J. A. Edmonds, Implications of limiting CO2 concentrations for land use and energy, *Science*, *324*, 1183-1186, 2009.
- Wu, T., Li, W., Ji, J., Xin, X., Li, L., Wang, Z., Zhang, Y., Li, J., Zhang, F., Wei, M., and Shi,
 X.: Global carbon budgets simulated by the Beijing Climate Center Climate
 System Model for the last century, Journal of Geophysical Research, 2013.
 118, 4326-4347, doi:10.1002/jgrd.50320, 2013.
- Zeng, F.-W., Collatz, G., Pinzon, J., and Ivanoff, A.: Evaluating and Quantifying the
 Climate-Driven Interannual Variability in Global Inventory Modeling and
 Mapping Studies (GIMMS) Normalized Difference Vegetation Index at Global
 Scales, Remote Sensing, 5, 3918-3950; doi: 3910.3390/rs508918, 2013.

Zhu, Z., Bi, J., Pan, Y., Ganguly, S., Anav, A., Xu, L., Samanta, A., Piao, S., Nemani, R., and Myneni, R.: Global data sets of vegetation leaf area index (LAI)3g and fraction of photosynthetically active radiation (FPAR)3g derived from global inventory modeling and mapping studies (GIMMS) Normalized difference vegetation index (NDVI3g) for the period 1981 to 2011, Remote Sensing, 5, 927-948, 2013.

Figure 3: Mean of all models for the annual mean change in LAI (m²/m²) over time relative to current climate (1981-2000) for 2081-2100 for RCP4.5. (a) The mean change (similar to Figure 1c), (b) the mean change across models normalized by the model standard deviation for 2081-2100 (similar to Figure 2c); and (c) the mean of all models for the percent of the time during which the annual mean LAI is considered "Low" (model projected annual mean LAI is less than one standard deviation of the current mean at each gridbox) (similar to Figure 5).

Supplemental material:

Sensitivity studies were conducted using the Community Land Model (CLM) (Lawrence et al., 2012), using online model derived meteorology in the CESM-BGC simulations (Lindsay et al., 2014), and using the CLM driven by observed meteorological winds (Qian et al., 2006; Harris et al., 2013). Contrasting these simulations suggest that correlations between LAI and temperature are robust to changing meteorological forcing (Figure S1a and S1b), but that the LAI relationship to precipitation is not robust to changing the input meteological driving data. (Figure S1c and d) This implies that errors in the simulations of the mean and variability in precipitation in the current climate, that are very difficult for ESMs to simulate well (e.g. Flato et al., 2014), are very important for the simulation of IAV in LAI. Thus we do not include analysis of the LAI to precipitation relationships in the evaluation of the current model simulation.

Supplemental tables

Table S1:Evaluation of LAI over globe. Metrics are described in text and Table 2, models in Table 1.

	Mean L	.AI	Seasonal		Std Dev IAV	LAI IAV correlations	
Models	Mod el/ob s	Corr.	Std Dev. Model /obs	Avg. Corr.	Model/ obs	LAI vs. Ts	LAI vs. time.
Obs.						0.11	0.21
bcc-csm1	1.74	0.70	1.28	0.54	1.64	-0.07	0.26
bcc-csm1-1	1.52	0.67	1.27	0.55	1.43	-0.13	0.28
BNU-ESM	2.12	0.56	1.47	0.48	1.79	0.27	0.32

Cornell University 1/7/2016 12:42 PM

Formatted: Font:Bold

CanESM2	1.05	0.66	0.75	0.40	1.15	0.02	0.17
CESM1-BGC	1.49	0.64	0.70	0.48	1.86	0.00	0.24
GFDL-ESM2G	2.27	0.45	0.78	0.18	1.64	-0.06	0.29
GFDL-ESM2M	2.35	0.39	0.78	0.18	1.93	-0.13	0.28
HadGEM2-CC	1.44	0.76	0.58	0.46	0.92	0.15	0.32
HadGEM2-ES	1.52	0.77	0.58	0.46	1.00	0.17	0.35
inmcm4	0.97	0.61	0.93	0.42	0.86	0.05	0.04
IPSL-CM5A-LR	1.44	0.67	0.98	0.49	1.21	0.03	0.08
IPSL-CM5A-MR	1.44	0.68	0.97	0.50	1.21	0.04	0.14
IPSL-CM5B-LR	1.33	0.60	0.95	0.50	1.36	0.02	0.17
MIROC-ESM	1.64	0.44	1.17	0.56	3.23	-0.08	0.11
MIROC-ESM- CHEM	1.62	0.44	1.11	0.53	3.23	-0.05	0.14
MPI-ESM-LR	1.32	0.59	0.83	0.45	0.85	-0.04	0.14
MPI-ESM-MR	1.36	0.60	0.86	0.26	0.85	0.02	0.20
NorESM1-ME	1.61	0.54	0.82	0.44	2.50	-0.05	0.17

Table S2a: Tropical LAI evaluation. As in Table S1, but for tropical region (<30°).

	Mean I		Seasona		Std Dev IAV	LAI IAV corre	
Models	Mod el/ob s	Corr.	Std. Dev. Model /obs	Avg. Corr.	Model/obs	LAI vs. Ts.	LAI vs. Preci p
Obs.						0.02	0.20
bcc-csm1	1.69	0.82	1.79	0.24	2.63	-0.45	0.38
bcc-csm1-1	1.44	0.78	1.66	0.27	2.06	-0.44	0.41
BNU-ESM	2.45	0.63	0.85	0.17	1.25	0.22	-0.04
CanESM2	1.23	0.54	0.74	0.13	1.81	-0.40	0.34
CESM1-BGC	1.72	0.72	0.91	0.38	2.69	-0.29	0.17
GFDL-ESM2G	1.80	0.64	0.96	0.17	2.06	-0.37	0.22
GFDL-ESM2M	1.74	0.63	0.92	0.16	2.50	-0.39	0.24
HadGEM2-CC	1.71	0.81	0.47	0.29	0.88	-0.08	0.28
HadGEM2-ES	1.76	0.81	0.47	0.28	0.94	-0.15	0.33
inmcm4	1.00	0.83	0.83	0.36	0.69	-0.19	0.68
IPSL-CM5A-LR	1.21	0.80	1.09	0.36	1.38	-0.25	0.39
IPSL-CM5A-MR	1.20	0.75	1.09	0.35	1.44	-0.24	0.41
IPSL-CM5B-LR	1.09	0.70	1.02	0.33	1.63	-0.19	0.36
MIROC-ESM	1.61	0.53	0.64	0.35	5.06	-0.37	0.14
MIROC- ESM_CHEM-	1.61	0.53	0.65	0.33	5.00	-0.38	0.15
MPI-ESM-LR	1.41	0.75	1.04	0.15	1.19	-0.51	0.46
MPI-ESM-MR	1.42	0.75	1.06	0.02	1.13	-0.50	0.45
NorESM1-ME	1.73	0.58	0.98	0.32	3.44	-0.34	0.24

Table S2b: Mid-latitude LAI evaluation. As in Table S1, but for mid-latitude region (between 30° and 60°).

	Mean L	-AI	Seasonal		Std Dev IAV	LAI IA\	/ correlations
Models	Mod el/ob s	Corr.	Std. Dev. Model/o bs	Avg. Corr.	Model/obs	LAI vs. Ts.	LAI vs. time.
Obs.						0.11	0.22
bcc-csm1	1.90	0.60	1.33	0.75	1.50	-0.14	0.23
bcc-csm1-1	1.61	0.52	1.34	0.74	1.50	-0.18	0.31
BNU-ESM	2.20	0.48	1.80	0.61	2.80	0.22	0.07
CanESM2	0.90	0.75	0.88	0.56	1.00	0.03	0.12
CESM1-BGC	1.86	0.73	0.82	0.51	2.80	-0.03	0.09
GFDL-ESM2G	1.37	0.52	1.46	0.17	2.20	-0.07	0.22
GFDL-ESM2M	1.36	0.42	1.37	0.20	2.40	-0.11	0.15
HadGEM2-CC	1.20	0.68	0.86	0.56	1.10	0.02	0.19
HadGEM2-ES	1.21	0.69	0.88	0.57	1.20	0.06	0.23
inmcm4	1.28	0.70	0.83	0.33	1.20	0.15	0.07
IPSL-CM5A-LR	1.68	0.71	1.05	0.53	1.90	0.00	0.05
IPSL-CM5A-MR	1.62	0.76	1.00	0.57	2.00	-0.04	0.09
IPSL-CM5B-LR	1.74	0.69	1.05	0.58	1.80	0.07	0.21
MIROC-ESM	1.89	0.44	1.79	0.71	2.80	-0.09	0.08
MIROC- ESM_CHEM-	1.88	0.46	1.75	0.69	2.80	-0.09	0.08
MPI-ESM-LR	1.42	0.41	0.94	0.71	0.70	0.11	0.07
MPI-ESM-MR	1.47	0.38	0.94	0.52	0.70	0.16	0.15
NorESM1-ME	2.19	0.68	1.09	0.50	3.80	-0.01	0.18

Table S2c. High-latitude LAI evaluation. As in Table S1, but for high-latitude region (> 60°).

	Mean LA	Al .	Seasonal		Std Dev IAV	LAI IAV correlat	ions
Models	Model /obs	Corr.	Std. Dev. Model/o bs	Avg. Corr.	Model/ obs	LAI vs. Ts.	LAI vs. time
Obs.						0.20	0.21
bcc-csm1	1.73	0.58	0.69	0.91	0.79	0.36	0.43
bcc-csm1-1	1.61	0.57	0.77	0.91	0.86	0.21	0.41
BNU-ESM	1.45	0.35	1.81	0.90	1.93	0.36	0.48
CanESM2	0.85	0.63	0.69	0.74	0.43	0.42	0.31
CESM1-BGC	0.84	0.48	0.35	0.69	0.50	0.31	0.44
GFDL-ESM2G	3.67	0.22	0.21	0.22	1.00	0.26	0.59
GFDL-ESM2M	4.03	0.21	0.23	0.24	1.14	0.11	0.54
HadGEM2-CC	1.13	0.61	0.39	0.70	0.85	0.48	0.58
HadGEM2-ES	1.31	0.60	0.40	0.72	0.92	0.57	0.66
inmcm4	0.71	0.14	1.21	0.79	0.93	0.21	0.08
IPSL-CM5A-	1.68	0.42	0.83	0.71	0.79	0.32	0.27
IPSL-CM5A-	1.74	0.44	0.82	0.72	0.64	0.36	0.31
IPSL-CM5B-	1.50	0.40	0.77	0.75	0.86	0.19	0.25
MIROC-ESM_	1.52	0.31	1.13	0.83	1.00	0.22	0.24
MIROC-ESM-	1.46	0.28	0.97	0.81	1.07	0.29	0.26
MPI-ESM-LR	1.09	0.49	0.56	0.75	0.36	0.32	0.37
MPI-ESM-MR	1.17	0.51	0.56	0.38	0.36	0.43	0.52
NorESM1-ME	0.95	0.38	0.42	0.66	0.71	0.20	0.19

Supplemental Figures

Figure S1: Rank correlation between model derived LAI and temperature (a and b) and precipitation (c and d) for the CESM-BGC (a and c) and for the CLM-obs (b and d). Both models have the same land model, but the difference is that the CESM-BGC meteorology is from the coupled climate model, while the CLM-obs is driven by datasets constrained by observations (Harris et al. 2013; Qian et al. 2006). **Figure S2**: Observed distributions of leaf area index (LAI) (units of m²/m²) from satellite (Zhu et al. 2013).

Figure S3: Mean of all models for the annual mean change in LAI (m²/m²) over time relative to current climate (1981-2000) for 2081-2100 for RCP4.5. (a) The mean change (similar to Figure 1c), (b) the mean change across models normalized by the model standard deviation for 2081-2100 (similar to Figure 2c); and (c) the mean of all models for the fraction of the time during which the annual mean LAI is considered "Low" (model projected annual mean LAI is less than one standard deviation of the current mean at each gridbox) (similar to Figure 5).

Figure S4: Scatter plot of the change in annual average surface temperature (Ts C) (x-axis) against the change in annual average LAI (m2/m2) (y-axis) for the global (a), tropics (b), mid-latitudes (c) and high-latitudes (d). Averages over four time periods are shown for each RCP: 1981-2000(with 0 changes), 2011-2030, 2041-2060 and 2081-2100, connected by a line. The final point (2081-2100) for RCP8.5 is a triangle, while RCP4.5 is a filled circle. The temperatures increase in all simulations with time, so increases in the x-axis indicate an increase in time. Note that there are 4 points along each line, and thus if there is no inflection point, the slope of the line is

constant across the 21st century. This figure is the same as figure 3, but includes RCP4.5.

Figure S5: Probability density function of the change in LAI between 2081-2100 at each grid box for each model for the globe (a), tropics (<30°) (b), mid-latitudes (between 30° and 50°) (c) and high-latitudes (>50°) (d). The probability density function indicates the fraction of the grid boxes with each LAI value. Probability density functions are smoothed using an Epanechnikov smoothing kernel. Models are show as colored lines, as indicated on legend in figure.

Figure S6: Rank correlation across models at every grid box of the mean model change in LAI (2081-2100 minus 1981-2000) for RCP4.5 against the model change over the same time period of temperature (a) and precipitation (b).

Figure S7: Rank correlation between observational-derived interannual variability in LAI and temperature (a) and year (b) at each grid-box. Correlations above an absolute value of 0.36 are significant at the 95% and are shown in darker colors.

Observations are derived from satellite retrievals (Zhu et al., 2013) for LAI and gridded datsasets GHCN-CAM (Fan and Dool, 2008) for temperature.