

## ***Interactive comment on “Future hydrological extremes: the uncertainty from multiple global climate and global hydrological models” by I. Giuntoli et al.***

**I. Giuntoli et al.**

ixg282@bham.ac.uk

Received and published: 5 February 2015

**Anonymous Referee 1** Received and published: 21 January 2015

**ERRATA CORRIGE / Specific Remarks** Due to Misplaced / Missing text, we provide an updated version of the ‘Specific Remarks’ section of the previous document - AC C17: ‘AC to Referee 1 comment C1: Review of the Manuscript’ from 04 Feb 2015. We apologize for the inconvenience. [In the previous document the 2nd comment was missing, and answer to the 1st comment was misplaced].

**Comment** Page 3 Line 6 mentions multi-model ensembles. However, the manuscript  
C29

continues with single model studies (e.g. Sheffield and Wood 2008). The reader expects a summary of multi-model simulations; maybe move this line to later in the manuscript when multi-model studies are discussed.

**Answer** We will address this in the manuscript by moving (in Page 3) the block of Lines 6-to-10 to Line 20, just before the introduction of the model inter-comparison projects.

**Comment** Page 4 Line 19-20 I believe the sentence should be ... all GCMs for each GCM averaged over all of the GCM or am I mistaken?

**Answer** As described by the authors (see Schewe et al. (2013), caption of Fig. 1, and Dankers et al. (2013), caption of Fig. 1), the GCM variance of the change is computed across all GCMs for each individual GHM, and then averaged over all of the GHMs.

**Comment** Page 7 Line 11-13: This  $SN2 > 1$  is rather arbitrary. Could there be a more statistically proper way to define this  $SN2$  threshold?

**Answer**  $SN2 > 1$  was chosen to facilitate the visualization of the model agreement on the maps. In our case, we wanted to remain consistent with Prudhomme et al. (2014) to facilitate comparisons.

**Comment** In general: Which dataset was used to computer the Koppen-Geiger climate classifications? I can assume they are different between the GCMs and different datasets. Please specify in the manuscript.

**Answer** The dataset we used for the Koppen-Geiger climate regions is based on present-day following the global data classification from Kottek et al. (2006) (a link is provided at the end of Table 1 in the manuscript - [http://koeppen-geiger.vu-wien.ac.at/pdf/kottek\\_et\\_al\\_2006\\_A4.pdf](http://koeppen-geiger.vu-wien.ac.at/pdf/kottek_et_al_2006_A4.pdf)). Therefore, there is no difference between models: all gridded model outputs have been assigned a Koppen-Geiger climate region from the Kottek et al. (2006) dataset. We will clarify this in the manuscript, in particular in the Data and Methods section (Page 8, Line 7). We are aware that region boundaries can change with time (as investigated by e.g. Wanders et al. (2014)); however, in our study we were seeking changes in projections with a present baseline as reference.

## References

Dankers, R., Arnell, N. W., Clark, D. B., Falloon, P. D., Fekete, B. M., Gosling, S. N., Heinke, J., Kim, H., Masaki, Y., Satoh, Y., Stacke, T., Wada, Y., and Wisser, D.: First look at changes in flood hazard in the Inter-Sectoral Impact Model Intercomparison Project ensemble, *P. Natl. Acad. Sci. USA*, 111, 1–5, 10.1073/pnas.1302078110, 2013.

Kottek, M., Grieser, J., Beck, C., Rudolf, B., and Rubel, F.: World Map of the Köppen-Geiger climate classification updated, *Meteorol. Z.*, 15, 259–263, 2006.

Prudhomme, C., Giuntoli, I., Robinson, E. L., Clark, D. B., Arnell, N. W., Dankers, R., Fekete, B. M., Franssen, W., Gerten, D., Gosling, S. N., Hagemann, S., Hannah, D. M., Kim, H., Masaki, Y., Satoh, Y., Stacke, T., Wada, Y., and Wisser, D.: Hydrological droughts in the 21st century, hotspots and uncertainties from a global multimodel ensemble experiment., *P. Natl. Acad. Sci. USA*, 111, 3262–3267, 10.1073/pnas.1222473110, 2014.

Schewe, J., Heinke, J., Gerten, D., Haddeland, I., Arnell, N. W., Clark, D. B., Dankers, R., Eisner, S., Fekete, B. M., Colón-González, F. J., Gosling, S. N., Kim, H., Liu, X., Masaki, Y., Portmann, F. T., Satoh, Y., Stacke, T., Tang, Q., Wada, Y., Wisser, D., Albrecht, T., Frieler, K., Piontek, F., Warszawski, L., and Kabat, P.: Multimodel assessment of water scarcity under climate change, *P. Natl. Acad. Sci. USA*, 111, 3245–3250, 10.1073/pnas.1222460110, 2013.

Wanders, N., Wada, Y., and Van Lanen, H. A. J.: Global hydrological droughts in the 21st century under a changing hydrological regime, *Earth Syst. Dynam. Discuss.*, 5, 649–681, doi:10.5194/esdd-5-649-2014, 2014.

---

Interactive comment on *Earth Syst. Dynam. Discuss.*, 6, 1, 2015.