

Young People's Burden: Requirement of Negative CO₂ Emissions

James Hansen,¹ Makiko Sato,¹ Pushker Kharecha,¹ Karina von Schuckmann,² David J. Beerling,³ Junji Cao,⁴ Shaun Marcott,⁵ Valerie Masson-Delmotte,⁶ Michael J. Prather,⁷ Eelco J. Rohling,^{8,9} Jeremy Shakun,¹⁰ Pete Smith,¹¹ Andrew Lacis,¹² Gary Russell,¹² Reto Ruedy^{12,13}

¹Climate Science, Awareness and Solutions, Columbia University Earth Institute, New York, NY 10115 ²Mercator Ocean, 10 Rue Hermes, 31520 Ramonville St Agne, France ³Leverhulme Centre for Climate Change Mitigation, University of Sheffield, Sheffield S10 2TN, UK ⁴Key Lab of Aerosol Chemistry and Physics, SKLLQG, Institute of Earth Environment, Xi'an 710061, China ⁵Department of Geoscience, 1215 W. Dayton St., Weeks Hall, University of Wisconsin-Madison, Madison, WI 53706 ⁶Institut Pierre Simon Laplace, Laboratoire des Sciences du Climat et de l'Environnement (CEA-CNRS-UVSQ) Université Paris Saclay, Gif-sur-Yvette, France ⁷Earth System Science Department, University of California at Irvine, CA ⁸Research School of Earth Sciences, The Australian National University, Canberra, 2601, Australia ⁹Ocean and Earth Science, University of Southampton, National Oceanography Centre, Southampton, SO14 3ZH, UK ¹⁰Department of Earth and Environmental Sciences, Boston College, Chestnut Hill, MA 02467 ¹¹Institute of Biological and Environmental Sciences, University of Aberdeen, 23 St Machar Drive, AB24 3UU, UK, ¹²NASA Goddard Institute for Space Studies, New York, NY 10025, ¹³SciSpace LLC, 2880 Broadway, New York, NY 10025

E-mail: jehl@columbia.edu

Keywords: climate change, carbon budget, intergenerational justice

Abstract

Global temperature is a fundamental climate metric highly correlated with sea level, which implies that keeping shorelines near their present location requires keeping global temperature within or close to its preindustrial Holocene range. However, global temperature excluding short-term variability now exceeds +1°C relative to the 1880-1920 mean and annual 2016 global temperature was almost +1.3°C. We show that global temperature has risen well out of the Holocene range and Earth is now as warm as it was during the prior (Eemian) interglacial period, when sea level reached 6-9 meters higher than today. Further, Earth is out of energy balance with present atmospheric composition, implying that more warming is in the pipeline, and we show that the growth rate of greenhouse gas climate forcing has accelerated markedly in the past decade. The rapidity of ice sheet and sea level response to global temperature is difficult to predict, but is dependent on the magnitude of warming. Targets for limiting global warming thus, at minimum, should aim to avoid leaving global temperature at Eemian or higher levels for centuries. Such targets now require “negative emissions”, i.e., extraction of CO₂ from the air. If phasedown of fossil fuel emissions begins soon, improved agricultural and forestry practices, including reforestation and steps to improve soil fertility and increase its carbon content, may provide much of the necessary CO₂ extraction. In that case, the magnitude and duration of global temperature excursion above the natural range of the current interglacial (Holocene) could be limited and irreversible climate impacts could be minimized. In contrast, continued high fossil fuel emissions today place a burden on young people to undertake massive technological CO₂ extraction if they are to limit climate change and its consequences. Proposed methods of extraction such as bioenergy with carbon capture and storage (BECCS) or air capture of CO₂ have minimal estimated costs of 89-535 trillion dollars this century and also have large risks and uncertain feasibility. Continued high fossil fuel emissions unarguably sentences young people to either a massive, implausible cleanup or growing deleterious climate impacts or both.




Figure 1. Fossil fuel (and cement manufacture) CO₂ emissions (note log scale) based on Boden et al. (2016) with BP data used to infer 2014-2015 estimates. Europe/Eurasia is Turkey plus the Boden et al. categories Western Europe and Centrally Planned Europe. Asia Pacific is sum of Centrally Planned Asia, Far East and Oceania. Middle East is Boden et al. Middle East less Turkey. Russia is Russian Federation since 1992 and 0.6 of USSR in 1850-1991. Ships/Air is sum of bunker fuels of all nations. Can+Aus is the sum of emissions from Canada and Australia.

1 Introduction

The United Nations 1992 Framework Convention on Climate Change (UNFCCC, 1992) stated its objective as “...stabilization of GHG concentrations in the atmosphere at a level that would prevent dangerous anthropogenic interference with the climate system.” The 15th Conference of the Parties (Copenhagen Accord, 2009) concluded that this objective required a goal to “...reduce global emissions so as to hold the increase of global temperature below 2°C....” The 21st Conference of the Parties (Paris Agreement, 2015), currently ratified by 148 nations, aims to strengthen the global response to the climate change threat by “[h]olding the increase in the global average temperature to well below 2°C above the pre-industrial levels and pursuing efforts to limit the temperature increase to 1.5°C above the pre-industrial levels.”

Global surface temperature has many merits as the principal metric for climate change, but additional metrics, such as atmospheric CO₂ amount and Earth’s energy imbalance, help refine targets for avoiding dangerous human-made climate change. Paleoclimate data and observations of Earth’s present energy imbalance led Hansen et al. (2008, 2013a, 2016) to recommend reducing CO₂ to less than 350 ppm, with the understanding that this target must be adjusted as CO₂ declines and empirical data accumulates. The 350 ppm CO₂ target is moderately stricter than the 1.5°C warming target. The near planetary energy balance anticipated at 350 ppm CO₂ implies a global temperature close to recent values, i.e., about +1°C relative to preindustrial.

We advocate pursuit of this goal within a century to limit the period with global temperature above that of the current interglacial period, the Holocene.¹ Limiting the period and magnitude of temperature excursion above the Holocene range is crucial to avoid strong stimulation of slow feedbacks. Slow feedbacks include ice sheet disintegration and thus sea level rise, which is probably the most threatening climate impact, and release of greenhouse gases (GHGs) via such mechanisms as thawing tundra and loss of soil carbon. Holocene climate stability allowed sea level to be stable for the past several millennia (Kopp et al., 2016) as civilizations developed. But there is now a danger that temperature rises so far above the Holocene range that slow feedbacks are activated to a degree that continuing climate change will be out of humanity’s control. Both the 1.5°C and 350 ppm targets require rapid phasedown of fossil fuel emissions.

¹ By Holocene we refer to the pre-industrial portion of the present interglacial period. As we will show, the rapid warming of the past century has brought temperature above the range in the prior 11,700 years of the Holocene.

Today, global fossil fuel emissions continue at rates that make these targets increasingly improbable (Fig. 1 and Appendix A1). On a per capita historical basis the U.S. is 10 times more accountable than China and 25 times more accountable than India for the increase of atmospheric CO₂ above its preindustrial level (Hansen and Sato, 2016). In response, a lawsuit [Juliana et al. vs United States, 2016, hereafter J et al. vs US, 2016] was filed against the United States asking the U.S. District Court, District of Oregon, to require the U.S. government to produce a plan to rapidly reduce emissions. The suit requests that the plan reduce emissions at the 6%/year rate that Hansen et al. (2013a) estimated as the requirement for lowering atmospheric CO₂ to a level of 350 ppm. At a hearing in Eugene Oregon on 9 March 2016 the United States and three interveners (American Petroleum Institute, National Association of Manufacturers, and the American Fuels and Petrochemical Association) asked the Court to dismiss the case, in part based on the argument that the requested rate of fossil fuel emissions reduction was beyond the court's authority. Magistrate Judge Coffin stated that he found "the remedies aspect of the plaintiff's complaint [to be] *troublesome*", in part because it involves "a separation of powers issue." But he also noted that some of the alleged climate change consequences, if accurate, could be considered "beyond the pale", and he rejected the motion to dismiss the case. Judge Coffin's ruling was certified, as required, by a second judge (Aiken, 2016) on 9 September 2016, and, barring a settlement that would be overseen by the court, the case is expected to proceed to trial in late 2017. It can be anticipated that the plausibility of achieving the emission reductions needed to stabilize climate will be a central issue at the remedy stage of the trial.

Urgency of initiating emissions reductions is well recognized (IPCC, 2013, 2014; Huntingford et al., 2012; Friedlingstein et al., 2014; Rogelj et al., 2016a) and was stressed in the paper (Hansen et al., 2013a) used in support of the lawsuit J et al. vs US (2016). It is also recognized that the goal to keep global warming less than 1.5°C likely requires negative net CO₂ emissions later this century if high global emissions continue in the near-term (Fuss et al., 2014; Anderson, 2015; Rogelj et al., 2015; Sanderson et al., 2016). The Intergovernmental Panel on Climate Change (IPCC) reports (IPCC 2013, 2014) do not address environmental and ecological feasibility and impacts of large-scale CO₂ removal, but recent studies (Smith et al., 2016; Williamson 2016) are taking up this crucial issue and raising the question of whether large-scale negative emissions are even feasible.

Our aim is to contribute to understanding of the required rate of CO₂ emissions reduction via an approach that is transparent to non-scientists. We consider potential drawdown of atmospheric CO₂ by reforestation and afforestation, the potential for improved agricultural practices to store more soil carbon, and potential reductions of non-CO₂ GHGs that could reduce human-made climate forcing². Quantitative examination reveals the merits of these actions to partly offset demands on fossil fuel CO₂ emission phasedown, but also their limitations, thus clarifying the urgency of government actions to rapidly advance the transition to carbon-free energies to meet the climate stabilization targets they have set.

We first describe the status of global temperature change and then summarize the principal climate forcings that drive long term climate change. We show that observed global warming is consistent with knowledge of changing climate forcings, Earth's measured energy imbalance,

² A climate forcing is an imposed change of Earth's energy balance, measured in W/m². For example, Earth absorbs about 240 W/m² of solar energy, so if the sun's brightness increases 1% it is a forcing of +2.4 W/m².

and the canonical estimate of climate sensitivity³, i.e., about 3°C global warming⁴ for doubled atmospheric CO₂. For clarity we make global temperature calculations with our simple climate model, which we show (Appendix A2) has a transient climate sensitivity near the midpoint of the sensitivity of models illustrated in Fig. 10.20a of IPCC (2013). The standard climate sensitivity and climate model do not include effects of “slow” climate feedbacks such as change of ice sheet size. There is increasing evidence that some slow feedbacks can be triggered within decades, so they must be given major consideration in establishing the dangerous level of human-made climate interference. We thus incorporate consideration of slow feedbacks in our analysis and discussion, even though precise specification of their magnitude and time scales is not possible. We present updates of GHG observations and find a notable acceleration during the past decade of the growth rate of GHG climate forcing. For future fossil fuel emissions we consider both the Representative Concentration Pathways (RCP) scenarios used in Climate Model Intercomparison Project 5 (CMIP5) IPCC studies, and simple emission growth rate changes that help evaluate the plausibility of needed emission changes. We use a Green’s function calculation of global temperature with canonical climate sensitivity for each emissions scenario, which yields the amount of CO₂ that must be extracted from the air – effectively the climate debt – to return atmospheric CO₂ to less than 350 ppm or limit global warming to less than 1.5°C above preindustrial levels. We discuss alternative extraction technologies and their estimated costs, and finally we consider the potential alleviation of CO₂ extraction requirements that might be obtained via special efforts to reduce non-CO₂ GHGs.

2 Global Temperature Change

The framing of human-caused climate change by the Paris Agreement uses global mean surface temperature as the metric for assessing dangerous climate change. We have previously argued the merits of additional metrics, especially Earth’s energy imbalance (Hansen et al., 2005; von Schuckmann et al., 2016) and atmospheric CO₂ amount (Hansen et al., 2008). Earth’s energy imbalance integrates over all climate forcings, known and unknown, and informs us where climate is heading, because it is this imbalance that drives continued warming. The CO₂ metric has merit because CO₂ is the dominant control knob on global temperature (Lacis et al., 2010, 2013), including paleo temperature change (cf. Fig. 28 of Hansen et al., 2016). Our present paper uses these alternative metrics to help sharpen determination of the dangerous level of global warming, and to quantify actions that are needed to stabilize climate. We here use global temperature as the principal metric because several reasons of concern are scaled to global warming (O’Neill et al., 2017), including specifically the potential for slow feedbacks such as ice sheet melt and permafrost thaw. The slow feedbacks, whose time scales depend on how strongly the climate system is being forced, will substantially determine the magnitude of climate impacts and affect how difficult the task of stabilizing climate will be.

Quantitative assessment of both ongoing and paleo temperature change is needed to define acceptable limits on human-made interference with climate, with paleo climate especially helpful

³ Climate sensitivity is the response of global average surface temperature to a standard forcing, with the standard forcing commonly taken to be doubled atmospheric CO₂, which is a forcing of about 4 W/m² (Hansen et al., 2005).

⁴ IPCC (2013) finds that 2×CO₂ equilibrium sensitivity is likely in the range 3 ± 1.5°C, as was estimated by Charney et al. (1979). Median sensitivity in recent model inter-comparisons is 3.2°C (Andrews et al., 2012; Vial et al., 2013).




Figure 2. Global surface temperature relative to 1880–1920 based on GISTEMP data (Appendix A3). (a) Annual and 5-year means since 1880, (b) 12- and 132-month running means since 1970. Blue squares in (b) are calendar year (Jan–Dec) means used to construct (a). (b) uses data through April 2017.

for characterizing long-term ice sheet and sea level response to temperature change. Thus we examine the modern period with near-global instrumental temperature data in the context of the current and prior (Holocene and Eemian) interglacial periods, for which less precise proxy-based temperatures have recently emerged. The Holocene, over 11,700 years in duration, had relatively stable climate, prior to the remarkable warming in the past half century. The Eemian, which lasted from about 130,000 to 115,000 years ago, was moderately warmer than the Holocene and experienced sea level rise to heights 6–9 m (20–30 feet) greater than today.

2.1 Modern Temperature

The several analyses of temperature change since 1880 are in close agreement (Hartmann et al., 2013). Thus we can use the current GISTEMP analysis (see Supporting Information), which is updated monthly and available (<http://www.columbia.edu/~mhs119/Temperature/>).

The popular measure of global temperature is the annual-mean global-mean value (Fig. 2a), which is publicized at the end of each year. However, as discussed by Hansen et al. (2010), the 12-month running mean global temperature is more informative and removes monthly “noise” from the record just as well as the calendar year average. For example, the 12-month running mean for the past 67 years (Fig. 2b) defines clearly the super-El Niños of 1997–98 and 2015–16 and the 3-year cooling after the Mount Pinatubo volcanic eruption in the early 1990s.

Global temperature in 2014, 2015 and 2016 reached successive record high levels for the period of instrumental data (Fig. 2). Temperature in the latter two years was partially boosted by the 2015–16 El Niño, but the recent warming is sufficient to remove the illusion of a hiatus of global surface warming after the 1997–98 El Niño (Appendix A4).

The present global warming rate, based on a linear fit for 1970–present (dashed line in Fig. 2b) is $+0.18^{\circ}\text{C}$ per decade⁵. The period since 1970 is the time with high growth rate of GHG climate forcing, which has been maintained at approximately $+0.4 \text{ W/m}^2/\text{decade}$ (see section 6 below)⁶ causing Earth to be substantially out of energy balance (Cheng et al., 2017). The energy

⁵ Extreme endpoints affect linear trends, but if the 2016 temperature is excluded the calculated trend ($0.176^{\circ}\text{C}/\text{decade}$) still rounds to $0.18^{\circ}\text{C}/\text{decade}$.

⁶ As forcing additions from chlorofluorocarbons (CFCs) and CH_4 declined, CO_2 growth increased (section 6).




Figure 3. Estimated average global temperature for (a) last interglacial (Eemian) period (Clark and Huybers, 2009; Turney and Jones, 2010; McKay et al., 2011; Hoffman et al, 2017), (b) centennially-smoothed Holocene (Marcott et al., 2013) temperature and the 11-year mean of modern data (Fig. 2), as anomalies relative to 1880-1920. Vertical downward arrows indicate likely overestimates (see text).

imbalance drives global warming, so unless and until there is substantial change in the rate of added climate forcing we expect the underlying warming to continue at a comparable rate.

Global temperature defined by the linear fit to temperature since 1970 now exceeds 1°C ⁷ relative to the 1880-1920 mean (Fig. 2b), where the 1880-1920 mean provides our best estimate of “preindustrial” temperature (Appendix A5). At the rate of $0.18^{\circ}\text{C}/\text{decade}$ the linear trend line of global temperature will reach $+1.5^{\circ}\text{C}$ in about 2040 and $+2^{\circ}\text{C}$ in the late 2060s. However, the warming rate can accelerate or decelerate, depending on policies that affect GHG emissions, developing climate feedbacks, and other factors discussed below.

2.2 Temperature during current and prior interglacial periods

Holocene temperature has been reconstructed at centennial-scale resolution from 73 globally distributed proxy temperature records by Marcott et al. (2013). This record shows a decline of 0.6°C from early Holocene maximum temperature to a “Little Ice Age” minimum in the early 1800s [that minimum being better defined by higher resolution data of Abram et al. (2016)]. Concatenation of the modern and Holocene temperature records (Fig. 3; Appendix A5) assumes that 1880-1920 mean temperature is 0.1°C warmer than the Little Ice Age minimum (Abram et al., 2016). The early Holocene maximum in the Marcott et al. (2013) data thus reaches $+0.5^{\circ}\text{C}$ relative to the 1880-1920 mean of modern data. The formal 95% confidence bounds to Holocene temperature (Marcott et al., 2013) are $\pm 0.25^{\circ}\text{C}$ (blue shading in Fig. 3b), but total uncertainty is larger. Specifically, Liu et al. (2014) points out a bias effect caused by seasonality in the proxy temperature reconstruction. Correction for this bias will tend to push early Holocene temperatures lower, increasing the gap between today’s temperature and early Holocene temperature (Marcott and Shakun, 2015).

We emphasize that comparisons of current global temperature with the earlier Holocene must bear in mind the centennial smoothing inherent in the Holocene data (Marcott et al. 2013). Thus the temperature in an anomalous single year such as 2016 is not an appropriate comparison. However, the temperature in 2016 based on the 1970-present linear trend (at least 1°C relative to the 1880-1920 mean) does provide a meaningful comparison. The trend line reduces the effect

⁷ It is 1.05°C for linear fit to 132-month running mean, but can vary by a few hundredths of a degree depending on the method chosen to remove short-term variability.

of interannual variability, but the more important point is that Earth's energy imbalance assures that this temperature will continue to rise unless and until the global climate forcing begins to decline. In other words, we know that mean temperature over the next several decades will not be lower than 1°C.

We conclude that the modern trend line of global temperature crossed the early Holocene (smoothed) temperature maximum (+0.5°C) in about 1985. This conclusion is supported by the accelerating rate of sea level rise, which approached 3 mm/year at about that date [Hansen et al. (2016) show a relevant concatenation of measurements in their Fig. 29]. Such a high rate of sea level rise, which is 3 meters per millennium, far exceeds the prior rate of sea level rise in the last six millennia of the Holocene (Lambeck et al., 2014). Note that near stability of sea level in the latter half of the Holocene as global temperature fell about 0.5°C, prior to rapid warming of the Modern Era (Fig. 3), is not inconsistent with that global cooling. Hemispheric solar insolation anomalies in the latter half of the Holocene favored ice sheet growth in the Northern Hemisphere and ice sheet decay in Antarctica (Fig. 27a, Hansen et al., 2016), but the Northern Hemisphere did not become cool enough to reestablish ice sheets on North America or Eurasia. There was a small increase of Greenland ice sheet mass (Larsen et al., 2015), but this was presumably at least balanced by Antarctic ice sheet mass loss (Lambeck et al., 2014).

The important point is that global temperature has risen above the centennially-smoothed Holocene range. Global warming is already having substantial adverse climate impacts (IPCC, 2014), including extreme events (NAS, 2016). There is widespread agreement that 2°C warming would commit the world to multi-meter sea level rise (Levermann et al., 2013; Dutton et al., 2015; Clark et al., 2016). Sea level reached 6-9 m higher than today during the Eemian (Dutton et al., 2015), so it is particularly relevant to know how global mean Eemian temperature compares to the preindustrial level and thus to today.

McKay et al. (2011) estimated peak Eemian annual global ocean SST as $+0.7^{\circ}\text{C} \pm 0.6^{\circ}\text{C}$ relative to late Holocene temperature, while models, as described by Masson-Delmotte et al. (2013), give more confidence to the lower part of that range. Hoffman et al. (2017) report the maximum Eemian annual global SST as $+0.5^{\circ}\text{C} \pm 0.3^{\circ}\text{C}$ relative to 1870-1889, which is $+0.65^{\circ}\text{C}$ relative to 1880-1920. The response of surface air temperature (SAT) over land is twice as large as the SST response to climate forcings in 21st century simulations with models (Collins et al., 2013), in good agreement with observed warming in the industrial era (Appendix A3 this paper, Fig. A3a). The ratio of land SAT change to SST change is reduced only to ~ 1.8 after 1000 years in climate models (Fig. A6, Appendix A6). This implies that, because land covers $\sim 30\%$ of the globe, SST warmings should be multiplied by 1.24-1.3 to estimate global temperature change. Thus the McKay et al. and Hoffman et al. data are equivalent to a global Eemian temperature of just under $+1^{\circ}\text{C}$ relative to the Holocene. Clark and Huybers (2009) and Turney and Jones (2010) estimated global temperature in the Eemian as 1.5-2°C warmer than the Holocene (Fig. 3), but Bakker and Renssen (2014) point out two biases that may cause this range to be an overestimate. Bakker and Renssen (2014) use a suite of models to estimate that the assumption that maximum Eemian temperature was synchronous over the planet overestimates Eemian temperature by $0.4 \pm 0.3^{\circ}\text{C}$ – a feature supported by a lack of synchronicity of warmest conditions in assessments with improved synchronization of records (Govin et al., 2015) – and that they also suggest that a possible seasonal bias of proxy temperature could make the total overestimate as large as $1.1 \pm 0.4^{\circ}\text{C}$. Given uncertainties in the corrections, it becomes a matter of expert judgment. Dutton et al. (2015) conclude that the best estimate for Eemian temperature




Figure 4. Estimated effective climate forcings [update through 2016 of Fig. 28b of Hansen et al. (2005), which are consistent with estimates of Myhre et al. (2013) in the most recent IPCC report (IPCC, 2013). Forcings are based on observations of each gas, except simulated CH₄-induced changes of O₃ and stratospheric H₂O included in the CH₄ forcing. Aerosols and surface albedo change are estimated from historical scenarios of emissions and land use. Oscillatory and intermittent natural forcings (solar irradiance and volcanoes) are excluded. CFCs include not only chlorofluorocarbons, but all Montreal Protocol Trace Gases (MPTGs) and Other Trace Gases (OTGs). Uncertainties (for 5-95% confidence) are 0.6 W/m² for total GHG forcing and 0.9 W/m² for aerosol forcing (Myhre et al., 2013).

is +1°C relative to preindustrial. Consistent with these estimates and the discussion of Masson-Delmotte et al. (2013), we assume that maximum Eemian temperature was +1°C relative to preindustrial with an uncertainty of at least 0.5°C.

These considerations raise the question of whether 2°C, or even 1.5°C, is an appropriate target to protect the well-being of young people and future generations. Indeed, Hansen et al. (2008) concluded “*If humanity wishes to preserve a planet similar to that on which civilization developed and to which life on Earth is adapted, ... CO₂ will need to be reduced ... to at most 350 ppm, but likely less than that.*” And further “*If the present overshoot of the target CO₂ is not brief, there is a possibility of seeding irreversible catastrophic effects.*”

A danger of 1.5°C or 2°C targets is that they are far above the Holocene temperature range. If such temperature levels are allowed to long exist they will spur “slow” amplifying feedbacks (Hansen et al., 2013b; Rohling et al., 2013; Masson-Delmotte et al., 2013), which have potential to run out of humanity’s control. The most threatening slow feedback likely is ice sheet melt and consequent significant sea level rise, as occurred in the Eemian, but there are other risks in pushing the climate system far out of its Holocene range. Methane release from thawing permafrost and methane hydrates is another potential feedback, for example, but the magnitude and time scale of this is unclear (O’Connor et al., 2010; Quicket, 2015).

Here we examine the fossil fuel emission reductions required to restore atmospheric CO₂ to 350 ppm or less, so as to keep global temperature close to the Holocene range, in addition to the canonical 1.5°C and 2°C targets. Quantitative investigation requires consideration of Earth’s energy imbalance, changing climate forcings, and climate sensitivity.

3 Global Climate Forcings and Earth’s Energy Imbalance

The dominant human-caused drivers of climate change are changes of atmospheric GHGs and aerosols (Fig. 4). GHGs absorb Earth’s infrared (heat) radiation, thus serving as a “blanket” warms Earth’s surface by reducing heat radiation to space. Aerosols, fine particles/droplets in the air that cause visible air pollution, both reflect and absorb solar radiation, but reflection of solar energy to space is their dominant effect, so they cause a cooling that partly offsets GHG




Figure 5. Ocean heat uptake in upper 2 km of ocean during 11 years 2005-2015 using analysis method of von Schuckmann and LeTraon (2011). Heat uptake in W/m^2 (0.5 and 0.7) refer to global (ocean + land) area, i.e., it is the contribution of the upper ocean to the heat uptake averaged over the entire planet.

warming. Estimated forcings (Fig. 4), an update of Fig. 28b of Hansen et al. (2005), are similar to those of Myhre et al. (2013) in the most recent IPCC report (IPCC, 2013).⁸

Climate forcings in Fig. 4 are the planetary energy imbalance that would be caused by the preindustrial-to-present change of each atmospheric constituent, if the climate were held fixed at its preindustrial state (Hansen et al., 2005). The CH_4 forcing includes its indirect effects, as increasing atmospheric CH_4 causes tropospheric ozone (O_3) and stratospheric water vapor to increase (Myhre et al., 2013). Uncertainties, discussed by Myhre et al. (2013), are typically 10-15% for GHG forcings. The aerosol forcing uncertainty, described by a probability distribution function (Boucher et al., 2013), is of order 50%. Our estimate of aerosol plus surface albedo forcing (-1.2 W/m^2) differs from the -1.5 W/m^2 of Hansen et al. (2005), as discussed below, but both are within the range of the distribution function of Boucher et al. (2013).

Positive net forcing (Fig. 4) causes Earth to be temporarily out of energy balance, with more energy coming in than going out, which drives slow global warming. Eventually Earth will become hot enough to restore planetary energy balance. However, because of the ocean's great thermal inertia (heat capacity), full atmosphere-ocean response to the forcing requires a long time: atmosphere-ocean models suggest that even after 100 years only 60-75% of the surface warming for a given forcing has occurred, the remaining 25-40% still being "in the pipeline" (Hansen et al., 2011; Collins et al., 2013). Moreover, we outline in the next section that global warming can activate "slow" feedbacks, such as changes of ice sheets or melting of methane hydrates, so the time for the system to reach a fully equilibrated state is even longer.

GHGs have been increasing for more than a century and Earth has partially warmed in response. Earth's energy imbalance is the portion of the forcing that has not yet been responded to. This imbalance thus defines additional global warming that will occur without further change of forcings. Earth's energy imbalance can be measured by monitoring ocean subsurface temperatures, because almost all excess energy coming into the planet goes into the ocean (von Schuckmann et al., 2016). Most of the ocean's heat content change occurs in the upper 2000 m (Levitus et al., 2012), which has been well measured since 2005 when the distribution of Argo

⁸ Our GHG forcings, calculated with formulae of Hansen et al. (2000), yield a CO_2 forcing 6.7% larger than the central IPCC estimate [Table 8.2 of Myhre et al. (2013)] for the CO_2 change from 1750-2011. For all well-mixed (long-lived) GHGs we obtain 3.03 W/m^2 , which is within the IPCC range $2.83 \pm 0.29 \text{ W/m}^2$.

floats achieved good global coverage (von Schuckmann and Le Traon, 2011). Here we update the von Schuckmann and Le Traon analysis with data for 2005-2015 (Fig. 5) finding a decade-average 0.7 W/m^2 heat uptake in the upper 2000 m of the ocean. Addition of the smaller terms raises the imbalance to $+0.75 \pm 0.25 \text{ W/m}^2$ averaged over the solar cycle (Appendix A7).

4 Climate Sensitivity and Feedbacks

Climate sensitivity has been a fundamental issue at least since the 19th century when Tyndall (1861) and Arrhenius (1896) stimulated interest in the effect of CO_2 change on climate.

Evaluation of climate sensitivity involves the full complexity of the climate system, as all components and processes in the system are free to interact on all time scales. Tyndall and Arrhenius recognized some of the most important climate feedbacks on both fast and slow time scales. The amount of water vapor in the air increases with temperature, which is an amplifying feedback because water vapor is a very effective greenhouse gas; this is a “fast” feedback, because water vapor amount in the air adjusts within days to temperature change. The area covered by glaciers and ice sheets is a prime “slow” feedback; it, too, is an amplifying feedback, because the darker surface exposed by melting ice absorbs more sunlight.

Diminishing climate feedbacks also exist. Cloud-cover changes, e.g., can either amplify or reduce climate change (Boucher et al., 2013). Thus it is not inherent that amplifying feedbacks should be dominant, but climate models and empirical data concur that amplifying feedbacks dominate on both short and long time scales, as we will discuss. Amplifying feedbacks lead to large climate change in response to even weak climate forcings such as ice age cycles caused by small perturbations of Earth’s orbit, and still larger climate change occurs on even longer time scales in response to gradual changes in the balance between natural sources and sinks of atmospheric CO_2 (Zachos et al., 2001; Royer et al., 2012; Franks et al., 2014).

4.1 Fast-Feedback Climate Sensitivity

Doubled atmospheric CO_2 , a forcing of $\sim 4 \text{ W/m}^2$, is a standard forcing in studies of climate sensitivity. Charney et al. (1979) concluded that equilibrium sensitivity, i.e., global warming after a time sufficient for the planet to restore energy balance with space, was $3^\circ\text{C} \pm 1.5^\circ\text{C}$ for $2\times\text{CO}_2$ or 0.75°C per W/m^2 forcing. The Charney analysis was based on climate models in which ice sheets and all long-lived GHGs (except for the specified CO_2 doubling) were fixed. The climate sensitivity thus inferred is the “fast-feedback” climate sensitivity. The central value found in a wide range of modern climate models (Flato et al., 2013) remains 3°C for $2\times\text{CO}_2$.

The possibility of unknown unknowns in models would keep the uncertainty in the fast-feedback climate sensitivity high, if it were based on models alone, but as discussed by Rohling et al. (2012a), paleoclimate data allow narrowing of the uncertainty. Ice sheet size and the atmospheric amount of long-lived GHGs (CO_2 , CH_4 , N_2O) under natural conditions change on multi-millennial time scales. These changes are so slow that the climate is in quasi-equilibrium with the changing surface condition and long-lived GHG amounts. Thus these changing boundary conditions, along with knowledge of the associated global temperature change, allow empirical assessment of the fast-feedback climate sensitivity. The central result agrees well with the model-based climate sensitivity estimate of 3°C for $2\times\text{CO}_2$ (Rohling et al., 2012b), with an uncertainty that is arguably 1°C or less (Hansen et al., 2013b).

385 The ocean has great heat capacity (thermal inertia), so it takes decades to centuries for Earth's surface temperature to achieve most of its fast-feedback response to a change of climate forcing (Hansen et al., 1985). Thus Earth has only partly responded to the human-made increase of GHGs in the air today, the planet must be out of energy balance with the planet gaining energy (via reduced heat radiation to space), and more global warming is "in the pipeline."

390 A useful check on understanding of ongoing climate change is provided by the consistency of the net climate forcing (Fig. 4), Earth's energy imbalance, observed global warming, and climate sensitivity. Observed warming since 1880-1920 is 1.05°C^9 based on the linear fit to the 132-month running mean (Fig. 2b), which limits bias from short-term oscillations. Global warming between 1700-1800 and 1880-1920 was $\sim 0.1^{\circ}\text{C}$ (Abram et al., 2016; Hawkins et al., 395 2017; Marcott et al., 2013), so 1750-2015 warming is $\sim 1.15^{\circ}\text{C}$. Taking climate sensitivity as 0.75°C per W/m^2 forcing, global warming of 1.15°C implies that $1.55 \text{ W}/\text{m}^2$ of the total $2.5 \text{ W}/\text{m}^2$ forcing has been "used up" to cause observed warming. Thus $0.95 \text{ W}/\text{m}^2$ forcing should remain to be responded to, i.e., the expected planetary energy imbalance is $0.95 \text{ W}/\text{m}^2$, which is reasonably consistent with the observed $0.75 \pm 0.25 \text{ W}/\text{m}^2$. If we instead take the aerosol + 400 surface albedo forcing as $-1.5 \text{ W}/\text{m}^2$, as estimated by Hansen et al. (2005), the net climate forcing is $2.2 \text{ W}/\text{m}^2$ and the forcing not responded to is $0.65 \text{ W}/\text{m}^2$, which is also within the observational error of Earth's energy imbalance.

4.2 Slow Climate Feedbacks

405 Large glacial-to-interglacial climate oscillations occur on time scales of tens and hundreds of thousands of years, with atmospheric CO_2 amount and the size of ice sheets (and thus sea level) changing almost synchronously on these time scales (Masson-Delmotte et al., 2013). It is readily apparent that these climate cycles are due to small changes in Earth's orbit and the tilt of its spin axis, which alter the geographical and seasonal distribution of sunlight striking Earth. The large 410 climate response is a result of two amplifying feedbacks: (1) atmospheric GHGs (mainly CO_2 but accompanied by CH_4 and N_2O), which increase as Earth warms and decrease as it cools (Ciais et al., 2013), thus amplifying the temperature change, and (2) the size of ice sheets, which shrink as Earth warms and grow as it cools, thus changing the amount of absorbed sunlight in the sense that also amplifies the climate change. For example, 20,000 years ago most of Canada and parts 415 of the United States were covered by an ice sheet, and sea level was about 130 m (~ 400 feet) lower than today. Global warming of $\sim 5^{\circ}\text{C}$ between the last glacial maximum and the Holocene (Masson-Delmotte et al., 2013) is accounted for almost entirely by radiative forcing caused by decrease in ice sheet area and increase of GHGs (Lorius et al., 1990; Hansen et al., 2007).

The glacial-interglacial time scale is set by the time scale of the weak orbital forcings. 420 Before addressing the crucial issue of the inherent time scale of slow feedbacks, we need to say more about the two dominant slow feedbacks, described above as ice sheets and GHGs.

The ice sheet feedback works mainly via the albedo (reflectivity) effect. A shrinking ice sheet exposes darker ground and warming darkens the ice surface by increasing the area and period with wet ice, thus increasing the ice grain size and increasing the surface concentration of

⁹ The IPCC (2013; p. 37 of Technical Summary) estimate of warming for 1880-2012 is 0.85°C [range 0.65 to 1.06°C]. While within that range, our value is higher because (1) use of 4-year longer period, (2) warming in the past few years eliminates the effect on the 1970-present trend from a seeming 1998-2012 warming hiatus, (3) the GISTEMP analysis has greater coverage of the large Arctic warming than the other analyses [Fig. TS.2, p. 39 of IPCC (2013)].

light-absorbing impurities (Tedesco et al., 2016). The ice albedo effect is supplemented by a change of surface albedo in ice-free regions due to vegetation changes. This vegetation albedo effect provides a significant amplification of warming as Earth's temperature increases from its present climate state, because dark forests tend to replace tundra or sparse low-level vegetation in large areas of Eurasia and North America (Lunt et al., 2010).

The GHG feedback on glacial-interglacial time scales is 75-80 percent from CO₂ change; N₂O and CH₄ account for 20-25 percent (Lorius et al., 1990, Hansen et al., 2007, Masson-Delmotte et al., 2013). In simple terms, the ocean and land release more of these gases as the planet becomes warmer. Mechanisms that control GHG release as Earth warms, and GHG drawdown as Earth cools, are complex, including many processes that affect the distribution of carbon, among the ocean, atmosphere, and biosphere (Yu et al., 2016; Ciais et al., 2013 and references therein). Release of carbon from methane hydrates and permafrost contributed to climate change in past warm periods (Zachos et al., 2008; DeConto et al., 2012) and potentially could have a significant effect in the future (O'Connor et al., 2010; Schädel et al., 2016).

Paleoclimate data help assess the possible time scale for ice sheet change. Ice sheet size, judged from sea level, varies almost synchronously with temperature for the temporal resolution available in paleoclimate records, but Grant et al. (2012) find that sea level change lags temperature change by 1-4 centuries. Paleoclimate forcing, however, is both weak and very slow, changing on millennial time scales. Hansen (2005, 2007) argues on heuristic grounds that the much faster and stronger human-made climate forcing projected this century with continued high fossil fuel emissions, equivalent to doubling atmospheric CO₂, would likely lead to substantial ice sheet collapse and multi-meter sea level rise on the time scale of a century. Modeling supports this conclusion, as Pollard et al. (2015) found that addition of hydro-fracturing and cliff failure to their ice sheet model not only increased simulated sea level rise from 2 m to 17 m in response to 2°C ocean warming, it also accelerated the time for multi-meter change from several centuries to several decades. Ice sheet modeling of Applegate et al. (2015) explicitly shows that the time scale for large ice sheet melt decreases dramatically as the magnitude of warming increases. Hansen et al. (2016), based on a combination of climate modeling, paleo data, and modern observations, conclude that continued high GHG emissions would likely cause multi-meter sea level rise within 50-150 years.

The GHG feedback plays a leading role in determining the magnitude of paleoclimate change and there is reason to suspect that it may already be important in modern climate. Rising temperatures increase the rate of CO₂ and CH₄ release from drying soils, thawing permafrost (Schädel et al., 2016; Schuur et al., 2015) and warming continental shelves (Kvenvolden, 1993; Judd et al., 2002), and affect the ocean carbon cycle as noted above. Crowther et al. (2016) synthesize results of 49 field experiments across North America, Europe and Asia, inferring that every 1°C global mean soil surface warming can cause a 30 PgC soil carbon loss and suggesting that continued high fossil fuel emissions might drive 2°C soil warming and a 55 PgC soil carbon loss by 2050. Although this analysis admits large uncertainty, such large soil carbon loss could wreak havoc with efforts to achieve the net soil and biospheric carbon storage that is likely necessary for climate stabilization, as we discuss in subsequent sections.

Recent changes of GHGs result mainly from industrial and agricultural emissions, but they also include any existing climate feedback effects. CO₂ and CH₄ are the largest forcings (Fig. 4), so it is especially important to examine their ongoing changes.




Figure 6. (a) Global CO₂ annual growth based on NOAA data (<http://www.esrl.noaa.gov/gmd/ccgg/trends/>). Dashed curve is for a single station (Mauna Loa). Red curve is monthly global mean relative to the same month of prior year; black curve is 12-month running mean of red curve. (b) CO₂ growth rate is highly correlated with global temperature, the CO₂ change lagging global temperature change by 7-8 months.

5 Observed CO₂ and CH₄ Growth Rates

Annual increase of atmospheric CO₂, averaged over a few years, grew from less than 1 ppm/year 50 years ago to more than 2 ppm/year today (Fig. 6), with global mean CO₂ now exceeding 400 ppm (Betts et al., 2016). Growth of atmospheric CO₂ is about half of fossil fuel CO₂ emissions as discussed in Appendix A8 and illustrated in Fig. A8. The large oscillations of annual growth are correlated with global temperature and with the El Niño/La Niña cycle, as discussed in Appendix A8. Recent global temperature anomalies peaked in February 2016, so as expected the CO₂ growth rate has been declining for the past several months (Fig. 6a).

Atmospheric CH₄ stopped growing between 1998 and 2006, indicating that its sources were nearly in balance with the atmospheric oxidation sink, but growth resumed in the past decade (Fig. 7). CH₄ growth averaged 10 ppb/year in 2014-2016, almost as fast as in the 1980s. Likely reasons for the recent increased growth of CH₄ are discussed in Appendix A8.

The continued growth of atmospheric CO₂ and reaccelerating growth of CH₄ raise important questions related to prospects for stabilizing climate. How consistent with reality are scenarios for phasing down climate forcing when tested by observational data? What changes to industrial and agricultural emissions are required to stabilize climate? We address these issues below.




Figure 7. Global CH₄ from Dlugokencky (2016), NOAA/ESRL (www.esrl.noaa.gov/gmd/ccgg/trends_ch4/). End months for three indicated slopes are January 1984, May 1992, August 2006, and February 2017.




Figure 8. GHG climate forcing annual growth rate (ΔFe) with historical data being 5-year running means, except 2015 is a 3-year mean. (a) includes scenarios used in IPCC AR3 and AR4 reports, and (b) has AR5 scenarios. GHG amounts are from NOAA/ESRL Global Monitoring Division. O_3 changes are not fully included, as they are not well-measured, but its tropospheric changes are partially included via the effective CH_4 forcing. Effective climate forcing (Fe), MPTGs and OTGs are defined in Fig. 4 caption.

6 GHG Climate Forcing Growth Rates and Emission Scenarios

Insight is obtained by comparing the growth rate of GHG climate forcing based on observed GHG amounts with past and present GHG scenarios. We examine forcings of IPCC Special Report on Emissions Scenarios (IPCC SRES, 2000) used in the 2001 AR3 and 2007 AR4 reports (Fig. 8a) and Representative Concentration Pathways scenarios (RCP: Moss et al., 2010; Meinshausen et al., 2011a) used in the 2013 IPCC AR5 report (Fig. 8b). We include the “alternative scenario” of Hansen et al. (2000) in which CO_2 and CH_4 emissions decline such that global temperature stabilizes near the end of the century.¹⁰ We use the same radiation equations for observed GHG amounts and scenarios, so errors in the radiation calculations do not alter the comparison. Equations for GHG forcings are from Table 1 of Hansen et al. (2000) with the CH_4 forcing using an efficacy factor 1.4 to include effects of CH_4 on tropospheric O_3 and stratospheric H_2O (Hansen et al., 2005).

The growth of GHG climate forcing peaked at $\sim 0.05 \text{ W/m}^2/\text{year}$ ($5 \text{ W/m}^2/\text{century}$) in 1978-1988, then falling to a level 10-25% below IPCC SRES (2000) scenarios during the first decade of the 21st century (Fig. 8a). The decline was due to (1) decline of the airborne fraction of CO_2 emissions (Fig. A8), (2) slowdown of CH_4 growth (Fig. 7), and (3) the Montreal Protocol, which initiated phase-out of the production of gases that destroy stratospheric ozone, primarily chlorofluorocarbons (CFCs).

¹⁰This scenario is discussed by Hansen and Sato (2004). CH_4 emissions decline moderately, producing a small negative forcing. CO_2 emissions (not captured and sequestered) are assumed to decline until in 2100 fossil fuel emissions just balance uptake of CO_2 by the ocean and biosphere. CO_2 emissions continue to decline after 2100.




Figure 9. Fossil fuel emission scenarios. (a) Scenarios with simple specified rates of emission increase or decrease. (b) IPCC (2013) RCP scenarios. Note: 1 ppm atmospheric CO₂ is ~2.12 GtC.

The 2013 IPCC RCP scenarios (Fig. 8b) use observed GHG amounts up to 2005 and diverge thereafter, fanning out into an array of potential futures driven by assumptions about energy demand, fossil fuel prices, and climate policy, chosen to be representative of an extensive literature on possible emissions trajectories (Moss et al., 2010; van Vuuren et al., 2011; Meinshausen et al., 2011a,b). Numbers on the RCP scenarios (8.5, 6.0, 4.5 and 2.6) refer to the GHG climate forcing (W/m²) in 2100.

Scenario RCP2.6 has the world moving into negative growth (net contraction) of GHG forcing 25 years from now (Fig. 8b), through rapid reduction of GHG emissions, along with CO₂ capture and storage. Already in 2015 there is a huge gap between reality and RCP2.6. Closing the gap (0.01 W/m²) between actual growth of GHG climate forcing in 2015 and RCP2.6 (Fig. 8b), with CO₂ alone, would require extraction from the atmosphere of more than 0.7 ppm of CO₂ or 1.5 PgC due to the emissions gap of a single year (2015). We discuss the plausibility and estimated costs of scenarios with CO₂ extraction in Section 9.

As a complement to RCP scenarios, we define scenarios with focus on the dominant climate forcing, CO₂, with its changes defined simply by percent annual emission decrease or increase. Below (Sec. 10.1 and Appendix A13) we conclude that efforts to limit non-CO₂ forcings could keep their growth small or even slightly negative, so a focus on long-lived CO₂ is appropriate. Thus for the non-CO₂ GHGs we use RCP6.0, a scenario with small changes of these gases. For CO₂ we consider rates -6%/year, -3%/year, constant emissions, and +2%/year; emissions stop increasing in the +2%/year case when they reach 25 Gt/year (Fig. 9a). Scenarios with decreasing emissions are preceded by constant emissions for 2015-2020, in recognition that some time is required to achieve policy change and implementation. Note similarity of RCP 2.6 with -3%/year, RCP 4.5 with constant emissions, and RCP 8.5 with +2%/year (Fig. 9).

7 Future CO₂ for Assumed Emission Scenarios

We must model Earth's carbon cycle, including ocean uptake of carbon, deforestation, forest regrowth and carbon storage in the soil, for the purpose of simulating future atmospheric CO₂ as a function of the fossil fuel emission scenario. Fortunately, the convenient dynamic-sink pulse-response function version of the well-tested Bern carbon cycle model (Joos et al., 1996) does a good job of approximating more detailed models, and it produces a good match to observed




Figure 10. (a) Atmospheric CO₂ for Fig. 9a emission scenarios. (b) Atmospheric CO₂ including effect of CO₂ extraction that increases linearly after 2020 (after 2015 in +2%/year case).

industrial-era atmospheric CO₂. Thus we use this relatively simple model, described elsewhere (Joos et al., 1996; Kharecha and Hansen, 2008, and references therein), to examine the effect of alternative fossil fuel use scenarios on the growth or decline of atmospheric CO₂. Assumptions about emissions in the historical period are given in Appendix A9.

Figure 10a shows the simulated atmospheric CO₂ for the baseline emission cases (Fig. 9a). These cases do not include active CO₂ removal. Five additional cases including CO₂ removal (Fig. 10b) achieve atmospheric CO₂ targets of either 350 ppm or 450 ppm in 2100, with cumulative removal amounts listed in parentheses (Fig. 10b). The rate of CO₂ extraction in all cases increases linearly from zero in 2010 to the value in 2100 that achieves the atmospheric CO₂ target (350 ppm or 450 ppm). The amount of CO₂ that must be extracted from the system exceeds the difference between the atmospheric amount without extraction and the target amount, e.g., constant CO₂ emissions and no extraction yields 547 ppm for atmospheric CO₂ in 2100, but to achieve a target of 350 ppm the required extraction is 328 ppm, not $547 - 350 = 197$ ppm. The well-known reason (Cao and Caldeira, 2010) is that ocean outgassing increases and vegetation productivity and ocean CO₂ uptake decrease with decreasing atmospheric CO₂, as explored in a wide range of Earth System models (Jones et al., 2016).

8 Simulations of Global Temperature Change

Analysis of future climate change, and policy options to alter that change, must address various uncertainties. One useful way to treat uncertainty is to use results of many models and construct probability distributions (Collins et al., 2013). Such distributions have been used to estimate the remaining budget for fossil fuel emissions for a specified likelihood of staying under a given global warming limit and to compare alternative policies for limiting climate forcing and global warming (Rogelj et al., 2016a,b).

Our aim here is a fundamental, transparent calculation that clarifies how future warming depends on the rate of fossil fuel emissions. We use best estimates for basic uncertain quantities such as climate sensitivity. If these estimates are accurate, actual temperature should have about equal chances of falling higher or lower than the calculated value. Important uncertainties in projections of future climate change include climate sensitivity, the effects of ocean mixing and dynamics on the climate response function discussed below, and aerosol climate forcing. We provide all defining data so that others can easily repeat calculations with alternative choices.

One clarification is important for our present paper. The climate calculations in this section include only fast-feedbacks, which is also true for most climate simulations by the scientific community for IPCC (2013). This is not a limitation for the past, i.e., for the period 1850-present, because we employ measured GHG changes, which include any GHG change due to slow feedbacks. Also we know that ice sheets did not change significantly in size in that period; there may have been some change in Greenland's albedo and expansion of forests in the Northern Hemisphere (Pearson et al., 2013), but those feedbacks so far have only a small global effect. However, this limitation to fast feedbacks may soon become important; it is only in the past few decades that global temperature rose above the prior Holocene range and only in the past two years that it shot far above that range. This limitation must be borne in mind when we consider the role of slow feedbacks in establishing the dangerous level of warming.

We calculate global temperature change T at time t in response to any climate forcing scenario using the Green's function (Hansen, 2008)

$$T(t) = \int_{1850}^t R(t-t') [dF(t')/dt'] dt' + F_v \times R(t - 1850) \quad (1)$$

where $R(t')$ is the product of equilibrium global climate sensitivity and the dimensionless climate response function (percent of equilibrium response), $dF(t')/dt'$ is the annual increment of the net forcing, and F_v is the negative of the average volcanic aerosol forcing during the few centuries preceding 1850. $F_v \times R(t)$ is a small correction term that prevents average volcanic aerosol activity from causing a long-term cooling, i.e., it accounts for the fact that the ocean in 1850 was slightly cooled by prior volcanoes. We take $F_v = 0.3 \text{ W/m}^2$, the average stratospheric aerosol forcing for 1850-2015. The assumed-constant pre-1850 volcanic aerosols caused a constant cooling up to 1850, which gradually decreases to zero after 1850 and is replaced by post-1850 time-dependent volcanic cooling; note that $T(1850) = 0^\circ\text{C}$. We use the "intermediate" response function in Fig. 5 of Hansen et al. (2011), which gives good agreement with Earth's measured energy imbalance. The response function is 0.15, 0.55, 0.75 and 1 at years 1, 10, 100 and 2000 with these values connected linearly in log (year). This defined response function allows our results to be exactly reproduced, or altered with alternative choices for climate forcings, climate sensitivity and response function. Forcings that we use are tabulated in Appendix A10.

We use equilibrium fast-feedback climate sensitivity 0.75°C per W/m^2 (3°C for $2\times\text{CO}_2$). This is consistent with climate models (Collins et al., 2013; Flato et al., 2013) and paleoclimate evidence (Rohling et al., 2012a; Masson-Delmotte et al., 2013; Bindoff and Stott, 2013). We use RCP6.0 for the non- CO_2 GHGs.

We take tropospheric aerosol plus surface albedo forcing as -1.2 W/m^2 in 2015, presuming the aerosol and albedo contributions to be -1 W/m^2 and -0.2 W/m^2 , respectively. We assume a small increase this century as global population rises and increasing aerosol emission controls in emerging economies tend to be offset by increasing development elsewhere, so aerosol + surface forcing is -1.5 W/m^2 in 2100. The temporal shape of the historic aerosol forcing curve (Table A10) is from Hansen et al. (2011), which in turn was based on the Novakov et al. (2003) analysis of how aerosol emissions have changed with technology change.




Figure 11. Climate forcings used in our climate simulations; F_e is effective forcing, as discussed in connection with Fig. 4. (a) Future GHG forcing uses four alternative fossil fuel emission growth rates. (b) GHG forcings are altered based on CO_2 extractions of Fig. 10.

Historic stratospheric aerosol data (Table A10, annual version), an update of Sato et al. (1993), include moderate 21st century aerosol amounts (Bourassa et al., 2012). Future aerosols, for realistic variability, include three volcanic eruptions in the rest of this century with properties of the historic Agung, El Chichon and Pinatubo eruptions, plus a background stratospheric aerosol forcing -0.1 W/m^2 . This leads to mean stratospheric aerosol climate forcing -0.3 W/m^2 for remainder of the 21st century, similar to the mean stratospheric aerosol forcing for 1850-2015 (Table A10). Reconstruction of historical solar forcing (Coddington et al., 2015; Kopp et al., 2016), based on data in Fig. A11, is extended with an 11-year cycle.

Individual and net climate forcings for the several fossil fuel emission reduction rates are shown in Fig. 11a,c. Scenarios with linearly growing CO_2 extraction at rates required to yield 350 or 450 ppm airborne CO_2 in 2100 are in Fig. 11b,d. These forcings and the assumed climate response function define expected global temperature for the entire industrial era considered here (Fig. 12). We extended the global temperature calculations from 2100 to 2200 by continuing the %/year change of CO_2 emissions. In the cases with CO_2 extraction we kept the GHG climate forcing fixed in the 22nd century, which meant that large CO_2 extraction continued in cases with continuing high emissions, e.g., the case with constant emissions that required extraction of 695 PgC during 2020-2100 required further extraction of $\sim 900 \text{ PgC}$ during 2100-2200. Even the cases with annual emission reductions $-6\%/year$ and $-3\%/year$ required small extractions to compensate for back-flux of CO_2 from the ocean that accumulated there historically.




Figure 12. Simulated global temperature for Fig. 11 forcings. Observations as in Fig. 2. Temperature zero-point is the 1880-1920 mean temperature for both observations and model. Gray area is 2σ (95% confidence) range for centennially-smoothed Holocene maximum, but there is further uncertainty about the magnitude of the Holocene maximum, as noted in the text and discussed by Liu et al. (2014).

A stark summary of alternative futures emerges from Fig. 12a. If emissions grow 2%/year, modestly slower than the 2.6%/year growth of 2000-2015, warming reaches $\sim 4^\circ\text{C}$ by 2100. Warming is about 2°C if emissions are constant until 2100. Furthermore, both scenarios launch Earth onto a course of more dramatic change well beyond the initial 2-3°C global warming, because: (1) warming continues beyond 2100 as the planet is still far from equilibrium with the climate forcing, and (2) warming of 2-3°C would unleash strong slow feedbacks, including melting of ice sheets and increases of GHGs.

The most important conclusion from Fig. 12a is the proximity of results for the cases with emission reductions of 6%/year and 3%/year. Although Hansen et al. (2013a) called for emission reduction of 6%/year to restore CO₂ to 350 ppm by 2100, that rate of reduction may have been regarded as implausibly steep by a federal court in 2012 when it declined to decide whether the U.S. was violating the public trust by causing or contributing to dangerous climate change (*Alec L v. Jackson*, 2012). Such a concern is less pressing for emission reductions of 3% per year. Note that reducing global emissions at a rate of 3%/year (or more steeply) maintains global warming at less than 1.5°C above preindustrial temperature.

However, end-of-century temperature still rises 0.5°C or more above the prior Holocene maximum with consequences for slow feedbacks that are difficult to foresee. Desire to minimize sea level rise spurs the need to get global temperature back into the Holocene range. That goal preferably should be achieved on the time scale of a century or less, because paleoclimate evidence indicates that the response time of sea level to climate change is 1-4 centuries (Grant et al., 2012, 2014) for natural climate change, and if anything the response should be faster to a stronger, more rapid human-made climate forcing. The scenarios that reduce CO₂ to 350 ppm succeed in getting temperature back close to the Holocene maximum by 2100 (Fig. 12b), but they require extractions of atmospheric CO₂ that range from 153 PgC in the scenario with 6%/year emission reductions to 1630 PgC in the scenario with +2%/year emission growth.

Scenarios ranging from constant emissions to +2%/year emissions growth can be made to yield 450 ppm in 2100 via extraction of 339-1270 PgC from the atmosphere (Fig. 10b). However, these scenarios still yield warming more than 1.5°C above the preindustrial level

(more than 1°C above the early Holocene maximum). Consequences of such warming and the plausibility of extracting such huge amounts of atmospheric CO₂ are considered below.

9 CO₂ Extraction: Estimated Cost and Alternatives

Extraction of CO₂ from the air, also called negative emissions or carbon dioxide removal (CDR), is required if large, long-term excursion of global temperature above its Holocene range is to be averted, as shown above. In estimating the cost and plausibility of CO₂ extraction we distinguish between (1) carbon extracted from the air by improved agricultural and forestry practices, and (2) additional “technological extraction” by intensive negative emission technologies.

We assume that improved practices will aim at optimizing agricultural and forest carbon uptake via relatively natural approaches, compatible with the land delivering a range of ecosystem services (Smith 2016; Smith et al., 2016). In contrast, proposed technological extraction and storage of CO₂ generally does not have co-benefits and remains unproven at relevant scales (NAS, 2015a). Improved practices have local benefits in agricultural yields and forest products and services (Smith et al., 2016), which may help minimize net costs. The Intended Nationally Determined Contributions (INDCs) submitted by 189 countries include carbon drawdown through land use plans (United Nations, 2016) with aggregate removal rate of ~2 PgCO₂/year (~0.55 PgC/year) after 2020. These targets are not the maximum possible drawdown, as they are only about a third of amounts Smith (2016) estimated as “realistic”.

Developed countries recognize a financial obligation to less developed countries that have done little to cause climate change (Paris Agreement 2015)¹¹. We suggest that at least part of developed country support should be channeled through agricultural and forestry programs, with continual evaluation and adjustment to reward and encourage progress (Bustamante et al., 2014). Efforts to minimize non-CO₂ GHGs can be included in the improved practices program.

Here, we do not estimate the cost of CO₂ extraction obtained via the “improved agricultural and forestry practices¹²,” because that would be difficult given the range of activities it is likely to entail, and because it is not necessary for reaching the conclusion that total CO₂ extraction costs will be high due to the remaining requirements for technological extraction. However, we do estimate the potential magnitude of CO₂ extraction that might be achievable via such improved practices, as that is needed to quantify the required amount of “technological extraction” of CO₂. Finally, we compare costs of extraction with estimated costs of mitigation measures that could limit the magnitude of required extraction, while admitting that there is large uncertainty in both extraction and mitigation cost estimates.

¹¹ Another conceivable source of financial support for CO₂ drawdown might be legal settlements with fossil fuel companies, analogous to penalties that courts have imposed on tobacco companies, but with the funds directed to the international “improved practices” programs.

¹² A comment is in order about the relation of “improved agricultural and forestry practices” with an increased role of biofuels in climate mitigation. Agriculture, forestry and other land use has potential for important contributions to climate change mitigation (Smith et al., 2014). However, first-generation biofuel production and use (which is usually based on edible portions of feedstocks, such as starch) is not inherently carbon neutral, indeed it is likely carbon-positive, as has been illustrated in specific quantitative analyses for corn ethanol in the United States (Searchinger et al., 2008; DeCicco et al., 2016). The need for caution regarding the role of biofuels in climate mitigation is discussed by Smith et al. (2014).

9.1 Estimated Cost of CO₂ Extraction

Hansen et al. (2013) suggested a goal of 100 PgC extraction in the 21st century, which would be almost as large as estimated net emissions from historic deforestation and land use (Ciais et al., 2013). Hansen et al. (2013a) assumed that 100 PgC was about as much as could be achieved via relatively natural reforestation and afforestation (Canadell and Raupach, 2008) and improved agricultural practices that increase soil carbon (Smith, 2016).

Here we first reexamine whether a concerted global effort on carbon storage in forests and soil might have potential to provide a carbon sink substantially larger than 100 PgC this century. Smith et al. (2016) estimate that reforestation and afforestation together have carbon storage potential of about 1.1 PgC/year. However, as forests mature, their uptake of atmospheric carbon decreases (termed “sink saturation”), thereby limiting CO₂ drawdown. Taking 50 years as the average time for tropical, temperate and boreal trees to experience sink saturation yields 55 PgC as the potential storage in forests this century.

Smith (2016) shows that soil carbon sequestration and soil amendment with biochar compare favorably with other negative emission technologies with less impact on land use, water use, nutrients, surface albedo, and energy requirements, but understanding of and literature on biochar are limited (NAS, 2015a). Smith estimates that soil carbon sequestration has potential to store 0.7 PgC/year. However, as with carbon storage in forest, there is a saturation effect. A commonly used 20-year saturation time (IPCC, 2006) would yield 14 PgC soil carbon storage, while an optimistic 50-year saturation time would yield 35 PgC. Use of biochar to improve soil fertility provides additional carbon storage of up to 0.7-1.8 PgC/year (Woolf et al., 2010; Smith 2016). Larger industrial-scale biochar carbon storage is conceivable, but belongs in the category of intensive negative emission technologies, discussed below, whose environmental impacts and costs require scrutiny. We conclude that 100 PgC is an appropriate ambitious estimate for potential carbon extraction via a concerted global-scale effort to improve agricultural and forestry practices with carbon drawdown as a prime objective.

Intensive negative emission technologies that could yield greater CO₂ extraction include (1) burning of biofuels, most commonly at power plants, with capture and sequestration of resulting CO₂ (Creutzig et al., 2015), and (2) direct air capture of CO₂ and sequestration (Keith, 2009; NAS, 2015a), and (3) grinding and spreading of minerals such as olivine to enhance geological weathering (Taylor et al., 2016). However, energy, land and water requirements of these technologies impose economic and biophysical limits on CO₂ extraction (Smith et al., 2016).

The popular concept of bioenergy with carbon capture and storage (BECCS) requires large areas, high fertilizer and water use, and may compete with other vital land use such as agriculture (Smith, 2016). Costs estimates are ~\$150-350/tC for crop-based BECCS (Smith et al., 2016).

Direct air capture has more limited area and water needs than BECCS and no fertilizer requirement, but it has high energy use, has not been demonstrated at scale, and cost estimates exceed those of BECCS (Socolow et al., 2011; Smith et al., 2016). Keith et al. (2006) have argued that, with strong research and development support and industrial-scale pilot projects sustained over decades, it may be possible to achieve costs ~\$200/tC, thus comparable to BECCS costs; however other assessments are higher, reaching \$1400-3700/tC (NAS, 2015a).

Enhanced weathering via soil amendment with crushed silicate rock is a candidate negative emission technology that also limits coastal ocean acidification as chemical products liberated by weathering increase land-ocean alkalinity flux (Kohler et al., 2010; Taylor et al., 2016). If two-thirds of global croplands were amended with basalt dust, as much as 1-3 PgC/year might be extracted, depending on application rate (Taylor et al., 2016), but energy costs of mining,

770 grinding and spreading likely reduce this by 10-25% (Moosdorf et al., 2014). Such large-scale enhanced weathering is speculative, but potential co-benefits for temperate and tropical agroecosystems could affect its practicality, and may put some enhanced weathering into the category of improved agricultural and forestry practices. Benefits include crop fertilization that increases yield and reduces use and cost of other fertilizers, increasing crop protection from 775 insect herbivores and pathogens thus decreasing pesticide use and cost, neutralizing soil acidification to improve yield, and suppression of GHG (N₂O and CO₂) emissions from soils (Edwards et al., 2017; Kantola et al., 2017). Against these benefits, we note potential negative impacts of air and water pollution caused by the mining, including downstream environmental consequences if silicates are washed into rivers and the ocean, causing increased turbidity, 780 sedimentation, and pH, with unknown impacts on biodiversity (Edwards et al., 2017). Cost of enhanced weathering might be reduced by deployment with reforestation and afforestation and with crops used for BECCS; this could significantly enhance the combined carbon sequestration potential of these methods.

For cost estimates, we first consider restoration of airborne CO₂ to 350 ppm in 2100 (Fig. 785 10b), which would keep global warming below 1.5°C and bring global temperature back close to the Holocene maximum by end-of-century (Fig. 12b). This scenario keeps the temperature excursion above the Holocene level small enough and brief enough that it has the best chance of avoiding ice sheet instabilities and multi-meter sea level rise (Hansen et al., 2016). If fossil fuel emission phasedown of 6%/year had begun in 2013, as proposed by Hansen et al. (2013a), this 790 scenario would have been achieved via the hypothesized 100 PgC carbon extraction from improved agricultural and forestry practices.

We examine here scenarios with 6%/year and 3%/year emission reduction starting in 2021, as well as scenarios with constant emissions and +2%/year emission growth starting in 2016 (Figs. 10b and 12b). The -6%/year and -3%/year scenarios leave a requirement to extract 153 795 and 237 PgC from the air during this century. Constant emission and +2%/year emission scenarios yield extraction requirements of 695 and 1630 PgC to reach 350 ppm CO₂ in 2100.

Total CO₂ extraction requirements for these scenarios are given in Fig. 10. Cost estimates here for extraction use amounts 100 Pg less than in Fig. 10 under assumption that 100 PgC can be stored via improved agricultural and forestry practices. Shortfall of this 100 PgC goal will 800 increase our estimated costs accordingly, as will the cost of the improved agricultural and forestry program.

Given a CO₂ extraction cost of \$150-350/tC for intensive negative emission technologies (Fig. 3f of Smith et al., 2016), the 53 PgC additional extraction required for the scenario with 6%/year emission reduction would cost \$8-18.5 trillion, thus \$100-230 billion per year if spread 805 uniformly over 80 years. We cannot rule out possible future reduction in CO₂ extraction costs, but given the energy requirements for removal and the already optimistic lower limit on our estimate, we do not speculate further about potential cost reduction.

In contrast, continued high emissions, between constant emissions and +2%/year, would require additional extraction of 595-1530 PgC (Fig. 10b) at a cost \$89-535 trillion or \$1.1-6.7 810 trillion per year over 80 years.¹³ Such extraordinary cost, along with the land area, fertilizer and water requirements (Smith et al., 2016) suggest that, rather than the world being able to buy its way out of climate change, continued high emissions would likely force humanity to live with climate change running out of control with all the consequences that would entail.

¹³ For reference, the United Nations global peacekeeping budget is about \$10B/year. National military budgets are larger: the 2015 USA military budget was \$596B and the global military budget was \$1.77 trillion (SIPRI 2016).

9.2 Mitigation Alternative

High costs of CO₂ extraction raise the question of how these costs compare to the alternative: taking actions to mitigate climate change by reducing fossil fuel CO₂ emissions. The Stern Review (Stern, 2006; Stern and Taylor, 2007) used expert opinion to produce an estimate for the cost of reducing emissions to limit global warming to about 2°C. Their central estimate was 1% of gross domestic product (GDP) per year, thus about \$800 billion per year. They argued that this cost was much less than likely costs of future climate damage if high emissions continue, unless we apply a high “discount rate” to future damage, which has ethical implications in its treatment of today’s young people and future generations. However, their estimated uncertainty of the cost is ±3%, i.e., the uncertainty is so large as to encompass GDP gain.

Hsu (2011) and Ackerman and Stanton (2012) argue that economies are more efficient if the price of fossil fuels better reflects costs to society, and thus GDP gain is likely with an increasing carbon price. Mankiw (2009) similarly suggests that a revenue-neutral carbon tax is economically beneficial. Hansen (2009, 2014) advocates an approach in which a gradually rising carbon fee is collected from the fossil fuel industry with the funds distributed uniformly to citizens. This approach provides incentives to business and the public that drive the economy toward energy efficiency, conservation, renewable energies and nuclear power. An economic study of this carbon-fee-and-dividend in the United States (Nystrom and Luckow, 2014) supports the conclusion that GDP increases as the fee rises steadily. These studies refute the common argument that environmental protection is damaging to economic prosperity.

We can also compare CO₂ extraction cost with the cost of carbon-free energy infrastructure. Global energy consumption in 2015 was 12.9 Gtoe,¹⁴ with coal providing 30% of global energy and almost 45% of global fossil fuel CO₂ emissions (BP, 2016). Most coal use, and its increases, are in Asia, especially China and India. Carbon-free replacement for coal energy is expected to be some combination of renewables (including hydropower) and nuclear power. China is leading the world in installation of wind, solar and nuclear power, with new nuclear power in 2015 approximately matching the sum of new solar and wind power (BP, 2016). For future decarbonization of electricity it is easiest to estimate the cost of the nuclear power component, because nuclear power can replace coal for baseload electricity without the need for energy storage or major change to national electric grids. Recent costs of Chinese and South Korean light water reactors are in the range \$2000-3000 per kilowatt (Chinese Academy of Engineering, 2015; Lovering et al., 2016). Although in some countries reactor costs stabilized or declined with repeated construction of the same reactor design, in others costs have risen for a variety of reasons (Lovering et al., 2016). Using \$2500 per kilowatt as reactor cost and assuming 85% capacity factor (percent uptime for reactors) yields a cost of \$10 trillion to produce 20% of present global energy use (12.9 Gtoe). Note that 20% of current global energy use is a huge amount (Fig. 13), exceeding the sum of present hydropower (6.8%), nuclear (4.4%), wind (1.4%), solar (0.4%), and other renewable energies (0.9%).

We do not suggest that new nuclear power plants on this scale will or necessarily should be built. Rather we use this calculation to show that mitigation costs are not large in comparison to costs of extracting CO₂ from the air. Renewable energy costs have fallen rapidly in the past 2-3 decades with the help of government subsidies, especially renewable portfolio standards that

¹⁴ Gtoe is gigatons oil equivalent. 1 Gtoe is 41.868 EJ (Exajoule = 10¹⁸ Joules) or 11,630 TWh (terawatt hours).




Figure 13. (a) Global fossil fuel emissions data from Boden et al. (2017) for 1751-2014 are extended to 2015 using BP (2016) data. (b) Global primary energy consumption data from BP (2016); energy accounting method is the substitution method (Macknick, 2011).

require utilities to achieve a specified fraction of their power from renewable sources. Yet fossil fuel use continues to be high, at least in part because fossil fuel prices do not include their full cost to society. Rapid and economic movement to non-fossil energies would be aided by a rising carbon price, with the composition of energy sources determined by competition among all non-fossil energy sources, as well as energy efficiency and conservation. Sweden provides a prime example: it cut per capita emissions by two-thirds since the 1990s while doubling per capita income in a capitalistic framework that embodies free-market principles (Pierrehumbert, 2016).

Mitigation of climate change deserves urgent priority. We disagree with assessments such as “The world will probably have only two choices if it wants to stay below 1.5°C of warming. It must either deploy carbon dioxide removal on an enormous scale or use solar geoengineering” (Parker and Geden, 2016). While we reject 1.5°C as a safe target – it is likely warmer than the Eemian and far above the Holocene range – Figure 12 shows that fossil fuel emission reduction of 3%/year beginning in 2021 yields maximum global warming ~1.5°C for climate sensitivity 3°C for 2×CO₂, with neither CO₂ removal nor geoengineering. These calculations show that mitigation – reduction of fossil fuel emissions – is very effective. We know no persuasive scientific reason to a priori reject as implausible a rapid phasedown of fossil fuel emissions.

10 Non-CO₂ GHGs, Aerosols and Purposeful Climate Intervention

10.1 Non-CO₂ GHGs

The annual increment in GHG climate forcing is growing, not declining. The increase is more than 20% in just the past five years (Fig. 8). Resurgence of CH₄ growth is partly responsible, but CO₂ is by far the largest contributor to growth of GHG climate forcing (Fig. 8). Nevertheless, given the difficulty and cost of reducing CO₂, we must ask about the potential for reducing non-CO₂ GHGs. Could realistic reductions of these other gases substantially alter the CO₂ abundance required to meet a target climate forcing?

We conclude, as discussed in Appendix A13, that a net decrease of climate forcing by non-CO₂ GHGs of perhaps −0.25 W/m² relative to today is plausible, but we must note that this is a dramatic change from the growing abundances, indeed accelerating growth, of these gases today. Achievement of this suggested negative forcing requires (i) successful completion of planned phase-out of MPTGs (−0.23 W/m²), (ii) absolute reductions of CH₄ forcing by 0.12 W/m² from

its present value, and (iii) N₂O forcing increasing by only 0.1 W/m². Achieving this net negative forcing of −0.25 W/m² for non-CO₂ gases would allow CO₂ to be 365 ppm, rather than 350 ppm, while yielding the same total GHG forcing. Absolute reduction of non-CO₂ gases is thus helpful, but does not alter the requirement for rapid fossil fuel emission reductions. Moreover, this is an optimistic scenario that is unlikely to occur in the absence of a reduction of CO₂, which is needed to limit global warming and thus avoid amplifying GHG feedbacks.

10.2 Aerosols and Purposeful Climate Intervention

Human-made aerosols today are believed to cause a large, albeit poorly measured, negative climate forcing (Fig. 4) of the order of −1 W/m² with uncertainty of at least 0.5 W/m² (Figure 7.19 Boucher et al., 2013). Fossil fuel burning is only one of several human-caused aerosol sources (Boucher et al., 2013). Given that human population continues to grow, and that human-caused climate effects such as increased desertification can lead to increased aerosols, we do not anticipate a large reduction in the aerosol cooling effect, even if fossil fuel use declines. Rao et al. (2017) suggest that future aerosol amount will decline due to technological advances and global action to control emissions. We are not confident of such a decline, as past controls have been at least matched by increasing emissions in developing regions, and global population continues to grow. However, to the extent that Rao et al. (2017) projections are borne out, they will only strengthen the conclusions of our present paper about the threat of climate change for young people and the burden of decreasing GHG amounts in the atmosphere.

Recognition that aerosols have a cooling effect, combined with the difficulty of restoring CO₂ to 350 ppm or less, inevitably raises the issue of purposeful climate intervention, also called geoengineering, and specifically solar radiation management (SRM). The cooling mechanism receiving greatest attention is injection of SO₂ into the stratosphere (Budyko, 1974; Crutzen, 2006), thus creating sulfuric acid aerosols that mimic the effect of volcanic aerosol cooling. That idea and others are discussed in a report of the U.S. National Academy of Sciences (NAS, 2015b) and references therein. We limit our discussion to the following summary comments.

Such purposeful intervention in nature, an attempt to mitigate effects of one human-made pollutant with another, raises additional practical and ethical issues. Stratospheric aerosols, e.g., could deplete stratospheric ozone and/or modify climate and precipitation patterns in ways that are difficult to predict with confidence, while doing nothing to alleviate ocean acidification caused by rising CO₂; we note that Keith et al. (2016) suggest alternative aerosols that would limit the impact on ozone. However, climate intervention also raises issues of global governance, and introduces the possibility of sudden global consequences if aerosol injection is interrupted (Boucher et al., 2013). Despite these issues, it is apparent that cooling by aerosols, or other methods that alter the amount of sunlight absorbed by Earth, could be effective more quickly than the difficult process of removing CO₂ from the air. Thus we agree with the NAS (2015b) conclusion that research is warranted to better define the climate, economic, political, ethical, legal and other dimensions of potential climatic interventions.

In summary, although research on climate interventions is warranted, the possibility of geoengineering can hardly be seen as alleviating the overall burden being placed on young people by continued high fossil fuel emissions. We concur with the assessment (NAS, 2015b) that such climate interventions are no substitute for the reduction of carbon dioxide emissions needed to stabilize climate and avoid deleterious consequences of rapid climate change.




Figure 14. Recent growth rate of total GHG effective climate forcing; points are 5-year running means, except 2015 point is a 3-year mean. See Fig. 8 for individual gases.

11. Discussion

Global temperature is now far above its range during the preindustrial Holocene, attaining at least the warmth of the Eemian period, when sea level reached +6-9 m relative to today. Also Earth is now out of energy balance, implying that more warming will occur, even if atmospheric GHG amounts are stabilized at today's level. Furthermore, the GHG climate forcing is not only still growing, the growth rate is actually accelerating, as shown in Fig. 14, which is extracted from data in our Fig. 8.

This summary, based on real world data for temperature, planetary energy balance, and GHG changes, differs from a common optimistic perception of progress toward stabilizing climate. That optimism may be based on the lowered warming target in the Paris Agreement (2015), slowdown in the growth of global fossil fuel emissions in the past few years (Fig. A1), and falling prices of renewable energies, but the hard reality of the climate physics emerges in Figs. 2, 5, 8 and 14. Although the scenarios employed in climate simulations for the most recent IPCC study (AR5) include cases with rapidly declining GHG growth, the scenarios do nothing to alter reality, which reveals that GHG growth rates not only remain high, they are accelerating.

The need for prompt action implied by these realities may not be a surprise to the relevant scientific community, because paleoclimate data revealed high climate sensitivity and the dominance of amplifying feedbacks. However, effective communication with the public of the urgency to stem human-caused climate change is hampered by the inertia of the climate system, especially the ocean and the ice sheets, which respond rather slowly to climate forcings, thus allowing future consequences to build up before broad public concern awakens. Some effects of human-caused global warming are now unavoidable, but is it inevitable that sea level rise of many meters is locked in, and, if so, on what time scale? Precise unequivocal answers to such questions are not possible. However, useful statements can be made.

First, the inertia and slow response of the climate system also allow the possibility of actions to limit the climate response by reducing human-caused climate forcing in coming years and decades. Second, the response time itself depends on how strongly the system is being forced; specifically, the response might be much delayed with a weaker forcing.

For example, studies suggesting multi-meter sea level rise in a century assume continued high fossil fuel emissions this century (Hansen et al., 2016) or at least a 2°C SST increase (DeConto and Pollard, 2016). Ice sheet response time decreases rapidly in models as the forcing increases, because processes such as hydrofracturing and collapse of marine-terminating ice cliffs spur ice sheet disintegration (Pollard et al., 2015). All amplifying feedbacks, including atmospheric water vapor, sea ice cover, soil carbon release and ice sheet melt could be reduced by rapid emissions phasedown. This would reduce the risk of climate change running out of humanity's control and provide time to assess the climate response, develop relevant technologies, and consider further purposeful actions to limit and/or adapt to climate change.

Concern exists that large sea level rise may be inevitable, because of numerous ice streams on Antarctica and Greenland with inward-sloping beds (beds that deepen upstream) subject to runaway marine ice sheet instability (Mercer, 1978; Schoof, 2007, 2010). Some ice stream instabilities may already have been triggered (Rignot et al., 2014), but the number of ice streams affected and the time scale of their response may differ strongly depending on the magnitude of the forcing (DeConto and Pollard, 2016). Sea level rise this century of say half a meter to a meter, which may be inevitable even if emissions decline, would have dire consequences, yet these are dwarfed by the humanitarian and economic disasters that would accompany sea level rise of several meters (McGranahan et al., 2007). Given the increasing proportion of global population living in coastal areas (Hallegatte et al., 2013), there is potential for forced migrations of hundreds of millions of people, dwarfing prior refugee humanitarian crises, challenging global governance (Biermann and Boas, 2010) and security (Gemenne et al., 2014).

Global temperature is a useful metric, because increasing temperature drives amplifying feedbacks. Global ocean temperature is a major factor affecting ice sheet size, as indicated by both model studies (Pollard et al., 2015) and paleoclimate analyses (Overpeck et al., 2006; Hansen et al., 2016). Eemian ocean warmth, probably not more than about +0.7°C warmer than pre-industrial conditions (McKay et al., 2011; Masson-Delmotte et al., 2013; Section 2.2 above), corresponding to global warmth about +1°C relative to preindustrial, led to sea level 6-9 m higher than today. This implies that, on the long run, the El Niño-elevated 2016 temperature of +1.3°C relative to preindustrial temperature, and even the (+1.05°C) underlying trend to date without the El Niño boost, are probably too high for maintaining our present coastlines.

We conclude that the world has already overshoot appropriate targets for GHG amount and global temperature, and we thus infer an urgent need for both (1) rapid phasedown of fossil fuel emissions, (2) actions that draw down atmospheric CO₂, and (3) actions that, at minimum, eliminate net growth of non-CO₂ climate forcings. These tasks are formidable and, with the exception of the Montreal Protocol agreement on HFCs that will halt the growth of their climate forcing (Appendix A13), they are not being pursued globally. Actions at citizen, city, state and national levels to reduce GHG emissions provide valuable experience and spur technical developments, but without effective global policies the impact of these local efforts is reduced by the negative feedback caused by reduced demand for and price of fossil fuels.

Our conclusion that the world has overshoot appropriate targets is sufficiently grim to compel us to point out that pathways to rapid emission reductions are feasible. Peters et al. (2013) note that Belgium, France and Sweden achieved emission reductions of 4-5%/year sustained over 10 or more years in response to the oil crisis of 1973. These rates were primarily a result of nuclear power build programs, which historically has been the fastest route to carbon-free energy (Fig. 2 of Cao et al., 2016). These examples are an imperfect analogue, as they were driven by a desire

for energy independence from oil, but present incentives are even more comprehensive. Peters et al. (2013) also note that a continuous shift from coal to natural gas led to sustained reductions of 1-2%/year in the UK in the 1970s and in the 2000s, 2%/year in Denmark in 1990-2000s, and 1.4%/year in the USA since 2005. Furthermore, these examples were not aided by the economy-wide effect of a rising carbon fee or tax (Hsu, 2011; Ackerman and Stanton, 2012; Hansen, 2014), which encourages energy efficiency and carbon-free energies.

In addition to CO₂ emission phase-out, large CO₂ extraction from the air is needed and a halt of growth of non-CO₂ climate forcings to achieve the temperature stabilization of our scenarios. Success of both CO₂ extraction and non-CO₂ GHG controls requires a major role for developing countries, given that they have been a large source of recent deforestation (IPCC, 2013) and have a large potential for reduced emissions. Ancillary benefits of the agricultural and forestry practices needed to achieve CO₂ drawdown, such as improved soil fertility, advanced agricultural practices, forest products, and species preservation, are of interest to all nations. Developed nations have a recognized obligation to assist nations that have done little to cause climate change yet suffer some of the largest climate impacts. If economic assistance is made partially dependent on verifiable success in carbon drawdown and non-CO₂ mitigation, this will provide incentives that maximize success in carbon storage. Some activities, such as soil amendments that enhance weathering, might be designed to support both CO₂ and other GHG drawdown.

Considering our conclusion that the world has overshoot the appropriate target for global temperature, and the difficulty and perhaps implausibility of negative emissions scenarios, we would be remiss if we did not point out the potential contribution of demand-side mitigation that can be achieved by individual actions as well as by government policies. Numerous studies (e.g. Hedenhus et al., 2014; Popp et al., 2010) have shown that reduced ruminant meat and dairy products is needed to reduce GHG emissions from agriculture, even if technological improvements increase food yields per unit farmland. Such climate-beneficial dietary shifts have also been linked to co-benefits that include improved sustainability and public health (Bajzelj et al., 2014; Tilman and Clark, 2014). Similarly, Working Group 3 of IPCC (2014) finds “robust evidence and high agreement” that demand-side measures in the agriculture and land use sectors, especially dietary shifts, reduced food waste, and changes in wood use have substantial mitigation potential, but they remain under-researched and poorly quantified.

There is no time to delay. CO₂ extraction required to achieve 350 ppm CO₂ in 2100 was ~100 PgC if 6%/year emission reductions began in 2013 (Hansen et al., 2013a). Required extraction is at least ~150 PgC in our updated scenarios, which incorporate growth of emissions in the past four years and assume that emissions will continue at approximately current levels until a global program of emission reductions begins in four years (in 2021 relative to 2020; see Figs. 9, 10 for reduction rates). The difficulty of stabilizing climate was thus markedly increased by a delay in emission reductions of eight years, from 2013 to 2021. Nevertheless, if rapid emission reductions are initiated soon, it is still possible that at least a large fraction of required CO₂ extraction can be achieved via relatively natural agricultural and forestry practices with other benefits. On the other hand, if large fossil fuel emissions are allowed to continue, the scale and cost of industrial CO₂ extraction, occurring in conjunction with a deteriorating climate and costly dislocations, may become unmanageable. Simply put, the burden placed on young people and future generations may become too heavy to bear.

Appendix A: Additional figures, tables and explanatory information




Figure A1. CO₂ emissions from fossil fuels and cement use based on Boden et al. (2017) through 2014, extended using BP (2016) energy consumption data. (a) is log scale and (b) is linear. Growth rates r in (a) for an n -year interval are from $(1+r)^n$ with end values being 3-year means to minimize noise.

A1. Fossil Fuel CO₂ Emissions

CO₂ emissions from fossil fuels in 2015 were only slightly higher than in 2014 (Fig. A1). Such slowdowns are common, usually reflecting the global economy. Given rising global population and the fact that nations such as India are still at early stages of development, the potential exists for continued emissions growth. Fundamental changes in energy technology are needed for the world to rapidly phase down fossil fuel emissions.

Emissions are growing rapidly in emerging economies; while growth slowed in China in the past two years, emissions remain high (Fig. 1). The Kyoto Protocol (1997), a policy instrument of the Framework Convention (UNFCCC, 1992), spurred emission reductions in some nations, and the collapse of the Soviet Union caused a large decrease of emissions by Russia (Fig. 1b). However, growth of international ship and air emissions (Fig. 1b) largely offset these reductions and the growth rate of global emissions actually accelerated from 1.5%/year in 1973-2000 to ~2.5%/year after 2000 (Fig. A1). China is now the largest source of fossil fuel emissions, followed by the U.S. and India, but on a per capita historical basis the U.S. is 10 times more accountable than China and 25 times more accountable than India for the increase of atmospheric CO₂ above its preindustrial level (Hansen and Sato, 2016). Tabular data for Figs. 1 and A1 are available on the web page www.columbia.edu/~mhs119/Burden.

A2. Transient Climate Response to cumulative CO₂ Emissions (TCRE)

The transient climate response (TCR), defined as the global warming at year 70 in response to a 1%/year CO₂ increase, for our simple Green's function climate model is 1.89°C with energy imbalance of 1.52 W/m² at that point; this TCR is in the middle of the range reported in the IPCC AR5 report (IPCC, 2013). We calculate the transient climate response to cumulative carbon emissions (TCRE) of our climate plus carbon cycle model as in Section 10.8.4 of IPCC (2013), i.e., $TCRE = TCR \times CAF/C_0$, where C_0 = preindustrial atmospheric CO₂ mass = 590 PgC and $CAF = C_{atm}/C_{sum}$, C_{atm} = atmospheric CO₂ mass minus C_0 and C_{sum} = cumulative CO₂ emissions (all evaluated at year 2100).




Figure A3a. Monthly (thin lines) and 12-month running mean (thick lines or filled colors for Niño 3.4) global, global land, sea surface, and Niño 3.4 temperatures. Temperatures are relative to 1951-1980 base period for the current GISTEMP analysis, which uses NOAA ERSST.v4 for sea surface temperature.

We find $\text{TCRE} = 1.54^{\circ}\text{C}$ per 1000 PgC at 2100 with constant emissions (which yields cumulative emissions of 1180 PgC at 2100, which is near the midpoint of the range assessed by IPCC, i.e., 0.8°C to 2.5°C per 1000 PgC (IPCC, 2013)). Our two cases with rapidly declining emissions never achieve 1000 PgC emissions, but TCRE can still be computed using the IPCC formulae, yielding $\text{TCRE} = 1.31$ and 1.25°C per 1000 PgC at 2100 for the cases of $-3\%/ \text{year}$ and $-6\%/ \text{year}$ respective emission reductions. As expected, the rapid emission reductions substantially reduce the temperature rise in 2100.

A3. Observed Temperature Data and Analysis Method

We use the current Goddard Institute for Space Studies global temperature analysis (GISTEMP), described by Hansen et al. (2010). The analysis combines data from: (1) meteorological station data of the Global Historical Climatology Network (GHCN) described by Peterson and Vose (1997) and Menne et al. (2012), (2) Antarctic research station data reported by the Scientific Committee on Antarctic Research (SCAR), (<http://www.antarctica.ac.uk/met/READER>), and (3) ocean surface temperature measurements from the NOAA Extended Reconstructed Sea Surface temperature (ERSST) (Smith et al., 2008; Huang et al., 2015).

Surface air temperature change over land is about twice SST change (Fig. A3a), and thus global temperature change is 1.3 times larger than the SST change. Note that the Arctic Ocean and parts of the Southern Ocean are excluded in the calculations because of inadequate data, but these regions are also not sampled in most paleo analyses and the excluded areas are small. Land area included covers 29% of the globe and ocean area included covers 65% of the globe.




Figure A3b. Global surface temperature relative to 1951-1980 in the GISTEMP analysis, comparing the current analysis using NOAA ERSST.v4 for sea surface temperature with results using ERSST.v3b.

The present analysis uses GHCN.v3.3.0 (Menne et al., 2012) for land data and ERSST.v4 for sea surface temperature (Huang et al., 2015). Update from GHCN.v2 used in our 2010 analysis to GHCN.v3 had negligible effect on global temperature change over the past century (see graph on http://www.columbia.edu/~mhs119/Temperature/GHCN_V3vsV2/). However, the adjustments to SST to produce ERSST.v4 have a noticeable effect, especially in the period 1939-1945, as shown by the difference between the two data sets (lower graph in Fig. A3b). This change is of interest mainly because it increases the magnitude of an already unusual global temperature fluctuation in the 1940s, making the 1939-1945 global temperature maximum even more pronounced than it was in ERSST.v3 data. Thompson et al. (2008) show that two natural sources of variability, the El Niño/Southern Oscillation and (possibly related) unusual winter Arctic warmth associated with advection over high Northern Hemisphere latitudes, partly account for global warmth of 1939-1945, and they suggest that the sharp cooling after 1945 is a data flaw, due to a rapid change in the mix of data sources (bucket measurements and engine room intake measurements) and a bias between these that is not fully accounted for.

Huang et al. (2015) justify the changes made to obtain version 4 of ERSST, the changes including more complete input data in ICOADS Release 2.5, buoy SST bias adjustments not present in version 3, updated ship SST bias adjustments using Hadley Nighttime Marine Air Temperature version 2 (HadNMat2), and revised low-frequency data filling in data sparse regions using nearby observations. ERSST.v4 is surely an improvement in the record during the past half century when spatial and temporal data coverages are best. On the other hand, the largest changes between v3 and v4 are in 1939-1945, coinciding with World War II and changes in the mix of data sources. Several hot spots appear in the Southern Hemisphere ocean during WWII in the v4 data, and then disappear after the war (Fig. A3c). These hot spots coincide with the locations of large SST changes between v3 and v4 (Fig. A3c), which leads us to suspect that




Figure A3c. Temperature anomalies in three periods relative to 1951-1980 comparing results obtained using ERSST.v3b (left column), ERSST.v4 (center column), and their difference (right column).

the magnitude of the 1940s global warming maximum (Fig. 2) is exaggerated; i.e., it is partly spurious. We suggest that this warming spike warrants scrutiny in the next version of the SST analysis. However, the important point is that these data adjustments and uncertainties are small in comparison with the long-term warming. Adjustments between ERSST.v3b and ERSST.v4 increase global warming over the period 1950-2015 by about 0.05°C, which is small compared with the ~1°C global warming during that period. The effect of the adjustments on total global warming between the beginning of the 20th century and 2015 is even smaller (Fig. A3b).

A4. Recent Global Warming Rate

Recent warming removes the illusion of a hiatus of global warming since the 1997-98 El Niño (Fig. 2). Several studies, including Trenberth and Fasullo (2013), England et al. (2014), Dai et al. (2015), Rajaratnam et al. (2015) and Medhaug et al. (2017), showed that temporary plateaus are consistent with expected long-term warming due to increasing atmospheric GHGs. Other analyses of the 1998-2013 plateau illuminate the roles of unforced climate variability and natural and human-caused climate forcings in climate change, with the Interdecadal Pacific Oscillation (a recurring pattern of ocean-atmosphere climate variability) playing a major role in the warming slowdown (Kosaka and Xie, 2013; Huber and Knutti, 2014; Meehl et al., 2014; Fyfe et al., 2016; Medhaug et al., 2017).

A5. Coincidence of 1880-1920 Mean and Preindustrial Global Mean Temperatures

The Framework Convention (UNFCCC, 1992) and Paris Agreement (2015) define goals relative to ‘preindustrial’ temperature, but do not define that period. We use 1880-1920, the earliest time with near global coverage of instrumental data, as the zero-point for temperature anomalies.

Although human-caused increases of GHGs would be expected to have caused a small warming by then, that warming was at least partially balanced by cooling from larger than average




Figure A6. 1000-year temperature response in two versions of GISS ModelE-R. (a) version used for CMIP5 simulations (Schmidt et al. 2014), which has higher resolution (40-layer atmosphere at $2^\circ \times 2.5^\circ$, 32-layer ocean at $1^\circ \times 1.25^\circ$), (b) version used by Hansen et al. (2016), which has coarse resolution (20-layer atmosphere at $4^\circ \times 5^\circ$, 12-layer ocean at $4^\circ \times 5^\circ$) and includes two significant improvements to small scale ocean mixing, cf. section 3.2 of Hansen et al. (2016).

volcanic activity in 1880-1920. Extreme Little Ice Age conditions may have been $\sim 0.1^\circ\text{C}$ cooler than the 1880-1920 mean (Abram et al., 2016), but the Little Ice Age is inappropriate to define preindustrial because the deep ocean temperature did not have time to reach equilibrium. Thus preindustrial global temperature has uncertainty of at least 0.1°C , and the 1880-1920 period, which has the merit of near-global data, yields our best estimate of preindustrial temperature.

A6. Land versus Ocean Warming at Equilibrium

Observations (Fig. A3a) show surface air temperature (SAT) over land increasing almost twice as much as sea surface temperature (SST) during the past century. This large difference is likely partly due to the thermal inertia of the ocean, which has not fully responded to the climate forcing due to increasing GHGs. However, land warming is heavily modulated by the ocean temperature, so land temperature too has not achieved its equilibrium response.

We use long climate model simulations to examine how much the ratio of land SAT change over ocean SST change (the observed quantities) is modified as global warming approaches its equilibrium response. This ratio is ~ 1.8 in years 901-1000 of doubled CO_2 simulations (Fig. A6) for two versions of GISS modelE-R (Schmidt et al., 2014; Hansen et al., 2016).

A7. Earth's Energy Imbalance

Hansen et al. (2011) inferred an Earth energy imbalance with the solar cycle effect removed of $+0.75 \pm 0.25 \text{ W/m}^2$, based on an imbalance of 0.58 W/m^2 during the 2005-2010 solar minimum, based on the analysis of von Schuckmann and Le Traon (2011) for heat gain in the upper 2 km of the ocean and estimates of small heat gains by the deep ocean, continents, atmosphere, and net melting of sea ice and land ice. The von Schuckmann and Le Traon analysis for 2005-2015 (Fig. 5) yields a decade-average 0.7 W/m^2 heat uptake in the upper 2 km of the ocean; addition of the smaller terms raises the imbalance to at least $+0.8 \text{ W/m}^2$ for 2005-2015, consistent with the recent estimate of $+0.9 \pm 0.1 \text{ W/m}^2$ by Trenberth et al. (2016) for 2005-2015. Other recent analyses including the most up-to-date corrections for ocean instrumental biases yield $+0.4 \pm 0.1$




Figure A8. Fossil fuel CO₂ emissions (left scale) and airborne fraction, i.e., the ratio of observed atmospheric CO₂ increase to fossil fuel CO₂ emissions.

W/m² by Cheng et al. (2017) for the period 1960-2015 and $+0.7 \pm 0.1$ W/m² by Dieng et al. (2017) for the period 2005-2013. We conclude that the estimate of $+0.75 \pm 0.25$ W/m² for the current Earth energy imbalance averaged over the solar cycle is still valid.

A8. CO₂ and CH₄ Growth Rates

Growth of airborne CO₂ is about half of fossil fuel CO₂ emissions (Fig. A8), the remaining portion of emissions being the net uptake by the ocean and biosphere (Ciais et al., 2013). Here we use the Keeling et al. (1973) definition of airborne fraction, which is the ratio of quantities that are known with good accuracy: the annual increase of CO₂ in the atmosphere and the annual amount of CO₂ injected into the atmosphere by fossil fuel burning. The data reveal that, even as fossil fuel emissions have increased by a factor of four over the past half century, the ocean and biosphere have continued to take up about half of the emissions (Fig. A8, right-hand scale). This seemingly simple relation between emissions and atmospheric CO₂ growth is not predictive as it depends on the growth rate of emissions being maintained, which is not true in cases with major changes in the emission scenario, so we use a carbon cycle model in Section 7 to compute atmospheric CO₂ as a function of emission scenario.

Oscillations of annual CO₂ growth are correlated with global temperature and with the El Niño/La Niña cycle¹⁵. Correlations (Fig. 6) are calculated for the 12-month running means, which effectively remove the seasonal cycle and monthly noise. Maxima of the CO₂ growth rate lag global temperature maxima by 7-8 months (Fig. 6b) and lag Niño3.4 [latitudes 5N-5S, longitudes 120-170W] temperature by ~10 months. These lags imply that the current CO₂ growth spike (Fig. 6 uses data through January 2017), associated with the 2015-16 El Niño, is well past its maximum, as Niño3.4 peaked in December 2015 and the global temperature anomaly peaked in February 2016.

CH₄ growth rate has varied over the past two decades, probably driven primarily by changes in emissions, as observations of CH₃CCl₃ show very little change in the atmospheric sink for

¹⁵ One mechanism for greater than normal atmospheric CO₂ growth during El Niños is the impoverishment of nutrients in equatorial Pacific surface water and thus reduced biological productivity that result from reduced upwelling of deep water (Chavez et al., 1999). However, the El Niño/La Niña cycle seems to have an even greater impact on atmospheric CO₂ via the terrestrial carbon cycle through effects on the water cycle, temperature, and fire, as discussed in a large body of literature (referenced, e.g., by Schwalm et al., 2011).

1235 CH₄ (Montzka et al., 2011; Holmes et al., 2013). Recent box-model inversions of the CH₄-CH₃-
CCl₃ system have argued for large fluctuations in the atmospheric sink over this period but there
is no identified cause for such changes (Rigby et al., 2017; Turner et al., 2017; Prather and
Holmes, 2017). Future changes in the sink could lead to increased atmospheric CH₄ separate
1240 from emission changes, but this effect is difficult to project and not included in the RCP
scenarios (Voulgarakis et al., 2013).

Carbon isotopes provide a valuable constraint (Saunois et al., 2016) that aids analysis of
which CH₄ sources¹⁶ contribute to the CH₄ growth resurgence in the past decade (Fig. 7).
Schaefer et al. (2016) conclude that the growth was primarily biogenic, thus not fossil fuel, and
located outside the tropics, most likely ruminants and rice agriculture. Such an increasing
1245 biogenic source is consistent with effects of increasing population and dietary changes (Tilman
and Clark, 2014). Nisbet et al. (2016) concur with Schaefer et al. (2016) that the CH₄ growth is
from biogenic sources, but from the latitudinal distribution of growth they conclude that tropical
wetlands¹⁷ have been an important contributor to the CH₄ increase. Their conclusion that
increasing tropical precipitation and temperature may be major factors driving CH₄ growth
1250 suggests the possibility that the slow climate-methane amplifying feedback might already be
significant. There is also concern that global warming will lead to a massive increase of CH₄
emissions from methane hydrates and permafrost (O'Connor et al., 2010), but as yet there is little
evidence for a substantial increase of emissions from hydrates or permafrost either now or over
the last 1,000,000 years (Berchet et al., 2016; Warwick et al., 2016; Quiquet et al., 2015).

1255 Schwietzke et al. (2016) use isotopic constraints to show that the fossil fuel contribution to
atmospheric CH₄ is larger than previously believed, but total fossil fuel CH₄ emissions are not
increasing. This conclusion is consistent with the above studies, and it does not contradict
evidence of increased fossil fuel CH₄ emissions at specific locations (Turner et al., 2016). A
recent inverse model study, however, contradicts the satellite studies and finds no evidence for
1260 increased US emissions (Bruhwiler et al., 2017). The recent consortium study of global CH₄
emissions finds with top-down studies that the recent increase is likely due to biogenic (natural
and human sources) sources in the tropics, but it is difficult to attribute the magnitude of the rise
to tropical wetlands alone (Saunois et al., 2017)

1265

A9. CO₂ Emissions in Historical Period

For land use CO₂ emissions in the historical period, we use the values labeled Houghton/2 by
Hansen et al (2008), which were shown in the latter publication to yield good agreement with
observed CO₂. We use fossil fuel CO₂ emissions data for 1850-2013 from Boden et al. (2016).
1270 BP (2016) fuel consumption data for 2013-2015 are used for the fractional annual changes of
each nation to allow extension of the Boden analysis through 2015. Emissions were almost flat
from 2014 to 2015, due to economic slowdown and increased use of low-carbon energies, but,
even if a peak in global emissions is near, substantial decline of emissions is dependent on
acceleration in the transformation of energy production and use (Jackson et al., 2016).

¹⁶ Estimated human-caused CH₄ sources (Ciais et al., 2013) are: fossil fuels (29%), biomass/biofuels (11%), Waste
and landfill (23%), ruminants (27%) and rice (11%)

¹⁷ Wetlands compose a majority of natural CH₄ emissions and are estimated to be equivalent to about 36% of the
anthropogenic source (Ciais et al., 2013)

A10. Tables of Effective Climate Forcings, 1850-2100

Table A10a. Effective forcings (W/m²) in 1850-2015 relative to 1850

	Year	CO ₂	^a CH ₄	^b CFCs	N ₂ O	^c O ₃	^d TA+SA	^e Volcano	Solar	Net
1280	1850	0.000	0.000	0.000	0.000	0.000	0.000	-0.083	0.000	-0.083
	1860	0.024	0.013	0.000	0.004	0.004	-0.029	-0.106	0.032	-0.058
	1870	0.048	0.027	0.000	0.008	0.009	-0.058	-0.014	0.048	0.068
	1880	0.109	0.041	0.000	0.011	0.014	-0.097	-0.026	-0.049	0.003
1285	1890	0.179	0.058	0.000	0.014	0.018	-0.146	-0.900	-0.070	-0.847
	1900	0.204	0.077	0.001	0.017	0.023	-0.195	-0.040	-0.063	0.024
	1910	0.287	0.115	0.002	0.022	0.026	-0.250	-0.072	-0.043	0.087
	1920	0.348	0.160	0.003	0.029	0.032	-0.307	-0.215	-0.016	0.034
1290	1930	0.425	0.206	0.004	0.037	0.036	-0.364	-0.143	0.014	0.215
	1940	0.494	0.247	0.005	0.043	0.045	-0.424	-0.073	0.037	0.374
	1950	0.495	0.291	0.009	0.052	0.056	-0.484	-0.066	0.055	0.408
	1960	0.599	0.365	0.027	0.061	0.078	-0.621	-0.106	0.102	0.505
1295	1970	0.748	0.461	0.076	0.075	0.097	-0.742	-0.381	0.093	0.427
	1980	0.976	0.568	0.185	0.097	0.115	-0.907	-0.108	0.169	1.095
	1990	1.227	0.659	0.303	0.125	0.117	-0.997	-0.141	0.154	1.447
	2000	1.464	0.695	0.347	0.150	0.117	-1.084	-0.048	0.173	1.814
1300	2005	1.619	0.651	0.356	0.162	0.123	-1.125	-0.079	0.019	1.770
	2010	1.766	0.710	0.364	0.177	0.129	-1.163	-0.082	0.028	1.929
	2015	1.927	0.730	0.373	0.195	0.129	-1.199	-0.100	0.137	2.192

Table A10b. Effective forcing (W/m²) in 2016-2100 relative to 1850

	Year	CO ₂	^a CH ₄	^b CFCs	N ₂ O	^c O ₃	^d TA+SA	^e Volcano	Solar	Net
1305	2016	1.942	0.698	0.367	0.192	0.130	-1.207	-0.100	0.097	2.119
	2020	2.074	0.702	0.373	0.201	0.130	-1.234	-0.100	-0.008	2.139
	2030	2.347	0.708	0.343	0.226	0.130	-1.296	-1.057	-0.008	1.393
	2040	2.580	0.735	0.301	0.254	0.123	-1.350	-0.100	0.027	2.569
1310	2050	2.803	0.766	0.267	0.288	0.117	-1.396	-0.100	0.062	2.807
	2060	3.017	0.791	0.243	0.322	0.111	-1.433	-1.208	0.097	1.940
	2070	3.222	0.804	0.229	0.358	0.105	-1.462	-0.100	0.132	3.289
	2080	3.421	0.792	0.215	0.391	0.098	-1.484	-0.100	0.167	3.500
1315	2090	3.614	0.722	0.199	0.427	0.091	-1.495	-1.240	0.167	2.484
	2100	3.801	0.619	0.191	0.456	0.085	-1.500	-0.100	0.167	3.719

^aCH₄: CH₄-induced changes of tropospheric O₃ and stratospheric H₂O are included

^bCFCs: This includes all GHGs except CO₂, CH₄, N₂O and O₃.

^cO₃: Half of troposphere O₃ forcing + stratosphere O₃ forcing from IPCC (2013).

^dTA+SA: tropospheric aerosols and surface albedo forcings combined.

^eVolc: volcanic forcing is zero when there are no stratospheric aerosols

Annual data are available at <http://www.columbia.edu/~mhs119/Burden/>.

CO₂, CH₄ and N₂O forcings are calculated with analytic formulae of Hansen et al. (2000). CH₄ forcing includes the factor 1.4 to convert adjusted forcing to effective forcing, thus incorporating the estimated effect of a CH₄ increase on tropospheric ozone and stratospheric water vapor. Our CH₄ adjusted forcing is significantly (~17%) higher than the values in IPCC (2013), but (~9%) smaller than values of Etminan (2017). Our factor 1.4 to convert direct radiative forcing to effective forcing is in the upper portion of the indirect effects discussed by Myhre et al. (2013), so our net CH₄ forcing agrees with Etminan et al. (2017) within uncertainties.




Figure A11. Solar irradiance and sunspot number in the era of satellite data. Left scale is the energy passing through an area perpendicular to Sun-Earth line. Averaged over Earth's surface the absorbed solar energy is $\sim 240 \text{ W/m}^2$, so the full amplitude of the measured solar variability is $\sim 0.25 \text{ W/m}^2$.

A11. Solar Irradiance

Solar irradiance has been measured from satellites since the late 1970s. Fig. A11 is a composite of several satellite-measured time series. Data through 28 February 2003 are an update of Frohlich and Lean (1998) obtained from Physikalisch Meteorologisches Observatorium Davos, World Radiation Center. Subsequent update is from University of Colorado Solar Radiation & Climate Experiment (SORCE). Historical total solar irradiance reconstruction is available at <http://lasp.colorado.edu/home/sorce/data/tsi-data/>. Data sets are concatenated by matching the means over the first 12 months of SORCE data. Monthly sunspot numbers support the conclusion that the solar irradiance in the current solar cycle is significantly lower than in the three preceding solar cycles.

The magnitude of the change of solar irradiance from the prior solar cycle to the current solar cycle is of the order of -0.1 W/m^2 , which is not negligible but is small compared with greenhouse gas climate forcing. On the other hand, the variation of solar irradiance from solar minimum to solar maximum is of the order of 0.25 W/m^2 , so the high solar irradiance in 2011 - 2015 contributes to the increase of Earth's energy imbalance between 2005-2010 and 2010-2015.

A12. Alternative Scenario

Simulated global temperature for the climate forcings of the "alternative scenario" discussed in Section 6 are shown in Fig. A12. The climate model, with sensitivity 3°C for doubled CO_2 , is the same as used for Fig. 12.

A13. Non- CO_2 GHGs

CO_2 is the dominant forcing in future climate scenarios. Growth of non- CO_2 GHG climate forcing is likely to be even smaller, relative to CO_2 forcing, than in recent decades (Fig. 8) if there is a strong effort to limit climate change. Indeed, recent agreement to use the Montreal




Figure A12. Simulated global temperature with historical climate forcings to 2000 followed by the alternative scenario. Historical climate forcings are discussed in the main text.

Protocol (2016) to phase down production of minor trace gases, the hydrofluorocarbons (HFCs), should cause annually added forcing of Montreal Protocol Trace Gases (MPTGs) + Other Trace Gases (OTGs) (red region in Fig. 8) to become near zero or slightly negative, thus at least partially off-setting growth of other non-CO₂ GHGs, especially N₂O.

Methane (CH₄) is the largest climate forcing other than CO₂ (Fig. 4). The CH₄ atmospheric lifetime is only about 10 years (Prather et al., 2012), so there is potential to reduce this climate forcing rapidly if CH₄ sources are reduced. Our climate simulations, based on the RCP6.0 non-CO₂ GHG scenarios, follow an optimistic path in which CH₄ increases moderately in the next few decades to 1960 ppb in 2070 and then decreases rapidly to 1650 ppb in 2100, yielding a forcing change of -0.1 W/m^2 . However, the IPCC (Kirtman et al., 2013) uses a more modern chemical model projection for the RCP anthropogenic emissions and gives a less beneficial view with a decrease to only 1734 ppb and a forcing change of -0.03 W/m^2 . RCP2.6 makes a more optimistic assumption: that CH₄ will decline monotonically to 1250 ppb in 2100, yielding a forcing of -0.3 W/m^2 (relative to today's 1800 ppb CH₄), but the IPCC projections of RCP2.6 reduce this to -0.2 W/m^2 (Kirtman et al., 2013).

Observed atmospheric CH₄ amount (Fig. A13a) is diverging on the high side of these optimistic scenarios. The downward offset (~ 20 ppb) of CH₄ scenarios relative to observations (Fig. A13a) is due to the fact that RCP scenarios did not include a data adjustment that was made in 2005 to match a revised CH₄ standard scale (E. Dlugokencky, pers. comm.), but observed CH₄ is also increasing more rapidly than in most scenarios. Reversal of CH₄ growth is made difficult by increasing global population, the diverse and widely distributed nature of agricultural sources, and global warming “in the pipeline,” as these trends create an underlying tendency for increasing CH₄. The discrepancy between observed and assumed CH₄ growth could also be due in part to increased natural sources or changes in the global OH sink (Dlugokencky et al., 2011; Turner et al., 2017). Evidence for increased natural sources in a warmer climate are suggested by glacial-interglacial CH₄ increases of the order of 300 ppb, and contributions to observed fluctuations cannot be ruled out on the basis of recent budgets (Ciais et al., 2013).

Methane emissions from rice agriculture and ruminants potentially could be mitigated by changing rice growing methods (Epule et al., 2011) and inoculating ruminants (Eckard et al., 2010; Beil, 2015)], but that would require widespread adoption of new technologies at the farmer level. California, in implementing a state law to reduce GHG emissions, hopes to dramatically cut agricultural CH₄ emissions (see www.arb.ca.gov/cc/scopingplan/scopingplan.htm), but




Figure A13. Comparison of observed CH₄ and N₂O amounts with RCP scenarios. RCP 6.0 and 4.5 scenarios for N₂O overlap. Observations are from NOAA/ESRL Global Monitoring Division. Natural sources and feedbacks not included in RCP scenarios may contribute to observed growth (see discussion).

California has one of the most technological and regulated agricultural sectors in the world. It is not clear that this level of management can occur in the top agricultural CH₄ emitters like China, India and Brazil. Methane leaks from fossil fuel mining, transportation and use can be reduced, indeed, percentage leakage from conventional fossil fuel mining and fuel use has declined substantially in recent decades (Schwietzke et al, 2016), but there is danger of increased leakage with expanded shale gas extraction (Caulton et al., 2014; Petron et al., 2013; Howarth, 2015; Kang et al., 2016).

Observed N₂O growth is exceeding all scenarios (Fig. A13b). Major quantitative gaps remain in our understanding of the nitrogen cycle (Kroeze and Bouwman, 2011), but fertilizers are clearly a principal cause of N₂O growth (Röckmann and Levin 2005; Park et al., 2012). More efficient use of fertilizers could reduce N₂O emissions (Liu and Zhang, 2011), but considering the scale of global agriculture, and the fact that fixed N is an inherent part of feeding people, there will be pressure for continued emissions at least comparable to present emissions. In contrast, agricultural CH₄ emissions are inadvertent and not core to food production. Given the current imbalance [emissions exceeding atmospheric losses by about 30% (Prather et al., 2012)] and the long N₂O atmospheric lifetime (116 ± 9 years; Prather et al., 2015) it is nearly inevitable that N₂O will continue to increase this century, even if emissions growth is checked. There can be no expectation of an N₂O decline that offsets the need to reduce CO₂.

The Montreal Protocol has stifled and even reversed growth of specific trace gases that destroy stratospheric ozone and cause global warming (Prather et al., 1996; Newman et al., 2009). The anticipated benefit over the 21st century is a drop in climate forcing of -0.23 W/m^2 (Prather et al., 2013). Protocol amendments that add other gases such as HFCs are important; forcings of these gases are small today, but without the Protocol their potential for growth is possibly as large as $+0.2 \text{ W/m}^2$ (Prather et al., 2013).

We conclude that a 0.25 W/m^2 decrease of climate forcing by non-CO₂ GHGs is plausible, but requires a dramatic change from the growing abundances of these gases today. Achievement requires (i) successful phase-out of MPTGs (-0.23 W/m^2), (ii) reduction of CH₄ forcing by 0.12 W/m^2 , and (iii) limiting N₂O increase to 0.1 W/m^2 . A net negative forcing of -0.25 W/m^2 for non-CO₂ gases would allow CO₂ to be 365 ppm, rather than 350 ppm, while yielding the same total GHG forcing. Thus potential reduction of non-CO₂ gases is helpful, but it does not alter the need for rapid fossil fuel emission reduction.

Acknowledgments

Support of the Climate Science, Awareness and Solutions program has been provided by the Durst family, the Grantham Foundation for Protection of the Environment, Jim and Krisann Miller, Gary Russell, Gerry Lenfest, the Flora Family Foundation, Elisabeth Mannschott, Alexander Totic and Hugh Perrine, which is gratefully acknowledged. DJB acknowledges funding through a Leverhulme Trust Research Centre Award (RC-2015-029). EJR acknowledges support from Australian Laureate Fellowship FL12 0100050

We appreciate the generosity, with data and advice, of Tom Boden, Ed Dlugokencky, Robert Howarth, Steve Montzka, Larissa Nazarenko and Nicola Warwick, the thoughtful reviews of the anonymous reviewers and several SC commenters, and assistance of the editor James Dyke.

The authors declare that they have no conflicts of interest. The first author (JH) notes that he is a plaintiff in the lawsuit Juliana et al. vs United States.

References

Abram, N. J., McGregor, H. V., Tierney, J. E., Evans, M. N., McKay, N. P., Kaufman, D. S., and PAGES 2k Consortium: Early onset of industrial-era warming across the oceans and continents, *Nature*, doi:10.1038/nature19082, 2016.

Ackerman, F. and Stanton, E. A.: Climate risks and carbon prices: revising the social cost of carbon, *Economics E-journal* 6 2012—10.5018/economics-ejournal.ja.2012—10, 2012.

Aiken, A.: Opinion and Order, Case No. 6:15-cv-01517-TC in United States District Court, District of Oregon, Eugene Division, 54 pp., 10 November 2016; Judge Aiken's Opinion and Order is available at: http://www.columbia.edu/~jeh1/mailings/2016/20161111_EmphaticRulingbyJudgeAiken.pdf

Alec L v. Jackson DDC, No. 11-CV-02235, 12/14/11, United States District Court, District of Columbia, May 31, 2012.

Anderson, K.: Talks in the city of light generate more heat, *Nature*, 528, 437, 2015.

Andrews, T., Gregory, J., Webb, M., and Taylor, K.: Forcing, feedbacks and climate sensitivity in CMIP5 coupled atmosphere-ocean climate models, *Geophys. Res. Lett.*, 39(9), L09712, 2012.

Applegate, P.J., Parizek, B.R., Nicholas, R.E., Alley, R.B., and Keller, K: Increasing temperature forcing reduces the Greenland Ice Sheet's response time scale, *Clim. Dyn.*, 45, 2001-2011, 2015.

Arrhenius, S.: On the influence of carbonic acid in the air upon the temperature on the ground, *Phil. Mag.*, 41, 237-276, 1896.

Bakker, P. and Renssen, H.: Last interglacial model-data mismatch of thermal maximum temperatures partially explained, *Clim. Past*, 10, 1633-1644, 2014.

Bajzelj, B., Richards, K. S., Allwood, J. M., Smith, P., Dennis, J. S., Curmi, E., and Gilligan, C. A.: Importance of food-demand management for climate mitigation, *Nature Clim. Change*, 4, 924-929, 2014.

Beil, L.: Getting creative to cut methane from cows, *Science News*, 188(11), 22-25, 2015.

Berchet, A., Bousquet, P., Pison, I., Locatelli, R., Chevallier, F., Paris, J.D., Dlugokencky, E.J., Laurila, T., Hataka, J., Viisanen, Y., Worthy, D.E.J., Nisbet, E.G., Fisher, R.E., France, J.L., Ivakhov, V., and Hermansen, O.: Atmospheric constraints on the methane emissions from the East Siberian Shelf, *Atmos. Chem. Phys.*, 16, 4147-4157, 2016.

Betts, R. A., Jones, C. D., Knight, J. R., Keeling, R. F., and Kennedy, J. J.: El Nino and a record CO₂ rise, *Nature Clim. Change*, 6, 806-810, 2016.

- 1475 Biermann, F. and Boas, I.: Preparing for a warmer world: towards a global governance system to protect climate refugees, *Global Environ. Politics*, 10, 60-88, 2010.
- Bindoff, N. L. and Stott, P. A.: Detection and attribution of climate change: from global to regional. In *Climate Change 2013: The Physical Science Basis*, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom, 2013.
- 1480 Boden, T. A., Marland, G., and Andres, R. J.: Global, Regional, and National Fossil-Fuel CO₂ Emissions, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee USA, doi 10.3334/CDIAC/00001_V2017, 2017.
- 1485 Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Fiengold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S. K., Sherwood, S., Stevens, B., and Zhang, X. Y.: Clouds and aerosols. in *Climate Change 2013: The Physical Science Basis*, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom, 2013.
- 1490 Bourassa, A. E., Robock, A., Randel, W. J., Deshler, T., Rieger, L. A., Lloyd, N. D., Llewellyn, E. J., and Degenstein, D. A.: Large volcanic aerosol load in the stratosphere linked to Asian Monsoon transport, *Science*, 337, 78-81, 2012.
- BP: 2016 BP Statistical Review of World Energy, available at: <http://www.bp.com/en/global/corporate/energy-economics/statistical-review-of-world-energy.html> (last access: 23 June 2016), 2016.
- 1495 Bruhwiler, L. M., Basu, S., Bergamaschi, P., Bousquet, P., Dlugokencky, E., Houweling, S., Ishizawa, M., Kim, H.-S., Locatelli, R., Maksyutov, S., Montzka, S., Pandey, S., Patra, P. K., Petron, G., Saunio, M., Sweeney, C., Schwietzke, S., Tans, P., and Weatherhead, E. C.: US CH₄ emissions from oil and gas production: Have recent large increases been detected?, *J Geophys Res-Atmos*, 122, 4070-4083, 10.1002/2016JD026157, 2017.
- Budyko, M.I.: *Izmeniya Klimata, Gidrometeoroizdat*, Leningrad, 1974.
- 1500 Bustamante, M., Robledo-Abad, C., Harper, R., Mbow, C., Ravindranath, N.H., Sperling, F., Haberl, H., de Siqueira Pinto, A. and Smith, P.: Co-benefits, trade-offs, barriers and policies for greenhouse gas mitigation in the Agriculture, Forestry and Other Land Use (AFOLU) sector, *Global Change Biology*, 20, 3270–3290, doi: 10.1111/gcb.12591, 2014.
- Canadell, J. G. and Raupach, M. R.: Managing forests for climate change mitigation, *Science*, 320, 1456, 2008.
- Cao, J., Cohen, A., Hansen, J., Lester, R., Peterson, P., and Xu, H.: China-U.S. cooperation to advance nuclear power, *Science*, 353, 547-548, 2016.
- 1505 Cao, L. and Caldeira, K.: Atmospheric carbon dioxide removal: long-term consequences and commitment, *Environ. Res. Lett.*, 5, doi:10.1088/1748-9326/5/2/024011, 2010.
- 1510 Caulton, D. R., Shepson, P. B., Santoro, R. L., Sparks, J. P., Howarth, R. W., Ingraffea, A. R., Cambaliza, M. O. L., Sweeney, C., Karion, A., Davis, K. J., Strim, B. H., Montzka, S. A., and Miller, B. R.: Toward a better understanding and quantification of methane emissions from shale gas development, *Proc. Natl. Acad. Sci. USA*, 111, 6237-6242, 2014.
- Charney, J., Arakawa, A., Baker, D., Bolin, B., Dickinson, R., Goody, R., Leith, C., Stommel, H., and Wunsch, C.: *Carbon Dioxide and Climate: A Scientific Assessment*, Natl. Acad. Sci. Press, Washington, DC, USA, 33p, 1979.
- 1515 Chavez, F. P., Strutton, P. G., Friederich, G. E., Feely, R. A., Feldman, G. C., Foley, D. G., and McPhaden, M. J.: Biological and chemical response of the Equatorial Pacific Ocean to the 1997-98 El Nino, *Science*, 286, 2126-2131, 1999.
- Cheng, L., Trenberth, K., Fasullo, J., Boyer, T., Abraham, J., and Zhu, J.: Improved estimates of ocean heat content from 1960-2015, *Sci. Advances*, 3, e1601545, 2017.

Chinese Academy of Engineering, Re-studying the nuclear development in China, Tsinghua University Press, edition 1, 2015. [in Chinese]

- 1520 Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Jones, C., Le Quéré, C., Myneni, R. B., Piao, S., and Thornton, P.: Carbon and other biogeochemical cycles. in *Climate Change 2013: The Physical Science Basis*, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom, 2013.
- 1525 Clark, P. U. and Huybers, P.: Interglacial and future sea level, *Nature*, 462, 856-857, 2009.

Clark, P.U., Shakun, J.D., Marcott, S.A., Mix, A.C., Eby, M., Kulp, S., Levermann, A., Milne, G.A., Pfister, P.L., Santer, B.D., Schrag, D.P., Solomon, S., Stocker, T.F., Strauss, B.H., Weaver, A.J., Winkelmann, R., Archer, D., Bard, E., Goldner, A., Lambeck, K., Pierrehumbert, R.T., and Plattner, G.K.: Consequences of twenty-first-century policy for multi-millennial climate and sea-level change, *Nature Clim. Change*, 6, 360-369, 2016.
- 1530 Coddington, O., Lean, J. L., Pilewskie, P., Snow, M., and Lindholm, D.: A solar irradiance climate data record, *Bull. Amer. Meteorol. Soc.*, doi:10.1175/BAMS-D-14-002265.12, 2015.

Collins, M., Knutti, R., Arblaster, J., Dufresne, J. L., Fichet, T., Friedlingstein, P., Gao, X., Gutowski, W. J., Johns, T., Krinner, G., Shongwe, M., Tebaldi, C., Weaver, A., and Wehner, M.: Long-term climate change: Projections, commitments and irreversibility, in: *Climate Change 2013: The Physical Basis*, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom, 2013.
- 1535 Copenhagen Accord 2009: United Nations Framework Convention on Climate Change, Draft decision available at: http://unfccc.int/meetings/copenhagen_dec_2009/items/5262.php, (last access: 9 September 2016), 2009.
- 1540 Creutzig, F., Ravindranath, N. H., Berndes, G., Bolwig, S., Bright, R., Cherubini, F., Chum, H., Corbera, E., Delucchi, M., Faaij, A., Fargione, J., Haberl, H., Heath, G., Lucon, O., Plevin, R., Popp, A., Robledo-Abad, C., Rose, S., Smith, P., Stromman, A., and Suh, S.: Bioenergy and climate change mitigation: an assessment, *GCB Bioenergy*, 7, 916-944, 2015.

Crowther, T.W. et al.: Quantifying global soil carbon losses in response to warming, *Nature*, 540, 104-108, 2016.
- 1545 Crutzen, P.J.: Albedo enhancement by stratospheric sulfur injections: A contribution to resolve a policy dilemma? *Clim. Change*, 77, 211-219, 2006.

Dai, A., Fyfe, J. C., Xie, S. P., and Dai, X.: Decadal modulation of global surface temperature by internal climate variability, *Nature Clim. Change*, published online 13 April doi:10.1038/nclimate2605, 2015.
- DeCicco, J.M., Liu, D.Y., Hoo, J., Krishnan, R., Kurthen, A., and Wang, L.: Carbon balance effects of U.S. biofuel production and use, *Clim. Change*, 138, 667-680, 2016.
- 1550 DeConto, R.M., Galeotti, S., Pagani, M., Tracy, D., Schaefer, K., Zhang, T., Pollard, D., and Beerling, D.J.: Past extreme warming events linked to massive carbon release from thawing permafrost, *Nature*, 484, 87-91, 2012.

DeConto, R. M. and Pollard, D.: Contribution of Antarctica to past and future sea-level rise, *Nature*, 531, 591-597, 2016.
- 1555 Dieng, H.B., Cazenave, A., Meyssignac, B., von Schuckmann, K., and Palanisamy, H.: Sea and land temperatures, ocean heat content, Earth's energy imbalance and net radiative forcing over the recent years, *International J. Climatol.*, doi:10.1002/joc.4996, 2017.
- Dlugokencky, E. J., Nisbet, E. G., Fisher, R., and Lowry, D.: Global atmospheric methane: budget, changes and dangers, *Phil. Trans. Roy. Soc. A*, 369, 2058-2072, 2011.
- 1560 Dlugokencky, E. J.: Global CH₄ data, NOAA/ESRL, available at: www.esrl.noaa.gov/gmd/ccgg/trends_ch4/, (last access: 13 September 2016), 2016.

- Dutton, A., Carlson, A. E., Long, A. J., Milne, G. A., Clark, P. U., DeConto, R., Horton, B. P., Rahmstorf, S., and Raymo, M. E.: Sea-level rise due to polar ice-sheet mass loss during past warm periods, *Science*, 349, 153 doi:10.1126/science.aaa4019, 2015.
- 1565 Eckard, R. J., Grainger, C., and de Klein, C. A. M.: Strategies to reduce methane emissions from farmed ruminants grazing on pasture, *Livestock Science*, 130, 47-56, 2010.
- Edwards, D. P., Lim, F., James, R. H., Pearce, C. R., Scholes, J., Freckleton, R. P., and Beerling, D. J.: Climate change mitigation: potential benefits and pitfalls of enhanced rock weathering in tropical agriculture, *Biol. Lett.*, 13, 20160715, <http://dx.doi.org/10.1098/rsbl.2016.0715>, 2017.
- 1570 England, M. H., McGregor, S., Spence, P., Meehl, G. A., Timmermann, A., Cai, W., Gupta, A. S., McPhaden, M. J., Purich, A., and Santoso, A.: Recent intensification of wind-driven circulation in the Pacific and the ongoing warming hiatus, *Nature Clim. Change*, published online 9 February, doi:10.1038/nclimate2106, 2014.
- Epule, E. T., Peng, C., and Mafany, N. M.: Methane emissions from paddy rice fields: strategies towards achieving a win-win sustainability scenario between rice production and methane reduction, *J. Sustain. Dev.*, 4, 188-196, 2011.
- 1575 Etminan, M., Myhre, G., Highwood, E.J., and Shine, K.P.: Radiative forcing of carbon dioxide, methane, and nitrous oxide: A significant revision of the methane radiative forcing, *Geophys. Res. Lett.*, 43, doi:10.1002/2016GL071930, 2017.
- 1580 Flato, G., Marotzke, J., Abiodun, B., Braconnot, P., Chou, S. C., Collins, W., Cox, P., Driouech, F., Emori, S., Eyring, V., Forest, C., Gleckler, P., Guilyardi, E., Jakob, C., Kattsov, V., Reason, C., and Rummukainen, M.: Evaluation of Climate Models. In: *Climate Change 2013: The Physical Science Basis*, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom, 2013.
- Franks, P.J., Royer, D.L., Beerling, D.J., Van de Water, P.K., Cantrill, D.J., Barbour, M.M., and Berry, J.A.: New constraints on atmospheric CO₂ concentration for the Phanerozoic, *Geophys. Res. Lett.*, 41, 4685-4694, 2014.
- 1585 Friedlingstein, P., Andrew, R. M., Rogelj, J., Peters, G. P., Canadell, J. G., Knutti, R., Luderer, G., Raupach, M. R., Schaeffer, M., van Vuuren, D. P., and Le Quéré, C.: Persistent growth of CO₂ emissions and implications for reaching climate targets, *Nat. Geosci.*, 1, 457-461, 2014.
- Frohlich, C. and Lean, J.: The Sun's total irradiance: cycles and trends in the past two decades and associated climate change uncertainties, *Geophys. Res. Lett.*, 25, 4377-4380, 1998.
- 1590 Fuss, S., Canadell, J. G., Peters, G. P., Tavoni, M., Andrew, R. M., Ciais, P., Jackson, R. B., Jones, C. D., Kraxner, F., Nakicenovic, N., Le Quere, C., Raupach, M. R., Sharifi, A., Smith, P., and Yamagata, Y.: Betting on negative emissions, *Nature Clim. Change*, 4, 850-853, 2014.
- Fyfe, J. C., Meehl, G. A., England, M. H., Mann, M. E., Santer, B. D., Flato, G. M., Hawkins, E., Gillett, N. P., Xie, S., Kosaka, Y., and Swart, N. C.: Making sense of the early-2000s warming slowdown, *Nature Clim. Change*, 6, 224-228, 2016.
- 1595 Gemenne, F., Barnett, J., Adger, W.N., and Dabelko, G.D.: Climate and security: evidence, emerging risks, and a new agenda, *Climatic Change*, 123(1), 1-9, 2014.
- 1600 Govin, A., Capron, E., Tzedakis, P.C., Verheyden, S., Ghaleb, B., Hillaire-Marcel, C., St-Onge, G., Stoner, J.S., Bassinot, F., Bazin, L., Blunier, T., Combourieu-Nebout, N., El Ouahabi, A., Gentry, D., Gersonde, R., Jimenez-Amat, P., Landais, A., Martrat, B., Masson-Delmotte, V., Parrenin, F., Seidenkrantz, M.S., Veres, D., Waelbroeck, C., and Zahn, R.: Sequence of events from the onset to the demise of the Last Interglacial: Evaluating strengths and limitations of chronologies used in climatic archives, *Quatern. Sci. Rev.*, 129, 1-36, 2015.
- Grant, K. M., Rohling, E. J., Bar-Matthews, M., Ayalon, A., Medina-Elizalde, M., Ramsey, C. B., Satow, C., and Roberts, A. P.: Rapid coupling between ice volume and polar temperature over the past 150,000 years, *Nature*, 491, 744-747, 2012.

- 1605 Grant, K. M., Rohling, E. J., Ramsey, C. B., Cheng, H., Edwards, R. L., Florindo, F., Heslop, D., Marra, F., Roberts, A. P., Tamisiea, M. E., and Williams, F.: Sea-level variability over five glacial cycles, *Nature Comm.*, 5, 5076, doi:10.1038/ncomms6076, 2014.
- Hallegatte, S., Green, C., Nicholls, R.J., and Corfee-Morlot, J.: Future flood losses in major coastal cities, *Nature Clim. Change*, 3, 802-806, 2013.
- 1610 Hansen, J., Russell, G., Lacis, A., Fung, I., Rind, D., and P. Stone, P.: [Climate response times: Dependence on climate sensitivity and ocean mixing](#). *Science*, **229**, 857-859, doi:10.1126/science.229.4716.857, 1985.
- Hansen, J., Sato, M., Ruedy, R., Lacis, A., and Oinas, V.: Global warming in the twenty-first century: An alternative scenario, *Proc. Natl. Acad. Sci. USA*, 97, 9875-9880, 2000.
- Hansen, J. and Sato, M.: Greenhouse gas growth rates, *Proc. Natl. Acad. Sci. USA*, 101, 16109-16114, 2004.
- 1615 Hansen, J.E.: [A slippery slope: How much global warming constitutes "dangerous anthropogenic interference"? An editorial essay](#). *Climatic Change*, **68**, 269-279, doi:10.1007/s10584-005-4135-0, 2005.
- Hansen, J., Sato, M., Ruedy, R., Nazarenko, L., Lacis, A., Schmidt, G. A., Russell, G., Aleinov, I., Bauer, M., Bauer, S., Bell, N., Cairns, B., Canuto, V., Chandler, M., Cheng, Y., Del Genio, A., Faluvegi, G., Fleming, E., Friend, A., Hall, T., Jackman, C., Kelley, M., Kiang, N. Y., Koch, D., Lean, J., Lerner, J., Lo, K., Menon, S., Miller, R. L., Minnis, P., Novakov, T., Oinas, V., Perlwitz, J.P., Perlwitz, J., Rind, D., Romanou, A., Shindell, D., Stone, P., Sun, S., Tausnev, N., Thresher, D., Wielicki, B., Wong, T., Yao, M., and Zhang, S.: [Efficacy of climate forcings](#), *J. Geophys. Res.*, 110, D18104, doi:10.1029/2005JD005776, 2005.
- 1620 Hansen, J.: Scientific reticence and sea level rise, *Environ. Res. Lett.*, 2, doi:10.1088/1748-9326/2/2/024002, 2007.
- Hansen, J., Sato, M., Kharecha, P., Russell, G., Lea, D.W., and Siddall, M.: [Climate change and trace gases](#). *Phil. Trans. Roy. Soc. A*, **365**, 1925-1954, doi:10.1098/rsta.2007.2052, 2007.
- 1625 Hansen, J., Sato, M., Kharecha, P., Beerling, D., Berner, R., Masson-Delmotte, V., Pagani, M., Raymo, M., Royer, D. L., and Zachos, J. C.: [Target atmospheric CO₂: Where should humanity aim?](#) *Open Atmos. Sci. J.*, 2, 217-231, 2008.
- Hansen, J.: Climate threat to the planet: implications for energy policy and intergenerational justice, Bjerknes lecture, American Geophysical Union, San Francisco, 17 December, available at: <http://www.columbia.edu/~jeh1/2008/AGUBjerknes20081217.pdf> (last access: 10 August 2016), 2008.
- 1630 Hansen, J.: *Storms of My Grandchildren*, New York, Bloomsbury, 304 pp., 2009.
- Hansen, J., Ruedy, R., Sato, M., and Lo, K.: [Global surface temperature change](#), *Rev. Geophys.*, 48, RG4004, 2010.
- Hansen, J., Sato, M., Kharecha, P., and von Schuckmann, K.: [Earth's energy imbalance and implications](#), *Atmos. Chem. Phys.*, 11, 13421-13449, doi:10.5194/acp-11-13421-2011, 2011.
- 1635 Hansen, J., Kharecha, P., Sato, M., Masson-Delmotte, V., Ackerman, F., Beerling, D. J., Hearty, P. J., Hoegh-Guldberg, O., Hsu, S., Parmesan, C., Rockstrom, J., Rohling, E. J., Sachs, J., Smith, P., Steffen, K., Van Susteren, L., von Schuckmann, K., and Zachos, J. C.: [Assessing "dangerous climate change": Required reduction of carbon emissions to protect young people, future generations and nature](#), *Plos One*, 8, e81648, 2013a.
- 1640 Hansen, J., Sato, M., Russell, G., and Kharecha, P.: Climate sensitivity, sea level and atmospheric carbon dioxide, *Phil Trans. Roy. Soc. A*, 371, 20120194, doi.org/10.1098/rsta.2012.0294, 2013b.
- Hansen, J. E.: *Environment and development challenges: the imperative of a carbon fee and dividend* The Oxford Handbook of the Macroeconomics of Global Warming editors Bernard, L. and Semmler, W. DOI:10.1093/oxfordhb/9780199856978,013.0026, Oxford University Press, 2014.
- 1645 Hansen, J. and Sato, M.: Regional climate change and national responsibilities, *Environ. Res. Lett.*, 11, published online 2 March, doi:10.1088/1748-9326/11/3/034009, 2016.

- 1650 Hansen, J., Sato, M., Hearty, P., Ruedy, R., Kelley, M., Masson-Delmotte, V., Russell, G., Tselioudis, G., Cao, J., Rignot, E., Velicogna, I., Tormey, B., Donovan, B., Kandiano, E., von Schuckmann, K., Kharecha, P., Legrande, A. N., Bauer, M., and Lo, K.-W.: Ice melt, sea level rise and superstorms: Evidence from paleoclimate data, climate modeling, and modern observations that 2°C global warming could be dangerous, Atmos. Chem. Phys., 16, 3761-3812, 2016.
- 1655 Hartmann, D.L., Klein Tank, A. M. G., Rusticucci, M., Alexander, L. V., Brönnimann, S., Charabi, Y., Dentener, F. J., Dlugokencky, E. J., Easterling, D. R., Kaplan, A., Soden, B. J., Thorne, P. W., Wild, M., and Zhai, P. M.: Observations: Atmosphere and Surface. In: Climate Change 2013: The Physical Science Basis, edited by: Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom, 2013.
- Hawkins, E., Ortega, P., Suckling, E., Schurer, A., Hegerl, G., Jones, P., Joshi, M., Osborn, T.J., Masson-Delmotte, V., Mignot, J., Thorne, P., and van Oldenborgh, G.J.: Estimating changes in global temperature since the pre-industrial period, Bull. Amer. Meteorol. Soc., <http://journals.ametsoc.org/doi/pdf/10.1175/BAMS-D-16-0007.1>, 2017.
- 1660 Hedenus, F., Wirsenius, S., and Johansson, D. J. A.: The importance of reduced meat and dairy consumption for meeting stringent climate change targets, Clim. Change, 124, 79, doi:10.1007/s10584-014-1104-5, 2014.
- Hoffman, J.S., Clark, P.U., Parnell, A.C., and He, F.: Regional and global sea-surface temperatures during the last interglaciation, Science, 355(6322), 276-279, 2017.
- 1665 Holmes, C. D., Prather, M. J., Sovde, O. A., and Myhre, G.: Future methane, hydroxyl, and their uncertainties: key climate and emission parameters for future predictions, Atmos. Chem. Phys., 13, 285-302, 2013.
- Howarth, R. W.: Methane emissions and climatic warming risk from hydraulic fracturing and shale gas development: implications for policy, Energy Emission Control Tech., 3, 45-54, 2015.
- Hsu, S. L.: The Case for a Carbon Tax, Island Press, Washington D.C., p 233, 2011.
- 1670 Huang, B., Banzon, V. F., Freeman, E., Lawrimore, J., Liu, W., Peterson, T. C., Smith, T. M., Thorne, P. W., Woodruff, S. D., and Zhang, H.-M.: Extended reconstructed sea surface temperature Version 4 (ERSST.v4). Part I: Upgrades and intercomparisons, J. Clim., 28, 911-930, 2015.
- Huber, M. and Knutti, R.: Natural variability, radiative forcing and climate response in the recent hiatus reconciled, Nature Geosci., 7(9), 651-656, 2014.
- 1675 Huntingford, C., Lowe, J. A., Gohar, L. K., Bowerman, N. H., Allen, M. R., Raper, S. C., and Smith, S. M.: The link between a global 2°C warming threshold and emissions in years 2020, 2050 and beyond, Environ. Res. Lett., 7, 014039, 2012.
- IPCC SRES (Intergovernmental Panel on Climate Change): Special Report on Emissions Scenarios, edited by: Nakicenovic, N. and Swart, R., Cambridge University Press, ISBN 0-521-80081-1, 978-052180081-5 (pb: 0-521-80493-0, 978-052180493-6), 2000.
- 1680 IPCC Guidelines for National Greenhouse Gas Inventories Volume 4 Agriculture, Forestry and Other Land Use, edited by: Eggleston, H. S., Buendia, L., Miwa, K., Ngara, T., and Tanabe, K., IGES Japan <http://www.ipcc-nggip.iges.or.jp/>, 2006.
- 1685 IPCC (Intergovernmental Panel on Climate Change): Climate Change 2013, edited by: Stocker, T., Qin, D., Q., Plattner, G. K., Tignor, M. M. B., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, 1535 pp., 2013.
- IPCC (Intergovernmental Panel on Climate Change): Climate Change 2014: Impacts, Adaptation, and Vulnerability, Field, C., Barros, V. R., Dokken, D. J., Mach, K.J., and Mastrandrea, M. D., Cambridge Univ. Press, 1132 pp, 2014.

- Jackson, R. B., Canadell, J. G., Le Quere, C., Andrew, R. M., Korsbakken, J. I., Peters, G. P., and Nakicenovic, N.: Reaching peak emissions, *Nature Clim. Change*, 6, 7-10, 2016.
- 1690 Jones, C.D., Ciais, P., Davis, S.J., Friedlingstein, Gasser, T., Peter, G.P., Rogelj, J., van Vuuren, D. P., Canadell, J. G., Cowie, A., Jackson, R. B., Jonas, M., Kriegl, E., Littleton, E., Lowe, J. A., Milne, J., Shrestha, G., Smith, P., Torvanger, A., and Wiltshire, A.: Simulating the Earth system response to negative emissions, *Environ. Res. Lett.*, 11(9), e095012, doi:10.1088/1748-9326/11/9/095012, 2016.
- 1695 Joos, F., Bruno, M., Fink, R., Stocker, T. F., Siegenthaler, U., Le, Quere. C., and Sarmiento, J. L.: An efficient and accurate representation of complex oceanic and biospheric models of anthropogenic carbon uptake, *Tellus*, 48B, 397-417, 1996.
- Judd, A. G., Hovland, M., Dimitrov, L. I., Garcia, Gil. S., and Jukes, V.: The geological methane budget at continental margins and its influence on climate change, *Geofluids*, 2, 109-126, 2002.
- 1700 Juliana et al. vs United States: Kelsey Cascadia Rose Juliana et al. v The United States of America; Barack Obama, in his official capacity as President of the United States, et al.; United States District Court, District of Oregon, Eugene Division Case no.: 6:15-cv-01517-TC, hearing held on 13 September 2016 on defendant's motion to dismiss case, decision pending, 2016.
- 1705 Kang, M., Christian, S., Celia, M.A., Mauzerall, D.L., Bill, M., Miller, A.R., Chen, Y., Conrad, M.E., Darrah, T.H., and Jackson, R.B.: Identification and characterization of high methane-emitting abandoned oil and gas wells, *Proc. Natl. Acad. Sci. USA*, doi:10.1073/pnas.1605913113, 2016.
- Kantola, I. B., McMasters, M. D., Beerling, D. J., Long, S. P., and DeLucia, E. H.: Potential of global croplands and bioenergy crops for climate change mitigation through deployment for enhanced weathering, *Biol. Lett.*, 13, 20160714, <http://dx.doi.org/10.1098/rsbl.20160714>, 2017.
- 1710 Keeling, C. D., Whorf, T. P., Wahlen, M., and van der Plicht, J.: Interannual extremes in the rate of rise of atmospheric carbon dioxide since 1980, *Nature*, 375, 666-670, 1993.
- Keith, D. W., Ha-Duong, M., and Stolaroff, J. K.: Climate strategy with CO₂ capture from the air, *Clim. Change*, 74, 17-45, 2006.
- Keith, D.: Why capture CO₂ from the atmosphere, *Science*, 325, 1654-1655, 2009.
- 1715 Keith, D.W., Weisenstein, D.K., Dykema, J.A., and Keutsch, F.N.: Stratospheric solar geoengineering without ozone loss, *Proc. Natl. Acad. Sci. USA*, doi/10.1073/pnas.1615572113, 2016.
- Kharecha, P. A. and Hansen, J. E.: Implications of “peak oil” for atmospheric CO₂ and climate, *Global Biogeochemical Cycles*, 22, GB3012 doi:10.1029/2007GB003142, 2008.
- 1720 Kirtman, B., S.B. Power, J.A. Adedoyin, G.J. Boer, R. Bojariu, I. Camilloni, F.J. Doblas-Reyes, A.M. Fiore, M. Kimoto, G.A. Meehl, M. Prather, A. Sarr, C. Schär, R. Sutton, G.J. van Oldenborgh, G. Vecchi and H.J. Wang, 2013: Near-term Climate Change: Projections and Predictability, in: *Climate Change 2013: The Physical Science Basis*, edited by Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom, 2013.
- Kohler, P., Hartman, J., and Wolf-Gladrow, D. A.: Geoengineering potential of artificially enhanced silicate weathering of olivine, *Proc. Natl. Acad. Sci. USA*, 107, 20228-20233, 2010.
- 1725 Kopp, G., Krivova, N., Lean, J., and Wu, C. J.: The Impact of the Revised Sunspot Record on Solar Irradiance Reconstructions, *Solar Physics*, doi:10.1007/s11207-016-0853-x, 2016.
- Kopp, R.E., Kemp, A.C., Bittermann, K., Horton, B.P., Donnelly, J.P., Gehrels, W.R., Hay, C.C., Mitrovica, J.X., Morrow, E.D., and Rahmstorf, S.: Temperature-driven global sea-level variability in the Common Era, *Proc. Natl. Acad. Sci. USA*, publ. online 22 February, E1434-E1441, 2016.

- 1730 Kosaka, Y. and Xie, S. P.: Recent global-warming hiatus tied to equatorial Pacific surface cooling, *Nature*, 501, 403-407, 2013.
- Kroeze, C. and Bouwman, L.: The role of nitrogen in climate change, *Curr. Opin. Envir. Sustain.*, 3, 279-280, 2011.
- Kvenvolden, K. A.: Gas hydrates – Geological perspective and global change, *Rev. Geophys.*, 31, 173-187, 1993.
- Kyoto Protocol: available at: http://unfccc.int/kyoto_protocol/items/2830.php (last access: 9 September 2016), 1997.
- 1735 Lacis, A.A, Schmidt, G.A., Rind, D., and Ruedy, R.A.: [Atmospheric CO₂: Principal control knob governing Earth's temperature](#). *Science*, **330**, 356-359, doi:10.1126/science.1190653, 2010.
- Lacis, A.A., Hansen, J.E., Russell, G.L., Oinas, V., and Jonas, J.: [The role of long-lived greenhouse gases as principal LW control knob that governs the global surface temperature for past and future climate change](#). *Tellus B*, **65**, 19734, doi:10.3402/tellusb.v65i0.19734, 2013.
- 1740 Lambeck, K., Rouby, H., Purcell, A., Sun, Y., and Sambridge, M.: Sea level and global ice volumes from the Last Glacial Maximum to the Holocene, *Proc. Natl. Acad. Sci. USA*, 111, 15296-15303, 2014.
- Levermann, A., Clark, P.U., Marzeion, B., Milne, G.A., Pollard, D., Radic, V., and Robinson, A.: The multimillennial sea-level commitment of global warming, *Proc. Natl. Acad. Sci. USA*, published online, doi:10.1073/pnas.1219414110, 2013.
- 1745 Levitus, S., Antonov, J. I., Boyer, T. P., Baranova, O. K., Garcia, H. E., Locarnini, R. A., Mishonov, A. V., Regan, J. R., Seidov, D., Yarosh, E. S., and Zweng, M. M.: World ocean heat content and thermocline sea level change (0-2000 m), 1955-2010, *Geophys. Res. Lett.*, 39, L10603 doi:10.1029/2012GL051106, 2012.
- Liu, X. and Zhang, F.: Nitrogen fertilizer induced greenhouse gas emissions in China, *Curr. Opin. Environ. Sustain.*, 3, 407-413, 2011.
- 1750 Liu, Z., Zhu, J., Rosenthal, Y., Zhang, X., Otto-Bliesner, B. L., Timmermann, A., Smith, R. S., Lohmann, G., Zheng, W., and Timm, O. E.: The Holocene temperature conundrum, *Proc. Natl. Acad. Sci. USA*, published online 11 August, doi:10.1073/pnas.1407229111, 2014.
- Lorius, C., Jouzel, J., Raynaud, D., Hansen, J., and Le Treut, H.: [The ice-core record: Climate sensitivity and future greenhouse warming](#). *Nature*, **347**, 139-145, doi:10.1038/347139a0, 1990.
- 1755 Lovering, J.R., Yip, A., and Nordhaus, T.: Historical construction costs of global nuclear power reactors, *Energy Policy*, 91, 371-382, 2016.
- Lunt, D.J., Haywood, A.M., Schmidt, G.A., Salzmann, U., Valdes, P.J., and Dowsett, H.J.: Earth system sensitivity inferred from Pliocene modelling and data, *Nature Geosci.*, 3, 60-64, 2010.
- Macknick, J.: Energy and CO₂ emission data uncertainties, *Carbon Management*, 2(2), 189-205, 2011.
- 1760 Mankiw, N.G.: Smart taxes: an open invitation to join the Pigou club, *Eastern Econ. J.*, 35, 14-23, 2009.
- Marcott, S. A., Shakun, J. D., Clark, P. U., and Mix, A. C.: A reconstruction of regional and global temperature for the last 11,300, *Science*, 339, 1198-1201, 2013.
- Marcott, S. A. and Shakun, J. D.: Holocene climate change and its context for the future, *PAGES*, 23(1), 28, 2015.
- 1765 Masson-Delmotte, V., Schulz, M., Abe-Ouchi, A., Beer, J., Ganopolski, A., Gonzalez Rouco, J. F., Jansen, E., Lambeck, K., Luterbacher, J., Naish, T., Osboorn, T., Otto-Bliesner, B., Quinn, T., Ramexh, R., Rojas, M. Shao, X., and Timmermann, A.: Information from paleoclimate archives, in: *Climate Change 2013: The Physical Science Basis*, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom, 2013.
- McGranahan, G., Balk, D., and Anderson, B.: The rising tide: assessing the risks of climate change and human settlements in low elevation coastal zones, *Environ. Urbanization*, 19, 17-37, 2007.
- 1770

- McKay, N. P., Overpeck, J. T., and Otto-Bliesner, B. L.: The role of ocean thermal expansion in Last Interglacial sea level rise, *Geophys. Res. Lett.*, 38, L14605, doi:10.1029/2011GL048280, 2011.
- Medhaug, I. Stolpe, M.B., Fischer, E.M. and Knutti, R.: Reconciling controversies about the ‘global warming hiatus’, *Nature*, 545, 41-47, 2017.
- 1775 Meehl, G. A., Teng, H., and Arblaster, J. M.: Climate model simulations of the observed early-2000s hiatus of global warming, *Nature Clim. Change*, published online 7, September doi:10.1038/nclimate2357, 2014.
- Meinshausen, M., Smith, S.J., Calvin, K., Daniel, J.S., Kainuma, M.L.T., Lamarque, J.F., Matsumoto, K., Montzka, S.A., Raper, S.C.B., Riahi, K., Thomson, A., Velders, G.J.M., and van Vuuren, D.P.P.: The RCP greenhouse gas concentrations and their extensions from 1765 to 2300, *Clim. Change*, 109, 213-241, 2011a.
- 1780 Meinshausen, M., Wigley, T. M. L., and Raper, S. C. B.: Emulating atmosphere-ocean and carbon cycle models with a simpler model, *MAGICC6 – Part 2: Applications, Atmos. Chem. Phys.*, 11, 1457-1471, 2011b.
- Menne, M. J., Durre, I., Vose, R. S., Gleason, B. E. and Houston, T. G.: An overview of the Global Historical Climatology Network-Daily database, *J. Atmos. Ocean Technol.*, 29, 897-910, 2012.
- Mercer, J.H.: West Antarctic ice sheet and CO₂ greenhouse effect: a threat of disaster, *Nature*, 271, 321-325, 1978.
- 1785 Montreal Protocol 2016: The Montreal Protocol on Substances that Deplete the Ozone Layer, available at: <http://www.ozone.unep.org/> (last access: 19 June 2016), 2016.
- Montzka, S. A., Krol, M., Dlugokencky, E., Hall, B., Jockel, P., and Lelieveld, J.: Small interannual variability of global atmospheric hydroxyl, *Science*, 331, 67-69, 2011.
- 1790 Moosdorf, N., Renforth, P., and Hartmann, J.: Carbon dioxide efficiency of terrestrial enhanced weathering, *Environ. Sci. Tech.*, 48, 4809-4816, 2014.
- Moss, R. H., Edmonds, J. A., Hibbard, K. A., Manning, M. R., Rose, S. K., van Vuuren, D. P., Carter, T. R., Emori, S., Kainuma, M., Kram, T., Meehl, G. A., Mitchell, J. F. B., Nakicenovic, N., Riahi, K., Smith, S. J., Stouffer, R. J., Thomson, A. M., Weyant, J. P., and Wilbanks, T. J.: The next generation of scenarios for climate change research and assessment, *Nature*, 463, 747-756, 2010.
- 1795 Myhre, G., Shindell, D., Breon, F., Collins, W., Fuglestad, J., Huang, J., Koch, D., Lamarque, J. F., Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura, T., and Zhang, H: Anthropogenic and natural climate forcing, in: *Climate Change 2013: The Physical Science Basis*, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom, 2013.
- 1800 National Academy of Sciences (NAS): Climate Intervention: Carbon Dioxide Removal and Reliable Sequestration, 154 pp., ISBN 978-0-309-30529-7, doi10.17226/18805, pdf at <http://nap.edu/18805>, 2015a.
- National Academy of Sciences (NAS): Climate Intervention: Reflecting Sunlight to Cool Earth, 234 pp., ISBN 978-0-309-31482-4, doi10.17226/18988, pdf at <http://nap.edu/18988>, 2015b.
- 1805 National Academy of Sciences (NAS): Attribution of Extreme Weather Events in the Context of Climate Change, National Academies Press, doi:10.17226/21852, 2016.
- Newman, P. A., Oman, L.D., Douglass, A. R., Fleming, E. L., Frith, S. M., Hurwitz, M. M., Kawa, S. R., Jackman, C. H., Krotkov, N. A., Nash, E. R., Nielsen, J. E., Pawson, S., Stolarski, R. S., and Velders, G. J. M.: What would have happened to the ozone layer if chlorofluorocarbons (CFCs) had not been regulated? *Atmos. Chem. Phys.*, 9, 2113-2128, 2009.
- 1810 Nisbet, E.G., Dlugokencky, E.J., Manning, M.R., Lowry, D., France, J.L., Michel, S.E., Miller, J.B., White, J.W.C., Vaughn, B., Bousquet, P., Pyle, J.A., Warwick, N.J., Cain, M., Brownlow, R., Zazzeri, G., Lanoiselle, M., Mannin, A.C., Gloor, E., Worthy, D.E.J., Brunke, E.G., Labuschagne, C., Wolff, E.W., and Ganesan, A.L.: Rising atmospheric methane: 2007-2014 growth and isotopic shift, *Global Biogeochem. Cycles*, 30, 1356-1370, 2016.

- 1815 Novakov, T., Ramanathan, V., Hansen, J. E., Kirchstetter, T. W., Sato, M., Sinton, J. E., and Sathaye, J. A.: Large historical changes of fossil-fuel black carbon aerosols, *Geophys. Res. Lett.*, 30, 1324, doi:10.1029/2005JD005977, 2003.
- Nystrom, S., and Luckow, M.S.: The economic, climate, fiscal, power, and demographic impact of a national fee-and-dividend carbon tax, Regional Economic Modeling, Inc., Washington, DC, <<http://bit.ly/1srHy3X>>, 126 pp., 2014.
- 1820 O'Connor, F. M., Boucher, O., Gedney, N., Jones, C. D., Folberth, G. A., Coppel, R., Friedlingstein, P., Collins, W. J., Chappellaz, J., Ridley, J., and Johnson, C. E.: Possible role of wetlands, permafrost, and methane hydrates in the methane cycle under future climate change, *Rev. Geophys.*, 48, RG4005 doi:10.1029/2010RG000326, 2010.
- 1825 O'Neill, B.C., Oppenheimer, M., Warren, R., Hallegatte, S., Kopp, R.E., Portner, H.O., Scholes, R., Birkmann, J., Foden, W., Licker, R., Mach, K.J., Marbaix, P., Mastrandrea, M.D., Price, J., Takahashi, K., van Ypersele, J.P., and Yohe, G.: IPCC reasons for concern regarding climate change risks, *Nature Clim. Change*, 7, 28-37, 2017.
- Overpeck, J.T., Otto-Bliesner, B.L., Miller, G.H., Muhs, D.R., Alley, R.B., and Kiehl, J.T.: Paleoclimatic evidence for future ice-sheet instability and rapid sea-level rise, *Science*, 311, 1747-1750, 2006.
- Paris Agreement 2015, UNFCCC secretariat, available at http://unfccc.int/paris_agreement/items/9485.php, (last access: 21 April 2017), 2015.
- 1830 Park, S., Croteau, P., Boering, K. A., Etheridge, D. M., Ferretti, D., Fraser, P. J., Kim, K. R., Krummel, P. B., Langenfelds, R. L., van Ommen, T. D., Steele, L. P., and Trudinger, C. M.: Trends and seasonal cycles in the isotopic composition of nitrous oxide since 1940, *Nature Geosci.*, 5, 261-265, 2012.
- Parker, A. and Geden, O.: No fudging on geoengineering, *Nature Geosci.*, www.nature.com/naturegeoscience, 2016.
- 1835 Pearson, R.G., Phillips, S.J., Loranty, M.M., Beck, P.S.A., Damoulas, T., Knight, S.J., and Goetz, S.J.: Shifts in Arctic vegetation and associated feedbacks under climate change, *Nature Clim. Change*, 3, 673-677, 2013.
- Peters, G. P., Andrew, R. M., Boden, T., Canadell, J. G., Ciais, P., Le Quéré, C., Marland, G., Raupach, M. R., and Wilson, C.: The challenge to keep global warming below 2°C, *Nature Clim. Change*, 3, 4-6, 2013.
- Peterson, T. C. and Vose, R.: An overview of the global historical climatology network temperature database, *Bull. Amer. Meteorol. Soc.*, 78, 2837-2849, 1997.
- 1840 Petron, G., Frost, G.J., Trainer, M.K., Miller, B.R., Dlugokencky, E.J., and Tans, P.: Reply to comment on "Hydrocarbon emissions characterization in the Colorado Front Range – A pilot study" by Michael A. Levi, *J. Geophys. Res. Atmos.*, 118, 236-242, doi:10.1029/2012JD018487, 2013.
- Pierrehumbert, R.: How to decarbonize? Look to Sweden, *Bull. Atomic Sci.*, 72, 105-111, 2016.
- 1845 Pollard, D., DeConto, R. M., and Alley, R. B.: Potential Antarctic ice sheet retreat driven by hydrofracturing and ice cliff failure, *Earth Planet. Sci. Lett.*, 412, 112-121, 2015.
- Popp, A., Lotze-Campen, H., and Bodirsky, B.: Food consumption, diet shifts and associated non-CO₂ greenhouse gases, *Global Environ. Change*, 20, 451-462, 2010.
- Prather, M., Midgley, P., Rowland, F. S., and Stolarski, R.: The ozone layer: the road not taken, *Science*, 381, 551-554, 1996.
- 1850 Prather, M. J., Holmes, C. D., and Hsu, J.: Reactive greenhouse gas scenarios: systematic exploration of uncertainties and the role of atmospheric chemistry, *Geophys. Res. Lett.*, 39, L09803, 2012.
- 1855 Prather, M., Flato, G., Friedlingstein, P., Jones, C., Lamarque, J.F., Liao, H. and Rasch P.: Annex II: Climate System Scenario Tables, in *Climate Change 2013: The Physical Science Basis*, edited by: Stocker, T., Qin, D., Q., Plattner, G. K., Tignor, M. M. B., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, pp. 1395-1445, 2013.

- Prather, M. J., Hsu, J., Deluca, N. M., Jackman, C. H., Oman, L. D., Douglass, A. R., Fleming, E. L., Strahan, S. E., Steenrod, S. D., Sovde, O. A., Isaksen, I. S. A., Froidevaux, L., and Funke, B.: Measuring and modeling the lifetime of nitrous oxide including its variability, *J. Geophys. Res. Atmos.*, 120, 5693-5705, 2015.
- 1860 Prather, M.J., and C. D. Holmes, Overexplaining or underexplaining methane's role in climate change, *PNAS* 114(21) 5324-5326, doi: 10.1073/pnas.1704884114, 2017.
- Quiquet, A., Archibald, A.T., Friend, A.D., Chappellaz, J., Levine, J.G., Stone, E.J., Telford, P.J., and Pyle, J.A.: The relative importance of methane sources and sinks over the Last Interglacial period and into the last glaciation. *Quaternary Sci. Rev.*, 112, 1-16, 2015.
- 1865 Rajaratnam, B., Romano, J., Tsiang, M., and Diffenbaugh, N. S.: Debunking the climate hiatus, *Clim. Change*, 133, 129-140, 2015.
- 1870 Rao, S., Klimont, Z., Smith, S.J., Van Dingenen, R., Dentener, F., Bouwman, L., Riahi, K., Amann, M., Bodirsky, B.L., van Vuuren, D.P., Reis, L.A., Calvin, K., Drouet, L., Fricko, O., Fujimori, S., Gernaat, D., Havlik, P., Harmsen, M., Hasegawa, T., Heyes, C., Hilaire, J., Luderer, G., Masui, T., Stehfest, E., Streffler, J., van der Sluis, S., and Tavoni, M.: Future air pollution in the Shared Socio-economic Pathways, *Global Environ. Change*, 42, 346-358, 2017.
- 1875 Rigby, M., Stephen A. Montzk, Ronald G. Prinn, James W. C. White, Dickon Younga, Simon O'Dohertya, Mark F. Lunta, Anita L. Ganesane, Alistair J. Manningf, Peter G. Simmondsa, Peter K. Salamehg, Christina M. Harthg, Jens Mühleg, Ray F. Weissg, Paul J. Fraserh, L. Paul Steeleh, Paul B. Krummelh, Archie McCullocha, and Sunyoung Park, Role of atmospheric oxidation in recent methane growth, *PNAS* 114(21): 5373–5377, doi: 10.1073/pnas.1616426114, 2017.
- Rignot, E., Mouginot, J., Morlighem, M., Seroussi, H., Scheuchl, B.: Widespread, rapid grounding line retreat of Pine Island, Thwaites, Smith, and Kohler glaciers, West Antarctica, from 1992 to 2011, *Geophys. Res. Lett.*, 41, 3502-3509, 2014.
- 1880 Rockmann, T. and Levin, I.: High-precision determination of the changing isotopic composition of atmospheric N₂O from 1990 to 2002, *J. Geophys. Res. Atmos.*, 110, D21304, 2005.
- Rogelj, J., Luderer, G., Pietzcker, R.C., Kriegler, E., Schaeffer, M., Krey, V., and Riahl, K.: Energy system transformations for limiting end-of-century warming to below 1.5°C, *Nature Clim. Change*, 5(6), 519-527, 2015.
- 1885 Rogelj, J., den Elzen, M., Hohne, N., Franzen, T., Fekete, H., Winkler, H., Schaeffer, R., Sha, F., Rihi, K., and Meinshausen, M.: Paris Agreement climate proposals need a boost to keep warming well below 2°C, *Nature*, 534, 631-639, 2016a.
- Rogelj, J., Schaeffer, M., Friedlingstein, P., Gillett, N. P., van Vuuren, D. P., Riahi, K., Allen, M., and Knutti, R.: Differences between carbon budget estimates unraveled, *Nature Clim. Change*, 534, 631-639, 2016b.
- 1890 Rohling, E. J., Sluijs, A., Dijkstra, H., Köhler, P., van de Wal, R.S., von der Heydt, A., Beerling, D., Berger, A., Bijl, P., Crucifix, M., DeConto, R., Drijfhout, S., Fedorov, A., Foster, G., Ganopolski, A., Hansen, J., Hönlisch, B., Hooghiemstra, H., Huber, M., Huybers, P., Knutti, R., Lea, D., Lourens, L.J., Lunt, D., Masson-Delmotte, V., Medina-Elizalde, M., Otto-Bliesner, B., Pagani, M., Pälike, H., Renssen, H., Royer, D., Siddall, M., Valdes, P., Zachos, J.C., and Zeebe, R.E.: Making sense of palaeoclimate sensitivity, *Nature*, 491, 683-691, doi:10.1038/nature11574, 2012a.
- 1895 Rohling, E. J., Medina-Elizade, M., Shepherd, J. G., Siddall, M., and Stanford, J. D.: Sea surface and high-latitude temperature sensitivity to radiative forcing of climate over several glacial cycles, *J. Clim.*, 25, 1635-1656, 2012b.
- Rohling, E. J., Haigh, I. D., Foster, G. L., Roberts, A. P., and Grant, K. M.: A geological perspective on potential future sea-level rise, *Scientific Reports*, 3, 3461 doi:10.1038/srep03461, 2013.
- Royer, D.L., Pagani, M., and Beerling, D.J.: Geobiological constraints on Earth system sensitivity to CO₂ during the Cretaceous and Cenozoic, *Geobiology*, 10, 298-310, 2012.

- 1900 Sanderson, B. M., O'Neill, B. C., and Tebaldi, C.: What would it take to achieve the Paris temperature targets? *Geophys. Res. Lett.*, 43, doi:10.1002/2016GL069563, 2016.
- Sato, M., Hansen, J. E., McCormick, M. P., and Pollack, J. B.: Stratospheric aerosol optical depths, 1850-1990, *J. Geophys. Res.*, 98, 22987-22994, doi:10.1029/93JD02553, 1993.
- 1905 Saunio, M., Jackson, R.B., Bousquet, P., Poulter, B., Canadell, J.G.: The growing role of methane in anthropogenic climate change, *Environ. Res. Lett.*, 11, 120207, doi:10.1088/1748-9326/11/12/120207, 2016.
- Saunio, M., Bousquet, P., Poulter, B., Peregon, A., Ciais, P., Canadell, J. G., Dlugokencky, E. J., Etiope, G., Bastviken, D., Houweling, S., Janssens-Maenhout, G., Tubiello, F. N., Castaldi, S., Jackson, R. B., Alexe, M., Arora, V. K., Beerling, D. J. Variability and quasi-decadal changes in the methane budget over the period 2000-2012, *Atmos. Chem. Phys. Discuss.*, doi:10.5194/acp-2017-296, 2017.
- 1910 Schädel, C., Bader, M. K.-F., Schuur, E. A. G., Biasi, C., Bracho, R., Čapek, P., De Baets, S., Diáková, K., Ernakovich, J., Estop-Aragones, C., Graham, D. E., Hartley, I. P., Iversen, C. M., Kane, E., Knoblauch, C., Lupascu, M., Martikainen, P. J., Natali, S. M., Norby, R. J., O'Donnell, J. A., Chowdhury, T. R., Šantrůčková, H., Shaver, G., Sloan, V. L., Treat, C. C., Turetsky, M. R., Waldrop, M. P., and Wickland, K. P.: Potential carbon emissions dominated by carbon dioxide from thawed permafrost soils, *Nature Clim. Change*, doi:10.1038/nclimate3054, 2016.
- 1915 Schaefer, H., Mikaloff Fletcher, S. E., Veidt, C., Lassey, K. R., Brailsford, G. W., Bromley, T. M., Dlugokencky, E. J., Michel, S. E., Miller, J. B., Levin, I., Lowe, D. C., Martin, R. J., Vaughn, B. H., White, J. W. C.: A 21st century shift from fossil-fuel to biogenic methane emissions indicated by ¹³CH₄, *Science*, 352, 80-84, 2016.
- Schmidt, G.A., Kelley, M., Nazarenko, L., Ruedy, R., Russell, G.L. Aleinov, I., Bauer, M. Bauer, S.E., Bhat, M.K., Bleck, R., Canuto, V., Chen, Y.-H., Cheng, Y., Clune, T.L. Del Genio, A., de Fainchtein, R., Faluvegi, G., Hansen, J.E., Healy, R.J., Kiang, N.Y., Koch, D., Lacis, A.A., LeGrande, A.N., Lerner, J., Lo, K.K., Matthews, E.E., Menon, S., Miller, R.L., Oinas, V., Olosio, A.O., Perlwitz, J.P., Puma, M.J., Putman, W.M., Rind, D., Romanou, A., Sato, M., Shindell, D.T., Sun, S., Syed, R.A., Tausnev, N., Tsigaridis, K., Unger, N., Voulgarakis, A., Yao, M.-S., and Zhang, J.: [Configuration and assessment of the GISS ModelE2 contributions to the CMIP5 archive](#). *J. Adv. Model. Earth Syst.*, 6, no. 1, 141-184, doi:10.1002/2013MS000265, 2014.
- 1920
- 1925 Schoof, C.: Ice sheet grounding line dynamics: steady states, stability, and hysteresis, *J. Geophys. Res.*, 112, F03S28, doi:10.1029/2006JF000664, 2007.
- Schoof, C.: Beneath a floating ice shelf, *Nature Geosci.*, 3, 450-451, 2010.
- Schuur, E. A. G., McGuire, A. D., Schädel, C., Grosse, G., Harden, J. W., Hayes, D. J., Hugelius, G., Koven, C. D., Kuhry, P., Lawrence, D. M., Natali, S. M., Olefeldt, D., Romanovsky, V. E., Schaefer, K., Turetsky, M. R., Treat, C. C., and Vonk, J. E.: Climate change and the permafrost carbon feedback, *Nature*, 520, 171-179, 2015.
- 1930
- Schwalm, C. R., Williams, C. A., Schaefer, K., Baker, I., Collatz, G. J., and Rödenbeck, C.: Does terrestrial drought explain global CO₂ flux anomalies induced by El Nino?, *Biogeosci.*, 8, 2493-2506, 2011.
- Schwietzke, S., Sherwood, O.A., Bruhwiler, L.M.P., Miller, J.B., Etiope, G., Dlugokencky, E.J., Michel, S.E., Arling, V.A., Vaughn, B.H., White, J.W.C., and Tans, P.P.: Upward revision of global fossil fuel methane emissions based on isotope database, *Nature*, 538, 88-91, 2016.
- 1935
- Searchinger, T., Heimlich, R., Houghton, R.A., Dong, F., Elobeid, A., Fabiosa, J., Tokgoz, S., Hayes, D., and Yu, T.-H.: Use of U.S. croplands for biofuels increases greenhouse gases through emissions from land-use change. *Science*, 319, 1238-40, 2008.
- SIPRI (Stockholm International Peace Research Institute): SIPRI Military Expenditure Database, available at: <https://www.sipri.org/databases/milex>, (last access: 17 September 2016), 2016.
- 1940
- Smith, P., Bustamante, M., Ahammad, H., Clark, H., Dong, H., Elsiddig, E.A., Haberl, H., Harper, R., House, J., Jafari, M., Masera, O., Mbow, C., Ravindranath, N.H., Rice, C.W., Abad, C.R., Romanovskaya, A., Sperling, F., and Tubiello, F.N.: Agriculture, Forestry and Other Land Use (AFOLU), in: *Climate Change 2014: Mitigation of Climate Change, Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel*

- 1945 on Climate Change, edited by: Edenhofer, O., Pichs-Madruga, R., Sokona, Y., Minx, J.C. Farahani, E., Kadner, S., Seyboth, K. Adler, A., Baum, I., Brunner, S., Eickemeier, P., Kriemann, B. Savolainen, J., Schlomer, S., von Stechow, C., and Zwickel, T., Cambridge University Press, Cambridge, United Kingdom, 2014.
- Smith, P.: Soil carbon sequestration and biochar as negative emission technologies, *Global Change Biology*, 22, 1315-1324, 2016.
- 1950 Smith, P., Davis, S. J., Creutzig, F., Fuss, S., Minx, J., Gavrielle, B., Kato, E., Jackson, R. B., Cowie, A., Kriegler, E., van Vuuren, D. P., Rogelj, J., Ciais, P., Milne, J., Canadell, J. G., McCollum, D., Peters, G., Andrew, R., Krey, V., Shrestha, G., Friedlingstein, P., Gasser, T., Grüber, A., Heidug, W. K., Jonas, M., Jones, C. D., Kraxner, F., Littleton, E., Lowe, J., Moreira, J. R., Nakicenovic, N., Obersteiner, M., Patwardhan, A., Rogner, M., Rubin, E., Sharifi, A., Torvanger, Asbjørn, Yamagata, Y., Edmonds, J., and Yongsung, C.: Biophysical and economic limits to negative CO₂ emissions, *Nature Clim. Change*, 6, 42-50, 2016.
- 1955 Smith, T. M., Reynolds, R. W., and Lawrimore, J.: Improvements to NOAA's historical merged land-ocean surface temperature analysis (1880-2006), *J. Clim.*, 21, 2283-2296, 2008.
- Socolow, R.: Direct air capture of CO₂ with chemicals: a technology assessment for the APS Panel on Public Affairs (American Physical Society), 2011.
- 1960 Stern, N.: The economics of Climate Change: The Stern Review, Cambridge University Press, Cambridge, 2006.
- Stern, N. and Taylor, C.: Climate change: risk, ethics, and the Stern review, *Science*, 317, 203-204, 2007.
- Taylor, L. L., Quirk, J., Thorley, R. M. S., Kharecha, P. A., Hansen, J., Ridgwell, A., Lomas, M. R., Banwart, S. A., and Beerling, D. J.: Enhanced weathering strategies for stabilizing climate and averting ocean acidification, *Nature Clim. Change*, 6, 402-406, 2016.
- 1965 Tedesco, M., Doherty, S., Fettweis, X., Alexander, P., Jeyaratnam, J., and Stroeve, J.: The darkening of the Greenland ice sheet: trends, drivers, and projections (1981-2100), *Cryosphere*, 10, 477-496, 2016.
- Thompson, D. W. J., Kennedy, J. J., Wallace, J. M., and Jones, P. D.: A large discontinuity in the mid-twentieth century in observed global-mean surface temperature, *Nature*, 453, 646-649 doi:10.1038/nature06982, 2008.
- Tilman, D. and Clark, M.: Global diets link environmental sustainability and human health, *Nature*, 515, 518-522, 2014.
- 1970 Trenberth, K. E. and Fasullo, J. T.: An apparent hiatus in global warming?, *Earth's Future*, doi:10.1002/2013EF000165, 2013.
- Trenberth, K., Fasullo, J., von Schuckmann, K., and Cheng, L.: Insights into Earth's energy imbalance from multiple sources, *J. Clim.*, 29(20), doi:10.1175/JCLI-D-16-0339.1 in press, 2016.
- 1975 Turner, A. J., Jacob, D. J., Benmergui, J., Wofsy, S. C., Maasakkers, J. D., Butz, A., Hasekamp, O., and Biraud, S. C.: A large increase in U.S. methane emissions over the past decade inferred from satellite data and surface observations, *Geophys. Res. Lett.*, 43, 2218–2224, doi:10.1002/2016GL067987, 2016.
- Turner, A.J., Frankenberg, C., Wennberg, P.O. and Jacob, D.J.: Ambiguity in the causes for decadal trends in atmospheric methane and hydroxyl, *Proc. Natl. Acad. Sci. USA*, 114, 1-6, 2017.
- 1980 Turney, C. S. M. and Jones, R. T.: Does the Agulhas Current amplify global temperatures during super-interglacials?, *J. Quatern. Sci.*, 25, 839-843, 2010.
- Tyndall, J.: On the absorption and radiation of heat by gases and vapours, and on the physical connexion of radiation, absorption, and conduction – The Bakerian Lecture, *Phil. Mag.*, 22, 273-285, 1861.
- 1985 United Nations: Framework Convention on Climate Change (UNFCCC), United Nations, New York, NY, available at: http://unfccc.int/essential_background/items/6036.php (last access: 3 March 2016), 1992.

United Nations: UNFCCC Technical Annex – Aggregate effect of the intended nationally determined contributions: an update. http://unfccc.int/focus/indc_portal/items/9240.php, 2016.

- 1990 van Vuuren, D. P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G. C., Kram, T., Krey, V., Lamarque, J.-F., Masui, T., Meinshausen, M., Nakicenovic, N., Smith, S. J., and Rose, S. K.: The representative concentration pathways: an overview, *Clim. Change*, 109, 5-31, 2011.
- Vial, J., Dufresne, J. L., and Bony, S.: On the interpretation of inter-model spread in CMIP5 climate sensitivity estimates, *Clim. Dyn.*, published online 23 March, doi:10.1007/s00382-013-1725-9, 2013.
- von Schuckmann, K. and Le Traon, P. Y.: How well can we derive Global Ocean Indicators from Argo data?, *Ocean Sci.*, 7, 783-791 doi:10.5194/os-7-783-2011, 2011.
- 1995 von Schuckmann, K., Palmer, M. D., Trenberth, K. E., Cazenave, A., Chamber, D., Champollion, N., Hansen, J., Josey, S. A., Loeb, N., Mathieu, P. P., Meyssignac, B., and Wild, M.: An imperative to monitor Earth's energy imbalance, *Nature Clim. Change*, 6, 138-144, 2016.
- 2000 Voulgarakis, A., Naik, V., Lamarque, J.-F., Shindell, D. T., Young, P. J., Prather, M. J., Wild, O., Field, R. D., Bergmann, D., Cameron-Smith, P., Cionni, I., Collins, W. J., Dalsøren, S. B., Doherty, R. M., Eyring, V., Faluvegi, G., Folberth, G. A., Horowitz, L. W., Josse, B., MacKenzie, I. A., Nagashima, T., Plummer, D. A., Righi, M., Rumbold, S. T., Stevenson, D. S., Strode, S. A., Sudo, K., Szopa, S., and Zeng, G.: Analysis of present day and future OH and methane lifetime in the ACCMIP simulations, *Atmos. Chem. Phys.*, 13, 2503-2587, 2013.
- 2005 Warwick, N.J., Cain, M.L., Fisher, R., France, J.L., Lowry, D., Michel, S.E., Nisbet, E.G., Vaughn, B.H., White, J.W.C. and Pyle, J.A.: Using $\delta^{13}\text{C}\text{-CH}_4$ and $\delta\text{D}\text{-CH}_4$ to constrain Arctic methane emissions, *Atmos. Chem. Phys.*, 16, 14891-14908, doi:10.5194/acp-16-14891-2016, 2016.
- Williamson, P.: Scrutinize CO₂ removal methods, *Nature*, 530, 153-155, 2016.
- Woolf, D., Amonette, J. E., Street-Perrott, F. A., Lehmann, J., and Joseph, S.: Sustainable biochar to mitigate global climate change, *Nature Comm.*, 1, 56 doi:10.1038/ncomms1053, 2010.
- 2010 Yu, J., Menviel, L., Jin, Z.D., Thornalley, D.J.R., Barker, S., Marino, G., Rohling, E.J., Cai, Y., Zhang, F., Wang, X., Dai, Y. Chen, P., and Broecker, W.S.: Sequestration of carbon in the deep Atlantic during the last glaciation, *Nature Geosci.*, 9, 319-324, 2016.
- Zachos, J., Pagani, M., Sloan, L., Thomas, E., and Billups, K.: Trends, rhythms, and aberrations in global climate 65 Ma to present, *Science*, 292, 686-693, 2001.
- 2015 Zachos, J.C., Dickens, G.R., and Zeebe, R.E.: An early Cenozoic perspective on greenhouse warming and carbon-cycle dynamics, *Nature*, 451, 279-283, 2008.