

# ***Interactive comment on “Changes in tropical cyclones under stabilized 1.5 °C and 2.0 °C global warming scenarios as simulated by the Community Atmospheric Model under the HAPPI protocols” by Michael F. Wehner et al.***

**Michael F. Wehner et al.**

mfwehner@lbl.gov

Received and published: 17 January 2018

Replies are in **red**

General comments: This paper describes changes in tropical cyclone (TC) activities simulated by the Community Atmospheric Model under the HAPPI protocols. The HAPPI protocol is a relatively smaller change in global mean temperature at 1.5 and 2.0 stabilizing levels compared to those mostly used for time slice experiments under global warming condition. One might have expected that the simulated signals of TCs

Printer-friendly version

Discussion paper



would be smaller or not be detected. However, in this paper, the simulation shows robust changes in TC activities similar to those obtained the existing literatures. In particular, the reduction of TC number, particularly that of Category 0, over the global domain is very robust and consistent. Projections of such extreme events as tropical cyclones under the condition of the HAPPI protocols are informative and useful for the society. This paper should be published soon with some minor revision suggested below.

The projection of the characteristics of tropical cyclones should also be tabulated. The data will be compiled later for comparison with other model results. Please refer to IPCC AR5 (Chapter 14, Supplementary Material). **We added tables to this effect to the Appendix.**

Specific comments: p. 10, Fig. 5: The curve for HAPPI 1.5 is not visible. Please clarify the figure. **Colors have been changed. The gray is now darker.**

p. 11, Fig. 6: This figure should be compared to observation. Add bars of the observational number of ACE. Why ACE in the South Indian Ocean is so high? Please add explanation. **We have added observational estimates of ACE according to Table 2 of Maue (2011). He did not calculate SIO ACE but did calculate southern Hemisphere. In the revised figure, the model and the observational estimate are quite consistent for the SH. However, we note that in the SH, TCs are probably not as well characterized as in the NH. Revised text added as follows:"Comparison with an observational estimate (Maue 2011) suggests that the model is overactive by this measure of tropical cyclone activity although differences in the methods that tracks and wind speeds are calculated could explain some of the biases shown in figure 6. "**

**Also note, there was a factor of 2 error in the original figure due to a miscommunication between the software engineer and the lead author. This has been corrected.**

p. 12: "Chavas et al. (2016)" (L8) is not included in the reference list, so that the definition of "Chavas radii" (L9) is not clear. We cannot understand the following sentences

Printer-friendly version

Discussion paper



in the paragraph and Fig. 8. Please explain why the radii is larger for the weaker tropical cyclones. We have added the missing citation and reworked this paragraph to replace "storm size" with "Chavas radii". The intent is to measure structural changes in tropical cyclones and we agree that having multiple "storm sizes" for a single storm is confusing. By referring this structural quantity to the Chavas radii, we hope it is clearer. This is a topic that we aim to provide a specialized paper on later this year.

p. 14, L16-18, “: : are not significantly altered in warmer climates, most notably the robust relationship between maximum wind speeds and central pressure minima”: The conclusion of the subtle behaviors is specific to this model. In particular, pressure-wind relation likely depends on the model resolution. Such remarks should be added. We expand the discussion on this point with the following sentences: "The peak wind speeds and central pressure minima relationship is controlled by the mechanical constraints of gradient wind balance, storm size and Coriolis force (Chavas 2017 and Chavas, private communication). The small poleward shift in the track density (figure 5) and subtle structural changes in wind speed radii discussed below are not large enough to change this relationship. Warmer temperatures do change the distribution of peak wind speeds and central pressure minima (figure 3) but does not appear to substantially change how they co-vary. We do note however that model resolution and structure may influence the simulation of this relationship..."

---

Interactive comment on Earth Syst. Dynam. Discuss., <https://doi.org/10.5194/esd-2017-101>, 2017.

Printer-friendly version

Discussion paper

