We want to thank the reviewers for their their constructive suggestions and comments. We have responded to their points below. In blue are the are the original responses suggesting how to revise the manuscript, and in red are our final responses, explaining what was done in the revised version of the manuscript.

Anonymous Referee #1

Received and published: 18 January 2018

This manuscript evaluates the quality of climate simulations with the NorESM1-Happi model, which is a slightly modified version of NorESM1-M (which was used in CMIP5). In addition, the authors discuss simulations with NorESM1-Happi that attempt to quantify the differential climate impacts between 1.5 and 2.0 degree global mean warming. Finally, the authors attempt to make inferences on the importance of coupled atmosphere-ocean-sea ice feedbacks in 1.5 and 2.0 degree global mean warming worlds, by comparison of AGCM-only and slab-ocean (SO) versions of NorESM1-Happi.

Unfortunately I have major concerns and cannot recommend publication of this manuscript at this time, due to lack of clarity, lack of focus and the subject matter being potentially out of scope for ESD.

Reply to overall response:

We understand the concerns of Referee #1 (as well as Referee #2) about the paper's scope for ESD, as well as the apparent lack of focus. Before we propose how to deal with this, please take into account that the manuscript was actually submitted to be considered for publication in GMD. It was intended for inclusion under the special collection of articles on the Norwegian Earth System Model (NorESM). This paper's version of the NorESM (NorESM1-Happi) is an update of NorESM1-M used for CMIP5. The AGCM-version of the model has been used to contribute to multi-model investigations in the HAPPI project, with AMIP-type experiments that address differences between a 1.5 degrees and a 2.0 degrees warmer world than the pre-industrial (1850). Our intentions with the present paper, when submitting it to GMD, were

- 1. to validate the fully coupled NorESM1-Happi which employs the atmosphere and land components that are used in the experiment in the HAPPI project, e.g. by comparing its performance to NorESM1-M, and
- 2. to apply a slab-ocean version (NorESM1-HappiSO) which includes the sea-ice model from the fully coupled NorESM1-Happi, to investigate how the polar amplification of the temperature signal may change the modelled difference between a 1.5 and a 2.0 degree world. In addition, results from the RCP2.6 and RCP4.5 runs with NorESM1-Happi complement the discussions, although these scenarios are not targeting the two temperature increments per se.

Obviously, the visibility of point (2) has suffered due to the lengthy and detailed discussions

over the first 4 sections. Point (1) was emphasized in the manuscript when designing it for GMD rather than ESD. The topic editor of GMD who was assigned for the paper almost a month after the submission, chose to reject the paper for GMD and recommended to transfer it to ESD. With the publication constraints for papers to be referred to in the scheduled Special Report from IPCC in mind, we chose to follow the recommendation without much discussion, although we suspected that the profile of the manuscript could produce confusion with the ESD editors and reviewers.

Now, with both the referees' comments in mind as well as our own concern about the paper's

profile for ESD, we propose to restructure the manuscript as follows, hoping to achieve stronger focus, better clarity, and a more suitable profile for ESD.

- We propose to reduce the amount of material that documents the standard validation of NorESM1-Happi as a global climate model, mainly by compressing much of the information given in the present text into tables, and collect them in a separate set of "supplementary material". A considerably shortened text will still be kept in one sub-section of the main manuscript, where the model design is described. Some validation results that are relevant for the polar amplification topic will be kept in the main text (feedback analysis, extratropical storm-tracks and blocking, sea-ice and aspects concerning the freshwater cycle).
- A clearer presentation of the reasons for running the slab ocean experiments to complement the AMIP-type experiments will be given, and in particular the replacement of a prescribed sea-ice cover in the AMIP-runs with a fully dynamic-thermodynamic sea-ice in the slab ocean model runs.
- We are underway with a selection of simulations with the fully coupled NorESM1-Happi model, which target the 1.5 degree and the 2.0 degrees warmer world (compared to 1850 pre-industrial). This will better complement the AMIP and SO-experiments, with specific focus on polar amplification than the runs based on RCP2.6 and 4.5 that are now included.

All in all, this should considerably strengthen the paper's focus on polar amplification, while still including the necessary results in support of the model validation as a secondary item.

With this, we believe the paper will be suitable for ESD, and in particular to be included in the

special issue about The Earth system at a global warming of 1.5°C and 2.0°C (https://www.earth-syst-dynam.net/special_issue909.html).

We have made substantial changes to the paper. It has been restructured and now is now focused on how ocean and sea-ice feedbacks affect the response to 1.5 degree and 2.0 degree warming. Details are given below:

• Following the suggestion of reviewer 2, we have moved the standard validation of NorESM1-Happi against NorESM1-M, observations and reanalysis out of the main manuscript and into a Supplement. The material has furthermore been condensed

substantially. A very short summary of the most important results are kept in the model section in the main text. We now only include some validation results for fields that are relevant for the discussion in the main text. We have also shortened the section describing the fully coupled NorESM1-Happi.

- We have designed and carried out new simulations with the fully-coupled NorESM1-Happi for 1.5 degree and the 2.0 degrees warming targets (compared to 1850 pre-industrial). Results from these simulations are compared to results from the SO and AMIP experiments, and replace the results from RCP2.6 and RCP4.5 in relevant cases.
- We have updated the slab-ocean simulations using Q-fluxes that are are calibrated using SST increments (difference between the PD SSTs and the 1.5C and 2.0C warmer SSTs) from the fully-coupled runs instead of the SST increments from the AMIP forcing data. The reason is that the increments from the fully-coupled simulations are more consistent with the model climate.
- We have added a new section dedicated to describing the set-up of the 1.5 degree and 2.0 degree warming experiments carried out with the AMIP version, the SO version, and the fully coupled version of NorESM1-Happi. This new section should make the differences between the AMIP, SO, and fully-coupled experiment more clear, but also accounts for the motivation behind running the fully-coupled and the SO experiments to complement the AMIP-type experiments.
- In addition to adding results from the fully-coupled simulations to compliment the already existing analysis of the AMIP and SO experiments, we now also consider changes in the upper-level and lower tropospheric equator-to-pole temperature gradients and changes in the upper-level and lower-level extratropical storm tracks. We also discuss how these results compare with the multi-model analysis carried out by Li et al. (2018).

The manuscript is now more clearly focused on the role of ocean and sea-ice feedbacks and polar amplification, while still including the necessary results for model validation in a Supplement.

We believe the revised paper is now well suited for ESD.

General comments:

1)

While the focus of the manuscript is claimed to be the differential climate impacts of 1.5 and 2.0 degree global mean warming, it feels more like a model evaluation paper. Most text and figures are dedicated to the evaluation of NorESM1-Happi compared to the CMIP5 model NorESM1-M. While this is a useful and necessary exercise, I am not sure if ESD it the right platform to report on this.

Reply:

As mentioned in the overall response, the paper was written for and submitted to GMD, but the assigned topical editor thought it was better suited for ESD. We now realize that the title of the manuscript may have given the GMD topical editor the impression that the paper is not a model evaluation paper (our fault). Since ESD has a special issue on "The Earth system at a global warming of 1.5°C and 2.0°C", we decided to follow the suggestion of the GMD topical editor (even though we did expect that the ESD-referees probably would comment on this).

We are therefore prepared to restructure the paper, and propose to include major parts of the pure model validation in a "supplementary material", reduce its volume, and only keep a shorter discussion in a subsection of the main text. Instead, the discussion of the differences in climate between a 1.5 and a 2.0 degrees warmer world will be more prominent, with the role of polar amplification as the paper's focus.

See changes listed under reply to general comments above.

2)

Often times, there is lack of clarity in the text. Perhaps the first-author is not a native English speaker, but this should be addressed as it makes it challenging to follow the discussion. Examples:

Reply:

We intend to seek help to copyedit the updated (or the very final) version of the manuscript by a native English speaker.

The manuscript has been largely re-written, both to change the scope from a model validation paper to a science paper and to improve the clarity of the text. The Supplement and parts of the manuscript have been proofread by a native English speaker.

p.3 I. 34 ('AA has been calculated with a strong response?),

Reply:

The sentence should read:

Even for the remote and regionally localized forcing caused by reduced European sulphate aerosols since the 1980s, the strongest amplification of warming is found in the Arctic (Acosta Navarro et al., 2016).

This sentence has been rewritten. See p3, L25-26.

p.11 I. 11 ('in reality?'),

Reply: Replace by "already", and in addition modify the entire paragraph to read:

NorESM1-Happi is a version of NorESM1-M with relatively minor updates. The most radical difference is a doubling of the horizontal resolution in the atmosphere and land models. As NorESM1-M is already thoroughly documented through CMIP5, this paper presents only selected features of the updated NorESM1-Happi.

This section has been moved to the Supplement and rewritten.

p. 11, I. 24 ('inaccurately calculated': was the calculation inaccurate or wrong?),

Reply:

"incorrectly" is the correct phrase.

This part of the text has been moved to the Supplement and rewritten.

p. 13, I. 6 ('calculated time-developments'),

Reply:

We propose to amend the sentence (if it will be used in the new manuscript) to become:

Fig. 5 shows the simulated time evolution from 1850 to 2100 of some of the quantities in Table 2.

Fig. 5 has been moved to the supplement (now Fig. S8) and the text describing it has been modified.

p. 14, l. 23 and Fig. 9 legend ('extension' -> 'edge'),

Reply:

will be changed to "edge".

Comment: "ice extent" will replace "ice extension". Unfortunately, in a few places the ice extent has been confused with ice area. Ice area takes into account the fractional ice-cover in any grid cell, while the ice extent measures the size of the entire domain where sea-ice is present. This will be corrected.

The second sentence in the Fig. 9 legend should read:

The solid black line shows the sea-ice edge estimated as the 15% iso-line of monthly sea-ice concentration in the OSI-SAF reprocessed data set (OSI-SAF, 2017).

This part of the text has been rewritten and moved to the Supplement along with Fig. 9. The second sentence in the legend of Fig. 9 (now Fig. S4) now reads: The solid black line shows the climatological 15 % concentration line for the same period from the OSI-SAF reprocessed data set (OSI-SAF, 2017).

p. 17 I. 29 ('describe' -> reproduce?').

Reply:

agreed. "reproduced" is better.

We use "reproduced" in the revised text. This part of the text has been moved to the Supplement (Figure S13).

Section 5.2 is really hard to follow:

Reply:

We are sorry for the sloppy and confusing language many places. This section will be re-written and expanded in a new manuscript.

This section has been rewritten to accommodate the new results and for clarity.

p. 22, I. 10 ('enhanced with the SO model compared to the AMIP': not sure what this means),

Reply:

see next point.

p. 22, I. 11 ('latter': not sure what this refers to).

Reply to both points:

Sorry! Yes, this is confusing. The sentences should read:

The PAF is considerably larger in the Arctic than in the Antarctic. Furthermore, the SO model produces stronger AA (by 18%) than the AMIP model.

This section has been rewritten to accommodate the new results and for clarity.

p.22 I. 15 (probably the tendency...: grammatically incorrect).

Reply:

see next point.

This part of the text has been rewritten.

p. 22, I 17-18 (unclear)

Reply to both points: The following amended text is proposed for the entire paragraph:

The SO model has a tendency to produce a colder winter climate than the AMIP model, consistent with the systematic cold bias in the fully coupled system. The differences in summer climate between the SO and AMIP simulations are much smaller. The SO model generally simulates less reduction in sea ice area than is prescribed in the AMIP simulations, with the exception of the NH summer response for 2.0 degrees - PD. These differences are also reflected in the temperatures and precipitation over land.

Reply: Please see the detailed replies above.

We suspect that missing clarity is also a consequence of the hectic situation caused by the deadline for submission on Nov. 1 st 2017 (for papers to be referred to in the IPCC's special report on 1.5 degrees). We will pay considerable attention to the clarity of language when updating the manuscript, and we intend to seek help by a native English speaker before finalization.

This part of the text has been rewritten to accommodate the new results and to make it more clear.

3)

The presentation should be improved. There are way too many numbers listed in the tables (only a small fraction is discussed in text). This is overwhelming and makes it hard for the reader to know which are the most relevant ones. I also suggest to have a more consistent lay-out: in evaluation of NorESM-happi, sometimes NorESM-M results are included, and sometimes not (e.g. Figs. 10 and 12). I suggest to always include NorESM-M results for consistency.

Also, the labelling in Fig. 13 is confusing, the x-axis in Fig. 14 is not legible (too small), and the different rcp's in Fig. 17b should have distinct colours.

Reply:

The many numbers are there to document basic properties in order to validate the model as a suitable tool for the study of processes in the climate system. We do not explicitly comment on every number in the text, although the overall model properties, e.g. concerning the simulation of a stable pre-industrial climate, climate sensitivity, and climate variability and change, are discussed. Such numbers are essential in a climate model validation. We understand, however, that a full model validation was not to be expected in an ESD-article, or from the title of the paper. We believe this will be considerably improved by moving many of the tables to a "supplementary material".

We think it can be a good idea to include NorESM1-M results in Figs. 10 and 12 as well, to the extent that they bring substantial additional information (there are already many figures). The details mentioned for Figs. 13, 14 and 17b will be taken into account. The figures will in any case be changed when re-organizing the manuscript.

To limit the number of figures in the Supplement, we prefer not to add versions of Figures 10 and 12 (now Figures S3 and S10) for NorESM1-M. We have added references to the figures in Bentsen et al. in the relevant captions to make it easier for the reader to locate the corresponding plots for NorESM1-M.

We have changed the labeling of Figure 13 (now Figure S11).

We have increased the font height of the labels for the x-axis in Figure 14 (now figure S12).

Figure Fig 17b has been updated and now show the time-evolution of the fully-coupled 1.5 degree and 2.0 degree warming experiments instead of the RCPs. We have kept the colors as we feel that adding more colors makes it more confusing.

4)

From the text I cannot derive what the major scientific advance is from the comparison between the AMIP and SO results. Aren't the SO sea ice and SST fields meant to mimic the AMIP sea ice and SST fields? Does the fact that there are differences between the SO and AMIP sea ice and SST fields mean that the 'target' SST and sea ice fields are not achieved in the SO runs? I'm concerned that this mismatch prevent a clean comparison of the climate impacts in the SO model versus the AMIP runs.

Reply:

We realize that the description of the differences between the SO-experiments and the AMIP-experiments is insufficient. We will expand the text accordingly as indicated below.

The referee is, quite understandably, concerned about the fact that the SO-model simulates sea-ice properties (and therefore also to some extent SSTs) which deviate considerably from the AMIP experiments, even though the calibration of the slab

ocean is constructed to mimic the prescribed ocean state used for the AMIP experiments.

Does this imply that the SO-model is profoundly wrong, and is not valid for the intended study of polar amplification? **The reply is no.** The explanation follows.

The SO model includes interactive sea-ice and snow processes which are missing in the AMIP-runs. In principle, the mean climatology for the SSTs used in the AMIP experiments should be reproduced by the SO-model experiments, provided the sea-ice properties (cover, thickness, concentration) are the same in the two setups. However, when calibrating the SO-model, we do not directly control the sea-ice properties. We define deep-ocean heat-fluxes (Q) in the SO in order to approximate the SSTs used in the AMIP experiments, but there is no similar relaxation of the sea-ice in the SO model towards the sea-ice which is prescribed in the AMIP runs. The Q-fluxes will indirectly constrain the SO-calculated sea-ice to some extent, but there is no reason why this sea-ice should be identical to the AMIP sea-ice.

We feel that this is a strength of the SO runs compared to the AMIP runs. The sea ice concentration prescribed in the present day AMIP experiment are based on satellite observations, but other properties of the ice and snow cover are more arbitrarily set as described in detail in Mitchell et al. (2017; doi:10.5194/gmd-10-571-2017). For example, the sea-ice thickness is 2 m over the entire Arctic sea ice extent and 1m in Antarctica, and the snow-cover is allowed to become unrealistically thick in many places. We will extend the description of this in an updated manuscript, thus clarifying why the SO-model with its more realistic feedback processes associated with sea-ice and snow cover, is useful.

The heat isolation and radiation properties of sea-ice thickness and snow cover are particularly important for e.g. the Arctic temperatures over extended winter seasons. In the SO-model, these properties are allowed to vary in accordance with physical processes represented in the sea-ice model, while the sea-ice thickness is constant in the AMIP runs. Hence, the SO-simulation should have a more realistic – or at least more physically based – representation of feedbacks that influence the Arctic amplification in ways not included in the AMIP runs.

Concerning SSTs, the SO-model should produce results close to the prescribed SST-fields used in AMIP, except in regions directly influenced by the sea-ice model results. By construction, the relaxation we use to define the Q-fluxes secures that the SO-model results hit the same global temperature targets as the AMIP runs. Deviations originate from the fact that the atmosphere in the SO-model "sees" a different sea-ice and snow cover than in the AMIP runs.

We will provide a comparison of the simulated sea-ice thickness changes in the 1.5C and 2.0C experiments that are underway with the fully coupled NorESM1-Happi model with those of the SO experiments, although the fully coupled experiments are not bias-corrected for present-day such as the SO-experiments.

In conclusion, and this will be better presented in an updated manuscript, the results from the SO-model should produce an Arctic amplification of the temperature response which differs from the AMIP-results. The differences are dominated by the changes in the sea-ice and snow cover in the SO-model with some regional influence on the SSTs. Otherwise, the contributions to AA from SST-feedbacks directly, should be included in the SST-fields prescribed for the AMIP experiments.

We have added a section describing the warming experiments carried out with the AMIP, SO, and fully-coupled version of NorESM1-Happi to make it more clear how the experiments with the different model versions compliment each other. We have also improved the section describing the SO model.

We now provide results for changes in sea ice for both the SO and the fully-coupled model.

Other comments:

1) p. 3, l. 5:

I'm not familiar with the term 'temperature ceiling', and find it a bit misleading. Better to use 'temperature target', as the temperature references in the Paris Agreement are generally believed to apply to long-term averages, and not to the maximum as the word 'ceiling' might suggest (see: doi: 10.1002/2017GL075612)

Reply:

We were concerned with the term "target". The target is in principle an upper bound of the global temperature change, not a target that one wishes to hit. However, if "ceiling" is a misleading word, we can change to "target", while also explaining that an under-shoot should not be considered a miss.

We do not use the term "temperature ceiling" in the revised manuscript.

2) Fig. 1: For a cleaner comparison between models and observations I suggest to sample the model only at the locations where and times when observations were made.

Reply:

In fact, model output was already sampled in grid volumes containing the observation points, and for dates in each of the ten years of the PD-period that coincide with the observation dates. Since the model is run in climate simulation mode without data-assimilation (or nudging), we cannot compare the exact observation times. We will update the figure legend to make this clear. The figure will be moved to the supplementary material section. We have updated the figure legend to make this clear. The figure has been moved to the Supplement (now Fig. S1).

3) p. 7, line 9: Do you really mean Fig. 11?

Reply:

No, it should be Fig. 9 (now Fig. S4). This part of the text has been rewritten and moved to the caption of Fig. S4.

4) p. 8, l. 6-8: 'changes in the major elements ...': unclear, please elaborate.

Reply:

More specifically: GHG, aerosols and land-use

This paragraph has been removed as we now dedicate a section to the set-up of the 1.5 degree and 2 degree warming experiments.

5) p. 10, l. 21:

what is the ACCESS version of the model? Please keep naming conventions/references to model versions consistent to avoid confusion

Reply:

Sorry about this. This sentence remained from an early version of the manuscript. It should read:

Before the piControl, the 1-degree version of the atmosphere and land model described in Seland and Debernard (2014) was spun up over 300 years, starting from....

We refer to the different model versions in a more consistent way in the revised manuscript.

6) p. 12:

maybe I missed it but did you discuss the trends in the AMOC and Drake Passage transport? These seem to be the quantities with the largest drift

Reply:

The trends in both quantities are mentioned on p12, I 3-11, and compared to the trends in the NorESM1-M model. They are not discussed in depth, and we will add a sentence in the manuscript stating that while the apparent drift in those indices indicates that the deep ocean is not in perfect balance, we do not think is a serious issue our study. The magnitude of the trends are comparable to those in NorESM1-M (Bentsen et al, 2013).

We have shortened this part of the text and moved it to the Supplement (caption of Table S3). We mention that the magnitude of the trends are comparable to those in NorESM1-M (Bentsen et al, 2013).

Otherwise, the strong AMOC leads to heating of the deep oceans and leaves less energy for surface heating and evaporation. We will add a figure in the planned supplementary material, which documents how heat penetrates into the deep ocean during simulated RCP scenario projections with NorESM1-Happi. On p13, I 14-30, we hypothesize a connection between the strong AMOC and the simulated speed of the atmospheric freshwater cycle, as well as the under-estimated cloud cover that can be related to the horizontal resolution of the atmosphere.

We have added figures in the manuscript showing the AMOC (Figure 4) and how heat penetrates into the deep ocean (Figure 3) in the fully-coupled simulations of the 1.5 degree and 2.0 degree warmer worlds.

7) p. 20, l. 21-22:

Very confusing to first list the numbers in the brackets and not mentioning what they mean until 5-10 sentences thereafter.

Reply:

Agreed. We will mention this in the beginning of the paragraph.

We have moved Table 6 and the paragraph discussing it to the Supplement (Table S6). The text has been rewritten for clarity and shortened.

p. 21, l. 1-10:

I don't think this discussion is accurate. Sanderson et al. (2017) used an emulator to construct emissions pathways that would lead to a 1.5 and 2.0 degree warmer world. Those emissions pathways were then prescribed to a coupled atmosphere-ocean model.

This is not a 'simplified method' as the authors suggest.

Reply:

Of course, this is wrongly stated in the paper and will be corrected.

We have corrected this in the revised manuscript.

Also, lines 2-4 are a bit misleading as well. It is now well established that the equilibrium climate response is determined by the accumulated carbon emissions. This implies that, in

reality and in climate models that include a carbon cycle, switching off emissions would lead to a rapid stabilization of global mean surface temperature (the delay in warming associated with ocean thermal inertia would be balanced by a decrease in greenhouse gas concentrations, see e.g. doi:10.1038/ngeo1047). My point is that it may not be as hard as the authors suggest to employ fully coupled atmosphere-ocean models to quantify the climate impacts of 1.5 and 2.0 degree global warming.

Reply:

We appreciate the referee's point.

We are now underway with some experiments with the fully coupled NorESM1-Happi model, targeting a 1.5 and a 2.0 degrees warmer world than the model simulated pre-industrial control run for 1850. These experiments are not emission-driven for GHGs, hence we do not base our calculations on the principle from Gillett et al. (2011, doi:10.1038/ngeo1047). Instead, we have designed GHG-based forcing with temperature targets based on amendments to RCP2.6 and 4.5, and our own estimates of the model's climate sensitivity.

We will not have time to produce an ensemble of model projections comparable to that of Sanderson et al (2017), but hopefully around 150 simulation years per temperature target. Statistics will replace those included for the RCP2.6 and RCP4.5 in the present manuscript, and thus better complement those of the SO- and AMIP-experiments.

We have conducted fully-coupled simulations for 1.5 and 2.0 degree warmer worlds. The set-up of the experiments is presented in Section 2.2 and the results are discussed alongside the results from the SO and AMIP experiments in the result sections.

9) p. 21, l. 13: which 'forcing data', please specify

Reply:

Forcing data here are prescribed atmospheric levels of greenhouse gases (GHGs). Details are given in Mitchell et al (2017).

This is specified in the new section describing the set-up of the experiments.

10) p. 21, l. 31: please give evidence that after 45 years a new equilibrium is indeed reached

Reply:

Below is a figure which shows the development of global temperature in the SO

simulations for present day 2005-2016 (SO-PD, black), the 1.5 (SO-15, blue), and the 2.0 (SO-20, red) degrees warmer world than pre-industrial. The numbering of years are arbitrary. All time series show considerable auto-correlation and multi-decadal variability. The weak, apparent trends from year 2161 to 2250 are not significant when data are sub-sampled based on the effective number of independent observations.



11) p. 22, l. 6: What is meant with 'single projections'?

Reply:

Single projections means that there is no ensemble of projections (e.g. with ensemble members starting from different initial states), but only one single estimate. We propose to remove "single".

We do not use the term single projections in the revised manuscript or in the Supplement.

12) p. 22, l. 7:

Table 7 does not show that the temperature targets (1.5 and 2.0 degrees above pre-industrial) are hit, but only show the temperatures relative to PD. What are the temperature relative to preindustrial?

Reply:

This is a consequence of the HAPPI-design (Mitchell et al, 2017), for which calculations are based on present-day (PD; 2006 – 2015) which is estimated independently at 0.8 o C above the pre-industrial. Since neither the SO- nor the AMIP-experiments are run explicitly for the pre-industrial situation, we will not present the pre-industrial numbers for these two sets of experiments. For the fully coupled experiments, however, we will add numbers for the pre-industrial situation.

The focus of the manuscript is on the changes in the 1.5 and 2.0 warming experiments relative to the PD climate as in the HAPPI project. This now clearly

stated several times in the revised manuscript. The PD climate is estimated to be 0.8 degrees warmer than the pre-industrial in the AMIP experiments. This is also now clearly stated in the manuscript. We show the time-evolution of the near-surface temperature response in the coupled warming experiments relative to pre-industrial conditions in Figure 2.

13)

As noted above, I found section 5.2 really hard to follow. What is particularly confusing is p. 22, I. 25-34. I think the sea ice area in the AMIP runs are compared to observations.

Questions I have here are:

1) why is this comparison not presented earlier,

2) why are there any differences if the prescribed data is based on observations?

3) why is sea ice extent in Table 4 compared with sea ice area in Table 8?

Reply:

There are obvious reasons for a reader to be confused here. In Table 4, sea-ice extent is presented, and not sea-ice area. In winter, the difference between area and extent (area being smaller than extent) may not be considerable, but still the comparison is not accurate. Nevertheless, we believe that the comparison indicates that there are some errors in the SO-model and probably also in the fully coupled NorESM1-Happi. The text in the paragraph will be reformulated and its emphasis will be reduced. This should have been done during the quality check before submission, and for this we apologize.

The text in the paragraph has been reformulated. Table 4 (now Table S5) has been moved to the Supplement and we no longer compare these results to Table 8 (now Table 4).

Anonymous Referee #2

Received and published: 30 January 2018

The authors look at a range of different versions of the NorESM1 model, and consider how those models hold up against reanalysis. They consider changes in many modes of variability, specifically related to key regional changes. Overall the paper was not what I

expected, from the title I expected the paper would be about NorESM1-Happi, Paris Agreement and Arctic Amplification. Very little of this was even mentioned until Figure 17!

As it stands the paper is a model description/validation paper, and a science paper. The science is completely lost due to the first part. Due to this, and a number of other major concerns, I recommend substantial corrections.

Major concerns

1.

In my view, the paper needs to be split into two. A paper focussing on the Arctic Amplification differences under Paris would be very welcome. So one suggestion is to put everything up to Figure 17 in online material, and just start the paper from there. Any reader that comes to this paper due to the title will be otherwise be completely lost in details of various models, and it will not be a productive read for them. More material would be needed for the science part though (see comments below). Alternatively, you could make this a model development only paper.

Reply:

As mentioned in the response to Referee #1, we agree that for the publication in ESD, the paper should be considerably restructured, and we propose to condense and move considerable parts of the sections that address pure validation of the NorESM1- Happi model, into a "supplementary material". In the main text, we will only keep those parts, which are directly relevant for the discussion of the difference between a 1.5 and a 2.0 degrees warmer world than pre-industrial, emphasizing the polar amplification of the temperature response.

Tentatively, we consider moving the following to supplementary material:

Tables 1, 2, 3, 4, and 6

Figures 1, 3, 4, 5, 6, 7, 8, 10, 11, 12, 15, and 16.

(Some of these figures will also be considered removed.).

There are also some numbers in the present main text that probably will be summarized in a new table. The text belonging to these items, which now fills up several pages in the manuscript will be compressed into one sub-section of the model description chapter. The text in the supplementary material will predominantly be written as extended Table headings and Figure legends.

What may remain in the main text are the discussions on Arctic amplification and the design of external forcing for reaching the given temperature targets with the fully

coupled NorESM1-Happi model. Furthermore, we will keep the discussion on the representation of sea-ice (Figure 9), extratropical cyclone activity (Figure 13) and blocking (Figure 14). Thus, the paper will be shorter and much more focused, while still documenting important aspects of NorESM1-Happi as a valid global climate model in the supplementary material.

We have shortened and moved the part of the paper concerned with the the validation of NorESM1-Happi against NorESM1-M, observations and reanalysis a Supplement. The paper is now focused on the role of ocean and sea-ice feedbacks under global warming of 1.5 and 2.0 degrees. We have expanded our analysis to include analysis of meridional temperature gradients and storm tracks at different levels. We have also added fully coupled simulations of the two warming targets and now present results from these simulations alongside results from the AMIP and SO simulations. The changes are listed in more detail under the reply to reviewer 1's overall response.

2.

The title makes it seem that NorESM1-Happi is the main model here, but actually it is not, the SO version is used the most, and the –M and –AMIP versions are used equally as much. I often got confused about which one was being used, as the paper jumped around a fair bit. It was not until half way through that I realised that the Happi version of the model did not have prescribed SSTs (as HAPPI is synonymous with prescribed SSTs).

Reply:

This confusion should be considerably reduced after the proposed paper reorganization. The definition of model versions (the AMIP, the slab-ocean (SO) and the fully coupled NorESM1-Happi) will be made already in the introduction. The NorESM1-M model is the CMIP5-version published in 2013, and is used only for documenting improvements in the NorESM1-Happi, and will thus be used for comparison in the supplementary material, and will not be prominent in the main text.

We consider to possibly use the name NorESM1-HappiCPL for the fully coupled model version, while NorESM1-HappiSO will be kept for the slab ocean version and NorESM1-HappiAMIP for the AGCM version.

This should be considerably less confusing in the revised manuscript, as we refer to the different model versions in a more consistent way. Also all the results for NorESM1-M are now in the Supplement, so the main text is now only concerned with the fully coupled, SO and AMIP versions of NorESM1-Happi. We have added a new section describing the set-up of the warming experiments with the different model versions, so it should be clear that the AMIP version has fixed SSTs and sea ice

3.

I was hoping to see more of a connection to Arctic Amplification here. Such as more of a

focus on latitude temperature gradients, changes in wave characteristics associated with this, and then relating this to blocking etc. This link was missing, and AA just seemed to be a 'hot topic' term. The authors should look at the recent work by Screen on this topic.

Reply:

It is mentioned in the introduction (p.3, I.26-20) that the intention of this paper's discussion of Arctic amplification is not to specifically address the potential impacts on planetary wave amplitudes and weather persistence.

We agree to extend the list of references to include publications by Screen, and we will mention the recently approved Polar Amplification MIP for CMIP6 (to which the NorESM-group plans to contribute). We should also mention that Figure 21 actually shows that there are more statistically significant changes in blocking occurrence in the NH in the slab ocean experiments than in the AMIP experiments. The AMIP experiment is not designed to produce a realistic Arctic amplification due to the way the sea-ice is prescribed (see our reply to pt. 4 of referee#1).

However, we disagree with the referee's statement that "AA just seemed to be a 'hot topic' term". Arctic amplification of the global warming is well established as a temperature signal, which deserves attention in its own right. Indeed, when discussing the difference between a "1.5 degree world" and a "2.0 degree world", this may prove to be the difference between an ice-free summer Arctic or not (e.g. Sanderson et al, 2017; and the results shown in our Figures 22 and 23), because of the amplified temperature response in the Arctic.

We have strengthened the results sections and now consider results from 1.5 degree and 2.0 degree warming experiments with AMIP, SO, and fully-coupled versions of the model. Our focus is on how ocean and sea-ice feedbacks in the SO and fully-coupled models affect climate response under 1.5 degree and 2.0 degree warming, i.e. how the AMIP experiments differ from the ones with ocean models. We have changed the title to reflect this.

We have added results showing changes in the upper-level and lower-level equator-to-pole temperature gradients in the Northern Hemisphere along with changes in the baroclinic-wave activity at similar levels. These results are moreover compared to the multi-model analysis in Li et al. (2018). The results for blocking activity mostly inconclusive due lack of consistency between the different models, but also the low statistical significance of the changes.

We have added references to studies by Screen and co-authors.

4.

A more comprehensive analysis seems to have already been done by the authors, in Li et al, (https://www.earth-syst-dynam-discuss.net/esd-2017-107/). Can the authors highlight what

their study adds?

Reply: While Li et al is a multi-model study on the response of selected features of the large-scale atmospheric dynamics entirely based on the HAPPI protocol of AMIP experiments, our paper's intention is to focus on the Arctic amplification of surface temperatures, which is not well represented in the HAPPI experimental set-up.

As emphasized in our reply to pt. 4 of referee#1, the reason for this is that some properties of the prescribed sea-ice in the AMIP experiments are not realistic. It is likely to misrepresent the Arctic surface temperature amplification, and therefore the differences between the 1.5 degree and the 2.0 degrees global warming in the Arctic, including sea-ice cover itself. There is a component of Arctic amplification in the HAPPI AMIP-experiments, but this is dominated by the prescribed SSTs and sea-ice concentrations in the 1.5 and 2.0 degrees warmer climate.

One consequence of this can be seen in the smaller response in NH extratropical storminess in our own AMIP results than in the results from both the slab ocean runs and the fully coupled NorESM1-Happi runs with RCP2.6 (Fig. 19). In pt.3 above, we have furthermore already mentioned the changes in NH blocking (Fig.21) and in Arctic sea-ice (Figs 22 and 23)

We have in the pipeline for the updated manuscript, longer simulations with the fully coupled model targeting the 1.5 degree and the 2.0 degrees warmer world.

We now compare our results to those in Li et al. (2018) both in the results sections and in the summary and disucssion. We also emphasize the difference between our study, which is focused on the NorESM1-Happi and now the AMIP experiments differ from similar experiments with active ocean components, and the study by Li et al., which is a multi-model study considering AMIP experiments from five different models (including the AMIP version NorESM1-Happi).

5.

I was very surprised by some of the differences between the SO model and the AMIP model. Surely the AMIP model will have smaller biases that the SO model (e.g. some Scaife papers could be referenced). It is not always clear that this is the case.

Reply:

(It would help if the referee#2 more specifically pointed to which results he/she is surprised to see.) The way we have calibrated and relaxed (i.e. bias corrected) the slab-ocean model for present-day conditions, we expect only small differences between the AMIP and SO model when comparing with e.g. re-analysed data for present-day conditions. This was also found for NH storminess and blocking (Figs. 18 and 20), and these results encouraged us to further employ the SO-model for the purpose of Arctic amplification on the target scenarios.

A more traditional way to calibrate the slab-ocean model (e.g. to study equilibrium climate sensitivity as mentioned in section 3.1), is likely to produce larger biases than in AMIP runs, where SSTs are prescribed from observationally based data.

The reviewer is correct, the AMIP model does have smaller biases than the SO model. This is for instance clear from the new figures showing the biases in the near-surface temperature (Figure 5) and baroclinic wave activity (Figure 11) and we point this out in the revised text.

At this stage, I am not sure minor comments are useful.

The "NorESM1-Happi" used for evaluating the role of ocean and sea-ice feedbacks under global warming of 1.5 °C and 2 °C

Lise S. Graff¹, Trond Iversen^{1,2}, Ingo Bethke³, Jens B. Debernard¹, Øyvind Seland¹, Mats Bentsen³, Alf Kirkevåg¹, Camille Li⁴, Dirk J. L. Olivié¹

⁵ ¹Norwegian Meteorological Institute, P.O.Box 43, Blindern, 0313 Oslo, Norway

²Dep. of Geosciences, University of Oslo, P.O. Box 1047 Blindern, 0315 Oslo, Norway

³Uni Research Climate, Bjerknes Centre for Climate Research, P.O. Box 7810, 5020 Bergen, Norway

⁴Geophysical Institute, University of Bergen, Bjerknes Centre for Climate Research, P.O. Box 7803, 5020 Bergen, Norway

10 Correspondence to: Lise S. Graff (<u>lise.s.graff@met.no</u>)

15

Abstract. Differences between a 1.5 K and a 2.0 K warmer climate than 1850 pre-industrial conditions are investigated using a suite of uncoupled (AMIP), fully coupled, and slab-ocean experiments performed with the NorESM1-Happi, an upgraded version of NorESM1-M. The AMIP-type runs with prescribed SSTs and sea ice from the NorESM1-Happi were provided to a multi-model project (HAPPI, <u>http://www.happimip.org/</u>). This paper compares the AMIP results to those from the fully coupled and the slab-ocean version of the model (NorESM1-HappiSO) in which SST and sea ice are allowed to respond to the warming, focusing on the role of ocean and sea-ice feedbacks and Arctic amplification of the global change signal.

The fully coupled and the slab-ocean runs generally show stronger responses than the AMIP runs in the warmer

worlds. Arctic amplification of the change in near-surface temperature is larger in the runs with active ocean models. Compared to the AMIP runs, the Arctic polar amplification factor is 54 % and 27 % stronger in fully coupled and SO 1.5 K warming runs relative to the present day climate, and 46 % and 19 % stronger with the additional 0.5 K warming. The low-level equator-to-pole temperature gradient consistently weakens more between the present-day and the 1.5 K warmer climate in the experiment with active ocean components. The magnitude of the upper-level equator-to-pole temperature gradient increase in a warmer climate, but is not systematically larger in the experiments with active ocean components. Implications for storm-tracks and blocking are investigated. There are considerable reductions in the Arctic sea-ice cover in the SO model; while ice-free summers are rare under 1.5 K warming, they are estimated to occur 18 % of the time under 2.0 K warming. The fully coupled model does not however reach ice-free conditions as it too cold and has too much ice in the present-day climate.

1 Introduction

5

15

In *The Paris Agreement*, the parties to the United Nations Framework Convention on Climate Change (UNFCCC) established a long-term temperature goal for climate protection of "holding the increase in the global average temperature to well below 2 °C above pre-industrial levels and pursuing efforts to limit the temperature increase to 1.5 °C above pre-industrial levels, recognising that this would significantly reduce the risks and impacts of climate change" (UNFCCC, 2015). This has triggered considerable attention from climate modelling groups and researchers alike (e.g. Hulme, 2016; Peters, 2016; Rogelj and Knutti, 2016; Mitchell et al., 2016; Anderson and Nevins, 2016; Boucher et al., 2016; Schleussner et al., 2016; and the special issue of

the electronic journal *Earth System Dynamics*: https://www.earth-syst-dynam.net/special_issue909.html). The

10 Special Report from the Intergovernmental Panel on Climate Change (IPCC) is to presented in October 2018 (<u>http://www.ipcc.ch/report/sr15/</u>).

In addressing differences in the climate impacts of the 1.5 K and 2 K global warming targest (we use the word "targets", although "upper bounds" would be more correct), there are two basic weaknesses of the available climate projections from the Coupled Model Intercomparison Project (CMIP) as reported in the assessment reports from the IPCC. There is a small body of research assessing impacts of 1.5 K warming compared to that for higher emission scenarios (James et al., 2017). The CMIP simulations are moreover generally designed on

- the basis of development scenarios that give rise to a certain top-of-the-model-atmosphere (TOA) radiative forcings, rather than selected temperature targets. Because different models simulate different responses of global, near-surface temperature to a given TOA radiative forcing, new types of model simulations are
 necessary to provide a scientifically-based evaluation of climate statistics for specific temperature targets.
- Under the acronym *HAPPI* (Half a degree additional warming, prognosis and projected impacts, http://www.happimip.org/), Mitchell et al. (2017) provided an experimental framework for model simulations of the present day climate and climates that are 1.5 K and 2.0 K warmer than the pre-industrial. The experiments are similar to those under the Atmospheric Model Inter-comparison Project (AMIP) protocol, employing active atmosphere and land components from state-of-the-art coupled ESMs and prescribed seasurface temperatures (SST) and sea ice. A multi-model ensemble with several hundred members was produced, enabling robust statistics for flow changes and rare events (e.g. Baker et al., 2018; Barcikowska et al., 2018; Li et al., 2018; Liu et al., 2018; Senerivatne et al., 2018; Wehner et al., 2018).
- Using a different approach, Sanderson et al. (2017) developed and applied an emulator to arrive at forcing scenarios that would produce global warming of 1.5 K and a 2 K above the pre-industrial levels in a model simulated stable climate. They carried out century-scale ensemble simulations with the fully coupled Community Earth System Model version 1 (CESM1; Hurrell et al., 2013). One striking result from this study is the strong increase in the probability of having an ice-free Arctic Ocean in the summer with the additional 0.5 K warming (the difference between the 1.5 K and 2 K warming scenarios). This aspect of the response to

the 1.5 K and 2.0 K warming was not evident in the HAPPI experiments because the sea ice is prescribed, but will be further addressed in the present paper.

We use various configurations of the Norwegian Earth System Model, *NorESM1-Happi*, which is an upgraded version of the NorESM1-M used in CMIP5 (Bentsen et al., 2013; Iversen et al., 2013, Kirkevåg et al., 2013).

- 5 The upgrades include double horizontal resolution and improved treatment of sea ice. The model was previously run in AMIP mode (*NorESM1-HappiAMIP*) to contribute a large ensemble of simulations to HAPPI. In order to study the role of feedbacks associated with the ocean and sea-ice, we here provide fully coupled simulations targeting quasi-sustained global warming levels of 1.5 K and a 2 K above pre-industrial levels. The forcings are constructed on the basis of those from the representative concentration pathway
- 10 scenarios (RCPs; van Vuuren et al., 2011) corresponding to an increased radiative forcing of 2.6 W m⁻² and 4.5 W m⁻² by the end of the 21st century (RCP2.6 and RCP4.5), but with important changes to the time evolution of the CO₂ concentrations. We also use a configuration where the full ocean model is replaced by a thermodynamic slab-ocean (SO) model (*NorESM1-HappiSO*). This configuration is applied as an intermediate option between the fully coupled (CPL) and the AMIP configurations, in order to partly correct for temperature biases in the CPL simulations, but still allowing for SST and sea ice feedbacks.

The role of Arctic amplification for specific warming levels (Arrhenius, 1896; Manabe and Stouffer, 1980, Holland and Bitz, 2003, Feldl et al., 2017) is relevant for the consequences of the Paris agreement. This is primarily due to the associated in-situ changes in the sea-ice and snow-cover, but also due to the potential triggering of irreversible feedbacks, such as changes in mid-latitude weather patterns and variability (Francis and Vavrus, 2012; Screen and Simmonds, 2013; Cohen et al., 2014; Screen, 2014; Barnes and Polvani, 2015; Screen and Francis, 2016; Screen, 2017a,b; Vihma, 2017; Screen et al., 2018; Cournou, et al., 2018).

20

Arctic amplification is predominantly driven by a positive regional lapse-rate feedback (negative at lower latitudes) in winter and a positive albedo feedback in summer (Winton, 2006; Pithan and Mauritsen, 2014). While the amplitude and pattern of Arctic amplification varies between models, it is nevertheless a robust

- 25 response to global warming. Even the remotely localized forcing caused by reduced European sulphate aerosols since the 1980s produces maximum warming in the Arctic (Acosta Navarro et al., 2016). Under the CMIP6 protocol, a Polar Amplification Model Intercomparison Project (PAMIP) is endorsed (Smith et al., 2018).
- In this paper, we focus on the Northern Hemisphere (NH) climate response to global warming of 1.5 K and
 2 K above pre-industrial levels, and the role of SST and sea-ice feedbacks. This includes changes in Arctic sea ice, midlatitude meridional temperature contrasts for different heights, and the storm tracks. We also consider blocking, although its representation in rather coarse resolution climate models is known to be of mixed quality (Dawson et al., 2012; Davini and D'Andrea, 2016; Woolings et al., 2018).

Section 2 describes the experiments in this paper. Section 3 provides an overview of the NorESM1-Happi, emphasizing the changes since NorESM1-M. Section 4 provides a description of the slab-ocean version NorESM1-HappiSO. Results are presented in Sect. 5–8. A summary and discussion are given at the end in Sect. 9. A Supplement to the paper contains an extensive validation of NorESM1-Happi in line with the CMIP5 protocol.

2 The 1.5 K and 2.0 K warming scenarios

5 2.1 The AMIP experiments

10

The "AMIP experiments" are those performed with NorESM1-Happi for the multi-model HAPPI project. The target of the experiments is to investigate the regional impacts of global warming under stabilisation scenarios that are 1.5 K and 2.0 K warmer than the 1850 climate. The three large ensemble experiments are: the present decade (PD; 2006–2015), a climate that is 1.5 K warmer than the pre-industrial (1850) climate, and a climate that is 2.0 K warmer. We refer to these as the AMIP-PD, the AMIP-15, and the AMIP-20 experiments.

- Designing a coupled model experimental protocol for 1.5 K and 2.0 K warming targets requires determining forcing conditions that will produce the target global-mean temperature change, and other characteristics of the warmer climate state. The same forcing conditions may however produce different temperature responses in different models. The CMIP5 models for instance display considerable spread in the near-surface
 15 temperature (2-m temperature) response to RCP2.6. While the multi-model mean response is very close to 2.0 K, the spread across the 95-5% range is approximately 1.5 K (see Fig. 2 in Mitchell et al., 2017). Fully coupled models are moreover computationally expensive because they require centuries or longer to approach new equilibria after sustained shifts in the TOA radiation balance.
- The experiments in the HAPPI project were therefore run with prescribed SSTs and sea ice. This constrains the climate state and makes it computationally feasible to run large ensembles. The experimental set-up resembles the AMIP protocol, thus we refer to the version of the NorESM1-Happi that follows the HAPPI protocol as NorESM1-HappiAMIP.

The construction of the input data for the HAPPI experiments is described in detail by Mitchell et al. (2017). The main points specific to our set-up are listed below:

In the AMIP-PD experiment, the SST and sea-ice fields are based on observations (Taylor et al., 2012).
 Sea-ice thickness is fixed at 2 m in the NH and 1 m in the Southern Hemisphere (SH). Anthropogenic greenhouse gas (GHG) concentrations (including CO₂, CH₄, N₂O, and CFCs), emissions of aerosols and their precursors, ozone concentrations, and land-use changes are taken from RCP8.5 for years 2006–2015, as it is common procedure to use RCP8.5 to extend the historical period beyond 2005 (van Vuuren et al., 2011).

In the AMIP-15 experiment, anthropogenic GHG and ozone concentrations, land-use and aerosols data are taken from RCP2.6 for year 2095. The SST increase relative to PD is the CMP5 multi-model mean difference between years 2091-2100 from RCP 2.6 and 2006-2015 from RCP8.5. Natural forcings are

as for PD. Sea-ice concentrations are estimates from a linear regression between observed anomalies of SST and sea ice (see Mitchell et al., 2017 p. 575 for details).

In the AMIP-20 experiment, the SST and sea-ice concentration differences are derived in a similar way, but using a weighted mean between RCP2.6 and RCP4.5 (0.41 for RCP2.6 and 0.59 for RCP4.5). The same weights are used for CO_2 (assuming a logarithmic relation). All other forcings are as for AMIP-15.

The NorESM1-HappiAMIP data set includes 125 ensemble members for each experiment, each of length 10 years (after a 1-year spin-up), giving 3750 years of data. To enable dynamical downscaling, output from 25 members of each experiment were stored with high temporal resolution. The data is available for download at http://portal.nersc.gov/c20c/data.html.

2.2 The fully coupled (CPL) experiments

5

10

15

One shortcoming of the AMIP-type simulations is that they neglect the effect of ocean and sea-ice related feedback mechanisms apart from the patterns included in the prescribed SST and sea ice fields. These feedbacks are particularly important in the Arctic, where albedo and lapse-rate feedbacks amplify the low-level temperature response relative to the global mean (Pithan and Mauritsen 2014; see also Fig. S14 in the Supplement).

To investigate the effect of such feedbacks, we have conducted 1.5 K and 2.0 K warming experiments (CPL-15 and CPL-20) with the fully coupled NorESM1-Happi. The forcings in the experiments are based on RCP2.6 and RCP4.5, but with important differences in the CO₂ concentration (Fig. 1). In CPL-15, the CO₂ concentration follows RCP2.6 from year 2000 to year 2095, after which it stays constant until year 2170, and then decreases following the pattern assumed in the original RCP2.6 from year 2095 onwards (i.e. the decrease is delayed 75 years compared to RCP2.6). In CPL-20, the CO₂ concentration follows RCP4.5 from year 2000 to year 2050, then stays constant until year 2170, after which it decays in the same fashion as CPL-15, but from the higher concentration level. The other GHGs and forcing-producing elements are as in RCP2.6.

25 The fully coupled present-day (CPL-PD) climate is represented by the 30-year time period 1991–2020 using output from the CMIP5 experiments carried out with NorESM1-Happi. We use the period 1991–2005 from three individual simulations of the historical climate (Hist1, Hist2, and Hist3; see Sect. 3.1 or Table S1 in the Supplement) and extend them with years 2006–2020 from three individual simulations of RCP8.5 (Sect. 3.1). Thus, CPL-PD, CPL-15, and CPL-20 are all sampled by 90 years of simulations with the fully coupled NorESM1-Happi.

The scenario runs CPL-15 and CPL-20 both start from simulation year 2005 of the Hist1 experiment. Figure 2 shows the change in near-surface temperature for Hist1 (1850–2005) and for CPL-15 and CPL-20 experiments (2006–2230) relative to the pre-industrial climate calculated under constant driving conditions valid for year 1850 (the piControl experiment, see Sect. 3.1 or Table S1). The global-mean temperature warms

rapidly between years 1960 and 2050, then the response flattens out over the next 150 years. In what follows, we study results from the 90-year periods 2111-2200 for which the mean temperature increase in CPL-15 and CPL-20 is 0.69 K and 1.15 K relative to CPL-PD (see discussion of Table 3 in Sect. 5.1).

- The experiments are, however, not entirely stabilized. By the end of the 22nd century, both CPL-15 and CPL-20 still have a positive radiative imbalance at the top of the model atmosphere (around 0.7 W m⁻², not shown) 5 and a positive heat flux into the ocean at depths below 200 m (Fig. 3). The net heat uptake in the upper ocean is, however, small at that point. The Atlantic meridional overturning circulation (AMOC) decreases with time over the first 100 years and is relatively stable over the last 150 years (Fig. 4).
- 10

Our fully coupled experiments differ from those in Sanderson et al. (2017), who first used a climate emulator to construct concentration scenarios, and then used these scenarios to produce stabilized 1.5 K and 2.0 K with the CESM1. The simulations presented in this study are far from reaching equilibrated climate states, but are quasi-stable over 90-year periods after spinning up for 100 years from present day. Full equilibration over several centuries is likely to produce different climate states (Gillet et al., 2011).

The slab ocean (SO) experiments 2.3

15 While results from the coupled simulations above will help us understand how 1.5 K and 2.0 K warming might manifest in the fully coupled earth system, CPL-15 and CPL-20 are not stabilized scenarios like the AMIP experiments. Moreover, Fig. 5 shows that the fully coupled PD experiments (panels a, d, g, and j) exhibits larger biases than the AMIP experiments (panels c, f, i, and l) relative to ERA-Interim (Dee et al., 2011) in all seasons. Prescribing the SSTs and sea ice to observationally-based fields constrains the climate in the AMIP-20 PD experiments, yielding smaller biases in the simulated climate. To be able to examine 1.5 K and 2.0 K warming experiments in a model which has smaller biases, but where the sea ice and SSTs are also free to respond, we have designed a slab-ocean configuration of NorESM1-Happi, NorESM1-HappiSO (see Sect. 4 for details).

We have conducted free-running SO experiments for the PD climate (SO-PD), and climates that are 1.5 K and 25 2.0 K warmer than the pre-industrial (SO-15 and SO-20). The SO model has been calibrated to mimic the three HAPPI experiments, using the same forcings for GHGs, aerosols, ozone, and land-use. In SO-PD, the SSTs are constrained to stay close to the observed values from AMIP-PD. The SST difference for SO-15 and SO-20 are based on the SST response in CPL-15 and CPL-20 relative to CPL-PD for consistency with the model climate in the NorESM1-Happi. The SO model and the set-up of the experiments are described in more detail in Sect. 4 and Table 2.

We carried out 150-year simulations for SO-PD, SO-15, and SO-20. After a spin-up of 60 years, a new quasiequilibrium is reached, leaving three equilibrated periods of 90 years each (270 years in total).

³⁰

The biases in the near-surface temperature for the present day climate are shown in Fig. 5b, e, h, and k (for the four seasons). While the biases are larger than those from AMIP-PD, they are still clearly reduced compared to CPL-PD. For instance, the global-mean bias in NH winter (December, January, and February; DJF) is reduced by 35 % in the SO and 64 % in AMIP model compared to the fully coupled model.

5 3 The model

10

20

30

In this section, we give a brief overview of the fully coupled NorESM1-Happi, which is NorESM1-M used for CMIP5 with some upgrades. A more exhaustive overview of the NorESM1 is given in Bentsen et al. (2013), Iversen et al. (2013) and Kirkevåg et al. (2013).

NorESM1-M is based on the fourth version of the Community Climate System Model (CCSM4) developed in the Community Earth System Model project at the US National Center for Atmospheric Research (NCAR) in collaboration with many partners (Gent et al., 2011).

The atmosphere component of the NorESM1-M and -Happi is the "Oslo" version of the CCSM4's Community Atmosphere Model version 4 (CAM4-Oslo). It is based on the CAM4 (Neale et al., 2010; Neale et al., 2014), but has a different aerosol module for aerosol lifecycle calculations and aerosol-cloud-radiation interactions (*Kirkeyåg* et al., 2013).

15 (Kirkevåg et al., 2013).

The ocean component is an elaborated version of the Miami Isopycnic Community Ocean Model (MICOM). This is an entirely different ocean component than the one used in the CCSM4. The MICOM version used in the NorESM1-M and -Happi has been adapted for multi-century simulations in coupled mode (Assmann et al., 2010; Otterå et al., 2010) and includes several extensions compared to the original MICOM (Bentsen et al., 2013).

The land and sea-ice component and the coupler are the same as in the CCSM4. The land component is the fourth version of the Community Land Model (CLM4; Oleson et al., 2010; Lawrence et al., 2011), including the SNow, ICe, and Aerosol Radiative model (SNICAR; Flanner and Zender, 2006). The sea-ice component is the fourth version of Los Alamos Sea Ice Model (CICE4) (Gent et al., 2011; Holland et al., 2012). The coupler is the *CPL7* (Craig et al., 2012).

coupler is the *CPL7* (Craig et al., 2012).

The ocean and sea-ice components of NorESM1-M and NorESM1-Happi were run with the standard CCSM4 land mask and ocean grid (the gx1v6) with 1.125 ° resolution along the equator and with the NH grid singularity located over Greenland. The atmosphere component, CAM4-Oslo was run with a horizontal resolution of 0.95 ° latitude by 1.25 ° longitude (in short: 1° resolution) in NorESM1-Happi and the double of the meshwidth (2 ° resolution) in NorESM1-M. In both versions, CAM4-Oslo has 26 vertical hybrid sigma-pressure levels and a model top at 2.194 hPa. The land component CLM4 employs the same horizontal grid as CAM4-Oslo, except for the river transport model which is configured on its own grid with a horizontal resolution of 0.5 ° in both versions.

Differences between NorESM1-Happi and NorESM1-M include finer horizontal resolution in the atmosphere and land, as described above, but also a few upgrades in the ocean, sea ice, and atmosphere components. In NorESM1-Happi, inertial-gravity waves are damped in shallow ocean regions in order to remove spurious oceanic variability in high-latitude shelf regions (Seland and Debernard, 2014). Wet snow albedo on sea ice is

5 reduced by increasing the wet snow grain size and by allowing a more rapid metamorphosis from dry to wet snow. This affects the Arctic sea ice more than the Antarctic, since the latter is less frequently influenced by mild and humid air (Seland and Debernard, 2014).

In the atmosphere, an error in the aerosol scheme (Kirkevåg et al., 2013) was found and rectified, resulting in faster condensation of secondary gas-phase matter on pre-existing particles. The changes in atmospheric residence time of aerosols compared to NorESM1-M are minor, except for the reductions for black carbon (BC) and organic matter due to more efficient wet deposition. Samset et al. (2013) and Allen and Landuyt (2014) indicated that NorESM1-M had too high upper-air concentrations of BC aerosols. This could cause overestimated absorption of solar radiation, suppressed upper-level cloudiness, and exaggerated static stability.

The increased efficiency of aerosol condensation in NorESM1-Happi enhances the scavenging efficiency of
BC compared to NorESM1-M. This is mainly affecting the upper-air BC concentrations (Fig. S1 in the Supplement) with minor impacts on surface temperatures, surface energy fluxes, and multi-decadal variability associated with the deep oceans (Sand et al., 2015; Stjern et al., 2017). To the extent that the observations from the HIPPO-campaign (Schwarz et al., 2013) is representative for the vertical distribution of BC in general, the model still mixes the BC too high up. A comprehensive discussion of the aerosols in a recently updated
NorESM version (NorESM1.2) is given in Kirkevåg et al. (2018).

3.1 Qualifying NorESM1-Happi: CMIP5 experiments

10

25

We performed a full range of CMIP5 experiments with NorESM1-Happi to document the performance of the model, and to obtain valid historical and RCP8.5 runs for the CPL-PD experiment (Sect. 2.2). The experiments are summarized in Table S1 in the Supplement. The set-up of the simulations follows that of the original CMIP5 simulations with NorESM1-M (Bentsen et al., 2013; Iversen et al., 2013; Kirkevåg et al., 2013).

The NorESM1-Happi with 1 ° resolution was spun up for 1850 conditions over 300 years, starting from model year 600 of the NorESM1-M spin-up with 2 ° resolution atmosphere and land. The ocean and sea ice were in both cases run with 1 ° resolution. The pre-industrial control experiment (piControl) was started from the end of the spin-up in model year 900. The three historical experiments start from the piControl in model years 930

30 (Hist1), 960 (Hist2) and 990 (Hist3). The code upgrades were introduced during the spin-up period, while the bug-fix in the aerosol scheme was introduced at the beginning of the piControl experiment, causing some adjustments over the first few years.

Here we briefly summarize the extensive model validation of NorESM1-Happi against NorESM1-M, observations and reanalysis given by tables and figures, which are commented, in the Supplement. The preindustrial control simulation is considerably more stable for NorESM1-Happi than for NorESM1-M, mainly because the control run started from a state closer to equilibrium in NorESM1-Happi. The NorESM1-Happi piControl experiment also deviates less from the World Ocean Atlas of 2009 (Locarnini et al., 2010; Antonov et al., 2010) than NorESM1-M. The increased horizontal resolution lead to reduced cloudiness in NorESM1-Happi, and along with this a cold bias, a faster atmospheric cycling of fresh water, and overestimated precipitation globally (Table S4 and Figure S5). The atmospheric residence time and ocean to continent transport of water-vapour appears satisfactory (Table S6). Also, the thermohaline forcing of the AMOC was strengthened, and is probably too strong (Figure S14).

NorESM1-Happi has a better representation of sea ice (Table S5 and Figure S4), improved NH extratropical cyclone (Figure S11) and blocking activity (Figure S12), and a fair representation of the Madden-Julian oscillation (Figure S10). The amplitude of the El Niño-Southern Oscillation (ENSO) signals is reduced and is too small, although the frequency is improved (Figure S13). NorESM1-Happi is less sensitive (3.34 K at CO₂ doubling) than NorESM1-M (3.50 K) and slightly more sensitive than CCSM4 (3.20 K; Table S7). The lapse-rate, albedo, and to a smaller extent the short-wave water vapour feedbacks contribute to Arctic amplification

in both model versions (Fig. S15).

4 Emulating the oceanic response with a slab ocean

NorESM1-HappiSO, the slab-ocean (SO) model version of the NorESM1-Happi, has the same atmosphere, land, and sea-ice components and coupler as the fully coupled model. The ocean component is however replaced by a SO model, which is a simplified 2-dimentional ocean model that represents a well-mixed surface mixed layer. Note that it allows for using the same sea-ice model as in the fully coupled model.

A SO model does not calculate ocean circulation and associated fluxes, but treats the upper-ocean mixed layer as a single layer which buffers heat-fluxes through the ocean surface, i.e. a thermodynamic "slab" governed by the equation

$$\rho_0 c_0 h_{mix} \frac{\partial SST}{\partial t} = F_{net} - Q_f - \alpha \left(SST - SST_{ext}\right)/\tau \tag{1}$$

where h_{mix} is the thickness of the slab which varies in space but not in time, ρ_0 and c_0 are the density and specific heat capacity of the sea-water, *SST* is (in this connection) the mixed-layer temperature, F_{net} is the net input of heat through the ocean surface from the atmosphere and sea ice, and Q_f is the net divergence of heat not accounted for by the explicit processes which are needed to maintain a stable climate with a predefined geographical distribution of SST. The last restoring term on the right-hand side could be used to estimate Q_f depending on the value of α . For free SO runs $\alpha = 0$.

30

25

5

10

15

The realism of the SO model climate depends on how Q_f is prescribed. In Bitz et al. (2012), Q_f is calculated using h_{mix} , SST, and F_{net} from a fully coupled stable control simulation, setting $\alpha = 0$. Both h_{mix} and SST should represent an assumed well-mixed layer in the vertical. With an annual mean (but still spatially variable) mixed-layer thickness, it is quite straightforward to obtain balance with the annual cycle of heat (Bitz et al.,

5 2012). This method gives a mean SST distribution from the SO model which is very similar to, and consistent with, the climate of the fully coupled model when external forcing is unchanged. Here, this method has been used when estimating the equilibrium climate sensitivity (ECS) for runs with abrupt CO₂ doubling ($\Delta T_{eq} =$ 3.31 K) and CO₂ quadrupling ($\Delta T_{eq4} = 6.74 K$), giving an average $\Delta T_{eq} = 3.34 K$ for doubling of the atmospheric CO₂ concentrations (Table S7 in the Supplement). The Q_f used in these experiments was 10 diagnosed from the 1850 fully coupled piControl experiment with NorESM1-Happi (Sect. 3.1), and kept

4.1 Calibration of NorESM1-HappiSO experiments

constant in the different SO runs.

One drawback with the method of Bitz et al. (2012) for quantifying Q_f is that biases in SST and the mean climate from the fully-coupled model are reflected in the SO model, which makes comparison with the AMIP 15 experiments difficult. Therefore, as an alternative, we also use a restoring method similar to Williams et al. (2001) and Knutson (2003), where a separate calibration run of the SO is done by setting $\alpha = 0$ in equation (1). SST_{ext} is an externally imposed SST field valid for some specific period (observation or model based) with τ as a prescribed time-scale for adjustment. After this run, the new Q_f is defined by adding the monthly climatology of the restoring flux to the Q_f used in the calibration run. Then, when used in a free SO run (setting 20 $\alpha = 0$), the new Q_f ensures a modelled SST climate close to the SST_{ext} fields imposed during the calibration. We have kept the sea-ice model free without any restoring or constraints to observed fields during the calibration. This increases the realism of the ice-ocean heat fluxes going into F_{net} , and ensures consistent changes in sea-ice mass and energy. As in Bitz et al. (2012), the sea ice in the SO set-up employs the full CICE4 dynamic and thermodynamic model, which is the same as used in the fully coupled NorESM1-M and 25 NorESM1-Happi. However, some tuning of snow albedo over sea ice has been done to increase the realism of sea-ice extent under PD conditions when using the restoring method for specifying Q_f .

In the present case, the purpose of the SO model is to emulate regional patterns of the climate response given a targeted global near-surface temperature change relative to the pre-industrial climate, considering the observed and analysed climate at PD (2006-2015). The experiments with NorESM1-HappiSO are designed to

30 be comparable to the NorESM1-HappiAMIP experiments, in which the SST and sea ice are prescribed (see Sect. 2.1). Three different calibrations of Q_f are therefore performed using the restoring method. For SO-PD we use 12-year averaged SSTs determined by the observationally based *Operational Sea Surface Temperature and Sea Ice Analysis* (OSTIA) for the years 2005–2016 (Donlon et al., 2012). In practice, this calibration also reduces biases. For SO-15 and SO-20, we determine new Q_f fields which adjust the model to an SST field which is consistent with 1.5 K and 2.0 K warming. These fluxes were obtained by adding SST increments based on the difference of the CPL-15 and CPL-20 runs relative to CPL-PD to the OSTIA PD SST field. The different Q_f -fields thus emulate the effects of oceanic circulation changes on the heat flux divergence in the mixed layer.

5 The Q_f -fields are determined for each month of the year, and the values used in the SO model at a given gridpoint and a given time are determined by linear interpolation between the former and the next monthly value. The same Q_f fields are used every year of the simulation. Fig. 6 shows annual averages for SO-PD together with the increments for the 1.5 K and the 2.0 K warmer worlds (SO-15 and SO-20). In addition, we use the same CO_2 levels, aerosols and precursor emissions, and other active forcing agents as in the AMIP 10 experiments. The Q_f for SO-PD (Fig. 6a), which includes bias corrections, is dominated by large negative values (hence SST increase) along the major currents in the North Pacific, North Atlantic, Southern Indian Ocean, and the Atlantic sector or the Arctic. Positive values are mainly seen along the equator and in some coastal upwelling zones. The increment patterns (Fig. 6b and c) appear largely independent of the level of the warming, with positive values (decreasing SST) over the Labrador Current, negative values (increasing SST) 15 south of Iceland, and values of both signs over the Southern Ocean.

20

25

Temperature response 5

In what follows we study results from the PD climate and the response to the warming in the 1.5 K experiment (with respect to PD) and the 0.5 K difference (between the 2.0 K and 1.5 K experiments) from three versions of the NorESM1-Happi: (1) NorESM1-HappiAMIP forced with prescribed SST and sea ice (Sect. 2.1); (2) NorESM-Happi which is fully coupled (Sect. 2.2); (3) NorESM1-HappiSO which employs slab ocean model (Sect. 2.3 and 4). The disadvantage with the AMIP model is that it does not capture any ocean and sea-ice feedbacks. The coupled model however has larger biases, for instance in the near-surface temperature (Fig. 5). The SO model offers an intermediate solution with smaller biases than the fully coupled model (Fig. 5), while still including feedbacks that are missing in the AMIP model. The AMIP experiments however comprise a much larger ensemble of experiments, which may enable statistical significance of smaller trends

5.1 Temperature targets and the polar amplification factor

The changes in the global-mean near-surface temperature for the 1.5 K and 2.0 K warmer worlds are given in Table 3. Note that these runs are designed to have temperature increases of 1.5 K and 2.0 K relative to preindustrial conditions, whereas we are comparing them to the PD climate, which is assumed to be 0.8 K warmer based on observations (Mitchell et al., 2017). Therefore, the ideal temperature increase between the PD experiments and the 1.5 K and 2.0 K warming experiments is 0.7 K and 1.2 K.

30

NorESM1-HappiAMIP hits the temperature targets of 0.7 K and 1.2 K with respect to the PD temperature quite accurately. The corresponding numbers are 0.56 K and 1.02 K for NorESM1-HappiSO and 0.69 K and 1.15 K for NorESM1-Happi. The warming with respect to the PD climate is thus somewhat too low in the SO model whereas it is closer to the targets in the fully coupled one. The difference between the 2.0 K and 1.5 K warming experiments is quite similar across the models: 0.49 K for NorESM-HappiAMIP, 0.43 K for NorESM-HappiSO, and 0.46 K for NorESM1-Happi. The smaller temperature response in the SO experiments

5 is caused by a cold bias over land (Table 4, see discussion below), which cannot be adequately controlled by the adjusted ocean Q_f -fluxes.

10

30

The time-evolution of the global-mean near-surface temperature response to 1.5 K and 2.0 K warming in NorESM1-Happi is shown alongside the response for the Arctic region (area poleward of 65 degN) in Fig. 7. The temperature response is clearly amplified over the Arctic compared to the global mean. The ratio of the polar to the global near-surface temperature response defines the polar amplification factor (PAF; Table 3). [The PAF is considerably larger in the Arctic than in the Antarctic, consistent with polar amplification being more pronounced in the NH. The Arctic amplification (NH-PAF) is furthermore stronger in the 1.5 K than in the 2.0 K warming scenarios.

The Arctic amplification is enhanced in the experiments with active ocean components. Compared to
15 NorESM1-HappiAMIP, the Arctic amplification is 27 % stronger in the 1.5 K warmer world in NorESM1-HappiSO and 54 % stronger in NorESM1-Happi. With the additional 0.5 K warming the Arctic amplification is moreover 19 % and 46 % stronger in the SO and the fully coupled model than in the AMIP model.

Table 4 shows similar statistics as Table 3, but for the NH extratropical (poleward of 20 °N) winter and summer land temperatures, land precipitation rates, and sea-ice area. The winter climate is colder over land in the coupled models than the AMIP model. The difference is -0.54 K for the fully coupled model and -0.57 K for the SO model with respect to the AMIP model. During summer, land temperatures are almost as high in the SO model as in the AMIP model, whereas the fully coupled model is 1.58 K colder. This is in line with the lager bias in the fully coupled model during this season (Fig. 5g–i).

The fully coupled model has the largest reduction in sea-ice area in the warmer climates during summer and
 winter. The SO model has larger changes than the AMIP model during summer and smaller changes during winter.

During summer, the SO and the fully coupled models have the largest changes in land temperatures and precipitation in the 1.5 K warming experiment, whereas the AMIP model has the largest changes with the additional 0.5 K warming. During winter, the AMIP model has the largest changes in precipitation and temperature in with the 1.5 K and the smallest changes in precipitation with the additional 0.5 K warming.

So far we have considered changes in surface fields, but changes are also occurring aloft. Figure 8 shows the zonal-mean temperature response to the 1.5 K warming relative to the PD climate for NH winter (DJF) and NH summer (June, July, and August; JJA). There is low-level warming in the Arctic and high-level warming in the tropics in all three models. The Arctic warming is strongest in the fully coupled model, consistent with

the PAF results in Table 3. The upper-level warming over the tropics appears to be more consistent between the seasons and is somewhat more pronounced in the AMIP model.

5.2 Equator-to-pole temperature gradients

coupled model has the smallest bias and the strongest contrasts.

The warming pattern in Fig. 8 is consistent with a sharpening of the upper-level equator-to-pole temperature
gradient and a weakening of the lower tropospheric gradient. Li et al. (2018) considered the multi-model changes in these gradients in five of the models contributing to the HAPPI project, including NorESM1-HappiAMIP. They found that the low-level gradient changes more with the initial 0.7 K warming (1.5K–PD) than with the additional 0.5 K warming (2.0 K–1.5 K) in all the models. The upper-level gradient on the other hand strengthens more with the additional 0.5 K warming than with the initial 0.7 K, except in NorESM1-HappiAMIP where the changes are more similar.

Figure 9 and 10 show the temperature gradients between the equator and the North Pole at 200 hPa and 850 hPa (e.g. Harvey et al., 2014) for the PD and the 1.5 K and 2.0 K warming experiments in NorESM1-Happi, NorESM1-HappiSO and NorESM1-HappiAMIP, and for each of the seasons.

- The magnitude of the PD gradients is weaker in the fully coupled than in the SO and AMIP models, except
 during summer when the low-level gradient is stronger in the fully coupled model. While the fully coupled model might seem like an outlier, the upper-level gradient is actually closer to the one in ERA-Interim (Dee et al., 2011), indicating that the SO and AMIP models overestimate the upper-level pole-to-equator temperature contrast (Fig. 9). At low-levels fully coupled model underestimates the gradient during winter and spring (March, April, and May; MAM), while the gradients in the SO and AMIP models are stronger and closer to the reanalysis (Fig. 10). During summer (JJA) and fall (September, August, and November; SON) the fully
 - In line with the zonal-mean response in Fig. 8, the high-level gradient generally increases with the warming (Fig. 9) while the low-level gradient decreases (Fig. 10). The low-level gradient decreases more with initial the 0.7 K warming than with the additional 0.5 K, consistent with Li et al. (2018). The decrease with the initial
- 25 0.7 K is moreover larger in the fully coupled and SO models than in the AMIP model, consistent the stronger Arctic amplification in these models (Table 3).

Changes in the upper-level gradient are less consistent across the experiments and seasons. During winter and spring, the gradient strengthens with the additional 0.5 K warming in all three models. There is however little change with the initial 0.7 K warming in the fully coupled and SO models. During summer and fall, the upper-

30 level gradient strengthens more with the initial 0.7 K warming than with the additional 0.5 K warming, like at low levels, only with no obvious differences between the model versions.

It is possible that the upper-level warming in the fully coupled and to SO experiments are affected by cold biases in the tropics. Both the fully coupled and the SO models are colder over land than the AMIP model (Table 4), and the fully coupled model additionally has cold biases over the tropical oceans (Fig. 5).

Extratropical storm-track activity 6

5 Changes in the temperature gradients are known to be associated with changes in the extratropical storm tracks, with stronger gradients being associated with poleward shifts and weaker gradients with equatorward shifts (Brayshaw et al., 2008; Graff and LaCasce, 2012; Harvey et al., 2014; Shaw et al. 2016).

Extratropical storm tracks can be defined as regions of growing and decaying baroclinic waves embedded in the zones of pronounced meridional temperature gradient and mean westerly wind currents. Here we represent the storm-track activity in terms of atmospheric fields, such as geopotential height, that have been bandpass filtered in time to isolate disturbances with timescales between 2.5 and 6 days (following Blackmon 1976 and Blackmon et al., 1977). The variability of the resulting fields is dominated by growing and decaying baroclinic waves, and the storm tracks are taken to be maxima in the bandpass-filtered variance fields (e.g. Blackmon et al., 1977; Chang et al. 2002; Chang et al., 2012).

15 Figure 11 shows the bias in the PD storm-track activity in terms of bandpass-filtered geopotential height at 500 hPa for NorESM1-Happi, NorESM1-HappiSO and NorESM1-HappiAMIP. The fully coupled model underestimates the variability in all seasons. The bias is largest over the North Atlantic during winter when the storm-track activity is underestimated on the equatorward side and over the Nordic Seas, consistent with the North Atlantic storm track being overly zonal (Iversen et al., 2013). The SO and AMIP models have both 20 positive and negative biases over the storm-track regions, and a North-Atlantic storm track which extends too far downstream over central Europe.

The storm-track biases are largest in the fully coupled model whereas they are substantially smaller in the SO and AMIP models. The area-averaged winter bias for the region shown in Figure 11 is for instance -4.24 m in the fully coupled model, 0.89 m in the SO model, and 0.51 m in the AMIP model.

25 Figure 12 shows the changes in upper-level storm-track activity in the 1.5 K warming experiments for the three models and all four seasons. The AMIP model has the most consistent changes with more storm-track activity at high latitudes and less at lower latitudes, consistent with a poleward shift, for all seasons. The exception is over the North Pacific where there is an equatorward shift near the North-American west coast region during winter and a more general equatorward shift of the whole storm track during summer. The results for the AMIP 30 model are in line with the multi-model mean results in Li et al. (2018).

10

Changes in the fully coupled and the SO model are less consistent. During summer and fall the changes resemble those in the AMIP model with more activity over the high latitudes and less over the low latitudes. The reductions on the equatorward side are however stronger for the fully coupled and the SO model. Changes during winter and spring are more complicated, and do not particularly resemble those in the AMIP model.

The upper-level storm-track response to the additional 0.5 K warming is shown in Fig. 13. Here the changes are more similar across the models and seasons, particularly over the North Atlantic where there tends to be more storm-track activity on the poleward side and less on the equatorward side. The poleward shifts are in line with changes in the upper-level temperature gradient, which strengthens with the 0.5 K warming for all

5 cases.

The white dots in Fig. 12 and 13 indicate that only the very strongest changes are significant in NorESM1-Happi and NorESM1-HappiSO whereas the changes in NorESM1-HappiAMIP are more generally significant. This could be caused by the smaller number of model years available for the fully coupled and SO model, but it could also reflect a larger spread between the decades/members. The similarity, or lack thereof, between the storm-track response in the two coupled models and the AMIP model does nonetheless increase, or reduce,

10

our confidence in the AMIP results. Li et al. (2018) found that while there is a poleward shift in upper-level storm-tracks activity with both the

initial 0.7 K and the additional 0.5 K warming in the HAPPI multi-model ensemble, the changes at low levels are less consistent. Figure 14 shows the changes in the low-level storm-track activity in the 1.5 K warming
experiment for NorESM1-Happi, NorESM1-HappiSO, and NorESM1-HappiAMIP during winter and summer. The low-level storm tracks are given in terms of the bandpass-filtered meridional eddy heat flux. As in Li et al., the response to the initial 0.7 K warming is generally a reduction in storm-track activity, here indicating that the storm-track eddies are transporting less heat poleward. The decrease over the North-Atlantic region is stronger in the fully coupled and the SO model than in the AMIP model. Changes during summer are weak.

The change in the low-level storm-track activity in response to the additional 0.5 K warming is shown in Fig. 15 for summer and winter. Again changes are weak during summer. During winter, the AMIP and SO models have an increase over the Nordic Season, but this is less pronounced (and not significant) in the fully coupled model. A similar increase is however present in the multi-model mean in Li et al. (2018).

25 7 Blocking frequency

Extratropical blocking is closely connected to persistent anticyclones, which can suppress precipitation at midlatitudes for periods of up to several weeks. The ability of climate models to simulate the occurrence of droughts at mid-latitudes in the present and in future climates is conditioned by the models ability to simulate blocking (e.g. Woolings et al., 2018).

30 Figure 16 shows the PD blocking frequency for NorESM1-Happi, NorESM1-HappiSO, and NorESM1-HappiAMIP for the winter and summer seasons. The blocking frequency is underestimated over the North Atlantic and western Europe during winter and over large parts of Eurasia during summer. The performance of the three models is generally similar, though some differences can be seen. The overestimation in NorESM1Happi at 120 °W is for instance not as pronounced in the other two models. The SO and AMIP models perform slightly better over the Pacific, but the blocking occurrence is still underestimated in the Atlantic sector.

It is well established that many global climate models have problems simulating the occurrence and duration of blocking in the Euro-Atlantic sector and that the systematic errors are particularly large during NH winter. Several studies tie these problems to poor horizontal resolution (Dawson et al., 2012; Davini and D'Andrea, 2016; Woolings et al., 2018)

The changes in the occurrence of winter and summer blocking in the 1.5 K warming experiment (relative to PD) and with the additional 0.5 K warming are shown in Fig. 17 for the NorESM1-Happi, NorESM1-HappiSO, and NorESM1-HappiAMIP. The magnitude of the response varies dramatically between the models, and although not shown, the same lack of consistency is also found for spring and fall. The magnitude of the changes is largest in the fully coupled model, but are almost as large in the SO model. Note that the AMIP response can be statistically significant relative to the internal variability in the AMIP model, even though the amplitude of the response is small.

There is, however, little consistency between the sign and significance (indicated by the asterisk) of the response for the different longitudes. There are indications of more consistent changes between the model versions with the 0.5 K incremental warming during NH summer, with increased blocking occurrence over parts of western Europe, the eastern Pacific, and the western Pacific. Changes are in these cases larger in the coupled models, but most significant in the AMIP model. Nevertheless, the results concerning NH blocking generally remain inconclusive.

20 8 Arctic sea-ice reduction

The extent[AK39], thickness and concentration of sea ice are important properties of the climate system. Figure 18 shows the [AK40] concentration of Arctic sea ice in March and September for NorESM1-Happi and NorESM1-HappiSO. For PD (Fig. 18a–d) the modelled concentrations are compared to remotely retrieved data from OSI-SAF (2017). The quality of the model data is better in March than in September, when the SO model seems to underestimate the concentration.

The sea-ice concentration is reduced in the warmer climates. In March, the largest changes occur along the edges of the ice (Fig. 18e–f, i–j). There is a larger reduction in the fully coupled than the SO model with the initial 0.7 K warming, whereas the changes are more similar with the additional 0.5 K. The changes occur over a larger fraction of the sea-ice covered area in September (Fig. 18g–h, k–l). While the changes again are larger

30

25

5

10

with the 0.7 K than with the additional 0.5 K warmings in the fully couple model, they are more comparable in the SO model.

While the sea-ice concentration is reduced more with the warming in the fully coupled model, ice-free summers are more likely in to SO model. Figure 19 shows histograms of the relative occurrence of NH September sea-

ice extent for NorESM1-Happi (Fig. 19a) and NorESM1-HappiSO (Fig. 19b). The sea-ice extent is shown for the observed and the modelled PD climate and the 1.5 K and 2.0 K warming experiments. For PD climate, the SO model produces too few cases with the largest sea-ice extent whereas the fully coupled model has too many. The overrepresentation in the latter case is likely caused by the cold bias in the model.

- 5 The probability of having an ice-free Arctic in September, i.e. having a sea-ice extent between 0 and 1×10^6 km², is practically zero for PD conditions in both models. The fully coupled model does not reach ice-free conditions with 1.5 K nor with 2.0 K warming (Fig. 19a). This is perhaps not surprising as the model is too cold and has too much ice in the PD climate. So even though there are larger reduction in the sea-ice concentration in the fully coupled model, it does not have an ice-free Arctic in September.
- 10 Results are different for the SO model which has smaller biases in temperature and sea-ice extent. While ice-free September conditions rather unlikely under 1.5K, but the probability increases substantially to about 18 % with the additional 0.5 K warming (Fig. 19b). The difference between the two temperature targets is therefore potentially very large for the Arctic sea ice in summer and fall, a result that was also found by Jahn (2018) and Sigmond et al. (2018).

15 9 Summary and discussion

This paper presents an evaluation of the importance of ocean and sea-ice feedbacks under global warming of 1.5 K and 2.0 K relative to pre-industrial conditions. We compare results from a fully coupled and a SO (slabocean) version of the NorESM1-Happi to results from the AMIP-style simulations that were carried out for the multi-model HAPPI project (Mitchell et al., 2017; <u>http://www.happimip.org/</u>).

20 Because the AMIP runs are forced with prescribed SSTs and sea ice they have small biases, but they also predefine aspects of the Arctic amplification. The fully coupled and the SO models allow for changes in SST and sea ice that can influence the surface albedo and atmospheric lapse rate, which are major elements in producing Arctic amplification (Pithan and Mauritsen, 2014). The motivation for using a SO model in addition to the fully coupled one is that the SO model has smaller biases, while still allowing the ocean and sea ice to 25 respond to the forcing in the warming runs.

We consider the PD (present day) climate, the response to the 0.7 K warming between the PD and the 1.5 K warming experiments (assuming 0.8 K warming between 1850 and PD), and the response to the 0.5 K warming between the 1.5 K experiment and the 2.0 K experiments.

Results show that Arctic amplification, as measured by the PAF (polar amplification factor) for the NH, is
larger in the models with active ocean components. In the fully coupled model, the PAF is 54 % stronger than in the AMIP model with the initial 0.7 K warming, and 46 % stronger with the additional 0.5 K warming. The difference is not as large for the SO model which has 27 % and 19 % stronger PAF for the same warmings.

Arctic amplification weakens the lower tropospheric equator-to-pole temperature gradient, and this decrease is larger with the initial 0.7 K warming than with the additional 0.5 K for all seasons. A similar result is also found in the AMIP runs from the five HAPPI models (including NorESM1-HappiAMIP) studied by Li et al. (2018). This study however shows that the changes with the initial 0.7 K warming is larger in the fully coupled and SO models than in the AMIP model, particularly during summer (JJA) and fall (SON).

5

35

The changes in the upper-level equator-to-pole gradients are less consistent. The gradients generally increase with the warming because the tropics are warming aloft (e.g. Collins et al., 2013). During summer and fall, the gradient changes more with the initial 0.7K warming, similar to with the low-level gradient. The magnitude of the response is however not systematically larger in the experiments with active ocean components. During winter and spring, the upper-level gradient changes very little with the initial 0.7 K warming in the coupled models and more with the additional 0.5 K, whereas the AMIP model has more similar changes with the 0.7 K and the 0.5 K warming. The changes in the upper-level gradient are also less consistent than those in the low-level gradient in Li et al. (2018); while the upper-level gradient changes more with the additional 0.5 C in the multi-model mean, there is considerable spread among the models.

15 Changes in temperature gradients are known to be associated with changes in the storm tracks, with the tracks shifting poleward with stronger gradients and equatorward with weaker ones (Brayshaw et al., 2018; Graff and LaCasce, 2012; Harvey et al., 2014; Shaw et al., 2016). Li et al. (2018) identified poleward shifts in the multi-model mean upper-level storm tracks with the initial 0.7 K warming and with the additional 0.5 K warming. We find that while the AMIP model displays consistent poleward shifts in the upper-level storm-track activity
20 with the initial 0.7 K warming for all seasons, the results from the coupled models are less consistent during winter and spring. The models agree more on the response to the additional 0.5 K. However, only the strongest changes in the fully coupled and the SO model are significant.

The low-level storm-track activity decreases with the initial 0.7 K warming. Changes with the additional 0.5 K warming are weak in the AMIP model, whereas the fully coupled and SO models have stronger reductions. All model versions have indications of more activity east of the British Isles, a response also seen in the multi-

- 25 All model versions have indications of more activity east of the British Isles, a response also seen in the multimodel mean in Li et al. (2018). These changes are however mostly not significant in the coupled models. To the extent that reduced low-level storm-track activity can be interpreted as slower propagation of cyclone waves in the westerlies, this can be associated with the reduced low-level temperature gradient associated with the high-latitude warming over the Arctic (e.g. Francis and Vavrus, 2012; Screen and Simmonds, 2013). The
- 30 results for blocking activity for the most remain inconclusive due lack of consistency between the model versions and the low statistical significance of the changes. Many aspects of blocking are also poorly simulated, likely because of relatively coarse model resolution (Woolings et al., 2018).

Our findings indicate that the storm-track response is not always very consistent between the model versions. There are moreover sizable biases in the storm tracks with respect to reanalysis, especially in the fully coupled model. Barcikowska et al. (2018) provided a study of the Euro-Atlantic winter storminess which showed that modelling the regional atmospheric circulation, extreme precipitation and winds with acceptable quality requires an atmospheric model with higher horizontal resolution $(0.25^{\circ} \text{ in their study})$ than that used in the present study and in CMIP5 models.

The SO model simulates considerable differences in the reduction of sea ice in the Arctic between a 1.5 K and a 2.0 K warmer world. Ice-free summer conditions in the Arctic are estimated to be rare under 1.5 K warming, while occurring 18 % of the time under 2.0 K warming. These results are consistent with other studies (Jahn, 2018; Sigmond et al., 2018; Notz and Strove, 2018). The fully coupled model is however too cold. It produces too much sea ice under PD conditions and is consequently not able to reach ice-free conditions in neither the 1.5 K nor the 2.0 K warming experiment.

- 10 This paper does not discuss practical or scientific challenges that must addressed in order to avoid exceeding certain temperature targets. Mathews et al. (2009) and Gillett et al. (2011) indicate that a constant equilibrium response in surface air temperature to anthropogenic CO₂ is determined by the accumulated carbon emissions. Hence, an ESM which calculates the atmospheric concentrations of CO₂ on-line from emissions, should produce quite rapid stabilization of the global mean surface temperature. This is enabled if the ocean thermal
- 15 inertia is balanced by decreasing atmospheric concentrations of CO₂ due to ocean uptake. NorESM1-Happi is not equipped with the possibility to run emission-driven GHG scenarios with on-line carbon-cycling. Instead, the atmospheric concentrations are prescribed.

10 Code and Data Availability

20

The source code for NorESM1-Happi is not open for everyone to download, because parts of the code is imported from several other code development centres. The code can be made available within the framework of an agreement. Data from the model experiments in this study can be made available as well, see e.g. NCC / NorESM1-HAPPI at http://portal.nersc.gov/c20c/data.html. Contacts: oyvindse@met.no and ingo.bethke@uni.no.

Acknowledgements

- 25 This work received financial support from the Norwegian Research Council, project no. 261821 (HappiEVA), and in-kind support from Geophysical Institute, Univ. of Bergen, UniResearch, Bergen, and from MET Norway. Jan Fuglestvedt, Cicero, Norway, Dann Mitchell, Univ. Bristol, UK, and Myles Allen, Univ. Oxford, UK, were instrumental in inspiring the initiating this work, and have strongly contributed in discussions. HPC-resources for the NorESM runs were provided in kind from Bjerknes Centre for Climate Research, MET
- 30 Norway, and Sigma2/Vilje (nn2345k). Storage for NorESM data was provided through Norstore/NIRD (ns9082k). The development of NorESM has been possible because of the granted early access to the later public versions of the CCSM4 and CESM1 by the US National Center for Atmospheric Research (NCAR). Changes in NorESM1-Happi compared to NorESM1-M were developed in the project ACCESS (Arctic Climate Change, Economy and Society) financed under the 7th Framework Programme of the European Union

(<u>http://www.access-eu.org/</u>) and in the Norwegian Research Council, project Earth system modelling of climate Variations in the Anthropocene1 (EVA, 229771/E10).

References

Acosta Navarro, J.C., Varma, V., Riipinen, I., Seland, Ø., Kirkevåg, A., Struthers, H., T. Iversen, Hansson, H.-C., and Ekman, A. M. L.: Amplification of Arctic warming by past air pollution reductions in Europe. Nature Geoscience, 03, DOI:10.1038/ngeo2673, 2016.

Allen, R. J., and Landuyt, W.: The vertical distribution of black carbon in CMIP5 models: Comparison to observations and the importance of convective transport, J. Geophys. Res. Atmos., 119, 4808–4835, doi:10.1002/2014JD021595, 2014.

10

25

Anderson, K. and Nevins, J.: Planting Seeds So Something Bigger Might Emerge: The Paris Agreement and the Fight Against Climate Change, Socialism and Democracy, 30, 209–218, 2016.

Assmann, K. M., Bentsen, M., Segschneider, J., and Heinze, C.: An isopycnic ocean carbon cycle model, Geosci. Model Dev., 3, 143–167, doi:10.5194/gmd-3-143-2010, 2010.

Arrhenius, S.: On the influence of carbonic acid in the air upon the temperature of the ground, Philos. Mag. J. Sci. 5, 237-276.

Baker, H. S., Millar, R. J., Karoly, D. J., Beyerle, U., Guillod, B. P., Mitchell, D., Shiogama, H., Sparrow, S.
Woollings, T. and Allen, M. R: Higher CO2 concentrations increase extreme event risk in a 1.5 °C world.
Nature Climate Change | VOL 8 | JULY 2018 | 604–608 |, doi.org/10.1038/s41558-018-0190-1, 2018.

20 Barcikowska, M. J., Weaver, S. J., Feser, F., Russo, S, Schenk, F., Stone, D. A., Wehner, M. F. and Zahn, M.: Euro-Atlantic winter storminess and precipitation extremes. Earth Syst. Dynam., 9, 679–699, https://doi.org/10.5194/esd-9-679-2018, 2018.

Barnes. E. A. and Polvani, L. M.: CMIP5 Projections of Arctic Amplification, of the North American/North Atlantic Circulation, and of Their Relationship. J. Climate, 28, 5254-5270. doi: 10.1175/JCLI-D-14-00589.1, 2015.

Bentsen, M., Bethke, I., Debernard, J. B., Iversen, T., Kirkevåg, A., Seland, Ø., Drange, H., Roelandt, C., Seierstad, I. A., Hoose, C., and Kristjánsson, J. E.: The Norwegian Earth System Model, NorESM1-M—Part 1: Description and basic evaluation of the physical climate. Geosci. Model Dev., 6, 687–720, doi:10.5194/gmd-6-687-2013, 2013.

Bitz, C. M., Shell, K. M., Gent, P. R., Bailey, D. A., Danabasoglu, G., Armour, K. C., M. M. Holland, J. T. Kiehl: Climate Sensitivity of the Community Climate System Model, Version 4. J. Climate, 25, 3053–3070. doi: <u>http://dx.doi.org/10.1175/JCLI-D-11-00290.1</u>. 2012.

Blackmon, M. L.: A climatological spectral study of the 500 mb geopotential height of the Northern Hemisphere. J. Atmos. Sci., 33, 1607–1623. 1976

Blackmon, M. L., Wallace, J. M., Lau, N., and Mullen, S. L.: An Observational Study of the Northern Hemisphere Wintertime Circulation. J. Atmos. Sci., 34, 1040\u20131053, <u>https://doi.org/10.1175/1520-</u>0469(1977)034<1040:AOSOTN>2.0.CO;2, 1977.

5

10

Boucher, O., Bellassen, V., Benveniste, H., Ciais, P., Criqui, P., Guivarch, C., Le Treut, H., Mathy, S., and Séférian, R.: Opinion: In the wake of Paris Agreement, scientists must embrace new directions for climate change research, P. Natl. Acad. Sci. USA, 113, 7287–7290, 2016.

Brayshaw, D. J., Hoskins, B., and Blackburn, M.: The Storm-Track Response to Idealized SST Perturbations in an Aquaplanet GCM. J. Atmos. Sci., 65, 2842–2860, <u>https://doi.org/10.1175/2008JAS2657.1</u>, 2008.

Chang, E. K., Lee, S., and Swanson, K. L.: Storm Track Dynamics. J. Climate, 15, 2163–2183, https://doi.org/10.1175/1520-0442(2002)015<02163:STD>2.0.CO;2, 2002.

Chang, E. K. M., Guo, Y., and Xia, X.: CMIP5 multimodel ensemble projection of storm track change under global warming, J. Geophys. Res., 117, D23118, doi: 10.1029/2012JD018578, 2012.

15 Cohen, J., Screen, J. A., Furtado, J. C., Barlow, M., Whittleston, D., Coumou, D., Francis, J., Dethloff, K., Entekhabi, D., Overland, J., and Jones, J.: Recent Arctic amplification and extreme mid-latitude weather, Nature Geoscience 7, 627–637, 2014.

Collins, M., Knutti, R., Arblaster, J., Dufresne, J.-L., Fichefet, T., Friedlingstein, P., Gao, X., Gutowski, W., Johns, T., Krinner, G., Shongwe, M., Tebaldi, C., Weaver, A., and Wehner, M.: Long-term Climate Change:

20 Projections, Commitments and Irreversibility, book section 12, p. 1029–1136, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, https://doi.org/10.1017/CBO9781107415324.024, 2013.

Coumou, D., Capua, G. Di, Vavrus, S., Wang, L., and Wang, S.: The influence of Arctic amplification on midlatitude summer circulation. Nature Comm., 9. doi: 10.1038/s41467-018-05256-8, 2018.

25 Davini P, D'Andrea F. Northern Hemisphere atmospheric blocking representation in global climate models: twenty years of improvements? J Clim., 29, 8823–40. doi.org/10.1175/JCLI-D-16-0242.s1, 2016.

Dawson, A., Palmer, T. N., and Corti, S.: Simulating regime structures in weather and climate prediction models. Geophys. Res. Lett., 39. L21805, doi:10.1029/2012GL053284, 2012.

- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M.
 A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-
 - K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N. and Vitart, F., The ERA-Interim reanalysis:

configuration and performance of the data assimilation system. Q.J.R. Meteorol. Soc., 137: 553–597. doi: <u>10.1002/qj.828</u>, 2011.

Donlon, C. J., Martin, M., Stark, J., Roberts-Jones, J., Fiedler, E., and Wimmer, W: The Operational Sea Surface Temperature and Sea Ice analysis (OSTIA). *Remote Sensing of the Environment*. doi: 10.1016/j.rse.2010.10.017, 2011.

5

15

25

Feldl, N., Anderson, B. T., and Bordoni, S.: Atmospheric eddies mediate lapse rate feedback and Arctic amplification. J. Climate., 30, 9213–9224, doi:10.1175/JCLI-D-16-0706.1, 2017.

Flanner, M. G., and Zender, C. S.: Linking snowpack micro-physics and albedo evolution. J. Geophys. Res., 111, D12208, doi:10.1029/2005JD006834, 2006.

10 Francis, J. A., and Vavrus, S. J.: Evidence linking Arctic amplification to extreme weather in mid-latitudes. Geophysical Research Letters, 39 (6), doi:10.1029/2012GL051000, 2012.

Gent, P. R., Danabasoglu, G., Donner, L. J., Holland, M. M., Hunke, E. C., Jayne, S. R., Lawrence, D. M., Neale, R. B., Rasch, P. J., Vertenstein, M., Worley, P. H., Yang, Z.-L., and Zhang, M.: The Community Climate System Model Version 4. J. Climate, 24, 4973–4991. doi: <u>http://dx.doi.org/10.1175/2011JCLI4083.1</u>, 2011.

Gillett, N. P., Arora, V. K., Zickfeld, K., Marshall, S. J. and Merryfield, W. J.: Ongoing climate change following a complete cessation of carbon dioxide emissions. Nature Geoscience, 4. doi: 10.1038/NGEO1047. 2011.

Graff, L. S. and LaCasce, J. H.: Changes in the Extratropical Storm Tracks in Response to Changes in SST in
an AGCM. J. Climate, 25, 1854–1870, https://doi.org/10.1175/JCLI-D-11-00174.1, 2012.

Harvey, B. J., Shaffrey, L. C., and Woollings, T. J.: Equator-to-pole temperature differences and the extratropical storm track responses of the CMIP5 climate models, Climate Dynamics, 43, 1171\u20131182, https://doi.org/10.1007/s00382-013-1883-9, 2014.

Holland, M. M., and Bitz, C. M.: Polar amplification of climate change in coupled models, Clim. Dyn., 21, 221–232, 2003.

Holland, M. M., Bailey, D. A., Briegleb, B. P., Light, B., and Hunke, E.: Improved sea ice shortwave radiation physics in CCSM4: the impact of melt ponds and aerosols on arctic sea ice, J. Climate, 25, 1413–1430, doi:10.1175/JCLI-D-11-00078.1, 2012.

Hulme, M.: 1.5 °C and climate research after the Paris Agreement, Nat. Clim. Change, 6, 222–224, 2016.

Hurrell, J. W., Holland, M. M., Gent, P. R., Ghan, S., Kay, J. E., Kushner, P. J., Lamarque, J.-F., Large, W. G., Lawrence, D., Lindsay, K., Lipscomb, W. H., Long, M. C., Mahowald, N., Marsh, D. R., Neale, R. B., Rasch, P., Vavrus, S., Vertenstein, M., Bader, D., Collins, W. D., Hack, J. J., Kiehl, J., and Marshall, S: The

Community Earth System Model: A Framework for Collaborative Research. Bull. Amer. Meteor. Soc., 94, 1339–1360, <u>https://doi.org/10.1175/BAMS-D-12-00121.1</u>, 2013

IPCC, 2013: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., Qin, D., Plattner, G.-

K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V. and Midgley, P. M. (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535pp, 2013.

Iversen, T., Bentsen, M., Bethke, I., Debernard, J. B., Kirkevåg, A., Seland, Ø., Drange, H., Kristjansson, J. E., Medhaug, I., Sand, M., and Seierstad, I. A.: The Norwegian Earth System Model, NorESM1-M – Part 2: Climate response and scenario projections, Geosci. Model Dev., 6, 389–415, doi:10.5194/gmd-6-389-2013, 2013.

10 201

15

Jahn, A.: Reduced probability of ice-free summers for 1.5 °C compared to 2 °C warming, Nature Climate Change, 5, 409-413. doi: 10.1038/s41558-018-0127-8, 2018

James, R., Washington, R., Schleussner, C.-F., Rogelj, J., and Conway, D.: Characterizing half-a-degree difference: a review of methods for identifying regional climate responses to global warming targets, WIREs Clim Change, 8:e457. doi: 10.1002/wcc.457doi:10.1002/wcc.457, 2017.

- Kirkevåg, A., Iversen, T., Seland, Ø., Hoose, C., Kristjánsson, J. E., Struthers, H., Ekman, A. M. L., Ghan, S., Griesfeller, J., Nilsson, E. D., and Schulz, M.: Aerosol–climate interactions in the Norwegian Earth System Model NorESM1-M, Geosci. Model Dev., 6, 207–244, doi:10.5194/gmd-6-207-2013, 2013.
- Kirkevåg, A., Grini, A., Olivié, D., Seland, Ø., Alterskjær, K., Hummel, M., Karset, I. H. H., Lewinschal, A.,
 Liu, X., Makkonen, R., Bethke, I., Griesfeller, J., Schulz, M., and Iversen, T.: A production-tagged aerosol module for earth system models, OsloAero5.3 extensions and updates for CAM5.3-Oslo, Geosci. Model Dev. Discuss., https://doi.org/10.5194/gmd-2018-46, accepted, 2018.

Knutson, T.: FMS slab ocean model technical documentation, 2003. [Visited online 2017 at http://www.gfdl.noaa.gov/fms-slab-ocean-model-technical-documentation.]

25 Lawrence, D. M., Oleson, K. W., Flanner, M. G., Thornton, P. E., Swenson, S. C., Lawrence, P. J., Zeng, X., Yang, Z.-L., Levis, S., Sakaguchi, K., Bonan, G. B., Slater, A. G.: Parameterization improvements and functional and structural advances in Version 4 of the Community Land Model, J. Adv. Model. Earth Syst., 3, M03001, doi: 10.1029/2011MS00045, 2011.

Li, C., Michel, C., Seland Graff, L., Bethke, I., Zappa, G., Bracegirdle, T. J., Fischer, E., Harvey, B. J., Iversen,
T., King, M. P., Krishnan, H., Lierhammer, L., Mitchell, D., Scinocca, J., Shiogama, H., Stone, D. A., and Wettstein, J. J.: Midlatitude atmospheric circulation responses under 1.5 and 2.0 °C warming and implications for regional impacts, Earth Syst. Dynam., 9, 359-382, https://doi.org/10.5194/esd-9-359-2018, 2018.

Liu, W., Lim, W. H., Sun, F., Mitchell, D., Wang, H., Chen, D.: Global freshwater availability below normal conditions and population impact under 1.5 and 2 °C stabilization scenarios. Geophys. Res. Lett., doi.org/10.1029/2018GL078789, 2018.

Manabe S., Stouffer R. J.: Sensitivity of a global climate model to an increase of CO2 concentration in the atmosphere. J Geophys Res 85: 5529–5554, 1980

Mitchell, D., James, R., Forster, P., Betts, R., Shiogama, H., and Allen, M.: Realizing the impacts of a 1.5 °C warmer world., Nat. Clim. Change, 6, 735–737, doi:10.1038/nclimate3055, 2016.

Mitchell, D., AchutaRao, K., Allen, M., Bethke, I., Forster, P., Fuglestvedt, J., Gillett, N., Haustein, K., Iverson, T., Massey, N., Schleussner, C.-F., Scinocca, J., Seland, Ø., Shiogama, H., Shuckburgh, E., Sparrow,

 S., Stone, D., Wallom, D., Wehner, M., and Zaaboul, R.: <u>Half a degree Additional warming, Projections,</u> <u>Prognosis and Impacts (HAPPI): Background and Experimental Design</u>, Geosci. Model Dev., 10, 571–583. doi:10.5194/gmd-10-571-2017, 2017.

Neale, B, R., Richter, J. H., Conley, A. J., Park, S., Lauritzen, P. H., Gettelman, A., Williamson, D. L., Rasch, P. J., Vavrus, S. J., Collins, W. D, Taylor, M. A., Zhang, M., and Lin, S.-J.: Description of the NCAR Community Atmosphere Model (CAM 4.0), NCAR TECHNICAL NOTE, 2010.

Neale, R. B., Richter, J. H., Park, S., Lauritzen, P. H., Vavrus, S. J., Rasch, P. J., Zhang, M.: The mean climate of the Community Atmosphere Model (CAM4) in forced SST and fully coupled experiments. J. Climate 26, 5150-5168. doi: 10.1175/JCLI-D-12-00236.1, 2013.

Notz, D., Stroeve, J.: The Trajectory Towards a Seasonally Ice-Free Arctic Ocean. Curr Clim Change Rep, doi: <u>10.1007/s40641-018-0113-2</u>, 2018

Oleson, K. W., and Coauthors: Technical Description of version 4.0 of the Community Land Model (CLM). NCAR Technical Note NCAR/TN-478+STR, doi:10.5065/D6FB50WZ, 2010.

OSI-SAF, 2017: EUMETSAT Ocean and Sea Ice Satellite Application Facility. Global sea ice concentration climate data record 1979-2015 (v2.0, 2017), [Online]. Norwegian and Danish Meteorological Institutes. doi: 10.15770/EUM_SAF_OSI_0008

Otterå, O. H., Bentsen, M., Drange, H., and Suo, L.: External focing as a metronome for Atlantic multidecadal variability, Nature Geosci., doi: 10.1038/NGEO955, 2010.

Pelly, J. L. and Hoskins, B. J.: A new perspective on blocking. J. Atmos. Sci., 60,743–755. 2003.

Peters, G. P.: The "best available science" to inform 1.5 _C policy choices, Nat. Clim. Change, 6, 646–649, doi:10.1038/nclimate3000, 2016.

30

5

15

20

25

Pithan, F. and Mauritsen, T.: Arctic amplification dominated by temperature feedbacks in contemporary climate models. Nature Geoscience, **7**, 181-184. doi:10.1038/ngeo2071, 2014.

Rogelj, J. and Knutti, R.: Geosciences after Paris, Nat. Geosci., 9, 187-189, 2016.

Samset, B. H., Myhre, G., Schulz, M., Balkanski, Y., Bauer, S., Berntsen, T. K., Bian, H., Bellouin, N., Diehl, T., Easter, R. C., Ghan, S. J.,

Iversen, T., Kinne, S., Kirkevåg, A., Lamarque, J.-F., Lin, G., Liu, X., Penner, J. E., Seland, Ø., Skeie, R. B., Stier, P., Takemura, T., Tsigaridis, K., and Zhang, K.: Black carbon vertical profiles strongly affect its radiative forcing uncertainty. Atmos. Chem. Phys., 13, 2423–2434, doi:10.5194/acp-13-2423-2013, 2013.

Sand, M., Iversen, T., Bohlinger, P., Kirkevåg, A., Seierstad, I. A., Seland, Ø., and Sorteberg, A.: A standardized global climate model study showing unique properties for the climate response to black carbon aerosols, J. Climate <u>http://dx.doi.org/10.1175/JCLI-D-14-00050.1</u>, 2015.

Sanderson, B. M., Xu, Y., Tebaldi, C., Wehner, M., O'Neill, B., Jahn, A., Pendergrass, A. G., Lehner, F., Strand, W. G., Lin, L., Knutti, R., and Lamarque, J. F.: Community climate simulations to assess avoided impacts in 1.5 and 2 C futures. Earth Syst. Dynam., 8, 827–847, https://doi.org/10.5194/esd-8-827-2017, 2017.

Schleussner C.-F., Rogelj, J., Schaeffer, M., Lissner, T., Licker, R., Fischer, E. M., Knutti, R., Levermann, A., Frieler, K., and Hare, W.: Science and policy characteristics of the Paris Agreement temperature goal, Nat. Clim. Change, 6, 827–835, doi:10.1038/nclimate3096, 2016.

Screen, J. A.: Arctic amplification decreases temperature variance in northern mid- to high-latitudes. Nature Clim. Change **4**, 577-582. 2014.

Screen, J. A.: Climate science: far-flung effects of Arctic warming. Nat. Geosci., 10, 253–254. 2017a.

Screen, J. A.: Simulated atmospheric response to regional and pan-Arctic sea ice loss, J. Clim., 30, 3945-3962, 2017b.

Screen, J. A. and Francis, J. A.: Contribution of sea-ice loss to Arctic amplification is regulated by Pacific
Ocean decadal variability. Nature Clim. Change, 6. doi: 10.1038/NCLIMATE3011. 2016.

Screen, J. A., and Simmonds, I.: Exploring links between Arctic amplification and mid-latitude weather, Geophys. Res. Lett., 40, 959–964, doi: 10.1002/grl.50174, 2013.

Screen, J. A., Deser, C., Smith, D. M., Zhang, X., Blackport, R., Kushner, P. J., Oudar, T., McCusker, K. E. and Sun, L.: Consistency and discrepancy in the atmospheric response to Arctic sea-ice loss across climate models, Nature Geoscience, doi:10.1038/s41561-018-0059-y, 2018.

30

5

15

Schwarz, J. P., Samset, B. H., Perring, A. E., Spackman, J. R., Gao, R. S., Stier, P., Schulz, M., Moore, F. L., Ray, E. A., and Fahey, D. W.: Global-scale seasonally resolved black carbon vertical profiles over the Pacific, Geophys. Res. Lett., 40, 5542–5547, doi:10.1002/2013gl057775, 2013.

Shaw, T. A., Baldwin, M., Barnes, E. A., Caballero, R., Garfinkel, C. I., Hwang, Y.-T., Li, C., O'Gorman, P. A., Rivière, G., Simpson, I. R., and Voigt, A.: Storm track processes and the opposing influences of climate change, Nat. Geosci., 9, 656–664, https://doi.org/10.1038/ngeo2783, 2016.

Sigmond, M., Fyfe, J. C., Swart, N. C.: Ice-free Arctic projections under the Paris Agreement. Nature Climate Change, 8, pp 404–408, doi: 10.1038/s41558-018-0124-y, 2018.

5

Seland, Ø. and Debernard, J. B., Sensitivities of Arctic Sea-ice in Climate Modelling, In: ACCESS Newsletter. No. 9, 10-13, 2014. <u>http://www.access-eu.org/en/publications/access_newsletter.html</u>.

Seneviratne S. I., R. Wartenburger, B. P. Guillod, A. L. Hirsch, M. M. Vogel, V. Brovkin, D. P. van Vuuren, N. Schaller, L. Boysen, K. V. Calvin, J. Doelman, P. Greve, P. Havlik, F. Humpenöder, T. Krisztin, D.

Mitchell, A. Popp, K. Riahi, J. Rogelj, C.-F. Schleussner, J. Sillmann and E. Stehfest: Climate extremes, land-climate feedbacks and land-use forcing at 1.5°C. Phil. Trans. R. Soc. A 376: 20160450. doi.org/10.1098/rsta.2016.0450, 2018.

Smith, D. M., Screen, J. A., Deser, C., Cohen, J., Fyfe, J. C., García-Serrano, J., Jung, T., Kattsov, V., Matei, D., Msadek, R., Peings, Y., Sigmond, M., Ukita, J., Yoon, J.-H., and Zhang, X.: The Polar Amplification

15 Model Intercomparison Project (PAMIP) contribution to CMIP6: investigating the causes and consequences of polar amplification, Geosci. Model Dev. Discuss., https://doi.org/10.5194/gmd-2018-82, in review. 2018.

Stjern, C. W., Samset, B. H., Myhre, G., Forster, P. M., Hodnebrog, Ø., Andrews, T., Boucher, O., Faluvegi, G., Iversen, T.,

Kasoar, M., Kharin, V., Kirkevåg, A., Lamarque, J.-F., Olivié, D., Richardson, T., Shawki, D., Shindell, D.,
Smith, C. J., Takemura, T., and Voulgarakis, A.: Rapid adjustments cause weak surface temperature response to increased black carbon concentrations., J. Geophys. Res., doi:10.1002/2017JD027326, 2017.

Taylor, K. E., Stouffer, R. J., and Meehl, G. A: An overview of CMIP5 and the experiment design, Bull. Am. Meteorol. Soc., 90, 485–498, doi:10.1175/BAMS-D-11-00094.1, 2012.

Tibaldi, S. and Molteni, F.: On the operational predictability of blocking. Tellus, 42A, 343–365, 1990.

25 van Vuuren, D. P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G. C., Kram, T., Krey, V., Lamarque, J.-F., Masui, T., Meinshausen, M., Nakicenovic, N., Smith, S. J., and Rose, S. K.: The representative concentration pathways: an overview. Clim. Change, 109, 5–31. DOI: 10.1007/s10584-011-0148-z, 2011.

Vihma, T.: Weather extremes linked to interaction of the Arctic and Midlatitudes. Clim. Extremes 226, 39– 30 50, 2017.

Wehner, M., Stone, D., Mitchell, D., Shiogama, H., Fischer, E., Graff, L. S., Kharin, V. V., Lierhammer, L., Sanderson, B., and Krishnan, H.: Changes in extremely hot days under stabilized 1.5 and 2.0 °C global

warming scenarios as simulated by the HAPPI multi-model ensemble. Earth Syst. Dynam., 9, 299–311, doi.org/10.5194/esd-9-299-2018, 2018.

Williams, K. D., Senior, C. A., and Mitchell, J.F.: Transient Climate Change in the Hadley Centre Models: The Role of Physical Processes. J. Climate, 14, 2659–2674, <u>https://doi.org/10.1175/1520-0442(2001)014<2659:TCCITH>2.0.CO;2</u>, 2001.

5

10

Winton, M.: Amplified Arctic climate change: What does surface albedo feedback have to do with it? Geophysical Research Letters, 33 (3), doi:10.1029/2005GL025244, 2006.

Woollings, T., Barriopedro, D., Methven, J., Son, S.-W., Martius, O., Harvey, B., Sillmann, J., Lupo, A. R., and Seneviratne, S.: Blocking and its Response to Climate Change. Current Climate Change Reports, 4, 287–300, doi.org/10.1007/s40641-018-0108-z, 2018.

Tables

Table 1: overview of the NorESM1-X versions referred to in this paper.

X =	Definition	Purpose	References
М	Fully coupled GCM for CMIP5 with concentration-driven GHGs: 2 ° atmosphere and land, 1 ° ocean and sea ice. 26 atmospheric levels, model top at 2.194 hPa.	Reference for model evaluation of NorESM1-Happi	Bentsen et al. (2013); Iversen et al. (2013)() Kirkevåg et al. (2013)
Наррі	Fully coupled GCM. Differences from NorESM1-M: 1 ° atmosphere and land; adjusted aging of snow on sea ice, with reduced albedo; bug-fix in the aerosol scheme, with faster removal of BC particles.	Basic GCM evaluation (Table S1); Coupled model scenarios targeting 1.5 K and 2.0 K above piControl	Seland and Debernard (2014)
HappiSO	Atmosphere, land and sea-ice models from NorESM1-Happi with slab-ocean replacing full ocean model.	Estimate equilibrium climate sensitivity (ECS); extend HAPPI AMIP-type runs which enables sea- ice response (Table 2)	
HappiAMIP	Atmosphere and land models from NorESM1-Happi with 1 ° resolution, set up with prescribed SST and sea ice.	Contribute to HAPPI: ensembles of AMIP-type runs with prescribed SST and sea ice, targeting present- day (2006–2015), 1.5 K, and 2.0 K above pre-industrial.	Mitchell et al. (2017); http://www.hap pimip.org/

Table 2: overview of the NorESM1-HappiSO experiments and their calibration. Q_f is the net divergence of heat not accounted for by the explicit processes, which is needed to maintain a stable climate with a predefined geographical distribution of SST. In SO-PD, SO-15, and SO-20, a restoring term – $(SST - SST_{ext})/\tau$ is included in Q_f , where $\tau = 30$ days is the applied time-scale of adjustment. Notice that sea ice is not restored except for via the indirect effect of the SST restoring term.

Name	Definition	Calibration	Length (years)
SO- piControl	Pre-industrial 1850 control run with constant external forcing.	Q_f calculated using h_{mix} , SST, and F_{net} (see Eq. 1) from a stable control simulation, piControl, for 1850 with NorESM1-Happi.	150
SO-4×CO ₂	Scenario-run with constant 4xCO ₂ mixing ratio.	As for SO-piControl	150
SO-PD	Present-day (2005–2016) equilibrium control.	 Q_f calculated with SST restored to 12- year averaged SST_{ext} determined by the Operational Sea Surface Temperature and Sea Ice Analysis (OSTIA) for 2005–2016 (Donlon et al., 2012), thus reducing SST biases. No restoring of sea ice. 	150
SO-15	Equilibrium climate change for an global surface air temperature response of 0.7 K above PD.	Forcing agents as in AMIP-15. Q_f calculated as for SO-PD by adding the CPL-15–CPL-PD increments to the OSTIA (2005–2016) climatology.	150
SO-20	Equilibrium climate change for an global surface air temperature response of 1.2 K above PD.	Forcing agents as in AMIP-20. Q_f calculated as for SO-15 using the CPL-20–CPL-PD increments.	150

Table 3: the NH and SH polar amplification factor (NH-PAF and SH-PAF) and global-mean near-surface temperature (T_{as}) in the PD experiments and differences associated with 1.5 K warming, 2.0 K warming, and the 0.5 K difference for NorESM1-Happi, NorESM1-HappiSO and NorESM1-HappiAMIP. PAF is defined as $\Delta T_{Polar}/\Delta T_{Global}$, where T is the near-surface temperature, and the Global and Polar (poleward of 60 °) subscripts indicate the averaging region.

	Period or Difference	NH-PAF	SH-PAF	T _{as} K
NorESM1-	AMIP-PD			287.30
HappiAMIP	AMIP-15-AMIP-PD	2.34	1.62	0.71
125×10	AMIP-20-AMIP-PD	2.17	1.35	1.20
years	AMIP-20-AMIP-15	1.93	0.95	0.49
	SO-PD			287.1 <mark>3</mark>
NorESM1- Happi <mark>SO</mark>	SO-15–SO-PD	2.98	-0.04	0. <mark>56</mark>
90 years	SO-20–SO-PD	2. <mark>68</mark>	0.30	1.02
·	SO-20–SO-15	2.2 <mark>9</mark>	0.77	0.43
	CPL-PD			286. <mark>72</mark>
NorESM1- Happi	CPL-15-CPL-PD	3. <mark>60</mark>	0.2 <mark>3</mark>	0. <mark>69</mark>
90 years	CPL-20-CPL-PD	2.99	0.5 <mark>6</mark>	1.15
	CPL-20-CPL-15	2.81	1.06	0.46

	Period or Difference	T _{Land} K	T _{Land} K	P _{Land} mm d ⁻¹	P _{Land} mm d ⁻¹	$\frac{\text{AREA}_{\text{Sealce}}^{\text{DJF}}}{10^6 \text{km}^2}$	AREA ^{JJA} 10 ⁶ km ²
NorESM1- HappiAMIP 125×10 years	AMIP-PD	265.87	292.62	1.214	2.532	11.26	5.81
	AMIP-15-AMIP-PD	+1.52	+0.84	+0.070	+0.104	-0.97	-0.54
	AMIP-20-AMIP-PD	+2.36	+1.65	+0.091	+0.139	-1.36	-0.86
	AMIP-20-AMIP-15	+0.83	+0.81	+0.021	+0.035	-0.39	-0.32
NorESM1- HappiSO 90 years	SO-PD	265.30	292.44	1.212	2.559	12.52	5.48
	SO-15–SO-PD	+1.46	+1.12	+0.041	+0.120	-0.65	-0.86
	SO-20–SO-PD	+2.19	+1.87	+0.078	+0.126	-1.02	-1.41
	SO-20–SO-15	+0.73	+0.75	+0.036	+0.006	-0.36	-0.55
NorESM1- Happi 90 years	CPL-PD	265.33	291.04	1.248	2.337	12.51	7.59
	CPL-15-CPL-PD	+1.44	+1.14	+0.048	+0.136	-1.41	-1.73
	CPL-20-CPL-PD	+2.41	+1.86	+0.073	+0.161	-1.93	-2.29
	CPL-20-CPL-15	+0.97	+0.71	+0.025	+0.025	-0.51	-0.56

Table 4: Similar as Table 3, but for near-surface temperature over land, precipitation on land, and sea-ice area in the NH (20 °N–90 °N) during winter (DJF) and summer (JJA).

Figures



Figure 1: Atmospheric CO₂ concentration for the 1.5 K and 2.0 K warming experiments with NorESM1-Happi. The 1.5 K experiment (black dotted line) initially follows RCP2.6 (blue solid line). At year 2095 the concentration deviates from RCP2.6, staying constant until year 2170, and decreases thereafter. The 2.0 K experiment (black dashed line) similarly follows RCP4.5 (red solid line) at first, but branches off at year 2050. The concentration is then constant until year 2170 before decreasing in the same fashion as in the 1.5 K experiment. Units are ppm.

10

15



Figure 2: Time-evolution of the global-mean near-surface temperature response in the Hist1 experiment (1850–2005; blue) and the CPL-15 (2006–2230; blue) and the CPL-20 experiment (2006–2230; red) relative to pre-industrial conditions (years 1850–1852). A three-year running average is used for both curves. Units are K.



Figure 3: Ocean heat uptake as a function of time in the CPL-15 (a) and CPL-20 (b) experiments. Shown is the heat uptake for depths 0–200 m (orange shading), 200–1000 m (green shading), 1000–2000 m (blue shading), 2000–3000 (pink shading), and below 3000 m (dark pink shading). Dashed vertical lines emphasize the time period analyzed in this study. Units are W m⁻².



Figure 4: Time-evolution of the maximum in the Atlantic meridional overturning circulation (AMOC) at 26.5 °N in Hist1 and RCP2.6 (black) and in the 1.5 K (red) and 2.0 K (blue) warming experiments with NorESM1-Happi. A 10-year running average is used for all curves. Units are Sv.



Figure 5: Near-surface temperature bias relative to ERA-Interim (colors) and near-surface temperature climatology (black contours; 260 to 350 K in increments of 10 K) for PD experiments from NorESM1-Happi (left), NorESM1-HappiSO (middle), and NorESM1-HappiAMIP (right). We use years 1986–2015 from ERA-Interim. The global-mean ensemble-mean bias is given in the upper-right corner of each panel. Units are K (a–1).



Figure 6: The annual-mean ocean heat flux Q_f needed in NorESM1-HappiSO to maintain a stable PD climate that is close to the observed SST used during calibration (a), and the change in Q_f for SO-15 (b) and SO-20 (c) compared to SO-PD. Negative values contribute to increasing SST (Eq. 1). Units are W m⁻² (a–c).



Figure 7: Time-evolution of global-mean near-surface temperature for Hist1 (1850–2005) and CPL-15 and CPL-20 (2005–2200) from NorESM1-Happi relative to the 1850–1899 average. Fields are shown for the global average (blue) and for an average taken over the area north of 65 °N (red), i.e. ca. 4.7 % of the global area. Units are K.



Figure 8: Zonal-mean temperature response (K) relative to PD (colors) and climatology (solid black contours; 210 K to 285 K in increments of 15 K) for the 1.5 K experiment from NorESM1-Happi (left; a, d), NorESM1-HappiSO (middle; b, e), and NorESM1-HappiAMIP (right; c, f). Fields are shown for DJF (top row; a–c) and JJA (bottom row; d–f). Units are K (a–f).



Figure 9: Upper tropospheric temperature contrast in the PD (grey), 1.5 K (blue), and 2.0 K (red) experiments from NorESM1-Happi (left column; a,d,g,j), NorESM1-HappiSO (middle column; b,e,h,k), and NorESM1-HappiAMIP (right column; e,f,i,l) for DJF (top row; a-c), MAM (middle row; d-f), JJA (third

5 row; g–i), and SON (bottom row; j–l). The upper-level temperature contrast $\Delta T200$ is defined as the 200 hPa temperature difference between an area over the tropics (30 °S-30 °N) and an area over the Arctic (poleward of 60 °N). The white lines within the boxes indicate the median, the boxes indicate the inter-quartile range, and the whiskers the full spread of the different decades in each experiment (9 in NorESM1-Happi, 9 in NorESM1-HappiSO, and 125 in NorESM1-HappiAMIP). The dashed horizontal lines emphasize the median values. Numbers are shown on the right side of each panel for ERA-Interim (top black), PD (second from

top, grey), the change with 1.5 K warming relative to PD (third from top, blue), and the change with the additional 0.5 K warming (bottom, red). Units are K (a–l).



Figure 10: As in Figure 9, but for the lower tropospheric temperature contrast at 850 hPa (ΔT 850).



Figure 11: Upper-level storm track bias relative to ERA-Interim (colors) and climatology (black contours; 8 m to 70 m in increments of 8 m) for the PD experiment from NorESM1-Happi (left, a, d, g, j), NorESM1-HappiSO (middle; b, e, g, k), and NorESM1-HappiAMIP (right; c, f, I, l) for DJF (top row; a–c), MAM

(second row; d–f), JJA (third row; g–i), and SON (bottom row; j–l). The bias is computed relative to ERA-Interim for years 1986–2015. The numbers in the upper-right corners of each plot give the mean bias for the area shown on the plot (latitudes poleward of 20 °N). Units are m (a–l).



Figure 12: Changes in upper-level storm-track activity relative to PD (colors) and climatology (black contours; 40 to 240 m² s⁻² in increments of 40 m² s⁻²) for the 1.5 K experiment from NorESM1-Happi (left; a, d, g, j), NorESM1-HappiSO (middle column; b, e, h, k), and NorESM1-HappiAMIP (right; e, f, i, l) for DJF (top row; a–c), MAM (middle row; d–f), JJA (third row; g–i), and SON (bottom; j–l). The storm tracks are represented in terms of bandpass-filtered EKE (eddy kinetic energy) at 250 hPa. The white dots indicate that the differences

are not significant at the 5 % level according to the Welch t-test. Units are m² s⁻² (a–l).



Figure 13: As in Figure 12, but for the upper-level storm track response to the additional 0.5 K warming (i.e. the difference between the respective 2.0 K and 1.5 K experiments).



Figure 14: Changes in the low-level storm-track activity relative to PD (colors) and PD climatology (black contours; -12 to 12 K m s⁻² in increments of 4 K m s⁻²) for the 1.5 K experiment from NorESM1-Happi (left; panels a and d), NorESM1-HappiSO (middle; panels b and e) and NorESM1-HappiAMIP (right; panels c and f) for DJF (top; panels a–c) and and JJA (bottom; panels d–f). The storm tracks are represented in terms of the bandpass-filtered eddy heat flux $\overline{v' T'}$ at 850 hPa. The white dots indicate that the differences are not significant at the 5 % level according to the Welch t-test. Units are K m s⁻² (a–f).



Figure 15: As in Figure 14, but for the low-level storm-track response to the additional 0.5 K warming (i.e. the difference between the 2.0 K and 1.5 K experiments).



Figure 16: PD climatology of blocking frequency from NorESM1-Happi (a-b), NorESM1-HappiSO (c-d), and NorESM1-HappiAMIP (e-f) for DJF (left; a, c, d) and JJA (right; b, d, f). Shown are the mean (solid black curve) and the spread (\pm one standard deviation) computed over the number of available decades (9 for 5 NorESM1-Happi and NorESM1-HappiSO, and 125 for NorESM1-HappiAMIP). Blocking frequency from ERA-Interim is shown for the period 1986–2015 (dotted black). The blocking events are identified using the the vTM index (Tibaldi and Molteni, 1990; Pelly and Hoskins, 2003), as in Iversen et al. (2013). It is based on the TM-index (Tibaldi and Molteni, 1990), which uses a persistent reversal of the meridional gradient of the 500-hPa geopotential height around the predefined central blocking latitude at 50 °N as an indicator for 10 blocking. The reversal must be present at 7.5 ° consecutive longitudes and persist for at least 5 days. In the vTM index the requirement of a predefined central blocking latitude is relaxed in order to reduce spurious detection (Pelly and Hoskins (2003). The central latitude is allowed to vary with longitude following the latitude of the maximum in the climatological storm track (using bandpass-filtered geopotential height at 500 hPa). To account for the seasonal cycle of the cyclone activity, the central latitude for a given month is 15 calculated as the climatological 3-month moving average centred on that month. Units are % (a-f).



Figure 17: Change in blocking frequency (solid black line with red and blue shading) in the 1.5 K experiment relative to PD (top three rows; a–f) and for the additional 0.5 K of warming (2.0 K–1.5 K; bottom three rows; g–l), shown along with the blocking climatology for the PD experiment (dotted black line). The fields are shown for the NorESM1-Happi (a, b, g, h), NorESM1-HappiSO (c, d, i, j), and NorESM1-HappiAMIP (e, f, k, l) during DJF (left; a, c, e, g, i, k) and JJA (right; b, d, f, h, j, l). The asterisks along the x-axis indicate where the changes at that longitude are statistically significant at the 5 % level according to the Welch t-test. Note that the left y-axis is for the difference field and the right y-axis is for the climatology. Units are % (a–l)



Figure 18: NH monthly-mean sea ice concentrations for PD (top; a–d), the 1.5 K warming relative to PD (second row; e–h), and the 0.5 K warming (bottom row; i–l) from NorESM1-Happi (first and third column; a, c, e, h, i, k) NorESM1-HappiSO (second and fourth column; b, d, f, h, j, l). Fields are shown for March (first and second column; a, b, e, f, i, j) and September (third and fourth column; c, d, g, h, k, l). The concentrations from the SO model are averaged over 90 years after 60 years of spin-up. The PD results (colors; top color bar) are shown together with observational estimates (OSI-SAF, 2017; solid black contours). Differences that are not statistically significant at the 5 % level according to the Mann-Whitney U test are marked with black dots. Units are % of ocean surface area (a–l).



Figure 19: The relative occurrence of NH monthly-mean sea-ice extent in September for observations (black bars; OSI-SAF, 2017), the PD experiments (blue bars), and the 1.5 K and 2.0 K warming experiments from NorESM1-Happi (a) and NorESM1-HappiSO. The sea-ice extent is binned in 1.0×106 km² increments. The observations are from 1996–2015 (20 values) in (a) and from 2005–2015 (11 values) in (b). The PD values from NorESM1-Happi and NorESM1-HappiSO is the default 90-year periods (Sect. 2).