



Climate sensitivity estimates – sensitivity to radiative forcing time series and observational data

Ragnhild Bieltvedt Skeie¹, Terje Berntsen^{1,2}, Magne Aldrin³, Marit Holden³, and Gunnar Myhre¹

¹CICERO-Center for International Climate and Environmental Research – Oslo, PB. 1129 Blindern, 0318 Oslo, Norway.

²Department of Geosciences, University of Oslo, PB. 1047 Blindern, 0316 OSLO, Norway.

³Norwegian Computing Center, PB. 114 Blindern, 0314 Oslo, Norway

Correspondence to: Ragnhild Bieltvedt Skeie (r.b.skeie@cicero.oslo.no)

Abstract. Inferred Effective Climate Sensitivity (ECS_{inf}) is estimated using a method combining radiative forcing (RF) time series and several series of observed ocean heat content (OHC) and near-surface temperature change in a Bayesian-framework using a simple energy balance model and a stochastic model. The model is updated compared to our previous analysis by using recent forcing estimates from IPCC, including OHC data for the deep ocean, and extending the time series to 2014. The mean value of the estimated ECS_{inf} is 2.0°C, with a 90% credible interval of 1.2-3.1°C. The mean estimate has recently been shown to be consistent with the higher values for the equilibrium climate sensitivity estimated by climate models. We show a strong sensitivity of the estimated ECS_{inf} to the choice of a priori RF time series, excluding pre-1950 data and the treatment of OHC data. Sensitivity analysis performed by merging the upper (0-700m) and the deep ocean OHC or using only one OHC data set (instead of four in the main analysis), both give an enhancement the mean ECS_{inf} by about 50% from our best estimate.

1 Introduction

A key question in climate science is how the global mean surface temperature (GMST) responds to changes in greenhouse gases or other forcings. The climate sensitivity is determined by complex feedbacks that operate on very different timescales and may depend on the transient climate state. The standard metric for climate sensitivity is the equilibrium climate sensitivity (ECS) (or Charney sensitivity) given as the change in temperature at equilibrium for a doubling of CO₂, neglecting long-term feedbacks associated with the vegetation changes, carbon feedbacks and ice sheet dynamics. Estimates of the ECS are either based on complex climate models or observations of past climate (Collins et al., 2013; Knutti et al., 2017). The Intergovernmental Panel on Climate Change (IPCC) presented a likely (>66% probability) range for ECS of 1.5 to 4.5°C (Collins et al., 2013).



Regarding the Earth as a climate laboratory and the changes in atmospheric composition and land use over the historical record as a perturbation experiment, observational based analysis of Earth's Energy Budget have been used to infer the climate sensitivity (Forster, 2016). Since the current climate is in a non-equilibrium state observationally based methods can only account for the feedbacks operating during the historical period. Thus, these estimates are often referred to as inferred or effective climate sensitivity (Armour, 2017;Forster, 2016) and are generally significantly lower than ECS estimates from Atmosphere-Ocean General Circulation Models (AOGCMs).

Since IPCCs fifth assessment report (AR5) there has been an improved understanding of the causes of the differences in estimates of climate sensitivity from climate models and observational based methods, directed to two main reasons. First, recent analysis of time-varying feedbacks in AOGCMs simulations from Coupled Model Intercomparison Project Phase 5 (CMIP5) (Proistosescu and Huybers, 2017;Armour, 2017;Andrews et al., 2015) have indicated that in most AOGCMs the net feedbacks become more positive over time as a new equilibrium is approached. This is most likely due to evolution of the pattern of sea surface temperature increase in the Pacific and Southern Ocean and associated cloud feedbacks. Whether this slow warming has manifested itself in the climate record used for the analysis is the difference between effective and equilibrium climate sensitivity (Armour, 2017;Knutti et al., 2017). Second, ECS formally refers to global near-surface air temperature ('tas' in CMIP5 nomenclature) and in observational based methods observed surface temperature records which are a blend of air temperature over land and sea surface temperature over ocean are used in the estimation. According to Richardson et al. (2016), there is a general bias in the surface temperature records since water heats slower than the air above and due to undersampling in fast warming regions (e.g. the Arctic). Taking both effects into account, Armour (2017) shows that previous estimates of ECS_{inf} of about $2.0^{\circ}C$ are consistent with estimates of ECS of $2.9^{\circ}C$ from climate models.

Although it is now established that the ECS estimated by the use of complex climate models and ECS_{inf} estimated by using historical observations would differ, there is still considerable spread in ECS estimates from models and between observationally based ECS_{inf} estimates. The observational based methods and using complex models are complementary approaches to quantify the net effect of the feedbacks that determines the climate sensitivity. Complex climate models include processes that are highly parameterized, in particular the representation of clouds, precipitation and convection, and associated feedbacks, which are crucial for estimating the ECS (Bony et al., 2015;Tan et al., 2016). There are also a large spread in observational based estimates (Knutti et al., 2017). Better understanding of the feedbacks in the complex models as well as improvements and understanding differences among the observational based methods are needed.

Observational estimates of climate sensitivity can be improved using longer data series of higher quality (e.g. correcting for observational biases in temperatures or better forcing estimates) (Urban et al., 2014). Estimates can also be improved by including observational data on other climate variables, which were not previously available. Several studies indicate that the temporary slowdown in GMST in the beginning of the millennium coexisted with increased accumulation of heat in the deep ocean (e.g. Meehl et al., 2011;Meehl et al., 2013;Balmaseda et al., 2013;Watanabe et al., 2013;Chen and Tung, 2014;Lyman and Johnson, 2013). Johansson et al. (2015) found that if OHC change below 700 meter over this period were included in their observational based methods the mean value of ECS_{inf} increased.



In this study we use our estimation model (cf. sect. 2, Aldrin et al., 2012; Skeie et al., 2014) with additional and extended observational datasets (including heating rates of the deep ocean) and new forcing time series to update our estimate of ECS_{inf} . Our method is more complex than the common energy balance based estimates (Forster, 2016) in that we embed a simple climate model treating the NH and SH separately and includes a vertical resolution of the ocean (40 layers). The Markov Chain Monte Carlo method used, estimates not only the ECS_{inf} but simultaneously also provides posterior estimates of the radiative forcing. We carry out a number of sensitivity experiments to investigate causes of differences in observational based ECS_{inf} estimates due to differences in the input data (observations of surface temperature, OHC and RF).

2 Data and methods

2.1 The model

Our full model consists of a simple climate model (SCM) with an idealized representation of the Earth's energy balance, a data model that describes how observations are related to the process states, and finally a parameter model that expresses our prior knowledge of the parameters (Aldrin et al., 2012).

The true state of some central characteristics (\mathbf{g}_t) of the climate system in year t with corresponding observations can be written as $\mathbf{g}_t = \mathbf{m}_t(\mathbf{x}_{1750:t}, \boldsymbol{\theta}) + \mathbf{n}_t$. Here, \mathbf{m}_t is the output from the deterministic SCM, whereas $\mathbf{x}_{1750:t}$ (the RF from 1750 until year t) and $\boldsymbol{\theta}$ are the true, but unknown, input values to the SCM. $\boldsymbol{\theta}$ is a vector of seven parameters, each with a physical meaning. One of these parameters is the climate sensitivity. Finally, \mathbf{n}_t is a stochastic process, with three terms, representing long-term and short-term internal variability and model error (Skeie et al., 2014).

For \mathbf{g}_t we consider the surface temperatures separately at the northern and southern hemispheres and the OHC separately for 0-700m and below 700m. Each of these elements of \mathbf{g}_t are associated with one or more corresponding observational-based data series, with individual error terms (Table 1). To gain as much information as possible, we use several data sets simultaneously (Aldrin et al., 2012; Skeie et al., 2014). Most of the data series are provided with corresponding yearly standard errors. However, these are often small compared to the differences between the data series, indicating that the errors reported by the data providers are too small. Therefore, we only use the temporal profiles of the reported errors and estimate their magnitudes within the model.

The unknown quantities are given prior distributions as presented in Skeie et al. (2014). We apply a Bayesian approach on calibration of computer models and use Markov Chain Monte Carlo (MCMC) techniques to sample from the posterior distribution.

2.1 Set up

The starting point is the main results from Skeie et al. (2014) (hereafter named Skeie14) with some modifications (see Appendix A). These modifications changed the mean ECS_{inf} value from 1.84 to 1.95°C and the 90% credible interval (C.I.)



from (0.92, 3.18) °C to (1.04, 3.41) °C (Fig. 1, case A). The transient climate response (TCR) is calculated by running the model with 1% per year increase in CO₂ using the joint posterior distribution of the model parameters, see Fig. S2.

In case A, we used four hemispheric pairs of observational based estimates of surface temperatures from about 1880 to 2010 and three series for OHC above 700m from about 1950 to 2010, and RF from Skeie et al. (2011,2014) (Table 1). We now include new and improved knowledge of the forcing time series and add new data for OHC below 700 meter (ORAS4), as well as ORAS4 OHC data above 700 meter, and observational data extended to 2014 (case B) (see Appendix B). Note that the deep ocean OHC is added as a separate dataset and not merged with the upper ocean. The vector θ contains several parameters that determine how heat is mixed into the ocean (the rate and vertical structure). This includes the mixed layer depth, the air-sea heat exchange coefficient, the vertical diffusivity in the ocean and the upwelling velocity (see Schlesinger et al. (1992) and Aldrin et al. (2012) for details). Including data on OHC in the deep ocean thus has the potential to better constrain θ , as well as the posterior estimates of the effective radiative forcing.

We perform three sensitivity experiments to investigate the effects of the use of OHC data (cases C and D, sect. 4.1) and how sensitive the results are to pre-1950 data (case E, sect. 4.2).

3 Improved estimate of inferred equilibrium climate sensitivity

Here we presents our revised estimate of ECS_{inf} by replacing the RF prior with IPCC data, including OHC data below 700 meter and extending the time series to 2014. We replaced the Forc_Skeie14 with the AR5 effective radiative forcing (ERF) estimates (Myhre et al., 2013), including the AR5 uncertainties. ERF includes rapid adjustments allowing the full influence on clouds except through surface temperature changes.

We consider the analysis using the IPCC forcing estimates, including deep ocean OHC and extending the length of the input data series as the most trustworthy and physical based case and thus regard it as our main estimate of the ECS_{inf}, with a mean of 2.0°C, and a 90% C.I. of 1.2-3.1°C. The mean value is similar while the 90% C.I. is narrower compared to the refined Skeie et al. (2014) estimate (Fig. 1). The mean value of TCR is 1.4°C with 90% C.I. of 0.9-2.0°C (Fig. S2). The GMST change is well reproduced (Fig. 2, case B), and less of the recent GMST change is attributed to long term internal variability (Fig. S5a-b) compared to the refined Skeie et al. (2014) estimate.

The rate of change in anthropogenic forcing is larger between 1940 and 1970 using Forc_AR5 compared to Forc_Skeie14 (Fig. 3). The fit to the GMST in the 1980s-1990s improved (Fig. 2 case B vs. A). The root mean square error between 1980 and 1999 decreased from 0.123 to 0.077. Parts of the increase in GMST over the last decades is explained as long-term internal variability, but the amplitude decreases in case B compared to case A (Fig. S5a-b).

The prior anthropogenic mean forcing in 2010 increased from 1.5 to 2.3 Wm⁻² from case A to case B when Forc_AR5 replaced Forc_Skeie14. The posterior is much closer to the prior in case B compared to case A (Fig. 3), even though the prior in case B is less constrained around the mean value compared to case A. The same holds for the total forcing (Fig. S4). The 90% C.I. for the posterior anthropogenic forcing was 1.3 to 2.8 Wm⁻² in case A compared to 1.3 to 3.4 Wm⁻² in case B.



The upper limit of the 90% C.I. is shifted to larger values. Using Forc_AR5 allows for stronger positive forcing than using Forc_Skeie14. The most uncertain part of the time series of ERF is associated with aerosols. The posterior estimated net ERF for aerosols change between case A and B (from -1.0 Wm^{-2} and -0.9 Wm^{-2} respectively), while the total ERF in 2010 is 0.2 Wm^{-2} higher in case B. A relative weak aerosol-cloud interaction as included in Forc_AR5 is consistent with the recent
5 findings in Malavelle et al. (2017) on how sulphate aerosols from volcanic emissions influences clouds.

The ERFs in AR5 are based on an assessment of several studies reflecting improved knowledge of the forcing mechanisms compared to the one-model RF results used in Skeie14. The new ERFs gave a better posterior estimate of GMST (Fig. 2) and reduced change from prior to posterior forcing (Fig. 3). Remark that the number of forcing time series that can be combined was 18 in Skeie14, including three time series for volcanic and eight for aerosol, compared to only one time series for each
10 of these forcing mechanisms in Forc_AR5 (Table S1). This gives less flexibility in the time development of the forcing in case B compared to case A, however the GMST change is better reproduced in the 1980s-1990s using Forc_AR5 compared to Forc_Skeie14.

Ultimately, global climate change is governed by the radiative imbalance at the top of the atmosphere (TOA) and modulated by the internal variability. Forcing by greenhouse gases and aerosols as well as albedo changes, feedback processes and the
15 radiative responses to temperature changes determine this imbalance. With a positive net imbalance at TOA energy accumulates in the Earth system, mainly as increasing OHC (Church et al., 2011). Since OHC is the dominant energy storage in the system, these data series have profound influence on the ECS_{inf} estimates (Tomassini et al., 2007; Skeie et al., 2014; Aldrin et al., 2012; Johansson et al., 2015). We have extended our use of OHC data, so in addition to the three OHC data series above 700m we now include the ORAS4 data above and below 700m (Table 1) as two separate data sources. The
20 deep ocean OHC data gives a stronger constraint on the overall accumulation of heat in the system, and the posterior estimates of the parameters of θ that determine the vertical transport of heat in the ocean, the effective diffusivity and the upwelling velocity increase by 44 and 31% respectively. Having separate data series for the two ocean layers also provides information that influence the balance between negative (by aerosols) and positive forcings, since these forcings have different evolution over time (cf. sect. 4.1).

In Fig. 4 the observed and fitted OHC for case A and B are shown. Including data on OHC change below 700 m increases the total heat uptake. The increase in OHC above 700 m over the last decade is larger in case B compared to case A. In case B the increase in OHC above 700 m is larger than the observational data, while below 700 meter, the observed OHC increase is underestimated (Fig. 4). This is to be expected since the parameters of θ do not change over time. Thus the observed rapid change in OHC below 700 meter over the last years with corresponding slower warming above 700 meter, is attributed to
30 long-term internal variability (a part of the n_t term) in the model (Figure S5c-d).

The update of the ECS_{inf} from case A to B (changing the forcing prior and including OHC data below 700 meters as well as updated and extended data up to and including 2014) were done stepwise (see Appendix B). The ECS_{inf} estimate was very sensitive to the forcing time series used. The new ERFs were first implemented. That resulted in a shift of the ECS_{inf} estimate to lower values, with an ECS_{inf} mean value of $1.5 \text{ }^\circ\text{C}$ and a 90% C.I. ranging from 0.9 to 2.3°C . Including also the OHC



change below 700m increased the total heat uptake and thus the mean value of ECS_{inf} from 1.5 to 1.7°C (see Fig. S1f-g). The last step to update the ECS_{inf} estimate from case A to case B was to extend the data series used. Using observational data up to 2014 as well as the most recent version of the data prior to 2010 (Table 1) increased the mean value of ECS_{inf} to 2.0°C and the reduction in ECS_{inf} in the first step is more or less counteracted.

5 4 Sensitivity tests – the use of input data

We now investigate possible causes of differences in observational based ECS_{inf} estimates due to the use of input data. We analyze the impacts of different usage of the OHC data (cases C and D) and the treatment of uncertainties in the GMST data (case E).

4.1 The role of the use of OHC data

10 The vertical transport of heat in the SCM (with 40 vertical layers) is quite simple. Turbulent diffusion mixes heat down from the surface, while down welling transports heat directly to the deepest layer, i.e. no detrainment to intermediate layers (Aldrin et al., 2012). Therefore, one may argue that it might be more appropriate to only constrain the model with OHC data for the total depth of the ocean. Merging the OHC above and below 700m (Case C, see Appendix D) results in a substantial decrease in the posterior ERF from 2.5 to 1.8 Wm^{-2} (Fig. S7b-c) and an increase in the ECS_{inf} estimate from a mean value of
15 2.0°C in case B to 3.2°C (Fig. 1). Without the separate constraint on the OHC above and below 700m, the posterior warming of the ocean increases faster (compared to case B) over the last 20 years (Fig. 5). This is mainly caused by enhanced warming in the upper 700m (Fig. S6). This allows for a stronger negative ERF estimate for aerosols. While the prior and posterior radiative forcing in Case B is similar, the posterior net forcing in case C is shifted towards lower values (Fig. S7c) and hence a higher estimated ECS_{inf} (Fig. 1) compared to case B. However, the observations show a stronger recent increase
20 in heat in the deep ocean (c.f. sect. 3), so this test might overestimate the aerosol forcing strength and hence overestimate the ECS_{inf} . Since the IPCC best estimate of $-0.9 Wm^{-2}$ was published in 2013 for aerosols ERF, studies point in different directions of either enhanced or weakened aerosol ERFs (Tan et al., 2016; Gordon et al., 2016; Malavelle et al., 2017), and hence no reason to refine the IPCC 2013 aerosol ERF best estimate jet.

A unique feature with our method is that we use data from more than one observational dataset. It is obvious that it is easier
25 to fit a model to one data series, thus giving less uncertainty in the posterior estimates. In case D we test the effect of using one alternative time series for OHC (Levitus2000 data, see Appendix D). As in case C, we do not separate the OHC data above and below 700m. Quite similar to case C, there is a more rapid increase in the posterior estimate of total OHC (Fig. 5) compared to case B, the increased warming is mostly in the upper 700 meter (Fig. S6) and the posterior forcing is shifted to lower values than in the prior (Fig. S7d). In case D the estimated mean ECS_{inf} is 2.8°C with a 90% C.I. of 1.5 to 4.6°C (Fig 1,
30 case F). This is higher than in case B, but lower than for case C.



The estimated total OHC has a narrower range when OHC above and below 700 meter are merged (Fig. 5, left panel). The range is also narrower in case D than in case C. As expected, using several data series for OHC (Case B: 5, Case C: 4, Case D: 1) increase the posterior observational error. Note that the magnitude of the observational errors are estimated (Aldrin et al., 2012; Skeie et al., 2014). In case D, the posterior standard deviation of the observed OHC is similar to the reported standard deviation, while using several OHC time series the posterior standard deviation is larger and arguably more correct than reported due to the large variability among the datasets. Hence, larger uncertainties in the observed OHC data result in larger uncertainties in the estimated OHC.

Johansson et al. (2015) used the same OHC data series as in our case D and a similar method, however their 90% C.I. for the OHC in the upper 2000 meter (their Fig. S5) is even narrower. This might not only be due to the use of one OHC dataset. While we estimate the magnitude of the observational error, Johansson et al. (2015) use the error given by the OHC data provider. In Johansson et al. (2015) the estimated uncertainties in OHC were smaller than the given observational uncertainties (their Fig. S5). The narrower ECS_{inf} range may primarily be because Johansson et al. (2015) assumed very small measurement errors in the most informative data (OHC), secondly that they ignored time correlation in observational errors and did not take into account long-term internal variability in the same degree as in our method.

To sum up, using several observational series (and estimate observational errors) increase the estimated observational errors to more realistic values, since data series are not well correlated, and hence increase the range of estimated OHC with implications on estimated ECS_{inf} .

4.2 The role of uncertainty estimates in the temperature series

The a priori standard deviation for the surface temperature data are quite different among the data sets (Fig. S8). The NCDC data has 3 to 5 times larger standard error prior to 1950 compared to after 1950, while it is more constant back to the 19th century for the three other data sets.

To investigate this, we re-estimated our model using data only after 1950, which is equivalent to assuming a very large uncertainty prior to 1950. The estimated magnitude of the ENSO-signal increases (Fig. S5a-b) since the data series are more correlated in the latter part of 20th century. For temperature, the model fits well to the observations of GMST, but with a larger 90% C.I. range (Fig. 2) and the observed NH and SH temperatures are well within the 90% C.I. of the model (Fig. S9). The mean ECS_{inf} increases from 2.0 to 2.2°C (Fig 1, case E vs. B) and the upper 90% C.I. limit increases from 3.1 to 3.8°C. Johansson et al. (2015) used only the NCDC data for GMST, thus the data prior to 1950 was given little weight when fitting the model. Our ENSO signal is now (case E) of similar magnitude as in Johansson et al. (2015) (their Fig. 1b). The ECS_{inf} uncertainty in this study is still larger and our mean value is slightly higher than their lower limit of 2°C.

Excluding data before 1950 also excludes the late 19th century period with a large volcanic eruption where the signal in the GMST data is small and quite uncertain (Santer et al., 2016). Santer et al. (2016) argued that the method in Johansson et al. (2015) down weights the volcanic forcing due to the small response of the Krakatau eruption in the temperature data. Johansson et al. (2016) responded that the observational uncertainty were large so the GMST data at that time will have a



limited effect. In our results, excluding observations before 1950, the GMST response following the Pinatubo eruption in 1991 increases (Fig. 2) and are similar to observations due to the larger ENSO signal and stronger posterior volcanic signal. In the early period, the aerosol forcing had a larger relative contribution to total ERF causing a more uncertain forcing trend in the early period. Uncertainty in the temporal trend of the forcing is not included, and better representation of forcing uncertainties than the scaling approach is needed (Tanaka et al., 2009). Omitting data before 1950 (case E), when the net forcing is more uncertain (Stevens, 2013), makes it easier to fit the model to observations but the uncertainty in estimated ECS_{inf}, GMST and TCR increases (Fig. 1, 2 and Fig. S2).

5. Discussions and conclusions

Causes of differences in observational based estimates of ECS_{inf} due to the use of input data are analyzed and an updated ECS_{inf} estimate is presented using our Bayesian estimation model. Adding observational data from 2011 to 2014, OHC data below 700 meter and replacing forcing data with IPCC AR5 ERFs, the ECS_{inf} posterior mean was 2.0°C with 90 % C.I. of 1.2 to 3.1°C. The mean value is similar and the range is slightly narrower than the refined Skeie14 estimated (Fig. 1 case B vs. A). The mean ECS_{inf} estimate is larger than in Skeie et al. (2014). Although the estimate in case A and B are quite similar, the ECS_{inf} estimate shifted to lower values when Forc_AR5 replaced Forc_Skeie14 and shifted to larger values when OHC data below 700m were included. The ECS_{inf} estimate was very sensitive to the forcing data used and we showed that the ECS_{inf} estimate was sensitive to the assumed uncertainties in the GMST data and how the OHC data were treated.

Bayesian methods have recently been reviewed by Annan (2015) and Bodman and Jones (2016) and limitation by assuming constant sensitivity over time, the role of the ECS_{inf} prior distribution and equal efficacy for different forcings have been discussed. Recently, studies have suggested that assuming equal efficacy for all forcings bias the ECS estimate low (Marvel et al., 2015; Shindell et al., 2015) even when ERFs are used (see also Appendix D). In our approach, the efficacy is implicitly included in the forcing uncertainty and thus accounted for.

The fit to the temperature data in the 1980s and 1990s improved using Forc_AR5 instead of Forc_Skeie14 indicating that the forcing trend over this period is better represented in Forc_AR5 compared to Forc_Skeie14. The trend in the forcing is more uncertain in the first half of the 20th century due to less dominance of CO₂, and in our method the same relative uncertainty for the prior forcing is used over the entire time period. A sensitivity simulation omitting observations before 1950, similar to making these observations very uncertain, gave better representation of the GMST in the latter part of the 20th century and an increased mean ECS_{inf}. Future work should include uncertainties in the temporal development of the forcing, and there is a clear need for an international effort to establish forcing time series, using a consistent forcing definition and allowing for uncertainties in emissions, to give a better representation of the temporal uncertainties.

Including OHC-data below 700 meter shifted the ECS_{inf} to higher values. The estimated ECS_{inf} was found to be very sensitive to how the OHC data were used. Including four OHC time series, but merging the data above and below 700 meter (case C), the ECS_{inf} mean value increased from 2.0 to 3.2°C. The probability of ECS_{inf} above 4.5°C increased to 13%, values



that are practically excluded in our main estimate (case B). Previous studies have used total column OHC data and due to the simple representation of the ocean one can argue that this is more appropriate. However, in case C most of the recent increase in OHC in the model occurred in the uppermost 700 meter allowing a stronger aerosol cooling (Fig. S7c) and hence a larger ECS_{inf} , while the observations indicate that the ocean is warming below 700 meter. Using only the total column
5 OHC might therefore overestimate the ECS_{inf} and the aerosol forcing strength. We recognize structural uncertainties in the model, and a multi-model intercomparison of observational methods using identical input data would be of great value to investigate these uncertainties.

Using only the Levitus2000 series for OHC for the total ocean column (case D), the ECS_{inf} 90% C.I. was shifted to lower values with a range from 1.5 to 4.6°C and the range shrunk compared to case C. The historical measurements of ocean
10 temperatures are sparse (Abraham et al., 2013), with large differences between the datasets. The temporal structure of the reported uncertainties differs, and the full uncertainties are often not assessed. Hence, relying on only one OHC series and its reported uncertainty may underestimate the observational uncertainties and hence overestimate the certainties in the estimated OHC with implications for the ECS_{inf} estimate.

Recent studies indicate that the upper-ocean warming is underestimated due to the gap-filling methods (Durack et al.,
15 2014;Li-Jing et al., 2015), in which case also the ECS_{inf} will be underestimated. Refining historical OHC estimates, not only the best value, but also the uncertainty is crucial for observational based ECS_{inf} estimation.

Other priorities is to improve the GMST series, including uncertainties, not only the recent trend (Karl et al., 2015;Cowtan and Way, 2014) but also for earlier time periods. Assuming a very large uncertainty prior to 1950, the GMST fit improved, ECS_{inf} mean increased while the estimated uncertainty ranges increased.

20 Our ECS_{inf} posterior mean was 2.0°C with 90 % C.I. of 1.2 to 3.1°C. This is consistent with a mean ECS of 2.9°C (Armour, 2017), which compares reasonably well with climate model estimates (Andrews et al., 2012;Forster et al., 2013). A final remark is that it is not obvious that the true ECS is a more relevant metric for the climate sensitivity than the ECS_{inf} in a policy context (i.e. the Paris agreement). The United Nations Framework Convention on Climate Change (UNFCCC) has not adopted a pre-defined definition of GMST and the stronger long-term feedbacks found in analysis of CMIP5 simulations
25 (Proistosescu and Huybers, 2017) operates on a time scale longer than the timescale for reaching 2°C.

Appendix A: Refinement of Skeie14

A few updates/corrections to Skeie14 (Fig. S1a) had to be made prior to the analyses presented in this study. In the Skeie14 study, the standard error of observed OHC above 700 meter for two out of the three series were constant in time, while for
30 the third dataset the standard error decreased with time. Due to the limited observational data back in history (e.g. Abraham et al., 2013), it is reasonable to assume that the shape of the standard error of observed global OHC increase back in time, as for the CSIRO series. Therefore, we now assume a common observational uncertainty temporal profile for OHC above 700



meter equal to CSIRO for all the OHC time series (Fig. S1b). Note that the magnitude of the observational errors are estimated in our approach (Aldrin et al., 2012; Skeie et al., 2014).

Another update of Skeie14 that was needed, was to use monthly volcanic RF data (Fig. S1c) compared to yearly data in Skeie14. In addition to the three global mean surface temperature (GMST) time series used in Skeie14, another time series
5 for GMST has been published recently (Cowtan and Way, 2014). This time series find a stronger increasing trend in temperature over the last decade compared to the HadCRUT4 data, due to their method of accounting for the unsampled regions in the world. This dataset is now included (Fig. S1d).

Our previous studies showed that the correlation between the observational errors in temperature data was almost uncorrelated with the observational errors in the OHC data. Therefore, to simplify the numerical computations, we from now
10 on assume that that these correlations are exactly zero (Fig. S1e).

The estimated ECS_{inf} for each step in the refinement of Skeie14 is presented in Fig. S1a-e.

Appendix B: The stepwise update of ECS_{inf}

The updated estimate of ECS_{inf} were done stepwise. Here these steps and the corresponding ECS_{inf} estimates are presented.

First, Forc_Skeie14 was replaced by Forc_AR5 in the estimation. With Forc_AR5 the prior anthropogenic mean forcing in
15 2010 increased from 1.5 to 2.3 Wm^{-2} , and the posterior forcing is much closer to the prior. The 90% C.I. for the posterior anthropogenic forcing was 1.3 to 2.8 Wm^{-2} using Forc_Skeie14 compared to 1.3 to 3.3 Wm^{-2} using Forc_AR5. The stronger forcing lead to a lower ECS estimate (Fig. S1f vs. e), reducing the mean ECS_{inf} to 1.5 °C (0.9 - 2.3°C). The fit to the GMST in the 1980s-1990s improved and the root mean square error between 1980 and 1999 decreased from 0.123 to 0.087 compared to case A.

20 So far, only OHC data in the upper 700 meter were used, leaving the model unconstrained with respect to the heating of the deeper ocean. We now include the ORAS4 data above and below 700m as two separate data sources. Similar to Johansson et al. (2015) we find that including OHC change below 700 m increases the total heat uptake and thus the mean value of ECS_{inf} (from 1.5 to 1.7°C, Fig. S1 g vs. f).

The last step to update the ECS_{inf} estimate is to extend the data series used. Some of the observational data series has been
25 updated, so first we use refined data up to 2010 (Fig. S1i) before we extend the data series to 2014 (cf. Appendix C). Using the refined data up to 2010, the estimated mean ECS_{inf} increased from 1.7 to 2.0°C, while extending the data up to 2014 slightly reduced the mean ECS_{inf} estimate (Fig. S1j). The upper bound of the 90% C.I. decreased from 3.3 to 3.1°C when the data series were extended from 2010 to 2014 (Fig. S1i and S1j) while the lower bound remained unchanged.

Appendix C: Extending data up to and including 2014

30 When extending the analysis from 2010 to 2014, not all the time series used in the estimation is available up to and including year 2014. Below is a description of how the different datasets are extended if not available up to 2014.



AR5 ERF: The end year for the forcing time series presented in AR5 is 2011 and has to be extended to 2014. For long-lived greenhouse gases the time series are extended using recent observations of global mean concentrations and the formulas relating concentrations and forcing used in Skeie et al. (2011). Tropospheric ozone, stratospheric ozone, aerosol ERF, land use change, BC on snow and volcanoes are kept constant 2011-2014. Stratospheric water vapor follow methane RF.

5 Contrails RF is extended using aircraft traffic data (<http://airlines.org/data/annual-results-world-airlines/>). Solar RF is extended using the Physikalisch-Meteorologisches Observatorium Davos (PMOD) composite (Frohlich and Lean, 2004). CSIRO: Data up to and including 2012 were downloaded. The time series were extended from 2012 to 2014 using the mean rate of change of the other OHC data. The uncertainty in 2014 and 2013 is set equal to the uncertainty in 2012.

ORAS4: Balmaseda et al. (2013) investigated the time evolution of global OHC at different depths of the ocean from 1958 to

10 2009 using the European Centre for Medium-Range Weather Forecasts ocean reanalysis system 4 (ORAS4). Five ensemble members of ORAS4 are generated that sample plausible uncertainties in the wind forcing, observation coverage, and the deep ocean. The ORAS4 system runs automatically in operations, with numerical weather prediction forcing and observations that are not manually quality controlled. The 1x1-degree Ocean potential temperature up to December 2014 are made available through the APDRC (http://apdrc.soest.hawaii.edu/datadoc/ecmwf_oras4.php) for one ensemble member.

15 The trend in OHC for the total depth and upper 700 meters from 2010 to 2014 based on the one ensemble member is used to extend the corresponding OHC data for all the five ensemble members from Balmaseda et al. (2013) up to 2014. The data after 2009 are based on the automatic ORAS4 system, and not quality controlled and the results in this paper using the data after 2009 should be interpreted by caution. The same method is used to extend the ORAS4 data from 2009 to 2010 (Fig. S1g-i). From the five ensemble members the estimate with uncertainty is calculated as the annual average and standard

20 deviation of OHC above and below 700 m. The standard deviations are modified by smoothing the curve (9-year moving average) since the curve was otherwise very static.

Appendix D: Sensitivity tests

Here we present in detail two of the sensitivity tests in section 4.

25 Case C: The OHC data used to constrain the model are the same series as in case B, but we do not separate the 0-700 meter from the deeper ocean. We use four data sets for total OHC by adding the ORAS4 700-2000 meter data to each of the four OHC above 700 meter estimates.

Case D: To test the effect of using one time series for the OHC we do a sensitivity test using only one OHC series and without separate constraint on the OHC above and below 700 meter. We choose to use the Levitus2000 time series, that is

30 the same OHC data as used in Johansson et al. (2015). The pentadal heat content are used from 1955 to 2012. The time series are extended to 2014 using the yearly OHC data for the upper 2000 meter from the same data source. We treat the pentadal observations as they were annual observations, and we use the OHC data for the upper 2000 meter as they were data for the total OHC. Observed energy stored below 2000 meter is not included in the estimation and hence the ECS might be



underestimated. Energy stored below 2000 meter is uncertain. Purkey and Johnson (2010) found that in the 1990s and 2000s there was an increase in OHC in the abyssal and deep Southern Ocean based on sparse observations from ships, but it is not clear if it is a long-term trend. Llovel et al. (2014) could not detect deep-ocean (below 2000 meter) contribution to sea level rise and energy budget between 2005 and 2013 using ocean observations and satellite measurements, however the
5 uncertainties are large.

Additional analyses:

Fig. S1h): Use the original ORAS4 standard error for OHC above and below 700 meter.

Fig. S1k): A sensitivity test using the same ENSO-index as in Johansson et al. (2015). Similar to a sensitivity test in
10 Johansson et al. (2015), the chosen ENSO-index had little effect on the ECS estimate.

Fig S1l): Shindell (2014) found that the transient climate sensitivity to historical aerosols and ozone is substantially greater than the transient climate sensitivity to CO₂ based on CMIP5 models. This enhanced sensitivity is primarily caused by more of the forcing being located in Northern Hemisphere where it triggers more rapid land responses and stronger feedbacks. Kummer and Dessler (2014) found that assuming larger efficacy (the amount of warming per unit global average forcing
15 divided by the warming per unit forcing from CO₂) for aerosol and ozone forcing increased the estimated equilibrium climate sensitivity. They suggested that forcing efficacy can explain the disagreement between estimates of climate sensitivity based on the twentieth century observational record and those based on climate models.

If we apply a efficacy of 1.5 for ozone, surface albedo, BC on snow and aerosols, which is the efficacy found in the analysis of Shindell (2014), the PDF of the ECS is shifted to larger values, with a 90% C.I. ranging from 1.2 to 3.7K. This is still
20 lower than the estimate by Kummer and Dessler (2014) of 3K and a 90 % C.I. of 1.9–6.8 when they included efficacy.

A recent study (Marvel et al., 2015) found that also when using ERF the TCR and ECR are biased low due to different response of the total forcing compared to CO₂ only. The efficacies presented is from one model only.

25 **Data availability.** Several publicly available data sets were used in this study. The specific references to the data sources are given in table 1. Model outputs are available upon request.

Author contributions. All authors designed the study and discussed the results; RBS and MH prepared the data; MH performed the simulations; RBS and MH made the figures; RBS prepared the paper with contributions from all co-authors.

Competing interests. The authors declare that they have no conflict of interest.

30 **Acknowledgements.** This research was supported by the Norwegian Research Council under the project EVA - Earth system modeling of climate Variations in the Anthropocene, grant number 229771. We kindly acknowledge the data providers listed in table 1 for providing the data for the analysis.



References

- Abraham, J. P., Baringer, M., Bindoff, N. L., Boyer, T., Cheng, L. J., Church, J. A., Conroy, J. L., Domingues, C. M., Fasullo, J. T., Gilson, J., Goni, G., Good, S. A., Gorman, J. M., Gouretski, V., Ishii, M., Johnson, G. C., Kizu, S., Lyman, J. M., Macdonald, A. M., Minkowycz, W. J., Moffitt, S. E., Palmer, M. D., Piola, A. R., Reseghetti, F., Schuckmann, K.,
- 5 Trenberth, K. E., Velicogna, I., and Willis, J. K.: A review of global ocean temperature observations: Implications for ocean heat content estimates and climate change, *Rev. Geophys.*, 51, 450-483, 10.1002/rog.20022, 2013.
- Aldrin, M., Holden, M., Guttorp, P., Skeie, R. B., Myhre, G., and Berntsen, T. K.: Bayesian estimation of climate sensitivity based on a simple climate model fitted to observations of hemispheric temperatures and global ocean heat content, *Environmetrics*, 23, 253-271, 10.1002/env.2140, 2012.
- 10 Andrews, T., Gregory, J. M., Webb, M. J., and Taylor, K. E.: Forcing, feedbacks and climate sensitivity in CMIP5 coupled atmosphere-ocean climate models, *Geophys. Res. Lett.*, 39, L09712, 10.1029/2012GL051607, 2012.
- Andrews, T., Gregory, J. M., and Webb, M. J.: The Dependence of Radiative Forcing and Feedback on Evolving Patterns of Surface Temperature Change in Climate Models, *J. Clim.*, 28, 1630-1648, 10.1175/jcli-d-14-00545.1, 2015.
- Annan, J. D.: Recent Developments in Bayesian Estimation of Climate Sensitivity, *Current Climate Change Reports*, 1-5,
- 15 10.1007/s40641-015-0023-5, 2015.
- Armour, K. C.: Energy budget constraints on climate sensitivity in light of inconstant climate feedbacks, *Nature Clim. Change*, 7, 331-335, 10.1038/nclimate3278, 2017.
- Balmaseda, M. A., Trenberth, K. E., and Källén, E.: Distinctive climate signals in reanalysis of global ocean heat content, *Geophys. Res. Lett.*, 40, 1754-1759, 10.1002/grl.50382, 2013.
- 20 Bodman, R. W., and Jones, R. N.: Bayesian estimation of climate sensitivity using observationally constrained simple climate models, *Wiley Interdisciplinary Reviews: Climate Change*, 7, 461-473, 10.1002/wcc.397, 2016.
- Bony, S., Stevens, B., Frierson, D. M. W., Jakob, C., Kageyama, M., Pincus, R., Shepherd, T. G., Sherwood, S. C., Siebesma, A. P., Sobel, A. H., Watanabe, M., and Webb, M. J.: Clouds, circulation and climate sensitivity, *Nature Geosci*, 8, 261-268, 10.1038/ngeo2398, 2015.
- 25 Chen, X., and Tung, K.-K.: Varying planetary heat sink led to global-warming slowdown and acceleration, *Science*, 345, 897-903, 10.1126/science.1254937, 2014.
- Church, J. A., White, N. J., Konikow, L. F., Domingues, C. M., Cogley, J. G., Rignot, E., Gregory, J. M., van den Broeke, M. R., Monaghan, A. J., and Velicogna, I.: Revisiting the Earth's sea-level and energy budgets from 1961 to 2008, *Geophys. Res. Lett.*, 38, L18601, 10.1029/2011gl048794, 2011.
- 30 Collins, M., R. Knutti, J. M. Arblaster, J.-L. Dufresne, T. Fichet, P. Friedlingstein, X. Gao, W. J. Gutowski, T. Johns, G. Krinner, M. Shongwe, C. Tebaldi, A. J. Weaver, and M. Wehner: Long-term Climate Change: Projections, Commitments and Irreversibility, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* edited by: Stocker, T. F., D. Qin, G.-K. Plattner, M.



- Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, and Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- Cowtan, K., and Way, R. G.: Coverage bias in the HadCRUT4 temperature series and its impact on recent temperature trends, *Q. J. R. Meteorol. Soc.*, 140, 1935-1944, 10.1002/qj.2297, 2014.
- 5 Durack, P. J., Gleckler, P. J., Landerer, F. W., and Taylor, K. E.: Quantifying underestimates of long-term upper-ocean warming, *Nature Clim. Change*, 4, 999-1005, 10.1038/nclimate2389, 2014.
- Forster, P. M., Andrews, T., Good, P., Gregory, J. M., Jackson, L. S., and Zelinka, M.: Evaluating adjusted forcing and model spread for historical and future scenarios in the CMIP5 generation of climate models, *J. Geophys. Res.*, 118, 1139-1150, 10.1002/jgrd.50174, 2013.
- 10 Forster, P. M.: Inference of Climate Sensitivity from Analysis of Earth's Energy Budget, *Annual Review of Earth and Planetary Sciences*, 44, 85-106, 10.1146/annurev-earth-060614-105156, 2016.
- Frohlich, C., and Lean, J.: Solar radiative output and its variability: evidence and mechanisms, *Astronomy and Astrophysics Review*, 12, 273-320, 2004.
- Gordon, H., Sengupta, K., Rap, A., Duplissy, J., Frege, C., Williamson, C., Heinritzi, M., Simon, M., Yan, C., Almeida, J.,
15 Tröstl, J., Nieminen, T., Ortega, I. K., Wagner, R., Dunne, E. M., Adamov, A., Amorim, A., Bernhammer, A.-K., Bianchi, F., Breitenlechner, M., Brilke, S., Chen, X., Craven, J. S., Dias, A., Ehrhart, S., Fischer, L., Flagan, R. C., Franchin, A., Fuchs, C., Guida, R., Hakala, J., Hoyle, C. R., Jokinen, T., Junninen, H., Kangasluoma, J., Kim, J., Kirkby, J., Krapf, M., Kürten, A., Laaksonen, A., Lehtipalo, K., Makhmutov, V., Mathot, S., Molteni, U., Monks, S. A., Onnela, A., Peräkylä, O., Piel, F., Petäjä, T., Praplan, A. P., Pringle, K. J., Richards, N. A. D., Rissanen, M. P., Rondo, L., Sarnela, N., Schobesberger,
20 S., Scott, C. E., Seinfeld, J. H., Sharma, S., Sipilä, M., Steiner, G., Stozhkov, Y., Stratmann, F., Tomé, A., Virtanen, A., Vogel, A. L., Wagner, A. C., Wagner, P. E., Weingartner, E., Wimmer, D., Winkler, P. M., Ye, P., Zhang, X., Hansel, A., Dommen, J., Donahue, N. M., Worsnop, D. R., Baltensperger, U., Kulmala, M., Curtius, J., and Carslaw, K. S.: Reduced anthropogenic aerosol radiative forcing caused by biogenic new particle formation, *PNAS*, 113, 12053-12058, 10.1073/pnas.1602360113, 2016.
- 25 Johansson, D. J. A., O'Neill, B. C., Tebaldi, C., and Haggstrom, O.: Equilibrium climate sensitivity in light of observations over the warming hiatus, *Nature Clim. Change*, 5, 449-453, 10.1038/nclimate2573, 2015.
- Johansson, D. J. A., O'Neill, B. C., Tebaldi, C., and Haggstrom, O.: Reply to 'Volcanic effects on climate', *Nature Clim. Change*, 6, 4-5, 10.1038/nclimate2861, 2016.
- Karl, T. R., Arguez, A., Huang, B., Lawrimore, J. H., McMahon, J. R., Menne, M. J., Peterson, T. C., Vose, R. S., and
30 Zhang, H.-M.: Possible artifacts of data biases in the recent global surface warming hiatus, *Science*, 348, 1469-1472, 10.1126/science.aaa5632, 2015.
- Knutti, R., Rugenstein, M. A. A., and Hegerl, G. C.: Beyond equilibrium climate sensitivity, 10, 727, 10.1038/ngeo3017, 2017.



- Kummer, J. R., and Dessler, A. E.: The impact of forcing efficacy on the equilibrium climate sensitivity, *Geophys. Res. Lett.*, 41, 2014GL060046, 10.1002/2014GL060046, 2014.
- Li-Jing, C., Jiang, Z., and Abraham, J.: Global Upper Ocean Heat Content Estimation: Recent Progress and the Remaining Challenges, *Atmos. Oceanic Sci. Lett.*, 8, 333-338, 10.3878/aosl20150031, 2015.
- 5 Llovel, W., Willis, J. K., Landerer, F. W., and Fukumori, I.: Deep-ocean contribution to sea level and energy budget not detectable over the past decade, *Nature Clim. Change*, 4, 1031-1035, 10.1038/nclimate2387, 2014.
- Lyman, J. M., and Johnson, G. C.: Estimating Global Ocean Heat Content Changes in the Upper 1800 m since 1950 and the Influence of Climatology Choice*, *J. Clim.*, 27, 1945-1957, 10.1175/JCLI-D-12-00752.1, 2013.
- Malavelle, F. F., Haywood, J. M., Jones, A., Gettelman, A., Clarisse, L., Bauduin, S., Allan, R. P., Karset, I. H. H.,
10 Kristjánsson, J. E., Oreopoulos, L., Cho, N., Lee, D., Bellouin, N., Boucher, O., Grosvenor, D. P., Carslaw, K. S., Dhomse, S., Mann, G. W., Schmidt, A., Coe, H., Hartley, M. E., Dalvi, M., Hill, A. A., Johnson, B. T., Johnson, C. E., Knight, J. R., O'Connor, F. M., Partridge, D. G., Stier, P., Myhre, G., Platnick, S., Stephens, G. L., Takahashi, H., and Thordarson, T.: Strong constraints on aerosol–cloud interactions from volcanic eruptions, *Nature*, 546, 485-491, 10.1038/nature22974, 2017.
- Marvel, K., Schmidt, G. A., Miller, R. L., and Nazarenko, L. S.: Implications for climate sensitivity from the response to
15 individual forcings, *Nature Clim. Change*, advance online publication, 10.1038/nclimate2888, 2015.
- Meehl, G. A., Arblaster, J. M., Fasullo, J. T., Hu, A., and Trenberth, K. E.: Model-based evidence of deep-ocean heat uptake during surface-temperature hiatus periods, *Nature Clim. Change*, 1, 360-364, 10.1038/nclimate1229, 2011.
- Meehl, G. A., Hu, A., Arblaster, J. M., Fasullo, J., and Trenberth, K. E.: Externally Forced and Internally Generated Decadal Climate Variability Associated with the Interdecadal Pacific Oscillation, *J. Clim.*, 26, 7298-7310, 10.1175/JCLI-D-12-
20 00548.1, 2013.
- Myhre, G., D. Shindell, F.-M. Bréon, W. Collins, J. Fuglestedt, J. Huang, D. Koch, J.-F. Lamarque, D. Lee, B. Mendoza, T. Nakajima, A. Robock, G. Stephens, T. Takemura, and Zhang, H.: Anthropogenic and Natural Radiative Forcing, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* edited by: Stocker, T. F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J.
25 Boschung, A. Nauels, Y. Xia, V. Bex, and Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- Proistosescu, C., and Huybers, P. J.: Slow climate mode reconciles historical and model-based estimates of climate sensitivity, *Science Advances*, 3, 2017.
- Purkey, S. G., and Johnson, G. C.: Warming of Global Abyssal and Deep Southern Ocean Waters between the 1990s and
30 2000s: Contributions to Global Heat and Sea Level Rise Budgets, *J. Clim.*, 23, 6336-6351, 10.1175/2010jcli3682.1, 2010.
- Richardson, M., Cowtan, K., Hawkins, E., and Stolpe, M. B.: Reconciled climate response estimates from climate models and the energy budget of Earth, *Nature Clim. Change*, 6, 931-935, 10.1038/nclimate3066, 2016.
- Santer, B., Solomon, S., Ridley, D., Fyfe, J., Beltran, F., Bonfils, C., Painter, J., and Zelinka, M.: Volcanic effects on climate, *Nature Clim. Change*, 6, 3-4, 10.1038/nclimate2859, 2016.



- Schlesinger, M. E., Jiang, X., and Charlson, R. J.: Implication of Anthropogenic Atmospheric Sulphate for the Sensitivity of the Climate System, in: *Climate Change and Energy Policy: Proceedings of the International Conference on Global Climate Change: Its Mitigation Through Improved Production and Use of Energy*, edited by: Rosen, L., and Glasser, R., American Institute of Physics, New York, 75-108, 1992.
- 5 Shindell, D. T.: Inhomogeneous forcing and transient climate sensitivity, *Nature Climate Change*, 4, 274-277, 10.1038/nclimate2136, 2014.
- Shindell, D. T., Faluvegi, G., Rotstayn, L., and Milly, G.: Spatial patterns of radiative forcing and surface temperature response, *J. Geophys. Res.*, 120, 5385-5403, 10.1002/2014JD022752, 2015.
- Skeie, R. B., Berntsen, T. K., Myhre, G., Tanaka, K., Kvalevåg, M. M., and Hoyle, C. R.: Anthropogenic radiative forcing
10 time series from pre-industrial times until 2010, *Atmos. Chem. Phys.*, 11, 11827-11857, 10.5194/acp-11-11827-2011, 2011.
- Skeie, R. B., Berntsen, T., Aldrin, M., Holden, M., and Myhre, G.: A lower and more constrained estimate of climate sensitivity using updated observations and detailed radiative forcing time series, *Earth Syst. Dynam.*, 5, 139-175, 10.5194/esd-5-139-2014, 2014.
- Stevens, B.: Climate science: Uncertain then, irrelevant now, *Nature*, 503, 47-48, 10.1038/503047a, 2013.
- 15 Tan, I., Storelvmo, T., and Zelinka, M. D.: Observational constraints on mixed-phase clouds imply higher climate sensitivity, *Science*, 352, 224-227, 2016.
- Tanaka, K., Raddatz, T., O'Neill, B. C., and Reick, C. H.: Insufficient forcing uncertainty underestimates the risk of high climate sensitivity, *Geophys. Res. Lett.*, 36, L16709, 10.1029/2009gl039642, 2009.
- Tomassini, L., Reichert, P., Knutti, R., Stocker, T. F., and Borsuk, M. E.: Robust Bayesian uncertainty analysis of climate
20 system properties using Markov chain Monte Carlo methods, *J. Clim.*, 20, 1239-1254, 10.1175/jcli4064.1, 2007.
- Urban, N. M., Holden, P. B., Edwards, N. R., Srivier, R. L., and Keller, K.: Historical and future learning about climate sensitivity, *Geophys. Res. Lett.*, 41, 2543-2552, 10.1002/2014gl059484, 2014.
- Watanabe, M., Kamae, Y., Yoshimori, M., Oka, A., Sato, M., Ishii, M., Mochizuki, T., and Kimoto, M.: Strengthening of ocean heat uptake efficiency associated with the recent climate hiatus, *Geophys. Res. Lett.*, 40, 3175-3179,
25 10.1002/grl.50541, 2013.




Figure 1: Posterior 90% C.I. for ECS_{inf} for the different analyses in this study. The estimated posterior mean is indicated by a dot. The IPCC AR5 likely range (>66% probability) for ECS is presented as gray shadings.




Figure 2: Observed and fitted (posterior mean) values for the GMST. The shaded areas show the 90% C.I. for fitted values i.e. the sum of the output from the deterministic SCM and the short-term internal variability excluding the terms for long-term internal variability and model error. Fig. S3 show three set of fitted values for the GMST for the main analysis that include the long-term internal variability and model error.

5




Figure 3: Posterior distribution of time series (a) and prior (dotted) and posterior (solid) probability density function (PDF) in 2010 (b) for anthropogenic forcing. The shaded areas in (a) represent the 90% C.I.




Figure 4: Observed and fitted (posterior mean) values for the OHC. The shaded areas indicate the 90% C.I. Left column: Upper 700 meter. Right column: Below 700 meter, if data included in the analysis.




Figure 5: Observed and fitted (posterior mean) total OHC using several OHC dataset (case B and C, left panel) and using only one dataset (case D, right panel). The shaded areas indicate the 90% C.I.

5 Table 1: List of data used in the estimation, the abbreviation used in the text, references and time of download. The months in parentheses are when data used in case A (see sect. 2.2) were downloaded.

Abbreviation	References	Downloaded
Surface temperature change:		
GISS	(Hansen et al., 2006; Hansen et al., 2010)	March 2015 (March 2011)
HadCRUT4	(Morice et al., 2012)	March 2015 (March 2011*)
NCDC	(Smith and Reynolds, 2005; Smith et al., 2008)	March 2015 (June 2011)
CowtanWay	(Cowtan and Way, 2014)	March 2015 (April 2014)
Ocean heat content upper 700 meter:		
Levitus	(Levitus et al., 2009)	March 2015 (March 2011)
CSIRO	(Domingues et al., 2008; Church et al., 2011)	April 2014 (October 2011)
Ishii and Kimoto	(Ishii and Kimoto, 2009)	March 2015 (October 2011)
ORAS4	(Balmaseda et al., 2013)	March 2015
Ocean heat content below 700 meter:		



ORAS4	(Balmaseda et al., 2013)	March 2015
Ocean heat content above 2000 meter:		
Levitus2000	(Levitus et al., 2012)	July 2015
SOI-index:		
SOI	Southern Oscillation index, Bureau of Meteorology, Australia http://www.bom.gov.au/climate/current/soihtml1.shtml	March 2015 (November 2011)
Forcing time series:		
Forc_Skeie14	(Skeie et al., 2011; Skeie et al., 2014)	
Forc_AR5	(Myhre et al., 2013)	

* HadCRUT3