Reply to Editor.

We very much thank the editor and 2 reviewers for the additional comments, which have helped improve the manuscript. We provide a response to the editor's comments below. Editors comments are in italics. Marked-up manuscript follows responses. Page and line numbers coincide with final

5 manuscript without mark-up.

Dear Authors

15

Thank you for your revised manuscript. I have decided to accept your manuscript subject to minor revisions that I will review.

10 Both reviewer reports are positive. Neither require corrections, but R2 provides a list of suggestions which I strongly recommend you consider. I have captured some of these and made additional recommendations below.

The Sustainable Development Goals may provide a useful framing for your methodology. How can we reduce hunger, provide clean water and reduce climate change impacts for example? This challenge seems to suggest an intergrated modelling approach. You briefly mention this at the end of the article, but it may serve as a more important framing near the start.

We have done this, see Page 2, Line 3 of the revised manuscript

P2 lin 25 Rockström et al., 2009, I suggest you use more recent Steffen et al (2015) as this also includes regional rather than global water use boundary.

20 We have included this reference

> P3 L15. You use climate modelling as a metaphor for model complexity. But to avoid any possible confusion, I suggest you mention that socioeconomic processes have come to be linked to biophysical models that include climate dynamics: IAMs. More generally, you are arguing for the need to develop socio-ecological modelling. That is a reasonably well established discipline or approach. It's a minor

25 issue, but you are setting the reader up in this introduction section so you need to be very clear how your approach is situated with existing work - and where and how it goes beyond it. Previous reviewer reports slightly misunderstood your motivations in places.

We have removed this metaphor and have made clearer that we are linking socioenvironmental (we prefer this term to socio-ecological as water is abiotic) dynamics with existing IAMs.

30 P5 L33 "However, the pathway to capturing these dynamics within global models is not obvious (Müller-Hansen et al., 2017)." this is a citation to a regional study. Verburg et al (2015) may offer a more productive global perspective. But it may not include sufficient discussion on water-energyclimate nexus.

We have included this useful citation.

35 P5 L34 "Thus, the second knowledge gap our framework aims to fill is to incorporate the complex dynamics associated with socioenvironmental systems within the IAM framework, without addina unwanted complexity to those models. We aim to achieve this by focusing on cities." You argue that cities represent a useful aggregation of the agents of interest, that You don't need to capture dynamics of individual farmers or farms for example. This is a centrally important modelling structure 40 decision. It needs to be defended or supported. You provide some high level data that suggests that a

focus on cities would represent sufficient capture of the relevant dynamics. Some readers may not be very convinced by that. How else can you effectively argue that resolving agents as cities is the best approach?

We have expanded on our reasoning for choosing cities as the focus of our study. Page 5, line 27-37

- 5 P8 L1 "However, we recommend tight coupling between dynamic vegetation models such as Lund-Potsdam-Jena managed Land (LPJ-ML) model and a complex hydrological model that captures water fluxes between the soil layer and groundwater reservoir, which is key for computing groundwater recharge rates (Bierkens and van Beek., 2009; Bondeau et al., 2007; Hanasaki et al., 2008b)." It is not clear if you have implemented this method, or have a route towards this method, or if this is an
- 10 aspiration or some sort of necessary condition for your method to be effective. In any event it is a large piece of work. I think the reader needs to get some understanding of what is involved here.

We have clarified that this is not yet done, to our knowledge and that it is a major undertaking. We have briefly explained what would be involved in achieving tight coupling between a DGVM and a hydrological model. Page 7 line 38 – Page 8 Line 9

15 P9 L7 "The upper level network consists of bilateral trade links among countries. Link weights among countries are calculated using a CGE model, which is coupled to certain IAMs (Stehfest et al., 2013)." More specificity here. Which "certain" IAMs?

We removed this statement. We discuss how a CGE is linked to the IMAGE IAM on Page 5 Line 1-3.

P9 L34 "This is a novel and potentially important step in capturing non-equilibrium dynamics within
pre-existing equilibrium approaches without adding unwanted complexity to CGEs or IAMs (Farmer and Geanakoplos, 2009)." I appreciate this is an important point. But, if we consider your multimodel framework to be a model (or meta-model), then this represents a very large increase in model complexity. On the one hand you argue that you don't want to add "unwanted complexity" to some kinds of models, on the other your entire approach is about capturing more dynamics and coupling

25 them within a much more complex modelling methodology. Can you be a bit clearer about your motivations here? I think you want to argue that your approach can fit in with the existing modelling 'ecosystem'. You won't be able to take models 'off the shelf' but you don't need to engineer entirely new IAMs. This is a very important point.

We have clarified this point on Page 10, line 5-11.

35

30 P11 The discussion on how intra city dynamics can impact inter city dynamics - up to the global scale - could be framed in terms of teleconnections. You are arguing that you can only capture such teleconnections in a complex social-ecological system by explicitly resolving agent dynamics and interactions.

We have made reference to teleconnections in our revised manuscript. Page 11, line 9. Page 4, line 17

L9 L21 "The main step required to make this framework a reality is the construction of a multi-agent network of city nodes and trade links." I agree! Which makes the lack of detail about such a model quite important. I think we need some framework for this element of the methodology. I would recommend you briefly review the ODD +D protocol. You may find that you are able to able to quickly

40 produce a schematic of your proposed multi-agent model. It's not much more than a list of entities and descriptions, but it will allow you to produce a concise table with which a reader will be able to quickly understand the nature - if not the details - of this central modelling component. We have added a table following the ODD +D protocol. Page 11, Line 30. Page 27 and 28 $\,$

References

Verburg, P.H., Dearing, J.A., Dyke, J.G., van der Leeuw, S., Seitzinger, S., Steffen, W. and Syvitski, J., 2016. Methods and approaches to modelling the Anthropocene. Global Environmental Change, 39, pp.328-340.

Steffen, W., Richardson, K., Rockström, J., Cornell, S.E., Fetzer, I., Bennett, E.M., Biggs, R., Carpenter, S.R., de Vries, W., de Wit, C.A. and Folke, C., 2015. Planetary boundaries: Guiding human development on a changing planet. Science, 347(6223), p.1259855.

3

10

5

15

A framework for modelling the complexities of food and water security under globalisation

5 Brian J. Dermody^{1,2,2}, Murugesu Sivapalan³Sivapalan⁴, Elke Stehfest⁴Stehfest⁵, Detlef P. van Vuuren^{1,2,4,5}, Martin J. Wassen¹, Marc F. P. Bierkens⁶Bierkens⁶, Stefan C. Dekker^{1,6,2}

¹Copernicus Institute of Sustainable Development, Faculty of Geosciences, Utrecht University, the Netherlands ²Centre for Complex Systems Studies, Utrecht University, the Netherlands

10 ³² Department of Science, University College Utrecht, the Netherlands

³ Department of Civil and Environmental Engineering, Department of Geography and Geographic Information Science, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA

⁴⁵PBL Netherlands Environmental Assessment Agency, The Hague, the Netherlands

15 ⁵⁶Department of Physical Geography, Faculty of Geosciences, Utrecht University, the Netherlands ⁷⁶Faculty of Management, Science and Technology, Department of Science. Open University, Heerlen, The Netherlands

Correspondence to: Brian J. Dermody (brianjdermody@gmail.comb.dermody@uu.nl)

20

25

30

Abstract

We present a new framework for modelling the complexities of food and water security under globalisation. The framework sets out a method to capture regional and sectoral interdependencies and cross scale feedbacks within the global food system that contribute to emergent water use patterns. The framework integrates aspects of existing models and approaches in the fields of hydrology and Integrated Assessment Modelling. The core of the framework is a multi-agent network of city agents connected by socioeconomic and infrastructural trade networks. Agents receive socioeconomic and ecohydrological environmental constraint information from existing IAMs and hydrological models respectively and simulate complex, socioenvironmental dynamics that operate within those constraints. The emergent changes in food and water resources are aggregated and fed back to the original models with minimal modification of the structure of those models. –It is our conviction that models built according to this the framework presented can form the basis for a new wave of decision tools that capture the complexities of complex socioenvironmental change within our globalised world. In doing so they will contribute to illuminating pathways towards a sustainable future for humans, ecosystems and the water they share.

35 Keywords: food and water security, globalisation, urbanization, complex_systemsity, hydrology, integratedassessment models Formatted: Font: (Default) Times New Roman, 10 pt Formatted: Default Paragraph Font, Font: (Default) +Body (Calibri), 11 pt

Formatted: Indent: First line: 0 cm

1 Introduction

5

Ensuring sustainable food and water security in our globalised world is an urgent and complex challenge (Shiklomanov, 2000; United Nations Water, 2015). The magnitude of the challenge is outlined in the Sustainable 10 Development Goals, which set a target of zero hunger globally whilst at the same time drastically reducing impacts of food production on aquatic and terrestrial ecosystems as well as the climate system (United Nations, 2015). Food Food security and water security are inextricably intertwined, with variability in agricultural production impacting water resource use and vice versa (Liu and Savenije, 2008). Trade, both international and domestic, plays a central role in determining water use, in the global food system, because when we trade food, we are also 15 trading the water embedded in the production of that food (Fader et al., 2013; Hoekstra and Chapagain, 2006; Hoekstra and Mekonnen, 2012). The global system of food production and trade is complex and drives interdependency among heterogeneous regions, meaning that socioeconomic or environmental changes in one part of the globe can have cascading impacts for food and water security throughout (Fig. 1b) (D'Odorico et al., 2010; J. Liu et al., 2015; Marchand et al., 2016; Young et al., 2006). Equally, there is interdependency across 20 sectors with the mechanisation of agriculture leading to a tighter coupling of the food, water and energy sectors in recent years, often referred as the food-water-energy nexus (Fig. 1a) (Bazilian et al., 2011)._-Within this interdependent system, interactions occur across spatial and temporal scales (d'Amour et al., 2016; Sivapalan and Blöschl, 2015) (Fig. 1c). For example, the combined effect of small-scale abstraction of ground and surface water to secure short-term food and water security can lead to large scale and long-term depletion of water resources, 25 which ultimately undermines food and water security (Sivapalan and Blöschl, 2015; Sophocleous, 2012).

Given the complexity of the food system, it is challenging to develop effective food and water management strategies because policies can leak across regions, sectors and scales (Eakin et al., 2009; Hejazi et al., 2015; Meyfroidt et al., 2013). Models and decision tools exist to inform policy makers on these issues. (Bierkens and van Beek, 2009; Hanasaki et al., 2008a; Lotze Campen et al., 2008; Sitch et al., 2005; Stehfest et

- 30 al., 2014), For example, Integrated Assessment Models have made significant progress in capturing sectoral interdependencies with the food-water-energy nexus (Bazilian et al., 2011; Lotze-Campen et al., 2008; Stehfest et al., 2014). In the hydrological sciences, water footprint studies capture regional interdependence whilst sociohydrological studies capture cross-scale sociohydrological interactions (Hoekstra and Chapagain, 2006; Sivapalan, 2012), however However to our knowledge, regional and sectoral interdependencies and cross-scale
- feedbacks associated with food and water security have not been captured in a single modelling framework. As a result, existing decision toolsapproaches miss important dynamics (Konar et al., 2016a; Srinivasan et al., 2017). It is urgent that we develop develop models and approachesdecision tools that capture these complexities within a single framework to assist decision makers in navigating the increasingly complex global food system and help ensure that we stay within the natural limits of our planet's boundaries for water resources (Steffen et al., 2015Rockström et al., 2009; Wagener et al., 2010).

Formatted: Indent: First line: 1.27 cm

Formatted: Font color: Text 1
Formatted: Font color: Text 1

Formatted: Font color: Text 1
Formatted: Font color: Text 1

Figure 1. Complex dynamics within the global food system that lead to emergent water use patterns.

In this paper, we present a modellingset out a framework to for eapture capturing regional and sectoral 5 interdependencies and cross-scale feedbacks in the global food system that contribute to emergent water use patterns. The framework builds upon and combines existing models and approachesapproaches in the fields of Integrated Assessment and hydrology and combines them via a multi-agent network of city agents and infrastructural trade links. In our framework city agents consist of an urban agglomeration and an associated hinterland, which we define as the agricultural area economically tied to a city. We focus on cities because they 10 are centres of food and water demand and important agents of change within the global food system, with 54% of the world's population urbanised, consuming a disproportionately large, 75% of the world's resources (United Nations, 2012; United Nations Environmental ProgrammeUNEP, 2013). Cities play a key role in driving regional interdependency via trade. In fact, cities and trade coevolved, with demand from cities determining the location of physical trade networks that constrain our ability to extract 15 food and water resources from the environment and redistribute them around the globe to meet demand (Barber et al., 2014; Barredo and Demicheli, 2003; Friedmann and Wolff, 1982; Fujita et al., 2001; Rees and Wackernagel, 2008) As city populations grow, infrastructural networks continue to upgrade capacity and extend further into natural systems, increasing our ability to extract natural resources, but also increasing regional interdependence (Ibisch et al., 2016; Laurance et al., 2015). The infrastructural networks that radiate from cities, serve to link

20 environmental resources in their city, hinterlands to global food markets, thus playing a key role in facilitating cross-scale socioenvironmental feedbacks within the global food system (Brenner, 1999; Güneralp et al., 2013; Harvey, 1990).
 <u>Cities are also crucibles for cross-sectoral socioeconomic interactions, Socioeconomic conditions across</u>

different sectors of the economy constrain cities demand for, and capacity to produce food (Stehfest, 2014; UNEP,
 2013). Socioeconomic conditions also constrain the ability of cities to trade food on regional and global markets (FAO, 2015). Trade cannot occur directly between two cities unless the socioeconomic links exist and not at all if the infrastructural links are missing (De Benedictis and Tajoli, 2011). Of course, physical infrastructural and

socioeconomic networks cannot be disentangled. Strengthening of socioeconomic ties invariably leads to strengthening of infrastructural links, whilst the cost of investment in that infrastructure serves to stabilise those
 socioeconomic ties (Khanna, 2016).

35

In our framework city agents consist of an urban agglomeration and an associated hinterland, which we define as the agricultural area economically tied to a city.Environmental conditions in the hinterlands of cities constrain the production of food and associated water use, whilst socioeconomic conditions across different sectors of the economy constrain the ability to exploit those environmental conditions for food production (Godfray et al., 2010). Socioeconomic conditions also constrain the ability of cities to trade food on regional and global markets (FAO, 2015). Trade cannot occur directly between two cities unless the socioeconomic links exist and not at all if the infrastructural links are missing (De Benedictis and Tajoli, 2011). Of course, physical infrastructural and socioeconomic networks cannot be disentangled. Strengthening of socioeconomic ties

Formatted: Font color: Text 1

Formatted: Font color: Text 1
Formatted: Indent: First line: 0 cm
Formatted: Font color: Text 1, English (United
Formatted: Font color: Text 1

invariably leads to strengthening of infrastructural links, whilst the cost of investment in that infrastructure serves to stabilise those socioeconomic ties (Khanna, 2016).

Capturing these different components of the global food system in a single framework is ambitious but has parallels to the discipline of elimate modelling, which began with models of sub-components of the elimate system. However, it became clear that these subcomponents were interdependent and it was necessary to capture these interdependencies in a single framework in order to understand the elimate system (Edwards, 2011). A similar approach is urgently required to understand how cross-scale and interdependent socioenvironmental interactions impact food and water security in our globalised and urbanised world (Liu et al., 2015). In the following section 2, we present an overview of existing models and approaches to understand food and water security. We highlight some important knowledge gaps in these approaches that our framework aims to fill. In section 3 we present our framework in detail, outlining how we aim to capture regional and sectoral

section 3 we present our framework in detail, outlining how we aim to capture regional and sectoral interdependencies and cross-scale feedbacks in a single framework. In section 4 we outline potential applications of a realised version of the framework. In section 5 we outline steps required to make this ambitious vision a reality.

15

2. <u>2. Challenges to capturing the complexity of food and water security using existing models</u><u>Challenges in</u> managing water within the global food system

Water resource use in the global food system emerges from complex, socioeconomic and environmental dynamics that are interdependent across regions, sectors and seales (Eakin et al., 2009; Hejazi et al., 2015; Meyfroidt et al., 2013).
 2013). However, to our knowledge, no current decision tools capture the two way interaction between socioeconomic and environmental processes across regions, sectors and scales within a single modelling framework. Given that many sustainability challenges are interlinked, it is urgent that we develop approaches that capture these interdependencies (Güneralp et al., 2013; Liu et al., 2015; Puma et al., 2015). In the following section, we outline the progress made in existing approaches that our framework aims to build upon as well as specific knowledge gaps that out framework aims to fill.

2.1 Regional Interdependence

30 among role ir water hydro soil la

35

40

becomes ever more important for food and water security, drives ing interdependency for food and water security among trading regions (FAO, 2015; Sartori and Schiavo, 2015). Water footprint studies have played an important role in quantifying the volume of water used in food production and embedded in traded food, known as virtual water trade (Allan, 1998; Fader et al., 2011; Hanasaki et al., 2010; Konar et al., 2016b). These studies apply hydrological models to estimate the amount of green (recently precipitated, rapidly replenished water in the upper soil layer) or blue water (slowly replenished ground and surface water resources) used in food production (Fader et al., 2011; Hoff et al., 2010). At a global scale, 84% of food production relies on green water, which has generally less negative impacts on the environment compared with blue water. In terms of internationally traded food, 16% of traded food comes from green water resources and 6% from blue water resources (Fader et al., 2011). As a result, at a global scale, international trade saves water (de Fraiture et al., 2004; Hanasaki et al., 2010; Konar et al., 2011). However, recent studies have revealed that an increasing proportion of traded food is produced from unsustainably abstracted blue water resources (Dalin et al., 2017; Wada et al., 2012, 2010). Owing to regional

As the food system becomes increasingly globalised and urbanised, trade, both domestic and international,

interdependency of the food system, these unsustainable practices threaten the future food and water security of trading regions (Gleeson et al., 2012; Wada and Bierkens, 2014) (Fig. 2a). The increase in unsustainable water abstraction is principally owing to increased demand arising from population growth, with climate change projected to exacerbate stress on food and water resources in the future (Gerten et al., 2011).

5

A knowledge gap in <u>these</u> water footprint studies is that there tends to be a focus on international bilateral trade, meaning the <u>impact ofmuch larger</u> domestic trade <u>on water resources isfluxes are</u> often neglected (Konar et al., 2016). In the developed world, almost all food reaches consumers through domestic and international trade networks-whilst in the developing world, the trade is also increasing in importance as people move to cities and <u>the</u> numbers of people involved in subsistence agriculture is also decreasinges, owing to rapid urbanisation trends

- 10 in these regions (Chen, 2007; IFPRI, 2017; Seto and Reenberg, 2014; United Nations, 2012). The lack of studies on domestic trade is in large part owing to the lack of data. However, increasingly there are attempts to quantify virtual water flows within countries (Fig. 2b) (Dalin et al., 2014; Dang et al., 2015) and the water footprints of cities (Hoff et al., 2014). Identifying the fine-scale networks that constrain where water resources are extracted from the environment to meet remote demand is essential for managing water under globalisation. By capturing 15 these fine scale virtual water flows, it will improve our ability to estimate how socioeconomic or environmental change in one part of the food system can be transmitted through the trade network (Karpiarz et al., 2014). This will improve our ability to diagnose the remote drivers of water resource use and understand how local water resource use may be teleconnected with socioeconomic-socioenvironmental and environmental changes may easeade through thechange in another part of the global trade networkfood system (d'Amour et al., 2016)(Puma 20 et al., 2015). Thus, the first knowledge gap our framework seeks to fill is to capture the fine-scale networks that constrain virtual water tradewater resource extraction and virtual water trade-within countries and among eities. We aim to close this knowledge gap by capturing spatially explicit demand and production at the city and hinterland scale and the infrastructure that constrains the redistribution of water resources to meet this demand (d'Amour et al., 2016).
- 25

Figure 2. Water footprint studies.

2.2 Sectoral Interdependence

30

35

40

Owing to the increased mechanisation of agriculture and the development of more diverse energy sources, water now sits at the centre of the the interdependencies among the food, water energy nexus sectors are stronger than ever (Bazilian et al., 2011; Kraucunas et al., 2015; United Nations Water, 2015). Currently 30% of energy produced is used within the food sector, with fluctuations in energy costs having direct impacts on agriculture and thus water resources (Frieler et al., 2015). Next to this, the energy sector itself requires considerable water resources (Bijl et al., 2016). Fossil fuel extraction such as coal mining and shale gas fracking are highly water intensive, whilst biofuel production competes with food production for land and water resources (Bonsch et al., 2016; Hejazi et al., 2015). Failure to capture sectoral interdependencies means we may fail to identify synergistic solutions, or worse still, policies in one sector may have unintended negative consequences in another. For example, investment in climate change mitigation measures such as biofuel production may lead to increased competition for water between the food and energy sectors and exacerbate unsustainable water resource use (Hejazi et al., 2015).

Integrated Assessment Models (IAMs) are powerful tools to explore sectoral interdependencies (Bazilian et al., 2011). Until now, IAMs have principally been focused on projecting greenhouse gas emissions, however increasingly, the agricultural sector is being captured within IAMs (Stehfest et al., 2013; Stehfest and Bouwman, 2006; Wise et al., 2009). For example, the IMAGE IAM contains an agriculture and land module which is internally coupled with an energy supply and demand module. IMAGE calculates changes in agriculture based on two-way interactions between the agriculture and energy sectors (Stehfest et al., 2014). IMAGE has also been coupled with the MAGNET agro-economic model, a Computable General Equilibrium (CGE) model that uses information from IMAGE on land availability and suitability, labour supply and, technological change to estimate international food trade, demand and supply (Stehfest et al., 2013; von Lampe et al., 2014; Woltjer et al., 2014). An integrated approach has also been applied by Bonsch et al. (2016) using MAgPIE (Lotze-Campen et al., 2008), a global land and water-use allocation model, to understand the trade-off between agricultural expansion and intensification via water abstraction to meet biofuel targets. Similarly to Hejazi et al. (2015), their integrated approach showed that changes in the energy sector can lead to competition for water resources within the food sector. The MAgPIE approach also underlines-underlined the importance of capturing dynamic vegetation and hydrological processes in the same modelling framework as the two are intrinisically intertwined with changes in vegetation impacting water resources and vice versa (Fader et al., 2011; Konar et al., 2013; Lotze-Campen et al., 2008).

5

10

15

Although IAM approaches capture important sectoral interdependencies, they cannot provide a representation of non-linear transformative change typical of socioenvironmental systems and the dynamics
associated with it (Filatova et al., 2013; Folke, 2006; Rockström et al., 2017). Incorporating these processes in IAMs would risk including too much complexity, possibly leading to a trade-off with transparency (van Vuuren et al., 2016). An alternative approach is to couple IAMs with models that capture these important processes, without changing the internal structure of IAMs. Some progress has already been made in this direction by coupling land use models with IAMs, where the IAM set regional socioeconomic constraints for the land use model. Based on these constraints, the and aggregating the emergent fine-scale patterns from the land use model are aggregated at the regional scale captured in the IAM. This approach has, which resulted in improved regional projections in IAMs as land allocation models capture important within-region heterogeneity (Hasegawa et al., 2017; Stehfest et al., 2013).

In order to model the complexity of socioenvironmental systems, it is necessary to capture dynamics
such as agency, emergence, non-linearities and feedbacks (Berger, 2001; Brown et al., 2005; Farmer and Foley, 2009; Farmer and Geanakoplos, 2009; Folke, 2006; Helbing, 2013; Sivapalan et al., 2012). However, the pathway to capturing these dynamics within global modelsat a global scale is not obvious (Müller-Hansen et al., 2017Verburg et al., 2016). For example, it is not practical from a computational or data perspective to represent farmers as agents at a global scale. Thus, the second knowledge gap our framework aims to fill is to incorporate the complex dynamics associated with socioenvironmental systems at a global scale within the IAM framework; without adding unwanted complexity to those models. We aim to achieve this by focusing on cities and infrastructural networks. As stated, Ccities are important agents key agents of environmental change and and lie at the intersection of scales within the global food system, whilst infrastructural networks constrain our ability to extract resources from the environment and redistribute them to meet demand (Brenner, 1999; Khanna, 2016; United Nations Water, 2015). An advantage of using cities as agents is that they have a number of important

features which are generalisable. Firstly, all cities, irrespective of where or when they existed, have a common utility function, which is to sustain the import of resources to maintain growth (Batty, 2008). Secondly, cities competitively interact for resources and services irrespective of whether they are in the same administrative region or not (Begg, 1999). Thirdly, the investment in infrastructure, which is a defining characteristic of cities, makes them highly path dependent (Khanna, 2016). This path dependency provides opportunities for narrowing the possibility space of future projections (Brown et al., 2005).sit at the intersection of scales within the global food system (Brenner, 1999; United Nations Water, 2015).

2.3 Cross-scale socioenvironmental feedbacks

10

5

Cross-scale socioenvironmental feedbacks describe a broad range of processes where small-scale or short-term actions bring about large-scale or long-term emergent change (Folke, 2006; Sivapalan and Blöschl, 2015). These emergent higher-level changes may feedback at the finer scale. In terms of water resources, sociohydrological studies have played an important role in helping to understand emergent water use patterns by capturing crossscale socioenvironmental feedbacks (Di Baldassarre et al., 2013b, 2013a; Montanari et al., 2013; Sivapalan et al., 15 2012; Troy et al., 2015). They have done so by explicitly considering bi-directional feedbacks between humans and the environment in hydrological basins. An example of emergent dynamics is a "pendulum swing", in communities that have alternated between water extraction for agriculture in the early stages of development. followed by subsequent efforts to mitigate or reverse the consequent degradation of the riparian ecosystems (Kandasamy et al., 2014; Liu et al., 2014). This has been explained by counteracting productive and restorative 20 forces, mediated via technology, environmental awareness and the intervention of governance institutions (Elshafei et al., 2015; Liu et al., 2015; van Emmerik et al., 2014).

However, sociohydrological studies maintain a disciplinary focus on water-and do not capture important, failing to capture important sectoral interdependencies as IAMs do (Troy et al., 2015). Equally, they have so far 25 globalised world, many different such entities may be linked through trade as demonstrated in water footprint studies (Konar et al., 2016a). Capturing linkages to regional and global markets is critical to understanding crossscale feedbacks, because the more connected water resources are to markets via trade, the more sensitive they are to cross-scale feedbacks (Eakin et al., 2009; Pande and Sivapalan, 2016). For example, the 2010 drought in Russia and Kazakhstan led to a spike in the price of wheat on global markets (Nelson et al., 2014). Food producers in other wheat producing regions of the world that were well connected to global markets via physical and 30 socioeconomic trade links were impacted by this price rise as they could sell their products to global markets at increased profit. Thus, environmental change in Russia impacted global markets which drove local water resource use in other parts of the world. There has been a recognition of the important role trade plays in facilitating crossscale feedbacks in literature related to land use change (Geist and Lambin, 2002; Lambin and Geist, 2008). These studies have highlighted that distant market demand drives local land use change and in order to predict land use 35 change we need to model land as an open with system with flows of resources coming in and out (Lambin and Meyfroidt, 2011). However, modelling of these issues in relation to water resources is still in its infancy (Konar et al., 2016a). Thus, the third challenge is to capture the crucial role of trade networks in facilitating cross-scale feedbacks between local water resources and regional and global markets.

3. Modelling Framework

3.1 The City Agent

5

3.1.1 City agent attributes

- The core structure of our framework is a multi-agent network of city nodes and trade links (Fig. 3). In our framework, cities are agents and comprise of an urban area and associated hinterland. City agents sit at the intersection of scales between <u>environmental ecohydrological</u> resources in their hinterlands and global markets via radiating trade networks. The <u>environmental ecohydrological</u> conditions in hinterlands of city agents are determined from a <u>grid based ecohydrologicalcoupled hydrological-vegetation</u> model for natural and managed land. Socioeconomic conditions in the hinterlands of cities are determined from an IAM and CGE. Socioeconomic
- 10 conditions constrain agent's ability to exploit environmental conditions within their hinterland for food production. City agents have a common utility function: to satisfy local and market demand for food, which in turn leads to emergent water use. City agents satisfy demand via food production and trade. How each city satisfies demand differs based on the heterogeneous socioeconomic and environmental conditions within in their hinterland and the socioeconomic and infrastructural networks that link them. The fine-scale interactions among cities and their
- 15 environment bring about higher-level emergent patterns which provide input conditions for the next simulation step of the ecohydrological model and the IAM and CGE. Thus, fine-scale interactions among heterogeneous city agents and their environment bring about emergent change which is incorporated within existing model structures with minimal change to the structure or complexity of those models.

Each city agent has an associated hinterland. The size definition of a hinterland varies in literature but of a

Formatted: Font color: Red

20 Figure 3. City agent attributes

3.1.2 The city and its hinterland

25

30

35

- hinterland-depends upon food resource demand in the city and on the ease of transportation between the city and the agricultural resource production area, which is determined by geographic, infrastructural and sociopolitical conomic factors (Billen et al., 2009). To capture these elements, we define city hinterlands based on the hierarchical overlay of supra-sub national administrative borders and <u>Ttheijess</u>sen polygon operation among cities based on the gravity equation of trade with mass equal to demand and distance equal to the cost-distance of trade via road, rail and inland water ways (Fig. 4) (Berthelon and Freund, 2008; Chaney, 2013). We base hinterlands on administrative borders so that the framework can capture the impact of heterogeneous policy or socioeconomic conditions at the administrative scale. For many cities, and depending on the food commodity, the effective hinterland will extend beyond these contiguous administrative boundaries and cities operate within these policy constraints. If policies stimulate free trade between administrative regions, then the effective hinterland of a city can expand (Knox and McCarthy, 2012). Therefore, the framework allows for the exploration of the impact of
- policy at different scales on food and water use and virtual water flows. By also basing the size of hinterlands on the cost-distance of trade via road, rail and shipping, we capture the key role that infrastructural networks play in linking demand in cities with environmental resources.

Hinterlands vary in terms of size and composition.

Formatted: Indent: First line: 1.27 cm

5

15

20

The hinterland of a city in Western Australia will be large, with a high proportion of natural landcover and low population density, whilst a hinterland in Eastern China will be smaller with a high proportion of agricultural and urban landcover and high population density. From a socioeconomic perspective, per capita demand will differ among hinterlands based on eulturediet, affluence etc. derived from the IAM. Equally, the agricultural intensification potential of a hinterland in the Netherlands will be greater compared with a sub-Saharan African region owing to factors such as mechanisation of agriculture, access to fertilizers etc. (Stehfest et al., 2014).

Figure 4. The city and its hinterland

10 3.2 Food production and water use

Within our framework, city agent food production decisions are constrained by socioeconomic and environmentalecohydrological conditions within their hinterlands. The To capture environmental ecohydrological conditions within the hinterlands of cities are determined using a process-based ecohydrological model for natural and managed land similar to the approach of Hejazi et al. (2015) and Lotze Campen et al. (2008). However, we recommend tight coupling between a dynamic vegetation models (DGVM) such as Lund-Potsdam-Jena managed Land (LPJ-ML) model and a complex hydrological model such as PC Raster Global Water Balance (PCR-GLOBWB) model (Bierkens and van Beek., 2009; Bondeau et al., 2007) (Fig. 3). In this way, the model would that captures the two way water fluxes between vegetation and water in water fluxes between the soil layer and and infiltration to the groundwater reservoir, which is key for computing groundwater recharge rates (Bierkens and van Beek., 2009; Bondeau et al., 2007; Hanasaki et al., 2008b). This is a considerable modelling challenge. Vegetative water demand and rooting depths from the DGVM would need to be sent to the hydrological model. Using the detailed soil water profile from the hydrological model, an improved estimate of water availability could

- be estimated. If water supply is lower than demand from the DGVM, then vegetation growth will be limited due to soil water stress. If water supply is higher than demand from the DGVM, then water infiltrates into the 25 groundwater reservoir in the hydrological model. Each city hinterland has a potential yield and associated water
 - resource usage based on these ecohydrological constraints.

Socioeconomic constraints on food production are taken from the IAM based on factors such as the mechanisation of agriculture, access to fertilizers etc. The demand for each food commodity in a city and its hinterland is equal to (per capita demand taken from IAM * population), with population based on spatially 30 explicit grid-scale population estimates (Bijl et al., 2017; Brinkhoff, 2016; Klein Goldewijk et al., 2011; Stehfest, 2014; UN Population Division, 2015). Cities and their hinterland are either in surplus or deficit for a crop type based on (local production - local demand) (Dermody et al., 2014). Cities with predominantly urbanised hinterlands will have a net demand for food resources whilst others, with large areas of agriculture within their hinterlands, will have a net surplus in food resources (Dermody et al., 2014). Demand is also based on trade demand calculated in the trade component of the framework (Section 3.3).

35

40

Cities adapt to changes in food demand through agricultural expansion/contraction, (de)intensification or trade. Agricultural expansion and intensification potential are estimated using land use algorithms. The algorithms used to calculate agricultural expansion and intensification potential in IAMs such as IMAGE may be extended to include detailed infrastructural data as well as more complex hydrology in determining agricultural suitability (Barber et al., 2014; Wada et al., 2012; Walker et al., 2013). In our framework, cities with increasing Formatted: Indent: First line: 1.27 cm

demand, low (high) agricultural intensity potential and high (low) expansion potential are likely to expand (intensify) agriculture (Fig. 5). Cities that are constrained from expanding or intensifying agriculture will increase imports to sustain growth. The agent decision process generates land-use maps that are prescribed to the ecohydrological model (Biemans et al., 2013; Wada et al., 2012).

5

10

The method for capturing food production outlined, blends macroscale interdependencies between the agricultural and energy sector with small-scale ecohydrologialecohydrological and demographic conditions to estimate land use change, similar to the approach by (Hasegawa et al.; (2017);- and Verstegen et al. (-2016)). Importantly, our approach also captures how market demand is transmitted heterogeneously within a region or country via socioeconomic and infrastructural networks (see section 3.3 for more details). This is a critical step in understanding how the critical role infrastructural networks play in determining water resource extraction and virtual water trade patterns regional interdependence for food and water security impacts local water resource use (Konar et al., 2016a; Lambin and Mevfroidt, 2011).

Figure 5. Food production decisions.

15

3.3 Virtual Water Trade

The trade network that links city agents is hierarchical. The network comprises a lower-level physical infrastructural network and an upper-level network with link weights reflecting bilateral socioeconomic trade links (Fig. 6). The lower level of the trade network is defined by physical trade infrastructure of roads, rail and shipping lanes (Berthelon and Freund, 2008; Karpiarz et al., 2014; Limão and Venables, 2001). The data required to build this network, such as open street maps and database of cargo ship movements, is openly available in digital format (Brinkhoff, 2016; Haklay and Weber, 2008; Kaluza et al., 2010). Each-In a realised version of the framework, each link type has aedge of the lower-level network has a different transport cost. For example, the cost of bulk trade is approximately 7-times less by ship than by road (Limão and Venables, 2001). In addition, 25 intermodal transport costs are applied for transferring goods between transport modes (Janic, 2007). Infrastructure development can be projected using infrastructure growth algorithms or manually added to explore the impacts of prospective plans such as the new trans-Eurasian Silk Road Economic Belt on virtual water trade and water use (Ahmed et al., 2013; Arima et al., 2008; Brugier, 2014; Walker et al., 2013). Because the infrastructural network constrains where resources can be extracted from the environment and redistributed to meet demand, all resource 30 flows travel via the lower level physical infrastructural network in a realised version of the model framework (Barber et al., 2014; Khanna, 2016). The upper level network consists of bilateral trade links among countries. Link-Edge weights among countries are calculated using a CGE model, which is coupled to certain IAMs. CGEs are constrained with historical data on bilateral trade balances, competitiveness (relative price developments) and trade policies to estimate trade patterns (Hertel et al., 2007; Hertel and Hertel, 1997; Woltjer et al., 2014). 35 (Stehfest et al., 2013). CGEs are constrained with historical data on bilateral trade balances, competitiveness (relative price developments) and trade policies to estimate trade patterns (Hertel et al., 2007; Hertel and Hertel, 1997; Woltier et al., 2014). If the CGE simulates high trade volumes between two countries, then the link-edge weight between those countries in our framework will reflect a high probability of bilateral trade.

Figure 6. Virtual Water Trade Network.

5

10

35

In a realised version of the framework, Fthe food produced in the hinterland of cities is either consumed locally or exported to meet market demand. Cities with a demand, import from cities with a surplus based on the cost-distance among cities (Berthelon and Freund, 2008). Thus, the probability of trade among cities decays with increasing cost-distance (Karpiarz et al., 2014; Limão and Venables, 2001). City agents within the same country trade with one another based solely on supply and demand for each food commodity and the cost-distance of trade among cities. Cities in different countries are also constrained by bilateral trade probabilities of the upper-level network (Fig. 6). For example, the probability of trade between Japan and American cities will be determined by

- the upper level network based on CGE output. However, it is more likely that the demand will be met from hinterlands on the West Coast of the United States owing to the lower cost-distance of trade. The emergent trade patterns are aggregated at the country and region scale of the CGE and fed back into it for the next simulation year. Because city agent trade decisions are stochastic, it allows for alternate emergent trade patterns within the
- 15 constraints of the same network. Thus, for given constraints there will be a range of possible solutions. The stronger (weaker) the constraints the narrower (wider) the solution space. In this way, the solution space for given environmental, socioeconomic and infrastructural constraints can be explored (An, 2012). Equally, if conditions change at the small scale, this will result in alternative emergent patterns at the large scale (Fig. 7b). In this way, agency, cross-scale processes, socioeconomic, environmental and infrastructural constraints are married with
- conventional general equilibrium modelling approaches. This is a novel and potentially important step in capturing non-equilibrium dynamics within pre-existing equilibrium modelling_approaches without_adding_unwanted eomplexity to CGEs or IAMs (Farmer and Geanakoplos, 2009). It also allows us to begin to uncover the important role of cross-scale feedbacks which in linking local water resources change to the dynamics of regional or global markets (Konar et al., 2016a). By capturing these cross-scale feedbacks, we can begin to see how heterogeneous water use can bring about emergent large-scale and long-term market change, which in turn feeds back to local scale water use.

Figure 7. Modelling Framework Workflow.

30 4. Applications of the framework

The framework is designed to be used with IAMs and CGEs, which are influential decision tools but are unable to capture non-linear, emergent change typical of socioenvironmental systems (Rockström et al., 2017; van Vuuren et al., 2016; Verburg et al. 2016). Our approach does not require IAMs or CGEs to be re-engineered, but rather simulates complex socioenvironmental processes within the constraints exogenously applied those models. The emergent patterns may be aggregated at the scale of IAMs and CGEs and used as input conditions for the next simulation step of those models. In this way, the framework blends bottom-up processes captured by the multi-agent network of cities with the top-down approach captured by pre-existing IAMs and CGEs. The framework is designed to be used with IAMs, which are influential decision tools but are unable to capture nonlinear, emergent change typical of socioenvironmental systems (Rockström et al., 2017; van Vuuren et al., 2016). Our approach blends agency and emergent processes captured by the multi-agent network of cities with the topdown approach captured by IAMs. Thus, a realised version of the framework has the potential to illuminate how the large scalehigher-level patterns demonstrated in IAMs and CGEs -emerge. In terms of informing policy, this is an essential important step as it allows policy makers to explore the impacts of policy changes at the different spatial and temporal scales at which policy is applied. In doing so, models based on the framework can help illuminate roadmaps, considering complex dynamics, to reach regional and long-term goals outlined in IAMs (Rockström et al., 2017). For example, it allows the exploration of how scenarios, such as the recently published Shared Socioeconomic Pathway (SSP), which outlines future environmental change according to different socioeconomic development storylines (O'Neill et al., 2015), can be implemented at finer scales and taking account of sectoral and regional interdependencies and cross-scale feedbacks. This can be a potentially important step in extending the usefulness of IAMs and CGEs whilst avoiding adding unwanted complexity to those modelss (van Vuuren et al., 2016)-.

The multi-agent network of cities at the core of the framework enables the exploration of the emergent solution space for realizing food and water security requirements based on socioeconomic and environmental these constraints. The emergent solution space illuminates how constrained regions or sectors are based on socioeconomic, environmental and infrastructural factors to transition to sustainable and secure water use pathways (Brown et al., 2005; Garud et al., 2010). Those regions or sectors that are locked into unsustainable or vulnerable water use pathways should be prioritised for intervention (Fig. 8) (Liebowitz and Margolis, 1995; Romero-Lankao et al., 2017; Sophocleous, 2012). Within the solution space, sustainable and secure development pathways can be explored. A common approach for exploring multi-dimensional solution space is to use optimality algorithms (Dermody et al., 2011; Konak et al., 2006; Vrugt et al., 2003). For example, optimality algorithms can be applied to find solutions that are optimal from a land saving and water saving perspective. Because the IAM captures cross-sectoral interdependencies, mMulti-objective optimality algorithms can be used to explore synergistically sustainable solutions across regions, sectors and scales. Such synergistic approaches are key to addressing the challenges set out in the UN Sustainable Development Goals (Costanza et al., 2016; Lu et al., 2015). In addition, our approach can uncover potential displacement of environmental impact of policies to other regions (Lambin and Meyfroidt, 2011). Thus, the model can be used to inform policies can be explored that are sustainable both in the administrative region, the economic sector where and the scale at which those policies are applied as well as interdependent regions, sectors and scales.

Figure 8. The solution space illustrates how constrained regions or sectors are along their current water use pathway.

35

30

5

10

15

20

25

Given the importance of cross-scale feedbacks for understanding local water use, it is essential that smaller scale studies on food and water security can be linked with regional-global scale models (Pande and Sivapalan, 2016; Sivapalan and Blöschl, 2015; Verburg and Overmars, 2009). Returning to the parallels with elimate modelling, small-scale studies use output from Global General Circulation Models as boundary conditions for local-regional elimate studies (Douglas et al., 2009). Equally, tThe model framework outlined presented can be used to understand how detailed changes within a city or catchment link to large-scale market or hydrological changes. Thus, models built according to our framework can contribute to, and benefit from, the rapid growth of within-city studies such as studies of urban metabolism studies or sociohydrological studies which focus on the

40

Formatted: Font color: Auto

catchment scale (Broto et al., 2012; Liu et al., 2014; van Emmerik et al., 2014; Zhang, 2013). <u>In</u>-In-the case of urban metabolism studies, detailed analysis is done of resources flows into and out of a city. Models based on our framework can illustrate how those-urban resource flows are <u>linked-teleconnected to-with</u> environmental change in other parts of the globe-and the feedback of those-global changes at the city-seale. <u>Equally, urban resource</u> <u>budgets can be used to test model performance</u>. In the case of sociohydrological studies, the framework outlined can help illustrate how small-scale changes across multiple catchments can <u>lead to</u> emergent market changes, which in turn feedback on water use within a single catchment.

5. Summary and recommendations

10 The complex and interdependent nature of food and water security within our globalised and urbanised world requires new models and approaches to inform policy makers. Models built using the framework outlined are ideally suited to informing policy on so-called "wicked problems" associated with water resource use <u>such as regional and sectoral interdependencies and cross-scale feedbacks</u> within the globalized food system (Dentoni et al., 2012; Duit and Galaz, 2008; Lach et al., 2005). The framework we present builds on the existing fields of hydrology and Integrated Assessment Modelling to understand emergent water use patterns. It is our conviction that models built according to the framework outlined will represent a new wave of decision tools that can help policy makers navigate the complexity of socioenvironmental interactions within our globalised world. Much of the groundwork for the framework already exists, however extra efforts are required to achieve an operational model.

20 IAMs and hydrological models provide the groundwork for the proposed framework. However, extra efforts are required to achieve an operational model based on the framework presented. The main step required to make this framework a reality-is the construction of a multi-agent network of city nodes and trade infrastructural links. A multi-agent network of city nodes and infrastructural links will provide a platform for a wide range of research questions on environmental change. The As outlined, cities and infrastructural trade networks play a key 25 role in environmental change. Cities are where the majority of resource demand is, whilst infrastructural networks constrain our ability to extract resources from the environment and redistribute them around the globe to meet this demand. Thus, a multi-agent network of cities and infrastructural links will provide a platform for a wide range of research questions related to environmental change. The resources required to build this network are openly available but need to be translated to a scalable and flexible network architecture. - The structure of this network 30 should be designed in collaboration with scientists from a wide range of disciplines to ensure the relevant elements are incorporated to meet a range of research questions. A summary description of a generalised framework for a multi-agent network of cities and infrastructural links is presented in Table 1.

<u>Table 1. Description of generalised framework of a multi-agent network of city nodes and infrastructural</u> links.

The structure of this network should be designed in collaboration with scientists from a wide range of different disciplines to ensure the relevant elements are incorporated to meet a range of research questions. In order to To facilitate collaboration in building such a multi-agent network, there is a need for interdisciplinary dialogue and collaboration. Indeed, this is essential for achieving the Sustainable Development Goals which

Formatted: Font: Bold
Formatted: Indent: First line: 0 cm
Formatted: Font: Bold

whic

5

35

present interdependent challenges across disciplines (Costanza et al., 2016; Lu et al., 2015). In our experience, the theories and methods associated with complex_systemsity science provide an ideal approach for tackling interdependent interdisciplinary sustainability challenges (Liu et al., 2015). Complexity Complexity is not just a suite of theories and methods, it delivers an intuitive way of understanding interdependent systems and provides a platform for deep interdisciplinarity collaboration that is required to meet today's sustainability challenges.

5

-Competing interests. The authors declare that they have no conflict of interest.

Special issue statement. This article is part of the special issue "Social dynamics and planetary boundaries in

10 Earth system modelling"

> Acknowledgements. The authors would like to thank the editor, James Dyke, and 3 anonymous reviewers for their considered and constructive comments, which helped improve the manuscript.

Competing interests. The authors declare that they have no conflict of interest.

15

45

References

Ahmed, S.E., Souza, C.M., Riberio, J., Ewers, R.M., 2013. Temporal patterns of road network development in the Brazilian Amazon. Reg. Environ. Change 13, 927-937. doi:10.1007/s10113-012-0397-z Allan, J. a., 1998. Virtual Water: A Strategic Resource Global Solutions to Regional Deficits. Ground Water 36, 20 545-546. doi:10.1111/j.1745-6584.1998.tb02825.x An, L., 2012. Modeling human decisions in coupled human and natural systems: Review of agent-based models. Ecol. Model., Modeling Human Decisions 229, 25-36. doi:10.1016/j.ecolmodel.2011.07.010 Arima, E.Y., Walker, R.T., Sales, M., Souza Jr, C., Perz, S.G., 2008. The Fragmentation of Space in the 25 Amazon Basin: Emergent Road Networks. Photogramm. Eng. Remote Sens. 74. Batty, M., 2008. The Size, Scale, and Shape of Cities. Science 319, 769-771. https://doi.org/10.1126/science.1151419 Barber, C.P., Cochrane, M.A., Souza Jr., C.M., Laurance, W.F., 2014. Roads, deforestation, and the mitigating effect of protected areas in the Amazon. Biol. Conserv. 177, 203-209. 30 doi:10.1016/j.biocon.2014.07.004 Barredo, J.I., Demicheli, L., 2003. Urban sustainability in developing countries' megacities: modelling and predicting future urban growth in Lagos. Cities 20, 297-310. doi:10.1016/S0264-2751(03)00047-7 Bazilian, M., Rogner, H., Howells, M., Hermann, S., Arent, D., Gielen, D., Steduto, P., Mueller, A., Komor, P., Tol, R.S.J., Yumkella, K.K., 2011. Considering the energy, water and food nexus: Towards an 35 integrated modelling approach. Energy Policy, Clean Cooking Fuels and Technologies in Developing Economies 39, 7896-7906. doi:10.1016/j.enpol.2011.09.039 Begg, I., 1999. Cities and Competitiveness. Urban Studies 36, 795-809. https://doi.org/10.1080/0042098993222 Berardy, A., Chester, M.V., 2017. Climate change vulnerability in the food, energy, and water nexus: concerns for agricultural production in Arizona and its urban export supply. Environ. Res. Lett. 12, 035004. 40 https://doi.org/10.1088/1748-9326/aa5e6d Berger, T., 2001. Agent-based spatial models applied to agriculture: a simulation tool for technology diffusion, resource use changes and policy analysis. Agric. Econ., Increasing Efficiency in Production, Research,

Markets and Environmental Management. Selected and edited papers presented during the XXIV Conference of the International Association of Agricultural Economists 25, 245-260. doi:10.1016/S0169-5150(01)00082-2

- Berthelon, M., Freund, C., 2008. On the conservation of distance in international trade. J. Int. Econ. 75, 310-320. doi:10.1016/j.jinteco.2007.12.005
- Biemans, H., Speelman, L.H., Ludwig, F., Moors, E.J., Wiltshire, A.J., Kumar, P., Gerten, D., Kabat, P., 2013. Future water resources for food production in five South Asian river basins and potential for adaptation 50 A modeling study. Sci. Total Environ., Changing water resources availability in Northern India with respect to Himalayan glacier retreat and changing monsoon patterns: consequences and adaptation 468-469, Supplement, S117-S131. doi:10.1016/j.scitotenv.2013.05.092

Formatted: Font: Not Bold Formatted: Font: Not Bold Formatted: Font: Not Bold

Formatted: Indent: First line: 0 cm

Bierkens, M.F.P., van Beek, L.P.H., 2009. Seasonal Predictability of European Discharge: NAO and Hydrological Response Time. J. Hydrometeorol. 10, 953–968. doi:10.1175/2009JHM1034.1

- Bijl, D.L., Bogaart, P.W., Dekker, S.C., Stehfest, E., de Vries, B.J.M., van Vuuren, D.P., 2017. A physicallybased model of long-term food demand. Glob. Environ. Change 45, 47–62. doi:10.1016/j.gloenvcha.2017.04.003
- Bijl, D.L., Bogaart, P.W., Kram, T., de Vries, B.J.M., van Vuuren, D.P., 2016. Long-term water demand for electricity, industry and households. Environ. Sci. Policy 55, Part 1, 75–86. doi:10.1016/j.envsci.2015.09.005

5

10

15

50

55

- Billen, G., Barles, S., Garnier, J., Rouillard, J., Benoit, P., 2009. The food-print of Paris: long-term reconstruction of the nitrogen flows imported into the city from its rural hinterland. Reg. Environ. Change 9, 13–24. doi:10.1007/s10113-008-0051-y
- Bondeau, A., Smith, P.C., Zaehle, S., Schaphoff, S., Lucht, W., Cramer, W., Gerten, D., Lotze-Campen, H., Müller, C., Reichstein, M., Smith, B., 2007. Modelling the role of agriculture for the 20th century global terrestrial carbon balance. Glob. Change Biol. 13, 679–706. doi:10.1111/j.1365-2486.2006.01305.x
- Bonsch, M., Humpenöder, F., Popp, A., Bodirsky, B., Dietrich, J.P., Rolinski, S., Biewald, A., Lotze-Campen, H., Weindl, I., Gerten, D., Stevanovic, M., 2016. Trade-offs between land and water requirements for large-scale bioenergy production. GCB Bioenergy 8, 11–24. doi:10.1111/gcbb.12226
- Brenner, N., 1999. Globalisation as Reterritorialisation: The Re-scaling of Urban Governance in the European
 Union. Urban Stud. 36, 431–451. doi:10.1080/0042098993466
- Brinkhoff, T., 2016. World city population database: Population Statistics for Countries, Administrative Areas, Cities and Agglomerations.

Broto, V.C., Allen, A., Rapoport, E., 2012. Interdisciplinary Perspectives on Urban Metabolism. J. Ind. Ecol. 16, 851–861. doi:10.1111/j.1530-9290.2012.00556.x

- 25 Brown, D.G., Page, S., Riolo, R., Zellner, M., Rand, W., 2005. Path dependence and the validation of agentbased spatial models of land use. Int. J. Geogr. Inf. Sci. 19, 153–174. doi:10.1080/13658810410001713399
 - Brugier, C., 2014. China's way: The new Silk Road (No. 14). EU Institute for Security Studies.
- Chaney, T., 2013. The Gravity Equation in International Trade: An Explanation (Working Paper No. 19285).
 National Bureau of Economic Research.
- Chen, J., 2007. Rapid urbanization in China: A real challenge to soil protection and food security. CATENA, Influences of rapid urbanization and industrialization on soil resource and its quality in China 69, 1–15. doi:10.1016/j.catena.2006.04.019

Costanza, R., Fioramonti, L., Kubiszewski, I., 2016. The UN Sustainable Development Goals and the dynamics
 of well-being. Front. Ecol. Environ. 14, 59–59. doi:10.1002/fee.1231

- d'Amour, C.B., Wenz, L., Kalkuhl, M., Steckel, J.C., Creutzig, F., 2016. Teleconnected food supply shocks. Environ. Res. Lett. 11, 035007. doi:10.1088/1748-9326/11/3/035007
- Dalin, C., Hanasaki, N., Qiu, H., Mauzerall, D.L., Rodriguez-Iturbe, I., 2014. Water resources transfers through Chinese interprovincial and foreign food trade. Proc. Natl. Acad. Sci. 111, 9774–9779.
 doi:10.1073/pnas.1404749111
- Dalin, C., Wada, Y., Kastner, T., Puma, M.J., 2017. Groundwater depletion embedded in international food trade. Nature 543, 700–704. doi:10.1038/nature21403
- Dang, Q., Lin, X., Konar, M., 2015. Agricultural virtual water flows within the United States. Water Resour. Res. 51, 973–986. doi:10.1002/2014WR015919
 De Benedictis, L., Tajoli, L., 2011. The World Trade Network. World Econ. 34, 1417–1454.
- 5 De Benedictis, L., Tajoli, L., 2011. The World Trade Network. World Econ. 34, 1417–1454. doi:10.1111/j.1467-9701.2011.01360.x
- de Fraiture, C., Cai, X., Amarasinghe, I., Rosegrant, M., Molden, D., 2004. Does international cereal trade save water?: the impact of virtual water trade on global water use. IWMI.
 - Dentoni, D., Hospes, O., Ross, R.B., others, 2012. Managing wicked problems in agribusiness: the role of multistakeholder engagements in value creation. Int. Food Agribus. Manag. Rev. 15.

Dermody, B.J., Tanner, C.J., Jackson, A.L., 2011. The evolutionary pathway to obligate scavenging in gyps vultures. PLoS ONE 6.

- Dermody, B.J., van Beek, R.P.H., Meeks, E., Klein Goldewijk, K., Scheidel, W., van der Velde, Y., Bierkens, M.F.P., Wassen, M.J., Dekker, S.C., 2014. A virtual water network of the Roman world. Hydrol Earth Syst Sci 18, 5025–5040. doi:10.5194/hess-18-5025-2014
- Di Baldassarre, G., Kooy, M., Kemerink, J.S., Brandimarte, L., 2013a. Towards understanding the dynamic behaviour of floodplains as human-water systems. Hydrol. Earth Syst. Sci. 17, 3235. doi: 10.5194/hess-17-3235-2013

- Di Baldassarre, G., Viglione, A., Carr, G., Kuil, L., Salinas, J.L., Blöschl, G., 2013b. Socio-hydrology: conceptualising human-flood interactions. Hydrol Earth Syst Sci 17, 3295-3303. doi:10.5194/hess-17-3295-2013
- D'Odorico, P., Laio, F., Ridolfi, L., 2010. Does globalization of water reduce societal resilience to drought? Geophys. Res. Lett. 37, L13403. doi:10.1029/2010GL043167 5
- Douglas, E.M., Beltrán-Przekurat, A., Niyogi, D., Pielke Sr., R.A., Vörösmarty, C.J., 2009. The impact of agricultural intensification and irrigation on land-atmosphere interactions and Indian monsoon precipitation -- A mesoscale modeling perspective. Glob. Planet. Change 67, 117-128. doi:10.1016/j.gloplacha.2008.12.007
- 10 Duit, A., Galaz, V., 2008. Governance and complexity-emerging issues for governance theory. Governance 21. 311-335.
- Eakin, H., Winkels, A., Sendzimir, J., 2009. Nested vulnerability: exploring cross-scale linkages and vulnerability teleconnections in Mexican and Vietnamese coffee systems. Environ. Sci. Policy, Special Issue: Food Security and Environmental ChangeFood Security and Environmental Change: Linking Science, Development and Policy for Adaptation 12, 398-412. doi:10.1016/j.envsci.2008.09.003 15
- Edwards, P.N., 2011. History of climate modeling. Wiley Interdiscip. Rev. Clim. Change 2, 128-139. doi:10.1002/wcc.95
- Elshafei, Y., Coletti, J.Z., Sivapalan, M., Hipsey, M.R., 2015. A model of the socio-hydrologic dynamics in a semiarid catchment: Isolating feedbacks in the coupled human-hydrology system. Water Resour. Res. 51, 6442-6471. doi:10.1002/2015WR017048 20
- Fader, M., Gerten, D., Krause, M., Lucht, W., Cramer, W., 2013. Spatial decoupling of agricultural production and consumption: quantifying dependences of countries on food imports due to domestic land and water constraints. Environ. Res. Lett. 8, 014046. doi:10.1088/1748-9326/8/1/014046
- Fader, M., Gerten, D., Thammer, M., Heinke, J., Lotze-Campen, H., Lucht, W., Cramer, W., 2011. Internal and external green-blue agricultural water footprints of nations, and related water and land savings through 25 trade. Hydrol Earth Syst Sci 15, 1641-1660. doi:10.5194/hess-15-1641-2011 Farmer, J.D., Foley, D., 2009. The economy needs agent-based modelling. Nature 460, 685-686.
 - doi:10.1038/460685a
- Farmer, J.D., Geanakoplos, J., 2009. The virtues and vices of equilibrium and the future of financial economics. Complexity 14, 11-38. doi:10.1002/cplx.20261 30
- Filatova, T., Verburg, P.H., Parker, D.C., Stannard, C.A., 2013. Spatial agent-based models for socio-ecological systems: Challenges and prospects. Environ. Model. Softw., Thematic Issue on Spatial Agent-Based Models for Socio-Ecological Systems 45, 1-7. doi:10.1016/j.envsoft.2013.03.017
- Folke, C., 2006. Resilience: The emergence of a perspective for social-ecological systems analyses. Glob. 35 Environ. Change 16, 253-267. doi: 10.1016/j.gloenvcha.2006.04.002
- Food and Agriculture Organisation of the United Nations, 2015. The State of Food Insecurity in the World. Rome
- Friedmann, J., Wolff, G., 1982. World city formation: an agenda for research and action. Int. J. Urban Reg. Res. 6, 309-344. doi:10.1111/j.1468-2427.1982.tb00384.x
- 40 Frieler, K., Levermann, A., Elliott, J., Heinke, J., Arneth, A., Bierkens, M.F.P., Ciais, P., Clark, D.B., Deryng, D., Döll, P., others, 2015. A framework for the cross-sectoral integration of multi-model impact projections: land use decisions under climate impacts uncertainties. Earth Syst. Dyn. 6, 447-460. Fujita, M., Krugman, P., Venables, A., 2001. The Spatial Economy: Cities, Regions, and International Trade
- (MIT Press Books). The MIT Press. Garud, R., Kumaraswamy, A., Karnøe, P., 2010. Path Dependence or Path Creation? J. Manag. Stud. 47, 760-45 774. doi:10.1111/j.1467-6486.2009.00914.x
- Geist, H.J., Lambin, E.F., 2002. Proximate Causes and Underlying Driving Forces of Tropical Deforestation Tropical forests are disappearing as the result of many pressures, both local and regional, acting in various combinations in different geographical locations. BioScience 52, 143-150. doi:10.1641/0006-50 3568(2002)052[0143:PCAUDF]2.0.CO;2
- Gerten, D., Heinke, J., Hoff, H., Biemans, H., Fader, M., Waha, K., 2011. Global Water Availability and Requirements for Future Food Production. J. Hydrometeorol. 12, 885-899. doi:10.1175/2011JHM1328.1
- Gleeson, T., Wada, Y., Bierkens, M.F.P., van Beek, L.P.H., 2012. Water balance of global aquifers revealed by groundwater footprint. Nature 488, 197-200. doi:10.1038/nature11295 55
 - Godfray, H.C.J., Beddington, J.R., Crute, I.R., Haddad, L., Lawrence, D., Muir, J.F., Pretty, J., Robinson, S., Thomas, S.M., Toulmin, C., 2010. Food Security: The Challenge of Feeding 9 Billion People. Science 327, 812-818. doi:10.1126/science.1185383
- Güneralp, B., Seto, K.C., Ramachandran, M., 2013. Evidence of urban land teleconnections and impacts on 60
 - hinterlands. Curr. Opin. Environ. Sustain. 5, 445-451. doi: 10.1016/j.cosust.2013.08.003

Haklay, M., Weber, P., 2008. OpenStreetMap: User-Generated Street Maps. IEEE Pervasive Comput. 7, 12–18. doi:10.1109/MPRV.2008.80

Hanasaki, N., Inuzuka, T., Kanae, S., Oki, T., 2010. An estimation of global virtual water flow and sources of water withdrawal for major crops and livestock products using a global hydrological model. J. Hydrol. 384, 232–244. doi:10.1016/j.jhydrol.2009.09.028

5

- Hanasaki, N., Kanae, S., Oki, T., Masuda, K., Motoya, K., Shirakawa, N., Shen, Y., Tanaka, K., 2008a. An integrated model for the assessment of global water resources – Part 2: Applications and assessments. Hydrol Earth Syst Sci 12, 1027–1037. doi:10.5194/hess-12-1027-2008
- Hanasaki, N., Kanae, S., Oki, T., Masuda, K., Motoya, K., Shirakawa, N., Shen, Y., Tanaka, K., 2008b. An integrated model for the assessment of global water resources–Part 1: Model description and input meteorological forcing. Hydrol. Earth Syst. Sci. 12, 1007–1025. doi: 10.5194/hess-12-1007-2008
 Harvey, D., 1990. Between Space and Time: Reflections on the Geographical Imagination1. Ann. Assoc. Am. Geogr. 80, 418–434. doi:10.1111/j.1467-8306.1990.tb00305.x
- Hasegawa, T., Fujimori, S., Ito, A., Takahashi, K., Masui, T., 2017. Global land-use allocation model linked to
 an integrated assessment model. Sci. Total Environ. 580, 787–796. doi:10.1016/j.scitotenv.2016.12.025
- Hejazi, M.I., Voisin, N., Liu, L., Bramer, L.M., Fortin, D.C., Hathaway, J.E., Huang, M., Kyle, P., Leung, L.R., Li, H.-Y., Liu, Y., Patel, P.L., Pulsipher, T.C., Rice, J.S., Tesfa, T.K., Vernon, C.R., Zhou, Y., 2015.
 21st century United States emissions mitigation could increase water stress more than the climate change it is mitigating. Proc. Natl. Acad. Sci. 112, 10635–10640. doi:10.1073/pnas.1421675112
- 20 Helbing, D., 2013. Globally networked risks and how to respond. Nature 497, 51–59. doi:10.1038/nature12047 Hertel, T., Hummels, D., Ivanic, M., Keeney, R., 2007. How confident can we be of CGE-based assessments of Free Trade Agreements? Econ. Model. 24, 611–635. doi:10.1016/j.econmod.2006.12.002 Hertel, Thomas W., Hertel, Thomas Warren, 1997. Global Trade Analysis: Modeling and Applications. Cambridge University Press.
- Hoekstra, A.Y., Chapagain, A.K., 2006. Water footprints of nations: Water use by people as a function of their consumption pattern. Water Resour. Manag. 21, 35–48. doi:10.1007/s11269-006-9039-x
 Hoekstra, A.Y., Mekonnen, M.M., 2012. The water footprint of humanity. Proc. Natl. Acad. Sci. 109, 3232–3237. doi:10.1073/pnas.1109936109
- Hoff, H., Döll, P., Fader, M., Gerten, D., Hauser, S., Siebert, S., 2014. Water footprints of cities indicators for sustainable consumption and production. Hydrol Earth Syst Sci 18, 213–226. doi:10.5194/hess-18-213-2014
 - Hoff, H., Falkenmark, M., Gerten, D., Gordon, L., Karlberg, L., Rockström, J., 2010. Greening the global water system. J. Hydrol., Green-Blue Water Initiative (GBI) 384, 177–186. doi:10.1016/j.jhydrol.2009.06.026
- 35 Ibisch, P.L., Hoffmann, M.T., Kreft, S., Pe'er, G., Kati, V., Biber-Freudenberger, L., DellaSala, D.A., Vale, M.M., Hobson, P.R., Selva, N., 2016. A global map of roadless areas and their conservation status. Science 354, 1423–1427. doi:10.1126/science.aaf7166
- IFPRI, 2017. Global Food Policy Report. International Food Policy Research Institute, Washington, DC, US. Janic, M., 2007. Modelling the full costs of an intermodal and road freight transport network. Transp. Res. Part
 Transp. Environ. 12, 33–44. doi:10.1016/j.trd.2006.10.004
- Kaluza, P., Kölzsch, A., Gastner, M.T., Blasius, B., 2010. The complex network of global cargo ship movements. J. R. Soc. Interface 7, 1093–1103. doi:10.1098/rsif.2009.0495
- Kandasamy, J., Sounthararajah, D., Sivabalan, P., Chanan, A., Vigneswaran, S., Sivapalan, M., 2014. Sociohydrologic drivers of the pendulum swing between agricultural development and environmental health:
 a case study from Murrumbidgee River basin, Australia. Hydrol Earth Syst Sci 18, 1027–1041. doi:10.5194/hess-18-1027-2014
 - Karpiarz, M., Fronczak, P., Fronczak, A., 2014. International Trade Network: Fractal Properties and Globalization Puzzle. Phys. Rev. Lett. 113, 248701. doi:10.1103/PhysRevLett.113.248701 Khanna, P., 2016. Connectography: Mapping the Global Network Revolution. Hachette UK.
- 50 Klein Goldewijk, K., Beusen, A., van Drecht, G., de Vos, M., 2011. The HYDE 3.1 spatially explicit database of human-induced global land-use change over the past 12,000 years. Glob. Ecol. Biogeogr. 20, 73–86. doi:10.1111/j.1466-8238.2010.00587.x
 - Knox, P.L., McCarthy, L., 2012. Urbanization: an introduction to urban geography. Pearson Boston, MA.
- Konak, A., Coit, D.W., Smith, A.E., 2006. Multi-objective optimization using genetic algorithms: A tutorial.
 Reliab. Eng. Syst. Saf., Special Issue Genetic Algorithms and ReliabilitySpecial Issue Genetic Algorithms and Reliability 91, 992–1007. doi:10.1016/j.ress.2005.11.018
 - Konar, M., Dalin, C., Suweis, S., Hanasaki, N., Rinaldo, A., Rodriguez-Iturbe, I., 2011. Water for food: The global virtual water trade network. Water Resour. Res. 47, W05520. doi:10.1029/2010WR010307

- Konar, M., Evans, T.P., Levy, M., Scott, C.A., Troy, T.J., Vörösmarty, C.J., Sivapalan, M., 2016a. Water resources sustainability in a globalizing world: who uses the water? Hydrol. Process. 30, 3330–3336. doi:10.1002/hyp.10843
- 5
- Konar, M., Hussein, Z., Hanasaki, N., Mauzerall, D.L., Rodriguez-Iturbe, I., 2013. Virtual water trade flows and savings under climate change. Hydrol Earth Syst Sci Discuss 10, 67–101. doi:10.5194/hessd-10-67-2013
- Konar, M., Reimer, J.J., Hussein, Z., Hanasaki, N., 2016b. The water footprint of staple crop trade under climate and policy scenarios. Environ. Res. Lett. 11, 035006. doi:10.1088/1748-9326/11/3/035006
- Kraucunas, I., Clarke, L., Dirks, J., Hathaway, J., Hejazi, M., Hibbard, K., Huang, M., Jin, C., Kintner-Meyer,
 M., Dam, K.K. van, Leung, R., Li, H.-Y., Moss, R., Peterson, M., Rice, J., Scott, M., Thomson, A., Voisin, N., West, T., 2015. Investigating the nexus of climate, energy, water, and land at decision-relevant scales: the Platform for Regional Integrated Modeling and Analysis (PRIMA). Clim. Change 129, 573–588. doi:10.1007/s10584-014-1064-9
- Lach, D., Rayner, S., Ingram, H., 2005. Taming the waters: strategies to domesticate the wicked problems of water resource management. Int. J. Water 3, 1–17. doi:10.1504/IJW.2005.007156
 Lambin, E.F., Geist, H.J., 2008. Land-Use and Land-Cover Change: Local Processes and Global Impacts.
 - Springer Science & Business Media. Lambin, E.F., Meyfroidt, P., 2011. Global land use change, economic globalization, and the looming land
- scarcity. Proc. Natl. Acad. Sci. 108, 3465–3472. doi:10.1073/pnas.1100480108
 Laurance, W.F., Peletier-Jellema, A., Geenen, B., Koster, H., Verweij, P., Van Dijck, P., Lovejoy, T.E., Schleicher, J., Van Kuijk, M., 2015. Reducing the global environmental impacts of rapid infrastructure
 - expansion. Curr. Biol. 25, R259–R262. doi:10.1016/j.cub.2015.02.050 Liebowitz, S.J., Margolis, S.E., 1995. Path Dependence, Lock-In, and History. J. Law Econ. Organ. 11, 205– 226.
- 25 Limão, N., Venables, A.J., 2001. Infrastructure, Geographical Disadvantage, Transport Costs, and Trade. World Bank Econ. Rev. 15, 451–479. doi:10.1093/wber/15.3.451
 - Liu, D., Tian, F., Lin, M., Sivapalan, M., 2015. A conceptual socio-hydrological model of the co-evolution of humans and water: case study of the Tarim River basin, western China. Hydrol Earth Syst Sci 19, 1035–1054. doi:10.5194/hess-19-1035-2015
- 30 Liu, J., Mooney, H., Hull, V., Davis, S.J., Gaskell, J., Hertel, T., Lubchenco, J., Seto, K.C., Gleick, P., Kremen, C., Li, S., 2015. Systems integration for global sustainability. Science 347, 1258832. doi:10.1126/science.1258832
 - Liu, J., Savenije, H.H.G., 2008. Food consumption patterns and their effect on water requirement in China. Hydrol Earth Syst Sci 12, 887–898. doi:10.5194/hess-12-887-2008
- 35 Liu, Y., Tian, F., Hu, H., Sivapalan, M., 2014. Socio-hydrologic perspectives of the co-evolution of humans and water in the Tarim River Basin, Western China: the Taije-Tire Model. Hydrol. Earth Syst. Sci. 18, 1289–1303. doi:10.5194/hess-18-1289-2014
- Lotze-Campen, H., Müller, C., Bondeau, A., Rost, S., Popp, A., Lucht, W., 2008. Global food demand, productivity growth, and the scarcity of land and water resources: a spatially explicit mathematical programming approach. Agric. Econ. 39, 325–338. doi:10.1111/j.1574-0862.2008.00336.x
 - Lu, Y., Nakicenovic, N., Visbeck, M., Stevance, A.-S., 2015. Policy: Five priorities for the UN Sustainable Development Goals. Nat. News 520, 432. doi:10.1038/520432a
- Marchand, P., Carr, J.A., Dell'Angelo, J., Fader, M., Gephart, J.A., Matti Kummu, Magliocca, N.R., Porkka, M., Puma, M.J., Ratajczak, Z., Rulli, M.C., Seekell, D.A., Suweis, S., Tavoni, A., D'Odorico, P., 2016.
 Reserves and trade jointly determine exposure to food supply shocks. Environ. Res. Lett. 11, 095009. doi:10.1088/1748-9326/11/9/095009
- Meyfroidt, P., Lambin, E.F., Erb, K.-H., Hertel, T.W., 2013. Globalization of land use: distant drivers of land change and geographic displacement of land use. Curr. Opin. Environ. Sustain., Human settlements and industrial systems 5, 438–444. doi:10.1016/j.cosust.2013.04.003
- Montanari, A., Young, G., Savenije, H.H.G., Hughes, D., Wagener, T., Ren, L.L., Koutsoyiannis, D., Cudennee, C., Toth, E., Grimaldi, S., Blöschl, G., Sivapalan, M., Beven, K., Gupta, H., Hipsey, M., Schaefli, B., Arheimer, B., Boegh, E., Schymanski, S.J., Baldassarre, G.D., Yu, B., Hubert, P., Huang, Y., Schumann, A., Post, D.A., Srinivasan, V., Harman, C., Thompson, S., Rogger, M., Viglione, A., McMillan, H., Characklis, G., Pang, Z., Belyaev, V., 2013. "Panta Rhei—Everything Flows": Change in hydrology and society—The IAHS Scientific Decade 2013–2022. Hydrol. Sci. J. 58, 1256–1275.
- doi:10.1080/02626667.2013.809088
 Müller, B., Bohn, F., Dreßler, G., Groeneveld, J., Klassert, C., Martin, R., Schlüter, M., Schulze, J., Weise, H., Schwarz, N., 2013. Describing human decisions in agent-based models – ODD + D, an extension of the ODD protocol. Environmental Modelling & Software 48, 37–48.

https://doi.org/10.1016/j.envsoft.2013.06.003

- Müller-Hansen, F., Schlüter, M., Mäs, M., Hegselmann, R., Donges, J.F., Kolb, J.J., Thonicke, K., Heitzig, J., 2017. How to represent human behavior and decision making in Earth system models? A guide to techniques and approaches. Earth Syst Dynam Discuss 2017, 1–53. doi:10.5194/csd-2017-18
- Nelson, G.C., Valin, H., Sands, R.D., Havlík, P., Ahammad, H., Deryng, D., Elliott, J., Fujimori, S., Hasegawa, T., Heyhoe, E., Kyle, P., Lampe, M.V., Lotze-Campen, H., d'Croz, D.M., Meijl, H. van, Mensbrugghe, D. van der, Müller, C., Popp, A., Robertson, R., Robinson, S., Schmid, E., Schmitz, C., Tabeau, A., Willenbockel, D., 2014. Climate change effects on agriculture: Economic responses to biophysical shocks. Proc. Natl. Acad. Sci. 111, 3274–3279. doi:10.1073/pnas.1222465110

5

10

20

- O'Neill, B.C., Kriegler, E., Ebi, K.L., Kemp-Benedict, E., Riahi, K., Rothman, D.S., van Ruijven, B.J., van Vuuren, D.P., Birkmann, J., Kok, K., others, 2015. The roads ahead: narratives for shared socioeconomic pathways describing world futures in the 21st century. Glob. Environ. Change.doi: 10.1016/j.gloenvcha.2015.01.004
- Pande, S., Sivapalan, M., 2016. Progress in socio-hydrology: a meta-analysis of challenges and opportunities. Wiley Interdiscip. Rev. Water e1193. doi:10.1002/wat2.1193
- 15 Puma, M.J., Bose, S., Chon, S.Y., Cook, B.I., 2015. Assessing the evolving fragility of the global food system. Environ. Res. Lett. 10, 024007. doi:10.1088/1748-9326/10/2/024007
 - Rees, W., Wackernagel, M., 2008. Urban ecological footprints: why cities cannot be sustainable—and why they are a key to sustainability, in: Urban Ecology. Springer, pp. 537–555.
 - Rockström, J., Gaffney, O., Rogelj, J., Meinshausen, M., Nakicenovic, N., Schellnhuber, H.J., 2017. A roadmap for rapid decarbonization. Science 355, 1269–1271. doi:10.1126/science.aah3443
 - Rockström, J., Steffen, W., Noone, K., Persson, Å., Chapin, F., Lambin, E., Lenton, T., Scheffer, M., Folke, C., Schellnhuber, H., Nykvist, B., Wit, C. de, Hughes, T., Leeuw, S. van der, Rodhe, H., Sörlin, S., Snyder, P., Costanza, R., Svedin, U., Falkenmark, M., Karlberg, L., Corell, R., Fabry, V., Hansen, J., Walker, B., Liverman, D., Richardson, K., Crutzen, P., Foley, J., 2009. Planetary Boundaries: Exploring the Safe Operating Space for Humanity. Inst. Sustain. Solut. Publ.
 - Romero-Lankao, P., McPhearson, T., Davidson, D.J., 2017. The food-energy-water nexus and urban complexity. Nature Climate Change 7, nclimate3260. https://doi.org/10.1038/nclimate3260
 - Sartori, M., Schiavo, S., 2015. Connected we stand: A network perspective on trade and global food security. Food Policy 57, 114–127. doi:10.1016/j.foodpol.2015.10.004
- 30 Seto, K.C., Reenberg, A., 2014. Rethinking Global Land Use in an Urban Era, in: Seto, K.C., Reenberg, A. (Eds.), Rethinking Global Land Use in an Urban Era. The MIT Press, pp. 1–7. Shiklomanov, I.A., 2000. Appraisal and Assessment of World Water Resources. Water Int. 25, 11–32. doi:10.1080/02508060008686794
- Sitch, S., Brovkin, V., von Bloh, W., van Vuuren, D., Eickhout, B., Ganopolski, A., 2005. Impacts of future
 land cover changes on atmospheric CO2 and climate. Glob. Biogeochem. Cycles 19.
- Sivapalan, M., Blöschl, G., 2015. Time scale interactions and the coevolution of humans and water. Water Resour. Res. 51, 6988–7022. doi:10.1002/2015WR017896
- Sivapalan, M., Savenije, H.H.G., Blöschl, G., 2012. Socio-hydrology: A new science of people and water. Hydrol. Process. 26, 1270–1276. doi:10.1002/hyp.8426
- 40 Sophocleous, M., 2012. Retracted: Conserving and Extending the Useful Life of the Largest Aquifer in North America: The Future of the High Plains/Ogallala Aquifer. Groundwater 50, 831–839. doi:10.1111/j.1745-6584.2012.00965.x
 - Srinivasan, V., Sanderson, M., Garcia, M., Konar, M., Blöschl, G., Sivapalan, M., 2017. Prediction in a sociohydrological world. Hydrol. Sci. J. 62, 338–345. doi:10.1080/02626667.2016.1253844
- 45 Steffen, W., Richardson, K., Rockström, J., Cornell, S.E., Fetzer, I., Bennett, E.M., Biggs, R., Carpenter, S.R., Vries, W. de, Wit, C.A. de, Folke, C., Gerten, D., Heinke, J., Mace, G.M., Persson, L.M., Ramanathan, V., Reyers, B., Sörlin, S., 2015. Planetary boundaries: Guiding human development on a changing planet. Science 347, 1259855. https://doi.org/10.1126/science.1259855
- Stehfest, E., 2014. Diet: Food choices for health and planet. Nature 515, 501–502.
 Stehfest, E., Berg, M. van den, Woltjer, G., Msangi, S., Westhoek, H., 2013. Options to reduce the environmental effects of livestock production Comparison of two economic models. Agric. Syst. 114, 38–53. doi:10.1016/j.agsy.2012.07.002
- Stehfest, E., Bouwman, L., 2006. N2O and NO emission from agricultural fields and soils under natural vegetation: summarizing available measurement data and modeling of global annual emissions. Nutr.
 Cycl. Agroecosystems 74, 207–228.
 - Stehfest, E., van Vuuren, D., Kram, T., Bouwman, L., Alkemade, R., Bakkenes, M., Biemans, H., Bouwman, A., den Elzen, M., Janse, J., Lucas, P., van Minnen, J., Muller, M., Gerdien Prins, A., 2014. Integrated Assessment of Global Environmental Change with IMAGE 3.0 - Model description and policy applications. Netherlands Environmental Assessment Agency, De Bilt, the Netherlands.

	 Troy, T.J., Konar, M., Srinivasan, V., Thompson, S., 2015. Moving sociohydrology forward: a synthesis across studies. Hydrol Earth Syst Sci 19, 3667–3679. doi:10.5194/hess-19-3667-2015 UN Population Division, 2015. World Population Prospects: the 2015 Revision. United Nations, Department of 	
5	Economic and Social Affairs, Population Division.	Formatted: Font: (Default) +Bo
	United Nations, 2012. World Urbanization Prospects. United Nations, Department of Economic and Social	Formatted: Normal
	Affairs, Population Division, New York. United Nations, 2015. Transforming our world: the 2030 agenda for sustainable development. New York, USA.	
10	United Nations Environmental Programme, 2013. City-level Decoupling: Urban Resource Flows and the Governance of Infrastructure Transitions. A Report of the Working Group on Cities of the International Resource Panel. Swilling M., Robinson B., Marvin S. and Hodson M.	
	United Nations Water, 2015. Annual Report. United Nations.	
	van Emmerik, T.H.M., Li, Z., Sivapalan, M., Pande, S., Kandasamy, J., Savenije, H.H.G., Chanan, A., Vigneswaran, S., 2014. Socio-hydrologic modeling to understand and mediate the competition for	
15	water between agriculture development and environmental health: Murrumbidgee River basin, Australia. Hydrol Earth Syst Sci 18, 4239–4259. doi:10.5194/hess-18-4239-2014	
	van Vuuren, D.P., Lucas, P.L., Häyhä, T., Cornell, S.E., Stafford-Smith, M., 2016. Horses for courses: analytical tools to explore planetary boundaries. Earth Syst Dynam 7, 267–279. doi:10.5194/esd-7-267-	
20	2016 Verburg, P.H., Dearing, J.A., Dyke, J.G., Leeuw, S. van der, Seitzinger, S., Steffen, W., Syvitski, J., 2016.	
20	Methods and approaches to modelling the Anthropocene. Global Environmental Change 39, 328-340.	
	https://doi.org/10.1016/j.gloenvcha.2015.08.007 Verburg, P.H., Overmars, K.P., 2009. Combining top-down and bottom-up dynamics in land use modeling:	
25	exploring the future of abandoned farmlands in Europe with the Dyna-CLUE model. Landsc. Ecol. 24,	
25	1167. doi:10.1007/s10980-009-9355-7 Verstegen, J.A., Hilst, F., Woltjer, G., Karssenberg, D., Jong, S.M., Faaij, A.P., 2016. What can and can't we	
	say about indirect land-use change in Brazil using an integrated economic-land-use change model?	
	Geb Bioenergy 8, 561–578. doi: 10.1111/gebb.12270 von Lampe, M., Willenbockel, D., Ahammad, H., Blanc, E., Cai, Y., Calvin, K., Fujimori, S., Hasegawa, T.,	
30	Havlik, P., Heyhoe, E., Kyle, P., Lotze-Campen, H., Mason d'Croz, D., Nelson, G.C., Sands, R.D., Schmitz, C., Tabeau, A., Valin, H., van der Mensbrugghe, D., van Meijl, H., 2014. Why do global	
	long-term scenarios for agriculture differ? An overview of the AgMIP Global Economic Model Intercomparison. Agric. Econ. 45, 3–20. doi:10.1111/agec.12086	
	Vrugt, J.A., Gupta, H.V., Bastidas, L.A., Bouten, W., Sorooshian, S., 2003. Effective and efficient algorithm for	
35	multiobjective optimization of hydrologic models. Water Resour. Res. 39. Doi: 10.1029/2002WR001746	
	Wada, Y., Beek, L.P.H. van, Kempen, C.M. van, Reckman, J.W.T.M., Vasak, S., Bierkens, M.F.P., 2010. Global depletion of groundwater resources. Geophys. Res. Lett. 37, 5 PP.	
40	doi:201010.1029/2010GL044571 Wada, Y., Bierkens, M.F.P., 2014. Sustainability of global water use: past reconstruction and future projections.	
40	Environ. Res. Lett. 9, 104003. doi:10.1088/1748-9326/9/10/104003	
	Wada, Y., van Beek, L.P.H., Bierkens, M.F.P., 2012. Nonsustainable groundwater sustaining irrigation: A global assessment. Water Resour. Res. 48, W00L06. doi:10.1029/2011WR010562	
45	Wagener, T., Sivapalan, M., Troch, P.A., McGlynn, B.L., Harman, C.J., Gupta, H.V., Kumar, P., Rao, P.S.C.,	
45	Basu, N.B., Wilson, J.S., 2010. The future of hydrology: An evolving science for a changing world. Water Resour. Res. 46, W05301. doi:10.1029/2009WR008906	
	Walker, R., Arima, E., Messina, J., Soares-Filho, B., Perz, S., Vergara, D., Sales, M., Pereira, R., Castro, W.,	
	2013. Modeling spatial decisions with graph theory: logging roads and forest fragmentation in the Brazilian Amazon. Ecol. Appl. 23, 239–254.	
50	Wise, M., Calvin, K., Thomson, A., Clarke, L., Bond-Lamberty, B., Sands, R., Smith, S.J., Janetos, A., Edmonds, J., 2009. Implications of Limiting CO ₂ Concentrations for Land Use and Energy. Science 324, 1183–1186. doi:10.1126/science.1168475	
	Woltjer, G.B., Kuiper, M., Kavallari, A., van Meijl, H., Powell, J.P., Rutten, M.M., Shutes, L.J., Tabeau, A.A.,	
55	2014. The MAGNET model: module description. LEI Wageningen UR. Young, O.R., Berkhout, F., Gallopin, G.C., Janssen, M.A., Ostrom, E., Van Der Leeuw, S., 2006. The	
55	Foung, O.K., Berkhout, F., Gaitopin, G.C., Janssen, M.A., Ostrom, E., Van Der Leeuw, S., 2006. The globalization of socio-ecological systems: An agenda for scientific research. Glob. Environ. Change 16, 304–316. doi:10.1016/j.gloenvcha.2006.03.004	
	Zhang, Y., 2013. Urban metabolism: A review of research methodologies. Environ. Pollut. 178, 463–473. doi:10.1016/j.envpol.2013.03.052	
	23	

ody (Calibri), 11 pt






Figure 1. Complex dynamics within the global food system that lead to emergent water use patterns. (A) Sectoral interdependence in water resources within the food-water-energy nexus (B) Regional interdependence in water resources owing to the the importance of importance of trade for food security under globalization and urbanization (C) Sectoral and regional interdependencies are linked via eCross-scale feedbacks. wherebyF_fine-scale interactions bring about emergent ehange-higher-level changeat larger spatial or longer spatial seales. Those-Higher-level, emergent changes may feed_back at the fine scale (Sivapalan and Bloschl, 2016).





Figure 2. Water footprint studies. Water footprint studies exploit hydrological models to estimate blue and green water use in agriculture. This data is combined with data on food trade to estimate the fluxes of virtual water embedded in food trade. A) The unsustainable water footprint of agriculture is shown in red, where groundwater abstraction exceeds aquifer recharge (*taken from* Wada and Bierkens, 2014). **B**) Virtual water flows within the United States. U.S. States are ranked according to the total trade volume and plotted clockwise in descending order. The size of the outer bar indicates the total virtual water trade volume of each State as a percentage of total U.S. trade. Destination volume is indicated with links emanating from the outer bar of the same colour. Origin volume is indicated with a white area separating the outer bar from Links of a different colour (*taken from* Dang et al., 2015).



JFigure 2. Water footprint studies. Water footprint studies exploit hydrological models to estimate blue and green water use in agriculture. This data is combined with data on food trade to estimate the fluxes of virtual water embedded in food trade. **A)** The unsustainable water footprint of agriculture is shown in red, where groundwater abstraction exceeds aquifer recharge (*taken from* Wada and Bierkens, 2014). **B**) Virtual water flows within the United States. States are ranked according to the total trade volume and plotted clockwise in descending order. The size of the outer bar indicates the total virtual water trade volume of each State as a percentage of total U.S. trade. Destination volume is indicated with links emanating from the outer bar of the same colour. Origin volume is indicated with a white area separating the outer bar from links of a different colour (*taken from* Dang et al., 2015). Formatted: Font: (Default) Times New Roman, 10 pt

5





Figure 3. City agent attributes. In our framework, city agents receive environmentalecohydrological conditions from and socioeconomic constraints from ecohydrological a tight coupling between a dynamic vegetation model and hydrological model. They receive socioeconomic constraint information from an and-Integrated Assessment Models. Ecohydrological conditions on natural and managed land determine food production potential and associated water use. Ecohydrological conditions are computed by a tight coupling between a dynamic vegetation model and hydrological model. The IAM captures heterogeneous socioeconomic conditions for food production. Apart from the multi-agent model, the other elements of the framework have already been developed. At the time of writing However, to our knowledge, a tight coupling between a dynamic vegetation model has yet to be coupled to aand a complex hydrological model that computes water exchange between soil and groundwaterhas yet to be implemented..



Formatted: Centered

Figure 4. The city and its hinterland. City agent hinterlands are defined by a hierarchical overlay of suprasub-national administrative boundaries and <u>Theisen-Thiessen</u> interpolation among city nodes based on the gravity equation of trade. Cities are linked by physical infrastructural network <u>where edges represent the costdistance of trade via of</u> roads, rail and shipping routes.

5




Figure 5. Food production decisions. City agents decide on agricultural expansion (left side) or intensification (right side) based on demand and spatially explicit crop production potential in their hinterlands.

Formatted Table





Figure 6. Virtual Water Trade Network. The trade network is hierarchical, containing an upper-level network that determines the strength of trade linkswith edge weights capturing the probability of trade among countries. The link weights of the upper level network are calculated using a CGE model based on factors such as historical trade patterns, trade agreements etc. and aThe lower level physical infrastructural network which captures the comprises an infrastructural network of roads, raid and shipping routes. Edge weights capture the -cost distance of trade via network of roads, rail and shipping-lanes. Because the infrastructural network constrains where resources can be extracted from the environment and redistributed to meet demand, all resource flows travel via the lower level physical infrastructural network in a realised version of the model framework-that link cities. The link weights of the upper level network are calculated using a CGE model based on factors such as historical trade patterns, trade agreements etc. Because infrastructure constrains trade, all virtual water trade flows via the physical trade network. City agents within the same country trade with one another based solely on supply and demand for each food commodity and the cost-distance of trade among eities. Cities in different countries are also constrained by bilateral trade probabilities of upper-level network.



Formatted: Indent: First line: 0 cm

Figure 7. Modelling Framework Workflow. The multi-agent network sits at the interface between the IAM and an ecohydrological model (A) The IAM, calculates per capita food demand at a regional scale, which is converted into spatially explicit demand using population maps. The IAM also calculates agricultural intensity constraints. Based on these constraints and demand, Based on agroeconomic and environmental constraints, aggents make food production decisions which change ecohydrological conditions in their catchment. The emergent crop production and water use patterns are aggregated and feedback to the IAM (B) A CGE trade model coupled to the IAM provides <u>link edge</u> weights for the upper level <u>trade</u> network. Virtual Water Trade among cities is calculated based on supply and demand and cost-distance of trade via the <u>multi-agent</u> infrastructural trade network. The emergent aggregated trade patterns feedback to the CGE as input for the next simulation year.




Figure 8. The solution space illustrates how constrained regions or sectors are along their current water **use pathway.** The solution space emerges from the multi-agent simulations and illuminates how locked-in regions or sectors are along their current water use pathway. The solution space can be explored using optimality algorithms to find pathways towards sustainable water futures, whilst taking account of regional and sectoral interdependencies and cross-scale feedbacks.

Table 1. Description of generalised framework for a multi-agent network of city nodes and infrastructural links. The description follows the ODD +D Protocol (Müller et al., 2013) for describing

agent-based models.				
	1.1 Purpose			
	A multi-agent network of cities and infrastructural links will provide a platform for a wide range of research questions related to global environmental			
	change. The framework is designed to be coupled with IAMs, CGEs and grid-based environmental models. The framework enables the simulation of regional			
	and sectoral interdependencies and cross-scale feedbacks within the constraints provided by these models.			
	1.2 State variables and Scales			
	The multi-agent network consists of 1 type of agent, the city. City agents interact with the environment within their own hinterland and with other city agents			
	by trading resources via the infrastructure network. Each agent has a geographic node location which is linked to a hinterland polygon from which agent			
Overview	attributes are derived. The distance among agents is based on the cost distance of trade via the infrastructural network. The network is scalable depending on			
	the detail of analysis. The scaling is based on infrastructure hierarchies and population. For example, at a global scale, only freeway roads are captured and			
	cities below a certain population threshold are aggregated into larger city nodes.			
1	1.3 Process Overview and Scheduling			
	Each year, cities receive socioeconomic constraint information from an IAM. Cities have a certain resource demand based on their internal demand and			
	demand from other cities in the network. Cities make resource use decisions based on demand, socioeconomic constraints and environmental conditions in			
	their hinterland. The output of the multi-agent model may be aggregated at the scale of IAMs or CGEs and used as input for the next iteration of those			
	models.			
	2.1 Theoretical and empirical background			
	The model is built to test the hypothesis that socioenvironmental interactions across regions, sectors and scales are mediated by cities and physical trade			
	infrastructure that link environmental resources with socioeconomic demand.			
	2.2 Individual Decision making			
	The decision model is based on the generalisable theory that cities competitively interact for environmental resources to sustain growth. City agent decisions			
	are individual but socioeconomic constraints on decision-making may be applied at a higher level, e.g. state level.			
	2.3 Learning Optimisation algorithms may be applied to explore solution space within exogenously applied socioeconomic and environmental constraints.			
	2.4 Individual Sensing			
	City agents sense exogenous socioeconomic constraint information from an IAM. City agents sense exogenous environmental conditions from an			
Design	environmental model. Cities sense endogenous resource demand from other agents.			
Design Concents	2.5 Individual Prediction Agents perceive temporal changes in resource demand.			
Concepts	2.6 Interaction Interactions among agents are determined by each agent's surplus or deficit for a resource and the cost distance among agents in the network.			
	2.7 Collectives Agents do not belong to collectives.			
	2.8 Heterogeneity			
	Agent attributes are heterogeneous and based upon socioeconomic constraint information, environmental conditions and population within their hinterland.			
	The local topology of the infrastructure trade network is also heterogeneous among agents.			
	2.9 Stochasticity			
	Socioeconomic constraint information is applied as probability distributions, meaning agent decisions are stochastic and based on these probability			
	distributions. Agent-Agent trade decisions are stochastic and based on supply and demand and the cost distance between agents in the infrastructure network.			
	The probability of trade decreases with increasing cost distance, assuming uniform supply and demand.			
1	2.10 Observation			

Formatted: Caption, Keep with next

Formatted: Width: 29.7 cm, Height: 21 cm

	Emergent agent decisions are sent to an environmental model as land-use change maps. The environmental model calculates environmental processes based on these land-use changes. The observable output is environmental change in forest cover, water resources, crop yields etc. The second observable output is resource consumption. Emergent resource consumption of agents is aggregated at a scale relevant for IAMs, e.g. the regional scale. The third observable output are trade patterns. Emergent trade patterns are aggregated at a scale relevant for CGEs, e.g. countries and regions.
Details	3.1 Implementation The framework is not yet implemented, and a software platform has not yet been chosen. 3.2 Initialisation It is envisaged that a model based on the framework presented can be initialised for past conditions and run until present to test the model's performance. The model can also be run for scenarios that are implemented within the IAM framework. Because the model is stochastic, an ensemble of simulations can be run for the same socioeconomic constraint information and used to explore different pathways for given constraints.
	3.3 Input Data The input data to define city agents are the global city database and population density maps. Administrative boundaries combined with Thiessen polygon analysis are used to define hinterlands of cities. The infrastructural trade network consists of road and rail networks and shipping lanes and ports. Other infrastructure relevant for resource extraction such as energy redistribution infrastructure could also be captured with the framework.