

1 **Structure and functioning of the acid-base system in the Baltic Sea.**

2 Karol Kuliński¹, Bernd Schneider², Beata Szymczycha¹, Marcin Stokowski¹

3 ¹Institute of Oceanology, Polish Academy of Sciences, IO PAN, ul. Powstańców Warszawy
4 55, 81-712 Sopot, Poland

5 ²Leibniz Institute for Baltic Sea Research, IOW, Seestrasse 15 Warnemünde, D-18119
6 Rostock, Germany

7 Correspondence to: Karol Kuliński (kroll@iopan.gda.pl)

8

9 Key words: pH, total alkalinity, CO₂ system, borate alkalinity, organic alkalinity, dissociation
10 constants, biomass production, mineralization

11

12 **Abstract**

13 The marine acid-base system is relatively well understood for oceanic waters. Its structure and
14 functioning is less obvious for the coastal and shelf seas due to a number of regionally specific
15 anomalies. In this review article we collect and integrate existing knowledge on the acid-base
16 system in the Baltic Sea. Hydrographical and biogeochemical characteristics of the Baltic Sea,
17 as manifested in horizontal and vertical salinity gradients, permanent stratification of the water
18 column, eutrophication, high organic matter concentrations and high anthropogenic pressure,
19 makes the acid-base system complex. We summarize in this study the general knowledge on
20 the marine acid-base system as well as describe the peculiarities identified and reported for the
21 Baltic Sea specifically. In this context we discuss issues such as: dissociation constants in
22 brackish water, different chemical alkalinity models including contributions by organic acid-
23 base systems, long term changes of total alkalinity, anomalies of borate alkalinity and the acid-
24 base effects of biomass production and mineralization. Finally, we identify research gaps and
25 specify bottlenecks concerning the Baltic Sea acid-base system.

26

27

28

29

1 **1. Introduction**

2 The acid-base system controlling the seawater pH is formed by a number of chemical
3 substances having acid-base properties (Dickson et al., 2007). The importance of each of these
4 substances (structure of the acid-base system) depends on both concentrations of individual
5 constituents and their dissociation constants. The control by physical conditions (temperature,
6 salinity) and biogeochemical processes (e.g. biomass production and mineralization) as well as
7 interrelationships between individual components determine the functioning of the acid-base
8 system. The understanding of the structure and functioning of the acid-base system is necessary
9 to investigate important issues that shape the Baltic Sea ecosystem and that are of interest in
10 present-day chemical oceanography like: ocean acidification, calcium carbonate (CaCO_3)
11 formation/dissolution and carbon dioxide (CO_2) exchange through the air-sea interface.

12 The steady increase of atmospheric CO_2 concentrations leads to enhanced dissolution of CO_2
13 in the ocean. Since CO_2 dissolved in seawater forms the diprotic carbonic acid, hydrogen ions
14 are released. Although the major fraction of the hydrogen ions are taken up by carbonate ions
15 (a buffering reaction), a significant fraction stays in the water column and thus causes a
16 decrease in pH. This phenomenon is known in the scientific literature as “ocean acidification”
17 although seawater does not really become acidic but only moves from its alkaline character
18 towards the acidic regime (Riebesell et al., 2010). Ocean acidification has been recognized as
19 one of the greatest threats for marine ecosystems not only by the scientific community (e.g.
20 Bates et al., 2014; Zeebe, 2012), but also in European Union (EU) legislation. EU Marine
21 Strategy Framework Directive (MSFD, 2008) explicitly points out that the EU Member States
22 should put more attention to ocean acidification, and emphasizes the necessity to include
23 measurements of pH and of the CO_2 partial pressure (pCO_2) as descriptors for the
24 environmental status of marine regions.

25 There is a general belief that the magnitude of ocean acidification can sufficiently be quantified
26 from the atmospheric pCO_2 levels and the CO_2 exchange between seawater and the atmosphere
27 (Zeebe, 2012; Riebesell et al., 2010; Caldeira and Wicket, 2003). This is approximately true
28 for oceanic waters. However, it is not the case for coastal seas because several other processes
29 are influencing the seawater pH such as: CaCO_3 formation and/or dissolution, eutrophication
30 or oligotrophication, total alkalinity (A_T) consumption and production, weathering,
31 contribution by organic substances and terrestrial inputs. The general structure and functioning
32 of the acid-base system in seawater are relatively well identified, however still some aspects
33 contain a lot of gaps and/or shortcomings, which can lead to wrong conclusions and

1 questionable predictions of the future pH development (Orr et al., 2015; Koeve and Oschlies,
2 2012; Hunt et al., 2011; Dickson et al., 2007; Cai et al; 1998).

3 Since the CO₂ system is an integral part of the ocean acid-base system, it is impossible to
4 understand the CO₂ system and to assess processes such as the CO₂ gas exchange or CaCO₃
5 dissolution/formation, without a clear idea about the structure and functioning of the whole
6 acid-base system. The saturation states of calcite and aragonite are of crucial importance for
7 organisms forming their exoskeletons from CaCO₃. Although pelagic calcifying organisms do
8 not occur in the Baltic Sea this aspect is still relevant as some benthic organisms can also build
9 CaCO₃ into their shells (Tyrrell et al., 2008). Moreover, calcifying processes not only depend
10 on the acid-base system structure but in turn can modify it by decreasing the concentration of
11 carbonate ions (CO₃²⁻) and thus the alkalinity.

12 Carbon dioxide is a major component of the global carbon cycle and transport of CO₂ links all
13 Earth's compartments. It is believed that the world oceans absorb about 25 % of anthropogenic
14 CO₂ emissions. However, there is still a debate on the role that the shelf seas play in this context
15 (Le Quéré, 2016). The direction and magnitude of the CO₂ exchange through the air-sea
16 interface depends largely on the pCO₂ difference between seawater and the atmosphere. The
17 level of seawater pCO₂ is mainly controlled by the structure of the acid-base system that is
18 influenced by the combined effect of biological activity (biomass production vs.
19 mineralization), CO₂ exchange with the atmosphere and temperature (Emerson and Hedges,
20 2008).

21 In this context the Baltic Sea can be considered as a very complex ecosystem. On one hand the
22 low buffer capacity (Beldowski et al., 2010) makes the seawater vulnerable to acidification in
23 most regions of the Baltic Sea, on the other hand the sea is exposed to various anthropogenic
24 influences which have the potential to change the acid-base system and thus also seawater pH
25 and all pH-related processes. This makes the Baltic Sea different from the oceans for which the
26 CO₂ system as part of the acid-base system has been well characterized during the last decades
27 (Bates et al., 2014; Riebesell et al., 2010; Dickson et al., 2007). Considerable research effort
28 was also undertaken in recent years to investigate the Baltic Sea CO₂ system and its
29 peculiarities. However, this was done from different perspectives and resulted in specific
30 problem-oriented and divers knowledge. Therefore, the goal of this review article is to collect
31 and integrate the existing knowledge on the structure and functioning of the acid-base system
32 in the Baltic Sea, to point out the research gaps and thus also to address challenges for the
33 future research in this field.

1

2 **2. The standard acid-base model for ocean water**

3 Four measurable variables exist to describe the acid-base system of seawater. These are: pCO₂,
4 pH, total inorganic carbon concentration (C_T, known also as DIC) and total alkalinity (A_T). C_T
5 is the total concentration of all inorganic carbon forms present in seawater (Eq. 1), where
6 [CO₂*] represents the sum of molecular dissolved CO₂ and undissociated carbonic acid
7 (H₂CO₃):

$$8 \quad C_T = [CO_2^*] + [HCO_3^-] + [CO_3^{2-}] \quad (1)$$

9 Measurements of C_T are usually based on acidification of the sample and coulometric or
10 infrared detection of the extracted CO₂ (Dickson et al., 2007). A_T is defined as the excess of
11 proton acceptors (bases formed from weak acids with a dissociation constant of $K \leq 10^{-4.5}$ at 25
12 °C) over proton donors (acids with $K > 10^{-4.5}$) and expressed as the hydrogen ion equivalent in
13 one kilogram of sample (Dickson, 1981):

$$14 \quad A_T = [HCO_3^-] + 2[CO_3^{2-}] + [B(OH)_4^-] + [OH^-] + [HPO_4^{2-}] + 2[PO_4^{3-}] + [SiO(OH)_3^-] + [NH_3] + \\ 15 \quad [HS^-] + [\text{minor bases}] - [H^+] - [HSO_4^-] - [HF] - [H_3PO_4] - [\text{minor acids}] \quad (2)$$

16 A_T can be determined by open- or closed-cell acidimetric titration. For both C_T and A_T certified
17 reference materials are provided by the Andrew Dickson laboratory, UC San Diego, USA
18 (Dickson et al., 2007). pH measurements in seawater are usually calibrated against TRIS buffer.
19 Its characteristics is well known at salinities of 20-35 (Mosley et al., 2004; DelValls and
20 Dickson, 1998). The efforts to improve the TRIS buffer characteristics for lower salinities are
21 ongoing within the BONUS PINBAL project ([https://www.io-warnemuende.de/pinbal-](https://www.io-warnemuende.de/pinbal-home.html)
22 [home.html](https://www.io-warnemuende.de/pinbal-home.html)). The use of TRIS buffer helps to calibrate the systems for pH measurements but
23 does not fully guarantee the quality of the pH results as reference materials in seawater matrix
24 still do not exist for pH measurements. In many cases pH is still measured potentiometrically
25 on the NBS scale although spectrophotometric measurements on the total scale (pH_{tot}), which
26 takes into account also HSO₄⁻ ions, are currently state of the art in the field of chemical
27 oceanography. The concentration of CO₂* in seawater is obtained from the partial pressure of
28 CO₂ in air in equilibrium with seawater. According to the standard operating procedure
29 (Dickson et al., 2007), measurements of pCO₂ require continuous flow of water, which makes
30 this parameter difficult to measure in discrete samples.

31 All four variables are interacting and control the pH as described by a set of equilibrium
32 constants and mass balance equations. In general the interrelationships between these four

1 parameters facilitate the calculation of any two variables, when the two others are known, e.g.
2 through measurements, and when the dissociation constants of the involved acid-base reactions
3 are known for the respective temperature and salinity. This fact is used in biogeochemical
4 models for simulations of the marine CO₂ system. All biogeochemical models are based on the
5 transport and transformations of A_T and C_T, because these variables are independent of
6 temperature and pressure and behave conservatively with respect to mixing (mass
7 conservation). Moreover, straightforward techniques for sampling and analysis for A_T and C_T
8 exist and are supported by the availability of reference materials (Dickson et al., 2007). Hence,
9 these two variables are very well suited for CO₂ system studies. However, the use of A_T and
10 C_T for the CO₂ system studies is not free from limitations. To obtain high accuracy for the
11 calculation of pH and pCO₂ from measured or modelled A_T and C_T, all dissociation constants
12 and total concentrations of all non-CO₂ components of the acid-base system must be known.
13 This requirement is approximately fulfilled with regard to ocean studies where the total
14 concentrations of non-CO₂ acid-base components are either negligible or can be approximated
15 as a function of salinity (Riebesell et al., 2010). However, this issue is more critical for coastal
16 and shelf regions, where the biogeochemical composition of seawater shows regional
17 peculiarities (e.g. Kuliński 2014; Hernandez-Ayon et al., 2007; Cai et al., 1998). This is the
18 case for the Baltic Sea, which shows various unique biogeochemical characteristics.


19

20 **3. Peculiarities of the Baltic Sea acid-base system**

21 **3.1. Hydrographic setting**

22 The Baltic Sea is one of the largest brackish ecosystems in the world. This is caused by both
23 limited inflows of saline oceanic water through the shallow and narrow Danish Straits and high
24 river runoff. The drainage basin of the Baltic Sea is almost four times larger than the area of
25 the sea itself, while the mean annual freshwater supply (428 km³) constitutes about 2 % of the
26 Baltic Sea water volume (22·10³ km³). The specific features of the Baltic Sea hydrology cause
27 clear horizontal and vertical salinity gradients. Salinity in the surface layer fluctuate from 2
28 in the northern Bothnian Bay to >20 observed in the Kattegat. Dense and highly saline water,
29 which irregularly enters the Baltic Sea, sinks and moves along the sea bottom. This deep water
30 is separated from the brackish surface water layer by a permanent halocline located at a depth
31 of 60-70 m, which is at the same time the maximum mixed layer depth in the Baltic Sea (Lass
32 and Matthäus, 2008; Winsor et al., 2001). The stratification limits ventilation of the deep water

1 masses. Simultaneously, the deeper water layers are supplied with large amounts of particulate
2 organic matter that either originated from terrestrial sources or derived from the biomass
3 production in the surface under the influence of high nutrient concentrations (eutrophication).
4 Some fraction of the sedimentary organic matter mineralizes releasing CO₂ and consuming
5 oxidants, which leads to hypoxia or even anoxia in the bottom waters (HELCOM, 2009).



6
7 Fig. 1. Map of the Baltic Sea showing its division into natural basins and sub-basins
8 (modified after Kuliński and Pempkowiak, 2011).

9
10 **3.2. Dissociation constants in brackish water**

11 If any variables of the marine CO₂ system are to be used for biogeochemical studies, it is
12 advisable to measure these directly and not to derive it from the measurements of other
13 variables of the CO₂ system (Dickson et al., 2007). However, sometimes direct measurements
14 may be prevented by technical reasons or routine analytical methods do not exist. The latter

1 refers for example to the determination of CO_3^{2-} ion concentrations which are important for the
2 assessment of the state of the CaCO_3 saturation and thus for the dissolution or formation of
3 CaCO_3 shells. In this case, calculations on the basis of other known variables such as C_T , A_T ,
4 pCO_2 or pH are indispensable and require knowledge about the equilibrium constants of the
5 CO_2 system. Here we examine the availability of the necessary constants for low salinity
6 brackish water. Three fundamental equations characterize the equilibria of the marine CO_2
7 system. The first refers to the solubility of gaseous CO_2 in seawater:

$$8 \quad [\text{CO}_2^*] = k_0 * f\text{CO}_2 \quad (3)$$

9 The solubility constant, k_0 , relates the concentration of CO_2^* in seawater to the CO_2 fugacity,
10 $f\text{CO}_2$. In contrast to the CO_2 partial pressure, pCO_2 , the fugacity accounts for the non-ideal
11 behaviour of CO_2 at atmospheric conditions. It differs only slightly from the pCO_2 and in many
12 cases can be replaced by the pCO_2 . The solubility and thus k_0 decreases with increasing
13 temperature and salinity and vice versa. Studies by Weiss (1974) that describe k_0 as a function
14 of temperature and salinity have received widespread acceptance and are almost exclusively
15 used in chemical oceanography. The situation is somewhat more complicated for the
16 determination of the dissociation constants for carbonic acid because it is a diprotic acid. Using
17 again CO_2^* as variable, the two dissociation equilibria are given by:

$$18 \quad K_1 = [\text{H}^+][\text{HCO}_3^-]/[\text{CO}_2^*] \quad (4)$$

$$19 \quad K_2 = [\text{H}^+][\text{CO}_3^{2-}]/[\text{HCO}_3^-] \quad (5)$$

20 In these formulas the terms for the hydrogen ions are given in concentration units which include
21 HSO_4^- ion concentration (Dickson, 1984).

22 Systematic studies concerning the dissociation constants of carbonic acid in seawater were
23 already performed during the twenties and thirties of the last century. One of the leading
24 scientists in this field was Kurt Buch from the former Finnish Institute for Marine Research,
25 whose research was mainly related to the Baltic Sea, and whose laboratory studies on the CO_2
26 system focused on brackish water with salinities down to zero. The results of his lab work and
27 of field studies performed during 1927 to 1936 are summarized in Buch (1945). The presented
28 dissociation constants must be considered as “hybrid” constants because the hydrogen ions are
29 represented by the hydrogen ion activity whereas the CO_2 constituents refer to concentrations.

30 With the increasing awareness of the importance of the oceans for the uptake of anthropogenic
31 CO_2 and thus for climate change scenarios, large-scale measurement programmes concerning

1 the state of the marine CO₂ system on an oceanic scale were performed. These efforts were
 2 accompanied by numerous laboratory studies aimed at the determination of improved
 3 dissociation constants. All these constants referred to the total hydrogen concentration scale
 4 that includes the contribution by HSO₄⁻ (Dickson, 1984). However, the validity of most of the
 5 constants was confined to salinities that are encountered in ocean water. Exceptions are the
 6 constants suggested by Roy et al. (1993), which were determined for salinities down to values
 7 of 5 and could be used accordingly for brackish water. However, with regard to research in the
 8 Baltic Sea where large areas, e.g. in the Gulf of Bothnia, have surface water salinities less than
 9 5, the situation was unsatisfactory. It took until 2006 when Millero et al. (2006) published
 10 dissociation constants that covered the salinity range from 0 to 50 and that were consistent with
 11 the constants for fresh water. An update of these constants was performed in 2010 (Millero,
 12 2010) and since then this set of dissociation constants is state of the art for CO₂ research in
 13 brackish waters. The salinity and the temperature dependency of K₁ and K₂ are presented in
 14 Fig. 2a and 2b, respectively.

15 A direct comparison between the dissociation constants that were used in the past for brackish
 16 water is difficult because the constants by Buch (1945) are based on a pH scale that is based
 17 on the hydrogen ion activity. Since it is difficult to convert the activities to total hydrogen ion
 18 concentrations, we use the ratio K₁/K₂ for a comparison. This ratio is obtained by dividing the
 19 equilibrium equations for the first dissociation step (Eq. 4) through that of the second
 20 dissociation step (Eq. 5):


$$21 \quad \frac{K_1}{K_2} = \frac{[HCO_3^-]^2}{[CO_2^*][CO_3^{2-}]} \quad (6)$$

22 Eq. 6 shows that K₁/K₂ is not connected to the hydrogen ion concentration and is thus not
 23 affected by the choice of the pH scale. Hence it is suited as an indicator for the quality of the
 24 early dissociation constants determined by Buch (1945). We used the logarithmic notation,
 25 log(K₁/K₂) which is equivalent to pK₁ – pK₂, for a comparison of the dissociation constants
 26 suggested by Buch (1945), Millero (2010) and Roy et al. (1993). The differences Δlog(K₁/K₂)
 27 referring to Millero (2010) and Buch (1945), and Millero (2010) and Roy et al. (1993) indicated
 28 as M-B and M-R, respectively, are shown as a function of salinity in Fig. 2c.

29 At salinities >7, M-R is less than 0.03, but the differences increase rapidly at lower salinities
 30 and amount to 0.06 and 0.10 already at S = 5.4 and S = 3.6, respectively. Such differences in
 31 log(K₁/K₂) are equivalent to a temperature change of 1 °C. The deviations of the Buch (1945)
 32 data from those of Millero (2010) (M-B in Fig. 2c) ranged below 0.10 at salinities below 20.

1 This is a surprisingly small difference in view of the limited technical capabilities that were
2 available to Buch and his co-workers in the twenties and thirties of the last century.

3



4

1 Fig. 2. Dissociation constants (a) pK_1 and (b) pK_2 as a function of salinity calculated according
2 to Millero (2010) for three different temperatures: 0, 10 and 20 °C; (c) differences in $\log(K_1/K_2)$
3 between the dissociation constants obtained from Millero (2010) and Roy et al. (1993) (M-R,
4 blue dots) and between Millero (2010) and Buch (1945) (M-B, purple dots) at different
5 salinities.

6

7 When alkalinity is also used for the calculation of any variables of the CO_2 system, the
8 contributions of non- CO_2 acid-base components to A_T (Eq. 2), which are generally low, must
9 be taken into account. This requires knowledge about the concentration of the individual acid-
10 base pairs and about the corresponding dissociation constants. In the deep basins of the central
11 Baltic Sea, ammonia and sulphide anions are released during organic matter mineralization at
12 anoxic conditions (see Section 3.4.2). Both these constituents are strong proton acceptors
13 forming mainly HS^- and NH_4^+ . Since their concentrations may be on the order of several tens
14 and up to a few hundred $\mu mol L^{-1}$ (Nausch et al., 2008), respectively, they have a strong effect
15 on alkalinity and pH. Therefore, any calculations concerning the CO_2 system at anoxic
16 conditions must account for the dissociation equilibria of H_2S-HS^- (Millero et al., 1988) and
17 $NH_4^+-NH_3$ (Clegg and Whitfield, 1995; Johansson and Wedborg, 1980). The concentrations of
18 S^{2-} at the pH of anoxic water are extremely small and can be neglected.


19

20 **3.3. The central role of alkalinity**

21 **3.3.1. Regional A_T vs. S distribution, consequences for pH distribution pattern**

22 There are several different A_T vs. S regimes in the Baltic Sea (Fig. 3, Beldowski et al., 2010).
23 They reflect different A_T concentrations in the respective rivers, which can be deduced from
24 extrapolation of regional A_T vs. S relationships to zero salinity. Low alkalinities are observed
25 in rivers entering the Gulf of Bothnia whereas rivers from south-eastern part of the Baltic
26 catchment, as shown by the A_T vs. S plot for the Gulf of Riga (Fig. 3), are rich in alkalinity.
27 These differences are a consequence of the geological conditions and weathering processes in
28 the respective catchment areas. As a result lower alkalinities and lower mean pH are observed
29 in the Gulf of Bothnia and Finland, whereas higher mean alkalinities and thus somewhat higher
30 pH are found in the Gulf of Riga and also in the Gdansk Bay (Kulinski et al., 2014; Beldowski
31 et al., 2010; Hjalmarsson et al., 2008).

1 The central Baltic Sea acts as a mixing chamber for the different water masses, including water
2 originating from the North Sea. This results in alkalinity in the surface water of the Baltic
3 Proper (salinity around 7) of about 1600-1700 $\mu\text{mol kg}^{-1}$.



4
5 Fig. 3. Different A_T vs. S regimes observed in the Baltic Sea (modified after Beldowski et al.,
6 2010)

7
8 An important, but still not well investigated, aspect of the A_T supply to the Baltic Sea is the A_T
9 seasonality in river water. As A_T is one of the variables used in biogeochemical models to study
10 the CO_2 system, well described spatial and temporal variability of A_T loads could improve the
11 characterization of the A_T distribution in the Baltic Sea and thus increase the accuracy of the
12 modelled pH and pCO_2 . This could be achieved by evaluation of the A_T monitoring data which
13 are regularly collected by some Baltic Sea states within the framework of the HELCOM
14 monitoring program. In some countries such as Sweden the monitoring data are publicly
15 accessible, in others still some limitations exist.


16 A_T can also enter the Baltic Sea by submarine groundwater discharge (SGD). The chemical
17 composition of groundwater, as in river water, depends to some extent on the geological
18 structure of the catchment through which the water flows. Thus, SGD entering the Baltic Sea
19 along the continental part of the coast can be rich in A_T . Szymczycha et al. (2014) noticed
20 significant C_T concentrations ($5400 \mu\text{mol kg}^{-1}$ on average) in SGD seeping to the southern

1 Baltic along the Polish coast. Although there might be some contribution from CO₂ to this high
2 C_T, it is very likely that SGD can play, at least locally, an important role for the A_T budget.
3 However, direct measurements of A_T concentrations and loads in SGD have not been reported
4 in the Baltic Sea so far.

5 At equilibrium with the atmospheric CO₂, A_T controls C_T and thus pH. Hence, pH may be
6 depicted as a function of A_T and pCO₂. A sensitivity study performed by Omstedt et al., (2010)
7 indicates that the pH of the Baltic Sea surface water that is at equilibrium with the atmosphere
8 (pCO₂ of ca. 400 μatm at 0 °C) can vary between 7.7 and 8.3 depending on the A_T (Fig. 4).
9 This range is significantly higher than that observed in the open ocean, where A_T oscillates
10 only in a narrow range: 2170-2460 μmol kg⁻¹ (Riebesell et al., 2010). The reason for that is the
11 high spatial variability of A_T in the surface Baltic Sea waters, from low A_T (below 1000 μmol
12 kg⁻¹) observed in the Bothnian Bay to the A_T-rich (more than 3000 μmol kg⁻¹) estuaries of the
13 large continental rivers. The diagram presented in Fig. 4 also shows that higher A_T reduces
14 shifts in seawater pH caused by changes in pCO₂.

15

16



1

2 Fig. 4. Distribution of pH_{tot} as a function of A_T and pCO_2 at salinity of 7.5 and temperature of
 3 0°C (modified after Omstedt et al., 2010). The A_T range represents findings by Beldowski et
 4 al. (2010). SSW is an abbreviation for surface seawater.

5

6 3.3.2. Long-term A_T changes, consequences for ocean acidification

7 Long-term high resolution pH measurements at station BATS (Bermuda Atlantic Time Series)
 8 in the sub-tropical North Atlantic have shown that ocean acidification occurs at a rate that is
 9 thermodynamically consistent with the increase of the atmospheric CO_2 (Bates, 2014). This
 10 gives reason to expect that the progression of ocean acidification can be assessed on the basis
 11 of CO_2 emission scenarios and the modelling of the effect on the atmospheric CO_2 . It also
 12 implies that changes in alkalinity will not occur in the future. On time scales of decades this

1 may be a reasonable assumption for the oceans which react very slowly to internal or external
2 changes. However, the situation is different for the Baltic Sea that is under the immediate
3 influence of natural and anthropogenic processes on land. The first hints of increasing alkalinity
4 in the surface water of the central Baltic Sea were reported by Schneider et al. (2015) who
5 argued that this may have considerably mitigated the acidification due to increasing
6 atmospheric CO₂. Müller et al. (2016) took up this idea and performed a thorough and
7 comprehensive statistical analysis of all alkalinity data that were available since the start of
8 CO₂ research in the Baltic Sea at the beginning of the last century. The authors focused on
9 high-quality data measured after 1995 and detected a distinct A_T trend that showed a clear
10 regional gradient. The highest trend was found in the Gulf of Bothnia (7.0 μmol kg⁻¹ yr⁻¹),
11 followed by 3.4 μmol kg⁻¹ yr⁻¹ in the central Baltic, whereas no trend could be detected in the
12 Kattegat. Müller et al. (2016) estimated that the changes in A_T have reduced the acidification
13 effect in the central Baltic Sea by about 50 % and that the Gulf of Bothnia was not subjected
14 at all to acidification during the last 20 years. Several reasons for the increasing A_T are
15 discussed by Müller et al. (2016), but a major player could not be identified and, hence, it
16 remains unclear whether the trend will continue in the future.

17

18 **3.3.3. Effect of organic alkalinity**

19 The Baltic Sea water contains 3-5 times more organic matter than open ocean waters. In the
20 surface water of the open Baltic Sea, concentrations of dissolved organic carbon (DOC) range
21 from about 260 to about 480 μmol C L⁻¹, while those in the surface water (top 100 m) of the
22 Atlantic Ocean are much lower and range between 50 and 80 μmol C L⁻¹ (Hoikkala et al., 2015;
23 Carlson et al., 2010). This is a result of both high inputs of terrestrial organic matter and
24 eutrophication driven by nutrient supply from land (Hoikkala et al., 2015; Kuliński and
25 Pempkowiak, 2011; Kuliński et al., 2011). Organic substances contain functional groups, some
26 of which (carboxylic, phenolic, amines) have acidic character and can dissociate in seawater
27 releasing protons (H⁺). This contributes to pH decrease. However, as most of these groups are
28 believed to act as weak acids (pK_a>4.5), their dissociation releases also an equivalent of organic
29 anions being strong bases. This, according to Eq. 2, causes no change in A_T concentration since
30 an equivalent amount of protons have been released. It affects the internal structure of A_T by
31 changing the contributions of different A_T components according to their individual
32 dissociation constants (Cai et al., 1998; Hunt et al., 2011; Kuliński et al., 2014). The lower the
33 pK_a of an organic acid added to the system, the greater are the shifts in the internal A_T

1 distribution. As a consequence bases forming the acid-base system are partially protonated and
2 their concentration decreases while the concentration of the corresponding undissociated acid
3 is increasing. In case of the carbonate alkalinity this means a reduction of CO_3^{2-} and an increase
4 of H_2CO_3 and thus of the pCO_2 .

5 Kuliński et al. (2014) estimated the organic alkalinity (A_{org}) in the Baltic Sea as an excess
6 alkalinity calculated from the difference between measured A_{T} and that calculated from C_{T} and
7 pH or pCO_2 . They found A_{org} of 25-35 $\mu\text{mol kg}^{-1}$ in the surface Baltic Sea water (from the
8 Arkona Basin to the Bothnian Bay) without any clear spatial distribution pattern. This A_{org}
9 contribution amounted from 1.5 % to 3.5 % of the measured A_{T} values. Even higher values of
10 more than 50 $\mu\text{mol kg}^{-1}$ were found by Hammer et al. (2017) in the Baltic Sea surface mixed
11 layer during the spring phytoplankton bloom. They reported also that A_{org} decreases with depth
12 and approaches zero below the permanent halocline due to pH decrease.

13 In sensitivity studies Kuliński et al. (2014) showed for the Baltic Sea that calculations
14 concerning the CO_2 system that are using measured A_{T} may lead to significant errors if A_{org} is
15 ignored. Highest deviations between calculated and observed values were found when A_{T} was
16 used together with C_{T} for computations of pCO_2 and pH. The pCO_2 values obtained in that way
17 were 27-56 % lower than the measured ones, while pH was overestimated by more than 0.4
18 units. These results are especially important as this combination (A_{T} and C_{T}) is used in
19 biogeochemical models (Edman and Omstedt, 2013; Kuznetsov and Neumann, 2013; Omstedt
20 et al., 2012 and 2009) because they are conservative variables. This means that they are
21 independent of pressure and temperature and follow the law of conservation of mass during
22 mixing, which are prerequisites for variables transported in models. The first attempts to
23 include an A_{org} parametrisation in biogeochemical models have been made by Gustafsson et
24 al., (2015) and Omstedt et al. (2015). However, the sensitivity of the models with regard to the
25 inclusion/negligence of A_{org} has not yet been reported.

26 The source of the calculation errors related to ignoring A_{org} is due to the fact that measurements
27 include all inorganic and organic contributors, while subsequent calculations using the standard
28 chemical A_{T} model and respective software, interpret the measured A_{T} only as inorganic
29 alkalinity. Including organic substances into the A_{T} model is, however, challenging as there are
30 number of organic substances having acidic functional groups and, according to the A_{T}
31 definition, for each of them the information on concentration and dissociation constants must
32 be available. Accounting for A_{org} in the chemical A_{T} model is less important for oceanic
33 research because the low concentrations of dissolved organic matter (DOM) in the oceans cause

1 only minor effects. As a first approximation of the acid-base properties of DOM for the Baltic
2 Sea water, Kuliński et al. (2014) proposed to use one single bulk dissociation constant, pK_{DOM}
3 and relate A_{org} to the concentration of dissolved organic carbon (DOC), a commonly measured
4 parameter. According to their studies, slightly corrected by Ulfsbo et al. (2015), pK_{DOM} in the
5 Baltic Sea amounts to 7.34. They also estimated experimentally that 12 % of DOC acts as a
6 carrier for acidic functional groups in the Baltic Sea DOM. The model studies by Ulfsbo et al.
7 (2015) showed that this experimentally derived share fits to the share estimated from the
8 hypothetical structure of the fulvic acids. They also suggested that the method proposed by
9 Kuliński et al. (2014) is the best available approach for representing organic alkalinity in
10 biogeochemical models at the current state of knowledge. They also showed that organic matter
11 (as represented by the fulvic acids) contains also some functional groups having pK_a already
12 below 4.5. According to the definition (Dickson, 1981) they should be considered as strong
13 acids. In experimental studies Hammer et al. (2017) also found that humic and/or fulvic
14 substances are more acidic than the bulk DOM naturally occurring in the Baltic Sea. This may
15 also be a source of uncertainty in the A_T measurements as the determination of the titration
16 end-point requires to titrate the samples through this pH region.

17

18 **3.3.4. Role of borate alkalinity**

19 The carbonate system plays a central role in the marine acid-base system. Thus, studies on
20 seawater pH and buffering capacity focus usually on the carbon species. Less attention is paid
21 to boron, though borates are, after bicarbonates and carbonates, the third most abundant
22 constituent of seawater A_T (Eq. 2). Boron exists in seawater in form of weak boric acid,
23 $B(OH)_3$, and the anion, $B(OH)_4^-$. The high pK_a (8.60 at salinity 35 and temperature 25 °C)
24 means that at seawater pH of about 8 the undissociated boric acid predominates. The boron
25 concentration, B_T , in seawater is approximated as a function of salinity or, for the historical
26 data, as a function of chlorinity ($S = 1.80655 * Cl$). First measurements of boron concentrations
27 in the Baltic Sea were reported by Buch (1945), who found that $B_T [mg\ kg^{-1}] = 0.133 * S$. This
28 finding was confirmed recently by Lee et al. (2010) for oceanic waters. Another relationship
29 linking B_T and S and often used in acid-base system studies is that by Uppström (1974), who
30 reported that $B_T [mg\ kg^{-1}] = 0.128 * S$. All these formulas suggest a fixed B_T/S ratio which is
31 only the case if the river water that enters the Baltic Sea contains no boron. However, the
32 studies by Kremling (1970, 1972) indicated that this is not the case and that a B_T concentration
33 anomaly exists in the Baltic Sea, where the experimentally determined B_T vs. S relationship

1 yielded a river water B_T (anomaly term) of $0.15 - 0.20 \text{ mg kg}^{-1}$ at $S = 0$. This anomaly is not
2 included in the chemical A_T model commonly used for numerical simulations of the CO_2
3 system and, similar to the effects of ignoring A_{org} , may lead to wrong conclusions. This can be
4 especially critical at low salinities, where the effect of the anomaly is the largest.

5

6 **3.4. Modulation of the acid-base system by organic matter production/mineralization** 7 **(concurrent A_T and C_T changes)**


8 **3.4.1. Biomass production**

9 In the case that pH changes refer to equilibrium with the atmosphere, it may be meaningful to
10 consider the pH at a given alkalinity as a function of $p\text{CO}_2$, because the atmospheric CO_2 is
11 then the only driver for any pH changes. However, when assessing pH changes as the
12 consequence of internal biogeochemical transformations, it is more appropriate to consider A_T
13 and C_T as the controls for pH and $p\text{CO}_2$. Biomass production primarily alters C_T because
14 phytoplankton (but also other plants) consume CO_2 in the course of photosynthesis (Fig. 5). As
15 a consequence it increases the pH and lowers the $p\text{CO}_2$ in the upper water layers and causes a
16 $p\text{CO}_2$ disequilibrium between seawater and the atmosphere. This leads to CO_2 uptake by the
17 seawater and thus counteracts the effect of biomass production and dampens the pH increase.
18 However, the CO_2 exchange through the air-sea interface is much slower than the effects of
19 biological processes. Therefore, the Baltic Sea water is in almost permanent $p\text{CO}_2$
20 disequilibrium with the atmosphere throughout the year. In spring and summer seawater is
21 undersaturated with respect to atmospheric CO_2 with two characteristic $p\text{CO}_2$ minima and two
22 pH maxima (Fig. 5) which reflect the spring bloom and the mid-summer nitrogen fixation
23 period. In autumn and winter the Baltic Sea surface water $p\text{CO}_2$ increases over the atmospheric
24 values as a consequence of less active production in the upper water column and transport of
25 deeper CO_2 -enriched water to the surface by mixing (Fig. 5; Schneider, 2011).

26 Another way in which biomass production influences seawater pH and $p\text{CO}_2$ is related to
27 nitrate consumption. As phytoplankton assimilate nitrate for growth an equivalent of H^+ is also
28 transported to the cells to keep the charge balance neutral. According to Eq. 2 this increases
29 the seawater A_T and thus raises the pH during biomass production (Brewer and Goldman,
30 1976). This assimilation also decreases the $p\text{CO}_2$ and therefore reinforces the drop in $p\text{CO}_2$ by
31 biomass production. Some small A_T changes can be also induced by phosphate consumption.

1 However, due to low phosphate concentration the effect is negligible (Wolf-Gladrow et al.,
2 2007).

3 Biological production also may consume CO_3^{2-} ions in the calcification processes. This causes
4 A_T reduction and in consequence pCO_2 increase and also pH decrease. However, this
5 mechanism has been recognized as less important in the Baltic Sea. According to studies by
6 Tyrrell et al. (2008) the Baltic Sea surface water is undersaturated in winter with respect to
7 both aragonite and calcite which are biogenic modifications of CaCO_3 . This possibly prevents
8 the growth of calcifying plankton and is the reason for the absence of coccolithophores in the
9 Baltic Sea.



10

11 Fig. 5. Typical seasonality of pCO_2 in surface water (blue line) and the atmosphere (green line)
12 in the Eastern Gotland Sea modified after Schneider (2011) and presented together with pH
13 calculated from the shown pCO_2 data and the mean A_T value.

14


15 3.4.2. Remineralization

16 In contrast to biomass production, CO_2 is released during the remineralization processes and
17 causes an increase in C_T . Consequently, this leads to a pH decrease and pCO_2 increase. Some
18 fraction of organic matter produced in situ undergoes remineralization in the upper water
19 layers. Particulate organic matter (POM) may be mineralized directly or via prior release of
20 dissolved organic matter (DOM). In the Baltic Sea terrigenous organic matter entering the sea
21 from rivers constitutes an important part of the organic matter. Kuliński and Pempkowiak

1 (2011) reported that the Baltic Sea receives annually 340 Gmol of organic carbon from land.
2 The model studies by Gustafsson et al. (2014a) suggested that as much as 39.5% of terrigenous
3 organic carbon that enters the Baltic Sea undergoes mineralization without distinguishing
4 between mineralization pathways. This makes that it is difficult to estimate all the
5 consequences of that terrestrial organic carbon loss for the Baltic Sea acid-base system. The
6 incubation experiments performed by Kuliński et al. (2016) indicated that ca. 20 % of terrestrial
7 dissolved organic carbon and 34 % of DOC present in the Baltic Sea is bioavailable. It is
8 important to mention here that the latter result refers likely to the mixture of terrestrial and
9 marine DOC.


10 Whereas the A_T in surface water is mainly controlled by mixing of different water masses, the
11 deep water A_T distribution depends additionally on the organic matter transformations by
12 various redox processes (Brenner et al., 2016; Krumins et al., 2013; Thomas et al., 2009; Schulz
13 and Zabel, 2006). A certain fraction of the organic matter produced in the euphotic zone is
14 exported to deeper water layers and to surface sediments, where it undergoes mineralization,
15 produces CO_2 , and changes the alkalinity. The change in A_T depends on the oxidant that is
16 required for the mineralization and may cover a wide range. Therefore, it is not possible to
17 predict the change of pH or pCO_2 during the mineralization process without knowledge of the
18 oxidant. In the presence of oxygen, mineralization takes place according to Eq. 7 which is
19 reversing the bulk photosynthesis reaction. However, in sediments and in deep water layers of
20 some basins of the central Baltic, where longer periods (years) of stagnation occur, oxygen
21 may be entirely depleted. Organic matter can then be mineralized in a certain
22 thermodynamically controlled sequence by other oxidants (Schulz and Zabel, 2006). First,
23 manganese dioxide takes over the role of oxygen (Eq. 8), followed by denitrification where
24 nitrate acts as oxidant (Eq. 9). After nitrate is consumed, the mineralization process may be
25 continued by iron (III) oxide (Eq. 10) before sulphate oxidizes the organic matter and generates
26 hydrogen sulphide (Eq. 11). In the Baltic Sea these processes may take place in the water
27 column where a pelagic redoxcline (an interface between oxic and anoxic conditions) can
28 develop during longer periods of stagnation. Only the final mineralization, that is an internal
29 oxidation and generates methane (methanogenesis, Eq. 12) after the sulphate concentration has
30 approached zero, is confined to deeper sediment layers only. Except oxic mineralization, all
31 other processes are connected with a simultaneous A_T increase caused either by release of
32 strong bases (S^{2-}) or by consumption of H^+ (see Eq. 2) (Ulfsbo et al., 2011; Schneider et al.,
33 2015; Lukawska-Matuszewska, 2016). The change in A_T (ΔA_T) for different mineralization

1 pathways is given in moles A_T per 106 moles of released CO_2 (Eqs. 7 – 12). It shows large
 2 differences and is highest when the insoluble oxides of iron (III) and manganese (IV) are
 3 reduced during the mineralization process. As a consequence of the higher A_T level, the pCO_2
 4 increase and pH decrease by increasing C_T during the mineralization is mitigated or may even
 5 be reversed. In the eastern Gotland Sea these mechanisms stabilize the pH in the deep, anoxic
 6 water layers at level of about 7 (Hammer et al., 2017).




8 Oxic mineralization, $\Delta A_T = -16$

9




12 Manganese reduction, $\Delta A_T = +472$

13




16 Denitrification, $\Delta A_T = +100.8$

17



20 Iron reduction, $\Delta A_T = +864$

21



24 Sulphate reduction, $\Delta A_T = +122$

25



27 Methanogenesis, $\Delta A_T = +16$

28

1 Generally, benthic processes linked to early diagenesis of organic matter can be an important
2 source for A_T to the water column (Brenner et al., 2016). This corresponds to the findings by
3 Gustafsson et al. (2014b), who found in a model study that external sinks and sources of A_T in
4 the Baltic Sea are imbalanced and cannot reproduce the observed A_T inventory, and that an
5 internal A_T source must exist in the Baltic Sea. However, it must be taken into account that
6 alkalinity released from sediments as sulphide or ammonia only influence the acid-base system
7 locally. Upon contact with oxic water most of the A_T generating processes – except
8 denitrification - are reversed and do not constitute a permanent A_T source. Some minor increase
9 of A_T can be also expected due to the burial of iron (II) minerals like pyrite or vivianite (Reed
10 et al., 2016).

11

12 **4. Challenges for future research.**

13 The general knowledge of the marine acid-base system was gained mainly from the oceanic
14 research. It has been integrated and documented in the form of a guide (Dickson et al., 2007)
15 and included in different software packages (Orr et al., 2015), that allow for computations of
16 two out of the four acid-base system parameters (pCO_2 , pH, A_T , C_T), when the other two are
17 known (e.g. by measurements). Although this works nicely for the open ocean waters, problems
18 appear when that knowledge is used for studying the acid-base system related processes in
19 coastal and shelf seas. This uncovers research gaps and bottlenecks but also define challenges
20 for marine biogeochemists working in these regions.

21 In this context the Baltic Sea can be considered as a model experimental field, where strong
22 horizontal and vertical salinity gradients, permanent stratification of the water column,
23 eutrophication, high organic matter concentrations and high anthropogenic pressure make the
24 acid-base system even more tangled. The challenges are related to analytical methods,
25 interpretation of the data and parametrization of the acid-base system related processes for the
26 use in numerical models. Great effort has recently been made to adapt spectrophotometric pH
27 measurements based on m-cresol purple to the Baltic Sea conditions. This was done by
28 Hammer et al. (2014) and was recently improved within a framework of the BONUS PINBAL
29 project. Further progress can be expected if pH reference materials for brackish water are made
30 available in the future. For the pCO_2 measurements the state-of-the-art method that is based on
31 measuring the CO_2 content in air equilibrated with the seawater is commonly used (e.g.
32 Schneider et al., 2014). Due to a relatively long response time, this method may, however, not
33 be fast enough to resolve steep horizontal pCO_2 gradients in the vicinities of river mouths. The

1 development of accurate and precise sensors for pCO₂, like optodes, could improve our
2 understanding of the CO₂ system in the transition area from the coast to the open sea.
3 Additionally, it would facilitate pCO₂ measurements in discrete samples, which are highly
4 desired for investigations of the organic matter mineralization processes in the water column,
5 but difficult to obtain with common underway measurement techniques. The determination of
6 A_T by titration with an acid (HCl) allows a high accuracy (+/-2 μmol kg⁻¹) for ocean water.
7 This is difficult to achieve for Baltic Sea water because the detection of the endpoint of the A_T
8 titration is more difficult due to the influence of organic acid-base constituents. The studies by
9 Ulfsbo et al. (2015) showed that fulvic and humic acids, substances widely spread in the Baltic
10 Sea, have relatively high concentrations of acid-base constituents, which have pK_a values close
11 to the pK_a of carbonic acid and may thus impede the correct detection of the titration endpoint.
12 C_T measurements are less critical since reference materials exist and the analytical procedure
13 is relatively straightforward. However, development of high accuracy underway methods for
14 C_T (but also A_T) would improve the horizontal coverage of the data.

15 Interpretations of the acid-base system data require appropriate parameterization of the related
16 processes and chemical reactions. With regard to the Baltic Sea some peculiarities of the
17 composition of the acid-base system must be taken into account which are neglectable in ocean
18 waters. An example is the influence of DOM on the acid-base system. In the oceans it is ignored
19 due to low DOM concentrations but in the Baltic Sea A_{org} becomes a noticeable component of
20 A_T (Kuliński et al., 2014). The challenge remains to include the contribution by DOM to the
21 chemical A_T model. The approach suggested by Kuliński et al. (2014) to use the so called bulk
22 dissociation constant, K_{DOM}, is only a first approximation, as K_{DOM} has no real thermodynamic
23 meaning. It also does not allow for distinguishing the influence of individual substances.
24 Respective improvements would require substantial progress in analytical techniques, as the
25 characterization of the structure and composition of DOM is still far from being satisfactory
26 (Nebbioso and Piccolo, 2013). Problems also arise due to the ion anomalies observed in the
27 Baltic Sea (Kremling 1970, 1972) that are not taken into account in the A_T model. An example
28 is boron, whose concentration is usually approximated by the assumption of a constant
29 boron/salinity ratio. However, the function linking both these parameters does not account for
30 river water as a source of boron. Furthermore, common software for CO₂ system calculation
31 also omits seawater constituents that are generated at anoxic conditions. These are mainly the
32 acid-base systems: H₂S-HS⁻-S²⁻ and NH₄⁺-NH₃ which require consideration in computational
33 models. All these shortcomings may cause uncertainties in any calculations of the CO₂ system

1 including those in biogeochemical models when A_T is used as a controlling variable. It is also
2 recommended to perform sensitivity studies in order to estimate the quantitative importance of
3 inadequate chemical characterization of A_T .

4 Furthermore, we promote the implementation of the acid-base system (or marine CO_2 system)
5 into biogeochemical models. This includes the simulation of surface water CO_2 partial
6 pressure, $p\text{CO}_2$. The $p\text{CO}_2$ is an ideal validation variable because on the one hand it can easily
7 be measured with high spatiotemporal resolution (e.g. on Voluntary Observing Ships, VOS),
8 and on the other hand it reflects the biogeochemical transformations such as biomass
9 production. Hence, agreement between simulated and measured $p\text{CO}_2$ data give an indication
10 of the quality of the biogeochemical process parameterization. Or, vice versa, unsatisfactory
11 agreement gives reason to rethink the description of the respective processes in the model.

12 The latter aspect was the focus of several model studies which initially failed to reproduce the
13 observed $p\text{CO}_2$ draw down during spring and summer (Fig. 5). This indicated that the
14 traditional parameterization of the biomass production that is based on the Redfield (1963)
15 C/N/P ratios and on nutrient concentrations at the start of the productive period, did not reflect
16 the reality. Omstedt et al. (2009) solved the problem by including a fractional nutrient release
17 into their model. This meant that nutrients that have been used for production are partly released
18 from the POM and reused for new production. In another study Kuznetsov et al. (2011)
19 introduced an additional cyanobacteria group and flexible C/N/P ratios into their model in order
20 to achieve approximate agreement between the simulated and measured $p\text{CO}_2$. A further study
21 by Gustafsson et al. (2014a) investigated the changes in the simulated CO_2 air-sea flux after
22 including organic alkalinity and changes in external inputs of carbon, alkalinity and nutrients.
23 Modelling the CO_2 system was also used to simulate the vertical distribution of total CO_2 and
24 alkalinity in stagnant waters of the deep basins (Edman and Omstedt, 2013) and to assess the
25 importance of internal alkalinity generation (Gustafsson et al., 2014b). Finally, model
26 calculations were used to estimate future changes in the Baltic Sea CO_2 system and in particular
27 in the development of the ocean acidification effect (Omstedt et al., 2012; Kuznetsov and
28 Neumann, 2013). Both simulations indicated that the increasing atmospheric CO_2 will mainly
29 control long-term changes in pH, of course, not taking into account the recently reported
30 current increase in alkalinity (see Chapter 3.3.2; Müller et al., 2016).

31

32 **5. Summary and conclusions**

1 With our review we have proposed that the Baltic Sea acid-base system cannot be explained
2 by dissolved CO₂ and a minor contribution of boric acid. Whereas acid-base components other
3 than CO₂ and borate may be ignored in ocean waters, this is not the case in the Baltic Sea and,
4 presumably, other marginal and semi-enclosed seas, which in a similar way are strongly
5 impacted by high biomass production and natural and anthropogenic processes in connected
6 catchment areas as well. The complexity of the Baltic Sea acid-base system is displayed in the
7 formulation of the alkalinity (Eq. 2), which is the central variable for the characterization of
8 the acid-base properties of seawater. The peculiarities of the Baltic Sea alkalinity system are
9 either caused by specific internal processes related to intense production/mineralization of
10 organic matter, or to the immediate effect of riverine input of acidic substances. This riverine
11 input refers mostly to dissolved organic matter which may strongly affect the composition of
12 the alkalinity in the Baltic Sea surface water (Kuliński et al., 2014), but also to boric acid
13 (borate) which in ocean water is linked to salinity by a constant ratio, but may show distinct
14 deviations from this ratio in some of the Baltic Sea estuaries because of boron input by river
15 water.

16 Internal biogeochemical processes such as organic matter production or mineralization also
17 have the potential to affect the acid-base system in the Baltic Sea. Alkalinity changes by
18 organic matter production caused by the removal of H⁺ during the uptake of nitrate are of minor
19 importance. This may be different if the production is associated with calcification, which
20 reduces the alkalinity, but the abundance of calcifying plankton in the Baltic Sea is restricted
21 to the high-salinity Kattegat region. Although the effect of organic matter production on
22 alkalinity is minor, it is the major control for the seasonal modulation of the acid-base
23 properties such as pH and pCO₂.

24 In contrast, organic matter mineralization taking place at anoxic conditions in the Baltic Sea
25 deep basins causes changes in the acid base system which are specific for marine systems with
26 a pelagic redoxcline. Anoxic mineralization of organic matter generates large amounts of
27 alkalinity by the formation of sulphide ions and the release of ammonia. This reduces the
28 increase of the pCO₂ in the deep water and stabilizes the pH at a value close to 7. Although
29 these processes are reversed upon re-oxidation during a deep water renewal event and do not
30 affect the acid-base system of the Baltic Sea as a whole, they do influence the deep water redox
31 chemistry which is partly controlled by the pH. On the other hand the existence of a pelagic
32 redoxcline has another more far-reaching importance. It is the medium for intense

1 denitrification, which increases the alkalinity not only locally, but may affect the alkalinity
2 budget of the entire Baltic Sea proper (Gustafsson et al., 2014).

3 Our compilation of the major features of the Baltic Sea acid-base system indicates that an exact
4 quantitative treatment of its properties is difficult to achieve. This refers to both the
5 biogeochemical modelling and the interpretation of measured alkalinity in terms of the
6 calculation of individual alkalinity contributions. An example is the calculation of the
7 carbonate alkalinity that is necessary for the full characterization of the CO₂ system based on
8 measurements of alkalinity together with another variable. This requires knowledge of the
9 dissociation constants and the determination of the total concentrations of the individual
10 compounds with acid-base properties. Regarding inorganic alkalinity contributions, it may be
11 realistic to obtain this information. But in view of our limited knowledge concerning the
12 composition of the dissolved organic matter, it is currently impossible to specify and
13 characterize the contributions of individual organic acid-base compounds to the alkalinity.
14 Hence we conclude that this is one of the greatest challenges for more adequate comprehensive
15 physico-chemical characterization of the acid-base system in the Baltic Sea and likely in other
16 coastal seas. Another important bottleneck, especially for the biogeochemical modelling, is
17 also an insufficient knowledge on short and long term development of alkalinity loads from
18 land caused by processes occurring in the catchment.

19

20 **6. Acknowledgements**

21 The study was completed thanks to funding provided by the National Science Centre, Poland,
22 grants no. 2015/19/B/ST10/02120 and 2014/13/D/ST10/01885. Significant financial support
23 was provided also by BONUS and the Polish National Centre for Research and Development
24 – sponsors of BONUS PINBAL project (grant no. BONUS-INNO-2012-03/2014). The
25 contribution of B. Schneider to this study was supported by the Foundation for Polish Science
26 within the Alexander von Humboldt Polish Honorary Research Scholarship. This study
27 comprises a part of the Baltic Earth — Earth System Science for the Baltic Sea Region. We
28 thank both anonymous reviewers for insightful and constructive comments and suggestions.

29

30 **References**

31 Bates, N. R., Astor, Y. M., Church, M. J., Currie, K., Dore, J. E., Gonzalez-Davila, M.,
32 Lorenzoni, L., Muller-Karger, F. E., Olafsson, J., and Santana-Casiano, J. M.: A time-

1 series view of changing surface ocean chemistry due to ocean uptake of anthropogenic
2 CO₂ and ocean acidification, *Oceanography*, 27, 126-141, 2014.

3 Beldowski, J., Loeffler, A., Schneider, B. and Joensuu, L.: Distribution and biogeochemical
4 control of total CO₂ and total alkalinity in the Baltic Sea, *J. Mar. Syst.*, 81, 252–259,
5 2010.

6 Brenner, H., Braeckman, U., Le Guitton, M. and Meysman, F.J.R.: The impact of sedimentary
7 alkalinity release on the water column CO₂ system in the North Sea., *Biogeosciences*,
8 13, 841-863, 2016.

9 Brewer, P. G., and Goldman, J. C.: Alkalinity changes generated by phytoplankton growth,
10 *Limnol. Oceanogr.*, 21, 108–117, 1976.

11 Buch, K.: Kolsyrejamvikten i Baltiska Havet., *Fennia*, 68, 5, 29–81, 1945.

12 Cai, W.J., Wang, Y.C. and Hodson, R.E.: Acid–base properties of dissolved organic matter in
13 the estuarine waters of Georgia, USA, *Geochim. Cosmochim. Ac.*, 62, 473–483, 1998.

14 Caldeira, K., Wickett, M. E., Anthropogenic carbon and ocean pH, *Nature*, 425, 365–365,
15 2003.

16 Carlson, C.A., Hansell, D.A., Nelson, N.B., Siegel, D.A., Smethie, W.M., Khatiwala, S.,
17 Meyers, M.M., Halewood, E., Dissolved organic carbon export and subsequent
18 remineralization in the mesopelagic and bathypelagic realms of the North Atlantic
19 basin, *Deep-Sea Res. II — Top. Stud. Oceanogr.*, 57, 1433–1445, 2010.

20 Clegg, S. L. and Whitfield, M.: Chemical model of seawater including dissolved ammonia and
21 the stoichiometric dissociation constant of ammonia in estuarine water and seawater
22 from -2 to 40°C, *Geochem. Cosmochim. Ac.*, 59, 2403-2421, 1995.

23 DelValls, T.A. and Dickson, A.G.: The pH of buffers based on 2-amino-2-hydroxymethyl-1,3-
24 propanediol ('tris') in synthetic sea water, *Deep Sea Res. Part 1 Oceanogr. Res. Pap.*,
25 45 (9), 1541–1554, 1998. Dickson, A.G., Sabine, C. L. and Christian, J. R.: Guide to
26 best practices for ocean CO₂ measurements, PICES Special Publication 3, North Pacific
27 Marine Science Organization, Sidney, British Columbia, 2007.

28 Dickson, A.G.: An exact definition of total alkalinity and a procedure for the estimation of
29 alkalinity and total inorganic carbon from titration data, *Deep Sea Res., Part A*, 28,
30 1981.

- 1 Dickson, A.G.: pH scales and proton-transfer reactions in saline media such as sea water,
2 *Geochim. Cosmochim. Acta*, 48(11), 2299-2308, 1984.
- 3 Edman, M., and Omstedt, A.: Modeling the dissolved CO₂ system in the redox environment of
4 the Baltic Sea, *Limnol. Oceanogr.*, 58, 2013.
- 5 Emerson, S. R. and Hedges, J. I. (Eds.): *Chemical Oceanography and the Marine Carbon*
6 *Cycle*, Cambridge University Press, Cambridge, 470, 2008.
- 7 Gustafsson, E., Deutsch, B., Gustafsson, B.G., Humborg, C. and Morth, C.M.: Carbon cycling
8 in the Baltic Sea - the fate of allochthonous organic carbon and its impact on air-sea
9 CO₂ exchange, *J. Mar. Syst.*, 129, 289–302, 2014a.
- 10 Gustafsson, E., Omstedt, A., Gustafsson, B.G., The air-water CO₂ exchange of a coastal sea-A
11 sensitivity study on factors that influence the absorption and outgassing of CO₂ in the Baltic
12 Sea, *Journal of Geophysical Research: Oceans*, 120, 5342–5357, 2015.
- 13 Gustafsson, E., Wällstedt, T., Humborg, Ch., Mörth, C. M., and Gustafsson, B. G.: External
14 total alkalinity loads versus internal generation: The influence of nonriverine alkalinity
15 sources in the Baltic Sea, *Glob. Biogeochem. Cycles* 28, 1358-1370, 2014b.
- 16 Hammer, K., Schneider, B., Kuliński, K. and Schulz-Bull, D.: Acid-base properties of Baltic
17 Sea dissolved organic matter, *J. Mar. Syst.*, 173, 114-121, 2017.
- 18 Hammer, K., Schneider, B., Kuliński, K. and Schulz-Bull, D.E.: Precision and accuracy of
19 spectrophotometric pH measurements at environmental conditions in the Baltic Sea,
20 *Estuar. Coast. Shelf S.*, 146, 24-32, 2014.
- 21 HELCOM: Eutrophication in the Baltic Sea, *Balt. Sea Environ. Proc.* 115B, 2009.
- 22 Hernandez-Ayon, J. M., Zirino, A., Dickson, A. G., Camiro-Vargas, T. and Valenzuela-
23 Espinoza, E.: Estimating the contribution of organic bases from microalgae to the
24 titration alkalinity in coastal seawaters, *Limnol. Oceanogr. Methods*, 5, 225–232, 2007.
- 25 Hjalmarsson, S., Wesslander, K., Anderson, L. G., Omstedt, A., Perttila, M., and Mintrop, L.:
26 Distribution, long-term development and mass balance calculation of total alkalinity in
27 the Baltic Sea, *Cont. Shelf Res.*, 28, 593–601, 2008.
- 28 Hoikkala, L., Kortelainen, P., Soinne, H. and Kuosa, H.: Dissolved organic matter in the Baltic
29 Sea, *J. Mar. Syst.*, 142, 47–61, 2015.

- 1 Hunt, C. W., Salisbury, J. E., and Vandemark, D.: Contribution of non-carbonate anions to
2 total alkalinity and overestimation of pCO₂ in New England and New Brunswick rivers,
3 *Biogeosciences*, 8, 3069–3076, 2011.
- 4 Johansson, O. and Wedborg, M.: The Ammonia-Ammonium equilibrium in sea water at
5 temperatures between 5-25°C, *J. Solution Chemistry*, 9, 37-44, 1980.
- 6 Koeve W. and Oschlies, A.: Potential impact of DOM accumulation on fCO₂ and carbonate
7 ion computations in ocean acidification experiments, *Biogeosciences*, 9, 3787–3798,
8 2012.
- 9 Kremling, K.: Untersuchungen über die chemische Zusammensetzung des Meerwassers aus
10 der Ostsee II. Frühjahr 1967 – Frühjahr 1968. *Kiel Meeresforsch*, 26, 1-20, 1970.
- 11 Kremling, K.: Untersuchungen über die chemische Zusammensetzung des Meerwassers aus
12 der Ostsee III. Frühjahr 1969 – Herbst 1970. *Kiel Meeresforsch*, 27, 99-118, 1972.
- 13 Krumins V., Gehlen M., Arndt S., Van Cappellen P. and Regnier P.: Dissolved inorganic
14 carbon and alkalinity fluxes from coastal marine sediments: model estimates for
15 different shelf environments and sensitivity to global change, *Biogeosciences*, 10, 371-
16 398, 2013.
- 17 Kuliński, K. and Pempkowiak, J.: The carbon budget of the Baltic Sea, *Biogeosciences*, 8,
18 3219-3230, 2011.
- 19 Kuliński, K., Hammer, K., Schneider, B. and Schulz-Bull, D.: Remineralization of terrestrial
20 dissolved organic carbon in the Baltic Sea, *Mar. Chem.*, 181, 10-17, 2016.
- 21 Kuliński, K., Schneider, B., Hammer, K., Machulik, U. and Schulz-Bull, D.: The influence of
22 dissolved organic matter on the acid-base system of the Baltic Sea. *J. Marine Syst.*, 132,
23 106-115, 2014.
- 24 Kuliński, K., She, J. and Pempkowiak, J.: Short and medium term dynamics of the carbon
25 exchange between the Baltic Sea and the North Sea, *Cont. Shelf Res.*, 31, 15, 1611–
26 1619, 2011.
- 27 Kuznetsov, I. and Neumann T.: Simulation of carbon dynamics in the Baltic Sea with a 3D
28 model, *J. Marine Syst.*, 111–112, 167–174, 2013.
- 29 Lass, H.-U. and Matthäus, W.: General Oceanography of the Baltic Sea, in: *State and Evolution*
30 *of the Baltic Sea, 1952-2005: Detailed 50-Year Survey of Meteorology and Climate*,

- 1 Physics, Chemistry, Biology and Marine Environment, Feistel, R., Nausch, G. and
2 Wasmund, N., John Wiley & Sons, Inc., USA, 2008.
- 3 Le Quéré, C., Andrew, R. M., Canadell, J. G., Sitch, S., Korsbakken, J. I., Peters, G. P.,
4 Manning, A. C., Boden, T. A., Tans, P. P., Houghton, R. A., Keeling, R. F., Alin, S.,
5 Andrews, O. D., Anthoni, P., Barbero, L., Bopp, L., Chevallier, F., Chini, L. P., Ciais,
6 P., Currie, K., Delire, Ch., Doney, S. C., Friedlingstein, P., Gkritzalis, T., Harris, I.,
7 Hauck, J., Haverd, V., Hoppema, M., Goldewijk, K. K., Jain, A. K., Kato, E.,
8 Körtzinger, A., Landschützer, P., Lefèvre, N., Lenton, A., Lienert, S., Lombardozzi, D.,
9 Melton, J. R., Metzl, N., Millero, F., Monteiro, P. M. S., Munro, D. R., Nabel, J. E. M.
10 S., Nakaoka, S., O'Brien, K., Olsen, A., Omar, A. M., Ono, T., Pierrot, D., Poulter, B.,
11 Rödenbeck, Ch., Salisbury, J., Schuster, U., Schwinger, J., Séférian, R., Skjelvan, I.,
12 Stocker, B. D., Sutton, A. J., Takahashi, T., Tian, H., Tilbrook, B., van der Laan-Luijkx,
13 I. T., van der Werf, G. R., Viovy, N., Walker, A. P., Wiltshire, A. J., Zaehle S.: Global
14 Carbon Budget 2016, *Earth Syst. Sci. Data*, 8, 605-649, 2016.
- 15 Lee, K., Kim, T.W., Byrne, R.H., Millero, F.J., Feely, R.A. and Liu, Y.M.: The universal ratio
16 of boron to chlorinity for the North Pacific and North Atlantic oceans, *Geochim.*
17 *Cosmochim. Ac.*, 74, 1801–1811, 2010.
- 18 Lukawska-Matuszewska, K.: Contribution of non-carbonate inorganic and organic alkalinity
19 to total measured alkalinity in pore waters in marine sediments (Gulf of Gdansk, S-E
20 Baltic Sea), *Mar. Chem.*, 186, 211-220, 2016.
- 21 Millero, F.J., Graham, T.B., Huang, F, Bustos-Serrano, H. and Pierrot, L.D.: Dissociation
22 constants of carbonic acid in seawater as a function of salinity and temperature, *Mar.*
23 *Chem*, 100, 1-2, 80-94, 2006.
- 24 Millero, F.J., Plese, T., Fernandez, M.: The dissociation of hydrogen sulfide in seawater (Note),
25 *Limnol. Oceanogr.*, 33, 2, 269-274, 1988.
- 26 Millero, F.J.: Carbonate constants for estuarine waters, *Mar. Freshwater Res.*, 61, 139–142,
27 2010.
- 28 Mosley, L.M., Husheer, S.L.G. and Hunter, K.A., Spectrophotometric pH measurement in
29 estuaries using thymol blue and m-cresol purple, *Mar. Chem.*, 91, 175-186, 2004.
- 30 MSFD, Directive 2008/56/EC of the European Parliament and of the Council (Marine Strategy
31 Framework Directive), 2008.

- 1 Müller, J.D., Schneider, B. and Rehder, G.: Long-term alkalinity trends in the Baltic Sea and
2 their implications for CO₂-induced acidification, *Limnol. Oceanogr.*, 61, 1984-2002,
3 2016.
- 4 Nausch, G., Nehring, D. and Nagel K.: Nutrients Concentrations, Trends and Their Relation to
5 Eutrophication, in: *State and Evolution of the Baltic Sea, 1952-2005: Detailed 50-Year*
6 *Survey of Meteorology and Climate, Physics, Chemistry, Biology, and Marine*
7 *Environment*, Feistel, R., Nausch, G. and Wasmund, N., John Wiley & Sons, Inc., USA,
8 2008.
- 9 Nebbioso, A. and Piccolo, A.: Molecular characterization of dissolved organic matter (DOM):
10 a critical review, *Anal. Bioanal. Chem.*, 405, 109–124, 2013.
- 11 Omstedt, A., Edman, M., Anderson, L. G. and Laudon H.: Factors influencing the acid–base
12 (pH) balance in the Baltic Sea: a sensitivity analysis, *Tellus B* 62, 280-295, 2010.
- 13 Omstedt, A., Edman, M., Claremar, B., Frodin, P., Gustafsson, E., Humborg, C., Hagg, H.,
14 Morth, M., Rutgersson, A., Schurgers, G., Smith, B., Wallstedt T. and Yurova, A.:
15 Future changes in the Baltic Sea acid–base (pH) and oxygen balances, *Tellus B* 64,
16 19586, 2012.
- 17 Omstedt, A., Edman, M., Claremar, B., Rutgersson, A., Modelling the contributions to marine
18 acidification from deposited SO_x, NO_x, and NH_x in the Baltic Sea: Past and present
19 situations, *Continental Shelf Research*, 111, 234–249, 2015.
- 20 Omstedt, A., Gustafsson, E. and Wesslander, K.: Modelling the uptake and release of carbon
21 dioxide in the Baltic Sea surface water, *Cont. Shelf Res.*, 29, 870-885, 2009.
- 22 Orr, J. C., Epitalon, J.-M. and Gattuso, J.-P.: Comparison of ten packages that compute ocean
23 carbonate chemistry, *Biogeosciences*, 12, 1483–1510, 2015.
- 24 Redfield, A. C., Ketchum, B. H. and Richards, F. A.: The influence of organisms on the
25 composition of sea water, in: *The Sea*, Hill, M. N. (Ed.), Interscience, New York, USA,
26 26–77, 1963.
- 27 Reed, D.C., Gustafsson, B.G., Slomp, C.P., Shelf-to-basin iron shuttling enhances vivianite
28 formation in deep Baltic Sea sediments, *Earth and Planetary Science Letters*, 434, 241–
29 251, 2016.

- 1 Riebesell, U., Fabry, V. J., Hansson, L. and Gattuso J.-P.: Guide to best practices for ocean
2 acidification research and data reporting, 260, Luxembourg, Publications Office of the
3 European Union, 2010.
- 4 Roy, R.N., Roy, L.N., Vogel, K.M., Porter-Moore, C., Pearson, T., Good, C.E., Millero, F.J.
5 and Campbell, D.M.: The dissociation constants of carbonic acid in seawater at
6 salinities 5 to 45 and temperatures 0 to 45°C: *Mar. Chem.*, 44, 2–4, 249–267, 1993.
- 7 Schneider, B., Eilola, K., Lukkari, K., Muller-Karulis, B. and Neumann T.: Environmental
8 Impacts – Marine Biogeochemistry, in: BACC II Author Team, Second Assessment
9 of Climate Change for the Baltic Sea Basin, Springer, Heidelberg, 337-361, 2015.
- 10 Schneider, B., Gülzow, W., Sadkowiak, B. and Rehder, G.: Detecting sinks and sources of CO₂
11 and CH₄ by ferrybox-based measurements in the Baltic Sea: Three case studies, *J.*
12 *Marine Syst.*, 140, 13-25, 2014.
- 13 Schneider, B.: The CO₂ system of the Baltic Sea : biogeochemical control and impact of
14 anthropogenic CO₂, in: *Global change and Baltic coastal zones*. Ed. by G. Schernewski,
15 J. Hofstede and T. Neumann. Dordrecht: Springer Science+Business Media (Coastal
16 research library): 33-49, 2011.
- 17 Schulz, H.D. and Zabel, M., *Marine geochemistry*, Springer-Verlag Berlin Heidelberg, 574,
18 2006.
- 19 Szymczycha, B., Maciejewska, A., Winogradow, A. and Pempkowiak, J.: Could submarine
20 groundwater discharge be a significant carbon source to the southern Baltic Sea?,
21 *Oceanologia*, 56, 2, 327-347, 2014.
- 22 Thomas H., Schiettecatte L.-S., Suykens K., Koné Y. J. M., Shadwick E. H., Prowe A. E. F.,
23 Bozec Y., de Baar H. J. W. and Borges A. V.: Enhanced ocean carbon storage from
24 anaerobic alkalinity generation in coastal sediments, *Biogeosciences*, 6, 267–274, 2009.
- 25 Tyrrell, T., Schneider, B., Charalampopoulou, A. and Riebesel, U.: Coccolithophores and
26 calcite saturation state in the Baltic and Black Seas, *Biogeosciences*, 5, 485–494, 2008.
- 27 Ulfsbo, A., Hulth S. and Anderson, L.G.: pH and biogeochemical processes in the Gotland
28 Basin of the Baltic Sea, *Mar. Chem.*, 127, 20-30, 2011.
- 29 Ulfsbo, A., Kuliński, K., Anderson, L.G. and Turner, D.R.: Modelling organic alkalinity in the
30 Baltic Sea using a Humic-Pitzer approach, *Mar. Chem.*, 168: 18-26, 2015.

- 1 Uppström, L.R.: The boron/chlorinity ratio of deep-sea water from the Pacific Ocean, Deep-
2 Sea Res., 21, 161–162, 1974.
- 3 Weiss, R.F.: Carbon dioxide in water and seawater: the solubility of a non-ideal gas, Mar.
4 Chem., 2, 203-215, 1974.
- 5 Winsor, P., Rodhe, J., Omstedt, A., 2001. Baltic Sea ocean climate: an analysis of 100 yr of
6 hydrographic data with focus on the freshwater budget. Climate Research 18, 5–15.
- 7 Wolf-Gladrow, D.A., Zeebe, R.E., Klaas, C., Körtzinger, A., Dickson, A.G., Total alkalinity:
8 The explicit conservative expression and its application to biogeochemical processes,
9 Marine Chemistry, 106, 287–300, 2007.
- 10 Zeebe, R.E., History of Seawater Carbonate Chemistry, Atmospheric CO₂, and Ocean
11 Acidification, Annual Review of Earth and Planetary Sciences, 40, 141–165, 2012.