

Dear Dr. Kleidon:

Please find attached a revision of the manuscript entitled “Recent changes of relative humidity: regional connection with land and ocean processes” to be considered for publication in Earth System Dynamics. In the revised manuscript, we have addressed all minor comments raised by the third reviewer; the existing typos have been removed, and the length of the manuscript has been noticeably reduced. We have removed 8 pages of text and three figures regarding the previous version of the manuscript.

We look forward to hearing from you at your earliest convenience, and should you have any questions please feel free to contact us.

Sincerely,

Sergio M. Vicente-Serrano and coauthors

1 | **Recent changes of relative humidity: regional ~~connection~~connections with land and**
2 | **ocean processes**

3 Sergio M. Vicente-Serrano¹, Raquel Nieto², Luis Gimeno², Cesar Azorin-Molina³,
4 Anita Drumond², Ahmed El Kenawy^{1,4}, Fernando Dominguez-Castro¹, Miquel Tomas-
5 Burguera⁵, Marina Peña-Gallardo¹

6 ¹ *Instituto Pirenaico de Ecología, Consejo Superior de Investigaciones Científicas*
7 *(IPE-CSIC), Zaragoza, Spain;* ² *Environmental Physics Laboratory, Universidade de*
8 *Vigo, Ourense, Spain.* ³ *Regional Climate Group, Department of Earth Sciences,*
9 *University of Gothenburg, Sweden.* ⁴ *Department of Geography, Mansoura University,*
10 *Mansoura, Egypt;* ⁵ *Estación Experimental Aula Dei, Consejo Superior de*
11 *Investigaciones Científicas (EEAD-CSIC), Zaragoza, Spain;*

12
13 * Corresponding author: svicen@ipe.csic.es
14

15 **Abstract.** We analyzed changes in surface relative humidity (RH) at the global scale
16 from 1979 to 2014 using both observations and ERA-Interim dataset. We compared the
17 variability and trends of RH with those of land evapotranspiration and ocean
18 evaporation in moisture source areas across a range of selected regions worldwide. The
19 sources of moisture for each particular region were identified by integrating different
20 observational data and model outputs into a lagrangian approach. The aim was to
21 account for the possible role of changes in air temperature over land, in comparison to
22 sea surface temperature (SST), but also the role of land evapotranspiration and the
23 ocean evaporation on RH variability. The results demonstrate that the patterns of the
24 observed trends in RH at the global scale cannot be linked to a particular individual
25 physical mechanism. Our results also stress that the different hypotheses that may
26 explain the decrease in RH under a global warming scenario could act together to
27 explain recent RH trends. Albeit with ~~the~~ uncertainty in establishing a direct causality
28 between RH trends and the different empirical moisture sources, we found that the
29 observed decrease in RH in some regions (~~e.g.~~) can be linked to lower water supply
30 from land evapotranspiration. In contrast, the empirical relationships also suggest that
31 RH trends in other target regions (~~e.g.~~) are mainly explained by the dynamic and
32 thermodynamic mechanisms related to the moisture supply from the oceanic source
33 regions. Overall, while this work gives insights ~~on~~into the ~~connection of~~connections
34 ~~between~~ RH trends ~~with~~and oceanic and continental processes at the global scale,
35 further investigation ~~is still desired~~ to assess the contribution of both dynamic and
36 thermodynamic factors to the evolution of RH over continental regions ~~at more detailed~~
37 ~~spatial scales is desired.~~

38
39 **Key-words:** Relative humidity; Evaporation; Evapotranspiration; Moisture; Trends;
40 Oceans.

41

42 **1. Introduction**

43 Relative Humidity (RH) is a key meteorological parameter that determines the
44 aerodynamic component of the atmospheric evaporative demand (AED) (Wang and
45 Dickinson, 2012; McVicar et al., 2012a). As such, changes in RH may impact
46 significantly the evolution of the AED (Vicente-Serrano et al., 2014a), with particular
47 implications for the intensity of the hydrological cycle (Sherwood, 2010), climate
48 aridity (Sherwood and Fu, 2014) as well as severity of drought events (Rebetez et al.,
49 2006; Marengo et al., 2008).

50 In a changing climate, temperature rise, as suggested by different climate scenarios,
51 may impact the atmospheric humidity. ~~According to the Clausius-Clapeyron (CC)~~
52 ~~relationship, a temperature rise of 1 °C is sufficient to increase the equilibrium amount~~
53 ~~of water vapor of the air by roughly 7%. Given the unlimited water availability in the~~
54 ~~oceans as well as the projected temperature rise, water vapor content is expected to~~
55 ~~increase, at least in the oceanic areas, in order to maintain RH constant in the future.~~

56 Particularly, there is empirical evidence on the increase in the water vapor content at
57 both the surface and upper tropospheric levels (Trenberth et al., 2005). In this context,
58 numerous studies have supported the constant RH scenario under global warming
59 conditions (e.g. Dai, 2006; Lorenz and Deweaver 2007; Willett et al., 2008; McCarthy
60 et al., 2009; Ferraro et al., 2015). In contrast, other studies supported the non-stationary
61 behavior of RH, not only in continental areas located far from oceanic humidity (e.g.
62 Pierce et al., 2013), but also in humid regions (e.g. Van Wijngaarden and Vincent,
63 2004). Assuming the stationary behavior of RH, the influence of RH on AED may- be
64 constrained, given that any possible change in AED would be mostly determined by
65 changes in other aerodynamic variables (e.g. air temperature and wind speed) (McVicar

66 et al., 2012a and b) or by changes in cloudiness and solar radiation (Roderick and
67 Farquhar, 2002; Fan and Thomas, 2013). However, a range of studies have supported
68 the non-stationary behavior of RH under global warming, giving insights on significant
69 changes in RH over the past decades. A representative example is Simmons et al.
70 (2010) who compared gridded observational and reanalysis RH data, suggesting a clear
71 dominant negative trend in RH over the Northern Hemisphere since 2000. Also, based
72 on a newly developed homogeneous gridded database that employed the most available
73 stations from the telecommunication system of the WMO, Willett et al. (2014) found
74 significant negative changes in RH, with strong spatial variability, at the global scale.
75 This global pattern was also confirmed at the regional scale, but with different signs of
76 change, including both negative (e.g. Vincent et al., 2007; Vicente-Serrano ~~et al.~~,
77 2014b; 2016; Zongxing et al., 2014) and positive trends (e.g. Shenbin, 2006; Jhajharia
78 et al., 2009; Hosseinzadeh Talaei et al., 2012).

79 There are different hypotheses that explain the non-stationary evolution of RH under
80 global warming conditions. One of these hypotheses is related to the slower warming of
81 oceans in comparison to continental areas (Lambert and Chiang, 2007; Joshi et al.,
82 2008). In particular, specific humidity of air advected from oceans to continents
83 increases more slowly than saturation specific humidity increases over land (Rowell and
84 Jones 2006; Fasullo 2010). This would decrease RH over continental areas, inducing an
85 increase in AED and aridity conditions (Sherwood and Fu, 2014). Some studies
86 employed global climate models (GCMs) to support this hypothesis under future
87 warming conditions (e.g. Joshi et al., 2008; O’Gorman and Muller, 2010; Byrne and
88 O’Gorman, 2013). ~~However, empirical studies that support this hypothesis using~~
89 ~~observational data are unavailable. Moreover, the observed decrease in RH over some~~

90 ~~coastal areas, which are adjacent to their sources of moisture, adds further uncertainty to~~
91 ~~this hypothesis (Vicente-Serrano et al., 2014b and 2016; Willet et al., 2014).~~

92 Another hypothesis to explain the non-stationary evolution of RH is associated with
93 land-atmosphere feedback processes. Different studies indicated that atmospheric
94 moisture and precipitation are strongly linked to moisture recycling in different regions
95 of the world (e.g. Rodell et al., 2015). Thus, evapotranspiration may contribute largely
96 to water vapor content and precipitation over land (Stohl and James, 2005; Bosilovich
97 and Chern, 2006; Trenberth et al., 2007; Dirmeyer et al., 2009; van der Ent et al., 2010).
98 Land-atmospheric feedbacks may also have marked influence on atmospheric humidity
99 (Seneviratne et al., 2006); given that soil drying can suppress evapotranspiration, reduce
100 RH and thus reinforce AED. All these processes would again reinforce soil drying
101 (Seneviratne et al., 2002; Berg et al., 2016).

102 Indeed, it is very difficult to determine which hypothesis can provide an understanding
103 of the observed RH trends at the global scale. Probably, the two hypotheses combined
104 together can be responsible for the observed RH trends in some regions of the world
105 (Rowell and Jones, 2006). In addition to the aforementioned hypotheses, some dynamic
106 forces, which are associated with atmospheric circulation processes, can explain the
107 non-stationary behavior of RH worldwide (e.g. Goessling and Reick, 2011). However,
108 defining the relative importance of these physical processes in different world regions is
109 quite challenging (Zhang et al., 2013; Laua and Kim, 2015).

110 The objective of this study is to compare the recent variability and trends of RH with
111 changes in the two types of fluxes that affect RH: i) vertical fluxes that were assessed
112 using land evapotranspiration and precipitation and ii) advection that was quantified
113 using oceanic evaporation from moisture source areas. The novelty of this work stems
114 from the notion that although different studies have already employed GCM's and

115 different scenarios to explain the possible mechanisms behind RH changes under
116 warming conditions, we introduce a new empirical approach that employs different
117 observational data sets, reanalysis fields and a lagrangian-based approach, not only for
118 identifying the continental and oceanic moisture areas for different target regions, but
119 also for exploring the relevance of the existing hypothesis to assess the magnitude, sign
120 and spatial patterns of RH trends in the past decades at the global scale.

121

122 **2. Data and methods**

123 2.1. Dataset description

124 *2.1.1. Observational RH dataset*

125 We employed the monthly RH HadISDH dataset, available through
126 <http://www.metoffice.gov.uk/hadobs/hadisdh> (Willet et al., 2014). ~~This dataset~~
127 ~~represents the most complete and accurate global dataset for RH, including~~
128 ~~observational data from a wide range of stations worldwide (Willet et al., W~~
129 ~~selected2014). Given that HadISDH includes some series with data gaps, our decision~~
130 ~~was to choose~~ only those series with no more than 20% of missing values over the
131 period 1979-2014. In order to fill these gaps, we created a standardized regional series
132 for each station using the most correlated series with each target series. ~~While this~~
133 ~~procedure maintains the temporal variance of the original data, it provides a low biased~~
134 ~~estimation of the missing values.~~ In order to avoid biases, mostly originated from
135 differences in the distribution parameters (mean and variance) between the candidate
136 and the objective data series, a bias correction was applied to the candidate data. ~~Thus,~~
137 ~~normal distribution was used for bias correction of RH.~~ The data of the candidate series
138 were re-scaled to match the statistical distribution of the observed series to be filled,
139 based on the overlapping period between them. Overall, a final dataset of 3462

165 | _____ (6), where)

166 | where

167 | _____ (7)

168 | _____ (8)

169 | and T is the 2 meters air temperature in °C

170 | e_s is obtained by substituting T_d by T .

171 |

172 | 2.1.3. Land precipitation and land air temperature

173 | We employed the gridded land precipitation and surface air temperature data (TS
174 | v.3.23), provided by the Climate Research Unit (CRU: UK), at a 0.5° spatial interval for
175 | the period 1979-2014 (Harris et al., 2014, 2014). ~~This product was developed using a~~
176 | ~~relatively high number of observational sites, which guarantees a robust representation~~
177 | ~~of climatic conditions across worldwide regions. Importantly, this product has been~~
178 | ~~carefully tested for potential data inhomogeneities as well as anomalous data.~~

179 |

180 | 2.1.4. Sea Surface Temperature (SST)

181 | We used the monthly SST data (HadSST3), compiled by the Hadley Centre for the
182 | common period 1979-2014 (<http://www.metoffice.gov.uk/hadobs/hadsst3/>). This dataset
183 | is provided at a 0.5° grid interval (Kennedy et al., 2011a and b).

184 |

185 | 2.1.5. Ocean evaporation and continental evapotranspiration data

186 | To quantify the temporal variability and trends of land evapotranspiration and oceanic
187 | evaporation, we employed two different datasets. First, the oceanic evaporation was
188 | quantified using the Objectively Analyzed air-sea Fluxes (OAFLUX) product (Yu et al.,
189 | 2008) from 1979 to 2014, which was used to analyze recent variability and changes in

190 evaporation from global oceans (Yu, 2007). To account for land evapotranspiration, we
191 employed the Global Land Evaporation Amsterdam Model (GLEAM) (Version 3.0a)
192 (<http://www.gleam.eu/>) (Miralles et al., 2011) from 1980 to 2014. This data set has been
193 widely validated using in situ measurements of surface soil moisture and evaporation
194 across the globe (Martens et al., 2016).

195

196 2.2. Methods

197 *2.2.1. Relative Humidity (RH) trends*

198 We assessed the seasonal (boreal cold season: October-March; boreal warm season:
199 April-September) and annual trends of RH for 1979-2014 using two different global
200 datasets (HadISDH and ERA-Interim). To quantify the magnitude of change in RH, we
201 used a linear regression analysis between the series of time (independent variable) and
202 RH series (dependent variable). ~~The slope of the regression indicates the amount of~~
203 ~~change (per year), with higher slope values indicating greater changes.~~ To assess the
204 statistical significance of the detectable changes, we applied the nonparametric Mann-
205 Kendall statistic, ~~which measures the degree to which a trend is consistently increasing~~
206 ~~or decreasing~~ (Zhang et al., 2001). ~~To~~ to account for any possible influence of serial
207 autocorrelation on the robustness of the defined trends, ~~we applied the modified Mann-~~
208 ~~Kendall trend test, which returns the corrected p-values after accounting for temporal~~
209 ~~pseudoreplication in RH series~~ (Hamed and Rao, 1998; Yue and Wang, 2004). The
210 statistical significance of the trend was tested at the 95% confidence level ($p < 0.05$).

211 Following the trend analysis results, we selected those regions that showed a high
212 agreement between HadISDH and ERA-Interim datasets in terms of the sign and
213 magnitude of RH changes. Nevertheless, we also extended our selection to some other
214 regions, with low station density in the HadISDH dataset. This decision was simply
215 motivated by the consistent changes found over these regions, as suggested by the ERA-

216 Interim dataset. For all the defined regions, we identified the oceanic and continental
217 moisture sources by means of the FLEXPART lagrangian model.

218
219 *2.2.2. Identification of continental and oceanic moisture sources*

220 We used the FLEXPART V9.0 particle dispersion model fed with the ERA-Interim
221 reanalysis data. According to this model, the atmosphere is divided homogeneously into
222 three-dimensional finite elements (hereafter “particles”); each represents a fraction of
223 the total atmospheric mass (Stohl and James, 2004). These particles may be advected
224 backward or forward in time using three-dimensional wind taken from the ERA-Interim
225 data every time step, with superimposed stochastic turbulent and convective motions.

226 The rates of increase (e) and decrease (p) of moisture (e-p) along the trajectory of each
227 particle were calculated via changes in the specific ~~moisture~~humidity (q) with time (e-p
228 = mdq/dt), where m is the mass of the particle. ~~Similar to the wind field, q is also taken~~
229 ~~from the meteorological data. FLEXPART allows identifying the particles affecting a~~
230 ~~particular region using information about the trajectories of these selected particles.~~ A
231 description of this methodology is detailed in Stohl and James (2004).

232 The FLEXPART dataset used in this study was provided by a global experiment in
233 which the entire global atmosphere was divided into approximately 2.0 million
234 “particles”. The tracks were computed using the ERA-Interim reanalysis data at 6 h
235 intervals, at a 1° horizontal resolution and at a vertical resolution of 60 levels from 0.1
236 to 1000 hPa. For each particular target region, all the particles were tracked backward in
237 time, and its position and specific humidity (q) were recorded every 6 h. ~~With this~~
238 ~~methodology, the evaporative sources and sink regions for the particles reaching the~~
239 ~~target region can be identified.~~ All areas where the particles gained humidity (E -P > 0)

Con formato: Espacio Después: 0 pto, Interlineado: Doble

240 along their trajectories towards the target region can be considered as “sources of
241 moisture”. In contrast, all areas with lost humidity ($E - P < 0$) are considered as “sinks”.

242 ~~A typical period used to track the particles backward in time is 10 days that is the~~
243 ~~average residence time of water vapor in the global atmosphere (Numaguti, 1999).~~
244 ~~However, we~~ We followed the methodology of Miralles et al (2016), where an optimal
245 lifetime of vapor in the atmosphere was calculated to reproduce the sources of moisture.
246 ~~As such, three steps were carried out in this order: i) all the particles that leave each~~
247 ~~target region were tracked back during 15 days and the “initial sources” at annual scale~~
248 ~~were defined as those areas with positive (E-P) values, ii) from these “initial sources”,~~
249 ~~all the particles were forward tracked during 1 to 15 days individually, and (E-P)<0 was~~
250 ~~calculated for these lifetime periods to estimate the precipitation contribution over the~~
251 ~~target region, iii) the optimal lifetime selected for each region was chosen according to~~
252 ~~the minimum absolute difference between the FLEXPART simulated precipitation and~~
253 ~~the CRU TS v.3.23 for each region, iv) and finally the backward tracking was~~
254 ~~recalculated during these optimal lifetimes.~~

255 We and we defined the climatological spatial extent of each source region
256 corresponding to a particular target region by applying a 95th percentile criterion
257 computed for the annual and seasonal (boreal summer and winter) positive (E-P) field
258 (Vazquez et al., 2016).— Then, for each year of the period, we estimated the total
259 moisture supply from each source region.

260 Also from FLEXPART simulations, we obtained the fractions of moisture from the
261 continental and oceanic sources annually and for each cold and warm season. The
262 purpose was to compare with the results obtained on the role of the land
263 evapotranspiration and ocean evaporation of RH variability and trends.

264

265 *2.2.3. Relationship between RH and the selected land/oceanic climate variables*

266 Based on defining the spatial extent of each moisture source region, we calculated
267 annual, warm and cold season regional series for ocean evaporation and land
268 evapotranspiration using the OAFLUX and GLEAM datasets, respectively. The
269 regional series of ocean evaporation and land evapotranspiration were created using a
270 weighted average based on the seasonal/annual fields of $(E-P) > 0$ (~~Section 2.2.2~~). This
271 approach allows creating a time series that better represents the interannual variability
272 of ocean evaporation and land evapotranspiration in the source(s) of moisture for each
273 defined region. Following the same approach, we also calculated the regional series of
274 SST corresponding to each oceanic moisture source region. Likewise, we calculated the
275 regional series of land precipitation and air temperature for each target region using
276 CRU TS v.3.23 dataset, and the ratio between air temperature in the target region and
277 SST in the source region.

278 For each target region, we related the regional series of seasonal and annual RH to the
279 corresponding regional times series of all aforementioned climatic variables. However,
280 to limit the possible influence of the trends presented in the data itself on the computed
281 correlations, we de-trended the series of the climate variables prior to calculating the
282 correlation. We also assessed changes in the regional series of the different variables;
283 their statistical significance was tested by means of the modified Mann-Kendall test at
284 the 95% level. For each target region, we summarized the results of the magnitude of
285 change in RH as well as other investigated variables at the seasonal and annual scales.
286 However, to facilitate the comparison among the different variables and the target
287 regions worldwide, we transformed the amount of change of each variable to
288 percentages.

289 Finally, we also computed the association between RH and land evapotranspiration at
290 the annual and seasonal scales using the available gridded evapotranspiration series.
291 While a pixel-to-pixel comparison does not produce a reliable assessment of the
292 possible contribution of land evapotranspiration to RH changes, given that the source of
293 moisture can apparently be far from the target region, we still believe that this
294 association can give insights on the possible relationship between land
295 evapotranspiration on RH changes.

296

297 **3. Results**

298 **3.1. Trends in Relative Humidity**

299 ~~Figure 1 shows the average seasonal and annual RH and the Vertically Integrated~~
300 ~~Moisture Flux (VIMF), which can be used to estimate regions~~There are regional
301 differences where the precipitation dominates (negative values) over the evaporation
302 (positive values), from the ERA-Interim dataset (Suppl. Fig. 1). Fig. 1 ~~RH shows higher~~
303 ~~average values over equatorial regions, Southeast Asia and the North Eurasia region.~~
304 ~~The lower values are recorded over tropical regions, mainly in the North Hemisphere.~~
305 ~~Spatial differences between the cold and warm seasons are very low. The annual pattern~~
306 ~~of the VIMF over continents shows that precipitation exceeds evaporation over the~~
307 ~~Intertropical Convergence Zone, Southeast Asia and the islands between the Pacific and~~
308 ~~Indian Oceans (Maritime continent), a great part of South America, Central America,~~
309 ~~Central Africa, and northward to 40°N in the Northern Hemisphere. Evaporation is~~
310 ~~higher than precipitation over the main area of Australia, the Pacific coast of North~~
311 ~~America, Northeast Brazil, areas around the Mediterranean Sea, eastern coast of Africa~~
312 ~~and southwest Asia. Seasonally it is evident the poleward movement of the ITCZ during~~
313 ~~the hemispheric summer, and the change of the pattern over North America and~~
314 ~~Eurasian continent.~~

315 | ~~Figure 21~~ summarizes the magnitude of change in RH for the boreal cold and warm
316 | seasons and at the annual scale, ~~calculated using the annual and seasonal (boreal~~
317 | ~~summer and winter)~~ for the period between 1979 and 2014. For HadISDH, it is noted
318 | that the available RH stations is unevenly distributed over the globe, with higher density
319 | in the mid-latitudes of the Northern Hemisphere. Nevertheless, the available stations
320 | show coherent and homogeneous spatial patterns of RH changes (~~Supplementary Figure~~
321 | ~~1). In the boreal cold season, the most marked decrease was observed in the Southwest~~
322 | ~~and areas of Northeast North America, central Argentina, the Fertile Crescent region in~~
323 | ~~western Asia, Kazakhstan, as well as in the eastern China and the Korea Peninsula. On~~
324 | ~~the other hand, the dominant RH increase was recorded in larger areas, including most~~
325 | ~~of Canada (mostly in the Labrador Peninsula), and large areas of North and central~~
326 | ~~Europe and India. While the density of complete and homogeneous RH series is low,~~
327 | ~~we found a dominant positive trend across the western Sahel and South Africa.~~~~Suppl.~~
328 | ~~Fig. 2).~~ The ERA-Interim dataset showed magnitudes of change close to those
329 | suggested by HadISDH. In addition, the ERA-Interim also provides information on RH
330 | changes in regions with low density of RH observations (e.g. ~~EastEastern~~ Amazonia,
331 | ~~eastEastern~~ Sahel and Iran), suggesting a dominant RH decrease across these regions.
332 | For the boreal warm season, a clear tendency towards a reduction in RH was observed
333 | in vast regions of the world, including (mostly the Iberian Peninsula, France, Italy,
334 | Turkey and ~~Morocco~~Morocco), Eastern Europe, and western part of Russia. ~~Based on~~
335 | ~~the available stations across central Asia, we also found a general reduction of RH; a~~
336 | ~~similar pattern was also observed in East Asia, including Mongolia, east China, north~~
337 | ~~Indonesia, South Japan and Korea.~~ This reduction was also noted South America, with a
338 | general homogeneous pattern over Peru, Bolivia and a strong decrease over central
339 | Argentina. On the other hand, the positive evolution of RH observed during the cold

340 season across Canada and Scandinavia was reinforced during the boreal warm season.

341 ~~In the Western Sahel and India, we found an upward trend of RH.~~ The ERA-Interim

342 also revealed a strong RH decrease over the whole Amazonian region and the ~~West~~

343 ~~Sahel, while a marked increase dominated over the Andean region between Colombia,~~

344 ~~Ecuador and North Peru. In Australia, the spatial patterns were more complex than~~

345 ~~those obtained using the available observatories.~~ Western Sahel. A wide range of these

346 regions exhibited statistically significant trends from 1979 to 2014 (Suppl. Fig. 3).

347 ~~The HadISDH dataset suggests a general decrease of RH over Southwest North~~

348 ~~America, Argentina, central Asia, Turkey, Mongolia and China, with a particular~~

349 ~~reduction over the Eastern Sahel, Iran, Mongolia and the eastern Asia. On the other~~

350 ~~hand, a dominant positive trend was observed across Canada, areas of North Southern~~

351 ~~America, the Western Sahel, South Africa (Namibia and Botswana), some areas of~~

352 ~~Kenia, India and the majority of Australia. A wide range of these regions exhibited~~

353 ~~statistically significant trends from 1979 to 2014. (Supplementary Figure 2). A~~

354 ~~statistically significant negative trend was observed at the seasonal and annual scales,~~

355 ~~not only in most of Southern America and Northern America, but in large regions of~~

356 ~~Africa, South Europe, central and East Asia as well. On the other hand, areas of~~

357 ~~complex topography in the Northern Hemisphere, Australia, India, Northern South~~

358 ~~America and Africa showed positive trends.~~

359 Albeit with these complex spatial patterns of RH changes, there is a global dominant

360 negative trend (~~Figure 3~~Fig. 2). This pattern was observed using both the HadISDH and

361 the ERA-Interim datasets, although there is marked spatial bias in data availability of

362 the HadISDH. ~~Figure 4 illustrates the relationship~~The relationships between the

363 magnitudes of change in RH, as suggested by the HadISDH dataset versus the ERA-

364 Interim dataset. ~~At the seasonal and annual scales, there is,~~ show a relatively high

365 correlation (mostly above 0.55). Given this high consistency between the HadISDH and
366 the ERA-Interim datasets in terms of both the magnitude and sign of change in RH
367 (~~Figures 2 and 3~~) and also in the interannual variations (~~Supplementary Figures 3~~
368 ~~and Suppl. Fig.s 4 to 6~~), we decided to restrict our subsequent analysis to the ERA-
369 Interim dataset, recalling its denser global coverage compared to the HadISDH.

370 As RH is mostly dependent on changes in specific humidity (q), there is a dominant
371 high correlation between the interannual variability of RH and q (~~Supplementary Figure~~
372 ~~5~~ ~~Suppl. Fig. 7~~). In accordance, the magnitude of observed change in these two
373 variables showed a strong agreement for 1979-2014. ~~Figure 5~~ ~~Fig. 3~~ summarizes the
374 magnitude of change in specific humidity (q) as well as changes in specific humidity
375 necessary to maintain RH constant as recorded in 1979. Specific humidity showed the
376 strongest decrease in Southwest North America, the Amazonian region, Southern South
377 America and the Sahel regions: a spatial pattern that is similar to RH pattern. Given the
378 evolution of air temperature between for 1979-2014, these regions exhibited a deficit of
379 water vapor on the order of -2 g/kg^{-1} in order to maintain RH constant.

380

381 **3.2. Spatial patterns of the dependency between RH and climate variables**

382 ~~Based on the high agreement between the HadISDH and the ERA-Interim datasets in~~
383 ~~reproducing consistent seasonal and annual trends in RH, we~~ We selected a range of
384 regions (N=14) worldwide (~~Figure 6~~ ~~Fig. 4~~). For these selected regions, we assessed the
385 connection between RH and some relevant climatic variables for the period 1979-2014.
386 In addition, we defined the oceanic and continental sources of moisture corresponding
387 to these regions using the FLEXPART model. We assessed the optimal lifetime for each
388 region: during 4 days in back for regions 1-5 and 7-11, during 5 days for regions 6, 12-
389 13, and during 7 days for region 5 (~~see section 2.2~~). ~~Figures. Figs.~~ 7-9 show some

390 examples of the dependency between RH and different climate variables at the annual
391 scale. Results for all regions at the seasonal and annual scales are presented in
392 ~~supplementary~~ Suppl. materials.

393

394 **3.2.1. Western Sahel**

395 ~~Figure 7~~ Fig. 5 (top) illustrates RH trends in the Western Sahel using the HadISDH and
396 ERA-Interim datasets. We also showed the distribution of the average annual moisture
397 sources (E-P in mm) over this region for 1979-2014. As illustrated, the atmospheric
398 moisture is mostly coming from the ~~western~~ Western Sahel region itself, in addition to
399 some oceanic sources located in the central eastern Atlantic Ocean. At the seasonal
400 scale, there are some differences in the location and the intensity of the moisture
401 sources, with more oceanic contribution during the boreal warm season. However, in
402 both cases, the continental moisture seems to be the key source of humidity in the
403 region (~~Suppl. Figures 21 and 35~~). ~~In other areas, e.g. the Western European region~~
404 ~~(Suppl. Figures 17 and 31), we observed marked differences in the location and the~~
405 ~~intensity of humidity sources between the boreal cold and warm seasons. Figure 7~~ Figs
406 23 and 37). Fig. 5 (central) shows different scatter plots summarizing the relationships
407 between the de-trended annual series of RH and those of relevant climate variables (e.g.
408 precipitation, air temperature and SST). ~~As illustrated, the~~ The interannual variability of
409 RH in the region is correlated to changes in the total annual precipitation and the total
410 annual land evapotranspiration in the continental source region. Specifically, the
411 correlation between the de-trended annual RH and precipitation and land
412 evapotranspiration is generally above 0.8 ($p < 0.05$). In contrast, RH shows negative
413 correlations with air temperature and SST ratio over the oceanic source. While the
414 correlation is statistically insignificant ($p > 0.05$), it suggests that higher differences

415 between air temperature and SST reinforce lower annual RH. At the seasonal scale, we
416 found similar patterns (~~Supplementary Figs. 21 and 35~~), ~~with RH being highly~~
417 ~~correlated with land evapotranspiration during the boreal cold and warm seasons. Suppl.~~
418 ~~Figs. 23 and 37~~). Nevertheless, in the warm season, a significant negative correlation
419 with air temperature and SST ratio was observed. This pattern concurs with the
420 significant increase in specific humidity (q) for 1979-2014; this is probably related to
421 the high increase in land evapotranspiration (19.5%, $p < 0.05$).

422

423 **3.2.2. La Plata region**

424 ~~Figure 8~~Fig. 6 summarizes the corresponding results, but for ~~the~~ La Plata region (South
425 America). Results indicate a general decrease in RH at the annual and seasonal scales
426 using both the HadISDH observational data and the ERA-Interim dataset. As depicted,
427 the main humidity sources are located in the same region, combined with some other
428 continental neighbor areas over South America. A similar finding was also observed on
429 the seasonal scale (~~Supplementary Suppl.~~ Figs. ~~2527~~ and ~~3941~~). Similar to the
430 ~~West Western~~ Sahel region, we found a significant association between the interannual
431 variations of RH and precipitation and the land evapotranspiration in the continental
432 source region. Similarly, we did not find any significant correlation between RH
433 changes and the interannual variability of the oceanic evaporation in the oceanic source
434 region as well as the ratio between air temperature in the continental target region and
435 SST in the oceanic source region. Again, we found a negative correlation between RH
436 and air temperature/SST ratio, though being statistically insignificant at the annual scale
437 ($p > 0.05$). In ~~the~~ La Plata region, we noted a strong decrease in RH (-6.21%/decade) for
438 1979-2014, which agrees well with the strong decrease in absolute humidity. ~~This~~

439 ~~region is strongly impacted by continental atmospheric moisture sources, with a general~~
440 ~~decrease in precipitation and land evapotranspiration during the analyzed period.~~

441

442 **3.2.3. Southwest North America**

443 Results for Southwest North America are also illustrated in ~~Figure 9~~Fig. 7. In
444 accordance with both previous studied examples (~~West~~Western Sahel and La Plata), this
445 region also exhibited a strong and positive relationship between the interannual
446 variability of RH and precipitation and land evapotranspiration. This pattern was also
447 recorded for the boreal warm and cold seasons (~~Supplementary Figures 28~~Suppl. Figs
448 30 and 4243). In this region, we found a strong negative trend of RH for 1979-2014,
449 which concurs with the significant decrease of absolute humidity. We noted a
450 significant increase in air temperature, air temperature and air temperature to SST ratio,
451 while a negative and statistically significant decrease in land evapotranspiration in the
452 continental sources of moisture was observed.

453

454 **3.2.4. Other regions**

455 Other regions of the world (see Supplementary Suppl. Material) also showed strong
456 dependency between the interannual variability of RH and that of land
457 evapotranspiration in the land moisture sources. Some examples include Western
458 Europe, Central-eastern Europe, Southeast Europe, Turkey, India and the ~~east Sahel.~~
459 ~~Nevertheless, the influence of land evapotranspiration was very different between the~~
460 ~~boreal warm and cold seasons (e.g. Scandinavia, Central East Europe and the~~
461 ~~Amazonian region).~~Eastern Sahel. In contrast, other regions showed a weak correlation
462 between the temporal variability of RH and land evapotranspiration in the moisture
463 source region. A representative example is China, which witnessed a strong decrease in

464 RH for 1979-2014. ~~This might be explained largely by the fact that relative interannual~~
465 ~~ET variations are just much weaker in China compared to other regions so that the~~
466 ~~signal to noise ratio is worse in China. In this region, RH changes correlated~~
467 ~~significantly with annual precipitation only: a variable that did not show significant~~
468 ~~changes from 1979 to 2014 (Supplementary ~~Nevertheless~~ Fig. 11). This annual pattern~~
469 ~~was also observed for the boreal cold and warm seasons (Supplementary Figs. 23 and~~
470 ~~37).~~

471 ~~Nevertheless, although the interannual variability of land evapotranspiration in the land~~
472 ~~moisture sources showed the highest correlation with RH variability in the majority of~~
473 ~~the analyzed regions, air temperature/SST ratio in the oceanic moisture sources also~~
474 exhibited negative correlations with RH in particular regions, including ~~West~~Western
475 Sahel, La Plata, West Coast of the USA, Central-Eastern Europe, India, central North
476 America and the Amazonian region. This finding suggests that higher differences
477 between air temperature in the target area and SST in the oceanic moisture region would
478 favor decreased RH.

479

480 **3.3. Land and ocean contribution to RH trends**

481 ~~It is well-recognized that establishing~~Establishing a direct influence of land
482 evapotranspiration on RH is a challenging task, including also any attempt to directly
483 compare these influences with the possible contribution from oceanic evaporation and
484 moisture transport. This is primarily because, apart from very humid regions, the
485 increase in land evapotranspiration could be driven by increased precipitation, which is
486 accompanied by anomalous RH conditions. -This dependency explains the correlation
487 found between precipitation and land evapotranspiration in some regions worldwide;
488 ~~although there are considerable spatial and seasonal differences (Supplementary Figures~~

489 ~~45 to 47). (Suppl. Figs 47 to 49).~~ In cold and humid regions, land evapotranspiration is
490 also related to the interannual variability of the AED (~~Supplementary Figures 48~~~~Suppl.~~
491 ~~Figs 50 to 5052).~~ Correspondingly, the magnitude of the oceanic evaporation may be
492 insufficient to explain RH anomalies in the target region.

493 Taken together, the transport of moisture to any target region is a fundamental process;
494 ~~in which atmospheric circulation configurations can contribute significantly to RH~~
495 ~~anomalies and their spatial variations. Hence, for better understanding of the possible~~
496 ~~contribution of oceanic evaporation and land evapotranspiration to RH. Hence,~~ we
497 assessed the contribution of land and ocean to precipitation, as represented by (E-P). ~~It~~
498 ~~may provide some clues on the possible connection between oceanic moisture and RH~~
499 ~~variability. Figures 9 and 10 illustrate the relationship between the interannual~~
500 ~~variability of RH in each target region and land-oceanic contribution to the annual~~
501 ~~precipitation.~~ Overall, results reveal important differences among the analyzed regions,
502 with- statistically insignificant correlations found between the interannual variations of
503 RH in some regions and ocean and land contribution to precipitation. ~~One example is~~
504 ~~the positive and significant correlation found between the annual RH (Figs 8 and the~~
505 ~~ocean E-P in regions 2 (Scandinavia), 5 (Western Sahel), 6 (India), 12 (West North~~
506 ~~America) and 13 (Amazonia9).~~ A similar pattern was observed at the seasonal scale,
507 albeit with greater contribution during the cold season, especially in the regions where
508 precipitation is mostly driven by western flows ~~during this season~~ (e.g. West of North
509 America, Western Europe and Scandinavia) (~~Supplementary~~~~Suppl.~~ Figs ~~51, 53 to 5456).~~
510 On the other hand, the land contribution ~~of land areas~~ to precipitation ~~in the analyzed~~
511 ~~target regions~~ is rather complex, with strong spatial differences. At the annual scale, a
512 positive and significant contribution to precipitation (E-P) is found in regions 3
513 (Central-East Europe), 6 (India), 7 (China), 9 (La Plata) and 11 (central US). ~~Notably, a~~

514 ~~wide range of regions exhibit a positive and significant correlation between land (E-P)~~
515 ~~and RH during the cold season, including Scandinavia, Central-East Europe, South-East~~
516 ~~Europe and Turkey, Western Sahel, India, North-East Asia, La Plata, Central USA,~~
517 ~~West North America and East Sahel. In contrast, during summertime, only regions 3~~
518 ~~(Central-East Europe) and 9 (La Plata) show a significant correlation between land~~
519 ~~contribution to precipitation (E-P) and annual RH. Given these noticeable spatial and~~
520 ~~seasonal differences, our findings suggest~~This suggests that there are no generalized
521 patterns in terms of the contribution of ocean and land to the interannual variability of
522 RH. ~~This complexity makes it quite difficult to attribute RH trends between 1979 and~~
523 ~~2014 at the global scale to a unique driver or process. The FLEXPART model outputs~~
524 ~~do not suggest consistent trends for ocean as well as land E-P. Irrespective of all these~~
525 ~~limitations, we believe that it is still possible to assess with a degree of confidence the~~
526 ~~evolution of land and ocean contribution (E-P) to precipitation of the target regions.~~
527 ~~Figure 12, making difficult to attribute RH trends to a unique driver.~~
528 Fig. 10 illustrates the evolution of the land contribution (E-P) to precipitation in the
529 different target regions. We noted positive and significant changes in the region 2
530 (Scandinavia), 3 (Central-East Europe), 5 (Western Sahel) and 14 (Eastern Sahel). A
531 contradictory behavior is observed for region 9 (La Plata), with a statistically significant
532 downward trend in land contribution (E-P) to precipitation. At the seasonal scale, results
533 suggest considerable differences (Supplementary Figures 54Suppl. Figs. 57 and 5558),
534 with no clear positive or negative trends.
535 ~~With respect to the different analysed variables, changes~~Changes in RH were more
536 associated with those of land evapotranspiration across the selected regions (Figure
537 13Fig. 11). In contrast, changes in annual RH did not correlate significantly with the
538 observed changes in precipitation, air temperature/SST and oceanic evaporation. The

539 observed patterns were similar for both the warm and the cold season
540 (~~Supplementary~~ Suppl. Figs. 5659 and 57). ~~Indeed, these~~ 60). These positive and
541 significant correlations do not imply causation between ~~these factors—land~~
542 evapotranspiration and RH variations over space and time. ~~This is simply evident in~~
543 different regions worldwide, where there are strong seasonal and spatial differences in
544 the contribution of land evapotranspiration to RH. Nevertheless, these findings also
545 confirm the suggest a role of land evapotranspiration in explaining the observed
546 variability of RH trends. Specifically, for many regions and at different temporal scales
547 (i.e. seasonal and annual), changes in land contribution to precipitation show
548 statistically significant positive correlation with changes in evapotranspiration and
549 precipitation (~~Supplementary Figure 58).~~ Suppl. Fig. 61). Again, this correlation does
550 not imply a true causal relationship between RH variability and evapotranspiration,
551 given the strong coupling between many of these controlling variables (e.g.
552 precipitation, RH, and land evapotranspiration). ~~This coupling is also spatially and~~
553 temporarily variable. However, this good The agreement between changes in the land
554 contribution to precipitation and changes in land evapotranspiration ~~emphasizes the role~~
555 of land evapotranspiration in explaining the complex spatial patterns of RH changes. In
556 , but also the fact that in many regions, land contribution (E-P) to precipitation is
557 evidently important, with (contributions close to 50% or even higher, including the
558 Western Sahel (54%), Eastern Sahel (61%) and North East Asia (64%)
559 (Supplementary Table 1). ~~), suggests the possible role of land evapotranspiration~~
560 in explaining the spatial patterns of RH changes. Nevertheless, there is some uncertainty
561 in this attribution since the good spatial agreement between RH, precipitation and land
562 evapotranspiration makes it difficult to accurately define the most dominant variable(s)
563 that control the temporal variability of RH (Suppl. Figs 62 - 64).

564 ~~Nevertheless, there is some uncertainty in attributing RH changes to land~~
565 ~~evapotranspiration. Figure 14 depicts the relationship between RH and land~~
566 ~~evapotranspiration seasonally and annually at the global scale. Note that these are local~~
567 ~~("pixel by pixel") correlations and the interpretation differs from the previous analysis~~
568 ~~The complex spatial~~ where RH in target regions is correlated with ET in corresponding
569 source regions. Results reveal strong positive and significant correlations in large areas
570 of the world. The strongest positive correlations were found in Central, West and
571 Southwest North America, Argentina, east Brazil, South Africa, the Sahel, central Asia
572 and the majority of Australia. Nevertheless, there are some exceptions, including large
573 areas of the Amazon, China, central Africa and the high latitudes of the Northern
574 Hemisphere, where the correlations were negative. In general, the areas with positive
575 and significant correlations between RH and land evapotranspiration corresponded to
576 those areas characterized by semiarid and arid climate characteristics, combined with
577 some humid areas (e.g. India and northwest North America). Nevertheless, at the global
578 scale, the correlation between RH and land evapotranspiration shows spatial patterns
579 consistent with those based on the correlation between RH and precipitation. Similar
580 spatial patterns are found also for the correlation between precipitation and
581 evapotranspiration at both seasonal and annual scales (Supplementary Figures 59 and
582 60). This high agreement makes it difficult to accurately define the most dominant
583 variable (s), which control the temporal variability of RH, given the good spatial
584 agreement between RH and different variables (e.g. precipitation, land
585 evapotranspiration). The complex spatial patterns of the observed trends for different
586 variables add another source of uncertainty to proper attribution of RH changes. **Figure**
587 **15** Fig. 12 illustrates the spatial distribution of the magnitude of change in annual and
588 seasonal land evapotranspiration at the global scale from 1979 to 2014. As depicted, the

589 spatial patterns of land evapotranspiration changes resemble those of RH in some
590 regions (refer to ~~Figure 2). For example, a positive trend in the annual land~~
591 ~~evapotranspiration dominated over Fig. 1), including -for example-~~ the Canadian region,
592 ~~which agrees well with the general increase in RH across the region. On the other hand,~~
593 ~~there was a dominant decrease in the annual land evapotranspiration across vast areas of~~
594 ~~North America, which concurs also with the strong decrease in RH. Similar to the~~
595 ~~pattern observed for land evapotranspiration, RH increased particularly over southwest~~
596 ~~North America. In South America, both variables also showed a dominant negative~~
597 ~~trend at the annual scale, but with some spatial divergences, mainly in the Amazonian~~
598 ~~region. Specifically, the western part of the basin showed the most important decrease~~
599 ~~in land evapotranspiration, whereas the most significant decrease in RH was observed in~~
600 ~~the eastern part. In the African continent, some areas showed good agreement between~~
601 ~~RH and land evapotranspiration changes, in terms of both the sign Western and~~
602 ~~magnitude. This can be clearly seen in the West and East Eastern Sahel, where a strong~~
603 ~~gradient in RH trend between the West (positive) and the East (negative) was observed.~~
604 Nevertheless, other ~~African~~ regions showed a divergent pattern between both variables:
605 ~~One example is (e.g. the Guinea Gulf in Nigeria and Cameroon),~~ where we noted a
606 strong increase in land evapotranspiration, as opposed to RH changes. ~~In Australia,~~
607 ~~although both variables showed a dominant positive trend, they did not match exactly in~~
608 ~~terms of the spatial pattern of the magnitude of change.~~ The Eurasian continent showed
609 the main divergences between both variables. ~~In the high latitudes of the continent,~~
610 ~~there was a dominant increase in both variables. For other regions (e.g. For example, in~~
611 ~~the Western Europe),~~ we noted a dominant RH decrease, which was not observed for
612 land evapotranspiration. A similar pattern was observed over east China, with a
613 dominant RH negative trend and a positive land evapotranspiration. Overall, the lack of

614 significant spatial association between the magnitude of trends in RH and the magnitude
615 of trends in evapotranspiration can be seen in the context of the strong spatial diversity
616 of trends of these two variables at both annual and seasonal scales (~~Supplementary~~
617 ~~Figure 61~~). ~~Similar patterns are found also for the trends of RH and precipitation, and~~
618 ~~the trends of precipitation and land evapotranspiration, although precipitation trends~~
619 ~~show a very different pattern, in comparison to land evapotranspiration trends~~
620 ~~(Supplementary Figure 62). Results~~ Suppl. Fig. 65). This complexity is similar to that
621 found for trends in precipitation (Suppl. Fig. 66). Thus, results suggest that while the
622 variability of precipitation, RH and land evapotranspiration show strong interannual
623 associations, their observed trends are completely decoupled over space. This high
624 spatial variability of trends at the global scale confirms that direct attribution of
625 observed RH changes to ~~oceanic~~/land contributions is a challenge and -quite a complex
626 task.

627 In relation to the ~~influence of the~~ ocean evaporation, ~~our results confirm that the global~~
628 ~~connection between oceanic evaporation and changes in RH is also complex. On one~~
629 ~~hand,~~ it is ~~difficult~~ also quite to establish ~~a pixel per pixel relationship since the sources~~
630 ~~of moisture may strongly differ at the global scale. On the other hand~~ this connection. In
631 particular, it is not feasible to identify the moisture sources and the ocean contribution
632 of the precipitation for each 0.5° land pixel at the global scale. However, we believe that
633 the analysis of the evolution of SST and oceanic evaporation for 1979-2014 and the
634 evolution of the oceanic contribution to precipitation can give indications on some
635 relevant patterns. ~~Figure 16~~ Fig. 13 illustrates the spatial distribution of the magnitude of
636 change of annual and seasonal SST and oceanic evaporation. ~~Supplementary~~ Suppl. Fig.
637 ~~6367~~ shows the spatial distribution of trend significance. As depicted, complex spatial
638 patterns and high variability of the trends were observed, particularly for oceanic

639 evaporation. Furthermore, the spatial distribution of the magnitude of change in annual
640 and seasonal oceanic evaporation was not related to the SST changes
641 (~~Supplementary Suppl. Fig. 68, Fig. 64. This finding suggests that oceanic evaporation~~
642 ~~is not only driven by changes in SST.~~ Thus, although some regions showed positive
643 changes in the oceanic evaporation, the amount of increase was much lower than that
644 found for SST, which suggests that only SST changes do not drive evaporation changes
645 (~~Supplementary Figure 65, Supplementary Suppl. Fig. 69, Suppl. Table 2).~~

646

647 **4. Discussion**

648 **4.1. Relative Humidity trends**

649 We assessed the temporal variability and trends of relative humidity (RH) at the global
650 scale using a dense observational network of meteorological stations (HadISDH) and
651 reanalysis data (ERA-Interim). Results revealed high agreement of the interannual
652 variability of RH using both datasets for 1979-2014. ~~This finding was also confirmed,~~
653 ~~even for the regions where the density of the HadISDH observatories was quite poor~~
654 ~~(e.g. the northern latitudes and tropical and equatorial regions).~~ Recent studies have
655 suggested dominant decrease in observed RH during the last decade (e.g. Simmons et
656 al., 2010; Willet et al., 2014). Our study suggests dominant negative trends of RH using
657 the HadISDH dataset. This decrease is mostly linked to the temporal evolution of RH
658 during the boreal warm season. ~~Nevertheless, other regions showed positive RH trends.~~
659 In accordance with the HadISDH dataset, the ERA-Interim revealed dominant negative
660 RH trends, albeit with a lower percentage of the total land surface compared to the
661 HadISDH dataset. These differences cannot be attributed to the selected datasets, given
662 that both mostly agree on the magnitude and sign of changes in RH.

663 Observed changes in RH were closely related to the magnitude and the spatial patterns
664 of specific humidity changes. Results demonstrate a general deficit of specific humidity
665 to maintain RH constant in large areas of the world, including the central and south
666 Northern America, the ~~Amazonas~~Amazonia and La Plata basins in South America and
667 the ~~East~~Eastern Sahel. In other regions, RH increased in accordance with higher specific
668 humidity. ~~Some studies suggested~~Studies suggest that changes in air temperature could
669 partly cancel the effects of the atmospheric humidity to explain RH changes (e.g.
670 McCarthy and Tuomi, 2004; Wright et al., 2010; Sherwood, 2010). Nevertheless,
671 although air temperature trends showed spatial differences at the global scale over the
672 past four decades (IPCC, 2013), our results ~~confirm~~show that air temperature is not the
673 main driver of the observed changes of RH globally. ~~The ERA-Interim dataset clearly~~
674 ~~showed~~There is a close resemblance between ~~RH~~relative and specific humidity trends
675 at the global scale. ~~This suggests, suggesting~~ that specific humidity is the main driver ~~of~~
676 ~~the observed changes~~ in the magnitude and spatial pattern of RH ~~during the past~~
677 ~~decade~~trends.

678

679 ***4.2. Contribution of continental areas to changes in RH***

680 Overall, there is an agreement between the interannual variability of precipitation ~~and~~,
681 land evapotranspiration in the continental moisture source and the interannual
682 variability of RH in different regions. Nevertheless, considering gridded datasets at the
683 global scale, we found that this good agreement is restricted only to the arid and
684 semiarid regions. In humid regions, soil moisture is not ~~a~~-constrained ~~variable; the~~
685 ~~variability of~~so land evapotranspiration variability is mostly driven by changes in the
686 AED (Stephenson, 1990). This makes ~~it~~ difficult to unravel the possible direct
687 contribution of land evapotranspiration to the variability of RH using statistical

688 approaches and empirical information, particularly with the strong coupling among
689 these variables. Land evapotranspiration is closely related to precipitation variability in
690 arid and semiarid regions; increased land evapotranspiration ~~thus~~ tends to be caused
691 primarily by increased precipitation, which is accompanied by corresponding RH
692 anomalies. Also, RH may affect land evapotranspiration, both in arid and humid
693 regions, given its important contribution to the aerodynamic component of the AED
694 (Wang et al., 2012; Vicente-Serrano et al., 2014a).

695 Nevertheless, although the interannual variability of these three variables ~~can be~~
696 ~~strongly~~are coupled in some regions, ~~the~~their long-term trends ~~in these variables~~ may
697 strongly differ; as a consequence of changes in precipitation, increasing influence of
698 ~~AED on land evapotranspiration~~the AED, and also changes in land and atmospheric
699 contribution to RH and precipitation. The fourteen analyzed regions, in which
700 FLEXPART was applied, show a relevant continental contribution (E-P) to precipitation
701 ~~in these areas~~. The average contribution is generally below 40% of the total
702 precipitation in some regions (e.g. Western Europe, Scandinavia or West North
703 America), ~~which~~but exceeds 50% in specific regions (e.g. Sahel and East China).

704 Therefore, it is reasonable to consider that changes in the contribution of
705 ~~continents~~continental areas to precipitation may affect land evapotranspiration processes
706 and ultimately affect RH variability ~~in these continental areas~~. Thus, our results suggest
707 an influence of land-atmosphere water feedbacks and recycling processes on RH trends.

708 This is simply because more available soil humidity under favorable atmospheric and
709 land conditions would result in more evapotranspiration and accordingly higher air
710 moisture (Eltahir and Bras, 1996; Domínguez et al., 2006; Kunstmann and Jung, 2007).

711 Recalling that the ocean surface ~~evaporates~~contributes about 84% of the water
712 evaporated over the Earth (Oki, 2005), ~~the~~ oceanic evaporation is highly important for

713 continental precipitation (Gimeno et al., 2010). However, the continental humidity
714 sources can be ~~also~~ more important than oceanic sources in some regions (e.g. the
715 Sahel) (Wei et al., 2016a). In this context, our results concur with previous works. For
716 example, ~~numerous model-based studies have supported an influence of land~~
717 ~~evaporation processes on air humidity and precipitation over land surfaces (e.g.~~
718 ~~Bosilovich and Chern, 2006; Dirmeyer et al., 2009; Goessling and Reick, 2011).~~
719 ~~Moisture~~ recycling is strongly important in some regions of the world, such as China
720 and central Asia, the western part of Africa and the central South America (Pfahl et al.,
721 2014; van der Ent et al., 2010).
722 ~~, and in general in semi-arid and desert areas worldwide (Miralles et al., 2016).~~ All these
723 studies assessed the role of continental evapotranspiration on average precipitation
724 conditions, with few studies focusing on the possible impacts of changes in soil
725 moisture/evapotranspiration on RH. ~~In Europe,~~ Rowell and Jones (2006) ~~analyzed~~
726 ~~different hypotheses to explain the projected summer drying conditions in Europe,~~
727 ~~suggesting that soil moisture decline and land sea contrast in lower tropospheric~~
728 ~~summer could be the key factors responsible for this drying. They concluded for future~~
729 ~~climate scenarios~~ that reduced evaporation in summer will drop RH and hence ~~reduced~~
730 continental rainfall. ~~These would impact soil moisture and evapotranspiration processes,~~
731 ~~inducing a reduction in RH and rainfall, through a range of atmospheric feedbacks. In~~
732 ~~the same context, the importance of moisture recycling processes for atmospheric~~
733 ~~humidity and precipitation has been recently identified in semi arid and desert areas of~~
734 ~~the world (Miralles et al., 2016).~~
735 Although our study was limited to specific regions across the world, results indicate that
736 humidity in the analyzed regions is substantially originated over continents. This
737 finding concurs with some regional studies that defined sources of moisture (e.g. Nieto

738 et al., 2014; Gimeno et al., 2010; Drumond et al., 2014; Ciric et al., 2016). Overall, the
739 spatial differences in the possible attribution of the observed changes in RH to changes
740 in land evapotranspiration are important. Nevertheless, in some regions ~~where RH is~~
741 ~~strongly correlated with land evapotranspiration, there is a significant correlation~~
742 ~~between land contributions to RH, suggesting our results suggest~~ a robust contribution of
743 land processes to the interannual variability of RH ~~in these regions~~. A representative
744 example is the La Plata ~~region~~, where a strong decrease (-6.6% from 1979 to 2014) in
745 RH is suggested by both observations and ~~reanalysis datasets. ERA-Interim~~. This region
746 did not exhibit a significant trend in precipitation, but conversely there is a significant
747 decrease in absolute humidity and land evapotranspiration. In the La Plata region, the
748 oceanic evaporation in the source region has shown a significant increase (6.33%) since
749 1979. However, this increase seems to be insufficient to favor an increase in RH values.
750 Herein, although the average oceanic contribution (E-P) to precipitation is slightly
751 higher (54%), compared to the continental contribution (46%), the interannual
752 variability of RH is positively correlated with the interannual variability of land E-P
753 rather than oceanic E-P. Moreover, this region exhibited a significant decrease in land
754 contribution to precipitation between 1979 and 2014.

755 ~~On the contrary, in other regions like~~In the Western US, the large decrease in RH (-
756 6.2%) corresponded to a large decrease in the absolute humidity (-0.58 g/kg). However,
757 it is difficult to attribute this association pattern to changes in land evapotranspiration,
758 given the low (37%) continental contribution to precipitation ~~in these regions~~.
759 ~~Moreover,~~ Wei et al. (2016b) ~~indicated~~showed that the transport of atmospheric
760 moisture from the Pacific is the main contributor to the interannual variability of
761 precipitation in the region. In this sense, we found a strong relationship between the
762 interannual variability of RH ~~in this region~~ and the oceanic E-P, albeit with insignificant

763 trends in the oceanic and continental contribution (E-P) to precipitation in the region.
764 Here, in the absence of significant changes in oceanic evaporation and contribution to
765 precipitation, it ~~could be~~ reasonable to consider that the decrease in absolute humidity
766 is linked to the atmospheric circulation that control moisture transport in the region
767 (Wei et al., 2016b). ~~The~~ but also the decrease in RH could be ~~also due to the positive~~
768 ~~and significant~~ avored by the trend toward higher differences between air temperature
769 in the land target region and the ~~oceanic temperature~~ SST in the source region.
770 Therefore, the relationships between RH, land evapotranspiration variability and
771 changes in the contribution of continental areas ~~to air moisture and precipitation in the~~
772 ~~target regions~~ can be extremely complex. Similarly, the relationships between RH and
773 land evapotranspiration are rather complex and cannot be easily interpreted.
774 Nonetheless, albeit with the strong uncertainty existing at the global scale, it is
775 reasonable to consider that ~~strong~~ changes in soil moisture budget, combined with land
776 evapotranspiration, could impact somehow the ~~complex~~ observed RH trends ~~observed at~~
777 ~~the global scale~~. In the same context, there is strong evidence that low levels of soil
778 moisture and land evapotranspiration are usually accompanied by a reinforcement of
779 low RH, particularly during drought episodes. Under these circumstances, the
780 suppression of the latent heat flows from the soil to the atmosphere would enhance soil
781 and vegetation warming and sensible heat, inducing air temperature rise. ~~Also, the lack~~
782 ~~of supply of water vapor to the atmosphere favors the decrease of RH~~ and the
783 reinforcement ~~of severity~~ of heat waves (Hirschi et al., 2011). Seneviratne et al. (2002)
784 showed that vegetation control on transpiration might contribute significantly to
785 enhancement of summer drying, particularly when soil water is limited. Other studies
786 confirmed this finding ~~for other regions worldwide~~, employing both observational data
787 (e.g. Hirschi et al., 2011) and model outputs (e.g. Seneviratne et al., 2006; Fischer et al.,

788 | 2007). ~~Our study suggests good spatial agreement between changes in RH and those of~~
789 | ~~continental contribution to precipitation as well as land evapotranspiration during~~
790 | ~~summertime. Although this finding is markedly evident for all the analyzed regions, it~~
791 | ~~should be seen with caution. This is mainly because physical processes driven soil~~
792 | ~~moisture are more active during the warm season (Vautard et al., 2007 and 2013;~~
793 | ~~Miralles et al., 2014), which adds difficulty to establish full causality between RH and~~
794 | ~~other driving forces during this season.~~

795

796 | **4.2. Contribution of oceans to changes in RH**

797 | ~~This study demonstrates~~We indicated strong differences among the fourteen
798 | ~~analysed~~analyzed regions in terms of the contribution of Oceanic water bodies to RH
799 | variability. In some regions (e.g. Western North America, India, Scandinavia and the
800 | Amazonian), the interannual variability of RH is closely related to oceanic E-P,
801 | indicating that changes in oceanic evaporation, combined with the processes of
802 | atmospheric moisture transport to target regions, play a main role in explaining changes
803 | in RH. ~~It is well recognized that t~~Moisture advection is the main driver of
804 | precipitation variability ~~in the majority of world regions~~ (Trenberth, 1999; Wei et al.,
805 | 2016a). Nevertheless, there is some uncertainty in recent trends of moisture advections
806 | from oceanic areas (Zhan and Allan, 2013). As such, it is difficult to determine -in
807 | general terms- whether the strong complexity of RH changes identified at the global
808 | scale ~~are~~is driven by changes in moisture advections from oceanic areas. ~~Nevertheless,~~
809 | given that - in the context of changing climate, - SST, oceanic evaporation, and oceanic
810 | contribution to precipitation in the target regions can jointly account for the possible
811 | influence of oceans on RH variability in some regions.

812 Different modelled climate studies suggested strong differences between land and ocean
813 RH trends, as a consequence of the different warming rates between oceanic and
814 continental areas (e.g. Joshi et al., 2008; Dessler and Sherwood, 2009; O’Gorman and
815 Muller, 2010). As the warming rates are generally slower over oceans, the specific
816 humidity of air advected from oceans to continents would increase more slowly than the
817 saturation specific humidity over land, causing a reduction in RH (Rowell and Jones
818 2006). Due to this effect, RH will not remain constant in areas located very far from
819 humidity sources, as warmer air temperatures under limited moisture humidity would
820 reduce RH (Pierce et al., 2013). This physical process could explain ~~the recent observed~~
821 ~~RH trends of RH~~ in some regions (e.g. ~~Amazonia~~), where around 68% of the average
822 atmospheric moisture originates over oceanic sources. ~~Moreover,~~ In this region RH is
823 correlated with the oceanic E-P, although there ~~are~~ is no ~~changes~~ change in the oceanic
824 contribution to precipitation from 1979 to 2014. ~~In Amazonia~~ Moreover, RH strongly
825 decreased at the annual scale (-7.7%), as a consequence of ~~the~~ a decrease ~~(-0.56 g/kg)~~
826 the absolute humidity; ~~(-0.56 g/kg)~~, accompanied by an increase ~~(1.09°C)~~
827 temperature; ~~(1.09°C)~~. While SST in the oceanic source region slightly increased by
828 0.33°C, other variables (i.e. oceanic evaporation, precipitation and land
829 evapotranspiration) did not ~~t~~ exhibit significant changes. Herein, we believe that the
830 mechanisms proposed by Sherwood and Fu (2014) to explain decreased RH are
831 applicable to our study. They attributed RH decrease to sub-saturated oceanic moisture
832 supply, which compensates ~~a strong~~ air temperature increase. In this study, this
833 mechanism is also capable of explaining RH variability, given that the difference in
834 warming rates between SST in the oceanic moisture source region and air temperature
835 of the Amazonia is increasing. This feature is also supported by the significant
836 correlation found between the interannual variability of RH and the ratio between air

837 | temperature and SST. ~~Again, this explanation does not guarantee full causality between~~
838 | ~~RH variability and oceanic moisture in the source regions. over land and SST.~~ A
839 | similar pattern is found for the Eastern Sahel, a region in which continental recycling is
840 | particularly important (Wei et al., 2016a). This region witnessed a strong decrease in
841 | RH, but not compensated by increased precipitation. Although there is no significant
842 | correlation between RH and the ratio between air temperature and SST, the latter
843 | variable shows a significant increase. This could reinforce the drying effect of the
844 | suggested trend toward a lower moisture supply from the oceanic source, especially
845 | with the significant negative trend in oceanic contribution to precipitation in this region.
846 | In other regions, there are no ~~coherent~~ empirical relationships that confirm the impact of
847 | oceanic moisture supply processes on RH changes. ~~Recalling the observed negative RH~~
848 | ~~trend at many coastal regions over the period 1979-2014, this study confirms that the~~
849 | ~~distance to oceanic humidity sources is not a key controller of the spatial patterns of RH~~
850 | ~~changes. In many instances, we found that continental regions, which are very far from~~
851 | ~~oceans (e.g. Canada, central China and Kazakhstan), recorded a positive RH trend.~~ A
852 | possible explanation of these contrasting findings could be related to the low differences
853 | in the warming rates between the oceanic sources and the majority of the continental
854 | target areas. We found that -in most of the cases- these differences were not strong
855 | enough to generate a clear effect at the global scale, particularly with the available
856 | number of observations. ~~In specific regions (e.g. Amazonia, East Sahel, Western North~~
857 | ~~America), there is a a positive trend in the difference between air temperature and SST~~
858 | ~~temperature in the source region. These processes could be the most reasonable physical~~
859 | ~~explanation of the strong observed RH trends.~~
860 | Although oceanic evaporation is decisive on moisture supply to continental regions
861 | (Gimeno et al., 2010), several processes, which are not considered in this study, may

862 strongly affect RH and precipitation in continental areas. ~~Under a generalized increase~~
863 ~~of SST at the global scale (e.g. Rayner et al., 2003; Deser et al., 2010), a higher~~
864 ~~moisture supply to the atmosphere and a strong decrease in RH can be expected in~~
865 ~~different regions. Some of these regions (e.g. Amazonia and Western North America)~~
866 ~~show an average high moisture supply from oceanic areas.~~ On one hand, a global
867 warming signal does not necessarily imply above-normal oceanic evaporation. Here,
868 we indicated that oceanic evaporation trends for 1979-2014 showed strong spatial
869 variability ~~at the global scale~~, with dominant positive trends. Nevertheless, large areas
870 also exhibited insignificant trends and even negative evaporation trends. While SST
871 increase is mainly associated with radiative processes, evaporation processes are mainly
872 controlled by a wide range of meteorological variables that impact the aerodynamic and
873 radiative components of the atmospheric evaporative demand (AED) rather than SST
874 alone (McVicar et al., 2012b). ~~This is consistent with the finding that global mean~~
875 ~~precipitation or evaporation does not scale with Clausius-Clapeyron (Held and Soden~~
876 ~~2006). Due to the unlimited water availability over oceans, air vapor pressure deficit is~~
877 ~~expected to be driven by the Clausius-Clapeyron relation. However~~ Thus, changes in
878 solar radiation and wind speed can also influence the evaporation evolution (Yu, 2007;
879 Kanamaru and Masunaga, 2013).

880 Due to the unlimited water availability over oceans, air vapor pressure deficit would be
881 expected to be driven by the Clausius-Clapeyron relation. Nevertheless, the observed
882 evolution indicates that the global mean precipitation/evaporation is not scaled by the
883 Clausius-Clapeyron relation (Held and Soden 2006). As such, given the slow oceanic
884 evaporation trends in large regions of the world, RH trends in some of the analyzed
885 regions could be seen in the context of a lower water supply to maintain RH constant,
886 particularly with air temperature rise.

887 Herein, we have not considered possible changes in other variables that could explain
888 the lowweak relationship between RH, oceanic evaporation and oceanic moisture
889 contribution. ~~Among these variables, we have not considered—for~~For example~~—the role~~
890 ~~of the “effectivity” of the oceanic moisture (Gimeno et al., 2012). This variables is of~~
891 ~~particular importance, because oceanic evaporation might not reach the target region,~~
892 ~~due to some geographical constraints (e.g. topography). Another relevant variable is,~~ the
893 transport mechanisms between the source and target regions, which ~~is~~are key
894 ~~variable~~variables in some regions like the Western North America (Wei et al., 2016b).
895 Moreover, moisture source regions are not stationary, as the intensity of humidity can
896 vary greatly from one year to another (Gimeno et al., 2013). This aspect could be
897 another source of uncertainty in the explanatory factors of current RH trends.
898 Furthermore, other different factors that control atmospheric humidity and RH have not
899 been approached in this study. Sherwood (1996) indicated that RH distributions are
900 strongly controlled by dynamical fields rather than local air temperatures. This suggests
901 that atmospheric circulation processes could largely affect the temporal variability and
902 trends of RH. A range of studies indicates noticeable RH changes ~~in RH~~, in response to
903 low-frequency atmospheric oscillations, such as the Atlantic Multidecadal Oscillation
904 (AMO) and El Niño-Southern Oscillation (e.g. McCarthy and Toumi, 2004; Zhang et
905 al., 2013), regional circulation (Wei et al., 2016a and 2016b), as well as changes in the
906 Hadley Cell (HC) (Hu and Fu, 2007). Wright et al. (2010) employed a global climate
907 model under double CO₂ concentrations to show that tropical and subtropical RH is
908 largely dependent on a poleward expansion of the Hadley cell: a deepening of the height
909 of convective detrainment, a poleward shift of the extratropical jets, and an increase in
910 the height of the tropopause. Also, Laua and Kim (2015) assessed changes in the HC
911 under CO₂ warming from the Coupled Model Intercomparison Project Phase-5 (CMIP5

912 model projections. They suggest that strengthening of the HC induces atmospheric
913 moisture divergence and reduces tropospheric RH in the tropics and subtropics. ~~This;~~
914 ~~this~~ spatial pattern resembles ~~the main~~ those areas ~~showing, with~~ negative RH trends in
915 ~~RH in~~ Northern as well as Southern hemisphere, ~~, as described in this study.~~

Con formato: Fuente: Sin Negrita

917 5. Conclusions

918 This study ~~analysed~~ analyzed relative humidity (RH) trends at the global scales using
919 observations and ERA-40 data. It extended further to link RH trends with a range of
920 variables, which can give indications on the possible oceanic and continental
921 contribution to RH trends. ~~As opposed to the widely accepted constant RH scenario~~
922 ~~under global warming, our~~ Our results suggest significant RH trends over many regions
923 worldwide, ~~but including: There are positive trends in RH over specific regions (e.g.~~
924 ~~high latitudes of the North Hemisphere, Northern South America, India, West Sahel);~~
925 ~~which is in contrast to the~~ and a generally dominant negative trend at the global scale.
926 ~~This decrease is mostly linked to the temporal evolution of RH during the boreal warm~~
927 ~~season.~~
928 ~~There is a~~ The strong diversity in the observed RH trends, ~~highlighting highlights~~ the
929 complex and divergent role of different physical processes ~~and drivers~~, including both
930 dynamic and thermodynamic ~~processes~~ components. In general the supply of specific
931 humidity is a main source of the observed RH trends since there is a high agreement
932 between RH and specific humidity trends ~~at the global scale, suggesting that moisture~~
933 ~~deficit contributes to RH variability, in opposition to atmospheric warming.~~ This
934 finding suggests that the evolution of specific humidity in vast areas of the world has
935 not provided the necessary humidity to maintain RH constant according to the observed
936 warming trends. ~~This feature is important, given its implications in terms of~~

937 ~~atmospheric evaporative demand and aridity conditions under the current climate~~
938 ~~change scenario.~~

939 This study also analyzed the possible contribution of continental and oceanic moisture
940 supply to explaining the magnitude and spatial patterns of RH trends. For this purpose,
941 fourteen regions were defined and the contribution of continental and oceanic sources to

942 RH ~~in these regions~~ was assessed using a Lagrangian scheme. Results
943 indicate that no single physical mechanism can be responsible for the observed trends in

944 RH at the global scale. ~~Globally, there are~~ Our findings stress that two ~~well-recognized~~
945 ~~hypotheses for explaining the possible decrease in RH under a global warming scenario:~~

946 (i) ~~the land water supply by means evapotranspiration processes~~, and (ii) the insufficient
947 oceanic moisture supply to maintain continental RH constant ~~under different warming~~

948 ~~rates between continental and oceanic regions. Our findings stress that these two~~
949 ~~hypotheses~~, could act together to explain recent RH trends. However, although it is

950 quite difficult to establish a direct causality between RH and different underlying
951 processes ~~and driven variables~~ using different empirical sources, the observed decrease

952 ~~in~~ RH in some regions (e.g. ~~the~~ La Plata) can be linked to the lower water supply from
953 land evapotranspiration. In other regions, the empirical relationships suggest dynamic

954 and thermodynamic mechanisms related to moisture supply from oceanic source regions
955 (e.g. Amazonia and Western North America). Taken together, these physical

956 mechanisms could coexist in some ~~analyzed~~ regions, given the strong relationship found
957 between precipitation, RH and land evapotranspiration. ~~This strong coupling among~~

958 ~~these variables makes it difficult to establish a direct physical attribution of RH~~
959 ~~variability.~~

960 Overall, this study confirms the strong complexity of determining a general physical
961 process that may explain the complex spatial patterns of RH trends, particularly at the

962 global scale. As such, further research is still needed to unravel the complex physical
963 factors driving the dominant RH negative trends over large continental regions. The
964 availability of long-term historical and ~~reanalysisreanalyses~~ data and the advancement
965 of modelling approaches ~~is-an-assetare potential assets~~ in any future research to explore
966 whether the land and oceanic processes drive the observed RH trends
967 ~~Understanding. This is important, given that understanding~~ current RH is relevant in
968 hydroclimatic research, due to its impacts on atmospheric evaporative demand, crop
969 development and yield, forest fire risk, bioclimatic comfort, besides other hydrological
970 processes. ~~This study provides the first comprehensive analysis of RH at the global~~
971 ~~scale based on empirical information, comprising state-of-the-art modelling approaches~~
972 ~~and foreing scenarios.~~

973

974 **Acknowledgements**

975 We thank Dr. H. F. Goessling and the two anonymous reviewers for their insightful and
976 constructive comments, which improved significantly the quality of the manuscript.
977 This work was supported by the EPhysLab (UVIGO-CSIC Associated Unit), PCIN-
978 2015-220, CGL2014-52135-C03-01, CGL2014-60849-JIN, ~~Red de variabilidad y~~
979 ~~cambio climático RECLIM (CGL2014-517221-REDT)~~ financed by the Spanish
980 Commission of Science and Technology and FEDER, IMDROFLOOD financed by the
981 Water Works 2014 co-funded call of the European Commission and INDECIS, which is
982 part of ERA4CS, an ERA-NET initiated by JPI Climate, and funded by FORMAS (SE),
983 DLR (DE), BMWFW (AT), IFD (DK), MINECO (ES), ANR (FR) with co-funding by
984 the European Union (Grant 690462).

985

986 **References**

- 987 Berg, A., Findell, K., Lintner, B., et al.: Land-atmosphere feedbacks amplify aridity
988 increase over land under global warming, *Nature Climate Change*, 6, 869-874,
989 2016.
- 990 Bosilovich, M.G., Chern, J.-D.: Simulation of water sources and precipitation recycling
991 for the MacKenzie, Mississippi, and Amazon River basins, *Journal of*
992 *Hydrometeorology*, 7, 312-329, 2006.

993 Byrne, M.P., O’Gorman, P.A.: Link between land-ocean warming contrasts and surface
994 relative humidities in ~~simulationssimulations~~ with coupled climate models,
995 Geophysical Research Letters, 40, 5223-5227, 2013.

996 Buck, A. L.: New equations for computing vapor pressure and enhancement factor, J.
997 Appl. Meteorol., 20, 1527–1532, 1981.

998 Ciric, D., Stojanovic, M., Drumond, A., Nieto, R., Gimeno, L.: Tracking the origin of
999 moisture over the ~~danube~~Danube river basin using a Lagrangian approach.
1000 Atmosphere, 7, 162, 2016.

1001 Dai, A.: Recent climatology, variability, and trends in global surface humidity. J Clim,
1002 19, 3589–3606, 2006.

1003 Dee, D. P., Uppala, S. M., Simmons, A. J. et al.: The ERA-Interim reanalysis:
1004 configuration and performance of the data assimilation system, Q. J. R.
1005 Meteorol. Soc., 137, 553–597, 2011.

1006 ~~Deser, C., Phillips, A.S., Alexander, M.A.: Twentieth century tropical sea surface~~
1007 ~~temperature trends revisited, Geophysical Research Letters, 37, L10701, 2010.~~

1008 Dessler, A. E., Sherwood, S. C.: Atmospheric Science: A Matter of Humidity, Science,
1009 323, 1020–1021, 2009.

1010 Dirmeyer, P.A., Schlosser, C.A., Brubaker, K.L.: Precipitation, recycling, and land
1011 memory: An integrated analysis, Journal of Hydrometeorology, 10, 278-288,
1012 2009.

1013 Dominguez, F., Kumar, P. Liang, X., Ting, M.: Impact of atmospheric moisture storage
1014 on precipitation recycling, J. Clim., 19, 1513–1530, 2006.

1015 Drumond, A., Marengo, J., Ambrizzi, T. et al.: The role of the Amazon Basin moisture
1016 in the atmospheric branch of the hydrological cycle: A Lagrangian analysis,
1017 Hydrology and Earth System Sciences, 18, 2577-2598, 2014.

1018 Eltahir, E.A.B., Bras, R.L.: Precipitation recycling, Reviews of Geophysics, 34, 367-
1019 378, 1996.

1020 Fan, Z.-X., Thomas, A.: Spatiotemporal variability of reference evapotranspiration and
1021 its contributing climatic factors in Yunnan Province, SW China, 1961-2004,
1022 Climatic Change, 116, 309-325, 2013.

1023 Ferraro, A.J., Lambert, F.H., Collins, M., Miles, G.M.: Physical mechanisms of tropical
1024 climate feedbacks investigated using temperature and moisture trends. Journal of
1025 Climate, 28, 8968-8987, 2015.

1026 Fischer, E.M., Seneviratne, S.I., Vidale, P.L., Lüthi, D., Schär, C.: Soil moisture-
1027 atmosphere interactions during the 2003 European summer heat wave, Journal of
1028 Climate, 20, 5081-5099, 2007.

1029 Gimeno, L., Nieto, R., Trigo, R.M., Vicente-Serrano, S.M., López-Moreno, J.I.: Where
1030 does the Iberian Peninsula moisture come from? An answer based on a
1031 Lagrangian approach, Journal of Hydrometeorology, 11, 421-436, 2010.

1032 ~~Gimeno, L., Stohl, A., Trigo, R.M., et al.: Oceanic and terrestrial sources of continental~~
1033 ~~precipitation, Reviews of Geophysics, 50, RG4003, 2012.~~

1034 ~~Gimeno, L.,~~ Nieto, R., Drumond, A., Castillo, R., Trigo, R.: Influence of the
1035 intensification of the major oceanic moisture sources on continental
1036 precipitation, Geophysical Research Letters, 40, 1443-1450, 2013.

1037 Goessling, H.F. and Reick, C.H., 2011. What do moisture recycling estimates tell us?
1038 Exploring the extreme case of non-evaporating continents. Hydrology and Earth
1039 System Sciences, 15, pp.3217-3235.

1040 Hamed, K. H., Rao, A. R.: A modified Mann Kendall trend test for autocorrelated data,
1041 J. Hydrol., 204, 182–196, 1998.

1042 Harris, I., Jones, P.D., Osborn, T.J., Lister, D.H.: Updated high-resolution grids of
1043 monthly climatic observations – the CRU TS3.10 Dataset, *International Journal*
1044 *of Climatology*, 34, 623–642, 2014.

1045 Held, I.M. and Soden, B.J., 2006. Robust responses of the hydrological cycle to global
1046 warming. *Journal of Climate*, 19: 5686-5699.

1047 Hirschi, M., Seneviratne, S.I., Alexandrov, V. et al.: Observational evidence for soil-
1048 moisture impact on hot extremes in southeastern Europe, *Nature Geoscience*, 4,
1049 17-21, 2011.

1050 Hosseinzadeh Talaei, P., Sabziparvar, A.A., Tabari, H.: Observed changes in relative
1051 humidity and dew point temperature in coastal regions of Iran, *Theoretical and*
1052 *Applied Climatology*, 110, 385-393, 2012.

1053 Hu, Y., Fu, Q.: Observed poleward expansion of the Hadley circulation since 1979,
1054 *Atmospheric Chemistry and Physics*, 7, 5229-5236, 2007.

1055 IPCC: *Climate Change 2013: The Physical Science Basis. Contribution of Working*
1056 *Group I to the Fifth Assessment Report of the Intergovernmental Panel on*
1057 *Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J.
1058 Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge
1059 University Press, Cambridge, United Kingdom and New York, NY, USA, 1535
1060 pp, 2013.

1061 Jensen, M. E., Burman, R. D., and Allen, R. G. (Eds.): *Evapotranspiration and Irrigation*
1062 *Water Requirements: ASCE Manuals and Reports on Engineering Practices No.*
1063 *70*, American Society of Civil Engineers, New York, 360 pp., 1990.

1064 Jhajharia, D., Shrivastava, S.K., Sarkar, D., Sarkar, S.: Temporal characteristics of pan
1065 evaporation trends under the humid conditions of northeast India, *Agricultural*
1066 *and Forest Meteorology*, 149, 763-770, 2009.

1067 Joshi, M.M., Gregory, J.M., Webb, M.J., Sexton, D.M.H., Johns, T.C.: Mechanisms for
1068 the land/sea warming contrast exhibited by simulations of climate change,
1069 *Climate Dynamics*, 30, 455-465, 2008.

1070 Kanamaru, K., Masunaga, H.: A satellite study of the relationship between sea surface
1071 temperature and column water vapor over tropical and subtropical oceans,
1072 *Journal of Climate*, 26, 4204-4218, 2013.

1073 Kennedy J.J., Rayner, N.A., Smith, R.O., Saunby, M., Parker, D.E.: Reassessing biases
1074 and other uncertainties in sea-surface temperature observations since 1850 part
1075 1: measurement and sampling errors, *J. Geophys. Res.*, 116, D14103, 2011a.

1076 Kennedy J.J., Rayner, N.A., Smith, R.O., Saunby, M. and Parker, D.E.: Reassessing
1077 biases and other uncertainties in sea-surface temperature observations since 1850
1078 part 2: biases and homogenisation. *J. Geophys. Res.*, 116, D14104, 2011b.

1079 Kunstmann, H., Jung, G.: Influence of soil-moisture and land use change on
1080 precipitation in the ~~volta~~Volta basin of West Africa, *International Journal of*
1081 *River Basin Management*, 5, 9-16, 2007.

1082 Lambert, F. H., Chiang, J.C.H.: Control of land-ocean temperature contrast by ocean
1083 heat uptake, *Geophys. Res. Lett.*, 34, L13704, 2007.

1084 Laua, W.K.M., Kim, K.-M.: Robust Hadley circulation changes and increasing global
1085 dryness due to CO2 warming from CMIP5 model projections. *Proceedings of*
1086 *the National Academy of Sciences of the United States of America*, 112, 3630-
1087 3635, 2015.

1088 Lorenz, D.J., DeWeaver, E.T.: The response of the extratropical hydrological cycle to
1089 global warming, *Journal of Climate*, 20, 3470-3484, 2007.

1090 Marengo, J.A., Nobre, C.A., Tomasella, J. et al.: The drought of Amazonia in 2005,
1091 *Journal of Climate*, 21, 495-516, 2008.

1092 Martens, B., Miralles, D.G., Lievens, H., van der Schalie, R., de Jeu, R.A.M.,
1093 Fernández-Prieto, D., Beck, H.E., Dorigo, W.A., and Verhoest, N.E.C.: GLEAM
1094 v3: satellite-based land evaporation and root-zone soil moisture, Geoscientific
1095 Model Development Discussions,
1096 McCarthy, M.P., Toumi, R.: Observed interannual variability of tropical troposphere
1097 relative humidity, *Journal of Climate*, 17, 3181-3191, 2004.
1098 McCarthy, M.P., Thorne, P.W., Titchner, H.A.: An analysis of tropospheric humidity
1099 trends from radiosondes, *Journal of Climate*, 22, 5820-5838, 2009.
1100 McVicar, T.R., Roderick, M.L., Donohue, R.J., Van Niel, T.G.: Less bluster ahead?
1101 Ecohydrological implications of global trends of terrestrial near-surface wind
1102 speeds, *Ecohydrology*, 5, 381–388, 2012a.
1103 McVicar, T.R., Roderick, M.L., Donohue, R.J., et al.: Global review and synthesis of
1104 trends in observed terrestrial near-surface wind speeds: implications for
1105 evaporation, *J. Hydrol.*, 416–417, 182–205, 2012b.
1106 Miralles, D.G., Holmes, T.R.H., de Jeu, R.A.M., Gash, J.H., Meesters, A.G.C.A.,
1107 Dolman, A.J.: Global land-surface evaporation estimated from satellite-based
1108 observations, *Hydrology and Earth System Sciences*, 15, 453–469, 2011.
1109 Miralles, D.G., Nieto, R., McDowell, N.G., Dorigo, W.A., Verhoest, N.E.C., Liu, Y.Y.,
1110 Teuling, A.J., Dolman, A.J., Good, S.P., Gimeno, L.: Contribution of water-
1111 limited ecoregions to their own supply of rainfall, *Environmental Research*
1112 *Letters*, 11, 124007, 2016.
1113 Nieto, R., Castillo, R., Drumond, A., Gimeno, L.: A catalog of moisture sources for
1114 continental climatic regions, *Water Resources Research*, 50, 5322-5328, 2014.
1115 ~~Numaguti, A.: Origin and recycling processes of precipitating water over the Eurasian~~
1116 ~~continent: experiments using an atmospheric general circulation model. *J.*~~
1117 ~~*Geophys. Res. Atmos.*, 104, 1957–1972, 1999.~~
1118 O’Gorman, P.A., Muller, C.J.: How closely do changes in surface and column water
1119 vapor follow Clausius-Clapeyron scaling in climate change simulations?
1120 *Environmental Research Letters*, 5, 025207, 2010.
1121 Oki, T.: The hydrologic cycles and global circulation. *Encyclopedia of Hydrological*
1122 *Sciences*, 13-22. 2005.
1123 Pfahl, S., Madonna, E., Boettcher, M., Joos, H., Wernli, H.: Warm conveyor belts in the
1124 ERA-Interim Dataset (1979-2010). Part II: Moisture origin and relevance for
1125 precipitation, *Journal of Climate*, 27, 27-40, 2014.
1126 Pierce, D.W., Westerling, A.L., Oyler, J.: Future humidity trends over the western
1127 United States in the CMIP5 global climate models and variable infiltration
1128 capacity hydrological modeling system. *Hydrology and Earth System Sciences*,
1129 17, 1833-1850, 2013.
1130 ~~Rayner, N.A., Parker, D.E., Horton, E.B. et al.: Global analyses of sea surface~~
1131 ~~temperature, sea ice, and night marine air temperature since the late nineteenth~~
1132 ~~century. *Journal of Geophysical Research D: Atmospheres*, 108, ACL 2-1–ACL~~
1133 ~~2-29, 2003.~~
1134 Rebetez, M., Mayer, H., Dupont, O., et al.: Heat and drought 2003 in Europe: A
1135 climate synthesis, *Annals of Forest Science*, 63, 569-577, 2006.
1136 Rodell, M., Beaudoin, H.K., L’Ecuyer, T.S. et al.: The observed state of the water
1137 cycle in the early twenty-first century, *Journal of Climate*, 28, 8289-8318,
1138 2015.
1139 Roderick, M., Farquhar, G.D.: The cause of decreased pan evaporation over the past 50
1140 years. *Science*, 15, 1410–1411, 2002.

1141 Rowell, D.P., Jones, R.G.: Causes and uncertainty of future summer drying over
1142 Europe, *Climate Dynamics*, 27, 281-299, 2006.

1143 Seneviratne, S.I., Pal, J.S., Eltahir, E.A.B., Schar, C.: Summer dryness in a warmer
1144 climate: A process study with a regional climate model, *Climate Dynamics*, 20,
1145 69-85, 2002.

1146 Seneviratne, S.I., Lüthi, D., Litschi, M., Schär, C.: Land-atmosphere coupling and
1147 climate change in Europe, *Nature*, 443, 205-209, 2006.

1148 Shenbin, C., Yunfeng, L., Thomas, A.: Climatic change on the Tibetan Plateau:
1149 Potential evapotranspiration trends from 1961-2000, *Climatic Change*, 76, 291-
1150 319, 2006.

1151 Sherwood, S.C.: Maintenance of the free-tropospheric tropical water vapor distribution.
1152 Part I: Clear regime budget, *Journal of Climate*, 9, 2903-2918, 1996.

1153 Sherwood, S.C.: Direct versus indirect effects of tropospheric humidity changes on the
1154 hydrologic cycle, *Environmental Research Letters*, 5, 025206, 2010.

1155 Sherwood, S., Fu, Q.: A drier future?, *Science*, 343, 737-739, 2014.

1156 Simmons, A.J., Willett, K.M., Jones, P.D., Thorne, P.W., Dee, D.P.: Low-frequency
1157 variations in surface atmospheric humidity, temperature, and precipitation:
1158 inferences from reanalyses and monthly gridded observational data sets. *J.*
1159 *Geophys. Res .D Atmos.*, 115, D01110, 2010.

1160 Stephenson, N.L. Climatic control of vegetation distribution: the role of the water
1161 balance. *Am. Nat.*, 135, 649-670, 1990.

1162 Stohl, A. and James, P.: A Lagrangian Analysis of the Atmospheric Branch of the
1163 GlobalWaterWater Cycle. Part I: Method Description, Validation and
1164 Demonstration for the August 2002 Flooding in Central Europe, *J.*
1165 *Hydrometeorol.*, 5, 656-678, 2004.

1166 Stohl, A., James, P.: A Lagrangian analysis of the atmospheric branch of the global
1167 water cycle. II. Moisture transports between Earth's ocean basins and river
1168 catchments, *J. Hydrometeorol.*, 6, 961-984, 2005.

1169 Trenberth, K.E. Atmospheric moisture recycling: Role of advection and local
1170 evaporation. *Journal of Climate* 12, 1368-1381. 1999.

1171 Trenberth, K.E., Fasullo, J., Smith, L.: Trends and variability in column-integrated
1172 atmospheric water vapor, *Climate Dynamics*, 24, 741-758, 2005.

1173 Trenberth, K.E., Smith, L., Qian, T., Dai, A., Fasullo, J.: Estimates of the global water
1174 budget and its annual cycle using observational and model data, *J.*
1175 *Hydrometeorol.*, 8, 758-769, 2007.

1176 Van Der Ent, R.J., Savenije, H.H.G., Schaeffli, B., Steele-Dunne, S.C. Origin and
1177 fate of atmospheric moisture over continents, *Water Resources Research*, 46,
1178 W09525, 2010.

1179 Van Wijngaarden, W.A., Vincent, L.A.: Trends in relative humidity in Canada from
1180 1953-2003, *Bulletin of the American Meteorological Society*, 4633-4636, 2004.

1181 ~~Vautard, R., Yiou, P., D'Andrea, F., et al. Summertime European heat and drought~~
1182 ~~waves induced by wintertime Mediterranean rainfall deficit ——— *Geophysical*~~
1183 ~~*Research Letters* 34(7),L07711. 2007.~~


1184 ~~Vautard, R., Gobiet, A., Jacob, D., et al. The simulation of European heat waves from~~
1185 ~~an ensemble of regional climate models within the EURO-CORDEX project.~~
1186 ~~*Climate Dynamics*, 41: 2555-2575. 2013.~~

1187 Vázquez, M., Nieto, R., Drumond, A., Gimeno, L.: Moisture transport into the Arctic:
1188 Source-receptor relationships and the roles of atmospheric circulation and
1189 evaporation, *Journal of Geophysical Research: Atmospheres*, 121, 493-509,
1190 2016.

- 1191 Vicente-Serrano, S.M., Azorin-Molina, C., Sanchez-Lorenzo, A., Revuelto, J., Morán-
 1192 Tejada, E., López-Moreno, J.I., Espejo, F.: Sensitivity of reference
 1193 evapotranspiration to changes in meteorological parameters in Spain (1961–
 1194 2011), *Water Resources Research*, 50, 8458–8480, 2014a.
- 1195 Vicente-Serrano, S.M., Azorin-Molina, C., Sanchez-Lorenzo, A., Morán-Tejada, E.,
 1196 Lorenzo-Lacruz, J., Revuelto, J., López-Moreno, J.I., Espejo, F.: Temporal
 1197 evolution of surface humidity in Spain: recent trends and possible physical
 1198 mechanisms. *Climate Dynamics*, 42, 2655–2674, 2014b.
- 1199 Vicente-Serrano, S.M., Azorin-Molina, C., Sanchez-Lorenzo, A., El Kenawy, A.,
 1200 Martín-Hernández, N., Peña-Gallardo, M., Beguería, S., Tomas-Burguera, M.:
 1201 Recent changes and drivers of the atmospheric evaporative demand in the
 1202 Canary Islands. *Hydrology and Earth System Science*, 20, 3393-3410, 2016.
- 1203 Vincent, L.A., van Wijngaarden, W.A., Hopkinson, R.: Surface temperature and
 1204 humidity trends in Canada for 1953-2005, *Journal of Climate*, 20, 5100-5113,
 1205 2007.
- 1206 Wang, K., Dickinson, R.E.: A review of global terrestrial evapotranspiration:
 1207 observation, modeling, climatology, and climatic variability, *Rev. Geophys.*, 50,
 1208 2012.
- 1209 Wang, K., Dickinson, R.E., Liang, S.: Global atmospheric evaporative demand over
 1210 land from 1973 to 2008, *J. Clim.*, 25, 8353–8361, 2012.
- 1211 Wei, J., Su, Hua, Zong-Liang Yang. Impact of moisture flux convergence and soil
 1212 moisture on precipitation: a case study for the southern United States with
 1213 implications for the globe. *Climate Dynamics* 46: 467–481. 2016a.
- 1214 Wei, J., Q. Jin, Z.-L. Yang, and Pand P. A. Dirmeyer. Role of ~~ocean~~evaporation
 1215 evaporation in California droughts ~~and floods~~and floods. *Geophys. Res. Lett.*, 43,
 1216 6554–6562, doi:10.1002/2016GL069386. 2016b.
- 1217 Willett, K.M., Jones, P.D., Gillett, N.P., Thorne, P.W.: Recent changes in surface
 1218 humidity: Development of the HadCRUH dataset, *Journal of Climate*, 21, 5364-
 1219 5383, 2008.
- 1220 Willett, K.M., Dunn, R.J.H., Thorne, P.W., Bell, S., de Podesta, M., Parker, D.E., Jones,
 1221 P.D., Williams Jr., C.N.: HadISDH land surface multi-variable humidity and
 1222 temperature record for climate monitoring, *Clim. Past*, 10, 1983-2006, 2014.
- 1223 Wright, J.S., Sobel, A., Galewsky, J.: Diagnosis of relative humidity changes in a
 1224 warmer climate, *J. Clim.*, 23, 4556-4569, 2010.
- 1225 Yu, L.: Global variations in oceanic evaporation (1958-2005): The role of the changing
 1226 wind speed. *J. Climate*, 20, 5376–5390, 2007.
- 1227 Yu, L., Jin, X., Weller, R.A.: Multidecade Global Flux Datasets from the Objectively
 1228 Analyzed Air-sea Fluxes (OAFlux) Project: Latent and sensible heat fluxes,
 1229 ocean evaporation, and related surface meteorological variables. Woods Hole
 1230 Oceanographic Institution, OAFlux Project Technical Report. OA-2008-01,
 1231 64pp. Woods Hole. Massachusetts. 2008.
- 1232 Yue, S., Wang, C.: The Mann-Kendall Test Modified by Effective Sample Size to
 1233 Detect Trend in Serially Correlated Hydrological Series, *Water Resour.*
 1234 *Manage.*, 18, 201–218, 2004.
- 1235 Zahn, M. and R. P. Allan, Quantifying present and projected future atmospheric
 1236 moisture transports onto land, *Water Resources Research*. 49, p.7266-
 1237 7277,doi:10.1002/2012WR013209. 2013.
- 1238 Zhang, X., Harvey, K. D., Hogg, W. D., and Yuzyk, T. R.: Trends in Canadian
 1239 streamflow, *Water Resour. Res.*, 37, 987–998, 2001.

Con formato: Inglés (Estados Unidos)

1240 Zhang, L., Wu, L., Gan, B.: Modes and mechanisms of global water vapor variability
1241 over the twentieth century, *Journal of Climate*, 26, 5578-5593, 2013.
1242 Zongxing, L., Qi, F., Wei, L., Tingting, W., Yan, G., Yamin, W., Aifang, C., Jianguo,
1243 L., Li, L.: Spatial and temporal trend of potential evapotranspiration and related
1244 driving forces in Southwestern China, during 1961-2009, *Quaternary*
1245 *International*, 336, 127-144, 2014.
1246
1247
1248
1249
1250
1251




1252

1253

1254

Figure 1. Annual and seasonal averages of RH and Vertically Integrated Moisture Flux (VIMF) based on ERA-Interim dataset.



1255

1256

Figure

2.




Fig. 1. Spatial distribution of the magnitude of change of RH (% per decade) over the period 1979-2014 from HadISDH (left) and ERA-Interim [dataset](#) (right). Results are provided for the boreal cold (October-March) and warm (April-September) seasons and annually.




Figure 2: Relative frequencies (%) of the RH magnitude of change in the HadISDH and ERA-Interim datasets. Color bar plots represent the percentage of stations (from HadISDH) and world regions (from ERA-Interim) with **positive and statistically significant (positive trends at $p < 0.05$) trends** (blue), **positive statistically insignificant positive trends** (cyan), **negative statistically insignificant negative trends** (orange) and **negative and statistically significant negative trends** (red).

Con formato: Fuente: 14 pto




Figure 4: Scatterplots showing the global relationship between the magnitude of change in RH with HadISDH stations and ERA-Interim dataset at the seasonal and annual scales. Colors represent the density of points, with red color showing the highest density of points.




Figure 3: Spatial distribution of the seasonal and annual magnitudes of change in specific humidity (g/kg^{-1}) (left) and the deficit/surplus of specific humidity to maintain the RH constant with the levels of 1979 according to the land air temperature evolution (from the CRU TS v.3.23 dataset) for 1979-2014.




Figure 4: Spatial distribution of the **selected** 14 world regions, **with based on the** high consistency in RH trends between the HadISDH and the ERA-Interim datasets. These regions were selected for the identification of the oceanic and land humidity sources by means of the FLEXPART scheme.





Figure 5: Top left: Annual RH humidity trends in the WestWestern Sahel (region 6), Top right: average (E-P) > 0 at the annual scale to identify the main humidity sources in the region (mm_{year}⁻¹). Center: Relationships between the de-trended annual RH and the de-trended annual variables for 1979-2014. Bottom: Annual evolution of the different variables corresponding to the WestWestern Sahel region. The

magnitude of change and their corresponding statistical significance of the trend is are indicated for each variable.




Figure 8 Fig. 6: The same as Fig. 75, but for the La Plata (region 9).




Figure 9 Fig. 7: The same as Fig. 75, but for the West North of Northern America (region 12).




Figure 10: Relationships between the annual oceanic contribution to annual precipitation (E-P) and the annual RH in the target regions.




Figure 11: Relationship between the annual land contribution to annual precipitation (E-P) and the annual RH in the target regions.




Figure 10: Evolution of the land contribution (%) to annual precipitation (%) in the different target regions




Figure 13: Relationships between the average annual **magnitudemagnitudes** of change in RH identified **infor** each one of the 14 analyzed regions and the annual **magnitudemagnitudes** of change in precipitation, the ratio between air temperature/SST, oceanic evaporation and land evapotranspiration.




Figure 14: Spatial distribution of the Pearson's r correlations between the detrended RH and land evapotranspiration series at the annual and seasonal time scales. The statistical significance of the correlations is also shown.




Figure 12: Spatial distribution of the magnitude of change in the annual and seasonal land evapotranspiration (1979-2014) and their corresponding statistical significance of trends.




Figure 16 Fig. 13: Annual and seasonal magnitude of change of SST and OAFLUX oceanic evaporation for 1979-2014.