

Interactive comment on “Thermodynamics of Saline and Fresh Water Mixing in Estuaries” by Zhilin Zhang and Hubert H. G. Savenije

Zhilin Zhang and Hubert H. G. Savenije

h.h.g.savenije@tudelft.nl

Received and published: 7 December 2017

We would like to thank referee #1 for the discussion. Below we reply to the first and third comment. The second comment about the derivation of the differential equation requires an illustration and equations, which we have put in the Supplement.

1) About the maximum and minimum. The theory in this paper follows the principle of Maximum Entropy Production which states that certain complex non-equilibrium thermodynamic systems can be successfully characterised as being in states in which the rate of thermodynamic entropy production is maximised (e.g., Kleidon, 2016). Mixing processes in estuaries continuously perform work by depleting salinity gradients. In doing so, these processes produce entropy, following the natural direction given by

C1

the second law of thermodynamics. So we assume the power to mix is maximized to maximize the entropy production in this system.

3) About the sloping or flat bed. In fact, the slope is small so the increase of depth due to the salinity difference (Δh) is trivial compared to the depth at the estuarine mouth (h) (but Δh is important for the mixing).

If, however, a downward slope were introduced in the picture, then we would have to include the bottom pressure as well: the water pressure over the additional depth near the downstream boundary would have to be balanced by the horizontal component of the pressure exercised on the water by the estuary bottom. This would make the sketch unnecessary complex.

Please also note the supplement to this comment:

<https://www.earth-syst-dynam-discuss.net/esd-2017-87/esd-2017-87-AC1-supplement.pdf>

Interactive comment on Earth Syst. Dynam. Discuss., <https://doi.org/10.5194/esd-2017-87>, 2017.

C2