

Interactive comment on “Interannual variability of the gravity wave drag – vertical coupling and possible climate links” by Petr Sacha et al.

Petr Sacha et al.

petr@pisoft.cz

Received and published: 21 February 2018

Responses to the referee’s comment on paper “*Interannual variability of the gravity wave drag - vertical coupling and possible climate links*” by Petr Sacha, Jiri Miksovsky, and Petr Pisoft.

We would like to thank the referee for taking the time to review our manuscript. We greatly appreciate the insightful and constructive comments, which we address in our responses below.

Printer-friendly version

Discussion paper



General comments:

1) *The study is focusing on orographic gravity wave drag, which is directly provided by the OGWD parameterization of the CMAM model. However, non-orographic sources such as convection or jet and storm sources are another important source of gravity wave drag. It is pointed out that the OGWD parameterization of the CMAM-sd simulation was "tuned" to obtain more realistic circulation patterns. Does this "tuning" overemphasize the role of orographic gravity waves compared with non-orographic sources? If non-orographic sources are neglected (as I understand), how does this affect the analysis presented in this paper?*

Thank you for this comment; it is a very good point. The settings of the OGWD and NOGWD parameterization scheme used in the CMAM-sd simulation are discussed in details in McLandress et al. (2013) and we will add a paragraph in the revised manuscript to make a clear summary discussing limitations connected to our results.

The dynamical role of the OGWD relative to the NOGWD at the vertical domain of our analysis is most likely overestimated compared to the current consensus on the GW impacts on the stratosphere. We will stress this out more in the revised version together with statement that our conclusions are directly applicable at the model atmosphere only and thus having indirect implications for the real atmosphere (as already stated in the current version of the discussion).

It would be very interesting to look at NOGWD variations connected with variability of jets, fronts etc. However, the CMAM NOGWD scheme (Scinocca, 2003) is based on launching a globally uniform isotropic NOGW spectrum in four cardinal horizontal directions at approximately 125 hPa. The aim is to produce reasonable seasonal evolution of the zonal mean zonal temperature and winds in the mesosphere and the zonal and merid-

ional asymmetry stems from propagation effects only. Regarding NOGWD, we have produced the same analysis as for the OGWD but due to the above mentioned reasons the resulting fields are highly zonally symmetrical and weaker in magnitude compared to the OGWD and so we decided not to show them in the manuscript. However, we attach selected figures as a supplement to this response (see Fig. 1 below).

2) It would be good if this work could be put better into the context of related work. There is a number of studies discussing global climatologies of gravity wave activity in the stratosphere from observations and models, e.g. ... Geller et al. (2013) showed that there are notable differences between momentum flux estimates from different models and observations. It might be good to provide more evidence that the results from the CMAM-sd simulation are realistic. Perhaps it might be helpful to also show gravity wave momentum flux distributions from the simulation, as this can be more easily compared to other studies.

We thank the referee for pointing out additional studies that can improve our assessment of realism of the CMAM-sd OGWD distribution and also we thank for the idea of comparing the momentum fluxes. We will adopt that and the results will be discussed in the revised version of the paper.

Specific comments:

1) Be more specific about what is meant by "lower tropospheric behavior"?

The OGWD variability is shown to be induced by lower tropospheric wind variations in a large part. There was also significant variability detected in near surface OGW momentum fluxes.

[Printer-friendly version](#)[Discussion paper](#)

2) *What is meant by "have a modified impact"? Do you mean "have a modifying impact on" or simply "have impact on"?*

We meant that there were modifications of the GWs impact. The statement will be adjusted in the following way.

We argue that the orographic gravity waves (OGWs) and GWs in general can be a quick mediator of the tropospheric variability into the stratosphere as the modifications of the OGWD distribution can result in different impacts on the stratospheric dynamics during different phases of the studied climate oscillations.

3) *This first sentence is quite long. The references to Plougonven and Zhang (2014) and Alexander et al. (2009) look a bit specific considering the broad statements made here.*

The statement will be adjusted in the following way.

Although the gravity wave (GW) sourcing (e.g. adjustment processes, Plougonven and Zhang, 2014), propagation and breaking is governed to some extent by processes in the stratosphere, there is a significant portion of the IGW spectra created in the troposphere (mostly orography and convection, Alexander et al., 2009). The highest amplitude upward propagating modes can break already in the troposphere and lower or middle stratosphere (Fritts et al., 2016).

- Fritts, D.C., R.B. Smith, M.J. Taylor, J.D. Doyle, S.D. Eckermann, A. Dörnbrack, M. Rapp, B.P. Williams, P. Pautet, K. Bossert, N.R. Criddle, C.A. Reynolds, P.A. Reinecke, M. Uddstrom, M.J. Revell, R. Turner, B. Kaifler, J.S. Wagner, T. Mixa, C.G. Kruse, A.D. Nugent, C.D. Watson, S. Gisinger, S.M. Smith, R.S. Lieberman, B. Laughman, J.J.

Printer-friendly version

Discussion paper



Moore, W.O. Brown, J.A. Haggerty, A. Rockwell, G.J. Stossmeister, S.F. Williams, G. Hernandez, D.J. Murphy, A.R. Klekociuk, I.M. Reid, and J. Ma, 2016: The Deep Propagating Gravity Wave Experiment (DEEPWAVE): An Airborne and Ground-Based Exploration of Gravity Wave Propagation and Effects from Their Sources throughout the Lower and Middle Atmosphere. Bull. Amer. Meteor. Soc., 97, 425–453, <https://doi.org/10.1175/BAMS-D-14-00269.1>

4) *How strong was the nudging? Does the CMAM-sd simulation closely follow the ERA-Interim winds and temperatures? Are the results of this study sensitive to the specific details/parameters of the nudging procedure?*

The nudging procedure is examined in detail in McLandress et al. (2014) and described in McLandress et al. (2013). Below 1 hPa, in spectral space (for horizontal scales with wavenumber lower than 21), CMAM is nudged to the 6 hourly horizontal winds and temperatures from ERA Interim. This would definitely be a problem, if we would analyze the GWD impact on the circulation. This impact is weakened by the relaxation. But, as we have only analyzed the OGWD interannual variability, the nudging procedure is very advantageous for us (compared to a free running model). Since the winds and temperatures in the lower atmosphere have a strong influence on the propagation and absorption of gravity waves the distribution of parameterized (and resolved) gravity wave fluxes in CMAM-sd can match the real fluxes (McLandress et al., 2013). This is true, of course, having the tuning and other specifics of GWD parameterization in mind.

5) *It might be worthwhile to briefly repeat/recap the definitions of the different indices?*

The definitions together with description of the utilized data will be added in the revised version of the paper.

Printer-friendly version

Discussion paper



6) *Is this exception of the Antarctic Peninsula due to its SW-NE orientation?*

We would not say an exception in general (only the strongest) but we agree with the referee that the topography orientation (relative to surface winds) is important. The orientation of the momentum flux in the OGWD parametrization in CMAM is a function of the near-surface wind direction relative to the orientation of the topography (McLandress et al., 2013). In Sacha et al. (2016) we highlighted the fact that the dynamical effects of the meridional GWD component do not receive a sufficient attention and are quite uncertain to date.

- Sacha, P., Lilienthal, F., Jacobi, C., and Pisoft, P.: Influence of the spatial distribution of gravity wave activity on the middle atmospheric dynamics, *Atmos. Chem. Phys.*, 16, 15755-15775, <https://doi.org/10.5194/acp-16-15755-2016>, 2016.

7) *The degree of correlation seems to decrease with height? Is this due to the stratospheric background affecting the propagation of the waves?*

Yes, this is due to the background. It is well manifested in the eastern Asia region where this hotspot is stronger at 50 hPa and when the stratospheric conditions are not advantageous for breaking at 50 hPa there is more breaking at 30 hPa. This results in the OGWD at 30hPa in this region having less correlation with tropospheric conditions because the stratospheric variability between 50 and 30 hPa is important.

But this is not the case for the Scandinavian hotspot that is climatologically strongest at 10 hPa and the fraction explained by near surface winds is lower at 50 and 30 hPa. At those levels the waves dissipate less frequently - predominantly in dependence on the stratospheric variability.

Printer-friendly version

Discussion paper



Based on the referee comment, we will add following statement to the manuscript:

p719 This is due to the stratospheric background affecting the critical line occurrence and propagation of the GWs between 50 and 30 hPa in the eastern Asia region.

8) *Reading this, I was wondering how well the CMAM-sd simulation itself captures the different climatological patterns (NAO, SO, QBO)?*

While there are some differences between the time series of CMAM-based and observational counterparts, their similarity is generally strong, and the Pearson correlation (i.e., a correlation measure best suited for quantifying the links explored by linear regression analysis) between them is high. See Fig. 2 below for an example of temporal variability of the CMAM-simulated and observational indices of the Southern Oscillation.

9) *This also triggers the question of how well the CMAM-sd simulation reflects reality?*

As with all model-based frameworks, the issue of reliable reproduction of the observed climate is quite complex and many statistics can be considered for validation. Our tests concentrated primarily on the ability of CMAM to realistically reproduce the spatial patterns of response of lower tropospheric characteristics to the phases of the oscillation considered in our analysis (NAO, SO/ENSO, QBO); the results suggested a high degree of match between the CMAM-based and observational data (see Fig. 3 below for a sample of the results, summarizing the 850 hPa temperature response to SO and NAO).

[Printer-friendly version](#)[Discussion paper](#)

10) *Recent studies showed that there might be notable gravity wave activity over remote islands in the Southern Ocean, e.g., South Georgia or Kerguelen Islands. Is the CMAM-sd simulation capable of capturing this?*

In the Figure 1, 50 hPa, DJF (!) we can see that there is some moderately strong significant OGWD over those remote islands. However, we must note that those are monthly mean values only, not reflecting the intermittent nature of the GWs. Our study is focused globally, but the fact that CMAM is able to get some pronounced OGWD above such small islands is good news for possible future research of individual hotspots.

- Alexander, M. J., and A. W. Grimsdell (2013), Seasonal cycle of orographic gravity wave occurrence above small islands in the Southern Hemisphere: Implications for effects on the general circulation, *J. Geophys. Res. Atmos.*, 118, 11,589–11,599, doi:10.1002/2013JD020526.
- Hoffmann, L., Grimsdell, A. W., and Alexander, M. J.: Stratospheric gravity waves at Southern Hemisphere orographic hotspots: 2003–2014 AIRS/Aqua observations, *Atmos. Chem. Phys.*, 16, 9381–9397, <https://doi.org/10.5194/acp-16-9381-2016>, 2016.

Interactive comment on Earth Syst. Dynam. Discuss., <https://doi.org/10.5194/esd-2018-1>, 2018.

Printer-friendly version

Discussion paper






Fig. 1. Response of the non-orographic GWD [m/s/s] at the 50 hPa level related to the activity of the Southern Oscillation (left), North Atlantic Oscillation (right) and Quasi-Biennial Oscillation (bottom)

[Printer-friendly version](#)[Discussion paper](#)




Fig. 2. Southern Oscillation index calculated as a normalized sea level pressure difference between Darwin and Tahiti, derived from CMAM data (red line) and from direct observations (blue; CRU)

[Printer-friendly version](#)[Discussion paper](#)




Fig. 3. 850 hPa temperature response to SO (top) and NAO (down) change from highly negative to highly positive phase, evaluated through multiple linear regression.