

Anonymous Referee #1

We thank the reviewer for all his/her comments which improved the manuscript substantially. We want to thank you for the references provided in the review. Please, find below the response to your comments, questions, and suggestions.

- In the abstract, you mention atmospheric rivers but then there is no further mention of them in the text. Is it really needed to mention them there? - In the abstract we had mentioned the atmospheric rivers (ARs) because together with low-level jets (LLJ) are the main structures that are associated with a transport of high atmospheric moisture and can trigger episodes of heavy rains or even floods. We agree with the reviewer that this may be misplaced and we removed the reference in the latest version of the manuscript.

- There have been studies looking at atmospheric rivers and precipitation focusing on specific events across the central United States. The papers by Moore et al. (2012) and Nayak et al. (2016) are likely worth mentioning. – Both references proposed by the reviewer will be added to the introduction.

- More broadly, there have been a growing body of work related to moisture transport over the central United States (e.g., Nakamura et al. 2013; Lavers and Villarini 2015; Steinschneider and Lall 2015, 2016; Nayak and Villarini 2017). – Most of the references proposed by the reviewer will be included in the latest version of the manuscript.


- Pg. 2, line 17: “Higgins et al. (1996)” – The typo has been corrected.

- Page 2, line 34: “which modulate a” – The typo has been corrected.

- Page 3, lines 17-18: “total amount of total precipitable water” seems a bit redundant. What about “amount of total precipitable water”? The same applies to other places in the text. – The text has been updated following the suggestion of the reviewer.

- Page 3, line 20: “as follows: in” – The typo has been corrected.

- Section 2.1: why is the focus only on July and not on June and August as well? Please clarify. - The Great Plains low-level jet (GPLLJ) is a phenomenon mostly related to the warm season. We have focused the study in the month of July because it is the month of the year when we find the highest frequency. Figure attached below shows the monthly distribution of GPLLJ detections.



- Page 3, line 24: “Rife et al. (2010)” – The typo has been corrected in the new version of the manuscript.

- Page 3, line 29: is there an impact on setting a threshold in terms of these differences? As of now, the only requirement is that the wind speed is higher at midnight compared to midday. What happens if you set a threshold, say 10% higher at midnight compared to midday? Would it be possible to have information related to the distribution of the differences between them?

The index is based on the temporal variation of the vertical wind structure. To obtain a day of GPLLJ it is necessary that two conditions are simultaneously met:

1. The wind speed is higher at midnight than at local noon.
2. The wind speed on the surface is higher than at high levels.

We do not believe it is adequate to apply a threshold in the detection of LLJ. The fact of establishing a threshold in one of the conditions would add subjectivity to the methodology used in the study. Nonetheless, if we set the arbitrary threshold of 10% in one of the conditions, the climatology however hardly changes. We have performed the calculation and applying the 10% threshold at midnight we identified 924 days of GPLLJ. However, without applying the threshold we get 931 days of GPLLJ (7 cases of difference). Have or not a day of GPLLJ is mainly due to the fulfilment of both initial conditions. Setting random thresholds only in one condition adds subjectivity to the methodology used in the study.

- Page 3, line 30: “than above it (~4km)” I still don’t think it is the right wording but it sounds a bit better than what is there. – Following the reviewer’s suggestion, the sentence now reads as follow: *The local midnight wind speed is higher at the surface (~500m) than above it (~4km).*

- Page 3, line 32: “Because of the jet”.

- Page 4, line 9: “variation of 5 the wind” is not clear.

- Page 4, line 17: “possible to calculate”.

- Page 4, line 30: “followed for 10”.


- Page 5, line 12: “above the 75th percentile”.

- Page 5, line 16: “is used to quantify the”.

The typos detected by the reviewer have been corrected in the latest version of the manuscript.

- Is it possible to show some results related to the validation of the WRF model with respect to observations?

We understand the reviewer is asking for some validation or comparison with observed precipitation, which “tricky” variable to solve by all models, including WRF. Figure attached below compares 11-days accumulated precipitation for WRF simulations versus CPC gauge-analysis observations (<https://www.esrl.noaa.gov/psd/data/gridded/data.unified.daily.conus.html>) throughout the same periods. As the reviewer can note, WRF tends to slightly overestimate the precipitation, but it is in all respects represented quite well, both in amount and field distribution. This figure will be added to the supplementary material.



- Page 6, line 1: the symbol ϕ for the instant flux of moisture was already used in equation 3. – “ ϕ ” has been replaced by “ σ ” in the latest version of the manuscript.

- Page 6, line 16: the use of the correlation coefficient is not appropriate. Please use the Lilliefors test to test whether the data can be described by a Gaussian distribution.

Another option is the Jarque-Bera test.

We agree with the referee on the fact that the correlation coefficient is not an appropriate indicator of the normality of the distribution. Thus, we have applied the Jarque-Bera test which provided a p-value equal to 0.0055. We understand that this p-value is low enough for considering the LLJ distribution as Gaussian.

Accordingly, “A clear peak around 11 m s^{-1} is observed together with a Gaussian behaviour ($R^2 = 0.95$, red line)” has been replaced by:

“A clear peak around 11 m s^{-1} is observed together with a Gaussian behaviour (Jarque-Bera test p-value=0.0055, which provides a confidence level close to 99.5 %, red line)”

- Page 6, line 19: “spans 11 days”.

- I would remove the equation from the caption of Figure 1.

- Pg. 7, line 5: shouldn't this be Figure 2 instead of Figure 1?

The manuscript has been updated following the Reviewer's suggestions.

- Page 7, line 27: why not computing the climatology using all the days, rather than based on just a handful? - The calculation of the climatology using every day of GPLLJ would increase the computational cost in an excessive way. The methodology used in this work, especially the Eulerian model WRF, has a high computational cost. Besides, this multiply the number of simulations making this work unaffordable. In addition, the aim to this study is to quantify the average transport of moisture in a general perspective of the GPLLJ's behaviour. Fig. 4 shows an approximation to this result. This figure is calculated based on the statistical weight of each simulation of the 5 GPLLJ events analysed. Thus, this figure can be understood as a climatological approach to the moisture transport associated with the GPLLJ.

References:

1. Lavers, D.A., and G. Villarini, The contribution of atmospheric rivers to precipitation in Europe and the United States, *Journal of Hydrology*, 522, 382-390, 2015.
2. Moore, B.J., P.J. Neiman, F.M. Ralph, and F.E. Barthold, Physical processes associated with heavy flooding rainfall in Nashville, Tennessee, and vicinity during 1–2 May 2010:

- The role of an atmospheric river and mesoscale convective systems, *Monthly Weather Review*, 140, 358–378, 2012.
3. Nakamura, J., U. Lall, Y. Kushnir, A.W. Robertson, and R. Seager, Dynamical structure of extreme floods in the U.S. Midwest and the United Kingdom, *Journal of Hydrometeorology*, 14, 485–504, 2013.
 4. Nayak, M.A., G. Villarini, and A.A. Bradley, Atmospheric rivers and rainfall during NASA's Iowa Flood Studies (IFloodS) campaign, *Journal of Hydrometeorology*, 17(1), 257–271, 2016.
 5. Nayak, M.A, and G. Villarini, A long-term perspective of the hydroclimatological impacts of atmospheric rivers over the central United States, *Water Resources Research*, 53, 1144–1166, 2017.
 6. Steinschneider, S., and U. Lall, Daily precipitation and tropical moisture exports across the Eastern United States: An application of archetypal analysis to identify spatiotemporal structure, *Journal of Climate*, 28(21), 8585–8602, 2015.
 7. Steinschneider, S., and U. Lall, Spatiotemporal structure of precipitation related to tropical moisture exports over the eastern United States and its relation to climate teleconnections, *Journal of Hydrometeorology*, 17(3), 897–913, 2016.

Anonymous Referee #2

We thank the reviewer for all his/her questions and suggestions, which improved the manuscript substantially. Please, find attached below the one-to-one reply to the comments.

Comment 1:

1) The FLEXPART analysis, to my understanding, is largely underutilized here. Is it only to justify the location of the wall line? I think the wall line location is fairly intuitive and I don't believe the authors would find much sensitivity to its location (within reasonable limits). As a minimum, I would encourage the authors to include the FLEXPART-derived moisture source regions for each case study in Supplemental Material.

We appreciate the comment and the figure of moisture sources for each case study will be added to the supplementary material. Answering the question, the lagrangian Flexpart model was used to objectively justify the position of the wall. Fig. RC2.1 (in this document) shows the 75th percentile the sources of moisture ($E - P > 0$) for the 6 cases of study. The purple line shows the position of the wall of moisture labelling. The vectors refer to the wind for each case study and are plotted at surface level (aprox. 500m). As it is observed, moisture enters the Great Plains in a channelized way through the wall from the Gulf of Mexico. In the Fig. RC2.1a (no-LLJ day) the vectors do not show the flow of the South. Therefore, the transport of moisture from the Gulf of Mexico will be weakened as is the case with simulation 0. In the remaining simulations we can observe a structure of LLJ with a maximum of moisture transport intensified to the south of the Great Plains responsible for advection of atmospheric moisture from the Gulf of Mexico.




Figure RC2.1. Moisture sources for the 6 study cases analysed. The vectors show the wind direction and intensity for each case study at surface level (aprox 500m). The purple line shows the position of the moisture labelling wall used in the WRF simulation.

2) The article focuses entirely on July GPLLJs with the logic that southerly GPLLJ frequency is highest for this month. Firstly, have the authors found this to be the case? In my own work, I have found May to be the month of highest frequency. The authors should include a figure or table of the ERA-INT-derived monthly GPLLJ climatology. Secondly, are July GPLLJs representative of the springtime LLJs that are predicted to increase in frequency and intensity (Ins 6-10, ph 3)? The authors could have designed their study to be better aligned with their motivations/stated best projections of a future GPLLJ.

We use the data from ERA-Interim and temporal period of 37 years (from 1980 to 2016). Examining the monthly variability, we find that July is the highest frequency month of GPLLJ. Understood the frequency as the largest number of days of GPLLJ (Fig. RC2.2, in this document). For the 37 years, we detected that the frequency increases during the months of May to October, with more than half of the days of GPLLJ. Nevertheless, we detected a peak in the month of July, with a frequency greater than 80%. Other authors, such as Rife et al. 2010 reported a frequency of GPLLJ of 78% also for the month of July. Despite this author used another dataset and a different time period than the one we use in our work.




Figure RC2.2 Monthly variability of the percentage of GPLLJ days.

Regarding the second question, the current literature foresees an increase both in the intensity and frequency of GPLLJ in spring. Nevertheless, the main goal of this work is to provide a first quantitative approach to moisture transported by the GPLLJ. Despite the numerous works related with GPLLJ, no work has objectively quantified the moisture associated with this structure. In the introduction we want to highlight the importance of studying these structures related to moisture transport. Changes in frequency and intensity will affect the sink regions and extreme events (droughts and floods). Therefore, it is of great interest to objectively estimate the moisture transported by the GPLLJ. However, in this paper we want to offer a first approximation in terms of average moisture transport. For this reason, we have objectively chosen objectively 5 cases of study based on the Gaussian distribution with the aim to quantify the average transport of moisture in a general perspective of the GPLLJ's behaviour.

Specific Comments:

Abstract: mention of ERA-INT and “southerly” GPLLJ needs to be made. - The word *southerly* was added and we mention the ERA-Interim reanalysis data used in the work.

Introduction: the work of Claudia Walters and Julie Winkler on GPLLJ (northerly and southerly) climatologies needs to be referenced here. There are several works from which to choose between 2001-present. - The work of Walters and Winkler, 2008 and other recent works have been added to the introduction.

Pg2,ln12 insert “southerly”. – The word “southerly” has been included in the latest version of the manuscript.

Pg2,ln21 specify whether Higgins et al (1997b) analysis was conditioned on GPLLJ occurrence -

Pg2,ln23 unclear meaning of “compared with the diurnal one”

Pg2,ln24 unclear if “this work” refers to Higgins or Mo reference

We consider the last three comments and the paragraph now read as follows:

Nevertheless, a large number of studies have documented the relationship between the major moisture transport and the GPLLJ. Higgins et al. (1996) studied the moisture budget over the central US in May employing NASA/DAO and NCEP/NCAR datasets, together with station observations, to evaluate the limitations of these products. Although both reanalyses overestimate daily mean precipitation rates, they accurately capture the basic temporal and structural characteristics of the GPLLJ. From the data, these authors calculated an increase in atmospheric moisture transport from the Gulf of Mexico during night-time of more than 50%. In a later work, Higgins et al. (1997) observed a well-defined nocturnal maximum of precipitation over the Great Plains in spring and summer by analysing station data. Particularly and linked to LLJ events this research found in this region an excess of 25% in nocturnal rainfall during summer when compared with the diurnal precipitation, associated with a rainfall decrease over the Gulf of Mexico. Additionally, Higgins et al. (1997) reported significant differences in precipitation pattern in coincidence (or not) with LLJ events. When a LLJ event occurs, the observations show an enhanced precipitation over the north-central United States and the Great Plains region, together with a decrease along the Gulf of Mexico and the western Atlantic. On the other hand, Mo and Juang (2003) found regional correlation at a distance between evaporation and precipitation, reflected in evaporation anomalies over the Great Plains along the trajectory of the GPLLJ, which are associated with downstream precipitation anomalies.

Pg2,ln27 suggest “found regional correlation at a distance between...” or similar – The text has been updated following the reviewer’s suggestion.

Pg2,lns30-32- one example of a “floating sentence” that needs to be grouped with another paragraph – Following the reviewer’s suggestion, the paragraph now reads as follows:

Otherwise, extreme rainfall events in the central US are related to an increase in moisture convergence downwind of the GPLLJ (Mo et al., 1997). A decisive factor that triggers heavy rains and floods is the presence of moisture advected by the GPLLJ from Gulf of Mexico and the Caribbean Sea. Moore et al. (2012) reported the physical processes related to the floods in May, 2010. A persistent southerly low-level jet associated with an atmospheric river (AR) enhanced the transport of moisture from the Gulf of Mexico into the heavy rainfall region. Thus, important socioeconomic impacts follow enhanced GPLLJ events, which modulate a large percentage of the local extreme precipitation events and flooding in warmer months (Mo et al., 1995, 1997; Beljaars et al., 1996; Trenberth and Guillemot, 1996; Arritt et al., 1997; Nakamura et al., 2013;

Nayak et al., 2016). All these results are consistent with the large-scale atmospheric moisture transport and support the marked influence of the GPLLJ over the central-eastern US, which has been shown to trigger more than 60% of the spring local precipitation there (Wang and Chen, 2009).

Pg2,ln31 meaning of “local” is unclear. Define local as opposed to non-local in this context. - The sentence has been rewrite as follow: *All these results are consistent with the large-scale atmospheric moisture transport and support the marked influence of the GPLLJ over the central-eastern US, which has been shown to trigger more than 60% of the spring precipitation over the Great Plains region (Wang and Chen, 2009).*

Pg3,ln4 suggest replacing “common” with “frequent” – “common” has been replaced by “frequent”.

Pg3,lns3-10 more floating sentences – The cited paragraph now reads as follows:

During the last decades, the GPLLJ has experienced a strengthening, accompanied by a northward migration causing a displacement of rainfall in the same direction. As a result, more frequent droughts have been observed in the southern Great Plains (Barandiaran et al., 2013). Besides, the increase in the number and intensity of GPLLJ events is also forecasted for future projections, which reveal an intensification of the GPLLJ during the spring season associated with global warming (Cook et al., 2008; Tang et al., 2017). As a result, increasing amounts of moisture transport and rainfall are expected, particularly from April to July, over the central US (Harding and Snyder, 2014). The same projections forecast a slight weakening of the GPLLJ from August to December, which could translate into increasing drought conditions.

Pg3,ln17 word “total” may be deleted – The word “total” has been removed from the manuscript.

Pg3,ln19 reword “and the at the point” - The sentence has been reworded as *a 37-year climatology was previously calculated at the point of maximum jet intensity...*

Pg3,ln27 on a monthly basis, I believe the max GPLLJ frequency is in May - We have calculated the monthly frequency for the 1980-2016 study period. As we show in figure RC2.2, we obtain the maximum frequency in the month of July.

Pg3,ln32 the native resolution of ERA-INT is closer to 0.75deg. How was it spatially interpolated (oversampled) to 0.25deg resolution? - The interim era data were downloaded directly from the ERA Interim web at a resolution of 0.25°. Although the original resolution is 0.75°, it was interim allowing the download to several horizontal resolutions. In our study, we obtained by downloading them at a resolution of 0.25. More information about the interpolation can be found in the following link: <https://confluence.ecmwf.int/display/CKB/ERA-Interim%3A+What+is+the+spatial+reference>

Pg5,ln29 delete “30” – The typo has been deleted.

Pg6,ln17 clarify for the reader whether these events were chosen from the NLLJ distribution at 32.75N,99W or for the regional distribution (w/I cyan outline) – The sentence: *The five case-studies were selected based on the Gaussian adjustment applied to the study.* has been included in the new version of the manuscript.

Pg6,ln25 should “LLJ” be “NLLJ”? – This typo has been corrected.

Pg7,ln22 “northeastern” – The typo has been corrected.

Pg8,ln8 clarify that this is done for a specific point (32.75N,99W) - The core (32.75°N,99°W) is used only to calculate the 37-yr climatology. We add the following sentence to the text to clarify

it: The target region used in FLEXPART was defined based on the 75th percentile of the index value.

Pg8,ln16 I do not believe it is true that GPLLJ occurs on more than 16/31 nights in July. Please quantify this using ERA-INT. - We use the ERA-Interim data for the detection of GPLLJ. As we discussed, we obtained a maximum frequency in July. Please, we refer you to the previous comment.


Pg8,ln20 “northeastern” – The typo has been corrected.

Pg8,ln21 synoptic and land preconditioning will impact ratio of GPLLJ TPW (Fig 3). – This clarification has been included in the text.

Pg8,ln26 replace “leaded” with “preceded” – “leaded” has been replaced by “preceded”.

Pg8,ln31 suggest “: : North America [using WRF-TT. Additional] simulations should...” – “WRF-TT” has been removed from the manuscript.

Fig. 1 the cyan color is hard to distinguish in my color print. Clarify whether these “frequency



distributions” are derived for the region contained in the cyan outline or for a single point (i.e., 32.75°N, 99°W). - The cyan color is replaced by the magenta. We think it differs a little better. The frequency of distribution was calculated at the point of maximum intensity. We have been modified the caption of the figure 2 to clarify it.

Figure 1. (a) Mean NLLJ index (shaded) and 500 m winds (arrows, in m s^{-1}) at local midnight in July (boreal summer) for 1980-2016, calculated from ERA-Interim reanalysis. The black cross at 32.75°N, 99°W shows the point of maximum NLLJ in the climatology. The cyan contour line surrounds the region containing points above the 75th percentile. (b) Frequency distribution of the GPLLJ for the months of July from 1980 to 2016 (blue bars). The red curve corresponds to the Gaussian fit (see table A2). Noted: The frequency distribution is calculated at the point of maximum intensity of NLLJ (at 32.75°N, 99°W, black cross in fig. 1a).

Figs 2-4. Lat/lon labels required on these figures.

Fig 4. Suggest adding state boundaries.

Lat and lon and the states boundaries have been included in the lasted version of the manuscript. An example figure is shown:




Fig5-6. The order of Fig5 and Fig6 should be switched. Would it also be informative to plot the vertical cross section of relative GPLLJ humidity? E.g., $q_{TR}:q$; $\phi_{TR}:\phi$? – Figure 5 and 6 will be swapped. Additionally, the vertical cross sections will be included and commented in the manuscript.

Table 1. Specify ERA-INT-derived as well as the lat/lon location or domain over which the frequency distribution was composed. - The title of the table has been rewritten as follows:

Table 1: Case-studies objectively selected based in the frequency distribution of the LLJ index to carry out WRF-TT simulations. μ is the mean of the distribution and σ its standard deviation. Noted: The frequency distribution is calculated at the point of maximum intensity of NLLJ at 32.75°N, 99°W (black cross in fig. 1a) using the ERA-Interim reanalysis dataset.

On the assessment of the moisture transport by the Great Plains low-level jet

Iago Algarra¹, Jorge Eiras-Barca¹, Gonzalo Miguez-Macho², Raquel Nieto¹, and Luis Gimeno¹

¹EPhysLab (Environmental Physics Laboratory), Facultade de Ciencias, Universidade de Vigo, Ourense, Galicia, Spain

²Non-Linear Physics Group, University of Santiago de Compostela, Galicia, Spain

Correspondence to: Iago Algarra (ialgarra@uvigo.es)

Abstract. Low-Level Jets (LLJs) can be defined as filamentous wind corridors of anomalously high wind speed values located within the first km of the troposphere. These structures, ~~together with atmospheric rivers (ARs),~~ are the major meteorological systems in the meridional transport of moisture on a global scale. ~~(together with atmospheric rivers).~~ In this work, we focus on the ~~southerly~~ Great Plains low-level jet, which plays an important role in the moisture transport balance over the central United States. The Gulf of Mexico is the main moisture source for the GPLLJ, which has been identified as a key factor for rainfall modulation over the eastern and central US.

The relationship between moisture transport from the Gulf of Mexico to the Great Plains and precipitation is well documented in previous studies. Nevertheless, a large uncertainty still remains in the quantification of the moisture amount actually carried by the GPLLJ. The main goal of this work is to address this question. For this purpose, a relatively new tool, the regional atmospheric Weather Research and Forecasting Model with 3D water vapour tracers (WRF-~~TFWVT~~, Insua-Costa and Miguez-Macho, 2018) is used together with the Lagrangian model FLEXPART to estimate the load of precipitable water advected within the GPLLJ. ~~Both models were fed with data from ERA Interim.~~ From a climatology of jet intensity over a 37-year period ~~(Rife et al., 2010)~~, which follows a Gaussian distribution, we select for study 5 cases representing the mean, and one and two standard deviations above and below it. Results show that the jet is responsible for roughly 70%-80% of the moisture transport occurring in the southern Great Plains when a jet event occurs. Furthermore, moisture transport by the GPLLJ extends to the northeast US, accounting for 50% of the total in areas near the Great Lakes. Vertical distributions show the maximum of moisture advected by the GPLLJ at surface levels and maximum values of moisture flux about 500 m above, in coincidence with the wind speed profile.

1 Introduction

It is well known that the Great Plains Low-Level Jet (hereafter, GPLLJ) plays an important role in the balance of the moisture transport over the central United States (Stensrud, 1996; Schubert et al., 1998). The atmospheric moisture is transported by the GPLLJ from tropical and subtropical latitudes (particularly the Gulf of Mexico and the Caribbean Sea) into the Great Plains

(Helfand and Schubert, 1995; Mo et al., 1997) where the jet is often responsible for nocturnal deep convective activity (Higgins et al., 1997; Pu et al., 2016). In the last decades, a large number of authors have shown the strong influence of the GPLLJ as a modulator of climate and rainfall over this region and even further east (Mo et al., 1995, 1997; Wu and Raman, 1998; Byerle and Paegle, 2003); for instance, throughout May and June it is estimated that at least one-third of the moisture penetrating into the continental US is carried by the GPLLJ (Helfand and Schubert, 1995).

Among the mechanisms which have been proposed as triggers of the GPLLJ are included a combination of inertial oscillations (Blackadar, 1957) and orographic forcing (Wexler, 1961; Byerle and Paegle, 2003; Pan et al., 2004; Ting and Wang, 2006). Particularly, the mechanism of Blackadar (1957) suggests that inertial oscillations near the friction layer can induce the formation of the GPLLJ (Wu and Raman, 1998). Nevertheless, orographic effects are also understood as a key factor in the maintenance of the GPLLJ. In this regard, Ting and Wang (2006) proved that, when the interaction with the orography is removed from numerical simulations, the GPLLJ vanishes, together with an important amount of the summer precipitation over the central and southern-southerly US.

The GPLLJ is a phenomenon confined within the first kilometres of the troposphere and is closely related to the warm season (Bonner, 1968). Besides, it is characterized by a strong diurnal oscillation, with a peak in strength during night hours (Augustin and Caracena, 1994). A long-term climatology of GPLLJ can be found in the work of Walters et al. (2008). The GPLLJ is a phenomenon extremely localized in time and space and its role in the continental moisture balance is difficult to study solely from observations.

Nevertheless, a large number of studies have documented the relationship between the major moisture transport and the GPLLJ. Higgins et al. (1996) studied the moisture budget over the central US in May employing NASA/DAO and NCEP/NCAR datasets, together with station observations, to evaluate the limitations of these products. Although both reanalyses overestimate daily mean precipitation rates, they accurately capture the basic temporal and structural characteristics of the GPLLJ. From the data, these authors calculated an increase in atmospheric moisture transport from the Gulf of Mexico during nighttime of more than 50%. In a later work, Higgins et al. (1997b) observed a well-defined nocturnal maximum of precipitation over the Great Plains in spring and summer by analysing station data. Particularly and linked to LLJ events they found in the region an excess of 25% in nocturnal rainfall during summer when compared with the diurnal precipitation, associated with a rainfall decrease over the Gulf of Mexico. Particularly, this research found over the region an excess of 25% in nocturnal precipitation during summer when compared with the diurnal one, associated with a rainfall decrease over the Gulf of Mexico. Additionally, Higgins et al. (1997) this work reveals reported significant differences in precipitation pattern in coincidence (or not) with LLJ events. When a LLJ event occurs, the observations show an enhanced precipitation over the north-central United States and the Great Plains region, together with a decrease along the Gulf of Mexico and the western Atlantic ~~(Mo et al., 1997)~~. On the other hand, Mo and Juang (2003) found a regional dependence-correlation at a distance between evaporation and precipitation, reflected in evaporation anomalies over the Great Plains along the trajectory of the GPLLJ, which are associated with downstream precipitation anomalies.

~~All these results are consistent with the large-scale atmospheric moisture transport and support the marked influence of the GPLLJ over the central-eastern US, which has been shown to trigger more than 60% of the spring local precipitation there (Wang and Chen, 2009).~~

Otherwise, extreme rainfall events in the central US are related to an increase in moisture convergence downwind of the GPLLJ (Mo et al., 1997). A decisive factor that triggers heavy rains and floods is the presence of moisture advected by the GPLLJ from the Gulf of Mexico and the Caribbean Sea. Moore et al. (2012) reported the physical processes related to the floods in May 2010, when a persistent southerly LLJ associated with an atmospheric river (AR) enhanced the transport of moisture from the Gulf of Mexico into the heavy rainfall region. Thus, important socioeconomic impacts follow enhanced GPLLJ events, which modulates a large percentage of the local extreme precipitation events and flooding in warmer months (Mo et al., 1995, 1997; Beljaars et al., 1996; Trenberth and Guillemot, 1996; Arritt et al., 1997; Nakamura et al., 2013; Nayak et al., 2016). All these results are consistent with the large-scale atmospheric moisture transport and support the marked influence of the GPLLJ over the central-eastern US, which has been shown to trigger more than 60% of the spring local precipitation over the Great Plains region~~there~~ (Wang and Chen, 2009).

During the last decades, the GPLLJ has experienced a strengthening, accompanied by a northward migration causing a displacement of rainfall in the same direction. As a result, more ~~frequent~~common droughts have been observed in the southern Great Plains (Barandiaran et al., 2013). Besides, ~~The~~ increase in the number and intensity of GPLLJ events is also forecasted for future projections, which reveal an intensification of the GPLLJ during the spring season associated with global warming (Cook et al., 2008; Tang et al., 2017). As a result, increasing amounts of moisture transport and rainfall are expected, particularly from April to July, over the central US (Harding and Snyder, 2014). The same projections forecast a slight weakening of the GPLLJ from August to December, which could translate into increasing drought conditions.

The knowledge about the GPLLJ, together with the insights on the relationship between the moisture transported by the GPLLJ and local precipitation patterns has increased considerably during the last decades. However, there are still unanswered questions about the quantification of such water vapour transport and specially about the estimation of the ratio of land to oceanic moisture sources associated with the GPLLJ. This estimate of the oceanic input to the moisture transport associated with the GPLLJ is essential to predict and understand the behaviour of the GPLLJ in future scenarios.

In this work, a new tool, the regional atmospheric Weather Research and Forecasting Model with 3D water vapour tracer diagnostics (WRF-~~TTWVT~~, Eiras-Barea et al., 2017; Insua-Costa and Miguez-Macho, 2018) is used to quantify the total amount of total precipitable water (TPW) transported by the GPLLJ. To show the differences between the transport of moisture on jet and non-jet days, a 37-year climatology was ~~calculated~~ previously calculated and the at the point of maximum jet intensity is obtained following the methodology by Rife et al., (2010). The structure of this work is: ~~as follows~~, in Section 2 we provide the methodology used, in Section 3 we show the results obtained, and finally in Section 4 we discuss the conclusions.

2 Data and methods

2.1 Detection of the Great Plains low-level jet

To objectively detect days with LLJ over the Great Plains, we applied the night-time index proposed in Rife et al., (2010), hereafter named as NLLJ. This index is based on the temporal variation of the wind's vertical structure and the fact that LLJs are most intense at local midnight. Because both frequency and intensity of GPLLJ are mostly associated with the warm season, we develop a 37-year climatology for the month of July (representative for the boreal summer). According to the NLLJs characteristics, and with the aim to define the index, two conditions should be met to consider a GPLLJ detection:

1. The wind speed is higher at local midnight than at midday.
2. The local midnight wind speed is higher at the surface ($\sim 500\text{m}$) than ~~in height~~above it ($\sim 4\text{km}$).

The index is calculated at each grid point over an area centred over the US using 6-hourly ERA-Interim reanalysis data (Dee et al., 2011) with a 0.25° horizontal resolution. ~~Due to~~Because of the jet core is located within of the first kilometre of the troposphere, it is necessary to take into account the elevation of the land, so sigma coordinates are used. The GPLLJ-climatology is developed for 37 years, from 1980 to 2016, and the NLLJ index can be defined as follows:

$$NLLJ = \lambda \varphi \sqrt{[(u_{00}^{L1} - u_{00}^{L2}) - (u_{12}^{L1} - u_{12}^{L2})]^2 + [(v_{00}^{L1} - v_{00}^{L2}) - (v_{12}^{L1} - v_{12}^{L2})]^2} \quad (1)$$

where u and v are the zonal and meridional wind components, respectively. $L1$ represents the winds at the surface at 53 sigma level (elevation near the jet core), approximately 500 m above ground level (AGL), while $L2$ corresponds to the wind at 42 sigma level (around 4000 m AGL). Numbers 00 and 12 refer to local midnight and local noon, respectively. λ and φ are binary multipliers representing the temporal and vertical variation of ~~the~~ the wind. Particularly, λ relates to the daily variation of the wind at 500 m and φ refers to the wind's vertical variation between 500 m and 4 km at midnight (Rife et al., 2010):

$$\lambda = \begin{cases} 0, & ws_{00}^{L1} \leq ws_{12}^{L1} \\ 1, & ws_{00}^{L1} > ws_{12}^{L1} \end{cases} \quad (2)$$

$$\varphi = \begin{cases} 0, & ws_{00}^{L1} \leq ws_{00}^{L2} \\ 1, & ws_{00}^{L1} > ws_{00}^{L2} \end{cases} \quad (3)$$

2.2 Identification of moisture sources associated with Great Plains low-level jet

For the objective identification of moisture sources associated with the GPLLJ, the Lagrangian backward trajectories from the FLEXPART v9.0 model are used (Stohl et al., 2005a). This model provides a large number of air parcel trajectories from which it is possible ~~to the calculation~~ of the evaporation minus precipitation budget, tracking all changes in the specific humidity of air parcels.

FLEXPART has been widely and successfully used to track moisture paths for the study of the atmospheric branch of the hydrologic cycle in different parts of the world (e.g., Hu et al., 2018; Sorí et al., 2018; Vázquez et al., 2016). Furthermore, this tool is able to infer the moisture sources for precipitation falling in a target region when backward trajectories are considered (eg., Sthol et al., 2008; Drumond et al., 2012; Gimeno et al., 2012; Wegmann et al., 2015; Ramos et al., 2016).

5 In this work we use the outputs of a global experiment in which FLEXPART v9.0 tracks approximately 2 million particles (air parcels) with constant mass distributed on the globe every time step during 37-year period (1980-2016). These air parcels are advected by the 3D wind field, and the variables of interest of each particle (such as position, height, specific humidity, temperature among many others) are saved at each time step. We perform a FLEXPART simulation fed with ERA-Interim reanalysis data at 1° horizontal resolution on 61 vertical levels from sea level pressure to 0.1 hPa and 6-hour time intervals (00, 10 06, 12 and 18 UTC). The model is run with a 3 h timestep, and linear interpolation is used to obtain data with this frequency from ERA-Interim. The backward trajectories are followed during-for 10 days, which is the average life time of water vapour in the atmosphere (Numaguti, 1999).

The changes in specific humidity (q) of each air parcel along its path can be expressed as follows:

$$15 \quad e - p = m \frac{dq}{dt} \quad (4)$$

where m is the mass of a particle (which remains constant in the simulation), q is the specific humidity, t the time step, and $e - p$ (evaporation minus precipitation) represents the water flux associated with the particle. To obtain the instantaneous values of the $E - P$ balance (E denotes evaporation and P the precipitation rate per unit area) in a given area (in this case, over one of 20 1-0° x 1-0 degrees in latitude and longitude), it is necessary to integrate the moisture changes for all particles present in the atmospheric column over such area (E denotes evaporation and P the precipitation rate per unit area) (Stohl and James 2004, 2005b). Thus, following this methodology, in a backward experiment, a moisture source is defined as those regions where $E - P$ is positive ($E - P > 0$), which implies that evaporation exceeds precipitation, while a moisture sink is defined as a region where ($E - P < 0$), meaning that precipitation is greater than evaporation.

25 In our study, backward trajectories were followed from the area composed of points with values of NLLJ above the percentile 75th percentile to found the main moisture source.-

2.3 The regional atmospheric Weather Research and Forecasting Model with 3D water vapour tracer diagnostics (WRF-~~TFWVT~~)

The mesoscale Weather Research and Forecast model (WRF 3.8.1) with the moisture tracers tool (WRF-~~TFWVT~~, Eiras Barea et al., 2017; Insua-Costa and Miguez-Macho, 2018) is used to carry out to quantify the total amount of total precipitable water (TPW) transported by the GPLLJ. In order to analyse the moisture transport associated with the GPLLJ avoiding the effects of other synoptic-scale transport events, we tag the moisture passing northward through a narrow wall located on the northern edge of the moisture source region identified using the FLEXPART model. When a particle of water (whether in liquid, solid

or gas state) crosses the wall, it is labeled for further analysis inside the simulation domain. We consider all water traversing the wall to be advected by the GPLLJ.

The horizontal resolution of the simulations is 20 km and the vertical column is divided into 38 levels. The simulation covers a time window of 11 days, starting 7 days prior to the day of interest. The model parameterizations together with the WRF-~~TT~~ **WVT** are set using the PBL Yonsei University (YSU) parametrization (Hu et al., 2013; Shin and Hong, 2011), the schemes of Kain-Fritsch for convection (Kain, 2004), the Dudhia one for short-wave radiation (Dudhia, 1989), the Rapid Radiative Transfer Model (RRTM) scheme for long-wave radiation (Mlawer et al., 1997), and the WRF Single-Moment 6-Class Microphysics Scheme (WSM6) (Hong and Lim, 2006).

In addition, we apply spectral nudging of waves longer than 1000 km above the boundary layer, with a relaxation time of 1h, to avoid ~~30~~-distortion of the large-scale circulation. This configuration has been validated and successfully applied several times with the WRF-~~TT~~-**WVT** in mid latitudes (e.g., Eiras-Barca et al., 2017). Spectral nudging ensures that the large-scale circulation is well captured in the simulations. ERA-Interim data provides lateral boundary and initial conditions for the runs (Dee et al., 2011). The variables of interest for the analysis of the GPLLJ event are computed as follows. Integrated Water Vapour (IWV), Eq. (5) is obtained by vertical integration of the specific humidity (q) in pressure (p) levels, where g represents the gravitational force. The instant flux of moisture ($\sigma\phi$) is calculated as stated in Eq. (6) and the conversion between (g) and the water vapour mixing ratio obtained from WRF is performed using Eq. (7), where u and v are the horizontal components of the wind field.

$$IWV_{(i,j)} = \frac{1}{g(k)} \int_{surface}^{top} q(i,j,k) dp \quad (5)$$

$$\sigma\phi(i,j,k) = |q \cdot (u,v)| \quad (6)$$

$$q = \frac{w}{w+1}, \text{ with } w \ll 1 \rightarrow q \approx w \quad (7)$$

3 Results

3.1 Characterization of the Great Plains low-level jet

As previously mentioned, the NLLJ index was calculated at each grid point over the North American region for the month of July over the period 1980-2016. July was found the month with maximum LLJs events (Figure S1 shows the monthly distribution of the GPLLJ days of detection). Fig. 1.a shows the climatological NLLJ index and the wind field at 500 hPa. The black cross indicates the point of maximum intensity of the index (8.8 m s⁻¹). At this point, located at 32.75°N-99°W, along the 37-years analysed, and for July, a total of 931 LLJ days are identified, that is, 81% of all days have a positive value of the index. On the point of maximum intensity showed in Fig1.a, Fig1.b displays the frequency distribution of the NLLJ for the period 1980-2016. A clear peak around 11 m s⁻¹ is observed together with a Gaussian behaviour ($R^2=0.95$, ~~red line~~ Jarque-

Bera test p-value=0.0055, which provides a confidence level close to 99.5%, red line). The latter has been used to select the five case studies to be ~~analyzed~~-analysed with WRF-~~TT~~-WVT and listed in Table 1. The five ~~events~~-case-studies were selected based on the Gaussian adjustment applied to the study. These five event~~s~~hosen correspond to μ , $\mu \pm 2\sigma$, $\mu \pm \sigma$ (where μ is the mean of the distribution and σ its standard deviation), and they provide a general perspective of the LLJ's behaviour. Since

5 each case-study WRF-~~TT~~-WVT simulation spans ~~for~~ 11 days, a condition of persistence of the index value for at least two days after the main jet day is applied. Additionally, we perform a sixth simulation with a non-jet day (simulation 0 in the Table 1). This non-jet day is selected from the developed climatology as the longest non-jet period, in order to avoid overlaps in moisture transport with jet days.

3.2 Moisture transport associated with the Great Plains low-level jet

10 In order to detect the main climatological oceanic source of moisture for the GPLLJ we used the FLEXPART trajectories outputs for 1980-2016. The area englobed in the 75th percentile of the NLLJ index values (eyan-magenta line in figure 1.a) was selected as the target region for the backward experiment (as it was explained in the methodology). Fig. 2 shows the source of moisture in red colour, obtained as the 75th percentile of the ($E - P > 0$) field. This area covers the southern Gulf of Mexico and extends into the Caribbean Sea, between 60°-98°W and 12°-28°N. Figure S2 in the supplementary material shows the

15 individual source of moisture for each case in study).

Although the flow originated in the source of moisture is advected in the low levels as a result of the strong intensity of the trade winds, a 3D-label wall (at 29-30°N and from 94.5°W to 100°W) was used in the WRF-~~TT~~-WVT simulations (orange line in fig. 2). The position of the sentinel wall was selected on the region where oceanic moisture associated with the GPLLJ landfalls. The wall remained constant in the WRF-~~TT~~-WVT simulations. ~~A~~The thin wall was used instead of the entire source

20 regions in order to avoid overlaps in the labelling of moisture caused by secondary synoptic scale mechanisms.

Fig. 3 shows the ratio of precipitable water transported by the GPLLJ to total precipitable water ($TPW_{tracers}=TPW$) for the six case-studies analysed. As mentioned earlier, $TPW_{tracers}$ represents the TPW that has crossed the "wall" highlighted in orange in Figure ~~4~~2, which we assume corresponds to moisture advected by the GPLLJ. Following the same behaviour of the GPLLJ itself, the moisture flux is initially in the northward direction and veers east as it penetrates into the Great Plains for all events

25 with positive NLLJ index values. As expected, the non-jet event with NLLJ value equal to zero (Sim 0) does not show remarkable moisture fluxes. For the jet events, ratios are close to one in regions near the tagging wall and extend for hundreds of km northward with significant values above 60%. Percentages between 70% and 80% are observed in the Great Plains. The large geographical reach of the moisture associated with the GPLLJ is evidenced in this figure, showing that for certain GPLLJ events it can occasionally explain more than 50% of TPW even in the north-east US. It is necessary to highlight that higher

30 values in the index value does not necessarily mean larger flows of moisture in the entire, as can be observed, for example, when SIM3 and SIM5 are compared. -Figure S3 in the supplementary material shows a comparison of accumulated precipitation at 11 days of WRF simulations versus CPC gauge-analysis observations (Chen et al., 2008).

As it was previously stated, the aim of this work is to study the general behaviour of the GPLLJ associated with its moisture transport. In the first simulated case of GPLLJ (fig.3 – Sim 1) it is observed that most of the precipitable water is concentrated on the Great Plains, exceeding ratios of 80% out of the total. In the second GPLLJ event simulated (fig.3 – Sim 2), the precipitable water extends northeast of the US and to the south of the Great Lakes and the GPLLJ, where explain close of the 50% of precipitable water. The third simulated case corresponds to the average behaviour of the GPLLJ (fig.3 – Sim 3) and evidences the influence of the GPLLJ in the northeast of US with ratios near 50% in the US East Coast. Nevertheless, in areas along the path of the GPLLJ, the advection of precipitable water is close to 80%. In the fourth and fifth simulations of GPLLJ (fig.3 – Sim 4 and 5), the plume of precipitable water is concentrated over the Great Plains. However, the water precipitable ratio is reduced as latitude increases, but values are still close to 50% in the northeastern areas of the US.

Fig. 4 shows the statistically weighted mean of the ratio shown in Fig. 3 for the five case studies with NLLJ > 0 considered in the analysis. The weights associated with each event are stated in Table 1 at the last column, and the objective criteria to assign them can be found in appendix A1. The aim of using weights in the analysis is to give greater importance to the event representing the mean value of the NLLJ and less relevance to the events in the tail of the distribution. Notwithstanding the limited number of simulations used in the analysis, this procedure allows us to interpret Figure 4 as a "climatology" of the moisture transport associated with the GPLLJ. Roughly 80-90% of the precipitable water in its core zone of influence over the Great Plains, in Texas and Oklahoma, is carried by the GPLLJ when a jet event occurs. With increased distance from that area, the ratio of precipitable water transported by the GPLLJ decreases; however, the contribution of moisture from the Gulf of Mexico to TPW is still of more than 50% as far north as the Great Lakes.

On the one hand, Fig. 5 shows the TPW and the cross sections for the main GPLLJ event (1992.07.11). On the other hand,

Fig. 5 shows the vertical distribution of tracer specific humidity (q_{TR}) and tracer water vapour flux (ϕ_{TR}) for cross sections at positions depicted in Fig. 6 for the main GPLLJ event (1992.07.11). Tracer moisture (a-c) has a maximum at surface levels, while the moisture flux (d- f) maximizes at 500 m AGL where the LLJ core is located. A significant presence of both tracer water vapour and tracer water vapour flux is restricted to the first 2 km AGL. Overall, as the latitude increases the water vapour plume from the Gulf of Mexico tends to rise in the vertical column and expand zonally along the GPLLJ path to the east of the

U.S. Equivalent conclusions can be obtained from the remaining events, which are shown in supplementary material S4(Figs. S4-S7). Figure 7 shows the steam ratio $q_{TR}:q$ for the vertical sections of Fig. 5 for the same event. The moisture pattern behaviour is similar to Fig. 6; for regions close to the Gulf of Mexico the moisture ratio is concentrated mostly at lower levels (Fig. 7a) and extends on the horizontal axis as it moves away from the source of moisture (Fig. 7b). Nevertheless, for remote regions the maximum moisture ratio continues at lower levels (Fig 7c). -The moisture flow has the same pattern, high moisture ratios that remain at low levels despite distancing longitudinal distance (Fig. 7d-7f). These results confirm that the GPLLJ transports a great concentrated quantify of moisture.

4 Conclusions and discussions

A combination of Lagrangian and Eulerian methods were used to identify and objectively quantify the moisture transport associated with the Great Plains Low-Level Jet (GPLLJ). First, the ~~L~~agrangian model FLEXPART was used to locate the GPLLJ moisture sources ~~in the Gulf of México~~ for the month of its maximum activity (July) throughout the period 1980-2016.

5 The target region used in the FLEXPART simulation to find the main source of moisture was defined based on the 75th percentile of the GPLLJ index value previously calculated based in Rife et al. (2010) method. Once the Gulf of Mexico was identified as the main source of moisture ($E - P > 0$) for the GPLLJ, we use a new tool known as eulerian 3D WRF-~~WTT~~
10 WVT (Eiras-Barea et al., 2017; Insua-Costa and Miguez-Macho, 2018) which was applied to track the moisture advected in six selected GPLLJ events based on the distribution of the GPLLJ index used previously to detect the GPLLJ (Rife et al., 2010). So, this work analysed the behaviour of the GPLLJ during the month of its maximum activity (July) for a long period, 1980-2016, and we select six representative cases for the WRF-WVT simulation-

The moisture transport analysis reveals the major role played by the GPLLJ in the water cycle of central North America, transporting large amounts of moisture from the Gulf of Mexico as far as the north-east US. Particularly, advection by the jet explains more than 80% of the precipitable water in the southern Great Plains when a jet event occurs, which, in July, is most
15 of the days. The Rocky Mountains blocks the circulation of GPLLJ and force it to turn to the east of the US, reaching even the eastern coast of the US. The moisture transport associated with the GPLLJ is in a plume of moisture where, the maximum transport occurs in the path of the GPLLJ. As expected, the ratio reduces as latitude increases, but values are still close to 50% in the north-eastern areas of the US.

We note that the extension of the GPLLJ is dependent on the synoptic conditions or land preconditioning, among other factors,
20 which may alter the ratio of the TPW. ~~are~~ Nevertheless, the analysis of these multiple factors is out of the scope of this paper. For example, the presence of a well-developed high pressure system in higher latitudes of North America may block the advection of the GPLLJ moisture to this region. Dong et al., (2011) related the drought of 2006 with an anomalous high over south-western U.S region and an anomalous low over the Great Lakes. This pattern inhibited the advection of moisture from the Gulf of Mexico contributing to the extreme dryness, and the lack precipitation was associated with a suppressed cyclonic
25 activity over the south-western US. However, the 2007 flood events were initially ~~preceded~~leaded by active synoptic weather patterns, linked to an active moisture transport from the Gulf of Mexico by the GPLLJ.

Besides, higher values in the NLLJ index mean larger differences between winds aloft and at the surface at the reference point, but do not necessarily mean stronger moisture transport.

Thus, results should be understood as a first approach to the quantification of the large extent of GPLLJ moisture advection
30 and its implications for the water budget in North America. More ~~WRF-TT~~ simulations should be conducted, and other months should be included in FLEXPART backward calculations to extend this work and produce a more comprehensive analysis.

Author contributions.

IA, RN and LG had the initial idea. GMM developed WRF-WVT tool. IA and JEB carried out the simulations and data analysis. IA and JEB wrote the original draf and reviewed the subsequent documents. RN, LG and GMM provided suggestions under the revision period, ~~commented and reviewed the manuscript before submission.~~

5 Competing interests.

The authors declare that they have no conflict of interest.

Acknowledgments.

~~The ECMWF ERA Interim reanalysis data were obtained from <https://www.ecmwf.int/en/forecasts/datasets/reanalysis-datasets/era-interim>.~~ Iago Algarra was financially supported by Spanish government (MINECO), (CGL2015-65141-R). Jorge Eiras-Barca has been financially supported by the EDB481B 2018/069 from the “Xunta de Galicia” and the Fullbrigh commission. This work was also partially supported by Xunta de Galicia under Project ED431C 2017/64-GRC ‘Programa de Consolidación e Estructuración de Unidades de Investigación Competitivas (Grupos de Referencia Competitiva)’. Iago Algarra would like to express his gratitude to ~~the~~ all the Non-Linear Physics Group for their kind support during his~~their~~ stay in the University of Santiago de Compostela.

15

References

- Arritt, R. W., Rink, T. D., Segal, M., Todey, D. P., Clark, C. A., Mitchell, M. J., Labas, K. M., Arritt, R. W., Rink, T. D., Segal, M., Todey, D. P., Clark, C. A., Mitchell, M. J. and Labas, K. M.: The Great Plains Low-Level Jet during the Warm Season of 1993, Mon. Weather Rev., 125(9), 2176–2192, doi:10.1175/1520-0493(1997)125<2176:TGPLLJ>2.0.CO;2, 1997.
- 20 Augustine, J. A. and Caracena, F.: Lower-tropospheric precursors to nocturnal MCS development over the central United States, Weather Forecast., 9(1), 116–135, doi:10.1175/1520-0434(1994)009<0116:LTPTNM>2.0.CO;2, 1994.
- Barandiaran, D., Wang, S.-Y. and Hilburn, K.: Observed trends in the Great Plains low-level jet and associated precipitation changes in relation to recent droughts, Geophys. Res. Lett., 40(23), 6247–6251, doi:10.1002/2013GL058296, 2013.
- Beljaars, A. C. M., Viterbo, P., Miller, M. J. and Betts, A. K.: The Anomalous Rainfall over the United States during July 1993: Sensitivity to Land Surface Parameterization and Soil Moisture Anomalies, Mon. Weather Rev., 124(3), 362–383, doi:10.1175/1520-0493(1996)124<0362:TAROTU>2.0.CO;2, 1996.
- 25 Blackadar, A. K.: Boundary layer wind maxima and their significance for the growth of nocturnal inversions, Bull. Amer. Meteor. Soc., 38, 283–290, doi:10.1175/1520-0477-38.5.283, 1957.

- Bonner, W. D.: Climatology of the low level jet, *Mon. Weather Rev.*, 96(12), 833–850, doi:10.1175/1520-0493(1968)096<0833:COTLLJ>2.0.CO;2, 1968.
- Byerle, L. A. and Paegle, J.: Modulation of the Great Plains low-level jet and moisture transports by orography and large-scale circulations, *J. Geophys. Res. D Atmos.*, 108(16), GCP 6-1-GCP 6-16, doi:10.1029/2002JD003005, 2003.
- 5 [Chen, M., Shi, W., Xie, P., Silva, V. B. S., Kousky, V. E., Higgins, R. W. and Janowiak, J. E.: Assessing objective techniques for gauge-based analyses of global daily precipitation, *J. Geophys. Res. Atmos.*, 113\(4\), 1–13, doi:10.1029/2007JD009132, 2008.](#)
- Cook, K. H., Vizu, E. K., Launer, Z. S. and Patricola, C. M.: Springtime intensification of the great plains low-level jet and midwest precipitation in GCM Simulations of the twenty-first century, *J. Clim.*, 21(23), 6321–6340, doi:10.1175/2008JCLI2355.1, 2008.
- 10
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N. and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q. J. R. Meteorol. Soc.*, 137(656), 553–597, doi:10.1002/qj.828, 2011.
- 15
- Dong, X., Xi, B., Kennedy, A., Feng, Z., Entin, J. K., Houser, P. R., Schiffer, R. A., L’Ecuyer, T., Olson, W. S., Hsu, K. L., Liu, W. T., Lin, B., Deng, Y. and Jiang, T.: Investigation of the 2006 drought and 2007 flood extremes at the Southern Great Plains through an integrative analysis of observations, *J. Geophys. Res. Atmos.*, 116(3), doi:10.1029/2010JD014776, 2011.
- 20
- Drumond, A., Nieto, R., Trigo, R., Ambrizzi, T., Souza, E. and Gimeno, L.: A lagrangian identification of the main sources of moisture affecting northeastern Brazil during its pre-rainy and rainy seasons, *PLoS One*, 5(6), 1–8, doi:10.1371/journal.pone.0011205, 2010.
- Dudhia, J.: Numerical Study of Convection Observed during the Winter Monsoon Experiment Using a Mesoscale Two-Dimensional Model, *J. Atmos. Sci.*, 46(20), 3077–3107, doi:10.1175/1520-0469(1989)046<3077:NSOCOD>2.0.CO;2, 1989.
- 25
- Eiras-Barca, J., Dominguez, F., Hu, H., Garaboa-Paz, D. and Miguez-Macho, G.: Evaluation of the moisture sources in two extreme landfalling atmospheric river events using an Eulerian WRF tracers tool, *Earth Syst. Dyn.*, 8(4), 1247–1261, doi:10.5194/esd-8-1247-2017, 2017.
- Gimeno, L., Stohl, A., Trigo, R. M., Dominguez, F., Yoshimura, K., Yu, L., Drumond, A., Durán-Quesada, A. M. and Nieto, R.: Oceanic and terrestrial sources of continental precipitation, *Rev. Geophys.*, 50(4), RG4003, doi:10.1029/2012RG000389, 2012.
- 30

- Harding, K. J. and Snyder, P. K.: Examining future changes in the character of central u.S. warm-season precipitation using dynamical downscaling, *J. Geophys. Res.*, 119(23), 13,113–116,136, doi:10.1002/2014JD022575, 2014.
- Helfand, H. M. and Schubert, S. D.: Climatology of the simulated Great Plains low-level jet and its contribution to the continental moisture budget of the United States, *J. Clim.*, 8(4), 784–806, doi:10.1175/1520-0442(1995)008<0784:COTSGP>2.0.CO;2, 1995.
- Higgins, R. W., Mo, K. C. and Schubert, S. D.: The moisture budget of the central United States in spring as evaluated in the NCEP/NCAR and the NASA/DAO reanalyses, *Mon. Weather Rev.*, 124(5), 939–963, doi:10.1175/1520-0493(1996)124<0939:TMBOTC>2.0.CO;2, 1996.
- Higgins, R. W., Yao, Y., Yarosh, E. S., Janowiak, J. E. and Mo, K. C.: Influence of the great plains low-level jet on summertime precipitation and moisture transport over the central United States, *J. Clim.*, 10(3), 481–507, doi:10.1175/1520-0442(1997)010<0481:IOTGPL>2.0.CO;2, 1997.
- Hong, S. and Lim, J.: The WRF single-moment 6-class microphysics scheme (WSM6), *J. Korean Meteorol. Soc.*, 42(2), 129–151, 2006.
- Hu, Q., Jiang, D. and Lang, X.: Sources of moisture for different intensities of summer rainfall over the Chinese Loess Plateau during 1979–2009, *Int. J. Climatol.*, 38, e1280–e1287, doi:10.1002/joc.5416, 2018.
- Hu, X. M., Klein, P. M. and Xue, M.: Evaluation of the updated YSU planetary boundary layer scheme within WRF for wind resource and air quality assessments, *J. Geophys. Res. Atmos.*, 118(18), 10490–10505, doi:10.1002/jgrd.50823, 2013.
- Insua-Costa, D. and Miguez-Macho, G.: A new moisture tagging capability in the Weather Research and Forecasting model: Formulation, validation and application to the 2014 Great Lake-effect snowstorm, *Earth Syst. Dyn.*, 9(1), 167–185, doi:10.5194/esd-9-167-2018, 2018.
- Kain, J. S.: The Kain–Fritsch Convective Parameterization: An Update, *J. Appl. Meteorol.*, 43(1), 170–181, doi:10.1175/1520-0450(2004)043<0170:TKCPAU>2.0.CO;2, 2004.
- Mlawer, E. J., Taubman, S. J., Brown, P. D., Iacono, M. J. and Clough, S. A.: Radiative transfer for inhomogeneous atmospheres: RRTM, a validated correlated-k model for the longwave, *J. Geophys. Res. Atmos.*, 102(D14), 16663–16682, doi:10.1029/97JD00237, 1997.
- Mo, K. C. and Juang, H.-M. H.: Relationships between soil moisture and summer precipitation over the Great Plains and the Southwest, *J. Geophys. Res. D Atmos.*, 108(16), GCP 5-1–GCP 5-17, doi:10.1175/1520-0469(1995)052<0879:PMOTSF>2.0.CO;2, 2003.
- Mo, K. C., Nogues-Paegle, J. and Paegle, J.: Physical mechanisms of the 1993 summer floods, *J. Atmos. Sci.*, 52(7), 879–895,

doi:10.1175/1520-0469(1995)052<0879:PMOTSF>2.0.CO;2, 1995.

Mo, K. C., Paegle, J. N. and Higgins, R. W.: Atmospheric processes associated with summer floods and droughts in the central United States, *J. Clim.*, 10(12), 3028–3046, doi:10.1175/1520-0442(1997)010<3028:APAWSF>2.0.CO;2, 1997.

5 Moore, B. J., Neiman, P. J., Ralph, F. M. and Barthold, F. E.: Physical Processes Associated with Heavy Flooding Rainfall in Nashville, Tennessee, and Vicinity during 1–2 May 2010: The Role of an Atmospheric River and Mesoscale Convective Systems*, *Mon. Weather Rev.*, 140(2), 358–378, doi:10.1175/MWR-D-11-00126.1, 2012.

Nakamura, J., Lall, U., Kushnir, Y., Robertson, A. W. and Seager, R.: Dynamical Structure of Extreme Floods in the U.S. Midwest and the United Kingdom, *J. Hydrometeorol.*, 14(2), 485–504, doi:10.1175/JHM-D-12-059.1, 2013.

10 Nayak, M. A., Villarini, G., Bradley, A. A., Nayak, M. A., Villarini, G. and Bradley, A. A.: Atmospheric Rivers and Rainfall during NASA’s Iowa Flood Studies (IFloodS) Campaign, *J. Hydrometeorol.*, 17(1), 257–271, doi:10.1175/JHM-D-14-0185.1, 2016.

Numaguti, A.: Origin and recycling processes of precipitating water over the Eurasian continent: Experiments using an atmospheric general circulation model, *J. Geophys. Res. Atmos.*, 104(D2), 1957–1972, doi:10.1029/1998JD200026, 1999.

15 Pan, Z., Segal, M. and Arritt, R. W.: Role of Topography in Forcing Low-Level Jets in the Central United States during the 1993 Flood-Altered Terrain Simulations, *Mon. Weather Rev.*, 132(1), 396–403, doi:10.1175/1520-0493(2004)132<0396:ROTIFL>2.0.CO;2, 2004.

Pu, B., Dickinson, R. E. and Fu, R.: Dynamical connection between Great Plains low-level winds and variability of central Gulf States precipitation, *J. Geophys. Res. Atmos.*, 121(7), 3421–3434, doi:10.1002/2015JD024045, 2016.

20 Ramos, A. M., Nieto, R., Tomé, R., Gimeno, L., Trigo, R. M., Liberato, M. L. R. and Lavers, D. A.: Atmospheric rivers moisture sources from a Lagrangian perspective, *Earth Syst. Dyn.*, 7(2), 371–384, doi:10.5194/esd-7-371-2016, 2016.

Rife, D. L., Pinto, J. O., Monaghan, A. J., Davis, C. A. and Hannan, J. R.: Global distribution and characteristics of diurnally varying low-level jets, *J. Clim.*, 23(19), 5041–5064, doi:10.1175/2010JCLI3514.1, 2010.

25 Schubert, S. D., Helfand, H. M., Wu, C. Y. and Min, W.: Subseasonal variations in warm-season moisture transport and precipitation over the central and eastern United States, *J. Clim.*, 11(10), 2530–2555, doi:10.1175/1520-0442(1998)011<2530:SIVIWSM>2.0.CO;2, 1998.

Shin, H. H. and Hong, S. Y.: Intercomparison of Planetary Boundary-Layer Parametrizations in the WRF Model for a Single Day from CASES-99, *Boundary-Layer Meteorol.*, 139(2), 261–281, doi:10.1007/s10546-010-9583-z, 2011.

Sorí, R., Marengo, J., Nieto, R., Drumond, A. and Gimeno, L.: The Atmospheric Branch of the Hydrological Cycle over the

- Negro and Madeira River Basins in the Amazon Region, *Water*, 10(6), 738, doi:10.3390/w10060738, 2018.
- Stensrud, D. J.: Importance of low-level jets to climate: A review, *J. Clim.*, 9(8), 1698–1711, doi:10.1175/1520-0442(1996)009<1698:IOLLJT>2.0.CO;2, 1996.
- Stohl, A., James, P., Stohl, A. and James, P.: A Lagrangian Analysis of the Atmospheric Branch of the Global Water Cycle. Part I: Method Description, Validation, and Demonstration for the August 2002 Flooding in Central Europe, *J. Hydrometeorol.*, 5(4), 656–678, doi:10.1175/1525-7541(2004)005<0656:ALAOTA>2.0.CO;2, 2004.
- Stohl, A. and James, P.: A Lagrangian Analysis of the Atmospheric Branch of the Global Water Cycle. Part II: Moisture Transports between Earth's Ocean Basins and River Catchments. *J. Hydrometeorol.*, 6, 961-984, doi:10.1175/JHM470.1, 2005a.
- 10 [Stohl, A.; Forster, C.; Frank, A.; Seibert, P.; Wotawa, G. Technical note: The Lagrangian particle dispersion model FLEXPART version 6.2. *Atmos. Chem. Phys.*, 5, 2461–2474, 2005b.](#)
- Stohl, A., Forster, C. and Sodemann, H.: Remote sources of water vapor forming precipitation on the Norwegian west coast at 60°N - A tale of hurricanes and an atmospheric river, *J. Geophys. Res. Atmos.*, 113(5), 1–13, doi:10.1029/2007JD009006, 2008.
- 15 [Tang, Y., Winkler, J., Zhong, S., Bian, X., Doubler, D., Yu, L. and Walters, C.: Future changes in the climatology of the Great Plains low-level jet derived from fine resolution multi-model simulations, *Sci. Rep.*, 7\(1\), doi:10.1038/s41598-017-05135-0, 2017.](#)
- Ting, M. and Wang, H.: The Role of the North American Topography on the Maintenance of the Great Plains Summer Low-Level Jet*, *J. Atmos. Sci.*, 63(3), 1056–1068, doi:10.1175/JAS3664.1, 2006.
- 20 Trenberth, K. E. and Guillemot, C. J.: Physical processes involved in the 1988 drought and 1993 floods in north America, *J. Clim.*, 9(6), 1288–1298, doi:10.1175/1520-0442(1996)009<1288:PPIITD>2.0.CO;2, 1996.
- Vázquez, M., Nieto, R., Drumond, A. and Gimeno, L.: Moisture transport into the Arctic: Source-receptor relationships and the roles of atmospheric circulation and evaporation, *J. Geophys. Res. Atmos.*, 121(22), 13,493-13,509, doi:10.1002/2016JD025400, 2016.
- 25 [Walters, C. K., Winkler, J. A., Shadbolt, R. P., van Ravensway, J. and Bierly, G. D.: A long-term climatology of southerly and northerly low-level jets for the central United States, *Ann. Assoc. Am. Geogr.*, 98\(3\), 521–552, doi:10.1080/00045600802046387, 2008.](#)
- Wang, S. Y. and Chen, T. C.: The late-spring maximum of rainfall over the U.S. central plains and the role of the low-level jet, *J. Clim.*, 22(17), 4696–4709, doi:10.1175/2009JCLI2719.1, 2009.

Wegmann, M., Orsolini, Y., Vázquez, M., Gimeno, L., Nieto, R., Bulygina, O., Jaiser, R., Handorf, D., Rinke, A., Dethloff, K., Sterin, A. and Brönnimann, S.: Arctic moisture source for Eurasian snow cover variations in autumn, *Environ. Res. Lett.*, 10(5), doi:10.1088/1748-9326/10/5/054015, 2015.

Wexler, H.: A Boundary Layer Interpretation of the Low-level Jet, *Tellus*, 13(3), 368–378, doi:10.1111/j.2153-3490.1961.tb00098.x, 1961.

Wu, Y. and Raman, S.: The summertime Great Plains low level jet and the effect of its origin on moisture transport, *Boundary-Layer Meteorol.*, 88(3), 445–466, doi:10.1023/A:1001518302649, 1998.

10

15

20

25

30

Captions

Figure 1: (a) Mean NLLJ index (shaded) and 500 m winds (arrows, in m s^{-1}) at local midnight in July (boreal summer) for 1980-2016, calculated from ERA-Interim reanalysis. The black cross at 32.75°N , 99°W shows the point of maximum NLLJ in the climatology. The cyan contour line surrounds the region containing points above the 75th percentile. (b) Frequency distributions of the GPLLJ for the months of July from 1980 to 2016 (blue bars). The red curve corresponds to the Gaussian fit: $y(x) = y_0 + A \exp(-\frac{(x-y)^2}{2\sigma^2})$. (see table A2). Noted: The frequency distributions are calculated at the point of maximum intensity of NLLJ (at 32.75°N , 99°W , black cross in fig. 1a).

Figure 2: Highlighted in red are moisture sources obtained with FLEXPART from backward trajectories originating in the region outlined in magenta in Fig 1.a. The orange line over the continent marks the position from where precipitable water is tagged in WRF-TFWVT, corresponding to the northern edge of the FLEXPART source region. All water vapour and condensate crossing through that line is considered as moisture advected by the GPLLJ for further analysis.

Figure 3: Ratio of tagged precipitable water transported by the GPLLJ to total for the six case-studies analysed.

Figure 4: Statistically weighted ratio of precipitable water transported by the GPLLJ for the five case studies with $\text{NLLJ} > 0$ considered in the analysis in Figure 3. Weights applied are stated in Table 1.

Figure 5: Tracer total precipitable water (TPW, g kg^{-1}) and positions of the cross sections along the central axis of the GPLLJ shown in Fig. 5, at latitudes 32°N (1), 35°N (2) and 38°N (3) for the main jet event of July 11, 2002-1992.

Figure 6: (a-c) $q\text{TR}$ in g kg^{-1} for the three vertical cross sections at the locations depicted with white lines in Fig. 5. (d-f) same as (a-c) but for ϕTR in g m (kg s)^{-1} .

Figure 7: (a-c) Ratio of $q\text{TR}:q$ for the three vertical cross sections at the locations depicted with white lines in Fig. 5. (d-f) same as (a-c) but for ratio of $\phi\text{TR}:\phi$.

Table 1: Case-studies objectively selected based in the frequency distribution of the LLJ index to carry out WRF-TFWVT simulations. μ is the mean of the distribution and σ its standard deviation. Noted: The frequency distribution is calculated at the point of maximum intensity of NLLJ at 32.75°N , 99°W (black cross in fig. 1a) using the ERA-Interim reanalysis dataset.




Figure 1

5

10

15

20

25




Figure 2

30

35

5

10

15

20

25




Figure 3

30

35

5

10

15

20

25




Figure 4

30

35


5

10

15

20

25



30 **Figure 5**

35




Figure 6




Figure 7

5

10

Table 1

15

Simulation	Gaussian	NLLJ value	Date	Stat. weight
0		0.00	2012-07-12	0
1	$\mu - 2\sigma$	1.49	1999-07-19	0.0623
2	$\mu - \sigma$	5.54	1983-07-23	0.2445
3	μ	10.19	1992-07-11	0.3864
4	$\mu + \sigma$	14.54	2002-07-28	0.2445
5	$\mu + 2\sigma$	18.89	2016-07-14	0.0623

20

25

Appendix A: Statistical weights in the analysis

Table A1. Events

Simulation	Gaussian Point
1	$\mu - 2\sigma$
2	$\mu - \sigma$
3	μ
4	$\mu + \sigma$
5	$\mu + 2\sigma$

Table A2. Gaussian fit $y(x) = y_0 + A \cdot \exp(-\frac{(x-y)^2}{2\sigma^2})$.

Simulation	Gaussian Point
y_0	0.03 ± 3.66
A	8.63 ± 4.39
μ	10.19 ± 0.19
σ	4.35 ± 0.30
R^2	0.95

5

$$StatWeight_1 = StatWeight_5 = \frac{\int_{\mu-2.5\sigma}^{\mu-1.5\sigma} y(x)dx}{\int_{\mu-2.5\sigma}^{\mu+2.5\sigma} y(x)dx} = 0.0623$$

$$StatWeight_2 = StatWeight_4 = \frac{\int_{\mu-1.5\sigma}^{\mu-0.5\sigma} y(x)dx}{\int_{\mu-2.5\sigma}^{\mu+2.5\sigma} y(x)dx} = 0.2446$$

10

$$StatWeight_3 = \frac{\int_{\mu-1.5\sigma}^{\mu+0.5\sigma} y(x)dx}{\int_{\mu-2.5\sigma}^{\mu+2.5\sigma} y(x)dx} = 0.38643$$

On the assessment of the moisture transport by the Great Plains low-level jet

Iago Algarra¹, Jorge Eiras-Barca¹, Gonzalo Miguez-Macho², Raquel Nieto¹, and Luis Gimeno¹

¹EPhysLab (Environmental Physics Laboratory), Facultade de Ciencias, Universidade de Vigo, Ourense, Galicia, Spain

²Non-Linear Physics Group, University of Santiago de Compostela, Galicia, Spain

Correspondence to: Iago Algarra (ialgarra@uvigo.es)

Abstract. Low-Level Jets (LLJs) can be defined as ~~filamentous~~-wind corridors of anomalously high wind speed values located within the first km of the troposphere. These structures, ~~together with atmospheric rivers (ARs),~~ are one of the major meteorological systems in the meridional transport of moisture on a global scale. In this work, we focus on the southerly Great Plains low-level jet, which plays an important role in the moisture transport balance over the central United States. The Gulf of Mexico is the main moisture source for the GPLLJ, which has been identified as a key factor for rainfall modulation over the eastern and central US.

The relationship between moisture transport from the Gulf of Mexico to the Great Plains and precipitation is well documented in previous studies. Nevertheless, a large uncertainty still remains in the quantification of the moisture amount actually carried by the GPLLJ. The main goal of this work is to address this question. For this purpose, a relatively new tool, the regional atmospheric Weather Research and Forecasting Model with 3D water vapour tracers (WRF-~~TTWVT~~, Insua-Costa and Miguez-Macho, 2018) is used together with the Lagrangian model FLEXPART to estimate the load of precipitable water advected within the GPLLJ. Both models were for ledfed with data from ERA Interim. From a climatology of jet intensity over a 37-year period (~~Rife et al., 2010~~), which follows a Gaussian distribution, we select for study 5 cases representing the mean, and one and two standard deviations above and below it. Results show that the jet is responsible for roughly 70%-80% of the moisture transport occurring in the southern Great Plains when a jet event occurs. Furthermore, moisture transport by the GPLLJ extends to the northeast US, accounting for 50% of the total in areas near the Great Lakes. Vertical distributions show the maximum of moisture advected by the GPLLJ at surface levels and maximum values of moisture flux about 500 m above, in coincidence with the wind speed profile.

1 Introduction

It is well known that the Great Plains Low-Level Jet (hereafter, GPLLJ) plays an important role in the balance of the moisture transport over the central United States (Stensrud, 1996; Schubert et al., 1998). The atmospheric moisture is transported by the GPLLJ from tropical and subtropical latitudes (particularly the Gulf of Mexico and the Caribbean Sea) into the Great Plains

(Helfand and Schubert, 1995; Mo et al., 1997) where the jet is often responsible for nocturnal deep convective activity (Higgins et al., 1997; Pu et al., 2016). In the last decades, a large number of authors have shown the strong influence of the GPLLJ as a modulator of climate and rainfall over this region and even further east (Mo et al., 1995, 1997; Wu and Raman, 1998; Byerle and Paegle, 2003); for instance, throughout May and June it is estimated that at least one-third of the moisture penetrating into the continental US is carried by the GPLLJ (Helfand and Schubert, 1995).

Among the mechanisms which have been proposed as triggers of the GPLLJ are included a combination of inertial oscillations (Blackadar, 1957) and orographic forcing (Wexler, 1961; Byerle and Paegle, 2003; Pan et al., 2004; Ting and Wang, 2006). Particularly, the mechanism of Blackadar (1957) suggests that inertial oscillations near the friction layer can induce the formation of the GPLLJ (Wu and Raman, 1998). Nevertheless, orographic effects are also understood as a key factor in the maintenance of the GPLLJ. In this regard, Ting and Wang (2006) proved that, when the interaction with the orography is removed from numerical simulations, the GPLLJ vanishes, together with an important amount of the summer precipitation over the central and southern-southerly US.

The GPLLJ is a phenomenon confined within the first kilometres of the troposphere and is closely related to the warm season (Bonner, 1968). Besides, it is characterized by a strong diurnal oscillation, with a peak in strength during night hours (Augustin and Caracena, 1994). [A long-term climatology of GPLLJ can be found in the work of Walters et al. \(2008\).](#) The GPLLJ is a phenomenon extremely localized in time and space and its role in the continental moisture balance is difficult to study solely from observations.

Nevertheless, a large number of studies have documented the relationship between the major moisture transport and the GPLLJ. Higgins et al. (1996) studied the moisture budget over the central US in May employing NASA/DAO and NCEP/NCAR datasets, together with station observations, to evaluate the limitations of these products. Although both reanalyses overestimate daily mean precipitation rates, they accurately capture the basic temporal and structural characteristics of the GPLLJ. From the data, these authors calculated an increase in atmospheric moisture transport from the Gulf of Mexico during nighttime of more than 50%. In a later work, Higgins et al. (1997b) observed a well-defined nocturnal maximum of precipitation over the Great Plains in spring and summer by analysing station data. [Particularly and linked to LLJ events they found in the region an excess of 25% in nocturnal rainfall during summer when compared with the diurnal precipitation, associated with a rainfall decrease over the Gulf of Mexico. Particularly, this research found over the region an excess of 25% in nocturnal precipitation during summer when compared with the diurnal one, associated with a rainfall decrease over the Gulf of Mexico.](#) Additionally, [Higgins et al. \(1997\) this work reveals reported](#) significant differences in precipitation pattern in coincidence (or not) with LLJ events. When a LLJ event occurs, the observations show an enhanced precipitation over the north-central United States and the Great Plains region, together with a decrease along the Gulf of Mexico and the western Atlantic (Mo et al., 1997). On the other hand, Mo and Juang (2003) found a regional [dependence-correlation at a distance](#) between evaporation and precipitation, reflected in evaporation anomalies over the Great Plains along the trajectory of the GPLLJ, which are associated with downstream precipitation anomalies.

~~All these results are consistent with the large-scale atmospheric moisture transport and support the marked influence of the GPLLJ over the central-eastern US, which has been shown to trigger more than 60% of the spring local precipitation there (Wang and Chen, 2009).~~

Otherwise, extreme rainfall events in the central US are related to an increase in moisture convergence downwind of the GPLLJ (Mo et al., 1997). A decisive factor that triggers heavy rains and floods is the presence of moisture advected by the GPLLJ from the Gulf of Mexico and the Caribbean Sea. Moore et al. (2012) reported the physical processes related to the floods in May 2010, when a persistent southerly LLJ associated with an atmospheric river (AR) enhanced the transport of moisture from the Gulf of Mexico into the heavy rainfall region. Thus, important socioeconomic impacts follow enhanced GPLLJ events, which modulates a large percentage of the local extreme precipitation events and flooding in warmer months (Mo et al., 1995, 1997; Beljaars et al., 1996; Trenberth and Guillemot, 1996; Arritt et al., 1997; Nakamura et al., 2013; Nayak et al., 2016). All these results are consistent with the large-scale atmospheric moisture transport and support the marked influence of the GPLLJ over the central-eastern US, which has been shown to trigger more than 60% of the spring local precipitation over the Great Plains region~~there~~ (Wang and Chen, 2009).

During the last decades, the GPLLJ has experienced a strengthening, accompanied by a northward migration causing a displacement of rainfall in the same direction. As a result, more ~~frequent~~~~common~~ droughts have been observed in the southern Great Plains (Barandiaran et al., 2013). Besides, ~~T~~the increase in the number and intensity of GPLLJ events is also forecasted for future projections, which reveal an intensification of the GPLLJ during the spring season associated with global warming (Cook et al., 2008; Tang et al., 2017). As a result, increasing amounts of moisture transport and rainfall are expected, particularly from April to July, over the central US (Harding and Snyder, 2014). The same projections forecast a slight weakening of the GPLLJ from August to December, which could translate into increasing drought conditions.

The knowledge about the GPLLJ, together with the insights on the relationship between the moisture transported by the GPLLJ and local precipitation patterns has increased considerably during the last decades. However, there are still unanswered questions about the quantification of such water vapour transport and specially about the estimation of the ratio of land to oceanic moisture sources associated with the GPLLJ. This estimate of the oceanic input to the moisture transport associated with the GPLLJ is essential to predict and understand the behaviour of the GPLLJ in future scenarios.

In this work, a new tool, the regional atmospheric Weather Research and Forecasting Model with 3D water vapour tracer diagnostics (WRF-~~TTWVT~~, ~~Eiras-Barea et al., 2017~~; Insua-Costa and Miguez-Macho, 2018) is used to quantify the total amount of total precipitable water (TPW) transported by the GPLLJ. To show the differences between the transport of moisture on jet and non-jet days, a 37-year climatology was ~~calculated~~ previously calculated ~~and the at the point~~ of maximum jet intensity is obtained following the methodology by Rife et al., (2010). The structure of this work is: ~~as follows~~, in Section 2 we provide the methodology used, in Section 3 we show the results obtained, and finally in Section 4 we discuss the conclusions.

2 Data and methods

2.1 Detection of the Great Plains low-level jet

To objectively detect days with LLJ over the Great Plains, we applied the night-time index proposed in Rife et al., (2010), hereafter named as NLLJ. This index is based on the temporal variation of the wind's vertical structure and the fact that LLJs are most intense at local midnight. Because both frequency and intensity of GPLLJ are mostly associated with the warm season, we develop a 37-year climatology for the month of July (~~representative for~~ representative for the boreal summer). According to the NLLJs characteristics, and with the aim to define the index, two conditions should be met to consider a GPLLJ detection:

1. The wind speed is higher at local midnight than at midday.
2. The local midnight wind speed is higher at the surface ($\sim 500\text{m}$) than ~~in height~~ above it ($\sim 4\text{km}$).

The index is calculated at each grid point over an area centred over the US using 6-hourly ERA-Interim reanalysis data (Dee et al., 2011) with a 0.25° horizontal resolution. ~~Due to~~ Because of the jet core is located within of the first kilometre of the troposphere, it is necessary to take into account the elevation of the land, so sigma coordinates are used. The GPLLJ-climatology is developed for 37 years, from 1980 to 2016, and the NLLJ index can be defined as follows:

$$NLLJ = \lambda \varphi \sqrt{[(u_{00}^{L1} - u_{00}^{L2}) - (u_{12}^{L1} - u_{12}^{L2})]^2 + [(v_{00}^{L1} - v_{00}^{L2}) - (v_{12}^{L1} - v_{12}^{L2})]^2} \quad (1)$$

where u and v are the zonal and meridional wind components, respectively. $L1$ represents the winds at the surface at 53 sigma level (elevation near the jet core), approximately 500 m above ground level (AGL), while $L2$ corresponds to the wind at 42 sigma level (around 4000 m AGL). Numbers 00 and 12 refer to local midnight and local noon, respectively. λ and φ are binary multipliers representing the temporal and vertical variation of ~~the~~ the wind. Particularly, λ relates to the daily variation of the wind at 500 m and φ refers to the wind's vertical variation between 500 m and 4 km at midnight (Rife et al., 2010):

$$\lambda = \begin{cases} 0, & ws_{00}^{L1} \leq ws_{12}^{L1} \\ 1, & ws_{00}^{L1} > ws_{12}^{L1} \end{cases} \quad (2)$$

$$\varphi = \begin{cases} 0, & ws_{00}^{L1} \leq ws_{00}^{L2} \\ 1, & ws_{00}^{L1} > ws_{00}^{L2} \end{cases} \quad (3)$$

2.2 Identification of moisture sources associated with Great Plains low-level jet

For the objective identification of moisture sources associated with the GPLLJ, the Lagrangian backward trajectories from the FLEXPART v9.0 model are used (Stohl et al., 2005a). This model provides a large number of air parcel trajectories from which it is possible ~~to the calculation of~~ to calculate the evaporation minus precipitation budget, tracking all changes in the specific humidity of air parcels.

FLEXPART has been widely and successfully used to track moisture paths for the study of the atmospheric branch of the hydrologic cycle in different parts of the world (e.g., Hu et al., 2018; Sorí et al., 2018; Vázquez et al., 2016). Furthermore, this tool is able to infer the moisture sources for precipitation falling in a target region when backward trajectories are considered (eg., Sthol et al., 2008; Drumond et al., 2012; Gimeno et al., 2012; Wegmann et al., 2015; Ramos et al., 2016).

5 In this work we use the outputs of a global experiment in which FLEXPART v9.0 tracks approximately 2 million particles (air parcels) with constant mass distributed on the globe every time step during 37-year period (1980-2016). These air parcels are advected by the 3D wind field, and the variables of interest of each particle (such as position, height, specific humidity, temperature among many others) are saved at each time step. We perform a FLEXPART simulation fed with ERA-Interim reanalysis data at 1° horizontal resolution on 61 vertical levels from sea level pressure to 0.1 hPa and 6-hour time intervals (00, 06, 12 and 18 UTC). The model is run with a 3 h timestep, and linear interpolation is used to obtain data with this frequency from ERA-Interim. The backward trajectories are followed [during-for](#) 10 days, which is the average life time of water vapour in the atmosphere (Numaguti, 1999).

The changes in specific humidity (q) of each air parcel along its path can be expressed as follows:

$$15 \quad e - p = m \frac{dq}{dt} \quad (4)$$

where m is the mass of a particle (which remains constant in the simulation), q is the specific humidity, t the time step, and $e - p$ (evaporation minus precipitation) represents the water flux associated with the particle. To obtain the instantaneous values of the $E - P$ balance ([E denotes evaporation and P the precipitation rate per unit area](#)) in a given area (in this case, over one of [1-0° x 1-0 degrees](#) in latitude and longitude), it is necessary to integrate the moisture changes for all particles present in the atmospheric column over such area ([E denotes evaporation and P the precipitation rate per unit area](#)) ([Stohl and James 2004, 2005b](#)). Thus, [following this methodology](#), in a backward experiment, a moisture source is defined as those regions where $E - P$ is positive ($E - P > 0$), which implies that evaporation exceeds precipitation, while a moisture sink is defined as a region where $E - P < 0$, meaning that precipitation is greater than evaporation.

25 In our study, backward trajectories were followed from the area composed of points with values of NLLJ above [the percentile 75th percentile to foundfind the main moisture source.-](#)

2.3 The regional atmospheric Weather Research and Forecasting Model with 3D water vapour tracer diagnostics (WRF-~~TTWVT~~)

The mesoscale Weather Research and Forecast model (WRF 3.8.1) with the moisture tracers tool (WRF-~~TTWVT~~, [Eiras Barea et al., 2017](#); Insua-Costa and Miguez-Macho, 2018) is used [to carry out](#) to quantify the ~~total~~ amount of total precipitable water (TPW) transported by the GPLLJ. In order to analyse the moisture transport associated with the GPLLJ avoiding the effects of other synoptic-scale transport events, we tag the moisture passing northward through a narrow wall located on the northern edge of the moisture source region identified using the FLEXPART model. When a particle of water (whether in liquid, solid

or gas state) crosses the wall, it is labeled for further analysis inside the simulation domain. We consider all water traversing the wall to be advected by the GPLLJ.

The horizontal resolution of the simulations is 20 km and the vertical column is divided into 38 levels. The simulation covers a time window of 11 days, starting 7 days prior to the day of interest. The model parameterizations together with the WRF-~~TT~~ ~~WVT~~ are set using the PBL Yonsei University (YSU) parametrization (Hu et al., 2013; Shin and Hong, 2011), the schemes of Kain-Fritsch for convection (Kain, 2004), the Dudhia one for short-wave radiation (Dudhia, 1989), the Rapid Radiative Transfer Model (RRTM) scheme for long-wave radiation (Mlawer et al., 1997), and the WRF Single-Moment 6-Class Microphysics Scheme (WSM6) (Hong and Lim, 2006).

In addition, we apply spectral nudging of waves longer than 1000 km above the boundary layer, with a relaxation time of 1h, to avoid ~~30~~-distortion of the large-scale circulation. This configuration has been validated and successfully applied several times with the WRF-~~TT~~-~~WVT~~ in mid latitudes (e.g., Eiras-Barca et al., 2017). Spectral nudging ensures that the large-scale circulation is well captured in the simulations. ERA-Interim data provides lateral boundary and initial conditions for the runs (Dee et al., 2011). The variables of interest for the analysis of the GPLLJ event are computed as follows. Integrated Water Vapour (IWV), Eq. (5) is obtained by vertical integration of the specific humidity (q) in pressure (p) levels, where g represents the gravitational force. The instant flux of moisture ($\sigma\phi$) is calculated as stated in Eq. (6) and the conversion between (g) and the water vapour mixing ratio obtained from WRF is performed using Eq. (7), where u and v are the horizontal components of the wind field.

$$IWV_{(i,j)} = \frac{1}{g(k)} \int_{surface}^{top} q(i,j,k) dp \quad (5)$$

$$\sigma\phi(i,j,k) = |q \cdot (u,v)| \quad (6)$$

$$q = \frac{w}{w+1}, \text{ with } w \ll 1 \rightarrow q \approx w \quad (7)$$

3 Results

3.1 Characterization of the Great Plains low-level jet

As previously mentioned, the NLLJ index was calculated at each grid point over the North American region for the month of July over the period 1980-2016. July was found to be the month with maximum LLJs events (Figure S1 shows the monthly distribution of the GPLLJ days of detection). Fig. 1.a shows the climatological NLLJ index and the wind field at 500 hPa. The black cross indicates the point of maximum intensity of the index (8.8 m s^{-1}). At this point, located at 32.75°N - 99°W , along the 37-years analysed, and for July, a total of 931 LLJ days are identified, that is, 81% of all days have a positive value of the index. On the point of maximum intensity ~~showed~~ shown in Fig1.a, Fig1.b displays the frequency distribution of the NLLJ for the period 1980-2016. A clear peak around 11 m s^{-1} is observed together with a Gaussian behaviour ($R^2=0.95$, ~~red line~~ Jarque-

Bera test p-value=0.0055, which provides a confidence level close to 99.5%, red line). The latter has been used to select the five case studies to be ~~analyzed~~ analysed with WRF-~~TT~~ WVT and listed in Table 1. The five ~~events~~ case-studies were selected based on the Gaussian adjustment applied to the study. These five events ~~chosen~~ correspond to μ , $\mu \pm 2\sigma$, $\mu \pm \sigma$ (where μ is the mean of the distribution and σ its standard deviation), and they provide a general perspective of the LLJ's behaviour. Selecting these case studies from the population of LLJ events decreases the computational expense. Since each case-study WRF-~~TT~~ WVT simulation spans ~~for~~ 11 days, a condition of persistence of the index value for at least two days after the main jet day is applied. Additionally, we perform a sixth simulation with a non-jet day (simulation 0 in the Table 1). This non-jet day is selected from the developed climatology as the longest non-jet period, in order to avoid overlaps in moisture transport with jet days.

10 3.2 Moisture transport associated with the Great Plains low-level jet

In order to detect the main climatological oceanic source of moisture for the GPLLJ we used the FLEXPART trajectories outputs for 1980-2016. The area ~~englobed-encompassed~~ in the 75th percentile of the NLLJ index values (~~cyan-magenta~~ line in figure 1.a) was selected as the target region for the backward experiment (as ~~it was~~ explained in the methodology). Fig. 2 shows the source of moisture in red colour, obtained as the 75th percentile of the $(E - P > 0)$ field. This area covers the southern Gulf of Mexico and extends into the Caribbean Sea, between 60°-98°W and 12°-28°N. Figure S2 in the supplementary material shows the individual sources of moisture for each case in study).

Although the flow originated in the source of moisture is advected in the low levels as a result of the strong intensity of the trade winds, a 3D-label wall (at 29-30°N and from 94.5°W to 100°W) was used in the WRF-~~TT~~ WVT simulations (orange line in fig. 2). The position of the sentinel wall was selected on the region where oceanic moisture associated with the GPLLJ ~~makes~~ landfalls. The wall remained constant in the WRF-~~TT~~ WVT simulations. ~~A~~The thin wall was used instead of the entire source regions in order to avoid overlaps in the labelling of moisture caused by secondary synoptic scale mechanisms.

Fig. 3 shows the ratio of precipitable water transported by the GPLLJ to total precipitable water ($TPW_{tracers}=TPW$) for the six case-studies analysed. As mentioned earlier, $TPW_{tracers}$ represents the TPW that has crossed the "wall" highlighted in orange in Figure ~~4~~2, which we assume corresponds to moisture advected by the GPLLJ. Following the same behaviour of the GPLLJ itself, the moisture flux is initially in the northward direction and veers east as it penetrates into the Great Plains for all events with positive NLLJ index values. As expected, the non-jet event with NLLJ value equal to zero (Sim 0) does not show ~~remarkable-significant~~ moisture fluxes. For the jet events, ratios are close to one in regions near the tagging wall and extend for hundreds of km northward with significant values above 60%. Percentages between 70% and 80% are observed in the Great Plains. The large geographical reach of the moisture associated with the GPLLJ is evidenced in this figure, showing that for certain GPLLJ events it can occasionally explain more than 50% of TPW even in the north-east US. It is necessary to highlight that higher values in the index value does not necessarily mean larger flows of moisture ~~in the entire~~, as can be observed, for example, when SIM3 and SIM5 are compared. -Figure S3 in the supplementary material shows a comparison of accumulated precipitation at 11 days of WRF simulations versus CPC gauge-analysis observations (Chen et al., 2008).

As ~~it was~~ previously stated, the aim of this work is to study the general behaviour of the GPLLJ associated with its moisture transport. In the first simulated case of GPLLJ (fig.3 – Sim 1) it is observed that most of the precipitable water is concentrated on the Great Plains, exceeding ratios of 80% out of the total. In the second GPLLJ event simulated (fig.3 – Sim 2), the precipitable water extends northeast of the US and to the south of the Great Lakes and the GPLLJ, where it explains close ~~of~~to the 50% of precipitable water. The third simulated case corresponds to the average behaviour of the GPLLJ (fig.3 – Sim 3) and evidences-shows the influence of the GPLLJ in the northeast of US with ratios near 50% in the US East Coast. Nevertheless, in areas along the path of the GPLLJ, the advection of precipitable water is close to 80%. In the fourth and fifth simulations of GPLLJ (fig.3 – Sim 4 and 5), the plume of precipitable water is concentrated over the Great Plains. However, the water precipitable ratio is reduced as latitude increases, but values are still close to 50% in the northeastern areas of the US.

Fig. 4 shows the statistically weighted mean of the ratio shown in Fig. 3 for the five case studies with NLLJ > 0 considered in the analysis. The weights associated with each event are stated in the last column of Table 1 ~~at the last column~~, and the objective criteria to assign them can be found in appendix A1. The aim of using weights in the analysis is to give greater importance to the event representing the mean value of the NLLJ and less relevance to the events in the tail of the distribution. Notwithstanding the limited number of simulations used in the analysis, this procedure allows us to interpret Figure 4 as a "climatology" of the moisture transport associated with the GPLLJ. Roughly 80-90% of the precipitable water in its core zone of influence over the Great Plains, in Texas and Oklahoma, is carried by the GPLLJ when a jet event occurs. With increased distance from that area, the ratio of precipitable water transported by the GPLLJ decreases; however, the contribution of moisture from the Gulf of Mexico to TPW is still ~~of~~ more than 50% as far north as the Great Lakes.

On the one hand, Fig. 5 shows the TPW and the cross sections for the main GPLLJ event (1992.07.11). On the other hand,

Fig. ~~5~~6 shows the vertical distribution of tracer specific humidity (q_{TR}) and tracer water vapour flux (ϕ_{TR}) for cross sections at positions depicted in Fig. ~~6-5~~for the main GPLLJ event (1992.07.11). Tracer moisture (a-c) has a maximum at surface levels, while the moisture flux (d- f) maximizes at 500 m AGL where the LLJ core is located. A significant presence of both tracer water vapour and tracer water vapour flux is restricted to the first 2 km AGL. Overall, as the latitude increases the water vapour plume from the Gulf of Mexico tends to rise in the vertical column and expand zonally along the GPLLJ path to the east of the

U.S. Equivalent conclusions can be obtained from the remaining events, which are shown in supplementary material ~~S4~~(Figs. S4-S7). Figure 7 shows the ~~steam~~water vapor ratio $q_{TR}:q$ for the vertical sections of Fig. 5 for the same event. The moisture pattern behaviour is similar to Fig. 6; for regions close to the Gulf of Mexico the moisture ratio is concentrated mostly at lower levels (Fig. 7a) and extends on the horizontal axis as it moves away from the source of moisture (Fig. 7b). Nevertheless, for remote regions the maximum moisture ratio continues at lower levels (Fig 7c). -The moisture flow has the same pattern, high moisture ratios that remain at low levels despite distancing longitudinal distance (Fig. 7d-7f). These results confirm that the GPLLJ transports a great concentrated quantity of moisture.

4 Conclusions and discussions

A combination of Lagrangian and Eulerian methods were used to identify and objectively quantify the moisture transport associated with the Great Plains Low-Level Jet (GPLLJ). First, the FLEXPART ~~L~~agrangian model ~~FLEXPART~~ was used to locate the GPLLJ moisture sources ~~in the Gulf of México~~ for the month of its maximum activity (July) throughout the period 1980-2016. The target region used in the FLEXPART simulation to find the main source of moisture was defined based on the 75th percentile of the GPLLJ index value previously calculated based in Rife et al. (2010) method. Once the Gulf of Mexico was identified as the main source of moisture ($E - P > 0$) for the GPLLJ, we use a new tool known as eulerian 3D WRF-~~WTT~~ WVT (~~Eiras-Barea et al., 2017;~~ Insua-Costa and Miguez-Macho, 2018) which was applied to track the moisture advected in six selected GPLLJ events based on the distribution of the GPLLJ index used previously to detect the GPLLJ (Rife et al., 2010). ~~Se~~Consequently, ~~F~~this work analysed the behaviour of the GPLLJ during the month of its maximum activity (July) for ~~thea long~~ period, 1980-2016, and we select six representative cases for the WRF-WVT simulation.

The moisture transport analysis reveals the major role played by the GPLLJ in the water cycle of central North America, transporting large amounts of moisture from the Gulf of Mexico as far as the north-east US. Particularly, advection by the jet explains more than 80% of the precipitable water in the southern Great Plains when a jet event occurs, which, in July, is most of the days. The Rocky Mountains blocks the circulation of GPLLJ and force it to turn to the east of the US, reaching even the eastern coast of the US. The moisture transport associated with the GPLLJ is in a plume of moisture where, the maximum transport occurs in the path of the GPLLJ. As expected, the ratio reduces as latitude increases, but values are still close to 50% in the ~~nort~~heastern areas of the US.

We note that the extension of the GPLLJ is dependent on the synoptic conditions or land preconditioning, among other factors, which may alter the ratio of the TPW. ~~are Nevertheless, the analysis of these multiple factors is out of the scope of this paper.~~ For example, the presence of a well-developed high pressure system in higher latitudes of North America may block the advection of the GPLLJ moisture to this region. Dong et al., (2011) related the drought of 2006 with an anomalous high over south-western U.S region and an anomalous low over the Great Lakes. This pattern inhibited the advection of moisture from the Gulf of Mexico contributing to the extreme dryness, and the lack precipitation was associated with a suppressed cyclonic activity over the south-western US. ~~However~~On the other hand, the 2007 flood events were initially ~~preceded~~leaded by active synoptic weather patterns, linked to an active moisture transport from the Gulf of Mexico by the GPLLJ. Nevertheless, the analysis of these multiple factors is out of the scope of this paper.

~~Besides,~~ hHigher values in the NLLJ index mean larger differences between winds aloft and at the surface at the reference point, but do not necessarily mean stronger moisture transport.

~~Thus,~~These results should be understood as a first approach to the quantification of the large extent of GPLLJ moisture advection and its implications for the water budget in North America. More ~~WRF TT~~ simulations should be conducted, and other months should be included in FLEXPART backward calculations to extend this work and produce a more comprehensive analysis.

Author contributions.

IA, RN and LG had the initial idea. GMM developed WRF-WVT tool. IA and JEB carried out the simulations and data analysis. [IA and JEB wrote the original draft and reviewed the subsequent documents.](#) RN, LG and GMM provided suggestions [under the revision period, commented and reviewed the manuscript before submission.](#)

5 Competing interests.

The authors declare that they have no conflict of interest.

Acknowledgments.

~~The ECMWF ERA Interim reanalysis data were obtained from <https://www.ecmwf.int/en/forecasts/datasets/reanalysis-datasets/era-interim>.~~ Iago Algarra was financially supported by Spanish government (MINECO), (CGL2015-65141-R). [Jorge Eiras-Barca has been financially supported by the EDB481B 2018/069 from the “Xunta de Galicia” and the Fullbrigh commission. This work was also partially supported by Xunta de Galicia under Project ED431C 2017/64-GRC ‘Programa de Consolidación e Estructuración de Unidades de Investigación Competitivas \(Grupos de Referencia Competitiva\).’](#) Iago Algarra would like to express his gratitude to ~~the~~ all [the](#) Non-Linear Physics Group for [their](#) kind support during ~~his~~[their](#) stay in the University of Santiago de Compostela.

15

References

- Arritt, R. W., Rink, T. D., Segal, M., Todey, D. P., Clark, C. A., Mitchell, M. J., Labas, K. M., Arritt, R. W., Rink, T. D., Segal, M., Todey, D. P., Clark, C. A., Mitchell, M. J. and Labas, K. M.: The Great Plains Low-Level Jet during the Warm Season of 1993, Mon. Weather Rev., 125(9), 2176–2192, doi:10.1175/1520-0493(1997)125<2176:TGPLLJ>2.0.CO;2, 1997.
- 20 Augustine, J. A. and Caracena, F.: Lower-tropospheric precursors to nocturnal MCS development over the central United States, Weather Forecast., 9(1), 116–135, doi:10.1175/1520-0434(1994)009<0116:LTPTNM>2.0.CO;2, 1994.
- Barandiaran, D., Wang, S.-Y. and Hilburn, K.: Observed trends in the Great Plains low-level jet and associated precipitation changes in relation to recent droughts, Geophys. Res. Lett., 40(23), 6247–6251, doi:10.1002/2013GL058296, 2013.
- Beljaars, A. C. M., Viterbo, P., Miller, M. J. and Betts, A. K.: The Anomalous Rainfall over the United States during July 1993: Sensitivity to Land Surface Parameterization and Soil Moisture Anomalies, Mon. Weather Rev., 124(3), 362–383, doi:10.1175/1520-0493(1996)124<0362:TAROTU>2.0.CO;2, 1996.
- 25 Blackadar, A. K.: Boundary layer wind maxima and their significance for the growth of nocturnal inversions, Bull. Amer. Meteor. Soc., 38, 283–290, doi:10.1175/1520-0477-38.5.283, 1957.

- Bonner, W. D.: Climatology of the low level jet, *Mon. Weather Rev.*, 96(12), 833–850, doi:10.1175/1520-0493(1968)096<0833:COTLLJ>2.0.CO;2, 1968.
- Byerle, L. A. and Paegle, J.: Modulation of the Great Plains low-level jet and moisture transports by orography and large-scale circulations, *J. Geophys. Res. D Atmos.*, 108(16), GCP 6-1-GCP 6-16, doi:10.1029/2002JD003005, 2003.
- 5 [Chen, M., Shi, W., Xie, P., Silva, V. B. S., Kousky, V. E., Higgins, R. W. and Janowiak, J. E.: Assessing objective techniques for gauge-based analyses of global daily precipitation, *J. Geophys. Res. Atmos.*, 113\(4\), 1–13, doi:10.1029/2007JD009132, 2008.](#)
- Cook, K. H., Vizy, E. K., Launer, Z. S. and Patricola, C. M.: Springtime intensification of the great plains low-level jet and midwest precipitation in GCM Simulations of the twenty-first century, *J. Clim.*, 21(23), 6321–6340, doi:10.1175/2008JCLI2355.1, 2008.
- 10
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N. and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q. J. R. Meteorol. Soc.*, 137(656), 553–597, doi:10.1002/qj.828, 2011.
- 15
- Dong, X., Xi, B., Kennedy, A., Feng, Z., Entin, J. K., Houser, P. R., Schiffer, R. A., L’Ecuyer, T., Olson, W. S., Hsu, K. L., Liu, W. T., Lin, B., Deng, Y. and Jiang, T.: Investigation of the 2006 drought and 2007 flood extremes at the Southern Great Plains through an integrative analysis of observations, *J. Geophys. Res. Atmos.*, 116(3), doi:10.1029/2010JD014776, 2011.
- 20
- Drumond, A., Nieto, R., Trigo, R., Ambrizzi, T., Souza, E. and Gimeno, L.: A lagrangian identification of the main sources of moisture affecting northeastern Brazil during its pre-rainy and rainy seasons, *PLoS One*, 5(6), 1–8, doi:10.1371/journal.pone.0011205, 2010.
- Dudhia, J.: Numerical Study of Convection Observed during the Winter Monsoon Experiment Using a Mesoscale Two-Dimensional Model, *J. Atmos. Sci.*, 46(20), 3077–3107, doi:10.1175/1520-0469(1989)046<3077:NSOCOD>2.0.CO;2, 1989.
- 25
- Eiras-Barca, J., Dominguez, F., Hu, H., Garaboa-Paz, D. and Miguez-Macho, G.: Evaluation of the moisture sources in two extreme landfalling atmospheric river events using an Eulerian WRF tracers tool, *Earth Syst. Dyn.*, 8(4), 1247–1261, doi:10.5194/esd-8-1247-2017, 2017.
- Gimeno, L., Stohl, A., Trigo, R. M., Dominguez, F., Yoshimura, K., Yu, L., Drumond, A., Durán-Quesada, A. M. and Nieto, R.: Oceanic and terrestrial sources of continental precipitation, *Rev. Geophys.*, 50(4), RG4003, doi:10.1029/2012RG000389, 2012.
- 30

- Harding, K. J. and Snyder, P. K.: Examining future changes in the character of central u.S. warm-season precipitation using dynamical downscaling, *J. Geophys. Res.*, 119(23), 13,113–116,136, doi:10.1002/2014JD022575, 2014.
- Helfand, H. M. and Schubert, S. D.: Climatology of the simulated Great Plains low-level jet and its contribution to the continental moisture budget of the United States, *J. Clim.*, 8(4), 784–806, doi:10.1175/1520-0442(1995)008<0784:COTSGP>2.0.CO;2, 1995.
- Higgins, R. W., Mo, K. C. and Schubert, S. D.: The moisture budget of the central United States in spring as evaluated in the NCEP/NCAR and the NASA/DAO reanalyses, *Mon. Weather Rev.*, 124(5), 939–963, doi:10.1175/1520-0493(1996)124<0939:TMBOTC>2.0.CO;2, 1996.
- Higgins, R. W., Yao, Y., Yarosh, E. S., Janowiak, J. E. and Mo, K. C.: Influence of the great plains low-level jet on summertime precipitation and moisture transport over the central United States, *J. Clim.*, 10(3), 481–507, doi:10.1175/1520-0442(1997)010<0481:IOTGPL>2.0.CO;2, 1997.
- Hong, S. and Lim, J.: The WRF single-moment 6-class microphysics scheme (WSM6), *J. Korean Meteorol. Soc.*, 42(2), 129–151, 2006.
- Hu, Q., Jiang, D. and Lang, X.: Sources of moisture for different intensities of summer rainfall over the Chinese Loess Plateau during 1979–2009, *Int. J. Climatol.*, 38, e1280–e1287, doi:10.1002/joc.5416, 2018.
- Hu, X. M., Klein, P. M. and Xue, M.: Evaluation of the updated YSU planetary boundary layer scheme within WRF for wind resource and air quality assessments, *J. Geophys. Res. Atmos.*, 118(18), 10490–10505, doi:10.1002/jgrd.50823, 2013.
- Insua-Costa, D. and Miguez-Macho, G.: A new moisture tagging capability in the Weather Research and Forecasting model: Formulation, validation and application to the 2014 Great Lake-effect snowstorm, *Earth Syst. Dyn.*, 9(1), 167–185, doi:10.5194/esd-9-167-2018, 2018.
- Kain, J. S.: The Kain–Fritsch Convective Parameterization: An Update, *J. Appl. Meteorol.*, 43(1), 170–181, doi:10.1175/1520-0450(2004)043<0170:TKCPAU>2.0.CO;2, 2004.
- Mlawer, E. J., Taubman, S. J., Brown, P. D., Iacono, M. J. and Clough, S. A.: Radiative transfer for inhomogeneous atmospheres: RRTM, a validated correlated-k model for the longwave, *J. Geophys. Res. Atmos.*, 102(D14), 16663–16682, doi:10.1029/97JD00237, 1997.
- Mo, K. C. and Juang, H.-M. H.: Relationships between soil moisture and summer precipitation over the Great Plains and the Southwest, *J. Geophys. Res. D Atmos.*, 108(16), GCP 5-1–GCP 5-17, doi:10.1175/1520-0469(1995)052<0879:PMOTSF>2.0.CO;2, 2003.
- Mo, K. C., Nogues-Paegle, J. and Paegle, J.: Physical mechanisms of the 1993 summer floods, *J. Atmos. Sci.*, 52(7), 879–895,

doi:10.1175/1520-0469(1995)052<0879:PMOTSF>2.0.CO;2, 1995.

Mo, K. C., Paegle, J. N. and Higgins, R. W.: Atmospheric processes associated with summer floods and droughts in the central United States, *J. Clim.*, 10(12), 3028–3046, doi:10.1175/1520-0442(1997)010<3028:APAWSF>2.0.CO;2, 1997.

5 [Moore, B. J., Neiman, P. J., Ralph, F. M. and Barthold, F. E.: Physical Processes Associated with Heavy Flooding Rainfall in Nashville, Tennessee, and Vicinity during 1–2 May 2010: The Role of an Atmospheric River and Mesoscale Convective Systems*, *Mon. Weather Rev.*, 140\(2\), 358–378, doi:10.1175/MWR-D-11-00126.1, 2012.](#)

[Nakamura, J., Lall, U., Kushnir, Y., Robertson, A. W. and Seager, R.: Dynamical Structure of Extreme Floods in the U.S. Midwest and the United Kingdom, *J. Hydrometeorol.*, 14\(2\), 485–504, doi:10.1175/JHM-D-12-059.1, 2013.](#)

10 [Nayak, M. A., Villarini, G., Bradley, A. A., Nayak, M. A., Villarini, G. and Bradley, A. A.: Atmospheric Rivers and Rainfall during NASA’s Iowa Flood Studies \(IFloodS\) Campaign, *J. Hydrometeorol.*, 17\(1\), 257–271, doi:10.1175/JHM-D-14-0185.1, 2016.](#)

Numaguti, A.: Origin and recycling processes of precipitating water over the Eurasian continent: Experiments using an atmospheric general circulation model, *J. Geophys. Res. Atmos.*, 104(D2), 1957–1972, doi:10.1029/1998JD200026, 1999.

15 Pan, Z., Segal, M. and Arritt, R. W.: Role of Topography in Forcing Low-Level Jets in the Central United States during the 1993 Flood-Altered Terrain Simulations, *Mon. Weather Rev.*, 132(1), 396–403, doi:10.1175/1520-0493(2004)132<0396:ROTIFL>2.0.CO;2, 2004.

Pu, B., Dickinson, R. E. and Fu, R.: Dynamical connection between Great Plains low-level winds and variability of central Gulf States precipitation, *J. Geophys. Res. Atmos.*, 121(7), 3421–3434, doi:10.1002/2015JD024045, 2016.

20 Ramos, A. M., Nieto, R., Tomé, R., Gimeno, L., Trigo, R. M., Liberato, M. L. R. and Lavers, D. A.: Atmospheric rivers moisture sources from a Lagrangian perspective, *Earth Syst. Dyn.*, 7(2), 371–384, doi:10.5194/esd-7-371-2016, 2016.

Rife, D. L., Pinto, J. O., Monaghan, A. J., Davis, C. A. and Hannan, J. R.: Global distribution and characteristics of diurnally varying low-level jets, *J. Clim.*, 23(19), 5041–5064, doi:10.1175/2010JCLI3514.1, 2010.

25 Schubert, S. D., Helfand, H. M., Wu, C. Y. and Min, W.: Subseasonal variations in warm-season moisture transport and precipitation over the central and eastern United States, *J. Clim.*, 11(10), 2530–2555, doi:10.1175/1520-0442(1998)011<2530:SIVIWSM>2.0.CO;2, 1998.

Shin, H. H. and Hong, S. Y.: Intercomparison of Planetary Boundary-Layer Parametrizations in the WRF Model for a Single Day from CASES-99, *Boundary-Layer Meteorol.*, 139(2), 261–281, doi:10.1007/s10546-010-9583-z, 2011.

Sorí, R., Marengo, J., Nieto, R., Drumond, A. and Gimeno, L.: The Atmospheric Branch of the Hydrological Cycle over the

- Negro and Madeira River Basins in the Amazon Region, *Water*, 10(6), 738, doi:10.3390/w10060738, 2018.
- Stensrud, D. J.: Importance of low-level jets to climate: A review, *J. Clim.*, 9(8), 1698–1711, doi:10.1175/1520-0442(1996)009<1698:IOLLJT>2.0.CO;2, 1996.
- Stohl, A., James, P., Stohl, A. and James, P.: A Lagrangian Analysis of the Atmospheric Branch of the Global Water Cycle. Part I: Method Description, Validation, and Demonstration for the August 2002 Flooding in Central Europe, *J. Hydrometeorol.*, 5(4), 656–678, doi:10.1175/1525-7541(2004)005<0656:ALAOTA>2.0.CO;2, 2004.
- Stohl, A. and James, P.: A Lagrangian Analysis of the Atmospheric Branch of the Global Water Cycle. Part II: Moisture Transports between Earth's Ocean Basins and River Catchments. *J. Hydrometeorol.*, 6, 961-984, doi:10.1175/JHM470.1, 2005a.
- 10 [Stohl, A.; Forster, C.; Frank, A.; Seibert, P.; Wotawa, G. Technical note: The Lagrangian particle dispersion model FLEXPART version 6.2. *Atmos. Chem. Phys.*, 5, 2461–2474, 2005b.](#)
- Stohl, A., Forster, C. and Sodemann, H.: Remote sources of water vapor forming precipitation on the Norwegian west coast at 60°N - A tale of hurricanes and an atmospheric river, *J. Geophys. Res. Atmos.*, 113(5), 1–13, doi:10.1029/2007JD009006, 2008.
- 15 [Tang, Y., Winkler, J., Zhong, S., Bian, X., Doubler, D., Yu, L. and Walters, C.: Future changes in the climatology of the Great Plains low-level jet derived from fine resolution multi-model simulations, *Sci. Rep.*, 7\(1\), doi:10.1038/s41598-017-05135-0, 2017.](#)
- Ting, M. and Wang, H.: The Role of the North American Topography on the Maintenance of the Great Plains Summer Low-Level Jet*, *J. Atmos. Sci.*, 63(3), 1056–1068, doi:10.1175/JAS3664.1, 2006.
- 20 Trenberth, K. E. and Guillemot, C. J.: Physical processes involved in the 1988 drought and 1993 floods in north America, *J. Clim.*, 9(6), 1288–1298, doi:10.1175/1520-0442(1996)009<1288:PPIITD>2.0.CO;2, 1996.
- Vázquez, M., Nieto, R., Drumond, A. and Gimeno, L.: Moisture transport into the Arctic: Source-receptor relationships and the roles of atmospheric circulation and evaporation, *J. Geophys. Res. Atmos.*, 121(22), 13,493-13,509, doi:10.1002/2016JD025400, 2016.
- 25 [Walters, C. K., Winkler, J. A., Shadbolt, R. P., van Ravensway, J. and Bierly, G. D.: A long-term climatology of southerly and northerly low-level jets for the central United States, *Ann. Assoc. Am. Geogr.*, 98\(3\), 521–552, doi:10.1080/00045600802046387, 2008.](#)
- Wang, S. Y. and Chen, T. C.: The late-spring maximum of rainfall over the U.S. central plains and the role of the low-level jet, *J. Clim.*, 22(17), 4696–4709, doi:10.1175/2009JCLI2719.1, 2009.

Wegmann, M., Orsolini, Y., Vázquez, M., Gimeno, L., Nieto, R., Bulygina, O., Jaiser, R., Handorf, D., Rinke, A., Dethloff, K., Sterin, A. and Brönnimann, S.: Arctic moisture source for Eurasian snow cover variations in autumn, *Environ. Res. Lett.*, 10(5), doi:10.1088/1748-9326/10/5/054015, 2015.

Wexler, H.: A Boundary Layer Interpretation of the Low-level Jet, *Tellus*, 13(3), 368–378, doi:10.1111/j.2153-
5 3490.1961.tb00098.x, 1961.

Wu, Y. and Raman, S.: The summertime Great Plains low level jet and the effect of its origin on moisture transport, *Boundary-Layer Meteorol.*, 88(3), 445–466, doi:10.1023/A:1001518302649, 1998.

10

15

20

25

30

Captions

Figure 1: (a) Mean NLLJ index (shaded) and 500 m winds (arrows, in m s^{-1}) at local midnight in July (boreal summer) for 1980-2016, calculated from ERA-Interim reanalysis. The black cross at 32.75°N , 99°W shows the point of maximum NLLJ in the climatology. The cyan contour line surrounds the region containing points above the 75th percentile. (b) Frequency distributions of the GPLLJ for the months of July from 1980 to 2016 (blue bars). The red curve corresponds to the Gaussian fit: $y(x) = y_0 + A \exp(-\frac{(x-y)^2}{2\sigma^2})$. (see table A2). Noted: The frequency distributions are calculated at the point of maximum intensity of NLLJ (at 32.75°N , 99°W , black cross in fig. 1a).

Figure 2: Highlighted in red are moisture sources obtained with FLEXPART from backward trajectories originating in the region outlined in magenta in Fig 1.a. The orange line over the continent marks the position from where precipitable water is tagged in WRF-TFWVT, corresponding to the northern edge of the FLEXPART source region. All water vapour and condensate crossing through that line is considered as moisture advected by the GPLLJ for further analysis.

Figure 3: Ratio of tagged precipitable water transported by the GPLLJ to total for the six case-studies analysed.

Figure 4: Statistically weighted ratio of precipitable water transported by the GPLLJ for the five case studies with $\text{NLLJ} > 0$ considered in the analysis in Figure 3. Weights applied are stated in Table 1.

Figure 5: Tracer total precipitable water (TPW, g kg^{-1}) and positions of the cross sections along the central axis of the GPLLJ shown in Fig. 5, at latitudes 32°N (1), 35°N (2) and 38°N (3) for the main jet event of July 11, 2002-1992.

Figure 6: (a-c) $q\text{TR}$ in g kg^{-1} for the three vertical cross sections at the locations depicted with white lines in Fig. 5. (d-f) same as (a-c) but for ϕTR in g m (kg s)^{-1} .

Figure 7: (a-c) Ratio of $q\text{TR}:q$ for the three vertical cross sections at the locations depicted with white lines in Fig. 5. (d-f) same as (a-c) but for ratio of $\phi\text{TR}:\phi$.

Table 1: Case-studies objectively selected based in the frequency distribution of the LLJ index to carry out WRF-TFWVT simulations. μ is the mean of the distribution and σ its standard deviation. Noted: The frequency distribution is calculated at the point of maximum intensity of NLLJ at 32.75°N , 99°W (black cross in fig. 1a) using the ERA-Interim reanalysis dataset.




Figure 1

5

10

15

20

25




Figure 2

30

35

5

10

15

20

25




Figure 3

30

35

5

10

15

20

25




Figure 4

30

35


5

10

15

20

25



30 **Figure 5**

35




Figure 6




Figure 7

5

10

Table 1

15

Simulation	Gaussian	NLLJ value	Date	Stat. weight
0		0.00	2012-07-12	0
1	$\mu - 2\sigma$	1.49	1999-07-19	0.0623
2	$\mu - \sigma$	5.54	1983-07-23	0.2445
3	μ	10.19	1992-07-11	0.3864
4	$\mu + \sigma$	14.54	2002-07-28	0.2445
5	$\mu + 2\sigma$	18.89	2016-07-14	0.0623

20

25

Appendix A: Statistical weights in the analysis

Table A1. Events

Simulation	Gaussian Point
1	$\mu - 2\sigma$
2	$\mu - \sigma$
3	μ
4	$\mu + \sigma$
5	$\mu + 2\sigma$

Table A2. Gaussian fit $y(x) = y_0 + A \cdot \exp(-\frac{(x-y)^2}{2\sigma^2})$.

Simulation	Gaussian Point
y_0	0.03 ± 3.66
A	8.63 ± 4.39
μ	10.19 ± 0.19
σ	4.35 ± 0.30
R^2	0.95

5

$$StatWeight_1 = StatWeight_5 = \frac{\int_{\mu-2.5\sigma}^{\mu-1.5\sigma} y(x)dx}{\int_{\mu-2.5\sigma}^{\mu+2.5\sigma} y(x)dx} = 0.0623$$

$$StatWeight_2 = StatWeight_4 = \frac{\int_{\mu-1.5\sigma}^{\mu-0.5\sigma} y(x)dx}{\int_{\mu-2.5\sigma}^{\mu+2.5\sigma} y(x)dx} = 0.2446$$

10

$$StatWeight_3 = \frac{\int_{\mu-1.5\sigma}^{\mu+0.5\sigma} y(x)dx}{\int_{\mu-2.5\sigma}^{\mu+2.5\sigma} y(x)dx} = 0.38643$$