



Changes in statistical distributions of sub-daily surface temperatures and wind speed

Robert J. H. Dunn¹, Kate M. Willett¹, and David E. Parker¹

¹Met Office Hadley Centre, FitzRoy Road, Exeter, EX1 3PB, UK

Correspondence: robert.dunn@metoffice.gov.uk

Abstract. With the ongoing warming of the globe, it is important to quantify changes in the recent behaviour of extreme events given their impacts on human health, infrastructure and the natural environment. We use the sub-daily, multi-variate, station-based HadISD dataset to study the changes in the statistical distributions of temperature, dewpoint temperature and wind speeds. Firstly, we use zonally averaged quantities to show that the lowest temperatures during both day and night are changing more rapidly than the highest, with the effect more pronounced in the northern high latitudes. Along with an increase in the zonally-averaged mean temperature, the standard deviation has decreased and the skew increased (increasing positive tail/decreasing negative tail) over the last 45 years, again with a stronger, more robust signal at higher latitudes. Changes in the distribution of dewpoint temperature are similar to those of temperature. However, changes in the distribution of wind speeds indicate a more rapid change at higher speeds than at lower.

Secondly, to assess in more detail the spatial distribution of changes as well as across seasons and hours of the day we study each station individually. For stations which show clear indications of change in the statistical moments, the higher the statistical moment, generally the more spatially heterogeneous the patterns of change. The standard deviation of temperatures are increasing in a band stretching from Europe through to China, but are decreasing across North America and in the high northern latitudes, indicating broadening and narrowing of the distributions respectively. Large seasonal differences are found in the change of standard deviations of temperatures over North America and eastern China. Temperatures in Eastern Asia also have increasing skew in the winter in contrast to the remainder of the year. The dewpoint temperatures show smaller variation in all of the moments, but similar patterns to the temperatures. For wind speeds, apart from the USA, standard deviations are decreasing across the world, indicating a decrease in variability.

Finally, we use quantile regression to show changes in the percentiles of distributions over time. These show an increase of high quantiles of temperature in eastern Europe during the summer, and also in northern Europe for low quantiles in the winter, also indicating broadening and narrowing of the distributions respectively. In North America, the largest changes are at the lower quantiles in northern latitudes for autumn and winter. Quantiles of dewpoint temperature are changing most in the autumn and winter, especially in the northern parts of Europe.



1 Introduction

25 The study of changes in the extremes of essential climate variables is vital given their impacts on human health, infrastructure, agriculture and the natural environment. The Intergovernmental Panel on Climate Change (IPCC) released a Special Report on Extremes (SREX, Field et al., 2012), and therein outlined three simple classes of the way changes in extremes occur; shifting the mean, increasing the variability, and changing the symmetry. Past studies have indicated that there is uncertainty around how changes in the occurrence or intensity of extreme values arise. Are extremes changing due to changes in the location of
30 the distribution mean with no change in the distribution shape (Griffiths et al., 2005; Simolo et al., 2011), or are changes in the shape of a distribution the primary driver of changes in extremes? If so, is the change in shape only the consequence of a change in the variance or does it also arise from changes in higher order moments (Della-Marta et al., 2007; Ballester et al., 2010)?

There are a whole host of ways to study climate extremes and determine how these have changed over the recent past.
35 One common approach is to use climate extremes indices to characterise moderate extremes in timeseries of station data, for example those developed by the World Meteorological Organisation (WMO) Commission for Climatology (CCI), World Climate Research Programme (WCRP), and Joint Technical Commission for Oceanography and Marine Meteorology (JCOMM) Expert Team in Climate Change Detection and Indices (ETCCDI, Alexander et al., 2006). A number of datasets comprising these indices now exist, allowing for detailed investigations of past changes and comparisons to model and reanalysis fields
40 (e.g. Caesar et al., 2006; Donat et al., 2013a, b). Another route uses Generalised Extreme Value theory (GEV) to model the tails of the distributions, and from these few points characterise the occurrence and intensity of extreme events, including those not yet observed in the modern data record (e.g. Brown et al., 2008; Christidis et al., 2011).

A further method characterises the complete distribution using all available data to establish what are the causes of changes in climate extremes. The advantage of this approach is that it uses all of the available information, and can easily be extended to
45 more or less extremal parts of the distribution. A number of the ETCCDI temperature indices use the 10th and 90th percentile values, which only probe moderate extremes. A recalculation of these indices at 1st and 99th percentile values is possible, but given the diverse sources of data in the HadEX datasets (Donat et al., 2013b), this would be a large undertaking by the international community. Furthermore, as the climate continues to warm, the relative intensity or duration of extreme events against warmer average conditions may be of interest; for example, how warm is the relatively warmest 10 per cent of days
50 now compared to the same fraction selected between 1961-90?

Hence, a number of recent studies have investigated past changes in the distributions of observed land-surface temperatures. Most have used the Global Historical Climate Network Daily (GHCND, Menne et al., 2012) or its gridded version (HadGHCND, Caesar et al., 2006, temperature only) as these provide daily maximum and minimum temperature values. Donat and Alexander (2012) compared two periods to show that the means of both temperature measures have shifted to warmer
55 values. Along with larger changes in the minimum than the maximum temperatures, this has resulted in increases in the skew of the distributions. However changes in standard deviation were less significant and more heterogeneous. Over North America, Shen et al. (2011) show that the standard deviation and skewness were decreasing, but that the kurtoses of the maximum



temperatures were increasing, in contrast to the decrease in the kurtoses of minimum temperatures. A subsequent study investigated seasonal changes in the distribution moments over 1950-2010 (Cavanaugh and Shen, 2014). McKinnon et al. (2016)
60 showed that for most stations between 1980 and 2015, over just the summer months, trends can be explained by a shift in the distribution with no change in shape. Reanalyses (e.g. Huntingford et al., 2013; Gross et al., 2018), or coupled models (e.g. Lewis and King, 2017) have also been used in the study of changes in the shapes of current and future distributions of essential climate variables (ECVs).

In this study, we use the sub-daily, station data from a single dataset, as opposed to the maximum and minimum daily
65 temperatures (either station based or gridded). We note that by using a single dataset, we have not investigated the effect of dataset choice on our findings, something which can have a large effect (Gross et al., 2018). In order to compare with a number of previous studies we perform three different assessments to see how the distributions of temperature, dewpoint temperature, and wind speed have changed in recent years. We outline the dataset and station selection criteria in Section 2. The three assessments are outlined in Sections 3 to 5, with a summary in Section 6.

70 2 Input Data

To study the changes in distributions related to climate extremes, we use the sub-daily HadISD dataset (Dunn et al., 2012, 2014, 2016; Dunn, 2019). This is updated annually, and now covers a period 1931-2018 inclusive. Improved station selection and merging processes as well as minor changes in the quality control tests are features of version 3.0.0.2018f over version 1.0.x. This version contains 8139 unique station locations, with some of these being composited from individual records in
75 the ISD (Integrated Surface Dataset, Smith et al., 2011). As a result of the long standing issues surrounding the free sharing of observational climate data (Thorne et al., 2017), the bulk of these stations are in the Northern Hemisphere, concentrated in North America and Europe. In contrast to some of the previous studies outlined in the Introduction (e.g. Donat and Alexander, 2012), we do not create a gridded form of the HadISD for this assessment, but retain the individual stations as input to the analyses, similar to e.g. Cavanaugh and Shen (2014); McKinnon et al. (2016).

80 HadISD undergoes a homogeneity assessment using the Pairwise Homogeneity Algorithm (PHA, Menne and Williams Jr, 2009) to identify the number and size of inhomogeneities in its four main ECVs (Dunn et al., 2014). Although no adjustments are made, the outputs of the PHA allow stations that have few or small inhomogeneities to be selected, i.e. those with the most homogeneous records (see Dunn et al., 2014). Therefore in all of the investigations below, we select those stations that have fewer than 10 inhomogeneities, allowing jumps of up to 1° C. This balances retaining sufficient stations to study the behaviour
85 across the globe with removing those with very large inhomogeneities.

Although the HadISD v3.0.x contains data from 1931 onwards, there is a large increase in the number of stations from 1973 onwards (the reason why HadISD v1.0.x started then, see Fig. 2 in Dunn et al., 2016). Therefore we restrict our analysis to the time period from 1st January 1973 unless otherwise stated. We also remove data from February 29th for ease of analysis.



3 Zonal Distribution Changes

90 We firstly look at the changes in distributions for all stations in a specified latitude band. By combining stations together, the quantity of observations available to define the distribution is increased. Donat and Alexander (2012) used the HadGHCND gridded daily temperature dataset (Caesar et al., 2006) to compare two 30-year periods of daily maximum and minimum temperatures (1951-80 and 1981-2010) globally, and for three smaller zonal regions (Northern Hemisphere, Tropics and Southern Hemisphere). Here we develop this method further using the sub-daily data available in the HadISD as well as narrower
95 10-degree zonal bands.

We select those stations that fall within each 10 degree band. An approximation for the whole-hour timezone of the station is calculated from the station longitude, enabling the Universal Time (UTC) observation time to be converted to local time. We then extract those portions of data that correspond to daytime and night-time, using the time periods 09-20 and 21-08 respectively, to ensure that the minimum temperatures are on the whole captured during the night-time portion, and the maximum
100 temperatures in the daytime. At high latitudes these definitions of day- and night-time will be vary in accuracy throughout the year, but should split the 24 hour day into the hours with most and least insolation. There are few stations at the highest latitudes where 24-hours of day or night occur.

Climatologies for stations for each day of the year for the day- and night-time data are calculated separately, requiring the equivalent of 15 years of 3-hourly observations over the 1981-2010 period. These climatologies are used to create anomalies
105 from the sub-daily data. The creation of climate anomalies also restricts this analysis to stations that have consistent records within this recent climatology period.

To investigate the changes in the distributions over time, we split the data up into nine intervals each of five years in length (1974-78, 1979-83, .., 2014-2018). By not including 1973, we can assess changes over a 45 year period. Furthermore, we only take those stations that have sufficient observations spread over the entire analysis period. In each interval, we require there to
110 be more observations than the number equivalent to hourly observations for 1/4 of the 5-year interval. Also, at least 3/4 of the intervals require sufficient data for the station to be incorporated into the final distribution. This is to ensure that non-uniform distributions of observations within a station record (both within a five-year interval and across the entire record) do not overly influence the final distributions. The stations which pass these selection criteria are shown in Fig 1.

We determine the values for the first four statistical moments (mean, standard deviation, skew and [excess] kurtosis) of the
115 resulting distributions. These in turn represent the central tendency, the spread, the asymmetry and the “peaked-ness” of the distribution. Distributions with non-zero skew or kurtosis indicate departures from a pure Gaussian shape. We note that the skew and kurtosis measures are not fully orthogonal (McKinnon et al., 2016), but will use these measures in this analysis because of their well-understood nature. The linear change in these values, along with 1σ range have been derived from the median of pairwise slopes method (e.g. Sen, 1968; Lanzante, 1996), and are shown for each latitude band in Table 1 and
120 Supplementary Information Table 1 & 2 for temperature, dewpoint temperature and wind speed respectively. We do not expect any changes in the parameters of the distributions to be linear, but it is a useful way to summarise their gross changes over time




Figure 1. Maps showing the locations of HadISD stations contributing to the zonal analysis along with the 10 degree latitude bands for TOP temperature, MIDDLE dewpoints, and BOTTOM windspeeds.




Figure 2. Dry bulb temperature distributions in latitudinal bands for LEFT day-, and RIGHT night-time observations for the Northern Hemisphere stations. The *y*-axis has been normalised for ease of comparison. The number of stations contributing to each set of distributions is shown in the top left of each sub-panel, with the latitude range in the top right. The short vertical bars indicate the location of the 5th and 95th percentile values for each distribution.

together with the relative certainty of these changes. Latitude bands where the range in trend values does not encompass zero are highlighted in bold, as for these we can be more confident that there is a true change in the value over time.

3.1 Temperatures

125 The results from day- and night-time temperature observations for the Northern Hemisphere are shown in Fig. 2 and Table 1. We can be more confident of changes that are seen in latitude bands containing a large number of stations than in those which



only contain a handful. The Southern Hemisphere is shown in Supplementary Information Fig. 1 and Table 1, as it includes considerably fewer stations. It is clearly evident from these figures and tables that there has been a shift in the distribution from the early period to the later period, with the most recent period being on average warmer than the earliest. This behaviour
130 agrees with the results presented in Donat and Alexander (2012) who shows increases in the mean of around 1°C between the early (1951-1980) and late (1981-2010) periods in their study (for all their regions and for both maximum and minimum temperatures). We note that their period of study is longer (60 years) compared to the period presented in this work (45 years).

In both day- and night-time observations, positive increases in the mean that are clearly different from zero are seen in all latitude bands (Table 1). The strongest increases are seen in the highest northern latitudes, with increases on the whole
135 becoming smaller when heading towards the equator for both halves of the day. Changes in the southern hemisphere show less variation with latitude, and are on the whole smaller than in the north, though still showing overall increases. There are few differences in the day- and night-time increases for each latitude band, in most cases the ranges overlap.

The northern mid-to-high latitudes also see a consistent decrease in the standard deviations in both day and night time. However, south of 20°N, the signal is less consistent. In fact, during day- and night-time most tropical and southern-latitude
140 bands have ranges that encompass zero, though the most bands suggest decreases. Although Donat and Alexander (2012) also found decreases in standard deviation in the extra-tropics, between 30°S-30°N they found an increase when using zonal averages.

Consistent changes in the skew and kurtosis for both day- and night-time or across latitude bands are not as apparent in Table 1. Slight increases in skew are seen for the daytime high latitudes (north and south). In contrast, during night-time, tropical
145 regions show decreases in skew. The kurtosis decreases in the tropics at both times of day, with a more mixed signal at the high latitudes. Donat and Alexander (2012) also found an increase in the skew in all regions except for T_{\max} in the tropics, which is broadly consistent with the changes shown in Table 1.

3.2 Dewpoint Temperatures

The changes in the dewpoint temperatures (Supplementary Information Table 1 and Figs. 2-3) show a slightly more complex
150 picture than for the dry-bulb temperatures. In general, the distribution mean again shows an increase in both day- and night-time values, with larger increases over time in the northern hemisphere high-latitudes, similar to the dry-bulb temperatures. In the southern hemisphere however, and especially at high latitudes, this increasing signal is less strong with few cases where the confidence intervals do not span zero. One southern latitude band even shows a clear decrease in mean dewpoint temperatures over the last 45 years. Also, trends close to the equator (10°S to 20°N) are greater than at latitudes immediately to the north
155 and south of this equatorial belt.

The tendency has been for standard deviations in many northern latitudes to decrease, but there are also a number of increases, especially in the southern hemisphere, though determined from few stations. For the skew, in the daytime, these increase in the north and decrease in the south, with a similar, but not as consistent pattern in the night-time. The changes in kurtosis do not show any clear pattern in comparison to the other moments.



Band	N-stations	Mean ($^{\circ}\text{C decade}^{-1}$)	St. Dev. ($^{\circ}\text{C decade}^{-1}$)	Skew (decade^{-1})	Kurtosis (decade^{-1})
Day					
70 → 60	173	0.40 (0.38 → 0.50)	−0.19 (−0.23 → −0.14)	0.04 (0.03 → 0.06)	0.06 (−0.02 → 0.14)
60 → 50	516	0.33 (0.28 → 0.41)	−0.09 (−0.14 → −0.05)	0.03 (0.02 → 0.05)	−0.03 (−0.13 → 0.03)
50 → 40	572	0.34 (0.33 → 0.40)	−0.06 (−0.09 → −0.03)	−0.00 (−0.02 → 0.02)	−0.02 (−0.07 → 0.00)
40 → 30	317	0.29 (0.27 → 0.31)	−0.11 (−0.14 → −0.05)	0.02 (0.01 → 0.05)	0.07 (0.02 → 0.11)
30 → 20	140	0.25 (0.23 → 0.29)	−0.05 (−0.07 → −0.00)	0.03 (0.01 → 0.06)	0.05 (0.03 → 0.07)
20 → 10	79	0.19 (0.17 → 0.23)	−0.01 (−0.03 → 0.02)	0.03 (0.02 → 0.05)	0.01 (−0.06 → 0.05)
10 → 0	25	0.15 (0.12 → 0.18)	0.01 (−0.00 → 0.02)	−0.00 (−0.02 → 0.01)	−0.10 (−0.11 → −0.07)
0 → −10	9	0.14 (0.14 → 0.17)	−0.01 (−0.01 → 0.01)	0.04 (0.04 → 0.06)	−0.15 (−0.17 → −0.13)
−10 → −20	30	0.15 (0.13 → 0.21)	0.07 (0.03 → 0.09)	0.02 (0.01 → 0.05)	−0.27 (−0.36 → −0.12)
−20 → −30	34	0.12 (0.11 → 0.14)	−0.08 (−0.09 → −0.07)	0.03 (0.01 → 0.06)	0.22 (0.20 → 0.25)
−30 → −40	29	0.21 (0.19 → 0.23)	−0.00 (−0.04 → 0.03)	0.07 (0.07 → 0.09)	0.13 (0.09 → 0.20)
−40 → −50	10	0.10 (0.05 → 0.15)	0.01 (−0.02 → 0.06)	0.06 (0.05 → 0.07)	0.01 (−0.03 → 0.03)
Night					
70 → 60	139	0.38 (0.33 → 0.46)	−0.17 (−0.20 → −0.12)	0.00 (−0.00 → 0.02)	0.02 (−0.03 → 0.11)
60 → 50	467	0.30 (0.25 → 0.40)	−0.08 (−0.11 → −0.04)	0.03 (0.01 → 0.06)	−0.10 (−0.18 → 0.01)
50 → 40	507	0.32 (0.31 → 0.35)	−0.10 (−0.11 → −0.06)	0.01 (−0.01 → 0.03)	−0.01 (−0.09 → 0.03)
40 → 30	256	0.31 (0.30 → 0.31)	−0.13 (−0.17 → −0.05)	0.05 (0.03 → 0.07)	0.03 (−0.01 → 0.12)
30 → 20	79	0.28 (0.27 → 0.30)	−0.06 (−0.08 → 0.00)	−0.01 (−0.02 → 0.01)	0.09 (0.07 → 0.13)
20 → 10	52	0.18 (0.17 → 0.24)	−0.03 (−0.05 → −0.02)	0.06 (0.03 → 0.08)	−0.30 (−0.33 → −0.23)
10 → 0	13	0.20 (0.17 → 0.23)	−0.00 (−0.00 → 0.03)	−0.07 (−0.08 → −0.04)	−0.41 (−0.48 → −0.38)
0 → −10	4	0.18 (0.14 → 0.22)	−0.03 (−0.08 → −0.01)	0.00 (−0.04 → 0.04)	−0.39 (−0.55 → −0.12)
−10 → −20	7	0.26 (0.23 → 0.32)	−0.02 (−0.05 → 0.02)	−0.06 (−0.09 → 0.02)	0.23 (0.17 → 0.40)
−20 → −30	18	0.16 (0.10 → 0.19)	−0.05 (−0.06 → −0.02)	0.02 (0.01 → 0.06)	0.08 (0.01 → 0.14)
−30 → −40	21	0.23 (0.22 → 0.25)	−0.07 (−0.08 → −0.01)	0.05 (0.03 → 0.06)	0.15 (0.12 → 0.17)
−40 → −50	9	0.10 (0.07 → 0.13)	−0.01 (−0.03 → 0.04)	0.01 (0.00 → 0.02)	−0.05 (−0.10 → −0.03)

Table 1. Linear change per decade in fits to parameters in zonal analysis for TOP day-, and BOTTOM night-time temperature observations over 1973–2017. Values in bold show parameters and bands where the 1σ range of the fitted trend does not include zero.



160 **3.3 Wind speeds**

We also assess changes in the wind speeds, but there are fewer stations where this variable has been sufficiently observed (see Supplementary Information Table 2 and Fig. 4). In fact, all latitude bands in the Southern Hemisphere have fewer than ten stations, and so we do not study this hemisphere. We also note that this variable is not normally distributed, unlike the temperatures investigated in previous sections. Despite this we use the same method, but note that the anomalies will be
165 truncated at the negative end by the zero-bounded nature of this variable.

For the northern hemisphere, most latitude bands show reductions in the mean and also the standard deviation over time for both day- and night-time observations, with confidence bounds distinct from zero. In the lower mid-latitudes, there are indications for a small increase in standard deviations, but at a lower magnitude than other changes. Tropical regions show decreases in the skewness over day- and night-time, but high latitudes an increase during the day.

170 These decreases in the mean wind speed over land have been noted in past studies (Roderick et al., 2007; McVicar et al., 2012), and monitoring studies have also shown decreases over time for the high and low wind speed values (e.g. Azorin-Molina et al., 2018). However, these recent regular global assessments also use the HadISD and so differences are not expected.

175 **3.4 Summary**

As can be seen in both Fig. 2 and Table 1 (as well as Fig. 1 in the Supplementary Information) there is a clear change in the
180 mean temperatures with time across all latitude bands and times of day. However, this is also combined with a decrease in the standard deviation over the northern hemisphere. The width of the distribution of temperatures is therefore narrowing. Also, the skew of the distribution is becoming more positive, which indicates that the positive tail is growing and the negative tail is shrinking. Together, these three changes indicate that there has been a greater change in the low tail than in the high tail of the temperature distribution. The changes in the kurtosis are mainly increases, indicating a change to more peaked distributions, and a reduction in both tails. Although Donat and Alexander (2012) found similar increases in the mean and skew of daily maxima and minima, they showed increases in the standard deviation for tropical regions not found here.

185 Markers indicating the location of the 5th and 95th percentile values in Fig. 2 track the changing tails of the distributions. Qualitatively for the northern hemisphere, the low tails have changed more than the high tails, as the vertical lines cover a greater spread, and more so for the higher latitudes than the tropical regions. These highlight changes in the standard deviation and also the skew on top of changes in the mean. However, there is no striking difference between the day- and night-time plots for temperature. The number of stations in the southern hemisphere is much smaller, but there are no systematic differences between the behaviours of the low and high tails (see Supplementary Information). A similar pattern is seen for the dewpoint temperatures, with a greater shift for the lower tail, and no striking difference between night- and daytime values. The wind speeds show little change in the lower tails (Supplementary Information Fig. 4), but decrease in the value of the 95th percentile over time. Further investigations into the changes in the 5th and 95th percentile values are presented in Section 5.

190 The intention of investigating the day and night time values separately was to echo analyses conducted in other studies of daily maximum and minimum temperatures. Many of these show that the minimum temperatures are changing faster than the



maxima, and so we expected to find a difference between day and night time, either for the parameters of the distributions or in the tail markers. However, the clearest difference remains between the upper and lower tails. This analysis shows increases
195 in the mean and skew, and decreases in the northern hemisphere standard deviations. These changes in the statistical moments of the temperature distributions indicate that the lower tails are shifting more rapidly than the upper tails. Hence, as well as a change in the location of the distribution, the shape has also changed, resulting in changes to occurrence of extremes. Although fewer stations contribute to the wind speed observations, these also show changes in location and shape, with larger changes in the upper tails caused by decreases in the mean and standard deviations. However, this approach removes any small scale,
200 regional changes in these and the other two characteristics and by focusing on the moments of the distributions makes it difficult to determine what happens to the tails and hence the climate extremes.

4 Station Distribution Changes

To investigate the geographical patterns of changes in these meteorological parameters in more detail than in Section 3, we study each station individually. This reduces the amount of data available to characterise the distribution at the gain of increased
205 spatial information. A similar approach was used by e.g. Cavanaugh and Shen (2014); McKinnon et al. (2016) in their investigations into the changing shape of temperature distributions on a daily basis from GHCND. However, the sub-daily nature of HadISD means we are able to investigate the distributions at different times throughout the day, rather than the changes of the maximum and minimum values independently. However, a large proportion of the stations in the HadISD only report every three hours, rather than hourly. Therefore we use the station longitude to adjust from UTC to local time, collating to the nearest
210 3-hourly time point (00:00, 03:00, ...).

For each station a climatology over 1981-2010 is calculated for each day of the year for each 3-hourly time point ($365 \times (24/3)$), requiring that at least 15 years of data are present. These 3-hourly climatologies are subtracted from the observations to create 3-hourly anomalies. The analysis is performed using observations over the entire year and also for standard three-month seasons.

215 The 45 years (1974-2018 inclusive) are split into five 9-year periods, and the anomalies within each period and at each three-hourly time point (0000, 0300, ... etc.), are combined, annually as well as seasonally. Using fewer periods of longer length than in Section 3 allows for better characterisation of the distributions at the individual station level, at the expense of the number of periods. This is in contrast to the previous analysis (Section 3) where combining stations together increases the number of observations contributing to a distribution. For the three-month seasons, using periods nine years in length means there are
220 maximally around 800 observations from which the distribution at a single time point can be characterised, if all observations are present (3 months $\times \sim 30$ days $\times 9$ years). The first four moments of the distributions are only calculated if there are more than 300 observations present within a 9-year period (roughly the equivalent of 1 months worth of observations at a given time point).

225 We again use the median of pairwise slopes method (Sen, 1968; Lanzante, 1996) to characterise the change in the moments over time, but only if there are sufficient data in at least four of the five periods. Again, we do not expect any changes to be




Figure 3. Trend over 1974-2018 in distribution parameters for temperature at 15:00 local time, showing the mean, standard-deviation, skew and kurtosis. Only the stations where the 1σ range of the fitted trend does not include zero are plotted (the number of these is shown in the top left hand corner of each plot). The total number of stations available is shown in parentheses.

linear, but this enables the convenient visual inspection of changes over time. Even if our data completeness criteria result in the inclusion of stations which have a non-uniform distribution of observations across the years and seasons, coherent, robust changes (if they are present) will still stand out across the stations.

4.1 Annual Temperature Changes

230 In Fig. 3 we show changes in the mean, standard deviation, skew and kurtosis for the temperature data for all seasons from
 HadISD at 15:00 and at 06:00 local time in Fig. 4, to correspond approximately with the times of maximum/daytime and
 minimum/nighttime temperature respectively. Other times are available in the Supplementary Material. Despite 8139 stations
 being available in the HadISD v3.0.0.2018f dataset, after selection for homogeneity and length of record, only around 3000
 235 have quantified changes in distributional parameters, the actual number of stations are shown in the top right-hand corner of
 each panel. Furthermore, only those stations where the 1σ range of the fitted trend does not include zero are plotted on each




Figure 4. Trend over 1974–2018 in distribution parameters for temperature at 06:00 local time, showing the mean ($^{\circ}\text{C}$), standard-deviation ($^{\circ}\text{C}$), skew and kurtosis. Only the stations where the 1σ range of the fitted trend does not include zero are plotted (the number of these is shown in the top left hand corner of each plot). The total number of stations available is shown in parentheses.

panel, as these were the ones where there is some confidence in there being a change over time. As is also clear from these figures, there are many fewer stations in the southern hemisphere than in the north. The highest concentration of stations is in Europe and eastern Asia, and this uneven distribution will have affected the zonal average calculations in Section 3.

As the highest station density is in the northern hemisphere, this is where any coherent changes are most clear. For the 240 distributions of daytime/maximum temperature (15:00, Fig. 3), it is immediately apparent that there are large and coherent changes in the mean and standard deviation over the period of record of these stations. Almost all changes in the mean are positive, with the strongest signal in Eurasia, and lower values in North America. Over 90 per cent (2806 plotted out of a total of 2956) of the stations have trends where the uncertainty range does not encompass zero. The majority of changes in standard deviation at high northern latitudes are negative, along with most for central North America. There are increases over Europe 245 and parts of China and southern Russia. Fewer stations (around 45 per cent, 1429/2956) have trends where the uncertainty range of the trends do not encompass zero.



Changes in skew and kurtosis have a more heterogenous signal, and again around half the stations exhibit changes where the uncertainty range does not include zero. However, the changes are not completely random, with consistent regional patterns in some areas, but these are smaller than for the standard deviation. The general pattern is for slight increases in the skew, 250 except in a belt around the tropics and in half of Europe. The kurtosis shows decreases in central Europe, and a band extending eastwards from the White Sea across to China and India, with a mixed signal elsewhere.

The night time observations (06:00) show very similar general patterns for changes in the mean and standard deviation, with quasi-global increases in the mean, and decreases in the standard deviation at high latitudes and increases for China, Europe and south-eastern Russia. However, the magnitude of the changes in the mean are lower than for the daytime (15:00) 255 observation, especially over Europe.

There is a stronger change between the skewness for the 06:00 observations and those from 15:00, with larger regions showing a coherent signal. Decreases are visible in more widely in Europe than at 15:00, and also in India and parts of south-east Asia; and increases in the high latitudes, as well as North America. Regions with a strong decrease in kurtosis are 260 mid-western North America, India through China and a band in north western Russia, but in contrast to 15:00, large parts of Europe show an increase. However, a smaller fraction of stations have trends where the uncertainty ranges do not encompass zero than for the higher moments.

The results from 15:00 and 06:00 can be compared to the maximum and minimum temperatures in the study by Donat and Alexander (2012) using HadGHCND between 1951 and 2010. Unsurprisingly, the increase in the mean temperatures is also 265 visible in the HadGHCND study, along with numerous other studies on recent climate change. However, HadGHCND shows stronger increases in the minimum temperatures (analogous to 06:00 here), whereas HadISD shows stronger increases in day-time (15:00) temperatures. The HadGHCND dataset uses inverse distance weighting to convert the daily station observations into a gridded dataset. One aspect of this process is that it smooths the data, and also can lead to very few stations contributing to large areas of interpolated values.

4.1.1 Temperature Changes Across the Day

270 In the Supplementary Information Fig. 5 we show the maps for changes in the mean for all eight of the 3-hourly time-stamps. All hours of the day show an increase in the mean, with the strongest signals during the daytime (09:00, 12:00, 15:00). It appears that Europe shows the largest trends, but this may be enhanced to the high station densities in this region compared to regions with a sparser distribution of stations.

The trends in standard deviation at 3-hourly intervals from HadISD show decreases at all hours in high northern latitudes, 275 North America, and southern Asia (Indian subcontinent and surroundings, Supplementary Information Fig. 6). Increases in the standard deviation are seen in a band, from Europe across to China. The skew from HadISD (Supplementary Information Fig. 7) shows increases in most regions except the majority of Europe and parts of Asia. The decreases in skew over Europe are strongest in the mid-morning (09:00) and weakest in the late evening (21:00). It also appears that the changes in standard deviation and skew are the somewhat inverse of each other, when comparing Figs. 6 and 7 of the Supplementary Information. 280 The decreasing (changing to more negative) skews indicate a shift from right to left-tailed distributions.



For HadGHCND in Donat and Alexander (2012), there are larger regions showing increases in standard deviation and a stronger increase around Hudson Bay for both maximum and minimum temperature values than for HadISD. For the skew, HadGHCND shows decreasing skew in Greenland, northern Africa, the coasts of eastern Asia and parts of Australia and increases over Europe, in contrast to the results of this study, which in some regions may be the result of low station densities
285 in HadISD. (Fig 1)

Although McKinnon et al. (2016) link the first four statistical moments derived from GHCND between 1980 and 2015 onto orthogonal basis functions, these still are closely tied to the original statistical moments. We also use their maximum and minimum temperatures to compare to the 15:00 and 06:00 local time values respectively. Over Europe, the geographical peak of trends in the daytime/maximum mean are both in eastern Europe. However, HadISD does not show such a strong
290 feature of increasing standard deviations across Asia, and GHCND does not show the decreasing daytime standard deviations in northern Europe, but those in eastern Asia do agree. The results from HadISD do not show daytime cooling in the eastern half of North America observed in GHCND. But the general decrease in variance is observed in both, though the increases in the central southern region of North America in GHCND is not present here. Over North America for the minimum/night-time temperatures, the agreement is better for the mean, standard deviation and also skew. And over Europe and Asia, the lower
295 magnitude trends for the mean also agree spatially, however the patterns for the higher moments do not.

4.2 Seasonal Temperature Changes

In the Supplementary Information Figs. 9-12 we show the changes for the first four moments from the 15:00 observations across the four seasons of the year (and Figs. 13-16 for the 06:00 observations). Again, the strongest variations are over Eurasia. Changes in the mean temperature at 15:00 are smallest in boreal winter¹ (DJF), and strongest over a wide area in spring (MAM).
300 However, summer (JJA) temperatures increase strongly with time around the Mediterranean and across continental Europe, and winter (DJF) has strong warming in the western part of Russia and Scandinavia. In North America, the largest trends are seen in SON (especially in higher latitudes), with little variation over the other seasons, especially spring and summer. A similar, but less intense pattern is observed at 06:00.

A smaller proportion of stations shows changes in the standard deviation where the uncertainty range does not encompass
305 zero, than for changes in the mean, but there are variations in the patterns of change over the year. Although increases in the standard deviation are seen across Asia in springtime and summer at 15:00, a more mixed pattern is present in this region in autumn; decreases are seen in the far east in winter. At 06:00, the pattern is somewhat similar, though with the Indian sub-continent and surrounding regions showing consistent decreases throughout the year. Over North America, standard deviations increase during spring, but mostly decrease during the other seasons, except in summer at 06:00. Further increases in the
310 standard deviation are also found around the Mediterranean and in south-eastern Europe for most of the year, with decreases

¹We will use the seasonal descriptions for the northern mid-latitudes, as the station density is highest and we are able to draw the clearest inferences. We note that these seasonal descriptions are not appropriate for all regions, and hence have defined them in the plots by months as well as the text where they are introduced.



in the north of the continent (British Isles and Scandinavia). Increases in the standard deviation are generally weaker and decreases more prevalent across all seasons at 06:00 then at 15:00.

Patterns of changes in the skew have some clear features, with decreases in China for all seasons except winter. Europe also shows decreases in the spring, but a more mixed pattern for the rest of the year. Increases in the skew at 15:00 are found in 315 northern high latitudes during spring, as well as India for most of the year. Over North America, decreases in skew are observed during spring and autumn in the southern half, but in the northern half during summer.

Comparing these seasonal plots with those showing changes in the distribution of average daily temperature from Cavanaugh and Shen (2014, 2015) shows generally good agreement for the changes in the mean. They find the strongest changes over 320 Europe and a clear “warming hole” over the US, especially in JJA and DJF. In HadISD, the strongest changes are over Europe, and the location of the largest values moves between the seasons; but the warming hole is only weakly apparent during JJA over North America. For standard deviation, both Cavanaugh and Shen (2014) and our study find coherent decreases in the Northern Hemisphere, as well as increases in MAM over North America. Over Europe, however, JJA shows the largest differences, with widespread increases in Cavanaugh and Shen (2014, 2015) but changes in both directions are clear in HadISD. McKinnon et al. (2016) also show the decrease in standard deviation over North America, except for a region of increase on the Gulf of 325 Mexico. Also across Eurasia, the patterns are similar in HadISD and McKinnon et al. (2016).

The positive trends in skewness found over North America (Cavanaugh and Shen, 2014, 2015) in GHCND are not as prominent in HadISD nor in McKinnon et al. (2016) also using GHCND, with contiguous regions of negative trends in all seasons. Part of these differences may of course be the result of the differing temporal periods used; HadISD using 1974-2018, and GHCND 1950-2010. We note that the number of stations reporting in HadISD prior to 1973 is much lower, restricting the 330 possibility of extending these analyses further into the past. Also, here we and McKinnon et al. (2016) do not combine the stations together and, unlike in Section 3 and in Cavanaugh and Shen (2014, 2015).

4.3 Dewpoint Temperature Changes

Comparing the changes in the dewpoint temperature moments across the day and the seasonal cycle (Supplementary Information Figs. 17 to 28) with those from the temperatures, shows that on the whole these two variables are quite closely aligned. 335 Despite fewer stations passing the selection criteria for this variable, there are still coherent patterns of change. Over the eight time points during the day, the changes in the mean dewpoints show less variation than the temperatures, most clearly apparent over Europe in the early afternoon. This suggests that the relative humidity over this region has decreased for this part of the day. A similar but less striking decrease in daytime relative humidity is also seen over North America. The changes in the higher moments of the dewpoint temperature are also reasonably constant through the day, but have a greater spatial heterogeneity 340 and lower magnitude than the temperatures. Notably, there is a clear increase in skew over south-east Asia during the daytime and early evening, which is not seen in the temperature results, indicating a change in shape away from a low-dewpoint tail towards a high-dewpoint tail.

Seasonally, the greatest change in the dewpoint mean is in SON and DJF over Europe in both the day- and night-time, contrasting with the temperatures of MAM and JJA for the day and also DJF at night. Again the standard deviations and



345 kurtoses are very similar for day and night across all seasons between the dewpoints and temperatures. For the skews, the increase in skew over south-east Asia noted above is most prominent during SON, during the daytime, but is also present at night and differs from the behaviour of the temperatures for all seasons.

4.4 Wind speed Changes

350 There are fewer stations with sufficient observations to assess changes in wind speed (Supplementary Information Figs. 29 to 33), and very few in the Southern Hemisphere. As a result we focus our analysis on North America and Eurasia. The proportion of stations where changes in the distribution could be assessed varies between each time step to a greater extent than for the two temperature variables. Despite this, regions with sufficient station densities show a diurnal cycle in the behaviour of wind speed trends. There are stronger trends for declining wind speeds during the daytime (09:00-18:00) than for the night-time hours. And there are also no large regions that show large increases (at the same magnitude as decreases) in the wind speeds at 355 any time point. However some individual stations do show some increases, for example in south-eastern Europe. For the other three moments (not shown) there are even smaller differences than for the mean for the annual changes over the day.

360 We show the seasonal changes at 15:00 for all four moments in Supplementary Information Figs. 30 to 33. For the mean there are no large changes across the seasons. The changes in standard deviation are also generally unvarying compared to the temperature measures. There are increases in the USA, with stronger values during spring and autumn. All other regions with sufficient station density have decreases in the standard deviation. Both skew and kurtosis show no strong seasonal variations. Skews are increasing over North America but there are no coherent, strong changes elsewhere. The kurtosis is mostly decreasing in south-eastern Europe, and also on the whole across Asia.

4.5 Summary

365 Across all seasons, times of day and locations, there are strong increases in the means of both temperature and dewpoint temperature with time. Although the geographical distribution of stations is not uniform, there appear to be larger increases in Europe, especially during the spring and summer, than elsewhere in the world. The largest increases in the high station density regions of Europe also occur during the daytime rather than night-time, which matches the behaviour seen in Table 1 40-60N (the latitudes corresponding to Europe). However, in most of the other latitude bands in Table 1 the night-time means are increasing faster than the daytime ones. The changes in the standard deviations show with stronger decreases in the 370 northern latitudes, but more regions with increases in the mid- and tropical latitudes in the northern hemisphere. Seasonally, North America shows decreases in the daytime and night-time standard deviation in winter and summer, but increases in spring (Supplementary Information Figs. 10 and 14), which was also observed in Cavanaugh and Shen (2015). This indicates a change towards less variability in winter and summer, but a change towards greater variability in spring for both day- and night-time temperatures.

375 We also note that in Section 3, standard deviations were decreasing more or less across the whole northern hemisphere, whereas in this study there are regions with clear increases. The zonal study combines the stations together, even if individually any change in the statistical moments over time would not show clear changes (1σ range of the trends encompassing zero). In



contrast, Figs. 3 and 4 only show those stations where the 1σ range of fitted trends do not include zero. As noted above, this is only around half of the total number of stations. Earlier versions of these figures plotted all stations, and suggested that many
380 of these exhibit decreases in the standard deviation. However, for clarity, we have decided to only plot those stations in which the trend is clearly different to zero here.

Changes in annual skew are different on either side of the North Atlantic, with predominantly increases in North America, and decreases in Europe across the day, and hence a rather small change in the zonally averaged results (Section 3). The seasonal results show increases at higher latitudes (Canada and Scandinavia), and decreases further south (USA and continental
385 Europe) in spring. Other seasons show a more heterogeneous pattern in Europe, but increases across central Asia in summer and autumn, similar to the patterns from GHCND in Cavanaugh and Shen (2015). In China which was not included in their study, all seasons except winter show a decreases in the skews. For the kurtoses, Cavanaugh and Shen (2015) found strong decreases in the southern parts of North America in the summer, which are not seen in HadISD. However, the strong decreases over parts of Europe in winter are present in HadISD.

390 5 Quantile Regression

Studies of the characteristics of a distribution are useful in describing the overall changes in a meteorological variable. However, they do not make the changes in extremes immediately clear, although they can be inferred. An alternative way to study the changes in the climate that show changes in past extreme events is to investigate the changes of the quantiles of a distribution themselves. This enables the question of “how much warmer are the warmest 10 per cent of days now than they were in
395 the past?” to be addressed. Quantile regression calculates trends in specified quantiles of a distribution, rather than trends in the mean (Koenker and Bassett Jr, 1978; Koenker and Hallock, 2001; Barbosa et al., 2011; Franzke, 2015). This allows, for example, cases where there is little or no mean trend but an increased variance over time to be assessed more completely. In the case of climate observations, and especially given the impact of extreme events on society and infrastructure, quantile regression allows the changes in these extremes over time to be made apparent.

400 The observations in HadISD naturally contain both an annual and a diurnal cycle. If run purely on these raw data, then the annual upper quantiles are likely dominated by summer daytime temperatures, and the lower by winter night-time temperatures. Although using standard three-month seasons would enable the more detailed study of winter and summer quantiles, those from the shoulder seasons (spring and autumn) would be dominated by the earlier or later parts of these seasons given the strong annual cycle. For this reason McKinnon et al. (2016) use daily maximum and minimum temperature data from GHCND over
405 July and August only to reduce the effect of the seasonal cycle in their study.

Our approach is to use de-seasonalised data, which removes the seasonal aspect described above and also the issue that quantile regression results can be biased when applied to discrete data (Machado and Silva, 2005; McKinnon et al., 2016). The observations in HadISD are at a finite numerical resolution (e.g. temperatures are reported to the nearest 0.1°C or 1°C). We use a daily climatology to create hourly anomalies of the HadISD data. Although within each calendar day, the numerical
410 resolution remains unchanged by the subtraction of a daily mean from the hourly observations (within-day differences will still




Figure 5. An example of the Quantile Regression analysis for a single station, with the trends in the quantiles in red, and the ordinary least squares (OLS) fit in yellow. This is for station 154810-99999, Constanța Mihail Kogălniceanu International Airport, Romania.

have 0.1°C or 1°C resolution). Between each calendar day, the different climatology value will result in a different offset, so that between-day differences would not result in discrete values. Hence, overall, the values passed to the quantile regression algorithm will be less discrete than the raw observations themselves.

Daily means were calculated for 1981-2010, requiring at least eight observations in each 24-hour period (equivalent to 415 3-hourly data). The daily climatology was calculated requiring at least 15 years of data to obtain an average. Finally, the climatology was smoothed using a 5-point binomial filter and subtracted from the sub-daily data for each day to make the climate anomalies. An example of the changes in the percentiles of a timeseries is shown in Fig. 5.

Both Europe and North America have high station densities (see Section 4) compared with other regions, and so our analysis is focussed on these two regions (see also McKinnon et al., 2016). Furthermore, we show the results using a space-filling 420 method; Voronoi tessellation. Voronoi tessellation (derived from Delaunay triangulation, Voronoï, 1908; Delaunay, 1934) uses the perpendicular bisectors of the lines joining each station to split the land surface up into polygons. These have then been coloured to show the results of the quantile regression. Stations for which the slope of the quantile shown is not significantly different from zero ($p > 0.05$ as determined by the quantile regression algorithm) are shown in grey, but we do no further significance testing on these trends. It should be noted that areas of high station density will naturally have smaller polygons, 425 but the human eye is drawn to the larger blocks of colour arising from sparse station networks. However, the advantage of this approach is that the underlying station information is retained, as opposed to it being smoothed when applying some form of gridding method.

We show the changes in the quantiles (0.01, 0.05, 0.1, 0.9, 0.95, and 0.99) over Europe for both JJA and DJF in Figs. 6 and 7 respectively. In some sub-regions large numbers of stations show similar patterns, suggesting that there is a coherent change in 430 the quantile in that area. However, within these there may be isolated stations/cells showing changes in the opposite direction.



Although we have selected those stations that do not have many or large breaks in their timeseries (as determined by PHA, see Section 2), and the HadISD has undergone a number of quality control checks, there will inevitably be a few data quality issues remaining.

The largest change in the summer (JJA) quantiles is observed in south-eastern Europe and western Asia (Fig. 6), focussed
435 on the area north and west of the Black Sea. The increase observed in these regions becomes progressively less strong towards lower quantiles and also with distance from this region. This indicates that the temperature distribution is broadening over time, with the upper tails changing more rapidly than the lower tails; i.e. the warmest N per cent of events are becoming warmer, more rapidly than the cool events. In contrast, the regions close to the North Sea and especially Scandinavia show slight increases for the lower quantiles that becomes less strong for higher quantiles. For the north and west of the UK in
440 general and Scandinavia except western Norway, there are even decreases for the highest quantiles.

Conversely, in the winter (DJF), the strongest increases are in the northern regions and the lower quantiles, with Scandinavia, the Baltic states along with Benelux, Denmark and northern Germany indicating a narrowing of the temperature distribution over time. In the higher quantiles, there are increases for eastern Europe but not as much as in the summer. Westernmost Europe shows very little change across all quantiles. We show the panels for the shoulder seasons (MAM and SON) in the
445 Supplementary Information Figs. 34 and 35. There is a general increase for all the percentiles, and the cold quantiles in the north east of the region shown are warming faster than the rest. But in contrast to Figs. 6 and 7 these increases are much slower and the regions are less coherent. Overall, these results show that warm extremes are becoming more common more quickly during the summer in south-eastern Europe, than elsewhere across the continent. Similarly, cool extremes are becoming less common more quickly during winter in northern Europe than elsewhere.

450 Franzke (2015) performed quantile regression on daily mean, maximum and minimum temperatures between 1950 and 2013. Mean temperatures, Franzke (2015) show strong regions of increase in both the 5th and 95th percentiles in eastern Europe and western Russia, north of the Black Sea, with the higher percentile region covering a larger area than for the results presented here. When using summer maxima, Franzke (2015) show a more uniform increase over all of Europe at the higher percentile, except Scandinavia, which behaves similarly in Fig. 6. The magnitudes of the change in the winter temperature percentiles
455 from HadISD are consistent with the studies of Barbosa et al. (2011); Franzke (2015). The behaviour over Turkey stands out in their study, but not in HadISD, suggesting a difference in the way these data have been processed or effects arising from the difference in temporal coverage or even station selection.

The difference in behaviour of the lower and upper quantiles indicate a change in the variance of the sub-daily temperatures over Europe, which has already been suggested by the results from Section 4. During the summer, the variance in the large
460 region to the north of the Black Sea, has increased. In contrast, during winter the variance over Scandinavia and northern Europe has decreased. Both of these changes are also clearly visible in Supplementary Information Fig. 10 showing the seasonal change of the standard deviation for each station. Franzke (2015) found a more widespread, uniform decrease in variance in the winter months over Europe, but with a stronger decrease over Scandinavia. However, Franzke (2015) found no increase in variance during the summer in eastern Europe.




Figure 6. Trend over 1973-2017 in quantiles of temperature ($^{\circ}\text{C decade}^{-1}$) at TOP 0.01, 0.05 and 0.10; and BOTTOM 0.90, 0.95, 0.99 over Europe during summer (JJA).

465 Over North America (Supplementary Information Figs. 36 to 39), the trends in quantiles are on the whole smaller than for Europe for most of the year. In the spring (MAM), there are moderate increases in the south-west, which are stronger for the higher quantiles. For the summer (JJA) the higher quantiles show decreases in the northern, central regions, but increases for the lower quantiles in the south west. In the autumn (SON) and winter (DJF) the strongest changes are in the north of the continent at the lower quantiles with more moderate increases in the south and the higher quantiles. Hence the most rapid 470 changes are in the cool extremes during autumn and winter in the north, with less rapid changes in the warm extremes in spring in the south-west.

475 Using GHCND, Rhines et al. (2017) show a similar pattern, with a strong increase in lower percentiles for northern latitudes in winter. In the spring their data show decreases in the lower percentiles in the north and increases in the south, almost the opposite in the autumn, and general warming for the minimum temperatures in the summer. We also find the strongest changes for the lower percentiles during winter, indicating a reduction in the range of temperatures during that season.

We also show changes in the dewpoint temperatures (Supplementary Information Figs. 40 to 43) and wind speeds (Fig. 44) over Europe. The dew point temperatures indicate small increases of the quantiles in most regions with no strong regional differences for MAM and JJA for any quantile. However, in SON and DJF, there are stronger increases in the lower quantiles




Figure 7. Trend over 1973-2017 in quantiles of temperature ($^{\circ}\text{C decade}^{-1}$) at TOP 0.01, 0.05 and 0.10; and BOTTOM 0.90, 0.95, 0.99 over Europe during winter (DJF).

for the north of the region (Scandinavia and north-western Russia), but small increases for other regions and quantiles. So,
 480 although the summer warm extremes are becoming more common in south-eastern Europe, there is no corresponding change
 in the dew point temperatures, suggesting a decrease in the relative humidity. However, in the northern parts, the dew point
 temperatures are changing roughly in step with the dry-bulb temperatures.

Only the annual changes in wind speeds are shown (Supplementary Information Fig. 44), with some blank tiles and regions
 because of the smaller number of stations with sufficient observations. There are no large changes in the lower quantiles
 485 but clear and larger decreases in the upper quantiles, which matches the behaviour observed in Section 3 and the decrease in
 variance observed in Section 4.4. The larger changes in the upper quantiles also suggest the skews are becoming more negative,
 with decreases in the number and intensities of extremely high wind speeds.

6 Summary

To investigate the changes in the shape of the distributions of temperatures and wind speeds in the HadISD dataset, we have
 490 performed three types of assessment. Firstly, by combining the data from the station based observations into zonal averages,



the resulting distributions are built from large numbers of observations at the expense of some spatial information. Using five-year bins from 1973 to 2017, we demonstrated the change in the distributions and percentile values. As expected, the mean temperatures increase for both day- and night-time. However, the standard deviations decrease for the northern hemisphere (where there are most stations). Although changes are seen in both tails of the distribution, they are greater in the lower tail.
495 This behaviour is consistent with both the more rapid change in low temperature extremes (Davy et al., 2017) as well as the decrease in diurnal temperature range (Alexander et al., 2006; Thorne et al., 2016b, a). The dewpoint temperatures show similar, if more heterogeneous, behaviour in this analysis.

In contrast, the wind speeds showed decreases in the mean, but also in the standard deviations, with larger shifts in the upper tail than the lower. A number of studies have investigated changes in global wind speeds, showing a steady decrease over the
500 past decades, termed “stilling” (e.g. Roderick et al., 2007; McVicar et al., 2012). The changes in distributions observed on a zonal level suggest this may be driven by changes in the upper tail.

To improve the spatial discrimination, we also assess changes in the distributions for each station individually, and show fitted linear trends. The sub-daily nature of HadISD allows these to be calculated across the hours of the local day as well as annually or seasonally. The temperature means show the expected increase over time, but with a daily cycle showing a
505 more rapid increase of mean temperatures during the early afternoon than in the early hours of the morning. However, the higher order moments are much less homogeneous, but have a smaller variation throughout the day. Trends in the standard deviations are on the whole negative, except for a band across central Asia. Conversely, trends in the skew of the temperatures are mainly positive, except over Europe and across into western Asia, indicating increasing warm extremes (positive tail) and decreasing cool extremes. These patterns are in some regions different from those reported in previous investigations by Donat
510 and Alexander (2012) and McKinnon et al. (2016) obtained using (derivatives of) GHCND (true maximum and minimum values). For the seasonal changes, which can be compared to Cavanaugh and Shen (2014), there are also differences to the patterns obtained from GHCND.

These disparities may result from a variety of differences in the datasets and methods. Firstly, the period used in this study is 1973-2017 compared to 1950-2010 (Donat and Alexander, 2012; Cavanaugh and Shen, 2014) and 1980-2015 (McKinnon
515 et al., 2016). Also, by using all of the sub-daily observations in HadISD, this work incorporates the effect of changes across the entire distribution, even if we are concentrating on either the moments or specific percentile values (e.g. all temperature observations rather than just the maxima and minima). The studies using GHCND use the true maximum and minimum values, which allow the focussed investigation of the momentary extremes. We also do not perform any spatial smoothing of the underlying stations, which results in less variable and less extreme fields compared to station data (Gross et al., 2018).

520 Finally, rather than study the moments of the distributions, we perform a quantile regression analysis to determine how the values of the percentiles have changed, and therefore how the warmest and coolest days feel in comparison to others. This approach is in contrast to the ETCCDI indices which use percentiles (e.g. TX90p), which capture the exceedence rate of a fixed percentile value. Here we show that over Europe, the upper percentiles are changing fastest in the large region north-west of the Black Sea in the summer, but the lower percentiles are changing faster in northern Europe and Scandinavia in the winter.
525 Hence the warmest N per cent of summer days are becoming warmer more rapidly than the coolest N per cent in south-eastern



Europe, and the reverse for winter in the north. Coherent changes in dewpoint temperatures are less clear, with increases in the lower percentiles in autumn and winter for the northern parts of Europe. For the wind speeds, as expected from the other analyses, the higher quantiles show a decline over Europe, with stronger decreases in the east.

All the studies presented herein show widespread increases in mean temperature, and less rapid rises in the dewpoints, as 530 expected from numerous other studies on the observed change in global temperatures. However, the changes in the higher moments are of more interest. Although in many cases (seasons and quantities) the patterns of change are very similar to prior studies (e.g. Donat and Alexander, 2012; Cavanaugh and Shen, 2014, 2015; McKinnon et al., 2016), there are still differences in the spatial patterns between all of the studies. These could arise from differences in the input data (daily extrema versus hourly 535 values) as well as the temporal and spatial coverage. Furthermore, we have used a single, station-based dataset, and differences between datasets (Gross et al., 2018) or the effect of gridding (Cavanaugh and Shen, 2015) can change the apparent behaviour of higher moments. However, in these investigations, we have shown that there are changes in the mean, the standard deviations and the skews of the distributions studied. And, when using a sub-daily dataset of temperatures, changes in the number and intensity of extremes appear to arise from the combination of changes in all three of these moments.

Data availability. The HadISD dataset is available under a non-commercial government licence at www.metoffice.gov.uk/hadobs/hadisd

540 *Author contributions.* Robert Dunn did the majority of the analysis, plotting and writing. Kate Willett and David Parker provided comments and guidance during this work. All authors have contributed text and edits to the main paper.

Competing interests. The authors declare there are no competing interests

545 *Copyright statement.* The works published in this journal are distributed under the Creative Commons Attribution 4.0 Licence. This licence does not affect the Crown Copyright work, which is re-usable under the Open Government Licence (OGL). The Creative Commons Attribution 4.0 Licence and the OGL are interoperable and do not conflict with, reduce or limit each other.

Acknowledgements. We thank Colin Morice for helpful discussions during the course of this work, and Elizabeth Good and Nick Rayner for useful suggestions on the text. This work was supported by the Met Office Hadley Centre Climate Programme funded by BEIS and Defra (GA01101).



References

550 Alexander, L., Zhang, X., Peterson, T., Caesar, J., Gleason, B., Tank, A. K., Haylock, M., Collins, D., Trewin, B., Rahimzadeh, F., et al.: Global observed changes in daily climate extremes of temperature and precipitation, *Journal of Geophysical Research: Atmospheres*, 111, 2006.

Azorin-Molina, C., Dunn, R. J. H., Mears, C., Berrisford, P., and McVicar, T.: Surface Wind Speed [in "State of the Climate in 2017"], *Bulletin of the American Meteorological Society*, 99, S41–42, <https://doi.org/10.1175/2018BAMSStateoftheClimate.1>, 2018.

555 Ballester, J., Giorgi, F., and Rodó, X.: Changes in European temperature extremes can be predicted from changes in PDF central statistics, *Climatic change*, 98, 277, 2010.

Barbosa, S., Scotto, M., and Alonso, A.: Summarising changes in air temperature over Central Europe by quantile regression and clustering, *Natural Hazards and Earth System Sciences*, 11, 3227–3233, 2011.

Brown, S., Caesar, J., and Ferro, C. A.: Global changes in extreme daily temperature since 1950, *Journal of Geophysical Research: Atmospheres*, 113, 2008.

560 Caesar, J., Alexander, L., and Vose, R.: Large-scale changes in observed daily maximum and minimum temperatures: Creation and analysis of a new gridded data set, *Journal of Geophysical Research: Atmospheres*, 111, 2006.

Cavanaugh, N. R. and Shen, S. S.: Northern Hemisphere climatology and trends of statistical moments documented from GHCN-daily surface air temperature station data from 1950 to 2010, *Journal of Climate*, 27, 5396–5410, 2014.

565 Cavanaugh, N. R. and Shen, S. S.: The effects of gridding algorithms on the statistical moments and their trends of daily surface air temperature, *Journal of Climate*, 28, 9188–9205, 2015.

Christidis, N., Stott, P. A., and Brown, S. J.: The role of human activity in the recent warming of extremely warm daytime temperatures, *Journal of Climate*, 24, 1922–1930, 2011.

Davy, R., Esau, I., Chernokulsky, A., Outten, S., and Zilitinkevich, S.: Diurnal asymmetry to the observed global warming, *International Journal of Climatology*, 37, 79–93, <https://doi.org/10.1002/joc.4688>, <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/joc.4688>, 2017.

570 Delaunay, B.: Sur la sphère vide, *Izv. Akad. Nauk SSSR, Otdelenie Matematicheskii i Estestvennyka Nauk*, 7, 1–2, 1934.

Della-Marta, P. M., Haylock, M. R., Luterbacher, J., and Wanner, H.: Doubled length of western European summer heat waves since 1880, *Journal of Geophysical Research: Atmospheres*, 112, 2007.

Donat, M., Alexander, L., Yang, H., Durre, I., Vose, R., Dunn, R., Willett, K., Aguilar, E., Brunet, M., Caesar, J., et al.: Updated analyses of 575 temperature and precipitation extreme indices since the beginning of the twentieth century: The HadEX2 dataset, *Journal of Geophysical Research: Atmospheres*, 118, 2098–2118, 2013a.

Donat, M. G. and Alexander, L. V.: The shifting probability distribution of global daytime and night-time temperatures, *Geophysical Research Letters*, 39, 2012.

Donat, M. G., Alexander, L. V., Yang, H., Durre, I., Vose, R., and Caesar, J.: Global land-based datasets for monitoring climatic extremes, 580 *Bulletin of the American Meteorological Society*, 94, 997–1006, 2013b.

Dunn, R.: HadISD version 3: Monthly Updates, Met Office Hadley Centre Technical Note, 2019.

Dunn, R., Willett, K., Thorne, P., Woolley, E., Durre, I., Dai, A., Parker, D., and Vose, R.: HadISD: a quality-controlled global synoptic report database for selected variables at long-term stations from 1973–2011, *Climate of the Past*, 8, 1649–1679, 2012.

Dunn, R., Willett, K., Morice, C., and Parker, D.: Pairwise homogeneity assessment of HadISD, *Climate of the Past*, 10, 1501–1522, 2014.



585 Dunn, R., Willett, K., Parker, D., and Mitchell, L.: Expanding HadISD: quality-controlled, sub-daily station data from 1931, *Geoscientific
Instrumentation, Methods and Data Systems*, 5, 473–491, 2016.

Field, C. B., Barros, V., Stocker, T. F., and Dahe, Q.: *Managing the risks of extreme events and disasters to advance climate change adaptation: special report of the intergovernmental panel on climate change*, Cambridge University Press, 2012.

Franzke, C. L.: Local trend disparities of European minimum and maximum temperature extremes, *Geophysical Research Letters*, 42, 6479–
590 6484, 2015.

Griffiths, G., Chambers, L., Haylock, M., Manton, M., Nicholls, N., Baek, H.-J., Choi, Y., Della-Marta, P., Gosai, A., Iga, N., et al.: Change
in mean temperature as a predictor of extreme temperature change in the Asia–Pacific region, *International Journal of Climatology: A
Journal of the Royal Meteorological Society*, 25, 1301–1330, 2005.

Gross, M. H., Donat, M. G., Alexander, L. V., and Sisson, S. A.: The sensitivity of daily temperature variability and extremes to dataset
595 choice, *Journal of Climate*, 31, 1337–1359, 2018.

Huntingford, C., Jones, P. D., Livina, V. N., Lenton, T. M., and Cox, P. M.: No increase in global temperature variability despite changing
regional patterns, *Nature*, 500, 327, 2013.

Koenker, R. and Bassett Jr, G.: Regression quantiles, *Econometrica: journal of the Econometric Society*, pp. 33–50, 1978.

Koenker, R. and Hallock, K. F.: Quantile regression, *Journal of economic perspectives*, 15, 143–156, 2001.

600 Lanzante, J. R.: Resistant, Robust and Non-Parametric techniques for the analysis of Climate Data: Theory and Examples, including Applications
to Historical Radiosonde Station Data, *International Journal of Climatology*, 16, 1197–1226, 1996.

Lewis, S. C. and King, A. D.: Evolution of mean, variance and extremes in 21st century temperatures, *Weather and climate extremes*, 15,
1–10, 2017.

Machado, J. A. F. and Silva, J. S.: Quantiles for counts, *Journal of the American Statistical Association*, 100, 1226–1237, 2005.

605 McKinnon, K. A., Rhines, A., Tingley, M. P., and Huybers, P.: The changing shape of Northern Hemisphere summer temperature distributions,
Journal of Geophysical Research: Atmospheres, 121, 8849–8868, 2016.

McVicar, T. R., Roderick, M. L., Donohue, R. J., Li, L. T., Van Niel, T. G., Thomas, A., Grieser, J., Jhajharia, D., Himri, Y., Mahowald,
N. M., et al.: Global review and synthesis of trends in observed terrestrial near-surface wind speeds: Implications for evaporation, *Journal
of Hydrology*, 416, 182–205, 2012.

610 Menne, M. J. and Williams Jr, C. N.: Homogenization of temperature series via pairwise comparisons, *Journal of Climate*, 22, 1700–1717,
2009.

Menne, M. J., Durre, I., Vose, R. S., Gleason, B. E., and Houston, T. G.: An overview of the global historical climatology network-daily
database, *Journal of Atmospheric and Oceanic Technology*, 29, 897–910, 2012.

Rhines, A., McKinnon, K. A., Tingley, M. P., and Huybers, P.: Seasonally resolved distributional trends of North American temperatures
615 show contraction of winter variability, *Journal of Climate*, 30, 1139–1157, 2017.

Roderick, M. L., Rotstayn, L. D., Farquhar, G. D., and Hobbins, M. T.: On the attribution of changing pan evaporation, *Geophysical Research
Letters*, 34, n/a–n/a, <https://doi.org/10.1029/2007GL031166>, <http://dx.doi.org/10.1029/2007GL031166>, 2007.

Sen, P. K.: Estimates of the Regression Coefficient Based on Kendall's Tau, *Journal of the American Statistical Association*, 63, pp. 1379–
1389, 1968.

620 Shen, S. S., Gurung, A. B., Oh, H.-S., Shu, T., and Easterling, D. R.: The twentieth century contiguous US temperature changes indicated by
daily data and higher statistical moments, *Climatic Change*, 109, 287–317, 2011.



Simolo, C., Brunetti, M., Maugeri, M., and Nanni, T.: Evolution of extreme temperatures in a warming climate, *Geophysical research letters*, 38, 2011.

Smith, A., Lott, N., and Vose, R.: The integrated surface database: Recent developments and partnerships, *Bulletin of the American Meteorological Society*, 92, 704–708, 2011.

625 Thorne, P., Donat, M., Dunn, R., Williams, C., Alexander, L., Caesar, J., Durre, I., Harris, I., Hausfather, Z., Jones, P., et al.: Reassessing changes in diurnal temperature range: Intercomparison and evaluation of existing global data set estimates, *Journal of Geophysical Research: Atmospheres*, 121, 5138–5158, 2016a.

Thorne, P., Menne, M., Williams, C., Rennie, J., Lawrimore, J., Vose, R., Peterson, T. C., Durre, I., Davy, R., Esau, I., et al.: Reassessing 630 changes in diurnal temperature range: A new data set and characterization of data biases, *Journal of Geophysical Research: Atmospheres*, 121, 5115–5137, 2016b.

Thorne, P. W., Allan, R. J., Ashcroft, L., Brohan, P., Dunn, R. J. H., Menne, M. J., Pearce, P. R., Picas, J., Willett, K. M., Benoy, M., 635 Bronnimann, S., Canziani, P. O., Coll, J., Crouthamel, R., Compo, G. P., Cuppett, D., Curley, M., Duffy, C., Gillespie, I., Guijarro, J., Jourdain, S., Kent, E. C., Kubota, H., Legg, T. P., Li, Q., Matsumoto, J., Murphy, C., Rayner, N. A., Rennie, J. J., Rustemeier, E., Slivinski, L. C., Slonosky, V., Squintu, A., Tinz, B., Valente, M. A., Walsh, S., Wang, X. L., Westcott, N., Wood, K., Woodruff, S. D., and Worley, S. J.: Toward an Integrated Set of Surface Meteorological Observations for Climate Science and Applications, *Bulletin of the American Meteorological Society*, 98, 2689–2702, <https://doi.org/10.1175/BAMS-D-16-0165.1>, 2017.

Voronoi, G.: Nouvelles applications des paramètres continus à la théorie des formes quadratiques. Deuxième mémoire. Recherches sur les parallélépédres primitifs., *Journal für die reine und angewandte Mathematik*, 134, 198–287, 1908.