

Interactive comment on “The Half-order Energy Balance Equation, Part 2: The inhomogeneous HEBE and 2D energy balance models” by Shaun Lovejoy

Anonymous Referee #2

Received and published: 13 July 2020

This second part reviewed here extends the approach of Part 1 to higher spatial dimension and inhomogeneous thermal models of the earth's response to radiative forcing. There is an appropriate summary of Part 1 that puts the new contribution into context. The full model considered here includes varying horizontal and vertical thermal diffusivities, thermal capacities, sensitivities and spatio-temporal forcing. By a heuristic method of Babenko, the author expands the inhomogeneous operator to give 2D energy balance equations that will be useful for studying spatio-temporal responses to forcing. The manuscript includes a number of appendices that examine horizontal structures, cross-correlations, space-time factorization of quantities such as autocorrelation and that extends the results from flat space to the sphere. The analysis seems

C1

to be carefully done, and care is taken to distinguish cases where there may not be a rigorous justification. I would be interested to see a bit more discussion of the “bottom boundary condition” $T=0$ at $z=-\infty$. I think it would also be useful to include some discussion of how atmosphere/ocean convection is/is not represented in the model.

Interactive comment on Earth Syst. Dynam. Discuss., <https://doi.org/10.5194/esd-2020-13>, 2020.

C2