

Regional variation in the effectiveness of methane-based and land-based climate mitigation options

Garry D. Hayman^{1*}, Edward Comyn-Platt¹, Chris Huntingford¹, Anna B. Harper², Tom Powell², Peter M. Cox², William Collins³, Christopher Webber³, Jason Lowe^{4,5}, Stephen Stith², Joanna I. House⁶, Jonathan C. Doelman⁷, Detlef P. van Vuuren^{7,8}, Sarah E. Chadburn², Eleanor Burke⁵, Nicola Gedney⁹.

¹ Centre for Ecology & Hydrology, Wallingford, OX10 8BB, U.K.

² University of Exeter, Exeter, EX4 4QF, U.K.

³ University of Reading, Reading, RG6 6BB, U.K.

⁴ University of Leeds, Leeds, LS2 9JT, U.K.

⁵ Met Office Hadley Centre, FitzRoy Road, Exeter, EX1 3PB, U.K.

⁶ Cabot Institute for the Environment, University of Bristol, Bristol, BS8 1SS, U.K.

⁷ Department of Climate, Air and Energy, Netherlands Environmental Assessment Agency (PBL), PO Box 30314, 2500 GH The Hague, Netherlands

⁸ Copernicus Institute of Sustainable Development, Utrecht University, Heidelberglaan 2, 3584 CS, the Netherlands

⁹ Met Office Hadley Centre, Joint Centre for Hydrometeorological Research, Wallingford, OX10 8BB, U.K.

Correspondence to: Garry Hayman (garr@ceh.ac.uk)

Abstract. Scenarios avoiding global warming greater than 1.5 or 2°C, as stipulated in the Paris Agreement, may require the combined mitigation of anthropogenic greenhouse gas emissions alongside enhancing negative emissions through approaches such as afforestation/reforestation (AR) and biomass energy with carbon capture and storage (BECCS). We use the JULES land-surface model coupled to an inverted form of the IMOGEN climate emulator to investigate mitigation scenarios that achieve the 1.5 or 2°C warming targets of the Paris Agreement. Specifically, within this IMOGEN-JULES framework, we focus on and characterise the global and regional effectiveness of land-based (BECCS and/or AR) and anthropogenic methane (CH₄) emission mitigation, separately and in combination, on the anthropogenic fossil fuel carbon dioxide (CO₂) emission budgets (AFFEBs) to 2100. We, using use consistent data and socio-economic assumptions from the IMAGE integrated assessment model, for the second Shared Socioeconomic Pathway (SSP2). The analysis includes the effects of the methane and carbon-climate feedbacks from wetlands and permafrost thaw, which we have shown previously to be significant constraints on the AFFEBs.

Globally, mitigation of anthropogenic CH₄ emissions has large impacts on the anthropogenic fossil fuel emission budgets, potentially offsetting (i.e. allowing extra) carbon dioxide emissions of 188-212 GtC. This is because of (a) the reduction in the direct and indirect radiative forcing of methane in response to the lower emissions and hence atmospheric concentration of methane; and (b) carbon-cycle changes leading to increased uptake by the land and ocean by CO₂-based fertilisation. Methane mitigation is beneficial everywhere, particularly for the major CH₄-emitting regions of India, USA and China. Land-based mitigation has the potential to offset 51-100 GtC globally, the large range reflecting assumptions and uncertainties associated with BECCS. The ranges for CH₄ reduction and BECCS implementation are valid for both the 1.5° and 2°C warming targets.

Commented [HGD1]: Response 7

Commented [HGD2]: Response 12

35 That is the mitigation potential of the CH₄ and of the land-based scenarios is similar for whether society aims for one or other
of the final stabilised warming levels. Further, both the effectiveness and the preferred land-management strategy (i.e., AR or
BECCS) have strong regional dependencies. Additional analysis shows extensive BECCS could adversely affect water security for
several regions. Although the primary requirement remains mitigation of fossil fuel emissions, our results highlight the
40 unrealised potential for the mitigation of CH₄ emissions to make the Paris climate targets more achievable.

Commented [HGD3]: Response 28

Commented [HGD4]: Response 8

1 Introduction

The stated aims of the Paris Agreement of the United Nations Framework Convention on Climate Change (UNFCCC, 2015) are “to hold the increase in global average temperature to well below 2°C and to pursue efforts to limit the increase to 1.5°C”. The global average surface temperature for the decade 2006-2015 was 0.87°C above pre-industrial levels and is likely
45 to reach 1.5°C between the years 2030 and 2052, if global warming continues at current rates (IPCC, 2018). The IPCC Special Report on Global Warming of 1.5°C (IPCC, 2018) gives the median remaining carbon budgets between 2018 and 2100 as 770 GtCO₂ (210 GtC) and 1690 GtCO₂ (~461 GtC) to limit global warming to 1.5°C and 2°C, respectively. These budgets represent ~20 and ~41 years at present-day emission rates. The actual budgets could however be smaller, as they exclude Earth system
50 feedbacks such as CO₂ released by permafrost thaw or CH₄ released by wetlands. Meeting the Paris Agreement goals will, therefore, require sustained reductions in sources of fossil carbon emissions, other long-lived anthropogenic greenhouse gases (GHGs) and some short-lived climate forcers (SLCFs) such as methane (CH₄), alongside increasingly extensive implementations of carbon dioxide removal (CDR) technologies (IPCC, 2018). Accurate information is needed on the range and efficacy of options available to achieve this.

Biomass energy with carbon capture and storage (BECCS) and afforestation/reforestation (AR) are among the most widely
55 considered CDR technologies in the climate and energy literature (Minx et al., 2018). For scenarios consistent with a 2°C warming target, the review by Smith et al. (2016) finds this may require (1) a median removal of 3.3 GtC yr⁻¹ from the atmosphere through BECCS by 2100 and (2) a mean CDR through AR of 1.1 GtC yr⁻¹ by 2100, giving a total CDR equivalent to 47% of present-day emissions from fossil fuel and other industrial sources (Le Quéré et al., 2018). Although there are fewer scenarios that look specifically at the 1.5°C pathway, BECCS is still the major CDR approach (Rogelj et al., 2018). For the
60 default assumptions in Fuss et al. (2018), BECCS would remove a median of 4 GtC yr⁻¹ by 2100 and a total of 41-327 GtC from the atmosphere during the twenty-first century, equivalent to about 4-30 years of current annual emissions. The land requirements for BECCS will be greater for the 1.5°C target within a given shared socio-economic pathway (e.g., SSP2), although published estimates are similar for the two warming targets, with between 380-700 Mha required for the 2°C target (Smith et al., 2016) and greater than 600 Mha for the 1.5°C target (van Vuuren et al., 2018). This is because the land
65 requirements for bioenergy production differ strongly across the different SSPs, depending on assumptions about the contribution of residues, assumed yields and yield improvement, start dates of implementation and the rates of deployment.

70 While the CDR figures assume optimism about the mitigation potential of BECCS, concerns have been raised about the potentially detrimental impacts of BECCS on food production, water availability and biodiversity, e.g., (Krause et al., 2017; Heck et al., 2018). Others note the risks and query the feasibility of large-scale deployment of BECCS e.g. (Anderson and Peters, 2016; Vaughan and Gough, 2016; Vaughan et al., 2018).

75 Harper et al. (2018) find the overall effectiveness of BECCS to be strongly dependent on the assumptions concerning yields, the use of initial above-ground biomass that is replaced and the calculated fossil-fuel emissions that are offset in the energy system. Notably, if BECCS involves replacing ecosystems that have higher carbon contents than energy crops, then AR and avoided deforestation can be more efficient than BECCS for atmospheric CO₂ removal over this century (Harper et al., 2018).

80 Mitigation of the anthropogenic emissions of non-CO₂ GHGs such as CH₄ and of SLCFs such as black carbon have been shown to be attractive strategies with the potential to reduce projected global mean warming by 0.22-0.5°C by 2050 (Shindell et al., 2012; Stohl et al., 2015). It should be noted that these were based on scenarios with continued use of fossil fuels. Through the link to tropospheric ozone (O₃), there are additional co-benefits of CH₄ mitigation for air quality, plant productivity and food production (Shindell et al., 2012) and carbon sequestration (Oliver et al., 2018). Control of anthropogenic CH₄ emissions leads to rapid decreases in its atmospheric concentration, with an approximately 9-year removal lifetime (and as such is an SLCF). Furthermore, many CH₄ mitigation options are inexpensive or even cost negative through the co-benefits achieved (Stohl et al., 2015), although expenditure becomes substantial at high levels of mitigation (Gernaat et al., 2015). The extra “allowable” carbon emissions from CH₄ mitigation can make a substantial difference to the feasibility or otherwise of achieving the Paris climate targets (Collins et al., 2018).

90 Some increases in atmospheric CH₄ are not related to direct anthropogenic activity, but indirectly to climate change triggering natural carbon and methane-climate feedbacks. These effects could act as positive feedbacks, and thus in the opposite direction to the mitigation of anthropogenic CH₄ sources. Wetlands are the largest natural source of CH₄ to the atmosphere and these emissions respond strongly to climate change (Melton et al., 2013; Gedney et al., 2019). A second natural feedback is from permafrost thaw. In a warming climate, the resulting microbial decomposition of previously frozen organic carbon is potentially one of the largest feedbacks from terrestrial ecosystems (Schuur et al., 2015). As the carbon and CH₄ climate feedbacks from natural wetlands and permafrost thaw could be substantial, this causes a reduction in anthropogenic CO₂ emission budgets compatible with climate change targets (Comyn-Platt et al., 2018; Gasser et al., 2018).

95 ~~For the first time, we combine these elements (land-based mitigation, anthropogenic CH₄ mitigation, and natural carbon and methane climate feedbacks) in a climate/Earth System modelling framework to quantify the unrealised potential from the mitigation of land-based options and anthropogenic CH₄ sources. In contrast to previous studies, we use a process-based land surface model to assess these mitigation options by region, yielding policy relevant information on the optimal mitigation strategy. This paper models the potential for mitigation of greenhouse gases to contribute to meeting the Paris targets of limiting global warming to 1.5°C and 2°C respectively. Specifically, we investigate the effectiveness of mitigation of anthropogenic methane emissions and land-based mitigation (e.g., implementation of BECCS and AR), combining results from three recent~~

Commented [HGD5]: Response 8

Commented [HGD6]: Response 1

105 papers (Collins et al., 2018;Comyn-Platt et al., 2018;Harper et al., 2018). We determine the effectiveness of these approaches in terms of their impact on the anthropogenic fossil fuel CO₂ emissions budget consistent with stabilising temperature at 1.5°C and 2° C of warming. The more effective the mitigation option, the larger the fossil fuel CO₂ emissions budget consistent with stabilisation at a given level. We estimate the impact of these mitigation scenarios relative to an existing scenario of greenhouse gas concentrations (based on the IMAGE SSP2 baseline), spanning uncertainties in both climate model projections (both global warming and regional climate change), process representation and the efficacy of BECCS. Sect. 2 provides a brief description of the models, the experimental set-up and the key datasets used in the model runs and subsequent analysis. Sect. 3 presents and discusses the results, starting with a global perspective before addressing the regional dimension. For BECCS, we additionally investigate the sensitivity to key assumptions and consider the implications for water security. Sect. 4 contains our conclusions.

2 Approach and Methodology

115 Figure 1 shows a schematic of our approach, the workflow and the prescribed data used. We use the Joint UK Land-Environment Simulator (JULES, Sect. 2.1) (Best et al., 2011;Clark et al., 2011), coupled with an inverted form of the “Integrated Model Of Global Effects of climatic aNomalies” (IMOGEN, Sect. 2.2) (Huntingford et al., 2010;Comyn-Platt et al., 2018). IMOGEN is an intermediate-complexity climate model, which uses “pattern-scaling” to emulate 34 models in the CMIP5 ensemble. In the inverted form used here, IMOGEN follows a prescribed temperature pathway (Sect. 2.2.2). We derive the overall radiative forcing consistent with this temperature pathway using an energy balance model, including a simplified model of ocean uptake (of energy and CO₂). Using a combination of calculated and prescribed time series of annual radiative forcings, we derive the atmospheric CO₂ radiative forcing and hence its concentration, taking account of any land and ocean feedbacks. For the mitigation scenarios considered (Sect 2.3), we use consistent and compatible time series of (a) anthropogenic CH₄ emissions, (b) prescribed land areas for crops and BECCS (where relevant) and (c) radiative forcings for SLCFs and non-CO₂ GHGs (except CH₄), from the IMAGE integrated assessment model. In a post-processing step (Sect. 2.4.1), we take the modelled carbon stores for land (=vegetation and soil carbon), atmosphere and oceans from the IMOGEN-JULES output and calculate the anthropogenic fossil fuel emission budgets (AFFEB) compatible with the warming pathway. For the land-based scenarios involving BECCS, we optimise the AFFEB by selecting the greater land carbon uptake from the ‘BECCS’ or the variant ‘natural’ (i.e., no BECCS) scenario, for those grid cells where BECCS is deployed (Sect. 2.4.2). Further, we investigate the sensitivity of the optimisation to the assumption made about BECCS productivity and carbon uptake (Sect. 2.4.3). Section 2.3.3 lists the model runs undertaken and the key assumptions and datasets used. Our overall modelling strategy is as follows. The starting point is the prescription of global temperature profiles that match the historical record, followed by a transition to a future stabilisation at either 1.5 or 2.0°C above pre-industrial levels. For these profiles, we then determine the related pathways in atmospheric radiative forcing by inversion of the global energy balance component of the IMOGEN impacts model. IMOGEN “Integrated Model Of Global Effects of climatic aNomalies” (Sect. 2.2) (Huntingford et

120
125
130
Commented [HGD7]: Response 1

al., 2010;Comyn-Platt et al., 2018) is an intermediate complexity climate model, which emulates 34 models in the CMIP5 climate model ensemble. Hence our radiative forcing (RF) trajectories have uncertainty bounds, reflecting the different thermal sensitivities of existing climate models.

For each radiative forcing pathway, we subtract the individual RF components for non-CO₂ and non-CH₄ radiatively-active gases that are perturbed by human activity, and for scenarios taken from the IMAGE integrated assessment model. Then, for CH₄, we use an understanding of its atmospheric lifetime to translate methane emissions into their atmospheric concentrations. The related RF for CH₄ is also subtracted from the overall value. Hence the remaining RF is that available for changes to atmospheric CO₂ concentration. The IMOGEN model uses pattern-scaling, again fitted to the same 34 climate models, to estimate local changes in near-surface meteorology. Combined with our global temperature pathways, these pattern-based changes (as well as atmospheric CO₂ concentration) drive the Joint UK Land-Environment Simulator land surface model (JULES, Sect. 2.1) (Best et al., 2011;Clark et al., 2011). JULES estimates atmosphere-land CO₂ exchange, and similarly, IMOGEN contains a single global description of oceanic CO₂ draw-down. These two estimates of carbon exchanges with the land and ocean respectively, in conjunction with atmospheric storage being linear in the CO₂ pathway, finally determine by simple summation compatible CO₂ emissions from fossil fuel burning. We call this the anthropogenic (CO₂) fossil fuel emission budgets (AFFEB) compatible with the warming pathway, subject to the assumptions made for non-CO₂ forcings.

Our numerical simulation structure allows us to investigate the implications of three different key changes on AFFEB, for stabilisation at both 1.5 and 2.0°C, and in a structure that captures features of a full set of climate models. First and maybe most importantly, we work to understand how regional reductions in CH₄ emissions allow higher values of AFFEB. Second, we consider how alternative scenarios of BECCS implementation alter atmosphere-land CO₂ exchanges, and again presented as the resultant implications for AFFEB. Third, we determine how the newer understanding of warming impacts on wetland methane emissions also affects AFFEB. Figure 1 captures the modelling framework, derivation of AFFEB, and our numerical experiments in a single overall schematic diagram.

Each of the scenarios investigated using the IMOGEN-JULES framework comprises 2 ensembles of 136 members, one ensemble for each of the warming targets. We make use of these ensembles to derive an “uncertainty” in the derived carbon budgets, specifically from climate change (as given by the 34 CMIP5 models) and from key land-surface processes (methane emissions from wetlands and the ozone vegetation damage). The climate change uncertainty comprises both the range of climate sensitivities of the CMIP5 models and the different regional patterns in the models. We use the median of the 136-member ensemble as the central value to derive the carbon budgets and the interquartile range (25-75%) for the uncertainty.

2.1 The JULES model

We use the JULES land surface model (Best et al., 2011;Clark et al., 2011), release version 4.8, but with a number of additions required specifically for our analysis:

Commented [HGD8]: Response 2 & 3

165 1. Land use: We adopt the approach used by Harper et al. (2018) and prescribe *managed* land-use and land-use change (LULUC). On land used for agriculture, C3 and C4 grasses are allowed to grow to represent crops and pasture. The land-use mask consists of an annual fraction of agricultural land in each grid cell. Historical LULUC is based on the HYDE 3.1 dataset (Klein Goldewijk et al., 2011), and future LULUC is based on two scenarios (SSP2 RCP-1.9 and SSP2 baseline), which were developed for use in the IMAGE integrated assessment model (IAM) (van Vuuren et al., 2017;Doelman et al., 2018) (see also Sect. 2.3).

170 Natural vegetation is represented by nine plant functional types (PFTs): broadleaf deciduous trees, tropical broadleaf evergreen trees, temperate broadleaf evergreen trees, needle-leaf deciduous trees, needle-leaf evergreen trees, C3 and C4 grasses, deciduous and evergreen shrubs (Harper et al., 2016). These PFTs are in competition for space in the non-agricultural fraction of grid cells, based on the TRIFFID (Top-down Representation of Interactive Foliage and Flora Including Dynamics) dynamic vegetation module within JULES (Clark et al., 2011). A further four PFTs are used to represent agriculture (C3 and C4 crops, and C3 and C4 pasture), and harvest is calculated separately for food and bioenergy crops (see Sect. 2.4.3, where we describe the modelling of carbon removed via bioenergy with CCS). When natural vegetation is converted to managed agricultural land, the vegetation carbon removed is placed into woody product pools that decay at various rates back into the atmosphere (Jones et al., 2011). Hence, the carbon flux from LULUC is not lost from the system. There are also four non-vegetated surface types: urban, water, bare soil and ice.

180 2. Soil carbon: Following Comyn-Platt et al. (2018), we also use a 14 layered soil column for both hydro-thermal (Chadburn et al., 2015) and carbon dynamics (Burke et al., 2017a). Burke et al. (2017b) demonstrated that modelling the soil carbon fluxes as a multi-layered scheme improves estimates of soil carbon stocks and net ecosystem exchange. In addition to the vertically discretised respiration and litter input terms, the soil-carbon balance calculation also includes a diffusivity term to represent cryoturbation/bioturbation processes. The freeze-thaw process of cryoturbation is particularly important in cold permafrost-type soils (Burke et al., 2017b). Following Burke et al. (2017a), we diagnose permafrost wherever the deepest soil layer is below 0°C (assuming that this layer is below the depth of zero annual amplitude, i.e., where seasonal changes in ground temperature are negligible (<0.1 °C)). Further, for permafrost regions, there is an additional variable to trace or diagnose “old” carbon and its release from permafrost as it thaws.

190 The multi-layered methanogenesis scheme improves the representation of high latitude CH₄ emissions, where previous studies underestimated production at cold permafrost sites during “shoulder seasons” (Zona et al., 2016). Figure 2 shows the annual cycle in the observed and modelled wetland CH₄ emissions at the Samoylov Island field site (panel a) and a comparison of observed and modelled annual mean fluxes at this and other sites (panel b). The range of uncertainty used in our study (JULES low Q₁₀ - JULES high Q₁₀) captures the range of uncertainty in the observations (In Fig. 2b, the error bars denote the lower and upper estimates from the low and high Q₁₀ simulations. The symbols represent the mean value between these estimates). Further, the layered methane scheme

Commented [HGD9]: Response 9

used in this work gives a better description of the shoulder season emissions when compared with the original, non-layered methane scheme in JULES. The multi-layered scheme allows an insulated sub-surface layer of active methanogenesis to continue after the surface has frozen. These model developments not only improve the seasonality of the emissions, but more importantly for this study capture the release of carbon as CH₄ from deep soil layers, including thawed permafrost. Further evaluation of the multi-layer scheme can be found in Chadburn et al. (2020).

3. Methane from wetlands: Following Comyn-Platt et al. (2018), we also use the multi-layered soil carbon scheme described in (2) above to give the local land-atmosphere CH₄ flux, E_{CH_4} (kg C m⁻² s⁻¹):

$$E_{CH_4} = k \cdot f_{wet} \cdot \sum_{i=1}^n C_s^{pools} \kappa_i \cdot \sum_{z=0m}^{3m} e^{-\gamma z} C_{s,i,z} \cdot Q_{10}(T_{soil,z})^{0.1(T_{soil,z}-T_0)} \quad (1)$$

where k is a dimensionless scaling constant such that the global annual wetland CH₄ emissions are 180 Tg CH₄ in 2000 (as described in Comyn-Platt et al. (2018)), z is the depth in soil column (in m), i is the soil carbon pool, f_{wet} (-) is the fraction of wetland area in the grid cell, κ_i (s⁻¹) is the specific respiration rate of each pool (Table 8 of Clark et al. (2011)), C_s (kg m⁻²) is soil carbon, T_{soil} (K) is the soil temperature. The decay constant γ (= 0.4 m⁻¹) describes the reduced contribution of CH₄ emission at deeper soil layers due to inhibited transport and increased oxidation through overlying soil layers. This representation of inhibition and of the pathways for CH₄ release to the atmosphere (e.g., by diffusion, ebullition and vascular transport) is a simplification. However, previous work which explicitly represented these processes showed little to no improvement when compared with in-situ observations (McNorton et al., 2016). We do not model CH₄ emissions from freshwater lakes (and oceans).

Comyn-Platt et al. (2018) varied Q_{10} in Eq. 1 to encapsulate a range of methanogenesis process uncertainty. They derive Q_{10} values for each GCM configuration to represent two wetland types identified in Turetsky et al. (2014) ('poor-fen' and 'rich-fen'). They also include a third 'low- Q_{10} ', which gives increased importance to high latitude emissions. Their ensemble spread was able to describe the magnitude and distribution of present-day CH₄ emissions from natural wetlands, according to the models used in the then-current global methane assessment (Saunois et al., 2016). Here, we use the 'low- Q_{10} ' value of Comyn-Platt et al. (2018) (=2.0) and adopt a 'high- Q_{10} ' value of ~4.8 from the rich-fen parameterisation. The two Q_{10} values used here still capture the full range of the methanogenesis process uncertainty.

4. Ozone vegetation damage: We use a JULES configuration including ozone deposition damage to plant stomata, which affects land-atmosphere CO₂ exchange (Sitch et al., 2007). JULES requires surface atmospheric ozone concentrations, O₃ (ppb), for the duration of the simulation period (1850-2100). As in Collins et al. (2018), we do not model tropospheric ozone production from CH₄ explicitly in IMOGEN. Instead, we use two sets of monthly near-surface O₃ concentration fields (January-December) from HADGEM3-A GA4.0 model runs, with the sets corresponding to low (1285 ppbv) and high (2062 ppbv) global mean atmospheric CH₄ concentrations (Stohl et al., 2015). We assume that the atmospheric O₃ concentration in each grid cell responds linearly to the atmospheric CH₄ concentration. We derive

230 separate linear relationships for each month and grid cell, and use these to calculate the surface O₃ concentration from the corresponding global atmospheric CH₄ concentration as it evolves during the IMOGEN run (Sect. 2.2.1). We use the CH₄ concentration profiles from the ~~prescribed-IMAGE~~ SSP2-Baseline and RCP-1.9-IMAGE scenarios (Sect. 2.3.1), adjusted for natural methane sources (see 3 above and Sect. 2.3.3). We undertake runs using both the ‘high’ and ‘low’ vegetation ozone-damage parameter sets (Sitch et al., 2007).

2.2 The IMOGEN intermediate complexity climate model

2.2.1 IMOGEN

235 The IMOGEN climate impacts model (Huntingford et al., 2010) uses “pattern-scaling” to estimate changes to the seven meteorological variables required to drive JULES. Huntingford et al. (2010) assume that changes in local temperature, precipitation, humidity, wind-speed, surface shortwave and longwave radiation and pressure are linear in global warming. Spatial patterns of each variable (based on the 34 GCM simulations in CMIP5 (Comyn-Platt et al., 2018)) are multiplied by the amount of global warming over land, ΔT_L , to give local monthly predictions of climate change. When using IMOGEN in forward mode, ΔT_L is calculated with an Energy Balance Model (EBM) as a function of the overall changes in radiative forcing, ΔQ (W m⁻²). ΔQ is the sum of the atmospheric greenhouse gas contributions (Eq. 2) (Etminan et al., 2016), which in the forward mode are either calculated (CO₂ and CH₄) or prescribed (for other atmospheric contributors) on a yearly time step.

$$\Delta Q(\text{total}) = \Delta Q(\text{CO}_2) + \Delta Q(\text{non CO}_2 \text{ GHGs}) + \Delta Q(\text{aerosols and other climate forcers}) \quad (2)$$

245 The EBM includes a simple representation of the ocean uptake of heat and CO₂ and uses a separate set of four parameters for each climate and Earth system model emulated (Huntingford et al., 2010): the climate feedback parameters over land and ocean, λ_l and λ_o (W m⁻² K⁻¹) respectively, the oceanic “effective thermal diffusivity”, κ (W m⁻¹ K⁻¹) representing the ocean thermal inertia and a land-sea temperature contrast parameter, ν , linearly relating warming over land, ΔT_l (K) to warming over ocean, ΔT_o (K), as $\Delta T_l = \nu \Delta T_o$. The climate feedback parameters (λ_l and λ_o) are calibrated using model-specific data for the top of the atmosphere radiative fluxes, the mean land and ocean surface temperatures, along with an estimate of the radiative forcing modelled for the CO₂ changes.

255 ~~Our simulations include a CH₄ feedback system that captures the climate impacts on CH₄ emissions from natural wetland sources. The approach used here follows Comyn-Platt et al. (2018) and Gedney et al. (2019), where the prescribed atmospheric CH₄ concentrations, which assume a constant annual wetland CH₄ emission (van Vuuren et al., 2017), are modified using the anomaly in the modelled annual wetland CH₄ emission. The increased/reduced atmospheric CH₄ concentration will have a corresponding faster/slower atmospheric decay rate than the prescribed concentration pathway. We account for this following the approach of Cubasch et al. (2001). Related changes in atmospheric radiative forcing, in response to altered atmospheric CH₄ concentrations, are calculated using the formulation from Etminan et al. (2016). We also include the indirect effect of these CH₄ emission changes on the forcing by tropospheric ozone and stratospheric water vapour by multiplying the CH₄~~

260 forcing by 1.65, based on Myhre et al. (2013). The atmospheric CH₄ concentrations available from the IMAGE database (see Sect. 2.3.1) assume a constant annual wetland CH₄ emission (van Vuuren et al., 2017). However, these emissions have interannual variability and a positive climate feedback (Comyn-Platt et al., 2018; Gedney et al., 2019), and their correct representation is a central part of our study. We follow the same approach that we used in our previous studies (Collins et al., 2018; Comyn-Platt et al., 2018; Gedney et al., 2019). As the IMOGEN-JULES modelling framework does not have an explicit representation of the atmospheric chemistry of methane, we represent the oxidation and hence loss of CH₄ by a single lifetime (τ).

$$\frac{d([CH_4] - [CH_4]_{IMAGE})}{dt} = C \{ \sum F [CH_4] - \sum F [CH_4]_{IMAGE} \} - \frac{[CH_4] - [CH_4]_{REF}}{\tau} \quad (3)$$

270 where [CH₄] and [CH₄]_{IMAGE} are the atmospheric methane concentrations using our new wetland-based, time varying (F[CH₄]) and the constant IMAGE (F[CH₄]_{IMAGE}) wetland emissions, respectively. Parameter C is a constant to convert from Tg CH₄ to a mixing ratio in parts per billion by volume (ppbv). Further, higher atmospheric concentrations of CH₄ and its oxidation product (carbon monoxide) lower the concentration of hydroxyl radicals, the major removal reaction for CH₄, thereby increasing the atmospheric lifetime of CH₄. Conversely, lower CH₄ concentrations will shorten its atmospheric lifetime. We take account of this feedback of CH₄ on its lifetime (τ), using Eq. 4 (Collins et al., 2018; Comyn-Platt et al., 2018; Gedney et al., 2019), as,

$$\ln(\tau/\tau_0) = s \cdot \ln([CH_4]/[CH_4]_0), \text{ i.e., } \tau = \tau_0 \cdot \exp(s [CH_4]/[CH_4]_0) \quad (4)$$

280 In Eq. 4, [CH₄]₀ and τ₀ are the contemporary atmospheric CH₄ concentration and lifetime, and s is the CH₄-OH feedback factor, defined by $s = \partial \ln(\tau) / \partial \ln(CH_4)$. We take values of τ₀ = 8.4 years, [CH₄]₀ = 1.745 ppbv and s = 0.28 from Ehhalt et al. (2001) (pages 248 and 250). In our earlier study on the climate-wetland methane feedback Gedney et al. (2019), we investigate the sensitivity to the methane lifetime and the feedback factor, in addition to an analysis of the main drivers on the wetland methane-climate feedback and the main sources of uncertainty. Gedney et al. (2019) conclude that the limited knowledge of contemporary global wetland emissions is a larger source of uncertainty than that from the projected climate spread of the 34 GCMs. We quantify this uncertainty in our experimental design by using two values of Q₁₀ (see Sect. 2.1).

285 In response to our dynamic interactive calculations of atmospheric CH₄ concentrations, we derive the related change in methane radiative forcing (RF). We use the formulation from Etmann et al. (2016), which accounts for the short-wave absorption by CH₄ and the overlap with N₂O. The atmospheric oxidation of methane (by the hydroxyl radical) leads to the production of tropospheric ozone and stratospheric water vapour. We calculate these indirect contributions of methane to the overall radiative forcing, following the approach for methane adopted in our previous work (Collins et al., 2018; Comyn-Platt et al., 2018; Gedney et al., 2019). Collins et al. (2018) represent the forcing contributions from O₃ and stratospheric water vapour as linear functions of the CH₄ mixing ratio, based on the analysis presented in IPCC AR5 Myhre et al. (2013). The indirect methane forcings amount to $2.36 \times 10^{-4} \pm 1.09 \times 10^{-4}$ W m⁻² per ppb CH₄ (i.e., 0.65 ± 0.3 times the CH₄ radiative

efficiency). Hence we incorporate the indirect effects of these CH₄ emission changes by an approximation, multiplying the CH₄ radiative forcing by 1.65.

In this study, we use the inverse version of IMOGEN, which follows prescribed temperature pathways (Fig. 3(a)), to derive the total radiative forcing (ΔQ [total]) and then the CO₂ radiative forcing (ΔQ [CO₂]), using Eq. 2. Comyn-Platt et al. (2018) describe the changes made to the EBM to create the inverse version. As each of the 34 GCMs that IMOGEN emulates has a different set of EBM parameters, each GCM has a different time-evolving radiative forcing (ΔQ) estimate for a given temperature pathway, $\Delta T_G(t)$. When IMOGEN is forced with a historical record of ΔT_G , the range of ΔQ for the near present day (year 2015) from the 34 GCMs is 1.13 W m⁻². To ensure a smooth transition to the modelled future, we require the historical period, 1850-2015, to match observations of both ΔT_G and atmospheric composition for all GCMs. As we have a model-specific estimate of the radiative forcing modelled for the CO₂ changes (see above), we, therefore, attribute the spread in ΔQ to the uncertainty in the non-CO₂ radiative forcing component, particularly the atmospheric aerosol contribution, which has an uncertainty range of -0.5 to -4 Wm⁻² (Stocker et al., 2013). Apart from our modelled CH₄ and CO₂ radiative forcings and the potential future balances between them, we use the projections from the IMAGE SSP2 baseline or RCP1.9 scenario for the radiative forcing of other atmospheric contributors (Fig. 3(b)).

2.2.2 Temperature Profile Formulation

Huntingford et al. (2017) define a framework to create trajectories of global temperature increase, based on two parameters, and which model the efforts of humanity to limit emissions of greenhouse gases and short-lived climate forcers, and, if necessary, capture atmospheric carbon. These profiles have the mathematical form of:

$$\Delta T(t) = \Delta T_0 + \gamma t - (1 - e^{-\mu(t)t})[\gamma t - (\Delta T_{Lim} - \Delta T_0)] \quad (35)$$

where $\Delta T(t)$ is the change in temperature from pre-industrial levels at year t , ΔT_0 is the temperature change at a given initial point (in this case $\Delta T_0 = 0.89^\circ\text{C}$ for 2015), ΔT_{Lim} is the final prescribed warming limit and

$$\mu(t) = \mu_0 + \mu_1 t \quad \text{and} \quad \gamma = \beta - \mu_0(\Delta T_{Lim} - \Delta T_0) \quad (46)$$

where β (= 0.00128) is the current rate of warming and μ_0 and μ_1 are tuning parameters which describe anthropogenic attempts to stabilise global temperatures (Huntingford et al., 2017). The parameter values used for the two profiles are: (a) 1.5°C profile: $\Delta T_{lim} = 1.5^\circ\text{C}$; $\mu_0 = 0.1$ and $\mu_1 = 0.0$; (b) 2°C profile: $\Delta T_{lim} = 2^\circ\text{C}$; $\mu_0 = 0.08$ and $\mu_1 = 0.0$.

2.3 Scenarios and model runs

We undertake a control run and other simulations with anthropogenic CH₄ mitigation or land-based mitigation, stabilising at either 1.5°C or 2.0°C warming without a temperature overshoot. We denote the control run as “CTL”, the anthropogenic CH₄ mitigation scenario, a land-based mitigation scenario using BECCS and a variant land-based scenario focussing on AR, as “CH₄”, as-“BECCS”, “Natural”, respectively. We also undertake runs combining the CH₄ and land-based mitigation

scenarios (coupled “BECCS+CH₄” and coupled “Natural+CH₄”) to determine if there are any non-linearities when we combine these mitigation scenarios. We summarise the key assumptions of these scenarios in [Table 1](#).

We use future projections of atmospheric CH₄ concentrations and LULUC (specifically, the areas assigned to agriculture and within that to BECCS) from the IMAGE SSP2 projections (Doelman et al., 2018) as input or prescribed data for both the methane and land-based mitigation strategies ([Table 1](#)). This ensures that all projections are consistent and based on the same set of IAM model and socio-economic pathway assumptions. The SSP2 socio-economic pathway is described as “middle of the road” (O’Neill et al., 2017), with social, economic, and technological trends largely following historical patterns observed over the past century. Global population growth is moderate and levels off in the second half of the century. The intensity of resource and energy use declines. We define the upper and lower limits of anthropogenic mitigation as the lowest (RCP1.9, denoted “IM-1.9”) and highest (“baseline”, denoted “IM-BL”) total radiative forcing pathways, respectively, within the IMAGE SSP2 ensemble (Riahi et al., 2017). [As described in Section 2.2.1, we modify the atmospheric concentrations of CH₄ in the IMOGEN-JULES modelling as the IMAGE scenarios assume constant natural and hence wetland methane emissions.](#)

Commented [HGD11]: Response 11

2.3.1 Methane: baseline and mitigation scenario

The anthropogenic CH₄ emission increase from 318 Tg yr⁻¹ in 2005 to 484 Tg yr⁻¹ in 2100 in the IMAGE SSP2 baseline scenario, but fall to 162 Tg yr⁻¹ in 2100 in the IMAGE SSP2 RCP1.9 scenario. The sectoral CH₄ emissions in 2005 (Energy Supply & Demand: 113; Agriculture: 136; Other Land Use (primarily burning): 18; Waste 52, all in Tg yr⁻¹) are in agreement with the latest estimates of the global methane cycle (Saunois et al., 2019). As summarised in Supplementary Information, Table SI.1, the reduction in CH₄ emissions from specific source sectors is achieved as follows: (a) [coal production](#) by maximising CH₄ recovery from underground mining of hard coal; (b) [oil/gas production & distribution](#), through control of fugitive emissions from equipment and pipeline leaks, and from venting during maintenance and repair; (c) [enteric fermentation](#), through change in animal diet and the use of more productive animal types; (d) [animal waste](#) by capture and use of the CH₄ emissions in anaerobic digesters; (e) [wetland rice production](#), through changes to the water management regime and to the soils to reduce methanogenesis; (f) [landfills](#) by reducing the amount of organic material deposited and by capture of any CH₄ released; (g) [sewage and wastewater](#), through using more wastewater treatment plants and also recovery of the CH₄ from such plants, and through more aerobic wastewater treatment. The levels of reduction vary between sectors, from 50% (agriculture) to 90% (fossil-fuel extraction and delivery). The abatement costs are between US\$ 300-1000 (1995 US\$) (Supplementary Information, Table SI.1). [Figure 4](#) presents the IMAGE baseline and RCP1.9 CH₄ emission pathways globally and for selected IMAGE regions, including the major-emitting regions of India, USA and China (Supplementary Information, Figure SI.1 shows the emission pathways for all 26 IMAGE regions). These two methane emission pathways ([IMAGE SSP2 baseline and RCP1.9](#)) define our “CTL” and “CH₄” scenarios, respectively.

2.3.2 Land-based mitigation: baseline, BECCS and Natural scenarios

355 For our land-based mitigation scenarios, we take time series of the annual areas assigned to agriculture (crops and pasture) and within that, the area allocated to bioenergy crops, from the IM-BL and IM-1.9 scenarios (defined at the start of Sect. 2.3). We use the dynamic vegetation module in JULES to calculate the evolution of the natural plant functional types and the non-vegetated surface on the remaining land area in the grid cell (see Land use in Sect. 2.1).

Commented [HGD12]: Response 18

360 The IM-BL LULUC scenario assumes (a) moderate land-use change regulation; (b) moderately effective land-based mitigation; (c) the current preference for animal products; (d) moderate improvement in livestock efficiencies; and (e) moderate improvement in crop yields (Table 1 in (Doelman et al., 2018)). It represents a control scenario within which agricultural land is accrued to feed growing populations associated with the SSP2 pathway and with no deployment of BECCS. Three types of land-based climate change mitigation are implemented in the IMAGE land use mitigation scenarios (Doelman et al., 2018): (1) bioenergy; (2) reducing emissions from deforestation and degradation (REDD or avoided deforestation); and (3) reforestation of degraded forest areas. For the IM-1.9 scenario, there are high levels of REDD and full reforestation. The scenario assumes a food-first policy (Daioglou et al., 2019) so that bioenergy crops are only implemented on land not required for food production (e.g., abandoned agricultural crop land, most notably, in central Europe, southern China and eastern USA, and on natural grasslands in central Brazil, eastern and southern Africa, and Northern Australia (Doelman et al., 2018)). The IM-1.9 scenario also requires bioenergy crops to replace forests in temperate and boreal regions (notably Canada and Russia). The demand for bioenergy is linked to the carbon price required to reach the mitigation target (Hoogwijk et al., 2009). In this scenario, the area of land used for bioenergy crops expands rapidly from 2030 to 2050, reaching a maximum of 550 Mha in 370 2060, and then declining to 430 Mha by 2100. Table 2 gives the maximum area of BECCS deployed in each IMAGE region for the IM-1.9 scenario. This defines the land use in the “BECCS” scenario.

375 We define a third LULUC pathway, which is identical to the “BECCS” scenario, except that any land allocated to bioenergy crops is allocated instead to natural vegetation, i.e., areas of natural land, which are converted to bioenergy crops, remain as natural vegetation, and areas, which are converted from food crops or pasture to bioenergy crops, return to natural vegetation. We make no allowance for any changes in the energy generation system, as this would require energy sector modelling that is beyond the scope of this study. We denote this scenario as “Natural”. Table 2 also summarises the main differences in land use between the BECCS and Natural scenarios for each IMAGE region.

380 Figure 5 presents time series of the land areas calculated for trees and prescribed for agriculture (including bioenergy crops) and bioenergy crops for the “BECCS” and “Natural” scenarios for the Russia and Brazil IMAGE regions, each as a difference to the baseline scenario (IM-BL). Supplementary Information, Figure SI.2 is equivalent to Fig. 5 for all the IMAGE regions.

2.3.3 Model runs

For each temperature pathway (1.5°C or 2.0°C) and for the baseline and each mitigation scenario, the set of factorial scenario runs comprises a 136-member ensemble (34 GCMs x 2 ozone damage sensitivities x 2 methanogenesis Q₁₀

385 temperature sensitivities). In all model runs, we include the effects of the methane and carbon-climate feedbacks from wetlands and permafrost thaw, which we have shown previously to be significant constraints on the AFFEBs (Comyn-Platt et al., 2018).

As shown in Fig. 1, we use a number of input or prescribed datasets: (a) time series of the annual area of land used for agriculture, including that for BECCS if appropriate; (b) time series of the global annual mean atmospheric concentrations of CH₄ (and N₂O for the radiative forcing calculations of CO₂ and CH₄); (c) time series of the overall radiative forcing by SLCFs and non-CO₂ GHGs (corrected for the radiative forcing of CH₄); and (d) time series of annual anthropogenic CH₄ emissions (used in the post-processing step). We take these from the IMAGE database for the relevant IMAGE SSP scenario (baseline or SSP2-1.9). Table 1 lists the main factorial-scenario runs, their key features and the prescribed datasets used (for agricultural land and BECCS, anthropogenic emissions and atmospheric concentrations of CH₄ and the non-CO₂ radiative forcing).

395 Figure 6 presents the effect of these scenarios on the modelled atmospheric CH₄ and CO₂ concentrations. We adjust the input atmospheric CH₄ concentrations to allow for the interannual variability in the wetland CH₄ emissions, as described in Sect. 2.2.1. The As we use the same input datasets for the two warming targets, the major control on the modelled atmospheric CH₄ concentrations is the CH₄ emission pathway followed, with the temperature pathway (1.5° versus 2°C warming) having a minor effect. For CO₂, on the other hand, the temperature and the CH₄ emission pathways both lead to increased atmospheric CO₂ concentrations, with the temperature pathway having a slightly larger effect.

2.4 Post-processing

2.4.1 Anthropogenic Fossil Fuel Emission Budget and Mitigation Potential

Following Comyn-Platt et al. (2018), we define the anthropogenic fossil fuel CO₂ emission budget (AFFEB) for scenario *i* as the change in carbon stores from present to the year 2100:

$$405 \quad AFFEB_i = [C^{land}(2100) - C^{land}(2015)]_i + [C^{ocean}(2100) - C^{ocean}(2015)]_i \\ + [C^{atmos}(2100) - C^{atmos}(2015)]_i + BECCS(2015:2100)_i \quad (57)$$

where $C^{land}(t)$, $C^{ocean}(t)$ and $C^{atmos}(t)$ are the carbon stored in the land, ocean and atmosphere, respectively, in year *t* and $BECCS(t_1:t_2)$ is the carbon sequestered via BECCS between the years *t*₁ and *t*₂. The atmospheric carbon store does not include CH₄. This is a reasonable approximation, however, given the relative magnitudes of the atmospheric concentrations of CH₄ (~2 ppmv at the surface) and CO₂ (400 ppmv).

410 Within the IMOGEN-JULES modelling framework, we use (a) the IMOGEN climate emulator to derive the changes in the ocean and atmosphere carbon stores, and (b) JULES for the changes in the land carbon store and carbon sequestered through BECCS. We discuss the changes in the carbon stores for the baseline and different mitigation scenarios in Sect. 3.1.

Commented [HGD13]: Response 13

Commented [HGD14]: Response 14

Commented [HGD15]: Response 15

For brevity in the subsequent discussion, we use the following shorthand where the terms on the RHS of Eq. 5-7 are equivalent to those on the RHS of Eq. 68:

$$AFFEB_i = \Delta C_i^{land} + \Delta C_i^{ocean} + \Delta C_i^{atmos} + BECCS_i \quad (68)$$

We define the mitigation potential (MP) for a mitigation strategy, j , as the difference between a control AFFEB ($AFFEB_{ctl}$) and the AFFEB resulting from applying the strategy i.e.:

$$MP_j = AFFEB_j - AFFEB_{ctl} \quad (79)$$

which can be broken down into its component parts as:

$$MP_j = MP_j^{land} + MP_j^{ocean} + MP_j^{atmos}$$

$$MP_j = (\Delta C_j^{land} - \Delta C_{ctl}^{land}) + (\Delta C_j^{ocean} - \Delta C_{ctl}^{ocean}) + (\Delta C_j^{atmos} - \Delta C_{ctl}^{atmos}) + BECCS_j \quad (810)$$

2.4.2 Optimisation of the land-based mitigation

Harper et al. (2018) find that the land-use pathways do not provide a clear choice for the preferred mitigation pathway. The key issue is that replacing natural vegetation with bioenergy crops often results in large emissions of soil carbon and the loss of the benefits of maintaining forest carbon stocks. In such circumstances, Harper et al. (2018) find that the loss of soil carbon in regions with high carbon density makes it difficult for BECCS to deliver a net negative emission of CO₂. Hence, to optimise the land-based mitigation (LBM), we compare the land-carbon stocks in the BECCS and Natural scenarios. We then select the optimum land-management option for each grid cell simulated as that, which maximises the $AFFEB$ by year 2100. That is:

$$AFFEB_{LBM} = \Delta DC_{\#M1 \rightarrow \#BECCS}^{atmos} + \Delta DC_{\#M1 \rightarrow \#BECCS}^{ocean} + \Delta DC_{LBM}^{land} \quad (911)$$

with

$$\Delta DC_{LBM}^{land} = \begin{cases} \sum_l^{grid\ cells} \Delta C_{\#M1 \rightarrow \#BECCS}^{land} + BECCS_{\#M1,9} & \text{where } \Delta C_{\#M1 \rightarrow \#BECCS}^{land} < \Delta C_{\#M1 \rightarrow \#BECCS}^{land} + BECCS_{\#M1,9} \\ \text{or} \\ \sum_l^{grid\ cells} \Delta C_{\#M1 \rightarrow \#Natural}^{land} & \text{where } \Delta C_{\#M1 \rightarrow \#Natural}^{land} > \Delta C_{\#M1 \rightarrow \#BECCS}^{land} + BECCS_{\#M1,9} \end{cases} \quad (1012)$$

where $\Delta C_{\#pathways\ scenario}^{store}$ is the change in carbon between 2015 and 2100 for the ‘store’ (= atmosphere, ocean or land) for the LULUC $\#pathways\ scenario$. We use the ocean and atmosphere contributions from the BECCS simulations as the changes in store size between the BECCS and Natural simulations are negligible (i.e. <2GtC).

2.4.3 Assumptions about BECCS efficiency

The efficacy of the BECCS scheme implemented in JULES is significantly lower than that of other implementations (Harper et al., 2018), reflecting the importance of assumptions about the efficiency of the BECCS process and bioenergy crop yields in determining their ability to contribute to climate mitigation. More specifically, there is (1) large uncertainty in carbon losses from farm to final storage (Harper et al. (2018) assumed a 40% loss compared to 13-52% loss found in other studies); and (2) a large range in potential productivity of second-generation lignocellulosic bioenergy crops, with JULES falling on the low end. JULES in this study and in Harper et al. (2018) simulated median average yields of ~4.8 and ~4.6 tDM ha⁻¹ yr⁻¹, respectively, compared to measured median of 11.5 tDM ha⁻¹ yr⁻¹ and simulated average of 15.8 tDM ha⁻¹ yr⁻¹ in IMAGE. The JULES yield of ~4.8 tDM ha⁻¹ yr⁻¹ corresponds to ~59 EJ yr⁻¹ of primary energy, using the maximum area for BECCS from Table 2 of 637.7 Mha and an energy yield of 19.5 GJ t DM⁻¹ (Daioglou et al., 2017). Bioenergy supplied 55.6 EJ yr⁻¹ or ~10% of primary energy requirement worldwide in 2017 (WBA, 2019). According to Smith et al. (2016), this would increase to ~170 EJ yr⁻¹ of primary energy in 2100, for negative emissions of 3.3 Gt Ceq yr⁻¹ from BECCS (as required for a 2°C warming target).

As both of these components are assumed to be diagnostics of the simulations, we can modify the contribution of BECCS to the AFFEB via a post-processing scaling factor, κ , which represents the efficiency of (1) and (2) with respect to the JULES parameterisation. That is, Eq. 10-12 becomes:

$$\Delta C_{LBM}^{land} = \begin{cases} \sum_l^{grid\ cells} \Delta C_{IM1.9BECCS}^{land} + \kappa BECCS_{IM1.9} & \text{where } \Delta C_{IM1.9Natural}^{land} < \Delta C_{IM1.9BECCS}^{land} + \kappa BECCS_{IM1.9} \\ \sum_l^{grid\ cells} \Delta C_{IM1.9Natural}^{land} & \text{where } \Delta C_{IM1.9Natural}^{land} > \Delta C_{IM1.9BECCS}^{land} + \kappa BECCS_{IM1.9} \end{cases} \quad (113)$$

Figure 7 presents maps of the scaling factor required for BECCS to be the preferable mitigation option, as opposed to natural land carbon uptake, for each grid cell for warming of 1.5°C or 2°C. There are large factors in the northern temperate and boreal regions, parts of Africa and Australia. As discussed in Harper et al. (2018), this follows from the loss of soil carbon in the tropics and at high northern latitude leading to long recovery or payback times (10-100+ years and >100 years, respectively, Fig. 6(c) in their paper). The payback time is however insignificant when bioenergy crops replace existing agriculture, for example in Europe and eastern North America.

Additionally, we define a threshold efficiency factor, κ^* , which represents the required BECCS efficiency for BECCS to be a preferable mitigation strategy for a given grid-cell, i.e.:

$$\kappa^* = \frac{\Delta C_{IM1.9Natural}^{land} - \Delta C_{IM1.9BECCS}^{land}}{BECCS_{IM1.9}} \quad (114)$$

This increased efficiency can be considered to be the additional bioenergy harvest (H) and/or the reduced carbon losses from farm to storage needed to pay back the carbon debt accrued due to land-use change (since carbon removed via BECCS

= ϵH , where ϵ is the assumed efficiency factor for farm to storage carbon conservation and H is the simulated biomass harvest). In addition, κ^* implies a new threshold (or break-even) level of BECCS:

$$BECCS^* = \kappa \kappa^* * BECCS_{IM1.9} \quad (1315)$$

In other words, $BECCS^*$ is equivalent to the carbon loss due to the land use change to grow the bioenergy crops. Our IMOGEN-JULES simulations assume a 40% carbon loss from farm to final storage, although other studies have assumed this to be as low as 13% (Harper et al., 2018). To assess the feasibility of meeting this break-even level of BECCS, we calculate the harvest (H^*) that would be needed if carbon losses are to be minimised, i.e. by increasing ϵ from 0.6 to 0.87, and assuming in Eq. 13-15 that:

$$BECCS^* = 0.87 H^* \text{ and } BECCS_{IM1.9} = 0.60 H$$

So:

$$H^* = \kappa \kappa^* * \frac{0.6}{0.87} * H \quad (1416)$$

We discuss this further in Sect. 3.2.

3 Results and Discussion

3.1 Global Perspective

We calculate the anthropogenic fossil fuel emission budget to limit global warming to a particular temperature target as the sum of the changes in the carbon stores of the atmosphere, land (vegetation and soil) and ocean between 2015 and 2100 (Sect. 2.4.1, Eq. 5-7 and 68). We use a BECCS scale factor (κ) of unity. We present in Fig. 8, we present the median and ensemble member spread of the AFFEB (as box and whiskers) from the 136-member ensemble, and the individual GCM/ESM contributions to the AFFEBs from the four carbon pools shown (points), for each of the factorial experiments main scenarios modelled using the IMOGEN-JULES or derived in the post-processing optimisation step (see Table 1 Table 1 for description of the scenarios-).

We find that there is increased uptake of atmospheric CO_2 in the land-based mitigation scenarios, although there is a reduction in land carbon from the land use changes in these scenarios. In all the scenarios apart from the BECCS scenario, there is an increase in the land carbon store (shown as positive changes for Coupled (Natural) and Coupled (Optimised) but as smaller negative changes for “ CH_4 ”, “Natural” and “Optimised” scenarios. In the “BECCS” scenario, the land carbon change becomes more negative than in the “CTL” scenario, as bioenergy crops replace ecosystems with higher carbon content. In the combined (‘coupled’) CH_4 and land-based mitigation scenarios, the reduction in the emissions and hence atmospheric concentrations of CH_4 allow increased atmospheric concentrations of CO_2 (Fig. 6). There is increased uptake of carbon by the

Commented [HGD16]: Response 16

Commented [HGD17]: Response 5

Commented [HGD18]: Responses 17 & 18

land, directly because of the increased atmospheric CO₂ concentration and indirectly through the reduction in O₃ damage. In the coupled “BECCS” scenario, this increased uptake of atmospheric CO₂ is again offset by, which is greater than the land carbon lost through land-use changes conversion of the land to bioenergy crops. We also find that there is increased uptake of CO₂ by the oceans for all scenarios. A further co-benefit of reducing the CH₄ emissions and allowing more CO₂ emissions is that the oceanic drawdown of CO₂ rises (although it eventually falls to zero under climate stabilisation and there would also be implications for ocean acidification). In Fig. 9(a), we compare the AFFEBs for both the 1.5°C and 2°C temperature pathways. We find that the absolute AFFEBs are 200-300 GtC larger for the 2°C target than the 1.5°C target. These budgets are in agreement with other estimates, which include corrections to the historical period (Millar et al., 2017). In both Figs. 8 and 9, it should be noted that the gain in the land carbon store for the “CH₄” mitigation option at -1.4 GtC (median of ensemble) is not visible in these figures. There has however been a net increase in the land carbon store in the “CH₄” scenario when compared to the land carbon store in the control scenario (shown as a reduction from -70.8 GtC, loss of land carbon in the control run to -1.4 GtC loss in the methane mitigation option (median of ensemble). This then explains the positive changes shown for the land carbon stores in the coupled “BECCS+ CH₄” and coupled “Natural+ CH₄” scenarios.

Commented [HGD19]: Responses 17 & 18

Figure 9(b) shows the mitigation potential of each strategy, calculated as the change in the AFFEB from the corresponding control simulation, for the two temperature pathways (Sect. 2.4.1, Eq. 7-9 and 810). Methane mitigation is a highly effective strategy; the AFFEBs are increased by 188-206 GtC and 193-212 GtC for the 1.5°C and 2°C scenarios, respectively, where the range represents the interquartile range from the 136-member ensemble (34 GCMs x 2 Q₁₀ x 2 ozone sensitivities). This AFFEB increase equates to roughly 20-24 years of emissions at current rates for the 1.5°C target. Land-based mitigation strategies also provide significant increases of 51-57 GtC and 56-62 GtC for the 1.5°C and 2°C AFFEB estimates, respectively. This is equivalent to 6-7 years of emissions at current rates. For our BECCS assumptions (see also below), we find that the BECCS contribution is small for the optimised land-based mitigation pathway and that AR are more effective land-based mitigation strategies (Fig. 9(b)). Although the primary challenge remains mitigation of fossil fuel emissions, these results highlight the unrealised potential of these mitigation options to make the Paris climate targets more achievable.

Commented [HGD20]: Response 23

Furthermore, the CH₄ and land-based mitigation strategies show little interaction and their potential can be summed to give a comparable result to the coupled simulation (coupled vs linear in Fig. 9(a) and (b)). This decoupling is despite the CH₄ emissions from the agricultural sector being influenced by land use choices. We can effectively treat the two mitigation strategies as independent, and their sum approximates the combined potential. Such linearity enables simpler and more direct comparisons.

Commented [HGD21]: Response 8

Despite the substantial differences in the absolute AFFEBs for the 1.5° and 2°C targets, the mitigation potential of the CH₄ and land-based strategies is similar for the two temperature scenarios-pathways considered. This similarity suggests that the mitigation strategies are robust to the target temperature; whether the international community aims for the 1.5° or 2°C target, afforestation, reforestation, reduced deforestation and CH₄ mitigation are beneficial mitigation approaches.

For both temperature pathways (i.e., 1.5°C or 2°C of warming), we investigate the contribution to the uncertainty range from ‘climate’ as represented by the 34 GCMs emulated and from the land processes investigated (Sect. 2.1). A GCM with

530 higher climate sensitivity will have a lower AFFEB for a specific warming target (and vice versa). In our post-processing steps,
we derive a number of statistical parameters from the complete 136-member or the 34-member GCM ensemble for the
individual factorial runs (low Q₁₀/low O₃, low Q₁₀/high O₃, high Q₁₀/low O₃ and high Q₁₀/high O₃), such as mean, standard
deviation, median, and various percentiles. Our focus is on the contribution different factors make to the overall standard
deviation of the 136-member ensemble (σ_{All}). By factoring out the climate variation (via their means), we calculate the standard
535 deviation for the land processes investigated (σ_{land}). With a knowledge of the overall standard deviation and that for land-only
processes, we derive the contribution from ‘climate’ ($\sigma_{climate}$) assuming that the variance are independent and can be summed
(Eq. 17). The contributions of uncertainty are by comparing ratios of σ_{land} to $\sigma_{climate}$.

$$\sigma_{all}^2 = \sigma_{climate}^2 + \sigma_{land}^2 \quad (17)$$

540 We present the results of this analysis in Table 3Table 3 for the Anthropogenic Fossil Field CO₂ Emission Budgets and
the Mitigation Potential (= scenario – “CTL”) for the 1.5°C temperature profile (Supplementary Information, Table SI.2 is
equivalent table for the 2°C temperature profile). Our overall finding is that the climate uncertainty dominates the uncertainty
of the AFFEBs. However, when considering different trade-offs between land uncertainty and mitigation options, the impact
of climate uncertainty is much weaker. Within the land uncertainty, the O₃ vegetation damage appears to make the greater
contribution (from the changes in the mean). Although there is some variation in the ratio ($\sigma_{climate}:\sigma_{land}$) between the scenarios
545 (0.32±0.13, mean ± standard deviation), this gives us confidence in the robustness of the uncertainty estimates derived, across
the scenarios and the 2 temperature profiles.

3.2 Sensitivity to BECCS Efficiency

550 The BECCS parameterisation used here makes BECCS less effective compared to those in other studies (van Vuuren et
al., 2018). Globally across the two temperature targets, our simulations imply a removal of 27-30 GtC from the active carbon
cycle via BECCS in the original “BECCS” scenario run, which is reduced to ~7-12 GtC after we optimise the land-use scenario.
555 These removal rates are significantly lower than other estimates based on the same land-use scenarios: 73 GtC in a similar
dynamic global vegetation model (LPJ-GUESS) and 130 GtC in IMAGE (Harper et al., 2018). We find that doubling the
carbon captured with BECCS in our simulations (Sect. 2.4.3, $\kappa=2$) has a relatively small impact on the total mitigation potential
in the optimised scenario (Fig. 10(a)). This low sensitivity is because the increased carbon removed by BECCS often
555 accompanies a comparable decrease in the carbon uptake from the “natural” vegetation that it replaces. It is only when setting
the BECCS carbon sequestration at 3-5 times its original value that there is a notable increase of the global AFFEB. Further,
as shown in Fig. 10(b), there is reduction in soil carbon in specific regions (e.g. Northern temperate and boreal regions), which
makes BECCS less effective for carbon sequestration than natural land management options (or there is a long payback time
as discussed in Harper et al. (2018)).

560 Increased carbon removal with BECCS could be realised through either (1) minimizing the loss of carbon from farm to final storage (ϵ in Sect. 2.4.3), or (2) maximizing the productivity of the bioenergy crop. Our IMOGEN-JULES simulations assume a 40% carbon loss from farm to final storage, although other studies have assumed this to be as low as 13% (Harper et al., 2018). The bioenergy crop yields in JULES (Fig. 10(c)) are lower than the median yield of Miscanthus (11.5 tons of dry matter ($\text{ton DM ha}^{-1} \text{ yr}^{-1}$), measured from 990 mostly European plots (Li et al., 2018), and are about half the productivity of those in the IMAGE simulations. We calculate for each IMOGEN grid cell the increase in carbon removed via BECCS and the associated increase in bioenergy crop yields (H^* in Sect. 2.4.3) required for BECCS to be the preferred mitigation option (Fig. 10(d)), rather than natural land carbon uptake, and assuming minimal amounts of carbon are lost during the BECCS lifecycle (13% carbon loss). In many places, we find that the required yield increases from <10 to 10-20 $\text{ton DM ha}^{-1} \text{ yr}^{-1}$ are achievable, but required yields of > 30 $\text{ton DM ha}^{-1} \text{ yr}^{-1}$ would be more difficult to realise, given the range of yields observed (Li et al., 2018). We provide additional information in the Supplementary Information, Tables SI.4a-SI.4d on the modelled bioenergy yields and the yields required for bioenergy crops to be the preferred land-based mitigation option by IMAGE region. The tables also show that area of bioenergy crops and carbon sequestered by BECCS increases, as expected, with the BECCS scale factor (κ).

575 We conclude that our uncorrected simulations are a lower estimate for the potential of carbon removal via BECCS. We provide a more optimistic estimate of the BECCS potential using $\kappa = 3$, which results from doubling the JULES yields and increasing the efficiency ϵ from 0.6 to 0.87 (i.e., $\kappa \sim 2 \times 0.87 / 0.6$). We now find the global land-based mitigation potential to be 88-100 GtC across the two temperature targets, as shown in Fig. 9(c) and (d). Supporting Information, Figure SI.3 shows the corresponding plots for the 2°C warming target. We use $\kappa = 3$ in the subsequent analysis of regional mitigation options and of BECCS water requirements.

580 3.3 Regional Analysis

We consider the sub-continental implications of CH_4 and land-based mitigation options, using the 26 regions of the IMAGE model (Stehfest et al., 2014). ~~Figure 11~~Figure 14 shows the contributions of the three mitigation options - CH_4 , carbon uptake through AR and BECCS - to the AFFEBs for each IMAGE region and for the temperature pathway stabilising at 1.5°C.

585 We estimate the regional land-based mitigation as the change in the land-carbon stores plus the carbon removal via BECCS for each IMAGE region in the IMOGEN-JULES model output. In this accounting, the region where the bioenergy crops are grown is credited with the carbon removal via BECCS. We assume a three-fold increase in carbon removal via BECCS compared to our default simulations ($\kappa=3$) to highlight regions where BECCS is potentially viable. Figure 12 shows the sensitivity of the global AFFEBs and Mitigation Potential for $\kappa = 1, 2$ and 3 for 1.5°C of warming (Supplementary Information, Figure SI.3 is the corresponding figure for 2°C of warming). For CH_4 , we ~~derive the regional contribution-use regional scale factors to the allocate~~ changes in the global atmospheric CH_4 concentration, and therefore the CH_4 mitigation potential, using

Commented [HGD22]: Response 19

595 regional fractions of the global difference in to each region, as shown in Supplementary Information, Table SI.3. To derive the regional scale factors, we separately sum the projected anthropogenic CH₄ emissions (between 2020- and 2100) between the IMAGE SSP2-Baseline and SSP2-1.9 scenarios (van Vuuren et al., 2017). We calculate the scale factor as the regional fraction of the global difference in the summed emissions (Supplementary Information, Table SI.3). These two CH₄ scenarios are consistent with the CH₄ concentration pathways considered in the CH₄ factorial-scenario simulations (Sect. 2.3). We use the scale factors to produce Fig. 11 and 12 (and Supplementary Information, Figures SI.3 and SI.4).

600 CH₄ mitigation is an effective mitigation strategy for all regions, and especially the major methane emitting regions: India, S. Africa, USA, China and Australasia. Figure 4 presented time series of the anthropogenic CH₄ emissions for selected IMAGE region from 2000 to 2100 (and Supplementary Information, Figure SI.1 presents emission time series for all IMAGE regions). The mitigation of CH₄ emissions from fossil-fuel production, distribution and use for energy is the largest contributor for India, S. Africa, USA, China and Australasia. The emissions from agriculture-cattle (for India, USA and China) and rice production (China and other Asian regions) make smaller contributions.

605 The impact of the land-based mitigation options links strongly to the managed land-use and land-use change (LULUC). As discussed in Sect. 2.3.2, we list in Table 2 the maximum area of BECCS deployed in each IMAGE region and the main differences in land use between the BECCS and Natural scenarios. Figure 5 presents time series of the land areas calculated for trees and prescribed for agriculture (including bioenergy crops) and bioenergy crops for the BECCS and Natural scenarios for the Russia and Brazil IMAGE regions, each as a difference to the baseline scenario (IM-BL) (see Supplementary Information, Figure SI.3 for all the IMAGE regions). The West Africa region shows the largest natural land carbon uptake (WAF in Fig. 12). Here, there is conversion of crop and pasture to forest, with little land used for bioenergy crops for BECCS. For Brazil (Fig. 5(a)) and the rest of South America, both bioenergy crops and forest expand at the expense of agricultural land. For many other regions, notably Canada, Russia, W. & C. Europe, China, Oceania, there is less carbon uptake from the 'land' in the optimised mitigation scenario, even though the overall carbon uptake has increased. For Canada and Russia, this results from the loss of forest in the BECCS land use scenario (see Fig. 5(b) and Supplementary Information, Figure SI.3). The carbon uptake by BECCS increases as κ increases from 1 to 3 because there are more grid cells where 'BECCS' is the preferred mitigation option in the optimisation process, as evidenced by the increase in area of bioenergy crops (Supplementary Information, Tables SI.3a and SI.3c). As κ only affects the 'BECCS' term (Sect. 2.4.3, Eq. 4+13), the increased carbon removed by BECCS is often accompanied by a decrease in the carbon uptake from the "natural" vegetation that it replaces. This can be seen more clearly in Fig. 12 (and Supplementary Information, Figure SI.3 for 2°C warming) and the Supplementary Information, Tables SI.3b and SI.3d. The version of JULES used in this study currently lacks a fire regime. There will be risks to long-term storage of carbon stored in vegetation in regions with significant areas of fire-dominated vegetation cover (e.g. savannah in Brazil and Africa). Further, this version of JULES does not include a nitrogen cycle, which has been implemented in more recent versions of the model. This will enable the impact of changes in land use and agriculture on N₂O emissions to be integrated into the assessments.

Commented [HGD23]: Response 20

625 There is relatively little difference in the additional allowable carbon emission budgets introduced by CH₄ and/or the land-
based mitigation between 2015 and 2100 for the two temperature [scenarios-pathways](#) considered (Supplementary Information,
Figure SI.4 for the contributions at 2°C of warming).

3.4 Water Resources

630 Smith et al. (2016) estimate the global water requirements for different negative emission technologies, including BECCS.
We also derive the water requirements from the carbon uptake by BECCS for our optimised land-based mitigation scenarios.
The IM-1.9 land use scenario (Sect. 2.3.2) assumes that bioenergy crops are grown sustainably and are rain-fed (Hoogwijk et
al., 2005; Daioglou et al., 2019). Our land surface modelling system explicitly accounts for this. We derive the additional water
requirements for BECCS, using $\kappa = 3$ and assuming (a) a marginal increase in water use of $80 \text{ m}^3 (\text{tC eq})^{-1} \text{ yr}^{-1}$ when replacing
the average short vegetation (i.e., C3/C4 grasses in JULES) by a biomass energy crop (Smith et al., 2016); and (b) $450 \text{ m}^3 (\text{tC}$
635 $\text{eq})^{-1} \text{ yr}^{-1}$ for the CCS component (Smith et al., 2016).

Following Postel et al. (1996), we derive the accessible runoff, using their assumptions that only 5% of the total runoff is
geographically and/or temporally accessible for the Brazil, Russia and Canada IMAGE regions, and 40% elsewhere. Our
present-day estimates of the global annual runoff ($43,000\text{--}44,200 \text{ km}^3 \text{ yr}^{-1}$) and the accessible runoff for human use ($11,400\text{--}$
 $11,720 \text{ km}^3 \text{ yr}^{-1}$) (see Fig. 13) are both in agreement with the values given in Postel et al. (1996), i.e., total and accessible
640 runoffs of $40,700$ and $12,500 \text{ km}^3 \text{ yr}^{-1}$, respectively.

We use the water withdrawals for each IMAGE region given in the IMAGE-SSP2-RCP2.6 scenario for the water demand
for agricultural irrigation (Rost et al., 2008) and for other human activities, such as energy generation, industry and domestic
usage (Bijl et al., 2016), between 2015 and 2100 (Table 4a and 4b). We assume the same water demands from these sectors
for both the 1.5° and 2°C warming targets.

645 Figure 14 compares the accessible water with the water demand for BECCS and other human activities for the regions
that produce a substantial amount of BECCS: Canada, USA, Brazil, Europe, Russia, China, Southern Africa and Oceania for
the optimised land-based mitigation. Table 4a and b show the additional water requirements of BECCS calculated for 2060
and 2100, respectively, for the 2°C warming target. We find that the additional demand for BECCS would lead to an
exceedence (or use >90%) of the available water for the Oceania and Rest of Southern Africa regions. We also find that the
650 additional demand for BECCS is greater than the total water withdrawals from anthropogenic activities for the Canada and
Brazil IMAGE regions. Our estimates represent a maximum possible water usage for BECCS as (i) the SSP2 scenario used
already accounts for the lower power generation efficiencies and hence higher water requirements in switching from fossil
fuels to bioenergy crops (which could be up to 20-25%) and (ii) the figure used for the CCS component does not allow for
future technological improvements in water use. For example, Fajardy and Mac Dowell (2017) indicate a 30-fold reduction in
655 water use when changing from a once-through to a recirculating cooling tower. Our results are less severe than other studies
considering BECCS water requirements (Séférián et al., 2018; Yamagata et al., 2018), because the carbon removed by BECCS

660 in this study (30 GtC) is already limited to regions where it is more beneficial to the AFFEB than forest-based mitigation options. We also note from Bijl et al. (2016) that the water demand for irrigation, derived using the coupled IMAGE-LPJmL models, is low compared to other estimates in the literature. Higher water demand for irrigation existing agriculture would be an additional constraint on the water available for BECCS. Nevertheless, our results indicate that the additional water demand for BECCS would have large impacts in half of the regions substantially invested in BECCS: Oceania, Rest of South Africa, Brazil and Canada.

4 Conclusions

665 Our paper brings together previous studies that looked separately into the potential of methane mitigation (Collins et al., 2018) and land-management options (especially forest conservation and BECCS) (Harper et al., 2018), into a single unified framework. Uniquely, this allows us to compare these options at local and regional scales. We utilise the detailed JULES land-surface model, which includes the temperature sensitivity of methanogenesis methane production from wetlands and permafrost thaw (Comyn-Platt et al., 2018) and the effect of CH₄ emissions on land carbon storage via ozone impacts on vegetation (Sitch et al., 2007), and also span the range of climate model projections using the IMOGEN ESM-emulator. For each temperature pathway and each of the three mitigation options, the set of factorial-scenario runs comprises a 136-member ensemble (34 670 GCMs x 2 ozone damage sensitivities x 2 methanogenesis Q₁₀ temperature sensitivities).

This analysis quantifies the regional differences in potential CH₄ and/or land-based strategies to aid mitigation of climate change. ~~Our~~ We present our findings are presented within a full probabilistic framework, capturing uncertainty in climate projections across the CMIP5 ensemble, as well as process uncertainties associated with the strength of natural CH₄ climate feedbacks from wetlands and ozone-induced vegetation damage. Globally, mitigation of anthropogenic CH₄ emissions and the optimised land-based mitigation can potentially offset (i.e. allow extra) fossil fuel carbon dioxide emissions of 188-212 GtC and 51-100 GtC, respectively. These bounds are almost independent of the eventual global-warming target, or the climate sensitivity of the climate models emulated. As shown in Sect. 3.1, the CH₄ and land-based mitigation strategies show little interaction and their potential can be summed to give a comparable result to the corresponding coupled simulation. This decoupling is despite the CH₄ emissions from the agricultural sector being influenced by land use choices. We can therefore treat the two mitigation strategies as independent, and sum their individual potentials. Such linearity enables simpler and more direct comparisons between the carbon budgets of methane and land-based mitigation strategies. Some caveats remain however. We acknowledge that ~~Land~~ land surface models still require refinement, alongside improved characterisation of the assumptions inherent in the socio-economic pathways and IAM modelling. Further, we do not allow for the reduced emissions from fossil fuel combustion due to the bioenergy crop being grown (or the converse when bioenergy crops are replaced in the Natural model run), as this would require energy sector modelling that is beyond the scope of this study.

685 For the "Natural" land-based scenario (see Table 1), we find a mitigation potential of 50-55 GtC (183-201 GtCO₂). The land-based mitigation estimates vary over wide ranges, partly related to different assumptions on land use and carbon pools.

Commented [HGD24]: Response 26

Commented [HGD25]: Response 26

Commented [HGD26]: Response 27

690 Our results are within the wide range of the overall deployment of CO₂ removal by Agriculture, Forestry and Other Land Use (including afforestation and reforestation) to 2100 of 200 [0-550] GtCO₂ (Page 2.40 in IPCC (2018)) and of estimates of the cumulative potential to 2100 from 80 to 260 GtCO₂ (Table 2) in Minx et al. (2018). In the “BECCS” scenario, we obtain a geological carbon storage via BECCS (27±1 GtC median, interquartile range) similar to that (30±1 GtC) derived by Harper et al. (2018), for the same land use scenario (IM-1.9). Our result is lower as we include the natural methane feedbacks from wetlands and permafrost thaw. Inclusion of this better process description leads to ~10% reduction in carbon budgets (Comyn-Platt et al., 2018). These estimates for the geological carbon storage via BECCS are much lower than the corresponding value derived by the IMAGE IAM (130 GtC). Harper et al. (2018) discuss this difference, identifying a number of reasons for the lower value: the use of initial above ground biomass harvested in boreal forests for BECCS, the replacement of fossil-fuel based emissions in the energy system, as well as specific assumptions about crop yields, conversion efficiency, use of residues, the proportion of bioenergy crops used with CCS. Estimates of the BECCS contribution in the literature vary over a wide range (from 178 to >1000 GtCO₂, according to Minx et al. (2018)), but in recent studies these result are typically revised downwards taking into account among others sustainability constraints (e.g. Fuss et al. (2018) suggests a potential of 0.5-5 GtCO₂ per year in 2050).

705 We investigate the efficacy of our “BECCS” scenario by increasing the productivity of BECCS (using a scale factor κ). From comparison with observed bioenergy crop yields, we argue that the scale factor could be between 1 and 3. We highlight how using this range of κ provides characterisation of an additional source of uncertainty on the land-based mitigation potential. In our optimised land-based mitigation scenario, which maximises the land carbon uptake (Sect 2.4.2, Eq. 10), the increased carbon removed by BECCS is often accompanied by a decrease in the carbon uptake from the “natural” vegetation that it replaces (as discussed in Sect. 3.3 and shown in Figure 12). This concern is equivalent to the statement in Harper et al. (2018) that the “use of BECCS in regions where bioenergy crops replace ecosystems with high carbon contents could easily result in negative carbon balance”. Hence the particularly novel feature of our paper is that our optimal approach accounts explicitly for that trade-off, only suggesting BECCS where there is a net gain. We quantify the sensitivity to the assumed productivity of bioenergy crops and the efficiency of the BECCS process. In consequence, our results for land-based mitigation strategies are nuanced, with considerable regional variations. For boreal forest regions there is a preference for avoided deforestation, whereas in tropical forest regions both AR and avoided deforestation offer significant potential. From a carbon sequestration perspective, growing bioenergy crops for BECCS is only preferable where it replaces existing agricultural land. BECCS has particular potential if productivities and power production efficiencies are towards the upper limit of expected photosynthetic capability, whilst noting the strong water demand of such crops requires consideration in the context of a growing population.

720 Stabilising the climate primarily requires urgent action to mitigate CO₂ emissions. However, we estimate that CH₄ mitigation has the potential to make the Paris targets more achievable by may offsetting up to 188-212 GtC of anthropogenic CO₂ emissions, while still meeting the same global-warming targets. This offset is a direct consequence of the reduced radiative forcing by methane and of carbon cycle gains. These balances and related flexibilities have the potential to make the Paris

Commented [HGD27]: Response 12

725 targets more achievable. Our range of additional CO₂ emissions broadly applies to both the 1.5° and 2°C warming targets, as
the mitigation potential of the CH₄ scenario is similar for the two temperature pathways considered. [Although there are
differences in the precise methane emission scenarios used, our mitigation potential is similar to that given in Collins et al.
(2018). That paper presents values of 155 or 235 GtC for offsetting CH₄ mitigation from a high to a medium or from a high to
a low emission scenario, respectively. Our value, and those of Collins et al. (2018), can be compared to the increase of 130
GtC in the carbon budget between a no and a stringent CH₄ emission mitigation scenario estimated by Rogelj et al. (2015).
More recently, Harmsen et al. (2020) have also investigated the mitigation potential of methane, although their results are
expressed in terms of changes in radiative forcing and temperature, rather than carbon budgets. An advantage of our analysis
remains the inclusion of climate response to altered radiative forcing, enabling understanding in terms of actual CO₂ emissions.

730 We conclude that CH₄ mitigation would be effective globally as a contribution to constraining global warming, and especially
so for the major CH₄-emitting regions of India, USA and China.

Code and Data Availability

735 The IMOGEN-JULES source code used in this work is available from the JULES code repository
(https://code.metoffice.gov.uk/trac/jules/browser/main/branches/dev/annaharper/r7971_vn4.8_IP5_DEGREES_CCS, at
JULES revision 14477, user account required). The rose suites used for the specific ~~factorial~~-IMOGEN-JULES runs are: u-
as624, u-at010, u-at011, u-at013, u-av005, u-av007, u-av008, u-av009, u-ax327, u-ax332, u-ax455, u-ax456, u-ax521, u-
ax523, u-ax524, u-ax525, u-bh009, u-bh023, u-bh046, u-bh081, u-bh084, u-bh098, u-bh103 and u-bh105. These can be found
740 at <https://code.metoffice.gov.uk/trac/roses-u/> (user account required).

The IMOGEN-JULES source code is also available as a zipped tarball from
https://github.com/GarryHayman/Regional_Mitigation_Paper, as are the python scripts used for post-processing. Relevant
outputs from the IMOGEN-JULES runs will be made available through a publically accessible data repository (tbc).

745 Requests for information about theAH code, data and parameterisations are available on request can be made to the
corresponding author.

Author Contributions

G.H., C.H., E.C-P., A.H., P.C., T.P., J.H., W.C., J.L. and S.C. designed the IMOGEN runs. All authors contributed to the
interpretation of the results and to the writing of or review of the paper. C.H. provided IMOGEN parameters calibrated against
the CMIP5 database, and E.C-P and C.H. led the development of the inverse IMOGEN model version. The following specific
750 contributions were also made: (a) E.B., S.C. and N.G.: code and expertise on permafrost, soil carbon and wetland methane
modelling, respectively; (b) A.H. and T.P.: land use change data; (c) W.C. and C.W.: ozone ancillary data; (d) D.P.vV. and

Commented [HGD28]: Response 26

Commented [HGD29]: Response 30

J.C.D.: IMAGE scenario data on land use, anthropogenic methane emissions and water consumption and withdrawals, and (e) S.S.: expertise on the ozone damage effects.

Competing interests

755 The authors declare no competing interests.

Acknowledgements

The work was undertaken as part of the UK Natural Environment Research Council's programme "Understanding the Pathways to and Impacts of a 1.5°C Rise in Global Temperature" through grants NE/P015050/1 CLIFFTOP (G.H., E.C-P, S.C.), NE/P014909/1, MOC1.5 (W.C., C.W., J.L., C.H., P.C., S.S.) and NE/P014941/1 CLUES (P.C., A.H., T.P., J.H.). We also acknowledge the support for: (a) G.H and E.C.P by NERC NE/N015746/1 The Global Methane Budget, MOYA; (b) A.H. through her EPSRC Fellowship "Negative Emissions and the Food-Energy-Water Nexus" (EP/N030141/1); (c) A.H. by NERC NE/P019951/1 FAB GGR, (d) W.C. from the Research Council of Norway, project no. 235548; (e) C.H. from CEH National Capability Funding; (f) E.B.. from the Joint UK BEIS/Defra Met Office Hadley Centre Climate Programme (GA01101); (g) E.B., D.P.vV. and J.C.D. from CRESCENDO (EU project 641816); and (h) NG from the Newton Fund through the Met Office Climate Science for Service Partnership Brazil (CSSP Brazil). All authors acknowledge the CMIP5 database, and its outputs from Earth System Models developed by climate research centres across the world. We also acknowledge Lars Kutzbach and David Holl, who kindly provided the methane emission data for the Samoylov Island field site. [We are grateful to the Editor and the two anonymous reviewers, whose comments have helped to improve the clarity of the paper.](#)

References

- 770 Anderson, K., and Peters, G.: The trouble with negative emissions, *Science*, 354, 182, <https://doi.org/10.1126/science.aah4567>, 2016.
- Best, M., Pryor, M., Clark, D., Rooney, G., Essery, R., Ménard, C., Edwards, J., Hendry, M., Porson, A., and Gedney, N.: The Joint UK Land Environment Simulator (JULES), model description–Part 1: energy and water fluxes, *Geoscientific Model Development*, 4, 677-699, <https://doi.org/10.5194/gmd-4-677-2011>, 2011.
- 775 Bijl, D. L., Bogaart, P. W., Kram, T., de Vries, B. J. M., and van Vuuren, D. P.: Long-term water demand for electricity, industry and households, *Environmental Science & Policy*, 55, 75-86, <https://doi.org/10.1016/j.envsci.2015.09.005>, 2016.
- Burke, E. J., Chadburn, S. E., and Ekici, A.: A vertical representation of soil carbon in the JULES land surface scheme (vn4.3_permafrost) with a focus on permafrost regions, *Geoscientific Model Development*, 10, 959, <https://doi.org/10.5194/gmd-10-959-2017>, 2017a.
- 780 Burke, E. J., Ekici, A., Huang, Y., Chadburn, S. E., Huntingford, C., Ciais, P., Friedlingstein, P., Peng, S., and Krinner, G.: Quantifying uncertainties of permafrost carbon–climate feedbacks, *Biogeosciences*, 14, 3051-3066, <https://doi.org/10.5194/bg-14-3051-2017>, 2017b.
- Chadburn, S., Burke, E., Essery, R., Boike, J., Langer, M., Heikenfeld, M., Cox, P., and Friedlingstein, P.: An improved representation of physical permafrost dynamics in the JULES land-surface model, *Geoscientific Model Development*, 8, 1493-1508, <https://doi.org/10.5194/gmd-8-1493-2015>, 2015.

785 Chadburn, S. E., Aalto, T., Aurela, M., Baldocchi, D., Biasi, C., Boike, J., Burke, E. J., Comyn-Platt, E., Dolman, A. J., Duran-Rojas, C., Fan, Y., Friborg, T., Gao, Y., Gedney, N., Göckede, M., Hayman, G. D., Holl, D., Hugelius, G., Kutzbach, L., Lee, H., Lohila, A., Parmentier, F.-J. W., Sachs, T., Shurpali, N. J., and Westermann, S.: Modeled Microbial Dynamics Explain the Apparent Temperature Sensitivity of Wetland Methane Emissions, *Global Biogeochemical Cycles*, 34, e2020GB006678, <https://doi.org/10.1029/2020GB006678>, 2020.

790 Clark, D., Mercado, L., Sitch, S., Jones, C., Gedney, N., Best, M., Pryor, M., Rooney, G., Essery, R., Blyth, E., Boucher, O., Harding, R., Huntingford, C., and Cox, P.: The Joint UK Land Environment Simulator (JULES), model description - Part 2: Carbon fluxes and vegetation dynamics, *Geoscientific Model Development*, 4, 701-722, <https://doi.org/10.5194/gmd-4-701-2011>, 2011.

Collins, W. J., Webber, C. P., Cox, P. M., Huntingford, C., Lowe, J., Sitch, S., Chadburn, S. E., Comyn-Platt, E., Harper, A. B., Hayman, G., and Powell, T.: Increased importance of methane reduction for a 1.5 degree target, *Environmental Research Letters*, 13, 054003, <https://doi.org/10.1088/1748-9326/aab89c>, 2018.

795 Comyn-Platt, E., Hayman, G., Huntingford, C., Chadburn, S. E., Burke, E. J., Harper, A. B., Collins, W. J., Webber, C. P., Powell, T., Cox, P. M., Gedney, N., and Sitch, S.: Carbon budgets for 1.5 and 2 °C targets lowered by natural wetland and permafrost feedbacks, *Nature Geoscience*, 11, 568-573, <https://doi.org/10.1038/s41561-018-0174-9>, 2018.

800 Cubasch, U., Meehl, G. A., Boer, G. J., Stouffer, R. J., Dix, M., Noda, A., Senior, C. A., Raper, S., and Yap, K. S.: Projections of Future Climate Change In: *Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change* [Houghton, J.T., Y. Ding, D.J. Griggs, M. Noguer, P.J. van der Linden, X. Dai, K. Maskell, and C.A. Johnson (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 881pp., Available from: <https://www.ipcc.ch/report/ar3/wg1/> (accessed November 2019), 2001.

Daioğlu, V., Doelman, J. C., Stehfest, E., Müller, C., Wicke, B., Faaij, A., and van Vuuren, D. P.: Greenhouse gas emission curves for advanced biofuel supply chains, *Nature Climate Change*, 7, 920-924, <https://doi.org/10.1038/s41558-017-0006-8>, 2017.

805 Daioğlu, V., Doelman, J. C., Wicke, B., Faaij, A., and van Vuuren, D. P.: Integrated assessment of biomass supply and demand in climate change mitigation scenarios, *Global Environmental Change*, 54, 88-101, <https://doi.org/10.1016/j.gloenvcha.2018.11.012>, 2019.

810 Doelman, J. C., Stehfest, E., Tabeau, A., van Meijl, H., Lassaletta, L., Gernaat, D. E. H. J., Hermans, K., Harmsen, M., Daioğlu, V., Biemans, H., van der Sluis, S., and van Vuuren, D. P.: Exploring SSP land-use dynamics using the IMAGE model: Regional and gridded scenarios of land-use change and land-based climate change mitigation, *Global Environmental Change*, 48, 119-135, <https://doi.org/10.1016/j.gloenvcha.2017.11.014>, 2018.

815 Ehhalt, D., Prather, M., Dentener, F., Derwent, R., Dlugokencky, E., Holland, E., Isaksen, I., Katima, J., Kirchhoff, V., P. Matson, Midgley, P., and Wang, M.: Atmospheric Chemistry and Greenhouse Gases, in: *Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Houghton, J. T., Y. Ding, D.J. Griggs, M. Noguer, P.J. van der Linden, X. Dai, K. Maskell and C.A. Johnson, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2001.

Etminan, M., G. Myhre, E. J. Highwood, and Shine, K. P.: Radiative forcing of carbon dioxide, methane, and nitrous oxide: A significant revision of the methane radiative forcing, *Geophysical Research Letters*, 43, <https://doi.org/10.1002/2016GL071930>, 2016.

820 Fajardy, M., and Mac Dowell, N.: Can BECCS deliver sustainable and resource efficient negative emissions?, *Energy & Environmental Science*, 10, 1389-1426, <https://doi.org/10.1039/C7EE00465F>, 2017.

Fuss, S., Lamb, W. F., Callaghan, M. W., Hilaire, J., Creutzig, F., Amann, T., Beringer, T., Garcia, W. d. O., Hartmann, J., Khanna, T., Luderer, G., Nemet, G. F., Rogelj, J., Smith, P., Vicente, J. L. V., Wilcox, J., M. del Mar Zamora, D., and Minx, J. C.: Negative emissions—Part 2: Costs, potentials and side effects, *Environmental Research Letters*, 13, 063002, <https://doi.org/10.1088/1748-9326/aabf9f>, 2018.

825 Gasser, T., Kechiar, M., Ciais, P., Burke, E. J., Kleinen, T., Zhu, D., Huang, Y., Ekici, A., and Obersteiner, M.: Path-dependent reductions in CO₂ emission budgets caused by permafrost carbon release, *Nature Geoscience*, <https://doi.org/10.1038/s41561-018-0227-0>, 2018.

Gedney, N., Huntingford, C., Comyn-Platt, E., and Wiltshire, A.: Significant feedbacks of wetland methane release on climate change and the causes of their uncertainty, *Environmental Research Letters*, 14, 084027, <https://doi.org/10.1088/1748-9326/ab2726>, 2019.

830 Gernaat, D. E. H. J., Calvin, K., Lucas, P. L., Luderer, G., Otto, S. A. C., Rao, S., Strefler, J., and van Vuuren, D. P.: Understanding the contribution of non-carbon dioxide gases in deep mitigation scenarios, *Global Environmental Change*, 33, 142-153, <https://doi.org/10.1016/j.gloenvcha.2015.04.010>, 2015.

- Harmsen, M., van Vuuren, D. P., Bodirsky, B. L., Chateau, J., Durand-Lasserve, O., Drouet, L., Fricko, O., Fujimori, S., Gernaat, D. E. H. J., Hanaoka, T., Hilaire, J., Keramidas, K., Luderer, G., Moura, M. C. P., Sano, F., Smith, S. J., and Wada, K.: The role of methane in future climate strategies: mitigation potentials and climate impacts, *Climatic Change*, 163, 1409-1425, [10.1007/s10584-019-02437-2](https://doi.org/10.1007/s10584-019-02437-2), 2020.
- 835 Harper, A. B., Cox, P. M., Friedlingstein, P., Wiltshire, A. J., Jones, C. D., Sitch, S., Mercado, L. M., Groenendijk, M., Robertson, E., Kattge, J., Bönsch, G., Atkin, O. K., Bahn, M., Cornelissen, J., Niinemets, Ü., Onipchenko, V., Peñuelas, J., Poorter, L., Reich, P. B., Soudzilovskaia, N. A., and Bodegom, P. V.: Improved representation of plant functional types and physiology in the Joint UK Land Environment Simulator (JULES v4.2) using plant trait information, *Geoscientific Model Development*, 9, 2415-2440, <https://doi.org/10.5194/gmd-9-2415-2016>, 2016.
- 840 Harper, A. B., Powell, T., Cox, P. M., House, J., Huntingford, C., Lenton, T. M., Sitch, S., Burke, E., Chadburn, S. E., Collins, W. J., Comyn-Platt, E., Daioglou, V., Doelman, J. C., Hayman, G., Robertson, E., van Vuuren, D., Wiltshire, A., Webber, C. P., Bastos, A., Boysen, L., Ciais, P., Devaraju, N., Jain, A. K., Krause, A., Poulter, B., and Shu, S.: Land-use emissions play a critical role in land-based mitigation for Paris climate targets, *Nature Communications*, 9, 2938, <https://doi.org/10.1038/s41467-018-05340-z>, 2018.
- Heck, V., Gerten, D., Lucht, W., and Popp, A.: Biomass-based negative emissions difficult to reconcile with planetary boundaries, *Nature Climate Change*, 8, 151-155, <https://doi.org/10.1038/s41558-017-0064-y>, 2018.
- 845 Hoogwijk, M., Faaij, A., Eickhout, B., de Vries, B., and Turkenburg, W.: Potential of biomass energy out to 2100, for four IPCC SRES land-use scenarios, *Biomass and Bioenergy*, 29, 225-257, <https://doi.org/10.1016/j.biombioe.2005.05.002>, 2005.
- Hoogwijk, M., Faaij, A., de Vries, B., and Turkenburg, W.: Exploration of regional and global cost–supply curves of biomass energy from short-rotation crops at abandoned cropland and rest land under four IPCC SRES land-use scenarios, *Biomass and Bioenergy*, 33, 26-43, <https://doi.org/10.1016/j.biombioe.2008.04.005>, 2009.
- 850 Huntingford, C., Booth, B. B. B., Sitch, S., Gedney, N., Lowe, J. A., Liddicoat, S. K., Mercado, L. M., Best, M. J., Weedon, G. P., Fisher, R. A., Lomas, M. R., Good, P., Zelazowski, P., Everitt, A. C., Spessa, A. C., and Jones, C. D.: IMOGEN: an intermediate complexity model to evaluate terrestrial impacts of a changing climate, *Geoscientific Model Development*, 3, 679-687, <https://doi.org/10.5194/gmd-3-679-2010>, 2010.
- 855 Huntingford, C., Yang, H., Harper, A., Cox, P. M., Gedney, N., Burke, E. J., Lowe, J. A., Hayman, G., Collins, W. J., Smith, S. M., and Comyn-Platt, E.: Flexible parameter-sparse global temperature time profiles that stabilise at 1.5 and 2.0 degrees C, *Earth System Dynamics*, 8, 617-626, <https://doi.org/10.5194/esd-8-617-2017>, 2017.
- IPCC: Global Warming of 1.5 °C, IPCC special report on the impacts of global warming of 1.5 °C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty, Available from: <http://www.ipcc.ch/report/sr15/> (accessed November 2019). 2018.
- 860 Jones, C., Hughes, J., Bellouin, N., Hardiman, S., Jones, G., Knight, J., Liddicoat, S., O'Connor, F., Andres, R. J., and Bell, C.: The HadGEM2-ES implementation of CMIP5 centennial simulations, *Geoscientific Model Development*, 4, 543, <https://doi.org/10.5194/gmd-4-543-2011>, 2011.
- Klein Goldewijk, K., Beusen, A., Van Drecht, G., and De Vos, M.: The HYDE 3.1 spatially explicit database of human-induced global land-use change over the past 12,000 years, *Global Ecology and Biogeography*, 20, 73-86, <https://doi.org/10.1111/j.1466-8238.2010.00587.x>, 2011.
- 865 Krause, A., Pugh, T. A. M., Bayer, A. D., Doelman, J. C., Humpenöder, F., Anthoni, P., Olin, S., Bodirsky, B. L., Popp, A., Stehfest, E., and Arneth, A.: Global consequences of afforestation and bioenergy cultivation on ecosystem service indicators, *Biogeosciences*, 14, 4829-4850, <https://doi.org/10.5194/bg-14-4829-2017>, 2017.
- 870 Le Quéré, C., Andrew, R. M., Friedlingstein, P., Sitch, S., Hauck, J., Pongratz, J., Pickers, P. A., Korsbakken, J. I., Peters, G. P., Canadell, J. G., Arneth, A., Arora, V. K., Barbero, L., Bastos, A., Bopp, L., Chevallier, F., Chini, L. P., Ciais, P., Doney, S. C., Gkritzalis, T., Goll, D. S., Harris, I., Haverd, V., Hoffman, F. M., Hoppema, M., Houghton, R. A., Hurtt, G., Ilyina, T., Jain, A. K., Johannessen, T., Jones, C. D., Kato, E., Keeling, R. F., Goldewijk, K. K., Landschützer, P., Lefèvre, N., Lienert, S., Liu, Z., Lombardozi, D., Metzl, N., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S. I., Neill, C., Olsen, A., Ono, T., Patra, P., Peregon, A., Peters, W., Peylin, P., Pfeil, B., Pierrot, D., Poulter, B., Rehder, G., Resplandy, L., Robertson, E., Rocher, M., Rödenbeck, C., Schuster, U., Schwinger, J., Séférian, R., Skjelvan, I., Steinhoff, T., Sutton, A., Tans, P. P., Tian, H., Tilbrook, B., Tubiello, F. N., van der Laan-Luijkx, I. T., van der Werf, G. R., Viovy, N., Walker, A. P., Wiltshire, A. J., Wright, R., Zaehle, S., and Zheng, B.: Global Carbon Budget 2018, *Earth Syst. Sci. Data*, 10, 2141-2194, <https://doi.org/10.5194/essd-10-2141-2018>, 2018.
- 875 Li, W., Ciais, P., Makowski, D., and Peng, S.: A global yield dataset for major lignocellulosic bioenergy crops based on field measurements, *Scientific Data*, 5, 180169, <https://doi.org/10.1038/sdata.2018.169>, 2018.

- 880 McNorton, J., Gloor, E., Wilson, C., Hayman, G. D., Gedney, N., Comyn-Platt, E., Marthews, T., Parker, R. J., Boesch, H., and
Chipperfield, M. P.: Role of regional wetland emissions in atmospheric methane variability, *Geophysical Research Letters*, 43, 11,433-
411,444, <https://doi.org/10.1002/2016gl070649>, 2016.
- Melton, J., Wania, R., Hodson, E., Poulter, B., Ringeval, B., Spahni, R., Bohn, T., Avis, C., Beerling, D., Chen, G., Eliseev, A., Denisov,
885 S., Hopcroft, P., Lettenmaier, D., Riley, W., Singarayer, J., Subin, Z., Tian, H., Zurcher, S., Brovkin, V., van Bodegom, P., Kleinen, T., Yu,
Z., and Kaplan, J.: Present state of global wetland extent and wetland methane modelling: conclusions from a model inter-comparison project
(WETCHIMP), *Biogeosciences*, 10, 753-788, <https://doi.org/10.5194/bg-10-753-2013>, 2013.
- Millar, R. J., Fuglestedt, J. S., Friedlingstein, P., Rogelj, J., Grubb, M. J., Matthews, H. D., Skeie, R. B., Forster, P. M., Frame, D. J.,
and Allen, M. R.: Emission budgets and pathways consistent with limiting warming to 1.5 °C, *Nature Geoscience*, 10, 741,
<https://doi.org/10.1038/ngeo3031>, 2017.
- 890 Minx, J. C., Lamb, W. F., Callaghan, M. W., Fuss, S., Hilaire, J., Creutzig, F., Amann, T., Beringer, T., de Oliveira Garcia, W.,
Hartmann, J., Khanna, T., Lenzi, D., Luderer, G., Nemet, G. F., Rogelj, J., Smith, P., Vicente Vicente, J. L., Wilcox, J., and del Mar Zamora
Dominguez, M.: Negative emissions—Part 1: Research landscape and synthesis, *Environmental Research Letters*, 13, 063001,
<https://doi.org/10.1088/1748-9326/aabf9b>, 2018.
- Myhre, G., Shindell, D., Bréon, F.-M., Collins, W., Fuglestedt, J. H., J., Koch, D., Lamarque, J.-F., Lee, D., Mendoza, B., Nakajima,
895 T., Robock, A., Stephens, G., Takemura, T., and Zhang, H.: Anthropogenic and Natural Radiative Forcing. In: *Climate Change 2013: The
Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*,
in: IPCC, 2013: *Climate Change 2013*, edited by: Stocker, T. F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y.
Xia, V. Bex and P.M. Midgley (eds.), Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA., 2013.
- O'Neill, B. C., Kriegler, E., Ebi, K. L., Kemp-Benedict, E., Riahi, K., Rothman, D. S., van Ruijven, B. J., van Vuuren, D. P., Birkmann,
900 J., Kok, K., Levy, M., and Solecki, W.: The roads ahead: Narratives for shared socioeconomic pathways describing world futures in the 21st
century, *Global Environmental Change*, 42, 169-180, <https://doi.org/10.1016/j.gloenvcha.2015.01.004>, 2017.
- Oliver, R. J., Mercado, L. M., Sitch, S., Simpson, D., Medlyn, B. E., Lin, Y. S., and Folberth, G. A.: Large but decreasing effect of
ozone on the European carbon sink, *Biogeosciences*, 15, 4245-4269, <https://doi.org/10.5194/bg-15-4245-2018>, 2018.
- Postel, S. L., Daily, G. C., and Ehrlich, P. R.: Human Appropriation of Renewable Fresh Water, *Science*, 271, 785,
905 <https://doi.org/10.1126/science.271.5250.785>, 1996.
- Riahi, K., van Vuuren, D. P., Kriegler, E., Edmonds, J., O'Neill, B. C., Fujimori, S., Bauer, N., Calvin, K., Dellink, R., Fricko, O., Lutz,
W., Popp, A., Cuaresma, J. C., Kc, S., Leimbach, M., Jiang, L., Kram, T., Rao, S., Emmerling, J., Ebi, K., Hasegawa, T., Havlik, P.,
Humpenöder, F., Da Silva, L. A., Smith, S., Stehfest, E., Bosetti, V., Eom, J., Gernaat, D., Masui, T., Rogelj, J., Strefler, J., Drouet, L., Krey,
910 V., Luderer, G., Harmsen, M., Takahashi, K., Baumstark, L., Doelman, J. C., Kainuma, M., Klimont, Z., Marangoni, G., Lotze-Campen, H.,
Obersteiner, M., Tabeau, A., and Tavoni, M.: The Shared Socioeconomic Pathways and their energy, land use, and greenhouse gas emissions
implications: An overview, *Global Environmental Change*, 42, 153-168, <https://doi.org/10.1016/j.gloenvcha.2016.05.009>, 2017.
- Rogelj, J., Meinshausen, M., Schaeffer, M., Knutti, R., and Riahi, K.: Impact of short-lived non-CO2 mitigation on carbon budgets for
stabilizing global warming, *Environmental Research Letters*, 10, 075001, 10.1088/1748-9326/10/7/075001, 2015.
- Rogelj, J., Popp, A., Calvin, K. V., Luderer, G., Emmerling, J., Gernaat, D., Fujimori, S., Strefler, J., Hasegawa, T., Marangoni, G.,
915 Krey, V., Kriegler, E., Riahi, K., van Vuuren, D. P., Doelman, J. C., Drouet, L., Edmonds, J., Fricko, O., Harmsen, M., Havlik, P.,
Humpenöder, F., Stehfest, E., and Tavoni, M.: Scenarios towards limiting global mean temperature increase below 1.5 °C, *Nature Climate
Change*, 8, 325-332, <https://doi.org/10.1038/s41558-018-0091-3>, 2018.
- Rost, S., Gerten, D., Bondeau, A., Lucht, W., Rohwer, J., and Schaphoff, S.: Agricultural green and blue water consumption and its
influence on the global water system, *Water Resources Research*, 44, <https://doi.org/10.1029/2007WR006331>, 2008.
- 920 Saunio, M., Bousquet, P., Poulter, B., Peregon, A., Ciais, P., Canadell, J. G., Dlugokencky, E. J., Etiope, G., Bastviken, D., Houweling,
S., Janssens-Maenhout, G., Tubiello, F. N., Castaldi, S., Jackson, R. B., Alexe, M., Arora, V. K., Beerling, D. J., Bergamaschi, P., Blake, D.
R., Brailsford, G., Brovkin, V., Bruhwiler, L., Crevoisier, C., Crill, P., Covey, K., Curry, C., Frankenberg, C., Gedney, N., Höglund-Isaksson,
L., Ishizawa, M., Ito, A., Joos, F., Kim, H. S., Kleinen, T., Krummel, P., Lamarque, J. F., Langenfelds, R., Locatelli, R., Machida, T.,
Maksyutov, S., McDonald, K. C., Marshall, J., Melton, J. R., Morino, I., Naik, V., O'Doherty, S., Parmentier, F. J. W., Patra, P. K., Peng,
925 C., Peng, S., Peters, G. P., Pison, I., Prigent, C., Prinn, R., Ramonet, M., Riley, W. J., Saito, M., Santini, M., Schroeder, R., Simpson, I. J.,
Spahni, R., Steele, P., Takizawa, A., Thornton, B. F., Tian, H., Tohjima, Y., Viovy, N., Voulgarakis, A., van Weele, M., van der Werf, G.
R., Weiss, R., Wiedinmyer, C., Wilton, D. J., Wilshire, A., Worthy, D., Wunch, D., Xu, X., Yoshida, Y., Zhang, B., Zhang, Z., and Zhu,
Q.: The global methane budget 2000–2012, *Earth Syst. Sci. Data*, 8, 697-751, <https://doi.org/10.5194/essd-8-697-2016>, 2016.

- 930 Saunois, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., Raymond, P. A., Dlugokencky, E. J., Houweling, S., Patra, P. K., Ciais, P., Arora, V. K., Bastviken, D., Bergamaschi, P., Blake, D. R., Brailsford, G., Bruhwiler, L., Carlson, K. M., Carrol, M., Castaldi, S., Chandra, N., Crevoisier, C., Crill, P. M., Covey, K., Curry, C. L., Etiopie, G., Frankenberg, C., Gedney, N., Heggin, M. I., Höglund-Isaksson, L., Hugelius, G., Ishizawa, M., Ito, A., Janssens-Maenhout, G., Jensen, K. M., Joos, F., Kleinen, T., Krummel, P. B., Langenfelds, R. L., Laruelle, G. G., Liu, L., Machida, T., Maksyutov, S., McDonald, K. C., McNorton, J., Miller, P. A., Melton, J. R., Morino, I., Müller, J., Murgia-Flores, F., Naik, V., Niwa, Y., Noce, S., O'Doherty, S., Parker, R. J., Peng, C., Peng, S., Peters, G. P., Prigent, C., Prinn, R., Ramonet, M., Regnier, P., Riley, W. J., Rosentretre, J. A., Segers, A., Simpson, I. J., Shi, H., Smith, S. J., Steele, P. L., Thornton, B. F., Tian, H., Tohjima, Y., Tubiello, F. N., Tsuruta, A., Viovy, N., Voulgarakis, A., Weber, T. S., van Weele, M., van der Werf, G. R., Weiss, R. F., Worthy, D., Wunch, D., Yin, Y., Yoshida, Y., Zhang, W., Zhang, Z., Zhao, Y., Zheng, B., Zhu, Q., Zhu, Q., and Zhuang, Q.: The Global Methane Budget 2000-2017, *Earth Syst. Sci. Data Discuss.*, 2019, 1-138, <https://doi.org/10.5194/essd-2019-128>, 2019.
- 940 Schuur, E. A. G., McGuire, A. D., Schadel, C., Grosse, G., Harden, J. W., Hayes, D. J., Hugelius, G., Koven, C. D., Kuhry, P., Lawrence, D. M., Natali, S. M., Olefeldt, D., Romanovsky, V. E., Schaefer, K., Turetsky, M. R., Treat, C. C., and Vonk, J. E.: Climate change and the permafrost carbon feedback, *Nature*, 520, 171-179, <https://doi.org/10.1038/nature14338>, 2015.
- Séférian, R., Rocher, M., Guivarch, C., and Colin, J.: Constraints on biomass energy deployment in mitigation pathways: the case of water scarcity, *Environmental Research Letters*, 13, 054011, <https://doi.org/10.1088/1748-9326/aabdc7>, 2018.
- 945 Shindell, D., Kuylenstierna, J. C. I., Vignati, E., van Dingenen, R., Amann, M., Klimont, Z., Anenberg, S. C., Müller, N., Janssens-Maenhout, G., Raes, F., Schwartz, J., Faluvegi, G., Pozzoli, L., Kupiainen, K., Hoglund-Isaksson, L., Emberson, L., Streets, D., Ramanathan, V., Hicks, K., Oanh, N. T. K., Milly, G., Williams, M., Demkine, V., and Fowler, D.: Simultaneously Mitigating Near-Term Climate Change and Improving Human Health and Food Security, *Science*, 335, 183-189, <https://doi.org/10.1126/science.1210026>, 2012.
- Sitch, S., Cox, P. M., Collins, W. J., and Huntingford, C.: Indirect radiative forcing of climate change through ozone effects on the land-carbon sink, *Nature*, 448, 791-794, <https://doi.org/10.1038/nature06059>, 2007.
- 950 Smith, P., Davis, S. J., Creutzig, F., Fuss, S., Minx, J., Gabrielle, B., Kato, E., Jackson, R. B., Cowie, A., Kriegler, E., van Vuuren, D. P., Rogelj, J., Ciais, P., Milne, J., Canadell, J. G., McCollum, D., Peters, G., Andrew, R., Krey, V., Shrestha, G., Friedlingstein, P., Gasser, T., Grübler, A., Heidug, W. K., Jonas, M., Jones, C. D., Kraxner, F., Littleton, E., Lowe, J., Moreira, J. R., Nakicenovic, N., Obersteiner, M., Patwardhan, A., Rogner, M., Rubin, E., Sharifi, A., Torvanger, A., Yamagata, Y., Edmonds, J., and Yongsung, C.: Biophysical and economic limits to negative CO₂ emissions, *Nature Climate Change*, 6, 42, <https://doi.org/10.1038/nclimate2870>, 2016.
- 955 Stehfest, E., van Vuuren, D., Kram, T., Bouwman, L., Alkemade, R., Bakkenes, M., Biemans, H., Bouwman, A., den Elzen, M., Jans, J., Lucas, P., van Minnen, J., Müller, C., and Prins, A.: Integrated Assessment of Global Environmental Change with IMAGE 3.0. Model description and policy applications, PBL Netherlands Environmental Assessment Agency, The Hague, Netherlands. , Available from: <http://www.pbl.nl/en/publications/integrated-assessment-of-global-environmental-change-with-IMAGE-3.0> (accessed November 2019), 2014.
- 960 Stocker, T., Qin, D., Plattner, G., Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, B., and Midgley, B.: The physical science basis. Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change, in: IPCC, 2013: Climate Change 2013, Cambridge University Press, 2013.
- 965 Stohl, A., Aamaas, B., Amann, M., Baker, L. H., Bellouin, N., Bernsten, T. K., Boucher, O., Cherian, R., Collins, W., Daskalakis, N., Dusinska, M., Eckhardt, S., Fuglestedt, J. S., Harju, M., Heyes, C., Hodnebrog, Ø., Hao, J., Im, U., Kanakidou, M., Klimont, Z., Kupiainen, K., Law, K. S., Lund, M. T., Maas, R., MacIntosh, C. R., Myhre, G., Myriokefalitakis, S., Orlivié, D., Quaas, J., Quennehen, B., Raut, J. C., Rumbold, S. T., Samset, B. H., Schulz, M., Seland, Ø., Shine, K. P., Skeie, R. B., Wang, S., Yttri, K. E., and Zhu, T.: Evaluating the climate and air quality impacts of short-lived pollutants, *Atmos. Chem. Phys.*, 15, 10529-10566, <https://doi.org/10.5194/acp-15-10529-2015>, 2015.
- 970 Turetsky, M. R., Kotowska, A., Bubier, J., Dise, N. B., Crill, P., Hornibrook, E. R. C., Minkinen, K., Moore, T. R., Myers-Smith, I. H., Nykänen, H., Olefeldt, D., Rinne, J., Saarnio, S., Shurpali, N., Tuittila, E.-S., Waddington, J. M., White, J. R., Wickland, K. P., and Wilming, M.: A synthesis of methane emissions from 71 northern, temperate, and subtropical wetlands, *Global Change Biology*, 20, 2183-2197, <https://doi.org/10.1111/gcb.12580>, 2014.
- UNFCCC: Adoption of the Paris Agreement, FCCC/CP/2015/L.9/Rev. 1, 2015.
- 975 van Vuuren, D. P., Stehfest, E., Gernaat, D. E. H. J., Doelman, J. C., van den Berg, M., Harmsen, M., de Boer, H. S., Bouwman, L. F., Daioglou, V., Edelenbosch, O. Y., Girod, B., Kram, T., Lassaletta, L., Lucas, P. L., van Meijl, H., Müller, C., van Ruijven, B. J., van der Sluis, S., and Tabeau, A.: Energy, land-use and greenhouse gas emissions trajectories under a green growth paradigm, *Global Environmental Change*, 42, 237-250, <https://doi.org/10.1016/j.gloenvcha.2016.05.008>, 2017.

van Vuuren, D. P., Stehfest, E., Gernaat, D. E. H. J., van den Berg, M., Bijl, D. L., de Boer, H. S., Daioglou, V., Doelman, J. C., Edelenbosch, O. Y., Harmsen, M., Hof, A. F., and van Sluisveld, M. A. E.: Alternative pathways to the 1.5 °C target reduce the need for negative emission technologies, *Nature Climate Change*, 8, 391-397, <https://doi.org/10.1038/s41558-018-0119-8>, 2018.

980 Vaughan, N. E., and Gough, C.: Expert assessment concludes negative emissions scenarios may not deliver, *Environmental Research Letters*, 11, 095003, <https://doi.org/10.1088/1748-9326/11/9/095003>, 2016.

Vaughan, N. E., Gough, C., Mander, S., Littleton, E. W., Welfle, A., Gernaat, D. E. H. J., and van Vuuren, D. P.: Evaluating the use of biomass energy with carbon capture and storage in low emission scenarios, *Environmental Research Letters*, 13, 044014, <https://doi.org/10.1088/1748-9326/aaaa02>, 2018.

985 WBA: Global Bioenergy Statistics 2019, World Bioenergy Association, Available from: https://worldbioenergy.org/uploads/191129%20WBA%20GBS%202019_HQ.pdf (accessed: November 2020), 2019.

Yamagata, Y., Hanasaki, N., Ito, A., Kinoshita, T., Murakami, D., and Zhou, Q.: Estimating water–food–ecosystem trade-offs for the global negative emission scenario (IPCC-RCP2.6), *Sustainability Science*, 13, 301-313, <https://doi.org/10.1007/s11625-017-0522-5>, 2018.

990 Zona, D., Gioli, B., Commene, R., Lindaas, J., Wofsy, S. C., Miller, C. E., Dinardo, S. J., Dengel, S., Sweeney, C., Karion, A., Chang, R. Y.-W., Henderson, J. M., Murphy, P. C., Goodrich, J. P., Moreaux, V., Liljedahl, A., Watts, J. D., Kimball, J. S., Lipson, D. A., and Oechel, W. C.: Cold season emissions dominate the Arctic tundra methane budget, *Proceedings of the National Academy of Sciences*, 113, 40-45, <https://doi.org/10.1073/pnas.1516017113>, 2016.






Figure 1 | Schematic of the modelling approach and the workflow. The coloured boxes and text show (a) the key components of the inverted IMOGEN-JULES model (blue), the prescribed and input data used in this study (orange) and the outputs (green).



(a)




(b)


Figure 2 | (a) Observed (circles) and modelled wetland methane emissions at the Samoylov Island field site. Modelled wetland methane emissions are shown for the standard JULES non-layered soil carbon configuration (green) and for the JULES layered soil carbon configurations with the low (blue line) and high (magenta line) Q_{10} temperature sensitivities; the low Q_{10} configuration gives higher methane emissions at high-latitude sites such as the Samoylov Island field site. The methane emission data is preliminary and was provided by Lars Kutzbach and David Holl. (b) Comparison of observed and modelled annual mean wetland CH_4 emission fluxes at a number of northern high-latitude and temperate sites. The error bars denote the lower and upper estimates from the low and high Q_{10} simulations. The symbols represent the mean value between these estimates.

Commented [HGD30]: Comment 21

1010 (a) Time series of the prescribed temperature pathways




1015 (b) Input time series of the input non-CO₂ radiative forcing



1015 Figure 3 | Time series of key datasets used in the study: (a) the historic temperature record (black) and the prescribed temperature profiles used to represent warming of 1.5°C (blue) and 2°C (orange); (b) the historic (black) and the projected non-CO₂ greenhouse gas radiative forcing (W m⁻²) for the control (orange) and methane mitigation (blue) scenarios.

Commented [HGD31]: Response 22




1020

Figure 4 | Time series of annual methane emissions between 2005 and 2100 from all and selected anthropogenic sources according to the IMAGE SSP2 Baseline (solid lines) and SSP2-RCP1.9 (dotted lines) scenarios, globally and for selected IMAGE regions, with total emissions in black, energy sector in red, agriculture-cattle in blue, agriculture-rice in green and waste in magenta. Note the y-axes have different scales for clarity.

1025

1026


(a) IMAGE Brazil Region



1027

1028


(b) IMAGE Russia Region



1029

1030

1031 **Figure 5 | Time series of the land areas (in Mha) calculated for trees and prescribed for agriculture (including bioenergy crops) and bioenergy crops for**
1032 **the 'BECCS' (orange) and 'Natural, (green), as a difference to the baseline scenario (IM-BL), for Brazil (panel a) and the Russia (panel b) IMAGE regions**
1033 **between 2000 and 2100. The dotted lines are the median and the spread the interquartile range for the 34 GCMs emulated and 4 factorial sensitivity**
1034 **simulations.**




1036

1037 **Figure 6 | (a, c, e)** Time series of the ensemble median atmospheric CH_4 concentrations (with interquartile range as spread) derived
 1038 for each temperature profile for the scenarios: (a) “CTL” and “ CH_4 ”, (c) “BECCS” and “BECCS+ CH_4 ”, (e) “Natural” and
 1039 “Natural+ CH_4 ”. (d, f, h) show the corresponding time series for the atmospheric CO_2 concentrations.


1040

1041 (a)



1042

1043 (b)



1044

1045 **Figure 7 | Scale factor required for BECCS to be the preferable mitigation option, as opposed to natural land carbon uptake. The**
1046 **data represents the median of the 136 member ensemble for the optimised land-based mitigation simulation. Panel (a) is for**
1047 **stabilisation at 1.5°C and panel (b) is for stabilisation at 2°C.**

1048





Figure 8 | The contribution to the allowable anthropogenic fossil fuel emission budget (AFFEBs, GtC) from the changes in the different carbon stores (atmosphere, ocean, land and BECCS) for the various control and mitigation scenarios, illustrated using the temperature pathways reaching for 1.5°C without overshoot of warming. The bars are the median of the component 136-member ensembles, with the individual members shown as points. The accompanying pink box and whisker plots to the right of each set of bars are for the AFFEBs (as the sum of the changes in the component carbon stores). The box and whisker plots show the median, interquartile range, minimum and maximum derived of the resulting AFFEB ensemble. The optimised land based and coupled mitigation options select the land use option, which maximises the AFFEB for each model grid cell. Note that the gain in the land carbon store for the CH₄ scenario is at -1.4 GtC (median of ensemble) is not visible, although the individual ensemble members can be seen as the green points shown as a reduction from -70.8 GtC in the control run to -1.4 GtC in the “CH₄” mitigation option (median of ensemble).


Commented [HGD32]: Response 23

1062 BECCS Scale Factor (κ) = 1



1063

1064 BECCS Scale Factor (κ) = 3




1065

1066 Figure 9 | Panels (a & c): The allowable anthropogenic fossil fuel emission budgets (AFFEBs; GtC) for the control (grey), CH₄
1067 mitigation (purple), land-based mitigation (green), coupled methane and land-based mitigation (orange) and the linearly summed
1068 methane and land-based mitigation (brown), for 2 temperature pathways asymptoting at 1.5°C (left) and 2.0°C (right). (b & d) The
1069 mitigation potential (GtC) as the increase in AFFEB from the corresponding control run. The breakdown of each AFFEB and
1070 mitigation potential by the changes in the carbon stores is also shown: atmosphere (pale yellow), ocean (light blue), land (dark green)
1071 and BECCS (gold) is included alongside each bar. Note that the gain in the land carbon store for the “CH₄-CH₄” scenario is at -1.4

|072 GtC (median of ensemble) is not visible. There has however been a net increase in the land carbon store in this scenario when
|073 compared to the land carbon store in the control run (shown as a reduction from -70.8 GtC₂ in the control run to -1.4 GtC in the
|074 "CH₄" mitigation option (median of ensemble)).

Commented [HGD33]: Response 23

1075
1076
1077
1078




1079
1080

1081 **Figure 10** (a) The total and component mitigation potential (GtC) for different mitigation options, involving methane and land use, as a function of the
1082 BECCS efficiency factor (κ , Sect. 2.4.3) for the temperature pathway reaching 1.5°C. The spread of the functions width of the lines represent the
1083 interquartile range of the sensitivity-136-member ensembles. Maps of (b) the change of the modelled soil carbon (kg-C m⁻²) between 2015 and 2099, as the
1084 difference between the scenario with BECCS and the natural land-management scenario; (c) the modelled mean bioenergy crop yield in the JULES
1085 simulations ($\kappa = 1$) and (d) the required bioenergy crop yield for BECCS to provide a larger carbon uptake than forest regrowth/afforestation (assuming
1086 $\kappa = \kappa^*$ and 87% efficiency of BECCS). Grid cells which do not exceed 1% BECCS cover for any year in the simulation are masked grey.


1087

Commented [HGD34]: Response 24




1088


1089 **Figure 11 |** The contribution to the allowable carbon emission budgets (GtC) between 2015 and 2100 for each of the 26 IMAGE IAM regions from methane
 1090 mitigation (purple bars) and land-based mitigation options (green: natural land uptake; yellow: BECCS with $\kappa = 3$), for the temperature pathway
 1091 stabilising at 1.5° warming without overshoot. The bars and error bars respectively show the median and the interquartile range, from the **34 GCMs**
 1092 **emulated and 4 factorial-run 136-member ensembles.**



(a)



(b)



(c)

1093

1094


1095

1096

1097 Figure 12 | Contribution of different mitigation options to the increase in allowable anthropogenic fossil fuel emission budgets by
 1098 IMAGE region to meet the 1.5°C target. The stacked bars represent the median methane mitigation potential (purple bars) and
 1099 median land-based mitigation potential (natural land uptake, green; BECCS, brown). Panel (a) is based on a BECCS scaling factor
 1100 of unity, (b) a BECCS scaling factor of 2 and (c) a BECCS scaling factor of 3. The total (pink) shows the median and interquartile
 1101 range for the 34 GCMs emulated and 4 factorial sensitivity simulations of the 136-member ensembles.

1102

1103




1104

1105 **Figure 13 | Global water availability (filled light blue bar) as a regionally dependent fraction of runoff (hollow light blue bar) for the**
1106 **year 2015. The water demand for irrigation (dark blue) and for other uses (i.e., energy generation, industry and domestic; yellow),**
1107 **are taken from the SSP2-RCP2.6-IMAGE database. Note there is very little BECCS additional water demand (green) in 2015.**

1108

1109

1110



1111

1112 **Figure 14 | Water availability (light blue), SSP2-IMAGE water demand estimates for irrigation (dark blue), other uses (i.e., energy**
 1113 **generation, industry and domestic; yellow) and the additional water demand from BECCS (green) for the years 2059-2060 and 2099-**
 1114 **2100 for the 2.0°C warming target, with a BECCS κ factor of 3. The points are the individual results from the 34 GCMs emulated**
 1115 **and 4 factorial runs 136-member ensembles, while the bars are the corresponding median values of the different GCM/factorial**
 1116 **ensembles.**

1117 **Tables**

1118 **Table 1 | The IMOGEN-JULES and post processing scenario runs, key features and the input and prescribed datasets used in the scenarios.**

1119 (a) IMOGEN-JULES modelling scenarios (Note 1)

	Scenario (Abbreviation) Key features of the Scenario	Scenario-specific input and prescribed datasets (Notes 2, 3)
1.	<p>Control (“CTL”)</p> <ul style="list-style-type: none"> • <u>Agricultural land accrued to feed growing populations associated with the SSP2 pathway</u> • <u>No deployment of BECCS</u> • <u>Anthropogenic CH₄ emissions rise from 318 Tg yr⁻¹ in 2005 to 484 Tg yr⁻¹ in 2100</u> • <u>Effects of the methane and carbon-climate feedbacks from wetlands and permafrost thaw included</u> 	<p><u>Scenario-specific input data</u></p> <ul style="list-style-type: none"> • <u>Time series of radiative forcing by non-CO₂ GHG and other non-CO₂ climate forcers, for the IMAGE SSP2 baseline scenario</u> • <u>Time series of annual global atmospheric concentrations of CH₄ and N₂O for the IMAGE SSP2 baseline scenario</u> <p><u>Scenario-specific prescribed data</u></p> <ul style="list-style-type: none"> • <u>Gridded annual time series of areas assigned to agriculture (crops & pasture), for the IMAGE SSP2 baseline scenario, converted into fractions of the IMOGEN-JULES grid cell</u>
2.	<p>Methane mitigation (“CH₄”)</p> <ul style="list-style-type: none"> • <u>Agricultural land-use as in Control (“CTL”) scenario</u> • <u>Anthropogenic CH₄ emissions decline from 318 Tg yr⁻¹ in 2005 to 162 Tg yr⁻¹ in 2100, from the IMAGE SSP2 RCPI.9 scenario</u> • <u>Effects of the methane and carbon-climate feedbacks from wetlands and permafrost thaw included</u> 	<p><u>Scenario-specific input data</u></p> <ul style="list-style-type: none"> • <u>Time series of radiative forcing by non-CO₂ GHG and other non-CO₂ climate forcers, for the IMAGE SSP2 RCPI.9 scenario</u> • <u>Time series of annual global atmospheric concentrations of CH₄ and N₂O for the IMAGE SSP2 RCPI.9 scenario</u> <p><u>Scenario-specific prescribed data</u></p> <ul style="list-style-type: none"> • <u>As 1, gridded annual time series of area assigned to agriculture (crops & pasture). Converted into fractions of the IMOGEN-JULES grid cell</u>
3.	<p>Land-based mitigation, including BECCS (“BECCS”)</p> <ul style="list-style-type: none"> • <u>Land use change based on the IMAGE SSP2 RCPI.9 scenario</u> • <u>High levels of REDD and full reforestation</u> • <u>Food-first policy so that bioenergy crops (BE) are only implemented on land not required for food production</u> • <u>Anthropogenic CH₄ emissions as in Control (“CTL”) scenario</u> • <u>Effects of the methane and carbon-climate feedbacks from wetlands and permafrost thaw included</u> 	<p><u>Scenario-specific input data</u></p> <ul style="list-style-type: none"> • <u>Time series of radiative forcing by non-CO₂ GHG and other non-CO₂ climate forcers, for the IMAGE SSP2 baseline scenario</u> • <u>Time series of annual global atmospheric concentrations of CH₄ and N₂O for the IMAGE SSP2 baseline scenario (as used in “CTL”)</u> <p><u>Scenario-specific prescribed data</u></p> <ul style="list-style-type: none"> • <u>Gridded annual time series of areas assigned to agriculture (crops & pasture) and within that the area for bioenergy crops, for the IMAGE SSP2 RCPI.9 scenario. Converted into a fraction of the IMOGEN-JULES grid cell</u>
4.	<p>Land-based mitigation with no BECCS (“Natural”)</p> <ul style="list-style-type: none"> • <u>Land use as 3, except any land area allocated to bioenergy crops is set to zero, allowing expansion of natural vegetation</u> • <u>Anthropogenic CH₄ emissions as in Control (“CTL”) scenario</u> • <u>Effects of the methane and carbon-climate feedbacks from wetlands and permafrost thaw included</u> 	<p><u>Scenario-specific input data</u></p> <ul style="list-style-type: none"> • <u>Time series of radiative forcing by non-CO₂ GHG and other non-CO₂ climate forcers, for the IMAGE SSP2 baseline scenario</u> • <u>Time series of annual global atmospheric concentrations of CH₄ and N₂O for the IMAGE SSP2 baseline scenario (as used in “CTL”)</u> <p><u>Scenario-specific prescribed data</u></p> <ul style="list-style-type: none"> • <u>Gridded annual time series of areas assigned to agriculture (crops & pasture). As 3, except any land allocated to bioenergy crops is set to zero. Converted into a fraction of the IMOGEN-JULES grid cell</u>

5.	<p><u>Combined methane & land-based mitigation “Coupled(CH₄+BECCS)”</u></p> <ul style="list-style-type: none"> Combines CH₄ mitigation of 2 with land-based mitigation scenario of 3 	<p><u>Scenario-specific input data</u></p> <ul style="list-style-type: none"> As 2, time series of radiative forcing by non-CO₂ GHG and other non-CO₂ climate forcers, for the IMAGE SSP2 RCP1.9 scenario As 2, time series of annual global atmospheric concentrations of CH₄ and N₂O for the IMAGE SSP2 RCP1.9 scenario <p><u>Scenario-specific prescribed data</u></p> <ul style="list-style-type: none"> As 3, gridded annual time series of areas assigned to agriculture (crops & pasture) and within that the area for bioenergy crops, for the IMAGE SSP2 RCP1.9 scenario. Converted into prescribed fractions of the IMOGEN-JULES grid cell
6.	<p><u>Combined methane & land-based mitigation with no BECCS “Coupled (CH₄+Natural)”</u></p> <ul style="list-style-type: none"> Combines CH₄ mitigation of 2 with land-based mitigation scenario of 4 	<p><u>Scenario-specific input data</u></p> <ul style="list-style-type: none"> As 2, time series of radiative forcing by non-CO₂ GHG and other non-CO₂ climate forcers, for the IMAGE SSP2 RCP1.9 scenario As 2, time series of annual global atmospheric concentrations of CH₄ and N₂O for the IMAGE SSP2 RCP1.9 scenario <p><u>Scenario-specific prescribed data</u></p> <ul style="list-style-type: none"> As 4, gridded annual time series of areas assigned to agriculture (crops & pasture). Converted into a fraction of the IMOGEN-JULES grid cell

1120

1121 (b) Post-processing scenarios (Note 1)

	<u>Scenario “Abbreviation”</u>	<u>Description of the Scenario</u>
7.	<u>Optimisation of land-based mitigation “Land-based mitigation: Optimised”</u>	<ul style="list-style-type: none"> Optimisation of scenarios 3 and 4 by selecting the scenario which has the larger carbon uptake, on a grid cell by grid cell basis
8.	<u>Optimisation of the combined methane & land-based mitigation “Coupled Optimised”</u>	<ul style="list-style-type: none"> Optimisation of scenarios 5 and 6 by selecting the scenario which has the larger carbon uptake, on a grid cell by grid cell basis

1122

1123 **Notes**

1124 1. Each scenario comprises two 136-member ensembles (34 GCMs x 2 ozone damage sensitivities x 2 methanogenesis Q₁₀ temperature sensitivities), one for the 1.5°C warming target
1125 and the second for the 2°C warming target.

1126 2. All of the above scenarios also use time series of (1) observed temperature changes between 1850 and 2015; (2) profiles of temperature change between 2015 and 2100 to achieve the
1127 1.5°C and the 2°C warming targets; and (3) the radiative forcing changes of non-CO₂ radiative forcing between 1850 and 2015.

1128 3. We define (a) a “prescribed” dataset as one that is used unchanged in the IMOGEN-JULES modelling; (b) an “input” dataset as one that provides the initial values that are subsequently
1129 changed.

1130 ~~The IMOGEN-JULES factorial runs, key features and IMAGE input/prescribed datasets.~~

Factorial Run	Abbreviation
Control	CTL

<p>IMAGE SSP2 baseline scenario</p> <p>Agricultural land accrued to feed growing populations associated with the SSP2 pathway. No deployment of BECCS</p> <p>Anthropogenic CH₄ emissions rise from 318 Tg yr⁻¹ in 2005 to 484 Tg yr⁻¹ in 2100</p> <p>IMAGE SSP2 baseline scenario for atmospheric concentrations of CH₄ and non-CO₂ radiative forcing</p>	
<p>Methane mitigation:</p> <p>IMAGE SSP2 RCP1.9 scenario for CH₄</p> <p>Agricultural land-use as in Control</p> <p>Anthropogenic CH₄ emissions decline from 318 Tg yr⁻¹ in 2005 to 162 Tg yr⁻¹ in 2100</p> <p>IMAGE SSP2 1.9 scenario for atmospheric concentrations of CH₄ and non-CO₂ radiative forcing</p>	CH₄
<p>Land-based mitigation, including BECCS:</p> <p>Land use from IMAGE SSP2 RCP1.9 scenario</p> <p>High levels of REDD and full reforestation</p> <p>Food first policy so that bioenergy crops are only implemented on land not required for food production</p> <p>Anthropogenic CH₄ emissions as in Control</p> <p>IMAGE SSP2 baseline scenario for atmospheric concentrations of CH₄ and non-CO₂ radiative forcing</p>	BECCS
<p>Land-based mitigation with no BECCS (Natural):</p> <p>As 3, except any land allocated to bioenergy crops is set to zero, allowing expansion of natural vegetation</p> <p>Anthropogenic CH₄ emissions as in Control</p>	Natural

IMAGE SSP2 baseline scenario for atmospheric concentrations of CH₄ and non-CO₂ radiative forcing	
Combined methane & land based mitigation: Combines CH₄ mitigation of 2 with land based mitigation of 3 IMAGE SSP2-1.9 scenario for atmospheric concentrations of CH₄ and non-CO₂ radiative forcing	Coupled (BECCS+CH₄)
Combined methane & land based mitigation with no BECCS (Natural) Combines CH₄ mitigation of 2 with land use scenario of 4 IMAGE SSP2-1.9 scenario for atmospheric concentrations of CH₄ and non-CO₂ radiative forcing	Coupled (Natural+CH₄)

1131

1132 ~~Note: Each factorial run comprises a 136 member ensemble: 34 GCMs x 2 ozone damage sensitivities x 2 methanogenesis Q₁₀ temperature sensitivities.~~

1133

1134 **Table 2 | IMAGE regions, the maximum area of BECCS deployed (Mha) and the main differences in land use between the BECCS**
1135 **and Natural scenarios.**

Region	Abbreviation	Max. area of bioenergy crops (Mha)	Main land-use difference between BECCS and Natural scenarios
Canada	CAN	65.9	Forest to BECCS in BECCS scenario
USA	USA	39.0	Agricultural land and forest to BECCS (BECCS). Agricultural land to forest (Natural)
Mexico	MEX	7.1	Agricultural land to BECCS and forest (BECCS). Agricultural land to forest (Natural)
Central America	RCAM	0.5	Little BECCS. Agricultural land to forests in both scenarios.
Brazil	BRA	27.8	Agricultural land to BECCS and forest (BECCS). Agricultural land to forest (Natural)
Rest of South America	RSAM	20.3	Agricultural land to BECCS and forest (BECCS). Agricultural land to forest (Natural)
Northern Africa	NAF	0.0	No BECCS. No real differences between scenarios
Western Africa	WAF	3.1	Little BECCS. Agricultural land to forests in both scenarios.
Eastern Africa	EAF	33.9	Agricultural land to BECCS and forest (BECCS). Agricultural land to forest (Natural)
South Africa	SAF	1.0	Little BECCS. Agricultural land to forests in both scenarios.
Rest of Southern Africa	RSAF	63.7	Agricultural land to BECCS and forest (BECCS). Agricultural land to forest (Natural)
Western Europe	WEU	23.6	Forest to BECCS in BECCS scenario
Central Europe	CEU	19.3	Forest to BECCS in BECCS scenario
Turkey	TUR	0.0	No BECCS. No real differences between scenarios
Ukraine Region	UKR	11.4	Forest to BECCS in BECCS scenario
Central Asia	STAN	0.7	Little BECCS. No real differences between scenarios
Russia Region	RUS	146.1	Forest to BECCS in BECCS scenario
Middle East	ME	0.0	No BECCS. No real differences between scenarios
India	INDIA	6.0	Forest to BECCS in BECCS scenario
Korea Region	KOR	4.3	Forest to BECCS in BECCS scenario
China	CHN	58.1	Forest to BECCS in BECCS scenario
South East Asia	SEAS	24.5	Forest to BECCS in BECCS scenario. Agricultural land to forest (Natural)
Indonesia	INDO	0.0	No BECCS. Agricultural land to forests in both scenarios.
Japan	JAP	2.7	Forest to BECCS in BECCS scenario
Rest of South Asia	RSAS	0.0	No BECCS. No real differences between scenarios
Oceania	OCE	78.7	Forest to BECCS in BECCS scenario

1136

1137

1138

1139 Table 3 | IMAGE regions, and the sum of the projected accumulated anthropogenic CH₄ emissions between (2020- and 2100) for the
 1140 SSP2-Baseline and SSP2-RCP1.9 scenarios, and the difference between these summed emissions. The regional scale factor is
 1141 calculated as the regional fraction of the global difference in anthropogenic CH₄ the summed emissions (2020-2100).

Region	Abbreviation	Projected Anthropogenic CH ₄ Emissions 2020-2100 (PgCH ₄)		Difference	Scale Factor
		SSP2-Baseline	SSP2-RCP1.9		
Canada	CAN	0.497	0.169	0.328	0.01471
USA	USA	3.281	1.573	1.708	0.07670
Mexico	MEX	0.542	0.320	0.222	0.00995
Central America	RCAM	0.312	0.195	0.117	0.00525
Brazil	BRA	2.502	1.638	0.865	0.03884
Rest of South America	RSAM	2.249	1.159	1.090	0.04896
Northern Africa	NAF	0.533	0.286	0.247	0.01110
Western Africa	WAF	2.035	1.128	0.907	0.04074
Eastern Africa	EAF	1.722	1.245	0.478	0.02146
South Africa	SAF	1.615	0.207	1.408	0.06324
Rest of Southern Africa	RSAF	1.883	0.924	0.959	0.04307
Western Europe	WEU	0.683	0.220	0.463	0.02081
Central Europe	CEU	0.409	0.219	0.190	0.00854
Turkey	TUR	0.387	0.128	0.259	0.01163
Ukraine Region	UKR	1.021	0.299	0.722	0.03241
Central Asia	STAN	1.743	0.514	1.228	0.05517
Russia Region	RUS	1.910	0.720	1.190	0.05343
Middle East	ME	4.873	1.788	3.085	0.13856
India	INDIA	0.170	0.081	0.089	0.00400
Korea Region	KOR	5.757	2.351	3.406	0.15296
China	CHN	1.923	0.908	1.015	0.04558
South East Asia	SEAS	1.005	0.457	0.547	0.02458
Indonesia	INDO	0.160	0.077	0.082	0.00369
Japan	JAP	1.316	0.460	0.856	0.03846
Rest of South Asia	RSAS	1.496	0.893	0.603	0.02710
Oceania	OCE	0.657	0.455	0.202	0.00907
World	World	40.680	18.415	22.265	1.00000

Commented [HGD35]: Response 20. Now in Supplementary Information

1142

For the 1.5°C temperature profile, the mean of the 34-GCM member ensembles for the “CTL” and mitigation scenarios for the different factorial runs (low Q₁₀/low O₃, low Q₁₀/high O₃, high Q₁₀/low O₃ and high Q₁₀/high O₃), the standard deviation of the full 136-member ensemble (GtC), the derived standard deviations for land processes (σ_{land}) and climate (σ_{climate}), as represented by the 34 GCMs) and the ratio of $\sigma_{\text{climate}}/\sigma_{\text{land}}$ for (a) the Anthropogenic Fossil Field CO₂ Emission Budgets and (b) the Mitigation Potential (= scenario – CTL).

(1) AFFEB

Scenario	Mean of 34-member Factorial Run (GtC)				Standard Deviation (GtC)			Ratio $\sigma_{\text{climate}}:\sigma_{\text{land}}$
	Low Q ₁₀ Low O ₃	Low Q ₁₀ High O ₃	High Q ₁₀ Low O ₃	High Q ₁₀ High O ₃	136-member Ensemble	Land σ_{land}	Climate σ_{climate}	
CTL	-9.66	-20.58	-18.91	-31.06	47.12	7.60	46.50	6.12
CH ₄	179.44	186.79	168.73	174.90	47.54	6.59	47.08	7.14
BECCS	6.49	3.42	-2.09	-5.80	47.45	4.76	47.21	9.91
Natural	42.57	24.60	35.00	16.05	48.95	10.07	47.90	4.75
Optimised Land-based	46.42	29.18	37.89	20.00	48.85	9.84	47.85	4.86
Linear BECCS+CH ₄	195.58	210.79	185.55	200.15	48.64	9.07	47.79	5.27
Linear Natural+CH ₄	231.67	231.97	222.64	222.00	48.70	4.76	48.47	10.19
Linear optimised	235.51	236.55	225.53	225.96	48.69	5.16	48.42	9.39
Coupled BECCS+CH ₄	199.69	214.62	189.50	203.94	48.48	9.01	47.64	5.29
Coupled Natural+CH ₄	237.83	238.95	228.72	228.91	48.60	4.80	48.36	10.07
Coupled optimised	241.50	243.29	231.35	232.60	48.60	5.27	48.31	9.17

(2) Mitigation Potential

Scenario	Mean of 34-member Factorial Run (GtC)				Standard Deviation (GtC)			Ratio $\sigma_{\text{climate}}:\sigma_{\text{land}}$
	Low Q ₁₀ Low O ₃	Low Q ₁₀ High O ₃	High Q ₁₀ Low O ₃	High Q ₁₀ High O ₃	136-member Ensemble	Land σ_{land}	Climate σ_{climate}	
CTL	-	-	-	-	-	-	-	-
CH ₄	189.10	207.37	187.64	205.96	9.28	9.18	1.39	0.15
BECCS	16.14	24.01	16.82	25.26	4.24	4.11	1.05	0.26
Natural	52.23	45.18	53.91	47.11	3.93	3.58	1.62	0.45
Optimised Land-based	56.07	49.76	56.80	51.06	3.44	3.06	1.57	0.51
Linear BECCS+CH ₄	205.24	231.38	204.46	231.21	13.39	13.23	2.09	0.16
Linear Natural+CH ₄	241.33	252.55	241.55	253.06	6.14	5.69	2.32	0.41
Linear optimised	245.17	257.13	244.44	257.02	6.55	6.14	2.28	0.37
Coupled BECCS+CH ₄	209.34	235.20	208.41	235.00	13.27	13.12	2.01	0.15
Coupled Natural+CH ₄	247.48	259.54	247.63	259.97	6.49	6.10	2.21	0.36
Coupled optimised	251.15	263.87	250.26	263.66	6.89	6.54	2.17	0.33

1149 **Table 4a | Comparison by IMAGE region of the modelled available water (km³ yr⁻¹), the projected water withdrawals (km³ yr⁻¹) for irrigation and for other anthropogenic activities**
1150 **(energy generation, industry, domestic) from the IMAGE SSP2-RCP2.6 scenario, and the additional water required for BECCS (km³ yr⁻¹ and as percentages of the net available water**
1151 **and of the water withdrawals for irrigation and other), for the year 2060. The percentage of runoff available for human use by IMAGE region is also included.**

Region	Abbreviation	% of Regional Runoff Available	Available Water (km ³ yr ⁻¹)	Water Demand			Total Demand as % of Available Water	BECCS Demand as % of Total Demand
				Irrigation (km ³ yr ⁻¹)	Other (km ³ yr ⁻¹)	BECCS (km ³ yr ⁻¹)		
Canada	CAN	40%	243.19	3.39	14.21	44.45	25.5%	71.6%
USA	USA	5%	1,010.82	149.55	96.07	44.55	28.7%	15.4%
Mexico	MEX	5%	75.89	76.58	25.56	24.48	166.8%	19.3%
Central America	RCAM	5%	185.92	8.16	15.49	2.28	13.9%	8.8%
Brazil	BRA	40%	310.65	12.24	34.44	73.12	38.6%	61.0%
Rest of South America	RSAM	5%	1,779.42	93.50	46.49	67.66	11.7%	32.6%
Northern Africa	NAF	5%	0.11	61.60	54.63	0.00	-	-
Western Africa	WAF	5%	1,962.47	28.29	118.83	0.39	7.5%	0.3%
Eastern Africa	EAF	5%	485.18	53.92	63.10	2.45	24.6%	2.1%
South Africa	SAF	5%	0.60	13.45	9.28	0.48	3868.3%	2.1%
Rest of Southern Africa	RSAF	5%	182.48	10.03	41.36	56.02	58.9%	52.2%
Western Europe	WEU	5%	642.34	78.72	82.01	56.22	33.8%	25.9%
Central Europe	CEU	5%	176.27	27.46	22.32	29.68	45.1%	37.4%
Turkey	TUR	5%	29.98	60.35	15.86	0.00	-	-
Ukraine Region	UKR	5%	67.47	11.73	25.90	12.28	74.0%	24.6%
Central Asia	STAN	5%	20.57	88.26	32.62	0.00	-	-
Russia Region	RUS	40%	270.32	42.30	51.60	103.87	73.2%	52.5%
Middle East	ME	5%	8.65	149.55	40.97	0.00	-	-
India	INDIA	5%	319.36	374.18	501.06	0.00	-	-
Korea Region	KOR	5%	42.85	6.20	9.75	12.64	66.7%	44.2%
China	CHN	5%	887.26	338.81	236.89	87.73	74.8%	13.2%
South East Asia	SEAS	5%	1,212.00	46.52	92.99	31.56	14.1%	18.4%
Indonesia	INDO	5%	1,293.05	8.18	113.87	0.00	-	-
Japan	JAP	5%	209.49	2.79	18.99	7.69	14.1%	26.1%
Rest of South Asia	RSAS	5%	74.57	259.95	154.42	0.00	-	-
Oceania	OCE	5%	85.46	24.99	8.91	48.06	95.9%	58.6%

1152

1153

1154 Table 4b | As Table 4a for 2100

Region	Abbreviation	% of Regional Runoff Available	Available Water (km ³ yr ⁻¹)	Water Demand			Total Demand as % of Available Water	BECCS Demand as % of Total Demand
				Irrigation (km ³ yr ⁻¹)	Other (km ³ yr ⁻¹)	BECCS (km ³ yr ⁻¹)		
Canada	CAN	40%	240.14	4.31	11.72	45.21	25.5%	73.8%
USA	USA	5%	993.09	148.57	81.35	45.45	27.7%	16.5%
Mexico	MEX	5%	72.79	77.27	23.78	11.14	154.1%	9.9%
Central America	RCAM	5%	182.12	8.74	13.96	0.66	12.8%	2.8%
Brazil	BRA	40%	307.53	12.31	30.80	54.89	31.9%	56.0%
Rest of South America	RSAM	5%	1,765.14	103.97	38.34	32.65	9.9%	18.7%
Northern Africa	NAF	5%	0.11	57.89	56.98	0.00	-	-
Western Africa	WAF	5%	1,953.10	37.23	262.07	0.62	15.4%	0.2%
Eastern Africa	EAF	5%	485.02	58.96	128.33	20.54	42.8%	9.9%
South Africa	SAF	5%	0.60	13.43	7.50	0.45	3563.3%	2.1%
Rest of Southern Africa	RSAF	5%	179.63	11.20	89.87	74.85	97.9%	42.5%
Western Europe	WEU	5%	637.68	80.39	118.64	45.25	38.3%	18.5%
Central Europe	CEU	5%	171.05	26.90	20.63	23.19	41.3%	32.8%
Turkey	TUR	5%	29.52	60.49	12.87	0.00	-	-
Ukraine Region	UKR	5%	66.45	10.40	19.58	8.62	58.1%	22.3%
Central Asia	STAN	5%	19.67	82.08	37.90	0.00	-	-
Russia Region	RUS	40%	266.36	40.25	43.82	58.40	53.5%	41.0%
Middle East	ME	5%	8.60	136.63	39.30	0.00	-	-
India	INDIA	5%	320.08	388.69	585.48	0.00	-	-
Korea Region	KOR	5%	42.73	7.41	5.47	0.00	-	-
China	CHN	5%	881.00	326.62	144.80	72.75	61.8%	13.4%
South East Asia	SEAS	5%	1,213.01	45.46	131.95	19.49	16.2%	9.9%
Indonesia	INDO	5%	1,291.53	15.08	114.33	0.00	-	-
Japan	JAP	5%	208.43	2.12	13.29	6.94	10.7%	31.1%
Rest of South Asia	RSAS	5%	74.19	245.78	227.85	0.00	0.0%	0.0%
Oceania	OCE	5%	85.46	30.57	8.77	62.96	136.5%	160.0%

Regional variation in the effectiveness of methane and land-based climate mitigation options

Garry D. Hayman^{1,*}, Edward Comyn-Platt¹, Chris Huntingford¹, Anna B. Harper², Tom Powell², Peter M. Cox², William Collins³, Christopher Webber³, Jason Lowe^{4,5}, Stephen Sitch², Joanna I. House⁶, Jonathan C. Doelman⁷, Detlef P. van Vuuren^{7,8}, Sarah E. Chadburn², Eleanor Burke⁵, Nicola Gedney⁹.

¹ Centre for Ecology & Hydrology, Wallingford, OX10 8BB, U.K.

² University of Exeter, Exeter, EX4 4QF, U.K.

³ University of Reading, Reading, RG6 6BB, U.K.

⁴ University of Leeds, Leeds, LS2 9JT, U.K.

⁵ Met Office Hadley Centre, FitzRoy Road, Exeter, EX1 3PB, U.K.

⁶ Cabot Institute for the Environment, University of Bristol, Bristol, BS8 1SS, U.K.

⁷ Department of Climate, Air and Energy, Netherlands Environmental Assessment Agency (PBL), PO Box 30314, 2500 GH The Hague, Netherlands

⁸ Copernicus Institute of Sustainable Development, Utrecht University, Heidelberglaan 2, 3584 CS, the Netherlands

⁹ Met Office Hadley Centre, Joint Centre for Hydrometeorological Research, Wallingford, OX10 8BB, U.K.

Correspondence to: Garry Hayman (garr@ceh.ac.uk)

Supplementary Information




Fig. SI.1 | Time series of annual methane emissions between 2005 and 2100 from all and selected anthropogenic sources according to the IMAGE SSP2 Baseline (solid lines) and SSP2-RCP1.9 (dotted lines) scenarios, globally and for each of the 26 IMAGE regions, with total emissions in black, energy sector in red, agriculture-cattle in blue, agriculture-rice in green and waste in magenta. Note the y-axes have different scales for clarity.

25





Fig. SI.1 (continued) | Time series of annual methane emissions between 2005 and 2100 from all and selected anthropogenic sources. Note the y-axes have different scales for clarity.



30

Fig. SI.1 (continued) | Time series of annual methane emissions between 2005 and 2100 from all and selected anthropogenic sources. Note the y-axes have different scales for clarity.

35 **Fig. SI.2** | Time series of the land areas (in Mha) calculated for trees and prescribed for
agriculture (including bioenergy crops) and bioenergy crops for the “BECCS” (orange) and
“NATURAL” (green) scenarios, as a difference to the baseline scenario (“CTL” = IM-BL),
for the 26 IMAGE regions between 2000 and 2100. The dotted lines are the median and the
spread the interquartile range for the 34 GCMs emulated and 4 factorial sensitivity
simulations.

40

- a) Canada
- b) USA
- c) Mexico
- d) Central America
- 45 e) Brazil
- f) Rest of South America
- g) Northern Africa
- h) Western Africa
- i) Eastern Africa
- 50 j) South Africa
- k) Rest of Southern Africa
- l) Western Europe
- m) Central Europe
- n) Turkey
- 55 o) Ukraine Region
- p) Central Asia
- q) Russia Region
- r) Middle East
- s) India
- 60 t) Rest of South Asia
- u) China
- v) Korea Region
- w) Japan
- x) South East Asia
- 65 y) Indonesia
- z) Oceania

Fig. SI.2a: Canada




Fig. SI.2b: USA




Fig. SI.2c: Mexico




Fig. SI.2d: Central America




Fig. SI.2e: Brazil




Fig. SI.2f: Rest of South America




Fig. SI.2g: Northern Africa




Fig. SI.2h: Western Africa




Fig. SI.2i: Eastern Africa




Fig. SI.2j: South Africa




Fig. SI.2k: Rest of Southern Africa




Fig. SI.2l: Western Europe




Fig. SI.2m: Central Europe




Fig. SI.2n: Turkey




Fig. SI.2o: Ukraine Region




Fig. SI.2p: Central Asia




Fig. SI.2q: Russia Region




Fig. SI.2r: Middle East




Fig. SI.2s: India




Fig. SI.2t: Rest of South Asia




Fig. SI.2u: China




Fig. SI.2v: Korea Region




Fig. SI.2w: Japan




Fig. SI.2x: South East Asia




Fig. SI.23y: Indonesia






Fig. SI.2z: Oceania






135

Fig. SI.3 | Contribution of different mitigation options to the increase in allowable anthropogenic fossil fuel emission budgets by IMAGE region to meet the 2°C target.

The stacked bars represent the median methane mitigation potential (purple bars) and median land-based mitigation potential (natural land uptake, green; BECCS, brown). Panel (a) is based on a BECCS scaling factor of unity, (b) a BECCS scaling factor of 2 and (c) a BECCS scaling factor of 3. The total (pink) shows the median and interquartile range for the 34 GCMs emulated and 4 factorial sensitivity simulations.

140



145 **Fig. SI.4 | Contribution of different mitigation options to the allowable anthropogenic carbon emission budgets by region.** The contribution to the allowable carbon emission budgets (GtC) between 2015 and 2100 for each of the 26 IMAGE IAM regions from methane mitigation (purple bars) and land-based mitigation options (green: natural land uptake; yellow: BECCS with $\kappa = 3$), for the temperature pathway stabilising at 2°C warming without overshoot. The bars and error bars respectively show the median and the interquartile range, from the 34 GCMs emulated and 4 factorial runs.

Table SI.1 | Mitigation options, estimated maximum reduction potential and the accompanying marginal price for mitigation of different anthropogenic methane source sectors for 2050 and 2100 [based on Lucas et al., 2007].

Source Sector	Mitigation option(s)	Max. possible reduction relative to baseline (%)	Marginal price of max. reduction (1995 US\$/tC _{eq})
Coal production	Maximising methane recovery from underground mining of hard coal	90 (2050) 90 (2100)	500 (2050) 500 (2100)
Oil/gas production & distribution	Control of fugitive emissions from equipment and pipeline leaks, and from venting during maintenance and repair.	75 (2050) 90 (2100)	300 (2050) 500 (2100)
Enteric fermentation	Change of animal diet and use of more productive animal types.	50 (2050) 60 (2100)	1000 (2050) 1000 (2100)
Animal waste	Capture and use of methane emissions through anaerobic digesters.	50 (2050) 60 (2100)	1000 (2050) 1000 (2100)
Wetland rice production	Changes to (1) the water management regime to reduce the period of anaerobic conditions in flooded fields; (2) the soils to reduce methanogenesis.	80 (2050) 90 (2100)	1000 (2050) 1000 (2100)
Landfills	(1) Reduced amount of organic material deposited in landfills; (2) capture of methane	90 (2050) 90 (2100)	500 (2050) 500 (2100)
Sewage and wastewater	(1) More wastewater treatment plants and also recovery of the methane from the plants; (2) More aerobic wastewater treatment.	80 (2050) 90 (2100)	500 (2050) 500 (2100)
Other anthropogenic sources	Note 1	-	-

Note: (1) These sources are either difficult to abate (e.g., land clearing for agricultural extension, and the use of traditional biomass for energy production and cooking) or are too small (e.g., methane emissions from industry, iron and steel production and the chemical sector).

Reference: Lucas, P. L., van Vuuren, D. P., Olivier, J. G. J. & den Elzen, M. G. J., 2007: Long-term reduction potential of non-CO₂ greenhouse gases. *Environmental Science & Policy* **10**, 85-103, doi: <https://doi.org/10.1016/j.envsci.2006.10.007>.

155

Table SI.2 | For the 2°C temperature profile, the mean of the 34-GCM member ensembles for the “CTL” and mitigation scenarios for the different factorial runs (low Q₁₀/low O₃, low Q₁₀/high O₃, high Q₁₀/low O₃ and high Q₁₀/high O₃), the standard deviation of the full 136-member ensemble (GtC), the derived standard deviations for land processes (σ_{land}) and climate (σ_{climate} , as represented by the 34 GCMs) and the ratio of $\sigma_{\text{climate}}/\sigma_{\text{land}}$ for (a) the Anthropogenic Fossil Field CO₂ Emission Budgets and (b) the Mitigation Potential (= scenario – CTL)

(1) AFFEB

Scenario	Mean of 34-member Factorial Run (GtC)				Standard Deviation (GtC)			Ratio $\sigma_{\text{climate}}:\sigma_{\text{land}}$
	Low Q ₁₀ Low O ₃	Low Q ₁₀ High O ₃	High Q ₁₀ Low O ₃	High Q ₁₀ High O ₃	136-member Ensemble	Land σ_{land}	Climate σ_{climate}	
CTL	229.67	230.28	213.87	213.23	86.80	8.22	86.41	10.51
CH ₄	423.93	443.83	405.57	424.31	89.03	13.53	88.00	6.50
BECCS	249.83	257.92	234.59	242.12	87.99	8.69	87.56	10.08
Natural	287.79	281.62	273.47	266.44	89.02	8.08	88.65	10.97
Optimised Land-based	291.34	285.95	276.06	270.11	88.94	8.28	88.55	10.69
Linear BECCS+CH ₄	444.09	471.47	426.30	453.20	90.63	16.29	89.15	5.47
Linear Natural+CH ₄	482.04	495.17	465.17	477.51	90.86	10.73	90.22	8.41
Linear optimised	485.60	499.50	467.76	481.19	90.84	11.33	90.13	7.95
Coupled BECCS+CH ₄	447.69	474.68	429.79	456.36	90.39	16.16	88.93	5.50
Coupled Natural+CH ₄	487.77	501.94	470.88	484.20	90.81	11.06	90.13	8.15
Coupled optimised	491.24	505.99	473.25	487.54	90.85	11.65	90.10	7.73

160

(2) Mitigation Potential

Scenario	Mean of 34-member Factorial Run (GtC)				Standard Deviation (GtC)			Ratio $\sigma_{\text{climate}}:\sigma_{\text{land}}$
	Low Q ₁₀ Low O ₃	Low Q ₁₀ High O ₃	High Q ₁₀ Low O ₃	High Q ₁₀ High O ₃	136-member Ensemble	Land σ_{land}	Climate σ_{climate}	
CTL	-	-	-	-	-	-	-	-
CH ₄	194.25	213.55	191.70	211.08	9.95	9.75	1.98	0.20
BECCS	20.16	27.64	20.72	28.89	4.21	3.94	1.48	0.37
Natural	58.11	51.34	59.60	53.20	4.22	3.40	2.50	0.74
Optimised Land-based	61.67	55.67	62.19	56.88	3.73	2.87	2.39	0.83
Linear BECCS+CH ₄	214.41	241.19	212.43	239.96	13.97	13.60	3.18	0.23
Linear Natural+CH ₄	252.37	264.89	251.30	264.28	7.55	6.39	4.02	0.63
Linear optimised	255.92	269.22	253.89	267.95	7.94	6.89	3.94	0.57
Coupled BECCS+CH ₄	218.02	244.40	215.93	243.13	13.76	13.42	3.03	0.23
Coupled Natural+CH ₄	258.10	271.66	257.01	270.96	7.94	6.89	3.94	0.57
Coupled optimised	261.56	275.71	259.38	274.31	8.30	7.33	3.90	0.53

Table SI.3 | IMAGE regions, the sum of the projected anthropogenic CH₄ emissions between 2020 and 2100 for the SSP2-Baseline and SSP2-RCP1.9 scenarios, and the difference between these summed emissions. The regional scale factor is calculated as the regional fraction of the global difference in the summed emissions.

<u>Region</u>	<u>Abbreviation</u>	<u>Sum of Projected Anthropogenic CH₄ Emissions 2020-2100 (PgCH₄)</u>		<u>Difference</u>	<u>Scale Factor</u>
		<u>SSP2-Baseline</u>	<u>SSP2-RCP1.9</u>		
		<u>Canada</u>	<u>CAN</u>		
<u>USA</u>	<u>USA</u>	<u>3.281</u>	<u>1.573</u>	<u>1.708</u>	<u>0.07670</u>
<u>Mexico</u>	<u>MEX</u>	<u>0.542</u>	<u>0.320</u>	<u>0.222</u>	<u>0.00995</u>
<u>Central America</u>	<u>RCAM</u>	<u>0.312</u>	<u>0.195</u>	<u>0.117</u>	<u>0.00525</u>
<u>Brazil</u>	<u>BRA</u>	<u>2.502</u>	<u>1.638</u>	<u>0.865</u>	<u>0.03884</u>
<u>Rest of South America</u>	<u>RSAM</u>	<u>2.249</u>	<u>1.159</u>	<u>1.090</u>	<u>0.04896</u>
<u>Northern Africa</u>	<u>NAF</u>	<u>0.533</u>	<u>0.286</u>	<u>0.247</u>	<u>0.01110</u>
<u>Western Africa</u>	<u>WAF</u>	<u>2.035</u>	<u>1.128</u>	<u>0.907</u>	<u>0.04074</u>
<u>Eastern Africa</u>	<u>EAF</u>	<u>1.722</u>	<u>1.245</u>	<u>0.478</u>	<u>0.02146</u>
<u>South Africa</u>	<u>SAF</u>	<u>1.615</u>	<u>0.207</u>	<u>1.408</u>	<u>0.06324</u>
<u>Rest of Southern Africa</u>	<u>RSAF</u>	<u>1.883</u>	<u>0.924</u>	<u>0.959</u>	<u>0.04307</u>
<u>Western Europe</u>	<u>WEU</u>	<u>0.683</u>	<u>0.220</u>	<u>0.463</u>	<u>0.02081</u>
<u>Central Europe</u>	<u>CEU</u>	<u>0.409</u>	<u>0.219</u>	<u>0.190</u>	<u>0.00854</u>
<u>Turkey</u>	<u>TUR</u>	<u>0.387</u>	<u>0.128</u>	<u>0.259</u>	<u>0.01163</u>
<u>Ukraine Region</u>	<u>UKR</u>	<u>1.021</u>	<u>0.299</u>	<u>0.722</u>	<u>0.03241</u>
<u>Central Asia</u>	<u>STAN</u>	<u>1.743</u>	<u>0.514</u>	<u>1.228</u>	<u>0.05517</u>
<u>Russia Region</u>	<u>RUS</u>	<u>1.910</u>	<u>0.720</u>	<u>1.190</u>	<u>0.05343</u>
<u>Middle East</u>	<u>ME</u>	<u>4.873</u>	<u>1.788</u>	<u>3.085</u>	<u>0.13856</u>
<u>India</u>	<u>INDIA</u>	<u>0.170</u>	<u>0.081</u>	<u>0.089</u>	<u>0.00400</u>
<u>Korea Region</u>	<u>KOR</u>	<u>5.757</u>	<u>2.351</u>	<u>3.406</u>	<u>0.15296</u>
<u>China</u>	<u>CHN</u>	<u>1.923</u>	<u>0.908</u>	<u>1.015</u>	<u>0.04558</u>
<u>South East Asia</u>	<u>SEAS</u>	<u>1.005</u>	<u>0.457</u>	<u>0.547</u>	<u>0.02458</u>
<u>Indonesia</u>	<u>INDO</u>	<u>0.160</u>	<u>0.077</u>	<u>0.082</u>	<u>0.00369</u>
<u>Japan</u>	<u>JAP</u>	<u>1.316</u>	<u>0.460</u>	<u>0.856</u>	<u>0.03846</u>
<u>Rest of South Asia</u>	<u>RSAS</u>	<u>1.496</u>	<u>0.893</u>	<u>0.603</u>	<u>0.02710</u>
<u>Oceania</u>	<u>OCE</u>	<u>0.657</u>	<u>0.455</u>	<u>0.202</u>	<u>0.00907</u>
<u>World</u>	<u>World</u>	<u>40.680</u>	<u>18.415</u>	<u>22.265</u>	<u>1.00000</u>

Table SI.4a | By IMAGE region for the 1.5°C temperature profile, the modelled bioenergy crop productivity (ton DM ha⁻¹ yr⁻¹) and that required to match the land carbon uptake in the variant “Natural” land-based mitigation scenario, the grid-cell scale-factors (Required/Modelled BECCS productivity), the maximum area of BECCS (Mha) and the area of bioenergy crops (Mha) in the optimised land-based scenario as a function of the BECCS scale factor κ , (all entries are median [10%, 90% percentiles] of the grid cells in the IMAGE region from the 136-member ensembles).

Region	Maximum area of BECCS (Mha)	BECCS Productivity (ton DM ha ⁻¹ yr ⁻¹)		Required Scale Factor	Area of BECCS (Mha) in optimised land-based scenario			
		Modelled	Required to match Natural		BECCS scale factor $\kappa = 1$	BECCS scale factor $\kappa = 2$	BECCS scale factor $\kappa = 3$	BECCS scale factor $\kappa = 4$
Canada	65.9	2.99 (0.00-4.75)	6.39 (0.00-20.16)	2.93 (0.81-24.94)	1.02 (0.74-1.24)	6.69 (4.84-11.41)	23.49 (14.23-30.93)	31.93 (25.29-35.80)
USA	39.0	5.40 (0.00-6.86)	3.36 (0.00-10.88)	1.16 (0.42-3.68)	10.42 (5.06-15.73)	23.71 (19.65-32.99)	29.88 (25.55-37.69)	35.98 (34.87-38.93)
Mexico	7.1	6.86 (2.12-9.30)	3.09 (0.98-5.84)	0.73 (0.31-1.26)	5.25 (5.25-5.81)	7.28 (7.28-7.28)	7.28 (7.28-7.28)	7.28 (7.28-7.28)
Central America	0.5	7.61 (0.07-9.60)	2.64 (0.05-4.63)	0.59 (0.03-0.92)	0.56 (0.56-0.56)	0.56 (0.56-0.56)	0.56 (0.56-0.56)	0.56 (0.56-0.56)
Brazil	27.8	8.21 (0.00-10.89)	3.46 (0.00-8.45)	0.79 (0.34-2.32)	14.64 (13.72-16.06)	26.21 (26.21-31.42)	33.27 (29.36-33.27)	33.27 (31.89-33.27)
Rest of South America	20.3	5.88 (0.01-9.98)	3.53 (0.01-11.08)	0.82 (0.40-6.29)	12.14 (11.21-14.65)	16.05 (16.03-16.96)	18.54 (17.62-18.90)	18.55 (17.62-18.90)
Northern Africa	=	=	=	=	=	=	=	=
Western Africa	3.1	0.02 (0.00-4.72)	0.00 (0.00-16.62)	1.29 (0.33-8.01)	1.96 (1.89-1.96)	2.10 (2.10-2.10)	2.10 (2.10-2.10)	2.10 (2.10-2.10)
Eastern Africa	33.9	4.67 (0.00-7.84)	3.27 (0.00-35.69)	2.43 (0.43-53.31)	2.98 (2.58-3.41)	5.05 (4.72-5.74)	8.17 (7.78-8.37)	8.37 (8.16-11.37)
South Africa	1.0	0.00 (0.00-3.16)	0.00 (0.00-2.71)	1.03 (0.44-1.40)	0.72 (0.72-1.02)	1.02 (0.96-1.02)	1.02 (1.01-1.02)	1.02 (1.02-1.02)
Western Europe	23.6	4.79 (0.00-5.71)	3.80 (0.00-7.40)	1.17 (0.74-2.50)	4.84 (1.61-6.98)	19.47 (19.47-19.47)	22.52 (21.54-23.49)	23.49 (23.49-23.49)
Rest of Southern Africa	63.7	5.38 (0.00-8.83)	4.42 (0.00-13.68)	1.31 (0.57-10.90)	13.17 (8.29-14.28)	24.76 (24.76-25.25)	24.76 (24.76-27.22)	24.76 (24.76-27.59)
Central Europe	19.3	5.05 (4.07-6.10)	4.60 (1.96-11.93)	1.32 (0.69-3.27)	3.56 (1.21-6.49)	9.60 (7.50-11.71)	13.55 (13.55-13.55)	17.14 (13.55-19.62)
Turkey	=	=	=	=	=	=	=	=
Ukraine Region	11.4	4.78 (3.63-5.38)	4.73 (2.87-41.12)	1.35 (0.82-13.81)	2.46 (0.00-5.20)	5.20 (5.20-5.20)	5.20 (5.20-5.20)	8.06 (5.20-8.06)
Central Asia	0.5	2.14 (0.00-4.58)	0.00 (0.00-0.00)	=	0.47 (0.47-0.47)	0.47 (0.47-0.47)	0.47 (0.47-0.47)	0.47 (0.47-0.47)
Russia region	146.1	3.39 (0.03-4.50)	6.52 (0.02-44.97)	3.34 (1.14-33.58)	2.51 (0.48-4.59)	24.36 (13.68-36.87)	49.21 (36.39-69.33)	75.28 (68.13-81.69)
Middle East	=	=	=	=	=	=	=	=
India	6.0	6.92 (6.72-7.22)	2.50 (1.38-9.07)	0.53 (0.29-1.85)	0.14 (0.14-6.08)	6.08 (6.08-6.08)	6.08 (6.08-6.08)	6.08 (6.08-6.08)
Korea region	4.3	6.09 (5.86-6.29)	4.42 (3.41-5.54)	1.07 (0.84-1.30)	2.15 (0.00-4.30)	4.30 (4.30-4.30)	4.30 (4.30-4.30)	4.30 (4.30-4.30)
China region	58.1	5.08 (0.05-7.06)	3.00 (0.04-7.64)	0.89 (0.48-4.41)	18.13 (13.20-22.02)	35.11 (32.61-36.85)	38.84 (37.77-39.79)	46.08 (41.53-48.81)
Southeastern Asia	24.5	7.19 (0.03-9.65)	4.22 (0.02-16.68)	0.83 (0.48-11.75)	6.92 (6.92-6.92)	7.30 (7.30-7.30)	7.30 (7.30-7.30)	7.30 (7.30-7.30)
Indonesia region	=	=	=	=	=	=	=	=
Japan	2.7	5.59 (1.61-6.27)	6.38 (3.33-23.76)	1.56 (0.87-18.64)	1.08 (0.00-2.46)	2.46 (2.46-2.46)	2.46 (2.46-2.46)	2.46 (2.46-2.46)
Rest of South Asia	=	=	=	=	=	=	=	=
Oceania	78.7	2.66 (0.00-7.52)	3.26 (0.00-10.71)	1.88 (0.79-4.67)	18.36 (17.71-19.58)	33.78 (30.20-39.76)	64.41 (59.14-67.85)	77.58 (74.43-81.72)

Table SI.4b | As Table SI.4a for the 2°C temperature profile.

Region	Maximum area of BECCS (Mha)	BECCS Productivity		Required Scale Factor	Area of BECCS (Mha) in optimised land-based scenario			
		Modelled	Required to match Natural		BECCS scale factor $\kappa = 1$	BECCS scale factor $\kappa = 2$	BECCS scale factor $\kappa = 3$	BECCS scale factor $\kappa = 4$
Canada	65.9	3.41 (0.00-5.26)	6.23 (0.00-19.88)	2.48 (0.81-18.40)	0.93 (0.74-1.14)	10.61 (4.89-18.12)	26.13 (17.68-34.09)	38.96 (30.97-44.66)
USA	39.0	5.93 (0.00-7.35)	3.66 (0.00-11.77)	1.15 (0.41-3.52)	9.98 (5.07-15.73)	24.77 (23.35-32.23)	33.65 (25.55-37.71)	36.34 (35.63-38.91)
Mexico	7.1	6.95 (1.78-9.63)	3.46 (1.01-6.38)	0.81 (0.33-1.36)	5.25 (4.70-5.25)	7.28 (7.05-7.28)	7.28 (7.28-7.28)	7.28 (7.28-7.28)
Central America	0.5	7.12 (0.07-9.95)	2.96 (0.05-5.06)	0.63 (0.08-1.09)	0.56 (0.41-0.56)	0.56 (0.56-0.56)	0.56 (0.56-0.56)	0.56 (0.56-0.56)
Brazil	27.8	8.25 (0.00-10.94)	3.83 (0.00-9.36)	0.88 (0.41-2.48)	14.01 (12.35-15.57)	27.68 (26.09-30.84)	29.36 (28.06-31.42)	33.27 (30.11-33.27)
Rest of South America	20.3	6.29 (0.02-10.21)	3.81 (0.01-11.79)	0.84 (0.41-5.60)	12.14 (11.19-13.08)	16.05 (16.03-16.96)	18.54 (17.62-18.90)	18.55 (17.77-18.90)
Northern Africa	=	=	=	=	=	=	=	=
Western Africa	3.1	0.02 (0.00-4.82)	0.00 (0.00-16.45)	1.50 (0.33-7.99)	1.89 (1.89-1.96)	2.10 (2.10-2.10)	2.10 (2.10-2.10)	2.10 (2.10-2.10)
Eastern Africa	33.9	4.87 (0.00-8.16)	3.64 (0.00-38.66)	2.39 (0.46-54.83)	2.98 (2.58-3.41)	4.77 (3.90-5.24)	8.17 (7.74-8.37)	8.38 (8.17-12.20)
South Africa	1.0	0.00 (0.00-3.42)	0.00 (0.00-2.95)	1.24 (0.65-1.36)	0.72 (0.72-0.72)	1.02 (1.02-1.02)	1.02 (1.02-1.02)	1.02 (1.02-1.02)
Western Europe	23.6	5.18 (0.00-6.15)	3.99 (0.00-7.80)	1.16 (0.71-2.40)	4.29 (2.37-8.77)	19.47 (19.47-19.47)	22.52 (21.54-23.49)	23.49 (23.49-23.49)
Rest of Southern Africa	63.7	5.45 (0.00-8.92)	4.85 (0.00-15.90)	1.44 (0.68-11.31)	6.59 (5.68-11.60)	24.76 (23.65-25.25)	25.12 (24.76-27.12)	25.25 (24.76-27.12)
Central Europe	19.3	5.50 (4.50-6.48)	4.70 (1.91-12.59)	1.28 (0.66-3.21)	2.65 (1.21-5.91)	8.42 (7.50-13.55)	13.55 (13.55-18.49)	19.05 (13.55-19.62)
Turkey	=	=	=	=	=	=	=	=
Ukraine Region	11.4	5.23 (3.94-5.91)	4.92 (3.02-46.34)	1.31 (0.79-14.56)	1.84 (0.00-5.20)	5.20 (5.20-5.20)	5.20 (5.20-5.20)	8.06 (5.20-8.06)
Central Asia	0.5	2.14 (0.00-4.61)	0.00 (0.00-0.00)	=	0.47 (0.47-0.47)	0.47 (0.47-0.47)	0.47 (0.47-0.47)	0.47 (0.47-0.47)
Russia region	146.1	3.70 (0.04-4.89)	6.60 (0.03-40.53)	3.04 (1.12-29.66)	2.69 (0.58-6.56)	25.92 (15.99-40.00)	60.45 (44.83-71.40)	78.97 (72.99-84.72)
Middle East	=	=	=	=	=	=	=	=
India	6.0	7.06 (6.84-7.71)	2.38 (1.43-14.27)	0.49 (0.30-2.70)	0.14 (0.14-6.08)	6.08 (0.14-6.08)	6.08 (6.08-6.08)	6.08 (6.08-6.08)
Korea region	4.3	6.65 (6.39-6.96)	4.82 (3.84-5.61)	1.05 (0.86-1.21)	2.15 (0.00-4.30)	4.30 (4.30-4.30)	4.30 (4.30-4.30)	4.30 (4.30-4.30)
China region	58.1	5.44 (0.06-7.42)	3.22 (0.04-8.05)	0.92 (0.48-4.28)	17.78 (13.89-21.67)	35.11 (32.97-36.89)	39.72 (38.84-41.01)	47.15 (42.31-49.46)
Southeastern Asia	24.5	7.01 (0.03-9.80)	4.82 (0.02-17.29)	0.98 (0.55-12.08)	6.82 (4.51-6.92)	7.30 (7.30-7.30)	7.30 (7.30-7.30)	7.30 (7.30-7.30)
Indonesia region	=	=	=	=	=	=	=	=
Japan	2.7	6.17 (2.19-6.82)	6.62 (3.58-29.58)	1.46 (0.85-16.68)	1.08 (0.00-2.46)	2.46 (2.46-2.46)	2.46 (2.46-2.46)	2.46 (2.46-2.46)
Rest of South Asia	=	=	=	=	=	=	=	=
Oceania	78.7	2.76 (0.00-7.81)	3.67 (0.00-12.17)	1.96 (0.76-5.44)	19.24 (15.74-20.25)	33.30 (30.28-35.15)	61.07 (59.03-67.85)	71.84 (63.39-77.58)

Table SI.4c | By IMAGE region for the 1.5°C temperature profile, the carbon uptake by BECCS (GtC) and the change in the land carbon store (GtC) compared to the control in the optimised land-based mitigation scenario. All entries shown as median (interquartile range).

Region	Carbon Uptake by BECCS (GtC)/Change in the land carbon store (GtC)							
	BECCS	Land	BECCS	Land	BECCS	Land	BECCS	Land
	BECCS scale factor $\kappa = 1$	BECCS scale factor $\kappa = 1$	BECCS scale factor $\kappa = 2$	BECCS scale factor $\kappa = 2$	BECCS scale factor $\kappa = 3$	BECCS scale factor $\kappa = 3$	BECCS scale factor $\kappa = 4$	BECCS scale factor $\kappa = 4$
Canada	0.02 (0.01,0.02)	0.15 (0.12,0.21)	0.77 (0.59,1.41)	-0.49 (-0.87,-0.40)	3.88 (2.58,4.61)	-2.63 (-2.89,-1.83)	6.61 (5.88,6.95)	-3.91 (-4.00,-3.52)
USA	0.70 (0.36,1.07)	1.30 (0.58,1.88)	2.76 (2.67,3.22)	0.25 (-0.37,0.44)	4.72 (4.53,5.44)	-0.30 (-0.68,0.01)	7.39 (7.26,7.51)	-1.01 (-1.15,-0.86)
Mexico	0.44 (0.41,0.48)	1.00 (0.83,1.05)	1.25 (1.21,1.32)	0.76 (0.71,0.80)	1.88 (1.82,1.98)	0.76 (0.71,0.79)	2.50 (2.42,2.64)	0.76 (0.71,0.79)
Central America	0.06 (0.05,0.06)	0.24 (0.20,0.28)	0.11 (0.11,0.12)	0.24 (0.20,0.28)	0.17 (0.16,0.18)	0.24 (0.20,0.28)	0.23 (0.22,0.24)	0.24 (0.20,0.28)
Brazil	1.72 (1.63,2.03)	5.84 (5.28,6.58)	5.21 (5.02,5.35)	4.84 (4.51,5.33)	8.05 (7.68,8.21)	4.66 (4.28,5.13)	10.80 (10.44,10.96)	4.63 (4.28,5.06)
Rest of South America	1.32 (1.20,1.47)	4.20 (3.73,4.56)	3.41 (3.29,3.55)	3.69 (3.41,3.94)	5.28 (5.10,5.48)	3.57 (3.28,3.84)	7.05 (6.81,7.32)	3.57 (3.27,3.83)
Northern Africa	=	=	=	=	=	=	=	=
Western Africa	0.02 (0.01,0.02)	17.63 (16.30,19.04)	0.06 (0.06,0.06)	17.62 (16.28,19.02)	0.09 (0.09,0.09)	17.62 (16.28,19.02)	0.12 (0.12,0.12)	17.62 (16.28,19.02)
Eastern Africa	0.12 (0.09,0.15)	1.97 (1.88,2.14)	0.50 (0.41,0.58)	1.76 (1.61,1.97)	1.11 (1.05,1.22)	1.44 (1.38,1.63)	1.48 (1.39,2.04)	1.40 (1.19,1.63)
South Africa	0.00 (0.00,0.01)	-0.13 (-0.14,-0.13)	0.03 (0.02,0.03)	-0.15 (-0.15,-0.13)	0.04 (0.04,0.04)	-0.15 (-0.15,-0.13)	0.05 (0.05,0.06)	-0.15 (-0.15,-0.13)
Western Europe	0.39 (0.10,0.59)	-0.27 (-0.43,-0.04)	3.13 (3.08,3.19)	-1.83 (-2.25,-1.61)	4.97 (4.90,5.05)	-2.15 (-2.46,-1.84)	6.69 (6.57,6.80)	-2.16 (-2.57,-1.84)
Rest of Southern Africa	1.49 (0.96,1.59)	4.87 (4.68,5.60)	4.84 (4.74,4.93)	3.57 (3.47,3.71)	7.27 (7.11,7.44)	3.57 (3.46,3.70)	9.70 (9.48,9.92)	3.56 (3.43,3.70)
Central Europe	0.24 (0.07,0.46)	-0.33 (-0.44,-0.22)	1.33 (1.09,1.61)	-0.90 (-0.96,-0.88)	2.62 (2.58,2.66)	-1.66 (-1.76,-1.07)	3.87 (3.52,4.16)	-1.71 (-1.76,-1.62)
Turkey	=	=	=	=	=	=	=	=
Ukraine Region	0.14 (0.00,0.31)	-0.13 (-0.25,-0.03)	0.64 (0.63,0.66)	-0.36 (-0.47,-0.25)	0.96 (0.95,0.99)	-0.44 (-0.47,-0.25)	1.57 (1.32,1.64)	-0.54 (-0.57,-0.47)
Central Asia	0.00 (0.00,0.00)	0.02 (0.01,0.03)	0.00 (0.00,0.00)	0.02 (0.01,0.03)	0.01 (0.01,0.01)	0.02 (0.01,0.03)	0.01 (0.01,0.01)	0.02 (0.01,0.03)
Russia region	0.15 (0.03,0.28)	0.84 (0.64,1.02)	3.15 (1.95,4.32)	-1.43 (-2.08,-0.52)	7.85 (6.43,9.22)	-3.81 (-4.38,-3.36)	13.05 (12.41,13.60)	-5.84 (-6.55,-5.49)
Middle East	=	=	=	=	=	=	=	=
India	0.00 (0.00,0.12)	0.18 (0.15,0.20)	0.25 (0.24,0.26)	0.02 (-0.01,0.12)	0.38 (0.37,0.39)	0.01 (-0.02,0.08)	0.51 (0.49,0.52)	0.01 (-0.02,0.08)
Korea region	0.13 (0.00,0.26)	-0.11 (-0.23,-0.01)	0.54 (0.53,0.55)	-0.29 (-0.36,-0.23)	0.81 (0.79,0.83)	-0.29 (-0.36,-0.23)	1.09 (1.05,1.11)	-0.29 (-0.36,-0.23)
China region	1.36 (1.03,1.61)	-0.23 (-0.43,-0.01)	5.42 (5.27,5.56)	-2.16 (-2.47,-2.07)	8.54 (8.42,8.65)	-2.63 (-3.06,-2.19)	12.10(11.61,12.49)	-3.18 (-3.23,-3.12)
Southeastern Asia	0.88 (0.86,0.90)	1.52 (1.33,1.65)	1.82 (1.77,1.85)	1.49 (1.31,1.61)	2.72 (2.66,2.78)	1.49 (1.31,1.61)	3.63 (3.55,3.71)	1.49 (1.31,1.61)
Indonesia region	=	=	=	=	=	=	=	=
Japan	0.10 (0.00,0.23)	-0.04 (-0.16,0.05)	0.47 (0.45,0.48)	-0.24 (-0.32,-0.16)	0.70 (0.68,0.72)	-0.24 (-0.32,-0.16)	0.93 (0.91,0.96)	-0.24 (-0.32,-0.16)
Rest of South Asia	=	=	=	=	=	=	=	=
Oceania	0.20 (0.19,0.27)	1.71 (1.60,1.88)	2.37 (1.62,2.94)	0.47 (-0.14,1.15)	6.17 (5.72,6.90)	-1.54 (-1.88,-1.38)	9.18 (8.51,10.26)	-2.52 (-2.81,-2.17)
Global	9.31 (7.19,11.93)	41.82 (36.12,46.59)	37.83 (35.74,41.73)	27.50 (23.51,31.04)	67.96 (65.96,71.25)	18.82 (16.30,21.23)	98.62(95.98,100.77)	12.61 (11.16,13.67)

Table SI.4d | As Table SI.4c for the 2°C temperature profile.

Region	Carbon Uptake (GtC)							
	BECCS	Land	BECCS	Land	BECCS	Land	BECCS	Land
	BECCS scale factor $\kappa = 1$	BECCS scale factor $\kappa = 1$	BECCS scale factor $\kappa = 2$	BECCS scale factor $\kappa = 2$	BECCS scale factor $\kappa = 3$	BECCS scale factor $\kappa = 3$	BECCS scale factor $\kappa = 4$	BECCS scale factor $\kappa = 4$
Canada	0.02 (0.02-0.02)	0.13 (0.09-0.17)	1.52 (0.66-2.27)	-1.03 (-1.49--0.44)	4.91 (3.69-5.73)	-3.09 (-3.46--2.62)	8.44 (7.37-9.00)	-4.65 (-4.89--4.47)
USA	0.82 (0.38-1.15)	1.42 (0.68-1.95)	3.18 (3.08-3.43)	0.16 (-0.25-0.32)	5.71 (4.97-6.04)	-0.50 (-0.83-0.04)	8.17 (8.03-8.32)	-1.06 (-1.20--0.91)
Mexico	0.40 (0.34-0.43)	1.13 (1.02-1.21)	1.27 (1.23-1.32)	0.85 (0.77-0.91)	1.91 (1.84-1.98)	0.85 (0.76-0.91)	2.55 (2.45-2.65)	0.85 (0.76-0.91)
Central America	0.05 (0.04-0.06)	0.30 (0.25-0.38)	0.11 (0.10-0.12)	0.30 (0.25-0.37)	0.17 (0.15-0.18)	0.30 (0.25-0.37)	0.23 (0.20-0.24)	0.30 (0.25-0.37)
Brazil	1.71 (1.56-1.94)	6.68 (6.14-7.48)	5.21 (5.01-5.38)	5.60 (5.21-6.15)	7.95 (7.64-8.17)	5.49 (5.09-6.06)	10.77 (10.41-11.05)	5.35 (4.99-5.88)
Rest of South America	1.30 (1.19-1.43)	4.65 (4.44-5.13)	3.52 (3.38-3.68)	4.16 (3.92-4.55)	5.46 (5.24-5.71)	4.00 (3.78-4.39)	7.29 (6.99-7.61)	3.99 (3.78-4.38)
Northern Africa	0.00 (0.00-0.00)	0.00 (0.00-0.00)	0.00 (0.00-0.00)	0.00 (0.00-0.00)	0.00 (0.00-0.00)	0.00 (0.00-0.00)	0.00 (0.00-0.00)	0.00 (0.00-0.00)
Western Africa	0.01 (0.01-0.02)	20.31 (19.00-21.68)	0.06 (0.06-0.06)	20.29 (18.99-21.66)	0.09 (0.09-0.09)	20.29 (18.99-21.66)	0.12 (0.12-0.12)	20.29 (18.99-21.66)
Eastern Africa	0.12 (0.09-0.15)	2.13 (2.04-2.28)	0.44 (0.35-0.53)	2.02 (1.77-2.12)	1.17 (1.05-1.45)	1.54 (1.43-1.73)	1.84 (1.40-2.21)	1.46 (1.13-1.65)
South Africa	0.00 (0.00-0.00)	-0.13 (-0.13--0.13)	0.03 (0.02-0.03)	-0.15 (-0.15--0.14)	0.04 (0.04-0.05)	-0.15 (-0.15--0.14)	0.06 (0.05-0.06)	-0.15 (-0.15--0.14)
Western Europe	0.37 (0.16-0.83)	-0.25 (-0.61--0.09)	3.39 (3.32-3.48)	-1.98 (-2.37--1.75)	5.40 (5.31-5.51)	-2.32 (-2.59--1.97)	7.26 (7.14-7.42)	-2.38 (-2.71--1.97)
Rest of Southern Africa	0.76 (0.71-0.94)	6.11 (5.42-6.30)	4.75 (4.40-4.94)	3.92 (3.77-4.27)	7.26 (7.07-7.47)	3.89 (3.76-3.97)	9.68 (9.43-9.98)	3.86 (3.74-3.97)
Central Europe	0.19 (0.08-0.48)	-0.28 (-0.47--0.24)	1.27 (1.18-1.82)	-0.97 (-1.11--0.92)	2.89 (2.78-3.32)	-1.76 (-1.84--1.61)	4.31 (3.80-4.55)	-1.81 (-1.87--1.76)
Turkey	0.00 (0.00-0.00)	-0.03 (-0.04--0.02)	0.00 (0.00-0.00)	-0.03 (-0.04--0.02)	0.00 (0.00-0.00)	-0.03 (-0.04--0.02)	0.00 (0.00-0.00)	-0.03 (-0.04--0.02)
Ukraine Region	0.12 (0.00-0.35)	-0.12 (-0.28--0.03)	0.71 (0.69-0.73)	-0.38 (-0.49--0.28)	1.07 (1.04-1.10)	-0.46 (-0.50--0.28)	1.72 (1.45-1.83)	-0.59 (-0.62--0.50)
Central Asia	0.00 (0.00-0.00)	0.02 (0.01-0.03)	0.00 (0.00-0.00)	0.02 (0.01-0.03)	0.01 (0.01-0.01)	0.02 (0.01-0.03)	0.01 (0.01-0.01)	0.02 (0.01-0.03)
Russia region	0.18 (0.03-0.45)	0.84 (0.52-1.01)	3.56 (2.30-4.87)	-1.67 (-2.48--0.67)	9.45 (8.11-10.49)	-4.79 (-5.18--4.38)	14.83 (14.13-15.22)	-6.67 (-7.29--6.05)
Middle East	0.00 (0.00-0.00)	0.00 (0.00-0.01)	0.00 (0.00-0.00)	0.00 (0.00-0.01)	0.00 (0.00-0.00)	0.00 (0.00-0.01)	0.00 (0.00-0.00)	0.00 (0.00-0.01)
India	0.00 (0.00-0.12)	0.21 (0.18-0.23)	0.26 (0.01-0.27)	0.18 (0.03-0.22)	0.40 (0.39-0.42)	-0.02 (-0.09-0.16)	0.54 (0.52-0.55)	-0.04 (-0.12-0.09)
Korea region	0.14 (0.00-0.29)	-0.12 (-0.26--0.01)	0.60 (0.59-0.61)	-0.32 (-0.37--0.26)	0.90 (0.88-0.92)	-0.32 (-0.37--0.26)	1.20 (1.17-1.22)	-0.32 (-0.37--0.26)
China region	1.43 (1.08-1.72)	-0.22 (-0.41-0.05)	5.78 (5.62-6.02)	-2.38 (-2.60--2.22)	9.19 (9.03-9.35)	-2.85 (-3.30--2.34)	12.95 (12.35-13.45)	-3.38 (-3.46--3.31)
Southeastern Asia	0.85 (0.63-0.89)	1.71 (1.47-1.91)	1.80 (1.66-1.86)	1.59 (1.44-1.73)	2.70 (2.49-2.79)	1.59 (1.44-1.73)	3.60 (3.32-3.72)	1.59 (1.44-1.73)
Indonesia region	0.00 (0.00-0.00)	0.90 (0.83-0.97)	0.00 (0.00-0.00)	0.90 (0.83-0.97)	0.00 (0.00-0.00)	0.90 (0.83-0.97)	0.00 (0.00-0.00)	0.90 (0.83-0.97)
Japan	0.11 (0.00-0.25)	-0.05 (-0.18-0.05)	0.52 (0.51-0.54)	-0.26 (-0.34--0.18)	0.78 (0.76-0.80)	-0.26 (-0.34--0.18)	1.04 (1.02-1.07)	-0.26 (-0.34--0.18)
Rest of South Asia	0.00 (0.00-0.00)	0.13 (0.12-0.14)	0.00 (0.00-0.00)	0.13 (0.12-0.14)	0.00 (0.00-0.00)	0.13 (0.12-0.14)	0.00 (0.00-0.00)	0.13 (0.12-0.14)
Oceania	0.21 (0.20-0.27)	1.96 (1.86-2.14)	2.13 (1.77-2.72)	0.75 (0.15-1.30)	6.34 (5.82-7.28)	-1.68 (-2.14--1.46)	9.22 (8.07-10.47)	-2.20 (-2.82--1.67)
Global	9.00 (6.95-11.62)	48.12 (41.99-52.11)	39.84 (37.27-43.80)	32.16 (27.36-35.64)	73.36 (70.87-76.54)	21.20 (18.99-23.39)	105.3 (101.7-108.7)	15.11 (13.67-16.45)