

Agricultural management effects on mean and extreme temperature trends

Aine M. Gormley-Gallagher¹, Sebastian Sterl^{1,2,3}, Annette L. Hirsch⁴, Sonia I. Seneviratne⁵, Edouard L. Davin⁵, Wim Thiery^{1,5}

- 5 ¹Department of Hydrology and Hydraulic Engineering, Vrije Universiteit Brussel, Brussels, Belgium
²Department of Earth and Environmental Sciences, KU Leuven, Leuven, Belgium
³Center for Development Research, University of Bonn, Bonn, Germany
⁴ARC Centre of Excellence for Climate Extremes, University of New South Wales, Sydney, Australia
⁵Institute for Atmospheric and Climate Science, ETH Zurich, Zurich, Switzerland

10

Correspondence to: Aine M. Gormley-Gallagher (a.gormley@ulster.ac.uk)

Abstract.

15 Understanding and quantifying land management impacts on local climate is important for distinguishing between the effects of land management and large-scale climate forcings. This study for the first time explicitly considers the radiative forcing resulting from realistic land management and offers new insights on the local land surface response to land management. Regression-based trend analysis is applied to observations and present-day ensemble simulations with the Community Earth System Model (CESM) version 1.2.2 to assess the impact of irrigation and conservation agriculture (CA) on warming trends using an approach that is less sensitive to temperature extremes. At the regional scale, an irrigation- and CA-induced acceleration of the annual mean near-surface air temperature (T_{2m}) warming trends and the annual maximum daytime temperature (TXx) warming trends were evident. Estimation of the impact of irrigation and CA on the spatial average of the warming trends indicated that irrigation and CA have a pulse cooling effect on T_{2m} and TXx, after which the warming trends increase at a greater rate than the control 20 simulations. This differed at the local (subgrid) scale under irrigation where surface temperature cooling and the dampening of warming trends were both evident. As the local surface warming trends, in contrast to regional trends, do not account for atmospheric (water vapour) feedbacks, their dampening confirms the importance of atmospheric feedbacks (water vapour forcing) in explaining the enhanced regional

Deleted: radiative

Deleted: at the top of the atmosphere

Deleted: if this model overestimates warming trends because

trends. At the land surface, the positive radiative forcing signal arising from enhanced atmospheric water vapour is too weak to offset the local cooling from the irrigation-induced increase in the evaporative fraction. Our results underline that agricultural management has complex and nonnegligible impacts on the local climate and highlight the need to carefully represent and evaluate land management in climate models.

- Deleted: s
- Deleted: account
- Deleted: for
- Deleted: to
- Deleted: climate projections

1 Introduction

According to observational and global climate model (GCM) data, temperatures associated with hot extremes have increased consistent with global anthropogenic climate change (Sillmann and Croci-Maspoli, 2009; Donat et al., 2013a, 2013b; Hartmann et al., 2013; Pendergrass and Hartmann, 2014; Fischer and Knutti, 2015). However, hot spots of accelerated warming in annual maximum daytime temperature (TXx) relative to local mean temperature (T_{2m}) simulated by climate models from phase 5 of the World Climate Research Programme's (WCRP) Coupled Model Intercomparison Project (CMIP5) are spatially inconsistent with observations (Donat et al., 2017). This is particularly the case over southeast China, South America, north America and parts of Australia and Europe. In these regions, the modelled TXx warming from the mid-twentieth century (1951–1980) to the late 20th/early 21st century (1981–2010) was greater than the modelled T_{2m} warming. In contrast to the models, the observations showed that TXx warmed at a slower rate than T_{2m} . Further analysis of the CMIP5 ensemble over central Europe by Vogel et al. (2018) highlighted that several GCMs overestimate the observed negative correlation between summer precipitation and TXx, resulting in too strong future drying and associated TXx increases under RCP8.5. This underlines the importance of a correct representation of land-atmosphere coupling for simulating changes in temperature extremes at regional scales. These discrepancies between multiple GCMs and observations raise the questions as to whether: (1) these model results can be used to reliably project changes in local temperature extremes; (2) the discrepancies remain if the rates at which warming occurs over a time period is examined, which less sensitive to outliers common in extreme temperature data than the absolute temperature difference between two time periods, as used in Donat et al. (2017); and (3) the inclusion of more processes that represent land-atmosphere coupling would enhance model skill.

Deleted: in TXx

Deleted: .

Deleted: the
Deleted: study

70 Agricultural land management techniques, including irrigation and conservation agriculture, can have a cooling effect on hot temperature extremes (Davin et al., 2014; Hirsch et al., 2017; Thiery et al., 2017, 2020; Chen and Dirmeyer, 2019; Hauser et al., 2019; Jia et al., 2019). Irrigation diverts surface and groundwater resources to agricultural land to increase crop production (Fereses and Soriano, 2007). The addition of this water to the land surface is balanced by the loss of water via runoff, deep percolation, soil storage and/or evapotranspiration (ET) (Fereses and Connor, 2004). Under drier conditions, less evaporative cooling leads to amplified warming because the energy budget becomes dominated by sensible heating instead of latent heating (Donat et al., 2017). If irrigation water is added to the surface, this increases soil moisture as well as latent heat flux over the summer months, leading to more evaporative cooling at the land surface. This irrigation-induced surface cooling, in turn, challenges the radiative forcing concept, which assumes that as radiative forcing increases (from enhanced atmospheric water vapour) so too does surface temperature (IPCC, 2001; Boucher et al., 2004).

85 Conservation agriculture (CA), which involves crop residue management, crop rotation (Carrer et al., 2018; Lombardozi et al., 2018) and minimal or no tillage (Kassam et al., 2015), can create climate feedbacks due to the presence of a crop residue over CA land that change both the radiative and hydrological properties at the surface (Davin et al., 2014). Hirsch et al. (2018) explored whether applying the no-till component of CA within the Community Earth System Model (CESM) improves the simulation of present-day climate. They found that the surface temperature response was influenced by three competing effects: (1) a surface albedo increase – which reduces the availability of energy for partitioning between the sensible and latent heat fluxes; (2) increased surface resistance (e.g. from mulch) – which reduces soil evaporation; and (3) increased soil moisture retention leading to enhanced transpiration. The local cooling response to CA was somewhat counteracted by grid-scale changes in climate over North America, Europe, and Asia because of negative atmospheric feedbacks. Grid-scale changes in climate counteracting local responses to land use change has also been demonstrated by Malyshev et al. (2015) who showed that the subgrid signal of land use change in near surface temperature was diminished by the averaging with undisturbed portions of the pixels. The importance of local-scale responses to land cover

Field Code Changed

Formatted: French

Formatted: French

Formatted: French

Field Code Changed

Field Code Changed

Field Code Changed

Deleted: Thiery et al. (2017) used the Community Earth System Model version 1.2.2 (CESM1.2.2, hereafter referred to as CESM) to investigate the influence of irrigation on temperature extremes on a global scale. They found that irrigation induced strong cooling on irrigated pixels – an average of -0.78 K compared to a no-irrigation case – during the hottest day of the year.

Formatted: Font colour: Text 1

Formatted: Font colour: Text 1

Formatted: Font colour: Text 1

Formatted: Font colour: Text 1

Formatted: Font colour: Text 1

Formatted: Font colour: Text 1

Formatted: Font colour: Text 1

Deleted: increase

Deleted: latent heat fluxes due to increased surface resistance to

Deleted: That is, the decrease in ET – both due to higher albedo and higher soil resistance – appeared to activate a decrease in cloud cover in the model that increased incoming radiation and therefore temperature. Grid-scale changes in climate counteracting local responses to land use change has also been demonstrated by Malyshev et al. (2015) who showed that the subgrid signal of land use change in near surface temperature was diminished by the averaging with undisturbed portions of the grid cells. The importance of local-scale responses to land cover change has also been indicated in observation-based studies (e.g., Mahmood et al., 2013; Li et al., 2016), yet few modelling studies offer insight on the local land surface response to land management (e.g., Paulot et al., 2018; Meier et al., 2018).

change has also been indicated in observation-based studies (e.g., Mahmood et al., 2014; Li et al., 2015), yet few global-scale modelling studies examine the local land surface response to land management (Paulot et al., 2018; Meier et al., 2018).

Using GCMs, such as CESM, to simulate land-atmosphere interactions for investigating the effects of irrigation and agricultural conversion has been criticized as insufficient (Niyogi et al., 2002). This is partly because their coarse resolution (e.g., of order 100 km) hampers their performances in describing the present-day climate at the regional scale (Jiang et al., 2016). Furthermore, economic, societal and water resource factors are ignored – a void that initiated the so-called ‘bottom-up’ approach to evaluating the effects of land-use change (Douglas et al., 2006). Regarding the applicability of the knowledge produced by GCMs, they do not provide the skill required at the spatial scale to offer practical responses at the infrastructure scale (Hossain et al., 2015). Despite these constraints, GCMs remain a prime tool for projecting changes in the climate system (Fajardo et al., 2020; Gupta et al., 2020; Hofer et al., 2020). Examples include the GCMs that are part of the latest Coupled Model Intercomparison Project (CMIP6) and used by the IPCC in consecutive assessment reports (Yazdandoost et al., 2021). However, these GCMs largely exclude agricultural management. In particular, no CMIP5 model incorporates irrigation or CA and only three CMIP6 models include irrigation, while none have CA. Pielke et al. (2011) suggested that landscape change is omitted from the CMIP5 models because the direct radiative impact of global landscape is a lower order than the radiative forcing from greenhouse gas emissions. This constitutes a reason to investigate their inclusion. That is, to distinguish between the effects of land management and other large-scale forcings such as rising CO₂ concentrations (Schultz et al., 2016), it is important to evaluate these processes in the GCMs and ultimately gain insight into the contrasts of impacts between regions under different climate regimes.

Considering the potential effects of irrigation and CA on climate (Thiery et al., 2017), it is possible that the discrepancies between climate models and observations regarding temperature changes (Donat et al., 2017) are because the models exclude the effect of agricultural management techniques on temperature.

Formatted: Font colour: Text 1

Deleted: Using GCMs, such as CESM, to simulate land-atmosphere interactions for investigating the effects of irrigation and agricultural conversion has been criticized as insufficient (Niyogi et al., 2002). This is partly because their coarse resolutions hamper their performances in describing the present-day climate at the regional scale (Jiang et al., 2016). Furthermore, economic, societal and water resource factors are ignored – a void that initiated the so-called ‘bottom-up’ approach to evaluating the effects of land-use change (Douglas et al., 2006). Regarding the applicability of the knowledge produced by GCMs, they do not provide the skill required at the spatial scale to offer practical responses at the infrastructure scale (Hossain et al., 2015). Despite these constraints, GCMs remain a prime tool for projecting changes in the climate system (Fajardo et al., 2020; Gupta et al., 2020; Hofer et al., 2020). Examples include the GCMs that are part of the latest Coupled Model Intercomparison Project (CMIP6) and used by the IPCC in consecutive assessment reports (Yazdandoost et al., 2020). However, these GCMs largely exclude agricultural management. In particular, no CMIP5 model incorporates irrigation or CA and only three CMIP6 models include irrigation, while none have CA. Pielke et al. (2011) suggested that landscape change is omitted from the CMIP5 models because the direct radiative impact of global landscape is a lower order than the radiative forcing from greenhouse gas emissions. This constitutes a reason to investigate their inclusion. That is, to distinguish between the effects of land management and other large-scale forcings such as a doubling of CO₂ (Schultz et al., 2016), it is important to evaluate these processes in the GCMs and ultimately gain insight into the contrasts of impacts between regions under different climate regimes

Deleted: Interactions between crop albedo, irrigation, cloud formation and temperature were explored in a previous study by Hirsch et al. (2017). They found that the CESM tends to produce large cloud feedbacks over Central Europe, Central North America, North Asia, and South Asia when more energy is reflected at the surface. Irrigation-induced increases in latent heat fluxes over the boreal summer months were noted and revealed more evaporative cooling at the surface. The increase in the latent heat flux led to more water vapor in the lower atmosphere, which generated low-level clouds (see also Sherwood et al., 2017). This limited shortwave radiation and hence the amount of energy available at the surface because the increased cloud cover reflected more downward shortwave radiation above the cloud layer, resulting in surface cooling. This was enhanced by a corresponding decrease in sensible heat fluxes, reflecting the decrease in the amount of energy available at the surface and of 1

Deleted: The importance of soil moisture-temperature coupling and its mechanisms controlling the regional amplification of extreme temperatures was assessed by Vogel et al. (2017) using models that contributed to the Global Land-Atmosphere Coupling Experiment (GLACE)-CMIP5 project, which aims to quantify soil moisture-climate interactions (Seneviratne et al., 2013). Vogel et al. (2017). [2]

Formatted: Font colour: Auto

Deleted: d

Deleted: cooling

Deleted: the

Deleted: overestimated warming trends

240 The goal of this study is thus to test the hypothesis that CESM version 1.2.2 overestimates warming trends in some regions because irrigation and CA are excluded. That is, warming rates are hypothesised to increase at a slower rate – showing signs of cooling, in irrigation- and CA-affected regions when climate models do account for a theoretical constant level of these land management practices. To realise this goal, the following objectives were formulated: (1) determine spatial warming rates using simulations that account for irrigation and CA and inspect whether CESM overestimates warming trends; (2) compare the observed rates of warming to the modelled rates of warming for irrigated and CA pixels, as well as non-irrigated and non-CA pixels; and (3) estimate the impact of irrigation on the spatial average of the warming rates over time for all land, selected regions, and irrigated and CA pixels. Within this framework, the novelty of this study lies in (i) an explicit focus on land management impacts on trends as opposed to the climatology; (ii) a comparison of the subgrid versus grid-scale response, offering important new insights on the local land surface land surface response to land management; and (iii) consideration of the radiative forcing resulting from realistic land management.

Deleted: It remains challenging to resolve this, however, because it is difficult to separate land management from other effects in GCMs – particularly natural climate variability (Cook et al., 2015). The question remains as to whether the inclusion of land management practices in GCMs will improve the estimation of warming rates trends.

Deleted: decline

Deleted: three

Deleted: D

Deleted: GCM

Deleted: C

Deleted: E

Deleted: (1981-2010)

Formatted: Font: 12 pt, Not Italic, Font colour: Text 1

Formatted: Font colour: Text 1

2 Materials and Methods

2.1 Irrigation and conservation agriculture implementation in CESM

255 To assess the influence of a theoretical constant level of either irrigation or CA on mean and extreme temperatures, we use the Community Earth System Model (CESM) version 1.2.2, which has contributed output to CMIP5 (Hurrell et al., 2013). The CESM atmospheric model was version 5.3 of the Community Atmosphere Model (CAM5.3) while the land surface model was version 4.0 of the Community Land Model (CLM4). Sea surface temperatures and sea ice fractions were prescribed from the data set described by Hurrell et al. (2008).

We analyse the control (1) and experimental (2) simulations presented in Thiery et al. (2017) for irrigation and in Hirsch et al. (2018) for CA. This set consists of three 5-member ensembles.

265 The first ensemble, the control (CTL), was set up to capture land-atmosphere components within a framework akin to that of the Atmospheric Model Intercomparison Project (AMIP). The period 1976-

2010 was simulated with a horizontal pixel resolution of 0.9° latitude \times 1.25° longitude. The first 5 years were discarded as spin-up, with trends evaluated for the period 1981-2010. On 1 January 1976, small random perturbations of 10^{-14} K were applied to the initial atmospheric temperature conditions. To focus on the influence of land-atmosphere interactions, rather than ocean-atmosphere feedbacks on the climate system, sea surface temperatures and sea ice fractions were prescribed from the data set described by Hurrell et al. (2008). Greenhouse gas concentrations were also prescribed from measurements, and satellite-based observations of vegetation phenology were imposed in CLM4.

The second ensemble, the irrigation (IRR) ensemble, follows an identical setup as the CTL experiment except that the interactive irrigation module in CLM4 was enabled. As described by Oleson et al. (2013), the irrigation parameterization in CLM4 divides the cropland area of each grid cell into non-irrigated and irrigated fractions corresponding to the portions that are equipped for irrigation – in accordance with Siebert et al.'s (2005) global map of irrigated areas (Figure 1a). The area of irrigated cropland in each grid cell is assigned as the smaller of the grid cells total cropland area and its area equipped for irrigation. What remains of the cropland area in the grid cell is regarded as non-irrigated cropland. It is important to note that implementation of transient irrigation was technically not possible in the CESM version 1.2, despite transient area equipped for irrigation data being available (Siebert et al., 2005), and therefore trends in the forcing are not considered.

The third ensemble, the CA ensemble, also follows the CTL experiment setup, but in this case the most likely distribution of CA was applied based on the CA dataset developed by Prestele et al. (2018). By splitting the existing CLM crop plant functional types (PFT) into a fraction under conservation agriculture and a fraction under conventional management, both forms of management are possible within a grid cell. Although the crop residue is assumed present all year, the implementation ensures that the increased soil albedo effect on the total surface albedo is dampened during the growing season by the inclusion of canopy cover (Hirsch et al., 2018). Implementation of transient CA, however, was not possible due to data limitation as only a static CA map was available; hence we study a theoretical constant level of CA.

310 To examine heterogeneous influences within grid cells, subgrid tiles that represent local physical, biogeochemical, and ecological characteristics – and therefore local (subgrid) influences of irrigation and CA – were evaluated against regional (grid-scale) influences. Up to 21 surface tiles may occur within one grid cell in CLM4, including glacier, wetland, lake, urban, bare soil and 16 PFTs. For subgrid irrigation influences, all tiles are placed on one single soil column, except for the irrigated crop tile. Separating the soil columns in this way allows the soils underneath irrigated and rainfed crop tiles to have individual
315 responses to atmospheric forcing (Schultz et al., 2016). Therefore, the subgrid-scale difference is the irrigated crop tile minus the rainfed crop tile. For subgrid CA influences, using the PFT-level outputs from CLM, it is possible to examine the subgrid-scale effect by subtracting the conventionally managed crop tiles from the CA crop tiles.

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Deleted: ¶

320 In addition, land masks were used to define and analyse: (1) all land pixels; (2) irrigated pixels only (where grid cells have a nonzero irrigated fraction); (3) CA pixels (the grid cells with a nonzero CA fraction) and (4) those regions of the Special Report on Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation (SREX) (IPCC, 2013) where irrigation and CA is extensive (Figure 1). The spatial points outside these masks as well as missing values in the observations
325 were excluded (as ‘NaN’ values). These masks were applied to the investigations undertaken in this study. As the observational datasets (see below) were remapped to the model grid, this meant the same land masks (excluding Antarctica) could be used for each dataset.

Formatted: Font: 12 pt, Font colour: Black, Pattern: Clear

Deleted: ¶

2.2 Observational datasets

330 For evaluation purposes, observational datasets for annual mean T_{2m} with a spatial resolution of $0.5^\circ \times 0.5^\circ$ for the same time period were obtained from the Climate Research Unit (CRU) (Harris et al., 2014). Annual mean TXx observational datasets were obtained from the daily Global Historical Climatology Network extremes data set (GHCNDEX) (Donat et al., 2013a) and the Hadley Centre extremes data set (HadEX2) (Donat et al., 2013b) with a spatial resolution of $2.5^\circ \times 2.5^\circ$. These observational products were
335 regrided to the CESM resolution using second-order conservative remapping (Jones, 1999). Thiery et al. (2017) and Hirsch et al. (2018) previously evaluated how the IRR and CA experiments alter the skill

Deleted: All

of CESM simulations (in terms of their agreement with observations). Thiery et al. (2017) demonstrated
340 that including irrigation has a small yet robust beneficial effect on the representation of TXx and T_{2m} in
CESM over irrigated and all land pixels. By including CA, Hirsch et al. (2018) showed a general
improvement in the simulation skill over MED for TXx and T_{2m} and enhanced skill for T_{2m} over WNA,
CNA, and CEU.

345 Observational data for the surface radiative temperature (T_s) at the subgrid scale were obtained from the
E-OBS European CDG dataset for 1981-2010 over MED pixels. As a regional dataset, it has a higher
spatial resolution and therefore enabled a skill of the models with respect to the local effects of land
management. The E-OBS data were regridded to the CESM resolution using bilinear remapping.

Formatted: Font colour: Text 1

Formatted: Font colour: Text 1, Subscript

Formatted: Font colour: Text 1

350 2.3 Statistical analysis

The warming rate β was calculated using Sen's slope approach (Sen, 1968) based on the time and
temperature values in each grid cell. This means that at each longitude and latitude point on land, there
are 30 time measurements (1981-2010) with an associated temperature measurement (for each annual
mean T_{2m} and TXx). Therefore, there are $30 \times 29/2$ possible pairs of sample points, rendering 435 pairs for
355 each location.

Annual TXx and T_{2m} values averaged across all land pixels and all irrigated pixels were computed for the
CTL, IRR and CA ensemble means, as well as the GHCNDEX (TXx), HadEX2 (TXx) and CRU (T_{2m})
observations. A Sen's slope regression analysis was then carried out on the spatial mean temperatures of
360 TXx and T_{2m} change over time (1981-2010) for (a) all pixels, (b) irrigated pixels and (c) CA pixels only,
for both observations and the model ensembles.

The spatial mean warming rate across all (land or irrigated) pixels was also calculated. Additionally, all
pixels within the SREX regions where irrigation is extensive (Thiery et al., 2017) –WNA, CNA, MED,
365 WAS, SAS, SEA and EAS – were selected and their spatial means determined and examined. The SREX

regions where CA is extensive (Hirsch et al., 2018) were also examined in greater detail. These include WNA, CNA, MED, SSA, CEU and SAU (Figure 1).

3 Results

3.1 Model Evaluation

370 First, we explore how the existing CESM climate simulation skill (i.e., how well the simulated and observed trends agree) is altered in IRR and CA relative to the skill obtained in the CTL. The model biases and spatial root mean square error (RMSE) values relative to the warming trends of the T_{2m} and TXx global observational products are provided in Table 1. For the IRR ensemble, T_{2m} warming trends are overestimated by $\sim 0.001 \text{ K yr}^{-1}$ across irrigated pixels, whereas over CA pixels T_{2m} warming trends are overestimated by $\sim 0.002\text{-}0.004 \text{ K yr}^{-1}$ in both the CA and CTL ensemble. On average, the CTL, IRR and CA ensembles overestimate TXx warming trends by $\sim 0.007\text{-}0.03 \text{ K yr}^{-1}$ over all land pixels. Over irrigated pixels, the CTL and IRR ensemble overestimate TXx by $\sim 0.008\text{-}0.013 \text{ K yr}^{-1}$. Over CA pixels, the CTL and CA ensemble overestimate TXx by $\sim 0.006\text{-}0.013 \text{ K yr}^{-1}$. This means that while T_{2m} warming rates have a slight low bias on average over all land and partially over irrigated areas, TXx warming trends are consistently too high over all land, irrigated and CA areas.

380 Second, to investigate how the uncertainty between the different irrigation and CA estimates of warming trends influences simulation skill, we examine the added value of including irrigation and CA for TXx and T_{2m} over the regions where irrigation and/or the CA extent is greatest, as well as over global land, global irrigated land and global CA land (Figure 2). The added value is evaluated by calculating the absolute change (experiment minus control) in the spatial RMSE. Accounting for irrigation improves the simulation skill for trends over MED, WAS and SAS for T_{2m} and over MED, WAS, SAS and SEA for TXx (with HadEX2 as reference product). For WNA, CNA and EAS, the added value is negative or limited for both temperature metrics. Accounting for CA improves the simulation skill over CNA, CEU and SAU for the T_{2m} and both TXx observational products and over the MED for the T_{2m} and the TXx HaxEX2 observational products. For WNA, skill is reduced for all CA estimates. If we consider the grid

Deleted: ¶
This section
Deleted: s

Deleted: The CTL ensemble underestimates the mean T_{2m} warming trends by $\sim 0.004 \text{ K yr}^{-1}$ over all land pixels and $\sim 0.003 \text{ K yr}^{-1}$ over irrigated pixels. In contrast, f

Formatted: Font: (Default) Cambria Math

Formatted: Font: (Default) Cambria Math

Deleted: CESM

Deleted: also

Deleted: s

Formatted: Font: (Default) Cambria Math

Deleted: over all land pixels in the three ensembles

Deleted: .

Formatted: Font: (Default) Cambria Math

Deleted: over irrigated pixels in the CTL and IRR ensembles and

Formatted: Font: (Default) Cambria Math

Deleted: over all CA pixels in the CTL and CA ensemble

Deleted: ¶

Deleted: T

cells where the land fraction within the CESM exceeds 50% (“all land”) or just the grid cells that have a nonzero irrigation (“Irrigated land”) is present, there is added value for T_{2m} observational product over all land and the grid cells where irrigation has been applied. There is limited skill improvement for the
410 TXx HadEx2 observational product. For the CA simulations, if we consider all land and the grid cells with a nonzero CA fraction (“CA land”), the model skill improves for the T_{2m} observational product.

415 Third, we explore how the CESM climate simulation skill is altered in the subgrid-scale irrigation (IRR_{SUB}) and CA crop tiles (CA_{SUB}) relative to the skill obtained in the conventionally managed (CM) and rainfed crop tiles (RAIN) in the MED region. The model biases and spatial RMSE values relative to the warming trends of the T_S observational product are provided in Table 2. For IRR_{SUB}, T_S warming trends are overestimated by $\sim 0.004 \text{ K yr}^{-1}$ across irrigated MED pixels, which is an improvement in terms of bias when compared to the subgrid-scale data that does not account for irrigation (i.e., RAIN). However, according to the change in the spatial RMSE, accounting for irrigation does not improve the
420 simulation skill for trends over MED irrigated pixels. This is likely because RMSE is more sensitive to outliers – whereas the bias is based on the spatial mean.

3.2 Impact of Irrigation and Conservation Agriculture on Mean and Extreme Warming Trends

425 Neither irrigation nor CA has a cooling effect on T_{2m} and TXx warming rates in irrigated/CA or non-irrigated/CA regions (Figure 3 and Table 3). The results suggest a slight irrigation- and CA-induced acceleration of the annual T_{2m} and TXx warming trends, rather than the hypothesised cooling. For instance, irrigation induced an increased T_{2m} warming rate of 0.0023 K yr^{-1} on average over land and 0.004 K yr^{-1} across all irrigated pixels. To put these increases into context, the mean T_{2m} CRU observed warming trend over irrigated pixels was 0.029 K yr^{-1} .

430 When the annual T_{2m} and TXx temperatures are spatially averaged for each ensemble, the IRR and CTL simulations both overestimate the observed values for irrigated pixels (Figure 4a and 4b), and the CA and CTL simulations both overestimate the observed values over CA pixels (Figure 4c and 4d). However, the

Formatted: Font colour: Auto, Pattern: Clear

Moved (insertion) [1]

Deleted: 2

Deleted: a and 3c

Deleted: I

Deleted: (Table 2)

Deleted: For

Deleted: As a result of CA, T_{2m} warming rates increased by 0.002 K yr^{-1} on average over land and 0.002 K yr^{-1} across all CA pixels (Table 2). The mean T_{2m} CRU observed warming trend over CA pixels was 0.024 K yr^{-1} .

Deleted: ¶

Deleted: Likewise, no cooling effect of irrigation or CA was evident on TXx warming rates in irrigated/CA or non-irrigated/CA regions, according to the warming rates generated (Figure 3b and 3d). Irrigation induced an increased TXx warming rate of 0.006 K yr^{-1} on average over land and 0.004 K yr^{-1} across all irrigated pixels (Table 2). The mean TXx observed warming trend over irrigated pixels was 0.022 K yr^{-1} for HadEX2 and 0.025 K yr^{-1} for GHCNDEX. Due to CA, TXx warming rates increased by 0.003 K yr^{-1} on average over land and 0.005 K yr^{-1} across all CA pixels (Table 2). The mean TXx observed warming trend over CA pixels was 0.033 K yr^{-1} for HadEX2 and 0.031 K yr^{-1} for GHCNDEX. The results suggest a slight irrigation- and CA-induced acceleration of the annual T_{2m} and TXx warming trends, rather than the hypothesised cooling.¶

Moved up [1]: The results suggest a slight irrigation- and CA-induced acceleration of the annual T_{2m} and TXx warming trends, rather than the hypothesised cooling.

Deleted: 3.3 Spatial average¶
CONSIDER CHANGING THE SECTIONS – SPATIAL AVERAGE IMP FOR ALL¶

The results (as shown in Table 1 especially) are difficult to reconcile with the statements made in the abstract and conclusions. Looking at Table 1, if the smallest RMSE (or the anomalies closest to zero) are considered, the Control simulation is better $\sim 2/3$ of the time. The abstract says, “our results underline... the need to account for land management in climate projections”. Surely the opposite is true, as the Control scenario does better by the measure most used to assess model skill. Even within the results, there appear to be contradictions. Line 218: “the impact of irrigation and CA on the modeled spatially averaged temperatures... is an overall cooling effect”. Line 223: “for the IRR and CA models... the spatially averaged T_{2m} and TXx warming rates are higher than those of the CTL model” .¶

impact of irrigation and CA on the modelled spatially averaged temperatures improves the closeness to that of the observations, i.e. there is an overall ~~there is an overall decrease in absolute temperature~~ (Figure 4a-d), which ~~aligns~~ with current theory (Kueppers et al., 2007; Saeed et al., 2009; Kueppers and Snyder, 2012; Thiery et al., 2017, 2020; Hirsch et al., 2018).

What these results show in addition is, for the IRR and CA models – for all land, irrigated and CA pixels, the spatially averaged T_{2m} and TXx warming rates (~~the slopes~~) are higher than those of the CTL model. Therefore, rather than continuous cooling, there is evidence in Figure 4 of a pulse cooling phase during the spin-up years (Smith et al., 1998), after which the T_{2m} and TXx warming trends increases at a greater rate than the control simulations.

In the case of CA, ~~because crop residue is more likely to be applied during the summer/dry season (when TXx is typically recorded) to reduce evaporation (Figure 4l and Figure 6f), energy is shifted to the sensible heat flux (SHF) (Figure 4h and Figure 6j), increasing TXx (Figure 4d). The SHF response is not always consistent with the decrease in the latent heat flux (LHF) (Figure 6h), with some increases over Eastern South America, Eastern North America, parts of Europe and Southeast Australia.~~

In the case of irrigation, the response also suggests two competing effects: (1) there is more water at the surface, so the energy budget shifts to the LHF (Figure 4i, Figure 6g and 6i), resulting in evaporative cooling (Figure 4a and 4b); and (2) because irrigation globally adds $418 \text{ km}^3 \text{ yr}^{-1}$ of moisture to the atmosphere (Thiery et al., 2017) and as water vapour acts as a greenhouse gas (GHG), it traps outgoing longwave radiation, radiating it back to the Earth's surface as downward longwave radiation (Figure 4m), resulting in increased T_{2m} and TXx warming trends (Figure 4a and 4b). The first effect appears more pronounced than the second due to the net cooling in Figure 4a and 4b. This means that despite the water vapour (acting as a GHG) increasing downward radiation and the overall energy budget thus increasing, most of it still goes to the latent heat flux leading to a net reduction in temperature (as compared to a situation without irrigation, where the sensible/latent ratio is more in favour of the latter). The limited warming effect of irrigation on atmospheric temperatures through water vapour forcing is

- Deleted: cooling-effect
- Deleted:
- Deleted: is consistent
- Field Code Changed
- Field Code Changed
- Deleted: ¶
- Deleted: the two paragraphs starting line 228 speculate that latent and sensible heat partitioning and changes in ET are responsible for the differences between CA and IRR, and the Control. But instead of exploring these and showing how the latent heat changes, there is just references to previous papers. I.e. it is not a result, it is a summary of previously published research.
- Deleted: ¶
- Formatted: Justified, Line spacing: 1.5 lines
- Deleted: the results indicate two opposing effects: (1) increased surface albedo associated with no tillage reduces the solar energy absorbed by the surface, inducing cooling (Davin et al., 2014); and (2) the presence of a crop residue over CA land can decrease soil evaporation, inhibiting energy partitioning to the latent heat flux, leading to surface warming (Hirsch et al., 2017). For TXx (Figure [3])
- Formatted: Not Highlight
- Formatted: Not Highlight
- Formatted: Not Highlight
- Formatted: Not Highlight
- Formatted: Not Highlight
- Formatted: Font: (Default) Times New Roman
- Formatted: Font: (Default) Times New Roman
- Formatted: Font: (Default) Times New Roman
- Formatted: Font: (Default) Times New Roman
- Formatted: Font: (Default) Times New Roman
- Formatted: Font: (Default) Times New Roman
- Formatted: Font: (Default) Times New Roman
- Formatted: ... [4]
- Deleted: ¶
- Deleted: latent instead of sensible heat flux
- Formatted: Not Highlight
- Formatted: Not Highlight
- Formatted: Not Highlight
- Deleted: c
- Formatted: Not Highlight
- Deleted: es
- Formatted: Not Highlight

consistent with earlier GCM studies inputting more than twice the amount of water vapour into the atmosphere through irrigation ($32500 \text{ m}^3 \text{ s}^{-1}$ or $1026 \text{ km}^3 \text{ yr}^{-1}$) and finding limited radiative forcing (Boucher et al., 2004; Sherwood et al., 2018).

We further investigate the potential warming of the Earth System irrigation-induced enhanced atmospheric water vapour by computing the top-of-atmosphere net radiation ($R_{n,TOA}$) in the CTL and IRR ensembles over the 1981-2010 period (Figure 5). As both ensembles employ prescribed, transient sea surface temperatures, the difference in $R_{n,TOA}$ is a measure of irrigation-induced radiative forcing. The area-weighted global average $R_{n,TOA}$ is 0.4961 W m^{-2} for the CTL ensemble (Figure 5a) and 0.5450 W m^{-2} for the IRR ensemble. The radiative forcing from irrigation is therefore 0.0489 W m^{-2} , at least an order of magnitude smaller compared to other combined anthropogenic forcings over this period (IPCC, 2013) and consistent with previous estimates (Boucher et al., 2004; Sherwood et al., 2018). The positive radiative forcing is mainly located over South Asia, and partially offset by negative forcing over central Asia, Greenland and Antarctica (Figure 5b). Breakdown of the irrigation-induced $R_{n,TOA}$ change into the shortwave and longwave components shows that the forcing is dominated by the longwave signal ($+0.0583 \text{ W m}^{-2}$), with the shortwave signal even showing signs of a slight albedo increase (-0.0094 W m^{-2}), presumably from enhanced low-level cloud cover (Sherwood et al., 2018). The additional water vapor in the atmosphere and associated longwave trapping in CESM can thus explain the small, positive radiative forcing contributing to Earth System warming and associated enhanced near-surface temperature trends in irrigated regions (Figure 4a-b), but at the land surface this signal is too weak to offset the local pulse cooling from the irrigation-induced increase in evaporative fraction.

3.4 Subgrid-Scale Impacts

Our results indicate a subgrid-scale cooling effect of irrigation on T_s warming trends that is more distinct and spatially consistent over irrigated pixels than grid-scale effects (Figure 6a versus Figure 7a). T_s warming trends on irrigated tiles are on average -0.008 K yr^{-1} (-24%) lower than their rainfed counterparts, whereas the trends are on average 0.001 K yr^{-1} (+11%) higher on the grid cell level over

Field Code Changed

Deleted: ¶

Formatted: Font: 12 pt, Font colour: Auto

Deleted: (Figure 5b)

Formatted: Font colour: Text 1

Deleted: c

Deleted: surface radiative temperature (

Deleted:)

Deleted: s

Deleted: 6

Deleted: and

Deleted: 6

Deleted: c

575 irrigated land (Table 3). The subgrid-scale influences of irrigation on ET rates over irrigated tiles were also pronounced as they increased by 0.653 mm yr^{-1} in comparison to rainfed tiles (Figure 7c and Table 3). The subgrid-scale influences of CA on T_s warming trends are smaller in comparison to irrigation, with only a 0.001 K yr^{-1} (-3%) dampening of warming trends and ET rates increased by 0.083 mm yr^{-1} (46%), relative to their conventionally managed counterparts (Figure 7b and 7d and Table 3).

580 The cooler warming trends from irrigation at the subgrid-scale (Figure 7a) occurs where the ET rate increases (Figure 7g) as well as the latent heat flux (Figure 7e), suggesting the cooling is due to an increase in the latent heat flux, which is consistent with Cook et al. (2015) and Thiery et al. (2017). The heightened grid-scale T_s warming trends (Figure 6a) generally align with a greater TMQ flux (Figure 6c) and 585 increased T_{2m} warming trends over irrigated pixels (Figure 3a), which signifies the longwave radiation trapping potential of the additional atmospheric water vapour. As the impact on trends is small (e.g. T_{2m} and T_s warming trends increased, respectively, by 0.004 K yr^{-1} and 0.001 K yr^{-1} across irrigated pixels), the finding is in agreement with Sherwood et al. (2018) who showed that additional water vapour has a small impact on global warming potential mainly because it rains out before reaching the altitudes needed to significantly contribute to the greenhouse effect. These findings thus support the concept of radiative forcing and the proviso that, at the land surface, the water vapour signal does not offset local cooling from the irrigation-induced increase in evaporative fraction, as described for Figure 4 and 5 and previously proposed by Boucher et al. (2004). However, because the subgrid-scale T_s trends, in contrast to grid-scale trends, are computed within the same ensemble and thus do not account for atmospheric (water vapour) 595 feedbacks, the sign reversal of irrigation-induced impact on grid-scale and subgrid-scale T_s trends confirms the importance of atmospheric feedbacks (water vapour forcing) in explaining the increased grid-scale T_s and T_{2m} trends.

600 When spatially averaged, over all pixels, the T_s warming trends at the subgrid-scale show no evidence of a pulse cooling phase due to irrigation (Figure 8c), which is in contrast the results over irrigated pixels – where there is both a cooling effect on T_s and a dampening of T_s warming trends (Figure 8a). This contrast is likely due to a combination of the remote effects of irrigation, the larger contribution of natural

- Deleted: 2
- Deleted: 6
- Deleted: c
- Deleted: 2
- Deleted: ¶
- Deleted: 6
- Deleted: 6
- Deleted: 2
- Deleted: 6a
- Deleted: 6g
- Formatted: Not Highlight
- Formatted: Not Highlight
- Deleted: 6c
- Formatted: Font colour: Text 1
- Deleted: total atmospheric column water (
- Deleted: e
- Deleted: 4a and

- Formatted: English (US)
- Deleted: ¶
- Formatted: Font: 1 pt, Bold, Font colour: Auto, Pattern: Clear
- Deleted: 7
- Deleted: c
- Deleted: 7

variability and an increased relative contribution of other components when considering all land pixels (Puma and Cook, 2010; Cook et al., 2015; De Vrese et al., 2016; Thiery et al., 2017).

625 Regarding CA, the slight overall warming of T_s temperatures (Figure 8b) as well as the increase in T_s warming trends over CA pixels for the MED region (Figure 8f) is possibly because of the decrease in soil evaporation as a result of crop residue over CA land (Figure 6f), inhibiting energy partitioning from the SHE (Table 3). The cooling of T_s temperatures over all land pixels (Figure 8d) and the slight decline in T_s warming trends over CA pixels (Figure 8b and Table 3), however, suggests that the effect of increasing surface albedo and thus reducing the solar energy absorbed by the surface is dominant. Additionally, the close correspondence between CA and CM (Figure 8b) may reflect that the temperature response spatially is both positive and negative depending on which mechanism dominates and therefore the spatial aggregation for all CA and all CM pixels globally loses this (Figure 8d).

635 4 Discussion

This study examined the hypothesis of whether excluding a theoretical constant level of irrigation and CA contributes to the overestimation of warming by an Earth System Model. A Sen's slope model was built and applied to ensemble simulations from the Community Earth System Model that include irrigation parameterization to determine if there are spatiotemporal patterns and why they exist. This unexpectedly showed that warming trends are not dampened due to either irrigation or CA, except for the subgrid-scale effect of irrigation on the warming trends of T_s .

The key findings of this investigation are a net cooling effect of irrigation and CA on the modelled spatially averaged T_{2m} and TXx , but, rather than continuous cooling, the warming trends showed a pulse cooling phase, after which the sensitivity to climatic change remains. Under irrigation, the opposing effects are the result of: (1) evaporative cooling; and (2) atmospheric water vapour strengthening the greenhouse effect. Under CA, the contrasting effects are due to: (1) cooling from a tillage-induced increase in surface albedo; and (2) reduced soil evaporation due to the presence of crop residue, limiting

- Formatted: Not Highlight
- Deleted: slight overall warming of
- Formatted: Not Highlight
- Formatted: Not Highlight
- Formatted: Not Highlight
- Deleted: temperatures
- Formatted: Not Highlight
- Formatted: Not Highlight
- Deleted: 7
- Deleted: b
- Deleted: to the latent heat flux
- Formatted: Not Highlight
- Formatted: Not Highlight
- Formatted: Not Highlight
- Deleted: 0
- Deleted: 7
- Formatted: Not Highlight
- Formatted: Not Highlight
- Deleted: the subgrid-scale response
- Deleted: 7
- Deleted: o
- Deleted: 7
- Deleted: and

energy partitioning to the latent heat flux. At the subgrid-scale, there was both a cooling effect on T_s and in the dampening of warming trends. This implies that enhanced evaporative cooling is the dominant driver of the subgrid-scale temperature trends.

665 Although this study was constructed with great care and built on a state-of-the-art modelling suite, several future developments could improve understanding of the impact of irrigation and CA on climate. Firstly, the quality of the model(s) could be improved by using transient irrigation and CA extents and new land cover datasets from the 6th phase of the Coupled Model Intercomparison Project (CMIP6) (Lawrence et al., 2016). In this study, a static irrigation map for the year 2000 was used for the whole simulation period.

Formatted: English (UK)

Formatted: English (UK)

Formatted: English (UK)

670 This likely contributes to our results being conservative. If, for instance, irrigation expands over time, the cooling effect may become stronger and thus affect the warming trends. Furthermore, the extent to which the increase in surface albedo (i.e. the first competing effect of CA) affects the sensible and latent heat fluxes partly depends on soil moisture, which too is not static. Also, CMIP6 experiments are based on annual emissions, whereas CMIP5 was based on decadal emissions and CMIP6 models were updated with irrigation-related features and land cover maps that incorporate irrigation and CA expansion over time (Goddard et al., 2013; Miao et al., 2014; Boer et al., 2016; Meinshausen et al., 2017; Stouffer et al., 2017). CMIP6 models may therefore improve the dynamics between irrigation, CA and climate change, provided that they represent these land management techniques in their surface schemes.

Formatted: English (UK)

Formatted: French

Formatted: English (UK)

680 The second consideration is that all simulations used in this study (5 control, 5 irrigation and 5 CA) were from a single model. Ensembles completed as such with the same model but different simulations (i.e. based on different initial conditions) characterise the uncertainty associated with internal climate variability only, while multi-model ensembles also account for the impact of model differences (Tebaldi and Knutti, 2007; Knutti et al., 2010). This limitation can impact cloud uncertainties. Hirsch et al. (2017) found that the CESM tends to produce large cloud feedbacks over Central Europe, Central North America, North Asia, and South Asia when more energy is reflected at the surface. Irrigation-induced increases in latent heat fluxes led to more water vapor in the lower atmosphere, which generated low-level clouds (see also Sherwood et al., 2017). This limited shortwave radiation and hence the amount of energy available

690 at the surface because the increased cloud cover reflected more downward shortwave radiation above the
cloud layer, resulting in surface cooling. This was enhanced by a corresponding decrease in sensible heat
fluxes, reflecting the decrease in the amount of energy available at the surface and/or the increase in latent
heating. The impact of cloud cover combined with land management change remains challenging to
resolve. Therefore, this study should ideally be repeated with other models. Donat et al. (2017), for
instance, conducted their study on 20 CMIP5 models, but these models did not incorporate irrigation and
695 CA.

Thirdly, irrigation and CA are the only agricultural management practices considered in this study (and
done so individually), whereas other agricultural management practices have been shown as impactful
(Luyssaert et al., 2014; Erb et al., 2016, 2018). Trend analysis of integrated land management practices
700 could affect the outcome if there is a lumped effect. Building an additional stochastic model could account
for variations in the distribution of the impact of land management practices on warming trends. This
would enable sensitivity analyses to ascertain the relative importance of irrigation and CA to the total
warming trends (based on all land management practices), as well as the relative contributions of the
uncertainty sources (model input, parameter, structure) to the total uncertainty in the model output.

705
The final consideration is whether regression-based models are suitable for analysing changes in highly
variable climate data, particularly annual extreme temperature data (von Storch, 2006). Essentially, the
regression slope blends forced temperature change and variability, to provide an estimation of the
temperature variation over time – within which variance can be lost due to noisy data. Whether the TXx
710 and T_{2m} temperatures were first spatially averaged and then the slope retrieved or if each slope was
estimated for each pixel and then the overall trends examined, the outcome remains. This is unsurprising
considering that in the spatial averaging the noise contributions are averaged out, while the individual
regression data suffers from the variance loss related to regression. However, when applied to over 60
years of observational data, the regression model used in this study showed similar trends to using the
715 difference between the past and the present average temperatures (not shown). This implies that the
irrigation and CA-inclusive climate system may require a longer timeframe (than the 30 years plus a 5-

Deleted: gridcell

year spin-up period used) for trends to overtake the natural variability. Additionally, rather than aggregating all months, trends during individual months or seasons could be examined. This can affect, for instance, the influence of irrigation on T_s , which has a clear seasonal pattern, with more cooling during the driest and/or hottest months (Thiery et al., 2017). A smaller magnitude in TXX response to CA at the subgrid-scale has also been noted during the summer season due to a larger leaf area index (LAI) reducing soil surface exposure and thus the contrast between CA and conventionally managed crops (Hirsch et al., 2018). Furthermore, the implementation of CA within CESM does not capture crop planting and harvesting cycling (Davin et al., 2014), which would affect the LAI of the crop and potentially the effect of CA on surface climate.

5 Conclusion

In this study the impact of a theoretical constant level of irrigation and CA on warming trends in global climate and climate extremes was assessed for the period of 1981–2010 using the Community Earth System Model. A Sen’s slope regression-based analysis was performed to compute spatial-explicit warming trends and spatially averaged warming trends. Insight into how modelled temperature is affected in its median by irrigation and CA over time was provided.

An irrigation- and CA-induced acceleration of the annual T_{2m} and TXX warming trends was evident. Estimating the impact of irrigation and CA on the spatial average of the warming trends indicated that irrigation and CA have a pulse cooling effect on T_{2m} and TXX, after which warming trends increased at a greater rate than the control simulations. This differed at the subgrid-scale under irrigation where surface temperature cooling and the dampening of warming trends were both evident. Therefore, irrigation-induced evaporative cooling is a more dominant effect at the local level than the strengthening of the greenhouse effect at regional scales as a result of enhanced atmospheric water vapour.

A model evaluation demonstrated that the simulations accounting for irrigation and CA satisfactorily reproduce observed warming trends in T_{2m} , but not the trends in temperature extremes of TXX. This signifies that the GCMs have more trouble representing the greater variability in the extreme

745 temperatures, compared to that of the mean annual temperature, and that the Sen's slope models are more suited to the blended variability inherent to annual mean temperatures.

The findings overall provide valuable context on how model complexity can impact the simulation of trends and emphasise the need for a more in-depth evaluation of the sensitivity of future climate projections to irrigation and CA-induced temperature changes. A sensitivity analysis, using transient irrigation and CA extents, as well as additional land management techniques, within coupled climate models based on CMIP6 output, is recommended. In this way, the variance can be approximated and the relative contributions of the uncertainty sources to the total uncertainty in the model output, as well as the relative importance of irrigation and CA to the total warming trends, can be quantified and compared. If the fundamental uncertainties relating to model structure dominate, then a more detailed analysis than the regression approach used in this study is suggested. This will support decision-making on the incorporation of agricultural management processes in future GCM projects.

Deleted: and

Deleted: model

Deleted: This will support decision-making when planning land management strategies that combine resource use efficiency with climate change adaptation and mitigation, enabling sustainable intensification of land management to meet mitigation targets and future demand for food, fuel, fibre, and water.

Acknowledgments

We thank Prof Piers Forster and Dr Chris Smith at the University of Leeds for their valuable discussions and insight on the theoretical outcomes of this project.

A. L. Hirsch is supported through funding from the Australian Research Council (ARC) Centre of Excellence for Climate Extremes (CE170100023).

Deleted: L

References

- Boer, G. J., Smith, D. M., Cassou, C., Doblus-Reyes, F., Danabasoglu, G., Kirtman, B., Kushnir, Y., Kimoto, M., Meehl, G. A., Msadek, R., Mueller, W. A., Taylor, K. E., Zwiers, F., Rixen, M., Ruprich-Robert, Y., and Eade, R.: The Decadal Climate Prediction Project (DCPP) contribution to CMIP6, Geoscientific Model Development, doi: 10.5194/gmd-9-3751-2016, 2016.
- Boucher, O., Myhre, G., and Myhre, A.: Direct human influence of irrigation on atmospheric water vapour and climate, *Climate Dynamics*, doi: 10.1007/s00382-004-0402-4, 2004.

- Carrer, D., Pique, G., Ferlicoq, M., Ceamanos, X., and Ceschia, E.: What is the potential of cropland albedo management in the fight against global warming? A case study based on the use of cover crops, *Environmental Research Letters*, 13 044030, doi: 10.1088/1748-9326/aab650, 2018.
- 780 [Chen, L. and Dirmeyer, P. A.: Global observed and modelled impacts of irrigation on surface temperature. *International Journal of Climatology*, 39: 2587-2600, 2019.](#)
- Cook, B. I., Shukla, S. P., Puma, M. J., and Nazarenko, L. S.: Irrigation as an historical climate forcing, *Climate Dynamics*, doi: 0.1007/s00382-014-2204-7, 2015.
- 785 Davin, E. L., Seneviratne, S. I., Ciais, P., Ollio, A., and Wang, T.: Preferential cooling of hot extremes from cropland albedo management, *Proceedings of the National Academy of Sciences*, doi: 10.1073/pnas.1317323111, 2014.
- De Vrese, P., Hagemann, S., and Claussen, M.: Asian irrigation, African rain: Remote impacts of irrigation, *Geophysical Research Letters*, doi: 10.1002/2016GL068146, 2016.
- 790 Donat, M. G., Alexander, L. V., Yang, H., Durre, I., Vose, R., and Caesar, J.: Global Land-Based Datasets for Monitoring Climatic Extremes, *Bulletin of the American Meteorological Society*, doi: 10.1175/BAMS-D-12-00109.1, 2013a.
- Donat, M. G., Alexander, L. V., Yang, H., Durre, I., Vose, R., Dunn, R. J. H., Willett, K. M., Aguilar, E., Brunet, M., Caesar, J., Hewitson, B., Jack, C., Klein Tank, A. M. G., Kruger, A. C., Marengo, J.,
- 795 Peterson, T. C., Renom, M., Oria Rojas, C., Rusticucci, M., Salinger, J., Elrayah, A. S., Sekele, S. S., Srivastava, A. K., Trewin, B., Villarroya, C., Vincent, L. A., Zhai, P., Zhang, X., and Kitching, S.: Updated analyses of temperature and precipitation extreme indices since the beginning of the twentieth century: The HadEX2 dataset, *Journal of Geophysical Research Atmospheres*, doi: 10.1002/jgrd.50150, 2013b.
- 800 Donat, M. G., Pitman, A. J., and Seneviratne, S. I.: Regional warming of hot extremes accelerated by surface energy fluxes, *Geophysical Research Letters*, doi: 10.1002/2017GL073733, 2017.
- [Douglas, E.M., D. Niyogi, S. Frolking, J.B. Yeluripati, R. A. Pielke Sr., N. Niyogi, C.J. Vörösmarty, and Mohanty, U. C.: Changes in moisture and energy fluxes due to agricultural land use and irrigation in the Indian Monsoon Belt. *Geophys. Res. Letts*, 33, doi:10.1029/2006GL026550, 2006.](#)

805

- Erb, K. H., Fetzel, T., Plutzer, C., Kastner, T., Lauk, C., Mayer, A., Niedertscheider, M., Körner, C., and Haberl, H.: Biomass turnover time in terrestrial ecosystems halved by land use, *Nature Geoscience*, doi: 10.1038/ngeo2782, 2016.
- Erb, K. H., Kastner, T., Plutzer, C., Bais, A. L. S., Carvalhais, N., Fetzel, T., Gingrich, S., Haberl, H.,
810 Lauk, C., Niedertscheider, M., Pongratz, J., Thurner, M., and Luysaert, S.: Unexpectedly large impact of forest management and grazing on global vegetation biomass, *Nature*, doi: 10.1038/nature25138, 2018.
- [Fajardo, J., Corcoran, D., Roehrdanz, P. R., Hannah, L. and Marquet, P. A.: GCM compareR: a web application to assess differences and assist in the selection of general circulation models for climate change research, *Methods in Ecology and Evolution*, doi: 10.1111/2041-210X.13360, 2020.](#)
815
- Fereres, E., and Connor, D.: Sustainable water management in agriculture, *Challenges of the New Water Policies for the XXI Century: Proceedings of the Seminar on Challenges of the New Water Policies for the 21st Century*, 2004.
- Fereres, E., and Soriano, M. A.: Deficit irrigation for reducing agricultural water use, *Journal of*
820 *Experimental Botany*, 58, 147-159, 2007.
- Fischer, E. M., and Knutti, R.: Anthropogenic contribution to global occurrence of heavy-precipitation and high-temperature extremes, *Nature Climate Change*, doi: 10.1038/nclimate2617, 2015.
- Goddard, L., Kumar, A., Solomon, A., Smith, D., Boer, G., Gonzalez, P., Kharin, V., Merryfield, W., Deser, C., Mason, S. J., Kirtman, B. P., Msadek, R., Sutton, R., Hawkins, E., Fricker, T., Hegerl, G.,
825 Ferro, C. A. T., Stephenson, D. B., Meehl, G. A., Stockdale, T., Burgman, R., Greene, A. M., Kushnir, Y., Newman, M., Carton, J., Fukumori, I., and Delworth, T.: A verification framework for interannual-to-decadal predictions experiments, *Climate Dynamics*, doi: 10.1007/s00382-012-1481-2, 2013.
- [Gupta, V., Singh, V. and Jain, M. K.: Assessment of precipitation extremes in India during the 21st century under SSP1-1.9 mitigation scenarios of CMIP6 GCMs. *Journal of Hydrology*, doi: 10.1016/j.jhydrol.2020.125422, 2020.](#)
830
- Harris, I., Jones, P. D., Osborn, T. J., and Lister, D.H.: Updated high-resolution grids of monthly climatic observations - the CRU TS3.10 Dataset, *International Journal of Climatology*, doi: 10.1002/joc.3711, 2014.

Formatted: Superscript

- Hartmann, D. L., Tank, A. M. G. K., Rusticucci, M., Alexander, L., Brönnimann, S., Charabi, Y.,
835 Dentener, F., Dlugokencky, E., Easterling, D., Kaplan, A., Soden, B., Thorne, P., Wild, M., and Zhai, P.
M.: Observations: Atmosphere and Surface Supplementary Material, Climate Change 2013 the Physical
Science Basis: Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental
Panel on Climate Change, 2013.
- Hauser, M., Thiery, W., and Seneviratne, S. I.: Potential of global land water recycling to mitigate local
840 temperature extremes, *Earth System Dynamics*, doi:10.5194/esd-10-157-2019, 2019.
- Hirsch, A. L., Prestele, R., Davin, E. L., Seneviratne, S. I., Thiery, W., and Verburg, P. H.: Modelled
biophysical impacts of conservation agriculture on local climates, *Global Change Biology*,
doi:10.1111/gcb.14362, 2018.
- Hirsch, A. L., Wilhelm, M., Davin, E. L., Thiery, W., and Seneviratne, S. I.: Can climate-effective land
845 management reduce regional warming? *Journal of Geophysical Research*, doi: 10.1002/2016JD026125,
2017.
- [Hofer, S., Lang, C., Amory, C., Kittel, C., Delhasse, A., Tedstrone, A. and Fettweis, X.: Greater
Greenland ice sheet contribution to global sea level rise in CMIP6, *Nature Communications*, 11: 6289,
2020.](#)
- 850 [Hossain, F., J. Arnold, E. Beighley, C. Brown, S. Burian, J. Chen, S. Madadgar, A. Mitra, D. Niyogi,
R.A. Pielke Sr., V. Tidwell, and Wegner, D.: Local-to-regional landscape drivers of extreme weather
and climate: Implications for water infrastructure resilience, *Journal of Hydrological Engineering*, doi:
\[10.1061/\\(ASCE\\)HE.1943-5584.0001210, 2015\]\(#\)](#)
- Hurrell, J. W., Holland, M. M., Gent, P. R., Ghan, S., Kay, J. E., Kushner, P. J., Lamarque, J. F., Large,
855 W. G., Lawrence, D., Lindsay, K., Lipscomb, W. H., Long, M. C., Mahowald, N., Marsh, D. R., Neale,
R. B., Rasch, P., Vavrus, S., Vertenstein, M., Bader, D., Collins, W. D., Hack, J. J., Kiehl, J., and
Marshall, S.: The community earth system model: A framework for collaborative research, *Bulletin of
the American Meteorological Society*, doi: 10.1175/BAMS-D-12-00121.1, 2013.
- IPCC.: Summary for policymakers. Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part
860 A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of
the Intergovernmental Panel on Climate Change, 2013.

Formatted: English (UK)

Field Code Changed

Formatted: Font: 12 pt, Not Bold

Formatted: Font: 12 pt, Not Bold

Jia, G., Shevliakova, E., Artaxo, P., De Noblet-Ducoudré, N., Houghton, R., House, J., Kitajima, K., Lennard, C., Popp, A., Sirin, A., Sukumar, R., and Verchot, L.: Land–climate interactions. In: Climate Change and Land: an IPCC special report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems [P.R. Shukla, J. Skea, E. Calvo Buendía, V. Masson-Delmotte, H.-O. Pörtner, D.C. Roberts, P. Zhai, R. Slade, S. Connors, R. van Diemen, M. Ferrat, E. Haughey, S. Luz, S. Neogi, M. Pathak, J. Petzold, J. Portugal Pereira, P. Vyas, E. Huntley, K. Kissick, M. Belkacemi, J. Malley, (eds.)], 2019.

865 [Jiang, D., Tian, Z. and Lang, X.: Reliability of climate models for China through the IPCC third to fifth assessment report, International Journal of Climatology, doi: 10.1002/joc.4406, 2015.](#)

870 Jones, P. W.: First- and second-order conservative remapping schemes for grids in spherical coordinates, Monthly Weather Review, doi: 10.1175/1520-0493(1999)127<2204:FASOCR>2.0.CO;2, 1999.

Kassam, A., Friedrich, T., Derpsch, R., and Kienzle, J.: Overview of the worldwide spread of conservation agriculture, Field Actions Science Report, 2015.

875 Knutti, R., Furrer, R., Tebaldi, C., Cermak, J., and Meehl, G.A.: Challenges in combining projections from multiple climate models, Journal of Climate, doi: 10.1175/2009JCLI3361.1, 2010.

Kueppers, L. M., and Snyder, M. A.: Influence of irrigated agriculture on diurnal surface energy and water fluxes, surface climate, and atmospheric circulation in California, Climate Dynamics, doi: 10.1007/s00382-011-1123-0, 2012.

880 Kueppers, L. M., Snyder, M. A., and Sloan, L. C.: Irrigation cooling effect: Regional climate forcing by land-use change, Geophysical Research Letters, doi: 10.1029/2006GL028679, 2007.

Lawrence, D. M., Hurtt, G. C., Arneth, A., Brovkin, V., Calvin, K. V., Jones, A. D., Jones, C. D., Lawrence, P. J., Noblet-Ducoudré, N. De., Pongratz, J., Seneviratne, S. I., and Shevliakova, E.: The Land Use Model Intercomparison Project (LUMIP) contribution to CMIP6: Rationale and experimental design, Geoscientific Model Development, doi: 10.5194/gmd-9-2973-2016, 2016.


885 [Li, Y., Zhao, M., Motesharrei, S., Mu, Q., Kalnay, E. and Li, S.: Local cooling and warming effects of forests based on satellite observations, Nature Communications, 6: 1-8, 2015.](#)

- 890 Lombardozzi, D. L., Bonan, G. B., Wieder, W., Grandy, A. S., Morris, C., and Lawrence, D. L.: Cover
crops may cause winter warming in snow-covered regions, *Geophysical Research Letters*, doi:
10.1029/2018GL079000, 2018.
- Luyssaert, S., Jammot, M., Stoy, P. C., Estel, S., Pongratz, J., Ceschia, E., Churkina, G., Don, A., Erb,
K., Ferlicoq, M., Gielen, B., Grünwald, T., Houghton, R. A., Klumpp, K., Knohl, A., Kolb, T.,
Kuemmerle, T., Laurila, T., Lohila, A., Loustau, D., McGrath, M. J., Meyfroidt, P., Moors, E. J.,
895 Naudts, K., Novick, K., Otto, J., Pilegaard, K., Pio, C. A., Rambal, S., Rebmann, C., Ryder, J., Suyker,
A. E., Varlagin, A., Wattenbach, M., and Dolman, A.J.: Land management and land-cover change have
impacts of similar magnitude on surface temperature, *Nature Climate Change*, doi:
10.1038/nclimate2196, 2014.
- [Malyshev, S., Shevliakova, E., Stouffer, R. J. and Pacala, S. W.: Contrasting local versus regional
900 effects of land-use-change-induced heterogeneity on historical climate: analysis with the GFDL Earth
System Model. *Journal of Climate*, doi: 10.1175/JCLI-D-14-00586.1, 2015.](#)
- [Mahmood, R., Pielke, R. A., Hubbard, K. G., Niyogi, D., Dirmeyer, P. A., McAlpine, C., Carleton, A.
M., Hale, R., Gameda, S., Beltran-Przekurat, A., Baker, B., McNider, R., Legates, D. R., Shepherd, M.,
905 Du, J., Blanken, P. D., Frauenfeld, O. W., Nair, U. S. and Fall, S.: Land cover changes and their
biogeophysical effects on climate. *International Journal of Climatology*, 34\(4\): 929-953, 2014.](#)
- [Meier, R., Davin, E. L., Lejeune, Q., Hauser, M., Li, Y., Martens, B., Schultz, N. M., Sterling, S., and
Thiery, W.: Evaluating and improving the Community Land Model's sensitivity to land cover.
Biogeosciences, 15, 4731–4757, <https://doi.org/10.5194/bg-15-4731-2018>, 2018.](#)
- Meinshausen, M., Vogel, E., Nauels, A., Lorbacher, K., Meinshausen, N., Etheridge, D. M., Fraser, P.
910 J., Montzka, S. A., Rayner, P. J., Trudinger, C. M., Krummel, P. B., Beyerle, U., Canadell, J. G.,
Daniel, J. S., Enting, I. G., Law, R. M., Lunder, C. R., O'Doherty, S., Prinn, R. G., Reimann, S.,
Rubino, M., Velders, G. J. M., Vollmer, M. K., Wang, R. H. J., and Weiss, R.: Historical greenhouse
gas concentrations for climate modelling (CMIP6), *Geoscientific Model Development*, doi:
10.5194/gmd-10-2057-2017, 2017.
- 915 Miao, C., Duan, Q., Sun, Q., Huang, Y., Kong, D., Yang, T., Ye, A., Di, Z., and Gong, W.: Assessment
of CMIP5 climate models and projected temperature changes over Northern Eurasia, *Environmental*

- Research Letters, doi: 10.1088/1748-9326/9/5/055007, 2014.
- [Niyogi, D., S., Xue, Y. and Raman, S.: Hydrological land surface response in a regime and a midlatitudinal regime, *Journal of Hydrometeorology*, 3: 39-56, 2002.](#)
- 920 Oleson, K. W., Lawrence, D. M., Gordon, B., Flanner, M. G., Kluzek, E., Peter, J., Levis, S., Swenson, S. C., Thornton, E., and Feddema, J.: Technical description of version 4.0 of the Community Land Model (CLM), NCAR/TN-503+STR NCAR Technical Note, 2013.
- [Paulot, F., Malyshev, S., Nguyen, T., Crouse, J. D., Shevliakova, E., and Horowitz, L. W.: Representing sub-grid scale variations in nitrogen deposition associated with land use in a global Earth system model: implications for present and future nitrogen deposition fluxes over North America, *Atmos. Chem. Phys.*, 18, 17963–17978, <https://doi.org/10.5194/acp-18-17963-2018>, 2018.](#)
- 925 Pendergrass, A. G., and Hartmann, D. L.: Changes in the distribution of rain frequency and intensity in response to global warming, *Journal of Climate*, doi: 10.1175/JCLI-D-14-00183.1, 2014.
- [Pielke, R. A., Sr., A. Pitman, D. Niyogi, R. Mahmood, C. McAlpine, F. Hossain, K. K. Goldewijk, U. Nair, R. Betts, S. Fall, M. Reichstein, P. Kabat, and De Noblet, N.: Land use/land cover changes and climate: modeling analysis and observational evidences, *WIREs Clim Change* 2011, 2, 828–850, doi:10.1002/wcc.144, 2011.](#)
- 930 Prestele, R., Hirsch, A. L., Davin, E. L., Seneviratne, S. I., and Verburg, P.H.: A spatially explicit representation of conservation agriculture for application in global change studies, *Global Change Biology*, doi: 10.1111/gcb.14307, 2018.
- Puma, M. J. and Cook, B. I.: Effects of irrigation on global climate during the 20th century, *Journal of Geophysical Research Atmospheres*, doi: 10.1029/2010JD014122, 2010.
- Saeed, F., Hagemann, S., and Jacob, D.: Impact of irrigation on the South Asian summer monsoon, *Geophysical Research Letters*, doi: 10.1029/2009GL040625, 2009.
- 940 Schultz, N. M., Lee, X., Lawrence, P. J., Lawrence, D. M., and Zhao, L.: Assessing the use of subgrid land model output to study impacts of land cover change, *Journal of Geophysical Research*, doi: 10.1002/2016JD025094, 2016.
- Sen, P. K.: Estimates of the Regression Coefficient Based on Kendall's Tau, *Journal of the American Statistical Association*, doi: 10.1080/01621459.1968.10480934, 1968.

- 945 Seneviratne, S. I., Corti, T., Davin, E. L., Hirschi, M., Jaeger, E. B., Lehner, I., Orlowsky, B., and Teuling, A. J.: Investigating soil moisture-climate interactions in a changing climate: A review, *Earth-Science Reviews*, 2010.
- Seneviratne, S. I., Wilhelm, M., Stanelle, T., Van Den Hurk, B., Hagemann, S., Berg, A., Cheruy, F., Higgins, M. E., Meier, A., Brovkin, V., Claussen, M., Ducharne, A., Dufresne, J. L., Findell, K. L.,
- 950 Ghattas, J., Lawrence, D. M., Malyshev, S., Rummukainen, M., and Smith, B.: Impact of soil moisture-climate feedbacks on CMIP5 projections: First results from the GLACE-CMIP5 experiment, *Geophysical Research Letters*, doi: 10.1002/grl.50956, 2013.
- Sherwood, J., Clabeaux, R., and Carbajales-Dale, M.: An extended environmental input-output lifecycle assessment model to study the urban food-energy-water nexus, *Environmental Research Letters*, doi: 10.1088/1748-9326/aa83f0, 2017.
- 955 Sherwood, S. C., Dixit, V., and Salomez, C.: The global warming potential of near-surface emitted water vapour, *Environmental Research Letters*, doi: 10.1088/1748-9326/aae018, 2018.
- Siebert, S., Döll, P., Hoogeveen, J., Faures, J. M., Frenken, K., and Feick, S.: Development and validation of the global map of irrigation areas, *Hydrology and Earth System Sciences*, doi: 10.5194/hess-9-535-2005, 2005.
- 960 Sillmann, J., and Croci-Maspoli, M.: Present and future atmospheric blocking and its impact on European mean and extreme climate, *Geophysical Research Letters*, doi: 10.1029/2009GL038259, 2009.
- Smith, R. A., Ditmire, T., and Tisch, J. W. G.: Characterization of a cryogenically cooled high-pressure gas jet for laser/cluster interaction experiments, *Review of Scientific Instruments*, doi: 10.1063/1.1149181, 1998.
- 965 Stouffer, R. J., Eyring, V., Meehl, G. A., Bony, S., Senior, C., Stevens, B., and Taylor, K. E.: CMIP5 scientific gaps and recommendations for CMIP6, *Bulletin of the American Meteorological Society*, doi: 10.1175/BAMS-D-15-00013.1, 2017.
- 970 Tebaldi, C., and Knutti, R.: The use of the multi-model ensemble in probabilistic climate projections, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 2007.

- 975 Thiery, W., Davin, E. L., Lawrence, D. M., Hirsch, A. L., Hauser, M., and Seneviratne, S. I.: Present-day irrigation mitigates heat extremes, *Journal of Geophysical Research*, doi: 10.1002/2016JD025740, 2017.
- Thiery, W., Visser, A. J., Fischer, E. M., Hauser, M., Hirsch, A. L., Lawrence, D. M., Lejeune, Q., Davin, E. L., and Seneviratne, S. I.: Warming of hot extremes alleviated by expanding irrigation, *Nature Communications*, doi: 10.1038/s41467-019-14075-4, 2020.
- 980 Vogel, M. M., Orth, R., Cheruy, F., Hagemann, S., Lorenz, R., van den Hurk, B. J. J. M., and Seneviratne, S. I.: Regional amplification of projected changes in extreme temperatures strongly controlled by soil moisture-temperature feedbacks, *Geophysical Research Letters*, doi:10.1002/2016GL071235, 2017.
- Vogel, M. M., Zscheischler, J., & Seneviratne, S. I.: Varying soil moisture-atmosphere feedbacks explain divergent temperature extremes and precipitation projections in central Europe. *Earth System Dynamics*, doi: 10.5194/esd-9-1107-2018, 2018.
- 985 von Storch H.: Response to Comment on “Reconstructing Past Climate from Noisy Data”, *Science*, doi: 10.1126/science.1121571, 2006.
- 990 [Yazdandoost, F., Moradian, S., Zakipour, M., Izadi, A. and Bavandpour, M.: Improving the precipitation forecasts of the North-American multi model ensemble \(NMME\) over Sistan basin, *Journal of Hydrology*, 590: 125263, 2021.](#)



995 Figure 1. (a) Percentage of each grid cell equipped for irrigation (%) (Siebert *et al.*, 2005). (b) Potential estimate of CA extent mapped
 000 to the CLM crop PFT (Prestele *et al.*, 2018). The red boxes in (a) denote the regional domains where irrigation is extensive and were
 thus examined in greater detail including Western North America (WNA), Central North America (CNA), south Europe and
 Mediterranean (MED), West Asia (WAS), South Asia (SAS), Southeast Asia (SEA), and East Asia (EAS). The red boxes in (b) denote
 the regional domains where CA is extensive and were thus examined in greater detail including WNA, CNA, MED, South-eastern
 South America (SSA), Central Europe (CEU) and Southern Australia (SAU).













Figure 2. Added value of including irrigation and CA in the simulated warming trends over 1981-2010. Absolute change in spatial root-mean-square error (RMSE) for the (a) IRR and (b) CA ensemble relative to the CTL ensemble over different regions (x axis) and with respect to 3 observational products (y axis). Considered regions are the SREX regions where irrigation is extensive (as highlighted in Figure 1a) and where CA is extensive (Figure 1b), in addition to global land, global irrigated land and global CA land. Observational products are for near-surface air temperature T_{2m} (CRU), annual maximum daytime temperature TXx (GHCNDEX and HadEX2). The spatial RMSEs are computed for the ensemble mean warming trend in every pixel, and subsequently averaged over the selected region. Regions with an observational coverage below 50% are marked in white.



030







Deleted: ¶

Formatted: Font: 6 pt

Formatted: Heading-Main



Deleted: 1 ... [6]



Deleted: 7 ... [7]

Deleted: over irrigated pixels

Deleted: and

Formatted: Font colour: Text 1

Formatted: Font colour: Text 1

Deleted: For irrigated pixels (a-b), CA pixels (c-d), and (e-f) all land pixels.

Formatted: Font colour: Text 1

Deleted: -

Formatted: Font colour: Text 1

Deleted: -


Formatted: Font colour: Text 1

Deleted: Legend: ...TL (red), IRR (blue), CA (cyan), CRU (purple), HadEX2 (green) ... [8]

Deleted: 1

040 Figure 4. Spatial average of the warming rates for T_{2m} (a, c and e), TXx (b, d and f), SHF (g and h), LHF (i and j) and ET (k and l) for the CESM ensembles and observations. Data points specify the mean T_{2m} and TXx temperatures, SHF and LHF and ET volumes for irrigated pixels (a, b, g, i and k), CA pixels (c, d, h, j and l), and (e-f) all land pixels. The slope was estimated using Sen's slope for the CTL (red), IRR (blue), CA (cyan), CRU (purple), HadEX2 (green), and GHCNDEX (black) temperatures.

045









Figure 5. Top-of-atmosphere (TOA) net radiation $R_{n,TOA}$ [$W m^{-2}$] in (a) the CTL ensemble, (b) Impact of irrigation on $R_{n,TOA}$. Difference map is based on the ensemble mean of each experiment for 1981–2010.


Deleted: and
 Deleted: the IRR ensemble. (c)






Deleted: b
 Deleted: b
 Deleted: Figure 6. Subgrid-scale differences between the irrigated and rainfed crop tile in the IRR ensemble (irrigated minus rainfed) (a and g) and between CA and conventionally managed (CM) crops (CA minus CM) (e and h). For T, (a-b) and ET (g-h). Grid-scale differences between the CTL and IRR ensemble (IRR minus CTL) (c, e and i) and between CA and conventionally managed (CM) crops (CA minus CM) (b, f and j).
 Formatted: Font: 20 pt, Not Bold

Deleted: b
 Deleted: b



Formatted: Font: 20 pt, Not Bold
 Deleted: Subgrid-scale differences between the irrigated and [12]
 Deleted: c
 Deleted: I, m and
 Deleted: q
 Deleted: and conventionally managed (CM) crops
 Deleted: CM
 Deleted: f
 Deleted: h
 Deleted: n
 Deleted: r
 Deleted: c
 Deleted: d
 Deleted: ET (i-j),
 Deleted: e
 Deleted: f
 Deleted: m
 Deleted: n
 Deleted: q
 Deleted: r

110 **Figure 6. Grid-scale differences between the CTL and IRR ensemble (IRR minus CTL) (a, c, e, g and i) and between the CTL and CA ensemble (CA minus CTL) (b, d, f, h and j). For T, (a-b), TMO (c-d), ET (e-f), LHF (g-h) and SHF (i-j), displayed over irrigated/CA pixels for comparative purposes. Differences are based on the ensemble mean warming trends of each experiment for 1981–2010. Hatching denotes less than 10% change induced by the model on mean warming trends of lumped ensemble members.**










Figure 7. Subgrid-scale differences between the irrigated and rainfed crop tile in the IRR ensemble (irrigated minus rainfed) (a, c, e and g) and between CA and conventionally managed (CM) crops (CA minus CM) (b, d, f and h). For T, (a-b), ET (c-d), LHF (e-f) and SHF (g-h), displayed over irrigated/CA pixels for comparative purposes. Differences are based on the ensemble mean warming trends of each experiment for 1981–2010. Hatching denotes less than 10% change induced by the model on mean warming trends of lumped ensemble members.

170

Formatted: Normal






Figure 8. Average of the subgrid-scale warming rates for TS (a-f), the SHF (g) and LHF (h) over (a and g) irrigated pixels for the irrigated and rainfed crop files; (b and h) CA pixels for the CA and CM crop files; (b) all pixels for the irrigated and rainfed crop files; (d) all pixels for the CA and CM crop files; (e) irrigated pixels over the MED SREX region; and (f) CA pixels over the MED SREX region. Data points are the mean TS, LHF and SHF values within the crop files and pixels specified. The slope was estimated using Sen's slope for the rainfed/CM (red), irrigated/CA (blue) experiments. For (a), (b), (c), (d), (e) and (f) the regions where less than 50% of the land pixels did not contain a value were excluded. For all land pixels (g and h), the minimum number of land pixels that needed to contain a value in order to be retained in the analysis was 15%.

Moved (insertion) [3]

Formatted: Normal, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Deleted: ¶

Deleted: ¶


Deleted: ¶

... [13]

Formatted: Normal

Moved (insertion) [2]

Formatted: Font: 20 pt



Moved up [3]:


Formatted: Font: 12 pt

Formatted

... [14]

Deleted: irrigated = 0.034, rain = 0.045, e-obs = 0.03 (BIAS= 0.004, 0.015)¶

... [15]



Moved up [2]:

Deleted: ¶

285 **Table 1. Bias and spatial RMSE of the ensemble mean warming trends (slopes) of the CTL, IRR and CA experiments versus the observational products for the years 1981-2010^a.**

| Physical (Units) | Quantity | All land bias | | | Irrigated land bias | | CA land bias | | All land RMSE | | | Irrigated land RMSE | | CA land RMSE | |
|-------------------------------------|----------|---------------|--------|--------|---------------------|-------|--------------|-------|---------------|-------|-------|---------------------|-------|--------------|-------|
| | | CTL | IRR | CA | CTL | IRR | CTL | CA | CTL | IRR | CA | CTL | IRR | CTL | CA |
| CRU T_{2m} (K yr ⁻¹) | | -0.006 | -0.004 | -0.004 | -0.003 | 0.001 | 0.002 | 0.004 | 0.027 | 0.025 | 0.027 | 0.020 | 0.019 | 0.018 | 0.017 |
| DHCNDEX TXx (K yr ⁻¹) | | 0.024 | 0.030 | 0.027 | 0.009 | 0.013 | 0.0 | 0.011 | 0.107 | 0.111 | 0.110 | 0.078 | 0.082 | 0.124 | 0.125 |
| | | | | | | | 06 | | | | | | | | |
| HadEX2 TXx (K yr ⁻¹) | | 0.007 | 0.013 | 0.010 | 0.008 | 0.012 | 0.008 | 0.013 | 0.135 | 0.135 | 0.136 | 0.121 | 0.122 | 0.086 | 0.087 |

- Deleted: S
- Deleted: E
- Deleted: M
- Deleted: W
- Deleted: T
- Deleted: S
- Deleted: E
- Deleted: V
- Deleted: the Warming Trends of
- Deleted: O
- Deleted: P

^aRegions with an observational coverage below 50% are excluded.

290 **Table 2. Bias and spatial RMSE of the subgrid-scale ensemble mean warming trends (slopes) of the RAIN, IRR_{SUB}, CA_{SUB} and CM Experiments Versus the E-OBS (K yr⁻¹) observational product in the MED region for the years 1981-2010.**

| All MED pixels bias | | | | Irrigated MED pixels bias | | | CA MED pixels bias | | All MED pixels RMSE | | | | Irrigated MED pixels RMSE | | CA MED pixels RMSE | |
|---------------------|-------|--------------------|-------------------|---------------------------|--------------------|-------|--------------------|-------|---------------------|--------------------|-------------------|-------|---------------------------|-------|--------------------|--|
| RAIN | CM | IRR _{SUB} | CA _{SUB} | RAIN | IRR _{SUB} | CM | CA _{SUB} | RAIN | CM | IRR _{SUB} | CA _{SUB} | RAIN | IRR _{SUB} | CM | CA _{SUB} | |
| 0.032 | 0.033 | 0.022 | 0.035 | 0.015 | 0.004 | 0.013 | 0.022 | 0.040 | 0.039 | 0.031 | 0.026 | 0.028 | 0.031 | 0.027 | 0.026 | |

Table 3. Impact of irrigation and CA on various climatological values (absolute slope differences calculated as IRR minus CTL and CA minus CTL for grid-scale, IRR_{SUB} minus RAIN and CA_{SUB} minus CM for subgrid-scale) for the years 1981-2010^a.

| | Physical Quantity (Units) | Irrigated Land | | | CA Land | | |
|----------------------------------|--|----------------|--------------------|---------------------------|---------------|-------------------|---------------------|
| | | CTL | IRR | ABS | CTL | CA | ABS |
| Grid-scale | T _{2m} (K yr ⁻¹) | 0.026 | 0.030 | 0.004 ^c | 0.026 | 0.028 | 0.002 ^c |
| | TXx (K yr ⁻¹) | 0.034 | 0.038 | 0.004 ^c | 0.039 | 0.044 | 0.005 ^c |
| | T _s (K yr ⁻¹) | 0.009 | 0.010 | 0.001 ^c | 0.016 | 0.015 | -0.001 ^c |
| | LHF (W/m ² yr ⁻¹) | <u>0.029</u> | <u>0.041</u> | <u>0.012</u> | <u>0.029</u> | <u>0.053</u> | <u>0.024</u> |
| | SHF (W/m ² yr ⁻¹) | <u>-0.010</u> | <u>-0.001</u> | <u>0.009</u> | <u>-0.010</u> | <u>0.004</u> | <u>0.014</u> |
| Subgrid-scale^b | Physical Quantity (Units) | RAIN | IRR _{SUB} | ABS | CM | CA _{SUB} | ABS |
| | T _s (K yr ⁻¹) | 0.038 | 0.030 | -0.008 ^c | 0.031 | 0.030 | -0.001 ^c |
| | ET (mm yr ⁻¹) | 0.286 | 0.939 | 0.653 ^c | 0.182 | 0.265 | 0.083 |
| | LHF (W/m ² yr ⁻¹) | <u>0.004</u> | <u>0.060</u> | <u>0.056</u> | <u>0.009</u> | <u>0.014</u> | <u>0.005</u> |
| | SHF (W/m ² yr ⁻¹) | <u>0.009</u> | <u>0.005</u> | <u>-0.004^c</u> | <u>0.056</u> | <u>0.060</u> | <u>0.004</u> |

^aABS denotes the absolute change of each given quantity.

^bRegions with a coverage below 25% are excluded. For grid-scale calculations, regions with a coverage below 50% are excluded.

^cThe changes significant at the 1% significance level (two-sided Wilcoxon signed rank test on ensemble mean slopes for irrigated/CA pixels).

- Deleted: 2
- Deleted: I
- Deleted: V
- Deleted: C
- Deleted: V
- Deleted: A
- Deleted: SI
- Deleted: D
- Deleted: C
- Deleted: M
- Deleted: M
- Deleted: G
- Deleted: S
- Deleted: M
- Deleted: M
- Deleted: S
- Deleted: S

310

315

Page 4: [1] Deleted Aine Gormley 19/12/2020 12:44:00

Page 4: [2] Deleted Aine Gormley 19/12/2020 08:31:00

Page 11: [3] Deleted Gormley, Aine 11/04/2021 21:48:00

Page 11: [4] Formatted Gormley, Aine 01/05/2021 18:24:00

Font: (Default) Times New Roman, Font colour: Black, Pattern: Clear

Page 28: [5] Deleted Gormley, Aine 16/04/2021 14:13:00

Page 31: [6] Deleted Gormley, Aine 14/04/2021 15:30:00

Page 31: [7] Deleted Gormley, Aine 14/04/2021 15:33:00

Page 31: [8] Deleted Aine Gormley 16/09/2020 12:47:00

Page 31: [8] Deleted Aine Gormley 16/09/2020 12:47:00

Page 33: [9] Deleted Gormley, Aine 15/04/2021 21:36:00

Page 34: [10] Deleted Gormley, Aine 15/04/2021 21:36:00

Page 34: [11] Deleted Gormley, Aine 15/04/2021 21:36:00

Page 34: [12] Deleted Gormley, Aine 15/04/2021 21:43:00

Page 38: [13] Deleted Gormley, Aine 15/04/2021 22:11:00

Page 38: [14] Formatted Gormley, Aine 15/04/2021 22:06:00

Font colour: Text 1, Not Highlight

Page 38: [14] Formatted Gormley, Aine 15/04/2021 22:06:00

Font colour: Text 1, Not Highlight

Page 38: [14] Formatted Gormley, Aine 15/04/2021 22:06:00

Font colour: Text 1, Not Highlight

Page 38: [15] Deleted Gormley, Aine 15/04/2021 22:14:00