

Response to Reviewer #3

The authors present a detailed analysis of multi-dynamic-ecosystem-model estimates of LULCC emissions under different initial and environmental conditions, with some comparison with recent bookkeeping approaches. A main focus is the contribution of the “Loss of Additional Sink Capacity” (LASC) to emissions, which can be best estimated by dynamic ecosystem models with transient environmental conditions. The authors conclude that LASC is considerable, and has some regional variation. Furthermore, they show that CO₂ effects dominate the contributions to LASC over climate effects. They also raise the question of whether LASC is appropriate for a harmonized attribution of LULCC emissions.

Overall review

This is an important contribution to LULCC and carbon/climate science and presents interesting results from the trendy model ensemble. There are a few main issues that need to be addressed, however, prior to publication. These are summarized here, with further detail following:

Thank you for this thorough review of our manuscript and the positive evaluation. We will follow your suggestions which will considerably improve the manuscript.

1) There seems to be a better way to calculate the CO₂ vs climate estimates. You have simulations that you can difference directly, rather than using the ratio method that has limitations.

To our knowledge, the proposed ratio method is the only possible option to derive climate vs CO₂ related alterations in f_{LULCC} based on available data (TRENDY v8). Because S0, S1, and S2 simulations have fixed pre-industrial LULCC forcing, their comparison does not allow to derive f_{LULCC} .

Additionally, S1 is the only simulation where only CO₂ but not climate conditions are transient while no corresponding simulation exists that includes LULCC. Therefore, the available TRENDY simulations do not allow to directly derive climate vs CO₂ related alteration in f_{LULCC} . We will better explain this in the revised manuscript. For the details, please refer to the answers given below (in particular the answer to comment on lines 220-240).

2) The results and discussion of regional LASC are confusing and not to the point. These can be better constrained and made more clear with main points in mind.

See below and responses to the other reviewers how we will improve the results and discussion. In particular, we will change the structure and wording in Sect. 3.1, 3.2, 3.2.1 and 3.2.2 (see response to your comment on lines 296-320 and the comment on lines 321-352).

3) The suggested “harmonized attribution” of LULCC emissions is not consistent with the paper findings and doesn’t differentiate between science and policy. The real issue here is full scientific estimates of emissions, as the authors clearly show require the inclusion of LASC, vs attribution of LULCC emissions for policy purposes, which is a political issue. This paper isn’t about harmonizing different model approaches, but rather clarifying and understanding their differences. This provides scientific bases for accurate estimation of emissions, and provides info for political decisions

regarding accounting for attribution policy, but doesn't show an answer to the "correct" accounting for policy or whether/how the different approaches should be "harmonized."

Thank you, we agree that we did not differentiate enough between science and policy relevance of our findings. Instead of referring to a 'correct' or 'harmonised attribution', we will now talk about an attribution that is more robust against choices of environmental drivers, historical timing of LULCCs and the accumulation period. With this in mind, we will incorporate your suggestion on the reference period (and a suggestion given by Reviewer #2) to give various possibilities for a more balanced estimation of f_{LULCC} and LASC (and highlight the need for future research, in particular, to investigate a suited reference period). Additionally, we will highlight the policy relevance where needed to enable a better differentiation for the reader. The revised manuscript, at several places, will include respective assignments (see responses below for more details, in particular answers given to your comment on lines 416-445 and comment on lines 467-472).

Specific comments and suggestions

Abstract

The abstract focuses more on the argument for LASC and DGVMs, than on the results of the study. The argument is better suited for the discussion, with a brief statement in the abstract stating that you use DGVMs to estimate LASC. Then you can add more findings/explanation to the abstract, such as temporal transitions in PTD that support the DGVM vs bookkeeping difference statements.

We will carefully rephrase the abstract in several places to avoid the impression that we make recommendations for policy and/or science. However, the LASC is one of the most important differences between DGVM and bookkeeping estimates and has never been analyzed over time and space and many models, which justifies its prominence in the abstract; we will clarify this as well through rephrasing and shorten the LASC description to also include the temporal transition in PTD around 1960 (though not namely, as the term is not introduced in the abstract):

NEW - 'The LASC results from the impact of environmental changes on land carbon storage potential of managed land compared to potential vegetation, and accumulates over time, which is not captured in bookkeeping models. f_{LULCC} from transient DGVM simulations, thus, strongly depends on the timing of land use and land cover changes mainly because LASC accumulation is cut off at the end of the simulated period. To estimate the LASC, f_{LULCC} from pre-industrial DGVM simulations, which is independent of changing environmental conditions, can be used. Additionally, DGVMs using constant present-day environmental forcing enable an approximation of bookkeeping estimates.'

NEW - 'Around 1960, the accumulating nature of the LASC causes global transient f_{LULCC} estimates to exceed estimates under present-day conditions, despite generally increased carbon stocks in the latter.'

line 8:

“arbitrary chosen” is unnecessary.

We will delete ‘arbitrary chosen’, see comment above.

lines 10-11:

Constant environmental forcing does not generate a condition independent of timing and legacy of lulcc. It generates a condition independent of changing environmental conditions, which is how this should be described.

Ensuring that the lulcc magnitude and trajectory are identical across simulations, along with constant environmental forcing, creates a condition almost independent of the timing and legacy of lulcc, as long as the transient dependence isn’t the question. Full independence in this case requires that the biogeochemical dynamics are also identical across simulations, such as may be possible when a single model is used. But this is not possible across different models because each model will respond differently to the same exact lulcc (which is also very difficult to achieve across models), even if the environmental forcing is identical.

I think you mean this in the context of a single-model experiment, with the exact same lulcc and environmental forcing, and not focused on how the LULCC trajectory determines emissions (i.e., path dependence), but this needs to be qualified here as it sounds more general.

Thank you for this interesting point. We agree that f_{LULCC} is not fully independent of timing of LULCCs and legacy effects due to different biogeochemical dynamics implemented in the different DGVMs. We will change the wording according to your suggestion to avoid misunderstanding (see comment above and further comments below).

lines 21-23:

I am not sure that your analysis calls for this, nor that your approach bridges the two. The issue is a full scientific accounting of emissions vs a partial accounting for policy purposes. While you have evidence for a full scientific accounting, you still have little basis for a partial-LASC approach.

Thank you, we agree that our original statement was overstated. We suggest to change the wording to better match the main outputs of this study, as follows:

NEW - ‘Our study unravels the strong dependence of f_{LULCC} estimates on the time a certain land use and land cover change event happened to occur, and on the chosen time period for the forcing of environmental conditions in the underlying simulations. We argue for an approach that provides an accounting of f_{LULCC} that is more robust against these choices, for example by estimating a mean DGVM-ensemble f_{LULCC} and LASC for a defined reference period and homogeneous environmental changes (CO₂-only).’

Introduction

lines 28-30:

awkward sentence. delete “e.g.” and use three clauses: deforestation...high lats, tropical deforestation, and recent forest expansion in high lats.

Thank you, we will change the wording according to your suggestion and a comment of Reviewer #2 as follows:

NEW - '[...] in particular through deforestation driven by early agricultural expansion in mid-latitudes, recent tropical deforestation, and recent forest expansion in mid- and high-latitudes (Klein Goldewijk et al., 2011).'

line 31:

“...contributed approximately one-third of global anthropogenic...”

Will be changed accordingly.

line 35:

delete “in line.” then: flulcc may also gain...

Will be changed accordingly.

line 42:

“...but also change in...”

Will be changed accordingly.

line 50:

delete “allow to”

Will be changed accordingly.

lines 69-72:

The language here starts to confuse the definition and example of LASC. It sounds like just having the potential vegetation or the foregone sink is part of LASC, in addition to the difference of environmental effects on the managed vs unmanaged vegetation. These two sentences should be reworded to clarify that “assuming potential vegetation” and “capture the foregone sinks a given LULCC event destroys” do not obfuscate the definition of LASC.

Thank you, we will reword this paragraph considering your comment and a comment of Reviewer #2 as follows:

NEW - 'Thus, even after the instantaneous emissions of the deforestation event may have ceased, deforestation continues to alter the f_{LULCC} since the reference simulation assumes the potential vegetation cover in the absence of LULCCs, and simulates its response to environmental changes. This example illustrates an interesting aspect of the LASC: It has been acknowledged that the LASC in its literal sense (a loss of carbon, positive LASC values) is an unrealized C uptake potential and is

not reflected in any real change in atmospheric CO₂ concentration (Pongratz et al., 2014). However, as the LASC captures the foregone sinks a given LULCC event destroys and likely accumulates even in absence of further LULCCs, it manifests in the budget of atmospheric CO₂ as compared to a reference excluding LULCCs (Pongratz et al., 2014). In contrast to the theoretical nature of positive LASC values, negative values counted towards the LASC, for example due to reforestation, depict realized C uptake which is theoretically observable (though observations in the field are highly complex due to co-occurrence of natural carbon fluxes).'

lines 71-72:

this statement is not correct. there is no guarantee that the change in environmental conditions is increasing LASC at the time of change. so "same direction" does not make sense. I think you are still referring to the example here, but this is a more general concluding statement.

it is more correct to state that these emissions "can" accumulate in the absence of further LULCC, depending on the environmental conditions. which you then give examples of in the next paragraph.

Thank you, we agree that the statement, referring to the example before, was not generally valid. We will change the wording according to your suggestion (see response to previous comment).

lines 98-99:

I don't agree that constant conditions create a condition independent of lulcc timing. regardless of the forcing, the time since disturbance makes a difference, especially in the context of DGVMs where ecosystems are not static. A subsequent LULCC has a different emission depending on the period between it and the previous LULCC. And an event early in time on 'pristine' land may affect a different biomass then one later in time, even if the conditions are identical. Even with a spun-up model, the biomass may not remain constant during a simulation, and depending on the resolution and application of LULCC, a later LULCC may or may not be applied consistently to 'pristine' or 'managed' area (and corresponding biomass), or some combination of the two.

I think this description should be changed to "independent of environmental trends" or something like this.

Thank you. We agree that, although based on fixed environmental forcings, f_{LULCC} derived under present-day and/or pre-industrial conditions is not totally time-independent given the reasons you stated. We will alter the description accordingly:

NEW - 'However, as f_{LULCC} quantities derived under constant present-day conditions are independent of long-term environmental trends [...]'

line 108:

delete "in line"

Thank you, we will delete 'in line'.

lines 109-110:

unclear: "and consequently into the natural land c sink." how is this related to 'included or excluded?'

We apologize for the unclear wording and will modify the sentence to make it clearer:

NEW - 'It needs to be decided whether the LASC should be included or excluded (as argued e.g. in Gasser and Ciais 2013; Gasser et al. 2020) as part of f_{LULCC} and consequently be counted towards the terrestrial C sink or not.'

Data and methods

lines 171-172 and 178-179 and 194-195:

These really seem like a single equation each.

Thank you. We will put all related equations into single lines.

line 188:

again, this isn't true. timing and legacy of LULCC are inherently critical and determining elements in any transient LULCC simulation. these simulations are indifferent to environmental trends or changing environmental conditions.

Thank you, we will improve the description according to your suggestion:

NEW - 'The latter two are derived under constant environmental forcing, meaning that both are indifferent to long-term environmental trends [...]'

line 219:

"...C stock changes..."

Thank you, we will change it.

lines 220-240:

Why did you do it this way, particularly for climate effects? You can get co_2+ndep -related emissions directly by subtracting S0 and S1, which would be interesting to compare with your estimate of co_2 only emissions using this ratio method.

And you can get the climate-related emissions directly by subtracting S1 and S2, including interactions with CO_2 , without using this indirect ratio method. and your assumption regarding zero interactions between climate and co_2 is not sound because S2 includes both at the same time, and S1 includes only co_2 . I suggest you use the actual nbp difference of these 2 sims to estimate the climate effects.

Indeed, subtracting NBP from S0 and S1 simulations enables the isolation of effects related to CO_2 concentration and nitrogen deposition on the biospheric carbon fluxes. Unfortunately, as the land

use/cover distribution is fixed to the pre-industrial state in both simulations, a derivation of the effects of CO₂ and nitrogen deposition effects on f_{LULCC} is not possible from these simulations. We will add a sentence to clarify this and slightly change the respective paragraph:

NEW - 'These simulations do not include transient LULCCs and can therefore not directly be used to estimate the climate vs CO₂ related alteration of f_{LULCC} . However, [...]'

Similarly, subtracting NBP from S1 and S2 enables isolation of climate effects on biospheric C fluxes, while f_{LULCC} alteration by climate can not directly be derived this way. In Figure A3 we show the relative change of C stocks in S1 vs S2 simulations for comparison.

It is true that the S2 simulation includes climate and CO₂ related effects on NBP at the same time. Thus, the statement 'Synergies [...] are assumed zero in this case.' was confusing. However, the statement is referring to our ratio approach, with the climate-induced f_{LULCC} share being derived as the total f_{LULCC} minus f_{LULCC_CO2} (the latter being the flux share if only CO₂ concentrations would have changed). In line with the response to a comment of Reviewer #2 we will add an explanation on the underlying assumption. Additionally, to eliminate confusion we will change the wording as follows:

NEW - 'Here we note, interacting effects of elevated CO₂ concentrations and temperature or precipitation on biomass productivity (observed under experimental setups; e.g. Obermeier et al. 2017) might obscure this attribution (Lombardozzi et al., 2018). Nevertheless, the assessment of the relative contribution as done by this approach seems valid as no significant interactions between these influencing factors on C stocks were observed within the TRENDY ensemble (Fernández-Martínez et al., 2019) nor within a fully coupled single model investigation (Devaraju et al., 2016).'

NEW - 'Note, by subtracting f_{LULCC_CO2} from the total f_{LULCC} to derive the climate-only flux shares, this approach assumes zero synergies between effects of CO₂ concentrations and climatic changes on NBP in the DGVMs.'

Additionally, we apologize, as this confusion might have been caused by an error in our formula for $f_{LULCC_Climate}$. The formula should read $f_{LULCC_Climate} = f_{LULCC_trans} - f_{LULCC_CO2}$ instead of $f_{LULCC_Climate} = f_{LULCC_Climate} - f_{LULCC_CO2}$.

Also, you don't report on the co2+no2 effects at all, so why is this simulation listed in the methods?

We agree that this information was not necessary and will remove it from Table 1 (columns *Nitrogen deposition* and *Nitrogen Fertilization*). However, high spread between models might partly result from 'co2+no2' effects since some of the used model include nitrogen fertilization while others do not. We will rephrase the respective wording in the appendix to highlight this as follows:

NEW - 'This pronounced model spread can be explained by intertwining issues, such as the low quality of historical LULCC data (with different data bases), the consideration or neglect of

relevant processes (e.g. nitrogen fertilisation), the simplified representation and uncertainty in the parameterization of management and natural processes, uncertainties in soil and vegetation C stocks, and the lack of observational constraints (Friedlingstein et al., 2019, Gasser et al., 2020, Lienert et al., 2018, Goll et al., 2015, Li et al., 2017).'

Results and discussion

lines 246-247:

unclear: "...to consistently use the same models for the flux and bias estimates on a spatio-temporal level..."

We apologize for the confusion and will change the sentence also in line with your next comment '[...] to consistently use the same models where possible [...]'].

line 254:

did you use the trendyv9 version of sdgvm throughout? if so, you need to state this in the methods.

Yes, we used TRENDY v9 output from SDGVM throughout the study. According to your suggestion, we suggest to add a sentence to section 2. Data and Methods and refer to it in the results section as follows:

NEW - 'Note, for SDGVM, model output from TRENDY v9 was used due to erroneous merging of land cover and LULCC datasets in earlier versions that caused a C loss over the period ~1900-1970 mainly in semi-arid regions.'

NEW - 'Slight differences ($<0.1 \text{ PgC yr}^{-1}$) between $f_{\text{LULCC_trans}}$ derived in this study and the DGVM-derived GCB2019 estimates are attributable to the fact that we used only a subset ($n=12$) of the models analyzed within the GCB2019 ($n=15$), to consistently use the same models where possible, and the inclusion of TRENDY v9 model output for SDGVM (for reasons refer to Sect. 2).'

NEW - 'Note, the LASC of 0.8 PgC yr^{-1} (0.84 PgC yr^{-1} with two decimal places) for 2009-2018 presented here is based on TRENDY v8 model output combined with the newer (TRENDY v9) SDGVM output. When consistently using TRENDY v8 output, the resulting LASC (usually rounded to one decimal place) becomes pronounced higher ($>10\%$) with 0.9 PgC yr^{-1} due to rounding of 0.85 PgC yr^{-1} .'

lines 256-257:

not necessarily. the single year pre-ind estimate is closer to bk estimates, and is equally close to gasser as the present-day for the decade.

Thank you, we suggest to rephrase the sentence as follows:

NEW - 'As expected (Sect. 1), $f_{\text{LULCC_pd}}$ is the DGVM-based f_{LULCC} estimate that is most similar to the bookkeeping mean in the GCB2019 when compared over multiple years.'

line 259:

you haven't shown the cumulative yet

Thank you, we suggest to add a reference to the cumulative estimates in the beginning of the results section:

NEW - 'A general overview of most recent annual and cumulative estimates of f_{LULCC} shows that our estimates are in good agreement to the published ones (Friedlingstein et al. 2019; Gasser et al. 2020; Tables 4 to 6).

line 260:

awkward. try "...in the 1950's and again at the end of the simulation..."

Thank you, we will change it according to your suggestion.

line 272:

"...comes into play..."

Thank you, we will change it.

lines 272-273:

awkward first half of sentence.

Thank you, we will change the wording of the respective sentence:

NEW - 'In general, beneficial environmental alterations for C sequestration widely increased the potential C stocks (Fig. 8), and thus, the LASC steadily increased (Fig. 2b,e), reaching about ~40% in recent annual and ~20% in cumulative contributions to f_{LULCC_trans} (Fig. 2c,f).'

line 277:

during this period (first half of 20th century)

Thank you, we will change the wording according to your suggestion:

NEW - '[...] still increase faster than the other estimates during this period (first half of 20th century; EED and PTD remain increasing) [...]

line 279:

"these peaks"

Thank you, we will change it.

lines 281-284:

maybe because the LULCC occurrence is the same between simulations. but post-peak carbon stocks might have been lowered enough differentially between the sims to reduce the post-peak difference because carbon reductions are generally implemented as fractions, in which case the same LULCC would reduce more carbon in the pd sim than in the pi sim because of higher carbon

stocks in the pd sim. and the eed peak is actually after the other peaks and stretched out. so is this really independent of the lulcc timing, as there is a peak in lulcc area that coincides with this emission peak.

We agree that the wording was misleading. We will change it accordingly as follows:

NEW - 'The latter is highlighted by the simultaneous peak in EED which in essence is the intersection of LULCCs with the difference in standing biomass and actual soil C stocks due to altered environmental conditions over the last hundred years (under pre-industrial vs present-day environmental conditions).'

line 293:

post 1950

Thank you, we will add 'post 1950'.

lines 296-320:

This section is difficult to follow, and it isn't entirely clear what the main point is, other than that LULCC dominates the EED pattern. But more importantly, LULCC seems to drive the fLULCC pattern also, since the EED and the fLULCC patterns also match. temporally, which is does not need a regional breakdown to be shown. And it would appear that the difference in biomass between Pi and Pd is a main driver of the corresponding difference in fLULCC. This boils down to biomass+LULCC controlling fLULCC estimates, with some regional variation in how this relationship contributes to the global estimates.

If the EED pattern is the main point, then take out the other regional plots (5 and 6), as figs 3 and 4 and 7 and 8 show this. The regional plots are difficult to interpret, and so the discussion is helpful, but it should be more clear how the regional patterns contribute to the global patterns, as figure 8 can show (and 7 can show magnitudes better if the scales are matched).

We agree that this section (Sect. 3.2) was not to the point and will change it according to your suggestion. Additionally, we will shift Figures 5-7 into the appendix and create new Figures 5 and 6 with unified y-scales, to ease the comparability. The new section will mainly discuss EED and the environmental effects on estimates of $f_{LULCC_{pi}}$ and $f_{LULCC_{pd}}$ and end with a passage shortly describing the higher complexity when comparing with $f_{LULCC_{trans}}$ (seen e.g. by trend reversal in PTD) and a connecting sentence to the following sections where the positive and negative LASC estimates are discussed as the underlying drivers:

NEW - 'A large sensitivity of cumulative f_{LULCC} towards choice of pre-industrial vs present-day environmental forcing is found in vast stretches across the globe: EED cumulated >8 PgC in the USA (mainly eastern parts), Brazil (mainly southern parts) and Southeast Asia, >5 PgC in Russia, China, Equatorial Africa, Southern Africa, and >2 PgC in Europe (mainly eastern parts), Southwest South America and South Asia from 1800 until 2018 (Figs. 10e and A12). Strikingly, the last decade saw the tropics to become more dominant in positive EED than other regions due to recent clearings

(Figs. 5 and 10f). All these reflect particularly forested areas where LULCCs caused highest f_{LULCC} quantities (Figs. 5 and 9a,c,e; compare increasing deviation of linear model from 1:1 line with higher values) due to the conversion of land with high NBP where positive changes in potential C stocks between 1800 and 2018 occurred (Fig. 8).

Conversely, very distinct regions of negative cumulative EED in Central Europe reflect relatively increased negative f_{LULCC_pd} due to early and widespread reforestation causing increased C uptake (Fig. 9e). Such a strong C uptake due to reforestation causes also globally wide-spread negative EED values in the last decade (Fig. 10f; with hotspots in northeastern Brazil, southern Africa and the Eurasian steppe zone), while the poor representation of recent large-scale reforestation programs with a concomitantly increased C sink in China (Lu et al., 2018; Chen et al., 2019) in the LUH2 data prevents EED (and also f_{LULCC} estimates) to become negative in this region.

Regions, for example remote rain forests, that were only affected little by LULCCs hardly show up in EED. The pattern of EED is thus dominated by the pattern of LULCC with variations due to ecosystem sensitivity (namely in NBP) to environmental conditions (see Fig. 7). This shows that the choice of pre-industrial vs present-day environmental conditions can play a substantial role in regional f_{LULCC} attribution.

As seen for the global estimates, the approach to derive f_{LULCC} under transient environmental conditions introduces even more complexity, as it includes the LASC and strongly depends on the timing of LULCCs. In line, PTD undergoes a trend reversal with widely negative values in the most recent period in many regions (Figs. 6c,d and 10d), which we discuss in detail in the next section.'




Figure 5. Regionwise smoothed multi-model mean annual f_{LULCC_trans} (a&b), f_{LULCC_pi} (c&d), and f_{LULCC_pd} (e&f) from 1800 to 2018, derived according to Eqs. 1 to 3. For discussion on individual models refer to Sect. A1 and Figures A7 to A9. f_{LULCC_pd} was not derived for CLM5.0, JULES, LPJ and OCN (Table 2).




Figure 6. Regionwise smoothed multi-model mean annual Loss of Additional Sink Capacity (LASC; a&b), difference between f_{LULCC} under present-day and pre-industrial environmental conditions (Environmental Equilibrium Difference, EED; c&d), and 'Present-day' vs 'Transient' environmental conditions Difference in f_{LULCC} (PTD; e&f) from 1800 to 2018, derived according to Eqs. 4 to 6. For discussion on individual models refer to Sect. A1 and Figures A4 to A6 and A10 to A12. PTD was not derived for CLM5.0, JULES, LPJ and OCN (Table 2).

Plots 9-11 are relevant, but they are not used to explain why the EED pattern dominates f_{LULCC} , but they can support the basic relationship between biomass, LULCC trajectory, and f_{LULCC} .

There are too many plots that are too difficult to read, and their referencing is difficult to follow also.

We agree and will change the figures in the main manuscript also according to your previous comment and next comments. Thus, there is one less figure in the main part with the two new figures containing only 6 panels (compared to 16 in the earlier version of the regionwise plots). In line, we will change the referencing to new Figures 5 and 6 and rearranged the occurrence of the figures.

lines 302-303:

The regional comparison plots (figures 5 - 7 and the supplemental) are difficult to compare because they are all on different scales. while it may make it more difficult to see some of the individual lines, putting them on the same scale is the best way to show the differences, which I think is the point.

You can split these plots into multiple figures also because they are so small, which also makes them difficult to read. If that is too many figures, then maybe all the regional plots should be supplemental.

Thank you for these suggestions. Accordingly, and to ease the understanding of the main story, we will create new Figures 5 and 6 to show regionwise multi-model mean emission estimates on unified y-scale for each hemisphere. We will shift all region-wise plots showing individual model performances into the appendix; individual model performances in the main script will now only be found for global estimates.

lines 321-352:

This section is difficult to follow, and it isn't clear what the point is of the comparison between LASC and EED. LASC is a component of f_{LULCC} estimation, and EED is a difference between two estimates of f_{LULCC} that don't include LASC. In general, the dynamics discussed here are still driven by biomass+LULLCC, since the forests are the high biomass areas and where LASC will have the most pronounced influence. While the regional breakdown and LASC discussion is somewhat informative, as the LASC dynamics are interesting and a key part of this study and since there is some regional variation, the relationships between LASC and EED seem to still be similar across regions, in that LASC tends to grow over time due to LULCC timing, while EED is driven mainly by biomass differences and the fixed LULCC trajectory. So is the point here to just discuss LASC dynamics, or to support the hypothesis that LASC is critical to f_{LULCC} estimates because static biomass conditions do not include temporal effects of changing conditions (and drive errors associated with these biomass conditions)? I don't think you need the EED stuff here, as your larger point about estimation methods is discussed elsewhere (section 4?).

It is true that LASC and EED are not directly related to each other. Nevertheless, we believe that a comparison of the two is helpful as the EED spans the 'realistic' range of possible f_{LULCC} estimates that can be derived within the historic period. Thus, one could assume that the f_{LULCC} estimates under transient simulations (where, in the end of the simulation, C densities approach the ones under present-day simulations) would never exceed the f_{LULCC} estimates under present-day conditions. However, given the accumulating nature of the LASC, transient f_{LULCC} is widely higher compared to present-day f_{LULCC} in particular for most recent estimates, which by definition results in higher LASC values compared to EED (and consequently a negative PTD).

We will clarify the purpose of Sect. 3.2.1 (and of the following) by rewording of Sect. 3.1 (see comment above) and restructure and modify the text in 3.2.1:

NEW - 'While EED is more relevant than the LASC for cumulative industrial-era emissions (change of sign in correlation in inlet Fig. 10e compared to Fig. 10f), the accumulating LASC heavily alters recent f_{LULCC} estimates – Fig. 10b shows which regions would be attributed much higher emissions when the LASC is included in the f_{LULCC} definition. Aggregated time series for the RECCAP2 regions reveal that the LASC started to increase ~1850 in the USA, Russia and Southeast- and South Asia, ~1900 in SW South- and Central- America and Southern Africa (Figs. 6a,b and A4). It then becomes even more pronounced ~1950 in Brazil, Equatorial Africa and China, with the latter two and Southeast Asia showing a particular strong increase after 2000 (Figs. 6a,b, 10b and A4). Overall, the LASC accumulated to more than 4 PgC in the USA, Brazil, Equatorial Africa and Southeast Asia, and

to 2–4 PgC in China, Russia, SW South- and Central- America, Southern Africa and South Asia (Figs. 10A and A10). As stated above, these high cumulative and annual LASC estimates mainly result from an initial high forest coverage and subsequent C losses in particular on areas where higher C stocks resulted from environmental changes over time (Sect. 3.4 and Fig. 8). Due to the different start of organized human agricultural, the forest clearings in the USA (mid of 19th century, with an early LASC initiation) though on forests with comparably low C stocks (Fig. A3) have caused similar cumulative sums as a much later onset of wide-spread LULCCs (beginning of 20th century) in Brazil, Equatorial Africa, China, and Southeast Asia (Fig. A10) due to rapidly increasing and pronounced higher vegetation C stocks in these regions (with strong response to CO₂ increase).

The widely negative PTD values across the globe for the period 2009–2018 indicate that the LASC causes recent f_{LULCC} estimates from the current DGVM approach (under transient conditions) to be higher compared to bookkeeping estimates (which are similar to f_{LULCC_pd}). However, small areas exist where EED remains larger than the LASC (i.e. positive PTD values) for the recent decade, here, more recent LULCCs caused even shorter period for the LASC to accumulate: in the tropics (mainly Brazil, Tanzania, Indonesia), sub-tropics (Eastern China, Southern Australia), and in the transition zones from temperate to boreal zone (Scandinavia, Russia). These regions would likely be attributed higher emissions by bookkeeping approaches than by f_{LULCC_trans} from DGVMs. This highlights another difficulty especially in regional f_{LULCC} attribution: as the LASC accumulates emissions caused by past LULCCs, recent LULCCs are given less weight in relative terms. This also applies to recent LULCCs reducing atmospheric CO₂ such as reforestation, which cannot quickly compensate for past LULCC in approaches including the LASC, while they could in f_{LULCC_pi} and f_{LULCC_pd} estimates.'

lines 353-377:

This discussion is a bit more clear as it focuses on LASC dynamics and doesn't mix in references to other calculations. I suggest combining this section and the previous section and focusing just on LASC dynamics. And the last sentence seems less a concluding sentence and more an evidence sentence to support the discussion.

Due to the outstanding importance of the LASC for f_{LULCC} estimation under transient environmental conditions, we aim to keep the structure of the manuscript. In Sect. 3.2.1 we discuss the well-known LASC and associated lost C sinks, for the first time on a spatially explicit level, with the aim to highlight regions whose f_{LULCC} estimates show strongest susceptibility to LASC inclusion and explain why the LASC can be greater than EED. In the following section (Sect. 3.2.2), we discuss the not so well-known occurrence of negative LASC values in some regions and particularly in most recent periods. To make this more clear we will rephrase several parts (compare also earlier comments) and add the following concluding sentence:

NEW - 'Comparing the different f_{LULCC} estimates over time and across space, previous discussion has shown that the choice of method to derive f_{LULCC} strongly impacts the estimated quantities. The effects of the interaction of the environmental forcing with the timing of the actual LULCC is

particularly pronounced for estimates under transient environmental forcing and where NBP was strongly altered by environmental changes.'

line 383-384:

What about LULCC effects due to abandonment of agricultural land? This also contributes to higher biomass in some places, and isn't trivial. You show some evidence of this in figure 12, where it can outweigh the climatic effects in even the long term, but your net results over the whole period do not show how regrowth has contributed more recently to increased biomass in places where this doesn't outweigh previous emissions.

Indeed, some areas experienced LULCCs that were beneficial for the carbon stocks, such as abandonment of agricultural land. However, for the derivation of CO₂- vs climate-related f_{LULCC} shares we used the total stock alterations of the S1 and S2 simulations, both of them have fixed pre-industrial LULCC forcing. Due to the fixed LULCC forcing, effects from LULCCs can be excluded, which we will explicitly state in the updated manuscript:

NEW - 'The mean simulated global vegetation C stock increased by ~23% from 664 PgC to 815 PgC from 1800 until today, in both the S1 and S2 simulation (which by protocol exclude effects from LULCCs; see Figs. 8 and A3 for maps and Fig. A1b for global estimates).'

lines 416-445:

This section seems to be referring to a political question for emission attribution to regions for policy, rather than a scientific question of what are LULCC emissions.

Thank you, we will correct the wording and highlight the political context of this discussion as follows:

NEW - 'From a policy standpoint and disregarding considerations from the natural land sink perspective, these results highlight the need for a f_{LULCC} estimate that is comparable over time and across space.'

NEW - 'Within the political context, if environmental effects on potential C stocks should be accounted for, we argue to include the LASC into regional budgets, thereby highlighting the need to use DGVMs for the assessment.'

So you should separate these two questions/recommendations. Is your scientific recommendation to discard some LULCC emissions? It doesn't seem to be so as you clearly show that LASC is an important component in "Accurate quantification of the net carbon flux from land use and land cover changes..." (line 447). Whether this makes sense for regional carbon policy is a different question.

From a policy standpoint, the question here is then why include only part of the total emissions from LULCC (i.e. LASC from second half of 21st century) while including some estimate of the directly effects of changes all the way back to 1850? If it isn't ok to include LASC back to 1850, why

should any effects of pre-1950 LULCC be included? Why not just start full attribution of emissions at 1950?

Thank you. We agree that the definition of the reference period only for the LASC was somehow arbitrary chosen. From a political standpoint, and in line with your suggestion, one could argue to derive not only the LASC but also f_{LULCC} for such a unified reference period. Considering your comment and a comment of Reviewer #2, we will name different possibilities and carefully rephrase the whole paragraph as follows (in mind that this is a political decision which we leave open):

NEW - 'To circumvent these issues, as could be desired in the political context, one could derive f_{LULCC} and the LASC based on simulations forced with the cycled climate and CO₂ conditions that occurred during the actual LULCC event. However, this would still result in differing accumulation periods and varying environmental conditions during and following a LULCC event. While the influence of the latter could be reduced using cycled pre-industrial or present-day environmental forcings, these neglect transient C stock changes. To consider the LASC but counteract spatial heterogeneity in f_{LULCC} differences resulting from synergistic effects of environmental conditions and the timing of LULCCs, one could derive f_{LULCC} and LASC from a defined reference period which is independent of the actual time that LULCCs occurred and shares the same reference conditions. For example, f_{LULCC} and LASC could always be modelled for the second half of the 21st century, as here the environmental C stock changes have been amplified due to the accelerating increase of atmospheric CO₂ concentrations (alternative start times are of course conceivable). By using such reference period, the LASC could fully be captured also for most recent LULCCs (may they act positive or negative on C stocks) and foregone sinks would be more equally counted (same length of accumulation period with similar environmental changes). Along these lines, it may be considered to calculate such adapted LASC based on CO₂-only simulations as here the impact of humans is more homogeneously distributed, while the spatially heterogeneous climate impact on f_{LULCC} , determined foremost by action outside the location of LULCCs, causes a questionable attribution of regional f_{LULCC} when compared across the globe (without even considering externalized f_{LULCC} s e.g. due to remote market demand of food and timber; Lambin and Meyfroidt 2011; Meyfroidt et al., 2013). To detach f_{LULCC} estimates from the timing of LULCCs and the spatially heterogeneous climate evolution, we argue to address the delineation of an adapted LASC in future studies, where, in particular, the reference to calculate the LASC should further be investigated. Such methodology could limit f_{LULCC} to locally determined factors (namely LULCCs) and reduce the dependence on the timing of LULCCs while still reflecting the foregone C sink capacity by human intervention.'

liens 467-472:

Here you clearly state that full accounting includes LASC. Then you vaguely backtrack and mention the derived LASC without context, which is the political context of attribution for policy. The "arbitrary length of their simulation period" is a given fact for all modeling and is handled by

explicit definition of the period in question. For example, climate change is defined in reference to the “pre-industrial” period (usually pre-1850 equilibrium).

Thank you, we apologize for the incorrect wording (in particular the misleading description of ‘full accounting’) and the missing link for the political context of attribution for policy. We will change the sentences as follows:

NEW – ‘Within the political context, if environmental effects on potential C stocks should be accounted for, we argue to include the LASC into regional budgets, thereby highlighting the need to use DGVMs for the assessment. As LASC values derived by the approach so far taken in the GCB are widely independent of locally determined environmental changes (rather depend on globally determined climatic changes) and strongly dependent on accumulation periods (defined by the timing of LULCCs and the end year of the simulations), we argue for a f_{LULCC} attribution that is more robust against choices of environmental drivers and accumulation period by using an adapted LASC, for example, based on a defined common reference period and homogeneously altered environmental conditions (such as only driven by CO₂ alterations).’

Figures

Your figures and supplemental figures are out of order with respect to references in the text.

We will change the order of the figures in the main text by adjusting the appearance and order of tables (Table 1 is changed with Table 2) and figures (shifting Figures 5-7 into the appendix, and creating new Figures 5 & 6). Additionally, we will add additional references in the text at several places and put additional labels in the figures to ease comparability.

Consider using the same scale across the regional comparison plots, both in the paper and in the supplemental.

Thank you for this suggestion. As highlighted before, we will create new Figures 5 and 6 showing region-wise multi-model mean emission estimates on unified y-scale for each hemisphere. However, since the emissions quantities of the different regions are extremely different (> factor of ten variation), a clear depiction of individual models on the same scale for each region is not feasible. Therefore, we suggest to keep the region-wise plots showing individual model performances on individual scales, but now assemble all of them in the appendix section.

Appendix A1

lines 484-485:

This statement appears incorrect based on the discussion that follows.

It appears that the variation may arise from models with different ecological and productivity responses to the cycled environmental conditions.

Thank you, we will shift the respective statement into Sect. 3.1 and rephrase it (in line with a comment from Reviewer #2) as follows:

NEW - 'Note, a high internal climate variability translated into a high interannual variability in NBP and consequently a high variability of f_{LULCC} estimates (Figs. 3, 5 and A13) and of their respective differences (Figs. 4 and 6). For the differences in f_{LULCC} estimates, some artefact might additionally arise due to comparison of simulations with different forcing cycles (e.g. on global scale, with periodic fluctuations in annual relative shares of EED to f_{LULCC_pd} in Fig. 2c and, on regional scale, in Figs. 6 and A4 to A6 with pronounced oscillations in some regions).'

#####

In addition to changes resulting from the reviewer comments, we suggest the following changes:

- change to capitalized journal abbreviations in references and added doi that where missing
- add new Table 3 with overview of different estimates

#####

New references:

Devaraju, N., Bala, G., Caldeira, K., and Nemani, R.: A model based investigation of the relative importance of CO₂-fertilization, climate warming, nitrogen deposition and land use change on the global terrestrial carbon uptake in the historical period, CLIM DYNAM, 47, 173–190, <https://doi.org/10.1007/s00382-015-2830-8>, 2016.

Fernández-Martínez, M., Sardans, J., Chevallier, F., Ciais, P., Obersteiner, M., Vicca, S., Canadell, J., Bastos, A., Friedlingstein, P., Sitch, S., et al.: Global trends in carbon sinks and their relationships with CO₂ and temperature, NAT CLIM CHANGE, 9, 73–79, <https://doi.org/10.1038/s41558-018-0367-7>, 2019.

Lombardozzi, D. L., Bonan, G. B., Levis, S., and Lawrence, D. M.: Changes in Wood Biomass and Crop Yields in Response to Projected CO₂, O₃, Nitrogen Deposition, and Climate, J GEOPHYS RES-BIOGEO, 123, 3262–3282, <https://doi.org/10.1029/2018JG004680>, 2018.

Obermeier, W. A., Lehnert, L. W., Kammann, C., Müller, C., Grünhage, L., Luterbacher, J., Erbs, M., Moser, G., Seibert, R., Yuan, N., et al.: Reduced CO₂ fertilization effect in temperate C3 grasslands under more extreme weather conditions, NAT CLIM CHANGE, 7, 137–141, <https://doi.org/10.1038/nclimate3191>, 2017.