



# Is time a variable like the others in multivariate statistical downscaling and bias correction?

Robin Yoann<sup>1,\*</sup> and Vrac Mathieu<sup>2,\*</sup>

<sup>1</sup>Météo-France, 42 avenue Gaspard-Coriolis, 31057, Toulouse, France

<sup>2</sup>Laboratoire des Sciences du Climat et de l'Environnement (LSCE-IPSL), CEA/CNRS/UVSQ, Université Paris-Saclay, Centre d'Etudes de Saclay, Orme des Merisiers, 91191 Gif-sur-Yvette, France

\*The two authors contributed equally to this study

**Correspondence:** Robin Y. (yoann.robin@meteo.fr), Vrac M. (mathieu.vrac@lsce.ipsl.fr)

**Abstract.** Bias correction and statistical downscaling are now regularly applied to climate simulations to make them more usable for impact models and studies. Over the last few years, various methods were developed to account for multivariate – inter-site or inter-variable – properties in addition to more usual univariate ones. Among such methods, temporal properties are either neglected or specifically accounted for, i.e., differently from the other properties. In this study, we propose a new multivariate approach called “Time Shifted Multivariate Bias Correction” (TSMBC), which targets to correct the temporal dependency in addition to the other marginal and multivariate aspects. TSMBC relies on considering the initial variables at various times (i.e., lags) as additional variables to correct. Hence, temporal dependencies (e.g., auto-correlations) to correct are viewed as inter-variable dependencies to be adjusted and an existing multivariate bias correction (MBC) method can then be used to answer this need. This approach is first applied and evaluated on synthetic data from a Vector Auto Regressive (VAR) process. In a second evaluation, we work in a “perfect model” context where a Regional Climate Model (RCM) plays the role of the (pseudo-) observations, and where its forcing Global Climate Model (GCM) is the model to be downscaled/bias corrected. For both evaluations, the results show a large reduction of the biases in the temporal properties, while inter-variable and spatial dependence structures are still correctly adjusted. However, increasing too much the number of lags to consider does not necessarily improve the temporal properties and a too strong increase in the number of dimensions of the dataset to correct can even imply some potential instability in the adjusted/downscaled results, calling for a reasoned use of this approach for large datasets.

## 1 Introduction

Climate and Earth System models (ESM) and their simulations are the main physical tools to investigate the potential future evolutions of the climate system (e.g., Flato et al., 2013; Kirtman et al., 2013). They are clearly indispensable to test how different scenarios of greenhouse gas emission trajectories might induce climate changes and, thus, to try to anticipate potential



impacts of those changes (e.g., Shukla et al., 2019). Although such elaborated models contain many relevant and complex processes characterizing the climate properties and their dependencies, the numerical simulations they generate are often tainted with biases and disagreements with respect to observations. Those can stem: (i) from the spatial resolution of the simulations – from 200km×200km for global climate models, GCMs, down to a few km for regional climate modes, RCMs –, usually too low compared to needs of impact models that may require very local or high-resolution input climate data, e.g., kilometre, hundreds of meters or below, down to the weather stations (e.g., Chen et al., 2011; Maraun and Widmann, 2018); and/or (ii) from inherent biases in the model simulations. Even with respect to observations or reanalysis data at the models’ spatial resolution, the simulations can present biases, i.e., disagreements on some statistical properties (e.g., mean, variability, distribution) that differ from reference data (e.g., Christensen et al., 2008).

Those issues make that impact models (such as for hydrology, energy, environment, etc.) cannot directly employ the climate simulations as input (e.g., Teutschbein and Seibert, 2012; Chen et al., 2013). Indeed, most impact models are calibrated on observational data and the use of biased climate data as input could result in inappropriate or even false impact projections. To overcome those spatial or inherent biases, various statistical post-processing methods have been developed for downscaling and/or bias correction of the climate simulations. The common idea is to statistically transform the numerical simulations – over a historical period – in such a way that some properties become equal or close to those of a chosen reference dataset (e.g., Gudmundsson et al., 2012). The statistical downscaling or correction transformation is then usually supposed valid under other climate conditions and applied to changed (e.g., future) climate. The obtained downscaled/bias corrected climate data can then serve as input into impact models.

Over the last two decades, many such post-processing methods were developed, either in a “Perfect prognosis” (PP) context, generally for downscaling (DS), or in a “Model Output Statistics” (MOS) one, generally for bias correction (BC) – see e.g., Vaittinada Ayar et al. (2015) or Maraun and Widmann (2018) for the differences between the two, or Vrac and Vaittinada Ayar (2017) for a combination of PP and MOS. Note that, in practice, BC methods (i.e., MOS) are often used to perform downscaling (see, e.g., Thrasher et al., 2012; Hempel et al., 2013; Frieler et al., 2017). Hence, in the following, we will simply refer to “bias correction” (hereafter BC) even for downscaling purposes. Up to recently, most of the BC methods were designed to work on “univariate” data, representing one climate variable at one given location. In such a univariate context, PP approaches include, e.g., linear regressions (Jeong et al., 2012), or non-linear ones such as polynomial regressions and artificial neural network (Xiaoli et al., 2008), or stochastic weather generators (Wilks, 2012). In a similar univariate context, MOS methods can vary from very simple methods correcting only the mean via anomalies adjustments (e.g., Xu, 1999, with the “anomaly” method), or the variance (e.g., Eden et al., 2012; Schmidli et al., 2006, with the “simple scaling” method) to more complete and widespread methods such as the “quantile-mapping” approach that corrects the whole univariate distribution and therefore all statistical moments (including mean and variance) of the variable of interest. Quantile-mapping (e.g., Haddad and Rosenfeld, 1997; Déqué, 2007) is certainly the most used 1d-BC method by practitioners and numerous variants have been developed, e.g., by Vrac et al. (2012) to account for non-stationarity, by Kallache et al. (2011) for modelling of extremes, by Cannon



et al. (2015) to ensure preservation of the climate change signal, or by Vrac et al. (2016) specifically for precipitation including occurrences, among many other variants.

60 However, the univariate correction of simulations (i.e., one variable at a time and one site at a time) may not be enough. Indeed, the use of several 1d-corrections separately for different physical variables and/or sites will not correct the dependencies between them (Vrac, 2018). The consequence is that, if the simulations are biased in their spatial and/or inter-variable correlations (or more generally in their dependencies), most of the 1d-BC methods will conserve those biases (Vrac, 2018; François et al., 2020). If an impact (e.g., hydrological) model needs realistic dependencies between its input climate variables, 65 the use of univariate BC may not provide corrections realistic enough (e.g. Boé et al., 2007). More generally in climate sciences, the accurate modelling of dependencies is a key aspect for proper assessments and projections of compound events and their associated risks (e.g. Leonard et al., 2014; Zscheischler et al., 2018; Bevacqua et al., 2019). This characteristics strongly motivated the development of BC methods accounting for multivariate links between the variables and/or the sites of interest. As detailed in Vrac (2018) or François et al. (2020), Multivariate BC methods can be categorized in three approaches: the 70 “marginal/dependence” correction approach – separating the correction of the marginal distributions and that of the dependence structure (e.g., Bárdossy and Pegram, 2012; Cannon, 2018; Vrac, 2018) –, the “successive conditional” correction one – applying successive corrections conditionally on the previously corrected variables (e.g., Piani and Haerter, 2012; Dekens et al., 2017) – and the “all-in-one” correction approach – correcting altogether the univariate marginal properties and their dependence structures (Robin et al., 2019). Those methodologies have been applied to correct either inter-variable dependence 75 only (i.e., multiple variables but spatial dependence not accounted for), spatial dependence only (one variable at multiple locations) or even both (see François et al., 2020, for a review and comparison of the methods).

One main conclusion provided in the MBC comparison study brought by François et al. (2020) is that if inter-site and inter-variable properties can be reasonably adjusted by multivariate bias correction (MBC) methods, the temporal properties are 80 usually not taken into account in most MBC procedures. As any MBC will necessarily modify the rank chronology of the simulations to perform the multivariate correction (Vrac, 2018), it results that temporal properties and dependencies issued from the MBC output are transformed with respect to the raw simulations but are not necessarily closer to those of the reference data, depending on the method itself, its setting and the variables of interest (François et al., 2020). Therefore, accounting for temporal properties (i.e., autocorrelations or even cross-autocorrelations) when performing MBC is needed to increase the 85 realism of the multivariate corrected time series. This is a major concern for impact studies: for example, hydrological models are very sensitive to the atmospheric forcing used as inputs and particularly, among others, to biases in the chronology of events (e.g., Raimonet et al., 2017; Bhuiyan et al., 2018). Some works tried to correct temporality properties in addition to (spatial or inter-variable) MBC. For example: Johnson and Sharma (2012) used a nesting 1d-BC model across various timescales; this was extended to include inter-site structures by Mehrotra and Sharma (2015) and then by Mehrotra and Sharma (2016) 90 to account also for inter-variable dependence. In those studies, time dependence was specifically modeled via (multivariate) autoregressive models with periodic parameters. Another recent study by Vrac and Thao (2020) proposed an MBC method



for both inter-site and inter-variable properties, where the temporal aspects are accounted for through an analog-based method applied to multivariate sequences of ranks.

95 All those approaches, although different, share the fact that they try to correct temporal properties by a (parametric or non-  
parametric) model that is specific to the variable “time”. In other words, they separate time from the other variables (variables  
at various locations) of interest. However, one can wonder if there is a real need for such a specificity. Indeed, let us take an  
example, in one dimension for sake of clarity (this can be easily generalised to  $n$  dimensions). Say we have a time series of  
a climate variable  $\mathbf{X}$  to be corrected (e.g., temperature) at one location, for  $t = 1, \dots, N$ :  $\mathbf{X}_{1:N} = (x_1, \dots, x_N)$ . If we apply a  
100 1d-BC method to the sample  $\mathbf{X}_{1:N}$ , most of the BC methods will preserve the rank correlation of  $\mathbf{X}$  (Vrac, 2018). Now, instead  
of applying a 1d-BC, if we duplicate and lag – say of one time step – the time series  $\mathbf{X}_{1:N}$ , we obtain a shifted time series  
noted  $\mathbf{X}_{1:N-1}^S = \mathbf{X}_{2:N} = (x_2, \dots, x_N)$ . If a multivariate bias correction method is able to correct the dependence between two  
univariate time series, say  $\mathbf{X}$  and  $\mathbf{Y}$ , when applied to the bivariate time series

$$(\mathbf{X}, \mathbf{X}^S)_{1:N-1} = \begin{pmatrix} x_1 & x_2 \\ x_2 & x_3 \\ \vdots & \vdots \\ x_{N-1} & x_N \end{pmatrix}, \quad (1)$$

105 it should then be able to correct the dependence between the univariate time series  $\mathbf{X}$  and  $\mathbf{X}^S$ , i.e., it should be able to correct  
the lag-1 temporal dependence of  $\mathbf{X}$ . As far as we know, this specific approach of considering time as any other statistical vari-  
able in an MBC procedure has never been investigated. This is therefore the goal of the present study. More specifically, the  
aim is to investigate (i) if this “time shifted multivariate bias correction” (TSMBC) approach does correct temporal properties  
(e.g., auto-correlations); (ii) if and how it impacts other properties (e.g., inter-site and inter-variable); (iii) if this TSMBC is  
110 able to work for more than lag-1.

To do so, the rest of the manuscript is organized as follows: Section 2 describes the climate and synthetic data used in this  
study and provides detailed explanations about the proposed TSMBC approach. Section 3 investigates the method based on  
artificially simulated multivariate dataset obtained from a Vector Autoregressive model (VAR). Then, TSMBC is applied to  
115 correct daily temperature and precipitation simulated from a climate model and the results are given in Section 4. Finally,  
conclusions are reminded and perspectives are discussed in Section 5.

## 2 Data and methodology

### 2.1 Data: Climate simulations and VAR processes

To apply any (M)BC method, it is necessary to dispose of a dataset of reference – i.e., supposed to be as close as possible to  
120 real observed climate – and a dataset of simulations (e.g., stemming from a GCM) that are biased with respect to the reference



dataset.

Here, the climate simulations to be corrected are daily temperature and precipitation times series over South-East of France, for the time period 1951-2010, extracted at a  $1.25^\circ \times 2.5^\circ$  spatial resolution from the IPSL-CM5A-MR global climate model (Marti et al., 2010; Dufresne et al., 2013) developed at “Institut Pierre Simon Laplace” (IPSL). The simulations are forced by historical conditions up to 2005 and then by RCP8.5 scenario conditions from 2006. The region of interest is presented in Fig 1 and corresponds to a zone where correlations between precipitation and temperature as well as autocorrelations are difficult to model due to various geographical constrains, such as Mediterranean influences and 3 mountain ranges (the Pyrenees, the Massif Central, the Alps), which can generate high precipitation events known as “Cevennol events” (see, e.g. Delrieu et al., 2005).

Regarding the reference dataset, regional climate simulations are used in this study, instead of observational or reanalysis data. Those are daily temperature and precipitation from the KNMI-RACMO22E regional climate model (KNMI, 2017) developed at “The Royal Netherlands Meteorological Institute” (KNMI) and forced by the larger-scale IPSL simulations. The same time period and region as for the IPSL simulations were extracted but with a  $0.11^\circ \times 0.11^\circ$  spatial resolution.

The GCM data are then interpolated with a nearest neighbors method to the  $0.11^\circ \times 0.11^\circ$  spatial resolution of the RCM to apply the TSMBC approaches. The (GCM and reference) data over the period 1951-1980 will serve to calibrate the proposed TSMBC method, while projections and evaluations will be performed over 1981-2010. Moreover, the calibration/projection steps are realized on a seasonal basis, i.e., for the 4 seasons separately to account for specific seasonal features of the biases. For sake of clarity, only the summer (JJA) results are shown in the manuscript, winter (DJF) results being provided as supplementary materials.

In addition to the evaluations that will be done based on those climate simulations, a preliminary analysis will first be performed on synthetic data, i.e., data artificially generated from statistical models. A “Vector autoregressive” (VAR) process is employed to generate multivariate time series  $(\mathbf{X}_t)_{t=1,\dots,N}$  where each  $\mathbf{X}_t$  is a vector of dimension  $d$  (e.g.,  $d = 2$  with temperature and precipitation at one location, or with temperature only at two locations), with prescribed autocorrelations up to a certain lag  $s$ , such that:

$$\mathbf{X}_n = \mathbf{b} + \sum_{l=1}^s \mathbf{A}_l \mathbf{X}_{n-l} + \boldsymbol{\epsilon} \quad (2)$$

where  $s$  is the order of the VAR process for which the autocorrelation is accounted for,  $\mathbf{b}$  is the  $d$ -dimensional intercept vector,  $\mathbf{A}_l$  are matrices of coefficients and  $\boldsymbol{\epsilon}$  is a  $d$ -dimensional noise following a multivariate Gaussian law with 0 mean vector and covariance matrix  $\boldsymbol{\Sigma}$ . Two VAR processes are generated: one used as the large-scale or biased simulations and the other one used as the reference.



To generate such synthetic data and analyse them in a comprehensive way, the dimension  $d$  has been chosen equal to 2 and lag  
 155  $s$  equal to 3. Also, in order to be in a realistic case, i.e., close to our climate simulations context, the parameters  $\mathbf{b}$ ,  $(\mathbf{A}_l)_{l=1\dots 3}$   
 and  $\Sigma$  have been estimated from a VAR fitted to temperature at two opposite grid points in summer (from the GCM or RCM,  
 for VARs representing biased or reference data respectively) in the region of interest. The parameters obtained for these two  
 VARs are then used to simulate 2500 synthetic data for each of the two VAR processes, approximately corresponding to the  
 number of summer days over the available calibration period. The univariate and bivariate probability density functions of  
 160 those simulations are plotted in Fig. 2(a), in red for VAR data to be corrected and in blue for the reference VAR.

## 2.2 The TSMBC approach: autocorrelations seen as correlations

The main philosophy of the proposed “multivariate time shifted bias correction” (TSMBC) approach has been briefly intro-  
 duced (with lag 1) in Eq. (1) of the introduction section. We now described the approach in a more general way and with  
 165 more details. Say we dispose of a  $d$ -dimensional time series (i.e., matrix)  $\mathbf{X}_{1:N} = (\mathbf{X}^1, \mathbf{X}^2, \dots, \mathbf{X}^d)_{1:N}$ , where each  $\mathbf{X}^i_{1:N}$  is a  
 $N$ -dimensional vector (i.e., time series of length  $N$ ):  $(x_1^i, \dots, x_N^i)^T$ , where the superscript  $T$  on a vector denotes its transpose.  
 The idea proposed and tested in this article consists in applying a multivariate bias correction onto the matrix  $\mathbf{M}(s)$  constituted  
 by the gathering of the initial  $\mathbf{X}_{1:N}$  time series and those obtained after shifting it by different lags up to lag  $s$ :

$$\mathbf{M}_{\mathbf{X}}(s) = \left( (\mathbf{X}^1, \mathbf{X}^2, \dots, \mathbf{X}^d)_{1:N-s}, (\mathbf{X}^1, \mathbf{X}^2, \dots, \mathbf{X}^d)_{2:N-s+1}, \dots, (\mathbf{X}^1, \mathbf{X}^2, \dots, \mathbf{X}^d)_{s+1:N} \right) \quad (3)$$

170 or equivalently:

$$\mathbf{M}_{\mathbf{X}}(s) = \left( \left( \begin{array}{cccc} x_1^1 & x_1^2 & \dots & x_1^d \\ x_2^1 & x_2^2 & \dots & x_2^d \\ \vdots & \vdots & \ddots & \vdots \\ x_{N-s}^1 & x_{N-s}^2 & \dots & x_{N-s}^d \end{array} \right), \left( \begin{array}{cccc} x_2^1 & x_2^2 & \dots & x_2^d \\ x_3^1 & x_3^2 & \dots & x_3^d \\ \vdots & \vdots & \ddots & \vdots \\ x_{N-s+1}^1 & x_{N-s+1}^2 & \dots & x_{N-s+1}^d \end{array} \right), \dots, \left( \begin{array}{cccc} x_{s+1}^1 & x_{s+1}^2 & \dots & x_{s+1}^d \\ x_{s+2}^1 & x_{s+2}^2 & \dots & x_{s+2}^d \\ \vdots & \vdots & \ddots & \vdots \\ x_N^1 & x_N^2 & \dots & x_N^d \end{array} \right) \right), \quad (4)$$

which can then be grouped by variable (dimension) and thus reordered as:

$$\mathbf{M}_{\mathbf{X}}(s) = \left( \left( \begin{array}{cccc} x_1^1 & x_2^1 & \dots & x_{s+1}^1 \\ x_2^1 & x_3^1 & \dots & x_{s+2}^1 \\ \vdots & \vdots & \ddots & \vdots \\ x_{N-s}^1 & x_{N-s+1}^1 & \dots & x_N^1 \end{array} \right), \left( \begin{array}{cccc} x_1^2 & x_2^2 & \dots & x_{s+1}^2 \\ x_2^2 & x_3^2 & \dots & x_{s+2}^2 \\ \vdots & \vdots & \ddots & \vdots \\ x_{N-s}^2 & x_{N-s+1}^2 & \dots & x_N^2 \end{array} \right), \dots, \left( \begin{array}{cccc} x_1^d & x_2^d & \dots & x_{s+1}^d \\ x_2^d & x_3^d & \dots & x_{s+2}^d \\ \vdots & \vdots & \ddots & \vdots \\ x_{N-s}^d & x_{N-s+1}^d & \dots & x_N^d \end{array} \right) \right). \quad (5)$$



Such a transformation can be made to create  $\mathbf{M}_Y(s)$  and  $\mathbf{M}_X(s)$ , the lagged matrices obtained respectively from the reference  
 175 dataset and the biased one to correct. An MBC method can thus be applied to correct the multivariate properties of  $\mathbf{M}_X(s)$   
 with respect to  $\mathbf{M}_Y(s)$  as reference. The result of the MBC is therefore a matrix  $\mathbf{M}_Z(s)$  of the same size as  $\mathbf{M}_X(s)$ . The ques-  
 tion to answer now is: “From this result MBC matrix, how to extract/reconstruct a  $N \times d$  matrix  $\mathbf{Z}$  that corresponds to proper  
 multivariate (time included) corrected data?”. There are indeed various possibilities for that and they may not be equivalent.

180 In this study, we propose a method based on a reconstruction *by rows*. For illustration, let us take this example of a bivariate  
 dataset  $\mathbf{X}$  and its associated matrix  $\mathbf{M}_X(1)$ , i.e., including a lag-1 shift:

$$\mathbf{X} = \begin{pmatrix} 1 & 10 \\ 2 & 20 \\ 3 & 30 \\ 4 & 40 \\ 5 & 50 \\ 6 & 60 \end{pmatrix}, \mathbf{M}_X(1) = \begin{pmatrix} \mathbf{1} & \mathbf{10} & \mathbf{2} & \mathbf{20} \\ 2 & 20 & 3 & 30 \\ \mathbf{3} & \mathbf{30} & \mathbf{4} & \mathbf{40} \\ 4 & 40 & 5 & 50 \\ \mathbf{5} & \mathbf{50} & \mathbf{6} & \mathbf{60} \end{pmatrix}$$

In this example, the bold rows of the matrix  $\mathbf{M}_X(1)$  (its rows 1, 3 and 5) can be used to reconstruct the original dataset  $\mathbf{X}$ . More  
 generally, a row  $s$  of  $\mathbf{M}_X(1)$  is a vector corresponding to a portion of  $\mathbf{X}$ , which continues at row  $s+2$ , and which continues at  
 185 row  $2s+3$ , etc. Concatenating these rows, we can re-construct the dataset  $\mathbf{X}$ . Applying this method to the matrix  $\mathbf{M}_Z(s)$ , we  
 can reconstruct a corrected dataset  $\mathbf{Z}$ . Because the same operation can be applied by starting at the second row (and any row  
 between 1 and  $s+1$ ), the reconstruction depends of the choice of the starting row. This leads us to wonder about the influence of  
 the choice of the starting row. This point will be investigated with the VAR processes. Our approach is summarized in the Alg. 1.

190 Note that a reconstruction “*by column*” could also be performed: each column of  $\mathbf{M}_X(1)$  being a sub-column of  $\mathbf{X}$ , it could  
 then be used to reconstruct the original dataset. However, preliminary analyses showed that this approach does not allow to  
 correct auto-correlations, as the temporal dependence structure in each row is not corrected and mostly corresponds to that of  
 the model to adjust (not shown). Hence, only the TSMBC approach “*by rows*” is investigated in the following.

195 Finally, because TSMBC uses an underlying MBC, potentially any MBC method can be used, as MBCn (Cannon, 2018)  
 or R2D2 (Vrac and Thao, 2020). Here, the MBC method used is the “dynamical Optimal Transport Correction” (dOTC)  
 developed by Robin et al. (2019). While most of the bias correction methods build a mapping between the biased and the  
 reference dataset, dOTC infers a probability distribution  $\mathbb{P}$  such that  $\mathbb{P}(\mathbf{x}, \mathbf{y})$  is *the probability that  $\mathbf{y}$  is the correction of  $\mathbf{x}$* .  
 This probability distribution is inferred – with optimal transport methods, see App. A and Villani (2008); Santambrogio (2015)  
 200 – between the biased and reference dataset in calibration period (representing the bias), and between the biased dataset in  
 calibration and projection periods (representing the evolution). The “evolution” distribution is then transferred along the “bias”  
 distribution to construct a correction in the reference world with an evolution similar to that of the biased data. Note that this



---

**Algorithm 1** Summary of the TSMBC method “by rows”

---

**Require:** A  $N_{\mathbf{X}} \times d$  biased dataset  $\mathbf{X}$ , where  $N_{\mathbf{X}}$  is the number of time steps for  $\mathbf{X}$ , and  $d$  the number of variables

**Require:** A  $N_{\mathbf{Y}} \times d$  reference dataset  $\mathbf{Y}$ , where  $N_{\mathbf{Y}}$  is the number of time steps for  $\mathbf{Y}$

**Require:** A multivariate bias correction method, dOTC in this paper

**Require:** A lag  $s$  and a starting row  $r$

Build the  $(N_{\mathbf{X}} - s) \times d(s + 1)$  matrix  $\mathbf{M}_{\mathbf{X}}(s)$  from  $\mathbf{X}$  with Eq. (5)

Build the  $(N_{\mathbf{Y}} - s) \times d(s + 1)$  matrix  $\mathbf{M}_{\mathbf{Y}}(s)$  from  $\mathbf{Y}$  with Eq. (5)

Apply the multivariate bias correction method between  $\mathbf{M}_{\mathbf{X}}(s)$  and  $\mathbf{M}_{\mathbf{Y}}(s)$ , generating the  $(N_{\mathbf{X}} - s) \times d(s + 1)$  matrix  $\mathbf{M}_{\mathbf{Z}}(s)$

Apply the reconstruction with the row method with starting row  $r$  to the matrix  $\mathbf{M}_{\mathbf{Z}}(s)$ . This generates the correction  $\mathbf{Z}$ .

**return** The  $N_{\mathbf{X}} \times d$  multivariate correction  $\mathbf{Z}$ .

---

method seeks to preserve the evolution of the model while reducing bias. All details about the dOTC method can be found in Robin et al. (2019).

## 205 3 Results: Synthetic VAR data

In this section we test our TSMBC method on synthetic data, generated from two VAR processes (see sub-section 2.1). A biased dataset and a reference one – noted respectively  $\mathbf{X}$  and  $\mathbf{Y}$  – are drawn from two VAR processes fitted from two time series of temperature of the GCM/RCM. TSMBC is applied to correct  $\mathbf{X}$  with respect to  $\mathbf{Y}$ , i.e., to generate a corrected dataset  $\mathbf{Z}$ .

210 Because the reconstruction step preserves the dependence structure, we propose to test which part of the correction is due to the underlying method (here dOTC), and which part is due to the reconstruction. To do so, a second underlying bias correction method is then used as a benchmark. It corresponds to a very naive method: the correction is randomly drawn from the reference dataset. Note that this naive method corresponds to dOTC with a transport plan given by the product probability distribution between the biased/reference dataset. We note this multivariate method a “Random Bias Correction” (RBC), and TSMBC is  
215 thus available in two versions for this evaluation: TSMBC(dOTC) and TSMBC(RBC). For these two methods, we test in a first part the influence of the choice of the starting row, and in a second part the influence of the choice of the lag.

### 3.1 Influence of the choice of the starting row

We fix the number of lags  $s = 10$  and we compute the correction for each of the two methods with a starting row  $r \in \{1, 3, 6, 9, 12\}$ . Note that, since here  $s = 10$ , TSMC applied with  $r = 12$  will provide the same results as for  $r = 1$  but will  
220 ignore the first row. The corrections are noted  $\mathbf{Z}_r$ .

To measure the similarity of the corrections from different starting rows, we compute the matrix of Pearson correlations between the pair  $\mathbf{Z}_{r_1}/\mathbf{Z}_{r_2}$ . We also compute the correlations with  $\mathbf{X}$  and  $\mathbf{Y}$  to compare the corrections with the biased and reference datasets. The results are presented in Fig. 3a for TSMBC(dOTC) and in Fig. 3b for TSMBC(RBC). Only the first





dimension is represented, the second dimension – which gives similar results – and the associated  $p$ -values are given in Fig. S1  
225 of the supplementary material. We can see for TSMBC(dOTC) that all corrections are highly correlated between them – with  
values close to 1 – whereas for TSMBC(RBC) no significant correlation appears. Remark that for TSMBC(dOTC) the correc-  
tions stay highly corrected with  $\mathbf{X}$ . It is an effect of the dOTC method, which preserves as much as possible the initial model  
simulations.

230 From this experiment we can conclude first that the choice of the starting row has only a very marginal influence on the cor-  
rection. Therefore, from now on, we use the integer part of  $r = \frac{s+1}{2}$  as starting row. Secondly, the TSMBC(RBC) method  
produces many different corrections, extremely weakly correlated between them. Even if, by construction, the TSMBC(RBC)  
is able to provide satisfactory inter-variable dependence structures and correlations, it is not able to provide proper temporal  
statistics, highlighting the importance of the choice of the underlying bias correction method.

235

### 3.2 Influence of the choice of the lag

In this section the starting row is fixed at  $r = \frac{s+1}{2}$ , and we vary the number of lags  $s \in \{3, 5, 10, 20, 100\}$ . This generate the  
corrections  $\mathbf{Z}_s$  for the two methods TSMBC(dOTC) and TSMBC(RBC).

As for the previous sub-section, the correlations between corrections, reference and biased dataset are computed, and repre-  
240 sented in Fig. 3c,d for the first variable, the second variable and  $p$ -values are given in Fig. S2. We can see for TSMBC(dOTC)  
that all corrections are highly correlated – and correlated with the biased dataset –, with a correlations decreasing with  $s$ . This  
shows that the corrections are similar, even when increasing the lag to a reasonable extent. As for the previous section, the  
method TSMBC(RBC) provides many different corrections, different from each other and thus difficult to link together.

Furthermore, we have added in Fig. 2a the density of the corrections with TSMBC(dOTC) and with TSMBC(RBC) in green  
245 and black respectively. The two methods adjust correctly the density and the dependence structure of the reference dataset.  
Therefore the differences seen in Fig. 3c,d come only from their capability to adjust the temporal structure.

The (cross) auto-correlations between the 2 dimensions for various lags are also given in Fig. 2b-e. On each panel, we can see  
that the results for the lag 0 – which corresponds to classic correlations – are closed to the references for all corrections, which  
is confirmed by the Fig. 2a. For non-zero lags, the (cross) auto-correlations of corrections with TSMBC(dOTC) are closed to  
250 the reference one, validating the method. For TSMBC(RBC), the (cross) auto-correlations become close to the reference if  $s$   
is large enough. This shows that the ability to correct the (cross) auto correlation when  $s$  is small comes from dOTC, and not  
only from the reconstruction part. When  $s$  increases, TSMBC(RBC) tends to adjust the temporal properties correctly but this  
is due to the reconstruction that replaces the biased dataset by the reference one for large  $s$ .

255 Generally, from the synthetic VAR dataset, we can see the ability of the TSMBC approach to correct the (cross) auto-  
correlations. The choice of the starting row has little influence on the final corrections, and we fix it now at  $\frac{s+1}{2}$ . We also



highlighted the importance of dOTC as underlying bias correction method (compared to the naive random approach), as the ability to properly correct the temporal structure does not come only from the reconstruction step of TSMBC.

#### 4 Results: DS/BC of temperature and precipitation

260 We now apply the TSMBC method with the underlying dOTC method to the bias correction/downscaling of the IPSL GCM simulations with respect to the RCM simulations taken as references. Following the strategy proposed by François et al. (2020), four kinds of corrections are applied and analysed:

- Each variable and grid point are corrected independently. This approach will be referred to as “L1V” (Local 1 Variable). The BC method employed here is dOTC in its univariate version (when  $s = 0$ );
- 265 – The dependence between temperature and precipitation (i.e., inter-variable dependence) is taken into account in the correction, but not the spatial dependence. This approach is noted “L2V” (Local 2 Variables), and employs the bivariate version of dOTC (when  $s = 0$ );
- The spatial dependence is corrected, but not the relations between temperature and precipitation. This approach is noted “S1V” (Spatial 1 Variable) and uses dOTC in a  $16$  (longitudes)  $\times$   $13$  (latitudes) =  $208$  dimensional configuration (when  
270  $s = 0$ );
- All dependencies (i.e., inter-variable and spatial) are corrected. This approach is noted “S2V” (Spatial 2 Variables), and dOTC has thus a  $2$  (variables)  $\times$   $16$  (longitudes)  $\times$   $13$  (latitudes) =  $416$ -dimensional configuration (when  $s = 0$ ).

Furthermore, for each of these approaches, we apply TSMBC to account for various lags, up to some maximum lags: 0-day (i.e., corresponding to dOTC, without any lag), 5-day and 10-day lags, noted dOTC, TSMBC-5 and TSMBC-10, respectively.  
275 Hence, we have finally 12 correction approaches, with dimensions varying from 1 (dOTC without any kind of dependence) to  $2 \times 16 \times 13 \times (10 + 1) = 4576$  (MSTBC-10 correcting spatial and inter-variable dependencies and temporal dependence up to lag 10-day). A summary of the dimension of each method is given in Tab. 1.

Recall that only the results for summer are given in the rest of this article (winter results are provided as supplementary materials), the calibration period is 1951-1980, and the validation/projection period is 1981-2010.  
280

##### 4.1 Bias reduction in marginal properties: mean, standard deviation, auto-correlation

We start by controlling the ability of the different methods to reduce the bias of the first two statistical moments: the mean (noted  $\mathbb{E}$ ) and the standard deviation (noted  $\sigma$ ). Noting  $\mathbf{Z}$  any correction, and  $\kappa$  the statistics of interest – such as the mean  $\mathbb{E}$ ,  
285 the standard deviation  $\sigma$ , or the lag- $s$  autocorrelation  $\rho_s$  –, we compute the following criterion  $BR_\kappa$  to characterize the bias



reduction:

$$BR_{\kappa} = 1 - \left| \frac{\kappa(\mathbf{Z}) - \kappa(\text{RCM})}{\kappa(\text{GCM}) - \kappa(\text{RCM})} \right|. \quad (6)$$

This criterion lives in the interval  $]-\infty, 1]$ . A value of 1 indicates a perfect correction. Boxplots of mean bias reduction ( $BR_{\bar{x}}$ ) and standard deviation bias reduction ( $BR_{\sigma}$ ) criteria from all grid points are presented in Fig. 4 for each variable and correction. For the calibration period (first column) we can see a bias reduction (both in means and standard deviations) between 0.95 and 1 for the temperature, and between 0.8 and 1 for precipitation. The bias reduction slightly decreases when the dimension increases, indicating a “curse of dimensionality” problem. For the projection period, the bias reduction in mean temperature stay reasonable, between 0.7 and 1. However, for the precipitation the results are more contrasted. For 75% of the grid points, the reduction of the mean precipitation bias lies between 0.2 and 0.95 and that of the standard deviation bias between 0.6 and 1. Interestingly, the use of a TSMBC approach relying on dOTC implies bias reductions in means and standard deviations equivalent to those provided by the dOTC method, i.e., without “time shifted” consideration. In other words, TSMBC does not degrade the basic marginal properties of the corrections from dOTC. Note that the ability of the methods to reduce the biases can be strongly affected by the evolution of the GCM variables between the calibration and validation period. Indeed, this evolution is different from that of the RCM (i.e., here, reference) variables. As dOTC preserves the evolution of the GCM to be corrected, the resulting corrections for the projection period have necessarily properties different from the reference over the same period.

The same boxplot are now represented in Fig 5 for the lag-2 auto-correlation bias reduction  $BR_{\rho_2}$ . Over the calibration period, the use of TSMBC clearly improves the (cross-) auto-correlation compared to dOTC. In the projection period, the TSMBC correction is better than the dOTC correction for the pr/pr couple. No clear improvement appears for the couples pr/tas and tas/pr, and for the couple tas/tas a degradation appears when dimension increases, related to the problems already described for the mean and standard deviation.

Figure 6 presents the maps of auto-correlations of precipitation at lag 1 (first two lines) and at lag 4 (last two lines). The calibration period corresponds to the first and third lines, and the projection period to the second and last lines. We focus here only on the L2V approach, but the results are equivalent for the others (not shown, except for temperature with L2V given in Fig. S4). Regarding the calibration period, it is clear that dOTC does not reproduce the RCM auto-correlation maps. In addition, the obtained auto-correlation maps are quite different from those of the GCM, justifying the need for TSMBC. Based on TSMBC-5 and TSMBC-10, the corrections are closer to the reference RCM. Using 10 lags instead of 5 does not show a clear improvement, and can sometimes even degrade the corrections. For the projection period, the situation is quite different and requires to compare the GCM, RCM and corrections, as well as their evolution between the calibration and projection periods. For the GCM, the lag-1 auto-correlation decreases between the two periods for the northern part, but increases in the



southern part. As dOTC mostly reproduces the evolution of the model (see Robin et al., 2019), the same feature appears in the  
320 corrections over the projection period. TSMBC-5 or TSMBC-10 using dOTC as underlying MBC method, the same conclusion  
holds. This result is also true for the lag 4, with auto-correlations more noisy for TSMBC-5 and TSMBC-10 due to a higher  
number of dimensions to consider.

Globally, TSMBC is able to reduce biases in means and standard deviations as well as dOTC but clearly improves the cor-  
325 rections of the auto-correlations. We now propose to further study the dependence structure of the corrections brought by  
TSMBC.

#### 4.2 Bias reduction in dependencies: the $\mathcal{W}$ -cross-auto-correlogram metric

The present sub-section targets the evaluation of the TSMBC corrections in terms of spatial structure of auto-correlations  
between variables and grid points. This requires a new tool: the  $\mathcal{W}$ -cross-auto-correlogram metric, based on the classic correl-  
330 ogram.

In order to evaluate spatial dependencies present in a univariate sample, correlograms (i.e., correlations expressed as function  
of the distance) are classically used (see, e.g. Vrac, 2018; François et al., 2020). To estimate it, the correlation between each  
pair of locations (or grid cells in the present study) is computed. Then the set of distance/correlation pairs (DCP) is divided into  
classes (e.g., 0-10km, 10-20km, etc) and the conditional mean correlation is calculated for each class. However, following an  
335 equivalent procedure, the calculation of correlation in a univariate context can be replaced by the auto-correlation for any lags,  
or, in a multivariate setting, by cross-auto-correlations between variables (e.g., correlation between temperature at time  $t$  for  
one location and precipitation at time  $t + \text{lag}$  for another location). This gives an auto-correlogram or a cross-auto-correlogram,  
respectively.

340 Figure 7 presents the scatter-plots of the DCP values, where correlations are cross-auto-correlations between temperature and  
precipitations for all pairs of grid points with lag 1 for the GCM data (panel a), the RCM (panel b), the corrections with dOTC  
(S2V, panel c) and with TSMBC-5 (S2V, panel d). In each panel, the black line is the estimated mean cross-auto-correlogram.  
It is clear that the cross-auto-correlograms – as mean conditional correlations given distances – are not quite representative  
of the DCP scatter-plot structures. Indeed, the DCP structures show large spreads that cannot be visible on simple lines.  
345 Moreover, those structures are different from one dataset to another, for example between the GCM data to be corrected and  
from the reference RCM data, while their associated cross-auto-correlogram (i.e., mean black lines) can appear relatively close  
to each other. However, the DCP scatter-plots of the dOTC (c) and TSMBC-5 (d) corrections exhibit a large improvement  
compared to the uncorrected GCM, with TSMBC-5 that seems better than dOTC, with more realistic spread and shape of the  
DCP set. To quantify this improvement, for the GCM, the RCM and all the corrections, all DCP sets are first computed for lags  
350 between 0 and 9, for auto-correlations (temperature/temperature and precipitation/precipitation) and for cross-auto-correlations  
(temperature/precipitation and precipitation/temperature). Then, the Wasserstein distance  $\mathcal{W}$  (see App. A) is calculated and  
used as a metric to measure the difference between the DCP sets from the RCM (i.e., here, the reference) and the DCP sets



of the GCM or the corrections. The Wasserstein distance (e.g., Santambrogio, 2015; Robin et al., 2019) is a distance between two multivariate distributions and can therefore be considered as an alternative to the Energy distance (Rizzo and Székely, 2016; François et al., 2020). Noting respectively  $DCP_{RCM}$ ,  $DCP_{GCM}$  and  $DCP_{\mathbf{Z}}$  these DCP sets, we propose the following indicator,  $BR_{\mathcal{W}}$ , based on the Wasserstein distance, to measure the bias reduction in dependence with respect to the raw GCM:

$$BR_{\mathcal{W}} = 1 - \left| \frac{\mathcal{W}(DCP_{\mathbf{Z}}, DCP_{RCM})}{\mathcal{W}(DCP_{GCM}, DCP_{RCM})} \right| \quad (7)$$

where  $\mathcal{W}(DCP_{\mathbf{Z}}, DCP_{RCM})$  is the Wasserstein distance between the DCP set from the corrections  $\mathbf{Z}$  and that from the reference RCM data; and  $\mathcal{W}(DCP_{GCM}, DCP_{RCM})$  the equivalent between the GCM and RCM DCP sets. A value of  $BR_{\mathcal{W}}$  close to 1 indicates that the chosen dependencies (correlation, auto-correlation or cross-auto-correlation) of the correction is close to the dependence structure of the reference RCM. We call the set of this indicator a  $\mathcal{W}$ -cross-auto correlogram.

As the Wasserstein metric is sensitive to the scale of the multivariate data (here, the DCP sets) it is applied to, two normalizations of the DCP sets are proposed before the computation of the  $BR_{\mathcal{W}}$  values. First, we normalize the correlation values of the DCP sets between  $-1$  and  $1$ , independently for each method and lag. This allows us to compare only the pattern of the DPC sets, by removing mean and scale biases. The results are shown in Fig. 8. Alternatively, a second type of normalization is performed, with a normalization common to all methods for a given lag but different for different lags. Hence, this normalization conditional to the lag allows us to compare the different bias correction methods and include both the intensities of the correlations and the DCP patterns while first normalization only accounts for the pattern of the DCP sets. The results are presented in Fig. 9, where only the  $BR_{\mathcal{W}}$  values coming from the same “column” (i.e., lag) can be compared.

Starting with the first normalization (for each method/lag separately), allowing to compare only the pattern of the DPC sets, in Fig. 8, we can first see that  $BR_{\mathcal{W}}$  matrices for calibration and projection periods are relatively similar, indicating some robustness of the TSMBC method over projections. Generally speaking, for a specific configuration of the method (i.e., L1V, L2V, S1V or S2V), TSMBC (5 or 10) is better than dOTC that does not account for temporal properties. This is true for almost all lags  $>0$  and any  $BR_{\mathcal{W}}$  matrix (tas/tas, tas/pr, pr/tas, pr/pr). However, although dOTC provides similar results for the L1V and L2V configurations (potentially of poor quality for various lags), the use of S1V and S2V approaches within dOTC – i.e., without specifically accounting for temporal dependence – strongly improves the  $BR_{\mathcal{W}}$  results. This indicates that imposing to account for spatial properties within dOTC can improve the correction of the auto-correlation and cross-auto-correlation patterns. Moreover, regarding the tas/tas and pr/pr  $BR_{\mathcal{W}}$  matrices, the spatial versions (i.e., S1V and S2V) of the method appear largely better than the local (i.e., L1V and L2V) ones, especially for tas/tas for most of the lags. This was somehow expected since the S1V and S2V configurations also correct spatial dependencies. Regarding the  $BR_{\mathcal{W}}$  matrices for tas/pr and pr/tas, results are better for L2V and S2V configurations than with L1V and S1V ones: accounting for correlations and cross-auto-correlations between temperature and precipitation allows improving the results over configurations based on temperature and precipitation considered separately. Furthermore, it is not clear that increasing the number of lags in TSMBC significantly



improves the results. Here, TSMBC-10 and TSMBC-5 provide quite similar  $BR_{\mathcal{W}}$  values whatever the couple of variables of interest, with some non-systematic variations along the lags. However, globally, the DCP patterns are clearly improved by the TSMBC corrections, as shown by the positive  $BR_{\mathcal{W}}$  values in Fig. 8.

390 Continuing with the  $BR_{\mathcal{W}}$  results obtained based on the second type of normalization in Fig. 9, the results are more contrasted, as negative values are now visible. This traduces the fact that, although DCP patterns are globally improved by the various configurations of dOTC and TSMBC (as seen previously in Fig. 8), biases are present in the intensities of correlations and, when accounted for, degrades the  $BR_{\mathcal{W}}$  values. However, as for the first normalization, the spatial configurations (S1V or S2V) seem to reduce biases more than the local (L1V or L2V) configurations; the L2V and S2V approaches are in general better than the  
395 L1V and S1V ones for the cross-correlations matrices (i.e., involving both tas and pr): and the S1V versions appear better than the S2V ones for matrices with singles variables (i.e., tas/tas and pr/pr). This latter remark is due to the fact that with the S2V approach, the complexity of the methods is obviously higher than with S1V: this is done at the expense of the quality of each variable separately. Hence, in such a case where only one variable is of interest in a spatial context, the S1V methods have to be favoured. Moreover, generally, the bias reduction in dependence is stronger (i.e., better) for the cross-auto-correlations  
400 between precipitation and temperature (matrices “pr/tas”) and between precipitation and precipitation (matrices “pr/pr”) than for the other couples of variables (“tas/tas” and “tas/pr”). This comes from the fact that the initial biases (i.e., Wasserstein distances) of the raw GCM data for pr/tas and pr/pr are larger than those from tas/tas and tas/pr. This is visible in Fig. S7 given as supplementary material, showing the values of the Wasserstein distances (based on the second normalization of the DCP sets) used to compute  $BR_{\mathcal{W}}$ . In addition, some negative  $BR_{\mathcal{W}}$  values in Fig. 9 are related to very small differences between  
405 Wasserstein distances very close to zero. Indeed, if the model simulations to be corrected have a DCP set already close to the reference, its Wasserstein distance will be near zero. If a corrected dataset has also its distance to the reference close to zero, the resulting ratio can be quite different from 1 (and thus induce a strongly negative  $BR_{\mathcal{W}}$  value) while the two  $\mathcal{W}$  values are only slightly different. For example among others, this is the case for TSMBC-5 (S1V) at lag 1 for the “tas/tas” matrix under projection context in Fig. S7, where the  $\mathcal{W}$  value is small but slightly higher than that of the raw IPSL simulations,  
410 implying a negative (red)  $BR_{\mathcal{W}}$  value in Fig. 9. Finally, when comparing TSMBC-5 and TSMBC-10, it appears that, whatever the configuration, increasing the dimension (i.e., 10 lags to be accounted for in the MBC, instead of 5) degrades the gain in correlations. The increase in the complexity (i.e., the number of dimensions) of the method is made at the expense of the quality of the results. This was not visible on previous Fig. 8 that only accounted for the shape of the DCPs. Thus, it indicates that this degradation is mostly due to biases in the marginals that are not fully removed by TSMBC in a high-dimensional setting.

415

## 5 Conclusions and discussion

The goal of bias correction (BC) is to transform biased climate simulations in order to make their statistical properties more similar to those from reference data. Over the last decades, many univariate BC methods were developed and applied, working



on one climate variable at a time and one location at a time. Over the last few years, various multivariate bias correction (MBC)  
420 methods were also designed to correct not only some marginal properties of the simulations (e.g., means, variances, distribu-  
tions) but also their dependencies (e.g., correlations), either in a multivariate context, inter-site context, or both. Some methods  
were even specifically developed to adjust the temporal properties (e.g., Mehrotra and Sharma, 2016) of the simulations. How-  
ever, the latter usually consider time-related properties and temporal (auto- or cross-) correlations specifically, i.e., differently  
425 like other variables – i.e., by adding to the (multivariate) data to be corrected some lagged time series of itself –, an MBC  
method is able to correct both the multivariate (inter-variable and inter-site) properties and the temporal properties. To test  
this, the “dynamical Optimal Transport Correction” (dOTC) method has been applied first to a synthetic (i.e., statistically  
generated) dataset and, second, to a dataset of daily temperature and precipitation over South of France, simulated from the  
IPSL global climate model. For evaluation, the reference dataset was extracted from higher-resolution regional climate simu-  
430 lations over the same region. dOTC was then been applied to correct the IPSL dataset where various lagged versions of those  
simulations have been added. This approach – performing an MBC on lagged time series in addition to the initial ones – has  
been called “Time Shifted Multivariate Bias Correction” (TSMBC). Furthermore, it has been tested based on 4 configurations:  
only one variable is corrected at a time, either for a given location (L1V) or for all locations at the same time (S1V); the 2  
variables are jointly corrected, for a single location (L2V) or for all locations (S2V).

435

From the synthetic data experiment, a comparison with a “reasonably naive” multivariate bias correction method, RBC, based  
on random sampling, has been proposed. The results showed that:

- TSMBC(dOTC) provides a clear improvement compared to TSMBC(RBC),
- the choice of the starting row only has a marginal influence on the corrections,
- 440 – for a relatively low number  $s$  of lags to account for in TSMBC – say  $s \leq 10$ -day –, the TSMBC- $s$  results are roughly  
equivalent, whatever  $s$ .

Those first conclusions indicates some robustness of the proposed TSMBC methodology that, despite some choices to make  
by the user – starting row, number of lags to include –, provides stable corrections.

445 In order to evaluate the results in a fully multivariate manner (i.e., intervariable, inter-site and temporal aspects), a new sta-  
tistical criterion has been proposed. It is based on the Wassertein distance between the set of distance/correlation pairs (DCP)  
from reference and that of a dataset (from corrections or simulations). This distance can be computed on lagged data, mul-  
tiple variables and at different locations, hence providing assessments of cross-auto-correlations, generalizing the traditional  
correlogram tool.

450 The results obtained by applying TSMBC to climate simulations provided the following conclusions:



- In terms of means and standard deviations, for both temperature and precipitation, the inclusion of lagged data does not strongly modify the results of the dOTC correction method. Although some evidences of degradation are present when the number of lags increase (e.g., for  $BR_\sigma$  in temperature), the bias reduction values with respect to the raw simulations are mostly positive, indicating an improvement over the raw GCM.
- 455 – This is also mostly the case for autocorrelation bias reductions  $BR_\rho$  but, this time, the increase in number of lags in TSMBC globally improves the results, even though TSMBC-10 does not clearly improve TSMBC-5.
- Moreover, the main spatio-temporal patterns of the TSMBC results are globally improving those from the raw GCM.
- However, biases in the intensities of the (intervariable, inter-site or temporal) correlations might remain. This is typically related to very small differences between two Wasserstein distances very close to zero: if the raw simulations already  
460 have a DCP set close to the reference, its Wasserstein distance will be near zero. Therefore, the relative reduction of bias  $BR$  can be strongly negative, even though the absolute difference is potentially very small.
- Finally, if the TSMBC methodology seems to reasonably adjust temporal (cross-auto) correlations, while still performing well on multivariate properties, when the number of lags increases (e.g., from 5 to 10 days), the gain in the quality of the corrections is not obvious and the latter can even be degraded. It is thus required to limit the temporal constrains to a few  
465 time steps, depending on the variable of interest. This would avoid to apply the MBC method in a too high dimensional context and then allow robust results.

Globally, the results of the different tests indicate that the proposed approach of “time shifted multivariate bias correction” – i.e., including lagged versions of the simulated and reference datasets in a multivariate correction procedure – can indeed be relevant to adjust temporal properties, in addition to more usual marginal and multivariate components.

470

Despite its promising results, the TSMBC approach can be further investigated and improved. For example, in the present study, only the dOTC multivariate method was used as correction technique. Other MBC methods exist. Hence, it would be interesting to test how those alternative MBCs – such as “ $R^2D^2$ ” (Vrac, 2018; Vrac and Thao, 2020), “MBCn” (Cannon, 2018), or “MRrec” (Bárdossy and Pegram, 2012), among others – would behave in this TSMBC framework.

475 Moreover, some adjustment methods were designed to account specifically for the correction of temporal properties (e.g., Johnson and Sharma, 2012; Mehrotra and Sharma, 2015, 2016). A comparison of TSMBC to such methods would then be of interest to understand if specific modelling is needed for adjusting temporal properties or if considering lagged data into MBCs (i.e., no specific modelling as in TSMBC) provides equivalent results.

Finally, the Wasserstein cross-auto-correlation based metric introduced in this study could be used more generally to compare  
480 various datasets and/or assess their diverse properties with respect to a reference. It can then be useful to make evaluations of climate simulations (adjusted or not) in a more holistic way.





## Appendix A: Optimal transport

Optimal transport and Wasserstein distance (see, e.g. Villani, 2008; Santambrogio, 2015) are used in this paper by the dOTC  
485 method to perform a bias correction, and by  $\mathcal{W}$ -cross-auto-correlation-based metric. Starting with two probability distributions  
 $\mathbb{P}_1$  and  $\mathbb{P}_2$  on  $\mathbb{R}^d$ , we consider the following set  $\Gamma$  of probability distributions on  $\mathbb{R}^d \times \mathbb{R}^d$ :

$$\Gamma = \left\{ \gamma : \mathbb{P}_1(A) = \int_{A \times \mathbb{R}^d} d\gamma, \mathbb{P}_2(B) = \int_{\mathbb{R}^d \times B} d\gamma \right\}.$$

Thus,  $\Gamma$  – called the set of transport plan – is the set of probability distributions  $\gamma$  such that the first projection is  $\mathbb{P}_1$ , and  
490 the second is  $\mathbb{P}_2$ . The underlying idea in the context of bias correction with dOTC is that  $\gamma(\mathbf{x}, \mathbf{y})$  is the probability that  $\mathbf{x}$  is  
transformed into  $\mathbf{y}$ , justifying the name of “transport plan”. Among all  $\gamma \in \Gamma$ , we consider  $\tilde{\gamma}$  minimizing the following cost  
function:

$$\tilde{\gamma} = \arg \min_{\gamma \in \Gamma} \int_{\mathbb{R}^d \times \mathbb{R}^d} \|\mathbf{x} - \mathbf{y}\|^2 d\gamma(\mathbf{x}, \mathbf{y}) \quad (\text{A1})$$

This cost function gives the “energy” needed to transform  $\mathbb{P}_1$  into  $\mathbb{P}_2$ , following the plan  $\tilde{\gamma}$ . So, the minimum is the plan with  
495 minimal energy, and we note it  $\tilde{\gamma}$ . This particular plan is used by the dOTC method to perform the bias correction, and define  
also a metric called the Wasserstein distance:

$$\mathcal{W}(\mathbb{P}_1, \mathbb{P}_2)^2 = \int_{\mathbb{R}^d \times \mathbb{R}^d} \|\mathbf{x} - \mathbf{y}\|^2 d\tilde{\gamma}(\mathbf{x}, \mathbf{y}) = \inf_{\gamma \in \Gamma} \int_{\mathbb{R}^d \times \mathbb{R}^d} \|\mathbf{x} - \mathbf{y}\|^2 d\gamma(\mathbf{x}, \mathbf{y}) \quad (\text{A2})$$

*Author contributions.* MV and YR had the idea of the method together. They designed the study and the experiments together. YR made the  
computations and plots. YR and MV jointly analysed the results. MV wrote the major part of the article with inputs from YR.

500 *Competing interests.* The authors declare no competing interest.

*Acknowledgements.* We acknowledge the World Climate Research Program’s Working Group on Coupled Modelling, which is responsible  
for CMIP, and we thank the climate modeling groups for producing and making available their model output. For CMIP the U.S. Department  
of Energy’s Program for Climate Model Diagnosis and Intercomparison provides coordinating support and led development of software



infrastructure in partnership with the Global Organization for Earth System Science Portals.

505

We acknowledge the World Climate Research Programme’s Working Group on Regional Climate, and the Working Group on Coupled Modelling, former coordinating body of CORDEX and responsible panel for CMIP5. We also thank the climate modelling groups for producing and making available their model output. We also acknowledge the Earth System Grid Federation infrastructure an international effort led by the U.S. Department of Energy’s Program for Climate Model Diagnosis and Intercomparison, the European Network for Earth System

510 Modelling and other partners in the Global Organisation for Earth System Science Portals (GO-ESSP).

MV has been supported by the CoCliServ project. MV and YR have been supported by the EUPHEME project. Both CoCliServ and EUPHEME are part of ERA4CS, an ERA-NET initiated by JPI Climate and cofunded by the European Union (grant no. 690462). MV has also been supported by project C3S 428J (“HR-CDFt”). YR has also been supported by project C3S 62 (“Prototype Extreme Events and

515 Attribution Service”).



## References

- Bárdossy, A. and Pegram, G.: Multiscale spatial recorrelation of RCM precipitation to produce unbiased climate change scenarios over large areas and small, *Water Resour. Res.*, 48, 9502–, <https://doi.org/10.1029/2011WR011524>, 2012.
- 520 Bevacqua, E., Maraun, D., Vousdoukas, M. I., Voukouvalas, E., Vrac, M., Mentaschi, L., and Widmann, M.: Higher probability of compound flooding from precipitation and storm surge in Europe under anthropogenic climate change, *Sci. Adv.*, 5, <https://doi.org/10.1126/sciadv.aaw5531>, <https://advances.sciencemag.org/content/5/9/eaaw5531>, 2019.
- Bhuiyan, M. A. E., Nikolopoulos, E. I., Anagnostou, E. N., Quintana-Seguí, P., and Barella-Ortiz, A.: A nonparametric statistical technique for combining global precipitation datasets: development and hydrological evaluation over the Iberian Peninsula, *Hydrol. Earth Syst. Sci.*, 22, 1371–1389, <https://doi.org/10.5194/hess-22-1371-2018>, <https://hess.copernicus.org/articles/22/1371/2018/>, 2018.
- 525 Boé, J., Terray, L., Habets, F., and Martin, E.: Statistical and dynamical downscaling of the Seine basin climate for hydro-meteorological studies, *Int. J. Climatol.*, 27, 1643–1655, <https://doi.org/10.1002/joc.1602>, 2007.
- Cannon, A. J.: Multivariate quantile mapping bias correction: an N-dimensional probability density function transform for climate model simulations of multiple variables, *Clim. Dyn.*, 50, 31–49, <https://doi.org/10.1007/s00382-017-3580-6>, <https://doi.org/10.1007/s00382-017-3580-6>, 2018.
- 530 Cannon, A. J., Sobie, S. R., and Murdock, T. Q.: Bias Correction of GCM Precipitation by Quantile Mapping: How Well Do Methods Preserve Changes in Quantiles and Extremes?, *J. Clim.*, 28, 6938–6959, <https://doi.org/10.1175/JCLI-D-14-00754.1>, 2015.
- Chen, J., Brissette, F. P., and Leconte, R.: Uncertainty of downscaling method in quantifying the impact of climate change on hydrology, *J. Hydrol.*, 401, 190–202, <https://doi.org/https://doi.org/10.1016/j.jhydrol.2011.02.020>, <https://www.sciencedirect.com/science/article/pii/S0022169411001351>, 2011.
- 535 Chen, J., Brissette, F. P., Chaumont, D., and Braun, M.: Finding appropriate bias correction methods in downscaling precipitation for hydrologic impact studies over North America, *Water Resour. Res.*, 49, 4187–4205, <https://doi.org/https://doi.org/10.1002/wrcr.20331>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/wrcr.20331>, 2013.
- Christensen, J. H., Boberg, F., Christensen, O. B., and Lucas-Picher, P.: On the need for bias correction of regional climate change projections of temperature and precipitation, *Geophys. Res. Lett.*, 35, <https://doi.org/https://doi.org/10.1029/2008GL035694>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2008GL035694>, 2008.
- 540 Dekens, L., Parey, S., Grandjacques, M., and Dacunha-Castelle, D.: Multivariate distribution correction of climate model outputs: A generalization of quantile mapping approaches: Multivariate distribution correction of climate model outputs, *Environmetrics*, 28, e2454, <https://doi.org/https://doi.org/10.1002/env.2454>, <https://onlinelibrary.wiley.com/doi/abs/10.1002/env.2454>, 2017.
- Delrieu, G., Nicol, J., Yates, E., Kirstetter, P.-E., Creutin, J.-D., Anquetin, S., Obled, C., Saulnier, G.-M., Ducrocq, V., Gaume, E., Payrastre, O., Andrieu, H., Ayrat, P.-A., Bouvier, C., Neppel, L., Livet, M., Lang, M., du Châtelet, J. P., Walpersdorf, A., and Wobrock, W.: The Catastrophic Flash-Flood Event of 8–9 September 2002 in the Gard Region, France: A First Case Study for the Cévennes–Vivarais Mediterranean Hydrometeorological Observatory, *Journal of Hydrometeorology*, 6, 34–52, <https://doi.org/10.1175/JHM-400.1>, <https://doi.org/10.1175/JHM-400.1>, 2005.
- 545 Déqué, M.: Frequency of precipitation and temperature extremes over France in an anthropogenic scenario: Model results and statistical correction according to observed values, *Global Planet. Change*, 57, 16 – 26, <https://doi.org/https://doi.org/10.1016/j.gloplacha.2006.11.030>, <https://www.sciencedirect.com/science/article/pii/S0921818106002748>, 2007.




- Dufresne, J.-L., Foujols, M.-A., Denvil, S., Caubel, A., Marti, O., Aumont, O., Balkanski, Y., Bekki, S., Bellenger, H., Benschila, R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Cheruy, F., Codron, F., Cozic, A., Cugnet, D., de Noblet, N., Duvel, J.-P., Ethé, C., Fairhead, L., Fichefet, T., Flavoni, S., Friedlingstein, P., Grandpeix, J.-Y., Guez, L., Guilyardi, E., Hauglustaine, D., Hourdin, F., Idelkadi, A., Ghattas, J., Joussaume, S., Kageyama, M., Krinner, G., Labetoulle, S., Lahellec, A., Lefebvre, M.-P., Lefevre, F., Levy, C., Li, Z. X., Lloyd, J., Lott, F., Madec, G., Mancip, M., Marchand, M., Masson, S., Meurdesoif, Y., Mignot, J., Musat, I., Parouty, S., Polcher, J., Rio, C., Schulz, M., Swingedouw, D., Szopa, S., Talandier, C., Terray, P., Viovy, N., and Vuichard, N.: Climate change projections using the IPSL-CM5 Earth System Model: from CMIP3 to CMIP5, *Clim. Dyn.*, 40, 2123–2165, <https://doi.org/https://doi.org/10.1007/s00382-012-1636-1>, 2013.
- 555
- Eden, J., Widmann, M., Grawe, D., and Rast, S.: Skill, Correction, and Downscaling of GCM-Simulated Precipitation, *J. Climate*, 25, 3970–3984, <https://doi.org/https://doi.org/10.1175/JCLI-D-11-00254.1>, 2012.
- 560
- Flato, G., Marotzke, J., Abiodun, B., Braconnot, P., Chou, S., Collins, W., Cox, P., Driouech, F., Emori, S., Eyring, V., Forest, C., Gleckler, P., Guilyardi, E., Jakob, C., Kattsov, V., Reason, C., and Rummukainen, M.: Evaluation of Climate Models, book section 9, pp. 741–866, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/CBO9781107415324.020>, [www.climatechange2013.org](http://www.climatechange2013.org), 2013.
- 565
- François, B., Vrac, M., Cannon, A., Robin, Y., and Allard, D.: Multivariate bias corrections of climate simulations: Which benefits for which losses?, *Earth Syst. Dyn.*, 11, 537–562, <https://doi.org/10.5194/esd-11-537-2020>, <https://esd.copernicus.org/articles/11/537/2020/>, 2020.
- Frieler, K., Lange, S., Piontek, F., Reyer, C. P. O., Schewe, J., Warszawski, L., Zhao, F., Chini, L., Denvil, S., Emanuel, K., Geiger, T., Halladay, K., Hurtt, G., Mengel, M., Murakami, D., Ostberg, S., Popp, A., Riva, R., Stevanovic, M., Suzuki, T., Volkholz, J., Burke, E., Ciais, P., Ebi, K., Eddy, T. D., Elliott, J., Galbraith, E., Gosling, S. N., Hattermann, F., Hickler, T., Hinkel, J., Hof, C., Huber, V., Jägermeyr, J., Krysanova, V., Marcé, R., Müller Schmied, H., Mouratiadou, I., Pierson, D., Tittensor, D. P., Vautard, R., van Vliet, M., Biber, M. F., Betts, R. A., Bodirsky, B. L., Deryng, D., Frolking, S., Jones, C. D., Lotze, H. K., Lotze-Campen, H., Sahajpal, R., Thonicke, K., Tian, H., and Yamagata, Y.: Assessing the impacts of 1.5 °C global warming – simulation protocol of the Inter-Sectoral Impact Model Intercomparison Project (ISIMIP2b), *Geosci. Model Dev.*, 10, 4321–4345, <https://doi.org/10.5194/gmd-10-4321-2017>, <https://gmd.copernicus.org/articles/10/4321/2017/>, 2017.
- 570
- 575
- Gudmundsson, L., Bremnes, J. B., Haugen, J. E., and Engen-Skaugen, T.: Technical Note: Downscaling RCM precipitation to the station scale using statistical transformations; a comparison of methods, *Hydrol. Earth Syst. Sci.*, 16, 3383–3390, <https://doi.org/10.5194/hess-16-3383-2012>, <https://hess.copernicus.org/articles/16/3383/2012/>, 2012.
- Haddad, Z. S. and Rosenfeld, D.: Optimality of empirical Z-R relations, *Q. J. R. Meteorol. Soc.*, 123, 1283–1293, <https://doi.org/https://doi.org/10.1002/qj.49712354107>, <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.49712354107>, 1997.
- 580
- Hempel, S., Frieler, K., Warszawski, L., Schewe, J., and Piontek, F.: A trend-preserving bias correction - the ISI-MIP approach, *Earth Syst. Dyn.*, 4, 219–236, <https://doi.org/10.5194/esd-4-219-2013>, <https://esd.copernicus.org/articles/4/219/2013/>, 2013.
- Jeong, D., St-Hilaire, A., Ouarda, T., and Gachon, P.: Multisite statistical downscaling model for daily precipitation combined by multivariate multiple linear regression and stochastic weather generator., *Clim. Change*, 114, 567–591, <https://doi.org/https://doi.org/10.1007/s10584-012-0451-3>, 2012.
- 585
- Johnson, F. and Sharma, A.: A nesting model for bias correction of variability at multiple time scales in general circulation model precipitation simulations, *Water Resour. Res.*, 48, <https://doi.org/https://doi.org/10.1029/2011WR010464>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2011WR010464>, 2012.




- Kallache, M., Vrac, M., Naveau, P., and Michelangeli, P.-A.: Non-stationary probabilistic downscaling of extreme precipitation, *J. Geophys. Res. D*, 116, <https://doi.org/https://doi.org/10.1029/2010JD014892>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2010JD014892>, 2011.
- Kirtman, B., Power, S., Adedoyin, J., Boer, G., Bojariu, R., Camilloni, I., Doblas-Reyes, F., Fiore, A., Kimoto, M., Meehl, G., Prather, M., Sarr, A., Schär, C., Sutton, R., van Oldenborgh, G., Vecchi, G., and Wang, H.: Near-term Climate Change: Projections and Predictability, book section 11, pp. 953–1028, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/CBO9781107415324.023>, [www.climatechange2013.org](http://www.climatechange2013.org), 2013.
- KNMI: cordex EUR-11 KNMI RACMO22E, <http://cera-www.dkrz.de/WDCC/ui/Compact.jsp?acronym=CXEU11KNRA>, 2017.
- Leonard, M., Westra, S., Phatak, A., Lambert, M., Hurk, B., McInnes, K., Risbey, J., Schuster, S., Jakob, D., and Stafford Smith, M.: A compound event framework for understanding extreme impacts, *Wiley Interdiscip. Rev. Clim. Change*, 5, 113–128, 2014.
- Maraun, D. and Widmann, M.: Statistical Downscaling and Bias Correction for Climate Research, Cambridge University Press, <https://doi.org/10.1017/9781107588783>, 2018.
- Marti, O., Braconnot, P., Dufresne, J., Bellier, J., R., B., Bony, S., Brockmann, P., Cadule, P., Caubel, A., Codron, F., de Noblet, N., Denvil, S., Fairhead, L., Fichefet, T., Foujols, M., Friedlingstein, P., Goosse, H., Grandpeix, J., Guilyardi, E., Hourdin, F., Idelkadi, A., Kageyama, M., Krinner, G., Lévy, C., Madec, G., Mignot, J., Musat, I., Swingedouw, D., and Talandier, C.: Key features of the IPSL ocean atmosphere model and its sensitivity to atmospheric resolution, *Clim. Dyn.*, 34, 1–26, <https://doi.org/https://doi.org/10.1007/s00382-009-0640-6>, 2010.
- Mehrotra, R. and Sharma, A.: Correcting for systematic biases in multiple raw GCM variables across a range of timescales, *J. Hydrol.*, 520, 214–223, <https://doi.org/https://doi.org/10.1016/j.jhydrol.2014.11.037>, <https://www.sciencedirect.com/science/article/pii/S0022169414009573>, 2015.
- Mehrotra, R. and Sharma, A.: A Multivariate Quantile-Matching Bias Correction Approach with Auto- and Cross-Dependence across Multiple Time Scales: Implications for Downscaling, *J. Clim.*, 29, 3519–3539, <https://doi.org/https://doi.org/10.1175/JCLI-D-15-0356.1>, <https://journals.ametsoc.org/view/journals/clim/29/10/jcli-d-15-0356.1.xml>, 2016.
- Piani, C. and Haerter, J.: Two dimensional bias correction of temperature and precipitation copulas in climate models, *Geophys. Res. Lett.*, 39, <https://doi.org/https://doi.org/10.1029/2012GL053839>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2012GL053839>, 2012.
- Raimonet, M., Oudin, L., Thieu, V., Silvestre, M., Vautard, R., Rabouille, C., and Le Moigne, P.: Evaluation of Gridded Meteorological Datasets for Hydrological Modeling, *J. Hydrometeorol.*, 18, 3027–3041, <https://doi.org/10.1175/JHM-D-17-0018.1>, [https://journals.ametsoc.org/view/journals/hydr/18/11/jhm-d-17-0018\\_1.xml](https://journals.ametsoc.org/view/journals/hydr/18/11/jhm-d-17-0018_1.xml), 2017.
- Rizzo, M. L. and Székely, G. J.: Energy distance, *Wiley Interdiscip. Rev. Comput. Stat.*, 8, 27–38, <https://doi.org/https://doi.org/10.1002/wics.1375>, <https://onlinelibrary.wiley.com/doi/abs/10.1002/wics.1375>, 2016.
- Robin, Y., Vrac, M., Naveau, P., and Yiou, P.: Multivariate stochastic bias corrections with optimal transport, *Hydrol. Earth Syst. Sci.*, 23, 773–786, <https://doi.org/10.5194/hess-23-773-2019>, <https://hess.copernicus.org/articles/23/773/2019/>, 2019.
- Santambrogio, F.: Optimal Transport for Applied Mathematicians: Calculus of Variations, PDEs, and Modeling, *Progress in Nonlinear Differential Equations and Their Applications*, Birkhäuser, Cham, 1 edn., <https://doi.org/https://doi.org/10.1007/978-3-319-20828-2>, 2015.
- Schmidli, J., Frei, C., and Vidale, P.: Downscaling from GCM precipitation: a benchmark for dynamical and statistical downscaling methods, *Int. J. Climatol.*, 26, 679–689, <https://doi.org/https://doi.org/10.1002/joc.1287>, <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/joc.1287>, 2006.



- Shukla, P., Skea, J., Calvo Buendia, E., Masson-Delmotte, V., Pörtner, H.-O., Roberts, D. C., Zhai, P., Slade, R., Connors, S., van Diemen, R., Ferrat, M., Haughey, E., Luz, S., Neogi, S., Pathak, M., Petzold, J., Portugal Pereira, J., Vyas, P., Huntley, E., Kissick, K., Belkacemi, M., and Malley, J.: Climate Change and Land: an IPCC special report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems, 2019.
- 630 Teutschbein, C. and Seibert, J.: Bias correction of regional climate model simulations for hydrological climate-change impact studies: Review and evaluation of different methods, *J. Hydrol.*, 456–457, 12–29, <https://doi.org/https://doi.org/10.1016/j.jhydrol.2012.05.052>, <https://www.sciencedirect.com/science/article/pii/S0022169412004556>, 2012.
- Thrasher, B., Maurer, E. P., McKellar, C., and Duffy, P. B.: Technical Note: Bias correcting climate model simulated daily temperature extremes with quantile mapping, *Hydrol. Earth Syst. Sci.*, 16, 3309–3314, <https://doi.org/10.5194/hess-16-3309-2012>, <https://hess.copernicus.org/articles/16/3309/2012/>, 2012.
- 635 Vaittinada Ayar, P., Vrac, M., Bastin, S., Carreau, J., Déqué, M., and Gallardo, C.: Intercomparison of statistical and dynamical downscaling models under the EURO- and MED-CORDEX initiative framework: Present climate evaluations, *Clim. Dyn.*, 46, 1301–1329, <https://doi.org/https://doi.org/10.1007/s00382-015-2647-5>, 2015.
- Villani, C.: Optimal Transport: Old and New, *Grundlehren der mathematischen Wissenschaften*, Springer-Verlag Berlin Heidelberg, 1 edn.,  
640 <https://doi.org/10.1007/978-3-540-71050-9>, 2008.
- Vrac, M.: Multivariate bias adjustment of high-dimensional climate simulations: the Rank Resampling for Distributions and Dependences (R2D2) bias correction, *Hydrol. Earth Syst. Sci.*, 22, 3175–3196, <https://doi.org/10.5194/hess-22-3175-2018>, <https://hess.copernicus.org/articles/22/3175/2018/>, 2018.
- Vrac, M. and Thao, S.: R<sup>2</sup>D<sup>2</sup> v2.0: accounting for temporal dependences in multivariate bias correction via analogue rank resampling, *Geosci. Model Dev.*, 13, 5367–5387, <https://doi.org/10.5194/gmd-13-5367-2020>, <https://gmd.copernicus.org/articles/13/5367/2020/>, 2020.
- 645 Vrac, M. and Vaittinada Ayar, P.: Influence of bias correcting predictors on statistical downscaling models, *J. Appl. Meteorol. Climatol.*, 56, 5–26, <https://doi.org/https://doi.org/10.1175/JAMC-D-16-0079.1>, <https://journals.ametsoc.org/view/journals/apme/56/1/jamc-d-16-0079.1.xml>, 2017.
- Vrac, M., Drobinski, P., Merlo, A., Herrmann, M., Lavaysse, C., Li, L., and Somot, S.: Dynamical and statistical downscaling of the French  
650 Mediterranean climate: uncertainty assessment, *Nat. Hazards Earth Syst. Sci.*, 12, 2769–2784, <https://doi.org/10.5194/nhess-12-2769-2012>, <https://nhess.copernicus.org/articles/12/2769/2012/>, 2012.
- Vrac, M., Noël, T., and Vautard, R.: Bias correction of precipitation through Singularity Stochastic Removal: Because occurrences matter, *J. Geophys. Res. D*, 121, 52U37–5258, <https://doi.org/https://doi.org/10.1002/2015JD024511>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2015JD024511>, 2016.
- 655 Wilks, D. S.: Stochastic weather generators for climate-change downscaling, part II: multivariable and spatially coherent multisite downscaling, *Wiley Interdiscip. Rev. Clim. Change*, 3, 267–278, <https://doi.org/https://doi.org/10.1002/wcc.167>, <https://onlinelibrary.wiley.com/doi/abs/10.1002/wcc.167>, 2012.
- Xiaoli, L., Coulibaly, P., and Evora, N.: Comparison of data-driven methods for downscaling ensemble weather forecasts, *Hydrol. Earth Syst. Sci.*, 12, 615–624, <https://doi.org/10.5194/hess-12-615-2008>, <https://hess.copernicus.org/articles/12/615/2008/>, 2008.
- 660 Xu, C.-y.: From GCMs to river flow: a review of downscaling methods and hydrologic modelling approaches, *Prog. Phys. Geogr.: Earth Environ.*, 23, 229–249, <https://doi.org/10.1177/030913339902300204>, <https://doi.org/10.1177/030913339902300204>, 1999.
- Zscheischler, J., Westra, S., Hurk, B., Seneviratne, S., Ward, P., Pitman, A., AghaKouchak, A., Bresch, D., Leonard, M., Wahl, T., and Zhang, X.: Future climate risk from compound events, *Nat. Clim. Chang.*, 8, 469–477, <https://doi.org/10.1038/s41558-018-0156-3>, 2018.




**Figure 1.** Left: Map of elevation (in m) over France. The region of interest lies in the south-east box. Right: Mean summer (JJA) temperature (in °C) and precipitation (in mm/day) for the region of interest (up: IPSL-CM5A-MR; bottom: KNMI-RACMO22E) over 1951-2010.




**Figure 2.** a) Probability densities and covariance matrices of the VAR processes and their corrections. b) Auto-correlation of the first coordinate. c) Cross-auto-correlation between the first and the second coordinates. d) Cross-auto-correlation between the second and the first coordinates. e) Auto-correlation of the second coordinate.







**Figure 3.** The first row of matrices is the correlations between corrections of the VAR with different choice of starting row for the reconstruction step. The second row is the correlations between corrections of the VAR with different choice of lag. The starting row for the reconstruction step is the middle value between 0 and lag of corrections. The first column is the correction with the method TSMBC with the underlying method dOTC, and the second column is also TSMBC, but with the underlying method RBC.




**Figure 4.** Boxplots of bias reduction in mean and standard deviation ( $BR_E$  and  $BR_\sigma$ ) in calibration (first column) and projection (second column) periods for temperature and precipitations in summer. The closer the boxplot is to 1, the closer the results are to the reference and therefore the better they are.




**Figure 5.** Boxplots of bias reduction in lag 2-(cross)-auto-correlations ( $BR_{\rho_2}$ ) in calibration (first column) and projection (second column) period for temperature and precipitations in summer. The closer the boxplot is to 1, the closer the results are to the reference and therefore the better they are.




**Figure 6.** Map of lag-1 (two first rows) and lag-4 (two last rows) auto-correlations of precipitation in summer. The first and third rows are for the calibration period, the second and fourth rows are for the projection period. The first (resp. second, third, fourth and last) column gives the maps of auto-correlations of the RCM to correct (resp. the GCM, the correction dOTC, the correction TSMBC-5 and the correction TSMBC-10 with method L2V). A key point of this figure is to compare the evolution between calibration and projection periods for the RCM, GCM and the corrections. The evolution of the TSMBC corrections is similar to the GCM evolution, which is different from the RCM evolution, leading to a fail of the correction in projection period.



**Figure 7.** Auto-correlogram (lag 1) between temperatures and precipitation in calibration period in summer for **a)** the model; **b)** the observations; **c)** the S2V dOTC correction and **d)** the S2V TSMBC-5 correction.



**Figure 8.** Bias reduction of dependence ( $BR_{WV}$ ) values, based on the Wasserstein distances computed on bivariate (correlation, distance) distributions between reference and the different BC datasets or model simulations in summer. The (correlation, distance) 2d-distributions come from the calculated correlograms (see figure 7 and text for details). The first line of matrices corresponds to the calibration period, and the second line to the projection period. The BC results and model simulations correspond to rows. The correlations are calculated between tas and tas (first matrix), tas and pr (second matrix), pr and tas (third matrix) and pr and pr (fourth matrix). For each matrix, the columns correspond to different lags and thus correlations indicate auto-correlations. Hence, the two central matrices (tas/pr and pr/tas) contain cross-correlations and cross-auto-correlations. In order to compare the shape of (correlation,distance) set, a normalization is performed separately for each cell (i.e., each couple method-lag) of each matrix. This normalization allows us characterizing the pattern of the distributions, and to get rid of the marginal properties. Hence, the comparison between different couples method-lag is possible but only to characterize the shape of the (distance, correlation) distributions. The closer the  $BR_{WV}$  value is to 1, the closer the results are to the reference and therefore the better they are.



**Figure 9.** Same as Fig. 8, but with a normalization by column (i.e., by given lag), before computing the  $BR_W$  values. Hence, the  $BR_W$  values of a given method for two different lags cannot be compared with this normalization. The closer the  $BR_W$  value is to 1, the closer the results are to the reference and therefore the better they are.



	dOTC	TSMBC-5	TSMBC-10
L1V	$1 \times 1 \times 1 \times 1 = 1$	$1 \times 1 \times 1 \times (5 + 1) = 6$	$1 \times 1 \times 1 \times (10 + 1) = 11$
L2V	$2 \times 1 \times 1 \times 1 = 2$	$2 \times 1 \times 1 \times (5 + 1) = 12$	$2 \times 1 \times 1 \times (10 + 1) = 22$
S1V	$1 \times 16 \times 13 \times 1 = 208$	$1 \times 16 \times 13 \times (5 + 1) = 1248$	$1 \times 16 \times 13 \times (10 + 1) = 2288$
S2V	$2 \times 16 \times 13 \times 1 = 416$	$2 \times 16 \times 13 \times (5 + 1) = 2496$	$2 \times 16 \times 13 \times (10 + 1) = 4576$

**Table 1.** Summary of the dimensions of the bias correction for each methods used in Sec. 4. For each cell, the first value corresponds to the dependence or not between temperature and precipitations. The second and third are the dimension of the grid of the RCM ( $16 \times 13$  grid cells). The last values is the dimension to correct until the lag 0 for the first column, the lag 5 for the second column and the lag 10 for the last column.