



# Space-time dependence of compound hot-dry events in the United States: assessment using a multi-site multi-variable weather generator

Manuela I. Brunner<sup>1</sup>, Eric Gilleland<sup>1</sup>, and Andrew W. Wood<sup>1</sup>

<sup>1</sup>Research Applications Laboratory, National Center for Atmospheric Research, 3450 Mitchell Ln, Boulder, CO 80301, USA

**Correspondence:** Manuela Brunner (manuela.i.brunner@gmail.com)

**Abstract.** Compound hot and dry events can lead to severe impacts whose severity may depend on their time scale and spatial extent. Despite their potential importance, the climatological characteristics of these joint events have received little attention regardless of growing interest in climate change impacts on compound events. Here, we ask how event time scale relates to (1) spatial patterns of compound hot-dry events in the United States, (2) the spatial extent of compound hot-dry events, and (3) the importance of temperature and precipitation as drivers of compound event occurrence. To study such rare spatial and multivariate events, we introduce a multi-site multi-variable weather generator (*PRSim.weather*), which enables generation of a large number of spatial compound hot-dry events. We show that the stochastic model realistically simulates distributional and temporal autocorrelation characteristics of temperature and precipitation at single sites, dependencies between the two variables, spatial correlation patterns, and spatial heat and drought indicators and their co-occurrence probabilities. The results of our compound event analysis demonstrate that (1) the Northwestern and Southeastern United States are most susceptible to compound hot-dry events independent of time scale and susceptibility decreases with increasing time scale, (2) the spatial extent and time scale of compound events are strongly related with sub-seasonal events (1–3 months) showing the largest spatial extents, and (3) the importance of temperature and precipitation as drivers of compound events varies with time scale where temperature is most important at short and precipitation at seasonal time scales. We conclude that time scale is an important factor to be considered in compound event assessments and suggest that climate change impact assessments should consider several instead of a single time scale when looking at future changes in compound event characteristics. The largest future changes may be expected for short compound events because of their strong relation to temperature.

## 1 Introduction

Compound hot and dry events, i.e. events that are extreme with respect to both temperature and precipitation, can lead to severe impacts in agriculture and other sectors as illustrated by the 2010 heatwave-drought in Russia or the 2012 heatwave-drought in the Central United States (US, Mo and Lettenmaier, 2015) which led to substantial reductions in crop yields (Wegren, 2011; Christian et al., 2020; Fuchs et al., 2012). The US has been shown to be affected by concurrent hot and dry events at different time scales including short and long events effective at weekly to monthly (Zhang et al., 2020) and seasonal to annual time scales (Alizadeh et al., 2020), respectively. The interest in these impactful compound events is reflected in



25 an increasing number of studies assessing changes in their frequency of occurrence. Substantial increases in the number of concurrent droughts and heat waves over the last few decades have been reported not just for the US (Alizadeh et al., 2020; Mazdiyasn and AghaKouchak, 2015; Tavakol et al., 2020) but also globally (Feng et al., 2020; Sarhadi et al., 2018) and for other regions of the world such as China (Wu et al., 2019; Zhou and Liu, 2018; Yu and Zhai, 2020) and Europe (Manning et al., 2019).

30 While frequency of occurrence is an important factor determining impacts, the severity of impacts related to compound events likely also depends on their spatial extent, i.e. how large the affected region is, and their time scale, i.e. whether they just last weeks or extend over a longer period of time. Indeed, the spatio-temporal behavior is a common target of analyses in general for drought, a related phenomenon, as in the multi-temporal severity-area-duration analyses presented by Andreadis et al. (2005). Despite their potential importance for understanding and projecting the physical manifestation and impacts of  
35 compound events, these spatio-temporal characteristics have received comparably little attention. Only recently, Alizadeh et al. (2020) and Wu et al. (2020) have shown that the area affected by concurrent hot-dry extremes has increased significantly over the past few decades in the US and globally for long, i.e. seasonal, time scales. However, it remains to be investigated how the time scale of compound events influences their characteristics and spatial extent.

This study aims to deepen our understanding of how the time scale of compound hot-dry events in the US relates to (1) spatial  
40 patterns of compound event affectedness, (2) spatial extents of compound events, and (3) the role of temperature and precipitation as drivers of compound events by focusing on multivariate and spatially compounding extreme events (Zscheischler et al., 2020). To answer the question of how time scale shapes compound event characteristics, we determine the probability, extent, and drivers of spatial compound heatwaves and meteorological drought over the conterminous US (CONUS) for different time scales ranging from weekly to annual events.

45 Studying such spatial compound events is challenging because they are rare in observational records (Zscheischler et al., 2018). This challenge can be tackled by developing stochastic simulation approaches to generate large data sets with similar statistical properties as the observations (Vogel and Stedinger, 1988). A stochastic approach to simulate spatial compound hot-dry events at different time scales needs to (1) represent spatial dependencies between sites to capture the spatial aspect, (2) represent dependencies between variables to capture dependencies between precipitation and temperature, and (3) be continuous to enable studying time scales from weeks to years. However, existing models often only fulfill one or two out of  
50 these three requirements. On the one hand, existing spatial models for simulating spatial extreme events such as the conditional exceedance model by Heffernan and Tawn (2004) are event-based (Keef et al., 2013; Diederer et al., 2019) and often applied to one variable, e.g. flood peaks. On the other hand, continuous stochastic approaches such as autoregressive moving average type models (Stedinger and Taylor, 1982) or bootstrap approaches (Rajagopalan et al., 2010) do not represent spatial dependencies  
55 well. Therefore, Brunner and Gilleland (2020) recently proposed a novel stochastic approach for simulating continuous stream-flow time series in multiple catchments based on the wavelet transform. The Phase Randomization Simulation using wavelets (*PRSim.wave*) model combines an empirical spatio-temporal model based on the wavelet transform and phase randomization with the flexible four-parameter kappa distribution and builds on an earlier, univariate version of the model (PRSim; Brunner



et al., 2019). It is able to simulate continuous, spatially consistent time series but has so far only been applied to one variable  
60 (streamflow).

We extend *PRSim.wave*, here, to multiple variables by proposing a multi-site multi-variable stochastic weather generator  
(*PRSim.weather*) that simulates long time series of spatially consistent temperature ( $T$ ) and precipitation ( $P$ ) time series.  
This multi-site multi-variable stochastic model reproduces local variable distributions using flexible distributions for  $T$  and  $P$   
and introduces spatio-temporal and variable dependence using the wavelet transform (Torrence and Compo, 1998) and phase  
65 randomization (Schreiber and Schmitz, 2000; Lancaster et al., 2018). Using this multi-site multi-variable generator to simulate  
a large set of spatial compound hot-dry events will help to shed light on the question of how time scale shapes compound  
event characteristics including spatial extent. Thus, this analysis will provide crucial information to increase preparedness and  
develop adaptation measures to potentially impactful spatial compound events.

## 2 Methods and Materials

70 We develop a multi-variable multi-site weather generator that stochastically simulates spatially consistent  $T$  and  $P$  time series  
for a large number of locations. We apply this model to a gridded  $T$  and  $P$  data set in the CONUS to generate a large sample  
of spatial compound hot-dry events. We subsequently use this sample to determine which regions in the US are susceptible to  
compound events and large spatial compound event extents at different time-scales. Last, we look at how the importance of  $T$   
and  $P$  for compound event development varies with time scale.

### 75 2.1 Study region and data

The analysis is performed using a gridded data set of  $T$  and  $P$  time series for 894 equally-spaced grid cells in the CONUS.  $T$   
and  $P$  data were obtained from the ERA5-Land reanalysis for the period 1981–2018 (ECMWF, 2019). ERA5-Land relies on  
atmospheric forcing from the ERA5 reanalysis (Hersbach et al., 2020) and provides variables at a spatial resolution of 9 km  
for the period 1981 to present. We chose a subset of regularly-spaced grid cells by sampling 1500 grid cells over the extent of  
80 the CONUS which resulted in 894 grid cells over land that are used for this analysis.

## 2.2 Methods

### 2.2.1 Stochastic multi-site multi-variable modeling

To study spatial compound hot-dry events, we develop a multi-site multi-variable weather generator *PRSim.weather* that en-  
ables simulation of large sets of spatially consistent compound hot-dry events. *PRSim.weather* combines an empirical spatio-  
temporal model based on the wavelet transform and phase randomization with two flexible distributions for  $T$  and  $P$ . It builds  
85 on the spatial stochastic model *PRSim.wave* (Phase Randomization Simulation using wavelets) proposed by Brunner and Gille-  
land (2020), which simulates continuous streamflow time series at multiple sites. We expand the functionality of *PRSim.wave* to




simulate multiple variables, i.e.  $T$  and  $P$ , at multiple sites. The weather generation procedure implemented in *PRSim.weather* consists of five main steps (Figure 1):

- 90 1. feed in observed daily  $T$  and  $P$  time series for multiple sites (here grid cells);
  2. fit monthly distributions to  $T$  and  $P$  time series at each site to capture seasonal variations in distribution parameters. For  $T$ , we use the flexible skewed exponential power (SEP) distribution with 4 parameters (Fernández and Steel, 1998), which generalizes the Gaussian distribution, can reproduce different skewness and kurtosis, and has been previously applied for multi-site temperature simulation (Evin et al., 2019). The parameters of the SEP distribution are estimated using  
95 L-moments (R-package *lmomco*; Asquith, 2020). For  $P$ , we use an extended Generalized Pareto distribution (E-GPD; Papastathopoulos and Tawn, 2013) with three parameters to model positive precipitation values. The E-GPD jointly models non-extreme and extreme values of  $P$  while bypassing the threshold selection problem as it enables smooth transitioning between a gamma-like distribution and a heavy-tailed Generalized Pareto distribution (GPD; Naveau et al., 2016). The E-GPD has been demonstrated to be valuable in multi-site precipitation modeling thanks to its flexibility (Evin et al.,  
100 2018). We combine the E-GPD with as many zero-values as in the observations to obtain the full  $P$  distribution;
  3. transform the  $T$  and  $P$  time series from the time to the frequency domain by decomposing the series into an amplitude and phase signal using a continuous wavelet transform with the Morlet wavelet (Torrence and Compo, 1998);
  4. generate a random time series using bootstrap resampling on the time series of one randomly sampled site by sampling years with replacement. Use the wavelet transform to also decompose this bootstrapped series in order to obtain a random  
105 phase signal;
  5. generate stochastic time series for  $T$  and  $P$  by applying the inverse wavelet transform to the observed amplitude signals and the randomly generated phases. Rank-transform the newly generated time series to the desired distribution for each month using the monthly distribution parameters derived in Step 2 (SEP parameters for  $T$  and E-GPD parameters for  $P$ ).
- 110 The spatial and variable dependencies are introduced in Step 5 by using the same random phases in the wavelet transform at all sites and for both variables.

The stochastic multi-site multi-variable model is evaluated with respect to the following characteristics: (1)  $T$  and  $P$  distributions (cdfs) at individual sites, (2) temporal autocorrelation of  $T$  and  $P$  (acfs) at individual sites, (3) spatial dependencies across sites for  $T$  and  $P$  (variograms), (4)  $T$ - $P$  variable dependencies (scatter plots), and (5) simulated spatial patterns of the  
115 standardized temperature index (STI), the standardized precipitation index (SPI), and the probability of concurrent high STI and low SPI anomalies at a 1-month aggregation level for moderate, severe and extreme events according to the empirical copula (see Section 2.2.2).

*PRSim.weather* is finally run  $n = 100$  times for the 894 grid cells in the US in order to substantially increase the sample size available for the assessment of spatial compound hot-dry events (28 years \* 100 = 2800 years).



**Figure 1.** Illustration of 5 working steps of *PRSim.weather*: (1) Feed in observed temperature ( $T$ ) and precipitation ( $P$ ) time series for multiple sites ( $1, \dots, n$ ); (2) fit monthly SEP distribution to  $T$  and E-GP distribution to  $P$  time series of all sites; (3) decompose  $T$  and  $P$  time series of all sites into an amplitude ( $A$ ) and phase ( $P$ ) signal using the wavelet transform; (4) generate one random time series using bootstrap resampling and decompose that random series into an amplitude and phase signal too, and (5) generate random  $T$  and  $P$  time series by combining the observed amplitude signal of each site and variable with the randomly generated phase signal and by backtransforming the signals to the time domain using the inverse wavelet transform. Rank-transform the newly generated signal to the desired distribution using the parameter estimates from Step 2.

## 120 2.2.2 Compound event analysis


While the focus is on the simulated series, compound events and their corresponding  $T$  and  $P$  characteristics are identified at different time scales in both the observed and stochastically simulated time series to assess the reliability of the stochastic model. To look at different time scales, we first convert the  $T$  and  $P$  series to weekly/monthly series using mean values and sums, respectively. We work with aggregation levels of 1 week to represent 'flash' compound events and of 1, 3, 6, and 12  
 125 months to represent sub-seasonal, seasonal, and annual time scales. In a second step, we transform the aggregated  $T$  and  $P$  series to series of standardized indices, which we will use to study relationships between the marginal behavior of compound events because they guarantee variable and site comparability. Standardized precipitation index (SPI) series (McKee et al., 1993) for each location are computed by transforming the  $P$  values to a standardized normal distribution (mean = 0 and sd = 1) using a site-specific Gamma distribution (Kolmogorov–Smirnov test did not not reject Gamma in over 80% of the grid  
 130 cells). Similarly, we compute standardized temperature index series (STI; Zscheischler et al., 2014) using the SEP distribution for transformation. Last, compound hot-dry events are identified for each time scale and grid cell using a bivariate empirical copula (Deheuvels, 1979; Genest and Favre, 2007), which describes the bivariate distribution of  $T$  (STI) and  $P$  (-SPI). We



change the sign of the SPI values to convert negative to positive anomalies as we are interested in events where both STI and -SPI are jointly exceeded. The empirical copula is described as:

$$135 \quad C_n(u, v) = \frac{1}{n} \sum_{i=1}^n 1\left(\frac{R_i}{n+1} \leq u, \frac{S_i}{n+1} \leq v\right), \quad (1)$$

where  $R_i$  and  $S_i$  represent pairs of ranks,  $n$  the sample size, and  $C_n(u, v)$  the rank-based estimator of the copula  $C(u, v)$ . An example of how the empirical copula (purple) is related to the margins STI (yellow) and -SPI (blue) is provided in Figure 2.



**Figure 2.** Illustration of the relationship between STI (yellow) and -SPI (blue) time series and their bivariate copula (purple) for one example grid cell. Concurrent high STI and -SPI events exceeding a threshold of 0.9 are highlighted by purple boxes.

Using the time series of empirical bivariate distribution values, we identify moderate, severe, and extreme droughts using three thresholds at 0.8, 0.9, and 0.95, respectively (see Figure 2 for an example with a threshold of 0.9).

140 To assess the spatial extent of compound events at different time scales, we determine the percentage of grid cells affected by each of the compound events identified at individual grid cells. Then, for each grid cell, we determine the median spatial extent of those events it is affected by.

To explain the role of the individual variables  $T$  and  $P$  in compound event occurrence, we compute Kendall's correlation between the median bivariate distribution (empirical copula) and the median standardized indices STI and SPI at different time scales. This correlation analysis is performed for nine hydro-climatic regions in the United States (Bukovsky; Bukovsky, 2011)  
 145 to quantify the regional spread in the role of STI and SPI for compound event development. We look at correlations for different time scales and event extremeness levels to assess to which degree these two factors influence STI and SPI importance.


### 3 Results

#### 3.1 Evaluating the weather generator

150 The multi-site multi-variable stochastic simulation approach *PRSim.weather* is capable of reproducing the observed statistical characteristics of  $T$  and  $P$  time series at multiple locations (Figure 3). The flexible SEP and E-GPD distributions capture the




local  $T$  and  $P$  distributions well as indicated by the good match of simulated with observed densities (Figure 3a,b). The temporal autocorrelation in both variables is realistically reproduced as shown by the good agreement of simulated with observed autocorrelation functions thanks to the observed frequency spectrum information used in the inverse wavelet transform (Figure 3c,d). The simulated time series well mimick the main temporal characteristics of the observed time series including seasonality and temporal event distribution/clustering as illustrated by three years of observed and simulated  $T$  and  $P$  data (Figure 3e,f). The  $T$ - $P$  variable dependence is also generally well captured thanks to the use of the same random phases for both variables when applying the inverse wavelet transform (Figure 3g,h). However, the number of high  $T$ -low  $P$  events at a daily scale is slightly underestimated. In addition to these local characteristics, spatial correlations are captured as illustrated by the similarity of observed and simulated variograms (Figure 4). However, the spatial correlation of  $T$  is slightly overestimated by the simulations. Achieving a 'perfect' joint representation of the three forms of dependence – temporal, spatial, and variable – is very challenging. The model is considered suitable for the analysis of compound hot-dry events because it has an acceptable performance with respect to all three aspects.




**Figure 3.** *PRSim.weather* evaluation for one example grid cell: (a, b) marginal distributions for observed and simulated  $T$  (orange) and  $P$  (blue), (c, d) temporal autocorrelation for observed and simulated  $T$  and  $P$ , (e, f) 3-year time series of  $T$  and  $P$  for observations and simulations, and (g, h) heat scatter plot of  $P$ - $T$  relationship in observations and simulations.





**Figure 4.** *PRSim.weather* evaluation for spatial dependence: (a) observed vs. simulated  $T$  (orange) variograms for 92 equally spaced grid cells (b) observed vs. simulated  $P$  (blue) variograms for 92 grid cells.


*PRSim.weather* enables simulation of a large sample of extreme events in terms of standardized temperature (STI) and precipitation indices (SPI), which reflect the spatial STI and SPI patterns detected in the observations for different levels of extremeness (Figure 5). While the simulated spatial STI and SPI patterns look similar as the observed ones, they are more expressed because of the larger sample size available. The spatial pattern for STI is rather weak with STI values being relatively homogeneously distributed except for the Pacific Northwest and along the West Coast where STI values are slightly higher than in the rest of the country. In contrast, the spatial pattern of median SPIs is expressed with substantially higher negative anomalies in the western than the eastern US and particularly strong negative anomalies in the Southwest.



**Figure 5.** Spatial distribution of median observed (upper panel) and simulated (lower panel) STIs (left panel) and SPIs (right panel) at a monthly time scale for three levels of extremeness: moderate (STI > 1 and SPI < -1), severe (STI > 1.5 and SPI < -1.5) and extreme (STI > 2 and SPI < -2). The darker the color, the more severe are median events.

The spatial STI and SPI patterns are reflected in the spatial distribution of the probability of concurrent hot-dry events, which is also realistically represented by *PRSim.weather* (Figure 6). The highest probability of concurrent hot-dry events at a monthly time scale are found in the Pacific Northwest, along the West coast, in the Rocky Mountains, and the Southeast, in particular in Texas. In contrast, concurrent hot-dry events are relatively rare in the Great Plains, the Midwest, and Florida. For the remainder of our analysis, we are focusing on the stochastic simulations because of their large sample size, which allows us to study rare spatial compound hot-dry events.






**Figure 6.** Spatial distribution of observed (upper panel) and simulated (lower panel) probability of occurrence of hot-dry events (number of compound events/total number of months, -) at a monthly time scale for three levels of extremeness: moderate ( $C_n > 0.8$ ), severe ( $C_n > 0.9$ ), and extreme ( $C_n : 0.95$ ). The darker the color, the more likely are concurrent hot-dry events.

### 3.2 Concurrent hot-dry events

The stochastically simulated compound hot-dry events reveal that the probability of co-occurring hot and dry periods is highest in the northwestern and southeastern US independently of the time scale considered (Figure 7). However, the probability of concurrent events decreases with increasing time scale, as can be expected due to the increasing aggregation of multiple weather events in longer periods, and event extremeness, as the events become more rare at more extreme thresholds. Still, there are spatial nuances depending on the time scale considered. For example, the high probabilities of concurrent events are located in the south for short time scales and move to the southeast as we move towards longer time scales. Time scale does not only affect local concurrence probabilities but also the size of the regions affected by concurrent hot-dry events, which decreases with increasing time scale and event extremeness. At an annual time scale, the probability of events at all extreme thresholds is negligible.




**Figure 7.** Probability of concurrent hot-dry events (number of compound events/total number of months, -) at different time scales (1 week, 1 month, 3 months, 6 months, 1 year) and for three levels of extremeness (moderate  $C_n > 0.8$ , severe  $C_n > 0.9$ , and extreme  $C_n > 0.95$ ) per grid cell. The darker the color, the higher the probability that a grid cell is affected by compound hot-dry events.


Different regions of the US differ not only in how susceptible they are to compound hot-dry event occurrence but also in how likely they are to be affected by a widespread (large spatial scale) compound event. The spatial occurrence patterns for spatially extensive compound hot-dry events vary by time scale (Figure 8). For moderate extremes, the Midwest is most affected by large events, with more prevalence in the upper Midwest at shorter time scales and the central to southern Midwest at longer time scales. For the severe category, the western and Southeastern regions are more affected, a similar spatial pattern to the probability of concurrent hot-dry events (Figure 7), although there are no large-scale events at short time scale. In addition, large spatial compound events generally become less likely as we move beyond the 3 month time scale and toward extreme



195 events (Figure 9). While  $\sim 20\%$  of the CONUS may be jointly affected by moderate and short compound events, spatial extents of compound events become small to non-existent for extreme and long-lasting (i.e. annual) compound events.




**Figure 8.** Spatial patterns of median spatial compound event extent per grid cell for different time scales and extremeness levels over nine hydro-climatic regions. The darker the color, the higher the median spatial extent of compound events a grid cell is affected by.



**Figure 9.** Relationship between spatial compound event extent, time scale, and event extremeness. Boxplots summarize the spread of median extent (percentage of overall area affected, -) across grid cells for weekly (red), monthly (orange), 3-monthly (purple), 6-monthly (blue), and annual (green) time scales and three levels of extremeness: moderate, severe, and extreme.

The importance of STI and SPI as drivers of compound event occurrence varies by time scale and level of extremeness (Figure 10).  $T$  is a particularly important driver at short time scales (Figure 10a). The importance of  $P$  as a driver of compound event occurrence increases with time scale up to event durations of 6 months but decreases with level of extremeness (Figure 10b). In summary, the longer the time scale, the more important  $P$  becomes as a driver compared to  $T$  (up to a seasonal time scale).

200



**Figure 10.** Importance of  $T$  and  $P$  as drivers of compound events across time scales and extremeness levels. Correlation of bivariate distribution (empirical copula) with (a) STI and (b) SPI for nine hydro-climatic (Bukovsky) regions (spread of boxplot) per time scale (color) and level of extremeness (hue).



#### 4 Discussion

The multi-site multi-variable stochastic model *PRSim.weather* proposed for the joint simulation of  $T$  and  $P$  at multiple sites has been shown to be suitable for the simulation of spatial compound hot-dry events. It reproduces the distributional and temporal autocorrelation characteristics of  $T$  and  $P$  at single sites, the dependence between the two variables, the spatial correlation of  $T$  and  $P$  across sites, and spatial patterns of STI, SPI, and their concurrence probabilities. Further model development should focus on how to improve the representation of dependencies in very high  $T$ -low  $P$  events at a daily scale. While the current application focuses on the two variables  $T$  and  $P$  in the US, the model can be adapted to other regions, other variables, and a multivariate context where more than two variables are of interest. Adapting the model to other regions and variables requires reconsidering distribution choices and extending it to a multivariate context necessitates adding more input variables, which are subsequently randomized in the same way as all other variables. Potential multivariate applications include the simulation of spatial concurrent pluvial, river, and coastal flooding by jointly modeling precipitation, discharge, and water levels or the joint simulation of wildfire drivers such as wind speed, temperature, and humidity. Please note that while the model will be able to retain the statistical dependencies between variables to some degree, individual simulated events may not necessarily be physically consistent if many variables are jointly simulated. If physical consistency is a requirement for a specific application, stochastic approaches may be combined with physical approaches as e.g. in the weather generator AWE-GEN-2 by Peleg et al. (2017).

The finding that the western and southeastern US are most likely to be affected by compound hot-dry events suggests that the likelihood of compound event occurrence is somehow related to precipitation seasonality with regions receiving most of their precipitation in winter or spring and comparably less in summer and fall (Finkelstein and Truppi, 1991) being most likely to be affected by compound events. In 'normal' years, both the western and southeastern US receive a large part of their precipitation through recurrent patterns such as atmospheric rivers (Rutz et al., 2015) or tropical cyclones (Kunkel et al., 2012), respectively. Anomalies can arise because of temporal shifts or a weakening of these patterns in specific seasons/years. In addition, the regions most likely to experience compound events are the regions found to be most susceptible to heatwaves in the US (Smith et al., 2013).

Our finding that spatial extents of compound events are largest for moderate events at subseasonal time scales implies that while these moderate events may have less severe impacts at a local scale, they may still be highly relevant at a regional scale. Compound events with large spatial extents represent a particular management challenge because they may preclude the transfer of resources and emergency supplies from one to another region. Consequently, the societal impacts of large-scale compound events can be amplified, since many coping strategies are predicated on some degree of resource transfers from less severely affected adjacent regions.

The finding that temperature is a comparably more important driver for short compound events only while precipitation is comparably more important at seasonal time scales corroborates the findings of previous studies about the importance of different hydro-meteorological drivers at different time scales. Zhang et al. (2020) have shown that temperature is the most important hydro-meteorological driver of short term concurrent hot-dry extremes, which aligns with our findings. In addition,



235 Tavakol et al. (2020) have shown that at long (i.e. annual) time scales, hot-dry-windy events co-occurred with major heat waves, which is in line with our finding that temperature is an important driver of extreme compound hot-dry events at seasonal to annual time scales.

Future changes in the frequency and severity of compound hot-dry events are expected because of both changes in temperature and precipitation and their interdependence. The importance of temperature as an important driver of short and extreme  
240 compound hot-dry events suggests that the increasing temperatures associated with climate change may induce future changes in the frequency and magnitude of short and extreme compound events. Such future increases have been projected globally (Wu et al., 2020) and regionally e.g. for China (Zhou and Liu, 2018). In addition, previous studies have shown that the number and intensity of compound hot-dry events may increase because temperature and precipitation may become increasingly  
245 coupled/correlated in summer (De Luca et al., 2020; Zscheischler and Seneviratne, 2017) possibly as a consequence of an intensification of land-atmosphere feedbacks (Seneviratne et al., 2010). As the number of compound events increases locally, the area exposed to compound hot-dry events is projected to increase with global warming (Vogel et al., 2019) continuing a trend that has been already observed during the past few decades (Alizadeh et al., 2020). How exactly future changes in compound event extents relate to changes in drought spatial extent (Brunner et al., 2020) and in heatwave spatial extent remains to be investigated.

## 250 5 Summary and Conclusions

We introduce the multi-variable multi-site stochastic model *PRSim.weather* to simulate continuous and spatially consistent multivariate time series. The model is shown to realistically simulate distributional and temporal autocorrelation characteristics of temperature and precipitation at single sites, dependencies between the two variables up to moderate extremes, spatial correlation patterns, and spatial heat and drought indicators and their co-occurrence probabilities for a gridded large-sample  
255 data set in the United States. However, future work is needed to improve the representation of very extreme hot-dry events. We apply the stochastic model to generate a large set of spatial and multivariate hot-dry events and use these simulated compound events to assess how event time scale and extremeness influence the spatial affectedness by compound hot-dry events over the United States, the spatial extent of compound events, and their main drivers temperature and precipitation. Our results show that (1) the Northwest and Southeast are most likely to be affected by compound hot-dry events independent of time scale; (2)  
260 the spatial extent of compound hot-dry events decreases with increasing event extremeness and time scale, i.e., the events with the largest spatial extents are typically short and only moderately extreme; and (3) temperature is an important driver of short compound events while precipitation is an important driver at seasonal time scales particularly for the moderately extreme events. These findings highlight that occurrences of compound events are strongly influenced by the time scales at which they are defined. Research to quantify current compound event risk and to project it into the future will need to take time scale  
265 into consideration, especially as it also influences the sensitivity to different climate drivers and their potential future changes. Considering space-time scales in compound event assessments will allow us to make nuanced statements about which types of compound events may be changing because of increasing temperatures in a warming world. For example, short compound



events and therefore events with large spatial extents may become more frequent with increasing temperatures, which will pose new regional management challenges.

270 *Code and data availability.* The ERA5-Land temperature and precipitation data used for this analysis can be downloaded from the Copernicus Climate Data Store: <https://cds.climate.copernicus.eu/cdsapp#!/dataset>. The stochastic weather generator *PRSim.weather* is implemented in the R-package *PRSim* under the function *PRSim.weather* and available for download: <https://cran.r-project.org/web/packages/PRSim/>.

*Author contributions.* MIB developed the study concept and stochastic simulation model, performed all the analyses, and wrote the first draft of the manuscript. EG provided methodological advice and revised and edited the manuscript. AW contributed to the interpretation of  
275 the results and revised and edited the manuscript.

*Competing interests.* The authors declare no competing interests.

*Acknowledgements.* This work was supported by the Swiss National Science Foundation via a PostDoc.Mobility grant (Number: P400P2\_183844, granted to MIB). We would like to acknowledge high-performance computing support from Cheyenne (doi:10.5065/D6RX99HX) provided by NCAR's Computational and Information Systems Laboratory, sponsored by the National Science Foundation.





## 280 References

- Alizadeh, M. R., Adamowski, J., Nikoo, M. R., AghaKouchak, A., Dennison, P., and Sadegh, M.: A century of observations reveals increasing likelihood of continental-scale compound dry-hot extremes, *Science Advances*, 6, 1–12, <https://doi.org/10.1126/sciadv.aaz4571>, 2020.
- Andreadis, K. M., Clark, E. A., Wood, A. W., Hamlet, A. F., and Lettenmaier, D. P.: Twentieth-century drought in the conterminous United States, *Journal of Hydrometeorology*, 6, 985–1001, <https://doi.org/10.1175/JHM450.1>, 2005.
- 285 Asquith, W.: Imomco: L-moments, censored L-moments, trimmed L-moments, L-comoments, and many distributions, <https://cran.r-project.org/web/packages/Imomco/index.html>, 2020.
- Brunner, M. I. and Gilleland, E.: Stochastic simulation of streamflow and spatial extremes: a continuous, wavelet-based approach, *Hydrology and Earth System Sciences*, 24, 3967–3982, <https://doi.org/10.5194/hess-24-3967-2020>, 2020.
- Brunner, M. I., Bárdossy, A., and Furrer, R.: Technical note: Stochastic simulation of streamflow time series using phase randomization, *Hydrology and Earth System Sciences*, 23, 3175–3187, <https://doi.org/10.5194/hess-23-3175-2019>, 2019.
- 290 Brunner, M. I., Swain, D. L., Gilleland, E., and Wood, A.: Increasing importance of temperature as a driver of streamflow drought spatial extent, *Environmental Research Letters*, p. in press, <https://doi.org/10.1088/1748-9326/abd2f0>, 2020.
- Bukovsky, M. S.: Masks for the Bukovsky regionalization of North America, <http://www.narccap.ucar.edu/contrib/bukovsky/>, 2011.
- Christian, J. I., Basara, J. B., Hunt, E. D., Otkin, J. A., and Xiao, S.: Flash drought development and cascading impacts associated with the 2010 Russian heatwave, *Environmental Research Letters*, 15, 094078, <https://doi.org/10.1088/1748-9326/ab9faf>, 2020.
- 295 De Luca, P., Messori, G., Faranda, D., Ward, P. J., and Coumou, D.: Compound warm-dry and cold-wet events over the Mediterranean, *Earth System Dynamics*, 11, 793–805, <https://doi.org/10.5194/esd-11-793-2020>, 2020.
- Deheuvels, P.: La fonction de dépendance empirique et ses propriétés. Un test non paramétrique d'indépendance, *Bulletin de la Classe des sciences*, 65, 274–292, <https://doi.org/10.3406/barb.1979.58521>, 1979.
- 300 Diederer, D., Liu, Y., Gouldby, B., Diermanse, F., and Vorogushyn, S.: Stochastic generation of spatially coherent river discharge peaks for continental event-based flood risk assessment, *Natural Hazards and Earth System Sciences*, 19, 1041–1053, <https://doi.org/10.5194/nhess-19-1041-2019>, 2019.
- ECMWF: ERA5-Land hourly data from 1981 to present, <https://doi.org/10.24381/cds.e2161bac>, 2019.
- Evin, G., Favre, A. C., and Hingray, B.: Stochastic generation of multi-site daily precipitation focusing on extreme events, *Hydrology and Earth System Sciences*, 22, 655–672, <https://doi.org/10.5194/hess-22-655-2018>, 2018.
- 305 Evin, G., Favre, A. C., and Hingray, B.: Stochastic generators of multi-site daily temperature: comparison of performances in various applications, *Theoretical and Applied Climatology*, 135, 811–824, <https://doi.org/10.1007/s00704-018-2404-x>, 2019.
- Feng, S., Wu, X., Hao, Z., Hao, Y., Zhang, X., and Hao, F.: A database for characteristics and variations of global compound dry and hot events, *Weather and Climate Extremes*, 30, 100299, <https://doi.org/10.1016/j.wace.2020.100299>, 2020.
- 310 Fernández, C. and Steel, M. F.: On bayesian modeling of fat tails and skewness, *Journal of the American Statistical Association*, 93, 359–371, <https://doi.org/10.1080/01621459.1998.10474117>, 1998.
- Finkelstein, P. L. and Truppi, L. E.: Spatial distribution of precipitation seasonality in the United States, *Journal of Climate*, 4, 373–385, 1991.
- Fuchs, B. A., Wood, D. A., and Ebbeka, D.: From too much to too little. How the central U.S. drought of 2012 evolved out of one of the most devastating floods in record in 2011, Tech. rep., National Drought Mitigation Center, Lincoln, [https://www.drought.gov/drought/sites/drought.gov.drought/files/media/reports/regional\\_outlooks/CentralRegion2012DroughtAssessment\\_1-5-15.pdf](https://www.drought.gov/drought/sites/drought.gov.drought/files/media/reports/regional_outlooks/CentralRegion2012DroughtAssessment_1-5-15.pdf), 2012.
- 315



- Genest, C. and Favre, A.-C.: Everything you always wanted to know about copula modeling but were afraid to ask, *Journal of Hydrologic Engineering*, 12, 347–367, [https://doi.org/10.1061/\(ASCE\)1084-0699\(2007\)12:4\(347\)](https://doi.org/10.1061/(ASCE)1084-0699(2007)12:4(347)), 2007.
- 320 Heffernan, J. E. and Tawn, J.: A conditional approach to modelling multivariate extreme values, *Journal of the Royal Statistical Society. Series B: Statistical Methodology*, 66, 497–546, <https://doi.org/10.1111/j.1467-9868.2004.02050.x>, 2004.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, 325 J.: The ERA5 Global Reanalysis, *Quarterly Journal of the Royal Meteorological Society*, pp. 1–51, <https://doi.org/10.1002/qj.3803>, 2020.
- Keef, C., Tawn, J. A., and Lamb, R.: Estimating the probability of widespread flood events, *Environmetrics*, 24, 13–21, <https://doi.org/10.1002/env.2190>, 2013.
- Kunkel, K. E., Easterling, D. R., Kristovich, D. A. R., Gleason, B., Stoecker, L., and Smith, R.: Meteorological causes of the secular variations in observed extreme precipitation events for the conterminous United States, *Journal of Hydrometeorology*, 13, 1131–1141, 330 <https://doi.org/10.1175/JHM-D-11-0108.1>, 2012.
- Lancaster, G., Iatsenko, D., Pidde, A., Ticcinelli, V., and Stefanovska, A.: Surrogate data for hypothesis testing of physical systems, *Physics Reports*, 748, 1–60, <https://doi.org/10.1016/j.physrep.2018.06.001>, 2018.
- Manning, C., Widmann, M., Bevacqua, E., Van Loon, A. F., Maraun, D., and Vrac, M.: Increased probability of compound long-duration dry and hot events in Europe during summer (1950–2013), *Environmental Research Letters*, 14, 094006, <https://doi.org/10.1088/1748-9326/ab23bf>, 2019.
- Mazdiyasn, O. and AghaKouchak, A.: Substantial increase in concurrent droughts and heatwaves in the United States, *Proceedings of the National Academy of Sciences of the United States of America*, 112, 11484–11489, <https://doi.org/10.1073/pnas.1422945112>, 2015.
- McKee, T. B., Doesken, N. J., and Kleist, J.: The relationship of drought frequency and duration to time scales, in: *Proceedings of the 8th Conference on Applied Climatology*, American Meteorological Society, January, pp. 179–184, Anaheim, California, <https://doi.org/citeulike-article-id:10490403>, 1993.
- 340 Mo, K. C. and Lettenmaier, D. P.: Heat wave flash droughts in decline, *Geophysical Research Letters*, 42, 2823–2829, <https://doi.org/10.1002/2015GL064018>, 2015.
- Naveau, P., Huser, R., Ribereau, P., and Hannart, A.: Modeling jointly low, moderate, and heavy rainfall intensities without a threshold selection, *Water Resources Research*, 52, 2753–2769, <https://doi.org/10.1002/2015WR018552>, 2016.
- 345 Papastathopoulos, I. and Tawn, J. A.: Extended generalised Pareto models for tail estimation, *Journal of Statistical Planning and Inference*, 143, 131–143, <https://doi.org/10.1016/j.jspi.2012.07.001>, 2013.
- Peleg, N., Fatichi, S., Paschalis, A., Molnar, P., and Burlando, P.: An advanced stochastic weather generator for simulating 2-D high-resolution climate variables, *Journal of Advances in Modeling Earth Systems*, 9, 1595–1627, <https://doi.org/10.1002/2013MS000282>, 2017.
- Rajagopalan, B., Salas, J. D., and Lall, U.: Stochastic methods for modeling precipitation and streamflow, in: *Advances in data-based approaches for hydrologic modeling and forecasting*, edited by Sivakumar, B. and Berndtsson, R., chap. 2, pp. 17–52, World Scientific, 2010.
- 350 Rutz, J. J., James Steenburgh, W., and Martin Ralph, F.: The inland penetration of atmospheric rivers over western North America: A Lagrangian analysis, *Monthly Weather Review*, 143, 1924–1944, <https://doi.org/10.1175/MWR-D-14-00288.1>, 2015.



- Sarhadi, A., Ausín, M. C., Wiper, M. P., Touma, D., and Diffenbaugh, N. S.: Multidimensional risk in a nonstationary climate: Joint probability of increasingly severe warm and dry conditions, *Science Advances*, 4, eaau3487, <https://doi.org/10.1126/sciadv.aau3487>, 2018.
- Schreiber, T. and Schmitz, A.: Surrogate time series, *Physica D: Nonlinear Phenomena*, 142, 346–382, [https://doi.org/10.1016/S0167-2789\(00\)00043-9](https://doi.org/10.1016/S0167-2789(00)00043-9), 2000.
- Seneviratne, S. I., Corti, T., Davin, E. L., Hirschi, M., Jaeger, E. B., Lehner, I., Orlowsky, B., and Teuling, A. J.: Investigating soil moisture-climate interactions in a changing climate: A review, *Earth-Science Reviews*, 99, 125–161, <https://doi.org/10.1016/j.earscirev.2010.02.004>, 2010.
- Smith, T. T., Zaitchik, B. F., and Gohlke, J. M.: Heat waves in the United States: definitions, patterns and trends, *Climatic Change*, 118, 811–825, <https://doi.org/10.1007/s10584-012-0659-2>, 2013.
- Stedinger, J. R. and Taylor, M. R.: Synthetic streamflow generation. 1. Model verification and validation, *Water Resources Research*, 18, 909–918, <https://doi.org/0733-9429>, 1982.
- Tavakol, A., Rahmani, V., and Harrington, J.: Temporal and spatial variations in the frequency of compound hot, dry, and windy events in the central United States, *Scientific Reports*, 10, 1–13, <https://doi.org/10.1038/s41598-020-72624-0>, 2020.
- Torrence, C. and Compo, G. P.: A practical guide to wavelet analysis, *Bulletin of the American Meteorological Society*, 79, 61–78, 1998.
- Vogel, M. M., Zscheischler, J., Wartenburger, R., Dee, D., and Seneviratne, S. I.: Concurrent 2018 hot extremes across northern hemisphere due to human-induced climate change, *Earth's Future*, 7, 692–703, <https://doi.org/10.1029/2019EF001189>, 2019.
- Vogel, R. M. and Stedinger, J. R.: The value of stochastic streamflow models in overyear reservoir design applications, *Water Resources Research*, 24, 1483–1490, <https://doi.org/10.1029/WR024i009p01483>, 1988.
- Wegren, S.: Food security and Russia's 2010 drought, *Eurasian Geography and Economics*, 52, 140–156, <https://doi.org/10.2747/1539-7216.52.1.140>, 2011.
- Wu, X., Hao, Z., Hao, F., Singh, V. P., and Zhang, X.: Dry-hot magnitude index: A joint indicator for compound event analysis, *Environmental Research Letters*, 14, <https://doi.org/10.1088/1748-9326/ab1ec7>, 2019.
- Wu, X., Hao, Z., Tang, Q., Singh, V. P., Zhang, X., and Hao, F.: Projected increase in compound dry and hot events over global land areas, *International Journal of Climatology*, pp. 1–11, <https://doi.org/10.1002/joc.6626>, 2020.
- Yu, R. and Zhai, P.: More frequent and widespread persistent compound drought and heat event observed in China, *Scientific Reports*, 10, 1–7, <https://doi.org/10.1038/s41598-020-71312-3>, 2020.
- Zhang, H., Wu, C., Yeh, P. J., and Hu, B. X.: Global pattern of short-term concurrent hot and dry extremes and its relationship to large-scale climate indices, *International Journal of Climatology*, 40, 5906–5924, <https://doi.org/10.1002/joc.6555>, 2020.
- Zhou, P. and Liu, Z.: Likelihood of concurrent climate extremes and variations over China, *Environmental Research Letters*, 13, 094 023, <https://doi.org/10.1088/1748-9326/aade9e>, 2018.
- Zscheischler, J. and Seneviratne, S. I.: Dependence of drivers affects risks associated with compound events, *Science Advances*, 3, 1–11, <https://doi.org/10.1126/sciadv.1700263>, 2017.
- Zscheischler, J., Michalak, A. M., Schwalm, C., Mahecha, M. D., Huntzinger, D. N., Reichstein, M., Berthier, G., Ciais, P., Cook, R. B., El-Masri, B., Huang, M., Ito, A., Jain, A., King, A., Lei, H., Lu, C., Mao, J., Peng, S., Poulter, B., Ricciuto, D., Shi, X., Tao, B., Tian, H., Viovy, N., Wang, W., Wei, Y., Yang, J., and Zeng, N.: Impact of large-scale climate extremes on biospheric carbon fluxes: An intercomparison based on MsTMIP data, *Global Biogeochemical Cycles*, 28, 585–600, <https://doi.org/10.1002/2014GB004826>, 2014.
- Zscheischler, J., Westra, S., Hurk, B. J. J. M. V. D., Seneviratne, S. I., Ward, P. J., Pitman, A., Aghakouchak, A., Bresch, D. N., and Leonard, M.: Future climate risk from compound events, *Nature Climate Change*, 8, 469–477, <https://doi.org/10.1038/s41558-018-0156-3>, 2018.



Zscheischler, J., Martius, O., Westra, S., Bevacqua, E., Raymond, C., Horton, R. M., Hurk, B. v. d., AghaKouchak, A., Jezequel, A., Mahecha, M. D., Maraun, D., Ramos, A. M., Ridder, N. N., Thiery, W., and Vignotto, E.: A typology of compound weather and climate events, *Nature Reviews Earth & Environment*, 1, 333–347, <https://doi.org/10.1038/s43017-020-0060-z>, 2020.