

Climate Change in the Baltic Sea Region: A Summary

H. E. Markus Meier^{1,2}, Madline Kniebusch¹, Christian Dieterich^{2,†}, Matthias Gröger¹, Eduardo Zorita³, Ragnar Elmgren⁴, Kai Myrberg^{5,6}, Markus ~~P.~~ Ahola⁷, Alena Bartosova², Erik Bonsdorff⁸, Florian Börgel¹, Rene Capell², Ida Carlén⁹, Thomas Carlund¹⁰, Jacob Carstensen¹¹, Ole B. Christensen¹², Volker Dierschke¹³, Claudia Frauen^{1,14}, Morten Frederiksen¹¹, Elie Gagel^{15,16}, Anders Galatius¹¹, Jari J. Haapala¹⁷, Antti Halkka¹⁸, Gustaf Hugelius¹⁹, Birgit Hünicke³, Jaak Jaagus²⁰, Mart Jüssi²¹, Jukka Käyhkö²², Nina Kirchner¹⁹, Erik Kjellström², Karol Kulinski²³, Andreas Lehmann²⁴, Göran Lindström², Wilhelm May²⁵, Paul A. Miller^{25,26}, Volker Mohrholz¹, Bärbel Müller-Karulis²⁷, Diego Pavón-Jordán²⁸, Markus Quante²⁹, Marcus Reckermann³⁰, Anna Rutgersson³¹, Oleg P. Savchuk²⁷, Martin Stendel¹², Laura Tuomi¹⁷, Markku Viitasalo³, Ralf Weisse³, Wenyan Zhang³

¹Department of Physical Oceanography and Instrumentation, Leibniz Institute for Baltic Sea Research Warnemünde, Rostock, Germany

²Research and Development Department, Swedish Meteorological and Hydrological Institute, Sweden

³Institute of Coastal Systems-Analysis and Modeling, Helmholtz-Zentrum Hereon, Geesthacht, Germany

⁴Department of Ecology, Environment and Plant Sciences, Stockholm University, Stockholm, Sweden

⁵Marine Research Centre, Finnish Environment Institute, Finland

⁶Marine Research Institute, University of Klaipeda, Lithuania

⁷Swedish Museum of Natural History, Stockholm, Sweden

⁸Environmental and Marine Biology, Åbo Akademi University, Finland

⁹Coalition Clean Baltic, Uppsala, Sweden

¹⁰Information and Statistics Department, Swedish Meteorological and Hydrological Institute, Norrköping, Sweden

¹¹Department of ~~EcoBio~~science, Aarhus University, Roskilde, Denmark

¹²National Centre for Climate Research, Danish Meteorological Institute, Copenhagen, Denmark

¹³Gavia EcoResearch, Winsen (Luhe), Germany

¹⁴Deutsches Klimarechenzentrum, Hamburg, Germany

¹⁵Department of Biology, University of Turku, Turku, Finland

¹⁶International Institute for Applied Systems Analysis (IIASA), Laxenburg, Austria

¹⁷Finnish Meteorological Institute, Helsinki, Finland

¹⁸Department of Biological and Environmental Sciences, University of Helsinki, Helsinki, Finland

¹⁹Department of Physical Geography, Stockholm University, Stockholm

²⁰Department of Geography, Institute of Ecology and Earth Sciences, University of Tartu, Tartu, Estonia

²¹Pro Mare, Estonia

²²Department of Geography and Geology, University of Turku, Finland

²³Institute of Oceanology, Polish Academy of Sciences, Gdansk, Poland

²⁴GEOMAR Helmholtz Centre for Ocean Research, Kiel, Germany

²⁵Centre for Environmental and Climate Science, Lund University, Lund, Sweden

²⁶Department of Physical Geography and Ecosystem Science, Lund University, Lund, Sweden

²⁷Baltic Sea Centre, Stockholm University, Stockholm, Sweden

²⁸Department of Terrestrial Ecology, Norwegian Institute for Nature Research (NINA), ~~P.O. Box 5685 Torgarden, N-7485~~ Trondheim, Norway

²⁹Institute of Coastal ~~Environmental~~ Chemistry, Helmholtz-Zentrum Hereon, Geesthacht, Germany

³⁰International Baltic Earth Secretariat, Helmholtz-Zentrum Hereon, Geesthacht, ~~21502~~, Germany

³¹Department of Earth Sciences, Uppsala University, Sweden

[†]Deceased

Correspondence to: H. E. Markus Meier (markus.meier@io-warnemuende.de)

Abstract. Based on the Baltic Earth Assessment Reports of this thematic issue in Earth System Dynamics and recent peer-reviewed literature, current knowledge ~~of about~~ the effects of global warming on past and future changes in climate of the Baltic Sea region is summarized and assessed. The study is an update of the Second Assessment of Climate Change (BACC II) published in 2015 and focusses on the atmosphere, land, cryosphere, ocean, sediments and the terrestrial and marine biosphere. Based on the summaries of the recent knowledge gained in paleo-, historical and future regional climate research, we find that the main conclusions from earlier assessments remain still valid. However, new long-term, homogenous observational records, e.g. for Scandinavian glacier inventories, sea-level driven saltwater inflows, so-called Major Baltic Inflows, and phytoplankton species distribution and new scenario simulations with improved models, e.g. for glaciers, lake ice and marine food web, have become available. In many cases, uncertainties can now be better estimated than before, because more models ~~were can be~~ included in the ensembles, especially for the Baltic Sea. With the help of coupled models, feedbacks between several components of the Earth System have been studied and multiple driver studies were performed, e.g. projections of the food web that include fisheries, eutrophication and climate change. New data sets and projections have led to a revised understanding of changes in some variables such as salinity. Furthermore, it has become evident that natural variability, in particular for the ocean on multidecadal time scales, is greater than previously estimated, challenging our ability to detect observed and projected changes in climate. In this context, the first paleoclimate simulations regionalized for the Baltic Sea region are instructive. Hence, estimated uncertainties for the projections of many variables increased. In addition to the well-known influence of the North Atlantic Oscillation, it was found that also other low-frequency modes of internal variability, such as the Atlantic Multidecadal Variability, have profound effects on the climate of the Baltic Sea region. Challenges were also identified, such as the systematic discrepancy between future cloudiness trends in global and regional models and the difficulty of confidently attributing large observed changes in marine ecosystems to climate change. Finally, we compare our results with other coastal sea assessments, such as the North Sea Region Climate Change Assessment (NOSCCA) and find that the effects of climate change on the Baltic Sea differ from those on the North Sea, since Baltic Sea oceanography and ecosystems are very different from other coastal seas such as the North Sea. While the North Sea dynamics is dominated by tides, the Baltic Sea is characterized by brackish water, a perennial vertical stratification in the southern sub-basins and a seasonal sea-ice cover in the northern sub-basins.

~~During the time in which this paper was prepared, shortly before submission, Christian Dieterich passed away (1964-2021). This sad event marked the end of the life of a distinguished oceanographer and climate scientist who made important contributions to the climate modeling of the Baltic Sea, North Sea and North Atlantic regions. This paper is dedicated to him.~~

1 Introduction

1.1 Overview

In this study, the results concerning climate change of the various articles of this thematic issue, the so-called Baltic Earth Assessment Reports (BEARs) coordinated by the Baltic Earth program¹ (Meier et al., 2014), and other relevant literature are summarized and assessed. We focus on the knowledge gained during 2013-2020 ~~of about~~ past, present and future climate changes in the Baltic Sea region. The methodology of all BEARs follows the earlier assessments of climate change in the Baltic Sea region (BACC Author Team, 2008; BACC II Author Team, 2015). The aim of this review is to inform and update scientists, policymakers and stakeholders about recent research results. The focus is on the atmosphere, hydrosphere, cryosphere, lithosphere and biosphere. In contrast to the earlier assessments, we do not investigate the impact of climate change on human society. We start (Section 1) with a summary of key messages from the earlier assessments of climate change in the Baltic Sea region, a description of the Baltic Sea region and its climate, a comparison of the Baltic Sea with other coastal seas and a

¹ <https://baltic.earth>

summary of current knowledge of global climate change assessed in the latest Intergovernmental Panel on Climate Change (IPCC) reports (see Table 1 of contents). In Section 2, the methods for the literature assessment, proxy data, instrumental measurements, climate model data and uncertainty estimates are outlined. In Section 3, the results of the assessment for selected variables (Table 24) under past (paleoclimate), present (historical period with instrumental data) and future (until 2100) climate conditions are presented, *inter alia* by summarizing the results in various papers of this special issue by Lehmann et al. (2021), Kuliński et al. (2021), Rutgersson et al. (2021), Weisse et al. (2021), Gröger et al. (2021a), Christensen et al. (2021), Meier et al. (2021a) and Viitasalo and Bonsdorff (2021) and by other relevant review studies. In Section 4, the interactions of climate with other anthropogenic drivers are summarized from Reckermann et al. (2021). As the adjacent North Sea has different physical characteristics and topographical features but is located in a similar climatic zone as the Baltic Sea, we compare the results of this assessment with the results of the North Sea Region Climate Change Assessment (NOSCCA; Quante and Colijn, 2016; Section 5). Knowledge gaps (Section 6), key messages (Section 7) and conclusions (Section 8) finalize the study. Acronyms used in this study are defined in Table 3.

1.2 The BACC and BEAR projects

This assessment is an update to the two BACC books, published as comprehensive textbooks in 2008 and 2015 (BACC Author Team, 2008; BACC II Author Team, 2015). The acronym BACC (BALTEX Assessment of Climate Change) refers to the Baltic Earth pre-cursor programme BALTEX (Baltic Sea Experiment; Reckermann et al., 2011). From the beginning, BALTEX tried to approach three basic questions: 1. What is the evidence for past and present regional climate change? 2. What are the model projections for future regional climate change? 3. Which impacts can we already observe in terrestrial and marine ecosystems?

First ideas for a comprehensive appraisal of the current knowledge on climate change and its impact on the Baltic Sea region evolved in 2004 as it became evident that there was a demand for this, in particular by the Baltic Marine Environment Protection Commission, the Helsinki Commission (HELCOM; BALTEX, 2005). A steering group of leading experts from the Baltic Sea region was enlisted, which elaborated a grand chapter structure at several preparatory workshops and meetings and also recruited a group of lead authors. In total, more than 80 scientists from 12 countries and all relevant scientific disciplines contributed to the first regional climate change assessment (BACC Author Team, 2008), which underwent a rigorous review process.

In 2011, a second edition of the BACC book was initiated as an update, but also as a complement to the first book, by including new topics like an overview of changes since the last glaciation, and a new section on regional drivers and attribution. The Second Assessment of Climate Change for the Baltic Sea Basin (BACC II Author Team, 2015) was published in 2015, used the same procedures and principles, but with a new steering and author group, and under the auspices of Baltic Earth, the successor of BALTEX. Close collaboration with HELCOM was envisaged from the very beginning, with HELCOM using material from both BACC assessments for their own climate change assessment reports (HELCOM, 2007; 2013b).

129 In 2018, the Baltic Earth Science Steering Group decided to produce a series of new assessment reports, the
 130 BEARs, on the current, six Baltic Earth Grand Challenges (1. Salinity dynamics in the Baltic Sea, 2. Land-Sea
 131 biogeochemical linkages in the Baltic Sea region, 3. Natural hazards and extreme events in the Baltic Sea region,
 132 4. Sea level dynamics in the Baltic Sea, 5. Regional variability of water and energy exchanges, 6. Multiple drivers
 133 for regional Earth system changes, see Baltic Earth, 2017), Earth System Models (ESMs) and projections for the
 134 Baltic Sea region. The BEARs are comprehensive, peer-reviewed review articles in journal format, and the
 135 update to BACC II (this article) is one of the ten envisaged contributions summarizing the current knowledge on
 136 regional climate change and its impacts, knowledge gaps and advice for future work. For further details about our
 137 knowledge on climate change, the reader is referred to the other BEARs. The close collaboration with HELCOM
 138 is continued in the joint HELCOM-Baltic Earth Expert Network of Climate Change (EN CLIME), which was
 139 assembled to produce a Baltic Earth – HELCOM Climate Change Fact Sheet for the Baltic Sea region².

140
 141 Hence, this thematic issue comprises nine BEARs and, in addition, this summary of the current knowledge about
 142 past, present and future climate changes for the Baltic Sea region ("BACC III"). Below a few key-words
 143 characterizing the BEARs' contents are listed:

- 144 1. Salinity dynamics of the Baltic Sea (Lehmann et al., 2021): water and energy cycles with focus on Baltic
 145 Sea salinity during past climate variability, meteorological patterns at various space and time scales and
 146 mesoscale variability in precipitation, variations in river runoff and various types of inflows of saline
 147 water, exchange of water masses between various sub-basins and vertical mixing processes. The paper
 148 also includes the observed trends of salinity during the last >100 years.
- 149 2. Baltic Earth Assessment Report on the biogeochemistry of the Baltic Sea (Kuliński et al., 2021): terrestrial
 150 biogeochemical processes and nutrient loads to the Baltic Sea, transformations of C, N, P in the coastal
 151 zone, organic matter production and remineralization, oxygen availability, burial and turnover of C, N, P
 152 in the sediments, the Baltic Sea CO₂ system and seawater acidification, role of specific microorganisms
 153 in Baltic Sea biogeochemistry, interactions between biogeochemical processes and chemical
 154 contaminants.
- 155 3. Natural hazards and extreme events in the Baltic Sea region (Rutgersson et al., 2021): extremes in
 156 wind, waves, and sea level, sea-effect snowfall, river floods, hot and cold spells in the atmosphere, marine
 157 heat waves, droughts, ice seasons, ice ridging, phytoplankton blooms and some implications of extreme
 158 events for society (including forest fires, coastal flooding, offshore wind mills and shipping).
- 159 4. Sea level dynamics and coastal erosion in the Baltic Sea region (Weisse et al., 2021): sea level
 160 dynamics and coastal erosion in past and future climates. The current knowledge about the diverse
 161 processes affecting mean and extreme sea level changes is assessed.
- 162 5. Coupled regional Earth system modelling in the Baltic Sea region (Gröger et al., 2021a): status report
 163 on coupled regional Earth system modeling with focus on the coupling between atmosphere and ocean,

² <https://helcom.fi/helcom-at-work/groups/state-and-conservation/en-clime/>,
<https://baltic.earth/projects/en-clime/index.php/en>, <https://helcom.fi/media/publications/Baltic-Sea-Climate-Change-Fact-Sheet-2021.pdf>

Formatiert: Schriftart: 10 Pt.

Formatiert: Absatz-Standschriftart, Englisch (USA)

Formatiert: Nicht unterstrichen, Schriftfarbe: Automatisch, Englisch (USA)

Formatiert: Absatz-Standschriftart, Englisch (USA)

Formatiert: Absatz-Standschriftart

Formatiert: Nicht unterstrichen, Schriftfarbe: Automatisch, Englisch (USA)

Formatiert: Absatz-Standschriftart, Englisch (USA)

Formatiert: Nicht unterstrichen, Schriftfarbe: Automatisch, Englisch (USA)

Formatiert: Absatz-Standschriftart, Englisch (USA)

Formatiert: Nicht unterstrichen, Schriftfarbe: Automatisch, Englisch (USA)

Formatiert: Absatz-Standschriftart, Englisch (USA)

Formatiert: Nicht unterstrichen, Schriftfarbe: Automatisch, Englisch (USA)

Formatiert: Absatz-Standschriftart, Englisch (USA)

atmosphere and land surface including dynamic vegetation, ocean, sea ice and waves and atmosphere and hydrological components to close the water cycle.

6. Atmospheric regional climate projections for the Baltic Sea ~~r~~Region until 2100 (Christensen et al., 2021): comparison of coupled and uncoupled regional future climate model projections. As the number of atmospheric scenario simulations of the ~~EURO~~EURO-CORDEX program (Kjellström et al., 2018; Teichmann et al., 2018; Jacob et al., 2018) is large, uncertainties can be better estimated and the effects of mitigation measures can be better addressed compared to earlier assessments.
7. Oceanographic regional climate projections for the Baltic Sea until 2100 (Meier et al., 2021a): new projections with a coupled physical-biogeochemical ocean model of future climate considering global sea level rise, regional climate change and nutrient input scenarios are compared with previous studies and the differences are explained by differing scenario assumptions and experimental setups.
8. Climate change and the Baltic Sea ecosystem: direct and indirect effects on species, communities and ecosystem function (Viitasalo ~~and~~ ~~Bonsdorff~~, 2021): impact of past and future climate changes on the marine ecosystem.
9. Human impacts and their interactions in the Baltic Sea region (Reckermann et al., 2021): interlinkages of factors controlling environmental changes. Changing climate is only one of the many anthropogenic and natural impacts that effect the environment. Other investigated factors are coastal processes, hypoxia, acidification, submarine groundwater discharge, marine ecosystems, non-indigenous species, land use and land cover (called natural) and agriculture and nutrient loads, aquaculture, fisheries, river regulations and restorations, offshore wind farms, shipping, chemical contaminants, unexploded and dumped warfare agents, marine litter and microplastics, tourism, and coastal management (called human-induced).

1.3 Summary of BACC I and II key messages

Quotation by the BACC II Author Team (2015):

“The key findings of the BACC I assessment were as follows:

- The Baltic Sea region is warming, and the warming is almost certain to continue throughout the twenty-first century.
- It is plausible that the warming is at least partly related to anthropogenic factors.
- So far, and as is likely to be the case for the next few decades, the signal is limited to temperature and to directly related variables, such as ice conditions.
- Changes in the hydrological cycle are expected to become obvious in the coming decades.
- The regional warming is almost certain to have a variety of effects on terrestrial and marine ecosystems—some will be more predictable (such as the changes in phenology) than others.

The key findings of the BACC II assessment [...] are as follows:

1. The results of the BACC I assessment remain valid.
2. Significant additional material has been found and assessed. Some previously contested issues have been resolved (such as trends in sea-surface temperature).

3. The use of multi-model ensembles seems to be a major improvement; there are first detection studies, but attribution is still weak.
4. Regional climate models still suffer from biases related to the heat and water balances. The effect of changing atmospheric aerosol load to date cannot be described; first efforts at describing the effect of land-use change have now been done.
5. Data homogeneity is still a problem and is sometimes not taken seriously enough.
6. The issue of multiple drivers on ecosystems and socioeconomics is recognized, but more efforts to deal with them are needed.
7. In many cases, the relative importance of different drivers of change, not only climate change, needs to be evaluated (e.g. atmospheric and aquatic pollution and eutrophication, overfishing, and changes in land cover).
8. Estimates of future concentrations and deposition of substances such as sulphur and nitrogen oxides, ammonia/ammonium, ozone, and carbon dioxide depend on future emissions and climate conditions. Atmospheric warming seems relatively less important than changes in emissions. The specification of future emissions is plausibly the biggest source of uncertainty when attempting to project future deposition or ocean acidification.
9. In the narrow coastal zone, the combination of climate change and land uplift acting together creates a particularly challenging situation for plant and animal communities in terms of adaptation to changing environmental conditions.
10. Climate change is a compounding factor for major drivers of changes in freshwater biogeochemistry, but evidence is still often based on small-scale studies in time and space. The effect of climate change cannot yet be quantified on a basin-wide scale.
11. Climate model scenarios show a tendency towards future reduced salinity, but due to the large bias in the water balance projections, it is still uncertain whether the Baltic Sea will become less or more saline.
12. Scenario simulations suggest that the Baltic Sea water may become more acidic in the future. Increased oxygen deficiency, increased temperature, changed salinity, and increased ocean acidification are expected to affect the marine ecosystem in various ways and may erode the resilience of the ecosystem.
13. When addressing climate change impacts on, for example, forestry, agriculture, urban complexes, and the marine environment in the Baltic Sea basin, a broad perspective is needed which considers not only climate change but also other significant factors such as changes in emissions, demographic and economic changes, and changes in land use.
14. Palaeoecological 'proxy' data indicate that the major change in anthropogenic land cover in the Baltic Sea catchment area occurred more than two thousand years ago. Climate model studies indicate that past anthropogenic land-cover change had a significant impact on past climate in the northern hemisphere and the Baltic Sea region, but there is no evidence that land cover change since AD 1850 was even partly responsible for driving the recent climate warming."

For comparison, the findings of this assessment study can be found in Section 8.

1.4 Baltic Sea region characteristics

1.4.1 Climate variability of the Baltic Sea Region

The Baltic Sea region (including the Kattegat) is located between maritime temperate and continental sub-arctic climate zones, in the latitude–longitude box 54°N – $66^{\circ}\text{N} \times 9^{\circ}\text{E}$ – 30°E (Fig. 1). The climate of the Baltic Sea region has a large variability due to the opposing effects of moist and relatively mild marine air flows from the North Atlantic Ocean and the Eurasian continental climate. The regional weather regimes vary depending on the exact location of the polar front and the strength of the westerlies, and both seasonal and interannual variations are considerable. The westerlies are particularly important in winter, when the temperature difference between the marine and continental air masses is large.

The southern and western parts of the Baltic Sea belong to the Central European mild climate zone in the westerly circulation. The northern part locates at the polar front and the winter climate is cold and dry due to cold arctic air outbreaks from the east. In terms of classical meteorology, during winter the polar front fluctuates over the Baltic Sea region but during summer it is located farther to the north. Depending on the particular year, the central part of the Baltic Sea can be either on the mild or the cold side of the polar front. The temperature difference between winter and summer is much larger in the north. During warm summers and cold winters the air pressure field is smooth and winds are weak, and blocking high pressure situations are common. During such periods, the weather can be very stable for several weeks.

The climate of the Baltic Sea region is strongly influenced by the large-scale atmospheric variability (e.g. Andersson, 2002; Tinz, 1996; Meier and Kauker, 2003; Omstedt and Chen, 2001; Zorita and Laine, 2000; Lehmann et al., 2002). In particular, the North Atlantic Oscillation (NAO), blocking and, on longer time scales, circulation patterns related to the Atlantic Multidecadal Oscillation (AMO) play important roles for the climate of the Baltic Sea region. ~~The AMO consists of an unforced component which is the result of atmosphere-ocean interactions (e.g. Wills et al., 2018) and a forced component such as volcanic eruptions (Mann et al., 2021; Mann et al., 2020).~~ The AMO consists of an unforced component which is the results of atmosphere-ocean interactions (e.g. Wills et al., 2018) and a forced component. It has been shown that external forcing such as solar activity, ozone, and volcanic and anthropogenic aerosols are also important drivers altering variance and phase of the AMO (Mann et al., 2020; Mann et al., 2021; Watanabe and Tatebe, 2019). However, the relative importance of its forced and unforced components is still debated (Mann et al., 2021).

The NAO is the dominant mode of near-surface pressure variability over the North Atlantic and its influence is strongest in winter (Hurrell et al., 2003), when it accounts for almost one-third of the sea level pressure (SLP) variance (e.g. Kauker and Meier, 2003). During the positive (negative) phase of the NAO the Icelandic Low and Azores High pressure systems are stronger (weaker), leading to a stronger (weaker) than normal westerly flow (Hurrell, 1995). Positive NAO phases are associated with mild temperatures and increased precipitation and storminess whereas negative NAO phases are characterized by warm summers, cold winters, and less precipitation (Hurrell et al., 2003). Increasing winter temperatures in the Baltic Sea have also been linked to an observed

northward shift in the storm tracks (BACC II Author Team, 2015). There is a large interannual to interdecadal variability in the NAO, reflecting interactions with and changes in surface properties, including sea surface temperature (SST) and sea ice cover. This makes it difficult to detect a possible long-term trend in the NAO.

Atmospheric blocking occurs when persistent high pressure systems interrupt the normally westerly flow over the middle and high latitudes, e.g. the North Atlantic. By redirecting the pathways of midlatitude cyclones, blockings lead to negative precipitation anomalies in the region of the blocking anticyclone and positive anomalies in the surrounding areas (Sousa et al., 2017). In this way, blockings can also be associated with extreme events such as heavy precipitation (Lenggenhager and Martius, 2019) or drought (Schubert et al., 2014).

The AMO describes fluctuations in North Atlantic sea surface temperature (SST) with a period of 50-90 years (Knight et al., 2006). Thus only a few distinct AMO phases have been observed in the 150-year instrumental record. However, a recent model study suggests that variations in the AMO may influence atmospheric circulation that leads to additional precipitation over the Baltic Sea region (Börgel et al., 2018). A recent model study suggested that variations in the AMO may influence atmospheric circulation that leads to additional precipitation during positive AMO phases over the Baltic Sea region (Börgel et al., 2018). However, the ensemble mean response of the CMIP6 control runs showed an increase in precipitation during negative AMO phases. Further, it was found that the AMO altered the zonal position of the NAO and affected the regional imprint of the NAO for the Baltic Sea region (Börgel et al., 2020).

1.4.2 A unique brackish water basin

The Baltic Sea is a unique brackish water basin in the World Ocean which has a salinity less than 24.7 g kg^{-1} in all areas (Leppäranta and Myrberg, 2009; Voipio, 1981; Magaard and Rheinheimer, 1974; Feistel et al., 2008; Omstedt et al., 2014). The sea is very shallow (with a mean depth of only 54 m), and can be characterized as a number of sub-basins (Fig. 2). The Baltic Sea has the only connection to the North Sea through the Danish straits (Fig. 2). The exchange of water between the Baltic Sea and North Sea through the narrow straits is quite limited. The Baltic Sea has a positive fresh-water balance with an average salinity of about 7.4 g kg^{-1} (e.g. Meier and Kauker, 2003) – this being only one-fifth the salinity of the World Ocean, thus water masses are brackish. The Baltic Sea is located between mild maritime and continental sub-arctic climate zones and partly ice-covered in every winter. However, it is completely frozen over only during extremely cold winters. The highly variable coastal geomorphology and the extended archipelago areas make the Baltic Sea unique (see Section 5).

The World Ocean has only four large brackish water basins (Leppäranta and Myrberg, 2009). These are from the largest to the smallest the Black Sea (Ivanov and Belokopytov, 2013) located between Europe and Asia Minor, the Baltic Sea, the Gulf of Ob in the Kara Sea (Volkov et al., 2002) and the Chesapeake Bay (Kjerfve, 1988), on the east coast of the United States of America. All these sea areas developed into brackish water basins during the Holocene. During the most recent (Weichselian) glaciation period the Black Sea was a freshwater lake, the Baltic Sea and the Gulf of Ob were under the Eurasian ice sheet, and the Chesapeake Bay was a river valley (Leppäranta and Myrberg, 2009). The mean depth of the Black Sea is 1200 m, and due to the strong salinity stratification and

extremely slow deep water renewal the water masses below 200 m are anoxic. The Sea of Azov in the north-eastern part of the Black Sea is often frozen during the winter. The Gulf of Ob is the long (800 km), narrow estuary of the River Ob in the Kara Sea in the Russian Arctic, and ice-covered in winter. Finally, Chesapeake Bay is a small, very shallow basin and a drowned river valley or ria, in the humid subtropical climate zone, with hot summers and ice formation in river mouths in some winters.

Table 42 gives basic information of the brackish water seas and other basins comparable with the Baltic Sea. Most similar to the brackish water seas is Hudson Bay (Roff and Legendre, 1986). It is an oceanic, semi-enclosed basin with a positive fresh-water balance, and a salinity of about 30 g kg⁻¹. In contrast, small Mediterranean seas with a negative fresh-water balance and salinities above 40 g kg⁻¹ are found in the tropical zone; e.g. the Red Sea and Persian Gulf. The largest lakes are comparable in size to the Baltic Sea, and the Caspian Sea is even larger in volume.

The Baltic Sea basin is a very old geomorphological depression. Prior to the Weichselian glaciation this basin contained the Eem Sea, which extended from the North Sea to the Barents Sea, making Fennoscandia an island. At the end of the Weichselian glaciation, 13,500 years ago, the Baltic Ice Lake was formed by glacier meltwater. During the Holocene fresh and brackish phases followed dictated by the balance of glacier retreats and progressions, land uplift and eustatic changes of the global sea level (Tikkanen and Oksanen, 2002). The present brackish phase commenced 7000 years ago, and since about 2000 years ago the salinity has been close to the present level. Postglacial land uplift has slowly changed the Baltic Sea landscape, making it possible to observe how land rises from the sea and how terrestrial life gradually takes over. People living in the region have adapted to this slow long-term change.

1.4.3 The Baltic Sea - a specific European sea

The basic features of the European seas reveal key differences, in areal extent, depth profile, salinity level, fresh water budget, climate, and tidal motions (Table 53). The Baltic Sea and the North Sea are shallow, with a mean depth of less than 100 m; the Baltic can be described as a “coastal sea”, with a mean depth of only 54 m. The Black Sea and the Mediterranean Sea are much deeper, with mean depths of approximately 1200 m and 1500 m, respectively, whereas the North-East Atlantic reaches the full oceanic depth of ca. 4 km, fringed by much shallower continental shelf areas, at about 400 m. These depth differences influence, among other things, the mixing of the water column, variability in temperature, and distribution of benthic ecosystems (Myrberg et al., 2019).

Among the European sSeas, the Baltic Sea physics stands out in terms of its small tidal amplitudes, low salinity, strong stratification and anoxic conditions. Additionally, frequent and spatially extensive upwelling and regular seasonal ice cover are typical of the Baltic Sea (Leppäranta and Myrberg, 2009). To summarize:

- The Baltic Sea is permanently stratified due to a large salinity (density) difference between the fresh upper layer and the more saline bottom layer. This limits ventilation, leading to oxygen deficiency in the bottom layer. For instance in autumn 2016, some 70 000 km² of the seabed experienced permanent hypoxia. Irregular Major Baltic Inflows (MBIs; Matthäus and Franck, 1992; Mohrholz, 2018) are the main

mechanism transporting oxygen-rich waters from the North Sea to Baltic Sea deeps. The associated salt transport in turn intensifies vertical stratification and eventually enlarges hypoxic area (Conley et al., 2002).

- In the small, semi-enclosed Baltic Sea, almost any winds are likely to blow parallel to some section of the coast and thus cause coastal upwelling. At the Swedish south-~~eastern~~~~western~~ coast, upwelling occurs 25-40 % of time (Lehmann et al., 2012). At times, about one third of the entire Baltic Sea may be under the influence of upwelling.
- Among European seas, ice is a unique feature of the Baltic Sea that strongly limits air-sea interaction and modifies the Baltic Sea ecosystem in many ways.

The salinity in the Baltic Sea is not only an oceanographic variable as in other more ventilated seas, but also integrates the complete water and energy cycles, with their specific Baltic Sea features. Baltic Sea salinity, and especially its low ~~mean~~~~basie~~ value and the large variations, is also an elementary factor controlling the marine ecosystem. The salinity dynamics is governed by several factors: net precipitation, river runoff, surface outflow of brackish Baltic Sea water and the compensating deep inflow of higher salinity water from the Kattegat. The latter is strongly controlled by the prevailing atmospheric forcing conditions. Due to freshwater supply from the Baltic Sea catchment area and due to the limited water exchange with the World Ocean, surface salinity varies from $> 20 \text{ g kg}^{-1}$ in Kattegat to $< 2 \text{ g kg}^{-1}$ in the Bothnian Bay and is close to zero at the mouth of the River Neva, in the easternmost end of the Gulf of Finland. In the vertical direction, the dynamics of the Baltic Sea is characterized by a permanent, two-layer system because of a pronounced, perennial vertical gradient in salinity. In summer, a shallow thermocline is also formed, complicating the vertical structure.

1.5 Global climate change

In the following, a brief overview is given of the latest global climate assessments, based on the IPCC Fifth (~~AR5~~; ~~IPCC, 2014b~~) and Sixth Assessment Report (~~AR6~~; IPCC, 2021~~44b~~), ~~including~~~~and~~ results so far available from the current Coupled Model Intercomparison Project (CMIP) phase 6 (Eyring et al., 2016). The focus is on large-scale changes in climate that are of particular relevance for the Baltic Sea region (mainly ~~in the~~ North Atlantic ~~and~~; Arctic ~~regions~~). Furthermore, whenever feasible, changes are described in terms of pattern scaling which relies on the fact that for many quantities the geographical change patterns are sufficiently consistent across models and scenarios to emerge from the background noise (IPCC, 2014a). Hence, changes in e.g. local temperatures can be scaled to changes per °C of global mean temperature change relative to ~~a defined historical period, e.g. 1986-2005~~ ~~for EURO-CORDEX~~ (Christensen et al., 2019).

Our future climate change assessment relies on the concentration driven scenarios RCP2.6, RCP4.5 and RCP8.5 from the CMIP5 suite (RCP = Representative Concentration Pathway), corresponding to changes in radiative forcing for the 21st century. Hence, policy targeted goals inspired by the United Nations Framework Convention on Climate Change (UNFCCC; United Nations Climate Change, 2015) to limit global mean warming below 2.0 or 1.5°C compared to preindustrial level, i.e. prior to the 20th century (the Paris Agreement, ~~PA~~), are not considered in many scenario simulations but referred to studies within the ~~EURO~~~~uro~~-CORDEX framework (Kjellström et al.,

2018; Teichmann et al., 2018; Jacob et al., 2018) and for a broader region. In order to achieve the goal of a significant reduction of the risks and impacts of climate change, the Paris Agreement commits the participating countries to aim “to reach global peaking of greenhouse gas emissions as soon as possible” and “to undertake rapid reductions thereafter in accordance with best available science, so as to achieve a balance between anthropogenic emissions by sources and removals by sinks of greenhouse gases in the second half of this century”. Furthermore, the countries “should take action to conserve and enhance, as appropriate, sinks and reservoirs of greenhouse gases [...], including forests”.

RCP8.5 is a totally unmitigated scenario and assumes a radiative forcing of +8.53 W m⁻² in year 2100, as compared to the preindustrial period. Assumptions for RCP8.5 are described in Riahi et al. (2011). RCP8.5 has been criticized because it assumes continued use of coal for energy production translating into too high greenhouse gas emissions (Hausfather and Peters, 2020). Moderate mitigation actions are reflected by RCP4.5 (Thomson et al., 2011), and RCP2.6 was developed for effective mitigation scenarios aiming at limiting global mean warming to ~+2°C (van Vuuren et al., 2011). With respect to global development, RCP2.6 and RCP8.5 might be unrealistic (Hausfather and Peters, 2020). However, both scenarios can be used as envelopes of plausible pathways of future greenhouse gas emissions- or as representing stronger feedbacks in the climate system than assumed in the default derivation of the scenarios.

Confidence levels expressing evidence and agreement are provided following the definitions of the IPCC (see Method Section 2.3).

1.5.1 Atmosphere

1.5.1.1 Surface air temperature

For the three considered scenarios, the IPCC AR5 (IPCC, 2014a; 2014b; Collins et al., 2013) reported a likely increase in global mean air temperature for the period 2081-2100 relative to 1986-2005 in the likely range (5th to 95th percentile of CMIP5 models) between 0.3 to 1.7°C (RCP2.6), 1.1 to 2.6°C (RCP4.5), and 2.6 to 4.8°C (RCP8.5). The corresponding mean changes are 1.0°C (RCP2.6), 1.8°C (RCP4.5) and 3.7°C (RCP8.5; IPCC, 2014b).

The large-scale geographical patterns of change remain stable among CMIP5 models and are consistent with the results of the IPCC AR4 (IPCC, 2007). The dominant feature is a strong warming of the Arctic north of 67.5 °N that exceeds global mean warming by a factor of 2.2 to 2.4 as a multi-model mean value. The Arctic warming is strongest for the winter season, when sea-ice retreat and reduced snow cover provide positive feedbacks (Arctic amplification), and weakest in summer, when melting sea ice consumes latent heat and the ice-free ocean absorbs heat (IPCC, 2014b). Besides these thermodynamic processes, the lateral transport of latent heat into the Arctic increases under global warming. Weakest warming is found over the Southern Ocean and in the North Atlantic south of Greenland with minimum values per degree global warming of about 0.25°C °C⁻¹ (Fig. 12.10 in IPCC, 2014b). This is partly due to a deeper ocean mixed layer that promotes vigorous oceanic heat uptake in these regions compared to others. Generally, land masses warm at a rate 1.4 to 1.7 times more than open ocean regions,

Formatiert: Englisch (Großbritannien)

leading to a pronounced land-sea pattern in the temperature anomaly ~~and indicating to a lower effective heat capacity of continents compared to the ocean.~~ The difference in heat capacity plays some role in the land-sea warming contrast during the transient phase of warming, but it is not its main reason. As first shown by Joshi et al. (2008), the overall land-sea contrast is to a large extent caused by the dryness of land surfaces, which makes it impossible for evaporation to increase as much in a warmer climate as it does over the oceans. The mechanistic pathway involves also atmospheric dynamics. For further details, the reader is referred to, for example, Byrne and O’Gorman (2018).

1.5.1.2 Precipitation

Projected global precipitation changes scale nearly linear with global mean temperature changes and range from +0.05 mm d⁻¹ or ~2% (RCP2.6) to 0.15 mm d⁻¹ or ~5% (RCP8.5; IPCC, 2014a). As a result of an accelerated global water cycle, the contrast between dry and wet regions in annual mean precipitation increases. Likewise, there is high confidence that the contrast between wet and dry seasons will become more pronounced (IPCC, 2014a). In the mid to high latitudes, yearly mean precipitation generally increases, with the strongest response over the Arctic, exceeding almost everywhere +12% °C⁻¹. Note that this normalization is by the global mean warming as already mentioned above.

Precipitation changes vary greatly among models. ~~Under RCP8.5,~~ High latitude land masses will likely get more precipitation, due to higher moisture content of the lower atmosphere and an increased moisture transport from the tropics (IPCC, 2014a). In the northern hemisphere the poleward branch of the Hadley Cell will expand further north, causing a northward expansion of the subtropical dry zone and reducing precipitation in affected regions. Further dynamical changes probably include a poleward shift of mid-latitude storm tracks (Seager et al., 2010; Scheff and Frierson, 2012) which is, however, of low confidence, especially for the North Atlantic region (IPCC, 2014a).

1.5.2 Cryosphere

The IPCC AR5 postulates a reduction of average February Arctic sea-ice extent ranging from 8% for RCP2.6 to 34% for RCP8.5. For the monthly mean summer minimum in September, reductions range from 43% for RCP2.6 to 94% for RCP8.5. These values are given medium confidence, because of biases in the simulation of present day trends and a large spread across models. For September, ice-free conditions are reached before 2090 in 90% of all CMIP5 models.

The permafrost area is projected to decrease in a likely range from 24 ± 16% for RCP2.6 to 69 ± 20% for RCP8.5.

Arctic autumn and spring snow cover are projected to decrease by 5–10%, under RCP2.6, and 20–35% under RCP8.5 (high confidence). In high mountain areas, projected decreases in mean winter snow depth are in a likely range of 10–40% for RCP2.6 and 50–90% for RCP8.5. The likely range of projected inland glacier mass reduction^s (ice sheets excluded) between 2015 and 2100 varies from 18 ± 7% for RCP2.6 to 36 ± 11% for RCP8.5. Regions with mostly smaller glaciers (e.g. ~~c~~Central Europe, Scandinavia) are projected to lose over 80% of their

current ice mass by 2100 under RCP8.5 (medium confidence), with many glaciers disappearing regardless of future emissions (very high confidence).

1.5.3 Ocean

1.5.3.1 Sea level

For 2081-2100, global mean sea level (GMSL) is projected to rise between 0.40 m under RCP2.6 (likely range 0.26-0.55 m) and 0.63 m under RCP8.5 (likely range 0.45-0.82 m) relative to 1986-2005 (IPCC, 2014^{ba}; their Chapter 13, Table 13.5). In all scenarios, thermal expansion gives the largest contribution to GMSL rise, accounting for about 30 to 55% of the projections. Glaciers are the next largest contributor, accounting for about 15-35%. By 2100, the Greenland Ice Sheet's projected contribution to GMSL rise is 0.07 m (likely range 0.04–0.12 m) under RCP2.6, and 0.15 m (likely range 0.08–0.27 m) under RCP8.5. The Antarctic Ice Sheet is projected to contribute 0.04 m (likely range 0.01–0.11 m) under RCP2.6, and 0.12 m (likely range 0.03–0.28 m) under RCP8.5. ~~The incomplete knowledge about~~~~Uncertainties concerning the~~ melting of ice sheets ~~is~~are, however, intensively discussed (Bamber et al., 2019).

Based on the same suite of model projections from CMIP5, the IPCC Special Report on the Ocean and Cryosphere in a Changing Climate (IPCC, 2019a) has updated these numbers by including new estimates of the contribution from Antarctica, for which new ice-sheet modelling results were available (Oppenheimer et al., 2019). While the differences in projected changes until 2100 are small for RCP2.6, projected changes for RCP8.5 increased by about 10 cm compared to AR5 (see Section 3.3.5.4).

It is important to note that sea level rise will continue in all RCP scenarios. This is made clear by a quote from IPCC's AR6 (IPCC, 2021): "In the longer term, sea level is committed to rise for centuries to millennia due to continuing deep ocean warming and ice sheet melt, and will remain elevated for thousands of years (high confidence). Over the next 2000 years, global mean sea level will rise by about 2 to 3 m if warming is limited to 1.5°C, 2 to 6 m if limited to 2°C and 19 to 22 m with 5°C of warming, and it will continue to rise over subsequent millennia (low confidence)."

1.5.3.2 Water temperature and salinity

By the end of the century, the projected global ocean warming ranges from about 1°C (RCP2.6) to more than 3°C (RCP8.5) at the surface and from 0.5°C (RCP2.6) to 1.5°C (RCP8.5) at a depth of 1 km. The subtropical waters of the Southern Ocean and the North Atlantic are projected to become saltier, whereas almost all other regions become fresher, in particular the northern North Atlantic (IPCC, 2014a). The freshening at high latitudes in the North Atlantic and Arctic basin is consistent with a weaker Atlantic Meridional Overturning Circulation (AMOC), and a decline in the volume of sea ice, as well as with the intensified water cycles (IPCC, 2019a).

By the end of the century, the annual mean stratification of the top 200 m (averaged between 60°S–60°N, relative to 1986–2005) is projected to increase in the very likely range of 1–9% for RCP2.6 and 12–30% for RCP8.5 (IPCC, 2019a).

1.5.3.3 Atlantic Meridional Overturning Circulation

Based on the CMIP5 models, the AMOC is estimated to be reduced by 11% (1 to 24%) under RCP2.6 and 34% (12 to 54%) under RCP8.5. There is low confidence in the projected evolution of the AMOC beyond the 21st century (IPCC, 2014a).

1.5.4 Marine biosphere

By 2081-2100, global net primary productivity relative to 2006-2015 will very likely decline by 4–11% for RCP8.5, due to the combined effects of warming and changes in stratification, light, nutrients and predation, with regional variations between low and high latitudes (IPCC, 2019a).

Globally, and relative to 2006-2015, the oxygen content of the ocean by 2081-2100 is very likely to decline by 1.6–2.0% for the RCP2.6 scenario, or by 3.2–3.7% for the RCP8.5 scenario (IPCC, 2019a). While warming is the primary driver of deoxygenation in the open ocean, eutrophication is projected to increase in estuaries due to human activities and due to intensified precipitation, which increase riverine nitrogen loads under both RCP2.6 and RCP8.5 scenarios, both by mid-century (2031–2060) and later (2071–2100; Sinha et al., 2017). Moreover, stronger stratification in estuaries due to warming is expected to increase the risk of hypoxia by reducing vertical mixing (IPCC, 2019a; Hallett et al., 2018; Warwick et al., 2018; Du et al., 2018).

1.5.5 Coupled Model Intercomparison Project 6

In this study, CMIP5 instead of the successor CMIP6 scenario simulation results have been used for the regionalization of climate change because dynamical downscaling experiments based on CMIP6 projections are still lacking, while the IPCC Sixth Assessment Report (AR6; IPCC, 2021) relies on the updated generation of ESMs from CMIP6 (Eyring et al., 2016). In CMIP6, RCP scenarios have been replaced by SSP (Shared Socioeconomic Pathway) scenarios. Future climate change assessments like the coming IPCC sixth Assessment Report AR6 (due 2021/2022), will rely on a new generation of Earth System Models (ESMs), developed during CMIP6, which offerings a wider range of scenarios than during CMIP5 (IPCC AR6 Technical Summary, p. 21-23). In particular, scenarios aiming to limit global warming to 1.5°C and 2.0°C and overshoot scenarios including negative emissions in the second part of the century are will be now available. The lowest of these scenarios, SSP1-1.9, which was designed to limit the global warming to 1.5°C above the preindustrial, has lower radiative forcing than RCP2.6. For the other SSP scenarios, the effective radiative forcing tends to be slightly higher than for the nominally corresponding RCP scenarios (e.g., SSP5-8.5 is slightly higher than RCP8.5).

A key result in IPCC AR6 (IPCC, 2021) was the narrowed uncertainty range of the estimated response to an instantaneous doubling of CO₂ (equilibrium climate sensitivity, ECS), as a consequence of the improved scientific understanding and accumulation of new data. Largely based on the review by Sherwood et al. (2020), the IPCC AR6 concluded a likely range of 2.5–4°C for the ECS.

A subset of current CMIP6 models have been shown to be more sensitive to greenhouse gases than previous generations of CMIP models. Thus, the ECS estimated response to an instantaneous doubling of CO₂ (equilibrium

Formatiert: Englisch (Großbritannien)

539 ~~climate sensitivity, ECS~~ is higher in CMIP6 models (1.8 – 5.6°C) than in CMIP5 models (1.5 – 4.5°C) and their
540 predecessors (Meehl et al., 2020). Indeed, the first transient simulations with the CMIP6 EC-Earth ESM found
541 stronger warming than with earlier versions, with about half of the increase attributed to differences between
542 CMIP5 and CMIP6 greenhouse gas forcing (Wyser et al., 2020).

543
544 However, it turns out that models with the highest projected warmings fail to capture past warming trends well,
545 and therefore recent studies argue that those models should not be used for climate assessments and policy
546 decisions (Forster et al., 2020; Nijse et al., 2019; Tokarska et al., 2020; Brunner et al., 2020). Furthermore,
547 systematic errors in many CMIP5 and CMIP6 models prevent the simulation of the observed 1951-2014 summer
548 warming trend in ~~w~~Western Europe, and ~~that~~ neither higher resolution nor better representation of the sea surface
549 is likely to improve this (Boé et al., 2020_b).

550
551 Despite the differences in model sensitivity between CMIP5 and CMIP6, CMIP6 results generally confirmed the
552 findings of CMIP5 but added value to the uncertainty assessment because of a larger number of participating
553 ESMs. Hence, from the dynamical downscaling of CMIP6 scenario simulations major changes on our conclusions
554 for the Baltic Sea region cannot be expected. To illustrate differences between CMIP5 and CMIP6, we compare,
555 with the help of the IPCC AR6 interactive atlas (Iturbide et al., 2021; Gutiérrez et al., 2021), applying the land
556 only mask, CMIP5 and CMIP6 results in two regions, Northern Europe (NEU) and Western and Central Europe
557 (WCE), which together comprise a large part of the Baltic Sea region. We assume that this analysis is relevant
558 although the two selected regions also include other adjacent areas.

559
560 The comparison of the two regions for all CMIP5 and all CMIP6 models under RCP8.5 and SSP5-8.5, respectively
561 (about 30 for both experiments), shows for seasonal and annual mean temperature increases at the same global
562 warming levels similar results (Table 6). Notably, the CMIP6 models appear to show a somewhat larger summer
563 warming signal than in CMIP5 and a lower increase in winter temperatures, in particular for higher warming levels.
564 Note that these comparisons are made for warming levels which means that differences between scenarios, i.e.
565 RCP8.5 and SSP5-8.5, are removed.

566
567 Also precipitation changes are similar. Differences in annual mean precipitation changes between CMIP5 and
568 CMIP6 are less than 3% (Table 6). Precipitation increase in northern Europe (NEU) is somewhat smaller in CMIP6
569 than in CMIP5. Also in central Europe (WCE) winter precipitation increase is somewhat smaller, in CMIP6. In
570 summer, WCE shows even stronger reduction in precipitation in CMIP6 compared to CMIP5 for all warming
571 levels apart from +1.5°C.

572
573 Also for various time slices, the differences between temperature and precipitation changes between CMIP5 and
574 CMIP6 models under RCP8.5 and SSP5-8.5, respectively, are relatively small (Table 7). The identified differences
575 can be attributed to differences in ESMs and emission scenarios. Largest differences between CMIP5 and CMIP6
576 in temperature changes between 1850-1900 and 2081-2100 are found in WCE during summer with CMIP6 models
577 showing more pronounced warming by 0.9 °C than the CMIP5 models. The larger climate change signals in CMIP6

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

578 compared to CMIP5 results are confirmed by a subset of ESMs from CMIP5 and CMIP6 analyzed by Coppola et
579 al. (2021). During all seasons, the precipitation increases in NEU are somewhat smaller in CMIP6 than in CMIP5,
580 in particular during summer, and the precipitation decrease in WCE during summer is larger (Table 7).

Formatiert: Englisch (USA)

581
582 Table 8 also includes scenario simulation results from EURO-CORDEX Regional Climate Models (RCMs), i.e.
583 47 simulations. Note that the changes are calculated relative to a different reference period (1986-2005) compared
584 to Tables 6 and 7 (1850-1900). Hence, the values of temperature and precipitation changes differ from the other
585 two tables. For EURO-CORDEX, the temperature changes are smaller than in CMIP5 and for precipitation there
586 is generally a smaller increase during the winter but more positive values during the summer (larger increase in
587 NEU or smaller decrease in WCE). That RCMs may change the results compared to the underlying GCMs also on
588 large regional scales is well-known and has previously been shown for different subsets of EURO-CORDEX
589 RCMs (e.g. Sørland et al., 2018; Coppola et al, 2021).

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

590
591 In the following, we list selected examples of publications comparing CMIP5 and CMIP6 results, with relevance
592 for the Baltic Sea region. Nie et al. (2019) compared the historical forcings in CMIP5 and CMIP6 simulations. As
593 more CMIP6 models include aerosol-cloud interactions than in CMIP5, the effect of the stronger aerosol forcing
594 results in an approximately 10% strengthening of the AMOC in the multimodel mean during 1850-1985, a larger
595 change than was seen in CMIP5 models before (Menary et al., 2020). Moreno-Chamarro et al. (2021) concluded
596 that the horizontal resolution of the ESMs is more important for the calculation of winter precipitation changes
597 over northern Europe than the differences between CMIP5 and CMIP6 models. Furthermore, there is robust
598 evidence that CMIP6 models simulate blocking frequency and persistence better than CMIP5 models, presumably
599 because of the higher resolution in CMIP6 models (Schiemann et al., 2020). Li et al. (2021) compared extreme
600 precipitation and drought changes between CMIP5 and CMIP6 models and found differing results in various
601 regions. According to Seneviratne and Hauser (2020) climate change results in extreme temperature and heavy
602 precipitation reveal close similarity between CMIP5 and CMIP6 ensembles in the regional climate sensitivity of
603 the projected multimodel mean change as a function of global warming although global (transient and equilibrium)
604 climate sensitivity in the two multimodel ensembles differ. Statistical-empirical downscaling was used by
605 Kreienkamp et al. (2020) to investigate temperature and precipitation changes for Germany using selected ESMs
606 of CMIP5 and CMIP6. The SSTs around the North Atlantic Subpolar Gyre are better simulated than their CMIP5
607 predecessors, largely due to the more accurate modelling of the influence of natural climate forcing factors
608 (Borchert et al., 2020). Finally, CMIP6 ESMs generally project greater ocean warming, acidification,
609 deoxygenation, and nitrate reductions but lesser primary production declines than those from CMIP5 under
610 comparable radiative forcing (Kwiatkowski et al., 2020). However, the inter-model spread in net primary
611 production changes in CMIP6 projections increase compared to CMIP5.

2 Methods

2.1 Assessment of literature

33 variables representing the components of the Earth system (atmosphere, land, terrestrial biosphere, cryosphere, ocean and sediment, marine biosphere) of the Baltic Sea region were selected (Table 24). ~~Scientific peer-reviewed publications and reports of scientific institutes since 2013 on past, present and future climate changes in these variables were assessed and with respect to past, present and future climate changes corresponding scientific peer-reviewed publications and reports of scientific institutes since 2013 were assessed by 478 experts (see Table 1 see the section about for the~~ author contributions). The year 2013 was chosen as a starting point ~~for the oldest publications~~ because earlier material was ~~already~~ included in the last assessment by the BACC II Author Team (2015). Information about climate change available in the BEARs (Section 12.1) was summarized and cross-references can be found in Table 24.

For the selected 33 variables and even ~~in~~ more general ~~terms~~, knowledge gaps (Section 6) and key messages (Section 7) as well as overall conclusions (Section 8) were formulated. Key messages, new compared to the results of the BACC II Author Team (2015), ~~awere~~ marked. The identified changes of the selected variables of the Earth system and their ~~estimated~~ uncertainties, following the definitions of the IPCC reports as outlined in Section 2.3, are summarized in Table 1540. The attribution of a changing variable to climate change, here the deterministic response to changes in external anthropogenic forcing such as greenhouse gas and aerosol emissions, is illustrated by Figure 3534. This study does not claim to be complete, neither with regard to the ~~importance of the limited~~ selection of variables, ~~characterising for~~ the Earth system, nor with regard to the discussed and assessed publications.

The assessment was done without influence from any political, economic or ideological group or party. The results of the BEARs including this summary about climate change impacts in the Baltic Sea region were used by the joint HELCOM-Baltic Earth Expert Network of Climate Change (EN CLIME) for the compilation of ~~thea~~ Climate Change Fact Sheet for the Baltic Sea region ~~(see Section 1.2)~~.

For further details about the assessment methods, the reader is referred to the BACC Author Team (2008) and the BACC II Author Team (2015).

2.2 Proxy data, instrumental measurements and climate model data

In addition to selected figures that are reproduced from the literature, for the assessment previously published datasets were analyzed and discussed.

2.2.1 Past climate

For the Holocene climate evolution, paleo-pollen data with a decadal resolution, reconstructing seasonal temperature and precipitation changes compared to preindustrial climate (Mauri et al., 2015), were analyzed (Fig. 3). More accurate tree-ring data, resolving annual summer mean temperatures, are available for the past

millennium (Luterbacher et al., 2016) and have been discussed here (Fig. 4). For further details, the reader is referred to Section 3.1.2.

2.2.2 Present climate

Historical station data of sea level pressure and ~~sea surface temperature~~ (SST) were used to calculate climate indices such as the NAO (sea level pressure differences, Fig. 5) and the AMO (SST anomalies, Fig. 6), describing decadal to multidecadal variability of the large-scale atmospheric circulation. Furthermore, selected records of variables such as air temperature (Fig. 8), river runoff (Fig. 10), land nutrient inputs (Fig. 11, Table 105), glacier masses (Fig. 12, Table 116), maximum sea ice extent (Fig. 14), ice thickness data (Figs. 15 and 16), length of the ice season (Fig. 17), sea level (Fig. 24) and gridded data sets of air temperature, e.g. the land-based CRUTEM4 data (Jones et al., 2012; Fig. 7, Table 95), and of precipitation, e.g. Copernicus data (Fig. 9), were analyzed.

For the Baltic Sea, intensive environmental monitoring started more than 100 years ago. Since 1898 an agreement between various Baltic Sea countries on simultaneous investigations on a regular basis at a few selected deep stations was signed and 1902 the International Council of the Exploration of the Sea (ICES) started its work. Examples from the national monitoring programs for water temperature (Figs. 18, 19, 20) and salinity (Figs. 21 and 22) are shown, illustrating climate variability and climate change of the Baltic Sea.

In addition, some institutes such as the Swedish Meteorological and Hydrological Institute (SMHI) provide environmental/climate indices, e.g. averaged sea level station data corrected for land uplift (Fig. 23) and hypoxic and anoxic areas (Fig. 25).

Since 1979 satellite data have become available, complementing traditional Earth observing systems and having the advantage of spatially high resolution (e.g. Karlsson and Devasthale, 2018).

Atmospheric reanalysis products, i.e. the combination of model data and observations (e.g. NCEP/NCAR, ERA40, ERA-Interim, ERA5, UERRA), were important for calculating water and energy budgets of the Baltic Sea region (BACC Author Team, 2008; BACC II Author Team, 2015). More recently, also ocean reanalysis products have been developed (e.g. Liu et al., 2017; Axell et al., 2019; Liu et al., 2019) and were, for instance, used for the evaluation of models (e.g. Placke et al., 2018).

Furthermore, various gridded datasets for North Sea SSTs exist and were compared (Fig. 343).

All data sets presented here are publicly online available. For further details on various datasets, the reader is referred to Rutgersson et al. (2021).

2.2.3 Future climate

For the BEARs, regionalisations ~~with the help of~~ ~~dynamically downscaling~~ of Global Climate Models (GCMs) or ESMs from CMIP3 and CMIP5 analyzed by ~~IPCC~~ (2014b) ~~and~~ ~~IPCC~~ (2019b) are assessed. The scenario

simulations of CMIP5 are driven by greenhouse gas concentration scenarios, the Representative Concentration Pathways, RCP2.6, 4.5 and 8.5 (see Section 1.5).

Uncoupled atmospheric regional climate simulations for the 21st century from the EURO-CORDEX framework, calculated with several ~~Regional Climate Models (RCMs)~~ and global ESMs were analyzed by Christensen et al. (2021) and conclusions are summarized here. The choice of working with regional climate model projections, downscaling a limited subset of all available CMIP5 GCMs, implies that the resulting ensemble may not represent all available data properly. Previous studies of parts of the 72-member EURO-CORDEX RCP8.5 ensemble (a sparsely filled GCM-RCM matrix with in total 11 RCMs downscaling 12 GCM projections) assessed here, and presented in more detail by Christensen et al. (2021), illustrated this hypothesis.

By investigating 18 of these RCM simulations (8 RCMs downscaling 9 GCMs), Kjellström et al. (2018) found that the 9-member GCM ensemble showed lower temperature response for northern and eastern Europe compared to the entire CMIP5 ensemble. In addition, it was found that the RCMs can – to some degree – alter the results of the driving GCMs (as also discussed by Sørland et al., 2018). In a more recent study, Coppola et al. (2021) investigated a 55-member ensemble with the same 11 RCMs downscaling the same 12 GCMs as assessed by Christensen et al. (2021). They compared the 55-member ensemble to the driving 12 GCMs and concluded that the RCMs modify the results. In their analysis, Coppola et al. (2021) also considered a set of 12 CMIP6 GCMs finding that these show a stronger warming signal than the 12 CMIP5 GCMs. This was related to the higher equilibrium climate sensitivity in several global models of the new generation.

Furthermore, coupled atmosphere – sea ice – ocean simulations for the Baltic Sea and North Sea regions with one, so-called Regional Climate System Model (RCSM, Dieterich et al., 2013; Bülow et al., 2014; Dieterich et al., 2019; Wang et al., 2015; Gröger et al., 2015; Gröger et al., 2019; Gröger et al., 2021b) driven by eight ESMs and three greenhouse gas concentration scenarios, i.e. RCP2.6, 4.5 and 8.5, were compared with atmosphere-only RCM simulations by (Christensen et al., 2021). In this study, we present figures of these internally consistent results from the coupled ~~atmosphere – sea ice – ocean~~atmosphere-ice-ocean scenario simulations, e.g. for air temperature and precipitation (Fig. 26, Tables 127 and 138), and for sea surface temperature (Fig. 3029, Table 149). The state-of-the-art of coupled modeling is discussed by Gröger et al. (2021a). For further details about the comparison between coupled and uncoupled scenario simulations, the reader is referred to Christensen et al. (2021).

Novel compared to the assessment by the BACC II Author Team (2015) are high-resolution projections of glacier masses including Scandinavian glaciers (Hock et al., 2019). These results are reproduced in Figure 28. In Figure 27 results are reproduced.

Oceanographic regional climate model projections for the Baltic Sea until 2100 driven by the atmospheric surface fields of the above mentioned RCSM by Dieterich et al. (2019) have been developed and analyzed by Saraiva et al. (2019a; 2019b) and Meier et al. (2021a; 2021b). In Meier et al. (2021b), global sea level rise was also considered, a driver of the Baltic Sea climate variability that was previously neglected (cf. Hordoir et al., 2015;

Arneborg, 2016; Meier et al., 2017). Here, we compare the latest scenario simulation results by Saraiva et al. (2019b) with previous projections by Meier et al. (2011a; 2011c) for, e.g. SST, sea surface and bottom salinities, sea level (Fig. 319), bottom oxygen concentration (Fig. 324), and Secchi depth (Fig. 332),

For further details about the latest oceanographic regional climate model projections for the Baltic Sea, the reader is referred to Meier et al. (2021a).

2.3 Uncertainty estimates

Uncertainties of future projections were estimated basically following the IPCC (2014a) guidance note for lead authors of the Fifth Assessment Report on consistent treatment of uncertainties (Mastrandrea et al., 2010). These uncertainty estimates are based upon a matrix of consensus and evidence reported in the literature. For the high confidence of a statement, high levels of both consensus and cases of evidence are required.

In this assessment, we applied a three-level confidence scale measuring low, medium and high confidence of identified climate changes (as defined in Section 2.1) of the selected 33 Earth system variables according to current knowledge (Table 1540). We assessed the sign of a change but not its magnitude. Only detected or projected changes undoubtedly attributed to climate change were considered and synthesized in Figure 354. Changes likely not caused by increasing greenhouse gas concentrations or changing aerosol emissions were not considered. Other external drivers of climate and environmental variability are internal “random” variations of the climate system, land use, eutrophication, contaminants, litter, river regulations, fishery, aquaculture, underwater noise, traffic, spatial planning, etc. (see Reckermann et al., 2021).

Note that our likelihood terminology of an outcome or a result differs from IPCC’s. We do not differentiate between probabilities such as virtually certain 99–100%, very likely 90–100%, likely 66–100%, about as likely as not 33–66%, unlikely 0–33%, very unlikely 0–10%, exceptionally unlikely 0–1%, as the IPCC assessment report does. For many variables the probability information does not exist at the regional scale because large ensembles of regional scenario simulations do not exist.

Key messages of this assessment that are new compared to the previous assessment by the BACC II Author Team (2015) are specially marked (Section 7).

3 Current state of knowledge

3.1 Past climate change

3.1.1 Key messages from previous assessments

Climate variations may be triggered by changes in drivers external to the climate system or may be due to internal processes that reflect the non-linear, chaotic interactions between the different components of the climate system. The analysis of past climate variations is, therefore, useful for two purposes. One is to estimate the

Formatiert: Nicht unterstrichen, Schriftfarbe: Automatisch, Englisch (USA)

Formatiert: Nicht unterstrichen, Schriftfarbe: Automatisch, Englisch (USA)

~~response~~reaction of the climate to changes in the external forcing. The second is to better understand the mechanisms of internal climate variations. Since future climate change will include a mixture of both types of climate variations, the analysis of past climate variations is also necessary for better estimations of future climate change.

The past climate of the Baltic Sea region can be reconstructed from paleo-pollen and dendroclimatological records, with different time resolutions and degrees of accuracy. Paleo-pollen in lake sediments give information about the dominant plant species of a certain period. Combining the environmental ranges of those species in terms of annual maximum temperatures, minimum temperatures and total annual precipitation allows an approximate reconstruction of past climate conditions over the past millennia, with time resolutions of a few decades (e.g. K \ddot{u} hl et al., 2002). Dendroclimatological data of tree ring widths, wood density and sometimes also carbon and oxygen isotopic composition in tree-rings can be dated as exactly as at annual scales.

As described by the BACC II Author Team (2015), the climate history of the Baltic Sea region during the Holocene, i.e. the last 12,000 years, involved very large climate changes, much larger than those during the 20th century. These climate changes were caused by strong changes in external forcing factors, in particular the Earth's orbit. These changes first brought about a warming that terminated the Last Ice Age about 13,000 years BP, then caused a period of very warm temperatures (~ 3°C above preindustrial levels) centered around 6,000 years BP (the Holocene Thermal Maximum), followed by a slow temperature decline towards preindustrial levels. During this long period, other shorter-lived climate events, with durations of a few centuries, caused abrupt drops of temperature. These events, e.g. the Younger Dryas (12,000 years BP) or the 8.2K event (8,200 years BP) were possibly related to abrupt changes in the North Atlantic circulation, when sudden melting of portions of the remnants of the North American ice-sheet disturbed the circulation of the North Atlantic Ocean and disrupted the poleward heat transport.

In general, annual precipitation is believed to have changed with the slow multicentennial-scale changes in temperature. Warmer periods, in particular the Mid-Holocene Optimum, tended to be wetter, although the regional heterogeneity may have been larger than for temperature.

Following the end of the last glaciation, the coastlines of the Baltic Sea underwent changes due to the interplay between the rising global sea-level and the local rebound of the Earth's crust after the disappearance of the Fennoscandian ice sheet. The weight of this ice sheet depressed Fennoscandia by about 500 meters, and its slow, viscous rebound continues until today, with a rate of about 10 mm year⁻¹ at the northern Baltic Sea coast. Due to this interplay, the Baltic Sea experienced periods of open or closed connections to the North Sea that governed the transport of salinity and heat and the nature of the Baltic Sea ecosystems (Groß et al., 2018).

The climate evolution during more recent historical times – the past 1,000 years (Section 3.1.3) - can be reconstructed with better accuracy and higher time resolution due to better dendrochronological data availability. These data show the imprint of the Medieval Warm Period (approx. 900-1350 AD), the Little Ice Age

(approx. 1550-1850 AD) and the Contemporary Warm Period (1850-present) on the Baltic Sea region. These periods were likely caused by [long-term internal climate variability](#) and changes in the external forcing (volcanic eruptions and solar radiation), and during the Contemporary Warm Period also by the increase in anthropogenic greenhouse gases (Hegerl et al., 2003).

In the Baltic Sea, this succession of warm-cold-warm temperatures was accompanied by changes in the deep water oxygen content, with low oxygen conditions in warmer periods (next section). The reasons for these oxygen variations are still not fully understood, but may be relevant for the future, should future warming also cause lower oxygen concentrations.

3.1.2 New paleoclimate reconstructions

Since the publication of the BACC II report (BACC II Author Team, 2015), new reconstructions of the evolution of the European climate over the Holocene and over the past millennium have become available. Like previous reconstructions, the new ones are based on paleo-pollen data and now comprise summer and winter temperatures and summer and winter precipitation. They are available for 1,000-year time segments (Mauri et al., 2015). The reconstructions of the late-spring-summer temperature evolution over the past millennium are based on dendroclimatological data, as the previous reconstructions, but they are now based on wood density measurements, which reflect the slow climate variations better than tree-ring width. These reconstructions are available for [Western Europe](#) from 755 AD onwards (Luterbacher et al., 2016). In this study, only the results for a regular geographical box approximately covering the Baltic Sea region are discussed.

In addition, new regional climate simulations since the publication of the BACC II report better demonstrate the connections between the Baltic Sea and North Atlantic climates on multidecadal timescales ([Schimanke et al., 2012](#); [Schimanke and Meier, 2016](#); Börgel et al., 2018; Börgel et al., 2020; Kniebusch et al., 2019a).

3.1.3 Holocene climate evolution

The picture of the Holocene climate evolution from the BACC II report (BACC II Author Team, 2015) is essentially confirmed, but the regional details are now clearer (Mauri et al., 2015). Between 7,000 and 5,000 years BP, the Baltic Sea region (especially the western Baltic Sea) experienced a period with summer temperatures about 2.5-3°C warmer than in the preindustrial reference period (before the 20th century). However, according to these reconstructions, the [Eastern Baltic Sea region](#) (Finland and the Baltic [states](#) [Republics](#), i.e. [Lithuania](#), [Latvia](#) and [Estonia](#)) did not experience a Mid-Holocene Optimum in summer, when temperatures were similar to the preindustrial period. In contrast, winter temperatures showed a clear Mid-Holocene Optimum over the whole Baltic Sea region, lasting about 8,000-4,000 BP, with winter temperatures roughly 3°C warmer than during the preindustrial period. In the eastern Baltic Sea, winter temperatures were even slightly higher, especially between 6,000-5,000 BP. As a result, annual mean temperatures during the millennia of the Mid-Holocene [Optimum](#), were generally warmer than in the preindustrial period. This warming was limited to the winter in the eastern Baltic Sea, where the amplitude of the annual temperature cycle was clearly lower than in the preindustrial period.

The warm temperatures in the Baltic Sea region during the Mid-Holocene ~~Optimum~~ are not surprising and basically agree with the previous review (BACC II Author Team, 2015). They also agree with evidence from regions ~~f~~urther north, indicating that the Arctic Ocean in summer may have been ice-free during this period (Jakobsson et al., 2010). These findings do not contradict the anthropogenic effect on climate observed during recent decades. During the Mid-Holocene Optimum, the orbital configuration of the Earth was different and favored warmer temperatures at northern high latitudes, especially in summer, as explained later. For the analysis of climate impacts on ecosystems it is relevant that high latitudes were exposed to warm temperatures and reduced sea-ice cover just a few millennia ago. However, at that time temperature changed at a much slower pace – around 2-3°C over several millennia – compared to present and projected rates of about 2°C in just a few decades.

For precipitation, the new reconstructions give a regionally more nuanced view of climate evolution during the Holocene. The BACC II report (BACC II Author Team, 2015) indicated that warmer climates were generally more humid. The new reconstructions (Mauri et al., 2015) modulate this vision and constrain the wetter conditions to the eastern Baltic Sea region, both in summer and winter seasons, with a stronger signal in winter. Precipitation anomalies in the eastern Baltic Sea region were of the order of + 1-2 mm month⁻¹ relative to preindustrial climate. In the western Baltic Sea region, the Mid-Holocene Optimum tended to be slightly drier than the preindustrial reference period both in summer and winter, with precipitation deficits of the order of 1-2 mm month⁻¹.

The main external forcing that drove the millennial climate evolution over the Holocene ~~p~~eriod is the changing orbital configuration of the Earth (the so-called Milanković cycles), as explained in the BACC-II report (BACC II Author Team, 2015), and especially the variation in the time of the year of the perihelion (when Earth is nearest to the sun). The perihelion is now at the beginning of January, but ~10,000 BP it was in July. This changes the seasonal distribution of solar insolation and determines the rate of melting of winter snow and its possible survival into the next winter. The solar insolation at 60°N at the top of the atmosphere during the Holocene, derived from Laskar et al. (2004) is depicted in Figure 3. The shift of the perihelion from summer to winter diminishes summer insolation - and in principle summer temperature - and increases winter insolation during the past few millennia. The long-term evolution of temperatures would, however, not be a linear response to the long-term evolution of the seasonal insolation. For instance, the presence or absence of ice-sheets may influence the timing of the response to increasing insolation during the early Holocene, delaying the Holocene temperature maximum with respect to the annual insolation maximum. In wintertime, the insolation is rather weak, so that its effect may be overwhelmed by other factors, such as changes in the large-scale atmospheric and oceanic heat transports.

For the ~~fifth last~~ IPCC ~~assessment~~ report (IPCC, 2014b), the ~~M~~id-Holocene climate was simulated with 14 global ~~ESMsEarth-System-models~~ within CMIP5 (Schmidt et al., 2011). These models were essentially the same as those used for future climate projections, although in some cases with a few simplifications required by limitations in computer power and by the long timescales involved. These models were driven by known external forcings, including the orbital forcing. The common evaluation of these simulations with reconstructions helps to interpret the reconstructions and sheds light on model limitations. An important aspect in this comparison was that the

874 spatial resolution of global models was relatively coarse, about 2 x 2 degrees longitude x latitude, so that smaller
875 details within the Baltic Sea region cannot be properly represented.

876
877 The simulations showed some agreements with the reconstructions, but also clear, not yet resolved disagreements
878 (Mauri et al., 2014). In summer, all models showed temperatures 2-3°C warmer during the Mid-Holocene
879 Optimum than in the preindustrial climate (Fig. 3). However, no model showed the gradient with clearer warming
880 in the western Baltic Sea, seen in the reconstructions. For wintertime, the disagreement was much clearer. Whereas
881 reconstructions show a clear warming over the whole region, the 14 models displayed widely varying patterns of
882 temperature change. Only three models agreed with the reconstructions. For precipitation, the models disagreed
883 with the west (wet) – east (dry) dipole shown by the reconstructions for summer and winter precipitation (Fig. 3).
884 Not a single simulation showed this pattern of summer precipitation change, and in general the simulated
885 precipitation deviations were much smaller than in the reconstructions. This disagreement regarding temperature
886 (especially in winter) and precipitation, known as the **M**id-Holocene conundrum, is not unique for the Baltic Sea
887 region, but was also found for the Mediterranean (Mauri et al., 2014; Liu et al., 2014). Errors in the applied external
888 (orbital) forcing can be ruled out, as this forcing can be accurately calculated during this period. The reasons for
889 the disagreement are still unknown. They may involve the influence of chaotic internal climate variations (unlikely
890 over such long time scales), model deficiencies, or reconstruction **inaccuracies****uncertainties**.

891 3.1.4 The past millennium

892 For shorter periods closer to the present, like the past one or two millennia, the data available for reconstructing
893 past climate are denser and more accurate. Abundant dendroclimatological information is available, dated to the
894 exact year, in contrast to the uncertain decadal-scale dating of paleo-pollen data. Recently, temperature
895 reconstructions for **w**estern Europe, spatially resolved and approximately covering the last 1,200 years, have
896 become available (Luterbacher et al., 2016) and are presented here in some detail for the Baltic Sea region. These
897 data are based on analysis of wood density in tree rings. Wood density is more sensitive to growing season
898 temperature than tree-ring width. In addition, tree-ring width variations usually contain too weak multidecadal-
899 scale variations, even when the year-to-year variations in temperature may be well captured. This makes wood
900 density a better proxy for temperature reconstructions at these latitudes.

901
902 Figure 4 shows the reconstructed growing-season temperature (spring-early summer) for the period 755-2000 AD,
903 averaged over the Baltic Sea region, based on the European reconstructions by Luterbacher et al. (2016). The
904 reconstructed **spring-early summer** temperature displays warmer conditions around 950 AD, confirmed also by
905 the previous pollen-based reconstructions (Mauri et al., 2015), colder conditions between 1200 and 1850 AD,
906 followed by the recent warming. This temperature evolution confirms that presented by the BACC II Author Team
907 (2015). ~~The relative temperature difference between the Medieval Warm Period and the Contemporary Warm~~
908 ~~Period (mid-20th century) agree within their respective uncertainties~~~~The temperatures in the Medieval Warm Period~~
909 ~~and the Contemporary Warm Period (mid 20th century) are similar within their respective error bars.~~ According to
910 these reconstructions, the Little Ice Age was on average about 0.8 °C colder than the 20th century.

911

There is no new analysis of the causes of this temperature evolution specific for the Baltic Sea region. For Europe as a whole, for which the reconstructions display a similar temporal pattern, the main identified forcings were volcanic activity - more intense during the Little Ice Age and weaker during the Medieval Warm Period - and solar activity, with roughly the reverse temporal signal (Luterbacher et al., 2016). With industrialization, greenhouse gases have become dominant.

The CMIP5 project also included simulations of the past millennium with ~~ESMs~~Earth-System-Models, although with fewer models than for the ~~M~~mid-Holocene. These simulations have been compared with the temperature reconstructions for Europe, in general yielding agreement. However, for the Baltic Sea region, the simulated temperature changes tend to be smaller ~~than those reconstructed~~, especially for the transition between the Medieval Warm Period and the Little Ice Age, with a modelled temperature difference of only $\sim 0.2^{\circ}\text{C}$ (compare with Figure 4 by Luterbacher et al., 2016).

Climate fluctuations are driven not only by the external forcings but also by chaotic internal dynamics of the Earth system. Regional climate simulations indicate that North Atlantic temperature variability influences Baltic Sea temperatures (Kniebusch et al., 2019a) and precipitation (Börger et al., 2018). North Atlantic temperatures tend to fluctuate internally at multidecadal timescales, the AMO, ~~which~~and influences the atmospheric circulation of the Baltic Sea region. Further, the interaction between internal modes of climate variability has recently been identified as a key driver for the state of the Baltic Sea. Internal fluctuations in the North Atlantic are likely to influence the spatial position of the NAO, affecting the regional importance of this climate mode for the Baltic Sea (Börger et al., 2020).

Climate simulations also indicate an impact of internally driven climate variability on the frequency of wind extremes. In the present climate, the wintertime wind regime in the Baltic Sea is linked to the NAO, but at the longer time scales of the preindustrial period, variations in wind extremes appear related neither to the mean wind conditions nor to the external climate forcings (Bierstedt et al., 2015). In the recent centuries, the main driver of trends of wind extremes over land appears to be land-use changes such as de- and reforestation (Bierstedt et al., 2015; Gröger et al., 2021a, and references therein).

An important question is how North Atlantic variations can influence the state of the Baltic Sea, especially its oxygen conditions, since freshwater input and water temperature (less strongly) affect the stratification of the water column, and therefore the exchange of oxygen between the surface and deeper layers. Temperature also modulates algal blooms and thus dissolved oxygen, when bacteria use oxygen to decompose dead algae. Analysis of sediment cores indicated that the ~~M~~mid-Holocene Optimum, the Roman Period (2000 BP), and the Medieval Warm Period were all periods of oxygen deficiency at the bottom of the Baltic Sea. Low oxygen conditions are also observed during the Contemporary Warm Period, unique in their extent on a thousand year perspective (Norbäck Ivarsson et al., 2019, and references therein). Hence, factors other than temperature, like nutrients input into the Baltic Sea, can also affect oxygen conditions, and thus the reasons for those hypoxic phases during the past millennia are not yet completely clear (Schimanke et al., 2012). It had been suggested that agricultural nutrient input was large

951 enough to influence oxygen conditions already during the Medieval Warm Period, perhaps also modulated by
952 changes in river runoff due to the described climate fluctuations (Zillén and Conley, 2010). However, a detailed
953 analysis of new sediments records find little evidence of anthropogenic eutrophication before the industrial period
954 (Norbäck Ivarsson et al., 2019; Ning et al., 2018; van Helmond et al., 2018). In view of the large temperature
955 increases projected for this region in the next decades, further study of the influence of climate on oxygen
956 conditions is warranted.

957 3.2 Present climate change

958 This section assesses our knowledge of Baltic Sea region climate variability during the past ~200 years, based on
959 instrumental records, model based reconstructions and reanalyses. We focus on changes in means, extremes, trends
960 and decadal to multidecadal climate variability.

961 3.2.1 Atmosphere

962 3.2.1.1 Large-scale atmospheric circulation

963 Long-term trends in NAO could not be detected (e.g. Deser et al., 2017; Marshall et al., 2020). For the period
964 1960-1990, a positive trend in NAO, with more zonal circulation, mild and wet winters and increased storminess
965 in central and northern Europe was found (Hurrell et al., 2003; Gillett et al., 2013; Ruosteenoja et al., 2020).
966 However, from after the mid-1990s to the early 2010s, there was a tendency towards more negative NAO indices,
967 i.e. a more meridional circulation and more cold spells in winter (Fig. 5).

968
969 There is no consensus on how strongly the interannual NAO variability is forced externally (Stephenson et al.,
970 2000; Feldstein, 2002; Rennert and Wallace, 2009). Several external forcing mechanisms have been proposed,
971 most prominently SST (Rodwell et al., 1999; Marshall et al., 2001) and sea ice in the Arctic (Strong and
972 Magnusdottir, 2011; Peings and Magnusdottir, 2016; Kim et al., 2014; Nakamura et al., 2015). However, other
973 authors (Screen et al., 2013; Sun et al., 2016; Boland et al., 2017) found no dependence on sea-ice extent.
974 Furthermore, the impact of changes in the Arctic on midlatitude dynamics isare still under debate (Dethloff et al.,
975 2006; Francis and Vavrus, 2012; Barnes, 2013; Cattiaux and Cassou, 2013; Vihma, 2017). However, Scaife and
976 Smith (2018) suggested that the atmospheric circulation in climate models might not be sensitive enough to
977 changes in, for instance, sea surface conditions,

978
979 A weakening of the zonal wind, eddy kinetic energy and amplitude of Rossby waves in summer (Coumou et al.,
980 2015) as well as an increased waviness of the jet stream associated with Arctic warming (Francis and Vavrus,
981 2015) in winter have been identified, which may be linked to an increase in blocking frequencies. Blackport and
982 Screen (2020) argued that previously observed correlations between surface temperature gradients and the
983 amplitude of Rossby waves have broken down in recent years. Therefore, previously observed correlations may
984 have to be reinterpreted as internal variability. On the other hand, it has been shown that observed trends in
985 blocking are sensitive to the choice of the blocking index, and that there is a huge natural variability that
986 complicates the detection of forced trends (Woollings et al., 2018), compromising the robustness of observed
987 changes in blocking.

Formatiert: Schriftart: (Standard) Times New Roman, Nicht unterstrichen

Formatiert: Schriftart: (Standard) Times New Roman, Nicht unterstrichen

Formatiert: Schriftart: (Standard) Times New Roman, Nicht unterstrichen

Formatiert: Schriftart: (Standard) Times New Roman, Nicht unterstrichen

Formatiert: Schriftart: Schriftfarbe: Schwarz

988
989 With ongoing global warming, the Arctic will warm faster than the rest of the [Earth](#). This decrease of the poleward
990 temperature gradient will tend to weaken the westerlies and increase the likelihood of blockings. On the other
991 hand, maximum warming (compared to other tropospheric levels) will occur just below the tropical tropopause
992 due to the enhanced release of latent heat, which tends to increase the poleward gradient, strengthen upper-level
993 westerlies and affect the vertical stability, thus altering the vertical shear in midlatitudes. It is not clear which of
994 these two factors will have the largest effect on the jet streams (Stendel et al., 2021).

995
996 The atmospheric circulation over Europe naturally varies significantly on decadal time scales (Dong et al., 2017;
997 Ravestein et al., 2018). Proposed drivers for these circulation changes include polar and tropical amplification,
998 stratospheric dynamics and the AMOC (Haarsma et al., 2015; Shepherd et al., 2018; Zappa and Shepherd, 2017).
999 The attribution of drivers is more straightforward for local changes, in particular for the soil-moisture feedback,
1000 for which an enhancement of heat waves due to a lack of soil moisture has been demonstrated (Seneviratne et al.,
1001 2013; Teuling, 2018; Whan et al., 2015). Räisänen (2019) found only a [weak](#) effect of circulation changes on the
1002 observed annual mean temperature trends [for 1979-2018](#) in Finland, but circulation changes have considerably
1003 modified the trends in individual months. In particular, circulation changes explain the lack of observed warming
1004 in June, the very modest warming in October in southern Finland, and about a half of the very large warming in
1005 December.

1006
1007 As part of its natural variability, the North Atlantic warmed from the late 1970s to 2014 (Fig. 6). Recently, the
1008 AMO began transitioning to a negative phase again (Frajka-Williams et al., 2017). Paleoclimate reconstructions
1009 and model simulations suggest that the AMO might change its dominant frequency over time (Knudsen et al.,
1010 2011; Wang et al., 2017). The impact of the AMO on climate is, however, independent of its frequency (Börgel et
1011 al., 2018; Börgel et al., 2020). Its influence on regional climate has been analyzed in several studies (Enfield et al.,
1012 2001; Knight et al., 2006; Sutton and Hodson, 2005; Ting et al., 2011; Casanueva et al., 2014; Ruprich-Robert et
1013 al., 2017; Peings and Magnusdottir, 2014), some dealing with the Baltic Sea (Börgel et al., 2018; Börgel et al.,
1014 2020; Kniebusch et al., 2019a). Kniebusch et al. (2019a) suggested that the influence of the AMO on [the warming](#)
1015 [of Baltic Sea SSTs temperature](#) during 1980-2008 might have been at least as strong as that induced by humans
1016 (IPCC, 2014b).

1017 3.2.1.2 Air temperature

1018 A significant increase in surface air temperature in the Baltic Sea region during the last century has been shown
1019 previously (e.g. BACC Author Team, 2008; Rutgersson et al., 2014; BACC II Author Team, 2015). The
1020 temperature increase was not monotonous but accompanied by large multidecadal variations that divided the 20th
1021 century into three main phases: (1) warming from the beginning of the century until the 1930s; (2) slight cooling
1022 until 1960s; and (3) a distinct warming during the last decades of the time series that has continued also during
1023 2014-2020 (Figs. 7 and 8 and Table [94](#)).

1024

1025 Linear trends of the annual mean temperature anomalies during 1878–2020 were $0.10\text{ }^{\circ}\text{C decade}^{-1}$ north of 60°N
1026 as well as south of 60°N in the Baltic Sea region. This is larger than the global mean temperature trend and slightly
1027 larger compared to the earlier BACC reports. Over the Baltic Sea surface air temperature trends were smaller than
1028 over land. During 1856–2005, surface air temperature over the Baltic Sea increased by 0.06 and $0.08\text{ }^{\circ}\text{C decade}^{-1}$
1029 in the central Baltic Sea and in the Bothnian Bay, respectively (Kniebusch et al., 2019a).

1031 There is a large variability in annual and seasonal mean temperatures, ~~in~~ particularly during winter, but the
1032 warming is seen for all seasons (being largest during spring in the northern part of the region).

1034 Both daily minimum and daily maximum temperatures have increased. A decrease in the daily temperature range
1035 (DTR) ~~has~~ been observed in many regions of the world, but there is no clear signal for the entire Baltic Sea
1036 region (see for example, Jaagus et al., 2014, for DTR analysis of the Baltic ~~states~~~~countries~~).

1038 These changes have also resulted in seasonality changes: the growing season has lengthened by about 5 days
1039 decade⁻¹ in the period 1965–2016 (Cornes et al., 2019). From this follows that the cold season has become shorter.

1041 Extreme air temperatures can be high or low, but extended periods of extreme temperatures (spells or waves) are
1042 often the most influential. Averaged over land areas, warm spell duration increased during recent decades
1043 (Rutgersson et al., 2021). For some regions, the ~~annual number of days defined to belong in average annual days~~
1044 ~~defined as~~ warm spells increased from 6–8 to 14 during recent decades. Along with more frequent and longer
1045 warm spells came decreases in the frequency, duration and severity of cold spells, based both on observations
1046 (Easterling et al., 2016) and model results. The length of the frost season and the annual number of frost days also
1047 decreased (Sillmann et al., 2013).

1048 3.2.1.3 Solar radiation and cloudiness

1049 Multidecadal variations of solar radiation at the Earth's surface, called “dimming” and “brightening”, have been
1050 observed in Europe and other parts of the world, particularly in the northern hemisphere (Wild et al., 2005; Wild,
1051 2012; Wild et al., 2017).

1053 One of the world's longest time series of global radiation, i.e. incoming solar radiation at the Earth's surface, is
1054 from Stockholm, where measurements started in 1922. Recently, a first attempt to homogenize this time series was
1055 made by Josefsson (2019). No significant trend was found over the whole time series, but there were large
1056 variations over one to three decades. Other long time series of global radiation in ~~n~~Northern Europe are from
1057 Potsdam, Germany, (Wild et al., 2021), and Tõravere, Estonia, (Russak, 2009). All three time series show a
1058 minimum in global radiation around the mid-1980s. Then a clear increase or “brightening” of about 5–8% followed,
1059 until at least 2005. Before the 1980s minimum there was a period of “dimming” at all stations but with differences
1060 in the details. ~~In Potsdam, there was rather stable dimming all the time from the late 1940s. In Tõravere, there was~~
1061 ~~a maximum dimming around mid-1960s, while Stockholm recorded high values both around 1950 and 1970, with~~
1062 ~~a minimum dimming in between. In Potsdam, there was rather stable dimming with decreasing solar radiation~~

Formatiert: Englisch (USA)

from the late 1940s until the mid-1980s. In Tõravere, the dimming period started around mid-1960s, while in Stockholm the dimming phase started around 1950 but with a temporary interruption with increased solar radiation around 1970. The data also suggest an early brightening in Stockholm time series, but this is still uncertain, especially before 1950.

Current twentieth century reanalyses models provide results for surface solar radiation. However, most of them fail to capture multidecadal surface radiation variability in central and southern Europe (Wohland et al., 2020). The ~~model~~ CERA20C reanalysis, which shows best results for central and southern Europe still gives questionable results over Scandinavia, showing a weak increase instead of a decrease in surface solar radiation during the presumed dimming period before 1980.

Satellite data allowing analyses of cloudiness and solar radiation at the Earth's surface are available since the early 1980s. For Europe, important work has been done within the EUMETSAT Satellite Application Facility on Climate Monitoring (CM SAF). Several satellite data records have been validated and used in climate studies (e.g. Urraca et al., 2017; Pfeifroth et al., 2018). At the highest latitudes of the Baltic Sea region there are, however, larger inaccuracies/uncertainties (Riihelä et al., 2015) or often no data at all, due to the low standing Sun and slant viewing geometry from the satellites.

The satellite data only cover the latest brightening period observed at ground-based stations in Europe. While the geographical patterns of global average cloud conditions agree well among several satellite cloud-data sets, there are clear differences in the distribution and size of trends (Karlsson and Devasthale, 2018). However, there seems to be consensus on a decreasing trend in total cloud fraction of about 1-2% per decade over the Baltic Sea region during 1984-2009.

Recent CM SAF satellite products on solar irradiance at the Earth's surface, the SARA-2 and CLARA-A2 datasets, both agree well with station data according to Pfeifroth et al. (2018). In many cases this holds both for climatological averages and for trend detection. The average trend for the period 1983-2015 is about $+3 \text{ W m}^{-2} \text{ decade}^{-1}$ both at the stations closest to the Baltic Sea and in the SARA-2 dataset. The three long-term stations mentioned above are all used as reference stations for the satellite data validation. For example, the on-going monitoring at stations spread over all Sweden show an average increase of about 8% (corresponding to $+4 \text{ W m}^{-2} \text{ decade}^{-1}$) from 1983 until 2005-2006 (SMHI, 2021). In later years the solar radiation leveled off, but *inter alia* the extremely sunny 2018 in Northern Europe contributed to keeping the trend increasing over time.

The multidecadal variations in the solar radiation at the Earth's surface were most probably caused by a combination of changes in cloudiness and in anthropogenic aerosols. Which of the two drivers is the largest contribution is still an open question, and might differ among regions. Aerosol concentrations over Northern Europe decreased during the brightening period from the mid-80s onwards (Ruckstuhl et al., 2008; Russak, 2009; Markowicz and Uscka-Kowalkowska, 2015; Glantz et al., 2019). Russak (2009) considered changes in cloudiness caused by variations in atmospheric circulation to be the most important factor in Estonia, but aerosol changes also

played a role. In an early study of the modern radiation measurements in Sweden the strong increase in solar radiation 1983-1997 was also accompanied by a clear decrease in total cloud cover, especially during the half-year of summer (Persson, 1999). The satellite datasets SARAH-2 and CLARA-A2 were both derived using an aerosol climatology as input. This underlines the important role of changes in cloudiness for surface solar radiation. Stjern et al. (2009) also stressed the importance of the contribution of clouds and the atmospheric circulation for dimming and brightening periods in Northern Europe.

Other studies, e.g. Ruckstuhl et al. (2008) and Wild et al. (2021), concluded that aerosol effects under clear skies is the main contributor to the multidecadal variations of solar radiation in central Europe. Aerosol-induced multidecadal variations in surface solar radiation could be expected also over oceans (Wild, 2016), but long-term measurements are lacking. The interaction between aerosols and clouds, the indirect aerosol effects, needs also be better understood and quantified.

3.2.1.4 Precipitation

During the twentieth century in the Baltic Sea region, changes in precipitation were spatially more variable than for temperature (BACC II Author Team, 2015). Irregularly distributed precipitation measurement stations make it difficult to determine statistically significant trends and regime shifts. Sweden shows an overall ~~increasing wetting~~ trend in precipitation since the 1900s, in particular since the mid 20th century (Chen et al., 2020). In Finland, the overall increase detected for 1961-2010 is neither regionally consistent nor always statistically significant (Aalto et al., 2016). The same holds for the Baltic States (Jaagus et al., 2018). In the south of the Baltic Sea region, changes were small and not significant. Nevertheless, precipitation averaged over the Baltic Sea catchment area has increased since 1950 due to an increase in winter (Fig. 9).

The number of heavy precipitation days is largest in summer. Compared to southern Europe, precipitation extremes in the Baltic Sea region are not as intense, ~~with the 95th percentile of wet-day precipitation with daily~~ amounts typically ranging from 8 to 20 mm (Cardell et al., 2020). Extreme precipitation intensity increased during the period 1960-2018. ~~An index for the annual maximum five-day precipitation~~ ~~An index for the maximum annual five consecutive days of precipitation~~ (Rx5d) shows significant increases of up to 5 mm per decade over the eastern part of the Baltic Sea catchment (EEA, 2019b). The change is more pronounced in winter than in summer.

3.2.1.5 Wind

In situ observations allow direct analysis of winds, in particular over sea (e.g. Woodruff et al., 2011). However, in situ measurements, especially over land, are often locally influenced, and inhomogeneities make the straightforward use of such data difficult, even for recent decades. Therefore, many studies use reanalyses rather than direct wind observations. But analysis of storm-track activity for longer periods using reanalysis data suffers necessarily from ~~inaccuracies~~ ~~uncertainties~~ associated with changing data assimilation and observations before and after the introduction of satellites, resulting in large variations of storm-track changes across assessments (Wang et al., 2016; Chang and Yau, 2016). Another concern about wind trends in reanalyses, especially over land, is related to appropriate consideration of the effects of land use changes on surface roughness.

1139

1140 Owing to inherent inhomogeneities and the large climate variability in the Baltic Sea region, it is unclear whether

1141 there is a general trend in wind speed in the recent climate. Results regarding changes or trends in the wind climate

1142 are strongly dependent on period and region considered (Feser et al., 2015). Due to the strong link to large-scale

1143 atmospheric variability over the North Atlantic, conclusions about changes over the Baltic Sea region are perhaps

1144 best made in a wider spatial context, considering *inter alia* the NAO.

1145

1146 Recent trend estimates for the total number of cyclones over the Northern Hemisphere extratropics during 1979-

1147 2010 revealed a large spread across the reanalysis products, strong seasonal differences, as well as decadal-scale

1148 variability (Tilinina et al., 2013; Wang et al., 2016; Chang et al., 2016; Matthews et al., 2016; Chang et al., 2012).

1149 Common to all reanalysis datasets is a weak upward trend in the number of moderately deep and shallow cyclones

1150 (~~7 to 11% per decade for both winter and summer~~), but a decrease in the number of deep cyclones, in particular

1151 for the period 1989-2010. Chang et al. (2016) reported a minor reduction in cyclone activity in the Northern

1152 Hemisphere summer due to a decrease in baroclinic instability as a consequence of Arctic temperatures rising

1153 faster than at low latitudes. Chang et al. (2012) also noticed that state-of-the-art models (CMIP5) generally

1154 underestimate this trend. In the Northern Hemisphere winter, recent studies reported a decrease in storm track

1155 activity related to Arctic warming (Ceppi and Hartmann, 2015; Shaw et al., 2016; Wills et al., 2019; Stendel et al.,

1156 2021).

1157

1158 Despite large decadal variations, there is still a positive trend in the number of deep cyclones (< 980 hPa) over the

1159 last six decades, which is consistent with results based on the NCEP reanalysis since 1958 over the northern North

1160 Atlantic Ocean (Lehmann et al., 2011). Using an analogue-based field reconstruction of daily pressure fields over

1161 central to ~~n~~Northern Europe (Schenk and Zorita, 2012), the increase in deep lows over the region might be

1162 unprecedented since 1850 (Schenk, 2015). However, for limited areas the conclusions were rather uncertain.

1163

1164 The effect of differential temperature trends on storm tracks has been recently addressed, both in terms of upper

1165 tropospheric tropical warming (Zappa and Shepherd, 2017) and lower tropospheric Arctic amplification (Wang et

1166 al., 2017), including the direct role of Arctic sea-ice loss (Zappa et al., 2018), and a possible interaction of these

1167 factors (Shaw et al., 2016). The remote and local SST influence has been further examined by Ciaso et al. (2016),

1168 who confirmed the sensitivity of the storm tracks to the SST trends generated by the models and suggested that

1169 the primary greenhouse gas influence on storm track changes was indirect, acting through its influence on SSTs.

1170 The importance of the stratospheric polar vortex for storm track changes has ~~recently also~~ received ~~more~~ attention

1171 (Zappa and Shepherd, 2017). ~~In an aqua planet simulation, Sinclair et al. (2020) found a decrease in the number~~

1172 ~~of extratropical cyclones and a poleward and downstream displacement due to an increase in diabatic heating.~~

1173 **3.2.1.6 Air pollution, air quality and atmospheric nutrient deposition**

1174 Air pollution continues to significantly impair the health of the European population, particularly in urban areas.

1175 Brandt et al. (2013) estimated the total number of premature deaths due to air pollution in Europe in the year 2000

1176 to be ~680 000 year⁻¹. Although this number was predicted to decrease to approximately 450 000 by 2020, it is

still a matter of ~~grave~~-concern. Particulate matter (PM) concentrations were reported to be the primary reason for adverse health effects. Estimates indicated that PM_{2.5} concentrations in 2016 were responsible for ~412 000 premature deaths in Europe, due to long-term exposure (EEA, 2019a).

The state of air pollution is often expressed as air quality, when human health is in focus. The ambient air quality in the Baltic Sea region is dominated by anthropogenic emissions, and natural emissions play only a minor role. These emissions show an overall decreasing trend in recent years (EEA, 2019a), as reflected in the ambient concentrations reported in the EMEP status report (EMEP, 2018). To quantify air quality, concentrations of certain gases and particulate matter are used as measures. The general conclusions in the field of air quality reported by the BACC II Author Team (2015) still hold today, i.e. that land-based emissions and concentrations of major constituents continue to decrease due to emission control measures, with the possible exception of certain emissions from the shipping sector. Sulphur emissions from shipping have continued to decrease strongly in the Baltic Sea from 2015, due to much lower limit values for the sulphur content of ship fuel in the emission control areas. A noticeable decrease in nitrogen emissions due to the newly (2021) implemented nitrogen emission control area (NECA) is expected in the next decade.

In Europe, the pollutants most harmful to human health are ~~particulate-matter~~-(PM), nitrogen dioxide (NO₂) and ground-level ozone (O₃). About 14% of the EU-28 urban population was exposed to O₃ concentrations above the EU target value threshold (EEA, 2019a). When compared to other European countries, air pollution was relatively low in Scandinavia, ~~excepti~~on for a few urban traffic hotspots, with annual mean NO₂ concentrations elevated to near the limit value (EEA, 2019b). In this comparison, ~~n~~Northern Germany is located in the lower mid-field, while northern Poland is among the more polluted countries, especially with PM. Biomonitoring samples analyzed for toxic metals by Schröder et al. (2016) tended to show lowest concentrations in ~~n~~Northern Europe.

Contributions to air pollution and pollutant deposition in coastal areas by shipping can be substantial. Major pollutants from shipping are SO₂, NO_x and PM (including black carbon). The BACC II assessment estimated emissions from shipping in the Baltic Sea region. Several studies have recently been published, including Jonson et al. (2015), Claremar et al. (2017), and Karl et al. (2019b), which used ~~chemistry~~ transport models to predict ~~ed~~ ambient concentrations from known emissions. They show ~~ed~~, as expected, that the highest air pollution concentrations due to ship exhaust are found near major shipping lanes and harbors, but also that considerable concentrations of NO₂ and PM reach populated land areas. This effect is pronounced in the south-western Baltic Sea area (Quante et al., 2021). Exact numbers from such modelling should still be interpreted with care, as shown by Karl et al. (2019a), who compared output from three state-of-the-art chemistry transport models for the Baltic Sea area.

The most important recent change in shipping emissions in the North ~~Sea~~ and Baltic ~~Seas~~ are due to the 2015 strengthening of the fuel sulphur content limit for the Sulphur Emission Control Areas (SECAs), by lowering the maximum allowed sulphur content from 1 to 0.1%. Model calculations indicated ~~large~~ reductions in sulphur deposition in countries bordering these two sea areas after the implementation of the lowered sulphur limit (Gauss

et al., 2017). Barregard et al. (2019) estimated the contribution of Baltic Sea shipping emissions to PM_{2.5} before 2014 and after 2016, when the new SECA regulation of marine fuel sulphur was implemented. These authors also estimated human exposure to PM_{2.5} from shipping and its health effects in the countries around the Baltic Sea. They concluded that PM_{2.5} emissions from Baltic Sea shipping, and resulting health impacts decreased substantially after the 2015 SECA regulation. Population exposure studies estimating the influence of shipping emissions for selected Baltic Sea harbor cities were published for Rostock, Riga and Gdańsk–Gdynia by Ramacher et al. (2019) and for Gothenburg by Tang et al. (2020). Ramacher et al. (2019) found that shipping emissions strongly influence NO₂ exposure in the port areas (50–80 %), while the average influence in home, work and other environments is lower (3–14 %) but still with strong influence close to the ports. It should, however, be noted that reduction of sulphur emissions to the atmosphere by the use of new cleaning techniques (e.g. open loop scrubbers) can increase the risk of acidification and marine pollution (Turner et al., 2017; 2018).

Johansson et al. (2020) published a first comprehensive assessment of emissions from leisure boats in the Baltic Sea. While the modeled NO_x and PM_{2.5} emissions from leisure boats are clearly lower than those from commercial shipping, these first estimates suggest that carbon monoxide (CO) emissions from leisure boats equal 70 % of the registered shipping emissions and non-methane volatile organic carbon (NMVOC) emissions equal 160 %. It should be noted that most of the leisure boat emissions occur in summer, and often occur near areas for nature conservation and tourism. Most of these emissions can be attributed to Swedish, Finnish and Danish leisure boats, but the leisure boat fleet has the potential for large future increases also in Russia, Estonia, Latvia, Lithuania and Poland.

Air pollution leads to environmental degradation by affecting natural ecosystems and biodiversity. Ground-level ozone (O₃) can damage crops, forests and other vegetation, impairing growth and reducing biodiversity. According to a recent study by Proietti et al. (2021), ~~assessed trends of the O₃ mean concentration in Northern Europe were not statistically significant for the time period from 2000 to 2014.~~ for the period 2000-2014 no statistically significant trend in the O₃ mean concentration in northern Europe could be identified, due to the large internal variability. The annual mean ozone concentration is reported to be slightly below 35 ppb, as compared to 43 to 45 ppb in the Mediterranean ~~r~~Region, for which a significant decreasing trend is found. The exposure index AOT40 (sum of the hourly exceedances above 40 ppb, for daylight hours during the growing season) significantly declined in all European regions except for ~~n~~Northern Europe, for which a positive but not significant trend is seen. On the nation level, among the six European countries showing a positive trend were Denmark, Germany, Sweden (Proietti et al., 2021). A clear difference in trends between rural sites and other station typologies is found for Europe for the period 2000 to 2017. ~~That is, i.e.~~ for traffic sites a substantial increase of annual mean O₃ concentration was observed, in contrast to rural stations, for which a slight decrease was found (Colette and Rouïl, 2020). Regarding the monitored population exposed to a large number of days with high ozone concentrations ~~population exposed to ozone~~ all countries in Europe showed a decrease from 2000 to 2014 (NDGT60 > 25 days per year; Fleming et al., 2018).

Formatiert: Nicht unterstrichen, Schriftfarbe: Automatisch, Tiefgestellt

Harmful exposure and impacts of air pollutants on ecosystems are assessed using the concept of critical loads (CLs; Nilsson and Grennfelt, 1988). The CL is the amount of pollutants that an ecosystem can tolerate without risking unacceptable damage. The most harmful air pollutants in terms of damage to ecosystems in addition to O₃ are ammonia (NH₃) and nitrogen oxides (NO_x). It is estimated that about 62% of the European ecosystem area is still exposed to high levels of NO_x, leading to exceedances of CLs for eutrophication of open water bodies in all European countries in all countries in 2016 (EEA, 2019a). Hotspots of exceedances of CLs for acidification in 2016 were the Netherlands and its borders with Germany and Belgium, southern Germany and also Czechia. However, most of Europe including the Baltic Sea region did not exceed the CLs for acidification (EEA, 2019a).

Since the 1980s, the total nitrogen deposition on the Baltic Sea has decreased substantially, due to an overall reduction of European emissions, but emission and deposition reductions have stalled since the mid-2000s (Colette et al., 2015; Gauss et al., 2021). Atmospheric phosphorus deposition remains highly uncertain in amount and trends (HELCOM, 2015; Kanakidou et al., 2018; Ruoho-Airola et al., 2012).

Air quality and climate interact in several ways. On the one hand, air pollutants can affect climate both directly and indirectly by changing the radiative balance of the atmosphere. On the other hand, climate change alters meteorological conditions, which may affect concentrations of air pollutants via several pathways, since air quality is strongly dependent on weather (Jacob and Winner, 2009). The effects of important meteorological and climate variables on surface O₃ and PM were discussed in a comprehensive review by Doherty et al. (2017). The connection between high temperatures and increased ground-level ozone concentrations is well established. Increases in temperature related to climate change (i.e. during heat waves) are expected to lead to higher ozone concentrations in certain regions with the required precursor concentrations. Other important meteorological factors influencing air pollution concentrations are a possible change in the number of midlatitude cyclones and in the number of occurrences and duration of stagnant weather conditions (Jacob and Winner, 2009).

3.2.2 Land

3.2.2.1 River discharge

The total river discharge to the Baltic Sea is approximately 14,000 m³ s⁻¹ (Bergström and Carlsson, 1994). This is substantially more than the direct net precipitation (precipitation minus evaporation) on the Baltic Sea itself, which has been estimated at 1,000-2,000 m³ s⁻¹ (Meier and Kauker, 2003; Meier and Döscher, 2002; Meier et al., 2019d), see also the discussion by Leppäranta and Myrberg, (2009). In other words, most of the fresh water entering the Baltic Sea comes from the terrestrial part of the catchment. Therefore, the fresh-water input to the Baltic Sea cannot be described entirely with only climatic parameters. Non-climatic drivers of runoff include river regulation by dams and reservoirs, land-use changes in the catchment, and water uptake for irrigation. Although dams are known to have altered the seasonality of discharge (e.g. McClelland et al., 2004; Adam et al., 2007; Adam and Lettenmaier, 2008), they do not seem to be responsible for annual discharge changes. In the long-term, net precipitation over the catchment area and river runoff are strongly correlated (Meier and Kauker, 2003).

Formatiert: Nicht unterstrichen, Schriftfarbe: Automatisch, Tiefgestellt

For the period 1850-2008, the total river discharge from the Baltic Sea catchment area, reconstructed from observations (Bergström and Carlsson, 1994; Cyberski et al., 2000; Hansson et al., 2011; Mikulski, 1986) and hydrological model results (Graham, 1999), showed no statistically significant trend but a pronounced multidecadal variability, with a period of about 30 years (Meier et al., 2019d). Furthermore, summed river flow observations in the period 1900-2018 (Lindström, 2019) and a historical reconstruction of the annual river discharge for the past 500 years showed no statistically significant trend either (Hansson et al., 2011). However, river runoff from northern Sweden, a part of the catchment area of the Bothnian Bay, significantly increased since the 1980s compared to 1911-2018 (Lindström, 2019).

There are indeed substantial regional and decadal variations in the river flow. Stahl et al. (2010) studied near-natural rivers of Europe over the period 1942-2004 and found a clear overall pattern of positive trends in annual streamflow in the northern areas. (Kniebusch et al., 2019b) also identified a statistically significant positive trend in the river discharge to the Bothnian Bay for 1921-2004. In Estonian rivers, regime shifts in annual specific runoff corresponded to the alternation of wet and dry periods (Jaagus et al., 2017). A dry period started in 1963/1964, followed by a wet period from 1978, with the latest dry period commencing at the beginning of the 21st century.

For the period 1920-2005, positive trends in stream flow at stations of a pan-Nordic dataset dominate annual mean, winter and spring figures, whereas summer trends are statistically not significant (Wilson et al., 2010). A clear signal of earlier snow-melt floods and a tendency towards more severe summer droughts in southern and eastern Norway were found.

~~The observed temperature increases have affected stream flow in the northern Baltic Sea region for 1920-2002 in a manner corresponding well to the projected consequences of a continued rise in global temperature (Hisdal et al., 2010). Regarding precipitation, however, the regional impacts of both the observed and projected changes on stream flow are still unclear.~~
The observed temperature increases have affected stream flow in the northern Baltic Sea region for 1920-2002 in a manner corresponding well to the projected consequences of a continued rise in global temperature in term of increasing winter time discharges (Hisdal et al., 2010). However, the regional impacts of precipitation change on both the observed and projected changes in stream flow are still unclear as the combined effects of changes in precipitation and temperature are still not well known (Stahl et al., 2010).

In the northern Baltic Sea region, all the way south to the Gulf of Finland, runoff is strongly linked to the climate indices air temperature, wind and rotational circulation components. In the southern region, runoff is associated more with the strength and torque of the cyclonic or anticyclonic pressure systems (Hansson et al., 2011).

In the Baltic states ~~(Lithuania, Latvia and Estonia)~~, changes in streamflow over the 20th century showed a redistribution of runoff over the year, with a significant increase in winter and a tendency for decreasing spring floods (Reihan et al., 2007; Sarauskiene et al., 2015; Jaagus et al., 2017). A similar winter trend was found also for the reconstructed river discharge to the entire Baltic Sea since the 1970s (Meier and Kauker, 2003).

For the period 1911-2010, a trend of observed annual maximum daily flows in Sweden could not be detected (Arheimer and Lindström, 2015). However, in particular the annual minimum daily flows in northern Sweden considerably increased in the period 1911-2018 (Lindström, 2019). Analyzing a pan-European database, Blöschl et al. (2017) showed that river floods over the past five decades occurred earlier in spring due to (1) an earlier spring snow melt in northeastern Europe, (2) delayed winter storms associated with polar warming around the North Sea, and (3) earlier soil moisture maxima in ~~w~~Western Europe.

3.2.2.2 Land nutrient inputs

The Baltic Sea catchment area of 1.7 million km², which is more than four times larger than the sea surface area ~~of the Baltic Sea~~ (cf. Fig. 1), is populated by over 84 million inhabitants. Stretching between 49° - 69°N and 10° - 38°E, the catchment exhibits significant gradients in both natural (precipitation, river discharge, temperature, etc.) and anthropogenic (population density and occupation, agricultural and industrial development, etc.) environmental factors. These factors change both in time (phenological changes, long-term trends and lags due to land cover processes) and space (north-south gradients in climate and land use, east-west gradients in socio-economic features and climate) thus determining heterogeneity and variation of land nutrient inputs that drive long-term eutrophication of the Baltic Sea (Savchuk, 2018, and references therein; Kuliński et al., 2021).

Estimates of nutrient inputs had been attempted since the 1980s (e.g. Larsson et al., 1985; Stålnacke et al., 1999) and are now being compiled within a permanent process of the HELCOM Pollution Load Compilation (PLC, e.g. HELCOM, 2019). However, these data officially reported to HELCOM by the participating riparian states have been and still are suffering from gaps and inconsistencies. Therefore, the “best available estimates” have been reconstructed in attempts to both fill in such gaps and correct possible sources of inconsistencies (Savchuk et al., 2012; ~~the present study based on~~ Svendsen and Gustafsson, 2020). For long-term studies of the Baltic Sea ecosystem, a historical reconstruction of nutrient inputs since 1850 is available (Gustafsson et al., 2012).

According to HELCOM (~~HELCOM~~, 2018a; Savchuk, 2018) and updated estimates (HELCOM, 2018c), substantial reductions of land nutrient inputs, comprising riverine inputs and direct point sources at the coast, have been achieved since the 1980s (Fig. 11, Table 105). Since there are no statistically significant trends in annual river discharge (Section 3.2.2.1), these reductions are attributed to socio-economic development, including expansion of the wastewater treatment and reduction of atmospheric nitrogen deposition (Gauss et al., 2021) over the entire Baltic Sea drainage basin, and not to climate-related effects (HELCOM, 2018a; Svendsen and Gustafsson, 2020). As an example, the coastal point sources of ~~total nitrogen (TN)~~ and ~~total phosphorus (TP)~~ decreased three- and ten-fold, respectively, comparing to the 1990s (Savchuk et al., 2012) and today contribute to the Baltic Sea less nutrients than they did in 1900 (Savchuk et al., 2008; Kuliński et al., 2021).

Agriculture is the main source of anthropogenic diffuse nutrient inputs, which comprise 47% of the riverine nitrogen and 36% of the riverine phosphorus inputs (HELCOM, 2018c). In turn, mineral fertilizer dominates the anthropogenic nutrient inputs to the Baltic Sea catchment, in particular in its intensely farmed southern part (Hong et al., 2017). However, during 2000-2010 only about 17% of the net anthropogenic nitrogen and only about 4.7%

of the net anthropogenic phosphorus input to the catchment ~~we are currently~~ exported with rivers to the sea (Hong et al., 2017). While denitrification might have removed part of the nitrogen applied in agriculture, the remaining phosphorus has accumulated in the drainage basin. A global budget estimated that agriculture has increased the soil storage of phosphorus in the drainage basin by 50 Mt during 1900-2010 (Bouwman et al., 2013) and a regional approach calculated an increase by 40 Mt during 1900-2013 (McCrackin et al., 2018). However, McCrackin et al. (2018) estimated that about 60% of these phosphorus inputs were retained in a stable pool and did not contribute noticeably to the riverine export. About 40% accumulated in a mobile pool with a residence time of 27 years. McCrackin et al. (2018) suggested that leakage from this mobile legacy pool is, though slowly declining, the dominant source of present riverine phosphorus inputs.

3.2.3 Terrestrial biosphere

Previous assessments

The comprehensive review of climate-related changes in terrestrial ecosystems in the first BACC report (BACC Author Team, 2008), Smith et al. (2008), concluded that climate change during the preceding 30-50 years had already caused measurable changes in terrestrial ecosystems in the Baltic Sea region, e.g. an advancement of spring phenological phases in some plants, upslope displacement of the alpine tree-line and increased land-surface greenness in response to improved growth conditions and a richer CO₂ supply. But as nearly all ecosystems in the region were managed to some extent, the climate impacts might be alleviated or intensified by human interventions, e.g. by choosing favorable tree species in forestry. The observed trends were expected to continue for at least several decades, assuming that continued future increases in the atmospheric CO₂ concentration will cause continued warming.

In the second BACC report (BACC II Author Team, 2015), climate effects on terrestrial ecosystems were less in focus, with a section related to forests and natural vegetation in the chapter on environmental impacts on coastal ecosystems, birds and forests (Niemelä et al., 2015) and as part of the chapter on socioeconomic impacts on forestry and agriculture (Krug et al., 2015). On the other hand, the second BACC report also considered anthropogenic land-cover changes as a driver of regional climate change (Gaillard et al., 2015). Niemelä et al. (2015) concluded that the observed positive effects of climate change on forest growth would continue, in particular for boreal forest stands that benefitted more than temperate forest stands. The species composition of natural vegetation in the Baltic Sea region was expected to undergo changes, with a predominantly northward shift of the hemiboreal and temperate mixed forests. Terrestrial carbon storage was likely to increase in the region, but land-use change could play an important modifying role, affecting this storage both positively and negatively. Krug et al. (2015) concluded that there were regional differences in how the vulnerability and productivity of forestry systems were affected by climate change, with the southern and eastern parts of the Baltic Sea region likely to experience reduced production and the northern and western parts increased production. Gaillard et al. (2015) found no indication that deforestation in the Baltic Sea region since 1850 could have been a major cause of the observed climate warming.

Acknowledging the importance of the land component in the climate system, the IPCC recently published its special report entitled 'Climate Change and Land' on climate change, desertification land degradation, sustainable

land management, food security, and greenhouse gas fluxes in terrestrial ecosystems (IPCC, 2019b). This is because land, including its water bodies, provides the basis for human livelihoods and well-being through primary productivity, the supply of food, fresh water, and multiple other ecosystem services.

Biophysical and biogeochemical interactions

The land surface and its terrestrial ecosystems in the Baltic Sea region interact with the atmosphere and are, thus, coupled to the local and regional climate. These interactions determine the exchanges of heat, water and momentum between the land surface and the atmosphere via biophysical processes, the exchange of greenhouse gases, e.g. CO₂, CH₄ and N₂O, and emissions of black carbon, from forest fires, aerosol precursors, e.g. biogenic volatile compounds, or organic carbon aerosols via biogeochemical processes, altering the atmospheric composition.

The nature of the biophysical and biogeochemical interactions between the land surface and the atmosphere and their effects on climate are studied by comparing the effects of forests with that of open land, e.g. grassland, pastures or cropland (e.g. Bonan, 2016). Details on the nature of these feedbacks are given in Gröger et al. (2021b).

On average, across the globe, forests absorb atmospheric CO₂ and, thus, reduce net radiation and have a cooling effect on climate. In equilibrium, forest ecosystems, are expected to be carbon neutral, with carbon loss through phenological turnover, mortality and decomposition over large areas on average matching plant productivity. CO₂ fertilization is considered a strong driver for the terrestrial carbon sink, but demographic recovery following past land use, e.g. afforestation and replanting of harvested forest stands, likely provides an equally important explanation for net carbon uptake by forests in industrialized regions of North America, Europe and Asia (Pugh et al., 2019). Deforestation, on the other hand, can lead to a release of carbon and to an increase in net radiation and, thus, has a warming effect on climate (Friedlingstein et al., 2020, and references therein). In contrast to the biophysical effects ~~described above~~ and some of the other biogeochemical interactions, the biogeochemical interactions associated with the carbon cycle have global impacts and operate at very long time scales.

Anthropogenic land-use and land cover changes

Using remote sensing data, Jin et al. (2019) investigated recent trends in springtime plant phenology in the Baltic Sea region and the sensitivities of phenological trends to temperature and precipitation, in spring, winter and summer. Considering the entire region and combining all vegetation types, the authors found an advancement of the growing season by 0.30 day year⁻¹ over the period 2000-2016. The advancement was particularly strong for evergreen needle-leaved forests (0.47 day year⁻¹) and weaker for cropland and grassland (0.14 day year⁻¹). Evergreen needle-leaved forests, together with deciduous broadleaf forests, dominate the northern part of the Baltic Sea region, while the southern part is mainly grassland and cropland. Jin et al. (2019) found that the most important driver of the advancement of the growing season is spring mean temperature, with an advancement rate of 2.47 day (°C)⁻¹ of spring warming, considering the entire area and all vegetation types. In this study, spring was defined as the three months preceding the start of the growing season. Spring drying could further increase the advancement rate by 0.18 day cm⁻¹ decrease in precipitation. While the sensitivity of the start of the growing season

Formatiert: Schriftart: Nicht Kursiv, Nicht unterstrichen, Schriftfarbe: Automatisch

Formatiert: Schriftart: Nicht Kursiv, Nicht unterstrichen, Schriftfarbe: Automatisch

Formatiert: Nicht unterstrichen, Schriftfarbe: Automatisch, Nicht Hochgestellt/ Tiefgestellt

Formatiert: Nicht unterstrichen, Schriftfarbe: Automatisch, Nicht Hochgestellt/ Tiefgestellt

to climate conditions in spring is comparable for the entire Baltic Sea region, the sensitivity to climate conditions in the previous summer and winter seasons differs between the northern and southern parts of the region. In both seasons an increase in the mean temperature was found to advance the growing season in the southern part of the Baltic Sea region but to delay the start of the growing season in the northern part, in contrast to the spring warming. These sensitivities were markedly stronger for summer than for winter. These trends in plant phenology in spring result in changes of the land cover early in the year, thus affecting climate through biophysical and biogeochemical interactions with the atmosphere.

As mechanisms for the delayed start of the growing season in Fennoscandia in relation to a warming in the preceding summer and winter seasons Jin et al. (2019) suggested the effects on plant dormancy. A winter warming could, for instance, prolong the chilling accumulation required to break winter dormancy of trees. Later summer temperatures, on the other hand, could affect bud dormancy initiation, while reduced soil moisture associated with higher summer and autumn temperatures could delay the leafing and flowering of plants. However, the details of the mechanisms involved have not yet been explored.

Observations reveal a local impact of changes in forest cover on near-surface temperatures, due to biophysical effects that depend on the geographical latitude, roughly separating the boreal regions and the temperate zone of the Northern Hemisphere. When investigating the effects of small-scale clearings at sites in the Americas and Asia, Zhang et al. (2014a) found on both continents that annual mean temperatures cooled over open land north of about 35°N and warmed south of this latitude. Changes in forest cover have, however, different effects on daily minimum and daily maximum temperatures. Zhang et al. (2014a) found that the warming effect over open land south of 35°N was related to an increase in daily maximum temperatures, with little change in daily minimum temperatures, while the cooling effect to the north was due to a decrease in daily minimum temperatures. Lee et al. (2011) showed consistent results for North America, where the cooling effect of $0.85 \pm 0.44^\circ\text{C}$ over non-forested areas north of 45°N was due to a decrease in daily minimum temperatures, associated with the reduced roughness length. At night, open land cools more than forests, regardless of geographical latitude. This is confirmed by Alkama and Cescatti (2016), who analyzed the impacts of recent losses in forest cover on near-surface and land-surface temperatures in the boreal zone. For both, the authors found cooling trends in daily minimum and warming trends in daily maximum temperatures in response to deforestation and opposite tendencies after afforestation. These effects were somewhat stronger for land-surface temperatures than air temperatures.

Regional Climate Models (RCMs) have been used to investigate the biophysical effects of changes in forest cover on climate in Europe. Strandberg and Kjellström (2019), for instance, used simulations with the Rossby Centre Atmosphere (RCA) RCM to assess the climate effect of maximal afforestation or deforestation in Europe, focusing on seasonal mean temperatures and precipitation, as well as daily temperature extremes. Maximum afforestation and deforestation were inferred from a simulation with the LPJ-GUESS dynamical vegetation model, providing a map for potential natural forest cover for Europe in equilibrium with present-day climate (Gröger et al., 2021a). To simulate effect maximum afforestation, present-day land cover classes, which represent considerable agricultural activity in Europe (particularly in western, central and southern Europe), were replaced by the potential

Formatiert: Schriftart: Nicht Kursiv, Nicht unterstrichen, Schriftfarbe: Automatisch

Formatiert: Schriftart: Nicht Kursiv, Nicht unterstrichen, Schriftfarbe: Automatisch

Formatiert: Schriftart: Nicht Kursiv, Nicht unterstrichen, Schriftfarbe: Automatisch

1477 natural forest cover. In the case of deforestation, on the other hand, the potential natural forest cover was converted
1478 to grassland in the model.

1479 The simulations indicated that afforestation in Europe generally increased evapotranspiration, which, in turn, led
1480 to colder near-surface temperatures. In western, central and southern Europe, the cooling in winter due to
1481 afforestation was between 0.5 and 2.5°C. The cooling effect was somewhat stronger in summer, exceeding 2.5°C
1482 in large parts of western and southeastern Europe. Deforestation had the opposite effect, warmer near-surface
1483 temperatures due to decreased evapotranspiration, typically in the range between 0.5 and 2°C in western and
1484 central Europe and reaching up to 3°C in southeastern Europe. In regions with low evapotranspiration, however,
1485 changes in the surface albedo were relatively more important for temperatures. During summer, warming by
1486 deforestation affected the entire Baltic Sea region (in the range between 0.5 and 1.5°C), while the cooling
1487 associated with afforestation only affected its southern part. Over parts of Scandinavia, afforestation actually
1488 resulted in a slight warming of about 0.5°C. In winter, the cooling effect of afforestation was only evident over the
1489 southern part of the Baltic Sea region, while deforestation had no effect on winter temperatures.

1490 Strandberg and Kjellström (2019) found relatively strong biophysical effects of afforestation or deforestation in
1491 Europe on daily maximum temperatures in summer. Deforestation markedly increased daily maximum
1492 temperatures over the entire Baltic Sea region (typically between 2 and 6°C), while afforestation lowered daily
1493 maximum temperatures in the southern part of the region by about 2 to 6°C and slightly increased over parts of
1494 Scandinavia. In contrast to its cooling effect on mean winter temperatures, afforestation lead to a warming of the
1495 daily minimum temperatures in the southern part of the Baltic Sea region in the range between 2 and 6°C.

1496 In a similar study with the Regional Model (REMO) RCM, Gálos et al. (2013) investigated the biophysical effects
1497 of afforestation in Europe on climate, and also compared these effects to the climatic changes expected from future
1498 global warming. Potential afforestation was implemented by specifying deciduous forest cover at all vegetated
1499 areas that were not covered by forests at the end of the 20th century, mainly in western, ~~c~~Central and ~~e~~Eastern
1500 Europe. Given the strong historical deforestation in ~~c~~Central Europe, there is potential for rather extensive
1501 afforestation in the southern part of the Baltic Sea region, but lesser potential in the northern part, i.e. Scandinavia
1502 and Finland. The results indicated a cooling effect of the re-established forests in boreal summer exceeding 0.3°C,
1503 mainly related to increased evapotranspiration from the trees, in combination with intensified fluxes of latent heat
1504 due to stronger vertical mixing ~~-(see above)~~. The stronger latent heat fluxes also enhanced precipitation by more
1505 than 10% in some regions. These effects of potential afforestation counteracted the projected future changes in
1506 climate, i.e. somewhat reduced the magnitude of the pronounced future warming and markedly reduced the future
1507 drying in the boreal summer. In some cases, such as in northern Germany, the enhanced precipitation was found
1508 to completely offset the drying effect of future warming. More recent results (Meier et al., 2021c) have used rain -
1509 gauge data to estimate precipitation changes induced by land cover change. Meier et al. (2021c) created a statistical
1510 model to show that reforestation of agricultural land can increase precipitation locally, especially in winter, and
1511 were able to separate the effects on both local and downwind precipitation regionally and seasonally. They also
1512 found that climate change induced summer precipitation reductions could be offset by reforestation, with a
1513 particularly strong effect in southwest Europe. ~~However, though~~ their analyses also indicate small precipitation

1514 increases in the Baltic Sea region, relative to a baseline scenario with no land cover change, consistent with the
1515 results of Gálos et al. (2013).

1516 In a more regionalized study, Gao et al. (2014) applied ~~an~~the-REMO RCM to investigate the biophysical effects
1517 of peatland forestation in Finland before (1920) and after drainage (2000s). In Finland, as in other northern
1518 European countries, vast areas of naturally tree-less or sparsely tree-covered peatland were drained for timber
1519 production in the second half of the 20th century. The total peatland area of Finland was estimated to be 9.7 million
1520 ha in the 1950s, but at the beginning of the 21th century the area of peatland drained for forestry was estimated to
1521 5.5 million ha. The authors found that the peatland forestation caused warming in spring, i.e. during the snow-melt
1522 season, and slight cooling in the growing season (May through October). The spring warming was mainly caused
1523 by decreased surface albedo and the cooling in the growing season by increased evapotranspiration.

1524 3.2.4 Cryosphere

1525 3.2.4.1 Snow

1526 In the Baltic Sea region, snow cover is an important feature that greatly affects cold season weather conditions. It
1527 is characterized by very high interannual and spatial variability. Snow cover is a sensitive indicator of climate
1528 change, and its variations are closely related to air temperature in many regions. General climate warming is
1529 expected to reduce snow cover. Several thaw periods now interrupt snow cover, making it less stable. Total winter
1530 snowfall in ~~n~~Northern Europe is projected to decrease, but is still expected to increase in mid-winter in the very
1531 coldest regions (Räsänen, 2016).

1532
1533 Previous climate change assessments demonstrated a number of snow-cover trends in recent decades in the Baltic
1534 Sea region (BACC Author Team, 2008; BACC II Author Team, 2015). A decrease in snow cover was observed
1535 in the south, while an increase in snow storage and duration of snow cover was detected in the north-east, and in
1536 the Scandinavian mountains. The spring snow melt has become earlier in most of the region. As a result, the spring
1537 maximum river discharge has become smaller and earlier, in many regions shifting from April to March.

1538
1539 Recent investigations confirm these results. Snow cover in the Northern Hemisphere has decreased since mid-20th
1540 century (IPCC, 2014a), as also shown by satellite measurements (Estilow et al., 2015). The largest decline in the
1541 extent of snow cover has been observed in March-April and also in summer. Using the satellite-based NOAA-
1542 CDR data for the period 1970–2019, it was shown that the annual snow cover fraction has reduced over most areas
1543 of the Northern Hemisphere by up to 2% decade⁻¹ (Zhu et al., 2021). Thereby, the annual snow cover area has
1544 reduced by $2 \times 10^5 \text{ km}^2 \text{ decade}^{-1}$.

1545
1546 In 1980–2008, snow-cover duration in ~~n~~Northern Europe decreased by about 3–7 days per decade, and the trend
1547 was significant at many stations (Peng et al., 2013). Most of the reduction happened in spring, with the end-date
1548 of snow cover five days earlier per decade, on average (Peng et al., 2013). Snow-cover variability over Europe is
1549 closely related to temperature fluctuations, which, in turn, are determined by large-scale atmospheric circulation
1550 during the cold season (Ye and Lau, 2017). A recent study of European snow-depth data in 1951–2017

Formatiert: Nicht unterstrichen, Schriftfarbe: Automatisch, Hochgestellt

1551 demonstrated an accelerated decrease after the 1980s (Fontrodona Bach et al., 2018), with an average decline,
 1552 excluding the coldest climates, of 12.2% per decade for mean snow depth and 11.4% per decade for its maximum.
 1553 A decreasing trend in snow density was detected in the eastern Baltic Sea region in 1966–2008 (Zhong et al.,
 1554 2014).

1555

1556 In Poland, rather large changes in snow cover parameters were found for 1952–2013 (Szwed et al., 2017). The
 1557 duration of snow cover decreased in almost the whole country but this change is mostly not statistically significant.
 1558 The total reduction in snow cover duration was 1–3 weeks over the 62 years, but the mean and maximum snow
 1559 depths did not change. The start date of snow cover has not changed, but the end date moved slightly earlier (Szwed
 1560 et al., 2017). A recent study found a statistically significant decreasing trends in snow cover duration as well as of
 1561 in snow depth, based on 40 Polish stations in 1967–2020 (Tomczyk et al., 2021). The trend values for the number
 1562 of days with snow cover were from -3.5 to -4.9 days per decade.

1563

1564 The snow-cover regime at 57 stations in the eastern Baltic Sea region (Lithuania, Latvia and Estonia) in 1961–
 1565 2015 was analyzed by Rimkus et al. (2018). The mean decrease in snow-cover duration was 3.3 days per decade,
 1566 and was statistically significant at 35% of the measuring sites, mostly in the southern part of the region. There
 1567 were no trends in maximum snow depth. An earlier study for Lithuania found similar results (Rimkus et al., 2014).

1568

1569 A detailed study of snow cover data at 22 stations in Estonia during the period 1950/51–2015/16 revealed
 1570 remarkable decreasing trends (Viru and Jaagus, 2020). Snow-cover duration decreased significantly at 16 stations,
 1571 and the mean decrease was 4 days per decade. Start dates for permanent snow cover had a non-significant tendency
 1572 to occur later. Permanent snow cover had a statistically significant trend to end earlier at almost all stations. There
 1573 were no overall trends in maximum snow depth in Estonia in 1951–2016 (Viru and Jaagus, 2020).

1574

1575 Significant decreases in snow depth parameters were found in Finland in recent decades (Aalto et al., 2016;
 1576 Luomaranta et al., 2019). Regional differences were substantial. In 1961–2014, the largest decrease in snow depth
 1577 occurred in the southern, western and central parts of Finland in late winter and early spring. In northern Finland,
 1578 a decrease in snow depth was most evident in spring, with no change in the winter months, even though the amount
 1579 of solid precipitation was found to increase in December–February (Luomaranta et al., 2019). Winter mean snow
 1580 depth (Jylhä et al., 2014) as well as the annual maximum snow depth (Lehtonen, 2015) has decreased significantly
 1581 at many stations in southern Finland. At the same time, the annual maximum snow depth has not changed in
 1582 Finnish Lapland (Lépy and Pasanen, 2017; Merkouriadi et al., 2017).

1583

1584 For the century 1909–2008, a general decrease was detected in many snow-cover parameters at three stations in
 1585 different parts of Finland (Irannezhad et al., 2016). A sharp decline in annual peak snow-water equivalent was
 1586 detected since 1959. The period of permanent snow cover shortened by 21–32 days per century. However, it
 1587 should be noted that the study of Irannezhad et al. (2016) was based on a temperature-index snowpack model using
 1588 daily temperature and precipitation as input, rather than directly on snow observations.

Formatiert: Schriftart: (Standard) Times New Roman, Nicht
 unterstrichen

1590 A decline in snow cover parameters was found also in Norway for 1961–2010 (Dyrrdal et al., 2012; Rizzi et al.,
1591 2017).

1592 3.2.4.2 Glaciers

1593 A recent basic inventory of Scandinavian glaciers is available through the Randolph Glacier Inventory (RGI
1594 Consortium, 2017; Pfeffer et al., 2014), a collection of digital outlines of the world's glaciers prepared to meet the
1595 needs of the Fifth IPCC Assessment Report (IPCC, 2014b; Vaughan et al., 2014). It has since been updated in
1596 support of the IPCC Special Report on the Ocean and Cryosphere in a Changing Climate (IPCC, 2019a; Hock et
1597 al., 2019). Of the 3,417 Scandinavian glaciers reported in the Randolph Glacier Inventory (v6.0), 365, with a
1598 combined area of c. 360 km², lie within the Baltic Sea Drainage Basin as defined by ~~(Vogt et al., 2007; Vogt et~~
1599 ~~al., 2008), all in the Scandinavian mountains). The combined glacier volume estimate (following Farinotti et al.,~~
1600 ~~2019) for the Scandinavian glaciers reported in Hock et al. (Hock et al., 2019) is 0.7 ± 0.2 mm global sea level~~
1601 ~~equivalent or 254 ± 72 Gt. The rate of mass loss for all Scandinavian glaciers (not only in the Baltic Sea Drainage~~
1602 ~~Basin) for the period 2006–2015 is 2 ± 1 Gt year⁻¹ corresponding to a negligible 0.01 ± 0.00 mm year⁻¹ of global~~
1603 ~~sea level rise equivalent, yet with potential importance for local streamflow (Hock et al., 2019; Zemp et al., 2019).~~
1604 With a 90–100% likelihood, atmospheric warming is the primary driver of glacier mass loss (Hock et al., 2019).

1605
1606 Most of the 365 glaciers in the Baltic Sea Drainage Basin are in Sweden (c. 72% by number, c. 75% by area), with
1607 the remaining ones in Norway. Both Sweden and Norway do nationally coordinated glacier monitoring, with the
1608 most recent results summarized in the Global Glacier Change Bulletin (GGCB) No. 3 (World Glacier Monitoring
1609 Service, ~~2020–2021~~; Zemp et al., 2020) as, among others, glacier mass balance changes. A glacier mass balance
1610 year usually covers the period from September 1st to August 31st in the subsequent year. The GGCB includes four
1611 Swedish glaciers, two of which (Rabots glaciär and Storglaciären in the Kebnekaise Massif, which has the world's
1612 longest continuous mass balance record, starting in 1945/46) are so-called reference glaciers. ~~This meansing~~
1613 ~~that~~ their dynamics are not dominated by non-climatically driven dynamics such as calving or surging, and that more
1614 than 30 years of ongoing measurements are available (Fig. 12). In Table ~~116~~, mass balances of the Swedish GGCB
1615 glaciers following World Glacier Monitoring Service (~~2020–2021~~) are summarized. None of the Norwegian
1616 GGCB glaciers are in the Baltic Sea drainage basin.

1617
1618 These with time increasingly negative mass balances coincide with globally increasing air temperatures, with the
1619 latest six years, 2015–2020, the warmest since instrumental recording began (World Meteorological Organization,
1620 2020). Regional and local deviations in mass loss from that ~~anticipated by expected from~~ long-term global warming
1621 ~~are is~~, however, expected. ~~For example, slightly positive mass balances were observed for Måmaglaciär,~~
1622 ~~Storglaciären and Riwojekna in 2016/2017 as a result of a cold summer. The slightly positive mass balances for~~
1623 ~~Måmaglaciär, Storglaciären and Riwojekna in 2016/2017 is the result of a cold summer, explained by the glacier~~
1624 ~~mass balance years starting on September 1, and that the summer months (June, July, August) included in the~~
1625 ~~2016/2017 balance are therefore June, July, and August 2017—a period during which average air temperature at~~
1626 ~~Tarfala Research Station was measured to 5.8°C, compared to 6.4°C and 7.4°C in the previous and subsequent~~

mass-balance-periods-(Swedish Infrastructure for Ecosystem Science, 2020; World Glacier Monitoring Service, 2020; 2021).

The ice summit of Kebnekaise Sydtopp (South Peak) lost its status as Sweden's highest in September 2019 and 2020 when due to melting of its ice-covered summit its elevation dropped below that of the rocky, non-ice-covered Kebnekaise Nordtopp (North Peak; Stockholm University - Department of Physical Geography, 2017, 2019, 2020).

3.2.4.3 Permafrost

The drainage basin of the Bothnian Bay is characterized by low mean annual air temperatures and includes boreal forest and mountain ecosystems as well as large peatland areas. In this region, permafrost — ground frozen for at least two consecutive years — exists in high alpine environments and in peatlands. While almost all Baltic Sea drainage basin permafrost is found in the Bothnian Bay catchment, some isolated occurrences of permafrost are found in the upper reaches of Alpine rivers further south (mainly Umeälven headwaters). Permafrost is a thermal state of ground, rock, soil or sediment, and occurs in regions with low mean annual air temperatures (MAAT). Temperature is the strongest control on permafrost, but thin winter snow depth also favors permafrost aggradation and stability. In alpine permafrost, insolation is an important factor. In the upper Torne River catchment, incoming shortwave summer radiation causes a difference in the altitude of alpine permafrost (permafrost $p > 0.8$) from 850 m a.s.l. on shaded slopes to 1100 m a.s.l. on south-facing slopes (Ridefelt et al., 2008). In lowland areas, permafrost is more likely to form and persist in peatlands, where the low thermal conductivity of peat insulates the ground from warm summer air (Seppälä, 1986). Mire complexes with palsas, elevated peat mounds with an ice-rich permafrost core, are the most common form of lowland permafrost in the Baltic Sea drainage basin (Luoto et al., 2004). Palsa mires in the Baltic Sea basin are predominantly found in regions with mean annual air temperature MAAT $< -3^{\circ}\text{C}$ and low mean annual precipitation (often < 450 mm), based on the 1961-1990 climate period (Fronzek et al., 2006).

Earlier maps of northern hemisphere permafrost extent showed relatively extensive Fennoscandian permafrost, especially in Alpine regions (Brown et al., 1997). Recent advances in permafrost modeling reveal a more nuanced picture, where permafrost is spatially patchy, persisting at high elevations and in lowland regions with low precipitation and large expanses of peat plateaus (Obu et al., 2019; Gislén et al., 2017). Consistent with observation of permafrost warming and thawing (Biskaborn et al., 2019), models indicate project substantial permafrost losses in recent decades. High resolution modelling (1 km pixels) driven by remotely sensed land surface temperature showed that c. 6,200 km² of permafrost in 1997, was reduced to 4,800 km² in 2018 (Obu et al., 2020). Most of this loss is modeled alpine permafrost (here defined as > 700 m a.s.l.) decreasing from 4,700 to 3,700 km², while lowland permafrost has decreased from 1,500 km² to 1,100 km² (Fig. 13).

3.2.4.4 Sea ice

Introduction

Sea ice is an essential indicator of climate change and variability in the Baltic Sea region. Not only does existence of sea ice indicate the general severity of the winter due to its close correlation with winter air temperature, but in addition parameters such as annual maximum sea-ice extent of the Baltic Sea (MIB), duration of ice season and maximum thickness of level ice have been monitored regularly in the Baltic Sea since the late 19th century. The MIB was even earlier observed.

The BACC II Author Team (2015) concluded that all sea-ice observations demonstrated large inter-annual variations, but with a long-term, statistically significant trend to milder ice conditions that is projected to continue in future (Fig. 14). In this section, we review recent ice-climate research in the Baltic Sea and provide updated figures on sea-ice trends and projections that largely confirm previous conclusions.

An important indicator of advancing climate change in the Baltic Sea region is that two of the latest ~~six~~five ice winters have been extremely mild (2015-2020). In winter 2015, the Bothnian Bay was never fully covered by ice (with a MIB of 51,000 km²), the first such extreme winter observed with certainty. The winter of 2020 was even milder, with a MIB of only 37,000 km², the lowest value in a time series that began in 1720.

Sea-ice conditions in the Baltic Sea

On recent average, the northern sea areas of the Baltic Sea are ice-covered every year, from December to May. During the mildest winters, only the Bothnian Bay (in a few years even only partially) and coastal zones of other basins are ice-covered. In the past, the entire Baltic Sea was ice-covered only in the most severe winters, e.g. 1940, 1942 and 1947 (Vihma and Haapala, 2009).

In the fast ice regions near the coast, sea ice grows by thermodynamic processes only. Maximum sea-ice thickness in the fast ice regions typically amounts to 40 – 70 cm in the Bothnian Bay and 10 – 40 cm in the Bothnian Sea, Gulf of Finland and Gulf of Riga. Ice thickness is regularly monitored at tens of fast ice sites.

Observations of sea-ice thickness in drift ice are much more limited. A recent study combined all airborne electromagnetic ice thickness measurements in the Bothnian Bay and derived the first estimate of basin-scale ice thickness distribution in the Baltic Sea (Fig. 15; Ronkainen et al., 2018). An important finding of that study was that mean ice thickness in drift ice regions is greater than the thickness of fast ice, and also greater than the ice thickness indicated on the ice charts. As expected, the data showed large inter-annual variability, but temporal and spatial coverage was not sufficient for conclusion on changes in drift ice thickness.

Individual ice ridges caused by compression and shearing of ice drift can be 30 meters thick. The largest gradients in ice motion are found in the coastal zone (Leppäranta et al., 2012), where mean ice thickness over several km² can be 1-3 meters (Ronkainen et al., 2018).

In some circumstances, sea ice can accumulate towards the coast and cause spectacular on-shore ridges or ride-up of several hundred meters from shore to land, causing damage to built structures (Leppäranta, 2013). Such events

have been observed in exposed coastal regions where the stability of fast ice can be overcome by combinations of storms, currents and water level (Leppäranta, 2013). In the Bothnian Bay, such events can occur regardless of the severity of ice seasons. In the southern Baltic Sea, on-shore ice has been common during severe winters. During the last ten years, such events were observed in 2010, 2011, 2012 and 2019 (Girjatowicz and Łabuz, 2020).

Observed changes

Long-term changes of the MIB (Seinä and Palosuo, 1996; Niskanen et al., 2009) are shown in Figure 14. The trend of the MIB during the last 100 years (1921-2020) is -6,400 km² per decade. This is almost twice the trend reported by the BACC II Author Team (2015), based on the period 1910-2011. Since 1987 no severe ~~and no extremely severe~~ ice winters and since 2012 only average, mild or extremely mild winters have been observed. The latter sea-ice conditions explain the accelerated trend after 2011.

The recent 30-year period (1991-2020) is definitely the mildest since 1720 (Uotila et al., 2015). The probability distribution of the MIB has shifted towards ~~low values~~~~zero~~, with severe winters very rare (Fig. 14). The 30-year mean MIB is now 139,000 ~~+0³~~-km² and the winter 2021 with a MIB of 127,000 km² on 15 February 2021 (Jouni Vainio, FMI, personal communication) was close to this mean. During the second mildest 30-year period (1909-1938), the average MIB was 184,000 ~~+0³~~-km² and during the last 100 years it was 182,000 ~~+0³~~-km². The shape of the MIB probability distribution has changed, also indicating a change in sea-ice extremes. According to the ice season classification (Seinä and Palosuo, 1996), the recent 30-year period includes only one ~~severe or extremely severe~~ ice winter (2011) and 13 mild ~~or extremely mild~~ ice winters.

Present ice conditions differ from the past to the extent that Rjazin and Pärn (2020) even suggested defining this change as a regime shift. They analyzed changes in sea-ice extent and air temperature in the Baltic Sea in 1982 – 2016, using a method of splitting the time series in two and concluded that a regime shift towards milder ice conditions occurred in 2006-2007.

Other studies complement these and BACC II conclusions. Kiani et al. (2018) examined the influence of atmospheric changes on ice roads between Oulu and Hailuoto. They used air temperature data to calculate freezing and thawing degree days and found that freezing degree days decreased and thawing degree days increased significantly during 1974 – 2009. As a consequence, the ice road season started later and ended earlier.

Merkouriadi and Leppäranta (2014) analyzed ice thickness and freezing and breakup dates collected at Tvärminne Zoological Station, at the entrance to the Gulf of Finland. They found a decrease of almost 30 days in the ice-covered period and a reduction of 8 cm in maximum annual ice thickness in the last 40 years. Laakso et al. (2018) used observations from the Utö Atmospheric and Marine Research station during 1914-2016 and concluded that the length of the ice season has decreased from 10-70 days before 1988 to 0-35 days after 1988 in the northern Baltic Sea proper. Figures 16 and 17 show level-ice thickness at Kemi and Loviisa and the length of the ice season at Kemi, Loviisa and Utö, ~~respectively~~. All graphs show statistically significant decreasing trends, except level-

ice thickness at station Kemi, which was probably be-influenced by snow cover or changes in measurement location.

3.2.4.5 Lake ice

The recent change in ice phenology is probably the single most important climatically induced alteration in lake environments within the Baltic Sea catchment. New literature demonstrates almost unanimously significant changes towards earlier ice break-up, later freeze-up, and shorter duration of ice cover across the Baltic Sea catchment, apart from the coldest climate regime in Lapland. The available centennial data indicate that the ice-cover duration has decreased by several days per century, whereas the intensified warming in recent decades has produced a similar change per decade (Efremova et al., 2013; Filazzola et al., 2020; Kļaviņš et al., 2016; Knoll et al., 2019; Korhonen, 2019; Lopez et al., 2019; Nöges and Nöges, 2014; O'Reilly et al., 2015; Ptak et al., 2020; Sharma et al., 2016; Sharma et al., 2020; Wrzeński et al., 2015). Some lakes have, however, responded only weakly to the warming trend, such as Lake Peipsi in Estonia, probably due to increasing snowfall. In individual years, a positive wintertime NAO seems to be an important factor causing a short ice-cover duration. Among the main properties that affect the ice cover of individual lakes are size, depth, and shoreline complexity.

3.2.5 Ocean and marine sediments

3.2.5.1 Water temperature

The main driver of annual mean water temperature variations and long-term changes is air temperature (Meier et al., 2019d; 2019c; Kniebusch et al., 2019a; Duthéil et al., 2021). Baltic Sea water temperature has risen fastest at the sea surface (Meier et al., 2021a). With time the heat spreads downward through different processes, such as lateral inflows, vertical down-welling and diffusion, and eventually the whole water column warms up, with smallest trends in the cold intermediate layer between the thermo- and halocline (Meier et al., 2021a).

Since the 1980s, marginal seas around the globe have warmed faster than the global ocean (Belkin, 2009), and the Baltic Sea has warmed the most (Belkin, 2009). Climate change and decadal variability led to an annual mean, area averaged increase in Baltic Sea SST of $+0.59^{\circ}\text{C decade}^{-1}$ for 1990-2018 (Siegel and Gerth, 2019) and of $+0.5^{\circ}\text{C decade}^{-1}$ for 1982-2013 (Stramska and Białogrodzka, 2015). Both figures were derived from satellite data. In accordance with earlier investigations (BACC Author Team, 2008; BACC II Author Team, 2015), SST variability in winter can be linked to the NAO (Stramska and Białogrodzka, 2015). However, the spatial maps of SST trends by Stramska and Białogrodzka (2015) differ from those by Lehmann et al. (2011), perhaps because of the differing horizontal resolution of the satellite data products. Linear trends for the Baltic Sea during 1982-2012 of $0.41^{\circ}\text{C decade}^{-1}$ are slightly larger than $0.37^{\circ}\text{C decade}^{-1}$ for the North Sea (Høyer and Karagali, 2016).

Using monitoring data, Liblik and Lips (2019) found that the upper layer has warmed by $0.3\text{--}0.6^{\circ}\text{C decade}^{-1}$ and the sub-halocline deep layer by $0.4\text{--}0.6^{\circ}\text{C decade}^{-1}$ in most of the Baltic Sea during 1982-2016. The total warming in the whole Baltic Sea at all monitoring stations was 1.07°C over 35 years, approximately twice that of the upper 100 m in the Atlantic Ocean.

During 1856–2005, the reconstructed Baltic Sea average, annual mean SST increased by 0.03 and 0.06°C decade⁻¹ in the northeastern and southwestern areas, respectively (Kniebusch et al., 2019a). The largest SST increase trends were found in the summer season in the northern Baltic Sea (Bothnian Bay). Bottom water temperature trends were smaller than SST trends, with the largest increase in the Bornholm Basin. Independent monitoring data support the results of the long-term reconstruction (Figs. 18 and 19), see also Meier et al. (2019c; 2019d). The largest SST warming occurred in summer (May to September), while trends in winter were smaller (Kniebusch et al., 2019a; Liblik and Lips, 2019).

During the more recent period of 1978–2007, the annual mean SST trend was tenfold higher, with a mean of 0.4°C decade⁻¹ (Kniebusch et al., 2019a). Trends increased more in the northeastern areas than in the southwestern, and exceeded the contemporary trends in air temperature. See also MARNET station data at Darss Sill and Arkona Deep in the southwestern Baltic Sea (Fig. 20).

The seasonal ice cover clearly plays an important role in the Baltic Sea by decoupling the ocean and the atmosphere in winter and spring. Hence, the large trends in air temperature in winter were not reflected by the SST trends because the air temperature was still below the freezing point. During the melting period, the ice-albedo feedback led to larger trends in SST than during the ice-covered period, because of a prolonged warming period of sea-water.

It has been suggested that the accelerated warming in 1982–2006 might partly be explained by a dominance of the positive phase of the AMO (Kniebusch et al., 2019a). Furthermore, historical eutrophication re-distributed the heat in the ocean by warming the surface layer more than the underlying layers, in particular during spring and summer, because the increased water turbidity caused an enhanced absorption of sunlight close to the sea surface. However, modeling studies suggest that the historical eutrophication had no impact on SST trends (Löptien and Meier, 2011).

The summer of 2018 was the warmest on instrumental record in Europe, and also the warmest summer in the past 30 years in the southern half of the Baltic Sea (Naumann et al., 2019), with surface-water temperatures 4–5°C above the 1990–2018 long-term mean. This heat wave was also observed in the bottom temperatures (Humborg et al., 2019). However, systematic studies on changes in Baltic Sea marine heat waves are not available.

3.2.5.2 Salinity and saltwater inflows

During the last decade, many new insights have been gained about the salt balance of the Baltic Sea and the dynamics of inflow and mixing processes. The Major Baltic Inflow (MBI) in December 2014, in particular, triggered new investigations and is by far the most intensively observed and modeled inflow event. Pathways and timing of the inflowing water were tracked by observations (Mohrholz et al., 2015), and numerical modelling (Gräwe et al., 2015) could reproduce the salt mass and volume of the inflow, calculated from observations. The inflow was found to be barotropic (pressure) controlled in the Danish straits but dominated by baroclinic (density stratification) processes on the further-pathway farther into the Baltic proper. At the Bornholm Gat and the Ślupsk Furrow the water exchange showed the clear two-layer flow pattern of an estuarine circulation. The inflow-related

1815 studies were underpinned by theoretical work based on the famous Knudsen Relation (Knudsen, 1900) for
 1816 estuarine exchange flow, and its extension to total exchange flow by Burchard et al. (2018). The contribution of
 1817 the inflowing saline water to the spatial distribution of salt and the total salt budget depends essentially on mixing
 1818 with ambient brackish waters. In the course of the inflow path, the character of the mixing process between the
 1819 deep salty layer and the brackish water above changes with increasing depth and decreasing current velocities. In
 1820 the entrance area the mixing is dominated by entrainment of brackish water into the eastward spreading saline
 1821 bottom water, due to turbulence generated by shear instability. The sills between the consecutive Baltic basins are
 1822 particular mixing hotspots (Neumann et al., 2017). In the deeper basins of the Baltic proper, boundary mixing
 1823 driven by the interaction of currents and internal waves with the topography, and mixing processes at sill overflows
 1824 contribute to the upward salinity flux (Reissmann et al., 2009). Mixing in the eastern Gotland Basin was
 1825 investigated during the Baltic Sea Tracer Experiment (BATRE). Using an inert tracer gas, the basin-scale vertical
 1826 diffusivities were estimated to $10^{-5} \text{ m}^2 \text{ s}^{-1}$, whereas the locally inferred diffusivities in the basins' interior were one
 1827 order of magnitude lower (Holtermann et al., 2012). This finding holds also for the inflow of saline water in course
 1828 of an MBI (Holtermann et al., 2017). The interior mixing is often controlled by double diffusive convection that
 1829 leads to a typical stair-case-like vertical stratification structure (Umlauf et al., 2018). The crucial role of boundary
 1830 mixing at the basin rim was confirmed by Holtermann and Umlauf (2012), and Lappe and Umlauf (2016). Both
 1831 studies identified near-boundary turbulence as the key processes for basin-scale mixing. Main energy sources for
 1832 boundary mixing are basin-scale topographic waves, deep rim currents, and near-inertial waves.

1833
 1834 The temporal statistics of barotropic saline inflows was reviewed by Mohrholz (2018). In contrast to earlier
 1835 investigations he found no long-term trend in inflow frequency, but a pronounced multidecadal variability of 25
 1836 to 30 years. Lehmann and Post (2015) and Lehmann et al. (2017), who studied the frequency and intensity of large
 1837 volume changes in the Baltic Sea due to inflows, likewise could not find a long-term trend. The distinction between
 1838 MBIs and smaller inflows is artificial and does not correspond to the frequency distribution of the inflows, which
 1839 shows an exponential decrease in frequency with increasing inflow intensity (Mohrholz, 2018). The classical MBIs
 1840 are only responsible for about 20% of the total salt input, while the rest is accounted for by medium and small
 1841 inflows with much less pronounced interannual variability.

1842
 1843 Paleoclimate simulations covering nearly the recent millennium (Schimanke and Meier, 2016) have provided new
 1844 insights into the long-term behavior of the mean salinity of the Baltic Sea. In accordance with previous historical
 1845 reconstruction studies (Schimanke and Meier, 2016; Meier and Kauker, 2003), Schimanke and Meier (2016)
 1846 identified river discharge, net precipitation and zonal winds as main drivers of the decadal variability in Baltic Sea
 1847 salinity. However, their relative contributions are not constant. Extreme periods with strong salinity decrease for
 1848 about 10 years occurred once per century. Thus, the long stagnation period from 1976 to 1992 was obviously a
 1849 rare but natural event, although its extreme duration might be caused by anthropogenic effects. The Baltic Sea
 1850 salinity also has a natural centennial variability. Based on the same numerical simulations, Börgel et al. (2018)
 1851 could show a strong coherence between the AMO climate mode and river runoff on timescales between 60 and
 1852 180 years. Accordingly, the Baltic Sea salinity and the AMO are correlated, probably due to the dominating impact
 1853 of river discharge on salinity. The river runoff leads salinity changes by about 20 years during the entire modeling

period of 850 years. Schimanke and Meier (2016) reported a similar lag of 15 years between river runoff and Baltic Sea salinity.

According to model results, multidecadal variations in runoff (Gailiusis et al., 2011; Meier et al., 2019d) explain about half the long-term variability of volume-averaged Baltic Sea salinity (Meier and Kauker, 2003). Radtke et al. (2020) found that the direct dilution effect was only responsible for about one fourth of the multidecadal variability and proposed a link between river runoff and inflow activity. Furthermore, they found that the influence of vertical turbulent mixing is small. Salt-water inflows contribute to the multidecadal salinity variability, in particular for the bottom layer salinity. The positive trend of river runoff in the northern catchment area led to a significant increase in the North-South salinity gradient in the Baltic Sea surface water layer (Kniebusch et al., 2019b). The causal relationship between the trends in runoff and horizontal salinity gradient was confirmed by numerically sensitivity experiments. Additionally, their model ~~simulationsbased study~~ revealed a multidecadal oscillation of salinity, river runoff and saltwater inflows of about 30 years, consistent with the long term observations.

From observations during 1982–2016, Liblik and Lips (2019) detected decreasing surface (see also Vuorinen et al., 2015) and increasing bottom salinities, but no long term trend in the total salt budget were found (cf. Fig. 21). Both temperature and salinity contribute to a strengthening of the vertical stratification. Enhanced freshwater fluxes combined with higher deep water salinities intensify the vertical density gradient throughout the year.

3.2.5.3 Stratification and overturning circulation

A direct consequence of increasing stratification is that mixing ~~weakens~~ between well ventilated surface waters and badly ventilated deep waters weakens, making the Baltic Sea vulnerable to deoxygenation of bottom waters (Conley et al., 2002). An increase in seasonal thermal stratification (e.g. Gröger et al., 2019) can additionally lower the vertical nutrient transport from deeper layers to the euphotic zone, thereby limiting nutrient supply and potentially affecting algal and cyanobacterial blooms, at least at the species level. The latter potential effect has not yet been thoroughly investigated. However, the hypothesis was supported by the results of Lips and Lips (2008) who found a correlation between cyanobacteria bloom intensity in the Gulf of Finland and the frequency of upwelling events along both coasts.

Since the start of regular salinity measurements at the end of the 19th century, the haline stratification has been dominated by sporadic inflows from the adjacent North Sea and variations in river discharge (Fig. 22). While no long-term trends could be demonstrated in Baltic Sea salinity during 1921–2004 (Kniebusch et al., 2019b) or in halocline depth during 1961–2007 (Väli et al., 2013), a trend towards increased horizontal salinity difference between the northern and southern Baltic Sea was found during 1921–2004 (Section 3.2.5.2~~Kniebusch et al., 2019b~~). Modeling studies by Kniebusch et al. (2019b) attributed this trend to increased river runoff from the northernmost catchment area. Furthermore, vertical sStratification increased in most of the Baltic Sea during 1982–2016, with the seasonal thermocline strengthening by 0.33–0.39 kg m⁻³ and the perennial halocline by 0.70–0.88 kg m⁻³ (Liblik and Lips, 2019).

1892
 1893 Sensitivity studies with a numerical model suggest~~ed~~ that the basin-wide overturning circulation will decrease if
 1894 the climate warms or when river runoff increases, but will tend to increase if global sea level rises (Placke et al.,
 1895 2021). However, historical multidecadal variations of the overturning circulation are mainly wind-driven.
 1896 Multidecadal variations in neither river runoff nor saltwater inflow had an impact, according to Placke et al. (2021).

1897 3.2.5.4 Sea level

1898 For the era of continuously operated satellite altimetry, absolute mean sea level (relative to the reference geoid)
 1899 increased in the Baltic Sea. Available estimates vary depending on the exact period considered, but are broadly
 1900 consistent with or slightly above the global average (3-4 mm year⁻¹; Oppenheimer et al., 2019; Nerem et al., 2018).
 1901 For the period 1992-2012, Stramska and Chudziak (2013) estimated an increase of 3.3 mm year⁻¹, and for the
 1902 period 1993-2015 Madsen et al. (2019a) an increase of 4 mm year⁻¹ in the Baltic Sea absolute mean sea level. For
 1903 the period 1886/1889-2018, the analysis of Swedish mareograph data suggest a sea level rise of about 1-2 mm
 1904 year⁻¹ (Figs. 23 and 24). Passaro et al. (2021) showed that the increase is not uniform across the Baltic Sea but
 1905 varies between about 2 mm year⁻¹ in the western Baltic Sea and more than 5 mm year⁻¹ in the Gulf of Bothnia for
 1906 the period 1995-2019. The acceleration of sea level rise in the Baltic Sea was studied by Hünicke and Zorita
 1907 (2016). They found that present acceleration is small and could only be detected through spatial averaging of
 1908 observations.

1909
 1910 Sea level changes relative to the coast are more complex, since land is rising in the northern Baltic Sea, by up to
 1911 about 108 mm year⁻¹, and sinking in the southern Baltic Sea, by about 0.5+ mm year⁻¹, relative to the Earth's centre
 1912 of mass (Hünicke et al., 2015; Groh et al., 2017). In addition to the global mechanisms (thermal expansion due to
 1913 warming and land-ice melting), sea level changes in the Baltic Sea are also affected by the changes in atmospheric
 1914 circulation, water inflow from the North Sea, and changes in the freshwater budget (river runoff, precipitation and
 1915 evaporation). Precipitation and river runoff are linked to westerly winds and affect salinity and the salinity gradient
 1916 across the Baltic Sea (Kniebusch et al., 2019b) and thus the sea-level height and its gradient. Stronger than normal
 1917 westerly winds are associated with increased transports across the Danish straits which leads to an increase in
 1918 Baltic mean sea level. The correlations between sea-level height and westerly wind are higher in the eastern and
 1919 northern parts and lower in the southern and western parts of the Baltic Sea. Westerly winds in the region became
 1920 more intense from the 1960s until the early 1990s, but have weakened somewhat thereafter (Feser et al., 2015).
 1921 Over longer periods, no significant long-term trend is detected (Feser et al., 2015).

1922
 1923 The Baltic mean sea level shows a pronounced seasonal cycle with a minimum in spring and maxima in late
 1924 summer (in 1900-1930) or winter (in 1970-1998). According to Hünicke and Zorita (2008), the amplitude of the
 1925 seasonal cycle increased over the 20th century. Other authors found different periods without systematic long-term
 1926 trends (Barbosa and Donner, 2016) or even regional decreases (Männikus et al., 2020).

1927
 1928 ~~Baltic sea level extremes are caused by strong atmospheric cyclones, or more seldom by wind-induced~~
 1929 ~~meteotsunamis (Pellikka et al., 2020) and seiches (Neumann, 1941; Wübbler and Krauss, 1979). Cyclones are~~

associated with strong winds that cause coastal storm surges over one or two days, and if their pathway is aligned along the west-east direction, cyclones may also increase the total volume of the Baltic Sea over one week by pushing in water masses from the North Sea into the Baltic Sea. Coastal storm surges can then reach 20 cm above the spatially averaged level (Weisse and Weidemann, 2017), in the eastern Gulf of Finland in Neva Bay even more. Extreme sea levels over a predefined threshold become more frequent with rising mean sea level (Pindsoo and Soomere, 2020). In addition, model results and analysis of observations indicate that atmospheric forcing is responsible for the long-term increases in storm surges in some localized areas of the eastern Baltic Sea (Ribeiro et al., 2014). The presence of sea-ice impedes the development of extreme sea levels by shielding the ocean surface from forcing by the wind, and coastal ice protects the coast from erosion by extreme sea levels. Baltic sea level extremes are caused by strong atmospheric cyclones sometimes in interaction with seiches (Neumann, 1941; Wübber and Krauss, 1979; Matthäus and Franck, 1992; Wolski et al., 2014; Soomere and Pindsoo, 2016; Girjatowicz, 2004; Leppäranta, 2013; Orviku et al., 2011), or more seldom and on smaller scales by wind-induced meteotsunamis (Pellikka et al., 2020). Cyclones associated with strong onshore winds can cause coastal storm surges over several hours to about one day (Wolski and Wisniewski, 2020). If the pathway of the cyclones is aligned along the west-east direction, they may also increase the volume of water in the Baltic Sea. Major inflow events are associated with volumes changes corresponding to an increase in Baltic sea level of about 24 cm (Matthäus and Franck, 1992) at time scales of about 10 days or longer (Soomere and Pindsoo, 2016). Under such conditions, even moderate wind and wind surges may lead to coastal sea level extremes (e.g., Weiss and Weidemann, 2017). Extreme sea levels over a predefined threshold become more frequent with rising mean sea level (Pindsoo and Soomere, 2020). In addition, model results and analysis of observations indicate that changes in atmospheric forcing are responsible for the long-term increases in storm surges in some localized areas of the eastern Baltic Sea (Ribeiro et al., 2014). The presence of sea-ice impedes the development of extreme sea levels by shielding the ocean surface from forcing by the wind. Coastal ice protects the coast from erosion by extreme sea levels, but pileup of ice and ice growth down to the sea bottom can also result in transport of bottom sediment when the ice drifts out to sea (Girjatowicz, 2004; Leppäranta, 2013; Orviku et al., 2011).

Storm surges caused by strong onshore winds represent a substantial hazard for the low-lying parts of the Baltic Sea coast, in particular, the southwestern parts (Wolski et al., 2014), the Gulf of Finland (e.g. Suursaar and Sooäär, 2016), the Gulf of Riga (e.g. Männikus et al., 2019), and the Gulf of Bothnia (Averkiev and Klevanny, 2010). Highest surges were reported for the Gulf of Finland (about 4 m in 1824 in St. Petersburg) and the western Baltic Sea (more than 3 m in 1871, Wolski and Wisniewski, 2020). For the Gulf of Riga and the western Baltic Sea values around 2 and 1-1.5 m are frequent, respectively (Wolski and Wisniewski, 2020). Hundred-year storm surges are higher (up to 2.4 m) at the inner end of the basins, farthest away from the Baltic proper, than in the center of the Baltic Sea (up to 1.2 m). No consistent long-term trend for an increase in extreme sea levels relative to the mean sea level of the Baltic Sea has been found, in agreement with earlier assessments (BACC II Author Team, 2015). This finding is supported by paleoclimate model studies that show no influence on extreme sea levels in the North Sea in warmer climate periods compared to colder periods (Lang and Mikolajewicz, 2019), and by recent studies of sea level records that suggest a pronounced decadal to multidecadal variability in storm surges relative to the mean sea level (Marcos et al., 2015; Marcos and Woodworth, 2017; Wahl and Chambers, 2016). Although

Formatiert: Schriftart: Nicht Kursiv, Nicht unterstrichen, Schriftfarbe: Automatisch

Formatiert: Schriftart: Nicht Kursiv, Nicht unterstrichen, Schriftfarbe: Automatisch

Formatiert: Schriftart: Nicht Kursiv, Nicht unterstrichen, Schriftfarbe: Automatisch

Formatiert: Schriftart:

1969 ~~Ribeiro et al.~~ (2014) argued for an increase in annual maximum sea level during 1916–2005, especially in the
 1970 northern Baltic Sea, and attributed the trends to changes in wind, these results were likely affected by the long-
 1971 term internal variability. Furthermore, extreme sea level in the Gulf of Finland, especially in Neva Bay, are very
 1972 sensitive to the position of storm tracks (Suursaar and Sooäär, 2007).

1973 3.2.5.5 Waves

1974 Instrumental wave measurements in the Baltic Sea have been made since the 1970s, first as measurement
 1975 campaigns and since the 1990s as continuous monitoring (e.g. Broman et al., 2006; Tuomi et al., 2011). As the
 1976 spatial coverage of wave measurements is still quite sparse, and there are long-term data from few locations only,
 1977 wave hindcasts have become a valuable tool for estimating the Baltic Sea wave climate (e.g. Björkqvist et al.,
 1978 2018). Lately, satellite altimeter measurements have been used to estimate the changes in the Baltic Sea wave
 1979 climate (Kudryavtseva and Soomere, 2017).

1980
 1981 Hindcast studies (e.g. Räämet and Soomere, 2010; Tuomi et al., 2011; Björkqvist et al., 2020) are in good
 1982 agreement with wave measurements, and estimate the annual mean significant wave height (SWH) ~~at 0.5–1.5 m~~ in
 1983 the open sea areas of the Baltic Sea at 0.5–1.5 m. Wave growth in the Baltic Sea is hampered by the shape and
 1984 small size of the basins. The highest mean values are recorded in the Baltic proper, with the longest and widest
 1985 fetches. The gulfs have less severe wave climates (Björkqvist et al., 2020). In addition, wave growth in the northern
 1986 Baltic Sea in winter is limited by the seasonal ice cover, leading to considerably lower mean and maximum values
 1987 of SWH in the northernmost Gulf of Bothnia and the easternmost Gulf of Finland.

1988
 1989 Although the measurement and hindcast periods have so far been quite short, some studies have also analyzed
 1990 trends in Baltic Sea SWH. For example, Soomere and Räämet (2011) and Kudryavtseva and Soomere (2017)
 1991 suggested ed an increasing trend in SWH since the 1990s, but results are site-specific and so far rather inconclusive.

1992
 1993 The seasonal wave climate is driven by the wind climate. The highest mean and maximum values of SWH are
 1994 reached in autumn and winter, while summer typically has the mildest wave climate. In sub-basins with long ice
 1995 season and large ice extent, such as the Bothnian Bay, the seasonal variation in the SWH is slightly different, since
 1996 waves are damped by the ice.

1997
 1998 So far the ~~n~~Northern Baltic proper ~~(NBP)~~ holds the record measured value of Baltic Sea SWH. In December 2004
 1999 a SWH of 8.2 m was measured by the northern Baltic proper~~NBP~~ wave buoy, with a highest individual wave of c.
 2000 14 m (Tuomi et al., 2011; Björkqvist et al., 2018). As the spatial coverage of the wave measurements has increased
 2001 and milder ice winters have allowed late autumn and even winter measurements also in the northern parts of the
 2002 Baltic Sea, 8 m SWH has been measured also in the Bothnian Sea, in January 2019. Björkqvist et al. (2020)
 2003 estimated a return period of 104 years for this event in the present climate. Hindcast statistics have suggested that
 2004 even higher maximum values between 9.5 – 10.5 m may occur in areas and times for which wave buoy
 2005 measurements are not available (Soomere et al., 2008; Tuomi et al., 2011; Björkqvist et al., 2018).

3.2.5.6 Sedimentation and coastal erosion

The Glacial Isostatic Adjustment and eustatic sea level change impose a first-order control on Baltic Sea coastal landscape change (Harff et al., 2007). In the subsiding southern Baltic Sea region, wind-driven coastal currents and waves are the major drivers for erosion and sedimentation, especially along the sandy and clayey sections of sandy beaches, dunes and soft moraine cliffs (Zhang et al., 2015; Harff et al., 2017).

Owing to spatial variation in aero- and hydrodynamic conditions (winds, waves and longshore currents) and underlying geological structure (lithology, sediment composition), a diversity of morphological patterns have developed along the Baltic Sea coast. Because of the dominant westerly winds that blow 60% of the year (Zhang et al., 2011) and a sheltering by land in the west, wind-waves are larger in the south-eastern Baltic Sea than in the south-western. As a result, sediment transport and dune development are more active and dynamic along the south-eastern coast. Thus, the largest coastal dunes are found along the Polish coast, with wave length >100 m and height > 20 m (Ludwig, 2017), while dunes along the German coast normally have wave length less than 60 m and height below 6 m (e.g. Lampe and Lampe, 2018). Under conditions favorable for wind-driven sand accumulation along sandy Baltic Sea coasts, a typical cross-shore profile features one or several foredune ridges, generally with a height of between 3 and 12 m above the mean sea level (Zhang et al., 2015; Łabuz et al., 2018). At the backshore behind the established foredune ridges, drifting or stabilized dunes in transgressive forms, mainly parabolic or barchanoid types, are commonly developed. The source of sediment for dune development includes fluvio-glacial sands from eroded cliffs, river-discharged sands, and older eroded dunes (Łabuz, 2015).

Because the wind-wave energy increases from west to east, so does erosion along the Baltic Sea coast. The mean annual erosional rate of the soft moraine cliffs and sandy dunes along the north side of the southwestern Baltic Sea coast (southern Sweden and Denmark) is 1-2 m year^{-1} , larger than 0.4-1 m year^{-1} along the south side of the southwestern Baltic Sea coast (Germany). Erosion along the southern Baltic Sea coast increases eastward, with a mean annual rate of 0.5-1.5 m year^{-1} in Poland, and 0.5-4 m year^{-1} in Latvia, Lithuania and Russia (BACC II Author Team, 2015). Severe coastal erosion in the Baltic Sea region is often caused by storms. The maximum storm-induced erosion increases eastward from 2-3 m year^{-1} at the southwestern Baltic Sea coast (southern Sweden, Denmark and Germany) to 3-6 m year^{-1} along the Polish coast, and ~10 m year^{-1} along the coast of Lithuania and Russia (Kaliningrad). Each storm can erode soft Latvian cliffs 3–6 m, with a maximum of up to 20–30 m locally. Many sandy beaches along the Gulf of Finland have recently been severely damaged by frequent storm surges, despite extensive protective measures (BACC II Author Team, 2015).

The prevailing wind-wave pattern controls the spatial variations of not only coastline change rates but also submarine morphologies (Deng et al., 2019). In the southwestern Baltic Sea coast where wind-wave energy is relatively small, nearshore submarine morphology is generally featured by smooth transition from beach-dunes/moraine cliffs to deeper water perturbed by one or two longshore bars. Morphological perturbations (e.g. the number and amplitude of longshore bars and rip current channels) become increasingly larger toward the east due to increased wind-wave energy. The wave incidence angle also impacts the nearshore submarine morphology, with in general a smaller angle leading to a larger morphological heterogeneity, i.e. a larger amplitude of

2045 perturbations. The amplitude of nearshore morphological perturbations may significantly affect coastal erosion
2046 because rip currents act as efficient conduit for offshore sediment transport, despite that they only occur
2047 sporadically along the Baltic Sea coast (Schönhofer and Dudkowska, 2021).

2048
2049 The sediments eroded from the soft moraine cliffs are composed of grain sizes from clay to pebbles. The fine-
2050 grained sediments are mostly transported outwards to the deeper seafloor (i.e. the Baltic Sea basins), either
2051 suspended in the water column or in a concentrated benthic fluffy layer (Emeis et al., 2002). Eroded fine-grained
2052 sediments from the moraine cliffs have been found to contribute to a major portion (40-70%) of the Holocene
2053 deposits in the muddy Baltic Sea basins (Porz et al., 2021). Coarser material, such as sands, stays mostly nearshore,
2054 partly in the water, partly transported onto the beach and the dunes (Deng et al., 2014; Zhang et al., 2015).

2055 3.2.5.7 Marine carbonate system and biogeochemistry

2056 Studies summarized in BACC II showed that nitrate and phosphate concentrations in winter surface water of the
2057 Baltic proper had increased by a factor of about three in the second half of the twentieth century, and reached a
2058 peak between 1980 and 1990. This change was consistent with the enhanced nutrients inputs to the Baltic Sea and
2059 caused eutrophication in the affected basin. Based on the available CO₂ system data, it has been estimated that the
2060 net ecosystem production in the Baltic Sea has increased since the 1930s by a factor of about 2.5. Increase in net
2061 ecosystem production and poor ventilation of the deep water layers due to the permanent stratification of the water
2062 column caused significant expansion of the anoxic and hypoxic areas in the Baltic Sea. Since the 1980s, nutrients
2063 inputs to the Baltic Sea have decreased. This led to a decrease in winter surface-water nitrate concentrations. No
2064 decrease was, however, observed for phosphate, due to the long residence time of P in the Baltic Sea and reduced
2065 P storage in oxygen-deficient sediments by binding to Fe-oxyhydroxides.

2066 3.2.5.7.1 Oxygen and nutrients

2067 In the Baltic Sea, hypoxia (oxygen deficiency) and even anoxia has expanded considerably since the first oxygen
2068 measurements in 1898 (Gustafsson et al., 2012; Carstensen et al., 2014a; Fig. 25). In 2016, the maximum hypoxia
2069 area was about 70,000 km², almost the combined area of Belgium and the Netherlands, whereas it was presumably
2070 very small or even absent 150 years ago (Carstensen et al., 2014b; Carstensen et al., 2014a; Meier et al., 2019c;
2071 2019d). Hypoxia was caused mainly by increasing land nutrient inputs and atmospheric deposition that led to
2072 eutrophication of the Baltic Sea (Andersen et al., 2017; Savchuk, 2018). The impacts of other drivers like observed
2073 warming and eustatic sea level rise were smaller, but still important (Carstensen et al., 2014a; Meier et al., 2019c).
2074 On annual to decadal time scales, halocline variations also had considerable influence on the hypoxic area (Conley
2075 et al., 2002; Väli et al., 2013).

2076
2077 Besides its detrimental effects on biota, hypoxia is responsible for the redox alterations of nitrogen and phosphorus
2078 integral stocks reaching in the Baltic proper hundreds of thousand tonnes annually: the **dissolved inorganic**
2079 **nitrogen (DIN)** pool is being depleted by denitrification, while the **dissolved inorganic phosphorus (DIP)** pool
2080 increases due to phosphate release in the water and sediment anoxic environments (e.g. Savchuk, 2010; 2018 and
2081 references therein). Resulting changes of nitrate and phosphate concentrations at the upper boundary of the

halocline affect also neighboring gulfs, exporting to the Bothnian Sea and Gulf of Finland waters with elevated phosphorus concentration (Rolff and Elfving, 2015; Lehtoranta et al., 2017; Savchuk, 2018).

Despite the decrease of land nutrient inputs after the 1980s, the extent of hypoxia in the Baltic Sea remains unaltered. This is due to the long response time of the system to reductions in N and P inputs. According to recent computations, the residence times for TN and TP in the water and sediments of the Baltic Sea combined are 9 and 49 years, respectively (Gustafsson et al., 2017; Savchuk, 2018). Furthermore, recently observed oxygen consumption rates in the Baltic Sea are higher than earlier observed, ~~counteracting shortening oxygen relieves from~~ vertical oxygen supply and natural ventilation by oxygen-rich saltwater intrusions from the North Sea (Meier et al., 2018b). Although sediments are still the most important sinks of oxygen in the Baltic Sea, the increased rates of oxygen consumption was largely driven by water column processes with, for instance, bacterial nitrification as the most prominent. Also zooplankton and higher trophic level respiration were suggested to contribute more to oxygen consumption than 30 years ago (Meier et al., 2018b). However, the importance of the latter processes is still unknown. The present total oxygen consumption rate in the water column below 60 m depth in the Baltic proper, Gulf of Riga and Gulf of Finland is estimated to be about five times that in the period 1850-1950 (Meier et al., 2018b).

Hypoxia remains an important problem also in the Baltic Sea coastal zone (Conley et al., 2011). Coastal hypoxia most often has episodic or temporary character and is driven by the seasonal variations in organic matter supply, advective transports and water column stratification (Carstensen and Conley, 2019). The latter is mostly caused by seasonal temperature changes, but may in some areas be due to occasional inflows of saltier water and lower winds during summer, changing the vertical stratification. The coastal regions most affected by hypoxia are estuaries in the Danish straits and parts of Swedish and Finnish archipelagos located in the Baltic proper and Gulf of Bothnia (Conley et al., 2011). In contrast, hypoxia is rare along the southern and south-eastern coastline (from Poland to Estonia) due to enhanced water circulation as well as in the less productive coastal zone of the northern Baltic Sea. Despite the recently reduced nutrient inputs, bottom water oxygen concentrations have improved only in a few coastal ecosystems that have experienced the largest reductions (for instance in the Stockholm Archipelago). In most of the 33 coastal sites, evaluated by Caballero-Alfonso et al. (2015), oxygen conditions have deteriorated, especially along the Danish and Finnish coasts. This finding was explained as a coupled effect of climate changes, especially warming, which reduces oxygen solubility in water and strengthens thermal stratification as well as a delay of the system in responding to nutrient reduction.

N and P are removed from the Baltic Sea by burial in sediments, but much N is also lost by denitrification (Gustafsson et al., 2017). Coastal regions constitute an efficient nutrient filter (Almroth-Rosell et al., 2016; Asmala et al., 2017) that remove about 16% of N (by denitrification) and as much as 53% of P (by burial) delivered from land (Asmala et al., 2017). The filter effect of the coastal zone is, however, highly diverse. Denitrification rates are highest in lagoons that receive large inputs of nitrate and labile organic material, while P is most efficiently buried in archipelagos (Carstensen et al., 2020). Additionally, Hoikkala et al. (2015) argued that dissolved organic

2120 matter (DOM) plays an important role for nutrient cycling in the Baltic Sea, since more than 25% of bioavailable
2121 nutrients in riverine inputs and surface waters can be in organic form.

2122
2123 Furthermore, the exchange of nutrients between the coastal zone and the open sea (Eilola et al., 2012) and the role
2124 of MBIs for the phosphorus cycling (Eilola et al., 2014) were analyzed. Eilola et al. (2014) concluded that the
2125 overall impact of MBIs on the annual uplift of nutrients from below the halocline to the surface waters is small
2126 because vertical transports are comparably large also during periods without MBIs. Instead, phosphorus released
2127 from the sediments between 60 and 100 m depth ~~under anoxic conditions in the eastern Gotland Basin~~ contributes
2128 to the eutrophication, ~~especially in the coastal regions of the eastern Baltic proper.~~

2129
2130 The cycling between nutrients and phytoplankton biomass was studied by Hieronymus et al. (2018) who found a
2131 regime shift between nutrient-limited phytoplankton variations before 1950 and a less nutrient-limited regime after
2132 1950, with a larger impact of other variations such as those in water temperature.

2133 3.2.5.7.2 Marine CO₂ system

2134 The marine CO₂ system in the Baltic Sea is greatly influenced by the production and remineralization of organic
2135 matter, as well as inputs of organic and inorganic carbon from land (Kuliński et al., 2017). The combination of all
2136 these factors makes Baltic Sea pH and partial pressure of CO₂ (pCO₂) highly variable in space and time (Carstensen
2137 and Duarte, 2019). The Baltic Sea surface water is in almost permanent pCO₂ disequilibrium with the atmosphere
2138 throughout the year (Schneider and Müller, 2018). In spring and summer, the surface seawater is undersaturated
2139 with respect to atmospheric CO₂, as a consequence of biological production and the shallowing mixed layer depth.
2140 Thus, seawater pCO₂ typically has two minima corresponding to the spring bloom and the mid-summer nitrogen
2141 fixation period. In autumn and winter, pCO₂ increases due to shifting balance between autotrophy and heterotrophy
2142 and entrainment of deeper CO₂-rich waters.

2143
2144 Remineralization of terrestrial organic matter plays an important role in shaping pCO₂ fields in the Baltic Sea.
2145 Kuliński et al. (2016) found that about 20% of the dissolved organic carbon (DOC) delivered from the Vistula and
2146 Odra rivers is bioavailable, while Gustafsson et al. (2014) even estimated that 56% of allochthonous (originating
2147 outside the Baltic Sea) DOC is remineralized in the Baltic Sea. High inputs of terrestrial organic matter that is
2148 subsequently partially remineralized in seawater turned the basins most affected by riverine runoff (Gulf of
2149 Bothnia, Gulf of Finland, Gulf of Riga) to net CO₂ sources to the atmosphere in the period 1980-2005. This
2150 outgassing was more than compensated by the high CO₂ uptake in the open Baltic proper. In 1980-2005, the whole
2151 Baltic Sea was found to be on average a minor sink for atmospheric CO₂, absorbing $4.3 \pm 3.9 \text{ g C m}^{-2} \text{ yr}^{-1}$, with
2152 the rate of atmospheric CO₂ exchange highly sensitive to the inputs of terrestrial organic matter (Gustafsson et al.,
2153 2014).

2154
2155 The high seasonal variability of pCO₂, enhanced by eutrophication, causes large seasonal fluctuations in surface
2156 water pH, amounting to about 0.5 in the central Baltic Sea (Kuliński et al., 2017) and even more, often exceeding
2157 1, in productive coastal ecosystems (Carstensen and Duarte, 2019; Stokowski et al., 2021). Furthermore, low total

alkalinity (A_T , measures buffer capacity), prominent in the northern basins, makes the Baltic Sea potentially vulnerable to Ocean Acidification (OA), i.e. pH decrease caused by rising pCO_2 in the atmosphere and thus also in seawater. However, Müller et al. (2016) showed that A_T in the Baltic Sea has increased over time, which may partly be due to increasing inputs from land (Duarte et al., 2013). The highest trend, $7.0 \mu\text{mol kg}^{-1} \text{yr}^{-1}$, found in the Gulf of Bothnia, almost entirely mitigates the pH drop expected from rising pCO_2 in the atmosphere alone. In the southern Baltic Sea, the A_T increase is lower ($3.4 \mu\text{mol kg}^{-1} \text{yr}^{-1}$) and reduces OA by about half. High seasonal pH variability, increasing A_T and variable productivity imply that OA is not measurable in the central and northern Baltic Sea. In the Danish Straits, where no A_T increase has been detected (Müller et al., 2016), a mean pH decrease of 0.004yr^{-1} was identified in coastal waters in the period of 1972-2016 (Carstensen et al., 2018), approximately twice the ocean trend.

Recent studies showed that the Baltic Sea CO_2 system functions differently from the open ocean waters. These differences include a large CO_2 input from remineralization of terrestrial organic matter (Gustafsson et al., 2014), a considerable contribution by organic alkalinity (Kuliński et al., 2014; Ulfsbo et al., 2015; Hammer et al., 2017), A_T generation under hypoxic and anoxic conditions (Gustafsson et al., 2019; Łukawska-Matuszewska and Graca, 2018), and a borate-alkalinity anomaly (Kuliński et al., 2018), making modeling a challenge. Due to the insufficient understanding of the processes involved, state-of-the-art biogeochemical models cannot yet reproduce the positive A_T trend in the Baltic Sea.

3.2.6 Marine biosphere

3.2.6.1 Pelagic habitats

3.2.6.1.1 Microbial communities

Microbial communities respond to increases in sea surface temperature and river runoff that enhance metabolism and augment the amount of substrate available for bacteria. By using long time-series from 1994 to 2006, increased input of riverine dissolved organic matter (DOM) in the Bothnian Bay and Bothnian Sea was shown to suppress phytoplankton biomass production and shift the carbon flow towards microbial heterotrophy (Wikner and Andersson, 2012; Paczkowska et al., 2020). Berner et al. (2018) presented further evidence of changes in marine microbial communities.

3.2.6.1.2 Phytoplankton and cyanobacteria

The phytoplankton growing season has become considerably prolonged in recent decades (Kahru et al., 2016; Groetsch et al., 2016; Hjerne et al., 2019; Wasmund et al., 2019). In the Baltic proper, the duration of the growing season arbitrarily indicated by a threshold of 3 mg Chl m^{-3} of a satellite-derived chlorophyll has doubled from approximately 110 days in 1998 to 220 days in 2013 (Kahru et al., 2016). In the western Baltic Sea, it now extends from February to December (Wasmund et al., 2019). Wasmund et al. (2019) analyzed data on chlorophyll a and microscopically determined biomass from 1988-2017 and found an earlier start of the growing season, which correlated with a slight increase in sunshine duration in spring, and a later end to the growing season, which correlated with warmer water in autumn. The shifts in the spring and autumn blooms led to a prolongation of the

summer biomass minimum. However, time series were rather short (30 years) and trends in irradiance might be caused by internal variability (see Section 3.2.1.3). The spring phytoplankton communities have shifted from a preponderance of early-blooming diatoms to dominance by later-blooming dinoflagellates (Wasmund, 2017; Wasmund et al., 2017) and the autotrophic ciliate *Mesodinium rubrum* (Klais et al., 2011; Hållfors et al., 2013; Hjerne et al., 2019), perhaps due to reduced ice thickness and increased winter wind-speed since the 1970s (Klais et al., 2013). Wasmund (2017) suggested that the decline in the ratio between diatom and dinoflagellate biomasses between 1984 and 1991 was caused by warmer winters. Confidence in these results is, however, low.

In summer, the amount of cyanobacteria has increased and the phytoplankton biomass maximum, which in the 1980s was in spring, is now in July-August. This shift has been explained by a complex interaction between warming, eutrophication and increased top-down pressure (Suikkanen et al., 2013). There are, however, different opinions concerning the relative effects of eutrophication and climate on changes in phytoplankton biomass and community composition. In the long-term data, results vary according to area and species group (Wasmund et al., 2011; Groetsch et al., 2016). Some studies saw evidence of eutrophication effects, modified by climate-induced variations in temperature and salinity (Hållfors et al., 2013; Olofsson et al., 2020). Others found no explanation for the gradual change in community composition, and concluded that the Baltic Sea phytoplankton community is not in a steady state (Olli et al., 2011; Griffiths et al., 2020).

Cyanobacteria accumulations derived from satellite data for 1979-2018 show both short-term (two to three year) oscillations and decadal-scale variations (Kahru and Elmgren, 2014; Kahru et al., 2018; Kahru et al., 2020). Cyanobacteria accumulations in the Baltic proper were common in the 1970s and early 1980s, but rare during 1985–1990. They increased again starting in 1991 and, especially since 1998. In the 1980s, the annual chlorophyll maximum in the Baltic proper was caused by the spring diatom bloom, but has in recent decades shifted to the summer cyanobacteria bloom in July; the timing of this bloom has also advanced by about 20 days, from the end to the beginning of July (Kahru et al., 2016).

In the Baltic proper, the hypoxia-induced decrease in N:P ratio and increase of phosphate pool left over in the surface layer after the spring bloom led to intensification of the “vicious circle” (Vahtera et al., 2007), further augmented by increasing water temperature, and resulted in conspicuous expansion of the surface diazotrophic cyanobacteria accumulations, covering in the 21st century 150–200 thousand square kilometers (e.g. Kahru and Elmgren, 2014; Savchuk, 2018, and references therein). Although mechanisms of the interannual oscillations of two to three years remain unexplained (Kahru et al., 2018), there is a strong correlation of the accumulations with hypoxia-related biogeochemical variables and water temperature at the decadal scale of five to twenty years (Kahru et al., 2020). In the Bothnian Sea, the decreased nitrogen import and increased phosphorus import from the Baltic proper has shifted the nutrient balance and made cyanobacteria accumulations a permanent feature (Kahru and Elmgren, 2014; Kuosa et al., 2017) and probably also increased production and sedimentation (Ahlgren et al., 2017; Kahru et al., 2018; Kahru et al., 2020; Kuosa et al., 2017; Lehtoranta et al., 2017; Rolff and Elfving, 2015; Savchuk, 2010; Vahtera et al., 2007).

Experimental evidence supports the idea that climate change can and will drive changes in the pelagic ~~primary~~ production ~~from both ecological and evolutionary perspectives~~ (Sommer et al., 2012; Hattich et al., 2021), and a thorough review of benthic-pelagic coupling in the Baltic Sea demonstrates ecosystem-wide consequences of altered pelagic primary production (Griffiths et al., 2017).

3.2.6.1.3 Zooplankton

Several studies have confirmed that marine copepod species have declined in abundance since the 1980s, while euryhaline or limnetic, often small species have increased (Hänninen et al., 2015; Suikkanen et al., 2013; Kortsch et al., 2021). The observed decline of marine taxa has been linked to the reduction in surface-water salinity since the 1980s (Vuorinen et al., 2015), whereas the increase of brackish-water taxa has been positively influenced by the temperature increase, directly or indirectly (Mäkinen et al., 2017). Small-scale effects on individual species may affect reproductive success, and hence influence both populations and communities (Möller et al., 2015).

3.2.6.2 Benthic habitats

3.2.6.2.1 Macroalgae and vascular plants

Long-term changes in Baltic Sea macroalgae and charophytes have been attributed to changes in salinity, wind exposure, nutrient availability and water transparency (Gubelit, 2015; Blindow et al., 2016; Rinne and Salovius-Laurén, 2020), and biotic interactions may also play a role (Haavisto and Jormalainen, 2014; Korpinen et al., 2007). The long-term decrease of water transparency from 1936 to 2017 has been estimated to have reduced sea floor areas in the northern Baltic Sea favorable for *Fucus* spp. by 45% (Sahla et al., 2020). Overall, it is expected that climate change and its interaction with other environmental factors (e.g. eutrophication) will cause complex responses and influence carbon storage in both macroalgae and vascular plants in the Baltic Sea (Jonsson et al., 2018; Takolander et al., 2017; Röhr et al., 2016; Perry et al., 2019; Salo et al., 2020; Bobsien et al., 2021).

3.2.6.2.2 Zoobenthos

Soft-sediment benthic communities depend on variables that are influenced by climatic variability. On the southwestern coast of Finland, amphipods have been replaced by Baltic clam *Limecola balthica* and the invasive polychaetes *Marenzelleria* spp., a change attributed to an increase in near-bottom temperature and fluctuations in salinity and oxygen (Rousi et al., 2013). Variations of zoobenthos in the Åland archipelago during 1983-2012 were associated with a salinity decline (Snickars et al., 2015), and effects related to climate change have acted as drivers for the long-term progression of zoobenthic communities (Rousi et al., 2019; Weigel et al., 2015; Ehrnsten et al., 2020).

3.2.6.3 Non-indigenous species

Numerous non-indigenous species have gained a stronghold in the Baltic Sea ecosystem during the past few decades, and in many cases these species have wider tolerance-ranges than the native ones, thus making them highly competitive under changing climate including warmer, and possibly less saline water, further impacted by other drivers such as eutrophication. The ecological impacts of these species may vary from filling vacant

ecological niches to potentially outcompeting native species, and thus influencing the entire food web structure and functioning (Weigel et al., 2015; Griffiths et al., 2017; Ojaveer et al., 2017).

3.2.6.4 Fish

Sprat and herring in the Baltic Sea are influenced by multiple factors, including fisheries, predation, food availability and climatic variations. Sprat has benefited from the seawater warming (Voss et al., 2011). In 1990–2020, sprat populations were affected both by climate and top-down control, i.e. fisheries and predation by cod (Eero et al., 2016). In the 1980s, overfishing and a partly climate change-induced decline in suitable spawning habitat, ‘reproductive volume’, interacted to drastically reduce the cod population (Hinrichsen et al., 2011; Casini et al., 2016), with cascade effects on its main prey, sprat and herring, as well as zooplankton (Casini et al., 2008).

The various effects of temperature and salinity on sprat and cod also resulted in a spatial mismatch between these species, which contributed to an increase of sprat stocks (Reusch et al., 2018). The freshening of the Baltic Sea surface water, with the associated decline in marine copepods (Hänninen et al., 2015), contributed to a halving of weight-at-age of 3-year old herring, from 50–70 g in the late 1970s to 25–30 g in 2000s (Dippner et al., 2019).

Among coastal fish, pikeperch (*Sander lucioperca*) has recently expanded its distribution northwards along the coasts of the Bothnian Sea, apparently aided by warmer waters (Pekcan-Hekim et al., 2011). For many coastal piscivores (perch, pike, pike-perch), as well as for cyprinids, coastal eutrophication is, however, equally or more important than climate (Bergström et al., 2016; Snickars et al., 2015). Long-term studies illustrate that it is hard to disentangle abiotic and biotic interactions, e.g. between fish and their food (benthos), and climate-related drivers thus appear significant on a multidecadal time-scale across a large spatial scale (Törnroos et al., 2019).

3.2.6.5 Marine mammals

~~The breeding distributions of the ice-breeding seals in the Baltic Sea have evolved with ice coverage, with the seals breeding where and when ice optimal for breeding occurs. Breeding ringed seals need ice throughout their relatively long lactation period (>6 weeks), and also use ice as moulting habitat. Ringed seals prefer compact or consolidated pack ice as it provides cavities and snowdrifts suitable for the construction of the lairs, most importantly the breeding lair (Sundqvist et al., 2012).~~

~~Sea ice changes, along with implementation of specific management and protection measures, have had a rapid influence on the populations of several Baltic Sea mammal populations, in particular seals (Reusch et al., 2018). Reusch et al. (2018) attributed these changes also to reduced exposure to harmful substances and increases in overall fish stocks as a consequence of eutrophication (including reduced stocks of several commercial fish species). Thus, specific climate change related impacts on seals are hard to establish, although reconstructions of distributional histories since the last glaciation have been attempted for some seal species (Ukkonen et al., 2014).~~

2302 The availability of suitable breeding ice for ringed seals in the Bothnian Bay is decreasing (Section 3.2.4.4). During
2303 the 20th century the marine mammals of the Baltic Sea experienced large declines in abundance because of hunting
2304 of seals, bycatch of porpoises and exposure to harmful substances causing reduced fertility.

Formatiert: Englisch (USA)

2306 The breeding distributions of the ice-breeding seals in the Baltic Sea have evolved with ice coverage, with the
2307 seals breeding where and when ice optimal for breeding occurs. Breeding ringed seals need ice throughout their
2308 relatively long lactation period (>6 weeks), and also use ice as moulting habitat. Ringed seals prefer compact or
2309 consolidated pack ice as it provides cavities and snowdrifts suitable for the construction of the lairs, most
2310 importantly the breeding lair (Sundqvist et al., 2012).

Formatiert: Nicht unterstrichen, Schriftfarbe: Automatisch, Englisch (USA)

Formatiert: Nicht unterstrichen, Schriftfarbe: Automatisch, Englisch (USA)

Formatiert: Englisch (USA)

2312 Implementation of specific management- and protection measures, have had a profound positive influence on the
2313 populations of several Baltic Sea mammal populations, in particular seals (Reusch et al., 2018). Reusch et al.
2314 (2018) attributed these changes also to reduced exposure to harmful substances and initial increases in overall fish
2315 stocks as a consequence of eutrophication, followed by a reduction of several commercial stocks due to
2316 overfishing.

Formatiert: Nicht unterstrichen, Schriftfarbe: Automatisch, Englisch (USA)

Formatiert: Englisch (USA)

2318 Specific climate change-related impacts on seals are difficult to detect, although reconstructions of historical
2319 habitats since the last glaciation have been attempted for some seal species (Ukkonen et al., 2014). Along with
2320 warmer winters the availability of suitable breeding ice for ringed seals in the Bothnian Bay is decreasing (Section
2321 3.2.4.4). The breeding success of ringed seal was probably reduced by the exceptionally mild winter of 2007-2008
2322 (Jüssi, 2012) and several similar or even milder ice-winters have followed (Ilmatieteen laitos, 2020). The winters
2323 2019–2020, 2007–2008 and 2014–2015 are the mildest in the annual ice cover statistics for the Baltic Sea (Uotila
2324 et al., 2015). The southern breeding populations of the ringed seal in the Gulf of Finland, the Gulf of Riga and
2325 Archipelago Sea are already facing the challenges of milder winters: the ice covered area during the breeding
2326 season has been reduced and overlying snow for breeding lairs has been absent from the southern areas of the
2327 ringed seal breeding range in most winters of the past decade (Ilmatieteen laitos, 2020). Thus, the only available
2328 breeding ice in the Gulf of Finland in 2020 was found very near St. Petersburg (Halkka, 2020).

Formatiert: Nicht unterstrichen, Schriftfarbe: Automatisch, Englisch (USA)

Formatiert: Nicht unterstrichen, Schriftfarbe: Automatisch, Englisch (USA)

Formatiert: Nicht unterstrichen, Schriftfarbe: Automatisch, Englisch (USA)

Formatiert: Nicht unterstrichen, Schriftfarbe: Automatisch, Englisch (USA)

2330 For grey seals, the lower availability of suitable breeding ice in its core distribution area has led to more breeding
2331 on land in areas where drift ice used to be found (Jüssi et al., 2008). Grey seals are known to gather to breed in
2332 certain sea areas regardless of the winter severity, so some land colonies may become overpopulated. As an
2333 example, in 2016, 3,000 grey seal pups were born on three islets in the northern Gulf of Riga.

2335 Flooding of seal haul-outs due to sea level rise will first occur in the southernmost Baltic Sea, where relative sea
2336 level rise will be most rapid (EEA, 2019c), and haul-out sites are mainly low sand or shingle banks. In Kattegat,
2337 relative sea level rise is estimated to be lower, and while the haul-outs in the western part are low sand and shingle
2338 banks similar to those in the southern Baltic Sea, haul-outs in eastern Kattegat are skerries with a higher profile.
2339 In the central and northern Baltic Sea, haul-outs are mainly skerries and here relative sea level rise is estimated to
2340 be low or even negative in the 21st century (EEA, 2019c). Yet, in recent history, winter storm surges have been

2341 observed to flood grey seal breeding colonies and push limited ice with ringed seal pups onto shore in the Gulf of
2342 Riga and Pärnu Bay. In the southern Baltic Sea and western Kattegat, increasing sea levels may turn parts of larger
2343 islands or previously inhabited islands into suitable seal haul-outs, but this is hard to project and depends, among
2344 other things, on the future management and protection of such areas.

2345
2346 Sea levels in the southern Baltic Sea have been rising at up to 3 mm year⁻¹ since the 1970s (EEA, 2019c) and the
2347 available haul-out areas have thus seen reductions already. However, during this time, the relevant harbour and
2348 grey seal populations have been recovering at high rates from past depletion (HELCOM, 2018b), with no
2349 documented or suspected effects of rising sea levels published.

2350
2351 The only cetacean resident in the Baltic Sea, the harbour porpoise (*Phocoena phocoena*), is a wide-spread species
2352 and seems to be rather tolerant of different temperatures as well as habitats. There are harbour porpoise populations
2353 in the waters around Greenland as well as along the coast of the Iberian Peninsula, Morocco, West Sahara and in
2354 the Black Sea. However, the Baltic proper harbour porpoise population has been shown to differ genetically (Lah
2355 et al., 2016) and morphologically (Galatius et al., 2012) from neighbouring populations, which may imply local
2356 adaptations that we are currently unaware of. Sea ice limits the range available to the Baltic proper harbour
2357 porpoise population since they need to come to the surface to breathe every 1-5 minutes. Hence, a decreasing ice
2358 cover is likely to increase the available range for the population. However, a change in the prey community
2359 resulting from climate-change related factors could potentially have serious effects on this critically endangered
2360 (Hammond et al., 2008) population of a small whale, which is dependent on constant access to prey (Wisniewska
2361 et al., 2016).

2362 **3.2.6.6 Waterbirds**

2363 The winter distribution of many waterbirds has extended northwards in response to the increase in temperature
2364 and the decreasing extent of sea ice cover. This can be observed as an overall increase in winter abundance of
2365 waterbirds, because part of the population of some species (mainly diving ducks) that formerly wintered further to
2366 the southwest now remain in the Baltic Sea (Pavón-Jordán et al., 2019). Many species show decreasing trends in
2367 abundance in the southern parts of their wintering ranges (typically in western and southern Europe) but increases
2368 near the northern edge of their distribution, typically the Baltic Sea region (MacLean et al., 2008; Skov et al., 2011;
2369 Aarvak et al., 2013; Lehikoinen et al., 2013; Pavón-Jordán et al., 2015; Nilsson and Haas, 2016; Marchowski et
2370 al., 2017; Fox et al., 2019). Similar shifts are seen in species that traditionally wintered in the Baltic Sea, but
2371 currently show declining wintering numbers there, as part of the population now winters in the White, Barents and
2372 Kara seas (Fox et al., 2019).

2373
2374 Although the community composition changes rapidly, the changes are not fast enough to track the thermal isocline
2375 shifts (Devictor et al., 2012; Gaget et al., 2021). How species respond to changes in winter temperature seems,
2376 however, to be highly species- or group-specific (Pavón-Jordán et al., 2019). Many species now winter closer to
2377 their breeding areas, shortening migration distances (Lehikoinen et al., 2006; Rainio et al., 2006; Gunnarsson et
2378 al., 2012).

2379
2380 Mainly owing to milder spring temperatures and related effects on vegetation and prey, many waterbirds migrate
2381 earlier in spring (Rainio et al., 2006), and hence arrive earlier in the breeding area (Vähätalo et al., 2004), and
2382 some also start breeding earlier (van der Jeugd et al., 2009). Delayed autumn migrations have also been noted, but
2383 their relation to climate change is less clear (Lehikoinen and Jaatinen, 2012).

2384
2385 Earlier loss of sea ice was found to improve pre-breeding body condition of female common eiders, leading to
2386 increasing fledging success in offspring (Lehikoinen et al., 2006). On the other hand, algal blooms promoted by
2387 higher seawater temperature have in some cases caused low quality in bivalve prey for common eiders, leading
2388 more birds to skip breeding (Larsson et al., 2014). Warmer seawater in winter also increases the energy expenditure
2389 of mussels, thus directly reducing their quality as prey for eiders (Waldeck and Larsson, 2013).

2390
2391 Most Baltic Sea waterbird species are migratory and affected by climate change also outside the Baltic Sea region,
2392 in the Arctic (breeding season) and in southern Europe and western Africa (wintering; Fox et al., 2015). This is
2393 important, given that climate warming is most pronounced in the Arctic and northern Eurasia and above average
2394 also in southern Europe and northern Africa (Allen et al., 2018).

2395 3.2.6.7 Marine food webs

2396 The entire marine food web of the Baltic Sea has been greatly impacted by climate change-related drivers that
2397 have altered the physical environment and the physiological tolerance limits of several species, by causing micro-
2398 evolution of Baltic Sea species, and by interactive effects of climate change with other environmental drivers, such
2399 as eutrophication and hypoxia/anoxia (Niiranen et al., 2013; Wikner and Andersson, 2012; Schmidt et al., 2020;
2400 Pecuchet et al., 2020).

2401
2402 ~~Integrated approaches encompassing all of the ecosystem components discussed above, are needed in order to~~
2403 ~~understand and manage the linkages between large-scale and long-term changes driven by synergistic impacts of~~
2404 ~~over-arching climate change-related physical and chemical drivers in combination with other factors such as~~
2405 ~~eutrophication, which may complicate human adaptation to the changing marine ecosystem (Blenckner et al.,~~
2406 ~~2015; Stenseth et al., 2020; Bonsdorff, 2021).~~Integrated approaches encompassing all of the ecosystem-
2407 components discussed above are needed in order to understand and manage the linkages among large-scale and
2408 long-term climate effects. These are driven by synergistic interactions of climate change-related physical and
2409 chemical drivers with other factors, such as eutrophication or large-scale fisheries, which complicate human
2410 adaptation to the changing marine ecosystem (Niiranen et al., 2013; Blenckner et al., 2015; Hyytiäinen et al., 2021;
2411 Stenseth et al., 2020; Bonsdorff, 2021).

Formatiert: Schriftart:

2412 3.3 Future climate change

2413 3.3.1 Atmosphere

2414 3.3.1.1 Large-scale atmospheric circulation

2415 ~~Continued greenhouse-gas induced warming is the key driver for future climate change and changes in atmospheric~~
2416 ~~circulation are relatively less important (IPCC, 2021). However, as the regional climate in the Baltic Sea region is~~
2417 ~~strongly governed by the large-scale circulation of the atmosphere it is important to also consider changes in it~~
2418 ~~when assessing future regional climate (e.g. Kjellström et al., 2018).~~

2419
2420 In the future, the NAO is very likely to continue to exhibit large natural variations, similar to those observed in the
2421 past. In response to global warming, it is likely to become slightly more positive on average (Knudsen et al., 2011).
2422 Trends in the intensity and persistence of blocking remain ~~yet to be determined~~~~uncertain~~ (IPCC, 2014b). The AMO
2423 is expected to be very sensitive even to weak global warming, shortening the time scale of its response and
2424 weakening in amplitude (Wu et al., 2018; Wu and Liu, 2020). This will likely reduce the decadal variability of
2425 SSTs in the Northern Hemisphere. Recent studies indicate a degree of decadal predictability for blocking and the
2426 NAO influenced by the AMO (Athanasiadis et al., 2020; Wills et al., 2018; Jackson et al., 2015).

2427 3.3.1.2 Air temperature

2428 Table ~~127~~ lists the air temperature changes over the Baltic Sea catchment area and the Baltic Sea calculated from
2429 an ensemble of regional coupled atmosphere-ocean simulations (Gröger et al., 2021b). Due to the ice/snow-albedo
2430 feedback, warming is larger in winter than in summer, and the land is warming faster than the Baltic Sea (Fig.
2431 26a). Due to its proximity to the Arctic, the Baltic Sea region including both land and sea is warming faster than
2432 the global mean figures (Section 1.5.1.1). The surface air temperature increase is expected to be largest in the
2433 northern Baltic Sea region especially in winter. These statements are true for both uncoupled atmosphere
2434 (Christensen et al., 2021) and coupled atmosphere-ocean regional climate simulations (Gröger et al., 2019; Gröger
2435 et al., 2021b).

2436
2437 For RCP2.6, the global annual mean surface air temperature change ~~between 1976–2005 and 2069–2098~~ averaged
2438 over the simulations is 1.0°C. The corresponding global changes for RCP4.5 and RCP8.5 are 1.9 and 3.5°C,
2439 respectively. Over land in the Baltic Sea region, the warming is larger in each of the three scenarios, amounting to
2440 1.5, 2.6 and 4.3°C, respectively (Table ~~127~~). Over the Baltic Sea, the increase is slightly smaller than over land
2441 (1.4, 2.4 and 3.9°C, respectively) but still larger than the corresponding global mean increase in surface air
2442 temperature. The latter result was expected and found in coupled atmosphere-ocean scenario simulations (Table
2443 ~~127~~), but not in all atmosphere-only runs (Christensen et al., 2021).

2444 *Extreme Air temperatures:*

2445
2446 ~~Changes in daily minimum and maximum temperatures have similar spatial patterns as the mean air temperature~~
2447 ~~changes, with the expected greater warming for minimum temperature (Christensen et al., 2021). According to~~
2448 ~~Christensen et al. (2021) and previous studies (BACC II Author Team, 2015), the latter result is explained by the~~

Formatiert: Schriftart:

reduced outgoing long-wave radiation under increased greenhouse gas concentrations. The long-wave radiation acts to cool the surface, especially when the ground is warmer than the air, e.g. during winter and during nights. Changes in daily minimum and maximum temperatures have similar spatial patterns as the mean air temperature changes, although with greater warming for minimum temperature (Fig. 27). Such a decrease of the diurnal temperature range can have a number of explanations. Based upon an ensemble of CMIP5 GCMs, Lindvall and Svensson (2015) suggested that increasing downwelling longwave radiation due to larger greenhouse gas concentrations, increased cloudiness at high latitudes, changes in the hydrological cycle and in changes in shortwave incoming radiation might be an explanation for the decreased diurnal temperature range. In addition to these drivers, the difference in diurnal temperature range in northern Europe is small in winter implying that differences between daily maximum and minimum temperatures also substantially depend on synoptic-scale variability and air mass origin. As a consequence, reduced temperature variability on synoptic time scales, resulting from reduced temperature gradients between the Atlantic Ocean and Eurasia, may be a reason for the decreased diurnal temperature range.

The number of hot spells ~~is~~^{are} projected to increase, in particular in the southern Baltic Sea region (Gröger et al., 2021b). In coupled atmosphere-ocean simulations, the strongest increases in the annual mean number of consecutive days of tropical nights and the annual maximum number of tropical nights (with temperature above 20°C all night) in the Baltic Sea region were projected to occur over the open sea (Gröger et al., 2021b). In contrast, projections of tropical nights with atmosphere-only models show no significant change (Gröger et al., 2021b; Meier et al., 2019a). Due to the sea ice/snow albedo feedback, the largest decline in the number of frost days was projected to occur over the northeastern Baltic Sea region, i.e. northern Scandinavia and adjacent northern Russia (Gröger et al., 2021b). In addition to the magnitude of the warming that is affected by the sea ice/snow albedo feedback, the baseline climate may also play a role. Further southwest, where the winters are milder, there are less frost days to start with, and therefore less room for a further decrease in the future.

3.3.1.3 Solar radiation and cloudiness

There are a few studies on projected future solar radiation over Europe. Global climate models of the CMIP5 generation indicated an increase in surface solar radiation, highest over southern Europe and decreasing towards north, but still with a slight increase over the Baltic Sea (Bartók et al., 2017; Müller et al., 2019). The RCP scenarios include even a decrease in aerosol emissions, which enhances the warming and the increase in solar radiation in the global climate models.

However, some regional climate models instead showed a decrease in surface solar radiation over the Baltic Sea area, in winter by about 10% over most of the catchment (Bartók et al., 2017; Christensen et al., 2021). This change was largely attributed to increasing future cloud-cover, due to a more zonal airflow, and was accompanied by increased winter precipitation. Thus, there are large differences in modelled surface solar radiation between global and regional models (Bartók et al., 2017). Unknown future aerosol emissions add to the uncertainty. In most RCM scenario simulations the effect of changing aerosol concentrations were not considered (Boé et al., 20020a).

2487 Global mean energy balance components have improved with every new climate model generation. For the latest
2488 CMIP6, models show good agreement for clear sky shortwave energy fluxes in today's climate, both between
2489 models and ~~between models and compared to~~ reference data (Wild et al., 2021). However, there are still substantial
2490 discrepancies among the various CMIP6 models in their representation of several of the global annual mean energy
2491 balance components, and the inter-model spread increases further on regional, seasonal and diurnal scales. Thus,
2492 future changes in solar radiation and cloudiness remain highly uncertain, not least on the regional scale.

2493 3.3.1.4 Precipitation

2494 Precipitation in winter and spring is projected to increase over the entire Baltic Sea catchment, while summer
2495 precipitation is projected to increase in the northern half of the basin only (Christensen et al., 2021). In the south,
2496 summer precipitation is projected to change very little, although with a large spread between different models
2497 including both increases and decreases. The projected increase in the north is a rather robust feature among the
2498 regional climate models but with a large spread in the amount. Ensemble mean precipitation changes from coupled
2499 atmosphere-ocean simulations are summarized in Table 138. For the Baltic Sea catchment area, projected annual
2500 mean precipitation changes for the three RCP scenarios amount to 5, 9 and 15% (Table 138) and are much larger
2501 than global averages (Section 1.5.1.2). Over the Baltic Sea, the changes are similar ~~to those than~~ over the land area
2502 (6, 8 and 16%).

2503
2504 ~~Expressed by the Clausius-Clapeyron equation, warming increases the potential for extreme precipitation due to~~
2505 ~~intensification of the hydrological cycle associated with the growth of atmospheric moisture content. For Northern~~
2506 ~~Europe, regional climate models indicate an overall increase in the frequency and intensity of heavy precipitation~~
2507 ~~events in all seasons and longer wet and dry spells (Christensen and Kjellström, 2018; Rajczak and Schär, 2017;~~
2508 ~~Christensen et al., 2021, and references therein). The largest increase in the number of high precipitation days is~~
2509 ~~projected for autumn. The number of drought events per year are expected to decrease, while their length is~~
2510 ~~expected to increase (Christensen and Kjellström, 2018). Changes in more extreme events, like 10-, 20- or 50-year~~
2511 ~~events, are less certain. Expressed by the Clausius-Clapeyron equation, warming increases the potential for extreme~~
2512 ~~precipitation due to intensification of the hydrological cycle associated with the growth of atmospheric moisture~~
2513 ~~content. For northern Europe, regional climate models indicate an overall increase in the frequency and intensity~~
2514 ~~of heavy precipitation events in all seasons (Christensen and Kjellström, 2018; Rajczak and Schär, 2017;~~
2515 ~~Christensen et al., 2021, and references therein). The largest increase in the number of high precipitation days is~~
2516 ~~projected for autumn. Changes in more extreme events, like 10-, 20- or 50-year events, are less certain.~~

2517
2518 ~~Changes in dry spells are another feature of an intensified hydrological cycle. Coppola et al. (2021) found~~
2519 ~~increasing number of consecutive dry days without precipitation for countries south of the Baltic Sea while no~~
2520 ~~changes were reported in the North.~~

2521 3.3.1.5 Wind

2522 In general, projected changes in wind speed over the Baltic Sea region are not robust among ~~ESMsEarth System~~
2523 ~~Models~~ (Kjellström et al., 2018; Gröger et al., 2021b). However, Ruosteenoja et al. (2019) found in CMIP5

2524 projections a slight but significant wind speed increase in autumn and a decrease in spring over Europe and the
2525 North Atlantic. Furthermore, over sea areas where the ice cover is projected to diminish on average, such as the
2526 Bothnian Sea and the eastern Gulf of Finland, the mean wind is projected to increase systematically because of a
2527 warmer sea surface and reduced stability of the planetary boundary layer (Meier et al., 2011c; Gröger et al., 2021b;
2528 Räisänen, 2017).

2529
2530 Projections of the future behavior of extratropical cyclones are inherently uncertain because changes in several
2531 drivers result in opposite effects on cyclone activity. With global warming, the lower troposphere temperature
2532 gradient between low and high latitudes decreases due to polar amplification. Near the tropopause and in the lower
2533 stratosphere, the opposite is true, thus implying competing lower and upper tropospheric effects on changes in
2534 baroclinicity implying changes in baroclinicity (Grise and Polvani, 2014; Shaw et al., 2016; Stendel et al., 2021).
2535 An increase in water vapour enhances diabatic heating and tends to increase the intensity of extratropical cyclones
2536 (Willison et al., 2015; Shaw et al., 2016) and contribute to their propagation further poleward (Tamarin-Brodsky
2537 and Kaspi, 2017; Tamarin and Kaspi, 2017). The opposite is true in parts of the North Atlantic region, e.g. south
2538 of Greenland. For this region the North-South temperature gradient is increasing, as the weakest warming in the
2539 entire Northern Hemisphere is over ocean areas south of Greenland. North of this local minimum the opposite is
2540 true. The increase in the North-South gradient over the North Atlantic may be responsible for some ESMsEarth
2541 System-Models showing an intensification of the low pressure activity and thereby higher wind speeds over a
2542 region from the British Isles and through parts of north-central Europe in winter (Leckebusch and Ulbrich, 2004;
2543 Ulbrich et al., 2008). These projections have been confirmed by (Harvey et al., 2012). They compared the ensemble
2544 storm track response of CMIP3 and CMIP5 model simulations and found that both projections show an increase
2545 in storm activity in the midlatitudes, with a smaller spread in the CMIP5 simulations. In contrast to CMIP3, the
2546 CMIP5 ensemble showed a significant decrease in cyclone track density north of 60°N. Hence, pre-CMIP3 and
2547 CMIP3 studies showed a clear poleward shift of the North Atlantic storm track (e.g. Fischer-Bruns et al., 2005;
2548 Yin, 2005; Bengtsson et al., 2009), whereas the CMIP5 ensemble predicts only an eastward extension of the North
2549 Atlantic storm track (Zappa et al., 2013). The newest generation of models from CMIP6 resulted in significant
2550 reduction of biases in storm track representation compared to CMIP3 and CMIP5, but the response to climate
2551 change is quite similar compared to the previous assessments (Harvey et al., 2020). The eastward extension of the
2552 North Atlantic storm track seems to be a robust result as it is found in pre-CMIP3, CMIP3 and CMIP5 simulations
2553 (Feser et al., 2015). The response of CMIP6 models is similar to CMIP5 models, but it is considerably larger,
2554 probably due to the larger climate sensitivity in the CMIP6 models (Harvey et al., 2020).

2555
2556 In summary, there is no clear consensus among climate change projections in how changes in frequency and/or
2557 intensity of extratropical cyclones will affect the Baltic Sea region (Räisänen, 2017). However, in future climate
2558 the frequency of severe wind gusts in summer associated with thunderstorms may increase (Rädler et al., 2019).

2559 3.3.1.6 Air pollution, air quality and atmospheric nutrient deposition

2560 The main conclusions by the BACC II Author Team (2015) concerning projections of air quality in the Baltic Sea
2561 region still hold. The main factor determining future air quality in the region is regional emissions of air pollutants,

not changes in meteorological factors related to climate change or in intercontinental pollution transport (see e.g. Langner et al., 2012; Hedegaard et al., 2013).

Recent post-BACC II air quality modelling studies for the Baltic Sea area are Colette et al. (2013), Varotsos et al. (2013), Colette et al. (2015), Hendriks et al. (2016), and Watson et al. (2016). They concentrate mainly on particulate matter (PM) and ground-level ozone (O₃), the pollutants most likely to be affected by changing climate parameters. They agree with current day air quality trends in that the Baltic Sea region in general is less exposed to air pollution than the rest of Europe.

Jacob and Winner (2009) showed that climate change is likely to increase ground-level ozone in central and southern Europe. In a meta-analysis, Colette et al. (2015) assessed the significance and robustness of the impact of climate change on European ground-level ozone based on 25 model projections, including some driven by SRES (Special Report on Emission Scenarios by Nakicenovic et al., 2000) and RCP scenarios. They indicate that an increase in ground-level ozone is not expected for the Baltic Sea region. A latitudinal gradient was found from increase in large parts of continental Europe (+ 5 ppbv), but a small decrease over Scandinavia (up to -1 ppbv). Studies that explicitly compared the magnitude of projected climate and anthropogenic emission changes (Langner et al., 2012; Colette et al., 2013; Varotsos et al., 2013) all confirmed that changes in emission of ozone precursors (NO_x, VOCs) had the larger effect. For Northern Europe, Varotsos et al. (2013) estimated that reductions in snow cover and solar radiation in a SRES A1B scenario lead to an ozone decrease of about 2 ppb by 2050, compared to present conditions.

Varotsos et al. (2013) stress the importance of future biogenic isoprene emissions for ozone concentrations. In the 2050 climate, increases in ozone concentrations are associated with increased biogenic isoprene emissions due to increased temperatures, whereas increased water vapour over the sea, as well as increased wind speeds, are associated with decreases. Hendriks et al. (2016) emphasise that isoprene emissions may increase significantly in coming decades if short-rotation coppice plantations are greatly expanded, to meet the increased biofuel demand resulting from the EU decarbonisation targets. They investigate the competing effects of anticipated trends in land use, anthropogenic emissions of ozone precursors and climate change on European ground-level ozone concentrations and related health and environmental effects by 2050. They found that increased ozone concentrations and associated health damage caused by a warming climate (+ 2 to 5°C across Europe in summer) might be more than the reduction that can be achieved by cutting emissions of anthropogenic ozone precursors in Europe. This is an example for the dominance of the climate change effect over a pollutant emission reduction effect, an exception to the statement made at the beginning of this section.

Orru et al. (2013, 2019) and Geels et al. (2015) studied the effect of climate change on ozone-related mortality in Europe. Orru et al. (2019) present their results on country level, including all Baltic Sea EU-countries. They conclude that although mortality related to ground-level ozone is projected to be lower in the future (mainly due to decreased precursor emissions), the reduction could have been larger, without climate change and an increasingly susceptible population.

2601
2602 In parts of the Baltic Sea region, a considerable air pollution is due to shipping. Ship traffic in the region is
2603 projected to increase over the coming decades, which could lead to larger emissions (i.e. NO_x and PM) than today,
2604 unless stricter air quality regulations counter this potential trend. For the Baltic Sea, a nitrogen emission control
2605 area (NECA) will become effective in 2021. Karl et al. (2019a) designed future scenarios to study the effect of
2606 current and planned regulations of ship emissions and the expected fuel efficiency development on air quality in
2607 the Baltic Sea region. They showed that in a business-as-usual scenario for 2040 (SECA-0.1% and fuel efficiency
2608 regulation effective starting in 2015), the introduction of the NECA will reduce NO_x emissions from ship traffic
2609 in the Baltic Sea by about 80% in 2040. The reduction in NO_x emissions from shipping translates to a ~60%
2610 decrease in NO₂ summer mean concentrations in a wide corridor around the ship routes. The coastal population of
2611 northern Germany, Denmark and western Sweden will be exposed to less NO₂ in 2040 due to the introduction of
2612 the NECA. With lower atmospheric NO_x levels, less ozone will be formed, and the estimated daily maximum O₃
2613 concentration over the Baltic Sea in summer 2040 will on average be 6% lower than without the NECA. Compared
2614 to today, the introduction of the NECA will also reduce ship-related PM_{2.5} emissions by 72% by 2040, compared
2615 to -48% without the NECA. Simulated nitrogen deposition on the Baltic Sea decreases 40-44% on average between
2616 2012 and 2040. A similar study by Jonson et al. (2019) estimated that the contributions of Baltic Sea shipping to
2617 NO₂ and PM_{2.5} concentrations, and to the deposition of nitrogen, will be reduced by 40-50 % from 2016 to 2030,
2618 mainly as a result of NECA.

2619
2620 The pollutant concentrations reported in this section may drop to yet not known lower values if the shipping sector
2621 is successful in meeting the International Maritime Organization (IMO) target in greenhouse gas emission
2622 reduction of 50% by 2050. This reduction is only possible if low-carbon alternative fuels will be introduced.
2623 Employing a high energy efficiency, as it is already considered in scenarios used by Karl et al. (2019a), will not
2624 be sufficient. The new fuels will also lead to altered emissions of pollutants.

Formatiert: Schriftart: Englisch (Großbritannien)

2625 3.3.2 Land

2626 3.3.2.1 River discharge

2627 Climate change ~~may have anis-likely-to-have-a-clear~~ influence on the seasonal river flow regime, as a direct
2628 response to changes in air temperature, precipitation and evapotranspiration (BACC II Author Team, 2015; Blöschl
2629 et al., 2017).

2630
2631 For areas in the northern Baltic Sea region presently characterized by spring floods due to snow melt, the floods
2632 are likely to occur earlier in the year and their magnitude is likely to decrease owing to less snowfall, shorter snow
2633 accumulation period, and repeated melting during winter. As a consequence, sediment transport and the risk of
2634 inundation are likely to decrease.

2635
2636 In the southern part of the Baltic Sea region, increasing winter precipitation is projected to result in increased river
2637 discharge in winter. In addition, groundwater recharge is projected to increase in areas where infiltration capacity
2638 is not currently exceeded, resulting in higher groundwater levels. Decreasing precipitation combined with rising

temperature and evapotranspiration during summer is projected to result in drying of the root zone, increasing demands for irrigation in the southern Baltic Sea region.

Projections with a process-oriented hydrological model suggested that, under the RCP4.5 and RCP8.5 scenarios, the total river flow during 2069-2098 relative to 1976-2005 will increase 1-21% and 6-20%, respectively, illustrating the large inherent uncertainty in hydrological projections (Saraiva et al., 2019a; Meier et al., 2021b). According to these and previous projections, the increase of river flow will mainly take place in the north, while total river flow to the south will decrease (Stonevičius et al., 2017; Šarauskienė et al., 2017). Winter flow will increase due to intermittent melting (Stonevičius et al., 2017). Projected discharge changes attributed to increasing air temperature are reflected in observed trends (Section 3.2.2.1), whereas changes attributed to increasing precipitation are necessarily not (Wilson et al., 2010).

Since the publication of BACC II (BACC II Author Team, 2015), ensemble sizes of scenario simulations with hydrological models have increased, enabling the estimate of uncertainties in projections (e.g. Roudier et al., 2016; Donnelly et al., 2017). Donnelly et al. (2014) focused on projecting changes in discharge to the Baltic Sea, by using a semi-distributed conceptual hydrological model for the drainage basin BSDB (Balt-HYPE), combined with a small ensemble of climate projections under the SRES A1B and A2 scenarios. Results showed an increased overall discharge to the Baltic Sea, with a seasonal shift towards higher winter and lower summer flows and diminished seasonal snow-melt peaks. Efforts were made to assess the uncertainty in the model chain, and the simulated changes were not larger in magnitude than the ensemble spread-change magnitudes were shown to be within the range of the overall uncertainty estimates, highlighting the importance of such uncertainty assessments in effect studies to frame the quantitative model results.

Arheimer and Lindström (2015) studied future changes in annual maximum and minimum daily flows. Their projections suggested that snow-driven spring floods in the northern-central part of Sweden may occur about one month earlier than today and rain-driven floods in the southern part of Sweden may become more frequent. The boundary between the two flood regimes is projected to shift northward.

Past observations (see Section 3.2.2.1) and future projections (e.g. Graham, 2004) suggest a temporal shift in the seasonality of the river discharge, with decreasing flow in spring/summer and increasing flow in winter. Global warming and river regulation due to hydropower production cause similar changes. However, in snow-fed rivers globally the impact of climate change is projected to be minor compared to river regulation (Arheimer et al., 2017).

3.3.2.2 Land nutrient inputs

Projected changes in riverine discharge and nutrient inputs from the Baltic Sea drainage basin (BSDB) to Baltic Sea-coastal waters have been studied using a number of modelling frameworks in recent years. Projecting the regional effects of future climate and environmental change on hydrology and nutrient turnover poses challenges in terms of (i) the complex nature of the modelled system, including human influence on riverine nutrient inputs and transport processes alike, which necessitates long projection model chains and leads to uncertainty in modelled

hydrologically driven responses, and (ii) the significance of changes in human behaviour, e.g. in terms of land management, population, or nutrient emissions from point sources, which adds complexity to the formulation of scenarios for future change, on top of the climate change signal. Hydrological impact studies in the [catchment areaBSDB](#) (and elsewhere) therefore often explicitly use simplifying assumptions in order to reduce complexity of the modelled system and to put focus on certain aspects of impacts of projected changes.

Hesse et al. (2015) ~~also~~ reported increasing discharges in a ~~similar~~ model study of the Vistula lagoon catchment, using a hydrological model (SWIM), which also allows for nutrient load assessment, and climate change impact modelling based on a climate model ensemble. On average, results showed decreasing trends for nitrogen and phosphorus inputs, but a wide range of projections with individual ensemble members.

Hägg et al. (2014) used a split model approach to project changes in TN and TP inputs to Baltic Sea sub-basins. Changes in discharge were estimated with a hydrological model (CSIM) combined with a climate projection ensemble, which sampled a range of climate model and emission scenario combinations. Inputs were then calculated with a statistical model, based on modelled discharges and population as a proxy for human nutrient emissions, combining population change assumptions with climate projections. Results showed a general trend towards higher nutrient inputs across the region as a result of climate change, and a significant (i.e. potentially trend-changing) influence of human ~~nutrient input reductionadaptation~~ scenarios, particularly in the southern half of the [Baltic Sea catchment basinBSDB](#).

Øygarden et al. (2014) used measurements in a number of small agricultural catchments to establish functional relationships between precipitation, runoff, and N losses from agricultural land, and qualitatively related their findings to projected precipitation changes across the [Baltic Sea drainage basinBSDB](#) under climate change scenarios, as well as to ~~anthropogenicmitigation~~ measures to counteract the climate-driven effects ~~on nutrients~~. The analyses showed a positive relationship between runoff and N losses as well as between rainfall intensity and N losses, but stressed the wide range of feedback loops possible between climate change effects and ~~anthropogenicadaptation~~ measures, through management or policy changes. Such data-driven approaches avoid uncertainties related to effect-model chains at the expense of direct ~~basinBSDB~~-wide quantitative effect projections.

The potential effects of ~~socio-economic-nutrient input abatement scenariosadaptation~~ under climate change conditions were investigated by Huttunen et al. (2015) in a study of Finnish catchments draining to the Baltic Sea. A national nutrient load model (VEMALA) was combined with a mini-ensemble of climate ~~modeleffects~~, and then a number of agricultural ~~adaptation~~-scenarios were derived, based on crop yield and policy changes, and an economic model (DREMFA) was used to translate scenario assumptions to changes in the nutrient load model for evaluation of effects. On average, increased precipitation led to increased annual discharge and a shift from spring to winter peaks, with ~~total-nitrogen-(TN)~~ and ~~total-phosphorus-(TP)~~ inputs increasing with the discharge. Here, adaptation scenarios had less effect than climate change, with some regional variation, but significantly

different load reductions were found among assessed adaptation strategies, leading to the conclusion that adaptation measures are important for overall climate change effect mitigation in the region.

The relative importance of management decisions for TN and TP load effects was studied also by Bartosova et al. (2019), using ~~at the~~ hydrological model (E-HYPE) ~~applied for the entire Baltic Sea drainage basin on the full BSDB.~~ The ensemble approach combined climate and socioeconomic pathways based on IPCC fifth assessment data (Zandersen et al., 2019), where socioeconomic changes were directly translated into changes of the effect model setup. The influence of ~~socioeconomic adaptation choices on~~ nutrient inputs ~~abatement strategies to the Baltic Sea~~ were shown to be in the same magnitude range as climate effects, thus indicating the importance of effective ~~nutrient input~~ mitigation strategies for the region. In order to increase this efficiency, Refsgaard et al. (2019) developed and explored the concept of spatially differentiated measures for TN load reductions in the ~~Baltic Sea drainage basin BSDB~~, based on the realization that measures are not uniformly efficient over large area, and should therefore not be uniformly applied either.

3.3.3 Terrestrial biosphere

In the following, we focus on the European drought in 2018, to study the impact of very warm conditions on the terrestrial ecosystem, and on projections for the terrestrial ecosystems in the Arctic, because of the particularly strong climate warming in the Arctic and potentially strong feedbacks from the release of CO₂ and CH₄ in the northernmost part of the Baltic Sea region. Finally, we discuss mitigation scenarios for land use and land-cover changes associated with the Paris Agreement.

Terrestrial ecosystems in the European drought year 2018

The summer of 2018 saw extremely anomalous weather conditions over Europe, with high temperatures everywhere, as well as low precipitation and high incoming radiation in western, central and ~~n~~Northern Europe (Peters et al., 2020). These extreme weather conditions resulted in severe drought (indicated by soil moisture anomalies) in western, central and ~~n~~Northern Europe, including the entire Baltic Sea region. The impacts of the severe drought and heatwave in Europe in 2018 were investigated in a series of papers, ranging from individual sites to the continental scale (Peters et al., 2020).

Graf et al. (2020) studied the effects of the 2018 drought conditions on the annual energy balance at the land surface, in particular the balance between sensible and latent heat fluxes, across different terrestrial ecosystems at various sites in Europe. Graf et al. (2020) found a 9% higher incoming solar radiation compared to their reference period across the drought-affected sites. The outgoing shortwave radiation mostly followed the incoming radiation, with an increase of 11.5%, indicating a small increase in surface albedo. The incoming longwave radiation, on the other hand, did not change significantly, indicating that effects of higher atmospheric temperatures and reduced cloudiness cancelled out, while outgoing longwave radiation increased by 1.3% as a result of higher land surface temperatures. Overall, the net radiation increased by 6.3% due to the extreme drought conditions. As for the non-radiative surface energy fluxes, the sensible heat flux showed a strong increase by 32%, while the latent heat fluxes did not change significantly on average. Graf et al. (2020) attributed the negligible effect on latent heat fluxes to

the opposing roles of increased grass reference evapotranspiration on the one hand and soil water depletion, stomatal closure and plant development on the other. Evapotranspiration increased where and when sufficient water was available and later decreased only where stored soil water was depleted. As a consequence, latent heat fluxes typically decreased at sites with a severe precipitation deficit, but often increased at sites with a comparable surplus of grass reference evapotranspiration but only a moderate precipitation deficit. Consistent with this, peatlands were identified as the only ecosystem with very strong increases in latent heat fluxes but insignificant changes in sensible heat fluxes under drought conditions. Crop sites, on the other hand, showed significant decreases in latent heat fluxes.

Lindroth et al. (2020) analysed the impact of the drought on Scandinavian forests, based on 11 forest ecosystem sites differing in species composition, i.e. spruce, pine, mixed and deciduous. Compared to their reference year, in 2018 the forest ecosystem showed a slight decrease in evaporation at two of the sites, was nearly unchanged at most sites and increased at two sites with pine forest. At the same time, the mean surface conductance during the growing season was reduced 40-60% and the evaporative demand increased 15-65% due to the warm and dry weather conditions. The annual net ecosystem productivity (NEP) decreased at most sites, but the reasons differed. At some sites, the NEP decrease was due to an increase in ecosystem respiration (RE), while at others both RE and the gross primary productivity (GPP) decreased, with the decrease in GPP exceeding that in RE. At six sites, the annual NEP decreased by over 50 g C m⁻² year⁻¹ in 2018. Across all sites considered, NEP anomalies varied from -389 to +74 g C m⁻² year⁻¹. A multi-linear regression analysis revealed that the anomalous NEP could to a very large extent (93%) be explained by anomalous heterotrophic respiration and reduced precipitation, with most of the variation (77%) due to the heterotrophic component.

Rinne et al. (2020) studied the effects of the drought on greenhouse gas exchange in five northern mire ecosystems in Sweden and Finland. Due to low precipitation and high temperatures, the water table sank in most of the mires. This led not only to a lower CO₂ uptake, but also to lower CH₄ emissions by the ecosystems. Three out of the five mires switched from sinks to sources of CO₂. Estimates of the radiative forcing expected from the drought-related changes in greenhouse gas fluxes indicated an initial cooling effect due to the reduced CH₄ emissions, lasting up to several decades, followed by a warming caused by the lower CO₂ uptake. However, it is unknown whether these results can be generalized to all wetlands of the Baltic Sea region.

Terrestrial ecosystems in the Arctic region

Climate warming has been particularly strong at high northern latitudes, and climate change projections indicate that this trend will continue, due to the anticipated increase in anthropogenic climate forcing. This strong warming is expected to have major consequences for terrestrial ecosystems in Arctic and sub-Arctic regions.

Zhang et al. (2013) used the Arctic version of a dynamic global vegetation model (LPJ-GUESS, Smith et al., 2001), forced with a regionalised climate scenario (A1B anthropogenic emission scenario), to investigate land surface feedbacks from vegetation shifts and biogeochemical cycling in terrestrial ecosystems under future climate warming. They found marked changes in vegetation by the second half of the 21st century (2051-2080), i.e. a poleward advance of the boundary between forests and tundra, expansion of tundra covered with tall shrubs and a

2787 shift from deciduous trees, e.g. birch, to evergreen boreal coniferous forest. These changes in vegetation were
2788 associated with decreases in surface albedo, particularly in winter due to the snow-masking effect, and with
2789 increases in evapotranspiration. The reduced surface albedo would tend to enhance the projected warming (positive
2790 feedback), while increased evapotranspiration would dampen it (negative effect). The terrestrial ecosystems
2791 continued to act as carbon sinks during the 21st century, but at diminished rates in the second half of the century.
2792 The initial increase in carbon sequestration, due to a longer growing season and CO₂ fertilisation, could be reduced
2793 and eventually reversed by increased soil respiration and greater CO₂ release from increased wildfires. Peatlands
2794 were identified as hotspots of CH₄ release, which would further enhance the projected warming (positive
2795 feedback).

2796 Using a regional ~~ESMEarth-System-Model~~ (RCA-GUESS; Smith et al., 2011) over the Arctic region, Zhang et al.
2797 (2014b) investigated the role that the biophysical effects of the projected future changes of the land surface play
2798 for the terrestrial carbon sink in the Arctic region under a future climate scenario based on a high emission scenario
2799 (RCP8.5). Two simulations were performed to determine the role of the biophysical interactions, one with and one
2800 without the biophysical feedbacks resulting from the simulated climatic changes to the terrestrial ecosystems in
2801 the model. In both simulations the Arctic terrestrial ecosystems continued to sequester carbon until the 2060-
2802 2070s, after which they were projected to turn into weak sources of carbon, due to increased soil respiration and
2803 biomass burning. The biophysical effects were found to markedly enhance the terrestrial ecosystem carbon sink,
2804 particularly in the tundra areas. Two opposing feedback mechanisms, mediated by changes in surface albedo and
2805 evapotranspiration, contributed to the additional carbon sequestration. The decreased surface albedo in winter and
2806 spring notably amplified warming in spring (positive feedback), while the increased evapotranspiration led to a
2807 marked cooling during summer (negative feedback). These feedbacks stimulated vegetation growth due to an
2808 earlier start of the growing season, leading to changes in woody plant species and the distribution of vegetation.
2809 In a later study, Zhang et al. (2018) found that these biophysical feedbacks play essential roles also in climate
2810 scenario simulations with weaker anthropogenic climate forcing.

2811 *Mitigation*

2812 The beneficial effects of carbon sequestration by forest ecosystems on climate change may be reinforced,
2813 counteracted or even offset by management-induced changes in surface albedo, land-surface roughness, emissions
2814 of biogenic volatile compounds, transpiration and sensible heat flux (see above). Luyssaert et al. (2018)
2815 investigated the trade-offs associated with using European forests to meet the climate objectives in the Paris
2816 Agreement. ~~A central argument of this study was that the agreement requires more than that forest management~~
2817 ~~should dampen the rise in atmospheric CO₂ and reduce the radiative imbalance at the top of the atmosphere. The~~
2818 ~~authors argued that a more comprehensive assessment of forest management as a strategy to achieve the goals of~~
2819 ~~the Paris Agreement should go beyond the reduction of atmospheric CO₂ and, thus, of the radiative imbalance at~~
2820 ~~the top of the atmosphere. They~~~~The authors~~ suggested two additional targets, that forest management should
2821 neither increase the near-surface temperature nor decrease precipitation, because climate effects arising from the
2822 changes in the terrestrial biosphere would make adaptation to climate change more demanding. Analysing different
2823 forest management portfolios in Europe designed to maximize the carbon sink, maximize the forest albedo or

Formatiert: Schriftart: Nicht Kursiv, Nicht unterstrichen,
Schriftfarbe: Automatisch

reduce near-surface temperatures, Luyssaert et al. (2018) found that only the portfolio designed to reduce near-surface temperatures accomplished two of the objectives, i.e. to dampen the rise in atmospheric CO₂ and to reduce near-surface temperatures. This portfolio featured a decrease in the area of coniferous forest in favour of a considerable increase in the area of deciduous forest in Northern Europe, from 130,000 to 480,000 km².

3.3.4 Cryosphere

3.3.4.1 Snow

~~There is agreement among models that the average amount of snow accumulated in winter will decrease by over 70% in most of the Baltic Sea region. The high Scandinavian mountains, where the warming temperature will not reach the freezing point as often as in lower-lying regions, are an exception (Christensen et al., 2021). The reduction in snow amount is slightly larger than in maps presented by the BACC II Author Team (2015), which is consistent with the stronger average warming projected in the RCP8.5 scenario, compared to the SRES A1B scenario analyzed by the BACC II Author Team (2015). Generally, the relative decrease in snow amount is projected to be smaller in the colder northern and eastern parts of the Baltic Sea region than in the milder southern and western parts (e.g. Räisänen, 2021). There is agreement among the EURO-CORDEX RCMs that the average amount of snow accumulated in winter will decrease by around 50% for land areas north of 60°N for the RCP8.5 scenario by 2071-2100 relative to 1981-2010 (Christensen et al., 2021). South of 60°N, the corresponding decrease is almost 80%. The reduction in snow amount is slightly larger than in figures presented by the BACC II Author Team (2015), which is consistent with larger average warming and, in the northern part of the area, smaller precipitation increase, projected in the RCP8.5 scenario compared to the SRES A1B scenario analyzed by the BACC II Author Team (2015).~~

For Poland, two additional downscaling experiments were made to produce reliable high-resolution climate projections of precipitation and temperature, using the RCP4.5 and RCP8.5 scenarios (Szwed et al., 2019). The results were used as input to a snow model (seNorge), to transform bias-adjusted daily temperature and precipitation into daily snow conditions. The snow model projected future snow depth to decrease in autumn, winter and spring, in both the near and far future. ~~The maximum snow depth was projected to decrease 15-20% by 2021-2050 and at least double that decrease by 2071-2100 (Szwed et al., 2019). The maximum snow depth was projected to decrease 15-25% by 2021-2050 in both scenarios. By 2071-2100, decreases under RCP4.5 and RCP8.5 were estimated to be 18-34% and 44-60% respectively (Szwed et al., 2019).~~

3.3.4.2 Glaciers

The Special Report on the Ocean and Cryosphere in a Changing Climate (IPCC, 2019a) provides the most recent assessment of future projected glacier mass reduction under various RCPs, and treats Scandinavian glaciers separately (Hock et al., 2019). Previous projections, summarized in the Fifth Assessment Report of the IPCC (IPCC, 2014a; Vaughan et al., 2014), did not specifically focus on Scandinavian glaciers.

By 2100, likely (i.e. with a likelihood of 66-100%) mass losses for high-mountain glaciers are 22-44% (RCP2.6) to 37-57% (RCP8.5) of their mass in 2015. These losses exceed global projections for glacier mass loss of 18 ±

7% for RCP2.6, and $36 \pm 11\%$ for RCP8.5 (likely ranges, IPCC, 2019a). Glaciers in Scandinavia will lose ~~more~~ ~~than ever~~ 80% of their current mass by 2100 under RCP8.5 (medium confidence), and many are projected to disappear, regardless of future emission scenarios (Fig. 287). Furthermore, river runoff from glaciers is projected to change regardless of emission scenario (high confidence), and to result in increased average winter runoff (high confidence) and in earlier spring peaks (high confidence; Hock et al., 2019).

Projections of future glacier mass loss depend crucially on climate projections providing surface air temperature and precipitation as forcing factors in process-based glacier models. For high mountain glaciers, such as those along the Scandinavian mountains that drain into the Baltic Sea, this is challenging, as the interplay of regional effects such as high-mountain meteorology and elevation-dependent warming (Wang et al., 2016; Qixiang et al., 2018) with global climate is poorly understood (Hock et al., 2019). Surface air temperatures in mountain regions are projected to increase at an average rate of $0.3 \pm 0.2^\circ\text{C}$ per decade until 2050 (very high confidence), i.e. faster than the present global average of $0.2 \pm 0.1^\circ\text{C}$ (Hock et al., 2019). ~~Beyond 2050, and depending on the emission scenario, air surface temperatures in high mountain regions are projected to either stabilize at the 2015–2050 rate or to increase further.~~ Beyond 2050, air surface temperatures in high mountain regions are projected to increase under a high emission scenario (RCP8.5), and to stabilize at 2015–2050 levels under a low emission scenario (RCP2.6) (IPCC, 2019a).

Projected changes in surface air temperature for the period 2071–2100 (compared to 1971–2000, under various emission scenarios) for the part of the Baltic Sea Drainage Basin that extends along the Scandinavian mountains (SMHI, 2020; Kjellström et al., 2016) will be of great importance for the assessment of future mass loss from glaciers draining into the Baltic Sea.

3.3.4.3 Permafrost

Due to recent warming more than 20% of the permafrost in the region was already lost in 1997–2018 (Figure 13; Obu et al., 2020). As warming increases, so will loss of permafrost (Section 1.5.2). Global projections show very limited permafrost in the region already at $+2^\circ\text{C}$ (Chadburn et al., 2017), but ~~the global projection~~ (including Chadburn et al., 2017) do not account for peatland permafrost, which can persist for centuries outside of its climate equilibrium (Osterkamp and Romanovsky, 1999). Much of the permafrost in Baltic Sea region was very close to its climatic boundary even before the recent acceleration of climate warming. Much of the lowland permafrost in palsas and peat plateaus in this region is very close to the 0°C thawing point, and is likely relict permafrost, persisting from the Little Ice Age (Sannel et al., 2016). Observations also show that lowland permafrost thaw has been going on for decades (Åkerman and Johansson, 2008). Preliminary analyses of permafrost loss in 1997–2018 suggests that this was roughly equally divided between alpine and lowland permafrost (22 and 24%, respectively, Figure 13), in agreement with projections of loss of all types of Baltic permafrost in the future.

Permafrost thaw by climate warming is known to affect river runoff and its loads of carbon, nutrients and contaminants, such as mercury (Schuster et al., 2018; Vonk et al., 2015). The local effect of permafrost thaw in alpine headwaters can be significant (Lyon et al., 2009), but in the Baltic Sea Basin alpine permafrost thaw will

Formatiert: Schriftart:

likely have limited influence on the characteristics of river transport at their mouths on the Baltic Sea. This is because the alpine permafrost in the Baltic Sea drainage basin mainly affects solid bedrock or regolith, with almost no soil organic matter stored in permafrost (Fuchs et al., 2015). Thaw of permafrost in peatlands affects soils with very large stocks of organic material, and has been suggested to cause large losses of peat carbon and nutrients into aquatic ecosystems (Hugelius et al., 2020). However, these projections are highly uncertain and based on studies of peatland thaw chronosequences in North America that may not be applicable to Fennoscandian permafrost peatlands (though see Tang et al., 2018).

The extent of permafrost in the Baltic Sea drainage basin may decrease significantly in this century, and depending on which warming trajectory the Earth takes, may disappear altogether in the coming century. The thaw of alpine permafrost will have little effect on flows of water, carbon and nutrients to the Baltic Sea. Thawing peatlands may increase the loads of carbon, nutrients and mercury to the Baltic Sea, but these projections remain highly uncertain.

3.3.4.4 Sea ice

Two new projections for sea ice in the Baltic Sea have been produced after BACC II (BACC II Author Team, 2015). Luomaranta et al. (2014) used simplified regression and analytical models to estimate changes in sea-ice extent (Fig. 298) and fast-ice thickness. Due to their less demanding computational approach, they could base estimates on 28 CMIP5 models. As in the Arctic Ocean (Section 1.5.2), maximum annual ice extent and thickness were both estimated to decline in the future, but some sea ice will still form every year, even by the end of the century, in agreement with earlier studies (e.g. Haapala et al., 2001; Meier, 2002; Meier et al., 2004a). Under the RCP4.5 and RCP8.5 scenarios, the modelled mean maximum ice thicknesses in Kemi were projected to be 60 cm and nearly 40 cm, respectively, in 2081-2090. However, under the RCP8.5 scenario, two models projected Kemi to be ice-free.

Höglund et al. (2017) used a more advanced approach to examine changes in sea ice conditions with a coupled ice-ocean model (Hordoir et al., 2019; Pemberton et al., 2017). They used downscaled atmospheric data from the EC-Earth and the Max Planck Institute ~~ESMs~~^{Earth System models} and simulated the response of the ice for the RCP4.5 and RCP8.5 projections. Average annual maximum ice extent at the end of the century was projected to be 90 – 100 10³ km² and 30 – 40 10³ km², for the medium and high emission scenarios, respectively, and ice thickness to decrease 3 – 6 cm decade⁻¹. Höglund et al. (2017) also projected the mobility of the ice to increase, but with little effect on future ridged ice production.

3.3.4.5 Lake ice

The latest model experiments demonstrate that the Baltic Sea catchment will experience a substantial reduction in lake ice cover in the future, with many lakes becoming ice covered only intermittently (Maberly et al., 2020; Sharma et al., 2019; Sharma et al., 2021; Shatwell et al., 2019). This change will commence in the south and move northwards gradually. Lithuanian and Latvian lakes will lose their ice cover after +2°C warming, and further warming will gradually move winter ice loss northwards, so that at +8°C warming, only lakes in northernmost

Lapland will retain a winter ice cover (Maberly et al., 2020; Sharma et al., 2019; Sharma et al., 2021; Shatwell et al., 2019).

3.3.5 Ocean and marine sediments

3.3.5.1 Water temperature

Ocean temperatures are rising at accelerating rates (IPCC, 2019a; Section 1.5.3.2). For the end of this century, scenarios for the Baltic Sea project a sea surface temperature increase of 1.1°C (0.8-1.6°C, RCP2.6) to 3.2°C (2.5-4.1°C, RCP8.5) compared to 1976-2005 (Gröger et al., 2019; Gröger et al., 2021b), see Table 149. In brackets, the ensemble spreads indicated by the 5th and 95th percentiles are listed. These changes are slightly larger than the projected global sea surface temperature changes (Section 1.5.3.2). Other ensembles than the one by Gröger et al. (2019) give similar results that vary between 1.9°C (RCP4.5) and 2.9°C (RCP8.5) for the ensemble mean temperature increase (Meier et al., 2021b; see also Meier and Saraiva, 2020). By the end of the century, sea surface temperature changes for the RCP8.5 scenarios significantly exceed natural variability. Largest open-sea warming is found in summer in the northern Baltic Sea, due to earlier melting of the sea ice (Figs. 3029 and 3130). Even higher warming of +2–6°C (the range denotes RCP2.6 and RCP8.5 scenarios) is projected for the Curonian Lagoon by the year 2100 (Jakimavičius et al., 2018). The north/south gradient and pronounced seasonality of SST trends would be reduced in the future due to vanishing sea ice (Dutheil et al., 2021).

The main driver of interannual variations of monthly mean sea surface temperature is air temperature, through the sensible heat fluxes (Meier et al., 2021a). The second most important drivers are cloudiness over the open sea and latent heat flux and meridional and zonal wind velocities over coastal areas, the latter probably because of upwelling (Meier et al., 2021a). In the vertical, the surface layer is warming more than the winter water, which is sandwiched between the surface layer and the halocline. Hence, the spring and summer thermoclines are getting more intense (Gröger et al., 2019). Water temperature trends in the deep water of those sub-basins such as Bornholm Basin and Gotland Basin that are sporadically ventilated by salt-water inflows originating from surface water are projected to be elevated as well (Meier et al., 2021a). Projected changes of the vertical water temperature distribution are similar ~~to~~ those observed since 1850 (Kniebusch et al., 2019a).

For extreme events, projections suggest, inter alia, more tropical nights over the Baltic Sea, increasing the risk of record-breaking water temperatures (Meier et al., 2019a).

3.3.5.2 Salinity and saltwater inflows

Future changes in salinity will depend on changes in the wind fields over the Baltic Sea region (Lass and Matthäus, 1996), river runoff from the Baltic Sea catchment (Schinke and Matthäus, 1998) and mean sea level rise relative to the seabed of the sills in the entrance area (Meier et al., 2017; Meier et al., 2021b). A projected increase in river runoff will tend to decrease salinity, but sea level rise will have the opposing effect of tending to increase salinity, because the water level above the sills at the Baltic Sea entrance would be higher, increasing the cross-sectional area of the Danish straits. As a result, saltwater imports from Kattegat would be larger. A 0.5 m higher sea level relative to the sill bottom at the end of the century would increase estimated Gotland Deep surface salinity by 0.7

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

g kg⁻¹ and bottom salinity by 0.9 g kg⁻¹ (Meier et al., 2017; Meier et al., 2021b). Furthermore, hypothetically increasing westerly wind will block the freshwater flow out of the Baltic Sea. Consequently, the saltwater inflow is reduced due to mass conservation (Meier and Kauker, 2003).

Due to the large uncertainty in projected changes in wind fields over the Baltic Sea region (Section 3.3.1.5), in changes of the freshwater supply from the catchment (section 3.3.2.1) and in global sea level rise (Section 3.3.5.4), salinity projections show a wide spread. No robust changes were identified because the two main drivers, river runoff and sea level rise, approximately compensate each other (Meier et al., 2021b). According to Saraiva et al. (2019b) river runoff would increase by about 1 to 21% at the end of the century depending on the climate model under both RCP4.5 and RCP8.5, in the ensemble mean causing a decrease in surface and bottom salinity at Gotland Deep of about 0.6-0.7 g kg⁻¹, with a large spread among the ensemble members. Assuming a negligible global sea level rise, the intensity and frequency of MBIs were projected to slightly increase due to changes in the wind fields (Schimanke et al., 2014). Hence, in ensemble studies that considered all potential drivers, no significant changes in salinity were projected as the ensemble mean (Meier et al., 2021b). In case of salinity, global climate model uncertainty was identified to be the largest of all uncertainties (Meier et al., 2021b).

3.3.5.3 Stratification and overturning circulation

Model based estimations of future stratification are still rare and depend critically on how well the models project changes in the three-dimensional distributions of temperature and salinity. A first systematic attempt using a high resolution coupled ocean - atmosphere model and five different global climate models (Gröger et al., 2019) explored future stratification under RCP8.5. They assumed a 10% increase in river runoff (approximately the ensemble mean in Saraiva et al., 2019b) and an unchanged mean sea level in the North Sea at the end of the century. The ensemble consistently indicated a basin-wide intensification of the pycnocline (by 9–35%) for nearly the whole Baltic Sea, and a shallowing of the pycnocline depth in most regions, except the Gulf of Bothnia (Gröger et al., 2019). The area with a pycnocline intensity > 0.05 kg m⁻³m⁻¹ increased 23-100%. The warm season thermocline likewise intensified in nearly the entire Baltic Sea (Gröger et al., 2019).

All ensemble members indicate a strengthening of the zonal, wind driven near-surface overturning circulation in the southwestern Baltic Sea towards the end of the 21st century, whereas the zonal overturning at depth is reduced by ~ 25% (Gröger et al., 2019). In the Baltic proper, the meridional overturning shows no clear climate change signal. However, three out of five ensemble members indicate at least a northward expansion of the main overturning cell. In the Bothnian Sea, all ensemble members show a significant weakening of the meridional overturning.

As the study by Gröger et al. (2019) and previous projections (e.g. Meier et al., 2006) do not consider global sea level rise, these scenario simulations are no longer considered plausible (Meier et al., 2021a; 2021b). Considering all drivers of changes in salinity in the Baltic Sea (wind, river runoff, global sea level rise), neither the haline induced stratification nor the overturning circulation is projected to change systematically among climate models (Meier et al., 2021a). It was found that under a RCP4.5 or RCP8.5 scenario a linearly rising mean sea level by the

3010 figures suggested by IPCC (2019b) would approximately counteract the effects of projected river runoff increases
3011 and wind changes on salinity.

3012 3.3.5.4 Sea level

3013 Global mean and thus Baltic Sea level will continue to rise at an increasing rate. During this century, melting ice
3014 sheets in Antarctica and Greenland are expected to contribute more to the total sea level than in the past (e.g.
3015 Mitrović et al., 2018). The fingerprints from melting ice sheets in Antarctica on sea level rise will be more
3016 pronounced in the northern hemisphere and introduce large uncertainties for Baltic sea level rise. On the other
3017 hand, ice melt in Greenland has a relatively modest effect on sea level in the Baltic Sea. Furthermore, the sea level
3018 in shelf seas such as the Baltic Sea will rise more strongly than one would expect from the thermostatic expansion
3019 of the local water column only, due to spill-over effects from the open ocean (Landerer et al., 2007; Bingham and
3020 Hughes, 2012). In addition, the long-term rate of coastal land rise is not easy to estimate accurately, due to the
3021 limited length of Global Positioning System (GPS) measurements, and frequently revised geological model values.

3022
3023 Estimates for the ensemble mean global sea level rise by 2100 ranged from 43 cm (RCP2.6) to 84 cm (RCP8.5),
3024 with likely ranges of 29-56 cm and 61-110 cm, respectively (IPCC, 2019a), cf. Section 1.5.3.1 (note that the figures
3025 in Section 1.5.3.1 refer to the mean changes between 1986-2005 and 2081-2100 and the previous IPCC AR5
3026 (2014b) with 10 cm lower changes in sea level for RCP8.5 compared to IPCC (2019a)). In particular for RCP8.5,
3027 sea level rise projections by the fifth IPCC assessment report (IPCC, 2014a) are somewhat lower than those
3028 from the more recent Special Report on the Ocean and Cryosphere in a Changing Climate (IPCC, 2019a) because
3029 of the updated contribution from Antarctica based upon new ice-sheet modeling.

3030
3031 The projected sea level rise relative to land in the Baltic Sea entrance area was estimated to about 80% of the
3032 global rate (Grinsted et al., 2015; Grinsted, 2015). These results were confirmed by other studies (e.g. Kopp et al.,
3033 2014) and summarized by Pellikka et al. (2020) who found a regional ensemble mean absolute sea level rise of
3034 87% of the global sea level rise. For the period 2090-2099 relative to 1980-1999 and based on the SRES A1B
3035 scenario, the projected absolute sea level rise in the Baltic Sea was estimated to be about 80% of the global increase
3036 (Grinsted et al., 2015). These results were confirmed by other studies for other scenarios and slightly different
3037 reference periods (e.g. Kopp et al. 2014; Grinsted, 2015), and summarized by Pellikka et al. (2020) who, for the
3038 period 2000-2100, documented an ensemble mean absolute sea level rise in the Baltic Sea of about 87% of the
3039 global mean sea level rise.

3040
3041 Altogether, considering land uplift and eustatic sea level rise, very likely ranges (5-95% probability) of relative
3042 sea level change between 2000 and 2100 under the most pessimistic IPCC emissions scenario (RCP8.5) were
3043 projected to, e.g. 29-162 cm in Copenhagen (median 68 cm), -13 -117 cm in Stockholm (median -25 cm), and 21-
3044 151cm in St. Petersburg (median 59 cm) (Grinsted et al., 2015). For coastal sites in the northern Baltic Sea, relative
3045 sea level changes in the Gulf of Finland in 2000–2100 were projected to be +29 cm (–22 to+92 cm), –5 cm (–66
3046 to+65 cm) for the Bothnian Sea, and –27 cm (–72 to +28 cm) for the Bothnian Bay, where the land uplift is larger
3047 (Johansson et al., 2014). The ranges in the latter study were estimated from the 5% and 95% cumulative

Formatiert: Schriftart: Nicht Kursiv, Nicht unterstrichen,
Schriftfarbe: Automatisch

Formatiert: Schriftart: Nicht Kursiv, Nicht unterstrichen,
Schriftfarbe: Automatisch

Formatiert: Schriftart: Nicht Kursiv

3048 probabilities considering several published scenarios from the third and fourth IPCC assessment reports. In a recent
3049 study based upon IPCC (2019a), Hieronymus and Kalén (2020) also estimated a sea-level fall in the northern
3050 Baltic Sea and a 70 cm rise in the south by 2100, [based upon a RCP8.5 scenario](#). These upper bounds of the sea
3051 level rise projections imply a very strong future acceleration of present rates. Current observations seem to show
3052 an acceleration, but its present magnitude is still small (Hünicke and Zorita, 2016; see Section 3.2.5.4).

3053
3054 Recent efforts since the IPCC AR5 report (IPCC, 2014a) that focused on the contribution of Antarctic ice sheets
3055 to global mean sea level rise have shown that ~~the interaction of~~ warming ocean water, melting the ice sheets from
3056 below can lead to instabilities in the ice sheet dynamics. The ice sheets flowing from land into the ocean are in
3057 contact with the ocean floor out to the grounding line. From there on outward the ocean is melting the ice from
3058 below and the ice sheets become thinner and lighter. If the weight of the ice sheet becomes less than the weight of
3059 the ocean water it replaces, it floats up and away. The grounding line retreats inland where the ice sheet is thicker
3060 and the ice flow larger and reinforces the ice loss (Mercer, 1978). This and related feedback loops could lead to an
3061 extra meter of sea level rise until the end of the century (e.g. Sweet et al., 2017). The most recent estimates based
3062 on expert judgement (Bamber et al., 2019) for global mean sea level rise in 2100 relative to 2000, including these
3063 potential contributions (including land water storage) are 69 cm and 111 cm for low and high sea level scenarios,
3064 respectively. For the high sea level scenario the likely range (5 to 95%) is between 62 cm and 238 cm.

3065
3066 Future changes in sea level extremes in the Baltic Sea depend on future changes in mean sea level and future
3067 developments in large-scale atmospheric conditions associated with changing wind patterns. Model projections
3068 disagree regarding atmospheric circulation changes and therefore their relevance for extreme future sea levels
3069 remains unclear (Räisänen, 2017). Absolute mean sea levels will continue to rise in the entire Baltic Sea, but exact
3070 rates remain uncertain and depend on models and greenhouse gas emission scenarios (Grinsted, 2015; Hieronymus
3071 and Kalén, 2020). Relative sea level changes will strongly vary across the Baltic Sea because of the existing spatial
3072 gradient in glacial isostatic adjustment and the spatial inhomogeneity associated with the uncertain relative
3073 contributions of melting from Antarctica and Greenland (e.g. Hieronymus and Kalén, 2020). For the Baltic Sea,
3074 changing mean sea levels are expected to have larger effects on future extremes than changing atmospheric
3075 circulation (Gräwe and Burchard, 2012). Sea ice loss in the future will further directly expose the northern Baltic
3076 coastline to stronger storm surges.

3077
3078 Recent projections of extreme sea levels along Europe's coasts have considered all drivers by linear superposition,
3079 i.e. absolute mean sea level rise and land uplift, tides (small in the Baltic Sea), storm surges and waves
3080 (Vousdoukas et al., 2016; Vousdoukas et al., 2017). The results suggest that extreme sea levels will increase more
3081 than the mean sea level, due to small changes in the large-scale atmospheric circulation, such as a northward shift
3082 of the Northern Hemisphere storm tracks and westerlies, and increases in the NAO/[Arctic Oscillation \(AO\)](#) (IPCC,
3083 2014b). These changes in the large-scale atmospheric circulation of the Baltic Sea region are, however, not robust
3084 among GCMs, giving the projections of extreme sea levels by Vousdoukas et al. (2016; 2017) low confidence.

3.3.5.5 Waves

The few existing wave climate projections for the Baltic Sea indicate an increase in the mean wave conditions, either in the whole area (Groll et al., 2017) or in its northern part (Bonaduce et al., 2019). This increase in the mean conditions has been linked to two main drivers: 1) increased wind speeds and 2) reduced seasonal ice cover.

Groll et al. (2017) projected wave climate at the end of 21st century, based on two different scenarios. They found a slight increase in the median wind speeds for most of the Baltic Sea area, which led to an increase of up to 15 % in median Significant Wave Height (SWH). Using only one climate scenario, Bonaduce et al. (2019) found that decreased wind speed in the southern Baltic Sea led to a decrease in mean SWH, whereas increased wind speeds in the north, especially in winter, led to increased mean SWH. As neither study used multi-model ensembles of scenario simulations (an exception for the western Baltic Sea is the work by Dreier et al. (2021)), and there is ~~large~~^{high} uncertainty in the projected wind speeds and directions, which is not attributed to the decline in ice cover, the results may not be representative. The projected changes in SWH estimates are therefore inconclusive.

Ruosteenoja et al. (2019) estimated based on ~~on~~ CMIP5 simulations that in future, mean and extreme scalar wind speeds are not likely to significantly change in the Baltic Sea area. Hence, mean wave conditions would not change. They also estimated that frequency of strong westerly winds will increase while strong easterly winds will become less common. These type of changes might have more significance on the frequency of extreme SWH values and their spatial patterns.

For extreme values, these studies give even less reliable results. The results of Groll et al. (2017), Suursaar et al. (2016) and Bonaduce et al. (2019) all indicated large spatial variability in how the projected extremes changed. In addition to the wind speed, extreme values are quite sensitive to wind direction, since fetch varies with direction due to the geometry of the Baltic Sea. Mäll et al. (2020) simulated how wave conditions during three historical Baltic Sea storms would change under climate change conditions. The results showed slight, but not significant changes in extreme SWH values during the storms.

Future changes in seasonal sea-ice conditions in the northern Baltic Sea are more reliable, and their effect on the wave climate easier to estimate (Rutgersson et al., 2021). Mild ice winters have already become common, and new records of lowest annual maximum ice extent have been recorded. In the Baltic Sea, the ice season partly overlaps with the seasons of the strongest winds, namely autumn and winter. The mean and extreme values of SWH are therefore expected to increase in areas like the Bothnian Sea, which now typically has ice cover in winter, but will have ~~lost~~^{lose} it in the future Baltic Sea climate.

3.3.5.6 Sedimentation and coastal erosion

As a consequence of the probably accelerating sea level rise, coastal erosion will increase regionally, to fill the increased underwater accommodation space. How much erosion will increase will depend not only on the rate of sea level rise, but also on the intensity of storms (Zhang et al., 2017).

Coastal erosion, accretion and alongshore sediment transport are primarily controlled by winds and wind-induced waves in the Baltic Sea. Projecting the future rate of coastal erosion or accretion in the Baltic Sea is highly uncertain because of a lack of consensus in the prediction of future storms. Neglecting potential change in future storms and assuming an intermediate sea level rise scenario (RCP4.5), an increment of 0.1 -0.3 m year⁻¹ in coastline erosion has been projected for some parts of the southern Baltic Sea coast (Zhang et al., 2017; Deng et al., 2015). Due to the prevailing westerly winds, the dominant sediment transport will continue to be eastwards along most of the southern Baltic Sea coast, but with high variability along coastal sections with a small incidence angle of incoming wind-waves (Dudzińska-Nowak, 2017). It has been found that even a minor climate-change-driven rotation of the predominant wind directions over the Baltic Sea may substantially alter the structural patterns and pathways of wave-driven transport along large sections of the coastline (Viška and Soomere, 2013).

The presence of sea ice is an important factor moderating coastal erosion. Storm surges and wave run-up on the beach are much higher in ice-free periods than when there is even partial ice cover. The hydrodynamic forces are particularly effective in reshaping the shoreline when there is no ice and sediment is mobile (Ryabchuk et al., 2011). Due to global warming, both the area and duration of ice cover in the Baltic Sea will be reduced in future (Section 3.3.4.4), thus increasing coastal erosion.

Foredunes will likely continue to form on prograding coasts, but at rates influenced by the accelerating sea-level rise (Zhang et al., 2017). Fore dunes may tend to become higher, but with reduced prograding rate and wavelength, if sea level rise is accelerated or storm frequency increased. If the wind-wave climate is stable, the height of coastal foredunes on a prograding coast remains stable or increases linearly with a low to intermediate rate (<1.5 mm year⁻¹) of sea level rise. An accelerating rate of sea level rise and/or changing storm frequency will lead to a nonlinear growth in height (following a quadratic or a higher power law; Zhang et al., 2017; Lampe and Lampe, 2018). The critical threshold that separates linear and non-linear foredune growth in response to sea level rise is likely to be reached before 2050 in the RCP8.5 scenario (Zhang et al., 2017).

~~Yet to be determined a~~ Anthropogenic influence ~~affects~~imposes further uncertainty in sediment transport and coastal erosion ~~as well~~. Sediment transport and coastal erosion are relevant for coastal management, construction and protection strategies. In general two main types of management strategies exist for the Baltic Sea coast: 1) coastal protection by soft or hard measures; and 2) adaptation to coastal change, accepting that in some places the coast would be left in its natural state (BACC II Author Team, 2015). However, administrative efforts for coastal protection differ among Baltic Sea countries, even between neighboring states or nations. It has been found that engineering structures (e.g. piers, seawalls) may influence coastline change at a much larger spatial scale than the dimension of the structure itself.

3.3.5.7 Marine carbonate system and biogeochemistry

The BACC II Author Team (2015) concluded that model simulations indicated that climate change has a potential to intensify eutrophication in the Baltic Sea. However, they also showed that the implementation of nutrient load reductions according to the Baltic Sea Action Plan (BSAP, HELCOM, 2013a) may not only mitigate this effect

but may ~~even slightly~~ decrease hypoxic and anoxic areas in the Baltic Sea. ~~Note that because~~ the nutrient load abatement ~~calculations strategy~~ of the BSAP did not take the effect of climate change into account. In contrast, the business as usual nutrients input scenario may increase the hypoxic area by about 30% ~~hypoxic area~~ and even more than double the area affected by anoxia by 2100.

As atmospheric CO₂ rises, so will the concentration of CO₂ in the Baltic Sea surface water. This will influence the mean future pH, while eutrophication and enhanced organic matter production/remineralization will increase the amplitude of daily and seasonal pH fluctuations without much affecting the mean values. Taking various nutrient load input scenarios into account, projections suggested ~~It was also shown~~ that pH in the Baltic Sea surface water will decrease, in the worst-case emission scenario (atmospheric pCO₂ of 850 ppm) the pH will drop by about 0.40 by 2100, while the decrease in a more optimistic emission scenario (550 ppm) will be smaller, i.e. about 0.26 (BACC II Author Team, 2015).

3.3.5.7.1 Oxygen and nutrients

Projected warming, increased precipitation and global mean sea level rise may worsen eutrophication and oxygen depletion in the Baltic Sea by reducing air-sea fluxes and vertical transports of oxygen in the water column, intensifying internal nutrient cycling, and increasing river-borne nutrient loads due to increased river runoff (Meier et al., 2011a; Meier et al., 2012b; Meier et al., 2012c). However, the future response of deep-water oxygen conditions will depend mainly on nutrient loads from land (Saraiva et al., 2019a, b; Meier et al., 2021b; cf. Fig. 3.24). In contrast to the global ocean (see Section 1.5.4), future nutrient supplies will have a relatively larger effect on oxygen conditions and primary production than warming. With high nutrient loads, the changing climate will have a considerable negative effect, but if loads are kept low, climate effects can be small or negligible. Scenario simulations suggest that full implementation of the nutrient load reductions required by BSAP will significantly improve the eutrophication status of the Baltic Sea, irrespective of the driving global climate model (Saraiva et al., 2019b; Meier et al., 2021b) and regional coupled climate-environmental model (Meier et al., 2018a). Despite large inherent uncertainties of future projections, modeling studies suggested that the future Baltic Sea ecosystem may unprecedentedly change compared to the past 150 years (Meier et al., 2012a).

By the end of the century (2069-2098), the ensemble mean hypoxic area is projected to change only slightly under reference (-14% for RCP4.5 and -5% for RCP8.5) and high (-2% for RCP4.5 and +5% for RCP8.5) nutrient load scenarios, compared to 1976-2005 (Saraiva et al., 2019b). Nutrient loads in the reference scenario are the average loads in 2010-2012. The high, or worst, scenario assumes changes caused by a 'fossil-fuelled development', coupled to increasing river runoff (Saraiva et al., 2019a). Changes in nitrogen and phosphorus loads were estimated from assumptions on regional population growth, changes in agricultural practices, such as land and fertilizer use, and developments in sewage treatment (Zandersen et al., 2019; Pihlainen et al., 2020). Under the BSAP scenario, the ensemble mean hypoxic area will be reduced by 50-60% at the end of the century, in comparison with 1976-2005 (Saraiva et al., 2019a). The relative reductions in hypoxic area may decrease with increasing sea level (Meier et al., 2021b).

3199 In the same model ensemble (Saraiva et al., 2019a), BSAP implementation is projected to reduce the water column
3200 phosphate pool in the Baltic Sea by 59% (RCP4.5) and 56% (RCP8.5) by the end of the century, and even the
3201 reference loads would lead to a decline by 24% (RCP4.5) and 18% (RCP8.5). Also, a larger ensemble (Meier et
3202 al., 2018a) of 8 biogeochemical models forced by outputs of 7 ESMs downscaled by 4 different RCMs projected
3203 that the BSAP reduced phosphate concentrations in the Baltic proper, Gulf of Finland and Bothnian Sea despite
3204 climate change, with largest reductions in surface concentrations by approximately 3 mmol m⁻³ in the Gulf of
3205 Finland. Present day nutrient loads led to small increases in surface phosphate concentration in the Baltic proper,
3206 a small decline in the Gulf of Finland and little change in Bothnian Sea and Bothnian Bay. Little change was
3207 predicted for DIN concentrations in the Baltic proper, whereas simulations showed an increase in the Gulf of
3208 Finland and the Bothnian Sea, regardless whether nutrient loads were kept at present level or whether loads were
3209 reduced.

3210
3211 Furthermore, future projections suggested that the sea-ice decline in the northern Baltic Sea may have considerable
3212 consequences for the marine biogeochemistry, because of changing underwater light conditions and wave climate
3213 (Eilola et al., 2013). Eilola et al. (2013) found that, by the end of the century, the spring bloom would start by up
3214 to one month earlier and winds and wave-induced resuspension would increase, causing an increased transport of
3215 nutrients from the productive coastal zone into the deeper areas.

3216
3217 For the Baltic proper, the internal nutrient cycling and exchanges between shallow and deeper waters were
3218 projected to be intensified, and the internal removal of phosphorus may become weaker in future climate (Eilola
3219 et al., 2012). These effects may counteract the efforts of planned nutrient input reductions.

3220
3221 Uncertainties in projections from Baltic Sea ecosystem models have recently been systematically assessed for the
3222 first time (Meier et al., 2018a; Meier et al., 2019b; Meier and Saraiva, 2020; Meier et al., 2021b). ~~One of the~~
3223 ~~L~~larger sources of uncertainty ~~are~~ ~~is~~ ~~biases in~~ global and regional climate model ~~uncertainties~~, in particular
3224 concerning global mean sea level rise and regional water cycling (Meier et al., 2019b). The mechanism behind the
3225 correlation between large-scale meteorological conditions in the different climate periods and oxygen conditions
3226 in the Baltic Sea is not well understood and subject to ongoing research. With respect to nutrient concentrations,
3227 also uncertainties in conditions at the North Sea boundary as well as difficulties in simulating the long-term
3228 response of the Baltic Sea biogeochemical system to changes in nutrient inputs, play a role.

3229
3230 Under the BSAP scenario, mean nitrogen fixation would decrease (Meier et al., 2021b) and record-breaking
3231 cyanobacteria blooms may no longer occur in the future, but record-breaking events may reappear at the end of
3232 the century in a business-as-usual nutrient load scenario (Meier et al., 2019a).

3233 3.3.5.7.2 Marine CO₂ system

3234 The rising atmospheric pCO₂ due to anthropogenic emissions will increase the mean pCO₂ of surface seawater and
3235 thus has the potential to lower the pH. However, the magnitude of pH changes will also depend on the development
3236 of total alkalinity concentrations (A_T; Omstedt et al., 2012). Future A_T changes in the Baltic Sea will be shaped by

3237 both external inputs (riverine runoff and inflows from the North Sea) and internal generation. The latter is due to
3238 biogeochemical processes of organic matter production and remineralization, especially under euxinic conditions.
3239 Kuznetsov and Neumann (2013), who used A_T as a tracer in a model (no internal processes included)
3240 ~~calculated~~~~showed~~ that on average A_T in surface Baltic Sea waters ~~would~~~~should~~ decrease by about $150 \mu\text{mol kg}^{-1}$
3241 by 2100, a change corresponding to an assumed decrease in salinity. Simulations by Gustafsson et al. (2019) that
3242 include most of the biogeochemical processes affecting A_T (except S burial and Fe-oxide availability) showed that
3243 A_T in the central Baltic Sea in the “business as usual” scenario will first increase, by about $100 \mu\text{mol kg}^{-1}$ by 2050,
3244 and then revert to present levels by 2100. If BSAP is implemented, A_T will decrease by about $150 \mu\text{mol kg}^{-1}$ in
3245 2100 from present levels. Irrespective of the nutrient load scenario, pH is eventually expected to decrease in the
3246 central Baltic Sea due to anthropogenic CO_2 emissions. Assuming the A1B CO_2 emission scenario (pCO_2 increase
3247 to $700 \mu\text{atm}$ by 2100), pH will drop to about 7.9 and 7.8 under “business as usual” and BSAP scenarios,
3248 respectively.

3249
3250 All available future projections of changes in alkalinity are highly uncertain, as they address only a few out of
3251 multiple factors determining the alkalinity pool in the Baltic Sea, i.e. changes in salinity, river runoff, weathering
3252 in the catchment, organic matter production (eutrophication), remineralization (especially at low redox conditions),
3253 and processes in sediments (including pyrite and vivianite formation) (Kulinski et al., 2021).

3254 3.3.6 Marine biosphere

3255 3.3.6.1 Pelagic habitats

3256 3.3.6.1.1 Microbial communities

3257 The effects of climate change on microbes and the functioning of the microbial loop have been studied by
3258 experiments in which temperature, salinity, dissolved organic matter (DOM), and ~~ocean acidification~~ (OA) were
3259 manipulated. In general, microbial activity and biomass increased with increasing DOM and temperature
3260 (Ducklow et al., 2009), but effects can be mixed. For instance, an increase in DOM in the northern Gulf of Bothnia
3261 enhanced the abundance of bacteria, whereas a temperature increase (from 12 to 15°C) decreased their abundance,
3262 probably due to a simultaneous increase of bacterivorous flagellates (Nydahl et al., 2013).

3263
3264 In the southern Baltic Sea the impact of OA was limited, and the bacterial community responded primarily to
3265 temperature and phytoplankton succession (Bergen et al., 2016). In experiments where CO_2 was increased and
3266 salinity decreased (from 6 to 3), heterotrophic bacteria declined (Wulff et al., 2018). In experiments with increasing
3267 temperature (from 16 to $18\text{--}20^\circ\text{C}$) and reduced salinity (from 6.9 to 5.9 g kg^{-1}), the Baltic proper microbial
3268 community also showed mixed responses, probably due to indirect food web effects (Berner et al., 2018).

3269 3.3.6.1.2 Phytoplankton and cyanobacteria

3270 The projected increase in precipitation is expected to increase nutrient loads, especially into the northern Baltic
3271 Sea (Huttunen et al., 2015), and together with increased internal loading of nutrients, several modelling studies

project an increased phytoplankton biomass by the end of the century (Meier et al., 2012b; Meier et al., 2012c; Skogen et al., 2014; Ryabchenko et al., 2016).

Several mesocosm studies have investigated the effects of warming on southern Baltic Sea phytoplankton communities. Warming accelerated the phytoplankton spring bloom and increased primary productivity (Sommer and Lewandowska, 2011; Lewandowska et al., 2012; Paul et al., 2015). The total phytoplankton biomass still decreased, due to negative effects of warming on nutrient flux (Lewandowska et al., 2012; Lewandowska et al., 2014).

~~Ocean acidification (OA)~~ may enhance phytoplankton productivity by increasing the CO₂ concentration in the water. The biomass of southern Baltic Sea autumn phytoplankton increased in mesocosms simulating OA (Sommer et al., 2015). In many experiments OA had, however, little effect on phytoplankton community composition, fatty acid composition or biovolume in spring or autumn (Paul et al., 2015; Bermúdez et al., 2016; Olofsson et al., 2019).

It has been suggested that climate change may increase the blooming of toxic species, such as the dinoflagellate *Alexandrium ostenfeldii* (Kremp et al., 2012; Kremp et al., 2016) and the cyanobacterium *Dolichospermum* sp. (Brutemark et al., 2015; Wulff et al., 2018). There are also contradictory results, indicating that OA and warming may decrease the biomass of *Nodularia* sp. and *Dolichospermum* sp. (Eichner et al., 2014; Berner et al., 2018). Several modelling studies project increases in cyanobacteria in the warmer and more stratified future Baltic Sea (Meier et al., 2011b; Andersson et al., 2015; Neumann et al., 2012; Chust et al., 2014; Hense et al., 2013), but other modelling studies project that the environmental state of the Baltic Sea will be significantly improved, and extreme cyanobacteria blooms will no longer occur if BSAP is fully implemented (Meier et al., 2018a; Meier et al., 2019a; Saraiva et al., 2019a; see Figure 332).

3.3.6.1.3 Zooplankton

The effects of increasing temperature and ~~ocean acidification (OA)~~ on zooplankton have been studied experimentally. In *Acartia* sp., a dominant copepod in the northern Baltic Sea, warming decreased egg viability, nauplii development and adult survival, and both warming and OA had negative effects on adult female size (Garzke et al., 2015; Vehmaa et al., 2016; Vehmaa et al., 2013).

In contrast, the effects of climate change on microzooplankton (MZP) seem to be mostly beneficial. Warming improved the growth rate of southern Baltic Sea MZP, which led to a reduced time-lag between phytoplankton and MZP maxima, improving the food supply to microzooplankton in warm conditions (Horn et al., 2015). ~~(Aberle et al., (2015)~~ showed that while protozooplankton escaped predation by slower growing copepods at low temperatures, at warmer temperatures small ciliates in particular became more strongly controlled by copepod predation.

3308 3.3.6.2 Benthic habitats

3309 3.3.6.2.1 Macroalgae and vascular plants

3310 The effects of climate change on bladder-wrack, *Fucus vesiculosus*, have been studied in a number of experiments.
3311 ~~Ocean acidification (OA)~~ appears to have a relatively small effect on macroalgae (Al-Janabi et al., 2016; Wahl et
3312 al., 2020), while temperature effects can be significant. The effects of increasing temperature are not linear,
3313 however. Growth or photosynthesis is not impaired under projected temperature increase (from 15 to 17.5°C) but
3314 at extreme temperatures (27 to 29°C), photosynthesis declines, growth ceases and necrosis starts (Graiff et al.,
3315 2015; Takolander et al., 2017). In very low salinity (2.5 g kg⁻¹), sexual reproduction of *F. vesiculosus* ceases
3316 (Rothäusler et al., 2018).

3317
3318 The direct and indirect effects of changes in temperature, salinity and pH may alter the geographic distribution of
3319 ~~many species in the Baltic Sea. Assuming a decline in salinity, a reduced penetration of marine species, such as~~
3320 ~~bladderwrack, eelgrass and blue mussel, into the Baltic Sea has been predicted (Vuorinen et al., 2015). A large~~
3321 ~~number of other species is affiliated with such keystone species, and species distribution modelling has indicated~~
3322 ~~that, e.g., many species in the Baltic Sea. Retreat of marine species has been predicted for bladder wrack, eelgrass~~
3323 ~~and blue mussel, and up to 50 other species affiliated to these keystone species (Vuorinen et al., 2015). Species~~
3324 ~~distribution modelling has indicated that~~ a decrease of bladder-wrack will have large effects on the biodiversity
3325 and functioning of the shallow-water communities of the northern Baltic Sea (Jonsson et al., 2018; Kotta et al.,
3326 2019). The responses of eelgrass, *Zostera marina*, to climate change and eutrophication mitigation have recently
3327 been modeled by Bobsien et al. (2021).

3328
3329 Experiments on climate change effects have been made also with other macroalgae and vascular plants. Thus, OA
3330 increased the growth of the opportunistic green alga *Ulva intestinalis* in the Gulf of Riga (Pajusalu et al., 2013;
3331 Pajusalu et al., 2016). Other studies showed that charophyte photosynthesis increased under high pCO₂, whereas
3332 eelgrass did not respond to the elevated pCO₂ alone (Pajusalu et al., 2015). Salinity decline is projected to decrease
3333 the distribution of *Z. marina* and the red alga *Furcellaria lumbricalis*, whereas warming will probably favour
3334 charophytes (Torn et al., 2020).

3335 3.3.6.2.2 Zoobenthos

3336 The effects of warming on invertebrates are non-linear. Respiration and growth of the isopod *Idotea balthica*
3337 increased up to 20°C, and then decreased at 25°C (Ito et al., 2019). Many marine invertebrates, including isopods,
3338 will also directly and indirectly suffer from decreasing salinity (Kotta et al., 2019; Rugiu et al., 2017), as well as
3339 ~~ocean acidification (OA)~~. The size and time to settlement of the pelagic larvae of the Baltic clam *Limecola balthica*
3340 (syn *Macoma balthica*) increased with OA, suggesting a developmental delay (Jansson et al., 2016), whereas OA
3341 had no effect on the isopod *Saduria entomon* (Jakubowska et al., 2013) or larvae of the barnacle *Amphibalanus*
3342 *improvisus* (Pansch et al., 2012).

3343

Formatiert: Schriftart:

3344 Several modelling studies have estimated the relative effects of hydrodynamics, oxygen and food availability on
3345 Baltic Sea zoobenthos ~~under various scenarios. As the assumptions about climate change and nutrient inputs differ,~~
3346 ~~a comparison between studies is impossible and final conclusions cannot be drawn.~~ In previously hypoxic areas,
3347 benthic biomass was projected to increase (until 2100) by up to 200% after re-oxygenating bottom waters, whereas
3348 in permanently oxygenated areas macrofauna may decrease by 35% due to lowered food supply to the benthic
3349 ecosystem (Timmermann et al., 2012). It has, however, been concluded that nutrient reductions will be a stronger
3350 driver for Baltic Sea ecosystem than climate change (Friedland et al., 2012; Niiranen et al., 2013; Ehrnsten et al.,
3351 2019). These studies suggest that benthic-pelagic coupling will weaken in a warmer and less eutrophic Baltic Sea,
3352 resulting in gradually decreasing benthic biomass (Ehrnsten et al., 2020).

3353 3.3.6.3 Non-indigenous species

3354 It is often suggested that climate change will favour invasions by non-indigenous species worldwide (Jones and
3355 Cheung, 2014). It has been shown that non-native benthic species typically occur in areas with reduced salinity,
3356 high temperatures, high proportion of soft seabed and low wave exposure, whereas most native species show an
3357 opposite pattern (Jänes et al., 2017). Modelled temperature and salinity scenarios suggest an increase of Ponto-
3358 Caspian cladocerans in the pelagic community, and an increase in dreissenid bivalves, amphipods and mysids in
3359 the benthos of coastal areas in the northern Baltic Sea by 2100 (Holopainen et al., 2016). Disentangling factors
3360 facilitating establishment of non-native species demands long-term surveys, and data from multiple environments
3361 in order to distinguish climate-related effects from other ecosystem-level drivers (Bailey et al., 2020). In addition,
3362 studies on changing connectivity are needed (e.g. Jonsson et al., 2020).

3363 3.3.6.4 Fish

3364 Climate change may affect Baltic Sea fish through ~~changes in effects on~~ water temperature, salinity, oxygen and
3365 pH, as well as nutrient loads, which indirectly affect food availability for fish. The responses of cod larvae to
3366 ~~OA ocean acidification~~ and warming have been studied experimentally. Some studies found no effect on hatching,
3367 survival or development rates of cod larvae (Frommel et al., 2013), while in others mortality of cod larvae doubled
3368 when exposed to high-end OA projections (RCP8.5). Several modelling studies however project low abundances
3369 of cod towards the end of the century, due to continued poor oxygen conditions (Niiranen et al., 2013; Wählström
3370 et al., 2020).

3371
3372 Climate change may also be positive for fish stocks. Warmer spring and summer temperatures have been projected
3373 to increase productivity of sprat (Voss et al., 2011; MacKenzie et al., 2012; Niiranen et al., 2013). For herring,
3374 results are more varied: both increase (Bartolino et al., 2014) and a short-term decrease (Niiranen et al., 2013)
3375 have been projected.

3376
3377 Multi-species modelling has also emphasized the role of climate for cod stocks. If fishing is intense but climate
3378 remains unchanged, cod declines, but not very dramatically, while if climate change proceeds as projected, cod
3379 disappeared in two models out of seven, even with the current low fishing effort (Gårdmark et al., 2013). Different
3380 scenarios yield very different outcomes, however. A medium CO₂ concentration scenario (RCP4.5), low nutrients

3381 and sustainable fisheries resulted in high numbers of cod and flounder, while high emissions (RCP8.5) and high
3382 nutrient loads resulted in high abundance of sprat (Bauer et al., 2018; Bauer et al., 2019). All these studies assumed
3383 a more or less pronounced decrease in salinity.

3384 **3.3.6.5 Marine mammals**

3385 *Ringed seal and grey seal – sea ice*

3386 Climate change is projected to drastically reduce the extent of seasonal sea ice in the Baltic Sea (Luomaranta et
3387 al., 2014; Meier, 2006; Meier, 2015; Meier et al., 2004b). At the end of the 21st century, ice will probably in most
3388 years be confined to the Bothnian Bay, the eastern Gulf of Finland, the Archipelago sea, and the Moonsund (Sound
3389 between the Estonian mainland and the offshore western islands Saaremaa, Hiiumaa, Muhu and Vormsi) and
3390 eastern parts of the Gulf of Riga such as Pärnu Bay (Meier et al., 2004b), with corresponding changes in the
3391 breeding and moulting distribution of ringed seals. Aside from these projections, ice cover has been even more
3392 limited in all the southern areas in recent years. Extirpation of one or more of the three southern breeding ringed
3393 seal populations is possible (Sundqvist et al., 2012; Meier et al., 2004b).

3394
3395 The ringed seal is an obligatory ice breeder that digs lairs in the snowdrifts on offshore ice for protection of the
3396 pup (e.g. Smith and Stirling, 1975). The Baltic grey seal prefers loose floes of drift ice (Hook et al., 1972), but can
3397 also breed on land (Jüssi et al., 2008). Overall pup survival in land breeding grey seals is probably lower than for
3398 ice breeders (Jüssi et al., 2008). Absence or low quality of sea ice will adversely affect pup survival and quality in
3399 ice-breeding seals. The effects can be seen by the end of the breeding season, and beyond (Jüssi et al., 2008). Grey
3400 and ringed seals are capital breeders, i.e., their pup quality depends on effective transfer of maternal energy (fatty
3401 milk) during a short, intensive lactation period. Timing of birth for both species is strongly adapted to the
3402 availability of the optimal breeding platform, sea ice. The height of the pupping season is around February-early
3403 March, when the extent and strength of the sea-ice is usually greatest. The immediate breeding success can be
3404 defined as survival and quality of the offspring at the end of the breeding season, but breeding conditions may
3405 have population consequences by affecting the survival and fitness of the pups throughout their lives (McNamara
3406 and Houston, 1996; Kahala and Kurkilahti, 2020). A warming climate with higher air and water temperatures
3407 will decrease the extent of ice-cover, the ice thickness and the overlaying snow-cover as well as the stability and
3408 duration of the ice.

3409
3410 Loss of habitat is critical for reproductive success of the ice-associated seals, especially the ringed seal, and can
3411 eventually lead to local population decreases and changes in breeding distribution, starting in the southernmost
3412 parts of its range. The ringed seal populations breeding in the Gulf of Finland, Gulf of Riga and Archipelago Sea
3413 (SW-Finland) are already small and vulnerable to any negative changes in habitat quality.

3414
3415 *Harbour seal and grey seal – flooding of haul-outs*
3416 Harbour seals and grey seals rely on undisturbed haul-out areas for key life cycle events such as breeding, moulting
3417 and resting (Allen et al., 1984; Thompson, 1989; Watts, 1996; Reder et al., 2003). In the southern Baltic Sea,
3418 relative sea levels have risen by 1 to 3 mm per year over the interval 1970-2016 (section 3.2.5.4), and increased

3419 rates of sea level rise are expected in the future (EEA, 2019c; Grinsted, 2015). A low emissions scenario for the
3420 21st century projects an additional sea-level rise of 0.29-0.59 m, a high emissions scenario an extra 0.61-1.10 m,
3421 but substantially higher values cannot be ruled out (Grinsted, 2015; IPCC, 2019a). A high emission scenario is
3422 thus likely to flood all current seal haul-outs in the southern Baltic Sea and many important localities in Kattegat,
3423 while under a low emission scenario, most haul-outs in the southern Baltic Sea will be flooded, while others will
3424 be reduced to small fractions of their current area. In the northern and central Baltic Sea and eastern Kattegat
3425 archipelago areas, seals will have alternative islets and skerries and are not likely to be affected to the same degree
3426 as in the south and in eastern Kattegat. In parts of the Gulf of Bothnia, relative sealevels may even fall, due to
3427 post-glacial rebound (EEA, 2019c).

3428

3429 *Harbour porpoise*

3430 There are no direct studies of the effects of climate change on harbour porpoises in the Baltic Sea, hence the
3431 following is based mostly on informed guesswork and on a few studies in other areas. There are a multitude of
3432 ways that changes in one parameter can affect others and we do not currently have the knowledge to predict the
3433 cumulative effects this might have on the Baltic Sea harbour porpoise population.

3434

3435 Harbour porpoises are present from Greenland to the African coast and the Black Sea and seem to have a rather
3436 wide thermal tolerance. Therefore, even though it is predicted that we will see a 1.2-3.2° increase in SST in the
3437 Baltic Sea (Section 3.3.5.1), it seems unlikely that this will directly affect harbour porpoise distribution, unless the
3438 Baltic Sea harbour porpoise is specifically adapted to colder temperatures. If this is the case, a northwards range
3439 shift might occur. With the expected future decrease in sea ice extent, the winter habitat available for the harbour
3440 porpoise in the northern Baltic Sea would increase.

3441

3442 Harbour porpoises are small cetaceans with limited capacity to store energy that mostly live in cold environments.
3443 Hence, they need to eat almost constantly (Read and Hohn, 1995; Wisniewska et al., 2016) and are therefore
3444 expected to be tightly dependent on their prey (Sveegaard et al., 2012). Their main prey species in the Baltic proper
3445 are cod (at least before the recent cod stock collapse), sprat, herring, gobies and sand eel (where present). Climate-
3446 induced changes in for example SST, fronts, stratification and to some degree currents will affect the distribution,
3447 abundance and possibly the quality of prey species, and in turn the harbour porpoise population. Their distribution
3448 may shift as they follow their prey, and potential food shortages might lead to starvation, with possible population
3449 effects.

3450

3451 It has been hypothesized that the susceptibility of marine mammals to disease may increase as temperature
3452 increases. Higher temperatures can increase pathogen development and survival rates, facilitate transmission
3453 among individuals and increase individual susceptibility to disease. The negative effects of disease as well as
3454 environmental contaminants on individual fitness will obviously worsen if the animal is also under nutritional
3455 stress.

3456

3457 *Seals and changes in the distribution of prey species*

Any large alteration of the ecosystem can affect the distribution of seals if there are climate-related changes in the abundance and distribution of their main prey species, such as herring, sprat and cod, as is possible with climate change. Such changes in top consumer distribution have been modeled in other sea areas, such as the UK continental shelf, where the current distribution of harbour seals did not match well the projected future distribution of their prey (Sadykova et al. 2020). There are large differences [between Baltic Sea climate projections of future in-salinity-between Baltic Sea models](#) (Saraiva et al., 2019b), and other factors such as temperature, eutrophication, predation and competition also affect fish distributions. Thus, future changes in abundance and distribution of seal prey species, such as herring and cod, are hard to predict (Dippner et al., 2008; Lindegren et al., 2010; Vuorinen et al., 2015; Dippner et al., 2019).

3.3.6.6 Waterbirds

Climate change scenarios agree in projecting a strong temperature increase in the Arctic and sub-Arctic. This will likely cause a northward expansion of species ranges, with colonization by new breeding and wintering species, as well as local species declines following migration of populations to ice-free northern waters (Pavón-Jordán et al., 2019; Fox et al., 2019).

If salinity in the Baltic Sea decreases, invertebrate species serving as prey for waterbirds (e.g. blue mussels for common eiders) are likely to change in distribution, body size and quality as food, with consequences for the distribution, reproduction and survival of the waterbirds that eat them (Fox et al., 2015). Predicting the consequences of climate change for piscivorous seabirds is complex, because effects are not uniform among Baltic Sea fish species. For example, expected increase of recruitment and abundance in an important prey species (sprat; MacKenzie et al., 2012; Lindegren et al., 2012) as well as declining numbers of large piscivorous fish (cod) may favour fish-eating birds, although management efforts to improve cod stocks may counteract the expected increase in sprat and lead to population declines of their main bird predator, the common guillemot (Kadin et al., 2019). Herring, another important prey species, is reported to be negatively affected by decreasing salinity (declining energy content; Rajasilta et al., 2018).

A rising sea level will reduce the area of saltmarshes available for the breeding of waders and foraging by geese (Clausen et al., 2013), and other coastal habitats would likewise be affected (Clausen and Clausen, 2014). Sea level rise in combination with storms may cause loss by erosion of current coastal breeding habitats, and flood breeding sites, thus affecting the breeding success of coastal waterbirds. Climate change can also be expected to affect waterbirds in the Baltic Sea by changing the incidence of diseases and parasites (Fox et al., 2015).

3.3.6.7 Marine food webs

Climate change and other anthropogenic environmental drivers are expected to change entire marine food webs, from coastal to off-shore, from shallow to deep, from pelagic to benthic (sedimentary), as species-distributions are impacted, and key nodes and linkages in the food webs are altered or lost (Lindegren et al., 2010; Niiranen et al., 2013; Leidenberger et al., 2015; Griffiths et al., 2017; Kotta et al., 2019; Gårdmark and Huss, 2020). These climate-driven changes will also, when combined with societal changes, affect aquatic ecosystem services, for instance

3495 future primary production (a supportive ecosystem service) and fish catches (a provisioning ecosystem service;
3496 Hyttiäinen et al., 2021).

3497
3498 Certain marine species, e.g., cod and bladderwrack, may decline in both distribution and abundance (Gårdmark et
3499 al., 2013; Takolander et al., 2017), whereas others, e.g. sprat and certain mainly costal freshwater fish, may
3500 increase (MacKenzie et al., 2012; Bergström et al., 2016). An increase in cyanobacteria blooms has also been
3501 projected, especially for the Central Baltic Sea (Meier et al., 2011b; Funkey et al., 2014), while increased flow of
3502 DOC may reduce both primary and secondary production in the northernmost low-saline areas with pronounced
3503 brown-water runoff (Wikner & Andersson, 2012; Figueroa et al., 2021). The responses also depend on human
3504 intervention, i.e. the success of nutrient reduction schemes, and are most probably non-linear (Hyttiäinen et al.,
3505 2021; Ehrnsten et al., 2020). However, it can be summarised that – if only climate change is accounted for – most
3506 studies tend to project a decline in the overall state of the ecosystem, and a long-term decline in the provision of
3507 associated ecosystem services to humans is likely if the climate change is not significantly mitigated.

Formatiert: Englisch (Großbritannien)

3508 4 Interactions of climate with other anthropogenic drivers

Formatiert: Schriftart:

3509 The term “driver” in this section is defined as something affecting a variable of the Earth system. A driver itself
3510 may be affected by other drivers, or being affected by another force. In this respect, climate is a force affecting
3511 other drivers, e.g. land use or shipping. On the other hand, (regional) climate may be affected by other drivers, e.g.
3512 land use or shipping. This section summarizes plausible two-way dependencies that have been described in the
3513 literature. For a deeper analysis, see Reckermann et al. (2021).

3514
3515 Climate change affects air and water temperature as well as precipitation, with a clear impact on land use and land
3516 cover. Growth conditions are affected by these changes, but also by political or management decisions, which may
3517 in turn be influenced by climate change (Yli-Pelkonen, 2008). Agriculture is the most important land use in the
3518 southern part of the Baltic Sea basin. Climate change strongly influences the choice of crops, as crops differ in
3519 their requirements for water availability and soil type (Fronzek and Carter, 2007; Smith et al., 2008). Still, socio-
3520 economic considerations may be even more important than climate in determining agricultural land use
3521 (Rounsevell et al., 2005; Pihlainen et al., 2020).

3522
3523 Land use and land cover can influence the regional climate, through geophysical (e.g. albedo) and biogeochemical
3524 (e.g. carbon sequestrationCO₂-drawdown) effects. Bright surfaces like agricultural fields reflect more solar
3525 radiationenergy than dark surfaces, like forests and open waters. Thus, the type of land cover may affect regional
3526 warming, but its relative contribution is disputed (Gaillard et al., 2015; Strandberg and Kjellström, 2019).
3527 Increasing droughts with lower river flow at certain times of the year may influence water management and
3528 shipping in regulated rivers, especially in the southern catchment basins. On the other hand, extreme rain events
3529 may lead to inundations (Kundzewicz et al., 2005).

3530

3531 Climate change will strongly affect coastal structures through sea level rise and intensified coastal erosion. Storm
3532 surges, which run up higher as sea level rises, as well as changed currents and sediment relocations will endanger
3533 levees, groynes and other coastal structures, and have to be handled by coastal management (Le Cozannet et al.,
3534 2017; Labuz, 2015).

3535
3536 We can expect a considerable increase in offshore wind energy production worldwide, in order to counteract
3537 climate warming. Although projections of future winds are highly uncertain, the number of off-shore wind farms
3538 can be expected to increase due to the politically driven shift to renewable energies, and the limited space and low
3539 acceptance for wind mills on land. Offshore wind farms may in turn affect the regional climate by absorbing
3540 atmospheric kinetic energy on the regional scale (Akhtar et al., 2021), but the magnitude of this effect is
3541 unknown (Lundquist et al., 2019).

3542
3543 Shipping is affected by climate change. Perils at sea for ships are all climate sensitive, ranging from storms, waves,
3544 currents, ice conditions, visibility to sea level affecting navigational fairways. Winter navigation will be facilitated
3545 as drastically decreasing winter sea-ice cover is projected, but search and rescue missions in winter may increase
3546 because engine power may in the future be adapted to the lower expected ice cover. Further aspects are a potential
3547 increase in leisure boating, a potentially temperature-dependent functioning of antifouling paints, and different
3548 noise propagation through warmer water. The efficiency of SO_x scrubbing depends on the temperature, salinity
3549 and pH of the seawater. Furthermore, highly acid scrubber water may and eventually ends up contaminating the
3550 Baltic Sea, depending on the type of scrubber in use (Turner et al., 2018). Shipping itself affects climate through
3551 combusting fossil fuels, although the emissions can be expected to be reduced with the expected increase of
3552 renewable energy within the European Union (European Commission, 1998, 2018a, b)C, 2021).

3553
3554 Coastal processes, e.g. erosion and the translocation of sediments through erosion, currents and accretion, are
3555 affected by climate change through sea level rise and changes in storm frequency, severity and tracks (Defeo et
3556 al., 2009).

3557
3558 Climate change affects the amount of nutrients entering the sea in precipitation and by land runoff, which in turn
3559 is affected by precipitation, air temperature and runoff pattern, e.g. Arheimer et al. (2012) and Bartosova et al.
3560 (2019). How fertilization practices, crops grown, and land use will change in response to climate change is yet to
3561 be determined ~~largely unknown~~. Climate-related changes in the Baltic Sea, like warmer temperatures, changed
3562 stratification and altered ecosystems and biogeochemical pathways may change the fate of nutrients in the sea, e.g.
3563 Kuliński et al. (2021).

3564
3565 There is yet little ~~not much~~ evidence of a direct climate influence on the quantity and quality of submarine
3566 groundwater discharge, but considering the driving forces (topography-driven flow, wave set-up, precipitation, sea
3567 level rise and convection), it is an effect is highly plausible that an effect will occur, but its possible magnitude and
3568 relevance is unknown (e.g. Taniguchi et al., 2019).

3569

3570 Fisheries are strongly affected by climate change through its effect on the resources, i.e. the commercially
3571 interesting fish populations in the Baltic Sea, mostly cod, sprat and herring (Möllmann, 2019). Climate affects
3572 salinity and temperature in the Baltic Sea, thereby influencing the productivity of several fish species (MacKenzie
3573 and Schiedek, 2007; Köster et al., 2016), and the resources that fisheries exploit. Growth of planktivorous species
3574 or life stages is also affected by climatic conditions that regulate zooplankton dynamics (Casini et al., 2011; Köster
3575 et al., 2016).

3576
3577 Climatic change is a plausible driver for the migration and occurrence of non-indigenous species, although there
3578 is little direct evidence during current climate change. Shipping has been identified as a major vector for the
3579 introduction of new marine species into the Baltic Sea ecosystem, through ballast water or attachment to hulls or
3580 elimination of physical barriers (e.g. through the construction of canals between water bodies; Ojaveer et al., 2017).
3581 However, a northward migration of terrestrial (Smith et al., 2008) and marine species, including fish, due to
3582 increasing temperature is documented and expected to continue (MacKenzie and Schiedek, 2007; Holopainen et
3583 al., 2016).

3584
3585 Climate change affects contaminants in the Baltic Sea through an array of processes, like partitioning between
3586 environmental phase-pairs such as air-water, air-aerosols, air-soil, air-vegetation, leading to a different distribution
3587 between environmental compartments (Macdonald et al., 2003). Atmospheric transport and air-water exchange
3588 can be influenced by changes in wind fields and wind speeds (Lamon et al., 2009; Kong et al., 2014). Changing
3589 precipitation patterns influence chemical transport via atmospheric deposition (rain dissolution and scavenging of
3590 particles, (Armitage et al., 2011) and runoff, transporting terrestrial organic carbon (Ripszám et al., 2015). As ice-
3591 cover of lakes and the sea decreases, more organic contaminants may volatilize to the atmosphere (Macdonald et
3592 al., 2003; Undeman et al., 2015).

3593
3594 Dumped military ammunition threaten the Baltic Sea in the future, as poisonous substances are expected to leak
3595 due to advanced corrosion of hulls and containers. This process may be affected by climate, as corrosion rates
3596 depend on temperature and oxygen, so that warming and good ventilation of dumping sites can be expected to
3597 enhance corrosion rates (Vanninen et al., 2020). This is an urgent problem since the location of the dumped military
3598 material is only partially known.

3599
3600 There is no evidence of degradation of marine litter or microplastics in the marine environment and~~There is no~~
3601 ~~evident direct impact of climate change on this driver~~marine litter or microplastics (Oberbeckmann and Labrenz,
3602 2020), but there may be a connection via increased temperature and photolysis-dependent rates of degradation
3603 and dissolution of microplastics, and on distribution by currents. Societal decisions about the use of plastics, in
3604 part influenced by efforts to decarbonize, will very likely affect future plastic pollution more than climate change.

5 Comparison with the North Sea region

5.1 The North Sea region

Like the Baltic Sea basin, the North Sea region is both a precious natural environment and a place for settlement and commerce for millions of people, with a rich cultural heritage. The North Sea is one of the world's richest fishing grounds as well as one of the busiest seas with respect to shipping and infrastructure for oil and gas extraction, and of enormous economic value. In recent years the area has also become a major site for wind energy, with many large offshore wind farms.

As climate change is expected to have profound effects on North Sea ecosystems and economic development, an independent, voluntary, international team of scientists from across the region compiled the North Sea Region Climate Change Assessment (NOSCCA; Quante and Colijn, 2016). The NOSCCA approach is similar to BACC in format and intention. The assessment provides a comprehensive overview of all aspects of a changing climate, discussing a wide range of topics including past, current and future climate change, and climate-related changes in marine, terrestrial and freshwater ecosystems. It also explores the impact of climate change on some socio-economic sectors, such as fisheries, agriculture, coastal zone management, coastal protection, urban climate, recreation/tourism, offshore activities/energy, and air pollution.

The North Sea is a semi-enclosed marginal sea of the North Atlantic Ocean, situated on the north-west European shelf. It opens widely into the Atlantic Ocean at its northern boundary, with a smaller connection to the Atlantic Ocean via the Dover Strait and English Channel in the south-west. To the east it connects to the Baltic Sea. The Kattegat, a transition zone between the North and Baltic seas, is located between the Skagerrak and the Danish straits. Comprehensive reviews of North Sea physical oceanography are provided by Otto et al. (1990), Rodhe (1998) and Sündermann and Pohlmann (2011). Physical-chemical-biological interaction processes within the North Sea are reviewed by Rodhe et al. (2006) and Emeis et al. (2015), and a description of the North Sea marine ecosystem was compiled by McGlade (2002).

Among the most striking differences between the North and Baltic seas is the wide, direct opening of the North Sea to the North Eastern Atlantic, allowing free exchange of matter, heat and momentum between the two seas. As a result the North Sea water has a much higher salinity than the Baltic Sea. The North Sea dynamics are greatly influenced by tides, while Baltic Sea tides are much weaker than in the North Sea, where tidal amplitudes vary spatially from a few decimeters to several meters. In addition to the wind-driven circulation, which dominates the mean cyclonic current system, North Sea tidal currents show non-vanishing residual currents (due to nonlinear processes), which cannot be neglected. Tidal currents cause strong mixing. Low pressure systems often travel from the Atlantic with minimum blockage and cause strong storm surges, which are the greatest potential natural hazards for coastal communities in the North Sea region.

Only selected examples from NOSCCA will be presented here. In general, the North Sea region already experiences a changing climate and projections indicate that further, partly accelerating, changes are to be expected

(warming of air and water, changing precipitation intensities and patterns, sea level rise, seawater acidification). Changes in ecosystems (marine, coastal, terrestrial) are observed, and are projected to strengthen, with degree depending on scenario. Observational as well as modelling studies have revealed a large natural variability in the North Sea region (from annual to multidecadal time scales), making it difficult to identify regional climate change signals and impacts for some parameters. Projecting regional climate change and impacts for the North Sea region is currently limited by the small number of regional coupled model runs available and the lack of consistent downscaling approaches, both for marine and terrestrial impacts. The wide spread in results from multi-model ensembles indicates the present uncertainty in the amplitude and spatial pattern of the projected changes in sea level, temperature, salinity and primary production. For moderate climate change, anthropogenic drivers such as changes in land use, agricultural practice, river flow management or pollutant emissions often seem more important for impacts on ecosystems than climate change.

3654 **5.2 A few selected and highly aggregated results from NOSCCA**

3655 *Atmosphere:* Observations reveal that the near-surface atmospheric temperature has increased everywhere in the
3656 North Sea region, especially in spring and in the north. The rise was faster over land than over the sea. Linear
3657 trends in the annual mean land temperature are about +0.39°C per decade for the period 1980–2010. Generally,
3658 more warm extremes and fewer cold ones were observed. A north-eastward shift in storm tracks was observed, in
3659 agreement with projections from climate models forced by increased greenhouse gas concentrations. Overall,
3660 precipitation has increased in the northern North Sea region and decreased in the south, summers have become
3661 warmer and drier and winters have become wetter. Heavy precipitation events have become more extreme. A
3662 marked further mean warming of 1.7–3.2°C is projected for the end of the 21st century (2071–2100, with respect
3663 to 1971–2000) for different scenarios (RCP4.5 and RCP8.5, respectively), with stronger warming in winter than
3664 in summer and particularly strong warming over southern Norway.

3665
3666 *North Sea:* There is strong evidence of surface warming in the North Sea, especially since the 1980s (Fig. 343).
3667 Warming is greatest in the south-east, exceeding 1°C since the end of the 19th century. Absolute mean sea level
3668 in the North Sea rose by about 1.6 mm/year over the past 100–120 years, in agreement with the global rise. The
3669 North Sea is a sink for atmospheric carbon dioxide (CO₂); this uptake declined over the last decade, due to lower
3670 pH and warmer water. Models consistently project the surface water to warm further by the end of the century (by
3671 about 1–3°C; A1B scenario). Exact numbers are not given due to differences in spatial averaging and reference
3672 periods from published studies. Coherent findings from published climate change studies include an overall rise
3673 in sea level, an increase in ocean acidification and a decrease in primary production. Uncertainties are large for
3674 projected changes in extreme sea level and waves as well as for decreases in net primary production, which range
3675 from 1 to 36 %.

3676
3677 *Rivers:* To date, no significant trends in response to climate change are apparent for most individual rivers
3678 discharging into the North Sea. Nevertheless, climate models project increased socio-economically important risks
3679 for the region, due to more intense hydrological extremes in the North Sea region, such as flooding along rivers,
3680 droughts and water scarcity. The exposure and vulnerability of cities in the North Sea region to changes in extreme

hydrometeorological and hydrological conditions are expected to increase, due to greater urban land use and rising urban populations.

Ecosystems: Long-term knowledge from exploitation of the North Sea indicates that climate affects marine biota in complex ways. Climate change influences the distribution of all taxa, but other factors (fishing, biological interactions) are also important. The distribution and abundance of many species have changed. Warm-water species have become more common and species richness has increased. Among coastal ecosystems, estuaries and most mainland marshes will survive sea-level rise, while back-barrier salt marshes with lower suspended sediment concentrations and tidal ranges are probably more vulnerable. Plant and animal communities can suffer habitat loss in dunes and salt marshes through high wave energy, and are affected by changes in temperature and precipitation and by atmospheric deposition of nitrogen. Lakes in the North Sea region have experienced a range of physical, chemical and biological changes due to climatic drivers over past decades. Lake temperatures have increased, ice-cover duration has decreased. For terrestrial ecosystems there is strong empirical evidence of changes in phenology in many plant and animal taxa and northward range expansions of mobile heat-loving animals. Climate change projections and effect studies suggest a northward shift of vegetation zones, with terrestrial net primary production likely to increase in the North Sea region, due to warmer conditions and longer growing seasons.

Socio-economic effects: The assessments of climate change effects on the different socio-economic sectors in the North Sea region find that adaptation measures are essential for all of them, e.g. for coastal protection and in agriculture. For North Sea fisheries, the rapid temperature rise is already being felt in terms of shifts in species distribution and variability in stock recruitment. In agriculture an increased risk of summer drought and associated effects will be a challenge, particularly in the South. In general, extreme weather events are likely to more often severely disrupt crop production. Offshore and onshore activities in the North Sea energy sector (dominated by oil, gas and wind) are highly vulnerable to extreme weather events, in terms of extreme wave heights, storms and storm surges. All coastal countries around the North Sea with areas vulnerable to flooding by storm surges are preparing for the challenges expected due to climate change, but coastal protection strategies differ widely from country to country. Due to ~~the inadequate knowledge uncertainty about concerning~~ the extent and timing of climate-driven impacts, current coastal zone adaptation plans focus on no-regret measures.

5.3 Some differences in climate change effects between the North Sea and Baltic Seas

Many of the climate change signals in the Baltic and North Seas show similar behaviours and trends. But there are also some notable differences between the two regions, which are listed below. They are based on findings reported in the appropriate chapters of the recent assessments BACC II (BACC II Author Team, 2015) and NOSCCA (Quante and Colijn, 2016).

- In recent decades, the surface air temperature in the Baltic Sea and North Sea regions rose in a similar way, on the order of 1 °C in the past century. Projections of the surface air temperature as obtained by EURO-CORDEX downscaling for a moderate scenario (RCP4.5) indicate a stronger winter and spring warming (> 1 °C) at the end of the century (2071-2100) relative to present day (1971-2000) for most

parts of the Baltic Sea region than for the western part of the North Sea region. In the summer and autumn months the projected warming is at the same level.

- The North Sea is vigorously ventilated by the Atlantic (overturning time ~1 to 4 years). Therefore, climate change signals from the Atlantic are rapidly transferred to the North Sea, while climate change in the North Sea can be expected to be damped by the large thermal inertia of the Atlantic Ocean. By contrast, the Baltic Sea is more prone to changes in mean meteorological conditions as its connection to the World Ocean is very narrow.
- Projected changes in seasonal mean precipitation show a distinctive difference between the two sea regions for the summer (JJA) and autumn months (SON). In the Baltic Sea region the mean precipitation for a RCP4.5 scenario is projected to increase for most land areas (5 to 25%), whereas no noticeable change (5 to -5%) is projected along the western and southern shores of the North Sea region.
- SST is currently rising and projected to rise further for both sea areas, but the spatial pattern of the SST increase is different. In the southern North Sea SST rises more than in the northern North Sea, while SST warming trends are higher in the north-eastern part of the Baltic Sea (Bothnian Sea and Gulf of Finland) than in the southern part. These spatial differences are explained by water depth (North Sea) and the ice-albedo feedback (Baltic Sea). In addition, the northern North Sea is affected by Atlantic water inflow at the western side of the Norwegian trench.
- The coastal regions of the North Sea experience increases in both mean sea level (MSL, as measured by satellites) and relative mean sea level (RMSL, as measured by tide gauges). Trends in RMSL vary significantly across the North Sea region due to the influence of vertical land movement (uplift in northern Scotland, Norway and Denmark, and subsidence elsewhere). But the trend of RMSL is still positive everywhere in the North Sea coasts. In contrast, sea levels relative to land along the northern Baltic Sea coast are sinking because land levels continue rising, due to post-glacial rebound since the last ice age. The northern Baltic Sea will experience considerable land rise also in future. As a result, the sea level will probably continue to decrease relative to land in this region. As positive trends in RMSL are more relevant for coastal protection, all countries around the North Sea with coastal areas vulnerable to flooding due to storm surges face similar challenges, while in the Baltic Sea region coastal protection is of greater concern for the countries in the south.
- The frequency of sea ice occurrence in the North Sea has decreased since about 1961, with a similar development in the western Baltic Sea. In contrast, ice still forms in the northern Baltic Sea, where it will remain a prominent feature for many years, covering about 50 to 200 x 10³ km², with high interannual variability, even though for the 21st century a linear trend of 2% decrease per decade is reported.

6 Knowledge gaps and research needs

Knowledge gaps and research needs have been intensively discussed within the grand challenge working groups of Baltic Earth and are summarized by the BEARs (Lehmann et al., 2021; Kuliński et al., 2021; Rutgersson et al., 2021; Weisse et al., 2021; Christensen et al., 2021; Gröger et al., 2021a; Meier et al., 2021a; Viitasalo and Bonsdorff, 2021; Reckermann et al., 2021).

Formatiert: Nicht unterstrichen, Schriftfarbe: Automatisch, Hochgestellt

3756
 3757 In summary, we conclude that the processes that control the variability of salinity in the Baltic Sea and its entire
 3758 water and energy cycles are still not fully understood (Lehmann et al., 2021). The time-dependence of the haline
 3759 stratification and its links to climate change are in special need of further study. Salinity dynamics is important for
 3760 its dominant role in stratification, concerning both mixing conditions and ecosystem composition and functioning.
 3761 The environmental and biological factors favoring certain biogeochemical pathways through complex interactions,
 3762 the pools of dissolved organic matter, and sediment biogeochemical processes are poorly understood (Kuliński et
 3763 al., 2021). Although initial studies on the coastal filter capacity have been made, coastal zone models for the entire
 3764 Baltic Sea and an overall estimate of bioavailable nutrients and carbon loads from land to the open sea do not exist
 3765 (Kuliński et al., 2021). Considering the large internal variability, investigations of changes in extremes are limited
 3766 because high-resolution observational time series are too short and model ensembles too small (Rutgersson et al.,
 3767 2021). Global mean sea level rise, land uplift and wind field changes control sea level of the Baltic. However, the
 3768 future evolution of these drivers, which are needed for projections, is rather uncertain (Weisse et al., 2021).
 3769 Furthermore, databases for coastline changes and erosion and basin-scale models of coastal change under sea-level
 3770 rise do not exist (Weisse et al., 2021).
 3771
 3772 Fully coupled regional ESMs for the Baltic Sea including the various compartments of the Earth system,
 3773 atmosphere, land, ocean, sea ice, waves, terrestrial and marine ecosystems are under development but are not yet
 3774 available for dynamical downscaling (Gröger et al., 2021a). The numerical estimation of water and energy cycles
 3775 suffers from both model deficiencies and natural variability. For climate projections, even if large ensembles of
 3776 high-resolution regional atmosphere models are becoming increasingly available, the coverage of the underlying
 3777 global climate model ensembles is still small implying that detailed conclusions on uncertainty and/or robustness
 3778 in details of future climate change and its impacts cannot easily be drawn. In addition, only one ensemble with 22
 3779 members utilized a coupled atmosphere-ice-ocean regional model (Christensen et al., 2021). For climate
 3780 projections, large ensembles of regional atmosphere models are available but only one ensemble with 22 members
 3781 utilized a coupled atmosphere-ice-ocean model (Christensen et al., 2021). Also in the past, the ocean Baltic Sea
 3782 ecosystem model ensembles have had too few members to address well the spread uncertainty related to the large
 3783 multidecadal variability in the ocean (Meier et al., 2021a). Furthermore, the global sea level rise needs to be
 3784 considered when making salinity projections which is rather uncertain (Meier et al., 2021b).
 3785
 3786 -The large inherent uncertainty in future projections of salinity fundamentally affects the projections of the marine
 3787 ecosystem (Viitasalo and Bonsdorff, 2021). The response of food web interactions to climate change isare largely
 3788 unknown. The uncertainties of scenario simulations with coupled physical-biogeochemical ocean models were
 3789 discussed by Meier et al. (2018a; 2019b; 2021b). They found that in addition to natural variability the largest
 3790 uncertainties are caused by (i) poorly known current bioavailable nutrient inputsloads from land and atmosphere
 3791 and yet to be determineduncertain assumptions about future inputsloads, (ii) uncertainties of models including
 3792 global sea level rise, and (iii) poorly known long-term future greenhouse gas emissions.

Formatiert: Schriftart:

3794 Finally, the regional Earth system is driven by multiple drivers, of which climate change is just one. Multi-driver
3795 studies are just beginning to be made and only a few have yet been published (Reckermann et al., 2021).

3796
3797 In the following, we list a few selected knowledge gaps related to the variables addressed by this study.

3798 6.1 Large-scale atmospheric circulation

3799 The interactions between atmospheric modes of variability of importance for the Baltic Sea region are still not
3800 well known. For instance, while climate models are able to simulate the main features of the NAO, the frequency
3801 of blocking over the Euro-Atlantic sector is still underestimated (IPCC, 2014b). Since observational records are
3802 relatively short, our understanding of the AMO and its possible changes depends largely on models, and these
3803 cannot be reliably evaluated for time scales longer than the AMO period (Knight, 2009). However, while possible
3804 changes in these climate phenomena do contribute to the inherent uncertainty in near-term climate projections,
3805 they are not the main driver of the projected warming over Europe by the end of the century (Cattiaux et al., 2013;
3806 IPCC, 2014b).

3807 6.2 Air temperature

3808 Air temperature and its extremes are to a large extent determined by the large-scale atmospheric circulation
3809 patterns. There is limited knowledge primarily concerning changes in these large-scale-atmospheric circulation
3810 patterns in a changing climate, as mirrored by climate model discrepancies.

3811
3812 Furthermore, the strong dependence of even local temperature changes on the evolution of greenhouse gas
3813 emissions and the feedbacks that determine the global climate sensitivity deserve attention. Another uncertainty,
3814 of unknown importance, is the extent to which larger-than-expected decreases in the AMOC could potentially
3815 counteract the effects of global warming in northern Europe. As shown by the IPCC (2014b, their Fig. 12.9), there
3816 in fact was one model with a cooling of northern Europe in the CMIP5 ensemble. The recent suggestion that the
3817 AMOC may be more sensitive to anthropogenic climate change than current climate models indicate may also be
3818 relevant for the Baltic Sea region (Boers, 2021).

3819
3820 Nevertheless, the heat cycle of the Baltic Sea region is probably better understood than the water cycle.

3821 6.3 Solar radiation and cloudiness

3822 Multidecadal variations in surface solar radiation (SSR) are generally not well captured by current climate model
3823 simulations (Allen et al., 2013; Storelvmo et al., 2018). The extent to which the observed variations in SSR are
3824 caused by natural variation in cloudiness induced by atmospheric dynamic variability (Stanhill et al., 2014; Parding
3825 et al., 2014), or by anthropogenic aerosol emissions (Wild, 2012; Ruckstuhl et al., 2008; Philipona et al., 2009;
3826 Storelvmo et al., 2018), or perhaps additional causes, is not understood. Future cloudiness trends in global and
3827 regional models differ in their sign (Bartók et al., 2017). Most RCM scenario simulations lack time varying aerosol
3828 forcing (Boé et al., 2020b).

Formatiert: Rechts: 0,03 cm

3829 **6.4 Precipitation**

3830 Even if climate scenarios are becoming more frequent and there is now a growing ensemble of relatively high-
3831 resolution regional climate scenarios for Europe, they still represent only a subset of the global climate model
3832 projections assessed by the IPCC. This means that the uncertainties of future climate change in the Baltic Sea
3833 region are not fully captured at the horizontal resolution needed for detailed studies of climate change effects in
3834 the region (Christensen et al., 2021). Very high-resolution so called “convective-permitting” climate models
3835 operating at grid spacing of 1-3 km are lacking for the Baltic Sea region. In other regions, such models have better
3836 agreed with observations of precipitation extremes and sometimes also given a larger climate change signal than
3837 the more traditional “high-resolution” models operating at c. 10 km grid spacing (Christensen et al., 2021). Land
3838 use change and cover, including changes in forests, can induce both local and downwind precipitation change
3839 (Meier et al., 2021c), and need to be included in projections.

3840 **6.5 Wind**

3841 Historical wind measurements suffer from inhomogeneity and records too short for detecting changes, considering
3842 the large internal variability in the Baltic Sea region. Projected changes are not robust among the few available
3843 downscaled ESMs.

3844 **6.6 Air pollution**

3845 The spatially and time resolved air quality status of a region is often assessed by means of model systems, typically
3846 with emission, meteorological and chemistry transport submodels. These model systems, used for the calculation
3847 of atmospheric concentrations and deposition of pollutants, need further developments and validation.
3848 Uncertainties are often connected to the emission segment of the modelling chain. Improvements of the
3849 implemented time profiles for the different emission sectors are especially necessary (Matthias et al., 2018). For
3850 projections of air quality with climate change models, more work is needed to establish a set of emission scenarios
3851 for air pollutants consistent with regional socio-economic pathways, like those developed by Zandersen et al.
3852 (2019). The shipping sector is currently a considerable source of air pollution in the Baltic Sea region. More
3853 research and development is needed on new fuel types and emission factors for air pollutants, relevant for
3854 politically and technologically driven abatement measures. To better address exposure and health impacts of
3855 shipping emissions more studies are required like those of Ramacher et al. (2019) and Barregard et al. (2019),
3856 especially at the harbour and city scale. Better knowledge and reduced uncertainties will improve quantification
3857 of air pollution as part of the environmental imprint of shipping in the Baltic Sea region, as developed by
3858 Moldanová et al. (2021).

3859 **6.7 River discharge**

3860 Precipitation from regional atmosphere models is biased and the bias correction methods applied for hydrological
3861 modeling affect the sensitivity of hydrological models to climate change (Donnelly et al., 2014). Natural variability
3862 and model uncertainties may explain the large spread in current river discharge projections (Roudier et al., 2016;
3863 Donnelly et al., 2017). The values of the parameters of a hydrological model are normally found through

calibration against historical data and are always associated with ~~inaccuracies~~uncertainty. These ~~is~~
~~inaccuracies~~uncertainty will translate into uncertainty in the projected changes.

6.8 Nutrient inputs from land

The time scales for exchange of the nutrient pools in soils are not well known (McCrackin et al., 2018). Long-term observations do not exist. Future projections of river discharge and nutrient inputs in the Baltic Sea drainage basin agree on key aspects (e.g. increased annual discharge), but also highlight the uncertainty of the projections. To improve assessments, studies should be designed to allow explicit semi-quantitative comparisons of the effects of the incorporated change factors, e.g. climate, land management, policy. In the case of nitrogen inputs, the effect of changes in anthropogenic atmospheric deposition should also be included in future projections.

6.9 Terrestrial biosphere

Terrestrial ecosystems in the Baltic Sea region are governed by human activities, both changes in climate due to anthropogenic climate forcing and anthropogenic changes in land use and land cover. In return, terrestrial ecosystems affect climate by altering the composition and the energy and water cycles of the atmosphere. Biophysical interactions between the land surface and the atmosphere have been incorporated into regional ESMs, in order to assess the impacts of changes in land use and land cover on regional climate and terrestrial ecosystems. Still, biogeochemical processes related to the carbon cycle are lacking, as are explicit forest management actions (Lindeskog et al., 2021), while explicit descriptions of some disturbances (e.g. wildfires, major storms, insect attacks) are under development in ESMs. Only when all these interactions are incorporated can the effects of national or international (e.g. in the European Union) climate policies on regional climate and terrestrial ecosystems be fully assessed for compliance with the goals of the Paris Agreement.

6.10 Snow

A general decrease in snow-cover duration in the Baltic Sea region is well documented, especially for the southern part. Changes in snow depth due to climate warming are much more unclear. Some evidence of increasing snow depth in recent decades have been reported from the northern part of the study region and from mountainous areas. However, these increases are not projected to continue, according to climate model projections. Whether or not there is a discrepancy between the observed and projected trends is not known.

-Changes in sea-effect snowfall events during present climate are unknown.

6.11 Glaciers

-It is presently not known how glacier-fed lakes react to competing environmental drivers, such as the general Arctic warming, and the simultaneous warming-triggered lake cooling caused by increased inflow of cold glacier meltwater, potentially carrying high sediment, nutrient, and organic matter loads. Understanding changing lake thermal regimes and vertical mixing dynamics as well as timing and duration of seasonal ice cover is important because ecological, biological, chemical processes, including carbon-cycling, will be affected (Lundin et al., 2015; Smol et al., 2005; Jansen et al., 2019). Since Scandinavian glaciers are predicted to decline 80% in volume by

2100 under RCP8.5, Scandinavian glacier-fed lakes could be used as natural observatories, where changes in processes, timescales, and effects in response to competing drivers can be studied before they occur at other glacial lake sites, where glaciers melt more slowly (Kirchner et al., 2021).

6.12 Permafrost

Thawing permafrost peatlands may potentially release large amounts of organic matter, nutrients and greenhouse gases to aquatic systems locally, but the timing and magnitude of such releases remain highly uncertain.

6.13 Sea ice

While the extent of the sea ice cover is well observed, observations of ice thickness are scarce. Ice thickness is regularly monitored only at a few coastal sites with fast ice. Long records of the various ice classes, such as ridged ice, do not exist. Sea ice models do not represent sea ice classes correctly. Since the last assessment by the BACC II Author Team (2015) only two new scenario simulation studies on sea ice were published (Luomaranta et al., 2014; Höglund et al., 2017).

6.14 Lake ice

Research is required to better understand the reasons for regional and temporal differences in the patterns of change in lake ice phenology and its relationship to large-scale climatic forcing. There is a need to better understand how loss of lake ice cover modifies gas exchange between lake and atmosphere, mixing of the water column, biogeochemical cycling, and ecosystem structure and function. The socioeconomic and cultural importance of winter ice also deserves further research.

6.15 Water temperature

The causes of the pronounced natural variability of Baltic Sea temperature and its connection to large-scale patterns of climate variability are not well known. The occurrence of marine heat waves is projected to increase. However, only a few studies of their impacts on the marine ecosystem exist. Furthermore, sea surface temperature trends also depend on coastal upwelling, which affects large areas of the Baltic Sea surface (Lehmann et al., 2012; Dutheil et al., 2021). Projected changes in upwelling are, however, very uncertain (Meier et al., 2021a).

6.16 Salinity and saltwater inflows

Salinity change depends on wind, river discharge, net precipitation over the sea and global sea level rise. Due to considerable uncertainty in all drivers and the different signs in the response of salinity to these drivers, the relative uncertainty in salinity projections is large, and larger ensembles of scenario simulations are needed (Meier et al., 2021b). This knowledge gap is also associated with compounded by the uncertainty of whether saltwater inflows from the North Sea will change (Schimanke et al., 2014). As salinity is a very important variable for the circulation in the Baltic Sea and for the marine ecosystem, projections for the Baltic Sea salinity are a priority.

6.17 Stratification and overturning circulation

Stratification depends on mixing as well as on gradients in water temperature and salinity, making changes in stratification highly uncertain. Mixing processes such as thermal and haline convection, entrainment, double diffusive convection or boundary mixing are not fully understood (Holtermann et al., 2012; 2017; Umlauf et al., 2018). Initial results on the sensitivity of the vertical overturning circulation rely on model studies only (Placke et al., 2021). Hence, more measurements on the fine-structure of horizontal and vertical turbulence are needed (Reissmann et al., 2009).

6.18 Sea level

The regional variability of processes which drive sea-level changes, along with their uncertainties and relative importance over different timescales, display long-term developments that still require an explanation and are a challenge to planning by coastal communities (Hamlington et al., 2020). For instance, the annual cycle in Baltic Sea mean sea level (winter maxima minus spring minima) shows a basin-wide widening in the period 1800-2000 (Hünicke and Zorita, 2008). The precise mechanisms responsible for this effect are not yet completely understood, although it seems strongly controlled by atmospheric forcing (Barbosa and Donner, 2016). Furthermore, at the longer time-scales relevant for anthropogenic climate change, Baltic Sea and North Atlantic sea levels are strongly affected by whether warming is allowed to proceed to the point of destabilising the very uncertain future melting of the Antarctic ice sheets. Current estimates are mostly based on heuristic expert knowledge, as models are still under development. This is probably the largest knowledge gap affecting projections of future Baltic sea-level rise (Bamber et al., 2019). Finally, long-term relative sea level trends are strongly affected by the vertical land movement due to glacial isostatic adjustment. This can be as large as, or even larger, than global sea-level rise. Currently, it is estimated from relatively short GPS measurements and from geo-elastic models. Both are inaccurate and uncertain, as point GPS measurements are strongly affected by other geological and anthropogenic effects on vertical land velocities and results from model geo-elastic models are often revised (Weisse et al., 2021). In addition, the glacial isostatic adjustment may affect the flow intensity of river runoff into the northern Baltic Sea (coastal regions rising relative to inland regions), the effects of which, e.g. on salinity and water levels, have not been explored.

6.19 Waves

The lack of long-term instrumental wave measurements and gaps in the data due to the ice season complicate the analysis of extreme values. Although wave hindcasts provide a good alternative, the accuracy naturally does not match that of measured data. Furthermore, Björkqvist et al. (2020) showed that the calculation of return periods of extreme events may depend on the sampling frequency. Adding sampling variability typical for in-situ measurements to simulated hindcast data, will result in consistently shorter estimates of return periods for high significant wave heights than using the original hindcast data.

6.20 Sedimentation and coastal erosion

We lack a comprehensive understanding of alongshore sediment transport and its associated spatial and temporal variability along the Baltic Sea coast. In general, an eastward transport dominates along most of the southern Baltic Sea coast due to the prevailing westerly winds. However, the intensity of secondary transport induced by easterly and northerly winds is much less understood. Its combination with storm surges will expose sand dunes and cliffs to the greatest erosional impact, further complicating understanding (Musielak et al., 2017). Due to the orientation of the coastline, transport along some parts of the Baltic Sea coastline is very sensitive to the angle of incidence of the waves. For example, the incidence angle of westerly wind-waves at the western part of the Wolin Island in Poland (Dudzińska-Nowak, 2017) and the coast of Lithuania and Latvia (Soomere et al., 2017) is very small and even a slight change in the wind direction (e.g. by 10 degrees) could lead to a reversal of the direction of alongshore transport. Coastline changes at these sections vary greatly, and will hence be extremely sensitive to future changes in wind-wave climate (Viška and Soomere, 2013). Another knowledge gap in understanding coastal erosion in response to future climate change concerns the impact of water levels and the submergence of the beach. Water level plays a key role in dune toe erosion and also limits aeolian sand transport on the beach. The relationship between the intensity of the forcing (wave energy, run-up) and the morphological response (erosion at the beach and dunes) during storms is not straightforward (Dudzińska-Nowak, 2017; Zhang et al., 2017). At some sites (e.g. Miedzyzdroje in Poland), dune erosion is well correlated with maximum storm surge level and storm frequency, but at others (e.g. Swinoujście), the beach morphology is more important in determining the effect of erosion than the storm surge level.

6.21 Oxygen and nutrients

There are significant knowledge gaps related to the identification and quantification of oxygen sinks and sources in the Baltic Sea. In particular, more understanding is required on the dynamics of seawater inflows from the North Sea, the role of mixing processes in the ventilation of the deep water, rates of oxygen consumption in water column and sediments and how they depend on climate change (Kulinski et al., 2021). Knowledge gaps also exist concerning the transport and transformations of DOM (including terrestrial DOM) and better quantification of the processes occurring in the microbial loop is needed to understand the nutrient (but also C and O) dynamics in the Baltic Sea.

The direct effects of climate change are likely to be detectable first in the coastal zone, e.g. indicated by increasing seasonal hypoxia due to warming. However, long-term records from the coastal zone are rare. More important could be the intensification of the proposed hypoxia-related “vicious circle” in the Baltic proper, due to the warming of both surface and deep-water layers in the Baltic proper (Savchuk, 2018; Meier et al., 2018b). The consequent expansion of cyanobacteria blooms and increased nitrogen fixation in the Baltic proper and neighboring basins could further counteract nitrogen load reductions and maintain hypoxia, with all its detrimental effects. However, there are still no biogeochemical and ecosystem models capable of producing reliable long-term scenario simulations of these processes, with sufficient confidence and precision (see Meier et al., 2018a; Meier et al., 2019b).

6.22 Marine CO₂ system

Due to the high spatial and temporal variability of air-sea carbon fluxes, it is not known whether the Baltic Sea as a whole is a net sink or a net source of CO₂. The source of the alkalinity increase observed in the Baltic Sea is still unclear. Plausible hypotheses indicate increased weathering in the catchment and processes related to anoxic remineralization of organic matter. There is high uncertainty in quantifying sediment/water fluxes of C, N and P, which are important bottlenecks for understanding the dynamics of the marine CO₂ system and the C, N, P and O₂ cycling generally, especially in the deep water layers. The lack of system understanding is particularly evident in the ~~Bothnian Sea and Bothnian Bay~~~~Gulf of Bothnia~~. Fransner et al. (2018) suggested that non-Redfieldian stoichiometry in phytoplankton production could explain pCO₂ fields in these ~~sub~~-basins, but confirmation by observations is still lacking.

6.23 Marine biosphere

6.23.1 Lower trophic levels

The summer cyanobacteria bloom in the Baltic proper, and increasingly in recent years in the Bothnian Sea, is considered one of the main problems of Baltic Sea eutrophication, and the nitrogen fixation it carries out is an important process in Baltic ecosystem models (Munkes et al., 2021). It has long been considered limited by the availability of phosphorus (Larsson et al., 1985; Granéli et al., 1990). It is therefore remarkable that it is not possible to predict inter-annual variations in cyanobacteria blooms observed by satellites from water chemistry (Kahru et al., 2020; ~~Hieronimus et al., 2021~~).

There are significant knowledge gaps related to the quantification of nitrogen fixation and the fate of the fixed nitrogen in the Baltic Sea pelagic zone. Direct nitrogen fixation measurements were until recently dogged by method problems, and even if these are now hopefully largely resolved (Klawonn et al., 2015), the enormous patchiness of cyanobacteria blooms remains a huge problem. The alternative approach of directly measuring the increase in total combined nitrogen during the bloom (Larsson et al., 2001) requires very high precision, also suffers from patchiness problems (Rolff et al., 2007), and has not been much used. Finally, the amount of nitrogen fixed can be estimated by modelling, based on uptake of CO₂ or phosphorus and assuming a Redfield N:P or C:N ratio. This theoretically highly attractive approach (Eggert and Schneider, 2015) is hampered by the possibility of non-Redfieldian ratios, and has made some biologists skeptical by predicting high nitrogen fixation in spring, when there are not sufficient known nitrogen-fixing autotrophs in the water to carry out this nitrogen fixation.

While total nitrogen in the water column clearly increases during the summer cyanobacterial bloom, just a couple of months later this increase seems largely to have disappeared, even though sediment traps find little evidence that ~~nitrogen~~~~it~~ has settled out of the upper mixed layer. ~~Are~~~~S~~ediment trap measurements ~~might be~~ gross underestimates, or ~~are~~ there ~~are~~ unidentified sites of denitrification or other overlooked nitrogen sinks in the water column?[?] Nitrogen fixation is a central process in Baltic ecosystem models, and better observationally based estimates of processes in the nitrogen cycle are required for assessing their credibility (Munkes et al., 2021).

6.23.2 Marine mammals

There is a great need for further research on the effects of climate change on marine mammals, especially on the critically endangered Baltic proper harbour porpoise as well as the Baltic ringed seal, given the multiple threats and cumulative impacts on these populations. Seal and porpoise foraging distribution and the relation of seals to haul-out sites is not well known. The requirement of sea ice for successful breeding of ringed seals has not been sufficiently assessed. Land-breeding of grey seals is not monitored regularly in most Baltic statescountries. The effects of interspecific competition on distributions are not known. Range contraction can be conceptualized as three stages (Bates et al., 2014): performance decline, population decrease and local extinction, all of which should be studied. For example, studies on performance decline, such as physiological conditions that reduce reproductive potential (Helle, 1980; Jüssi et al., 2008; Kauhala et al., 2017; Kauhala et al., 2019) are important (Bates et al., 2014). Breeding success of Baltic Sea ringed seals in normal winters is poorly known, as the lairs in pack ice snowdrifts are rarely found. Likewise, observations of the effects of poor ice-conditions on breeding success of ringed seals in mild winters are very limited, but the lack of protection from breeding lairs against harsh weather and predators is assumed to be highly negative.

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Schriftart:

6.23.3 Waterbirds

The complex interaction between many primary parameters affected by climate change makes it hard to identify which environmental changes are actually causing changes in waterbird populations. It is currently not known in detail how shifts in distribution and timing of migration match the availability and quality of food, and thus the importance of potential temporal mismatches between food availability and requirements is unknown. In addition, changes in waterbird distribution are likely to alter inter- and intraspecific competition. Resolving these issues requires investigation of effects at other levels of the food web (e.g. loss of bivalves from areas of reduced salinity, species and size-class composition of fish communities) and their consequences for waterbirds. So far, knowledge of climate change effects on waterbirds in the Baltic Sea are mostly restricted to ducks (including diving and dabbling ducks), with much less known for other quantitatively important components of the waterbird community, i.e. divers, grebes, waders, gulls and auks. Interactions between fish and piscivorous waterbirds in particular need more attention. Responses to climate change are likely to vary between waterbird species and groups. There is still little information on which species are most affected (negatively or positively) by changes in climatic conditions and the uncertainty is therefore large on how species in future waterbird assemblages will interact and the consequences for the functioning of the Baltic Sea. To gain a better understanding on how single species (or groups with similar ecology, often closely-related) will respond to climate change is critical for projecting effects of climate change on waterbirds around the Baltic Sea.

6.23.4 Marine food webs

Some changes observed in marine food webs have been partly attributed to warming, brightening and sea-ice decline on long time-scales. Other drivers, such as eutrophication or fisheries, may however predominate and many records are too short to allow attributing the observed changes to climate change. Although effects of warming, ocean acidification and dissolved organic matter on some ecosystem functions have been identified in mesocosm experiments, changing food-web interactions are still impossible to project. It is, however, important to include

4073 the marine biosphere in management-strategies for tackling the complex interactive aspects of climate change-
4074 related effects on the marine ecosystem and human adaptations to them (Andersson et al., 2015; Stenseth et al.,
4075 2020).

4076 **6.24 Summary**

4077 The following gaps in knowledge are rated as the most serious. Overall, changes in the heat cycle of the Baltic Sea
4078 region are better understood than changes in the water, momentum or carbon cycles. Effects of climate change
4079 induced warming on the latter three cycles are less clear than the effect on the terrestrial and marine heat cycles
4080 including the cryosphere (Fig. 35). The uncertainty in salinity projections substantially limits our understanding
4081 of the marine ecosystem response to anthropogenic climate change. Furthermore, detected trends in observational
4082 records are often caused by internal variability, e.g. in storms, extreme sea level, rather than anthropogenic climate
4083 change because many records are too short. The response of the biosphere to climate change is highly uncertain
4084 because of unknown food-web interactions.

Formatiert: Überschrift 3

Formatiert: Schriftart:

4085 **7 Key messages**

4086 The following lists selected key messages from this assessment that either confirm the conclusions of previous
4087 assessments or are novel (marked with NEW). The estimated level of confidence based upon agreement and
4088 evidence (see Section 2.3) of each key message refers to whether a systematic change in the considered variable
4089 was detected and attributed to climate change. Climate change is here defined as the change in climate due to
4090 human impact only (BACC II Author Team, 2015; see Section 2.1). Key messages referring to observed or
4091 simulated changes in the Baltic Sea region caused by other drivers than climate change (e.g. afforestation,
4092 eutrophication, fisheries, etc.) are not classified by a confidence level. A summary of all key messages related to
4093 climate change is presented in Table 1540 and Figure 354.

4094 **7.1 Past climate changes**

- 4095 • Large-scale circulation: The AMO has undergone frequency changes, but its influence on climate
4096 variability in the Baltic Sea region remained similar, independent of the dominant frequency [NEW].
- 4097 • Air temperature: During the Holocene, The Baltic Sea region experienced periods as warm as the 20th
4098 century, such as the Mid-Holocene Optimum and the Medieval Warm Period. The implied rate of
4099 change was, however, much slower than the present. The past warming signal was regionally markedly
4100 heterogeneous, mostly along a west-east gradient [NEW].
- 4101 • Oxygen: The previous warm periods were accompanied by oxygen deficiency in the deeper waters of the
4102 Baltic Sea, which cannot be attributed to eutrophication and was likely a result of climate forcing [NEW].

4103 **7.2 Present climate changes**

- 4104 • Large-scale atmospheric circulation: Systematic changes in large-scale atmospheric circulation related to
4105 climate change could not be detected [low confidence]. The AMO is an important driver of climate

variability in the Baltic Sea region, affecting *inter alia* the correlation of regional climate variables with the NAO.

- Air temperature: Linear trends of the annual mean temperature anomalies during 1876–2018 were $+0.10\text{ }^{\circ}\text{C decade}^{-1}$ north of 60°N and $+0.09\text{ }^{\circ}\text{C decade}^{-1}$ south of 60°N in the Baltic Sea region [high confidence]. This is larger than the global mean temperature trend and slightly larger than estimated in the earlier BACC reports [NEW]. The warm spell duration index has increased during 1950–2018 [medium confidence]. Statistically significant decreases in winter cold spell duration index across the period 1979–2013 have been widespread in Norway and Sweden, but less prevalent in eastern Finland, while changes in summer cold spells have been small in general [medium confidence].
- Solar radiation and cloudiness: Various satellite data products suggest a small but robust decline in cloudiness over the Baltic Sea region since the 1980s [low confidence, NEW]. However, whether this signal is an indicator of a changing climate or due to internal variability is unknown.
- Precipitation: Since 1950, annual mean precipitation has generally increased in the northern part of the Baltic Sea region. There is some evidence of a long-term trend during 1950–2018 [low confidence]. However, long-term records suffer from inhomogeneity due to the increasing number of rain gauges. Frequency and intensity of heavy precipitation events have increased [medium confidence]. Drought frequency has increased across southern Europe and most of central Europe since 1950, but decreased in many parts of northern Europe [low confidence].
- Wind: Owing to the large internal variability, it is unclear whether there is an overall trend in mean wind speed. There has been an increase in the number of deep cyclones over central and northern Europe since the late 1950s, but no evidence for a long-term trend [low confidence].
- Air pollution: The influence of climate change on air pollution is small and undetectable, given the dominance of other human activities [low confidence]. Land-based emissions are declining due to emission control measures, but some emissions from the shipping sector may be increasing.
- River discharge: For the period 1900–2008, no trend in total river discharge was found, but there was a pronounced 30-year variability. Data for some rivers in the northern Baltic Sea catchment indicate a long-term positive trend during 1921–2004, but the confidence in these reconstructions is low. Since the 1970s, the total river winter discharge is increasing, perhaps due to warming or river regulations [low confidence]. Due to earlier snow-melt, driven by temperature increases in the region and a decreasing frequency of arctic air mass advection, high flow events in the Baltic Sea region shifted from late March to February occurred about a month earlier. In Sweden, trends in the magnitude of high flow events over the past 100 years are not statistically significant [low confidence, NEW].
- Riverine nutrient loads: The effect of changing climate on riverine nutrient loads is small and not detectable [low confidence].
- Terrestrial biosphere: Combining all vegetation types in the entire Baltic Sea region, satellite observations suggest an advancement of the growing season by 0.30 day/year over the period 2000–2016. The most important driver of the advancement of the growing season is spring mean temperature, with an advancement rate of $2.47\text{ day/}^{\circ}\text{C}$ of spring warming [medium confidence, NEW]. Observations and

Formatiert: Schriftart: Nicht Kursiv, Nicht unterstrichen,
Schriftfarbe: Automatisch

model results suggest cooling trends in daily minimum and warming trends in daily maximum temperatures in response to deforestation, and the opposite tendencies for afforestation. [NEW]

- Snow: The decrease in snow cover has accelerated in recent decades, except in the mountain areas and the north-eastern part of the Baltic Sea region [high confidence, NEW]. On average, the number of days with snow cover has declined by 3–5 days per decade, [high confidence]. Mean and maximum snow depth has also decreased, most clearly in the southern and central part of the region [high confidence]. Whether sea-effect snowfall events have changed is unknown [low confidence].
- Glaciers: Inventories of all Scandinavian glaciers, available only since 2006, show that they have lost 20 Gt of ice (~8% of their total mass) during 2006–2015. Atmospheric warming is very likely the primary driver of glacier mass loss [high confidence]. [NEW]
- Permafrost: Recent warming has caused losses of over 20% of the original 6200 km² of permafrost in the Baltic Sea catchment area during 1997–2018 [medium confidence, NEW].
- Sea ice: Long-term decreases in sea ice in the Baltic Sea have exceeded the large natural climatological variability and can only be attributed to global climate change [high confidence]. In addition, unprecedented mild ice seasons have occurred in the last ten years, and 100-year trends in sea ice cover showed an accelerated decline in 1921–2020 compared to 1910–2011 [high confidence, NEW].
- Lake ice: Warming in the Baltic Sea catchment during recent decades has resulted in earlier ice break-up, later freeze-up, and hence shorter ice cover duration on the lakes in the region. [high confidence, NEW]
- Water temperature: Monitoring data, satellite data and model-based historical reconstructions indicate an increase in annual mean sea surface temperature averaged over the Baltic Sea of 0.4–0.6 °C decade⁻¹ or ~1 – 2 °C since the 1980s [high confidence]. During 1856–2005, reconstructed SSTs increased by 0.03 and 0.06 °C decade⁻¹ in northeastern and southwestern areas, respectively. Hence, recent warming trends have accelerated tenfold [NEW]. Long-term measurements at Tvärminne, on the north coast of the Gulf of Finland, indicate that marine heat waves have increased since 1926 [low confidence].
- Salinity and saltwater inflows: The record of major Baltic inflows (MBIs) has been revised and the earlier reported decreasing trend is now seen as artifactual. On centennial time-scales, there are no statistically significant trends in salinity averaged over the Baltic Sea (1920–2008) or in MBIs (1887–2017), but pronounced multidecadal variability, with a period of about 30 years. Model results suggest that a decade of decreasing salinity, like the 1983–1992 stagnation, happens about once a century due to natural variability. Due to increased river runoff in the northern catchment, the North-South gradient in sea surface salinity likely increased in 1900–2008 [low confidence, NEW].
- Stratification and overturning circulation: No long-term trend in stratification was detected, but during 1982–2016 stratification increased in most of the Baltic Sea, with the seasonal thermocline and the perennial halocline strengthening by 0.33–0.39 and 0.70–0.88 kg m⁻³, respectively [low confidence, NEW].
- Sea level: Since 1886 the mean sea level in the Baltic Sea relative to the geoid has increased by about 1–2 mm per year, similar to the global mean rate [high confidence]. However, in the northern Baltic Sea rapid land uplift causes a relative sea level decrease [high confidence]. Although an acceleration of the

mean sea level rise at individual stations could not yet be detected, the all-station-average-record showed an almost statistically significant acceleration [~~medium~~low confidence, NEW]. Basin-wide, no statistically significant, long-term changes in extreme sea levels relative to the mean sea level of the Baltic Sea could be documented [low confidence, NEW].

- Waves: Wave hindcasts and observations are too short for studies of climate-relevant trends [low confidence].
- Sedimentation and coastal erosion: Dominance of mobile sediments makes the southern and eastern coasts more vulnerable to wind-wave induced transport than other Baltic Sea coasts [high confidence]. Prevailing westerly winds lead to mainly west-east sediment transport and an alternation of glacial till cliffs (sources), sandy beaches and spits (sinks). No statistically significant, long-term changes were found [low confidence, NEW].
- Oxygen and nutrients: Reconstructions of oxygen conditions in the Baltic Sea for the period 1898-2012 suggest a tenfold increase of the hypoxic area, with current values of up to 70,000 km². This increase was attributed mainly to increased nutrient loads, with a minor contribution from climate warming [low confidence, NEW]. Furthermore, recently estimated oxygen consumption rates in the Baltic Sea are higher than observed before, reducing the duration of improved oxygen conditions after natural ventilation events by oxygen-enriched saltwater inflows.
- Marine CO₂ system – air-sea exchange: In the period 1980-2005, sub-basins affected by high riverine runoff and related high loads of terrestrial organic matter (e.g. Gulf of Bothnia) were found to be on average a source of CO₂ to the atmosphere. This outgassing was more than compensated by the high CO₂ uptake by the open waters of the Baltic proper [medium confidence, NEW].
- Marine CO₂ system – alkalinity: During 1900-2015, a long-term trend in alkalinity was observed, with largest increases in the Gulf of Bothnia, where it almost entirely cancelled the pH decrease expected from rising atmospheric pCO₂. The smaller alkalinity increase in the southern Baltic Sea compensated ocean acidification by about 50%. Due to the high seasonal variability in pH, large interannual variability in productivity and the identified alkalinity trend, no acidification was measurable in the central and northern Baltic Sea [medium confidence, NEW].
- Microbial communities: Long-term time series from 1994 to 2006 show that increased riverine dissolved organic matter suppresses phytoplankton biomass production and shifts the carbon flow towards heterotrophic microbes [low confidence, NEW].
- Phytoplankton and cyanobacteria: The growing season for phytoplankton and cyanobacteria has lengthened significantly in the past few decades [medium confidence] and the ratio between diatom and dinoflagellate biomasses declined during the past century, probably due to warmer winters [low confidence, NEW]. The annual chlorophyll maximum, in the 1980s associated with the spring diatom bloom, has shifted to coincide with the summer cyanobacteria bloom [low confidence, NEW]. Although inter-annually oscillating, surface cyanobacteria accumulations became a recurrent summer feature of the southern Bothnian Sea in the 2010s [medium confidence, NEW].

- 4220 • Macroalgae: Long-term changes in Baltic Sea macroalgae and charophytes have been attributed to
4221 changes in salinity, wind exposure, nutrient availability and water transparency as well as biotic
4222 interactions [low confidence, NEW]. However, the role of climate change is unclear.
- 4223 • Zoobenthos: Increasing near-bottom temperature may partially explain the spreading of non-indigenous
4224 species, such as polychaetes of the genus *Marenzelleria*. The effects on zoobenthos are primarily
4225 synergistic, through e.g. eutrophication and hypoxia [low confidence, NEW].
- 4226 • Fish: Changes in temperature, salinity and species interactions can affect the stocks of cod, sprat and
4227 herring. However, the dominant driver is the fishery. For coastal fish, the distribution of pikeperch
4228 expanded northwards along the coasts of the Bothnian Sea, apparently due to the warming waters. For
4229 many coastal fish species eutrophication is, however, equally or more important than climate change [low
4230 confidence, NEW].
- 4231 • Marine mammals: Populations of ice-breeding seals, especially southern populations of the ringed seal,
4232 have likely suffered from the sea ice decline [medium confidence]. However, this is based on occasional
4233 ringed seal moult counts that indicat~~ing~~ing no population growth, while monitoring data on reproductive
4234 success are missing.
- 4235 • Waterbirds: Many waterbird species have shifted their wintering range northwards [high confidence].
4236 They now migrate earlier in spring [medium confidence]. Effects of warming sea temperature are
4237 inconsistent, because both positive and negative effects on foraging conditions and food quality have
4238 been found [low confidence]. Most migrating Baltic Sea waterbirds are also affected by climate change
4239 outside the Baltic Sea [medium confidence].
- 4240 • Marine food webs: Significant alterations in food web structure and functioning such as the shift from
4241 early diatom to later dinoflagellate dominated blooms have been observed. However, the causes of these
4242 changes are unknown [low confidence].

4243 7.3 Future climate changes

- 4244 • Large-scale circulation: Projections suggest a more zonal flow over n~~N~~orthern Europe and a northward
4245 shift in the mean summer position of the westerlies at the end of the century [low confidence].
- 4246 • Air temperature: Coupled atmosphere-ocean regional climate models project an increase in annual mean
4247 air temperature by between 1.5 and 4.3°C over the Baltic Sea catchment area at the end of the century.
4248 The range indicates ensemble mean values for RCP2.6 and RCP8.5 scenarios. On average, air over
4249 surrounding land will warm about 0.1 to 0.4°C more than the air over the Baltic Sea [high confidence,
4250 NEW]. A bias-adjusted median estimate of increase in warm spell duration index in Scandinavia for the
4251 period 2071-2100, compared to 1981-2010, was about 15 days under RCP8.5, with an uncertainty range
4252 of about 5-20 days [medium confidence]. The cold spell duration index in n~~N~~orthern Europe is projected
4253 to decrease in the future, with a likely range of from -5 to -8 days per year by 2071-2100, compared to
4254 1971-2000 [medium confidence].
- 4255 • Solar radiation and cloudiness: Projections for solar radiation and cloudiness differ systematically in sign
4256 between global and regional climate models, indicating high uncertainty [low confidence, NEW].

- Precipitation: Annual mean precipitation is projected to increase over the entire Baltic Sea catchment at the end of the century [medium confidence]. The signal is robust for winter among the various regional climate models but is highly uncertain for summer in the south. The intensity and frequency of heavy rainfall events are projected to increase. These increases are even larger for convection-resolving models [high confidence, NEW]. Projections show that the number of dry days in the southern and central parts of the Baltic Sea basin increases mainly in summer [low confidence].
- Wind: Changes in wind over the Baltic Sea region are highly uncertain [low confidence]. Over sea areas where the average ice cover is projected to diminish, such as the Bothnian Sea and the eastern Gulf of Finland, the mean wind is projected to increase because of a warmer sea surface and reduced stability of the planetary boundary layer [low confidence].
- Air pollution: The impact of climate change on air quality and atmospheric deposition is smaller than the assumed impact of future changes in emissions [low confidence].
- River discharge: River runoff is projected to increase 2–22% in RCP4.5 and 7–22% in RCP8.5. River discharge is projected to increase to the northern and decrease to the southern sub-basins [low confidence]. High flows are projected to decrease in spring and increase in autumn and winter due to earlier snow melt and more winter rain. Over much of continental Europe, an increase in intensity of high flow events is projected with increasing temperature [low confidence].
- Land nutrient inputs: The impact of climate change on land nutrient inputs is smaller than the impact of changes in land management, populations and nutrient point-source releases. In any given river, larger runoff would lead to larger nutrient inputs [medium~~low~~ confidence].
- Terrestrial growing season: Projections suggested that decreasing surface albedo in the Arctic region in winter and spring will notably amplify the future warming in spring (positive feedback), while the increased evapotranspiration will lead to a marked cooling during summer (negative feedback). These feedbacks will stimulate vegetation growth, due to an earlier start of the growing season, leading to compositional changes in woody plants and the distribution of vegetation. Arctic terrestrial ecosystems could continue to sequester carbon until the 2060–2070s, after which the terrestrial ecosystems are projected to turn into weak sources of carbon due to increased soil respiration and biomass burning [low confidence, NEW].
- Terrestrial carbon sequestration: Mitigation scenarios that decrease the fraction of coniferous forest in favour of deciduous forest, and increase the area of deciduous forest in nNorthern Europe from 130,000 to 480,000 km², were projected to reduce near-surface temperatures and give maximum carbon sequestration. [NEW]
- Snow: Projections under RCP8.5 suggest a reduction of the average snow amount between 1981–2010 and 2071–2100 by more 70% for most areas, with the exception of the high Scandinavian mountains, where the warming temperature does not reach the freezing point as often as in lower-lying regions [high confidence]. Sea-effect snowfall events in future climate have not been investigated yet.
- Glaciers: Scandinavian glaciers will lose more than 80% of their current mass by 2100 under RCP8.5, and many are projected to disappear, regardless of future emission scenarios [high confidence, NEW].

Furthermore, river runoff from glaciers is also projected to change regardless of the emission scenario, and to result in increased average winter runoff and in earlier spring peaks [high confidence, NEW].

- Permafrost: In the future climate, the on-going loss of permafrost in the Baltic Sea catchment will very likely accelerate [high confidence].
- Sea ice: Regional climate projections consistently project shrinking and thinning of Baltic Sea ice cover [high confidence], but still estimate that some ice will be formed even in mildest future winters. However, those estimates are based on a limited number of ensembles [members](#) and may not represent future climate variability correctly.
- Lake ice: The observed trends of earlier ice break-up, later freeze-up, and shorter ice cover duration on lakes in the region are projected to continue with future warming, and lakes with intermittent winter ice will consequently become increasingly abundant [high confidence, NEW].
- Water temperature: Coupled atmosphere-ocean regional climate models project an increase in annual mean SST of between 1.2 and 3.2°C, averaged for the Baltic Sea [in](#) the end of the century. The range indicates ensemble mean values for RCP2.6 and RCP8.5 scenarios. Warming will be largest in summer in the northern Baltic Sea [high confidence]. Under both RCP4.5 and RCP8.5, record-breaking summer mean SSTs were projected to increase at the end of the century [medium confidence, NEW]. However, due to the pronounced internal variability there might be decades in the near future without record-breaking events.
- Salinity and saltwater inflows: An increase in river runoff or westerly winds will tend to decrease salinity, but a global sea level rise will tend to increase it, because an enlarged cross-sectional area of the Danish Straits will increase the saltwater imports from the Kattegat. Due to the large uncertainty in projected river runoff, wind and global sea level rise, salinity projections show a wide spread, from increasing to decreasing salinities, and no robust changes were identified [low confidence, NEW].
- Stratification and overturning circulation: Considering all potential drivers of changes in salinity in the Baltic Sea (wind, river runoff, net precipitation, global sea level rise), neither the haline-induced stratification nor the overturning circulation is projected to change [low confidence, NEW]. Projections consistently show that the seasonal thermocline during summer will intensify across nearly the whole Baltic Sea [high confidence, NEW].
- Sea level: Future absolute sea level in the Baltic Sea will continue to rise with the global mean sea level [high confidence]. Its regional manifestation is, however, modulated by the future melting of Antarctica, which affects the Baltic Sea more strongly than the melting of Greenland [low confidence]. Using current estimates, the regional mean sea level is projected to rise by about 87% of the global mean sea level. Land uplift is roughly known but difficult to estimate accurately in practice, as many regional geological factors blur the signature of the glacial isostatic adjustment. Trends in sea level extremes will be determined by the changing mean sea level and possible future changes in storminess. The uncertainty in the latter driver is very large [low confidence].
- Waves: The projected decrease in seasonal sea ice cover will have considerable effects on the wave climate in the northernmost Baltic Sea [high confidence, NEW]. Otherwise, there are no conclusive

4333 results on possible changes in the wave climate and wave extremes, because of the uncertainty about
 4334 changes in wind fields [low confidence].

- 4335 • Sedimentation and coastal erosion: Changes in sea level, wind, waves and sea ice all affect sediment
 4336 transport and coastal erosion. Hence, available projections are highly uncertain [low confidence, NEW].
- 4337 • Oxygen and nutrients: The future response of deep water oxygen conditions will mainly depend on future
 4338 nutrient inputs from land [medium confidence]. However, coastal hypoxia might increase due to warming
 4339 of the water in shallow areas [medium confidence]. Implementation of the BSAP will lead to declining
 4340 phosphorus concentrations [medium confidence, NEW].
- 4341 • Marine CO₂ system: Due to anthropogenic emissions, atmospheric pCO₂ will rise, and consequently also
 4342 the mean pCO₂ of Baltic surface seawater, which has the potential to lower pH [high confidence].
 4343 However, the magnitude of the pH change also depends on alkalinity trends, which are highly uncertain
 4344 [low confidence]. Hence, projections for the Baltic Sea are different from the global ocean.
- 4345 • Microbial communities: The impact of climate change on microbes and the functioning of the microbial
 4346 loop have been studied experimentally. In the northern Gulf of Bothnia, adding DOM increased the
 4347 abundance of bacteria, whereas a temperature increase (from 12 to 15°C) reduced their abundance [low
 4348 confidence, NEW].
- 4349 • Phytoplankton and cyanobacteria: The effect of climate change on phytoplankton and cyanobacteria
 4350 blooms is larger under high nutrient concentrations, but nutrient loads are the dominant driver. If the
 4351 BSAP is fully implemented, the projected environmental status of the Baltic Sea will be significantly
 4352 improved, and extreme cyanobacteria blooms will be rare or absent [low confidence, NEW].
- 4353 • Zooplankton: Experimental studies suggested improved conditions for microzooplankton due to warming
 4354 but negative effects on some larger zooplankton species [low confidence, NEW].
- 4355 • Macroalgae and vascular plants: The direct and indirect effects of changes in temperature, salinity and
 4356 pH are likely to change the geographic distribution of Baltic Sea macrophytes. However, neither
 4357 experimental studies nor past observed changes provide conclusive projections for the effects of climate
 4358 change [low confidence, NEW].
- 4359 • Zoobenthos: In a warmer and less eutrophic Baltic Sea, benthic-pelagic coupling will be weaker, resulting
 4360 in decreasing benthic biomass [low confidence, NEW].
- 4361 • Non-indigenous species: Climate change may favour invasions of non-indigenous species. However, it is
 4362 impossible to project which species may enter the Baltic Sea in future [low confidence].
- 4363 • Fish: Projected changes in temperature and salinity will affect the stocks of cod, sprat and herring.
 4364 However, nutrient loads and especially fishing mortality are also important drivers. Although multi-driver
 4365 modeling studies have been performed, the impact of climate change is ~~unknown~~ ~~uncertain~~ [low confidence,
 4366 NEW].
- 4367 • Marine mammals: Mild winters are known to negatively affect Baltic ringed seals (*Phoca hispida*
 4368 *botnica*) because without their sea ice lair, the pups are more vulnerable to weather and predators, and it
 4369 has been projected that the growth rates of ringed seal populations will decline in the next 90 years. Also
 4370 for grey seals (*Halichoerus grypus*), it has been suggested that reduced ice cover in combination with

(partly climate-driven) changes in the food web, may affect their body condition and birth rate [low confidence].

- Waterbirds: The northward distributional shifts of waterbirds are expected to continue [medium confidence]. Effects on waterbird food will be manifold, but consequences are difficult to predict [low confidence]. The rising sea level and erosion are expected to reduce the availability of breeding habitats [low confidence].
- Marine food webs: Significant alterations in food web structure and functioning can be expected, since species distributions and abundances are expected to change with warming sea-water. The consequences are difficult to project, as research into the long-term dynamics of food webs is still scarce [low confidence].

8 Concluding remarks

We found that

1. The overall conclusions of the BACC I and BACC II assessments remain valid.
2. However, new coupled models (atmosphere-ice-ocean, atmosphere-land), larger ensembles of scenario simulations (CORDEX), new mesocosm experiments (warming, ocean acidification, and dissolved organic matter), extended monitoring (glaciers, satellite data) and homogenized records of observations (MBIs) have led to new insights into past and future climate variability.
3. Improved paleoclimate simulations of the Holocene, new dendroclimatological reconstructions of the past 1000 years and new climate regionalizations have added regional details (east-west gradients over the Baltic Sea region) and improved our understanding of internal variability (sea level extremes, stagnation periods) and the remote impact of low-frequency North Atlantic variability on the Baltic Sea region (AMO, Baltic Sea salinity). New sediment cores suggest that hypoxia during the Medieval Climate Anomaly was caused by climate variability, rather than by human influence, as claimed earlier.
4. Natural variability of many variables of the Earth system is larger than previously realized, requiring larger model ensembles for convincing future projections. Although the first, relatively large ensemble of scenario simulations utilizing a regional coupled atmosphere-ice-ocean model has become available, uncertainty estimates are still incomplete.
5. New regional ESMs including additional components of the Earth system are under development. However, the simulated water cycle is still biased.
6. The first complex multiple-driver study with focus on present and future climates addressing for instance eutrophication of the Baltic Sea, fisheries and climate change has become available and an overall assessment of the various drivers in the Baltic Sea region is part of the BEARs. However, further research on the interplay between drivers is needed.
7. More research on changing extremes was performed, acknowledging that the impact of changing extremes may be more important than that of changing means. However, most observational records are either too short or too heterogeneous for statistical studies of extremes.

- 4407 8. The climate change signal is still confined to increases in observed air and water temperatures, to
 4408 decreases in sea and lake ice, snow cover, permafrost and glacier mass, to the rise in mean sea level, and
 4409 to variables directly related to temperature and the cryosphere, such as ringed seal habitats. Compared to
 4410 the previous BACC report, changes in air temperature, sea ice, snow cover and sea level were shown to
 4411 have accelerated.
- 4412 9. Intensive research on the land-sea interface focussing on the coastal filter has been performed and nutrient
 4413 retention in the coastal zone was estimated for the first time. ~~Uncertainty concerning The~~ unknown
 4414 bioavailability of nutrient ~~inputs~~loads was identified as one of the foremost challenges for marine
 4415 biogeochemistry. However, a model for the entire Baltic Sea coastal zone is still missing and the effect
 4416 of climate change on the coastal filter capacity is still unknown.
- 4417 10. In contradiction to earlier results, observed MBIs have no declining trend. Due to the uncertainties in
 4418 projections of the regional wind, regional precipitation and evaporation, river discharge and global mean
 4419 sea level rise, projections of salinity in the Baltic Sea are inherently uncertain and it remains unknown
 4420 whether the Baltic Sea will become less or more salty. As salinity is a crucial variable for the marine
 4421 ecosystem and for Baltic circulation, projections for the Baltic Sea as a whole are regarded as highly
 4422 uncertain.
- 4423 11. The Baltic Sea may become more acidic in the future, but the decrease in pH may partly be compensated
 4424 by an alkalinity increase, as in the past. Hence, past changes in Baltic carbonate chemistry were different
 4425 from the global ocean acidification, and pH changes may differ also in future.
- 4426 12. Large marine food web changes were observed, which could partly be attributed to warming, brightening
 4427 and sea ice decline. However other factors also play important roles, and many records are too short for
 4428 attribution studies.

4429 **Author contributions**

4430 H.E.M.M. coordinated the assessment and writing of the review article and edited the manuscript. All co-authors
 4431 contributed with either text, figures, tables or comments as detailed in Table 1.

Chapter	Title	Authors
1	Introduction	
1.1	Overview	H.E.M. Meier
1.2	The BACC process	M. Reckermann, H.E.M. Meier
1.3	Summary of BACC I and BACC II key messages	H.E.M. Meier
1.4	Baltic Sea Region characteristics	K. Myrberg
1.5	Global climate change	M. Gröger
2	Methods	
2.1	Assessment of literature	H.E.M. Meier
2.2	Climate model data	H.E.M. Meier

Formatiert: Standard

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

2.3	Uncertainty estimates	H.E.M. Meier
3	Current state of knowledge	
3.1	Past climate change	E. Zorita
3.2	Present climate change	
3.2.1	Present climate change — Atmosphere	
3.2.1.1	Large-scale circulation	M. Stendel, C. Frauen, F. Börgel, H.E.M. Meier, M. Kniebuseh
3.2.1.2	Air temperature	A. Rutgersson
3.2.1.3	Solar radiation	T. Carlund, A. Rutgersson
3.2.1.4	Precipitation	J. Käyhkö, E. Kjellström
3.2.1.5	Wind	M. Stendel
3.2.1.6	Air pollution, air quality and atmospheric — nutrient deposition	M. Quante
3.2.2	Present climate change — Land	
3.2.2.1	River discharge	J. Käyhkö, G. Lindström
3.2.2.2	Land nutrient inputs	O.P. Savchuk, B. Müller-Karulis
3.2.3	Terrestrial biosphere	W. May, P.A. Miller
3.2.4	Present climate change — Cryosphere	
3.2.4.1	Snow	J. Jaagus
3.2.4.2	Glaciers	N. Kirehner
3.2.4.3	Permafrost	G. Hugelius
3.2.4.4	Sea ice	J.J. Haapala
3.2.4.5	Lake ice	J. Käyhkö
3.2.5	Present climate change — Ocean and marine sediments	
3.2.5.1	Water temperature	C. Dieterich, H.E.M. Meier
3.2.5.2	Salinity and saltwater inflows	V. Mohrholz, H.E.M. Meier, K. Myrberg, A. Lehmann
3.2.5.3	Stratification and overturning circulation	M. Gröger, H.E.M. Meier, K. Myrberg
3.2.5.4	Sea level	B. Hünicke, E. Zorita, C. Dieterich, R. Weisse
3.2.5.5	Waves	L. Tuomi
3.2.5.6	Sedimentation and coastal erosion	W. Zhang

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

3.2.5.7	Marine carbonate and biogeochemistry	K. Kulinski, J. Carstensen, B. Müller-Karulis, O. Sævehuk
3.2.6	Marine biosphere	M. Viitasalo, E. Bonsdorff, R. Elmgren, A. Galatius, M. Ahola, V. Dierschke, I. Carlen, M. Frederiksen, E. Gaget, A. Halkka, M. Jüssi, D. Pavon-Jordan
3.3	Future climate change	All
3.3.1	Future climate change Atmosphere	
3.3.1.1	Large-scale circulation	A. Rutgersson, F. Börgel, C. Frauen
3.3.1.2	Air temperature	A. Rutgersson, E. Kjellström, O.B. Christensen
3.3.1.3	Solar radiation	A. Rutgersson, T. Carlund
3.3.1.4	Precipitation	E. Kjellström, O.B. Christensen
3.3.1.5	Wind	M. Stendel, H.E.M. Meier
3.3.1.6	Air pollution, air quality and atmospheric nutrient deposition	M. Quante
3.3.2	Future climate change—Land	▲
3.3.2.1	River discharge	J. Käyhkö
3.3.2.2	Riverine nutrient loads	R. Capell, A. Bartosova
3.3.3	Terrestrial biosphere	W. May, P.A. Miller
3.3.4	Future climate change Cryosphere	
3.3.4.1	Snow	O.B. Christensen
3.3.4.2	Glaciers	N. Kirchner
3.3.4.3	Permafrost	G. Hugelius
3.3.4.4	Sea-ice	J.J. Haapala
3.3.4.5	Lake-ice	J. Käyhkö
3.3.5	Future climate change—Ocean and marine sediments	
3.3.5.1	Water temperature	▲ C. Dieterich, H.E.M. Meier, M. Gröger
3.3.5.2	Salinity and saltwater inflows	H.E.M. Meier
3.3.5.3	Stratification and overturning circulation	▲ M. Gröger, C. Dieterich, H.E.M. Meier
3.3.5.4	Sea-level	B. Hünicke, E. Zorita, C. Dieterich, R. Weisse
3.3.5.5	Waves	L. Tuomi
3.3.5.6	Sedimentation and coastal erosion	W. Zhang

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

3.3.5.7	Marine carbonate and biogeochemistry	K. Kulinski, J. Carstensen, B. Müller-Karulis, O. Saveluk
3.3.6	Marine biosphere	M. Viitasalo, E. Bonsdorff, R. Elmgren, A. Galatius, M. Ahola, V. Dierschke, I. Carlen, M. Frederiksen, E. Gaget, A. Halkka, M. Jüssi, D. Pavon-Jordan
4	Interactions with other drivers	M. Reckermann
5	Comparison with the North Sea region	M. Quante
6	Knowledge gaps	H.E.M. Meier and All
7	Key messages	H.E.M. Meier and All
8	Concluding remarks	H.E.M. Meier and All
Figures and Tables	Analysis of observed time series	M. Kniebusch
Figures and Tables	Analysis of scenario simulations	M. Kniebusch, H.E.M. Meier, C. Dieterich, M. Gröger, O.P. Saveluk, G. Lindström, N. Kirchner, E. Zorita, G. Hugelius


Formatiert: Englisch (USA)

Acknowledgements

During the time in which this paper was prepared, shortly before submission, Christian Dieterich passed away (1964-2021). This sad event marked the end of the life of a distinguished oceanographer and climate scientist who made important contributions to the climate modeling of the Baltic Sea, North Sea and North Atlantic regions. This paper is dedicated to him.

The research presented in this study is part of the Baltic Earth Assessment Reports project of the Baltic Earth program (Earth System Science for the Baltic Sea region, see <http://www.baltic.earth>). Glacier mass loss data were obtained from the SITES Data Portal (<https://data.fieldsites.se/portal/>; see World Glacier Monitoring Service (-, 2021) and Swedish Infrastructure for Ecosystem Science (SITES, 2021a, b, c). We thank Berit Recklebe for technical support and preparation of the reference list.

4444 **Figures**



4445
4446 **Figure 1:** The Baltic Sea and its catchment area, showing climatological mean salinity (in g kg^{-1}) and river runoff
4447 (in mm year^{-1}). (Source: Meier et al., 2014)
4448





Figure 2: Bottom topography of the Baltic Sea and locations of the monitoring stations Arkona Deep (BY2), Bornholm Deep (BY5), Gdansk Deep (BMPL1), Gotland Deep (BY15), Northern Deep (OMTF 0286), Landsort Deep (BY15), and Åland Sea (F64). The Baltic proper comprises the Arkona Basin, Bornholm Basin and Gotland Basin. (Source: H.E. Markus Meier, Leibniz Institute for Baltic Sea Research Warnemünde)






Figure 3: Orbital forcing (irradiance) at 60°N in January and July (derived from Laskar et al., 2004) and the anomalies of reconstructed seasonal temperature and precipitation compared to preindustrial climate (Mauri et al., 2015) in the Baltic Sea region at the Mid-Holocene Optimum (6,000 BPbefore-present). (Source: Eduardo Zorita, Helmholtz-Zentrum Hereon)





Figure 4: Reconstructed spring-early-summer air temperature in the Baltic Sea region (land areas in the box 0-40°E x 55-70°N, deviations from the 20th century mean) derived from Luterbacher et al. (2016). The record is smoothed by a 30-year low-pass filter. The approximate uncertainty range has been estimated here from the data provided by the original publication at interannual and grid-cell scale. (Source: Eduardo Zorita, Helmholtz-Zentrum Hereon)




Figure 5: Normalized winter (December through March; DJFM) mean NAO index during 1821/22-2018/19. Red: positive, blue: negative, black: 10-year running mean. Normalization: $(\text{data} - \text{mean}(\text{data})) / \text{standard deviation}(\text{data})$. (Data source: <https://crudata.uea.ac.uk/cru/data/nao/nao.dat>, compiled by Madline Kniebusch, Leibniz Institute for Baltic Sea Research Warnemünde)




Figure 6: Normalized annual mean AMO index during 1882-2018. Red: positive, blue: negative, black: 10-year running mean. Normalization: $(\text{data} - \text{mean}(\text{data})) / \text{standard deviation}(\text{data})$. (Data source: https://climexp.knmi.nl/data/iamo_hadsst_ts.dat, compiled by Madline Kniebusch, Leibniz Institute for Baltic Sea Research Warnemünde)




Figure 7: Annual and seasonal mean near-surface air temperature anomalies for the Baltic Sea basin for 1871–2020, taken from the CRUTEM4v dataset (Jones et al., 2012), compiled by Anna Rutgersson, Uppsala University. Baseline period is 1961-1990. Blue, red: Baltic Sea basin region north and south, respectively, of 60°N. Dots: individual years. Smoothed curves: variability on timescales longer than 10 years.




Figure 8: Homogenized annual and seasonal mean temperature in Stockholm during 1756-2018-measured at Bolin Centre, Stockholm. Each colored bar show the annual mean temperature, in red or blue, depending on whether the temperature is above or below the average during the reference period 1961-1990. The black curve represents smoothed 10-year mean temperatures. (Source: <https://bolin.su.se/data/stockholm-historical-temps-monthly>)




Figure 9: Mean annual precipitation over land in mm year⁻¹ in the Baltic Sea catchment area during 1950-2018. Blue: whole catchment area, green: North of 59°N, Red: South of 59°N. Bars: annual sum, black: 10-year running mean. (Data source: http://surfobs.climate.copernicus.eu/dataaccess/access_eobs.php#datafiles, compiled by Madline Kniebusch, Leibniz Institute for Baltic Sea Research Warnemünde). Trends: 1.44 mm year⁻¹ (Total), 1.51 mm year⁻¹ (North), 1.37 mm year⁻¹ (South), significant on 99% using the phase-scrambling method (Kniebusch et al., 2019b).




Figure 10: Area weighted river runoff anomalies relative to 1960-1990 (in %) from Sweden to the Baltic Sea. The black solid curve denotes Gaussian filtered data with a standard deviation of three years. (Source: Göran Lindström, Swedish Meteorological and Hydrological Institute):




Figure 11: Long-term dynamics (1970–2017) of annual nitrogen (A) and phosphorus (B) ~~riverineland~~ inputs to the major Baltic Sea basins: BB - Bothnian Bay; BS – Bothnian Sea; BP - Baltic proper; GF - Gulf of Finland; GR - Gulf of Riga; DS – Danish straits; KT – Kattegat. Time (in years) is on the horizontal axis. (Source: O.P. Savchuk, Stockholm University)




Figure 12: Cumulative mass loss for four Swedish glaciers: Storglaciären (since 1946), Rabots glaciär (since 1982, no data for 2004 and 2007 and hence interpolated), Riukojietna (since 1986, data for 2004 interpolated), [andoe](https://data.fieldsites.se/portal/) Mårmaglaciär (since 1990, no data for 2020). Data are accessible from the SITES Data Portal, <https://data.fieldsites.se/portal/> (World Glacier Monitoring Service, 2021; Swedish Infrastructure for Ecosystem Science, 2021a, c, b). (Source: Nina Kirchner, Stockholm University)





Figure 13: Modeled permafrost extent of alpine (> 700 m a.s.l.) and lowland permafrost for the years 1997-2018 in the Baltic Sea drainage basin. Permafrost data from Obu et al. (2020), extent of catchment from Hannerz and Destouni (2006) and elevation data from USGS Global Multi-resolution Terrain Elevation Data 2010 (GMTED2010). Analyses performed at 1 km resolution in an equal area projection. (Source: Gustav Hugelius, Stockholm University)



Formatiert: Schriftart: Nicht unterstrichen, Schriftfarbe: Automatisch




Figure 14: Left: Annual maximum sea-ice extent of the Baltic Sea (MIB) in km^2 during 1720-2020. Blue bars: annual, red: 15-year running mean. The dashed bars represent the error range of the early observations (Vihma and Haapala, 2009). The error range of the 30-year moving average is indicated by two red curves, converging into one when high quality data became available. Right: 30-year distribution functions of MIB during 1909-1938 and 1991-2020. (Data sources: https://www.eea.europa.eu/data-and-maps/daviz/maximum-extent-of-ice-cover-3#tab-chart_1, website Finnish Meteorological Institute: <https://en.ilmatieteenlaitos.fi/ice-season-in-the-baltic-sea>). (Source: Jari Haapala, Finnish Meteorological Institute)




Figure 15: An average sea ice thickness distribution in the Bay of Bothnia. Statistics is based on helicopter electromagnetic measurements conducted in winters 2003, 2004, 2005, 2007 and 2011. Sea-ice thickness distribution in the Bothnian Bay estimated from helicopter electromagnetic measurements during February-March in five winters ((a)-(e)) and it's five-winter average (f), also shown as a red line in (a)-(e). (Source: Ronkainen et al., 2018):





Figure 16: Level-ice thickness at Kemi, Finland and Loviisa, Finland during 1912-2019. **Dots:** annual mean values, **lines:** linear trend with 95% confidence intervals. (Data source: Jari Haapala, Finnish Meteorological Institute).

4572



4573

4574

4575

4576

4577

Figure 17: Length of the ice season in days at Kemi, Loviisa and Utö (Finland) during 1890-2019. **Dots** annual mean, lines: linear trend with 95% confidence intervals. (Data source: Jari Haapala, Finnish Meteorological Institute).




Figure 18: Annual mean values of de-seasonalized daily sea surface (left) and bottom (right) temperature (red dots) at seven monitoring stations during 1877-2018. For the location of the stations see Figure 2. The grey lines show the period when every station has data for every year (1954-2018). For Figures 18, 19, 21 and 22, ICES data (<https://ocean.ices.dk/HydChem/>) for temperature and salinity (bottle data, i.e. from specific depths) were used. Post processing of the data was done following Radtke et al. (2020) in order to overcome possible seasonal biases due to missing values in the observations. Therefore, gaps were statistically filled using a GAMM model (general additive mixed models) taking the seasonality into account. (Source: Madline Kniebusch, Leibniz Institute for Baltic Sea Research Warnemünde)




Figure 19: Seasonal mean sea surface and bottom temperature values during 1877-2018 at Gotland Deep (BY15). Blue: winter, green: spring, yellow: summer and orange: autumn. The grey lines show the period when every station has data for every year (1954-2018). (Source: Madline Kniebusch, Leibniz Institute for Baltic Sea Research Warnemünde)




Figure 20: Annual mean sea surface temperature anomalies to the reference period 2002-2018 of de-seasonalized measurements at the monitoring station BY2 (for the location see Figure 2) and the MARNET stations Darss Sill and Arkona Basin during 1980-2018. (Source: Madline Kniebusch, Leibniz Institute for Baltic Sea Research Warnemünde)




Figure 21: Annual mean values of de-seasonalized daily sea surface (left) and bottom (right) salinity (blue dots) at seven important stations during 1877-2018. The grey lines show the period when every station has data for every year (1954-2018). (Source: Madline Kniebusch, Leibniz Institute for Baltic Sea Research Warnemünde)




Figure 22: Difference between bottom and surface salinity as a measure for the vertical stratification (blue dots) during 1877-2018. Only time steps when both values were available are considered. The grey lines show the period when every station has data for every year (1954-2018). (Source: Madline Kniebusch, Leibniz Institute for Baltic Sea Research Warnemünde)




Figure 23: Annual mean sea level changes in centimeters for 14 Swedish mareographs since 1886. The data are corrected for land uplift. The grey line shows a smoothed curve. (Source: Swedish Meteorological and Hydrological Institute)




Figure 24: Annual mean sea level in Stockholm during 1774-2018. Grey bars: historic time series from Ekman (1988), red: SMHI data (RH2000, 1889-2018), blue: trend computed for 1774-1884 (estimated land uplift: 4.9 mm year⁻¹) and extrapolated until 2018. The SMHI data were adjusted to the historical data, has been bias corrected considering (the mean difference during the overlapping time period.) to make both time series comparable. Sea level rise of SMHI data corrected by estimated land uplift amounts 1.13 mm year⁻¹ during 1889-2018. (Source: Madline Kniebusch, Leibniz Institute for Baltic Sea Research Warnemünde)





Figure 25: Extent of hypoxic ($< 2 \text{ mL O}_2 \text{ L}^{-1}$) and anoxic ($< 0 \text{ mL O}_2 \text{ L}^{-1}$) bottom water (in 10^4 km^2) in the Baltic proper, Gulf of Finland and Gulf of Riga during regular cruises in August-October 1960-2020. (Source: Swedish Meteorological and Hydrological Institute)

Formatiert: Nicht unterstrichen, Schriftfarbe: Automatisch, Hochgestellt



T2m (2070-2099) minus (1970-1999)




Figure 26: (a) Ensemble mean 2 m air temperature change (°C) between 1970-1999 and 2070-2099 and 1970-1999 for winter (December through February/January, upper panels) and summer (June through August, lower panels) under RCP2.6, RCP4.5 and RCP8.5. (b) as (a) but for precipitation change (%mm-day⁻¹). Eight different dynamically downscaled Earth System Model (ESM) simulations are used. (Data source: Gröger et al., 2021b)

Formatiert: Nicht unterstrichen, Schriftfarbe: Automatisch, Nicht Hochgestellt/ Tiefgestellt





Figure 27: From left to right, daily temperature range (DTR, in °C) during historical (1981-2010) and future (2071-2100) periods, and changes in daily temperature range, daily minimum and daily maximum temperatures in RCP8.5 according to EURO-CORDEX scenario simulations (Christensen et al., 2021). Upper and lower panels show winter (December to February, DJF) and summer (June to August, JJA) means, respectively. (Data source: Christensen et al., 2021)

Formatiert: Schriftart: Fett

4643

Formatiert: Schriftart:






Figure 287: Mean projected glacier mass evolution between 2015 and 2100 relative to each region's glacier mass in 2015 (in %) and ± 1 standard deviation under RCP2.6, RCP4.5, and RCP8.5. (Source: Hock et al., 2019; their Figure CB6.1)




Figure 298: Median, 5th and 95th percentiles of the annual maximum sea-ice extent of the Baltic Sea (in 10³ km²) estimated from 28 CMIP5 models. The right-hand side vertical axis shows upper class limits for mild and average ice winters. (a) RCP4.5, (b) RCP8.5. (Source: Luomaranta et al., 2014; their Figure 5 distributed under the terms of the Creative Commons CC-BY 4.0 License, <http://creativecommons.org/licenses/by/4.0/>).


Formatiert: Hochgestellt

Formatiert: Hochgestellt


Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Schriftart:




SST (2070-2099) minus (1970-1999)



Standard deviation

Figure 3029: (a) Ensemble mean sea surface temperature change (°C) between 1970-1999 and 2070-2099 and 1970-1999 for winter (December through February/January, upper panels) and summer (June through August, lower panels) under RCP2.6, RCP4.5, and RCP8.5. (b) as (a) but for the standard deviation of the change, i.e. the ensemble spread (°C). Eight different dynamically downscaled Earth System Model (ESM) simulations are used. (Data Source: Gröger et al., 2019)







Figure 3130: From left to right changes of summer (June to August) mean sea surface temperature (SST; °C), annual mean sea surface salinity (SSS; g kg⁻¹), annual mean bottom salinity (BS; g kg⁻¹), and winter (December to February) mean sea level (SL; cm) between 1978-2007 and 2069-2098 are shown. From top to bottom results of the ensembles by Meier et al. (2011a) (ECOSUPPORT) under the A1B/A2 greenhouse gas emission scenario (white background), and by Saraiva et al. (2019b) (BalticAPP), under RCP-4.5 (grey background) and RCP-8.5 (grey background) are depicted. (Source: Meier et al., 2021a)







Figure 3234: Ensemble mean summer (June to August) bottom dissolved oxygen concentration changes (mL L^{-1}) between 1978-2007 and 2069-2098. From left to right results of the nutrient load scenarios Baltic Sea Action Plan (BSAP), Reference (REF) and Business-As-Usual (BAU) are shown. From top to bottom results of the ensembles by Meier et al. (2011a) ([ECOSUPPORT](#)) under the A1B/A2 greenhouse gas emission scenario (white background), and by Saraiva et al. (2019b) ([BalticAPP](#)) under RCP-4.5 (grey background) and RCP-8.5 (grey background) are depicted. (Source: Meier et al., 2021a)






Figure 3332: As Figure 3230 but for annual mean Secchi depth changes (m). Secchi depth changes indicate changes in water transparency caused by phytoplankton and detritus concentration changes. (Source: Meier et al., 2021a)





Figure 3433: Sea surface temperature anomaly in the Greater North Sea region from 1870 to 2016 (relative to the mean 1971 to 2000), according to different data sets. (Source: Huthnance et al., 2016); updated by Elizabeth Kent, Southampton)




Figure 3534: Synthesis of the knowledge on present and future climate changes. Shown are the anthropogenic climate changes in 33 Earth system variables (bubbles) of the atmosphere (yellow), land surface (brown), terrestrial biosphere (dark green), cryosphere (grey), ocean and sediment (blue), and marine biosphere (light green). The abbreviation NIS stands for non-indigenous species. The sign of a change (plus/minus) is shown together with the level of confidence denoted by the number of signs, i.e. one to three signs correspond to low, medium and high confidence levels, as the result of the literature assessment reflecting consensus and evidence following the IPCC definitions (Section 2.3). Sign colours indicate the direction of past (black) and future (red) changes following

4702 Table [1549](#). Uncertain changes (+/-) are not displayed. Investigated external anthropogenic drivers of the Earth
4703 system are shown as red squares, i.e. greenhouse gases, in particular CO₂, and aerosol emissions. [The prevailing](#)
4704 [climate change attribution relationships with sufficiently high confidence are shown by arrows \(black: heat cycle,](#)
4705 [blueorange: water cycle, orange: momentum cycle including sea level changes, pink: carbon cycle\).](#) Projections
4706 of carbon sequestration of Arctic terrestrial ecosystems for the 21st century showed first increased uptake and later
4707 a carbon source (Section 3.3.3), denoted by 1. + 2. -. [Future changes in mean sea level will be dominated by](#)
4708 [thermal expansion of the global ocean and melting of ice sheets outside the Baltic Sea region. Probably the impact](#)
4709 [of regional changes are small.](#)
4710

4711

4712 **Tables**4713 **Table 1:** Table of contents and contributing authors.

Chapter	Title	Authors
<u>1</u>	<u>Introduction</u>	
<u>1.1</u>	<u>Overview</u>	<u>H.E.M. Meier</u>
<u>1.2</u>	<u>The BACC and BEAR projects</u>	<u>M. Reckermann, H.E.M. Meier</u>
<u>1.3</u>	<u>Summary of BACC I and BACC II key messages</u>	<u>H.E.M. Meier</u>
<u>1.4</u>	<u>Baltic Sea region characteristics</u>	
<u>1.4.1</u>	<u>Climate variability of the Baltic Sea region</u>	<u>K. Myrberg</u>
<u>1.4.2</u>	<u>A unique brackish water basin</u>	<u>K. Myrberg</u>
<u>1.4.3</u>	<u>The Baltic Sea – a specific European sea</u>	<u>K. Myrberg</u>
<u>1.5</u>	<u>Global climate change</u>	▲
<u>1.5.1</u>	<u>Atmosphere</u>	<u>M. Gröger</u>
<u>1.5.1.1</u>	<u>Surface air temperature</u>	<u>M. Gröger</u>
<u>1.5.1.2</u>	<u>Precipitation</u>	<u>M. Gröger</u>
<u>1.5.2</u>	<u>Cryosphere</u>	<u>M. Gröger</u>
<u>1.5.3</u>	<u>Ocean</u>	<u>M. Gröger</u>
<u>1.5.3.1</u>	<u>Sea level</u>	<u>M. Gröger</u>
<u>1.5.3.2</u>	<u>Water temperature and salinity</u>	<u>M. Gröger</u>
<u>1.5.3.3</u>	<u>Atlantic Meridional Overturning Circulation</u>	<u>M. Gröger</u>
<u>1.5.4</u>	<u>Marine biosphere</u>	<u>M. Gröger</u>
<u>1.5.5</u>	<u>Coupled Model Intercomparison Project 6</u>	<u>H.E.M. Meier, M. Gröger, E. Kjellström</u>
<u>2</u>	<u>Methods</u>	▲
<u>2.1</u>	<u>Assessment of literature</u>	<u>H.E.M. Meier</u>
<u>2.2</u>	<u>Proxy data, instrumental measurements and climate model data</u>	
<u>2.2.1</u>	<u>Past climate</u>	<u>H.E.M. Meier</u>

Formatiert: Deutsch (Deutschland)

Formatiert: Links

Formatiert: Links

Formatiert: Links

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

<u>2.2.2</u>	<u>Present climate</u>	<u>H.E.M. Meier</u>
<u>2.2.3</u>	<u>Future climate</u>	<u>H.E.M. Meier</u>
<u>2.3</u>	<u>Uncertainty estimates</u>	<u>H.E.M. Meier</u>
<u>3</u>	<u>Current state of knowledge</u>	
<u>3.1</u>	<u>Past climate change</u>	
<u>3.1.1</u>	<u>Key messages from previous assessments</u>	<u>E. Zorita</u>
<u>3.1.2</u>	<u>New paleoclimate reconstructions</u>	<u>E. Zorita</u>
<u>3.1.3</u>	<u>Holocene climate evolution</u>	<u>E. Zorita</u>
<u>3.1.4</u>	<u>The past millennium</u>	<u>E. Zorita</u>
<u>3.2</u>	<u>Present climate change</u>	
<u>3.2.1</u>	<u>Atmosphere</u>	
<u>3.2.1.1</u>	<u>Large-scale atmospheric circulation</u>	<u>M. Stendel, C. Frauen, F. Börgel, H.E.M. Meier, M. Kniebusch</u>
<u>3.2.1.2</u>	<u>Air temperature</u>	<u>A. Rutgersson</u>
<u>3.2.1.3</u>	<u>Solar radiation and cloudiness</u>	<u>T. Carlund, A. Rutgersson</u>
<u>3.2.1.4</u>	<u>Precipitation</u>	<u>J. Käyhkö, E. Kjellström</u>
<u>3.2.1.5</u>	<u>Wind</u>	<u>M. Stendel</u>
<u>3.2.1.6</u>	<u>Air pollution, air quality and atmospheric nutrient deposition</u>	<u>M. Quante</u>
<u>3.2.2</u>	<u>Land</u>	
<u>3.2.2.1</u>	<u>River discharge</u>	<u>J. Käyhkö, G. Lindström</u>
<u>3.2.2.2</u>	<u>Land nutrient inputs</u>	<u>O.P. Savchuk, B. Müller-Karulis</u>
<u>3.2.3</u>	<u>Terrestrial biosphere</u>	<u>W. May, P.A. Miller</u>
<u>3.2.4</u>	<u>Cryosphere</u>	
<u>3.2.4.1</u>	<u>Snow</u>	<u>J. Jaagus</u>
<u>3.2.4.2</u>	<u>Glaciers</u>	<u>N. Kirchner</u>
<u>3.2.4.3</u>	<u>Permafrost</u>	<u>G. Hugelius</u>
<u>3.2.4.4</u>	<u>Sea ice</u>	<u>J.J. Haapala</u>
<u>3.2.4.5</u>	<u>Lake ice</u>	<u>J. Käyhkö</u>
<u>3.2.5</u>	<u>Ocean and marine sediments</u>	
<u>3.2.5.1</u>	<u>Water temperature</u>	<u>C. Dieterich, H.E.M. Meier</u>
<u>3.2.5.2</u>	<u>Salinity and saltwater inflows</u>	<u>V. Mohrholz, H.E.M. Meier, K. Myrberg</u>
<u>3.2.5.3</u>	<u>Stratification and overturning circulation</u>	<u>M. Gröger, H.E.M. Meier, K. Myrberg</u>
<u>3.2.5.4</u>	<u>Sea level</u>	<u>B. Hünicke, E. Zorita, C. Dieterich, R. Weisse</u>

Formatiert: Schriftart: Nicht Fett

Formatiert: Schriftart: Nicht Fett

Formatiert: Schriftart: Nicht Fett

Formatiert: Schriftart: Nicht Fett

3.2.5.5	Waves	L. Tuomi
3.2.5.6	Sedimentation and coastal erosion	W. Zhang
3.2.5.7	Marine carbonate system and biogeochemistry	▲
3.2.5.7.1	Oxygen and nutrients	K. Kulinski, J. Carstensen, B. Müller-Karulis, O. Savchuk
3.2.5.7.2	Marine CO₂ system	K. Kulinski, J. Carstensen, B. Müller-Karulis, O. Savchuk
3.2.6	Marine biosphere	
3.2.6.1	Pelagic habitats	M. Viitasalo, E. Bonsdorff, R. Elmgren
3.2.6.1.1	Microbial communities	M. Viitasalo, E. Bonsdorff, R. Elmgren
3.2.6.1.2	Phytoplankton and cyanobacteria	M. Viitasalo, E. Bonsdorff, R. Elmgren
3.2.6.1.3	Zooplankton	M. Viitasalo, E. Bonsdorff, R. Elmgren
3.2.6.2	Benthic habitats	M. Viitasalo, E. Bonsdorff, R. Elmgren
3.2.6.2.1	Macroalgae and vascular plants	M. Viitasalo, E. Bonsdorff, R. Elmgren
3.2.6.2.2	Zoobenthos	M. Viitasalo, E. Bonsdorff, R. Elmgren
3.2.6.3	Non-indigenous species	M. Viitasalo, E. Bonsdorff, R. Elmgren
3.2.6.4	Fish	M. Viitasalo, E. Bonsdorff, R. Elmgren
3.2.6.5	Marine mammals	A. Galatius, M. Ahola, I. Carlen, A. Halkka, M. Jüssi
3.2.6.6	Waterbirds	V. Dierschke, M. Frederiksen, E. Gaget, D. Pavon-Jordan
3.2.6.7	Marine food webs	E. Bonsdorff, M. Viitasalo, R. Elmgren
3.3	Future climate change	
3.3.1	Atmosphere	
3.3.1.1	Large-scale atmospheric circulation	A. Rutgersson, F. Börgel, C. Frauen
3.3.1.2	Air temperature	A. Rutgersson, E. Kjellström, O.B. Christensen
3.3.1.3	Solar radiation	A. Rutgersson, T. Carlund
3.3.1.4	Precipitation	E. Kjellström, O.B. Christensen
3.3.1.5	Wind	M. Stendel, H.E.M. Meier
3.3.1.6	Air pollution, air quality and atmospheric nutrient deposition	M. Quante
3.3.2	Land	

Formatiert: Englisch (USA)

Formatiert: Tiefgestellt

Formatiert: Schwedisch (Schweden)

Formatiert: Schwedisch (Schweden)

Formatiert: Schwedisch (Schweden)

3.3.2.1	River discharge	J Käyhkö
3.3.2.2	Riverine nutrient loads	R. Capell, A. Bartosova
3.3.3	Terrestrial biosphere	W. May, P.A. Miller
3.3.4	Cryosphere	
3.3.4.1	Snow	O.B. Christensen
3.3.4.2	Glaciers	N. Kirchner
3.3.4.3	Permafrost	G. Hugelius
3.3.4.4	Sea ice	J.J. Haapala
3.3.4.5	Lake ice	J. Käyhkö
3.3.5	Ocean and marine sediments	
3.3.5.1	Water temperature	C. Dieterich, H.E.M. Meier, M. Gröger
3.3.5.2	Salinity and saltwater inflows	H.E.M. Meier
3.3.5.3	Stratification and overturning circulation	M. Gröger, C. Dieterich, H.E.M. Meier
3.3.5.4	Sea level	B. Hünicke, E. Zorita, C. Dieterich, R. Weisse
3.3.5.5	Waves	L. Tuomi
3.3.5.6	Sedimentation and coastal erosion	W. Zhang
3.3.5.7	Marine carbonate system and biogeochemistry	▲
3.3.5.7.1	Oxygen and nutrients	K. Kulinski, J. Carstensen, B. Müller-Karulis, O. Savchuk
3.3.5.7.2	Marine CO2 system	K. Kulinski, J. Carstensen, B. Müller-Karulis, O. Savchuk
3.3.6	Marine biosphere	
3.3.6.1	Pelagic habitats	M. Viitasalo, E. Bonsdorff, R. Elmgren
3.3.6.1.1	Microbial communities	M. Viitasalo, E. Bonsdorff, R. Elmgren
3.3.6.1.2	Phytoplankton and cyanobacteria	M. Viitasalo, E. Bonsdorff, R. Elmgren
3.3.6.1.3	Zooplankton	M. Viitasalo, E. Bonsdorff, R. Elmgren
3.3.6.2	Benthic habitats	M. Viitasalo, E. Bonsdorff, R. Elmgren
3.3.6.2.1	Macroalgae and vascular plants	M. Viitasalo, E. Bonsdorff, R. Elmgren
3.3.6.2.2	Zoobenthos	M. Viitasalo, E. Bonsdorff, R. Elmgren
3.3.6.3	Non-indigenous species	M. Viitasalo, E. Bonsdorff, R. Elmgren
3.3.6.4	Fish	M. Viitasalo, E. Bonsdorff, R. Elmgren
3.3.6.5	Marine mammals	A. Galatius, M. Ahola, I. Carlen, A. Halkka, M. Jüssi

Formatiert: English (USA)

<u>3.3.6.6</u>	<u>Water birds</u>	<u>V. Dierschke, M. Frederiksen, E. Gaget, D. Pavon-Jordan</u>
<u>3.3.6.7</u>	<u>Marine food webs</u>	<u>E. Bonsdorff, M. Viitasalo, R. Elmgren</u>
<u>4</u>	<u>Interactions of climate with other anthropogenic drivers</u>	<u>M. Reckermann</u>
<u>5</u>	<u>Comparison with the North Sea region</u>	
<u>5.1</u>	<u>The North Sea region</u>	<u>M. Quante</u>
<u>5.2</u>	<u>A few selected and highly aggregated results from NOSCCA</u>	<u>M. Quante</u>
<u>5.3</u>	<u>Some differences in climate change effects between the North Sea and Baltic Sea</u>	<u>M. Quante</u>
<u>6</u>	<u>Knowledge gaps</u>	<u>H.E.M. Meier and All</u>
<u>7</u>	<u>Key messages</u>	
<u>7.1</u>	<u>Past climate changes</u>	<u>H.E.M. Meier and All</u>
<u>7.2</u>	<u>Present climate changes</u>	<u>H.E.M. Meier and All</u>
<u>7.3</u>	<u>Future climate changes</u>	<u>H.E.M. Meier and All</u>
<u>8</u>	<u>Concluding remarks</u>	<u>H.E.M. Meier and All</u>
<u>Figures and Tables</u>	<u>Analysis of observed, reconstructed and proxy-data time series</u>	<u>M. Kniebusch, J.J. Haapala, G. Hugelius, G. Lindström, N. Kirchner, M. Quante, A. Rutgersson, O.P. Savchuk, E. Zorita</u>
<u>Figures and Tables</u>	<u>Analysis of scenario simulations</u>	<u>O.B. Christensen, C. Dieterich, M. Gröger, E. Kjellström, H.E.M. Meier</u>

Formatiert: Schriftart: Nicht Fett

Formatiert: Schriftart: Nicht Fett

Formatiert: Schriftart: Nicht Fett

Formatiert: Schriftart: Nicht Fett

Formatiert: Schriftart: Nicht Fett

Formatiert: Schriftart: Nicht Fett

Formatiert: Deutsch (Deutschland)

Formatiert: Deutsch (Deutschland)

Table 2: Variables of this assessment and further references (1: Lehmann et al., 2021; 2: Kuliński et al., 2021; 3: Rutgersson et al., 2021; 4: Weisse et al., 2021; 5: Reckermann et al., 2021; 6: Gröger et al., 2021a; 7: Christensen et al., 2021; 8: Meier et al., 2021a; 9: Viitasalo and Bonsdorff, 2021)

Number	Variable	Past and present climates		Future climate	
Atmosphere					
1	Large-scale <u>atmospheric</u> circulation	3.2.1.1	3	3.3.1.1	3, 7
2	Air temperature Warm spell Cold spell	3.1.2, 3.1.3, 3.1.4 3.2.1.2	 3 3	3.3.1.2	7 3 3
3	Solar radiation and cloudiness	3.2.1.3		3.3.1.3	7
4	Precipitation Heavy precipitation Drought	3.1.2, 3.1.3, 3.1.4 3.2.1.4	 3 3	3.3.1.4	7 3 3
5	Wind Storm	3.2.1.5	3	3.3.1.5	7 3
6	Air pollution, air quality and atmospheric deposition	3.2.1.6		3.3.1.6	
Land					
7	River discharge High flow	3.2.2.1	3	3.3.2.1	8 3
8	Land nutrient inputs	3.2.2.2		3.3.2.2	8
Terrestrial biosphere					
9	Land cover (forest, crops, grassland, peatland, mires)	3.2.3	6	3.3.3	
10	Carbon sequestration			3.3.3	
Cryosphere					
11	Snow Sea-effect snowfall	3.2.4.1	3	3.3.4.1	7 3
12	Glaciers	3.2.4.2		3.3.4.2	

Formatiert: Schriftart: Fett, Nicht unterstrichen, Schriftfarbe: Automatisch, Deutsch (Deutschland)

Formatiert: Deutsch (Deutschland)

13	Permafrost	3.2.4.3		3.3.4.3	
14	Sea ice	3.2.4.4		3.3.4.4	8
	Extreme mild winter		3		3
	Severe winter		3		3
	Ice ridging		3		3
15	Lake ice	3.2.4.5		3.3.4.5	
Ocean and marine sediments					
16	Water temperature	3.2.5.1		3.3.5.1	8
	Marine heat wave		3		3
17	Salinity and saltwater inflows	3.2.5.2	1	3.3.5.2	8
18	Stratification and overturning circulation	3.2.5.3	1	3.3.5.3	8
19	Sea level	3.2.5.4	4	3.3.5.4	8
	Sea level extreme		3		3
20	Waves	3.2.5.5	4	3.3.5.5	
	Extreme waves		3		3
21	Sedimentation and coastal erosion	3.2.5.6	4	3.3.5.6	
22	Oxygen and nutrients	3.1.4			
		3.2.5.7.1	2	3.3.5.7.1	8
23	Marine CO ₂ system	3.2.5.7.2	2	3.3.5.7.2	
Marine biosphere					
24	Pelagic habitats: Microbial communities	3.2.6.1.1	2, 9	3.3.6.1.1	9
25	Pelagic habitats: Phytoplankton and cyanobacteria	3.2.6.1.2	2, 3, 9	3.3.6.1.2	3, 9
26	Pelagic habitats: Zooplankton	3.2.6.1.3	9	3.3.6.1.3	9
27	Benthic habitats: Macroalgae and vascular plants	3.2.6.2.1	9	3.3.6.2.1	9
28	Benthic habitats: Zoobenthos	3.2.6.2.2	9	3.3.6.2.2	9

4719

29	Non-indigenous species	3.2.6.3	9	3.3.6.3	9
30	Fish	3.2.6.4	9	3.3.6.4	9
31	Marine mammals	3.2.6.5	9	3.3.6.5	9
32	Waterbirds	3.2.6.6	9	3.3.6.6	9
33	Marine food web	3.2.6.7	9	3.3.6.7	9

Table 3: List of acronyms (in alphabetical order), their definitions, comments and references

Acronym	Definition	Comment	Reference	Formatiert
AMO	Atlantic Multidecadal Oscillation	Mode of climate variability	Knight et al. (2005)	Formatiert
AMOC	Atlantic Meridional Overturning Circulation	Large-scale ocean circulation pattern, part of the global conveyor belt	Buckley and Marshall (2016)	Formatiert
AO	Arctic Oscillation	Teleconnection pattern	Deser (2000), Ambaum et al. (2001)	Formatiert
BACC	Assessment of climate change for the Baltic Sea basin	Regional climate change assessment	BACC Author Team (2008), BACC II Author Team (2015)	Formatiert
BALTEX	Baltic Sea Experiment	Earth System Science for the Baltic Sea Region programme, predecessor of Baltic Earth	https://www.baltex-research.eu , https://baltic.earth	Formatiert
BalticAPP	Well-being from the Baltic Sea: applications combining natural science and economics	Climate modelling project for the Baltic Sea	Saraiva et al. (2019a)	Formatiert
BEAR	Baltic Earth assessment reports	Regional climate change assessment	https://baltic.earth	Formatiert
BSAP	Baltic Sea Action Plan	Nutrient load abatement strategy for the Baltic Sea	HELCOM (2013b)	Formatiert
BSAP, REF, BAU, WORST	Baltic Sea Action Plan, Reference, Business-As-Usual and WORST	Nutrient load scenarios	Meier et al. (2012a), Saraiva et al. (2019b)	Formatiert
CMIP	Coupled Model Intercomparison Project of the World Climate Research Programme	GCM/ESM results from CMIP3, CMIP5 and CMIP6 were assessed	https://www.wcrp-climate.org/wgcm-cmip	Formatiert
DTR	Daily temperature range	Climate index of air temperature		Formatiert
DOC	Dissolved organic carbon	Environmental variable		Formatiert
DOM	Dissolved organic matter	Environmental variable		Formatiert
ECOSUPPORT	Advanced modelling tool for scenarios of the Baltic Sea Ecosystem to SUPPORT decision making	Climate modelling project for the Baltic Sea	Meier et al. (2014)	Formatiert
EN CLIME	Expert Network on Climate Change	Joint HELCOM - Baltic Earth expert network	https://helcom.fi/helcom-at-work/groups/state-and-conservation/en-clime/	Formatiert

<u>ESM</u>	<u>Earth System Model</u>	<u>Model applied for global climate simulations including the carbon cycle</u>	<u>Heavens et al. (2013)</u>	Formatiert
				Formatiert
				Formatiert
<u>EURO-CORDEX</u>	<u>Coordinated Downscaling Experiment: European Domain</u>	<u>High-resolution climate change projections for European impact research</u>	<u>Jacob et al. (2014), https://euro-cordex.net/</u>	Formatiert
				Formatiert
				Formatiert
<u>GCM</u>	<u>General Circulation Model</u>	<u>Model applied for global climate simulations</u>	<u>Meehl et al. (2004)</u>	Formatiert
				Formatiert
<u>GGCB</u>	<u>Global Glacier Change Bulletin</u>		<u>Zemp et al. (2020)</u>	Formatiert
<u>HELCOM</u>	<u>Helsinki Commission</u>	<u>Consists of the Baltic Sea countries and the European Union</u>	<u>https://helcom.fi</u>	Formatiert
				Formatiert
				Formatiert
<u>ICES</u>	<u>International Council of the Exploration of the Sea</u>	<u>Marine science organisation</u>	<u>https://www.ices.dk</u>	Formatiert
				Formatiert
<u>IPCC</u>	<u>Intergovernmental Panel of Climate Change</u>	<u>Assessment reports (AR) of past and future changes in 1990, 1995, 2001, 2008, 2013, 2021, <i>inter alia</i> based upon CMIP results</u>	<u>http://www.ipcc.ch</u>	Formatiert
				Formatiert
				Formatiert
				Formatiert
<u>MBI</u>	<u>Major Baltic Inflow</u>	<u>Barotropic inflow of saltwater into the Baltic Sea</u>	<u>Matthäus and Frank (1992), Schinke and Matthäus (1998)</u>	
<u>MIB</u>	<u>Annual maximum sea-ice extent of the Baltic Sea</u>	<u>Climate index</u>	<u>https://www.eea.europa.eu/data-and-maps/data/external/the-classification-of-the-maximum-https://en.ilmatiiteenlaitos.fi/ice-season-in-the-baltic-sea</u>	
<u>MSL, GMSL, RMSL</u>	<u>Mean sea level, global mean sea level, relative mean sea level</u>	<u>Reference sea level</u>		Formatiert
<u>NAO</u>	<u>North Atlantic Oscillation</u>	<u>Mode of climate variability</u>	<u>Hurrell (1995)</u>	Formatiert
<u>NOSCCA</u>	<u>North Sea Region Climate Change Assessment</u>	<u>Regional climate change assessment</u>	<u>Quante and Colijn (2016)</u>	Formatiert
				Formatiert
<u>OA</u>	<u>Ocean acidification</u>	<u>Effect of rising atmospheric CO₂</u>	<u>Doney et al. (2009), Havenhand (2012)</u>	Formatiert
				Formatiert
<u>PM</u>	<u>Particulate matter</u>	<u>Environmental variable</u>		Formatiert
				Formatiert: Tiefgestellt

4721

4722
4723

Table 42: Comparison of the Baltic Sea with other intra-continental seas and large lakes.

Basin	Area	Mean depth	Mean salinity	Fresh-water budget	Ice cover on average	Location Centre
Unit	10 ³ km ²	m	g kg ⁻¹			Lat Long

Formatierte Tabelle

4723
4724

Baltic Sea	393	54	7.414	+	Half	60°N 20°E
Black Sea	436	1197	1820	+	Northeast	43°N 35°E
Gulf of Ob	41	12	5	+	All	73°N 74°E
Chesapeake Bay	12	6	15	+	Shores	38°N 76°W
Hudson Bay	1232	128	30	+	All	58°N 85°W
Red Sea	438	491	40	–	None	22°N 38°E
Persian Gulf	239	25	40	–	None	27°N 52°E
Caspian Sea	374	211	12	0	North	43°N 50°E
Lake Superior	82	149	< 0.1	0	All	48°N 88°W

Formatierte Tabelle

Table 53: The main characteristics of the physical features of the European Seas (Leppäranta and Myrberg, 2009; Sündermann and Pohlmann, 2011; <http://www.ospar.org>; British Oceanographic Data Centre). Greater North Sea – being the neighbouring sea area to the Baltic Sea - is shown as a sub-region of the NE-Atlantic, but other European sub-regions are not listed (from Myrberg et al., 2019).

Basin	Area 10 ³ km ²	Mean depth m	Mean salinity g kg ⁻¹	Fresh—water budget	Ice cover on average	Tides	Water residence time (years)
Baltic Sea	393	54	7.4	Pos.	37 % ¹⁾	Weak	40
Black Sea	43622	1 1972 00	18	Pos.	Northeast only	Weak	3 000
Greater North Sea	750	80	34–35	Pos.	No	Strong	Not applicable
Mediterranean Sea	2 970	1 500	38	Neg.	No	Weak/ Moderate	80-100
NE Atlantic shelf	13 500 ²⁾	1 500	34–35	Not applicable	No	Strong	Not applicable
1) Mean maximal ice cover between 2000-2017, see Fig. 143.2.4.4.1							
2) defined as the OSPAR convention area, incl. the Greater North Sea							

Table 6: Ensemble mean changes in temperature (°C) and precipitation (%) at selected future global mean surface temperature (GMST) levels (1.5 to 4.0°C) compared to preindustrial (PI) conditions during 1850-1900. Listed are annual, winter (December to January, DJF) and summer (June to August, JJA) means from CMIP5, CMIP6 and their differences for Northern Europe (NEU) and Western and Central Europe (WCE). N is the number of considered members in the two ensembles. Note that CMIP5 and CMIP6 are driven under RCP8.5 and SSP5-8.5 scenarios, respectively. (Source: IPCC AR6 interactive atlas (Iturbide et al., 2021; Gutiérrez et al., 2021) assessed on 25.11.2022, Erik Kjellström, Swedish Meteorological and Hydrological Institute)

GMST vs PI		CMIP5 N=33				CMIP6 N=34				CMIP6-CMIP5		
		ANN	DJF	JJA		ANN	DJF	JJA		ANN	DJF	JJA
Temp (°C)	NEU	NEU				NEU				NEU		
	1.5	2.3	3.1	1.8		2.3	3	2		0	-0.1	0.2
	2.0	2.9	3.8	2.4		3	3.9	2.7		0.1	0.1	0.3
	3.0	4.3	5.6	3.6		4.2	5.2	3.8		-0.1	-0.4	0.2
	4.0	5.6	7.1	4.7		5.5	6.4	5.2		-0.1	-0.7	0.5
	WCE	WCE				WCE				WCE		
	1.5	2	2.5	1.9		2.1	2.6	2.1		0.1	0.1	0.2
	2.0	2.6	3.1	2.6		2.8	3.3	3		0.2	0.2	0.4
	3.0	4	4.6	4		4	4.6	4.4		0	0	0.4
	4.0	5.2	6	5.6		5.3	5.7	6		0.1	-0.3	0.4
		CMIP5 N=28				CMIP6 N=33				CMIP6-CMIP5		
		ANN	DJF	JJA		ANN	DJF	JJA		ANN	DJF	JJA
Precip (%)	NEU	NEU				NEU				NEU		
	1.5	7	10.9	2.7		6.3	10.7	1.3		-0.7	-0.2	-1.4
	2.0	8.8	13.6	3.6		7.7	13	1.1		-1.1	-0.6	-2.5
	3.0	12.3	21.3	3.1		10	18.4	0.1		-2.3	-2.9	-3.0
	4.0	16.7	29.3	2.5		13.7	23.3	-0.9		-3.0	-6.0	-3.4
	WCE	WCE				WCE				WCE		
	1.5	2.9	8.5	-4.4		3.1	7.7	-3.8		0.2	-0.8	0.6
	2.0	3.3	10.8	-6.7		2.7	9.7	-8		-0.6	-1.1	-1.3
	3.0	3.4	15.7	-12		3.5	15.1	-13		0.1	-0.6	-1.6
	4.0	3.6	21.7	-18		2.8	19.4	-18		-0.8	-2.3	-0.4

Formatiert: Schriftart: Nicht Fett

Formatiert: Schriftart: Nicht Fett

Formatiert: Schriftart: Nicht Fett

Table 7: As Table 6 but ensemble mean changes in temperature (°C) and precipitation (%) at selected future time slices (2021-2040, 2041-2060 and 2081-2100) compared to preindustrial (PI) conditions during 1850-1900. (Source: IPCC AR6 interactive atlas (Iturbide et al., 2021; Gutiérrez et al., 2021) assessed on 25.11.2022, Erik Kjellström, Swedish Meteorological and Hydrological Institute)

GMST vs PI	CMIP5 N=33			CMIP6 N=34			CMIP6-CMIP5		
	ANN	DJF	JJA	ANN	DJF	JJA	ANN	DJF	JJA
Temp (°C)	NEU			NEU			NEU		
2021-2040	2.5	3.3	2	2.6	3.3	2.3	0.1	0	0.3
2041-2060	3.6	4.7	3	3.6	4.5	3.3	0	-0.2	0.3
2081-2100	6.1	7.6	5.3	6.3	7.5	5.9	0.2	-0.1	0.6
	WCE			WCE			WCE		
2021-2040	2.2	2.7	2.2	2.3	2.8	2.4	0.1	0.1	0.2
2041-2060	3.3	3.9	3.3	3.4	4	3.7	0.1	0.1	0.4
2081-2100	5.8	6.5	6.2	6.3	6.7	7.1	0.5	0.2	0.9
Precip (%)	CMIP5 N=28			CMIP6 N=33			CMIP6-CMIP5		
	ANN	DJF	JJA	ANN	DJF	JJA	ANN	DJF	JJA
	NEU			NEU			NEU		
2021-2040	7	10.9	2.7	6.3	10.7	1.3	-0.7	-0.2	-1.4
2041-2060	8.8	13.6	3.6	7.7	13	1.1	-1.1	-0.6	-2.5
2081-2100	12.3	21.3	3.1	10	18.4	0.1	-2.3	-2.9	-3.0
	WCE			WCE			WCE		
2021-2040	3.5	9.8	-4.1	3.1	8.3	-5	-0.4	-1.5	-0.9
2041-2060	3.2	12.8	-9.3	3.7	13	-9.3	0.5	0.2	0.0
2081-2100	4	23	-19.6	3.3	24.4	-23.8	-0.7	1.4	-4.2

Formatiert: Schriftart: Nicht Fett

Formatiert: Schriftart: Nicht Fett

Formatiert: Schriftart: Nicht Fett

Table 8: As Table 6 ensemble mean changes in temperature (°C) and precipitation (%) at selected future global mean surface temperature (GMST) levels (1.5 to 4.0°C) compared to preindustrial (PI) conditions during 1986-2005. In addition to CMIP5 and CMIP6, also EURO-CORDEX ensemble mean scenario simulations are shown. Note that the regional climate models of EURO-CORDEX are driven bei different ESMs than the presented ESMs of the CMIP5 ensemble. (Source: IPCC AR6 interactive atlas (Iturbide et al., 2021; Gutiérrez et al., 2021) assessed on 25.11.2022, Erik Kjellström, Swedish Meteorological and Hydrological Institute)

GMST vs PI		CMIP5 N=33				EURO N=47				CMIP6 N=34				EURO-CMIP5				CMIP6-CMIP5		
		ANN	DJF	JJA		ANN	DJF	JJA		ANN	DJF	JJA		ANN	DJF	JJA		ANN	DJF	JJA
	Temp (°C)	NEU				NEU				NEU				NEU				NEU		
1.5		1.3	1.6	1.1		1.3	1.3	1		1.4	1.6	1.3		0	-0.3	-0.1		0.1	0	0.2
2.0		2	2.4	1.7		1.9	2.3	1.5		2.1	2.5	2		-0.1	-0.1	-0.2		0.1	0.1	0.3
3.0		3.4	4.2	2.9		3	3.4	2.5		3.3	3.9	3.1		-0.4	-0.8	-0.4		-0.1	-0.3	0.2
4.0		4.5	5.6	4		4.1	4.9	3.4		4.5	4.9	4.5		-0.4	-0.7	-0.6		0	-0.7	0.5
		WCE				WCE				WCE				WCE				WCE		
1.5		1.3	1.3	1.3		1	0.9	1		1.5	1.6	1.8		-0.3	-0.4	-0.3		0.2	0.3	0.5
2.0		1.9	1.9	2		1.5	1.6	1.4		2.2	2.3	2.6		-0.4	-0.3	-0.6		0.3	0.4	0.6
3.0		3.2	3.4	3.5		2.7	2.8	2.6		3.6	3.6	4		-0.5	-0.6	-0.9		0.4	0.2	0.5
4.0		4.5	4.7	4.9		3.8	4.2	3.6		4.7	4.6	5.5		-0.7	-0.5	-1.3		0.2	-0.1	0.6
		CMIP5 N=28				EURO N=48				CMIP6 N=33				EURO-CMIP5				CMIP6-CMIP5		
		ANN	DJF	JJA		ANN	DJF	JJA		ANN	DJF	JJA		ANN	DJF	JJA		ANN	DJF	JJA
	Precip (%)	NEU				NEU				NEU				NEU				NEU		
1.5		4	4.7	2.5		3.5	2.6	3.3		4.1	6.2	1.3		-0.5	-2.1	0.8		0.1	1.5	-1.2
2.0		5.7	7.2	3.4		5.9	6.9	5.7		5.5	8.4	1.1		0.2	-0.3	2.3		-0.2	1.2	-2.3
3.0		9.2	14.4	3		8.4	10.3	5.1		8	13.5	0.1		-0.8	-4.1	2.1		-1.2	-0.9	-2.9
4.0		13.3	22.5	2.3		13.6	17.6	9.8		11.2	18.2	-0.1		0.3	-4.9	7.5		-2.1	-4.3	-2.4
		WCE				WCE				WCE				WCE				WCE		
1.5		2.5	4.4	-1.5		2.5	5	0.5		2	4.9	-3.6		0	0.6	2		-0.5	0.5	-2.1
2.0		2.9	6.5	-3.8		4.5	7.7	2.1		1.6	6.6	-7.8		1.6	1.2	5.9		-1.3	0.1	-4.0
3.0		3	11.3	-9		5.4	11.1	-0.6		2.3	12	-13.2		2.4	-0.2	8.4		-0.7	0.7	-4.2
4.0		3.3	16.9	-16		7.3	16.3	-1.8		1.4	14.8	-18		4	-0.6	13.7		-1.9	-2.1	-2.5

Formatiert: Schriftart: Nicht Fett

Formatiert: Schriftart: Nicht Fett

Formatiert: Schriftart: Nicht Fett

Table 94: Linear surface air temperature trends (K decade^{-1}) for the period 1878–2020 over the northern ($>60^\circ\text{N}$) and southern ($<60^\circ\text{N}$) Baltic Sea basin (1878–2020 is selected for comparison with (Rutgersson et al., 2014), with an equally long time period). Bold: significance at $p < 0.05$. Data from the updated CRUTEM4v dataset (Jones et al., 2012). (Source: Anna Rutgersson, Uppsala University)

	Annual	Winter	Spring	Summer	Autumn
North	0.10	0.11	0.14	0.08	0.08
South	0.10	0.13	0.10	0.09	0.09

Table 105: Average (2013-2017) riverine and coastal nutrient inputs (10^3 t N (P) yr^{-1}) to the major basins of the Baltic Sea (Source: Oleg P. Savchuk, Stockholm University). For abbreviated basin names see Figure 11. TN and TP are total nitrogen and total phosphorus inputs, respectively. (Source: Oleg P. Savchuk, Stockholm University)

	BB	BS	BP	GF	GR	DS	KT	Entire BS
TN river	48	40	251	85	73	32	46	575
TN coast	3.6	4.3	6.5	9.3	0.5	2.8	2.0	29
TP river	2.4	1.7	11.8	3.4	2.0	1.1	1.3	24.0
TP coast	0.1	0.2	0.5	0.4	0.1	0.2	0.1	1.6

Table 116: Mass balances for the Swedish glaciers Storglaciären, Rabots glaciär, Mårmaglaciär, and Riuekojietna. General references are given as footnotes in connection with balance years and long-term monitoring intervals, respectively. Selected specific references are given as footnotes in connection with the glacier names, and include also neighboring glaciers, Kårsa glaeier and Kebnepakteglaeiär. (Source: Nina Kirchner, Stockholm University)

	Recent mass balance years. Gains and losses in mm w.e. (millimeter water equivalent). Note that the unit mm w.e. is interchangeable with the unit kg m ⁻²				Long-term mass balance, losses per year in mm w.e.	
	2015/ 2016 ³	2016/ 2017 ³	2017/ 2018 ^{4,3}	2018/2019 ⁴	1980-2010 ⁵	1985-
Storglaciären ^{7,8,9}	-240	+470	-1600	-310	-113	-153
Rabots glaciär ^{10,11}	-650	-170	-1590	-650	-394	-465
Mårmaglaciär	-370	+260	-1370	-910	-430	-460
Riuekojietna	-1060	+150	-1400	-610	-592	-592

Formatiert: Nicht unterstrichen, Schriftfarbe: Automatisch, Hochgestellt

³ World Glacier Monitoring Service, 2020

⁴ World Glacier Monitoring Service, 2021

⁵ Blunden and Arendt, 2015

⁶ Hartfield et al., 2018

⁷ Mercer, 2016

⁸ Holmlund and Holmlund, 2019

⁹ Kirchner et al., 2019

¹⁰ Brugger and Pankratz, 2015

¹¹ Williams et al., 2016

Formatiert: Schriftart: 10 Pt.

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Table 127: Air temperature (T_{2m}) changes ($^{\circ}\text{C}$) between 1976 - 2005 and 2069 - 2098 averaged over each season and annual mean over the Baltic Sea catchment area and over the Baltic Sea calculated from [nine dynamically downscaled ESM simulations](#) ~~nine regionalized ESMs~~ (Data source: [Gröger et al., 2021b](#), compiled by [Christian Dieterich, Swedish Meteorological and Hydrological Institute](#)). In addition to the ensemble mean change, the 5th and 95th percentiles indicating the ensemble spread are listed (in brackets). (Data source: [Gröger et al., 2021b](#), compiled by [Christian Dieterich, Swedish Meteorological and Hydrological Institute](#))

	Annual	Winter	Spring	Summer	Autumn
Total land					
RCP2.6	1.5 (1.2, 2.0)	2.1 (1.5, 3.3)	1.5 (1.2, 2.0)	1.3 (0.8, 2.1)	1.3 (0.9, 1.8)
RCP4.5	2.6 (1.6, 3.2)	3.2 (2.1, 4.2)	2.4 (1.5, 3.3)	2.1 (1.3, 3.1)	2.3 (1.4, 2.8)
RCP8.5	4.3 (3.5, 5.2)	5.0 (3.4, 6.3)	3.8 (3.1, 4.5)	3.7 (2.5, 5.0)	3.8 (2.6, 4.8)
Land north of 60°N					
RCP2.6	1.7 (1.4, 2.4)	2.5 (1.9, 3.1)	1.7 (1.2, 2.3)	1.4 (0.8, 2.3)	1.5 (1.1, 2.1)
RCP4.5	2.9 (2.0, 3.7)	4.0 (2.9, 5.0)	2.8 (1.8, 3.8)	2.3 (1.3, 3.4)	2.5 (1.7, 3.2)
RCP8.5	4.9 (3.9, 5.9)	6.0 (4.2, 7.5)	4.2 (3.5, 5.1)	3.9 (2.8, 5.1)	4.2 (2.9, 5.3)
Land south of 60°N					
RCP2.6	1.4 (1.0, 1.8)	1.7 (1.1, 3.4)	1.3 (0.9, 1.7)	1.3 (0.9, 1.9)	1.2 (0.7, 1.6)
RCP4.5	2.2 (1.3, 2.8)	2.6 (1.5, 4.0)	2.2 (1.3, 2.9)	2.0 (1.1, 3.0)	2.1 (1.2, 2.7)
RCP8.5	3.9 (3.2, 4.7)	4.2 (2.9, 5.7)	3.4 (2.9, 4.0)	3.5 (2.2, 4.9)	3.5 (2.3, 4.5)
Baltic Sea					
RCP2.6	1.4 (1.2, 1.9)	1.9 (1.3, 2.8)	1.5 (1.1, 1.9)	1.2 (0.6, 1.8)	1.2 (0.9, 1.7)
RCP4.5	2.4 (1.4, 2.9)	2.9 (1.8, 3.7)	2.5 (1.5, 3.1)	2.0 (1.2, 2.7)	2.1 (1.2, 2.7)
RCP8.5	3.9 (3.1, 4.8)	4.6 (3.2, 5.8)	3.9 (3.0, 4.9)	3.5 (2.4, 4.6)	3.6 (2.6, 4.6)

Table 138: Relative precipitation changes (%) between 1976 - 2005 and 2069 - 2098 averaged over each season and annual mean over the Baltic Sea catchment area and over the Baltic Sea calculated from [nine dynamically downscaled ESM simulations](#)~~nine regionalized ESMs~~ (Data source: [Gröger et al., 2021b](#), compiled by [Christian Dieterich, Swedish Meteorological and Hydrological Institute](#)). In addition to the ensemble mean change, the 5th and 95th percentiles indicating the ensemble spread are listed (in brackets). (Data source: [Gröger et al., 2021b](#), compiled by [Christian Dieterich, Swedish Meteorological and Hydrological Institute](#))

	Annual	Winter	Spring	Summer	Autumn
Total land					
RCP2.6	5 (2, 14)	7 (1, 22)	8 (2, 12)	3 (-2, 13)	4 (-4, 12)
RCP4.5	9 (6, 14)	12 (4, 24)	13 (8, 17)	4 (1, 11)	6 (-5, 12)
RCP8.5	15 (11, 22)	22 (11, 38)	20 (7, 26)	5 (-4, 15)	13 (-1, 18)
Land north of 60°N					
RCP2.6	6 (2, 15)	7 (2, 23)	8 (0, 13)	5 (1, 17)	5 (-5, 14)
RCP4.5	11 (7, 18)	13 (6, 27)	15 (2, 21)	9 (4, 14)	8 (-3, 17)
RCP8.5	19 (12, 30)	22 (12, 41)	24 (7, 35)	13 (-1, 30)	17 (1, 26)
Land south of 60°N					
RCP2.6	5 (0, 13)	7 (-1, 22)	7 (3, 13)	2 (-5, 10)	3 (-7, 11)
RCP4.5	7 (4, 11)	12 (1, 22)	12 (6, 20)	1 (-5, 11)	4 (-8, 11)
RCP8.5	12 (8, 18)	21 (9, 35)	18 (7, 26)	-1 (-14, 9)	9 (-3, 17)
Baltic Sea					
RCP2.6	6 (0, 15)	5 (-3, 15)	4 (-1, 8)	8 (0, 22)	5 (-3, 13)
RCP4.5	8 (3, 13)	9 (-4, 20)	11 (1, 17)	6 (-1, 16)	6 (-3, 15)
RCP8.5	16 (8, 23)	18 (3, 31)	19 (-3, 32)	10 (-9, 22)	15 (4, 26)

Table 149: Sea surface temperature (SST) changes (°C) between 1976 - 2005 and 2069 - 2098 averaged over each season and annual mean over the Baltic Sea calculated from nine dynamically downscaled ESM simulations~~nine regionalized ESMs~~ (Data source: Gröger et al., 2021b, compiled by Christian Dieterich, Swedish Meteorological and Hydrological Institute). In addition to the ensemble mean change, the 5th and 95th percentiles indicating the ensemble spread are listed (in brackets). (Data source: Gröger et al., 2021b, compiled by Christian Dieterich, Swedish Meteorological and Hydrological Institute)

Baltic Sea	Annual	Winter	Spring	Summer	Autumn
RCP2.6	1.1 (0.8, 1.6)	1.0 (0.9, 1.4)	1.1 (0.9, 1.6)	1.2 (0.6, 1.7)	0.9 (0.7, 1.6)
RCP4.5	1.8 (1.1, 2.5)	1.7 (1.0, 2.3)	1.9 (1.2, 2.6)	2.0 (1.2, 2.6)	1.8 (1.1, 2.4)
RCP8.5	3.2 (2.5, 4.1)	3.0 (2.3, 3.8)	3.2 (2.5, 3.9)	3.4 (2.4, 4.5)	3.1 (2.4, 4.1)

4785 **Table 1510:** Summary of key messages about the impact of global warming on selected variables. The sign of a
4786 change (plus/minus) is listed together with the level of confidence denoted by the number of signs, i.e. one to three
4787 signs correspond to low, medium and high confidence levels. +/- means no detected or projected change due to
4788 climate change. Key messages of this assessment that are new compared to the previous assessment by BACC II
4789 Author Team (2015) are marked and a brief explanation is provided in the neighboring column. (NA = North
4790 Atlantic)

Number	Variable	Present climates		Future climate	
Atmosphere					
1	Large-scale circulation	+/-	Remote influence of the multi-decadal variability in the NA on the Baltic Sea	+/-	Impact of warming Arctic with declining sea ice might be relevant
2	Air temperature Warm spell Cold spell	+++ ++ --	Accelerated warming	+++ ++ --	Greater confidence due to increased ensemble size, coupled atmosphere-ocean models
3	Solar radiation	+	Comparison between various satellite products	+/-	GCM and RCM systematically differ
4	Precipitation Heavy precipitation Drought north (south) of 59°N	+ ++ -(+)		++ +++ -(+)	convection-resolving models became available
5	Wind Number of deep cyclones	+/- +		+ +/-	Small systematic increase in winter in the northern Baltic where the sea ice will melt
6	Air pollution, air quality and atmospheric deposition	+/-		+/-	
Land					
7	River discharge High flow ¹² in the north (south)	+/- +/- (+/-)	Dataset of observed time series for the past century from Sweden merged with	+ - (+)	Changing seasonality (decrease of river discharge in spring, increase in winter)

¹² Based upon annual maximum river discharges of daily data for Sweden with 10- and 100-year repeat periods (Roudier et al., 2016) and for Finland with 100-year repeat period (Veijalainen et al., 2010)

			high-resolution dynamic model projections of the upcoming century		may affect the occurrence of floods ^{13,14}
8	Land nutrient inputs	+/-		+	
Terrestrial biosphere					
9	Growing season in the Baltic Sea region	++	Study based on satellite data available	+/-	No new study
10	Carbon sequestration in northern terrestrial ecosystems	+/-		+, later -	First increasing sinks. Weak sources of carbon after 2060-2070s due to increased soil respiration and biomass burning
Cryosphere					
11	Snow Sea-effect snowfall	--- +/-		--- +/-	
12	Ice mass of glaciers	---	Since 2006 inventories of all Scandinavian glaciers have become available	---	High-resolution projections of Scandinavian glaciers available
13	Permafrost	--	High-resolution modeling	---	
14	Sea ice cover Extreme mild winter Extreme severe winter Ice ridging	--- +++ --- +/-		--- +++ --- -	
15	Lake ice	---	Systematic assessment available	---	Projections for global lake ice available
Ocean and marine sediments					
16	Water temperature Marine heat wave	+++ +	Accelerated warming	+++ +++	Increasing number of record-breaking summer

¹³ Roudier et al., 2016

¹⁴ Veijalainen et al., 2010

					mean SST events and number of heat waves
17	Salinity and saltwater inflows	+/-	Homogenous data of saltwater inflows, north-south salinity gradient has increased	+/-	Uncertainty sources of salinity due to wind, river discharge and global sea level rise changes were assessed
18	Stratification and overturning circulation	+/-	Systematic study of monitoring data since the 1980s	+/-	Intensified seasonal thermoclines during summer but no change of the halocline and overturning circulation
19	Absolute sea level Storm surge relative to the mean sea level	+++ +/-	Paleoclimate study on sea level extremes did not show systematic changes in changing climate, dissensus in the literature ^{15,16}	+++ +/-	Dissensus in the literature ^{17,18}
20	Waves Extreme waves	+/- +/-		+ +	Small increase in winter in the northern Baltic Sea
21	Sedimentation and coastal erosion	+/-		+/-	First modeling studies available, <u>but large uncertainty</u>
22	Hypoxic area	+	Warming contributed to the historical spread of hypoxia in the deep water and in the coastal zone, sediment cores suggest that changing climate	++	Oxygen decline in the coastal zone due to warming

¹⁵ Ribeiro et al., 2014

¹⁶ Marcos and Woodworth, 2017

¹⁷ Vousdoukas et al., 2016

¹⁸ Vousdoukas et al., 2017

			caused hypoxia during the Medieval Climate Anomaly instead of agriculture		
23	CO ₂ uptake pH southern (northern) Baltic Sea	++ --(+/-)	New observations and modeling, positive alkalinity trends identified	+/- +/-	
Marine biosphere					
24	Microbial communities	+	In the northern Baltic Sea increased riverine dissolved organic matter suppressed phytoplankton biomass production and shifts the carbon flow towards microbial heterotrophy	+/-	Increase of dissolved organic matter and temperature will enhance and decrease the abundance of bacteria, respectively
25	growing season of phytoplankton (cyanobacteria) cyanobacteria biomass ratio between diatom and dinoflagellate biomasses since 1901	++ +/- -	 new indicator for the environmental status developed	+/- + +/-	Warming causes prolonged and intensified cyanobacteria blooms but the nutrient control is dominating
26	Zooplankton	+/-		+/-	Increasing microzooplankton biomass
27	Macroalgae and vascular plants	+/-	Systematic studies on benthic ecosystems	+/-	

28	Zoobenthos	+/-	Systematic studies on benthic ecosystems, spreading of non-indigenous such as polychaete <i>Marenzelleria</i> spp.	+/-	Weaker benthic-pelagic coupling and decreasing benthic biomass in a warmer and less eutrophic Baltic
29	Non-indigenous species	+		++	
30	Fish	+/-	Food web modeling including fisheries	+/-	Multi-driver (climate change, eutrophication, fisheries) food web projections were performed
31	Populations of marine mammals	+/-		-	
32	Waterbird migration	+++	Northward shift of the wintering range of waterbirds	++	Controlled by food availability
33	Marine food web	+/-		+/-	

4791

4792 **References**

- 4793 Aalto, J., Pirinen, P., and Jylhä, K.: New gridded daily climatology of Finland: Permutation-based uncertainty
4794 estimates and temporal trends in climate, *J. Geophys. Res-Atmos.*, 121, 3807-3823,
4795 <https://doi.org/10.1002/2015JD024651>, 2016.
- 4796 Aarvak, T., Jostein Øien, I., Krasnov, Y. V., Gavrilov, M. V., and Shavykin, A. A.: The European wintering
4797 population of Steller's Eider *Polysticta stelleri* reassessed, *Bird Conserv. Int.*, 23, 337-343,
4798 <https://doi.org/10.1017/S0959270912000251>, 2013.
- 4799 Aberle, N., Malzahn, A. M., Lewandowska, A. M., and Sommer, U.: Some like it hot: the protozooplankton-
4800 copepod link in a warming ocean, *Mar. Ecol. Prog. Ser.*, 519, 103-113, <https://doi.org/10.3354/meps11081> 2015.
- 4801 Adam, J. C., Haddeland, I., Su, F., and Lettenmaier, D. P.: Simulation of reservoir influences on annual and
4802 seasonal streamflow changes for the Lena, Yenisei, and Ob' rivers, *J. Geophys. Res-Atmos.*, 112,
4803 <https://doi.org/10.1029/2007JD008525>, 2007.
- 4804 Adam, J. C., and Lettenmaier, D. P.: Application of New Precipitation and Reconstructed Streamflow Products
4805 to Streamflow Trend Attribution in Northern Eurasia, *J. Climate*, 21, 1807-1828,
4806 <https://doi.org/10.1175/2007jcli1535.1>, 2008.
- 4807 Ahlgren, J., Grimvall, A., Omstedt, A., Rolff, C., and Wikner, J.: Temperature, DOC level and basin interactions
4808 explain the declining oxygen concentrations in the Bothnian Sea, *J. Mar. Sys.*, 170, 22-30,
4809 <https://doi.org/10.1016/j.jmarsys.2016.12.010>, 2017.
- 4810 Åkerman, H. J., and Johansson, M.: Thawing permafrost and thicker active layers in sub-arctic Sweden,
4811 *Permafrost Periglac.*, 19, 279-292, <https://doi.org/10.1002/ppp.626>, 2008.
- 4812 Akhtar, N., Geyer, B., Rockel, B., Sommer, P. S., and Schrum, C.: Accelerating deployment of offshore wind
4813 energy alter wind climate and reduce future power generation potentials, *Sci. Rep.-UK*, 11, 11826,
4814 <https://doi.org/10.1038/s41598-021-91283-3>, 2021.
- 4815 Al-Janabi, B., Kruse, I., Graiff, A., Karsten, U., and Wahl, M.: Genotypic variation influences tolerance to
4816 warming and acidification of early life-stage *Fucus vesiculosus* L. (Phaeophyceae) in a seasonally fluctuating
4817 environment, *Mar. Biol.*, 163, 14, <https://doi.org/10.1007/s00227-015-2804-8>, 2016.
- 4818 Alkama, R., and Cescatti, A.: Biophysical climate impacts of recent changes in global forest cover, *Science*, 351,
4819 600-604, <https://doi.org/10.1126/science.aac8083>, 2016.
- 4820 Allen, M. R., Dube, O. P., Solecki, W., Aragón-Durand, F., Cramer, W., Humphreys, S., Kainuma, M., Kala, J.,
4821 Mahowald, N., Mulugetta, Y., Perez, R., Wairiu, M., and Zickfeld, K.: Framing and Context, in: *Global*
4822 *Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial*
4823 *levels and related global greenhouse gas emission pathways, in the context of strengthening the global response*
4824 *to the threat of climate change, sustainable development, and efforts to eradicate poverty*, edited by: Masson-
4825 Delmotte, V., Zhai, P., Pörtner, H.-O., Roberts, D., Skea, J., Shukla, P. R., Pirani, A., Moufouma-Okia, W.,
4826 Péan, C., Pidcock, R., Connors, S., Matthews, J. B. R., Chen, Y., Zhou, X., Gomis, M. I., Lonnoy, E., Maycock,
4827 T., Tignor, M., and Waterfield, T., <https://www.ipcc.ch/sr15/chapter/chapter-1/>, 2018.
- 4828 Allen, R. J., Norris, J. R., and Wild, M.: Evaluation of multidecadal variability in CMIP5 surface solar radiation
4829 and inferred underestimation of aerosol direct effects over Europe, China, Japan, and India, *J. Geophys. Res-
4830 Atmos.*, 118, 6311-6336, <https://doi.org/10.1002/jgrd.50426>, 2013.
- 4831 Allen, S. G., Ainley, D. G., Page, G. W., and Ribic, C. A.: The effect on distribution on harbor seal haul out
4832 patterns at Bolinas Lagoon, California, in: *Fish. Bull.*, edited by: Richards, W. J., Collette, B. B., Houde, E. D.,
4833 Ingham, M. C., Lasker, R., Malins, D. C., Pella, J. J., Quast, J. C., Sindermann, C. J., and Fukuyama, M. S.,
4834 National Marine Fisheries Service, NOAA, Seattle, WA, USA, 493-500, 1984.
- 4835 Almroth-Rosell, E., Edman, M., Eilola, K., Meier, H. E. M., and Sahlberg, J.: Modelling nutrient retention in the
4836 coastal zone of an eutrophic sea, *Biogeosciences*, 13, 5753-5769, <https://doi.org/10.5194/bg-13-5753-2016>,
4837 2016.

Formatiert: Englisch (USA)

4838 [Ambaum, M. H., Hoskins, B. J., and Stephenson, D. B.: Arctic oscillation or North Atlantic oscillation?. *Journal of*](#)
4839 [Climate, 14\(16\), 3495-3507, \[https://doi.org/10.1111/brv.12221\]\(https://doi.org/10.1175/1520-0442\(2001\)014<3495:AOONAO>2.0.CO;2, 2001.</p>
<p>4840 Andersen, J. H., Carstensen, J., Conley, D. J., Dromph, K., Fleming-Lehtinen, V., Gustafsson, B. G., Josefson,

4841 A. B., Norkko, A., Villnäs, A., and Murray, C.: Long-term temporal and spatial trends in eutrophication status of

4842 the Baltic Sea, <i>Biol. Rev.</i>, 92, 135-149, <a href=\), 2017.](#)

4843 Andersson, A., Meier, H. E. M., Ripszám, M., Rowe, O., Wikner, J., Haglund, P., Eilola, K., Legrand, C.,
4844 Figueroa, D., Paczkowska, J., Lindehoff, E., Tysklind, M., and Elmgren, R.: Projected future climate change and
4845 Baltic Sea ecosystem management, *Ambio* 44, 345-356, <https://doi.org/10.1007/s13280-015-0654-8>, 2015.

4846 Andersson, H. C.: Influence of long-term regional and large-scale atmospheric circulation on the Baltic sea level,
4847 *Tellus A*, 54, 76-88, <https://doi.org/10.3402/tellusa.v54i1.12125>, 2002.

4848 Arheimer, B., Dahné, J., and Donnelly, C.: Climate change impact on riverine nutrient load and land-based
4849 remedial measures of the Baltic Sea Action Plan, *Ambio*, 41, 600-612, [https://doi.org/10.1007/s13280-012-0323-](https://doi.org/10.1007/s13280-012-0323-0)
4850 0, 2012.

4851 Arheimer, B., and Lindström, G.: Climate impact on floods: changes in high flows in Sweden in the past and the
4852 future (1911-2100), *Hydrol. Earth Syst. Sci.*, 19, 771-784, <https://doi.org/10.5194/hess-19-771-2015>, 2015.

4853 Arheimer, B., Donnelly, C., and Lindström, G.: Regulation of snow-fed rivers affects flow regimes more than
4854 climate change, *Nat. Commun.*, 8, 62, <https://doi.org/10.1038/s41467-017-00092-8>, 2017.

4855 Armitage, J. M., Quinn, C. L., and Wania, F.: Global climate change and contaminants—an overview of
4856 opportunities and priorities for modelling the potential implications for long-term human exposure to organic
4857 compounds in the Arctic, *J. Environ. Monitor.*, 13, 1532-1546, <https://doi.org/10.1039/C1EM10131E>, 2011.

4858 Arneborg, L.: Comment on “Influence of sea level rise on the dynamics of salt inflows in the Baltic Sea” by R.
4859 Hordoir, L. Axell, U. Löptien, H. Dietze, and I. Kuznetsov, *J. Geophys. Res-Oceans*, 121, 2035-2040,
4860 <https://doi.org/10.1002/2015JC011451>, 2016.

4861 Asmala, E., Carstensen, J., Conley, D. J., Slomp, C. P., Stadmark, J., and Voss, M.: Efficiency of the coastal
4862 filter: Nitrogen and phosphorus removal in the Baltic Sea, *Limnol. Oceanogr.*, 62,
4863 <https://doi.org/10.1002/lno.10644>, 2017.

4864 Athanasiadis, P. J., Yeager, S., Kwon, Y.-O., Bellucci, A., Smith, D. W., and Tibaldi, S.: Decadal predictability
4865 of North Atlantic blocking and the NAO, *npj Clim. Atmos. Sci.*, 3, 20, [https://doi.org/10.1038/s41612-020-0120-](https://doi.org/10.1038/s41612-020-0120-6)
4866 6, 2020.

4867 Averkiev, A. S., and Klevanny, K. A.: A case study of the impact of cyclonic trajectories on sea-level extremes
4868 in the Gulf of Finland, *Cont. Shelf Res.*, 30, 707-714, <https://doi.org/10.1016/j.csr.2009.10.010>, 2010.

4869 Axell, L., Liu, Y., Jandt, S., Lorkowski, I., Lindenthal, A., Verjovkina, S., and Schwichtenberg, F.: Baltic Sea
4870 Production Centre BALTICSEA_REANALYSIS_BIO_003_012 COPERNICUS Marine Environment
4871 Monitoring Service, Quality Information Document, 51pp, 2019.

4872 BACC Author Team: Assessment of climate change for the Baltic Sea basin, *Regional Climate Studies*, Springer
4873 Science & Business Media, Berlin, Heidelberg, 473 pp., 2008.

4874 BACC II Author Team: Second Assessment of Climate Change for the Baltic Sea Basin, *Regional Climate*
4875 *Studies*, Springer International Publishing, Cham, 2015.

4876 Bailey, S. A., Brown, L., Campbell, M. L., Canning-Clode, J., Carlton, J. T., Castro, N., Chainho, P., Chan, F.
4877 T., Creed, J. C., Curd, A., Darling, J., Fofonoff, P., Galil, B. S., Hewitt, C. L., Inglis, G. J., Keith, I., Mandrak, N.
4878 E., Marchini, A., McKenzie, C. H., Occhipinti-Ambrogi, A., Ojaveer, H., Pires-Teixeira, L. M., Robinson, T. B.,
4879 Ruiz, G. M., Seaward, K., Schwindt, E., Son, M. O., Theriault, T. W., and Zhan, A.: Trends in the detection of
4880 aquatic non-indigenous species across global marine, estuarine and freshwater ecosystems: A 50-year
4881 perspective, *Divers. Distrib.*, 26, 1780-1797, <https://doi.org/10.1111/ddi.13167>, 2020.

4882 BALTEX: Minutes of 17th Meeting of the BALTEX Science Steering Group held at Institute of Meteorology
4883 and Water Management (IMGW), Poznan, Poland, 24 - 26 November 2004, International BALTEX Secretariat
4884 Publication No. 32, 100pp, 2005.

Formatiert: Englisch (USA)

Formatiert: Schriftart: Nicht Kursiv

Formatiert: Schriftart: +Textkörper (Calibri)

4885 [Baltic Earth: Baltic Earth Science Plan 2017, International Baltic Earth Secretariat Publication No. 11,](#)
4886 [Helmholtz-Zentrum Geesthacht GmbH, Germany, 34pp, 2017.](#)
4887 Bamber, J. L., Oppenheimer, M., Kopp, R. E., Aspinall, W. P., and Cooke, R. M.: Ice sheet contributions to
4888 future sea-level rise from structured expert judgment, *P. Natl. Acad. Sci. USA*, 116, 11195,
4889 <https://doi.org/10.1073/pnas.1817205116>, 2019.

4890 Barbosa, S. M., and Donner, R. V.: Long-term changes in the seasonality of Baltic sea level, *Tellus A*, 68,
4891 30540, <https://doi.org/10.3402/tellusa.v68.30540>, 2016.

4892 Barnes, E. A.: Revisiting the evidence linking Arctic amplification to extreme weather in midlatitudes, *Geophys.*
4893 *Res. Lett.*, 40, 4734-4739, <https://doi.org/10.1002/grl.50880>, 2013.

4894 Barregard, L., Molnár, P., Jonson, J. E., and Stockfelt, L.: Impact on Population Health of Baltic Shipping
4895 Emissions, *Int. J. Environ. Res. Public Health*, 16, 1954, <https://doi.org/10.3390/ijerph16111954>, 2019.

4896 Bartók, B., Wild, M., Folini, D., Lüthi, D., Kotlarski, S., Schär, C., Vautard, R., Jerez, S., and Imecs, Z.:
4897 Projected changes in surface solar radiation in CMIP5 global climate models and in EURO-CORDEX regional
4898 climate models for Europe, *Clim. Dyn.*, 49, 2665-2683, <https://doi.org/10.1007/s00382-016-3471-2>, 2017.

4899 Bartolino, V., Margonski, P., Lindegren, M., Linderholm, H. W., Cardinale, M., Rayner, D., Wennhage, H., and
4900 Casini, M.: Forecasting fish stock dynamics under climate change: Baltic herring (*Clupea harengus*) as a case
4901 study, *Fish. Oceanogr.*, 23, 258-269, <https://doi.org/10.1111/fog.12060>, 2014.

4902 Bartosova, A., Capell, R., Olesen, J. E., Jabloun, M., Refsgaard, J. C., Donnelly, C., Hyytiäinen, K., Pihlainen,
4903 S., Zandersen, M., and Arheimer, B.: Future socioeconomic conditions may have a larger impact than climate
4904 change on nutrient loads to the Baltic Sea, *Ambio*, 48, 1325-1336, <https://doi.org/10.1007/s13280-019-01243-5>,
4905 2019.

4906 Bates, A. E., Pecl, G. T., Frusher, S., Hobday, A. J., Wernberg, T., Smale, D. A., Sunday, J. M., Hill, N. A.,
4907 Dulvy, N. K., Colwell, R. K., Holbrook, N. J., Fulton, E. A., Slawinski, D., Feng, M., Edgar, G. J., Radford, B.
4908 T., Thompson, P. A., and Watson, R. A.: Defining and observing stages of climate-mediated range shifts in
4909 marine systems, *Glob. Environ. Change*, 26, 27-38, <https://doi.org/10.1016/j.gloenvcha.2014.03.009>, 2014.

4910 Bauer, B., Meier, H. E. M., Casini, M., Hoff, A., Margoński, P., Orio, A., Saraiva, S., Steenbeek, J., and
4911 Tomczak, M. T.: Reducing eutrophication increases spatial extent of communities supporting commercial
4912 fisheries: a model case study, *ICES J. Mar. Sci.*, 75, 1306-1317, <https://doi.org/10.1093/icesjms/fsy023>, 2018.

4913 Bauer, B., Gustafsson, B. G., Hyytiäinen, K., Meier, H. E. M., Müller-Karulis, B., Saraiva, S., and Tomczak, M.
4914 T.: Food web and fisheries in the future Baltic Sea, *Ambio*, <https://doi.org/10.1007/s13280-019-01229-3>, 2019.

4915 Belkin, I. M.: Rapid warming of large marine ecosystems, *Prog. Oceanogr.*, 81, 207-213,
4916 <https://doi.org/10.1016/j.pocean.2009.04.011>, 2009.

4917 Bengtsson, L., Hodges, K. I., and Keenlyside, N.: Will Extratropical Storms Intensify in a Warmer Climate?, *J.*
4918 *Climate*, 22, 2276-2301, <https://doi.org/10.1175/2008jcli2678.1>, 2009.

4919 Bergen, B., Endres, S., Engel, A., Zark, M., Dittmar, T., Sommer, U., and Jürgens, K.: Acidification and
4920 warming affect prominent bacteria in two seasonal phytoplankton bloom mesocosms, *Environ. Microbiol.*, 18,
4921 4579-4595, <https://doi.org/10.1111/1462-2920.13549>, 2016.

4922 Bergström, L., Heikinheimo, O., Svirsgden, R., Kruze, E., Ložys, L., Lappalainen, A., Saks, L., Minde, A.,
4923 Dainys, J., Jakubavičiūtė, E., Ådjers, K., and Olsson, J.: Long term changes in the status of coastal fish in the
4924 Baltic Sea, *Estuar. Coast. Shelf Sci.*, 169, 74-84, <https://doi.org/10.1016/j.ecss.2015.12.013>, 2016.

4925 Bergström, S., and Carlsson, B.: River runoff to the Baltic Sea - 1950-1990, *Ambio*, 23, 280-287, 1994.

4926 Bermúdez, R., Winder, M., Stühr, A., Almén, A. K., Engström-Öst, J., and Riebesell, U.: Effect of ocean
4927 acidification on the structure and fatty acid composition of a natural plankton community in the Baltic Sea,
4928 *Biogeosciences*, 13, 6625-6635, <https://doi.org/10.5194/bg-13-6625-2016>, 2016.

4929 Berner, C., Bertos-Fortis, M., Pinhassi, J., and Legrand, C.: Response of Microbial Communities to Changing
4930 Climate Conditions During Summer Cyanobacterial Blooms in the Baltic Sea, *Front. Microbiol.*, 9,
4931 <https://doi.org/10.3389/fmicb.2018.01562>, 2018.

Formatiert: Schriftart: (Standard) Times New Roman, 10 Pt.,
Englisch (USA)

Formatiert: Abstand Nach: 0 Pt.

Formatiert: Rechtschreibung und Grammatik nicht prüfen

4932 Bierstedt, S. E., Hünicke, B., and Zorita, E.: Variability of wind direction statistics of mean and extreme wind
4933 events over the Baltic Sea region, *Tellus A*, 67, 29073, <https://doi.org/10.3402/tellusa.v67.29073>, 2015.

4934 Bingham, R. J., and Hughes, C. W.: Local diagnostics to estimate density-induced sea level variations over
4935 topography and along coastlines, *J. Geophys. Res-Oceans*, 117, <https://doi.org/10.1029/2011jc007276>, 2012.

4936 Biskaborn, B. K., Smith, S. L., Noetzli, J., Matthes, H., Vieira, G., Streletskiy, D. A., Schoeneich, P.,
4937 Romanovsky, V. E., Lewkowicz, A. G., Abramov, A., Allard, M., Boike, J., Cable, W. L., Christiansen, H. H.,
4938 Delaloye, R., Diekmann, B., Drozdov, D., Etzelmüller, B., Grosse, G., Guglielmin, M., Ingeman-Nielsen, T.,
4939 Isaksen, K., Ishikawa, M., Johansson, M., Johannsson, H., Joo, A., Kaverin, D., Kholodov, A., Konstantinov, P.,
4940 Kröger, T., Lambiel, C., Lanckman, J.-P., Luo, D., Malkova, G., Meiklejohn, I., Moskalenko, N., Oliva, M.,
4941 Phillips, M., Ramos, M., Sannel, A. B. K., Sergeev, D., Seybold, C., Skryabin, P., Vasiliev, A., Wu, Q.,
4942 Yoshikawa, K., Zheleznyak, M., and Lantuit, H.: Permafrost is warming at a global scale, *Nat. Commun.*, 10,
4943 264, <https://doi.org/10.1038/s41467-018-08240-4>, 2019.

4944 Björkqvist, J.-V., Lukas, I., Alari, V., van Vledder, G. P., Hulst, S., Pettersson, H., Behrens, A., and Männik, A.:
4945 Comparing a 41-year model hindcast with decades of wave measurements from the Baltic Sea, *Ocean Eng.*, 152,
4946 57-71, <https://doi.org/10.1016/j.oceaneng.2018.01.048>, 2018.

4947 Björkqvist, J. V., Rikka, S., Alari, V., Männik, A., Tuomi, L., and Pettersson, H.: Wave height return periods
4948 from combined measurement–model data: A Baltic Sea case study, *Nat. Hazard. Earth Syst. Sci. Discuss.*, 2020,
4949 1-25, <https://doi.org/10.5194/nhess-2020-190>, 2020.

4950 Blackport, R., and Screen, J. A.: Insignificant effect of Arctic amplification on the amplitude of midlatitude
4951 atmospheric waves, *Sci. Adv.*, 6, eaay2880, <https://doi.org/10.1126/sciadv.aay2880>, 2020.

4952 Blenckner, T., Österblom, H., Larsson, P., Andersson, A., and Elmgren, R.: Baltic Sea ecosystem-based
4953 management under climate change: Synthesis and future challenges, *Ambio* 44, 507-515,
4954 <https://doi.org/10.1007/s13280-015-0661-9>, 2015.

4955 Blindow, I., Dahlke, S., Dewart, A., Flügge, S., Hendreschke, M., Kerkow, A., and Meyer, J.: Long-term and
4956 interannual changes of submerged macrophytes and their associated diaspore reservoir in a shallow southern
4957 Baltic Sea bay: influence of eutrophication and climate, *Hydrobiologia*, 778, 121-136,
4958 <https://doi.org/10.1007/s10750-016-2655-4>, 2016.

4959 Blöschl, G., Hall, J., Parajka, J., Perdigão, R. A. P., Merz, B., Arheimer, B., Aronica, G. T., Bilibashi, A.,
4960 Bonacci, O., Borga, M., Canjevac, I., Castellarin, A., Chirico, G. B., Claps, P., Fiala, K., Frolova, N.,
4961 Gorbachova, L., Gül, A., Hannaford, J., Harrigan, S., Kireeva, M., Kiss, A., Kjeldsen, T. R., Kohnová, S.,
4962 Koskela, J. J., Ledvinka, O., Macdonald, N., Mavrova-Guirguinova, M., Mediero, L., Merz, R., Molnar, P.,
4963 Montanari, A., Murphy, C., Osuch, M., Ovcharuk, V., Radevski, I., Rogger, M., Salinas, J. L., Sauquet, E., Šraj,
4964 M., Szolgay, J., Viglione, A., Volpi, E., Wilson, D., Zaimi, K., and Živković, N.: Changing climate shifts timing
4965 of European floods, *Science*, 357, 588, <https://doi.org/10.1126/science.aan2506>, 2017.

4966 Blunden, J., and Arndt, D. S.: State of the Climate in 2014, *Bull. Amer. Meteor. Soc.*, 96, ES1-ES32,
4967 <https://doi.org/10.1175/2015BAMSStateoftheClimate.1>, 2015.

4968 Bobsien, I. C., Hukriede, W., Schlamkow, C., Friedland, R., Dreier, N., Schubert, P. R., Karez, R., and Reusch,
4969 T. B. H.: Modeling eelgrass spatial response to nutrient abatement measures in a changing climate, *Ambio*, 50,
4970 400-412, <https://doi.org/10.1007/s13280-020-01364-2>, 2021.

4971 Boé, J., Somot, S., Corre, L., and Nabat, P.: Large discrepancies in summer climate change over Europe as
4972 projected by global and regional climate models: causes and consequences, *Clim. Dyn.*, 54, 2981-3002,
4973 <https://doi.org/10.1007/s00382-020-05153-1>, 2020a.

4974 Boé, J., Terray, L., Moine, M.-P., Valcke, S., Bellucci, A., Drijfhout, S., Haarsma, R., Lohmann, K., Putrasahan,
4975 D. A., Roberts, C., Roberts, M., Scoccimarro, E., Seddon, J., Senan, R., and Wyser, K.: Past long-term summer
4976 warming over western Europe in new generation climate models: role of large-scale atmospheric circulation,
4977 *Environ. Res. Lett.*, 15, 084038, <https://doi.org/10.1088/1748-9326/ab8a89>, 2020b.

4978 Boers, N.: [Observation-based early-warning signals for a collapse of the Atlantic Meridional Overturning](https://doi.org/10.1038/s41558-021-01097-4)
4979 [Circulation](https://doi.org/10.1038/s41558-021-01097-4), *Nat. Clim. Change*, 11, 680-688, <https://doi.org/10.1038/s41558-021-01097-4>, 2021.

Formatiert: Schriftart: (Standard) Times New Roman, 10 Pt.,
Englisch (USA)

Formatiert: Englisch (USA)

4981
4982 Boland, E. J. D., Bracegirdle, T. J., and Shuckburgh, E. F.: Assessment of sea ice-atmosphere links in CMIP5
4983 models, *Clim. Dyn.*, 49, 683-702, <https://doi.org/10.1007/s00382-016-3367-1>, 2017.

4984 Bonaduce, A., Staneva, J., Behrens, A., Bidlot, J.-R., and Wilcke, R. A. I.: Wave Climate Change in the North
4985 Sea and Baltic Sea, *J. Mar. Sci. Eng.*, 7, 166, <https://doi.org/10.3390/jmse7060166>, 2019.

4986 Bonan, G. B.: Forests, Climate, and Public Policy: A 500-Year Interdisciplinary Odyssey, *Annu. Rev. Ecol.*
4987 *Evol. S.*, 47, 97-121, <https://doi.org/10.1146/annurev-ecolsys-121415-032359>, 2016.

4988 Bonsdorff, E.: Eutrophication: Early warning signals, ecosystem-level and societal responses, and ways forward,
4989 *Ambio*, <https://doi.org/10.1007/s13280-020-01432-7>, 2021.

4990 [Borchert, L. F., Menary, M. B., Swingedouw, D., Sgubin, G., Hermanson, L., and Mignot, J.: Improved Decadal
4991 Predictions of North Atlantic Subpolar Gyre SST in CMIP6, 48, e2020GL091307,
4992 <https://doi.org/10.1029/2020GL091307>, 2021.](https://doi.org/10.1029/2020GL091307)

4993 Börgel, F., Frauen, C., Neumann, T., Schimanke, S., and Meier, H. E. M.: Impact of the Atlantic Multidecadal
4994 Oscillation on Baltic Sea Variability, *Geophys. Res. Lett.*, 45, 9880-9888,
4995 <https://doi.org/10.1029/2018GL078943>, 2018.

4996 Börgel, F., Frauen, C., Neumann, T., and Meier, H. E. M.: The Atlantic Multidecadal Oscillation controls the
4997 impact of the North Atlantic Oscillation on North European climate, *Environ. Res. Lett.*,
4998 <https://doi.org/10.1088/1748-9326/aba925>, 2020.

4999 [Börgel, F., Meier, H. E. M., Gröger, M., Rhein, M., Dutheil, C., and Kaiser, J. M.: Atlantic Multidecadal
5000 Variability and the implications for North European Climate using CMIP6 control simulations, *Environ. Res.*
5001 *Lett.*, under review, 2022.](https://doi.org/10.1029/2020GL091307)

5002 Bouwman, L., Goldewijk, K. K., Van Der Hoek, K. W., Beusen, A. H. W., Van Vuuren, D. P., Willems, J.,
5003 Rufino, M. C., and Stehfest, E.: Exploring global changes in nitrogen and phosphorus cycles in agriculture
5004 induced by livestock production over the 1900–2050 period, *P. Natl. Acad. Sci. USA*, 110, 20882,
5005 <https://doi.org/10.1073/pnas.1012878108>, 2013.

5006 Brandt, J., Silver, J. D., Christensen, J. H., Andersen, M. S., Bønløkke, J. H., Sigsgaard, T., Geels, C., Gross, A.,
5007 Hansen, A. B., Hansen, K. M., Hedegaard, G. B., Kaas, E., and Frohn, L. M.: Assessment of past, present and
5008 future health-cost externalities of air pollution in Europe and the contribution from international ship traffic
5009 using the EVA model system, *Atmos. Chem. Phys.*, 13, 7747-7764, <https://doi.org/10.5194/acp-13-7747-2013>,
5010 2013.

5011 Broman, B., Hammarklint, T., Rannat, K., Soomere, T., and Valdmann, A.: Trends and extremes of wave fields
5012 in the north-eastern part of the Baltic Proper, *Oceanologia*, 48, 2006.

5013 Brown, J., Ferrians Jr, O. J., Heginbottom, J. A., and Melnikov, E. S.: Circum-Arctic map of permafrost and
5014 ground-ice conditions, U.S. Geological Survey, U.S.A., Report 45, <https://doi.org/10.3133/cp45>, 1997.

5015 Brugger, K. A., and Pankratz, L.: Changes in the geometry and volume of rabots glaciär, sweden, 2003–2011:
5016 recent accelerated volume loss linked to more negative summer balances, *Geogr. Ann. A*, 97, 265-278,
5017 <https://doi.org/10.1111/geoa.12062>, 2015.

5018 Brunner, L., Pendergrass, A. G., Lehner, F., Merrifield, A. L., Lorenz, R., and Knutti, R.: Reduced global
5019 warming from CMIP6 projections when weighting models by performance and independence, *Earth Syst.*
5020 *Dynam. Discuss.*, 2020, 1-23, <https://doi.org/10.5194/esd-2020-23>, 2020.

5021 Brutemark, A., Engström-Öst, J., Vehmaa, A., and Gorokhova, E.: Growth, toxicity and oxidative stress of a
5022 cultured cyanobacterium (*Dolichospermum* sp.) under different CO₂/pH and temperature conditions, *Phycol.*
5023 *Res.*, 63, 56-63, <https://doi.org/10.1111/pre.12075>, 2015.

5024 [Buckley, M. W., and Marshall, J.: Observations, inferences, and mechanisms of the Atlantic Meridional
5025 Overturning Circulation: A review, *Reviews of Geophysics* 54.1, 5–63, 2016.](https://doi.org/10.1111/pre.12075)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Rechtschreibung und Grammatik nicht prüfen

5026 Bülow, K., Dietrich, C., Elizalde, A., Gröger, M., Heinrich, H., Hüttl-Kabos, S., Klein, B., Mayer, B., Meier, H.
5027 E. M., and Mikolajewicz, U.: Comparison of three regional coupled ocean atmosphere models for the North Sea
5028 under today's and future climate conditions (KLIWAS Schriftenreihe; KLIWAS-27/2014), Bundesanstalt für
5029 Gewässerkunde, 2014.

5030 Burchard, H., Bolding, K., Feistel, R., Gräwe, U., Klingbeil, K., MacCready, P., Mohrholz, V., Umlauf, L., and
5031 van der Lee, E. M.: The Knudsen theorem and the Total Exchange Flow analysis framework applied to the Baltic
5032 Sea, *Prog. Oceanogr.*, 165, 268-286, <https://doi.org/10.1016/j.pocean.2018.04.004>, 2018.

5033 [Byrne, M. P., and O'Gorman, P. A.: Trends in continental temperature and humidity directly linked to ocean](#)
5034 [warming, P. Natl. Acad. Sci. USA, 115, 4863, <https://doi.org/10.1073/pnas.1722312115>, 2018.](#)

5035 Caballero-Alfonso, A. M., Carstensen, J., and Conley, D. J.: Biogeochemical and environmental drivers of
5036 coastal hypoxia, *J. Mar. Syst.*, 141, 190-199, <https://doi.org/10.1016/j.jmarsys.2014.04.008>, 2015.

5037 Cardell, M. F., Amengual, A., Romero, R., and Ramis, C.: Future extremes of temperature and precipitation in
5038 Europe derived from a combination of dynamical and statistical approaches, *Int. J. Climatol.*, 40, 4800-4827,
5039 <https://doi.org/10.1002/joc.6490>, 2020.

5040 Carstensen, J., Andersen, J. H., Gustafsson, B. G., and Conley, D. J.: Deoxygenation of the Baltic Sea during the
5041 last century, *P. Natl. Acad. Sci. USA*, 111, 5628-5633, <https://doi.org/10.1073/pnas.1323156111>, 2014a.

5042 Carstensen, J., Conley, D. J., Bonsdorff, E., Gustafsson, B. G., Hietanen, S., Janas, U., Jilbert, T., Maximov, A.,
5043 Norkko, A., Norkko, J., and others: Hypoxia in the Baltic Sea: Biogeochemical cycles, benthic fauna, and
5044 management, *Ambio*, 43, 26-36, <https://doi.org/10.1007/s13280-013-0474-7>, 2014b.

5045 Carstensen, J., Chierici, M., Gustafsson, B. G., and Gustafsson, E.: Long-Term and Seasonal Trends in Estuarine
5046 and Coastal Carbonate Systems, *Global Biogeochem. Cy.*, 32, 497-513, <https://doi.org/10.1002/2017gb005781>,
5047 2018.

5048 Carstensen, J., and Conley, D. J.: Baltic Sea Hypoxia Takes Many Shapes and Sizes, *Limnol. Oceanogr. Bull.*,
5049 28, 125-129, <https://doi.org/10.1002/lob.10350>, 2019.

5050 Carstensen, J., and Duarte, C. M.: Drivers of pH Variability in Coastal Ecosystems, *Environ. Sci. Technol.*, 53,
5051 4020-4029, <https://doi.org/10.1021/acs.est.8b03655>, 2019.

5052 Carstensen, J., Conley, D. J., Almroth-Rosell, E., Asmala, E., Bonsdorff, E., Fleming-Lehtinen, V., Gustafsson,
5053 B. G., Gustafsson, C., Heiskanen, A.-S., Janas, U., Norkko, A., Slomp, C., Villnäs, A., Voss, M., and Zilius, M.:
5054 Factors regulating the coastal nutrient filter in the Baltic Sea, *Ambio*, 49, 1194-1210,
5055 <https://doi.org/10.1007/s13280-019-01282-y>, 2020.

5056 Casanueva, A., Rodríguez-Puebla, C., Frías, M. D., and González-Reviriego, N.: Variability of extreme
5057 precipitation over Europe and its relationships with teleconnection patterns, *Hydrol. Earth Syst. Sci.*, 18, 709 -
5058 725, <https://doi.org/10.5194/hess-18-709-2014>, 2014.

5059 Casini, M., Lövgren, J., Hjelm, J., Cardinale, M., Molinero, J.-C., and Kornilovs, G.: Multi-level trophic
5060 cascades in a heavily exploited open marine ecosystem, *P. Roy. Soc. B-Biol. Sci.*, 275, 1793-1801,
5061 <https://doi.org/10.1098/rspb.2007.1752> 2008.

5062 Casini, M., Kornilovs, G., Cardinale, M., Möllmann, C., Grygiel, W., Jonsson, P., Raid, T., Flinkman, J., and
5063 Feldman, V.: Spatial and temporal density dependence regulates the condition of central Baltic Sea clupeids:
5064 compelling evidence using an extensive international acoustic survey, *Popul. Ecol.*, 53, 511-523,
5065 <https://doi.org/10.1007/s10144-011-0269-2>, 2011.

5066 Casini, M., Käll, F., Hansson, M., Plikshs, M., Baranova, T., Karlsson, O., Lundström, K., Neuenfeldt, S.,
5067 Gårdmark, A., and Hjelm, J.: Hypoxic areas, density-dependence and food limitation drive the body condition of
5068 a heavily exploited marine fish predator, *Roy. Soc. Open Sci.*, 3, 160416, <https://doi.org/10.1098/rsos.160416>,
5069 2016.

5070 Cattiaux, J., and Cassou, C.: Opposite CMIP3/CMIP5 trends in the wintertime Northern Annular Mode
5071 explained by combined local sea ice and remote tropical influences, *Geophys. Res. Lett.*, 40, 3682-3687,
5072 <https://doi.org/10.1002/grl.50643>, 2013.

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Feldfunktion geändert

Formatiert: Englisch (USA)

Formatiert: Rechtschreibung und Grammatik nicht prüfen

5073 Cattiaux, J., Douville, H., and Peings, Y.: European temperatures in CMIP5: origins of present-day biases and
5074 future uncertainties, *Clim. Dyn.*, 41, 2889-2907, <https://doi.org/10.1007/s00382-013-1731-y>, 2013.

5075 Ceppi, P., and Hartmann, D. L.: Connections Between Clouds, Radiation, and Midlatitude Dynamics: a Review,
5076 *Curr. Clim. Chang. Rep.*, 1, 94-102, <https://doi.org/10.1007/s40641-015-0010-x>, 2015.

5077 Chadburn, S. E., Burke, E. J., Cox, P. M., Friedlingstein, P., Hugelius, G., and Westermann, S.: An observation -
5078 based constraint on permafrost loss as a function of global warming, *Nat. Clim. Change*, 7, 340-344,
5079 <https://doi.org/10.1038/nclimate3262>, 2017.

5080 Chang, E. K. M., Guo, Y., and Xia, X.: CMIP5 multimodel ensemble projection of storm track change under
5081 global warming, *J. Geophys. Res.-Atmos.*, 117, <https://doi.org/10.1029/2012JD018578>, 2012.

5082 Chang, E. K. M., Ma, C.-G., Zheng, C., and Yau, A. M. W.: Observed and projected decrease in Northern
5083 Hemisphere extratropical cyclone activity in summer and its impacts on maximum temperature, *Geophys. Res.
5084 Lett.*, 43, 2200-2208, <https://doi.org/10.1002/2016GL068172>, 2016.

5085 Chang, E. K. M., and Yau, A. M. W.: Northern Hemisphere winter storm track trends since 1959 derived from
5086 multiple reanalysis datasets, *Clim. Dyn.*, 47, 1435-1454, <https://doi.org/10.1007/s00382-015-2911-8>, 2016.

5087 Chen, D., Zhang, P., Seftigen, K., Ou, T., Giese, M., and Barthel, R.: Hydroclimate changes over Sweden in the
5088 twentieth and twenty-first centuries: a millennium perspective, *Geogr. Ann. A*, 1-29,
5089 <https://doi.org/10.1080/04353676.2020.1841410>, 2020.

5090 Christensen, J. H., Larsen, M. A. D., Christensen, O. B., Drews, M., and Stendel, M.: Robustness of European
5091 climate projections from dynamical downscaling, *Clim. Dyn.*, 53, 4857-4869, <https://doi.org/10.1007/s00382-019-04831-z>, 2019.

5093 Christensen, O. B., and Kjellström, E.: Projections for Temperature, Precipitation, Wind, and Snow in the Baltic
5094 Sea Region until 2100, *Oxford Research Encyclopedia of Climate Science*, Oxford University Press,
5095 <https://doi.org/10.1093/acrefore/9780190228620.013.695>, 2018.

5096 Christensen, O. B., Kjellström, E., Dieterich, C., Gröger, M., and Meier, H. E. M.: Atmospheric regional climate
5097 projections for the Baltic Sea Region until 2100, *Earth Syst. Dynam.*, 2021.

5098 Chust, G., Allen, J. I., Bopp, L., Schrum, C., Holt, J., Tsiaras, K., Zavatarelli, M., Chifflet, M., Cannaby, H.,
5099 Dadou, I., Daewel, U., Wakelin, S. L., Machu, E., Pushpadas, D., Butenschon, M., Artioli, Y., Petihakis, G.,
5100 Smith, C., Garçon, V., Goubanova, K., Le Vu, B., Fach, B. A., Salihoglu, B., Clementi, E., and Irigoien, X.:
5101 Biomass changes and trophic amplification of plankton in a warmer ocean, *Global Change Biol.*, 20, 2124-2139,
5102 <https://doi.org/10.1111/gcb.12562>, 2014.

5103 Ciasto, L. M., Li, C., Wettstein, J. J., and Kvamstø, N. G.: North Atlantic Storm-Track Sensitivity to Projected
5104 Sea Surface Temperature: Local versus Remote Influences, *J. Climate*, 29, 6973-6991,
5105 <https://doi.org/10.1175/jcli-d-15-0860.1>, 2016.

5106 Claremar, B., Haglund, K., and Rutgersson, A.: Ship emissions and the use of current air cleaning technology:
5107 contributions to air pollution and acidification in the Baltic Sea, *Earth Syst. Dynam.*, 8, 901-919,
5108 <https://doi.org/10.5194/esd-8-901-2017>, 2017.

5109 Clausen, K. K., Stjernholm, M., and Clausen, P.: Grazing management can counteract the impacts of climate
5110 change-induced sea level rise on salt marsh-dependent waterbirds, *J. Appl. Ecol.*, 50, 528-537,
5111 <https://doi.org/10.1111/1365-2664.12043>, 2013.

5112 Clausen, K. K., and Clausen, P.: Forecasting future drowning of coastal waterbird habitats reveals a major
5113 conservation concern, *Biol. Conserv.*, 171, 177-185, <https://doi.org/10.1016/j.biocon.2014.01.033>, 2014.

5114 Colette, A., Bessagnet, B., Vautard, R., Szopa, S., Rao, S., Schucht, S., Klimont, Z., Menut, L., Clain, G.,
5115 Meleux, F., CURCI, G., and Rouil, L.: European atmosphere in 2050, a regional air quality and climate
5116 perspective under CMIP5 scenarios, *Atmos. Chem. Phys.*, 13, 7451-7471, [https://doi.org/10.5194/acp-13-7451-](https://doi.org/10.5194/acp-13-7451-2013)
5117 2013, 2013.

5118 Colette, A., Andersson, C., Baklanov, A., Bessagnet, B., Brandt, J., Christensen, J. H., Doherty, R., Engardt, M.,
5119 Geels, C., Giannakopoulos, C., Hedegaard, G. B., Katragkou, E., Langner, J., Lei, H., Manders, A., Melas, D.,

5120 Meleux, F., Rouïl, L., Sofiev, M., Soares, J., Stevenson, D. S., Tombrou-Tzella, M., Varotsos, K. V., and Young,
5121 P.: Is the ozone climate penalty robust in Europe?, *Environ. Res. Lett.*, 10, 084015, [https://doi.org/10.1088/1748-](https://doi.org/10.1088/1748-9326/10/8/084015)
5122 9326/10/8/084015, 2015.

5123 Colette, A., and Rouïl, L.: Air Quality Trends in Europe: 2000-2017, European Environment Agency, Kjeller,
5124 Norway, Eionet Report - ETC/ATNI 2019/16, 36pp, 2020.

5125 Collins, M., Knutti, R., Arblaster, J., Dufresne, J.-L., Fichet, T., Friedlingstein, P., Gao, X., Gutowski, W. J.,
5126 Johns, T., Krinner, G., and others: Long-term climate change: projections, commitments and irreversibility, In:
5127 IPCC Chapter 12, Climate Change 2013: The Physical Science Basis. IPCC Working Group I Contribution to
5128 IPCC AR5; Cambridge University Press, Cambridge, UK and New York, USA, 1029-1136, 2013.

5129 Conley, D. J., Humborg, C., Rahm, L., Savchuk, O. P., and Wulff, F.: Hypoxia in the Baltic Sea and basin-scale
5130 changes in phosphorus biogeochemistry, *Environ. Sci. Technol.*, 36, 5315-5320,
5131 <https://doi.org/10.1021/es025763w>, 2002.

5132 Conley, D. J., Carstensen, J., Aigars, J., Axe, P., Bonsdorff, E., Eremina, T., Haahti, B. -M., Humborg, C.,
5133 Jonsson, P., Kotta, J., Lännegren, C., Larsson, U., Maximov, A., Medina, M. R., Lysiak-Pastuszek, E.,
5134 Remeikaitė-Nikienė, N., Walve, J., Wilhelms, S., and Zillén, L.: Hypoxia Is Increasing in the Coastal Zone of
5135 the Baltic Sea, *Environ. Sci. Technol.*, 45, 6777-6783, <https://doi.org/10.1021/es201212r>, 2011.

5136 Cornes, R. C., van der Schrier, G., and Squintu, A. A.: A reappraisal of the thermal growing season length across
5137 Europe, *Int. J. Climatol.*, 39, 1787-1795, <https://doi.org/10.1002/joc.5913>, 2019.

5138 Coppola, E., Nogherotto, R., Ciarlo', J. M., Giorgi, F., van Meijgaard, E., Kadygrov, N., Iles, C., Corre, L.,
5139 Sandstad, M., Somot, S., Nabat, P., Vautard, R., Levavasseur, G., Schwingshackl, C., Sillmann, J., Kjellström,
5140 E., Nikulin, G., Aalbers, E., Lenderink, G., Christensen, O. B., Boberg, F., Sørland, S. L., Demory, M.-E.,
5141 Bülow, K., Teichmann, C., Warrach-Sagi, K., and Wulfmeyer, V.: Assessment of the European Climate
5142 Projections as Simulated by the Large EURO-CORDEX Regional and Global Climate Model Ensemble, J.
5143 Geophys. Res-Atmos., 126, e2019JD032356, <https://doi.org/10.1029/2019JD032356>, 2021.

5144 Coumou, D., Lehmann, J., and Beckmann, J.: The weakening summer circulation in the Northern Hemisphere
5145 mid-latitudes, *Science*, 348, 324, <https://doi.org/10.1126/science.1261768>, 2015.

5146 Cyberski, J., Wróblewski, A., and Stewart, J.: Riverine water inflows and the Baltic Sea water volume 1901 -
5147 1990, *Hydrol. Earth Syst. Sci.*, 4, 1-11, <https://doi.org/10.5194/hess-4-1-2000>, 2000.

5148 Defeo, O., McLachlan, A., Schoeman, D. S., Schlacher, T. A., Dugan, J., Jones, A., Lastra, M., and Scapini, F.:
5149 Threats to sandy beach ecosystems: A review, *Estuar. Coast. Shelf Sci.*, 81, 1-12,
5150 <https://doi.org/10.1016/j.ecss.2008.09.022>, 2009.

5151 Deng, J., Zhang, W., Harff, J., Schneider, R., Dudzinska-Nowak, J., Terefenko, P., Giza, A., and Furmanczyk,
5152 K.: A numerical approach for approximating the historical morphology of wave-dominated coasts—A case study
5153 of the Pomeranian Bight, southern Baltic Sea, *Geomorphology*, 204, 425-443,
5154 <https://doi.org/10.1016/j.geomorph.2013.08.023>, 2014.

5155 Deng, J., Harff, J., Schimanke, S., and Meier, H. E. M.: A method for assessing the coastline recession due to the
5156 sea level rise by assuming stationary wind-wave climate, *Oceanol. Hydrobiol. Stud.*, Vol. 44, No. 3, 362-380,
5157 <https://doi.org/10.1515/ohs-2015-0035>, 2015.

5158 Deng, J., Wu, J., Zhang, W., Dudzinska-Nowak, J., and Harff, J.: Characterising the relaxation distance of
5159 nearshore submarine morphology: A southern Baltic Sea case study, *Geomorphology*, 327, 365-376,
5160 <https://doi.org/10.1016/j.geomorph.2018.11.018>, 2019.

5161 Deser, C.: On the teleconnectivity of the "Arctic Oscillation". *Geophysical Research Letters*, 27(6),
5162 779-782, <https://doi.org/10.1029/1999GL010945>, 2000.

5163 Deser, C., Hurrell, J. W., and Phillips, A. S.: The role of the North Atlantic Oscillation in European climate
5164 projections, *Clim. Dyn.*, 49, 3141-3157, <https://doi.org/10.1007/s00382-016-3502-z>, 2017.

5165 Dethloff, K., Rinke, A., Benkel, A., Koltzow, M., Sokolova, E., Kumar Saha, S., Handorf, D., Dorn, W., Rockel,
5166 B., von Storch, H., Haugen, J. E., Røed, L. P., Roeckner, E., Christensen, J. H., and Stendel, M.: A dynamical

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Feldfunktion geändert

Formatiert: Englisch (USA)

Formatiert: Rechtschreibung und Grammatik nicht prüfen

Formatiert: Englisch (USA)

Formatiert: Schriftart: Nicht Kursiv, Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Schriftart: Nicht Kursiv, Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Schriftart: (Standard) +Textkörper (Calibri), 11 Pt.

link between the Arctic and the global climate system, *Geophys. Res. Lett.*, 33,
<https://doi.org/10.1029/2005GL025245>, 2006.

Devictor, V., van Swaay, C., Brereton, T., Brotons, L., Chamberlain, D., Heliölä, J., Herrando, S., Julliard, R., Kuussaari, M., Lindström, Å., Reif, J., Roy, D. B., Schweiger, O., Settele, J., Stefanescu, C., Van Strien, A., Van Turnhout, C., Vermouzek, Z., WallisDeVries, M., Wynhoff, I., and Jiguet, F.: Differences in the climatic debts of birds and butterflies at a continental scale, *Nat. Clim. Change*, 2, 121-124, <https://doi.org/10.1038/nclimate1347>, 2012.

Dieterich, C., Schimanke, S., Wang, S., Väli, G., Liu, Y., Hordoir, R., Höglund, A., and Meier, H. E. M.: Evaluation of the SMHI coupled atmosphere-ice-ocean model RCA4-NEMO, SMHI, Norrköping, Sweden, 80, 2013.

Dieterich, C., Wang, S., Schimanke, S., Gröger, M., Klein, B., Hordoir, R., Samuelsson, P., Liu, Y., Axell, L., and Höglund, A.: Surface heat budget over the North Sea in climate change simulations, *Atmosphere*, 10, 272, <https://doi.org/10.3390/atmos10050272>, 2019.

Dippner, J. W., Vuorinen, I., Daunys, D., Flinkman, J., Halkka, A., Köster, F. W., Lehikoinen, E., MacKenzie, B. R., Möllmann, C., Møhlenberg, F., Olenin, S., Schiedek, D., Skov, H., and Wasmund, N.: Climate-related Marine Ecosystem Change, in: *Assessment of Climate Change for the Baltic Sea Basin*, Springer Berlin Heidelberg, Berlin, Heidelberg, 309-377, https://doi.org/10.1007/978-3-540-72786-6_5, 2008.

Dippner, J. W., Fründt, B., and Hammer, C.: Lake or Sea? The Unknown Future of Central Baltic Sea Herring, *Front. Ecol. Evol.*, 7, <https://doi.org/10.3389/fevo.2019.00143>, 2019.

Doherty, R. M., Heal, M. R., and O'Connor, F. M.: Climate change impacts on human health over Europe through its effect on air quality, *Environ. Health-Glob.*, 16, 118, <https://doi.org/10.1186/s12940-017-0325-2>, 2017.

Doney, S. C., Fabry, V. J., Feely, R. A., and Kleypas, J. A.: Ocean acidification: the other CO₂ problem. *Annual review of marine science*, 1, 169-192, <https://doi.org/10.1146/annurev.marine.010908.163834>, 2009.

Dong, B., Sutton, R. T., and Shaffrey, L.: Understanding the rapid summer warming and changes in temperature extremes since the mid-1990s over Western Europe, *Clim. Dyn.*, 48, 1537-1554, <https://doi.org/10.1007/s00382-016-3158-8>, 2017.

Donnelly, C., Yang, W., and Dahné, J.: River discharge to the Baltic Sea in a future climate, *Clim. Change*, 122, 157-170, <https://doi.org/10.1007/s10584-013-0941-y>, 2014.

Donnelly, C., Greuell, W., Andersson, J., Gerten, D., Pisacane, G., Roudier, P., and Ludwig, F.: Impacts of climate change on European hydrology at 1.5, 2 and 3 degrees mean global warming above preindustrial level, *Clim. Change*, 143, 13-26, <https://doi.org/10.1007/s10584-017-1971-7>, 2017.

Dreier, N., Nehlsen, E., Fröhle, P., Rechid, D., Bouwer, L. M., and Pfeifer, S.: Future Changes in Wave Conditions at the German Baltic Sea Coast Based on a Hybrid Approach Using an Ensemble of Regional Climate Change Projections, *Water*, 13, <https://doi.org/10.3390/w13020167>, 2021.

Du, J., Shen, J., Park, K., Wang, Y. P., and Yu, X.: Worsened physical condition due to climate change contributes to the increasing hypoxia in Chesapeake Bay, *Sci. Total Environ.*, 630, 707-717, <https://doi.org/10.1016/j.scitotenv.2018.02.265>, 2018.

Duarte, C. M., Hendriks, I. E., Moore, T. S., Olsen, Y. S., Steckbauer, A., Ramajo, L., Carstensen, J., Trotter, J. A., and McCulloch, M.: Is Ocean Acidification an Open-Ocean Syndrome? Understanding Anthropogenic Impacts on Seawater pH, *Estuar. Coast.*, 36, 221-236, <https://doi.org/10.1007/s12237-013-9594-3>, 2013.

Ducklow, H. W., Morán, X. A. G., and Murray, A. E.: Bacteria in the Greenhouse: Marine Microbes and Climate Change, in: *Environ. Microbiol.*, edited by: Mitchell, R., and Gu, J. D., Wiley-Blackwell, Hoboken, N.J., USA, 1-31, <https://doi.org/10.1002/9780470495117.ch1>, 2009.

Dudzińska-Nowak, J.: Morphodynamic Processes of the Swina Gate Coastal Zone Development (Southern Baltic Sea), in: *Coastline Changes of the Baltic Sea from South to East: Past and Future Projection*, edited by:

Formatiert: Schriftart: Nicht Kursiv, Englisch (USA)

Formatiert: Schriftart: +Textkörper (Calibri)

5213 Harff, J., Furmańczyk, K., and von Storch, H., Springer International Publishing, Cham, 219-255,
5214 https://doi.org/10.1007/978-3-319-49894-2_11, 2017.

5215 Dutheil, C., Meier, H.E.M., Gröger, M., and Börgel, F.: Understanding past and future sea surface temperature
5216 trends in the Baltic Sea, *Climate Dynamics*, <https://doi.org/10.1007/s00382-021-06084-1>, 2021.

5217 Dyrndal, A. V., Saloranta, T., Skaugen, T., and Stranden, H. B.: Changes in snow depth in Norway during the
5218 period 1961–2010, *Hydrol. Res.*, 44, 169-179, <https://doi.org/10.2166/nh.2012.064>, 2012.

5219 Easterling, D. R., Kunkel, K. E., Wehner, M. F., and Sun, L.: Detection and attribution of climate extremes in
5220 the observed record, *Weather. Clim. Extremes*, 11, 17-27, <https://doi.org/10.1016/j.wace.2016.01.001>, 2016.

5221 Eero, M., Andersson, H. C., Almroth-Rosell, E., and MacKenzie, B. R.: Has eutrophication promoted forage fish
5222 production in the Baltic Sea?, *Ambio*, 45, 649-660, <https://doi.org/10.1007/s13280-016-0788-3>, 2016.

5223 Efremova, T., Palshin, N., and Zdorovenkov, R.: Long-term characteristics of ice phenology in Karelian lakes,
5224 *Est. J. Earth Sci.*, 62, 33-41, <https://doi.org/10.3176/earth.2013.04>, 2013.

5225 Eggert, A., and Schneider, B.: A nitrogen source in spring in the surface mixed-layer of the Baltic Sea: evidence
5226 from total nitrogen and total phosphorus data, *Journal of Marine Systems*, 148, 39-47,
5227 <https://doi.org/10.1016/j.jmarsys.2015.01.005>, 2015.

5228 Ehrnsten, E., Bauer, B., and Gustafsson, B. G.: Combined Effects of Environmental Drivers on Marine Trophic
5229 Groups – A Systematic Model Comparison, *Front. Mar. Sci.*, 6, <https://doi.org/10.3389/fmars.2019.00492>, 2019.

5230 Ehrnsten, E., Norkko, A., Müller-Karulis, B., Gustafsson, E., and Gustafsson, B. G.: The meagre future of
5231 benthic fauna in a coastal sea—Benthic responses to recovery from eutrophication in a changing climate, *Global*
5232 *Change Biol.*, 26, 2235-2250, <https://doi.org/10.1111/gcb.15014>, 2020.

5233 Eichner, M., Rost, B., and Kranz, S. A.: Diversity of ocean acidification effects on marine N₂ fixers, *J. Exp.*
5234 *Mar. Biol. Ecol.*, 457, 199-207, <https://doi.org/10.1016/j.jembe.2014.04.015>, 2014.

5235 Eilola, K., Rosell, E. A., Dieterich, C., Fransner, F., Höglund, A., and Meier, H. E. M.: Modeling Nutrient
5236 Transports and Exchanges of Nutrients Between Shallow Regions and the Open Baltic Sea in Present and Future
5237 Climate, *Ambio*, 41, 586-599, <https://doi.org/10.1007/s13280-012-0322-1>, 2012.

5238 Eilola, K., Mårtensson, S., and Meier, H. E. M.: Modeling the impact of reduced sea ice cover in future climate
5239 on the Baltic Sea biogeochemistry, *Geophys. Res. Lett.*, 40, 149-154, <https://doi.org/10.1029/2012GL054375>,
5240 2013.

5241 Eilola, K., Almroth-Rosell, E., and Meier, H. E. M.: Impact of saltwater inflows on phosphorus cycling and
5242 eutrophication in the Baltic Sea: a 3D model study, *Tellus A*, 66, 23985,
5243 <https://doi.org/10.3402/tellusa.v66.23985>, 2014.

5244 Emeis, K.-C., van Beusekom, J., Callies, U., Ebinghaus, R., Kannen, A., Kraus, G., Kröncke, I., Lenhart, H.,
5245 Lorkowski, I., Matthias, V., Möllmann, C., Pätsch, J., Scharfe, M., Thomas, H., Weisse, R., and Zorita, E.: The
5246 North Sea — A shelf sea in the Anthropocene, *J. Mar. Syst.*, 141, 18-33,
5247 <https://doi.org/10.1016/j.jmarsys.2014.03.012>, 2015.

5248 Emeis, K., Christiansen, C., Edelvang, K., Jähmlich, S., Kozuch, J., Laima, M., Leipe, T., Löffler, A., Lund-
5249 Hansen, L. C., Miltner, A., Pazdro, K., Pempkowiak, J., Pollehne, F., Shimmield, T., Voss, M., and Witt, G.:
5250 Material transport from the near shore to the basinal environment in the southern Baltic Sea: II: Synthesis of data
5251 on origin and properties of material, *J. Mar. Syst.*, 35, 151-168, [https://doi.org/10.1016/S0924-7963\(02\)00127-6](https://doi.org/10.1016/S0924-7963(02)00127-6),
5252 2002.

5253 EMEP: Transboundary particulate matter, photo-oxidants, acidifying and eutrophying components, Norwegian
5254 Meteorological Institute, Oslo, Norway, EMEP Status Report 1/2018, 204pp, 2018.

5255 Enfield, D. B., Mestas-Núñez, A. M., and Trimble, P. J.: The Atlantic Multidecadal Oscillation and its relation to
5256 rainfall and river flows in the continental U.S, *Geophys. Res. Lett.*, 28, 2077-2080,
5257 <https://doi.org/10.1029/2000gl012745>, 2001.

Formatiert: Englisch (USA)

Formatiert: Schriftart: Nicht Kursiv

5258 Estilow, T. W., Young, A. H., and Robinson, D. A.: A long-term Northern Hemisphere snow cover extent data
 5259 record for climate studies and monitoring, *Earth Syst. Sci. Data*, 7, 137–142, [https://doi.org/10.5194/essd-7-137-](https://doi.org/10.5194/essd-7-137-2015)
 5260 2015, 2015.

5261 [European Commission: DIRECTIVE 98/70/EC OF THE EUROPEAN PARLIAMENT AND OF THE](#)
 5262 [COUNCIL of 13 October 1998 relating to the quality of petrol and diesel fuels and amending Council](#)
 5263 [Directive 93/12/EEC, 1998.](#)

5264 [European Commission: DIRECTIVE \(EU\) 2018/2001 OF THE EUROPEAN PARLIAMENT AND OF THE](#)
 5265 [COUNCIL of 11 December 2018 on the promotion of the use of energy from renewable sources \(recast\) \(Text](#)
 5266 [with EEA relevance\), 2018a.](#)

5267 [European Commission: REGULATION \(EU\) 2018/1999 OF THE EUROPEAN PARLIAMENT AND OF THE](#)
 5268 [COUNCIL of 11 December 2018 on the Governance of the Energy Union and Climate Action, amending](#)
 5269 [Regulations \(EC\) No 663/2009 and \(EC\) No 715/2009 of the European Parliament and of the Council,](#)
 5270 [Directives 94/22/EC, 98/70/EC, 2009/31/EC, 2009/73/EC, 2010/31/EU, 2012/27/EU and 2013/30/EU of the](#)
 5271 [European Parliament and of the Council, Council Directives 2009/119/EC and \(EU\) 2015/652 and repealing](#)
 5272 [Regulation \(EU\) No 525/2013 of the European Parliament and of the Council \(Text with EEA relevance\), 2018b.](#)

5273 European Environment Agency: Air quality in Europe - 2019 report, Publications Office of the European Union,
 5274 Luxembourg, EEA Report No. 10/2019, 99pp, <https://doi.org/10.2800/822355>, 2019a.

5275 European Environment Agency: Heavy precipitation in Europe. Indicator Assessment CLIM 004. IND -92-en:
 5276 https://www.eea.europa.eu/ds_resolveuid/998fbc113cc84e9a978becd87079f874, access: 24-03-2021, 2019b.

5277 European Environment Agency: Global and European sea-level rise: [https://www.eea.europa.eu/data-and-](https://www.eea.europa.eu/data-and-maps/indicators/sea-level-rise-6/assessment)
 5278 [maps/indicators/sea-level-rise-6/assessment](https://www.eea.europa.eu/data-and-maps/indicators/sea-level-rise-6/assessment), access: 08-12-2020, 2019c.

5279 Eyiring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the
 5280 Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geosci. Model*
 5281 *Dev.*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.

5282 Farinotti, D., Huss, M., Fürst, J. J., Landmann, J., Machguth, H., Maussion, F., and Pandit, A.: A consensus
 5283 estimate for the ice thickness distribution of all glaciers on Earth, *Nat. Geosci.*, 12, 168–173,
 5284 <https://doi.org/10.1038/s41561-019-0300-3>, 2019.

5285 Feistel, R., Nausch, G., and Wasmund, N.: State and evolution of the Baltic Sea, 1952–2005: a detailed 50-year
 5286 survey of meteorology and climate, physics, chemistry, biology, and marine environment, John Wiley & Sons,
 5287 Hoboken, NJ, 2008.

5288 Feldstein, S. B.: The Recent Trend and Variance Increase of the Annular Mode, *J. Climate*, 15, 88–94,
 5289 [https://doi.org/10.1175/1520-0442\(2002\)015<0088:Trtavi>2.0.Co;2](https://doi.org/10.1175/1520-0442(2002)015<0088:Trtavi>2.0.Co;2), 2002.

5290 [Ferreira, J. A., & Soares, C. G.: Modelling distributions of significant wave height, *Coastal Engineering*, 40\(4\),](#)
 5291 [361–374, 2000.](#)

5292 Feser, F., Barcikowska, M., Krueger, O., Schenk, F., Weisse, R., and Xia, L.: Storminess over the North Atlantic
 5293 and northwestern Europe—A review, *Q. J. Roy. Meteor. Soc.*, 141, 350–382, <https://doi.org/10.1002/qj.2364>,
 5294 2015.

5295 [Figueroa, D., Capo, E., Lindh, M. V., Rowe, O. F., Paczkowska, J., Pinhassi, J., and Andersson, A.: Terrestrial](#)
 5296 [dissolved organic matter inflow drives temporal dynamics of the bacterial community of a subarctic estuary](#)
 5297 [\(northern Baltic Sea\), *Environ. Microbiol.*, 23, 4200–4213, <https://doi.org/10.1111/1462-2920.15597>, 2021.](#)

5298 Filazzola, A., Blagrove, K., Imrit, M. A., and Sharma, S.: Climate Change Drives Increases in Extreme Events
 5299 for Lake Ice in the Northern Hemisphere, *Geophys. Res. Lett.*, 47, e2020GL089608,
 5300 <https://doi.org/10.1029/2020GL089608>, 2020.

5301 Fischer-Bruns, I., Storch, H. v., González-Rouco, J. F., and Zorita, E.: Modelling the variability of midlatitude
 5302 storm activity on decadal to century time scales, *Clim. Dyn.*, 25, 461–476, [https://doi.org/10.1007/s00382-005-](https://doi.org/10.1007/s00382-005-0036-1)
 5303 0036-1, 2005.

Formatiert: Schriftart: (Standard) Times New Roman, 10 Pt., Englisch (USA)

Formatiert: Rechtschreibung und Grammatik nicht prüfen

Formatiert: Schriftart: Nicht Kursiv

Formatiert: Schriftart: Nicht Kursiv

Formatiert: Schriftart: +Textkörper (Calibri)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Rechtschreibung und Grammatik nicht prüfen

Feldfunktion geändert

5304 Fleming, Z. L., Doherty, R. M., von Schneidmesser, E., Malley, C. S., Cooper, O. R., Pinto, J. P., Colette, A.,
5305 Xu, X., Simpson, D., Schultz, M. G., Lefohn, A. S., Hamad, S., Moolla, R., Solberg, S., and Feng, Z.:
5306 Tropospheric Ozone Assessment Report: Present-day ozone distribution and trends relevant to human health,
5307 *Elementa-Sci. Anthropol.*, 6, <https://doi.org/10.1525/elementa.273>, 2018.

5308 Fontrodona Bach, A., van der Schrier, G., Melsen, L. A., Klein Tank, A. M. G., and Teuling, A. J.: Widespread
5309 and Accelerated Decrease of Observed Mean and Extreme Snow Depth Over Europe, *Geophys. Res. Lett.*, 45,
5310 12,312-312,319, <https://doi.org/10.1029/2018GL079799>, 2018.

5311 Forster, P. M., Maycock, A. C., McKenna, C. M., and Smith, C. J.: Latest climate models confirm need for
5312 urgent mitigation, *Nat. Clim. Change*, 10, 7-10, <https://doi.org/10.1038/s41558-019-0660-0>, 2020.

5313 Fox, A. D., Jónsson, J. E., Aarvak, T., Bregnballe, T., Christensen, T. K., Clausen, K. K., Clausen, P., Dalby, L.,
5314 Holm, T. E., Pavón-Jordan, D., Laursen, K., Lehtikoinen, A., Lorentsen, S.-H., Møller, A. P., Nordström, M.,
5315 Öst, M., Söderquist, P., and Roland Therkildsen, O.: Current and Potential Threats to Nordic Duck Populations
5316 — A Horizon Scanning Exercise, *Annal. Zool.*, 52, 193-220, 128, 2015.

5317 Fox, A. D., Nielsen, R. D., and Petersen, I. K.: Climate-change not only threatens bird populations but also
5318 challenges our ability to monitor them, *Ibis*, 161, 467-474, <https://doi.org/10.1111/ibi.12675>, 2019.

5319 Frajka-Williams, E., Beaulieu, C., and Duchez, A.: Emerging negative Atlantic Multidecadal Oscillation index in
5320 spite of warm subtropics, *Sci. Rep.-UK*, 7, 11224, <https://doi.org/10.1038/s41598-017-11046-x>, 2017.

5321 Francis, J. A., and Vavrus, S. J.: Evidence linking Arctic amplification to extreme weather in mid-latitudes,
5322 *Geophys. Res. Lett.*, 39, <https://doi.org/10.1029/2012GL051000>, 2012.

5323 Francis, J. A., and Vavrus, S. J.: Evidence for a wavier jet stream in response to rapid Arctic warming, *Environ.*
5324 *Res. Lett.*, 10, 014005, <https://doi.org/10.1088/1748-9326/10/1/014005>, 2015.

5325 Fransner, F., Gustafsson, E., Tedesco, L., Vichi, M., Hordoir, R., Roquet, F., Spilling, K., Kuznetsov, I., Eilola,
5326 K., Mörrth, C.-M., Humborg, C., and Nycander, J.: Non-Redfieldian Dynamics Explain Seasonal pCO₂
5327 Drawdown in the Gulf of Bothnia, *J. Geophys. Res-Oceans*, 123, 166-188,
5328 <https://doi.org/10.1002/2017JC013019>, 2018.

5329 Friedland, R., Neumann, T., and Schernewski, G.: Climate change and the Baltic Sea action plan: model
5330 simulations on the future of the western Baltic Sea, *J. Mar. Syst.*, 105-108, 175-186,
5331 <https://doi.org/10.1016/j.jmarsys.2012.08.002>, 2012.

5332 Friedlingstein, P., O'Sullivan, M., Jones, M. W., Andrew, R. M., Hauck, J., Olsen, A., Peters, G. P., Peters, W.,
5333 Pongratz, J., Sitch, S., Quéré, C. L., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S., Aragão, L. E. O. C.,
5334 Arneth, A., Arora, V., Bates, N. R., Becker, M., Benoit-Cattin, A., Bittig, H. C., Bopp, L., Bultan, S., Chandra,
5335 N., Chevallier, F., Chini, L. P., Evans, W., Florentie, L., Forster, P. M., Gasser, T., Gehlen, M., Gilfillan, D.,
5336 Gkritzalis, T., Gregor, L., Gruber, N., Harris, I., Hartung, K., Haverd, V., Houghton, R. A., Ilyina, T., Jain, A.
5337 K., Joetzjer, E., Kadono, K., Kato, E., Kitidis, V., Korsbakken, J. I., Landschützer, P., Lefèvre, N., Lenton, A.,
5338 Lienert, S., Liu, Z., Lombardozzi, D., Marland, G., Metzl, N., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-I.,
5339 Niwa, Y., O'Brien, K., Ono, T., Palmer, P. I., Pierrot, D., Poulter, B., Resplandy, L., Robertson, E., Rödenbeck,
5340 C., Schwinger, J., Séférian, R., Skjelvan, I., Smith, A. J. P., Sutton, A. J., Tanhua, T., Tans, P. P., Tian, H.,
5341 Tilbrook, B., Werf, G. v. d., Vuichard, N., Walker, A. P., Wanninkhof, R., Watson, A. J., Willis, D., Wiltshire,
5342 A. J., Yuan, W., Yue, X., and Zaehle, S.: Global carbon budget 2020, *Earth Syst. Sci. Data*, 12, 3269-3340,
5343 <https://doi.org/10.5194/essd-12-3269-2020>, 2020.

5344 Frommel, A. Y., Schubert, A., Piatkowski, U., and Clemmesen, C.: Egg and early larval stages of Baltic cod,
5345 *Gadus morhua*, are robust to high levels of ocean acidification, *Mar. Biol.*, 160, 1825-1834,
5346 <https://doi.org/10.1007/s00227-011-1876-3>, 2013.

5347 Fronzek, S., Luoto, M., and Carter, T. R.: Potential effect of climate change on the distribution of palsa mires in
5348 subarctic Fennoscandia, *Clim. Res.*, 32, 1-12, <https://doi.org/10.3354/cr032001>, 2006.

5349 Fronzek, S., and Carter, T. R.: Assessing uncertainties in climate change impacts on resource potential for
5350 Europe based on projections from RCMs and GCMs, *Clim. Change*, 81, 357-371,
5351 <https://doi.org/10.1007/s10584-006-9214-3>, 2007.

5352 Fuchs, M., Kuhry, P., and Hugelius, G.: Low below-ground organic carbon storage in a subarctic Alpine
5353 permafrost environment, *Cryosphere*, 9, 427–438, <https://doi.org/10.5194/tc-9-427-2015>, 2015.

5354 Gaget, E., Pavón-Jordán, D., Johnston, A., Lehtikainen, A., Hochachka, W. M., Sandercock, B. K., Soutan, A.,
5355 Azafzaf, H., Bendjedda, N., Bino, T., Božič, L., Clausen, P., Dakki, M., Devos, K., Domsa, C., Encarnação, V.,
5356 Erciyas-Yavuz, K., Faragó, S., Frost, T., Gaudard, C., Gosztonyi, L., Haas, F., Hornman, M., Langendoen, T.,
5357 Ieronymidou, C., Kostyushin, V. A., Lewis, L. J., Lorentsen, S.-H., Luijckx, L., Meissner, W., Mikuska, T.,
5358 Molina, B., Musilová, Z., Natykanets, V., Paquet, J.-Y., Petkov, N., Portolou, D., Ridzoň, J., Sayoud, S., Šćiban,
5359 M., Sniauksta, L., Stipniece, A., Strebel, N., Teufelbauer, N., Topić, G., Uzunova, D., Vizi, A., Wahl, J.,
5360 Zenatello, M., and Brommer, J. E.: Benefits of protected areas for nonbreeding waterbirds adjusting their
5361 distributions under climate warming, *Conserv. Biol.*, n/a, <https://doi.org/10.1111/cobi.13648>, 2021.

5362 Gailiusis, B., Kriaučiūnienė, J., Jakimavičius, D., and Šarauskienė, D.: The variability of long-term runoff series
5363 in the Baltic Sea drainage basin, *Baltica*, 24, 45–54, 2011.

5364 Gaillard, M.-J., Kleinen, T., Samuelsson, P., Nielsen, A. B., Bergh, J., Kaplan, J., Poska, A., Sandström, C.,
5365 Strandberg, G., Trondman, A.-K., and Wramneby, A.: Causes of Regional Change—Land Cover, in: Second
5366 Assessment of Climate Change for the Baltic Sea Basin, edited by: BACC II Author Team, Springer
5367 International Publishing, Cham, 453–477, https://doi.org/10.1007/978-3-319-16006-1_25, 2015.

5368 Galatius, A., Kinze, C. C., and Teilmann, J.: Population structure of harbour porpoises in the Baltic region:
5369 evidence of separation based on geometric morphometric comparisons, *J. Mar. Biol. Assoc. UK*, 92, 1669–1676,
5370 <https://doi.org/10.1017/S0025315412000513>, 2012.

5371 Gálos, B., Hagemann, S., Hänsler, A., Kindermann, G., Rechid, D., Sieck, K., Teichmann, C., and Jacob, D.:
5372 Case study for the assessment of the biogeophysical effects of a potential afforestation in Europe, *Carbon Bal.*
5373 *Manage.*, 8, 3, <https://doi.org/10.1186/1750-0680-8-3>, 2013.

5374 Gao, Y., Markkanen, T., Backman, L., Henttonen, H. M., Pietikäinen, J. P., Mäkelä, H. M., and Laaksonen, A.:
5375 Biogeophysical impacts of peatland forestation on regional climate changes in Finland, *Biogeosciences*, 11,
5376 7251–7267, <https://doi.org/10.5194/bg-11-7251-2014>, 2014.

5377 Gårdmark, A., Lindegren, M., Neuenfeldt, S., Blenckner, T., Heikinheimo, O., Müller-Karulis, B., Niiranen, S.,
5378 Tomczak, M. T., Aro, E., Wikström, A., and Möllmann, C.: Biological ensemble modeling to evaluate potential
5379 futures of living marine resources, *Ecol. Appl.*, 23, 742–754, <https://doi.org/10.1890/12-0267.1>, 2013.

5380 Gårdmark, A., and Huss, M.: Individual variation and interactions explain food web responses to global
5381 warming, *Philos. T. Roy. Soc. B.*, 375, 20190449, <https://doi.org/10.1098/rstb.2019.0449>, 2020.

5382 Garzke, J., Ismar, S. M. H., and Sommer, U.: Climate change affects low trophic level marine consumers:
5383 warming decreases copepod size and abundance, *Oecologia*, 177, 849–860, [https://doi.org/10.1007/s00442-014-](https://doi.org/10.1007/s00442-014-3130-4)
5384 3130-4, 2015.

5385 Gauss, M., Jonson, J. E., and Nyíri, Á.: Emissions from international shipping, The Norwegian Meteorological
5386 Institute Oslo, Norway, 129–133, 2017.

5387 Gauss, M., Bartnicki, J., Jalkanen, J.-P., Nyíri, A., Klein, H., Fagerli, H., and Klimont, Z.: Airborne nitrogen
5388 deposition to the Baltic Sea: Past trends, source allocation and future projections, *Atmos. Environ.*, 253, 118377,
5389 <https://doi.org/10.1016/j.atmosenv.2021.118377>, 2021.

5390 Geels, C., Andersson, C., Hänninen, O., Lansø, A. S., Schwarze, P. E., Skjøth, C. A., and Brandt, J.: Future
5391 Premature Mortality Due to O₃, Secondary Inorganic Aerosols and Primary PM in Europe — Sensitivity to
5392 Changes in Climate, Anthropogenic Emissions, Population and Building Stock, *Int. J. Env. Res. Pub. He.*, 12,
5393 2837–2869, 2015.

5394 Gillett, N. P., Arora, V. K., Matthews, D., and Allen, M. R.: Constraining the Ratio of Global Warming to
5395 Cumulative CO₂ Emissions Using CMIP5 Simulations, *J. Climate*, 26, 6844–6858, [https://doi.org/10.1175/jcli-d-](https://doi.org/10.1175/jcli-d-12-00476.1)
5396 12-00476.1, 2013.

5397 [Girjatowicz, J. P.: Ice thrusts and piles on the shores of the Southern Baltic Sea coast \(Poland\) lagoons, Baltic](#)
5398 [Coastal Zone. Journal of Ecology and Protection of the Coastline, 8, 5-22, 2004.](#)

Formatiert: Englisch (USA)

Formatiert: Rechtschreibung und Grammatik nicht prüfen

5399 Girjatowicz, J. P., and Łabuz, T. A.: Forms of piled ice at the southern coast of the Baltic Sea, *Estuar. Coast.*
5400 *Shelf Sci.*, 239, 106746, <https://doi.org/10.1016/j.ecss.2020.106746>, 2020.

5401 Gisnäs, K., Etzelmüller, B., Lussana, C., Hjort, J., Sannel, A. B. K., Isaksen, K., Westermann, S., Kuhry, P.,
5402 Christiansen, H. H., Frampton, A., and Åkerman, J.: Permafrost Map for Norway, Sweden and Finland,
5403 *Permafrost Periglac.*, 28, 359-378, <https://doi.org/10.1002/ppp.1922>, 2017.

5404 Glantz, P., Freud, E., Johansson, C., Noone, K. J., and Tesche, M.: Trends in MODIS and AERONET derived
5405 aerosol optical thickness over Northern Europe, *Tellus B*, 71, 1554414,
5406 <https://doi.org/10.1080/16000889.2018.1554414>, 2019.

5407 Graf, A., Klosterhalfen, A., Arriga, N., Bernhofer, C., Bogen, H., Bornet, F., Brüggemann, N., Brümmer, C.,
5408 Buchmann, N., Chi, J., Chipeaux, C., Cremonese, E., Cuntz, M., Dušek, J., El-Madany, T. S., Fares, S., Fischer,
5409 M., Foltýnová, L., Gharun, M., Ghiasi, S., Gielen, B., Gottschalk, P., Grünwald, T., Heinemann, G., Heinesch,
5410 B., Heliasz, M., Holst, J., Hörtnagl, L., Ibrom, A., Ingwersen, J., Jurasinski, G., Klatt, J., Knohl, A., Koebsch, F.,
5411 Konopka, J., Korkiakoski, M., Kowalska, N., Kremer, P., Kruijt, B., Lafont, S., Léonard, J., Ligne, A. D.,
5412 Longdoz, B., Loustau, D., Magliulo, V., Mammarella, I., Manca, G., Mauder, M., Migliavacca, M., Mölder, M.,
5413 Neiryneck, J., Ney, P., Nilsson, M., Paul-Limoges, E., Peichl, M., Pitacco, A., Poyda, A., Reibmann, C., Roland,
5414 M., Sachs, T., Schmidt, M., Schrader, F., Siebicke, L., Šigut, L., Tuittila, E.-S., Varlagin, A., Vendrame, N.,
5415 Vincke, C., Völksch, I., Weber, S., Wille, C., Witzemann, H.-D., Zeeman, M., and Vereecken, H.: Altered energy
5416 partitioning across terrestrial ecosystems in the European drought year 2018, *Philos. T. Roy. Soc. B.*, 375,
5417 20190524, <https://doi.org/10.1098/rstb.2019.0524>, 2020.

5418 Graham, L. P.: Climate Change Effects on River Flow to the Baltic Sea, *Ambio* 33, 235-241, 237,
5419 <https://doi.org/10.1579/0044-7447-33.4.235>, 2004.

5420 Graham, P.: Modeling runoff to the Baltic Sea, *Ambio*, 28, 328-334, 1999.

5421 Graiff, A., Liesner, D., Karsten, U., and Bartsch, I.: Temperature tolerance of western Baltic Sea *Fucus*
5422 *vesiculosus* – growth, photosynthesis and survival, *J. Exp. Mar. Biol. Ecol.*, 471, 8-16,
5423 <https://doi.org/10.1016/j.jembe.2015.05.009>, 2015.

5424 Granéli, E., Kerstin, W., Larsson, U., Granéli, W., and Elmgren, R.: Nutrient Limitation of Primary Production
5425 in the Baltic Sea Area, *Ambio*, 19, 142-151, 1990.

5426 Gräwe, U., and Burchard, H.: Storm surges in the Western Baltic Sea: the present and a possible future, *Clim.*
5427 *Dyn.*, 39, 165-183, <https://doi.org/10.1007/s00382-011-1185-z>, 2012.

5428 Gräwe, U., Naumann, M., Mohrholz, V., and Burchard, H.: Anatomizing one of the largest saltwater inflows into
5429 the Baltic Sea in December 2014, *J. Geophys. Res-Oceans*, 120, 7676-7697,
5430 <https://doi.org/10.1002/2015JC011269>, 2015.

5431 Griffiths, J. R., Kadin, M., Nascimento, F. J. A., Tamelander, T., Törnroos, A., Bonaglia, S., Bonsdorff, E.,
5432 Brüchert, V., Gårdmark, A., Järnström, M., Kotta, J., Lindegren, M., Nordström, M. C., Norkko, A., Olsson, J.,
5433 Weigel, B., Žydelis, R., Blenckner, T., Niiranen, S., and Winder, M.: The importance of benthic–pelagic
5434 coupling for marine ecosystem functioning in a changing world, *Global Change Biol.*, 23, 2179-2196,
5435 <https://doi.org/10.1111/gcb.13642>, 2017.

5436 Griffiths, J. R., Lehtinen, S., Suikkanen, S., and Winder, M.: Limited evidence for common interannual trends in
5437 Baltic Sea summer phytoplankton biomass, *PLoS ONE*, 15, e0231690,
5438 <https://doi.org/10.1371/journal.pone.0231690>, 2020.

5439 Grinsted, A.: Projected Change - Sea Level, in: Second Assessment of Climate Change for the Baltic Sea Basin,
5440 edited by: BACC II Author Team, *Regional Climate Studies*, Springer International Publishing, Cham, 253-263,
5441 https://doi.org/10.1007/978-3-319-16006-1_14, 2015.

5442 Grinsted, A., Jevrejeva, S., Riva, R. E. M., and Dahl-Jensen, D.: Sea level rise projections for northern Europe
5443 under RCP8.5, *Clim. Res.*, 64, 15-23, <https://doi.org/10.3354/cr01309> 2015.

5444 Grise, K. M., and Polvani, L. M.: The response of midlatitude jets to increased CO₂: Distinguishing the roles of
5445 sea surface temperature and direct radiative forcing, *Geophys. Res. Lett.*, 41, 6863-6871,
5446 <https://doi.org/10.1002/2014GL061638>, 2014.

5447 Groetsch, P. M. M., Simis, S. G. H., Eleveld, M. A., and Peters, S. W. M.: Spring blooms in the Baltic Sea have
5448 weakened but lengthened from 2000 to 2014, *Biogeosciences*, 13, 4959–4973, [https://doi.org/10.5194/bg-13-](https://doi.org/10.5194/bg-13-4959-2016)
5449 4959-2016, 2016.

5450 Gröger, M., Dieterich, C., Meier, H. E. M., and Schimanke, S.: Thermal air-sea coupling in hindcast simulations
5451 for the North Sea and Baltic Sea on the NW European shelf, *Tellus A*, 67, 26911,
5452 <https://doi.org/10.3402/tellusa.v67.26911>, 2015.

5453 Gröger, M., Arneborg, L., Dieterich, C., Höglund, A., and Meier, H. E. M.: Summer hydrographic changes in the
5454 Baltic Sea, Kattegat and Skagerrak projected in an ensemble of climate scenarios downscaled with a coupled
5455 regional ocean–sea ice–atmosphere model, *Clim. Dyn.*, <https://doi.org/10.1007/s00382-019-04908-9>, 2019.

5456 Gröger, M., Dieterich, C., Ho-Hagemann, H. T. M., Hagemann, S., Jakacki, J., May, W., Meier, H. E. M.,
5457 Miller, P. A., Rutgersson, A., and Wu, L.: Coupled regional Earth system modelling in the Baltic Sea region,
5458 *Earth Syst. Dynam. Discuss.*, 1–50, <https://doi.org/10.5194/esd-2021-14>, 2021a.

5459 Gröger, M., Dieterich, C., and Meier, H. E. M.: Is interactive air sea coupling relevant for simulating the future
5460 climate of Europe?, *Clim. Dyn.*, 56, 491–514, <https://doi.org/10.1007/s00382-020-05489-8>, 2021b.

5461 Groh, A., Richter, A., and Dietrich, R.: Recent Baltic Sea Level Changes Induced by Past and Present Ice
5462 Masses, in: *Coastline Changes of the Baltic Sea from South to East: Past and Future Projection*, edited by: Harff,
5463 J., Furmańczyk, K., and von Storch, H., Springer International Publishing, Cham, 55–68,
5464 https://doi.org/10.1007/978-3-319-49894-2_4, 2017.

5465 Groll, N., Grabemann, I., Hünicke, B., and Meese, M.: Baltic Sea wave conditions under climate change
5466 scenarios, *Boreal Environ. Res.*, 22, 1–12, 2017.

5467 Groß, D., Zander, A., Boethius, A., Dreibrodt, S., Grøn, O., Hansson, A., Jessen, C., Koivisto, S., Larsson, L.,
5468 Lübke, H., and Nilsson, B.: People, lakes and seashores: Studies from the Baltic Sea basin and adjacent areas in
5469 the early and Mid-Holocene, *Quaternary Sci. Rev.*, 185, 27–40, <https://doi.org/10.1016/j.quascirev.2018.01.021>,
5470 2018.

5471 Gubelit, Y. I.: Climatic impact on community of filamentous macroalgae in the Neva estuary (eastern Baltic
5472 Sea), *Mar. Pollut. Bull.*, 91, 166–172, <https://doi.org/10.1016/j.marpolbul.2014.12.009>, 2015.

5473 Gunnarsson, G., Waldenström, J., and Fransson, T.: Direct and indirect effects of winter harshness on the
5474 survival of Mallards *Anas platyrhynchos* in northwest Europe, *Ibis*, 154, 307–317, [https://doi.org/10.1111/j.1474-](https://doi.org/10.1111/j.1474-919X.2011.01206.x)
5475 919X.2011.01206.x, 2012.

5476 Gustafsson, B. G., Schenk, F., Blenckner, T., Eilola, K., Meier, H. E. M., Müller-Karulis, B., Neumann, T.,
5477 Ruoho-Airola, T., Savchuk, O. P., and Zorita, E.: Reconstructing the development of Baltic Sea eutrophication
5478 1850–2006, *Ambio*, 41, 534–548, <https://doi.org/10.1007/s13280-012-0318-x>, 2012.

5479 Gustafsson, E., Deutsch, B., Gustafsson, B. G., Humborg, C., and Mörrh, C. M.: Carbon cycling in the Baltic Sea
5480 — The fate of allochthonous organic carbon and its impact on air–sea CO₂ exchange, *J. Mar. Syst.*, 129, 289–
5481 302, <https://doi.org/10.1016/j.jmarsys.2013.07.005>, 2014.

5482 Gustafsson, E., Savchuk, O. P., Gustafsson, B. G., and Müller-Karulis, B.: Key processes in the coupled carbon,
5483 nitrogen, and phosphorus cycling of the Baltic Sea, *Biogeochemistry*, 134, 301–317,
5484 <https://doi.org/10.1007/s10533-017-0361-6>, 2017.

5485 Gustafsson, E., Hagens, M., Sun, X., Reed, D. C., Humborg, C., Slomp, C. P., and Gustafsson, B. G.:
5486 Sedimentary alkalinity generation and long-term alkalinity development in the Baltic Sea, *Biogeosciences*, 16,
5487 437–456, <https://doi.org/10.5194/bg-2018-313>, 2019.

5488 [Gutiérrez, J.M., R.G. Jones, G.T. Narisma, L.M. Alves, M. Amjad, I.V. Gorodetskaya, M. Grose, N.A.B. Klutse,](https://www.ipcc.ch/interactive-atlas)
5489 [S. Krakovska, J. Li, D. Martínez-Castro, L.O. Mearns, S.H. Mernild, T. Ngo-Duc, B. van den Hurk, and J.-H.](https://www.ipcc.ch/interactive-atlas)
5490 [Yoon, 2021: Atlas. In Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to](https://www.ipcc.ch/interactive-atlas)
5491 [the Sixth Assessment Report of the Intergovernmental Panel on Climate Change \[Masson-Delmotte, V., P. Zhai,](https://www.ipcc.ch/interactive-atlas)
5492 [A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell,](https://www.ipcc.ch/interactive-atlas)
5493 [E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou \(eds.\)\]. Available](https://www.ipcc.ch/interactive-atlas)
5494 [from <http://interactive-atlas.ipcc.ch/>, access: 25.11.2021.](https://www.ipcc.ch/interactive-atlas)

5495 Haapala, J., Meier, H. E. M., and Rinne, J.: Numerical investigations of future ice conditions in the Baltic Sea,
5496 *Ambio*, 30, 237-244, <https://doi.org/10.1579/0044-7447-30.4.237>, 2001.

5497 Haarsma, R. J., Selten, F. M., and Drijfhout, S. S.: Decelerating Atlantic meridional overturning circulation main
5498 cause of future west European summer atmospheric circulation changes, *Environ. Res. Lett.*, 10, 094007,
5499 <https://doi.org/10.1088/1748-9326/10/9/094007>, 2015.

5500 Haavisto, F., and Jormalainen, V.: Seasonality elicits herbivores' escape from trophic control and favors induced
5501 resistance in a temperate macroalga, *Ecology*, 95, 3035-3045, <https://doi.org/10.1890/13-2387.1>, 2014.

5502 Hägg, H. E., Lyon, S. W., Wällstedt, T., Mörh, C.-M., Claremar, B., and Humborg, C.: Future Nutrient Load
5503 Scenarios for the Baltic Sea Due to Climate and Lifestyle Changes, *Ambio* 43, 337-351,
5504 <https://doi.org/10.1007/s13280-013-0416-4>, 2014.

5505 Halkka, A.: Changing climate and the Baltic region biota, Doctoral dissertation, Faculty of Biological and
5506 Environmental Sciences, University of Helsinki, Helsinki, Finland, 52 pp., 2020.

5507 Hallett, C. S., Hobday, A. J., Tweedley, J. R., Thompson, P. A., McMahon, K., and Valesini, F. J.: Observed and
5508 predicted impacts of climate change on the estuaries of south-western Australia, a Mediterranean climate region,
5509 *Reg. Environ. Change*, 18, 1357-1373, <https://doi.org/10.1007/s10113-017-1264-8>, 2018.

5510 Hällfors, H., Backer, H., Leppänen, J.-M., Hällfors, S., Hällfors, G., and Kuosa, H.: The northern Baltic Sea
5511 phytoplankton communities in 1903–1911 and 1993–2005: a comparison of historical and modern species data,
5512 *Hydrobiologia*, 707, 109-133, <https://doi.org/10.1007/s10750-012-1414-4>, 2013.

5513 Hamlington, B. D., Gardner, A. S., Ivins, E., Lenaerts, J. T. M., Reager, J. T., Trossman, D. S., Zaron, E. D.,
5514 Adhikari, S., Arendt, A., Aschwanden, A., Beckley, B. D., Bekaert, D. P. S., Blewitt, G., Caron, L., Chambers,
5515 D. P., Chandanpurkar, H. A., Christianson, K., Csatho, B., Cullather, R. I., DeConto, R. M., Fasullo, J. T.,
5516 Frederikse, T., Freymueller, J. T., Gilford, D. M., Giroto, M., Hammond, W. C., Hock, R., Holschuh, N., Kopp,
5517 R. E., Landerer, F., Larour, E., Menemenlis, D., Merrifield, M., Mitrovica, J. X., Nerem, R. S., Nias, I. J.,
5518 Nieves, V., Nowicki, S., Pangaluru, K., Piecuch, C. G., Ray, R. D., Rounce, D. R., Schlegel, N.-J., Seroussi, H.,
5519 Shirzaei, M., Sweet, W. V., Velicogna, I., Vinogradova, N., Wahl, T., Wiese, D. N., and Willis, M. J.:
5520 Understanding of Contemporary Regional Sea-Level Change and the Implications for the Future, *Rev. Geophys.*,
5521 58, e2019RG000672, <https://doi.org/10.1029/2019RG000672>, 2020.

5522 Hammer, K., Schneider, B., Kuliński, K., and Schulz-Bull, D. E.: Acid-base properties of Baltic Sea dissolved
5523 organic matter, *J. Mar. Syst.*, 173, 114-121, <https://doi.org/10.1016/j.jmarsys.2017.04.007>, 2017.

5524 Hammond, P., Bearzi, G., Bjørge, A., Forney, K., Karczmarski, L., Kasuya, T., Perrin, W., Scott, M., Wang, J.,
5525 and Wells, R.: *Phocoena phocoena* - Baltic Sea subpopulation (errata version published in 2016)(No. e.
5526 T17031A98831650), The IUCN Red List of Threatened Species, 2008.

5527 Hannerz, F., and Destouni, G.: Spatial Characterization of the Baltic Sea Drainage Basin and Its Unmonitored
5528 Catchments, *Ambio* 35, 214-219, 216, <https://doi.org/10.1579/05-A-022R.1>, 2006.

5529 Hänninen, J., Vuorinen, I., Rajasilta, M., and Reid, P. C.: Response of the Baltic and North Seas to river runoff
5530 from the Baltic watershed – Physical and biological changes, *Prog. Oceanogr.*, 138, 91-104,
5531 <https://doi.org/10.1016/j.pocean.2015.09.001>, 2015.

5532 Hansson, D., Eriksson, C., Omstedt, A., and Chen, D.: Reconstruction of river runoff to the Baltic Sea, AD
5533 1500–1995, *Int. J. Climatol.*, 31, 696-703, <https://doi.org/10.1002/joc.2097>, 2011.

5534 Harff, J., Deng, J., Dudzińska-Nowak, J., Fröhle, P., Groh, A., Hünicke, B., Soomere, T., and Zhang, W.: What
5535 Determines the Change of Coastlines in the Baltic Sea?, in: *Coastline Changes of the Baltic Sea from South to*
5536 *East: Past and Future Projection*, edited by: Harff, J., Furmańczyk, K., and von Storch, H., Springer International
5537 Publishing, Cham, 15-35, https://doi.org/10.1007/978-3-319-49894-2_2, 2017.

5538 Hartfield, G., Blunden, J., and Arndt, D. S.: State of the Climate in 2017, *Bull. Amer. Meteor. Soc.*, 99, Si-S310,
5539 <https://doi.org/10.1175/2018BAMSStateoftheClimate.1>, 2018.

5540 Harvey, B. J., Shaffrey, L. C., Woollings, T. J., Zappa, G., and Hodges, K. I.: How large are projected 21st
5541 century storm track changes?, *Geophys. Res. Lett.*, 39, <https://doi.org/10.1029/2012GL052873>, 2012.

5542 Harvey, B. J., Cook, P., Shaffrey, L. C., and Schiemann, R.: The Response of the Northern Hemisphere Storm
5543 Tracks and Jet Streams to Climate Change in the CMIP3, CMIP5, and CMIP6 Climate Models, J. Geophys. Res -
5544 Atmos., 125, e2020JD032701, <https://doi.org/10.1029/2020JD032701>, 2020.

5545 Hattich, G. S. I., Listmann, L., Govaert, L., Pansch, C., Reusch, T. B. H., and Matthiessen, B.:
5546 Experimentally decomposing phytoplankton community change into ecological and evolutionary
5547 contributions. Functional Ecology, <https://doi.org/10.1111/1365-2435.13923>, 2021.

5548 Hausfather, Z., and Peters, G. P.: Emissions - the 'business as usual' story is misleading, Nature, 577, 618-620,
5549 <https://doi.org/10.1038/d41586-020-00177-3>, 2020.

5550 Havenhand, J. N.: How will ocean acidification affect Baltic Sea ecosystems? An assessment of plausible
5551 impacts on key functional groups. Ambio, 41(6), 637-644, 2012.

5552 Hedegaard, G. B., Christensen, J. H., and Brandt, J.: The relative importance of impacts from climate change vs.
5553 emissions change on air pollution levels in the 21st century, Atmos. Chem. Phys., 13, 3569-3585,
5554 <https://doi.org/10.5194/acp-13-3569-2013>, 2013.

5555 Hegerl, G. C., Crowley, T. J., Baum, S. K., Kim, K.-Y., and Hyde, W. T.: Detection of volcanic, solar and
5556 greenhouse gas signals in paleo-reconstructions of Northern Hemispheric temperature, Geophys. Res. Lett., 30,
5557 <https://doi.org/10.1029/2002GL016635>, 2003.

5558 HELCOM: Climate Change in the Baltic Sea Area - HELCOM Thematic Assessment in 2007, Helsinki, Finland,
5559 Baltic Sea Environment Proceedings No. 111, 54pp, 2007.

5560 HELCOM: HELCOM Copenhagen Ministerial Declaration - Taking Further Action to Implement the Baltic Sea
5561 Action Plan - Reaching Good Environmental Status for a healthy Baltic Sea, Copenhagen, Denmark, HELCOM
5562 Ministerial Meeting, 2013a.

5563 HELCOM: Climate Change in the Baltic Sea Area - HELCOM Thematic Assessment in 2013, Helsinki, Finland,
5564 Baltic Sea Environment Proceedings No. 137, 70pp, 2013b.

5565 HELCOM: Updated Fifth Baltic Sea Pollution Load Compilation (PLC-5.5), Helsinki, Finland, Baltic Sea
5566 Environment Proceedings No. 145, 143pp, 2015.

5567 HELCOM: HELCOM thematic assessment of eutrophication 2011-2016., Helsinki, Finland, Baltic Sea
5568 Environment Proceedings No. 156, 83pp, 2018a.

5569 HELCOM: Population trends and abundance of seals, HELCOM core indicator report, 34pp, 2018b.

5570 HELCOM: The Sixth Pollution Load Compilation (PLC-6), Helsinki, Finland, 15pp, 2018c.

5571 Helle, E.: Reproduction, size and structure of the Baltic ringed seal population of the Bothnian Bay., PhD, Acta
5572 Universitatis Ouluensis series A Scientiae rerum naturalium No. 106 Biologica No. 11, 47 pp., 1980.

5573 Hendriks, C., Forsell, N., Kiesewetter, G., Schaap, M., and Schöpp, W.: Ozone concentrations and damage for
5574 realistic future European climate and air quality scenarios, Atmos. Environ., 144, 208-219,
5575 <https://doi.org/10.1016/j.atmosenv.2016.08.026>, 2016.

5576 Hense, I., Meier, H. E. M., and Sonntag, S.: Projected climate change impact on Baltic Sea cyanobacteria, Clim.
5577 Change, 119, 391-406, <https://doi.org/10.1007/s10584-013-0702-y>, 2013.

5578 Hesse, C., Krysanova, V., Stefanova, A., Bielecka, M., and Domnin, D. A.: Assessment of climate change
5579 impacts on water quantity and quality of the multi-river Vistula Lagoon catchment, Hydrol. Sci. J., 60, 890-911,
5580 <https://doi.org/10.1080/02626667.2014.967247>, 2015.

5581 Hieronymus, J., Eilola, K., Hieronymus, M., Meier, H. E. M., Saraiva, S., and Karlson, B.: Causes of simulated
5582 long-term changes in phytoplankton biomass in the Baltic proper: a wavelet analysis, Biogeosciences, 15, 5113-
5583 5129, <https://doi.org/10.5194/bg-15-5113-2018>, 2018.

5584 Hieronymus, M., and Kalén, O.: Sea-level rise projections for Sweden based on the new IPCC special report:
5585 The ocean and cryosphere in a changing climate, Ambio, <https://doi.org/10.1007/s13280-019-01313-8>, 2020.

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Schriftart: (Standard) +Textkörper (Calibri), 11 Pt.

Formatiert: Schriftart: Nicht Kursiv

Formatiert: Schriftart: Nicht Kursiv

Formatiert: Schriftart: +Textkörper (Calibri)

5586 Hieronymus, J., Eilola, K., Olofsson, M., Hense, I., Meier, H. E. M., and Almroth-Rosell, E.: Modeling
 5587 cyanobacteria life cycle dynamics and historical nitrogen fixation in the Baltic Sea, *Biogeosciences*, 18, 6213-
 5588 6227, <https://doi.org/10.5194/bg-18-6213-2021>, 2021.

5589 Hinrichsen, H.-H., Huwer, B., Makarchouk, A., Petereit, C., Schaber, M., and Voss, R.: Climate-driven long-
 5590 term trends in Baltic Sea oxygen concentrations and the potential consequences for eastern Baltic cod (*Gadus*
 5591 *morhua*), *ICES J. Mar. Sci.*, 68, 2019-2028, <https://doi.org/10.1093/icesjms/fsr145>, 2011.

5592 Hisdal, H., Holmqvist, E., Jonsdottir, J. F., Jonsson, P., Kuusisto, E., Lindstroem, G., and Roald, L. A.: Has
 5593 streamflow changed in the Nordic countries?, Norwegian Water Resources and Energy Directorate (NVE),
 5594 NorwayNVE-1/2010, 28pp, 2010.

5595 Hjerne, O., Hajdu, S., Larsson, U., Downing, A. S., and Winder, M.: Climate Driven Changes in Timing,
 5596 Composition and Magnitude of the Baltic Sea Phytoplankton Spring Bloom, *Front. Mar. Sci.*, 6,
 5597 <https://doi.org/10.3389/fmars.2019.00482>, 2019.

5598 Hock, R., Rasul, G., Adler, C., Cáceres, B., Gruber, S., Hirabayashi, Y., Jackson, M., Kääb, A., Kang, S.,
 5599 Kutuzov, S., Milner, A., Molau, U., Morin, S., Orlove, B., and Steltzer, H.: High Mountain Areas, in: IPCC
 5600 Special Report on the Ocean and Cryosphere in a Changing Climate, edited by: Pörtner, H.-O., Roberts, D. C.,
 5601 Masson-Delmotte, V., Zhai, P., Tignor, M., Poloczanska, E., Mintenbeck, K., Alegría, A., Nicolai, M., Okem,
 5602 A., Petzold, J., Rama, B., and Weyer, N. M., 2019.

5603 Höglund, A., Pemberton, P., Hordoir, R., and Schimanke, S.: Ice conditions for maritime traffic in the Baltic Sea
 5604 in future climate, *Boreal Environ. Res.*, 22, 245-265, 2017.

5605 Hoikkala, L., Kortelainen, P., Soinne, H., and Kuosa, H.: Dissolved organic matter in the Baltic Sea, *J. Mar.*
 5606 *Syst.*, 142, 47-61, <https://doi.org/10.1016/j.jmarsys.2014.10.005>, 2015.

5607 Holmlund, E. S., and Holmlund, P.: Constraining 135 years of mass balance with historic structure-from-motion
 5608 photogrammetry on Storglaciären, Sweden, *Geogr. Ann. A*, 101, 195-210,
 5609 <https://doi.org/10.1080/04353676.2019.1588543>, 2019.

5610 Holopainen, R., Lehtiniemi, M., Meier, H. E. M., Albertsson, J., Gorokhova, E., Kotta, J., and Viitasalo, M.:
 5611 Impacts of changing climate on the non-indigenous invertebrates in the northern Baltic Sea by end of the twenty-
 5612 first century, *Biol. Invasions*, 18, 3015-3032, <https://doi.org/10.1007/s10530-016-1197-z>, 2016.

5613 Holtermann, P. L., and Umlauf, L.: The Baltic Sea Tracer Release Experiment: 2. Mixing processes, *J. Geophys.*
 5614 *Res-Oceans*, 117, <https://doi.org/10.1029/2011JC007445>, 2012.

5615 Holtermann, P. L., Umlauf, L., Tanhua, T., Schmale, O., Rehder, G., and Waniek, J. J.: The Baltic Sea Tracer
 5616 Release Experiment: 1. Mixing rates, *J. Geophys. Res-Oceans*, 117, <https://doi.org/10.1029/2011JC007439>,
 5617 2012.

5618 Holtermann, P. L., Prien, R., Naumann, M., Mohrholz, V., and Umlauf, L.: Deepwater dynamics and mixing
 5619 processes during a major inflow event in the central Baltic Sea, *J. Geophys. Res-Oceans*, 122, 6648-6667,
 5620 <https://doi.org/10.1002/2017JC013050>, 2017.

5621 Hong, B., Swaney, D. P., McCrackin, M., Svanbäck, A., Humborg, C., Gustafsson, B., Yershova, A., and
 5622 Pakhomau, A.: Advances in NANI and NAPI accounting for the Baltic drainage basin: spatial and temporal
 5623 trends and relationships to watershed TN and TP fluxes, *Biogeochemistry*, 133, 245-261,
 5624 <https://doi.org/10.1007/s10533-017-0330-0>, 2017.

5625 Hook, O., Johnels, A. G., and Matthews, L. H.: The breeding and distribution of the grey seal (*Halichoerus*
 5626 *grypus* Fab.) in the Baltic Sea, with observations on other seals of the area, *P. Roy. Soc. B-Biol. Sci.*, 182, 37-58,
 5627 <https://doi.org/10.1098/rspb.1972.0065>, 1972.

5628 Hordoir, R., Axell, L., Löptien, U., Dietze, H., and Kuznetsov, I.: Influence of sea level rise on the dynamics of
 5629 salt inflows in the Baltic Sea, *J. Geophys. Res-Oceans*, 120, 6653-6668, <https://doi.org/10.1002/2014JC010642>,
 5630 2015.

5631 Hordoir, R., Axell, L., Höglund, A., Dieterich, C., Fransner, F., Gröger, M., Liu, Y., Pemberton, P., Schimanke,
 5632 S., Andersson, H., Ljungemyr, P., Nygren, P., Falahat, S., Nord, A., Jönsson, A., Lake, I., Döös, K.,
 5633 Hieronymus, M., Dietze, H., Löptien, U., Kuznetsov, I., Westerlund, A., Tuomi, L., and Haapala, J.: Nemo-

5634 Nordic 1.0: a NEMO-based ocean model for the Baltic and North seas – research and operational applications,
5635 Geosci. Model Dev., 12, 363-386, <https://doi.org/10.5194/gmd-12-363-2019>, 2019.

5636 Horn, H. G., Boersma, M., Garzke, J., Löder, M. G. J., Sommer, U., and Aberle, N.: Effects of high CO₂ and
5637 warming on a Baltic Sea microzooplankton community, ICES J. Mar. Sci., 73, 772-782,
5638 <https://doi.org/10.1093/icesjms/fsv198>, 2015.

5639 Høyer, J. L., and Karagali, I.: Sea Surface Temperature Climate Data Record for the North Sea and Baltic Sea, J.
5640 Climate, 29, 2529-2541, <https://doi.org/10.1175/jcli-d-15-0663.1>, 2016.

5641 Hugelius, G., Loisel, J., Chadburn, S., Jackson, R. B., Jones, M., MacDonald, G., Marushchak, M., Olefeldt, D.,
5642 Packalen, M., Siewert, M. B., Treat, C., Turetsky, M., Voigt, C., and Yu, Z.: Large stocks of peatland carbon and
5643 nitrogen are vulnerable to permafrost thaw, P. Natl. Acad. Sci. USA, 117, 20438-20446,
5644 <https://doi.org/10.1073/pnas.1916387117>, 2020.

5645 Humborg, C., Geibel, M. C., Sun, X., McCrackin, M., Mörtz, C.-M., Stranne, C., Jakobsson, M., Gustafsson, B.,
5646 Sokolov, A., Norkko, A., and Norkko, J.: High Emissions of Carbon Dioxide and Methane From the Coastal
5647 Baltic Sea at the End of a Summer Heat Wave, Front. Mar. Sci., 6, <https://doi.org/10.3389/fmars.2019.00493>,
5648 2019.

5649 Hünicke, B., and Zorita, E.: Trends in the amplitude of Baltic Sea level annual cycle, Tellus A, 60, 154-164,
5650 <https://doi.org/10.1111/j.1600-0870.2007.00277.x>, 2008.

5651 Hünicke, B., Zorita, E., Soomere, T., Madsen, K. S., Johansson, M., and Suursaar, Ü.: Recent Change—Sea
5652 Level and Wind Waves, in: Second Assessment of Climate Change for the Baltic Sea Basin, edited by: BACC II
5653 Author Team, Springer International Publishing, Cham, 155-185, 10.1007/978-3-319-16006-1_9, 2015.

5654 Hünicke, B., and Zorita, E.: Statistical Analysis of the Acceleration of Baltic Mean Sea-Level Rise, 1900–2012,
5655 Front. Mar. Sci., 3, <https://doi.org/10.3389/fmars.2016.00125>, 2016.

5656 Hurrell, J. W.: Decadal trends in the North Atlantic Oscillation: regional temperatures and precipitation, Science-
5657 AAAS-Weekly Paper Edition, 269, 676-679, <https://doi.org/10.1126/science.269.5224.676> 1995.

5658 Hurrell, J. W., Kushnir, Y., Ottersen, G., and Visbeck, M.: An overview of the North Atlantic oscillation,
5659 Geophys. Monogr. Ser., 134, 1-36, 2003.

5660 Huthnance, J., Weisse, R., Wahl, T., Thomas, H., Pietrzak, J., Souza, A. J., van Heteren, S., Schmelzer, N., van
5661 Beusekom, J., Colijn, F., Haigh, I., Hjøllø, S., Holford, J., Kent, E. C., Kühn, W., Loewe, P., Lorkowski, I.,
5662 Mork, K. A., Pätsch, J., Quante, M., Salt, L., Siddorn, J., Smyth, T., Sterl, A., and Woodworth, P.: Recent
5663 Change—North Sea, in: North Sea Region Climate Change Assessment, edited by: Quante, M., and Colijn, F.,
5664 Springer International Publishing, Cham, 85-136, https://doi.org/10.1007/978-3-319-39745-0_3, 2016.

5665 Huttunen, I., Lehtonen, H., Huttunen, M., Piirainen, V., Korppoo, M., Veijalainen, N., Viitasalo, M., and
5666 Vehviläinen, B.: Effects of climate change and agricultural adaptation on nutrient loading from Finnish
5667 catchments to the Baltic Sea, Sci. Total Environ., 529, 168-181, <https://doi.org/10.1016/j.scitotenv.2015.05.055>,
5668 2015.

5669 Hyytiäinen, K., Bauer, B., Bly Joyce, K., Ehrnsten, E., Eilola, K., Gustafsson, B. G., Meier, H. E. M., Norkko,
5670 A., Saraiva, S., Tomczak, M., and Zandersen, M.: Provision of aquatic ecosystem services as a consequence of
5671 societal changes: The case of the Baltic Sea, Popul. Ecol., 63, 61-74, <https://doi.org/10.1002/1438-390X.12033>,
5672 2021.

5673 Ilmatieteen laitos: Jäätalvi Itämerellä: <https://www.ilmatieteenlaitos.fi/jaatalvi-itamerella>, The ice cover of the
5674 Baltic 1961-2018. Public record of Finnish Meteorological Institute, access: 10.12.2020, 2020.

5675 [IPCC: Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth](#)
5676 [Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Solomon, S., Qin, D.,](#)
5677 [Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tigno, M., and Miller, H. L., Cambridge University Press,](#)
5678 [United Kingdom and New York, NY, USA, Cambridge, 2007.](#)

5679 IPCC: Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth
5680 Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Team, C. W., Pachauri, R. K.,
5681 and Meyer, L. A., Geneva, 2014a.

5682 IPCC: Climate Change 2013 – The Physical Science Basis: Working Group I Contribution to the Fifth
5683 Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Stocker, T. F., Qin, D.,
5684 Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M.,
5685 Cambridge University Press, Cambridge, 2014b.

5686 IPCC: IPCC Special Report on the Ocean and Cryosphere in a Changing Climate, edited by: Pörtner, H.-O.,
5687 Roberts, D. C., Masson-Delmotte, V., Zhai, P., Tignor, M., Poloczanska, E., Mintenbeck, K., Alegría, A.,
5688 Nicolai, M., Okem, A., Petzold, J., Rama, B., and Weyer, N. M., 2019a.

5689 IPCC: Climate Change and Land: an IPCC special report on climate change, desertification, land degradation,
5690 sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems, edited by:
5691 Shukla, P. R., Skea, J., Buendia, E. C., Masson-Delmotte, V., Pörtner, H.-O., Roberts, D. C., Zhai, P., Slade, R.,
5692 Connors, S., Diemen, R. v., Ferrat, M., Haughey, E., Luz, S., Neogi, S., Pathak, M., Petzold, J., Pereira, J. P.,
5693 Vyas, P., Huntley, E., Kissick, K., Belkacemi, M., and Malley, J., 2019b.

5694 IPCC: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth
5695 Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Masson-Delmotte, V., Zhai,
5696 P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M.,
5697 Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B.,
5698 Cambridge University Press, Cambridge, 2021.

5699 Irannezhad, M., Ronkanen, A.-K., and Kløve, B.: Wintertime climate factors controlling snow resource decline
5700 in Finland, *Int. J. Climatol.*, 36, 110-131, <https://doi.org/10.1002/joc.4332>, 2016.

5701 Ito, M., Scotti, M., Franz, M., Barboza, F. R., Buchholz, B., Zimmer, M., Guy-Haim, T., and Wahl, M.: Effects
5702 of temperature on carbon circulation in macroalgal food webs are mediated by herbivores, *Mar. Biol.*, 166, 158,
5703 <https://doi.org/10.1007/s00227-019-3596-z>, 2019.

5704 Iturbide, M., Fernández, J., Gutiérrez, J. M., Bedia, J., Gimadevilla, E., Díez-Sierra, J., Manzanar, R., Casanueva,
5705 A., Baño-Medina, J., Milovac, J., Herrera, S., Cofiño, A. S., San Martín, D., García-Díez, M., Hauser, M., Huard,
5706 D., Yelekçi, Ö. (2021) Repository supporting the implementation of FAIR principles in the IPCC-WG1 Atlas.
5707 Zenodo, <https://doi.org/10.5281/zenodo.3691645>. Available from: <https://github.com/IPCC-WG1/Atlas>, access:
5708 25.11.2021.

5709 Ivanov, V. A., and Belokopytov, V. N.: Oceanography of the Black Sea, National Academy of Sciences of
5710 Ukraine, Marine Hydrophysical Institute, Sevastopol, Ukraine, 2013.

5711 Jaagus, J., Briede, A., Rimkus, E., and Remm, K.: Variability and trends in daily minimum and maximum
5712 temperatures and in the diurnal temperature range in Lithuania, Latvia and Estonia in 1951–2010, *Theor. Appl.*
5713 *Climatol.*, 118, 57-68, <https://doi.org/10.1007/s00704-013-1041-7>, 2014.

5714 Jaagus, J., Sepp, M., Tamm, T., Järvet, A., and Mõisja, K.: Trends and regime shifts in climatic conditions and
5715 river runoff in Estonia during 1951–2015, *Earth Syst. Dynam.*, 8, 963-976, [https://doi.org/10.5194/esd-8-963-](https://doi.org/10.5194/esd-8-963-2017)
5716 2017, 2017.

5717 Jaagus, J., Briede, A., Rimkus, E., and Sepp, M.: Changes in precipitation regime in the Baltic countries in
5718 1966–2015, *Theor. Appl. Climatol.*, 131, 433-443, <https://doi.org/10.1007/s00704-016-1990-8>, 2018.

5719 Jackson, L. C., Kahana, R., Graham, T., Ringer, M. A., Woollings, T., Mecking, J. V., and Wood, R. A.: Global
5720 and European climate impacts of a slowdown of the AMOC in a high resolution GCM, *Clim. Dyn.*, 45, 3299-
5721 3316, <https://doi.org/10.1007/s00382-015-2540-2>, 2015.

5722 Jacob, D., Kotova, L., Teichmann, C., Sobolowski, S. P., Vautard, R., Donnelly, C., Koutroulis, A. G., Grillakis,
5723 M. G., Tsanis, I. K., Damm, A., Sakalli, A., and van Vliet, M. T. H.: Climate Impacts in Europe Under +1.5°C
5724 Global Warming, *Earths Future*, 6, 264-285, <https://doi.org/10.1002/2017ef000710>, 2018.

5725 Jacob, D. J., and Winner, D. A.: Effect of climate change on air quality, *Atmos. Environ.*, 43, 51-63,
5726 <https://doi.org/10.1016/j.atmosenv.2008.09.051>, 2009.

5727 Jakimavičius, D., Kriaučiūnienė, J., and Šarauskienė, D.: Impact of climate change on the Curonian Lagoon
5728 water balance components, salinity and water temperature in the 21st century, *Oceanologia*, 60, 378-389,
5729 <https://doi.org/10.1016/j.oceano.2018.02.003>, 2018.

Formatiert: Englisch (USA)

Formatiert: Rechtschreibung und Grammatik nicht prüfen

5730 Jakobsson, M., Long, A., Ingólfsson, Ó., Kjær, K. H., and Spielhagen, R. F.: New insights on Arctic Quaternary
5731 climate variability from palaeo-records and numerical modelling, *Quaternary Sci. Rev.*, 29, 3349-3358,
5732 <https://doi.org/10.1016/j.quascirev.2010.08.016>, 2010.

5733 Jakubowska, M., Jerzak, M., Normant, M., Burska, D., and Drzazgowski, J.: Effect of Carbon Dioxide-Induced
5734 Water Acidification on the Physiological Processes of the Baltic Isopod *Saduria entomon*, *J. Shellfish Res.*, 32,
5735 825-834, 810, <https://doi.org/10.2983/035.032.0326>, 2013.

5736 Jänes, H., Herkül, K., and Kotta, J.: Environmental niche separation between native and non-native benthic
5737 invertebrate species: Case study of the northern Baltic Sea, *Mar. Environ. Res.*, 131, 123-133,
5738 <https://doi.org/10.1016/j.marenvres.2017.08.001>, 2017.

5739 Jansen, J., Thornton, B. F., Jammet, M. M., Wik, M., Cortés, A., Friberg, T., MacIntyre, S., and Crill, P. M.:
5740 Climate-Sensitive Controls on Large Spring Emissions of CH₄ and CO₂ From Northern Lakes, *J. Geophys. Res.-*
5741 *Biogeo.*, 124, 2379-2399, <https://doi.org/10.1029/2019JG005094>, 2019.

5742 Jansson, A., Lischka, S., Boxhammer, T., Schulz, K. G., and Norkko, J.: Survival and settling of larval *Macoma*
5743 *balthica* in a large-scale mesocosm experiment at different fCO₂ levels, *Biogeosciences*, 13, 3377-3385,
5744 <https://doi.org/10.5194/bg-13-3377-2016>, 2016.

5745 Jin, H., Jönsson, A. M., Olsson, C., Lindström, J., Jönsson, P., and Eklundh, L.: New satellite-based estimates
5746 show significant trends in spring phenology and complex sensitivities to temperature and precipitation at
5747 northern European latitudes, *Int. J. Biometeorol.*, 63, 763-775, <https://doi.org/10.1007/s00484-019-01690-5>,
5748 2019.

5749 Johansson, L., Ytreberg, E., Jalkanen, J. P., Fridell, E., Eriksson, K. M., Lagerström, M., Maljutenko, I.,
5750 Raudsepp, U., Fischer, V., and Roth, E.: Model for leisure boat activities and emissions – implementation for the
5751 Baltic Sea, *Ocean Sci.*, 16, 1143-1163, <https://doi.org/10.5194/os-16-1143-2020>, 2020.

5752 Johansson, M. M., Pellikka, H., Kahma, K. K., and Ruosteenoja, K.: Global sea level rise scenarios adapted to
5753 the Finnish coast, *J. Mar. Syst.*, 129, 35-46, <https://doi.org/10.1016/j.jmarsys.2012.08.007>, 2014.

5754 Jones, M. C., and Cheung, W. W. L.: Multi-model ensemble projections of climate change effects on global
5755 marine biodiversity, *ICES J. Mar. Sci.*, 72, 741-752, <https://doi.org/10.1093/icesjms/fsu172>, 2014.

5756 Jones, P. D., Lister, D. H., Osborn, T. J., Harpham, C., Salmon, M., and Morice, C. P.: Hemispheric and large-
5757 scale land-surface air temperature variations: An extensive revision and an update to 2010, *J. Geophys. Res.-*
5758 *Atmos.*, 117, <https://doi.org/10.1029/2011jd017139>, 2012.

5759 Jonson, J. E., Jalkanen, J. P., Johansson, L., Gauss, M., and van der Gon, H. A. C. D.: Model calculations of the
5760 effects of present and future emissions of air pollutants from shipping in the Baltic Sea and the North Sea,
5761 *Atmos. Chem. Phys.*, 15, 783-783, 2015.

5762 Jonson, J. E., Gauss, M., Jalkanen, J.-P., and Johansson, L.: Effects of strengthening the Baltic Sea ECA
5763 regulations, *Atmos. Chem. Phys.*, 19, 2019.

5764 Jönsson, P. R., Kotta, J., Andersson, H. C., Herkül, K., Virtanen, E., Sandman, A. N., and Johannesson, K.: High
5765 climate velocity and population fragmentation may constrain climate-driven range shift of the key habitat former
5766 *Fucus vesiculosus*, *Divers. Distrib.*, 24, 892-905, <https://doi.org/10.1111/ddi.12733>, 2018.

5767 Jönsson, P. R., Moksnes, P.-O., Corell, H., Bonsdorff, E., and Nilsson Jacobi, M.: Ecological coherence of
5768 Marine Protected Areas: New tools applied to the Baltic Sea network, *Aquat. Conserv.*, 30, 743-760,
5769 <https://doi.org/10.1002/aqc.3286>, 2020.

5770 Josefsson, W.: Long-term global radiation in Stockholm, 1922-2018, SMHI (Swedish Meteorological and
5771 Hydrological Institute), Norrköping, Sweden, 92pp, 2019.

5772 [Joshi, M. M., Gregory, J. M., Webb, M. J., Sexton, D. M. H., and Johns, T. C.: Mechanisms for the land/sea](https://doi.org/10.1007/s00382-007-0306-1)
5773 [warming contrast exhibited by simulations of climate change, *Clim. Dyn.*, 30, 455-465,](https://doi.org/10.1007/s00382-007-0306-1)
5774 <https://doi.org/10.1007/s00382-007-0306-1>, 2008.

5775 Jüssi, M., Härkönen, T., Helle, E., and Jüssi, I.: Decreasing Ice Coverage Will Reduce the Breeding Success of
5776 Baltic Grey Seal (*Halichoerus grypu*) Females, *Ambio* 37, 80-85, 86, 2008.

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Feldfunktion geändert

Formatiert: Englisch (USA)

Formatiert: Rechtschreibung und Grammatik nicht prüfen

5777 Jüssi, M.: Living on an edge: land-locked seals in changing climate, 2012.

5778 Jylhä, K., Laapas, M., Ruosteenoja, K., Arvola, L., Drebs, A., Kersalo, J., Saku, S., Gregow, H., Hannula, H.-R.,
5779 and Pirinen, P. J. B. E. R.: Climate variability and trends in the Valkea-Kotinen region, southern Finland:
5780 comparisons between the past, current and projected climates, *Boreal Environ. Res.*, 19, 4-30, 2014.

5781 Kadin, M., Frederiksen, M., Niiranen, S., and Converse, S. J.: Linking demographic and food-web models to
5782 understand management trade-offs, *Ecol. Evol.*, 9, 8587-8600, <https://doi.org/10.1002/ece3.5385>, 2019.

5783 Kahru, M., and Elmgren, R.: Multidecadal time series of satellite-detected accumulations of cyanobacteria in the
5784 Baltic Sea, *Biogeosciences*, 11, 3619-3633, <https://doi.org/10.5194/bg-11-3619-2014>, 2014.

5785 Kahru, M., Elmgren, R., and Savchuk, O. P.: Changing seasonality of the Baltic Sea, *Biogeosciences*, 13, 1009-
5786 1018, <https://doi.org/10.5194/bg-13-1009-2016>, 2016.

5787 Kahru, M., Elmgren, R., Di Lorenzo, E., and Savchuk, O.: Unexplained interannual oscillations of
5788 cyanobacterial blooms in the Baltic Sea, *Sci. Rep.-UK*, 8, 6365, <https://doi.org/10.1038/s41598-018-24829-7>,
5789 2018.

5790 Kahru, M., Elmgren, R., Kaiser, J., Wasmund, N., and Savchuk, O.: Cyanobacterial blooms in the Baltic Sea:
5791 Correlations with environmental factors, *Harmful Algae*, 92, 101739, <https://doi.org/10.1016/j.hal.2019.101739>,
5792 2020.

5793 Kanakidou, M., Myriokefalitakis, S., and Tsigaridis, K.: Aerosols in atmospheric chemistry and biogeochemical
5794 cycles of nutrients, *Environ. Res. Lett.*, 13, 063004, <https://doi.org/10.1088/1748-9326/aabddb>, 2018.

5795 Karl, M., Bieser, J., Geyer, B., Matthias, V., Jalkanen, J.-P., Johansson, L., and Fridell, E.: Impact of a nitrogen
5796 emission control area (NECA) on the future air quality and nitrogen deposition to seawater in the Baltic Sea
5797 region, *Atmos. Chem. Phys.*, 19, 1721-1752, <https://doi.org/10.5194/acp-2018-1107>, 2019a.

5798 Karl, M., Jonson, J. E., Uppstu, A., Aulinger, A., Prank, M., Sofiev, M., Jalkanen, J. P., Johansson, L., Quante,
5799 M., and Matthias, V.: Effects of ship emissions on air quality in the Baltic Sea region simulated with three
5800 different chemistry transport models, *Atmos. Chem. Phys.*, 19, 7019-7053, <https://doi.org/10.5194/acp-19-7019-2019>,
5801 2019b.

5802 Karlsson, K.-G., and Devasthale, A.: Inter-comparison and evaluation of the four longest satellite-derived cloud
5803 climate data records: CLARA-A2, ESA Cloud CCI V3, ISCCP-HGM, and PATMOS-x, *Remote Sens.-Basel*, 10,
5804 1567, <https://doi.org/10.3390/rs10101567>, 2018.

5805 Kauhala, K., Bäcklin, B.-M., Raitaniemi, J., and Harding, K. C.: The effect of prey quality and ice conditions on
5806 the nutritional status of Baltic grey seals of different age groups, *Mammal Res.*, 62, 351-362,
5807 <https://doi.org/10.1007/s13364-017-0329-x>, 2017.

5808 Kauhala, K., Bergenius, M., Isomursu, M., and Raitaniemi, J.: Reproductive rate and nutritional status of Baltic
5809 ringed seals, *Mammal Res.*, 64, 109-120, 2019.

5810 Kauhala, K., and Kurkilahti, M.: Delayed effects of prey fish quality and winter temperature during the birth
5811 year on adult size and reproductive rate of Baltic grey seals, *Mammal Res.*, 65, 117-126,
5812 <https://doi.org/10.1007/s13364-019-00454-1>, 2020.

5813 Kauker, F., and Meier, H. E. M.: Modeling decadal variability of the Baltic Sea: 1. Reconstructing atmospheric
5814 surface data for the period 1902-1998, *J. Geophys. Res.-Oceans*, 108, <https://doi.org/10.1029/2003JC001797>,
5815 2003.

5816 Kiani, S., Irannezhad, M., Ronkanen, A.-K., Moradkhani, H., and Kløve, B.: Effects of recent temperature
5817 variability and warming on the Oulu-Hailuoto ice road season in the northern Baltic Sea, *Cold Reg. Sci.*
5818 *Technol.*, 151, 1-8, <https://doi.org/10.1016/j.coldregions.2018.02.010>, 2018.

5819 Kim, B.-M., Son, S.-W., Min, S.-K., Jeong, J.-H., Kim, S.-J., Zhang, X., Shim, T., and Yoon, J.-H.: Weakening
5820 of the stratospheric polar vortex by Arctic sea-ice loss, *Nat. Commun.*, 5, 4646,
5821 <https://doi.org/10.1038/ncomms5646>, 2014.

5822 Kirchner, N., Noormets, R., Kutteneuler, J., Erstorp, E. S., Holmlund, E. S., Rosqvist, G., Holmlund, P.,
5823 Wennbom, M., and Karlin, T.: High-resolution bathymetric mapping reveals subaqueous glacial landforms in the
5824 Arctic alpine lake Tarfala, Sweden, *J. Quat. Sci.*, 34, 452-462, <https://doi.org/10.1002/jqs.3112>, 2019.

5825 Kirchner, N., Kutteneuler, K., Rosqvist, G., Hancke, M., Granebeck, A., Weckström, J., Weckström, K.,
5826 Schenk, F., Korhola, A., and Eriksson, P.: A first continuous three-year temperature record from the dimictic
5827 arctic-alpine Lake Tarfala, northern Sweden, *Arct. Antarct. Alp. Res.*, 2021.

5828 Kjellström, E., Barring, L., Nikulin, G., Nilsson, C., Persson, G., and Strandberg, G.: Production and use of
5829 regional climate model projections – A Swedish perspective on building climate services, *Clim. Serv.*, 2-3, 15-
5830 29, <https://doi.org/10.1016/j.cliser.2016.06.004>, 2016.

5831 Kjellström, E., Nikulin, G., Strandberg, G., Christensen, O. B., Jacob, D., Keuler, K., Lenderink, G., van
5832 Meijgaard, E., Schär, C., and Somot, S.: European climate change at global mean temperature increases of 1.5
5833 and 2 degrees C above pre-industrial conditions as simulated by the EURO-CORDEX regional climate models,
5834 *Earth Syst. Dynam.*, 9, 459-478, <https://doi.org/10.5194/esd-9-459-2018>, 2018.

5835 Kjerfve, B.: Oceanography of Chesapeake Bay, in: *Hydrodynamics of Estuaries*, edited by: Kjerfve, B., Taylor
5836 & Francis group, CRC Press, 1988.

5837 Klais, R., Tamminen, T., Kremp, A., Spilling, K., and Olli, K.: Decadal-scale changes of dinoflagellates and
5838 diatoms in the anomalous Baltic Sea spring bloom, *PLOS ONE* 6, e21567,
5839 <https://doi.org/10.1371/journal.pone.0021567>, 2011.

5840 Klais, R., Tamminen, T., Kremp, A., Spilling, K., An, B. W., Hajdu, S., and Olli, K.: Spring phytoplankton
5841 communities shaped by interannual weather variability and dispersal limitation: Mechanisms of climate change
5842 effects on key coastal primary producers, *Limnol. Oceanogr.*, 58, 753-762,
5843 <https://doi.org/10.4319/lo.2013.58.2.0753>, 2013.

5844 Kļaviņš, M., Avotniece, Z., and Rodinovs, V.: Dynamics and Impacting Factors of Ice Regimes in Latvia Inland
5845 and Coastal Waters, *Proceedings of the Latvian Academy of Sciences. Section B. Natural, Exact, and Applied*
5846 *Sciences.*, 70, 400-408, <https://doi.org/10.1515/prolas-2016-0059>, 2016.

5847 Klawonn, I., Lavik, G., Böning, P., Marchant, H., Dekaezemacker, J., Mohr, W., and Ploug, H.: Simple approach
5848 for the preparation of 15–15N₂-enriched water for nitrogen fixation assessments: evaluation, application and
5849 recommendations, *Front. Microbiol.*, 6, <https://doi.org/10.3389/fmicb.2015.00769>, 2015.

5850 Kniebusch, M., Meier, H. E. M., Neumann, T., and Börgel, F.: Temperature Variability of the Baltic Sea Since
5851 1850 and Attribution to Atmospheric Forcing Variables, *J. Geophys. Res.-Oceans*, 124, 4168-4187,
5852 <https://doi.org/10.1029/2018jc013948>, 2019a.

5853 Kniebusch, M., Meier, H. E. M., and Radtke, H.: Changing Salinity Gradients in the Baltic Sea As a
5854 Consequence of Altered Freshwater Budgets, *Geophys. Res. Lett.*, 46, 9739-9747,
5855 <https://doi.org/10.1029/2019GL083902>, 2019b.

5856 Knight, J. R., Folland, C. K., and Scaife, A. A.: Climate impacts of the Atlantic Multidecadal Oscillation,
5857 *Geophys. Res. Lett.*, 33, <https://doi.org/10.1029/2006gl026242>, 2006.

5858 Knight, J. R.: The Atlantic Multidecadal Oscillation Inferred from the Forced Climate Response in Coupled
5859 General Circulation Models, *J. Climate*, 22, 1610-1625, <https://doi.org/10.1175/2008jcli2628.1>, 2009.

5860 Knoll, L. B., Sharma, S., Denfeld, B. A., Flaim, G., Hori, Y., Magnuson, J. J., Straile, D., and Weyhenmeyer, G.
5861 A.: Consequences of lake and river ice loss on cultural ecosystem services, *Limnol. Oceanogr. Lett.*, 4, 119-131,
5862 <https://doi.org/10.1002/lol2.10116>, 2019.

5863 Knudsen, M.: Ein hydrographischer lehrsatz, *Ann. Hydr. u. mar. Meteor.*, 28, 316-320, 1900.

5864 Knudsen, M. F., Seidenkrantz, M.-S., Jacobsen, B. H., and Kuijpers, A.: Tracking the Atlantic Multidecadal
5865 Oscillation through the last 8,000 years, *Nat. Commun.*, 2, 178, <https://doi.org/10.1038/ncomms1186>, 2011.

5866 Kong, D., MacLeod, M., and Cousins, I. T.: Modelling the influence of climate change on the chemical
5867 concentrations in the Baltic Sea region with the POPCYCLING-Baltic model, *Chemosphere*, 110, 31-40,
5868 <https://doi.org/10.1016/j.chemosphere.2014.02.044>, 2014.

5869 Kopp, R. E., Horton, R. M., Little, C. M., Mitrovica, J. X., Oppenheimer, M., Rasmussen, D. J., Strauss, B. H.,
5870 and Tebaldi, C.: Probabilistic 21st and 22nd century sea-level projections at a global network of tide-gauge sites,
5871 *Earths Future*, 2, 383-406, <https://doi.org/10.1002/2014EF000239>, 2014.

5872 Korhonen, J.: Long-term changes and variability of the winter and spring season hydrological regime in Finland,
5873 Faculty of Science, Institute for Atmospheric and Earth System Research, University of Helsinki, Finland, 2019.

5874 Korpinen, S., Honkanen, T., Vesakoski, O., Hemmi, A., Koivikko, R., Loponen, J., and Jormalainen, V.:
5875 Macroalgal Communities Face the Challenge of Changing Biotic Interactions: Review with Focus on the Baltic
5876 Sea, *Ambio* 36, 203-211, 209, [https://doi.org/10.1579/0044-7447\(2007\)36\[203:MCFTCO\]2.0.CO;2](https://doi.org/10.1579/0044-7447(2007)36[203:MCFTCO]2.0.CO;2), 2007.

5877 Kortsch, S., Frelat, R., Pecuchet, L., Olivier, P., Putnis, I., Bonsdorff, E., Ojaveer, H., Jurgensone, I., Sträke, S.,
5878 Rubene, G., Krüze, Ē., and Nordström, M. C.: Disentangling temporal food web dynamics facilitates
5879 understanding of ecosystem functioning, *J. Anim. Ecol.*, <https://doi.org/10.1111/1365-2656.13447>, 2021.

5880 Köster, F. W., Huwer, B., Hinrichsen, H.-H., Neumann, V., Makarchouk, A., Eero, M., Dewitz, B. V., Hüsey,
5881 K., Tomkiewicz, J., Margonski, P., Temming, A., Hermann, J.-P., Oesterwind, D., Dierking, J., Kotterba, P., and
5882 Plikshs, M.: Eastern Baltic cod recruitment revisited—dynamics and impacting factors, *ICES J. Mar. Sci.*, 74, 3-
5883 19, <https://doi.org/10.1093/icesjms/fsw172>, 2016.

5884 Kotta, J., Vanhatalo, J., Jänes, H., Orav-Kotta, H., Rugiu, L., Jormalainen, V., Bobsien, I., Viitasalo, M.,
5885 Virtanen, E., Sandman, A. N., Isaeus, M., Leidenberger, S., Jonsson, P. R., and Johannesson, K.: Integrating
5886 experimental and distribution data to predict future species patterns, *Sci. Rep.-UK*, 9, 1821,
5887 <https://doi.org/10.1038/s41598-018-38416-3>, 2019.

5888 [Kreienkamp, F., Lorenz, P., and Geiger, T.: Statistically downscaled CMIP6 projections show stronger warming
5889 for Germany, *Atmosphere*, 11, 1245, <https://doi.org/10.3390/atmos11111245>, 2020.](#)

5890 Kremp, A., Godhe, A., Egardt, J., Dupont, S., Suikkanen, S., Casabianca, S., and Penna, A.: Intraspecific
5891 variability in the response of bloom-forming marine microalgae to changed climate conditions, *Ecol. Evol.*, 2,
5892 1195-1207, <https://doi.org/10.1002/ece3.245>, 2012.

5893 Kremp, A., Oja, J., LeTortorec, A. H., Hakanen, P., Tahvanainen, P., Tuimala, J., and Suikkanen, S.: Diverse
5894 seed banks favour adaptation of microalgal populations to future climate conditions, *Environ. Microbiol.*, 18,
5895 679-691, <https://doi.org/10.1111/1462-2920.13070>, 2016.

5896 Krug, J., Eriksson, H., Heidecke, C., Kellomäki, S., Köhl, M., Lindner, M., and Saikkonen, K.: Socio-economic
5897 Impacts—Forestry and Agriculture, in: Second Assessment of Climate Change for the Baltic Sea Basin, edited
5898 by: BACC II Author Team, Springer International Publishing, Cham, 399-409, https://doi.org/10.1007/978-3-319-16006-1_21, 2015.

5900 Kudryavtseva, N., and Soomere, T.: Satellite altimetry reveals spatial patterns of variations in the Baltic Sea
5901 wave climate, *Earth Syst. Dynam.*, 8, 697-706, <https://doi.org/10.5194/esd-8-697-2017>, 2017.

5902 Köhl, N., Gebhardt, C., Litt, T., and Hense, A.: Probability Density Functions as Botanical-Climatological
5903 Transfer Functions for Climate Reconstruction, *Quat. Res.*, 58, 381-392, <https://doi.org/10.1006/qres.2002.2380>,
5904 2002.

5905 Kuliński, K., Schneider, B., Hammer, K., Machulik, U., and Schulz-Bull, D.: The influence of dissolved organic
5906 matter on the acid–base system of the Baltic Sea, *J. Mar. Syst.*, 132, 106-115,
5907 <https://doi.org/10.1016/j.jmarsys.2014.01.011>, 2014.

5908 Kuliński, K., Hammer, K., Schneider, B., and Schulz-Bull, D.: Remineralization of terrestrial dissolved organic
5909 carbon in the Baltic Sea, *Mar. Chem.*, 181, 10-17, <https://doi.org/10.1016/j.marchem.2016.03.002>, 2016.

5910 Kuliński, K., Schneider, B., Szymczycha, B., and Stokowski, M.: Structure and functioning of the acid–base
5911 system in the Baltic Sea, *Earth Syst. Dynam.*, 8, 1107-1120, <https://doi.org/10.5194/esd-8-1107-2017>, 2017.

5912 Kuliński, K., Szymczycha, B., Kozirowska, K., Hammer, K., and Schneider, B.: Anomaly of total boron
5913 concentration in the brackish waters of the Baltic Sea and its consequence for the CO₂ system calculations, *Mar.*
5914 *Chem.*, 204, 11-19, <https://doi.org/10.1016/j.marchem.2018.05.007>, 2018.

Formatiert: Standard, Abstand Nach: 12 Pt.

5915 Kuliński, K., Rehder, G., Asmala, E., Bartosova, A., Carstensen, J., Gustafsson, B., Hall, P. O. J., Humborg, C.,
5916 Jilbert, T., Jürgens, K., Meier, H. E. M., Müller-Karulis, B., Naumann, M., Olesen, J. E., Savchuk, O., Schramm,
5917 A., Slomp, C. P., Sofiev, M., Sobek, A., Szymczycha, B., and Undeman, E.: Baltic Earth Assessment Report on
5918 the biogeochemistry of the Baltic Sea, *Earth Syst. Dynam. Discuss.*, 1-93, <https://doi.org/10.5194/esd-2021-33>,
5919 2021.

5920 Kundzewicz, Z. W., Ulbrich, U., Brücher, T., Graczyk, D., Krüger, A., Leckebusch, G. C., Menzel, L., Pińskwar,
5921 I., Radziejewski, M., and Szwed, M.: Summer Floods in Central Europe – Climate Change Track?, *Nat. Hazards*,
5922 36, 165-189, <https://doi.org/10.1007/s11069-004-4547-6>, 2005.

5923 Kuosa, H., Fleming-Lehtinen, V., Lehtinen, S., Lehtiniemi, M., Nygård, H., Raateoja, M., Raitaniemi, J.,
5924 Tuimala, J., Uusitalo, L., and Suikkanen, S.: A retrospective view of the development of the Gulf of Bothnia
5925 ecosystem, *J. Mar. Syst.*, 167, 78-92, <https://doi.org/10.1016/j.jmarsys.2016.11.020>, 2017.

5926 Kuznetsov, I., and Neumann, T.: Simulation of carbon dynamics in the Baltic Sea with a 3D model, *J. Mar.*
5927 *Syst.*, 111-112, 167-174, <https://doi.org/10.1016/j.jmarsys.2012.10.011>, 2013.

5928 [Kwiatkowski, L., Torres, O., Bopp, L., Aumont, O., Chamberlain, M., Christian, J. R., Dunne, J. P., Gehlen, M.,](#)
5929 [Ilyina, T., John, J. G., Lenton, A., Li, H., Lovenduski, N. S., Orr, J. C., Palmieri, J., Santana-Falcón, Y.,](#)
5930 [Schwinger, J., Séférian, R., Stock, C. A., Tagliabue, A., Takano, Y., Tjiputra, J., Toyama, K., Tsujino, H.,](#)
5931 [Watanabe, M., Yamamoto, A., Yool, A., and Ziehn, T.: Twenty-first century ocean warming, acidification,](#)
5932 [deoxygenation, and upper-ocean nutrient and primary production decline from CMIP6 model projections,](#)
5933 [Biogeosciences](#), 17, 3439-3470, <https://doi.org/10.5194/bg-17-3439-2020>, 2020.

5934 Laakso, L., Mikkonen, S., Drebs, A., Karjalainen, A., Pirinen, P., and Alenius, P.: 100 years of atmospheric and
5935 marine observations at the Finnish Utö Island in the Baltic Sea, *Ocean Sci.*, 14, 617-632,
5936 <https://doi.org/10.5194/os-14-617-2018>, 2018.

5937 Łabuz, T. A.: Environmental Impacts—Coastal Erosion and Coastline Changes, in: Second Assessment of
5938 Climate Change for the Baltic Sea Basin, edited by: BACC II Author Team, Springer International Publishing,
5939 Cham, 381-396, https://doi.org/10.1007/978-3-319-16006-1_20, 2015.

5940 Łabuz, T. A., Grunewald, R., Bobykina, V., Chubarenko, B., Česnulevičius, A., Bautrėnas, A., Morkūnaitė, R.,
5941 and Tõnisson, H.: Coastal Dunes of the Baltic Sea Shores: A Review, *Quaest. Geogr.*, 37, 47-71,
5942 <https://doi.org/10.2478/quageo-2018-0005>, 2018.

5943 Lah, L., Trense, D., Benke, H., Berggren, P., Gunnlaugsson, Þ., Lockyer, C., Öztürk, A., Öztürk, B., Pawliczka,
5944 I., Roos, A., Siebert, U., Skóra, K., Víkingsson, G., and Tiedemann, R.: Spatially Explicit Analysis of Genome-
5945 Wide SNPs Detects Subtle Population Structure in a Mobile Marine Mammal, the Harbor Porpoise, *PLoS ONE*,
5946 11, e0162792, <https://doi.org/10.1371/journal.pone.0162792>, 2016.

5947 Lamon, L., von Waldow, H., MacLeod, M., Scheringer, M., Marcomini, A., and Hungerbühler, K.: Modeling the
5948 Global Levels and Distribution of Polychlorinated Biphenyls in Air under a Climate Change Scenario, *Environ.*
5949 *Sci. Technol.*, 43, 5818-5824, <https://doi.org/10.1021/es900438j>, 2009.

5950 Lampe, M., and Lampe, R.: Evolution of a large Baltic beach ridge plain (Neudarss, NE Germany): A
5951 continuous record of sea-level and wind-field variation since the Homeric Minimum, *Earth Surf. Process. Landf.*,
5952 43, 3042-3056, <https://doi.org/10.1002/esp.4468>, 2018.

5953 Landerer, F. W., Jungclauss, J. H., and Marotzke, J.: Regional dynamic and steric sea level change in response to
5954 the IPCC-A1B scenario, *J. Phys. Oceanogr.*, 37, 296-312, <https://doi.org/10.1175/JPO3013.1>, 2007.

5955 Lang, A., and Mikolajewicz, U.: The long-term variability of extreme sea levels in the German Bight, *Ocean*
5956 *Sci.*, 15, 651-668, 2019.

5957 Langner, J., Engardt, M., Baklanov, A., Christensen, J. H., Gauss, M., Geels, C., Hedegaard, G. B., Nuterman,
5958 R., Simpson, D., and Soares, J.: A multi-model study of impacts of climate change on surface ozone in Europe,
5959 *Atmos. Chem. Phys.*, 12, 10423-10440, 2012.

5960 Lappe, C., and Umlauf, L.: Efficient boundary mixing due to near-inertial waves in a nontidal basin:
5961 Observations from the Baltic Sea, *J. Geophys. Res-Oceans*, 121, 8287-8304,
5962 <https://doi.org/10.1002/2016JC011985>, 2016.

5963 Larsson, K., Hajdu, S., Kilpi, M., Larsson, R., Leito, A., and Lyngs, P.: Effects of an extensive *Prymnesium*
5964 *polylepis* bloom on breeding eiders in the Baltic Sea, *J. Sea Res.*, 88, 21-28,
5965 <https://doi.org/10.1016/j.seares.2013.12.017>, 2014.

~~5966 Larsson, U., Elmgren, R., and Wulff, F.: Eutrophication and the Baltic Sea: causes and consequences;~~
5967 ~~L'eutrophisation et la mer Baltique: causes et conséquences, *Ambio*, 14, 9-14, 1985-Larsson, U., Elmgren, R.,~~
5968 ~~and Wulff, F.: Eutrophication and the Baltic Sea: causes and consequences, *Ambio*, 14, 9-14, 1985.~~

5969 Larsson, U., Hajdu, S., Walve, J., and Elmgren, R.: Baltic Sea nitrogen fixation estimated from the summer
5970 increase in upper mixed layer total nitrogen, *Limnol. Oceanogr.*, 46, 811-820,
5971 <https://doi.org/10.4319/lo.2001.46.4.0811>, 2001.

5972 Laskar, J., Robutel, P., Joutel, F., Gastineau, M., Correia, A. C. M., and Levrard, B.: A long-term numerical
5973 solution for the insolation quantities of the Earth, *Astron. Astrophys.*, 428, 261-285,
5974 <https://doi.org/10.1051/0004-6361:20041335>, 2004.

5975 Lass, H. U., and Matthäus, W.: On temporal wind variations forcing salt water inflows into the Baltic Sea, *Tellus*
5976 *A*, 48, 663-671, <https://doi.org/10.1034/j.1600-0870.1996.t01-4-00005.x>, 1996.

5977 Le Cozannet, G., Nicholls, R. J., Hinkel, J., Sweet, W. V., McInnes, K. L., Van de Wal, R. S. W., Slangen, A. B.
5978 A., Lowe, J. A., and White, K. D.: Sea Level Change and Coastal Climate Services: The Way Forward, *J. Mar.*
5979 *Sci. Eng.*, 5, <https://doi.org/10.3390/jmse5040049>, 2017.

5980 Leckebusch, G. C., and Ulbrich, U.: On the relationship between cyclones and extreme windstorm events over
5981 Europe under climate change, *Global Planet. Change*, 44, 181-193,
5982 <https://doi.org/10.1016/j.gloplacha.2004.06.011>, 2004.

5983 Lee, X., Goulden, M. L., Hollinger, D. Y., Barr, A., Black, T. A., Bohrer, G., Bracho, R., Drake, B., Goldstein,
5984 A., Gu, L., Katul, G., Kolb, T., Law, B. E., Margolis, H., Meyers, T., Monson, R., Munger, W., Oren, R., Paw U,
5985 K. T., Richardson, A. D., Schmid, H. P., Staebler, R., Wofsy, S., and Zhao, L.: Observed increase in local
5986 cooling effect of deforestation at higher latitudes, *Nature*, 479, 384-387, <https://doi.org/10.1038/nature10588>,
5987 2011.

5988 Lehtikoinen, A., Kilpi, M., and Öst, M.: Winter climate affects subsequent breeding success of common eiders,
5989 *Global Change Biol.*, 12, 1355-1365, <https://doi.org/10.1111/j.1365-2486.2006.01162.x>, 2006.

5990 Lehtikoinen, A., and Jaatinen, K.: Delayed autumn migration in northern European waterfowl, *J. Ornithol.*, 153,
5991 563-570, 2012.

5992 Lehtikoinen, A., Jaatinen, K., Vähätalo, A. V., Clausen, P., Crowe, O., Deceuninck, B., Hearn, R., Holt, C. A.,
5993 Hornman, M., Keller, V., Nilsson, L., Langendoen, T., Tománková, I., Wahl, J., and Fox, A. D.: Rapid climate
5994 driven shifts in wintering distributions of three common waterbird species, *Global Change Biol.*, 19, 2071-2081,
5995 <https://doi.org/10.1111/gcb.12200>, 2013.

5996 Lehmann, A., Krauß, W., and Hinrichsen, H.-H.: Effects of remote and local atmospheric forcing on circulation
5997 and upwelling in the Baltic Sea, *Tellus A*, 54, 299-316, <https://doi.org/10.3402/tellusa.v54i3.12138>, 2002.

5998 Lehmann, A., Getzlaff, K., and Harlaß, J.: Detailed assessment of climate variability in the Baltic Sea area for
5999 the period 1958 to 2009, *Clim. Res.*, 46, 185-196, <https://doi.org/10.3354/cr00876>, 2011.

6000 Lehmann, A., Myrberg, K., and Höflich, K.: A statistical approach to coastal upwelling in the Baltic Sea based
6001 on the analysis of satellite data for 1990–2009, *Oceanologia*, 54, 369-393, <https://doi.org/10.5697/oc.54-3.369>,
6002 2012.

6003 Lehmann, A., and Post, P.: Variability of atmospheric circulation patterns associated with large volume changes
6004 of the Baltic Sea, *Adv. Sci. Res.*, 12, 219-225, <https://doi.org/10.5194/asr-12-219-2015>, 2015.

6005 Lehmann, A., Höflich, K., Post, P., and Myrberg, K.: Pathways of deep cyclones associated with large volume
6006 changes (LVCs) and major Baltic inflows (MBIs), *J. Mar. Syst.*, 167, 11-18,
6007 <https://doi.org/10.1016/j.jmarsys.2016.10.014>, 2017.

6008 Lehmann, A., Myrberg, K., Post, P., Chubarenko, I., Dailidienė, I., Hinrichsen, H.-H., Hüsey, K., Liblik, T.,
6009 Lips, U., Meier, H. E. M., and Bukanova, T.: Salinity dynamics of the Baltic Sea, *Earth Syst. Dynam. Discuss.*,
6010 1-36, <https://doi.org/10.5194/esd-2021-15>, 2021.

6011 Lehtonen, I.: Four consecutive snow-rich winters in Southern Finland: 2009/2010–2012/2013, *Weather* 70, 3-8,
6012 <https://doi.org/10.1002/wea.2360>, 2015.

6013 Lehtoranta, J., Savchuk, O. P., Elken, J., Dahlbo, K., Kuosa, H., Raateoja, M., Kauppila, P., Räske, A., and
6014 Pitkänen, H.: Atmospheric forcing controlling inter-annual nutrient dynamics in the open Gulf of Finland, *J.*
6015 *Mar. Syst.*, 171, 4-20, <https://doi.org/10.1016/j.jmarsys.2017.02.001>, 2017.

6016 Leidenberger, S., De Giovanni, R., Kulawik, R., Williams, A. R., and Bourlat, S. J.: Mapping present and future
6017 potential distribution patterns for a meso-grazer guild in the Baltic Sea, *J. Biogeogr.*, 42, 241-254,
6018 <https://doi.org/10.1111/jbi.12395>, 2015.

6019 Lenggenhager, S., and Martius, O.: Atmospheric blocks modulate the odds of heavy precipitation events in
6020 Europe, *Clim. Dyn.*, 53, 4155-4171, <https://doi.org/10.1007/s00382-019-04779-0>, 2019.

6021 [Leppäranta, M.: Land-ice interaction in the Baltic Sea, *Est. J. Earth Sci.*, 62, 2013.](#)

6022 Leppäranta, M., and Myrberg, K.: Physical oceanography of the Baltic Sea, Springer Science & Business Media,
6023 Berlin, 2009.

6024 Leppäranta, M., Oikonen, A., Shirasawa, K., and Fukamachi, Y.: A treatise on frequency spectrum of drift ice
6025 velocity, *Cold Reg. Sci. Technol.*, 76, 83-91, <https://doi.org/10.1016/j.coldregions.2011.12.005>, 2012.

6026 Leppäranta, M.: Land-ice interaction in the Baltic Sea, *Est. J. Earth Sci.*, 62, 2013.

6027 Lépy, É., and Pasanen, L.: Observed Regional Climate Variability during the Last 50 Years in Reindeer Herding
6028 Cooperatives of Finnish Fell Lapland, *Climate*, 5, 81, <https://doi.org/10.3390/cli5040081>, 2017.

6029 Lewandowska, A. M., Breithaupt, P., Hillebrand, H., Hoppe, H.-G., Jürgens, K., and Sommer, U.: Responses of
6030 primary productivity to increased temperature and phytoplankton diversity, *J. Sea Res.*, 72, 87-93,
6031 <https://doi.org/10.1016/j.seares.2011.10.003>, 2012.

6032 Lewandowska, A. M., Boyce, D. G., Hofmann, M., Matthiessen, B., Sommer, U., and Worm, B.: Effects of sea
6033 surface warming on marine plankton, *Ecol. Lett.*, 17, 614-623, <https://doi.org/10.1111/ele.12265>, 2014.

6034 [Li, J., Huo, R., Chen, H., Zhao, Y., and Zhao, T.: Comparative Assessment and Future Prediction Using CMIP6
6035 and CMIP5 for Annual Precipitation and Extreme Precipitation Simulation, 9,
6036 <https://doi.org/10.3389/feart.2021.687976>, 2021.](#)

6037 Liblik, T., and Lips, U.: Stratification Has Strengthened in the Baltic Sea – An Analysis of 35 Years of
6038 Observational Data, *Front. Earth Sci.*, 7, <https://doi.org/10.3389/feart.2019.00174>, 2019.

6039 Lindegren, M., Möllmann, C., Nielsen, A., Brander, K., MacKenzie, B. R., and Stenseth, N. C.: Ecological
6040 forecasting under climate change: the case of Baltic cod, *P. Roy. Soc. B-Biol. Sci.*, 277, 2121-2130,
6041 <https://doi.org/10.1098/rspb.2010.0353>, 2010.

6042 Lindegren, M., Blenckner, T., and Stenseth, N. C.: Nutrient reduction and climate change cause a potential shift
6043 from pelagic to benthic pathways in a eutrophic marine ecosystem, *Global Change Biol.*, 18, 3491-3503,
6044 <https://doi.org/10.1111/j.1365-2486.2012.02799.x>, 2012.

6045 Lindeskog, M., Lagergren, F., Smith, B., and Rammig, A.: Accounting for forest management in the estimation
6046 of forest carbon balance using the dynamic vegetation model LPJ-GUESS (v4.0, r9333): Implementation and
6047 evaluation of simulations for Europe, *Geosci. Model Dev. Discuss.*, 2021, 1-42, <https://doi.org/10.5194/gmd-2020-440>, 2021.

6049 Lindroth, A., Holst, J., Linderson, M.-L., Aurela, M., Biermann, T., Heliasz, M., Chi, J., Ibrom, A., Kolari, P.,
6050 Klemetsson, L., Krasnova, A., Laurila, T., Lehner, I., Lohila, A., Mammarella, I., Mölder, M., Löfvenius, M.
6051 O., Peichl, M., Pilegaard, K., Soosaar, K., Vesala, T., Vestin, P., Weslien, P., and Nilsson, M.: Effects of drought
6052 and meteorological forcing on carbon and water fluxes in Nordic forests during the dry summer of 2018, *Philos.*
6053 *T. Roy. Soc. B.*, 375, 20190516, <https://doi.org/10.1098/rstb.2019.0516>, 2020.

Formatiert: Englisch (USA)

Formatiert: Rechtschreibung und Grammatik nicht prüfen

6054 Lindström, G.: Hydrologiska aspekter på åtgärder mot vattenbrist och torka inom avrinningsområden02837722
6055 (ISSN), 60pp, 2019.

6056 [Lindvall, J., and Svensson, G.: The diurnal temperature range in the CMIP5 models, *Clim. Dyn.*, 44, 405-421,
6057 <https://doi.org/10.1007/s00382-014-2144-2>, 2015.](https://doi.org/10.1007/s00382-014-2144-2)

6058 Lips, I., and Lips, U.: Abiotic factors influencing cyanobacterial bloom development in the Gulf of Finland
6059 (Baltic Sea), *Hydrobiologia*, 614, 133-140, <https://doi.org/10.1007/s10750-008-9449-2>, 2008.

6060 Liu, Y., Meier, H. E. M., and Eilola, K.: Nutrient transports in the Baltic Sea – results from a 30-year physical–
6061 biogeochemical reanalysis, *Biogeosciences*, 14, 2113-2131, <https://doi.org/10.5194/bg-14-2113-2017>, 2017.

6062 Liu, Y., Axell, L., Jandt, S., Lorkowski, I., Lindenthal, A., Verjovkina, S., and Schwichtenberg, F.: Baltic Sea
6063 Production Centre BALTICSEA_REANALYSIS_PHY_003_012 COPERNICUS Marine Environment
6064 Monitoring Service, Quality Information Document, 35pp, 2019.

6065 Liu, Z., Zhu, J., Rosenthal, Y., Zhang, X., Otto-Bliesner, B. L., Timmermann, A., Smith, R. S., Lohmann, G.,
6066 Zheng, W., and Elison Timm, O.: The Holocene temperature conundrum, *Proceedings of the National Academy
6067 of Sciences*, 111, E3501, <https://doi.org/10.1073/pnas.1407229111>, 2014.

6068 Lopez, L. S., Hewitt, B. A., and Sharma, S.: Reaching a breaking point: How is climate change influencing the
6069 timing of ice breakup in lakes across the northern hemisphere?, *Limnol. Oceanogr.*, 64, 2621-2631,
6070 <https://doi.org/10.1002/lno.11239>, 2019.

6071 Löptien, U., and Meier, H. E. M.: The influence of increasing water turbidity on the sea surface temperature in
6072 the Baltic Sea: a model sensitivity study, *J. Mar. Syst.*, 88, 323-331,
6073 <https://doi.org/10.1016/j.jmarsys.2011.06.001>, 2011.

6074 Ludwig, J.: Climate Signals in Coastal Deposits, Staats- und Universitätsbibliothek Hamburg, Hamburg, 2017.

6075 Łukawska-Matuszewska, K., and Graca, B.: Pore water alkalinity below the permanent halocline in the Gdańsk
6076 Deep (Baltic Sea) - Concentration variability and benthic fluxes, *Mar. Chem.*, 204, 49-61,
6077 <https://doi.org/10.1016/j.marchem.2018.05.011>, 2018.

6078 Lundin, E. J., Klaminder, J., Bastviken, D., Olid, C., Hansson, S. V., and Karlsson, J.: Large difference in carbon
6079 emission – burial balances between boreal and arctic lakes, *Sci. Rep.-UK*, 5, 14248,
6080 <https://doi.org/10.1038/srep14248>, 2015.

6081 Lundquist, J. K., DuVivier, K. K., Kaffine, D., and Tomaszewski, J. M.: Costs and consequences of wind turbine
6082 wake effects arising from uncoordinated wind energy development, *Nat. Energy*, 4, 26-34,
6083 <https://doi.org/10.1038/s41560-018-0281-2>, 2019.

6084 Luomaranta, A., Ruosteenoja, K., Jylhä, K., Gregow, H., Haapala, J., and Laaksonen, A.: Multimodel estimates
6085 of the changes in the Baltic Sea ice cover during the present century, *Tellus A*, 66, 22617,
6086 <https://doi.org/10.3402/tellusa.v66.22617>, 2014.

6087 Luomaranta, A., Aalto, J., and Jylhä, K.: Snow cover trends in Finland over 1961–2014 based on gridded snow
6088 depth observations, *Int. J. Climatol.*, 39, 3147-3159, <https://doi.org/10.1002/joc.6007>, 2019.

6089 Luoto, M., Fronzek, S., and Zuidhoff, F. S.: Spatial modelling of palsa mires in relation to climate in northern
6090 Europe, *Earth Surface Processes and Landforms: The Journal of the British Geomorphological Research Group*,
6091 29, 1373-1387, <https://doi.org/10.1002/esp.1099>, 2004.

6092 Luterbacher, J., Werner, J. P., Smerdon, J. E., Fernández-Donado, L., González-Rouco, F. J., Barriopedro, D.,
6093 Ljungqvist, F. C., Büntgen, U., Zorita, E., Wagner, S., Esper, J., McCarroll, D., Toreti, A., Frank, D., Jungclauss,
6094 J. H., Barriendos, M., Bertolin, C., Bothe, O., Brázdil, R., Camuffo, D., Dobrovolný, P., Gagen, M., García-
6095 Bustamante, E., Ge, Q., Gómez-Navarro, J. J., Guiot, J., Hao, Z., Hegerl, G. C., Holmgren, K., Klimenko, V. V.,
6096 Martín-Chivelet, J., Pfister, C., Roberts, N., Schindler, A., Schurer, A., Solomina, O., von Gunten, L., Wahl, E.,
6097 Wanner, H., Wetter, O., Xoplaki, E., Yuan, N., Zanchettin, D., Zhang, H., and Zerefos, C.: European summer
6098 temperatures since Roman times, *Environ. Res. Lett.*, 11, 024001, [https://doi.org/10.1088/1748-
6099 9326/11/2/024001](https://doi.org/10.1088/1748-9326/11/2/024001), 2016.

Formatiert: Englisch (USA)

Feldfunktion geändert

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

6100 Luysaert, S., Marie, G., Valade, A., Chen, Y.-Y., Njakou Djomo, S., Ryder, J., Otto, J., Naudts, K., Lansø, A.
6101 S., Ghattas, J., and McGrath, M. J.: Trade-offs in using European forests to meet climate objectives, *Nature*, 562,
6102 259-262, <https://doi.org/10.1038/s41586-018-0577-1>, 2018.

6103 Lyon, S. W., Destouni, G., Giesler, R., Humborg, C., Mörtz, M., Seibert, J., Karlsson, J., and Troch, P. A.:
6104 Estimation of permafrost thawing rates in a sub-arctic catchment using recession flow analysis, *Hydrol. Earth*
6105 *Syst. Sci.*, 13, 595-604, <https://doi.org/10.5194/hess-13-595-2009>, 2009.

6106 Maberly, S. C., O'Donnell, R. A., Woolway, R. I., Cutler, M. E. J., Gong, M., Jones, I. D., Merchant, C. J.,
6107 Miller, C. A., Politi, E., Scott, E. M., Thackeray, S. J., and Tyler, A. N.: Global lake thermal regions shift under
6108 climate change, *Nat. Commun.*, 11, 1232, <https://doi.org/10.1038/s41467-020-15108-z>, 2020.

6109 Macdonald, R. W., Mackay, D., Li, Y. F., and Hickie, B.: How Will Global Climate Change Affect Risks from
6110 Long-Range Transport of Persistent Organic Pollutants?, *Hum. Ecol. Risk Assess.*, 9, 643-660,
6111 <https://doi.org/10.1080/713609959>, 2003.

6112 MacKenzie, B. R., and Schiedek, D.: Daily ocean monitoring since the 1860s shows record warming of northern
6113 European seas, *Global Change Biol.*, 13, 1335-1347, <https://doi.org/10.1111/j.1365-2486.2007.01360.x>, 2007.

6114 MacKenzie, B. R., Meier, H. E. M., Lindegren, M., Neuenfeldt, S., Eero, M., Blenckner, T., Tomczak, M. T.,
6115 and Niiranen, S.: Impact of Climate Change on Fish Population Dynamics in the Baltic Sea: A Dynamical
6116 Downscaling Investigation, *Ambio* 41, 626-636, <https://doi.org/10.1007/s13280-012-0325-y>, 2012.

6117 MacLean, I. M. D., Austin, G. E., Rehfish, M. M., Blew, J., Crowe, O., Delany, S., Devos, K., Deceuninck, B.,
6118 Günther, K., Laursen, K., von Roomen, M., and Wahl, J.: Climate change causes rapid changes in the
6119 distribution and site abundance of birds in winter, *Global Change Biol.*, 14, 2489-2500,
6120 <https://doi.org/10.1111/j.1365-2486.2008.01666.x>, 2008.

6121 Magaard, L., and Rheinheimer, G.: *Meereskunde der Ostsee*, Springer-Verlag, New York, N.Y., 1974.

6122 Mäkinen, K., Vuorinen, I., and Hänninen, J.: Climate-induced hydrography change favours small-bodied
6123 zooplankton in a coastal ecosystem, *Hydrobiologia*, 792, 83-96, <https://doi.org/10.1007/s10750-016-3046-6>,
6124 2017.

6125 Mäll, M., Nakamura, R., Suursaar, Ü., and Shibayama, T.: Pseudo-climate modelling study on projected changes
6126 in extreme extratropical cyclones, storm waves and surges under CMIP5 multi-model ensemble: Baltic Sea
6127 perspective, *Nat. Hazards*, 102, 67-99, <https://doi.org/10.1007/s11069-020-03911-2>, 2020.

6128 Mann, M. E., Steinman, B. A., Brouillette, D. J., and Miller, S. K.: Multidecadal climate oscillations during the
6129 past millennium driven by volcanic forcing, *Science*, 371, 1014, <https://doi.org/10.1126/science.abc5810>, 2021.

6130 Männikus, R., Soomere, T., and Viška, M.: Variations in the mean, seasonal and extreme water level on the
6131 Latvian coast, the eastern Baltic Sea, during 1961–2018, *Estuar. Coast. Shelf Sci.*, 245, 106827,
6132 <https://doi.org/10.1016/j.ecss.2020.106827>, 2020.

6133 Marchowski, D., Jankowiak, Ł., Wysocki, D., Ławicki, Ł., and Girjatowicz, J.: Ducks change wintering patterns
6134 due to changing climate in the important wintering waters of the Odra River Estuary, *PeerJ*, 5, e3604,
6135 <https://doi.org/10.7717/peerj.3604>, 2017.

6136 Marcos, M., Calafat, F. M., Berihuete, Á., and Dangendorf, S.: Long-term variations in global sea level
6137 extremes, *J. Geophys. Res-Oceans*, 120, 8115-8134, <https://doi.org/10.1002/2015jc011173>, 2015.

6138 Marcos, M., and Woodworth, P. L.: Spatiotemporal changes in extreme sea levels along the coasts of the North
6139 Atlantic and the Gulf of Mexico, *J. Geophys. Res-Oceans*, 122, 7031-7048,
6140 <https://doi.org/10.1002/2017JC013065>, 2017.

6141 Markowicz, K. M., and Uscka-Kowalkowska, J.: Long-term and seasonal variability of the aerosol optical depth
6142 at Mount Kasprowy Wierch (Poland), *J. Geophys. Res-Atmos.*, 120, 1865-1879,
6143 <https://doi.org/10.1002/2014JD022580>, 2015.

6144 Marshall, G. J., Jylhä, K., Kivinen, S., Laapas, M., and Dyrørdal, A. V.: The role of atmospheric circulation
6145 patterns in driving recent changes in indices of extreme seasonal precipitation across Arctic Fennoscandia, *Clim.*
6146 *Change*, 162, 741-759, <https://doi.org/10.1007/s10584-020-02747-w>, 2020.

6147 Marshall, J., Johnson, H., and Goodman, J.: A Study of the Interaction of the North Atlantic Oscillation with
6148 Ocean Circulation, *J. Climate*, 14, 1399-1421, [https://doi.org/10.1175/1520-](https://doi.org/10.1175/1520-0442(2001)014<1399:ASOTIO>2.0.CO;2)
6149 [0442\(2001\)014<1399:ASOTIO>2.0.CO;2](https://doi.org/10.1175/1520-0442(2001)014<1399:ASOTIO>2.0.CO;2), 2001.

6150 Mastrandrea, M. D., Field, C. B., Stocker, T. F., Edenhofer, O., Ebi, K. L., Frame, D. J., Held, H., Kriegler, E.,
6151 Mach, K. J., Matschoss, P. R., Plattner, G.-K., Yohe, G. W., and Zwiers, F. W.: Guidance Note for Lead Authors
6152 of the IPCC Fifth Assessment Report on Consistent Treatment of Uncertainties, Intergovernmental Panel on
6153 Climate Change (IPCC), <http://www.ipcc.ch>, 2010.

6154 Matthäus, W., and Franck, H.: Characteristics of major Baltic inflows—a statistical analysis, *Cont. Shelf Res.*,
6155 12, 1375-1400, [https://doi.org/10.1016/0278-4343\(92\)90060-W](https://doi.org/10.1016/0278-4343(92)90060-W), 1992.

6156 Matthews, T., Murphy, C., Wilby, R. L., and Harrigan, S.: A cyclone climatology of the British-Irish Isles 1871–
6157 2012, *Int. J. Climatol.*, 36, 1299-1312, <https://doi.org/10.1002/joc.4425>, 2016.

6158 Matthias, V., Arndt, J. A., Aulinger, A., Bieser, J., Denier van der Gon, H., Kranenburg, R., Kuenen, J.,
6159 Neumann, D., Pouliot, G., and Quante, M.: Modeling emissions for three-dimensional atmospheric chemistry
6160 transport models, *J. Air Waste Manag.*, 68, 763-800, <https://doi.org/10.1080/10962247.2018.1424057>, 2018.

6161 Mauri, A., Davis, B. A. S., Collins, P. M., and Kaplan, J. O.: The influence of atmospheric circulation on the
6162 mid-Holocene climate of Europe: a data-model comparison, *Clim. Past*, 10, 1925-1938,
6163 <https://doi.org/10.5194/cp-10-1925-2014>, 2014.

6164 Mauri, A., Davis, B. A. S., Collins, P. M., and Kaplan, J. O.: The climate of Europe during the Holocene: a
6165 gridded pollen-based reconstruction and its multi-proxy evaluation, *Quaternary Sci. Rev.*, 112, 109-127,
6166 <https://doi.org/10.1016/j.quascirev.2015.01.013>, 2015.

6167 McClelland, J. W., Holmes, R. M., Peterson, B. J., and Stieglitz, M.: Increasing river discharge in the Eurasian
6168 Arctic: Consideration of dams, permafrost thaw, and fires as potential agents of change, *J. Geophys. Res.-Atmos.*,
6169 109, <https://doi.org/10.1029/2004JD004583>, 2004.

6170 McCrackin, M. L., Muller-Karulis, B., Gustafsson, B. G., Howarth, R. W., Humborg, C., Svanbäck, A., and
6171 Swaney, D. P.: A Century of Legacy Phosphorus Dynamics in a Large Drainage Basin, *Global Biogeochem.*
6172 *Cy.*, 32, 1107-1122, <https://doi.org/10.1029/2018GB005914>, 2018.

6173 McGlade, J. M.: 12 The North Sea Large Marine Ecosystem, in: *Large Marine Ecosystems*, edited by: Sherman,
6174 K., and Skjoldal, H. R., Elsevier, 339-412, [https://doi.org/10.1016/S1570-0461\(02\)80064-7](https://doi.org/10.1016/S1570-0461(02)80064-7), 2002.

6175 McNamara, J. M., and Houston, A. I.: State-dependent life histories, *Nature*, 380, 215-221,
6176 <https://doi.org/10.1038/380215a0>, 1996.

6177 Meehl, G. A., Senior, C. A., Eyring, V., Flato, G., Lamarque, J.-F., Stouffer, R. J., Taylor, K. E., and Schlund,
6178 M.: Context for interpreting equilibrium climate sensitivity and transient climate response from the CMIP6 Earth
6179 system models, *Sci. Adv.*, 6, eaba1981, <https://doi.org/10.1126/sciadv.aba1981>, 2020.

6180 Meier, H. E. M.: Regional ocean climate simulations with a 3D ice–ocean model for the Baltic Sea. Part 2:
6181 results for sea ice, *Clim. Dyn.*, 19, 255-266, <https://doi.org/10.1007/s00382-001-0225-5>, 2002.

6182 Meier, H. E. M., and Döscher, R.: Simulated water and heat cycles of the Baltic Sea using a 3D coupled
6183 atmosphere-ice-ocean model, *Boreal Environ. Res.*, 7, 327-334, 2002.

6184 Meier, H. E. M., and Kauker, F.: Modeling decadal variability of the Baltic Sea: 2. Role of freshwater inflow and
6185 large-scale atmospheric circulation for salinity, *J. Geophys. Res.-Oceans*, 108,
6186 <https://doi.org/10.1029/2003JC001799>, 2003.

6187 Meier, H. E. M., Broman, B., and Kjellström, E.: Simulated sea level in past and future climates of the Baltic
6188 Sea, *Clim. Res.*, 27, 59-75, <https://doi.org/10.3354/cr027059>, 2004a.

6189 Meier, H. E. M., Döscher, R., and Halkka, A.: Simulated distributions of Baltic Sea-ice in warming climate and
6190 consequences for the winter habitat of the Baltic ringed seal, *Ambio*, 33, 249-256, [https://doi.org/10.1579/0044-](https://doi.org/10.1579/0044-7447-33.4.249)
6191 [7447-33.4.249](https://doi.org/10.1579/0044-7447-33.4.249), 2004b.

6192 Meier, H. E. M.: Baltic Sea climate in the late twenty-first century: a dynamical downscaling approach using two
6193 global models and two emission scenarios, *Clim. Dyn.*, 27, 39–68, <https://doi.org/10.1007/s00382-006-0124-x>,
6194 2006.

6195 Meier, H. E. M., Kjellström, E., and Graham, L. P.: Estimating uncertainties of projected Baltic Sea salinity in
6196 the late 21st century, *Geophys. Res. Lett.*, 33, L15705, <https://doi.org/10.1029/2006GL026488>, 2006.

6197 Meier, H. E. M., Andersson, H. C., Eilola, K., Gustafsson, B. G., Kuznetsov, I., Müller-Karulis, B., Neumann,
6198 T., and Savchuk, O. P.: Hypoxia in future climates: A model ensemble study for the Baltic Sea, *Geophys. Res.*
6199 *Lett.*, 38, L24608, <https://doi.org/10.1029/2011GL049929>, 2011a.

6200 Meier, H. E. M., Eilola, K., and Almroth, E.: Climate-related changes in marine ecosystems simulated with a 3-
6201 dimensional coupled physical-biogeochemical model of the Baltic Sea, *Clim. Res.*, 48, 31–55,
6202 <https://doi.org/10.3354/cr00968> 2011b.

6203 Meier, H. E. M., Höglund, A., Döscher, R., Andersson, H., Löptien, U., and Kjellström, E.: Quality assessment
6204 of atmospheric surface fields over the Baltic Sea from an ensemble of regional climate model simulations with
6205 respect to ocean dynamics, *Oceanologia*, 53, 193–227, <https://doi.org/10.5697/oc.53-1-TI.193>, 2011c.

6206 Meier, H. E. M., Andersson, H. C., Arheimer, B., Blenckner, T., Chubarenko, B., Donnelly, C., Eilola, K.,
6207 Gustafsson, B. G., Hansson, A., Havenhand, J., and others: Comparing reconstructed past variations and future
6208 projections of the Baltic Sea ecosystem—first results from multi-model ensemble simulations, *Environ. Res.*
6209 *Lett.*, 7, 34005, <https://doi.org/10.1088/1748-9326/7/3/034005>, 2012a.

6210 Meier, H. E. M., Hordoir, R., Andersson, H. C., Dieterich, C., Eilola, K., Gustafsson, B. G., Höglund, A., and
6211 Schimanke, S.: Modeling the combined impact of changing climate and changing nutrient loads on the Baltic
6212 Sea environment in an ensemble of transient simulations for 1961–2099, *Clim. Dyn.*, 39, 2421–2441,
6213 <https://doi.org/10.1007/s00382-012-1339-7>, 2012b.

6214 Meier, H. E. M., Müller-Karulis, B., Andersson, H. C., Dieterich, C., Eilola, K., Gustafsson, B. G., Höglund, A.,
6215 Hordoir, R., Kuznetsov, I., Neumann, T., and others: Impact of climate change on ecological quality indicators
6216 and biogeochemical fluxes in the Baltic Sea: a multi-model ensemble study, *Ambio*, 41, 558–573,
6217 <https://doi.org/10.1007/s13280-012-0320-3>, 2012c.

6218 Meier, H. E. M., Rutgersson, A., and Reckermann, M.: An Earth System Science Program for the Baltic Sea
6219 Region, *Eos Trans. AGU*, 95, 109–110, <https://doi.org/10.1002/2014EO130001>, 2014.

6220 Meier, H. E. M.: Projected Change—Marine Physics, in: Second Assessment of Climate Change for the Baltic
6221 Sea Basin, edited by: BACC II Author Team, Springer International Publishing, Cham, 243–252,
6222 https://doi.org/10.1007/978-3-319-16006-1_13, 2015.

6223 Meier, H. E. M., Höglund, A., Eilola, K., and Almroth-Rosell, E.: Impact of accelerated future global mean sea
6224 level rise on hypoxia in the Baltic Sea, *Clim. Dyn.*, 49, 163–172, <https://doi.org/10.1007/s00382-016-3333-y>,
6225 2017.

6226 Meier, H. E. M., Edman, M. K., Eilola, K. J., Placke, M., Neumann, T., Andersson, H. C., Brunnabend, S.-E.,
6227 Dieterich, C., Frauen, C., Friedland, R., Gröger, M., Gustafsson, B. G., Gustafsson, E., Isaev, A., Kniebusch, M.,
6228 Kuznetsov, I., Müller-Karulis, B., Omstedt, A., Ryabchenko, V., Saraiva, S., and Savchuk, O. P.: Assessment of
6229 Eutrophication Abatement Scenarios for the Baltic Sea by Multi-Model Ensemble Simulations, *Front. Mar. Sci.*,
6230 5, <https://doi.org/10.3389/fmars.2018.00440>, 2018a.

6231 Meier, H. E. M., Väli, G., Naumann, M., Eilola, K., and Frauen, C.: Recently accelerated oxygen consumption
6232 rates amplify deoxygenation in the Baltic Sea, *J. Geophys. Res.-Oceans*, 123, 3227–3240,
6233 <https://doi.org/10.1029/2017JC013686>, 2018b.

6234 Meier, H. E. M., Dieterich, C., Eilola, K., Gröger, M., Höglund, A., Radtke, H., Saraiva, S., and Wählström, I.:
6235 Future projections of record-breaking sea surface temperature and cyanobacteria bloom events in the Baltic Sea,
6236 *Ambio*, 48, 1362–1376, <https://doi.org/10.1007/s13280-019-01235-5>, 2019a.

6237 Meier, H. E. M., Edman, M., Eilola, K., Placke, M., Neumann, T., Andersson, H. C., Brunnabend, S.-E.,
6238 Dieterich, C., Frauen, C., Friedland, R., Gröger, M., Gustafsson, B. G., Gustafsson, E., Isaev, A., Kniebusch, M.,
6239 Kuznetsov, I., Müller-Karulis, B., Naumann, M., Omstedt, A., Ryabchenko, V., Saraiva, S., and Savchuk, O. P.:

6240 Assessment of Uncertainties in Scenario Simulations of Biogeochemical Cycles in the Baltic Sea, *Front. Mar.*
6241 *Sci.*, 6, <https://doi.org/10.3389/fmars.2019.00046>, 2019b.

6242 Meier, H. E. M., Eilola, K., Almroth-Rosell, E., Schimanke, S., Kniebusch, M., Höglund, A., Pemberton, P., Liu,
6243 Y., Väli, G., and Saraiva, S.: Correction to: Disentangling the impact of nutrient load and climate changes on
6244 Baltic Sea hypoxia and eutrophication since 1850, *Clim. Dyn.*, 53, 1167–1169, [https://doi.org/10.1007/s00382-](https://doi.org/10.1007/s00382-018-4483-x)
6245 018-4483-x, 2019c.

6246 Meier, H. E. M., Eilola, K., Almroth-Rosell, E., Schimanke, S., Kniebusch, M., Höglund, A., Pemberton, P., Liu,
6247 Y., Väli, G., and Saraiva, S.: Disentangling the impact of nutrient load and climate changes on Baltic Sea
6248 hypoxia and eutrophication since 1850, *Clim. Dyn.*, 53, 1145–1166, <https://doi.org/10.1007/s00382-018-4296-y>,
6249 2019d.

6250 Meier, H. E. M., and Saraiva, S.: Projected Oceanographical Changes in the Baltic Sea until 2100, *Oxford*
6251 *Research Encyclopedia of Climate Science*, <https://doi.org/10.1093/acrefore/9780190228620.013.699>, 2020.

6252 Meier, H. E. M., Börgel, F., Bøssing-Christensen, O., Dieterich, C., Dutheil, C., Gröger, M., and Kjellström, E.:
6253 Oceanographic regional climate projections for the Baltic Sea until 2100, *Earth Syst. Dynam.*, 2021a.

6254 Meier, H. E. M., Dieterich, C., and Gröger, M.: Natural variability is a large source of uncertainty in future
6255 projections of hypoxia in the Baltic Sea, *Commun. Earth Environ.*, 2, 50, [https://doi.org/10.1038/s43247-021-](https://doi.org/10.1038/s43247-021-00115-9)
6256 00115-9, 2021b.

6257 Meier, R., Schwaab, J., Seneviratne, S. I., Sprenger, M., Lewis, E., and Davin, E. L.: Empirical estimate of
6258 forestation-induced precipitation changes in Europe, *Nat. Geosci.*, 14, 473–478, [https://doi.org/10.1038/s41561-](https://doi.org/10.1038/s41561-021-00773-6)
6259 021-00773-6, 2021c.

6260 [Menary, M. B., Robson, J., Allan, R. P., Booth, B. B. B., Cassou, C., Gastineau, G., Gregory, J., Hodson, D.,](#)
6261 [Jones, C., Mignot, J., Ringer, M., Sutton, R., Wilcox, L., and Zhang, R.: Aerosol-Forced AMOC Changes in](#)
6262 [CMIP6 Historical Simulations](#), 47, e2020GL088166, <https://doi.org/10.1029/2020GL088166>, 2020.

6263 Mercer, A.: A DEM of the 2010 surface topography of Storglaciären, Sweden, *J. Maps*, 12, 1112–1118,
6264 <https://doi.org/10.1080/17445647.2015.1131754>, 2016.

6265 Mercer, J. H.: West Antarctic ice sheet and CO₂ greenhouse effect: a threat of disaster, *Nature*, 271, 321–325,
6266 1978.

6267 Merkouriadi, I., and Leppäranta, M.: Long-term analysis of hydrography and sea-ice data in Tvärminne, Gulf of
6268 Finland, Baltic Sea, *Clim. Change*, 124, 849–859, <https://doi.org/10.1007/s10584-014-1130-3>, 2014.

6269 Merkouriadi, I., Leppäranta, M., and Järvinen, O.: Interannual variability and trends in winter weather and snow
6270 conditions in Finnish Lapland, *Est. J. Earth Sci.*, 66, 47–57, <https://doi.org/10.3176/earth.2017.03>, 2017.

6271 Mikulski, Z.: Inflow from drainage basin, *Water Balance of the Baltic Sea-Baltic Sea Environment Proceedings*,
6272 16, 24–34, Baltic Marine Environment Protection Commission, Helsinki, Finland, 1986.

6273 Mitrovica, J. X., Hay, C. C., Kopp, R. E., Harig, C., and Latychev, K.: Quantifying the Sensitivity of Sea Level
6274 Change in Coastal Localities to the Geometry of Polar Ice Mass Flux, *J. Climate*, 31, 3701–3709,
6275 <https://doi.org/10.1175/jcli-d-17-0465.1>, 2018.

6276 Mohrholz, V., Naumann, M., Nausch, G., Krüger, S., and Gräwe, U.: Fresh oxygen for the Baltic Sea—An
6277 exceptional saline inflow after a decade of stagnation, *J. Mar. Syst.*, 148, 152–166,
6278 <https://doi.org/10.1016/j.jmarsys.2015.03.005>, 2015.

6279 Mohrholz, V.: Major Baltic Inflow Statistics – Revised, *Front. Mar. Sci.*, 5,
6280 <https://doi.org/10.3389/fmars.2018.00384>, 2018.

6281 Moldanová, J., Hassellöv, I.-M., Matthias, V., Fridell, E., Jalkanen, J.-P., Ytreberg, E., Quante, M., Tröltzsch, J.,
6282 Maljutenko, I., Raudsepp, U., and Eriksson, K. M.: Framework for the environmental impact assessment of
6283 operational shipping, *Ambio*, <https://doi.org/10.1007/s13280-021-01597-9>, 2021.

6284 Möller, K. O., Schmidt, J. O., St.John, M., Temming, A., Diekmann, R., Peters, J., Floeter, J., Sell, A. F.,
6285 Herrmann, J.-P., and Möllmann, C.: Effects of climate-induced habitat changes on a key zooplankton species, *J.*
6286 *Plankton Res.*, 37, 530-541, <https://doi.org/10.1093/plankt/fbv033>, 2015.

6287 Möllmann, C.: Effects of Climate Change and Fisheries on the Marine Ecosystem of the Baltic Sea, Oxford
6288 Research Encyclopedia of Climate Science, <https://doi.org/10.1093/acrefore/9780190228620.013.682>, 2019.

6289 [Moreno-Chamarro, E., Caron, L. P., Ortega, P., Loosveldt Tomas, S., and Roberts, M. J.: Can we trust CMIP5/6](#)
6290 [future projections of European winter precipitation?](#), *Environ. Res. Lett.*, 16, 054063,
6291 <https://doi.org/10.1088/1748-9326/abf28a>, 2021.

6292 Müller, J., Folini, D., Wild, M., and Pfenninger, S.: CMIP-5 models project photovoltaics are a no-regrets
6293 investment in Europe irrespective of climate change, *Energy*, 171, 135-148,
6294 <https://doi.org/10.1016/j.energy.2018.12.139>, 2019.

6295 Müller, J. D., Schneider, B., and Rehder, G.: Long-term alkalinity trends in the Baltic Sea and their implications
6296 for CO₂-induced acidification, *Limnol. Oceanogr.*, 61, 1984-2002, <https://doi.org/10.1002/lno.10349>, 2016.

6297 Munkes, B., Löptien, U., and Dietze, H.: Cyanobacteria blooms in the Baltic Sea: a review of models and facts,
6298 *Biogeosciences*, 18, 2347-2378, <https://doi.org/10.5194/bg-18-2347-2021>, 2021.

6299 Musielak, S., Furmańczyk, K., and Bugajny, N.: Factors and Processes Forming the Polish Southern Baltic Sea
6300 Coast on Various Temporal and Spatial Scales, in: *Coastline Changes of the Baltic Sea from South to East: Past*
6301 *and Future Projection*, edited by: Harff, J., Furmańczyk, K., and von Storch, H., Springer International
6302 Publishing, Cham, 69-85, https://doi.org/10.1007/978-3-319-49894-2_5, 2017.

6303 Myrberg, K., Korpinen, S., and Uusitalo, L.: Physical oceanography sets the scene for the Marine Strategy
6304 Framework Directive implementation in the Baltic Sea, *Mar. Policy*, 107, 103591,
6305 <https://doi.org/10.1016/j.marpol.2019.103591>, 2019.

6306 Nakamura, T., Yamazaki, K., Iwamoto, K., Honda, M., Miyoshi, Y., Ogawa, Y., and Ukita, J.: A negative phase
6307 shift of the winter AO/NAO due to the recent Arctic sea-ice reduction in late autumn, *J. Geophys. Res.-Atmos.*,
6308 120, 3209-3227, <https://doi.org/10.1002/2014JD022848>, 2015.

6309 Nakicenovic, N., Alcamo, J., Grubler, A., Riahi, K., Roehrl, R. A., Rogner, H.-H., and Victor, N.: Special report
6310 on emissions scenarios (SRES), a special report of Working Group III of the intergovernmental panel on climate
6311 change, IPCC Special Report, 2000.

6312 Naumann, M., Gräwe, U., Mohrholz, V., Kuss, J., Siegel, H., Wanek, J. J., and Schulz-Bull, D. E.:
6313 Hydrographic-hydrochemical assessment of the Baltic Sea 2018, *Meereswiss. Ber.*, Warnemünde, 110,
6314 <https://doi.io-warnemuende.de/10.12754/msr-2019-0110>, 2019.

6315 Nerem, R. S., Beckley, B. D., Fasullo, J. T., Hamlington, B. D., Masters, D., and Mitchum, G. T.: Climate-
6316 change-driven accelerated sea-level rise detected in the altimeter era, *Proceedings of the National Academy of*
6317 *Sciences*, 115, 2022, <https://doi.org/10.1073/pnas.1717312115>, 2018.

6318 Neumann, G.: *Eigenschwingungen der Ostsee*, Archiv der Deutschen Seewarte und des Marineobservatoriums,
6319 Hamburg, 1941.

6320 Neumann, T., Eilola, K., Gustafsson, B., Müller-Karulis, B., Kuznetsov, I., Meier, H. E. M., and Savchuk, O. P.:
6321 Extremes of temperature, oxygen and blooms in the Baltic Sea in a changing climate, *Ambio*, 41, 574-585,
6322 <https://doi.org/10.1007/s13280-012-0321-2>, 2012.

6323 Neumann, T., Radtke, H., and Seifert, T.: On the importance of Major Baltic Inflows for oxygenation of the
6324 central Baltic Sea, *J. Geophys. Res.-Oceans*, 122, 1090-1101, <https://doi.org/10.1002/2016JC012525>, 2017.

6325 [Nie, Y., Li, L., Tang, Y., and Wang, B.: Impacts of Changes of External Forcings from CMIP5 to CMIP6 on](#)
6326 [Surface Temperature in FGOALS-g2, SOLA, 15, 211-215, https://doi.org/10.2151/sola.2019-038, 2019.](#)

6327 Niemelä, P., Tolvanen, H., Rönkä, M., Kellomäki, S., Krug, J., Schurgers, G., Lehtikoinen, E., and Kalliola, R.:
6328 Environmental Impacts—Coastal Ecosystems, Birds and Forests, in: *Second Assessment of Climate Change for*
6329 *the Baltic Sea Basin*, edited by: BACC II Author Team, Springer International Publishing, Cham, 291-306,
6330 https://doi.org/10.1007/978-3-319-16006-1_16, 2015.

6331 Niiranen, S., Yletyinen, J., Tomczak, M. T., Blenckner, T., Hjerne, O., MacKenzie, B. R., Müller-Karulis, B.,
6332 Neumann, T., and Meier, H. E. M.: Combined effects of global climate change and regional ecosystem drivers
6333 on an exploited marine food web, *Global Change Biol.*, 19, 3327-3342, <https://doi.org/10.1111/gcb.12309>, 2013.

6334 Nijse, F. J. M. M., Cox, P. M., Huntingford, C., and Williamson, M. S.: Decadal global temperature variability
6335 increases strongly with climate sensitivity, *Nat. Clim. Change*, 9, 598-601, [https://doi.org/10.1038/s41558-019-](https://doi.org/10.1038/s41558-019-0527-4)
6336 0527-4, 2019.

6337 Nilsson, J., and Grennfelt, P.: Critical loads for sulphur and nitrogen, Report from Skokloster Workshop.
6338 Skokloster. Sweden, 1988.

6339 Nilsson, L., and Haas, F.: Distribution and numbers of wintering waterbirds in Sweden in 2015 and changes
6340 during the last fifty years, *Ornis Svecica*, 26, 3-54-53-54, 2016.

6341 Ning, W., Nielsen, A. B., Ivarsson, L. N., Jilbert, T., Åkesson, C. M., Slomp, C. P., Andrén, E., Broström, A.,
6342 and Filipsson, H. L.: Anthropogenic and climatic impacts on a coastal environment in the Baltic Sea over the last
6343 1000 years, *Anthropocene*, 21, 66-79, <https://doi.org/10.1016/j.ancene.2018.02.003>, 2018.

6344 Niskanen, T., Vainio, J., Eriksson, P., and Heiler, I.: Maximum extent of the Baltic sea ice recalculated for the
6345 period 1971-2008, Report Series in Geophysics, 164, 2009.

6346 Nöges, P., and Nöges, T.: Weak trends in ice phenology of Estonian large lakes despite significant warming
6347 trends, *Hydrobiologia*, 731, 5-18, <https://doi.org/10.1007/s10750-013-1572-z>, 2014.

6348 Norbäck Ivarsson, L., Andrén, T., Moros, M., Andersen, T. J., Lönn, M., and Andrén, E.: Baltic Sea Coastal
6349 Eutrophication in a Thousand Year Perspective, *Front. Environ. Sci.*, 7,
6350 <https://doi.org/10.3389/fenvs.2019.00088>, 2019.

6351 Nydahl, A., Panigrahi, S., and Wikner, J.: Increased microbial activity in a warmer and wetter climate enhances
6352 the risk of coastal hypoxia, *FEMS Microbiol. Ecol.*, 85, 338-347, <https://doi.org/10.1111/1574-6941.12123>,
6353 2013.

6354 O'Reilly, C. M., Sharma, S., Gray, D. K., Hampton, S. E., Read, J. S., Rowley, R. J., Schneider, P., Lenters, J.
6355 D., McIntyre, P. B., Kraemer, B. M., Weyhenmeyer, G. A., Straile, D., Dong, B., Adrian, R., Allan, M. G.,
6356 Anneville, O., Arvola, L., Austin, J., Bailey, J. L., Baron, J. S., Brookes, J. D., de Eyto, E., Dokulil, M. T.,
6357 Hamilton, D. P., Havens, K., Hetherington, A. L., Higgins, S. N., Hook, S., Izmet'seva, L. R., Joehnk, K. D.,
6358 Kangur, K., Kasprzak, P., Kumagai, M., Kuusisto, E., Leshkevich, G., Livingstone, D. M., MacIntyre, S., May,
6359 L., Melack, J. M., Mueller-Navarra, D. C., Naumenko, M., Noges, P., Noges, T., North, R. P., Plisnier, P.-D.,
6360 Rigosi, A., Rimmer, A., Rogora, M., Rudstam, L. G., Rusak, J. A., Salmaso, N., Samal, N. R., Schindler, D. E.,
6361 Schladow, S. G., Schmid, M., Schmidt, S. R., Silow, E., Soylu, M. E., Teubner, K., Verburg, P., Voutilainen, A.,
6362 Watkinson, A., Williamson, C. E., and Zhang, G.: Rapid and highly variable warming of lake surface waters
6363 around the globe, *Geophys. Res. Lett.*, 42, 10,773-710,781, <https://doi.org/10.1002/2015GL066235>, 2015.

6364 Oberbeckmann, S., and Labrenz, M.: Marine Microbial Assemblages on Microplastics: Diversity, Adaptation,
6365 and Role in Degradation, *Annu. Rev. Mar. Sci.*, 12, 209-232, [https://doi.org/10.1146/annurev-marine-010419-](https://doi.org/10.1146/annurev-marine-010419-010633)
6366 010633, 2020.

6367 Obu, J., Westermann, S., Bartsch, A., Berdnikov, N., Christiansen, H. H., Dashtseren, A., Delaloye, R.,
6368 Elberling, B., Etzelmueller, B., Kholodov, A., Khomutov, A., Kääh, A., Leibman, M. O., Lewkowicz, A. G.,
6369 Panda, S. K., Romanovsky, V., Way, R. G., Westergaard-Nielsen, A., Wu, T., Yamkhin, J., and Zou, D.:
6370 Northern Hemisphere permafrost map based on TTOP modelling for 2000-2016 at 1 km2 scale, *Earth Sci. Rev.*,
6371 193, 299-316, <https://doi.org/10.1016/j.earscirev.2019.04.023>, 2019.

6372 Obu, J., Westermann, S., Barboux, C., Bartsch, A., Delaloye, R., Grosse, G., Heim, B., Hugelius, G., Irrgang, A.,
6373 Kääh, A. M., Kroisleitner, C., Matthes, H., Nitze, I., Pellet, C., Seifert, F. M., Strozzi, T., Wegmüller, U.,
6374 Wiczorek, M., and Wiesmann, A.: ESA Permafrost Climate Change Initiative (Permafrost_cci): Permafrost
6375 extent for the Northern Hemisphere, v2.0., Centre for Environmental Data Analysis,
6376 <http://dx.doi.org/10.5285/28e889210f884b469d7168fde4b4e54f>, 2020.

6377 Ojaveer, H., Olenin, S., Naršcius, A., Florin, A.-B., Ezhova, E., Gollasch, S., Jensen, K. R., Lehtiniemi, M.,
6378 Minchin, D., Normant-Saremba, M., and Stråke, S.: Dynamics of biological invasions and pathways over time: a

Formatiert: Englisch (USA)

Feldfunktion geändert

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Rechtschreibung und Grammatik nicht prüfen

6379 case study of a temperate coastal sea, *Biol. Invasions*, 19, 799-813, <https://doi.org/10.1007/s10530-016-1316-x>,
6380 2017.

6381 Olli, K., Klais, R., Tamminen, T., Ptacnik, R., and Andersen, T.: Long term changes in the Baltic Sea
6382 phytoplankton community, *Boreal Environ. Res.*, 16 (suppl. A), 3-14, 2011.

6383 Olofsson, M., Torstensson, A., Karlberg, M., Steinhoff, F. S., Dinasquet, J., Riemann, L., Chierici, M., and
6384 Wulff, A.: Limited response of a spring bloom community inoculated with filamentous cyanobacteria to elevated
6385 temperature and pCO₂, *Bot. Mar.*, 62, 3-16, <https://doi.org/10.1515/bot-2018-0005>, 2019.

6386 Olofsson, M., Suikkanen, S., Kobos, J., Wasmund, N., and Karlson, B.: Basin-specific changes in filamentous
6387 cyanobacteria community composition across four decades in the Baltic Sea, *Harmful Algae*, 91, 101685,
6388 <https://doi.org/10.1016/j.hal.2019.101685>, 2020.

6389 Omstedt, A., and Chen, D.: Influence of atmospheric circulation on the maximum ice extent in the Baltic Sea, *J.*
6390 *Geophys. Res-Oceans*, 106, 4493-4500, <https://doi.org/10.1029/1999JC000173>, 2001.

6391 Omstedt, A., Edman, M., Claremar, B., Frodin, P., Gustafsson, E., Humborg, C., Hägg, H., Mörtz, M.,
6392 Rutgersson, A., Schurgers, G., and others: Future changes in the Baltic Sea acid-base (pH) and oxygen balances,
6393 *Tellus B*, 64, 19586, <https://doi.org/10.3402/tellusb.v64i0.19586>, 2012.

6394 Omstedt, A., Elken, J., Lehmann, A., Leppäranta, M., Meier, H. E. M., Myrberg, K., and Rutgersson, A.:
6395 Progress in physical oceanography of the Baltic Sea during the 2003–2014 period, *Prog. Oceanogr.*, 128, 139-
6396 171, <https://doi.org/10.1016/j.pocean.2014.08.010>, 2014.

6397 Oppenheimer, M., Glavovic, B. C., Hinkel, J., Wal, R. v. d., Magnan, A. K., Abd-Elgawad, A., Cai, R.,
6398 Cifuentes-Jara, M., DeConto, R. M., Ghosh, T., Hay, J., Isla, F., Marzeion, B., Meyssignac, B., and Sebesvari,
6399 Z.: Sea Level Rise and Implications for Low-Lying Islands, Coasts and Communities, in: *IPCC Special Report*
6400 *on the Ocean and Cryosphere in a Changing Climate*, edited by: Pörtner, H.-O., Roberts, D. C., Masson-
6401 Delmotte, V., Zhai, P., Tignor, M., Poloczanska, E., Mintenbeck, K., Alegria, A., Nicolai, M., Okem, A.,
6402 Petzold, J., Rama, B., and Weyer, N. M., 2019.

6403 Orru, H., Andersson, C., Ebi, K. L., Langner, J., Åström, C., and Forsberg, B.: Impact of climate change on
6404 ozone-related mortality and morbidity in Europe, *Eur. Respir. J.*, 41, 285-294,
6405 <https://doi.org/10.1183/09031936.00210411>, 2013.

6406 Orru, H., Åström, C., Andersson, C., Tamm, T., Ebi, K. L., and Forsberg, B.: Ozone and heat-related mortality in
6407 Europe in 2050 significantly affected by changes in climate, population and greenhouse gas emission, *Environ.*
6408 *Res. Lett.*, 14, 074013, 2019.

6409 [Orviku, K., Jaagus, J., and Tõnisson, H.: Sea ice shaping the shores, *J. Coastal Res.*, 681-685, 2011.](#)

6410 Osterkamp, T., and Romanovsky, V.: Evidence for warming and thawing of discontinuous permafrost in Alaska,
6411 *Permafrost Periglac.*, 10, 17-37, [https://doi.org/10.1002/\(SICI\)1099-1530\(199901/03\)10:1<17::AID-PPP303>3.0.CO;2-4](https://doi.org/10.1002/(SICI)1099-1530(199901/03)10:1<17::AID-PPP303>3.0.CO;2-4), 1999.

6412

6413 Otto, L., Zimmerman, J. T. F., Furnes, G. K., Mork, M., Saetre, R., and Becker, G.: Review of the physical
6414 oceanography of the North Sea, *Neth. J. Sea Res.*, 26, 161-238, [https://doi.org/10.1016/0077-7579\(90\)90091-T](https://doi.org/10.1016/0077-7579(90)90091-T),
6415 1990.

6416 Øygarden, L., Deelstra, J., Lagzdins, A., Bechmann, M., Greipsland, I., Kyllmar, K., Povilaitis, A., and Iital, A.:
6417 Climate change and the potential effects on runoff and nitrogen losses in the Nordic–Baltic region, *Agric.*
6418 *Ecosyst. Environ.*, 198, 114-126, <https://doi.org/10.1016/j.agee.2014.06.025>, 2014.

6419 Paczkowska, J., Brugel, S., Rowe, O., Lefebure, R., Brutemark, A., and Andersson, A.: Response of Coastal
6420 Phytoplankton to High Inflows of Terrestrial Matter, *Front. Mar. Sci.*, 7,
6421 <https://doi.org/10.3389/fmars.2020.00080>, 2020.

6422 Pajusalu, L., Martin, G., Pöllumäe, A., and Paalme, T.: Results of laboratory and field experiments of the direct
6423 effect of increasing CO₂ on net primary production of macroalgal species in brackish-water ecosystems, *P. Est.*
6424 *Acad. Sci.*, 62, 148, <https://doi.org/10.3176/proc.2013.2.09>, 2013.

6425 Pajusalu, L., Martin, G., Pöllumäe, A., Torn, K., and Paalme, T.: Direct effects of increased CO₂ concentrations
6426 in seawater on the net primary production of charophytes in a shallow, coastal, brackish-water ecosystem, *Boreal*
6427 *Environ. Res.*, 20, 413-422, 2015.

6428 Pajusalu, L., Martin, G., Paalme, T., and Pöllumäe, A.: The effect of CO₂ enrichment on net photosynthesis of
6429 the red alga *Furcellaria lumbricalis* in a brackish water environment, *PeerJ*, 4, e2505,
6430 <https://doi.org/10.7717/peerj.2505>, 2016.

6431 Pansch, C., Nasrolahi, A., Appelhans, Y. S., and Wahl, M.: Impacts of ocean warming and acidification on the
6432 larval development of the barnacle *Amphibalanus improvisus*, *J. Exp. Mar. Biol. Ecol.*, 420-421, 48-55,
6433 <https://doi.org/10.1016/j.jembe.2012.03.023>, 2012.

6434 Parding, K., Olseth, J. A., Dagestad, K. F., and Liepert, B. G.: Decadal variability of clouds, solar radiation and
6435 temperature at a high-latitude coastal site in Norway, *Tellus B*, 66, 25897,
6436 <https://doi.org/10.3402/tellusb.v66.25897>, 2014.

6437 Paul, C., Matthiessen, B., and Sommer, U.: Warming, but not enhanced CO₂ concentration, quantitatively and
6438 qualitatively affects phytoplankton biomass, *Mar. Ecol. Prog. Ser.*, 528, 39-51,
6439 <https://doi.org/10.3354/meps11264> 2015.

6440 Pavón-Jordán, D., Fox, A. D., Clausen, P., Dagys, M., Deceuninck, B., Devos, K., Hearn, R. D., Holt, C. A.,
6441 Hornman, M., Keller, V., Langendoen, T., Lawicki, L., Lorentsen, S. H., Luigujõe, L., Meissner, W., Musil, P.,
6442 Nilsson, L., Paquet, J.-Y., Stipniece, A., Stroud, D. A., Wahl, J., Zenatello, M., and Lehikoinen, A.: Climate-
6443 driven changes in winter abundance of a migratory waterbird in relation to EU protected areas, *Divers. Distrib.*,
6444 21, 571-582, <https://doi.org/10.1111/ddi.12300>, 2015.

6445 Pavón-Jordán, D., Clausen, P., Dagys, M., Devos, K., Encarnação, V., Fox, A. D., Frost, T., Gaudard, C.,
6446 Hornman, M., Keller, V., Langendoen, T., Lawicki, L., Lewis, L. J., Lorentsen, S.-H., Luigujõe, L., Meissner,
6447 W., Molina, B., Musil, P., Musilova, Z., Nilsson, L., Paquet, J.-Y., Ridzon, J., Stipniece, A., Teufelbauer, N.,
6448 Wahl, J., Zenatello, M., and Lehikoinen, A.: Habitat- and species-mediated short- and long-term distributional
6449 changes in waterbird abundance linked to variation in European winter weather, *Divers. Distrib.*, 25, 225-239,
6450 <https://doi.org/10.1111/ddi.12855>, 2019.

6451 Pecuchet, L., Lindegren, M., Kortsch, S., Całkiewicz, J., Jurgensone, I., Margonski, P., Otto, S. A., Putnis, I.,
6452 Stråke, S., and Nordström, M. C.: Spatio-temporal dynamics of multi-trophic communities reveal ecosystem-
6453 wide functional reorganization, *Ecography*, 43, 197-208, <https://doi.org/10.1111/ecog.04643>, 2020.

6454 Peings, Y., and Magnusdottir, G.: Forcing of the wintertime atmospheric circulation by the multidecadal
6455 fluctuations of the North Atlantic ocean, *Environ. Res. Lett.*, 9, 034018, [https://doi.org/10.1088/1748-](https://doi.org/10.1088/1748-9326/9/3/034018)
6456 9326/9/3/034018, 2014.

6457 Peings, Y., and Magnusdottir, G.: Wintertime atmospheric response to Atlantic multidecadal variability: effect of
6458 stratospheric representation and ocean-atmosphere coupling, *Clim. Dyn.*, 47, 1029-1047,
6459 <https://doi.org/10.1007/s00382-015-2887-4>, 2016.

6460 Pekcan-Hekim, Z., Urho, L., Auvinen, H., Heikinheimo, O., Lappalainen, J., Raitaniemi, J., and Söderkultalahti,
6461 P.: Climate Warming and Pikeperch Year-Class Catches in the Baltic Sea, *Ambio*, 40, 447-456,
6462 <https://doi.org/10.1007/s13280-011-0143-7>, 2011.

6463 Pellikka, H., Särkkä, J., Johansson, M., and Pettersson, H.: Probability distributions for mean sea level and storm
6464 contribution up to 2100 AD at Forsmark, Swedish Nuclear Fuel and Waste Management Company (SKB), 49pp,
6465 2020.

6466 Pemberton, P., Löptien, U., Hordoir, R., Höglund, A., Schimanke, S., Axell, L., and Haapala, J.: Sea-ice
6467 evaluation of NEMO-Nordic 1.0: a NEMO-LIM3. 6-based ocean-sea-ice model setup for the North Sea and
6468 Baltic Sea, *Geosci. Model Dev.*, 10, 3105, <https://doi.org/10.5194/gmd-10-3105-2017>, 2017.

6469 Peng, S., Piao, S., Ciais, P., Friedlingstein, P., Zhou, L., and Wang, T.: Change in snow phenology and its
6470 potential feedback to temperature in the Northern Hemisphere over the last three decades, *Environ. Res. Lett.*, 8,
6471 014008, <https://doi.org/10.1088/1748-9326/8/1/014008>, 2013.

6472 Perry, D., Staveley, T., Deyanova, D., Baden, S., Dupont, S., Hernroth, B., Wood, H., Björk, M., and Gullström,
6473 M.: Global environmental changes negatively impact temperate seagrass ecosystems, *Ecosphere*, 10, e02986,
6474 <https://doi.org/10.1002/ecs2.2986>, 2019.

6475 Persson, T.: Solar radiation climate in Sweden, *Physics and Chemistry of the Earth, Part B: Hydrology, Oceans*
6476 *and Atmosphere*, 24, 275-279, [https://doi.org/10.1016/S1464-1909\(98\)00050-1](https://doi.org/10.1016/S1464-1909(98)00050-1), 1999.

6477 Peters, W., Bastos, A., Ciais, P., and Vermeulen, A.: A historical, geographical and ecological perspective on the
6478 2018 European summer drought, *Philos. T. Roy. Soc. B.*, 375, 20190505, <https://doi.org/10.1098/rstb.2019.0505>,
6479 2020.

6480 Pfeffer, W. T., Arendt, A. A., Bliss, A., Bolch, T., Cogley, J. G., Gardner, A. S., Hagen, J.-O., Hock, R., Kaser,
6481 G., Kienholz, C., Miles, E. S., Moholdt, G., Mölg, N., Paul, F., Radić, V., Rastner, P., Raup, B. H., Rich, J., and
6482 Sharp, M. J.: The Randolph Glacier Inventory: a globally complete inventory of glaciers, *J. Glaciol.*, 60, 537 -
6483 552, <https://doi.org/10.3189/2014JoG13J176>, 2014.

6484 Pfeifroth, U., Sanchez-Lorenzo, A., Manara, V., Trentmann, J., and Hollmann, R.: Trends and Variability of
6485 Surface Solar Radiation in Europe Based On Surface- and Satellite-Based Data Records, *J. Geophys. Res-*
6486 *Atmos.*, 123, 1735-1754, <https://doi.org/10.1002/2017JD027418>, 2018.

6487 Philipona, R., Behrens, K., and Ruckstuhl, C.: How declining aerosols and rising greenhouse gases forced rapid
6488 warming in Europe since the 1980s, *Geophys. Res. Lett.*, 36, <https://doi.org/10.1029/2008GL036350>, 2009.

6489 Pihlainen, S., Zandersen, M., Hyytiäinen, K., Andersen, H. E., Bartosova, A., Gustafsson, B., Jabloun, M.,
6490 McCrackin, M., Meier, H. E. M., Olesen, J. E., Saraiva, S., Swaney, D., and Thodsen, H.: Impacts of changing
6491 society and climate on nutrient loading to the Baltic Sea, *Sci. Total Environ.*, 138935,
6492 <https://doi.org/10.1016/j.scitotenv.2020.138935>, 2020.

6493 Pindsoo, K., and Soomere, T.: Basin-wide variations in trends in water level maxima in the Baltic Sea, *Cont.*
6494 *Shelf Res.*, 193, 104029, <https://doi.org/10.1016/j.csr.2019.104029>, 2020.

6495 Placke, M., Meier, H. E. M., Gräwe, U., Neumann, T., Frauen, C., and Liu, Y.: Long-Term Mean Circulation of
6496 the Baltic Sea as Represented by Various Ocean Circulation Models, *Front. Mar. Sci.*, 5,
6497 <https://doi.org/10.3389/fmars.2018.00287>, 2018.

6498 Placke, M., Meier, H. E. M., and Neumann, T.: Sensitivity of the Baltic Sea Overturning Circulation to Long-
6499 Term Atmospheric and Hydrological Changes, *J. Geophys. Res-Oceans*, 126, e2020JC016079,
6500 <https://doi.org/10.1029/2020JC016079>, 2021.

6501 Porz, L., Zhang, W., and Schrum, C.: Density-driven bottom currents control development of muddy basins in
6502 the southwestern Baltic Sea, *Mar. Geol.*, 438, 106523, <https://doi.org/10.1016/j.margeo.2021.106523>, 2021.

6503 Proietti, C., Fornasier, M. F., Sicard, P., Anav, A., Paoletti, E., and De Marco, A.: Trends in tropospheric ozone
6504 concentrations and forest impact metrics in Europe over the time period 2000–2014, *J. For. Res.*, 32, 543-551,
6505 <https://doi.org/10.1007/s11676-020-01226-3>, 2021.

6506 Ptak, M., Sojka, M., and Nowak, B.: Effect of climate warming on a change in thermal and ice conditions in the
6507 largest lake in Poland – Lake Śniardwy, *J. Hydrol. Hydromech.*, 68, 260-270, <https://doi.org/10.2478/johh-2020-0024>, 2020.

6509 Pugh, T. A. M., Lindeskog, M., Smith, B., Poulter, B., Arneth, A., Haverd, V., and Calle, L.: Role of forest
6510 regrowth in global carbon sink dynamics, *P. Natl. Acad. Sci. USA*, 116, 4382,
6511 <https://doi.org/10.1073/pnas.1810512116>, 2019.

6512 Qixiang, W., Wang, M., and Fan, X.: Seasonal patterns of warming amplification of high-elevation stations
6513 across the globe, *Int. J. Climatol.*, 38, 3466-3473, <https://doi.org/10.1002/joc.5509>, 2018.

6514 Quante, M., and Colijn, F.: North Sea region climate change assessment, *Regional Climate Studies*,
6515 SpringerOpen, Cham, 2016.

6516 Räämet, A., and Soomere, T.: The wave climate and its seasonal variability in the northeastern Baltic Sea, *Est. J.*
6517 *Earth Sci.*, 59, <https://doi.org/10.3176/earth.2010.1.08>, 2010.

6518 Rädler, A. T., Groenemeijer, P. H., Faust, E., Sausen, R., and Púčik, T.: Frequency of severe thunderstorms
6519 across Europe expected to increase in the 21st century due to rising instability, *npj Clim. Atmos. Sci.*, 2, 30,
6520 <https://doi.org/10.1038/s41612-019-0083-7>, 2019.

6521 Radtke, H., Brunnabend, S. E., Gräwe, U., and Meier, H. E. M.: Investigating interdecadal salinity changes in the
6522 Baltic Sea in a 1850–2008 hindcast simulation, *Clim. Past*, 16, 1617–1642, [https://doi.org/10.5194/cp-16-1617-](https://doi.org/10.5194/cp-16-1617-2020)
6523 2020, 2020.

6524 Rainio, K., Laaksonen, T., Ahola, M., Vähätalo, A. V., and Lehikoinen, E.: Climatic responses in spring
6525 migration of boreal and arctic birds in relation to wintering area and taxonomy, *J. Avian Biol.*, 37, 507–515,
6526 <https://doi.org/10.1111/j.0908-8857.2006.03740.x>, 2006.

6527 Räisänen, J.: Twenty-first century changes in snowfall climate in Northern Europe in ENSEMBLES regional
6528 climate models, *Clim. Dyn.*, 46, 339–353, <https://doi.org/10.1007/s00382-015-2587-0>, 2016.

6529 Räisänen, J.: Effect of atmospheric circulation on recent temperature changes in Finland, *Clim. Dyn.*, 53, 5675–
6530 5687, <https://doi.org/10.1007/s00382-019-04890-2>, 2019.

6531 [Räisänen, J.: Snow conditions in northern Europe: the dynamics of interannual variability versus projected long -](https://doi.org/10.5194/tc-15-1677-2021)
6532 [term change, *Cryosphere*, 15, 1677–1696, <https://doi.org/10.5194/tc-15-1677-2021>, 2021.](https://doi.org/10.5194/tc-15-1677-2021)

6533 Rajasilta, M., Hänninen, J., Laaksonen, L., Laine, P., Suomela, J.-P., Vuorinen, I., and Mäkinen, K.: Influence of
6534 environmental conditions, population density, and prey type on the lipid content in Baltic herring (*Clupea*
6535 *harengus membras*) from the northern Baltic Sea, *Can. J. Fish. Aquat. Sci.*, 76, 576–585,
6536 <https://doi.org/10.1139/cjfas-2017-0504>, 2018.

6537 Rajczak, J., and Schär, C.: Projections of Future Precipitation Extremes Over Europe: A Multimodel Assessment
6538 of Climate Simulations, *J. Geophys. Res-Atmos.*, 122, 10,773–710,800, <https://doi.org/10.1002/2017JD027176>,
6539 2017.

6540 Ramacher, M. O. P., Karl, M., Bieser, J., Jalkanen, J.-P., and Johansson, L.: Urban population exposure to
6541 NO₂ emissions from local shipping in three Baltic Sea harbour cities - a generic approach, *Atmos. Chem.*
6542 *Phys.*, 19, 9153, 2019.

6543 Ravestein, P., van der Schrier, G., Haarsma, R., Scheele, R., and van den Broek, M.: Vulnerability of European
6544 intermittent renewable energy supply to climate change and climate variability, *Renew. Sust. Energ. Rev.*, 97,
6545 497–508, <https://doi.org/10.1016/j.rser.2018.08.057>, 2018.

6546 Read, A. J., and Hohn, A. A.: LIFE IN THE FAST LANE: THE LIFE HISTORY OF HARBOR PORPOISES
6547 FROM THE GULF OF MAINE, *Mar. Mammal Sci.*, 11, 423–440, [https://doi.org/10.1111/j.1748-](https://doi.org/10.1111/j.1748-7692.1995.tb00667.x)
6548 7692.1995.tb00667.x, 1995.

6549 Reckermann, M., Langner, J., Omstedt, A., von Storch, H., Keevallik, S., Schneider, B., Arheimer, B., Meier, H.
6550 E. M., and Hünicke, B.: BALTEX—an interdisciplinary research network for the Baltic Sea region, *Environ.*
6551 *Res. Lett.*, 6, 045205, <http://doi.org/10.1088/1748-9326/6/4/045205>, 2011.

6552 Reckermann, M., Omstedt, A., Soomere, T., Aigars, J., Akhtar, N., Beldowski, J., Brauwer, C. P.-S. d., Cronin,
6553 T., Czub, M., Eero, M., Hyttiäinen, K., Jalkanen, J.-P., Kiessling, A., Kjellström, E., Larsén, X. G., McCrackin,
6554 M., Meier, H. E. M., Oberbeckmann, S., Parnell, K., Poska, A., Saarinen, J., Szymczycha, B., Undeman, E.,
6555 Viitasalo, M., Wörman, A., and Zorita, E.: Human impacts and their interactions in the Baltic Sea region, *Earth*
6556 *Syst. Dynam. Discuss.*, 1–127, <https://doi.org/10.5194/esd-2021-54>, 2021.

6557 Reder, S., Lydersen, C., Arnold, W., and Kovacs, K. M.: Haulout behaviour of High Arctic harbour seals (*Phoca*
6558 *vitulina vitulina*) in Svalbard, Norway, *Polar Biol.*, 27, 6–16, <https://doi.org/10.1007/s00300-003-0557-1>, 2003.

6559 Refsgaard, J. C., Hansen, A. L., Højberg, A. L., Olesen, J. E., Hashemi, F., Wachniew, P., Wörman, A.,
6560 Bartosova, A., Stelljes, N., and Chubarenko, B.: Spatially differentiated regulation: Can it save the Baltic Sea
6561 from excessive N-loads?, *Ambio*, 48, 1278–1289, <https://doi.org/10.1007/s13280-019-01195-w>, 2019.

6562 Reihan, A., Koltsova, T., Kriaciuniene, J., Lizuma, L., and Meilutyte-Barauskiene, D.: Changes in water
6563 discharges of the Baltic states rivers in the 20th century and its relation to climate change, *Hydrol. Res.*, 38, 401–
6564 412, <https://doi.org/10.2166/nh.2007.020>, 2007.

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Feldfunktion geändert

Formatiert: Englisch (USA)

Formatiert: Rechtschreibung und Grammatik nicht prüfen

6565 Reissmann, J. H., Burchard, H., Feistel, R., Hagen, E., Lass, H. U., Mohrholz, V., Nausch, G., Umlauf, L., and
6566 Wiecezorek, G.: Vertical mixing in the Baltic Sea and consequences for eutrophication—A review, *Prog.*
6567 *Oceanogr.*, 82, 47–80, <https://doi.org/10.1016/j.pocean.2007.10.004>, 2009.

6568 Rennert, K. J., and Wallace, J. M.: Cross-Frequency Coupling, Skewness, and Blocking in the Northern
6569 Hemisphere Winter Circulation, *J. Climate*, 22, 5650–5666, <https://doi.org/10.1175/2009jcli2669.1>, 2009.

6570 Reusch, T. B. H., Dierking, J., Andersson, H. C., Bonsdorff, E., Carstensen, J., Casini, M., Czajkowski, M.,
6571 Hasler, B., Hinsby, K., Hyytiäinen, K., Johannesson, K., Jomaa, S., Jormalainen, V., Kuosa, H., Kurland, S.,
6572 Laikre, L., MacKenzie, B. R., Margonski, P., Melzner, F., Oesterwind, D., Ojaveer, H., Refsgaard, J. C.,
6573 Sandström, A., Schwarz, G., Tonderski, K., Winder, M., and Zandersen, M.: The Baltic Sea as a time machine
6574 for the future coastal ocean, *Sci. Adv.*, 4, <https://doi.org/10.1126/sciadv.aar8195> 2018.

6575 RGI Consortium: Randolph Glacier Inventory – A Dataset of Global Glacier Outlines: Version 6.0: Technical
6576 Report, Global Land Ice Measurements from Space, Colorado, USA. Digital Media, [https://doi.org/10.7265/N5-](https://doi.org/10.7265/N5-RGI-60)
6577 [RGI-60](https://doi.org/10.7265/N5-RGI-60), 2017.

6578 Riahi, K., Rao, S., Krey, V., Cho, C., Chirkov, V., Fischer, G., Kindermann, G., Nakicenovic, N., and Rafaj, P.:
6579 RCP 8.5—A scenario of comparatively high greenhouse gas emissions, *Clim. Change*, 109, 33,
6580 <https://doi.org/10.1007/s10584-011-0149-y>, 2011.

6581 Ribeiro, A., Barbosa, S. M., Scotto, M. G., and Donner, R. V.: Changes in extreme sea-levels in the Baltic Sea,
6582 *Tellus A*, 66, 20921, <https://doi.org/10.3402/tellusa.v66.20921>, 2014.

6583 Ridefelt, H., Etzelmüller, B., Boelhouwers, J., and Jonasson, C.: Statistic-empirical modelling of mountain
6584 permafrost distribution in the Abisko region, sub-Arctic northern Sweden, *Norsk Geografisk Tidsskrift-*
6585 *Norwegian Journal of Geography*, 62, 278–289, <https://doi.org/10.1080/00291950802517890>, 2008.

6586 Riihelä, A., Carlund, T., Trentmann, J., Müller, R., and Lindfors, A. V.: Validation of CM SAF Surface Solar
6587 Radiation Datasets over Finland and Sweden, *Remote Sens.-Basel*, 7, 6663–6682,
6588 <https://doi.org/10.3390/rs70606663>, 2015.

6589 Rimkus, E., Kažys, J., Valiukas, D., and Stankūnavičius, G.: The atmospheric circulation patterns during dry
6590 periods in Lithuania, *Oceanologia*, 56, 223–239, <https://doi.org/10.5697/oc.56-2.223>, 2014.

6591 Rimkus, E., Briede, A., Jaagus, J., Stonevicius, E., Kilpys, J., and Viru, B.: Snow-cover regime in Lithuania,
6592 Latvia and Estonia and its relationship to climatic and geographical factors in 1961–2015, *Boreal Environ. Res.*,
6593 23, 193–208, 2018.

6594 Rinne, H., and Salovius-Laurén, S.: The status of brown macroalgae *Fucus* spp. and its relation to environmental
6595 variation in the Finnish marine area, northern Baltic Sea, *Ambio*, 49, 118–129, [https://doi.org/10.1007/s13280-](https://doi.org/10.1007/s13280-019-01175-0)
6596 [019-01175-0](https://doi.org/10.1007/s13280-019-01175-0), 2020.

6597 Rinne, J., Tuovinen, J.-P., Klemetsson, L., Aurela, M., Holst, J., Lohila, A., Weslien, P., Vestin, P., Łakomiec,
6598 P., Peichl, M., Tuittila, E.-S., Heiskanen, L., Laurila, T., Li, X., Alekseychik, P., Mammarella, I., Ström, L.,
6599 Crill, P., and Nilsson, M. B.: Effect of the 2018 European drought on methane and carbon dioxide exchange of
6600 northern mire ecosystems, *Philos. T. Roy. Soc. B.*, 375, 20190517, <https://doi.org/10.1098/rstb.2019.0517>, 2020.

6601 Ripszám, M., Paczkowska, J., Figueira, J., Veenaas, C., and Haglund, P.: Dissolved Organic Carbon Quality and
6602 Sorption of Organic Pollutants in the Baltic Sea in Light of Future Climate Change, *Environ. Sci. Technol.*, 49,
6603 1445–1452, <https://doi.org/10.1021/es504437s>, 2015.

6604 Rizzi, J., Nilsen, I. B., Stagge, J. H., Gislås, K., and Tallaksen, L. M.: Five decades of warming: impacts on
6605 snow cover in Norway, *Hydrol. Res.*, 49, 670–688, <https://doi.org/10.2166/nh.2017.051>, 2017.

6606 Rjazin, J., and Pärn, O.: Determining the Regime Shift of the Baltic Sea Ice Seasons during 1982–2016, *NAŠE*
6607 *MORE: znanstveni časopis za more i pomorstvo*, 67, 53–59, 2020.

6608 Rodhe, J.: The Baltic and North Seas: A process-oriented review of the physical oceanography. Coastal segment.
6609 , in: *The Sea, Volume 11: The Global Coastal Ocean: Regional Studies and Syntheses*, edited by: Robinson, A.
6610 R., and Brink, K. H., John Wiley & Sons, Inc., 699–732, 1998.

- 6611 Rodhe, J., Tett, P., and Wulff, F.: The Baltic and North Seas: A Regional Review of some important Physical-
6612 Chemical-Biological Interaction Processes, in: *The Sea, Volume 14B: The Global Coastal Ocean*, edited by:
6613 Robinson, A. R., and Brink, K. H., Harvard University Press, Cambridge, MA, USA, 1033-1075, 2006.
- 6614 Rodwell, M. J., Rowell, D. P., and Folland, C. K.: Oceanic forcing of the wintertime North Atlantic Oscillation
6615 and European climate, *Nature*, 398, 320-323, <https://doi.org/10.1038/18648>, 1999.
- 6616 Roff, J. C., and Legendre, L.: Chapter 14 Physico-Chemical and Biological Oceanography of Hudson Bay, in:
6617 *Elsevier Oceanography Series*, edited by: Martini, I. P., Elsevier, 265-292, [https://doi.org/10.1016/S0422-](https://doi.org/10.1016/S0422-9894(08)70907-3)
6618 9894(08)70907-3, 1986.
- 6619 Röhr, M. E., Boström, C., Canal-Vergés, P., and Holmer, M.: Blue carbon stocks in Baltic Sea eelgrass (*Zostera*
6620 *marina*) meadows, *Biogeosciences*, 13, 6139-6153, <https://doi.org/10.5194/bg-13-6139-2016>, 2016.
- 6621 Rolff, C., Almesjö, L., and Elmgren, R.: Nitrogen fixation and abundance of the diazotrophic cyanobacterium
6622 *Aphanizomenon* sp. in the Baltic Proper, *Mar. Ecol. Prog. Ser.*, 332, 107-118,
6623 <https://doi.org/10.3354/meps332107>, 2007.
- 6624 Rolff, C., and Elfving, T.: Increasing nitrogen limitation in the Bothnian Sea, potentially caused by inflow of
6625 phosphate-rich water from the Baltic Proper, *Ambio*, 44, 601-611, <https://doi.org/10.1007/s13280-015-0675-3>,
6626 2015.
- 6627 Ronkainen, I., Lehtiranta, J., Lensu, M., Rinne, E., Haapala, J., and Haas, C.: Interannual sea ice thickness
6628 variability in the Bay of Bothnia, *Cryosphere*, 12, 3459-3476, <https://doi.org/10.5194/tc-12-3459-2018>, 2018.
- 6629 Rothäusler, E., Rugiu, L., and Jormalainen, V.: Forecast climate change conditions sustain growth and
6630 physiology but hamper reproduction in range-margin populations of a foundation rockweed species, *Mar.*
6631 *Environ. Res.*, 141, 205-213, <https://doi.org/10.1016/j.marenvres.2018.09.014>, 2018.
- 6632 Roudier, P., Andersson, J. C. M., Donnelly, C., Feyen, L., Greuell, W., and Ludwig, F.: Projections of future
6633 floods and hydrological droughts in Europe under a +2°C global warming, *Clim. Change*, 135, 341-355,
6634 <https://doi.org/10.1007/s10584-015-1570-4>, 2016.
- 6635 Rounsevell, M. D. A., Ewert, F., Reginster, I., Leemans, R., and Carter, T. R.: Future scenarios of European
6636 agricultural land use: II. Projecting changes in cropland and grassland, *Agric. Ecosyst. Environ.*, 107, 117-135,
6637 <https://doi.org/10.1016/j.agee.2004.12.002>, 2005.
- 6638 Rousi, H., Korpinen, S., and Bonsdorff, E.: Brackish-Water Benthic Fauna Under Fluctuating Environmental
6639 Conditions: The Role of Eutrophication, Hypoxia, and Global Change, *Front. Mar. Sci.*, 6,
6640 <https://doi.org/10.3389/fmars.2019.00464>, 2019.
- 6641 Ruckstuhl, C., Philipona, R., Behrens, K., Collaud Coen, M., Dürr, B., Heimo, A., Mätzler, C., Nyeki, S.,
6642 Ohmura, A., Vuilleumier, L., Weller, M., Wehrli, C., and Zelenka, A.: Aerosol and cloud effects on solar
6643 brightening and the recent rapid warming, *Geophys. Res. Lett.*, 35, L12708,
6644 <https://doi.org/10.1029/2008GL034228>, 2008.
- 6645 Rugiu, L., Manninen, I., Sjöroos, J., and Jormalainen, V.: Variations in tolerance to climate change in a key
6646 littoral herbivore, *Mar. Biol.*, 165, 18, <https://doi.org/10.1007/s00227-017-3275-x>, 2017.
- 6647 Ruoho-Airola, T., Eilola, K., Savchuk, O. P., Parviainen, M., and Tarvainen, V.: Atmospheric Nutrient Input to
6648 the Baltic Sea from 1850 to 2006: A Reconstruction from Modeling Results and Historical Data, *Ambio* 41, 549-
6649 557, <https://doi.org/10.1007/s13280-012-0319-9>, 2012.
- 6650 Ruosteenoja, K., Vihma, T., and Venäläinen, A.: Projected Changes in European and North Atlantic Seasonal
6651 Wind Climate Derived from CMIP5 Simulations, *J. Climate*, 32, 6467-6490, [https://doi.org/10.1175/JCLI-D-19-](https://doi.org/10.1175/JCLI-D-19-0023.1)
6652 0023.1, 2019.
- 6653 Ruosteenoja, K., Markkanen, T., and Räisänen, J.: Thermal seasons in northern Europe in projected future
6654 climate, *Int. J. Climatol.*, 40, 4444-4462, <https://doi.org/10.1002/joc.6466>, 2020.
- 6655 Ruprich-Robert, Y., Msadek, R., Castruccio, F., Yeager, S., Delworth, T., and Danabasoglu, G.: Assessing the
6656 Climate Impacts of the Observed Atlantic Multidecadal Variability Using the GFDL CM2.1 and NCAR CESM1
6657 Global Coupled Models, *J. Climate*, 30, 2785-2810, <https://doi.org/10.1175/jcli-d-16-0127.1>, 2017.

6658 Russak, V.: Changes in solar radiation and their influence on temperature trend in Estonia (1955–2007), *J.*
6659 *Geophys. Res.-Atmos.*, 114, <https://doi.org/10.1029/2008JD010613>, 2009.

6660 Rutgersson, A., Jaagus, J., Schenk, F., and Stendel, M.: Observed changes and variability of atmospheric
6661 parameters in the Baltic Sea region during the last 200 years, *Clim. Res.*, 61, 177–190,
6662 <https://doi.org/10.3354/cr01244>, 2014.

6663 Rutgersson, A., Kjellström, E., Haapala, J., Stendel, M., Danilovich, I., Drews, M., Jylhä, K., Kujala, P., Larsén,
6664 X. G., Halsnæs, K., Lehtonen, I., Luomaranta, A., Nilsson, E., Olsson, T., Särkkä, J., Tuomi, L., and Wasmund,
6665 N.: Natural Hazards and Extreme Events in the Baltic Sea region, *Earth Syst. Dynam. Discuss.*, 1–80,
6666 <https://doi.org/10.5194/esd-2021-13>, 2021.

6667 Ryabchenko, V. A., Karlin, L. N., Isaev, A. V., Vankevich, R. E., Eremina, T. R., Molchanov, M. S., and
6668 Savchuk, O. P.: Model estimates of the eutrophication of the Baltic Sea in the contemporary and future climate,
6669 *Oceanology+*, 56, 36–45, <https://doi.org/10.1134/S0001437016010161>, 2016.

6670 Ryabchuk, D., Kolesov, A., Chubarenko, B., Spiridonov, M., Kurennoy, D., and Soomere, T.: Coastal erosion
6671 processes in the eastern Gulf of Finland and their links with geological and hydrometeorological factors, *Boreal*
6672 *Environ. Res.*, 16 (Suppl. A), 117–137, 2011.

6673 Sahla, M., Tolvanen, H., Ruuskanen, A., and Kurvinen, L.: Assessing long term change of *Fucus* spp.
6674 communities in the northern Baltic Sea using monitoring data and spatial modeling, *Estuar. Coast. Shelf Sci.*,
6675 245, 107023, <https://doi.org/10.1016/j.ecss.2020.107023>, 2020.

6676 Salo, T., Mattila, J., and Eklöf, J.: Long-term warming affects ecosystem functioning through species turnover
6677 and intraspecific trait variation, *Oikos*, 129, 283–295, <https://doi.org/10.1111/oik.06698>, 2020.

6678 Sannel, A. B. K., Hugelius, G., Jansson, P., and Kuhry, P.: Permafrost Warming in a Subarctic Peatland – Which
6679 Meteorological Controls are Most Important?, *Permafrost Periglac.*, 27, 177–188,
6680 <https://doi.org/10.1002/ppp.1862>, 2016.

6681 Saraiva, S., Meier, H. E. M., Andersson, H., Höglund, A., Dieterich, C., Gröger, M., Hordoir, R., and Eilola, K.:
6682 Baltic Sea ecosystem response to various nutrient load scenarios in present and future climates, *Clim. Dyn.*, 52,
6683 3369–3387, <https://doi.org/10.1007/s00382-018-4330-0>, 2019a.

6684 Saraiva, S., Meier, H. E. M., Andersson, H., Höglund, A., Dieterich, C., Gröger, M., Hordoir, R., and Eilola, K.:
6685 Uncertainties in Projections of the Baltic Sea Ecosystem Driven by an Ensemble of Global Climate Models,
6686 *Front. Earth Sci.*, 6, <https://doi.org/10.3389/feart.2018.00244>, 2019b.

6687 Sarauskiene, D., Kriauciuniene, J., Reihan, A., and Klavins, M.: Flood pattern changes in the rivers of the Baltic
6688 countries, *J. Environ. Eng. Landsc. Manag.*, 23, 28–38, <https://doi.org/10.3846/16486897.2014.937438>, 2015.

6689 Šarauskiene, D., Akstinas, V., Kriauciūnienė, J., Jakimavičius, D., Bukantis, A., Kažys, J., Povilaitis, A., Ložys,
6690 L., Kesminas, V., Virbickas, T., and Pliauraitė, V.: Projection of Lithuanian river runoff, temperature and their
6691 extremes under climate change, *Hydrol. Res.*, 49, 344–362, <https://doi.org/10.2166/nh.2017.007>, 2017.

6692 Savchuk, O. P., Wulff, F., Hille, S., Humborg, C., and Pollehne, F.: The Baltic Sea a century ago — a
6693 reconstruction from model simulations, verified by observations, *J. Mar. Syst.*, 74, 485–494,
6694 <https://doi.org/10.1016/j.jmarsys.2008.03.008>, 2008.

6695 Savchuk, O. P., Gustafsson, B. G., Rodriguez Medina, M., Sokolov, A. V., and Wulff, F.: External nutrient loads
6696 to the Baltic Sea, 1970–2006, *Baltic Nest Institute Techn. Rep.*, 22, 2012.

6697 Savchuk, O. P.: Large-Scale Nutrient Dynamics in the Baltic Sea, 1970–2016, *Front. Mar. Sci.*, 5, 95,
6698 <https://doi.org/10.3389/fmars.2018.00095>, 2018.

6699 Scaife, A. A., and Smith, D.: A signal-to-noise paradox in climate science, *npj Clim. Atmos. Sci.*, 1, 28,
6700 <https://doi.org/10.1038/s41612-018-0038-4>, 2018.

6701 Scheff, J., and Frierson, D. M. W.: Robust future precipitation declines in CMIP5 largely reflect the poleward
6702 expansion of model subtropical dry zones, *Geophys. Res. Lett.*, 39, <https://doi.org/10.1029/2012GL052910>,
6703 2012.

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Feldfunktion geändert

Formatiert: Englisch (USA)

Formatiert: Rechtschreibung und Grammatik nicht prüfen

6704 Schenk, F., and Zorita, E.: Reconstruction of high resolution atmospheric fields for Northern Europe using
6705 analog-upscaling, *Clim. Past Discuss.*, 8, 819-868, <https://doi.org/10.5194/cp-8-1681-2012>, 2012.

6706 Schenk, F.: The analog-method as statistical upscaling tool for meteorological field reconstructions over
6707 Northern Europe since 1850, Doctoral Dissertation, University of Hamburg, Hamburg, 2015.

6708 [Schiemann, R., Athanasiadis, P., Barriopedro, D., Doblas-Reyes, F., Lohmann, K., Roberts, M. J., Sein, D. V.,](#)
6709 [Roberts, C. D., Terray, L., and Vidale, P. L.: Northern Hemisphere blocking simulation in current climate](#)
6710 [models: evaluating progress from the Climate Model Intercomparison Project Phase 5 to 6 and sensitivity to](#)
6711 [resolution, *Weather Clim. Dynam.*, 1, 277-292, <https://doi.org/10.5194/wcd-1-277-2020>, 2020.](#)

6712 Schimanke, S., Meier, H. E. M., Kjellström, E., Strandberg, G., and Hordoir, R.: The climate in the Baltic Sea
6713 region during the last millennium simulated with a regional climate model, *Clim. Past*, 8, 1419,
6714 <https://doi.org/10.5194/cp-8-1419-2012>, 2012.

6715 Schimanke, S., Dieterich, C., and Meier, H. E. M.: An algorithm based on sea-level pressure fluctuations to
6716 identify major Baltic inflow events, *Tellus A*, 66, 23452, <http://dx.doi.org/10.3402/tellusa.v66.23452>, 2014.

6717 Schimanke, S., and Meier, H. E. M.: Decadal-to-Centennial Variability of Salinity in the Baltic Sea, *J. Climate*,
6718 29, 7173-7188, <https://doi.org/10.1175/JCLI-D-15-0443.1>, 2016.

6719 Schinke, H., and Matthäus, W.: On the causes of major Baltic inflows - an analysis of long time series, *Cont.*
6720 *Shelf Res.*, 18, 67-97, [https://doi.org/10.1016/S0278-4343\(97\)00071-X](https://doi.org/10.1016/S0278-4343(97)00071-X), 1998.

6721 Schmidt, G. A., Jungclauss, J. H., Ammann, C. M., Bard, E., Braconnot, P., Crowley, T. J., Delaygue, G., Joos,
6722 F., Krivova, N. A., Muscheler, R., Otto-Bliesner, B. L., Pongratz, J., Shindell, D. T., Solanki, S. K., Steinhilber,
6723 F., and Vieira, L. E. A.: Climate forcing reconstructions for use in PMIP simulations of the last millennium
6724 (v1.0), *Geosci. Model Dev.*, 4, 33-45, <https://doi.org/10.5194/gmd-4-33-2011>, 2011.

6725 Schmidt, K., Birchill, A. J., Atkinson, A., Brewin, R. J. W., Clark, J. R., Hickman, A. E., Johns, D. G., Lohan,
6726 M. C., Milne, A., Pardo, S., Polimene, L., Smyth, T. J., Tarran, G. A., Widdicombe, C. E., Woodward, E. M. S.,
6727 and Ussher, S. J.: Increasing picocyanobacteria success in shelf waters contributes to long-term food web
6728 degradation, *Global Change Biol.*, 26, 5574-5587, <https://doi.org/10.1111/gcb.15161>, 2020.

6729 Schneider, B., and Müller, J. D.: Biogeochemical Transformations in the Baltic Sea. Observations Through
6730 Carbon Dioxide Glasses. , 110, Springer International Publishing, Cham, 2018.

6731 Schönhofer, J., and Dudkowska, A.: Rip currents in the southern Baltic Sea multi-bar nearshore zone, *Cont.*
6732 *Shelf Res.*, 212, 104324, <https://doi.org/10.1016/j.csr.2020.104324>, 2021.

6733 Schröder, W., Nickel, S., Schönrock, S., Meyer, M., Wosniok, W., Harmens, H., Frontasyeva, M. V., Alber, R.,
6734 Aleksiyenak, J., Barandovski, L., Carballeira, A., Danielsson, H., de Temmermann, L., Godzik, B., Jeran, Z.,
6735 Karlsson, G. P., Lazo, P., Leblond, S., Lindroos, A.-J., Liiv, S., Magnússon, S. H., Mankovska, B., Martínez-
6736 Abaigar, J., Piispanen, J., Poikolainen, J., Popescu, I. V., Qarri, F., Santamaria, J. M., Skudnik, M., Špirić, Z.,
6737 Stafilov, T., Steinnes, E., Stihl, C., Thöni, L., Uggerud, H. T., and Zechmeister, H. G.: Spatially valid data of
6738 atmospheric deposition of heavy metals and nitrogen derived by moss surveys for pollution risk assessments of
6739 ecosystems, *Environ. Sci. Pollut. Res.*, 23, 10457-10476, <https://doi.org/10.1007/s11356-016-6577-5>, 2016.

6740 Schubert, S. D., Wang, H., Koster, R. D., Suarez, M. J., and Groisman, P. Y.: Northern Eurasian Heat Waves
6741 and Droughts, *J. Climate*, 27, 3169-3207, <https://doi.org/10.1175/jcli-d-13-00360.1>, 2014.

6742 Schuster, P. F., Schaefer, K. M., Aiken, G. R., Antweiler, R. C., Dewild, J. F., Gryziec, J. D., Gusmeroli, A.,
6743 Hugelius, G., Jafarov, E., Krabbenhoft, D. P., Liu, L., Herman-Mercer, N., Mu, C., Roth, D. A., Schaefer, T.,
6744 Striegl, R. G., Wickland, K. P., and Zhang, T.: Permafrost Stores a Globally Significant Amount of Mercury,
6745 *Geophys. Res. Lett.*, 45, 1463-1471, <https://doi.org/10.1002/2017GL075571>, 2018.

6746 Screen, J. A., Simmonds, I., Deser, C., and Tomas, R.: The Atmospheric Response to Three Decades of
6747 Observed Arctic Sea Ice Loss, *J. Climate*, 26, 1230-1248, <https://doi.org/10.1175/jcli-d-12-00063.1>, 2013.

6748 Seager, R., Naik, N., and Vecchi, G. A.: Thermodynamic and Dynamic Mechanisms for Large-Scale Changes in
6749 the Hydrological Cycle in Response to Global Warming*, *J. Climate*, 23, 4651-4668,
6750 <https://doi.org/10.1175/2010JCLI3655.1>, 2010.

6751 Seinä, A., and Palosuo, E.: The classification of the maximum annual extent of ice cover in the Baltic Sea 1720–
6752 1995, *Meri*, 27, 79–91, 1996.

6753 Seneviratne, S. I., Wilhelm, M., Stanelle, T., van den Hurk, B., Hagemann, S., Berg, A., Cheruy, F., Higgins, M.
6754 E., Meier, A., Brovkin, V., Claussen, M., Ducharne, A., Dufresne, J.-L., Findell, K. L., Ghattas, J., Lawrence, D.
6755 M., Malyshev, S., Rummukainen, M., and Smith, B.: Impact of soil moisture-climate feedbacks on CMIP5
6756 projections: First results from the GLACE-CMIP5 experiment, *Geophys. Res. Lett.*, 40, 5212–5217,
6757 <https://doi.org/10.1002/grl.50956>, 2013.

6758 [Seneviratne, S. I., and Hauser, M.: Regional Climate Sensitivity of Climate Extremes in CMIP6 Versus CMIP5
6759 Multimodel Ensembles, 8, e2019EF001474, <https://doi.org/10.1029/2019EF001474>, 2020.](#)

6760 Seppälä, M.: The origin of palsas, *Geogr. Ann. A*, 68, 141–147, <https://doi.org/10.2307/521453>, 1986.

6761 Sharma, S., Magnuson, J. J., Batt, R. D., Winslow, L. A., Korhonen, J., and Aono, Y.: Direct observations of ice
6762 seasonality reveal changes in climate over the past 320–570 years, *Sci. Rep.-UK*, 6, 25061,
6763 <https://doi.org/10.1038/srep25061>, 2016.

6764 Sharma, S., Blagrove, K., Magnuson, J. J., O'Reilly, C. M., Oliver, S., Batt, R. D., Magee, M. R., Straile, D.,
6765 Weyhenmeyer, G. A., Winslow, L., and Woolway, R. I.: Widespread loss of lake ice around the Northern
6766 Hemisphere in a warming world, *Nat. Clim. Change*, 9, 227–231, <https://doi.org/10.1038/s41558-018-0393-5>,
6767 2019.

6768 Sharma, S., Meyer, M. F., Culpepper, J., Yang, X., Hampton, S., Berger, S. A., Brouil, M. R., Fradkin, S. C.,
6769 Higgins, S. N., Jankowski, K. J., Kirillin, G., Smits, A. P., Whitaker, E. C., Yousef, F., and Zhang, S.:
6770 Integrating Perspectives to Understand Lake Ice Dynamics in a Changing World, *J. Geophys. Res.-Biogeo*, 125,
6771 e2020JG005799, <https://doi.org/10.1029/2020JG005799>, 2020.

6772 Sharma, S., Blagrove, K., Filazzola, A., Imrit, M. A., and Hendricks Franssen, H.-J.: Forecasting the Permanent
6773 Loss of Lake Ice in the Northern Hemisphere Within the 21st Century, *Geophys. Res. Lett.*, 48,
6774 e2020GL091108, <https://doi.org/10.1029/2020GL091108>, 2021.

6775 Shatwell, T., Thiery, W., and Kirillin, G.: Future projections of temperature and mixing regime of European
6776 temperate lakes, *Hydrol. Earth Syst. Sci.*, 23, 1533–1551, <https://doi.org/10.5194/hess-23-1533-2019>, 2019.

6777 Shaw, T. A., Baldwin, M., Barnes, E. A., Caballero, R., Garfinkel, C. I., Hwang, Y. T., Li, C., O'Gorman, P. A.,
6778 Rivière, G., Simpson, I. R., and Voigt, A.: Storm track processes and the opposing influences of climate change,
6779 *Nat. Geosci.*, 9, 656–664, <https://doi.org/10.1038/ngeo2783>, 2016.

6780 Shepherd, T. G., Boyd, E., Calel, R. A., Chapman, S. C., Dessai, S., Dima-West, I. M., Fowler, H. J., James, R.,
6781 Maraun, D., Martius, O., Senior, C. A., Sobel, A. H., Stainforth, D. A., Tett, S. F. B., Trenberth, K. E., van den
6782 Hurk, B. J. J. M., Watkins, N. W., Wilby, R. L., and Zenghelis, D. A.: Storylines: an alternative approach to
6783 representing uncertainty in physical aspects of climate change, *Clim. Change*, 151, 555–571,
6784 <https://doi.org/10.1007/s10584-018-2317-9>, 2018.

6785 [Sherwood, S. C., Webb, M. J., Annan, J. D., Armour, K. C., Forster, P. M., Hargreaves, J. C., Hegerl, G., Klein,
6786 S. A., Marvel, K. D., Rohling, E. J., Watanabe, M., Andrews, T., Braconnot, P., Bretherton, C. S., Foster, G. L.,
6787 Hausfather, Z., von der Heydt, A. S., Knutti, R., Mauritsen, T., Norris, J. R., Proistosescu, C., Rugenstein, M.,
6788 Schmidt, G. A., Tokarska, K. B., and Zelinka, M. D.: An Assessment of Earth's Climate Sensitivity Using
6789 Multiple Lines of Evidence, *Rev. Geophys.*, 58, e2019RG000678, <https://doi.org/10.1029/2019RG000678>, 2020.](#)

6790 Siegel, H., and Gerth, M.: Sea Surface Temperature in the Baltic Sea 2018, HELCOM Baltic Sea Environment
6791 Fact Sheets 2019, 7pp, [https://helcom.fi/wp-content/uploads/2020/07/BSEFS-Sea-Surface-Temperature-in-the-
6792 Baltic-Sea-2018.pdf](https://helcom.fi/wp-content/uploads/2020/07/BSEFS-Sea-Surface-Temperature-in-the-Baltic-Sea-2018.pdf), 2019.

6793 Sillmann, J., Kharin, V. V., Zwiers, F. W., Zhang, X., and Bronaugh, D.: Climate extremes indices in the CMIP5
6794 multimodel ensemble: Part 2. Future climate projections, 118, 2473–2493, <https://doi.org/10.1002/jgrd.50188>,
6795 2013.

6796 Sinclair, V. A., Rantanen, M., Haapanala, P., Räisänen, J., and Järvinen, H.: The characteristics and structure of
6797 extra-tropical cyclones in a warmer climate, *Weather Clim. Dynam.*, 1, 1–25, 10.5194/wcd-1-1-2020, 2020.

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Feldfunktion geändert

Formatiert: Englisch (USA)

Formatiert: Rechtschreibung und Grammatik nicht prüfen

6798 Sinha, E., Michalak, A. M., and Balaji, V.: Eutrophication will increase during the 21st century as a result of
6799 precipitation changes, *Science*, 357, 405, <https://doi.org/10.1126/science.aan2409>, 2017.

6800 Skogen, M. D., Eilola, K., Hansen, J. L. S., Meier, H. E. M., Molchanov, M. S., and Ryabchenko, V. A.:
6801 Eutrophication status of the North Sea, Skagerrak, Kattegat and the Baltic Sea in present and future climates: A
6802 model study, *J. Mar. Syst.*, 132, 174-184, <https://doi.org/10.1016/j.jmarsys.2014.02.004>, 2014.

6803 Skov, H., Heinänen S., Žydelis R., Bellebaum J., Bzoma S., Dagys M., Durinck J., Garthe S., Grishanov G., Hario M,
6804 Kieckbusch JJ, Kube J, Kuresoo A, Larsson K, Luigujoe L, Meissner W, Nehls HW, Nilsson L, Petersen IK,
6805 Roos MM, Pihl S, Sonntag N, Stock A, and A, S.: Waterbird populations and pressures in the Baltic Sea, Nordic
6806 Council of Ministers, Copenhagen, 2011.

6807 Smith, B., Prentice, I. C., and Sykes, M. T.: Representation of Vegetation Dynamics in the Modelling of
6808 Terrestrial Ecosystems: Comparing Two Contrasting Approaches within European Climate Space, *Global Ecol.*
6809 *Biogeogr.*, 10, 621-637, 2001.

6810 Smith, B., Aasa, A., Ahas, R., Blenckner, T., Callaghan, T. V., de Chazal, J., Humborg, C., Jönsson, A. M.,
6811 Kellomäki, S., Kull, A., Lehtikoinen, E., Mander, Ü., Nöges, P., Nöges, T., Rounsevell, M., Sofiev, M.,
6812 Tryjanowski, P., and Wolf, A.: Climate-related Change in Terrestrial and Freshwater Ecosystems, in:
6813 Assessment of Climate Change for the Baltic Sea Basin, edited by: BACC Author Team, Springer Berlin
6814 Heidelberg, Berlin, Heidelberg, 221-308, https://doi.org/10.1007/978-3-540-72786-6_4, 2008.

6815 Smith, B., Samuelsson, P., Wramneby, A., and Rummukainen, M.: A model of the coupled dynamics of climate,
6816 vegetation and terrestrial ecosystem biogeochemistry for regional applications, *Tellus A*, 63, 87-106,
6817 <https://doi.org/10.1111/j.1600-0870.2010.00477.x>, 2011.

6818 Smith, T. G., and Stirling, I.: The breeding habitat of the ringed seal (*Phoca hispida*). The birth lair and
6819 associated structures, *Can. J. Zool.*, 53, 1297-1305, <https://doi.org/10.1139/z75-155>, 1975.

6820 Smol, J. P., Wolfe, A. P., Birks, H. J. B., Douglas, M. S. V., Jones, V. J., Korhola, A., Pienitz, R., Rühland, K.,
6821 Sorvari, S., Antoniades, D., Brooks, S. J., Fallu, M.-A., Hughes, M., Keatley, B. E., Laing, T. E., Michelutti, N.,
6822 Nazarova, L., Nyman, M., Paterson, A. M., Perren, B., Quinlan, R., Rautio, M., Saulnier-Talbot, É., Siitonen, S.,
6823 Solovieva, N., and Weckström, J.: Climate-driven regime shifts in the biological communities of arctic lakes, *P.*
6824 *Natl. Acad. Sci. USA*, 102, 4397, <https://doi.org/10.1073/pnas.0500245102>, 2005.

6825 Snickars, M., Weigel, B., and Bonsdorff, E.: Impact of eutrophication and climate change on fish and
6826 zoobenthos in coastal waters of the Baltic Sea, *Mar. Biol.*, 162, 141-151, [https://doi.org/10.1007/s00227-014-](https://doi.org/10.1007/s00227-014-2579-3)
6827 2579-3, 2015.

6828 Sommer, U., and Lewandowska, A.: Climate change and the phytoplankton spring bloom: warming and
6829 overwintering zooplankton have similar effects on phytoplankton, *Global Change Biol.*, 17, 154-162,
6830 <https://doi.org/10.1111/j.1365-2486.2010.02182.x>, 2011.

6831 Sommer, U., Aberle, N., Lengfellner, K., and Lewandowska, A.: The Baltic Sea spring phytoplankton bloom in
6832 a changing climate: an experimental approach, *Mar. Biol.*, 159, 2479-2490, [https://doi.org/10.1007/s00227-012-](https://doi.org/10.1007/s00227-012-1897-6)
6833 1897-6, 2012.

6834 Sommer, U., Paul, C., and Moustaka-Gouni, M.: Warming and Ocean Acidification Effects on Phytoplankton—
6835 From Species Shifts to Size Shifts within Species in a Mesocosm Experiment, *PLoS ONE*, 10, e0125239,
6836 <https://doi.org/10.1371/journal.pone.0125239>, 2015.

6837 Soomere, T., Behrens, A., Tuomi, L., and Nielsen, J. W.: Wave conditions in the Baltic Proper and in the Gulf of
6838 Finland during windstorm Gudrun, *Natural Hazards and Earth System Sciences*, 8, 37-46,
6839 <https://doi.org/10.5194/nhess-8-37-2008>, 2008.

6840 Soomere, T., and Pindsoo, K.: Spatial variability in the trends in extreme storm surges and weekly-scale high
6841 water levels in the eastern Baltic Sea, *Cont. Shelf Res.*, 115, 53-64, <https://doi.org/10.1016/j.csr.2015.12.016>,
6842 2016.

6843 Soomere, T., and Räämet, A.: Spatial patterns of the wave climate in the Baltic Proper and the Gulf of
6844 Finland**This study was supported by the Estonian Science Foundation (grant No. 7413), targeted financing by
6845 the Estonian Ministry of Education and Research (grants SF0140077s08 and SF0140007s11) and partially
6846 performed within the framework of the BalticWay project, which is supported by funding from the European

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Feldfunktion geändert

Formatiert: Englisch (USA)

Formatiert: Rechtschreibung und Grammatik nicht prüfen

Community's Seventh Framework Programme (FP/2007-2013) under grant agreement No. 217246 made with the joint Baltic Sea research and development programme BONUS, *Oceanologia*, 53, 335-371, <https://doi.org/10.5697/oc.53-1-TI.335>, 2011.

Soomere, T., Viška, M., and Pindsoo, K.: Retrieving the Signal of Climate Change from Numerically Simulated Sediment Transport Along the Eastern Baltic Sea Coast, in: *Coastline Changes of the Baltic Sea from South to East: Past and Future Projection*, edited by: Harff, J., Furmańczyk, K., and von Storch, H., Springer International Publishing, Cham, 327-361, https://doi.org/10.1007/978-3-319-49894-2_15, 2017.

Sørland, S. L., Schär, C., Lüthi, D., and Kjellström, E.: Bias patterns and climate change signals in GCM-RCM model chains, *Environ. Res. Lett.*, 13, 074017, <https://doi.org/10.1088/1748-9326/aacc77>, 2018.

Sousa, P. M., Trigo, R. M., Barriopedro, D., Soares, P. M. M., Ramos, A. M., and Liberato, M. L. R.: Responses of European precipitation distributions and regimes to different blocking locations, *Clim. Dyn.*, 48, 1141-1160, <https://doi.org/10.1007/s00382-016-3132-5>, 2017.

Stahl, K., Hisdal, H., Hannaford, J., Tallaksen, L. M., van Lanen, H. A. J., Sauquet, E., Demuth, S., Fendekova, M., and Jódar, J.: Streamflow trends in Europe: evidence from a dataset of near-natural catchments, *Hydrol. Earth Syst. Sci.*, 14, 2367-2382, <https://doi.org/10.5194/hess-14-2367-2010>, 2010.

Stålnacke, P., Grimvall, A., Sundblad, K., and Tonderski, A.: Estimation of riverine loads of nitrogen and phosphorus to the Baltic Sea, 1970-1993, *Environ. Monit. Assess.*, 58, 173-200, <https://doi.org/10.1023/a:1006073015871>, 1999.

Stanhill, G., Achiman, O., Rosa, R., and Cohen, S.: The cause of solar dimming and brightening at the Earth's surface during the last half century: Evidence from measurements of sunshine duration, *J. Geophys. Res.-Atmos.*, 119, 9102-9109, <https://doi.org/10.1002/2013JD021308>, 2014.

Stendel, M., Francis, J., White, R., Williams, P. D., and Woollings, T.: Chapter 15 - The jet stream and climate change, in: *Climate Change (Third Edition)*, edited by: Letcher, T. M., Elsevier, 327-357, <https://doi.org/10.1016/B978-0-12-821575-3.00015-3>, 2021.

Stenseth, N. C., Payne, M. R., Bonsdorff, E., Dankel, D. J., Durant, J. M., Anderson, L. G., Armstrong, C. W., Blenckner, T., Brakstad, A., Dupont, S., Eikeset, A. M., Goksøyr, A., Jónsson, S., Kuparinen, A., Våge, K., Österblom, H., and Paasche, Ø.: Attuning to a changing ocean, *P. Natl. Acad. Sci. USA*, 117, 20363, <https://doi.org/10.1073/pnas.1915352117>, 2020.

Stephenson, D. B., Pavan, V., and Bojariu, R.: Is the North Atlantic Oscillation a random walk?, *Int. J. Climatol.*, 20, 1-18, [https://doi.org/10.1002/\(SICI\)1097-0088\(200001\)20:1<1::AID-JOC456>3.0.CO;2-P](https://doi.org/10.1002/(SICI)1097-0088(200001)20:1<1::AID-JOC456>3.0.CO;2-P), 2000.

Stjern, C. W., Kristjánsson, J. E., and Hansen, A. W.: Global dimming and global brightening—an analysis of surface radiation and cloud cover data in northern Europe, *Int. J. Climatol.*, 29, 643-653, <https://doi.org/10.1002/joc.1735>, 2009.

Stockholm University - Department of Physical Geography: Kebnekaise's south peak still highest in Sweden: <https://www.su.se/english/research/kebnেকেise-s-south-peak-still-highest-in-sweden-1.343483>, access: 14.12.2020, 2017.

Stockholm University - Department of Physical Geography: Southern peak of Kebnekaise, lowest height ever measured: <https://www.natgeo.su.se/english/research/resarch-news/southern-peak-of-kebnেকেise-lowest-height-ever-measured-1.452263>, access: 14.12.2020, 2019.

Stockholm University - Department of Physical Geography: Tarfala Research Station: <https://www.natgeo.su.se/english/tarfala-research-station>, access: 14.12.2020, 2020.

Stokowski, M., Winogradow, A., Szymczycha, B., Carstensen, J., and Kuliński, K.: The CO₂ system dynamics in the vicinity of the Vistula River mouth (the southern Baltic Sea): A baseline investigation, *Estuar. Coast. Shelf Sci.*, 258, 107444, <https://doi.org/10.1016/j.ecss.2021.107444>, 2021.

Stonevičius, E., Rimkus, E., Štaras, A., Kažys, J., and Valiūškevičius, G.: Climate change impact on the Nemunas River basin hydrology in the 21st century, *Boreal Environ. Res.*, 22, 49-65, 2017.

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Feldfunktion geändert

Formatiert: Englisch (USA)

Formatiert: Schriftart: +Textkörper (Calibri), Rechtschreibung und Grammatik nicht prüfen

6893 Storelvmo, T., Heede, U. K., Leirvik, T., Phillips, P. C. B., Arndt, P., and Wild, M.: Lethargic Response to
6894 Aerosol Emissions in Current Climate Models, *Geophys. Res. Lett.*, 45, 9814-9823,
6895 <https://doi.org/10.1029/2018GL078298>, 2018.

6896 Stramska, M., and Białogrodzka, J.: Spatial and temporal variability of sea surface temperature in the Baltic Sea
6897 based on 32-years (1982–2013) of satellite data, *Oceanologia*, 57, 223-235,
6898 <https://doi.org/10.1016/j.oceano.2015.04.004>, 2015.

6899 Strandberg, G., and Kjellström, E.: Climate Impacts from Afforestation and Deforestation in Europe, *Earth*
6900 *Interact.*, 23, 1-27, <https://doi.org/10.1175/ei-d-17-0033.1>, 2019.

6901 Strong, C., and Magnúsdóttir, G.: Dependence of NAO variability on coupling with sea ice, *Clim. Dyn.*, 36,
6902 1681-1689, <https://doi.org/10.1007/s00382-010-0752-z>, 2011.

6903 Suikkanen, S., Pulina, S., Engström-Öst, J., Lehtiniemi, M., Lehtinen, S., and Brutemark, A.: Climate Change
6904 and Eutrophication Induced Shifts in Northern Summer Plankton Communities, *PLoS ONE*, 8, e66475,
6905 <https://doi.org/10.1371/journal.pone.0066475>, 2013.

6906 Sun, L., Perlwitz, J., and Hoerling, M.: What caused the recent “Warm Arctic, Cold Continents” trend pattern in
6907 winter temperatures?, *Geophys. Res. Lett.*, 43, 5345-5352, <https://doi.org/10.1002/2016GL069024>, 2016.

6908 Sündermann, J., and Pohlmann, T.: A brief analysis of North Sea physics, *Oceanologia*, 53, 663-689,
6909 <https://doi.org/10.5697/oc.53-3.663>, 2011.

6910 Sundqvist, L., Harkonen, T., Svensson, C. J., and Harding, K. C.: Linking Climate Trends to Population
6911 Dynamics in the Baltic Ringed Seal: Impacts of Historical and Future Winter Temperatures, *Ambio* 41, 865-872,
6912 <https://doi.org/10.1007/s13280-012-0334-x>, 2012.

6913 Sutton, R. T., and Hodson, D. L. R.: Atlantic Ocean Forcing of North American and European Summer Climate,
6914 *Science*, 309, 115-118, <https://doi.org/10.1126/science.1109496>, 2005.

6915 Suursaar, Ü., and Sooäär, J.: Decadal variations in mean and extreme sea level values along the Estonian coast of
6916 the Baltic Sea, *Tellus A*, 59, 249-260, <https://doi.org/10.1111/j.1600-0870.2006.00220.x>, 2007.

6917 Suursaar, Ü., Tõnisson, H., Alari, V., Raudsepp, U., Rästas, H., and Anderson, A.: Projected Changes in Wave
6918 Conditions in the Baltic Sea by the end of 21st Century and the Corresponding Shoreline Changes, *J. Coastal*
6919 *Res.*, 1012-1016, <https://doi.org/10.2112/si75-203.1>, 2016.

6920 Sveegaard, S., Andreasen, H., Mouritsen, K. N., Jeppesen, J. P., Teilmann, J., and Kinze, C. C.: Correlation
6921 between the seasonal distribution of harbour porpoises and their prey in the Sound, Baltic Sea, *Mar. Biol.*, 159,
6922 1029-1037, <https://doi.org/10.1007/s00227-012-1883-z>, 2012.

6923 Svendsen, L. M., and Gustafsson, B.: Waterborne nitrogen and phosphorus inputs and water flow to the Baltic
6924 Sea 1995-2018, *HELCOM Baltic Sea Environmental Fact Sheet*, 25pp., 2020.

6925 Swedish Infrastructure for Ecosystem Science: Meteorological data from Tarfala Research Station:
6926 <https://data.fieldsites.se/portal/>, access: 14.12.2020, 2020.

6927 Swedish Infrastructure for Ecosystem Science: Glacier data - surface mass balance from Rabots Glaciär, 1982–
6928 2020, <https://meta.fieldsites.se/objects/KI06fLDs-YUzlvq2GmVKTNg>, 2021a.

6929 Swedish Infrastructure for Ecosystem Science: Glacier data - surface mass balance from Måraglaciären, 1986–
6930 2020, <https://meta.fieldsites.se/objects/EXzEgEOQbn9ofJJrTX2fMHH>, 2021b.

6931 Swedish Infrastructure for Ecosystem Science: Glacier data - surface mass balance from Riukojietna, 1986–
6932 2020, <https://meta.fieldsites.se/objects/BZvbLykTx4cnEVAEuMSFVQeK>, 2021c.

6933 SMHI - Swedish Meteorological and Hydrological Institute: Climate scenarios:
6934 <http://www.smhi.se/en/climate/future-climate/climate-scenarios/>, access: 14.12.2020, 2020.

6935 SMHI - Swedish Meteorological and Hydrological Institute: Klimatindikator – Globalstrålning:
6936 <https://www.smhi.se/klimat/klimatet-da-och-nu/klimatindikatorer/stralning-1.17841>, access: 12.02.2021, 2021.

6937 Sweet, W., Kopp, R. E., Weaver, C. P., Obeysekera, J. T. B., Horton, R. M., Thieler, E. R., and Zervas, C. E.:
6938 Global and regional sea level rise scenarios for the United States, <https://doi.org/10.7289/v5/tr-nos-coops-083>,
6939 2017.

6940 Szwed, M., Pińskwar, I., Kundzewicz, Z. W., Graczyk, D., and Mezghani, A.: Changes of snow cover in Poland,
6941 *Acta Geophys.*, 65, 65-76, <https://doi.org/10.1007/s11600-017-0007-z>, 2017.

6942 Szwed, M., Dobler, A., Mezghani, A., and Saloranta, T. M.: Change of maximum snow cover depth in Poland -
6943 Trends and projections, *Időjárás*, 123, 487-500, <https://doi.org/10.28974/idojaras.2019.4.5>, 2019.

6944 Takolander, A., Cabeza, M., and Leskinen, E.: Climate change can cause complex responses in Baltic Sea
6945 macroalgae: A systematic review, *J. Sea Res.*, 123, 16-29, <https://doi.org/10.1016/j.seares.2017.03.007>, 2017.

6946 Tamarin-Brodsky, T., and Kaspi, Y.: Enhanced poleward propagation of storms under climate change, *Nat.*
6947 *Geosci.*, 10, 908-913, <https://doi.org/10.1038/s41561-017-0001-8>, 2017.

6948 Tamarin, T., and Kaspi, Y.: The poleward shift of storm tracks under global warming: A Lagrangian perspective,
6949 *Geophys. Res. Lett.*, 44, 10,666-610,674, <https://doi.org/10.1002/2017GL073633>, 2017.

6950 Tang, J., Yurova, A. Y., Schurgers, G., Miller, P. A., Olin, S., Smith, B., Siewert, M. B., Olefeldt, D., Pilesjö, P.,
6951 and Poska, A.: Drivers of dissolved organic carbon export in a subarctic catchment: Importance of microbial
6952 decomposition, sorption-desorption, peatland and lateral flow, *Sci. Total Environ.*, 622-623, 260-274,
6953 <https://doi.org/10.1016/j.scitotenv.2017.11.252>, 2018.

6954 Tang, L., Ramacher, M. O. P., Moldanová, J., Matthias, V., Karl, M., Johansson, L., Jalkanen, J. P., Yaramenka,
6955 K., Aulinger, A., and Gustafsson, M.: The impact of ship emissions on air quality and human health in the
6956 Gothenburg area – Part 1: 2012 emissions, *Atmos. Chem. Phys.*, 20, 7509-7530, <https://doi.org/10.5194/acp-20-7509-2020>, 2020.

6958 Taniguchi, M., Dulai, H., Burnett, K. M., Santos, I. R., Sugimoto, R., Stieglitz, T., Kim, G., Moosdorf, N., and
6959 Burnett, W. C.: Submarine Groundwater Discharge: Updates on Its Measurement Techniques, Geophysical
6960 Drivers, Magnitudes, and Effects, *Front. Environ. Sci.*, 7, <https://doi.org/10.3389/fenvs.2019.00141>, 2019.

6961 Teichmann, C., Bülow, K., Otto, J., Pfeifer, S., Rechid, D., Sieck, K., and Jacob, D.: Avoiding Extremes:
6962 Benefits of Staying below +1.5 °C Compared to +2.0 °C and +3.0 °C Global Warming, *Atmosphere*, 9, 115,
6963 <https://doi.org/10.3390/atmos9040115>, 2018.

6964 Teuling, A. J.: A hot future for European droughts, *Nat. Clim. Change*, 8, 364-365,
6965 <https://doi.org/10.1038/s41558-018-0154-5>, 2018.

6966 Thompson, P. M.: Seasonal changes in the distribution and composition of common seal (*Phoca vitulina*) haul-
6967 out groups, *J. Zool.*, 217, 281-294, <https://doi.org/10.1111/j.1469-7998.1989.tb02488.x>, 1989.

6968 Thomson, A. M., Calvin, K. V., Smith, S. J., Kyle, G. P., Volke, A., Patel, P., Delgado-Arias, S., Bond-
6969 Lamberty, B., Wise, M. A., Clarke, L. E., and Edmonds, J. A.: RCP4.5: a pathway for stabilization of radiative
6970 forcing by 2100, *Clim. Change*, 109, 77, <https://doi.org/10.1007/s10584-011-0151-4>, 2011.

6971 Tikkanen, M., and Oksanen, J.: Late Weichselian and Holocene shore displacement history of the Baltic Sea in
6972 Finland, *Fennia - International Journal of Geography*, 180, 9,20, 2002.

6973 Tilinina, N., Gulev, S. K., Rudeva, I., and Koltermann, P.: Comparing Cyclone Life Cycle Characteristics and
6974 Their Interannual Variability in Different Reanalyses, *J. Climate*, 26, 6419-6438, <https://doi.org/10.1175/jcli-d-12-00777.1>, 2013.

6976 Timmermann, K., Norkko, J., Janas, U., Norkko, A., Gustafsson, B. G., and Bonsdorff, E.: Modelling
6977 macrofaunal biomass in relation to hypoxia and nutrient loading, *J. Mar. Syst.*, 105-108, 60-69,
6978 <https://doi.org/10.1016/j.jmarsys.2012.06.001>, 2012.

6979 Ting, M., Kushnir, Y., Seager, R., and Li, C.: Robust features of Atlantic multi-decadal variability and its
6980 climate impacts, *Geophys. Res. Lett.*, 38, <https://doi.org/10.1029/2011GL048712>, 2011.

6981 Tinz, B.: On the relation between annual maximum extent of ice cover in the Baltic Sea and sea level pressure as
6982 well as air temperature field, *Geophysica*, 32, 319-341, 1996.

6983 Tokarska, K. B., Stolpe, M. B., Sippel, S., Fischer, E. M., Smith, C. J., Lehner, F., and Knutti, R.: Past warming
6984 trend constrains future warming in CMIP6 models, *Sci. Adv.*, 6, eaaz9549,
6985 <https://doi.org/10.1126/sciadv.aaz9549>, 2020.

6986 Tomczyk, A. M., Bednorz, E., and Szyga-Pluta, K.: Changes in Air Temperature and Snow Cover in Winter in
6987 Poland, *Atmosphere*, 12, 68, <https://doi.org/10.3390/atmos12010068>, 2021.

6988 Torn, K., Peterson, A., and Herkül, K.: Predicting the Impact of Climate Change on the Distribution of the Key
6989 Habitat-Forming Species in the Ne Baltic Sea, *J. Coastal Res.*, 95, 177-181, 175, [https://doi.org/10.2112/S195-](https://doi.org/10.2112/S195-035.1)
6990 035.1, 2020.

6991 Törnroos, A., Pecuchet, L., Olsson, J., Gårdmark, A., Blomqvist, M., Lindegren, M., and Bonsdorff, E.: Four
6992 decades of functional community change reveals gradual trends and low interlinkage across trophic groups in a
6993 large marine ecosystem, *Global Change Biol.*, 25, 1235-1246, <https://doi.org/10.1111/gcb.14552>, 2019.

6994 Tuomi, L., Kahma, K. K., and Pettersson, H.: Wave hindcast statistics in the seasonally ice-covered Baltic Sea,
6995 *Boreal Environ. Res.*, 16, 451-472, 2011.

6996 Turner, D. R., Hassellöv, I.-M., Ytreberg, E., and Rutgersson, A.: Shipping and the environment: Smokestack
6997 emissions, scrubbers and unregulated oceanic consequences, *Elementa-Sci. Anthropol.*, 5,
6998 <https://doi.org/10.1525/elementa.167>, 2017.

6999 Turner, D. R., Edman, M., Gallego-Urrea, J. A., Claremar, B., Hassellöv, I.-M., Omstedt, A., and Rutgersson,
7000 A.: The potential future contribution of shipping to acidification of the Baltic Sea, *Ambio*, 47, 368-378,
7001 <https://doi.org/10.1007/s13280-017-0950-6>, 2018.

7002 Ukkonen, P., Aaris-Sørensen, K., Arppe, L., Daugnora, L., Halkka, A., Lõugas, L., Oinonen, M. J., Pilot, M.,
7003 and Storå, J.: An Arctic seal in temperate waters: History of the ringed seal (*Pusa hispida*) in the Baltic Sea and
7004 its adaptation to the changing environment, *Holocene*, 24, 1694-1706,
7005 <https://doi.org/10.1177/0959683614551226>, 2014.

7006 Ulbrich, U., Pinto, J. G., Kupfer, H., Leckebusch, G. C., Spangehl, T., and Reyers, M.: Changing Northern
7007 Hemisphere Storm Tracks in an Ensemble of IPCC Climate Change Simulations, *J. Climate*, 21, 1669-1679,
7008 <https://doi.org/10.1175/2007jcli1992.1>, 2008.

7009 Ulfssbo, A., Kuliński, K., Anderson, L. G., and Turner, D. R.: Modelling organic alkalinity in the Baltic Sea
7010 using a Humic-Pitzer approach, *Mar. Chem.*, 168, 18-26, <https://doi.org/10.1016/j.marchem.2014.10.013>, 2015.

7011 Umlauf, L., Holtermann, P. L., Gillner, C. A., Prien, R. D., Merckelbach, L., and Carpenter, J. R.: Diffusive
7012 Convection under Rapidly Varying Conditions, *J. Phys. Oceanogr.*, 48, 1731-1747, [https://doi.org/10.1175/jpo-](https://doi.org/10.1175/jpo-d-18-0018.1)
7013 d-18-0018.1, 2018.

7014 Undeman, E., Gustafsson, B. G., Humborg, C., and McLachlan, M. S.: Application of a novel modeling tool
7015 with multistressor functionality to support management of organic contaminants in the Baltic Sea, *Ambio* 44,
7016 498-506, <https://doi.org/10.1007/s13280-015-0668-2>, 2015.

7017 United Nations Climate Change: Paris Agreement, 27pp, <https://unfccc.int/documents/37107>, 2015.

7018 Uotila, P., Vihma, T., and Haapala, J.: Atmospheric and oceanic conditions and the extremely low Bothnian Bay
7019 sea ice extent in 2014/2015, *Geophys. Res. Lett.*, 42, 7740-7749, <https://doi.org/10.1002/2015GL064901>, 2015.

7020 Urraca, R., Gracia-Amillo, A. M., Koubli, E., Huld, T., Trentmann, J., Riihelä, A., Lindfors, A. V., Palmer, D.,
7021 Gottschalg, R., and Antonanzas-Torres, F.: Extensive validation of CM SAF surface radiation products over
7022 Europe, *Remote Sens. Environ.*, 199, 171-186, <https://doi.org/10.1016/j.rse.2017.07.013>, 2017.

7023 Vähätalo, A. V., Rainio, K., Lehtikoinen, A., and Lehtikoinen, E.: Spring arrival of birds depends on the North
7024 Atlantic Oscillation, *J. Avian Biol.*, 35, 210-216, <https://doi.org/10.1111/j.0908-8857.2004.03199.x>, 2004.

7025 Vahtera, E., Conley, D. J., Gustafsson, B. G., Kuosa, H., Pitkänen, H., Savchuk, O. P., Tamminen, T., Viitasalo,
7026 M., Voss, M., Wasmund, N., and others: Internal ecosystem feedbacks enhance nitrogen-fixing cyanobacteria
7027 blooms and complicate management in the Baltic Sea, *Ambio*, 36, 186-194, [https://doi.org/10.1579/0044-](https://doi.org/10.1579/0044-7447(2007)36[186:IEFENC]2.0.CO;2)
7028 7447(2007)36[186:IEFENC]2.0.CO;2, 2007.

7029 Väli, G., Meier, H. E. M., and Elken, J.: Simulated halocline variability in the Baltic Sea and its impact on
7030 hypoxia during 1961-2007, *J. Geophys. Res.-Oceans*, 118, 6982-7000, <https://doi.org/10.1002/2013JC009192>,
7031 2013.

7032 van der Jeugd, H. P., Eichhorn, G., Litvin, K. E., Stahl, J., Larsson, K., van der Graaf, A. J., and Drent, R. H.:
7033 Keeping up with early springs: rapid range expansion in an avian herbivore incurs a mismatch between
7034 reproductive timing and food supply, *Global Change Biol.*, 15, 1057-1071, [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2486.2008.01804.x)
7035 2486.2008.01804.x, 2009.

7036 van Helmond, N. A. G. M., Jilbert, T., and Slomp, C. P.: Hypoxia in the Holocene Baltic Sea: Comparing
7037 modern versus past intervals using sedimentary trace metals, *Chem. Geol.*, 493, 478-490,
7038 <https://doi.org/10.1016/j.chemgeo.2018.06.028>, 2018.

7039 van Vuuren, D. P., Stehfest, E., den Elzen, M. G. J., Kram, T., van Vliet, J., Deetman, S., Isaac, M., Klein
7040 Goldewijk, K., Hof, A., Mendoza Beltran, A., Oostenrijk, R., and van Ruijven, B.: RCP2.6: exploring the
7041 possibility to keep global mean temperature increase below 2°C, *Clim. Change*, 109, 95,
7042 <https://doi.org/10.1007/s10584-011-0152-3>, 2011.

7043 van Vuuren, D. P., Kriegler, E., O'Neill, B. C., Ebi, K. L., Riahi, K., Carter, T. R., ... & Winkler, H.: A new scenario
7044 framework for climate change research: scenario matrix architecture, *Climatic Change*, 122(3), 373-386, 2014.

7045 Vanninen, P., Östin, A., Beldowski, J., Pedersen, E. A., Söderström, M., Szubska, M., Grabowski, M.,
7046 Siedlewicz, G., Czub, M., Popiel, S., Nawala, J., Dziedzic, D., Jakacki, J., and Pączek, B.: Exposure status of
7047 sea-dumped chemical warfare agents in the Baltic Sea, *Mar. Environ. Res.*, 161, 105112,
7048 <https://doi.org/10.1016/j.marenvres.2020.105112>, 2020.

7049 Varotsos, K. V., Giannakopoulos, C., and Tombrou, M.: Assessment of the Impacts of Climate Change on
7050 European Ozone Levels, Water, Air & Soil Pollution, 224, 1596, <https://doi.org/10.1007/s11270-013-1596-z>,
7051 2013.

7052 Vaughan, D. G., Comiso, J. C., Allison, I., Carrasco, J., Kaser, G., Kwok, R., Mote, P., Murray, T., Paul, F., Ren,
7053 J., Rignot, E., Solomina, O., Steffen, K., and Zhang, T.: Observations: Cryosphere, in: *Climate Change 2013 –*
7054 *The Physical Science Basis: Working Group I Contribution to the Fifth Assessment Report of the*
7055 *Intergovernmental Panel on Climate Change*, edited by: Intergovernmental Panel on Climate, C., Cambridge
7056 University Press, Cambridge, 317-382, <https://doi.org/10.1017/CBO9781107415324.012>, 2014.

7057 Vehmaa, A., Hogfors, H., Gorokhova, E., Brutemark, A., Holmborn, T., and Engström-Öst, J.: Projected marine
7058 climate change: effects on copepod oxidative status and reproduction, *Ecol. Evol.*, 3, 4548-4557,
7059 <https://doi.org/10.1002/ece3.839>, 2013.

7060 Vehmaa, A., Almén, A. K., Brutemark, A., Paul, A., Riebesell, U., Furuhausen, S., and Engström-Öst, J.: Ocean
7061 acidification challenges copepod phenotypic plasticity, *Biogeosciences*, 13, 6171-6182,
7062 <https://doi.org/10.5194/bg-13-6171-2016>, 2016.

7063 Veijalainen, N., Lotsari, E., Vehviläinen, B., Alho, P., and Käyhkö, J.: Changes in floods in Finland due to
7064 climate change: General assessment on national scale, 6th Alexander von Humboldt International Conference on
7065 Climate Change, 2010.

7066 Vihma, T., and Haapala, J.: Geophysics of sea ice in the Baltic Sea: A review, *Prog. Oceanogr.*, 80, 129-148,
7067 <https://doi.org/10.1016/j.pocean.2009.02.002>, 2009.

7068 Vihma, T.: Weather Extremes Linked to Interaction of the Arctic and Midlatitudes, in: *Climate Extremes*, edited
7069 by: Wang, S. Y. S., Yoon, J. H., Funk, C. C., and Gillies, R. R., 39-50,
7070 <https://doi.org/10.1002/9781119068020.ch2>, 2017.

7071 Viitasalo, M., and Bonsdorff, B.: Climate change and the Baltic Sea ecosystem: direct and indirect effects on
7072 species, communities and ecosystem function *Earth Syst. Dynam.*, 2021.

7073 Viru, B., and Jaagus, J.: Spatio-temporal variability and seasonal dynamics of snow cover regime in Estonia,
7074 *Theor. Appl. Climatol.*, 139, 759-771, <https://doi.org/10.1007/s00704-019-03013-5>, 2020.

7075 Viška, M., and Soomere, T.: Simulated and observed reversals of wave-driven alongshore sediment transport at
7076 the eastern Baltic Sea coast, *Baltica*, 26, 145-156, 2013.

Formatiert: Schwedisch (Schweden)

Formatiert: Schriftart: Nicht Kursiv

Formatiert: Schriftart: Nicht Kursiv

Formatiert: Schriftart: +Textkörper (Calibri)

7077 Vogt, J., Soille, P., Jager, A. d., Rimaviciute, E., Mehl, W., Foisneau, S., Bódis, K., Dusart, J., Paracchini, M. L.,
7078 Haastrop, P., and Bamps, C.: A pan-European River and Catchment Database, European Commission - Joint
7079 Research Centre - Institute for Environment and Sustainability, Ispra, Italy, 124pp, 2007.

7080 Vogt, J., Rimaviciute, E., and Jager, A. d.: CCM2 River and Catchment Database for Europe Version 2.1
7081 Release Notes, European Commission - Joint Research Centre - Institute for Environment and Sustainability,
7082 Ispra, Italy, 5pp, 2008.

7083 Voipio, A.: The Baltic Sea, Elsevier Oceanographic Series, Amsterdam, 1981.

7084 Volkov, V. A., Johannessen, O. M., Borodachev, V. E., Voinov, G. N., Pettersson, L. H., Bobylev, L. P., and
7085 Kouraev, A. V.: Polar seas oceanography: an integrated case study of the Kara Sea, Springer Science & Business
7086 Media, Berlin, Heidelberg, 2002.

7087 Vonk, J. E., Tank, S. E., Bowden, W. B., Laurion, I., Vincent, W. F., Alekseychik, P., Amyot, M., Billet, M.,
7088 Canário, J., and Cory, R. M.: Reviews and syntheses: Effects of permafrost thaw on Arctic aquatic ecosystems,
7089 Biogeosciences, 12, 7129-7167, <https://doi.org/10.5194/bg-12-7129-2015>, 2015.

7090 Voss, R., Hinrichsen, H.-H., Quaas, M. F., Schmidt, J. O., and Tahvonen, O.: Temperature change and Baltic
7091 sprat: from observations to ecological-economic modelling, ICES J. Mar. Sci., 68, 1244-1256,
7092 <https://doi.org/10.1093/icesjms/fsr063>, 2011.

7093 Vousdoukas, M. I., Voukouvalas, E., Annunziato, A., Giardino, A., and Feyen, L.: Projections of extreme storm
7094 surge levels along Europe, Clim. Dyn., 47, 3171-3190, <https://doi.org/10.1007/s00382-016-3019-5>, 2016.

7095 Vousdoukas, M. I., Mentaschi, L., Voukouvalas, E., Verlaan, M., and Feyen, L.: Extreme sea levels on the rise
7096 along Europe's coasts, Earths Future, 5, 304-323, <https://doi.org/10.1002/2016EF000505>, 2017.

7097 Vuorinen, I., Hänninen, J., Rajasilta, M., Laine, P., Eklund, J., Montesino-Pouzols, F., Corona, F., Junker, K.,
7098 Meier, H. E. M., and Dippner, J. W.: Scenario simulations of future salinity and ecological consequences in the
7099 Baltic Sea and adjacent North Sea areas—implications for environmental monitoring, Ecol. Indic., 50, 196-205,
7100 <https://doi.org/10.1016/j.ecolind.2014.10.019>, 2015.

7101 Wahl, M., Werner, F. J., Buchholz, B., Raddatz, S., Graiff, A., Matthiessen, B., Karsten, U., Hiebenthal, C.,
7102 Hamer, J., Ito, M., Gülzow, E., Rilov, G., and Guy-Haim, T.: Season affects strength and direction of the
7103 interactive impacts of ocean warming and biotic stress in a coastal seaweed ecosystem, Limnol. Oceanogr., 65,
7104 807-827, <https://doi.org/10.1002/lno.11350>, 2020.

7105 Wahl, T., and Chambers, D. P.: Climate controls multidecadal variability in U. S. extreme sea level records, J.
7106 Geophys. Res-Oceans, 121, 1274-1290, <https://doi.org/10.1002/2015JC011057>, 2016.

7107 Wählström, I., Höglund, A., Almroth-Rosell, E., MacKenzie, B. R., Gröger, M., Eilola, K., Plikshs, M., and
7108 Andersson, H. C.: Combined climate change and nutrient load impacts on future habitats and eutrophication
7109 indicators in a eutrophic coastal sea, Limnol. Oceanogr., 65, 2170-2187, <https://doi.org/10.1002/lno.11446>,
7110 2020.

7111 Waldeck, P., and Larsson, K.: Effects of winter water temperature on mass loss in Baltic blue mussels:
7112 Implications for foraging sea ducks, J. Exp. Mar. Biol. Ecol., 444, 24-30,
7113 <https://doi.org/10.1016/j.jembe.2013.03.007>, 2013.

7114 Wang, J., Yang, B., Ljungqvist, F. C., Luterbacher, J., Osborn, Timothy J., Briffa, K. R., and Zorita, E.: Internal
7115 and external forcing of multidecadal Atlantic climate variability over the past 1,200 years, Nat. Geosci., 10, 512-
7116 517, <https://doi.org/10.1038/ngeo2962>, 2017.

7117 Wang, Q., Fan, X., and Wang, M.: Evidence of high-elevation amplification versus Arctic amplification, Sci.
7118 Rep.-UK, 6, 19219, <https://doi.org/10.1038/srep19219>, 2016.

7119 Wang, S., Dieterich, C., Döscher, R., Höglund, A., Hordoir, R., Meier, H. E. M., Samuelsson, P., and
7120 Schimanke, S.: Development and evaluation of a new regional coupled atmosphere-ocean model in the North
7121 Sea and Baltic Sea, Tellus A, 67, 24284, <https://doi.org/10.3402/tellusa.v67.24284>, 2015.

7122 Warwick, R. M., Tweedley, J. R., and Potter, I. C.: Microtidal estuaries warrant special management measures
7123 that recognise their critical vulnerability to pollution and climate change, *Mar. Pollut. Bull.*, 135, 41–46,
7124 <https://doi.org/10.1016/j.marpolbul.2018.06.062>, 2018.

7125 Wasmund, N., Tuimala, J., Suikkanen, S., Vandepitte, L., and Kraberg, A.: Long-term trends in phytoplankton
7126 composition in the western and central Baltic Sea, *J. Mar. Syst.*, 87, 145–159,
7127 <https://doi.org/10.1016/j.jmarsys.2011.03.010>, 2011.

7128 Wasmund, N.: The Diatom/Dinoflagellate Index as an Indicator of Ecosystem Changes in the Baltic Sea. 2.
7129 Historical Data for Use in Determination of Good Environmental Status, *Front. Mar. Sci.*, 4,
7130 <https://doi.org/10.3389/fmars.2017.00153>, 2017.

7131 Wasmund, N., Kownacka, J., Göbel, J., Jaanus, A., Johansen, M., Jurgensone, I., Lehtinen, S., and Powilleit, M.:
7132 The Diatom/Dinoflagellate Index as an Indicator of Ecosystem Changes in the Baltic Sea 1. Principle and
7133 Handling Instruction, *Front. Mar. Sci.*, 4, <https://doi.org/10.3389/fmars.2017.00022>, 2017.

7134 Wasmund, N., Nausch, G., Gerth, M., Busch, S., Burmeister, C., Hansen, R., and Sadkowiak, B.: Extension of
7135 the growing season of phytoplankton in the western Baltic Sea in response to climate change, *Mar. Ecol. Prog.
7136 Ser.*, 622, 1–16, <https://doi.org/10.3354/meps12994>, 2019.

7137 Watanabe, M., and Tatebe, H.: Reconciling roles of sulphate aerosol forcing and internal variability in Atlantic
7138 multidecadal climate changes, *Clim. Dyn.*, 53, 4651–4665, <https://doi.org/10.1007/s00382-019-04811-3>, 2019.

7139 Watson, L., Lacressonnière, G., Gauss, M., Engardt, M., Andersson, C., Josse, B., Maréchal, V., Nyiri, A.,
7140 Sobolowski, S., Siour, G., Szopa, S., and Vautard, R.: Impact of emissions and +2 °C climate change upon future
7141 ozone and nitrogen dioxide over Europe, *Atmos. Environ.*, 142, 271–285,
7142 <https://doi.org/10.1016/j.atmosenv.2016.07.051>, 2016.

7143 Watts, P.: The diel hauling-out cycle of harbour seals in an open marine environment: correlates and constraints,
7144 *J. Zool.*, 240, 175–200, <https://doi.org/10.1111/j.1469-7998.1996.tb05494.x>, 1996.

7145 Weigel, B., Andersson, H. C., Meier, H. E. M., Blenckner, T., Snickars, M., and Bonsdorff, E.: Long-term
7146 progression and drivers of coastal zoobenthos in a changing system, *Mar. Ecol. Prog. Ser.*, 528, 141–159,
7147 <https://doi.org/10.3354/meps11279>, 2015.

7148 Weisse, R., and Weidemann, H.: Baltic Sea extreme sea levels 1948–2011: Contributions from atmospheric
7149 forcing, *Proc. IUTAM*, 25, 65–69, <https://doi.org/10.1016/j.piutam.2017.09.010>, 2017.

7150 Weisse, R., Dailidiene, I., Hünicke, B., Kahma, K., Madsen, K., Omstedt, A., Parnell, K., Schöne, T., Soomere,
7151 T., Zhang, W., and Zorita, E.: Sea Level Dynamics and Coastal Erosion in the Baltic Sea Region, *Earth Syst.
7152 Dynam. Discuss.*, 1–40, <https://doi.org/10.5194/esd-2021-6>, 2021.

7153 Whan, K., Zscheischler, J., Orth, R., Shongwe, M., Rahimi, M., Asare, E. O., and Seneviratne, S. I.: Impact of
7154 soil moisture on extreme maximum temperatures in Europe, *Weather. Clim. Extremes*, 9, 57–67,
7155 <https://doi.org/10.1016/j.wace.2015.05.001>, 2015.

7156 Wikner, J., and Andersson, A.: Increased freshwater discharge shifts the trophic balance in the coastal zone of
7157 the northern Baltic Sea, *Global Change Biol.*, 18, 2509–2519, <https://doi.org/10.1111/j.1365-2486.2012.02718.x>,
7158 2012.

7159 Wild, M., Gilgen, H., Roesch, A., Ohmura, A., Long, C. N., Dutton, E. G., Forgan, B., Kallis, A., Russak, V.,
7160 and Tsvetkov, A.: From Dimming to Brightening: Decadal Changes in Solar Radiation at Earth's Surface,
7161 *Science*, 308, 847, <https://doi.org/10.1126/science.1103215>, 2005.

7162 Wild, M.: Enlightening Global Dimming and Brightening, *Bull. Amer. Meteor. Soc.*, 93, 27–37,
7163 <https://doi.org/10.1175/bams-d-11-00074.1>, 2012.

7164 Wild, M.: Decadal changes in radiative fluxes at land and ocean surfaces and their relevance for global warming,
7165 *Wires. Clim. Change*, 7, 91–107, <https://doi.org/10.1002/wcc.372>, 2016.

7166 Wild, M., Ohmura, A., Schär, C., Müller, G., Folini, D., Schwarz, M., Hakuba, M. Z., and Sanchez-Lorenzo, A.:
7167 The Global Energy Balance Archive (GEBA) version 2017: a database for worldwide measured surface energy
7168 fluxes, *Earth Syst. Sci. Data*, 9, 601–613, <https://doi.org/10.5194/essd-9-601-2017>, 2017.

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Feldfunktion geändert

Formatiert: Englisch (USA)

Formatiert: Rechtschreibung und Grammatik nicht prüfen

7169 Wild, M., Wacker, S., Yang, S., and Sanchez-Lorenzo, A.: Evidence for Clear-sky Dimming and Brightening in
7170 Central Europe, *Geophys. Res. Lett.*, n/a, e2020GL092216, <https://doi.org/10.1029/2020GL092216>, 2021.

7171 Williams, C. N., Carrivick, J. L., Evans, A. J., and Rippin, D. M.: Quantifying uncertainty in using multiple
7172 datasets to determine spatiotemporal ice mass loss over 101 years at kårsagläciären, sub-arctic sweden, *Geogr.*
7173 *Ann. A*, 98, 61-79, <https://doi.org/10.1111/geoa.12123>, 2016.

7174 Willison, J., Robinson, W. A., and Lackmann, G. M.: North Atlantic Storm-Track Sensitivity to Warming
7175 Increases with Model Resolution, *J. Climate*, 28, 4513-4524, <https://doi.org/10.1175/jcli-d-14-00715.1>, 2015.

7176 Wills, R. C. J., Armour, K. C., Battisti, D. S., and Hartmann, D. L.: Ocean–Atmosphere Dynamical Coupling
7177 Fundamental to the Atlantic Multidecadal Oscillation, *J. Climate*, 32, 251-272, [https://doi.org/10.1175/jcli-d-18-](https://doi.org/10.1175/jcli-d-18-0269.1)
7178 [0269.1](https://doi.org/10.1175/jcli-d-18-0269.1), 2018.

7179 Wills, R. C. J., White, R. H., and Levine, X. J.: Northern Hemisphere Stationary Waves in a Changing Climate,
7180 *Curr. Clim. Chang. Rep.*, 5, 372-389, <https://doi.org/10.1007/s40641-019-00147-6>, 2019.

7181 Wilson, D., Hisdal, H., and Lawrence, D.: Has streamflow changed in the Nordic countries? – Recent trends and
7182 comparisons to hydrological projections, *J. Hydrol.*, 394, 334-346, <https://doi.org/10.1016/j.jhydrol.2010.09.010>,
7183 2010.

7184 Wisniewska, Danuta M., Johnson, M., Teilmann, J., Rojano-Doñate, L., Shearer, J., Sveegaard, S., Miller,
7185 Lee A., Siebert, U., and Madsen, Peter T.: Ultra-High Foraging Rates of Harbor Porpoises Make Them
7186 Vulnerable to Anthropogenic Disturbance, *Curr. Biol.*, 26, 1441-1446,
7187 <https://doi.org/10.1016/j.cub.2016.03.069>, 2016.

7188 Wohland, J., Brayshaw, D., Bloomfield, H., and Wild, M.: European multidecadal solar variability badly
7189 captured in all centennial reanalyses except CERA20C, *Environ. Res. Lett.*, 15, 104021,
7190 <https://doi.org/10.1088/1748-9326/aba7e6>, 2020.

7191 Wolski, T., Wiśniewski, B., Giza, A., Kowalewska-Kalkowska, H., Boman, H., Grabbi-Kaiv, S., Hammarklint,
7192 T., Holfort, J., and Lydeikaitė, Ž.: Extreme sea levels at selected stations on the Baltic Sea coast**This work was
7193 financed by the Polish National Centre for Science research project No. 2011/01/B/ST10/06470, *Oceanologia*,
7194 56, 259-290, <https://doi.org/10.5697/oc.56-2.259>, 2014.

7195 Woodruff, S. D., Worley, S. J., Lubker, S. J., Ji, Z., Eric Freeman, J., Berry, D. I., Brohan, P., Kent, E. C.,
7196 Reynolds, R. W., Smith, S. R., and Wilkinson, C.: ICOADS Release 2.5: extensions and enhancements to the
7197 surface marine meteorological archive, *Int. J. Climatol.*, 31, 951-967, <https://doi.org/10.1002/joc.2103>, 2011.

7198 Woollings, T., Barriopedro, D., Methven, J., Son, S.-W., Martius, O., Harvey, B., Sillmann, J., Lupo, A. R., and
7199 Seneviratne, S.: Blocking and its Response to Climate Change, *Curr. Clim. Chang. Rep.*, 4, 287-300,
7200 <https://doi.org/10.1007/s40641-018-0108-z>, 2018.

7201 World Glacier Monitoring Service: Fluctuations of Glaciers (FoG) Database:
7202 https://wgms.ch/data_databaseversions/, access: ~~14.12.2020~~22.11.2021, 2021.

7203 World Meteorological Organization: State of the Global Climate 2020, 38pp, 2020.

7204 Wrzesiński, D., Chosiński, A., Ptak, M., and Skowron, R.: Effect of the North Atlantic Oscillation on the Pattern
7205 of Lake Ice Phenology in Poland, *Acta Geophys.*, 63, 1664-1684, <https://doi.org/10.1515/acgeo-2015-0055>,
7206 2015.

7207 Wu, S., Liu, Z.-Y., Cheng, J., and Li, C.: Response of North Pacific and North Atlantic decadal variability to
7208 weak global warming, *Adv. Clim. Chang. Res.*, 9, 95-101, <https://doi.org/10.1016/j.accre.2018.03.001>, 2018.

7209 Wu, S., and Liu, Z.-Y.: Decadal Variability in the North Pacific and North Atlantic under Global Warming: The
7210 Weakening Response and Its Mechanism, *J. Climate*, 33, 9181-9193, <https://doi.org/10.1175/jcli-d-19-1012.1>,
7211 2020.

7212 Wübbert, C., and Krauss, W.: The two dimensional seiches of the Baltic Sea, *Oceanol. Acta*, 2, 435-446, 1979.

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Feldfunktion geändert

Formatiert: Englisch (USA)

Formatiert: Rechtschreibung und Grammatik nicht prüfen

7213 Wulff, A., Karlberg, M., Olofsson, M., Torstensson, A., Riemann, L., Steinhoff, F. S., Mohlin, M., Ekstrand, N.,
7214 and Chierici, M.: Ocean acidification and desalination: climate-driven change in a Baltic Sea summer
7215 microplanktonic community, *Mar. Biol.*, 165, 63, <https://doi.org/10.1007/s00227-018-3321-3>, 2018.

7216 Wyser, K., van Noije, T., Yang, S., von Hardenberg, J., O'Donnell, D., and Döscher, R.: On the increased
7217 climate sensitivity in the EC-Earth model from CMIP5 to CMIP6, *Geosci. Model Dev.*, 13, 3465-3474,
7218 <https://doi.org/10.5194/gmd-13-3465-2020>, 2020.

7219 Ye, K., and Lau, N.-C.: Influences of surface air temperature and atmospheric circulation on winter snow cover
7220 variability over Europe, *Int. J. Climatol.*, 37, 2606-2619, <https://doi.org/10.1002/joc.4868>, 2017.

7221 Yin, J. H.: A consistent poleward shift of the storm tracks in simulations of 21st century climate, *Geophys. Res.*
7222 *Lett.*, 32, <https://doi.org/10.1029/2005GL023684>, 2005.

7223 Yli-Pelkonen, V.: Ecological information in the political decision making of urban land-use planning, *J. Environ.*
7224 *Plan. Manag.*, 51, 345-362, <https://doi.org/10.1080/09640560801977224>, 2008.

7225 Zandersen, M., Hyytiäinen, K., Meier, H. E. M., Tomczak, M. T., Bauer, B., Haapasaari, P. E., Olesen, J. E.,
7226 Gustafsson, B. G., Refsgaard, J. C., Fridell, E., Pihlainen, S., Le Tissier, M. D. A., Kosenius, A.-K., and Van
7227 Vuuren, D. P.: Shared socio-economic pathways extended for the Baltic Sea: exploring long-term environmental
7228 problems, *Reg. Environ. Change*, 19, 1073-1086, <https://doi.org/10.1007/s10113-018-1453-0>, 2019.

7229 Zappa, G., Shaffrey, L. C., Hodges, K. I., Sansom, P. G., and Stephenson, D. B.: A Multimodel Assessment of
7230 Future Projections of North Atlantic and European Extratropical Cyclones in the CMIP5 Climate Models, *J.*
7231 *Climate*, 26, 5846-5862, <https://doi.org/10.1175/jcli-d-12-00573.1>, 2013.

7232 Zappa, G., and Shepherd, T. G.: Storylines of Atmospheric Circulation Change for European Regional Climate
7233 Impact Assessment, *J. Climate*, 30, 6561-6577, <https://doi.org/10.1175/jcli-d-16-0807.1>, 2017.

7234 Zappa, G., Pithan, F., and Shepherd, T. G.: Multimodel Evidence for an Atmospheric Circulation Response to
7235 Arctic Sea Ice Loss in the CMIP5 Future Projections, *Geophys. Res. Lett.*, 45, 1011-1019,
7236 <https://doi.org/10.1002/2017GL076096>, 2018.

7237 Zemp, M., Huss, M., Thibert, E., Eckert, N., McNabb, R., Huber, J., Barandun, M., Machguth, H., Nussbaumer,
7238 S. U., Gärtner-Roer, I., Thomson, L., Paul, F., Maussion, F., Kutuzov, S., and Cogley, J. G.: Global glacier mass
7239 changes and their contributions to sea-level rise from 1961 to 2016, *Nature*, 568, 382-386,
7240 <https://doi.org/10.1038/s41586-019-1071-0>, 2019.

7241 Zemp, M., Gärtner-Roer, I., Nussbaumer, S. U., Bannwart, J., Rastner, P., Paul, F., Hoelzle, M., and [eds.]:
7242 Global Glacier Change Bulletin No. 3 (2016–2017) World Glacier Monitoring Service, Zürich,
7243 <https://doi.org/10.5167/uzh-191797>, 2020.

7244 Zhang, M., Lee, X., Yu, G., Han, S., Wang, H., Yan, J., Zhang, Y., Li, Y., Ohta, T., Hirano, T., Kim, J.,
7245 Yoshifuji, N., and Wang, W.: Response of surface air temperature to small-scale land clearing across latitudes,
7246 *Environ. Res. Lett.*, 9, 034002, <https://doi.org/10.1088/1748-9326/9/3/034002>, 2014a.

7247 Zhang, W., Harff, J., and Schneider, R.: Analysis of 50-year wind data of the southern Baltic Sea for modelling
7248 coastal morphological evolution – a case study from the Darss-Zingst Peninsula, *Oceanologia*, 53, 489-518,
7249 <https://doi.org/10.5697/oc.53-1-TI.489>, 2011.

7250 Zhang, W., Miller, P. A., Smith, B., Wania, R., Koenigk, T., and Döscher, R.: Tundra shrubification and tree -
7251 line advance amplify arctic climate warming: results from an individual-based dynamic vegetation model,
7252 *Environ. Res. Lett.*, 8, 034023, <https://doi.org/10.1088/1748-9326/8/3/034023>, 2013.

7253 Zhang, W., Jansson, C., Miller, P. A., Smith, B., and Samuelsson, P.: Biogeophysical feedbacks enhance the
7254 Arctic terrestrial carbon sink in regional Earth system dynamics, *Biogeosciences*, 11, 5503-5519,
7255 <https://doi.org/10.5194/bg-11-5503-2014>, 2014b.

7256 Zhang, W., Schneider, R., Kolb, J., Teichmann, T., Dudzinska-Nowak, J., Harff, J., and Hanebuth, T. J. J.:
7257 Land–sea interaction and morphogenesis of coastal foredunes — A modeling case study from the southern Baltic
7258 Sea coast, *Coast. Eng.*, 99, 148-166, <https://doi.org/10.1016/j.coastaleng.2015.03.005>, 2015.

- 7259 Zhang, W., Schneider, R., Harff, J., Hünicke, B., and Fröhle, P.: Modelling of Medium-Term (Decadal) Coastal
7260 Foredune Morphodynamics- Historical Hindcast and Future Scenarios of the Świna Gate Barrier Coast
7261 (Southern Baltic Sea), in: *Coastline Changes of the Baltic Sea from South to East: Past and Future Projection*,
7262 edited by: Harff, J., Furmańczyk, K., and von Storch, H., Springer International Publishing, Cham, 107-135,
7263 https://doi.org/10.1007/978-3-319-49894-2_7, 2017.
- 7264 Zhang, W., Miller, P. A., Jansson, C., Samuelsson, P., Mao, J., and Smith, B.: Self-Amplifying Feedbacks
7265 Accelerate Greening and Warming of the Arctic, *Geophys. Res. Lett.*, 45, 7102-7111,
7266 <https://doi.org/10.1029/2018GL077830>, 2018.
- 7267 Zhong, X., Zhang, T., and Wang, K.: Snow density climatology across the former USSR, *Cryosphere*, 8, 785-
7268 799, <https://doi.org/10.5194/tc-8-785-2014>, 2014.
- 7269 Zhu, X., Lee, S.-Y., Wen, X., Wei, Z., Ji, Z., Zheng, Z., and Dong, W.: Historical evolution and future trend of
7270 Northern Hemisphere snow cover in CMIP5 and CMIP6 models, *Environ. Res. Lett.*, 16, 065013,
7271 <https://doi.org/10.1088/1748-9326/ac0662>, 2021.
- 7272 Zillén, L., and Conley, D. J.: Hypoxia and cyanobacteria blooms-are they really natural features of the late
7273 Holocene history of the Baltic Sea?, *Biogeosciences*, 7, 2567, <https://doi.org/10.5194/bg-7-2567-2010>, 2010.
- 7274 Zorita, E., and Laine, A.: Dependence of salinity and oxygen concentrations in the Baltic Sea on large -scale
7275 atmospheric circulation, *Clim. Res.*, 14, 25-41, <https://doi.org/10.3354/cr014025>, 2000.