



1 Climate Change in the Baltic Sea Region: A Summary

2 H. E. Markus Meier^{1,2}, Madline Kniebusch¹, Christian Dieterich^{2,†}, Matthias Gröger¹, Eduardo
 3 Zorita³, Ragnar Elmgren⁴, Kai Myrberg^{5,6}, Markus Ahola⁷, Alena Bartosova², Erik Bonsdorff⁸,
 4 Florian Börgel¹, Rene Capell², Ida Carlén⁹, Thomas Carlund¹⁰, Jacob Carstensen¹¹, Ole B.
 5 Christensen¹², Volker Dierschke¹³, Claudia Frauen^{1,14}, Morten Frederiksen¹¹, Elie Gaget^{15,16},
 6 Anders Galatius¹¹, Jari J. Haapala¹⁷, Antti Halkka¹⁸, Gustaf Hugelius¹⁹, Birgit Hünicke³, Jaak
 7 Jaagus²⁰, Mart Jüssi²¹, Jukka Käyhkö²², Nina Kirchner¹⁹, Erik Kjellström², Karol Kulinski²³,
 8 Andreas Lehmann²⁴, Göran Lindström², Wilhelm May²⁵, Paul A. Miller^{25,26}, Volker Mohrholz¹,
 9 Bärbel Müller-Karulis²⁷, Diego Pavón-Jordán²⁸, Markus Quante²⁸, Marcus Reckermann²⁹,
 10 Anna Rutgersson³⁰, Oleg P. Savchuk²⁷, Martin Stendel¹², Laura Tuomi¹⁷, Markku Viitasalo⁵,
 11 Ralf Weisse³, Wenyan Zhang³

12 ¹Department of Physical Oceanography and Instrumentation, Leibniz Institute for Baltic Sea Research
 13 Warnemünde, Rostock, Germany

14 ²Research and Development Department, Swedish Meteorological and Hydrological Institute, Sweden

15 ³Institute of Coastal Systems-Analysis and Modeling, Helmholtz-Zentrum Hereon, Geesthacht, Germany

16 ⁴Department of Ecology, Environment and Plant Sciences, Stockholm University, Stockholm, Sweden

17 ⁵Marine Research Centre, Finnish Environment Institute, Finland

18 ⁶Marine Research Institute, University of Klaipeda, Lithuania

19 ⁷Swedish Museum of Natural History, Stockholm, Sweden

20 ⁸Environmental and Marine Biology, Åbo Akademi University, Finland

21 ⁹Coalition Clean Baltic, Uppsala, Sweden

22 ¹⁰Information and Statistics Department, Swedish Meteorological and Hydrological Institute, Norrköping, Sweden

23 ¹¹Department of Bioscience, Aarhus University, Roskilde, Denmark

24 ¹²National Centre for Climate Research, Danish Meteorological Institute, Copenhagen, Denmark

25 ¹³Gavia EcoResearch, Winsen (Luhe), Germany

26 ¹⁴Deutsches Klimarechenzentrum, Hamburg, Germany

27 ¹⁵Department of Biology, University of Turku, Turku, Finland

28 ¹⁶International Institute for Applied Systems Analysis (IIASA), Laxenburg, Austria

29 ¹⁷Finnish Meteorological Institute, Helsinki, Finland

30 ¹⁸Department of Biological and Environmental Sciences, University of Helsinki, Helsinki, Finland

31 ¹⁹Department of Physical Geography, Stockholm University, Stockholm

32 ²⁰Department of Geography, Institute of Ecology and Earth Sciences, University of Tartu, Tartu, Estonia

33 ²¹Pro Mare, Estonia

34 ²²Department of Geography and Geology, University of Turku, Finland

35 ²³Institute of Oceanology, Polish Academy of Sciences, Gdansk, Poland

36 ²⁴GEOMAR Helmholtz Centre for Ocean Research, Kiel, Germany

37 ²⁵Centre for Environmental and Climate Science, Lund University, Lund, Sweden

38 ²⁶Department of Physical Geography and Ecosystem Science, Lund University, Lund, Sweden

39 ²⁷Baltic Sea Centre, Stockholm University, Stockholm, Sweden

40 ²⁸Department of Terrestrial Ecology, Norwegian Institute for Nature Research (NINA), P.O. Box 5685 Torgarden,
 41 N-7485 Trondheim, Norway

42 ²⁹International Baltic Earth Secretariat, Helmholtz-Zentrum Hereon, Geesthacht, 21502, Germany

43 ³⁰Department of Earth Sciences, Uppsala University, Sweden

44 [†]Deceased

46 Correspondence to: H. E. Markus Meier (markus.meier@io-warnemuende.de)



Abstract. Based on the Baltic Earth Assessment Reports of this thematic issue in Earth System Dynamics and recent peer-reviewed literature, current knowledge about the effects of global warming on past and future changes in climate of the Baltic Sea region is summarized and assessed. The study is an update of the Second Assessment of Climate Change (BACC II) published in 2015 and focusses on the atmosphere, land, cryosphere, ocean, sediments and the terrestrial and marine biosphere. Based on the summaries of the recent knowledge gained in paleo-, historical and future regional climate research, we find that the main conclusions from earlier assessments remain still valid. However, new long-term, homogenous observational records, e.g. for Scandinavian glacier inventories, sea-level driven saltwater inflows, so-called Major Baltic Inflows, and phytoplankton species distribution and new scenario simulations with improved models, e.g. for glaciers, lake ice and marine food web, have become available. In many cases, uncertainties can now be better estimated than before, because more models can be included in the ensembles, especially for the Baltic Sea. With the help of coupled models, feedbacks between several components of the Earth System have been studied and multiple driver studies were performed, e.g. projections of the food web that include fisheries, eutrophication and climate change. New data sets and projections have led to a revised understanding of changes in some variables such as salinity. Furthermore, it has become evident that natural variability, in particular for the ocean on multidecadal time scales, is greater than previously estimated, challenging our ability to detect observed and projected changes in climate. In this context, the first paleoclimate simulations regionalized for the Baltic Sea region are instructive. Hence, estimated uncertainties for the projections of many variables increased. In addition to the well-known influence of the North Atlantic Oscillation, it was found that also other low-frequency modes of internal variability, such as the Atlantic Multidecadal Variability, have profound effects on the climate of the Baltic Sea region. Challenges were also identified, such as the systematic discrepancy between future cloudiness trends in global and regional models and the difficulty of confidently attributing large observed changes in marine ecosystems to climate change. Finally, we compare our results with other coastal sea assessments, such as the North Sea Region Climate Change Assessment (NOSCCA) and find that the effects of climate change on the Baltic Sea differ from those on the North Sea, since Baltic Sea oceanography and ecosystems are very different from other coastal seas such as the North Sea. While the North Sea dynamics is dominated by tides, the Baltic Sea is characterized by brackish water, a perennial vertical stratification in the southern sub-basins and a seasonal sea ice cover in the northern sub-basins.

During the time in which this paper was prepared, shortly before submission, Christian Dieterich passed away (1964-2021). This sad event marked the end of the life of a distinguished oceanographer and climate scientist who made important contributions to the climate modeling of the Baltic Sea, North Sea and North Atlantic regions. This paper is dedicated to him.

1 Introduction

1.1 Overview

In this study, the results concerning climate change of the various articles of this thematic issue, the so-called Baltic Earth Assessment Reports (BEARs) coordinated by the Baltic Earth program¹ (Meier et al., 2014), and other relevant literature are summarized and assessed. We focus on the knowledge gained during 2013-2020 about past, present and future climate changes in the Baltic Sea region. The methodology of all BEARs follows the earlier assessments of climate change in the Baltic Sea region (BACC Author Team, 2008; BACC II Author Team, 2015). The aim of this review is to inform and update scientists, policymakers and stakeholders about recent research results. The focus is on the atmosphere, hydrosphere, cryosphere, lithosphere and biosphere. In contrast to the earlier assessments, we do not investigate the impact of climate change on human society. We start (Section 1) with a summary of key messages from the earlier assessments of climate change in the Baltic Sea region, a description of the Baltic Sea region and its climate, a comparison of the Baltic Sea with other coastal seas and a summary of current knowledge on global climate change assessed in the latest Intergovernmental Panel on Climate Change (IPCC) reports. In Section 2, the methods for the literature assessment, climate model data and uncertainty

¹ <https://baltic.earth>



estimates are outlined. In Section 3, the results of the assessment for selected variables (Table 1) under past (paleoclimate), present (historical period with instrumental data) and future (until 2100) climate conditions are presented, *inter alia* by summarizing the results in various papers of this special issue by Lehmann et al. (2021), Kuliński et al. (2021), Rutgersson et al. (2021), Weisse et al. (2021), Gröger et al. (2021a), Christensen et al. (2021), Meier et al. (2021a) and Viitasalo (2021) and by other relevant review studies. In Section 4, the interactions of climate with other anthropogenic drivers are summarized from Reckermann et al., 2021. As the adjacent North Sea has different physical characteristics and topographical features but is located in a similar climatic zone as the Baltic Sea, we compare the results of this assessment with the results of the North Sea Region Climate Change Assessment (NOSCCA; Quante and Colijn, 2016; Section 5). Knowledge gaps (Section 6), key messages (Section 7) and conclusions (Section 8) finalize the study.

1.2 The BACC and BEAR projects

This assessment is an update to the two BACC books, published as comprehensive textbooks in 2008 and 2015 (BACC Author Team, 2008; BACC II Author Team, 2015). The acronym BACC (**B**ALTEX **A**ssessment of **C**limate **C**hange) refers to the Baltic Earth pre-cursor programme BALTEX (Baltic Sea Experiment; Reckermann et al., 2011). From the beginning, BALTEX tried to approach three basic questions: 1. What is the evidence for past and present regional climate change? 2. What are the model projections for future regional climate change? 3. Which impacts can we already observe in terrestrial and marine ecosystems?

First ideas for a comprehensive appraisal of the current knowledge on climate change and its impact on the Baltic Sea region evolved in 2004 as it became evident that there was a demand for this, in particular by the Baltic Marine Environment Protection Commission, the Helsinki Commission (HELCOM; BALTEX, 2005). A steering group of leading experts from the Baltic Sea region was enlisted, which elaborated a grand chapter structure at several preparatory workshops and meetings and also recruited a group of lead authors. In total, more than 80 scientists from 12 countries and all relevant scientific disciplines contributed to the first regional climate change assessment (BACC Author Team, 2008), which underwent a rigorous review process.

In 2011, a second edition of the BACC book was initiated as an update, but also as a complement to the first book, by including new topics like an overview of changes since the last glaciation, and a new section on regional drivers and attribution. The Second Assessment of Climate Change for the Baltic Sea Basin (BACC II Author Team, 2015) was published in 2015, used the same procedures and principles, but with a new steering and author group, and under the auspices of Baltic Earth, the successor of BALTEX. Close collaboration with HELCOM was envisaged from the very beginning, with HELCOM using material from both BACC assessments for their own climate change assessment reports (HELCOM, 2007; 2013b).

In 2018, the Baltic Earth Science Steering Group decided to produce a series of new assessment reports, the BEARs, on the current Baltic Earth Grand Challenges, Earth System models and projections for the Baltic Sea Region. The BEARs are comprehensive, peer-reviewed review articles in journal format, and the update to BACC II (this article) is one of the ten envisaged contributions summarizing the current knowledge on regional climate change and its impacts, knowledge gaps and advice for future work. For further details about our knowledge on climate change, the reader is referred to the other BEARs. The close collaboration with HELCOM is continued in



the joint HELCOM-Baltic Earth Expert Network of Climate Change (EN CLIME), which was assembled to produce a Baltic Earth – HELCOM Climate Change Fact Sheet for the Baltic Sea region².

135

Hence, this thematic issue comprises nine BEARs and, in addition, this summary of the current knowledge about past, present and future climate changes for the Baltic Sea region (“BACC III”). Below a few key-words characterizing the BEARs’ contents are listed:

1. Salinity dynamics of the Baltic Sea (Lehmann et al., 2021): water and energy cycles with focus on Baltic Sea salinity during past climate variability, meteorological patterns at various space and time scales and mesoscale variability in precipitation, variations in river runoff and various types of inflows of saline water, exchange of water masses between various sub-basins and vertical mixing processes. The paper also includes the observed trends of salinity during the last >100 years.
2. Baltic Earth Assessment Report on the biogeochemistry of the Baltic Sea (Kuliński et al., 2021): terrestrial biogeochemical processes and nutrient loads to the Baltic Sea, transformations of C, N, P in the coastal zone, organic matter production and remineralization, oxygen availability, burial and turnover of C, N, P in the sediments, the Baltic Sea CO₂ system and seawater acidification, role of specific microorganisms in Baltic Sea biogeochemistry, interactions between biogeochemical processes and chemical contaminants.
3. Natural Hazards and Extreme Events in the Baltic Sea region (Rutgersson et al., 2021): extremes in wind, waves, and sea level, sea-effect snowfall, river floods, hot and cold spells in the atmosphere, marine heat waves, droughts, ice seasons, ice ridging, phytoplankton blooms and some implications of extreme events for society (including forest fires, coastal flooding, offshore wind mills and shipping).
4. Sea Level Dynamics and Coastal Erosion in the Baltic Sea Region (Weisse et al., 2021): sea level dynamics and coastal erosion in past and future climates. The current knowledge about the diverse processes affecting mean and extreme sea level changes is assessed.
5. Coupled regional Earth system modelling in the Baltic Sea Region (Gröger et al., 2021a): status report on coupled regional Earth system modeling with focus on the coupling between atmosphere and ocean, atmosphere and land surface including dynamic vegetation, ocean, sea ice and waves and atmosphere and hydrological components to close the water cycle.
6. Atmospheric regional climate projections for the Baltic Sea Region until 2100 (Christensen et al., 2021): comparison of coupled and uncoupled regional future climate model projections. As the number of atmospheric scenario simulations of the Euro-CORDEX program (Kjellström et al., 2018; Teichmann et al., 2018; Jacob et al., 2018) is large, uncertainties can be better estimated and the effects of mitigation measures can be better addressed compared to earlier assessments.
7. Oceanographic regional climate projections for the Baltic Sea until 2100 (Meier et al., 2021a): new projections with a coupled physical-biogeochemical ocean model of future climate considering global sea level rise, regional climate change and nutrient input scenarios are compared with previous studies and the differences are explained by differing scenario assumptions and experimental setups.

² <https://helcom.fi/helcom-at-work/groups/state-and-conservation/en-clime/>,
https://baltic.earth/projects/en_clime/index.php/en



- 170 8. Climate change and the Baltic Sea ecosystem: direct and indirect effects on species, communities and
 171 ecosystem function (Viitasalo, 2021): impact of past and future climate changes on the marine ecosystem.
- 172 9. Human impacts and their interactions in the Baltic Sea region (Reckermann et al., 2021): interlinkages of
 173 factors controlling environmental changes. Changing climate is only one of the many anthropogenic and
 174 natural impacts that effect the environment. Other investigated factors are coastal processes, hypoxia,
 175 acidification, submarine groundwater discharge, marine ecosystems, non-indigenous species, land use
 176 and land cover (called natural) and agriculture and nutrient loads, aquaculture, fisheries, river regulations
 177 and restorations, offshore wind farms, shipping, chemical contaminants, unexploded and dumped warfare
 178 agents, marine litter and microplastics, tourism, and coastal management (called human-induced).

179 1.3 Summary of BACC I and II key messages

180 Quotation by the BACC II Author Team (2015):

181 “The key findings of the BACC I assessment were as follows:

- 182 • The Baltic Sea region is warming, and the warming is almost certain to continue throughout the twenty-first
 183 century.
- 184 • It is plausible that the warming is at least partly related to anthropogenic factors.
- 185 • So far, and as is likely to be the case for the next few decades, the signal is limited to temperature and to directly
 186 related variables, such as ice conditions.
- 187 • Changes in the hydrological cycle are expected to become obvious in the coming decades.
- 188 • The regional warming is almost certain to have a variety of effects on terrestrial and marine ecosystems—some
 189 will be more predictable (such as the changes in phenology) than others.

190

191 The key findings of the BACC II assessment [...] are as follows:

- 192 1. The results of the BACC I assessment remain valid.
- 193 2. Significant additional material has been found and assessed. Some previously contested issues have been
 194 resolved (such as trends in sea-surface temperature).
- 195 3. The use of multi-model ensembles seems to be a major improvement; there are first detection studies, but
 196 attribution is still weak.
- 197 4. Regional climate models still suffer from biases related to the heat and water balances. The effect of
 198 changing atmospheric aerosol load to date cannot be described; first efforts at describing the effect of
 199 land-use change have now been done.
- 200 5. Data homogeneity is still a problem and is sometimes not taken seriously enough.
- 201 6. The issue of multiple drivers on ecosystems and socioeconomics is recognized, but more efforts to deal
 202 with them are needed.
- 203 7. In many cases, the relative importance of different drivers of change, not only climate change, needs to
 204 be evaluated (e.g. atmospheric and aquatic pollution and eutrophication, overfishing, and changes in land
 205 cover).
- 206 8. Estimates of future concentrations and deposition of substances such as sulphur and nitrogen oxides,
 207 ammonia/ammonium, ozone, and carbon dioxide depend on future emissions and climate conditions.
- 208 Atmospheric warming seems relatively less important than changes in emissions. The specification of



- future emissions is plausibly the biggest source of uncertainty when attempting to project future deposition or ocean acidification.
9. In the narrow coastal zone, the combination of climate change and land uplift acting together creates a particularly challenging situation for plant and animal communities in terms of adaptation to changing environmental conditions.
 10. Climate change is a compounding factor for major drivers of changes in freshwater biogeochemistry, but evidence is still often based on small-scale studies in time and space. The effect of climate change cannot yet be quantified on a basin-wide scale.
 11. Climate model scenarios show a tendency towards future reduced salinity, but due to the large bias in the water balance projections, it is still uncertain whether the Baltic Sea will become less or more saline.
 12. Scenario simulations suggest that the Baltic Sea water may become more acidic in the future. Increased oxygen deficiency, increased temperature, changed salinity, and increased ocean acidification are expected to affect the marine ecosystem in various ways and may erode the resilience of the ecosystem.
 13. When addressing climate change impacts on, for example, forestry, agriculture, urban complexes, and the marine environment in the Baltic Sea basin, a broad perspective is needed which considers not only climate change but also other significant factors such as changes in emissions, demographic and economic changes, and changes in land use.
 14. Palaeoecological ‘proxy’ data indicate that the major change in anthropogenic land cover in the Baltic Sea catchment area occurred more than two thousand years ago. Climate model studies indicate that past anthropogenic land-cover change had a significant impact on past climate in the northern hemisphere and the Baltic Sea region, but there is no evidence that land cover change since AD 1850 was even partly responsible for driving the recent climate warming.”
- For comparison, the findings of this assessment study can be found in Section 8.

1.4 Baltic Sea region characteristics

1.4.1 Climate variability of the Baltic Sea Region

The Baltic Sea region (including the Kattegat) is located between maritime temperate and continental sub-arctic climate zones, in the latitude–longitude box 54°N – $66^{\circ}\text{N} \times 9^{\circ}\text{E}$ – 30°E (Fig. 1). The climate of the Baltic Sea region has a large variability due to the opposing effects of moist and relatively mild marine air flows from the North Atlantic Ocean and the Eurasian continental climate. The regional weather regimes varies depending on the exact location of the polar front and the strength of the westerlies, and both seasonal and interannual variations are considerable. The westerlies are particularly important in winter, when the temperature difference between the marine and continental air masses is large.

The southern and western parts of the Baltic Sea belong to the Central European mild climate zone in the westerly circulation. The northern part locates at the polar front and the winter climate is cold and dry due to cold arctic air outbreaks from the east. In terms of classical meteorology, during winter the polar front fluctuates over the Baltic Sea region but during summer it is located farther to the north. Depending on the particular year, the central part of the Baltic Sea can be either on the mild or the cold side of the polar front. The temperature difference between winter and summer is much larger in the north. During warm summers and cold winters the air pressure field is



smooth and winds are weak, and blocking high pressure situations are common. During such periods, the weather can be very stable for several weeks.

250

The climate of the Baltic Sea region is strongly influenced by the large-scale atmospheric variability (e.g. Andersson, 2002; Tinz, 1996; Meier and Kauker, 2003; Omstedt and Chen, 2001; Zorita and Laine, 2000; Lehmann et al., 2002). In particular, the North Atlantic Oscillation (NAO), blocking and, on longer time scales, circulation patterns related to the Atlantic Multidecadal Oscillation (AMO) play important roles for the climate of the Baltic Sea region. The AMO consists of an unforced component which is the result of atmosphere-ocean interactions (e.g. Wills et al., 2018) and a forced component such as volcanic eruptions (Mann et al., 2021; Mann et al., 2020). However, the relative importance of its forced and unforced components is still debated (Mann et al., 2021).

259

The NAO is the dominant mode of near-surface pressure variability over the North Atlantic and its influence is strongest in winter (Hurrell et al., 2003), when it accounts for almost one-third of the sea level pressure (SLP) variance (e.g. Kauker and Meier, 2003). During the positive (negative) phase of the NAO the Icelandic Low and Azores High pressure systems are stronger (weaker), leading to a stronger (weaker) than normal westerly flow (Hurrell, 1995). Positive NAO phases are associated with mild temperatures and increased precipitation and storminess whereas negative NAO phases are characterized by warm summers, cold winters, and less precipitation (Hurrell et al., 2003). Increasing winter temperatures in the Baltic Sea have also been linked to an observed shift in the storm tracks (BACC II Author Team, 2015). There is a large interannual to interdecadal variability in the NAO, reflecting interactions with and changes in surface properties, including sea surface temperature (SST) and sea ice cover. This makes it difficult to detect a possible long-term trend in the NAO.

270

Atmospheric blocking occurs when persistent high pressure systems interrupt the normally westerly flow over the middle and high latitudes, e.g. the North Atlantic. By redirecting the pathways of midlatitude cyclones, blockings lead to negative precipitation anomalies in the region of the blocking anticyclone and positive anomalies in the surrounding areas (Sousa et al., 2017). In this way, blockings can also be associated with extreme events such as heavy precipitation (Lenggenhager and Martius, 2019) or drought (Schubert et al., 2014).

276

The AMO describes fluctuations in North Atlantic sea surface temperature (SST) with a period of 50-90 years (Knight et al., 2006). Thus only a few distinct AMO phases have been observed in the 150-year instrumental record. However, a recent model study suggests that variations in the AMO may influence atmospheric circulation that leads to additional precipitation over the Baltic Sea region (Börgel et al., 2018). Further, it was found that the AMO altered the zonal position of the NAO and affected the regional imprint of the NAO for the Baltic Sea region (Börgel et al., 2020).

1.4.2 A unique brackish water basin

The Baltic Sea is a unique brackish water basin in the World Ocean which has a salinity less than 24.7 g kg⁻¹ in all areas (Leppäranta and Myrberg, 2009; Voipio, 1981; Magaard and Rheinheimer, 1974; Feistel et al., 2008; Omstedt et al., 2014). The sea is very shallow (with a mean depth of only 54 m), and can be characterized as a number of sub-basins (Fig. 2). The Baltic Sea has the only connection to the North Sea through the Danish straits



(Fig. 2). The exchange of water between the Baltic Sea and North Sea through the narrow straits is quite limited. The Baltic Sea has a positive fresh water balance with an average salinity of about 7.4 g kg^{-1} – this being only one-fifth the salinity of the World Ocean, thus water masses are brackish. The Baltic Sea is located between mild maritime and continental sub-arctic climate zones and partly ice-covered in every winter. However, it is completely frozen over only during extremely cold winters. The highly variable coastal geomorphology and the extended archipelago areas make the Baltic Sea unique (see Section 5).

The World Ocean has only four large brackish water basins (Leppäranta and Myrberg, 2009). These are from the largest to the smallest the Black Sea (Ivanov and Belokopytov, 2013) located between Europe and Asia Minor, the Baltic Sea, the Gulf of Ob in the Kara Sea (Volkov et al., 2002) and the Chesapeake Bay (Kjerfve, 1988), on the east coast of the United States of America. All these sea areas developed into brackish water basins during the Holocene. During the most recent (Weichselian) glaciation period the Black Sea was a freshwater lake, the Baltic Sea and the Gulf of Ob were under the Eurasian ice sheet, and the Chesapeake Bay was a river valley (Leppäranta and Myrberg, 2009). The mean depth of the Black Sea is 1200 m, and due to the strong salinity stratification and extremely slow deep water renewal the water masses below 200 m are anoxic. The Sea of Azov in the north-eastern part of the Black Sea is often frozen during the winter. The Gulf of Ob is the long (800 km), narrow estuary of the River Ob in the Kara Sea in the Russian Arctic, and ice-covered in winter. Finally, Chesapeake Bay is a small, very shallow basin and a drowned river valley or ria, in the humid subtropical climate zone, with hot summers and ice formation in river mouths in some winters.

Table 2 gives basic information of the brackish water seas and other basins comparable with the Baltic Sea. Most similar to the brackish water seas is Hudson Bay (Roff and Legendre, 1986). It is an oceanic, semi-enclosed basin with a positive fresh water balance, and a salinity of about 30 g kg^{-1} . In contrast, small Mediterranean seas with a negative fresh water balance and salinities above 40 g kg^{-1} are found in the tropical zone; e.g. the Red Sea and Persian Gulf. The largest lakes are comparable in size to the Baltic Sea, and the Caspian Sea is even larger in volume.

The Baltic Sea basin is a very old geomorphological depression. Prior to the Weichselian glaciation this basin contained the Eem Sea, which extended from the North Sea to the Barents Sea, making Fennoscandia an island. At the end of the Weichselian glaciation, 13,500 years ago, the Baltic Ice Lake was formed by glacier meltwater. During the Holocene fresh and brackish phases followed dictated by the balance of glacier retreats and progressions, land uplift and eustatic changes of the global sea level (Tikkanen and Oksanen, 2002). The present brackish phase commenced 7000 years ago, and since about 2000 years ago the salinity has been close to the present level. Postglacial land uplift has slowly changed the Baltic Sea landscape, making it possible to observe how land rises from the sea and how terrestrial life gradually takes over. People living in the region have adapted to this slow long-term change.

1.4.3 The Baltic Sea - a specific European sea

The basic features of the European seas reveal key differences, in areal extent, depth profile, salinity level, fresh water budget, climate, and tidal motions (Table 3). The Baltic Sea and the North Sea are shallow, with a mean depth of less than 100 m; the Baltic can be described as a “coastal sea”, with a mean depth of only 54 m. The Black



328 Sea and the Mediterranean Sea are much deeper, with mean depths of approximately 1200 m and 1500 m,
 329 respectively, whereas the North-East Atlantic reaches the full oceanic depth of ca. 4 km, fringed by much shallower
 330 continental shelf areas, at about 400 m. These depth differences influence, among other things, the mixing of the
 331 water column, variability in temperature, and distribution of benthic ecosystems (Myrberg et al., 2019).

332

333 Among the European Seas, the Baltic Sea physics stands out in terms of its small tidal amplitudes, low salinity,
 334 strong stratification and anoxic conditions. Additionally, frequent and spatially extensive upwelling and regular
 335 seasonal ice cover are typical of the Baltic Sea (Leppäranta and Myrberg, 2009). To summarize:

- 336 • The Baltic Sea is permanently stratified due to a large salinity (density) difference between the fresh upper
 337 layer and the more saline bottom layer. This limits ventilation, leading to oxygen deficiency in the bottom
 338 layer. For instance in autumn 2016, some 70 000 km² of the seabed experienced permanent hypoxia.
 339 Irregular Major Baltic Inflows (MBIs; Matthäus and Franck, 1992; Mohrholz, 2018) are the main
 340 mechanism transporting oxygen-rich waters from the North Sea to Baltic Sea deeps. The associated salt
 341 transport in turn intensifies vertical stratification and eventually enlarges hypoxic area (Conley et al.,
 342 2002).
- 343 • In the small, semi-enclosed Baltic Sea, almost any winds are likely to blow parallel to some section of
 344 the coast and thus cause coastal upwelling. At the Swedish south-western coast, upwelling occurs 25-40
 345 % of time (Lehmann et al., 2012). At times, about one third of the entire Baltic Sea may be under the
 346 influence of upwelling.
- 347 • Among European seas, ice is a unique feature of the Baltic Sea that strongly limits air-sea interaction and
 348 modifies the Baltic Sea ecosystem in many ways.

349

350 The salinity in the Baltic Sea is not only an oceanographic variable as in other more ventilated seas, but also
 351 integrates the complete water and energy cycles, with their specific Baltic Sea features. Baltic Sea salinity, and
 352 especially its low basic value and the large variations, is also an elementary factor controlling the marine
 353 ecosystem. The salinity dynamics is governed by several factors: net precipitation, river runoff, surface outflow of
 354 brackish Baltic Sea water and the compensating deep inflow of higher salinity water from the Kattegat. The latter
 355 is strongly controlled by the prevailing atmospheric forcing conditions. Due to freshwater supply from the Baltic
 356 Sea catchment area and due to the limited water exchange with the World Ocean, surface salinity varies from > 20
 357 g kg⁻¹ in Kattegat to < 2 g kg⁻¹ in the Bothnian Bay and is close to zero at the mouth of the River Neva, in the
 358 easternmost end of the Gulf of Finland. In the vertical direction, the dynamics of the Baltic Sea is characterized
 359 by a permanent, two-layer system because of a pronounced, perennial vertical gradient in salinity. In summer, a
 360 shallow thermocline is also formed, complicating the vertical structure.

361 1.5 Global climate change

362 In the following, a brief overview is given of the latest global climate assessments, based on the IPCC Fifth
 363 Assessment Report (AR5; IPCC, 2014b) and results so far available from the current Coupled Model
 364 Intercomparison Project (CMIP) phase 6 (Eyring et al., 2016). The focus is on large-scale changes in climate that
 365 are of particular relevance for the Baltic Sea region (mainly North Atlantic, Arctic). Furthermore, whenever
 366 feasible, changes are described in terms of pattern scaling which relies on the fact that for many quantities the
 367 geographical change patterns are sufficiently consistent across models and scenarios to emerge from the



background noise (IPCC, 2014a). Hence, changes in e.g. local temperatures can be scaled to changes per °C of global mean temperature change relative to 1981-2005 (Christensen et al., 2019).

370

Our future climate change assessment relies on the concentration driven scenarios RCP2.6, RCP4.5 and RCP8.5 from the CMIP5 suite (RCP = Representative Concentration Pathway), corresponding to changes in radiative forcing for the 21st century. Hence, policy targeted goals inspired by the United Nations Framework Convention on Climate Change (UNFCCC; United Nations Climate Change, 2015) to limit global mean warming below 2.0 or 1.5°C compared to preindustrial level, i.e. prior to the 20th century (the Paris Agreement, PA), are not considered in many scenario simulations but referred to studies within the Euro-CORDEX framework (Kjellström et al., 2018; Teichmann et al., 2018; Jacob et al., 2018) and for a broader region. In order to achieve the goal of a significant reduction of the risks and impacts of climate change, the Paris Agreement commits the participating countries to aim “to reach global peaking of greenhouse gas emissions as soon as possible” and “to undertake rapid reductions thereafter in accordance with best available science, so as to achieve a balance between anthropogenic emissions by sources and removals by sinks of greenhouse gases in the second half of this century”. Furthermore, the countries “should take action to conserve and enhance, as appropriate, sinks and reservoirs of greenhouse gases [...], including forests”.

384

RCP8.5 is a totally unmitigated scenario and assumes a radiative forcing of +8.3 W m⁻² in year 2100, as compared to the preindustrial period. Assumptions for RCP8.5 are described in Riahi et al. (2011). RCP8.5 has been criticized because it assumes continued use of coal for energy production translating into too high greenhouse gas emissions. Moderate mitigation actions are reflected by RCP4.5 (Thomson et al., 2011), and RCP2.6 was developed for effective mitigation scenarios aiming at limiting global mean warming to ~+2°C (van Vuuren et al., 2011). With respect to global development, RCP2.6 and RCP8.5 might be unrealistic (Hausfather and Peters, 2020). However, both scenarios can be used as envelopes of plausible pathways of future greenhouse gas emissions.

392

Confidence levels expressing evidence and agreement are provided following the definitions of the IPCC (see Method Section 2.3).

1.5.1 Atmosphere

1.5.1.1 Surface air temperature

For the three considered scenarios, the IPCC AR5 (IPCC, 2014a; 2014b; Collins et al., 2013) reported a likely increase in global mean air temperature for the period 2081-2100 relative to 1986-2005 in the likely range (5th to 95th percentile of CMIP5 models) between 0.3 to 1.7°C (RCP2.6), 1.1 to 2.6°C (RCP4.5), and 2.6 to 4.8°C (RCP8.5). The corresponding mean changes are 1.0°C (RCP2.6), 1.8°C (RCP4.5) and 3.7°C (RCP8.5; IPCC, 2014b).

402

The large-scale geographical patterns of change remain stable among CMIP5 models and are consistent with the results of the IPCC AR4. The dominant feature is a strong warming of the Arctic north of 67.5 °N that exceeds global mean warming by a factor 2.2 to 2.4. The Arctic warming is strongest for the winter season, when sea ice retreat and reduced snow cover provide positive feedbacks (Arctic amplification), and weakest in summer, when melting sea ice consumes latent heat and the ice free ocean absorbs heat (IPCC, 2014b). Besides these



thermodynamic processes, the lateral transport of latent heat into the Arctic increases under global warming. Weakest warming is found over the Southern Ocean and in the North Atlantic south of Greenland with minimum values per degree global warming of about $0.25^{\circ}\text{C }^{\circ}\text{C}^{-1}$ (Fig. 12.10 in IPCC, 2014b). This is partly due to a deeper ocean mixed layer that promotes vigorous oceanic heat uptake in these regions compared to others. Generally, land masses warm at a rate 1.4 to 1.7 times more than open ocean regions, leading to a pronounced land-sea pattern in the temperature anomaly and indicating to a lower effective heat capacity of continents compared to the ocean.

1.5.1.2 Precipitation

Projected global precipitation changes scale nearly linear with global mean temperature changes and range from $+0.05 \text{ mm d}^{-1}$ or $\sim 2\%$ (RCP2.6) to 0.15 mm d^{-1} or $\sim 5\%$ (RCP8.5; IPCC, 2014a). As a result of an accelerated global water cycle, the contrast between dry and wet regions in annual mean precipitation increases. Likewise, there is high confidence that the contrast between wet and dry seasons will become more pronounced (IPCC, 2014a). In the mid to high latitudes, yearly mean precipitation generally increases, with the strongest response over the Arctic, exceeding almost everywhere $+12\% ^{\circ}\text{C}^{-1}$.

Precipitation changes vary greatly among models. Under RCP8.5, high latitude land masses will likely get more precipitation, due to higher moisture content of the lower atmosphere and an increased moisture transport from the tropics (IPCC, 2014a). In the northern hemisphere the poleward branch of the Hadley Cell will expand further north, causing a northward expansion of the subtropical dry zone and reducing precipitation in affected regions. Further dynamical changes probably include a poleward shift of mid-latitude storm tracks (Seager et al., 2010; Scheff and Frierson, 2012) which is, however, of low confidence, especially for the North Atlantic region (IPCC, 2014a).

1.5.2 Cryosphere

The IPCC AR5 postulates a reduction of average February Arctic sea ice extent ranging from 8% for RCP2.6 to 34% for RCP8.5. For the monthly mean summer minimum in September, reductions range from 43% for RCP2.6 to 94% for RCP8.5. These values are given medium confidence, because of biases in the simulation of present day trends and a large spread across models. For September, ice free conditions are reached before 2090 in 90 % of all CMIP5 models.

The permafrost area is projected to decrease in a likely range from $24 \pm 16\%$ for RCP2.6 to $69 \pm 20\%$ for RCP8.5.

Arctic autumn and spring snow cover are projected to decrease by 5–10%, under RCP2.6, and 20–35% under RCP8.5 (high confidence). In high mountain areas, projected decreases in mean winter snow depth are in a likely range of 10–40 % for RCP2.6 and 50–90% for RCP8.5. The likely range of projected inland glacier mass reductions (ice sheets excluded) between 2015 and 2100 varies from $18 \pm 7\%$ for RCP2.6 to $36 \pm 11\%$ for RCP8.5. Regions with mostly smaller glaciers (e.g. Central Europe, Scandinavia) are projected to lose over 80% of their current ice mass by 2100 under RCP8.5 (medium confidence), with many glaciers disappearing regardless of future emissions (very high confidence).



445 1.5.3 Ocean

446 1.5.3.1 Sea level

447 For 2081-2100, global mean sea level (GMSL) is projected to rise between 0.40 m under RCP2.6 (likely range
 448 0.26-0.55m) and 0.63 m under RCP8.5 (likely range 0.45-0.82 m) relative to 1986-2005 (IPCC, 2014a; their
 449 Chapter 13, Table 13.5). In all scenarios, thermal expansion gives the largest contribution to GMSL rise,
 450 accounting for about 30 to 55% of the projections. Glaciers are the next largest contributor, accounting for about
 451 15-35%. By 2100, the Greenland Ice Sheet's projected contribution to GMSL rise is 0.07 m (likely range 0.04–
 452 0.12 m) under RCP2.6, and 0.15 m (likely range 0.08–0.27 m) under RCP8.5. The Antarctic Ice Sheet is projected
 453 to contribute 0.04 m (likely range 0.01–0.11 m) under RCP2.6, and 0.12 m (likely range 0.03–0.28 m) under
 454 RCP8.5. Uncertainties concerning the melting of ice sheets are, however, intensively discussed (Bamber et al.,
 455 2019).

456

457 Based on the same suite of model projections from CMIP5, the IPCC Special Report on the Ocean and Cryosphere
 458 in a Changing Climate (IPCC, 2019a) has updated these numbers by including new estimates of the contribution
 459 from Antarctica, for which new ice-sheet modelling results were available (Oppenheimer et al., 2019). While the
 460 differences in projected changes until 2100 are small for RCP2.6, projected changes for RCP8.5 increased by about
 461 10 cm compared to AR5 (see Section 3.3.5.4).

462 1.5.3.2 Water temperature and salinity

463 By the end of the century, the projected global ocean warming ranges from about 1°C (RCP2.6) to more than 3°C
 464 (RCP8.5) at the surface and from 0.5°C (RCP2.6) to 1.5°C (RCP8.5) at a depth of 1km. The subtropical waters of
 465 the Southern Ocean and the North Atlantic are projected to become saltier, whereas almost all other regions
 466 become fresher, in particular the northern North Atlantic (IPCC, 2014a). The freshening at high latitudes in the
 467 North Atlantic and Arctic basin is consistent with a weaker Atlantic Meridional Overturning Circulation (AMOC),
 468 and a decline in the volume of sea ice, as well as with the intensified water cycles (IPCC, 2019a).

469

470 By the end of the century, the annual mean stratification of the top 200 m (averaged between 60°S–60°N, relative
 471 to 1986–2005) is projected to increase in the very likely range of 1–9% for RCP2.6 and 12–30% for RCP8.5
 472 (IPCC, 2019a).

473 1.5.3.3 Atlantic Meridional Overturning Circulation

474 Based on the CMIP5 models, the AMOC is estimated to be reduced by 11% (1 to 24%) under RCP2.6 and 34%
 475 (12 to 54%) under RCP8.5. There is low confidence in the projected evolution of the AMOC beyond the 21st
 476 century (IPCC, 2014a).

477 1.5.4 Marine biosphere

478 By 2081-2100, net primary productivity relative to 2006-2015 will very likely decline by 4–11% for RCP8.5, due
 479 to the combined effects of warming, stratification, light, nutrients and predation, with regional variations between
 480 low and high latitudes (IPCC, 2019a).

481



Globally, and relative to 2006–2015, the oxygen content of the ocean by 2081–2100 is very likely to decline by 1.6–2.0% for the RCP2.6 scenario, or by 3.2–3.7% for the RCP8.5 scenario (IPCC, 2019a). While warming is the primary driver of deoxygenation in the open ocean, eutrophication is projected to increase in estuaries due to human activities and due to intensified precipitation, which increase riverine nitrogen loads under both RCP2.6 and RCP8.5 scenarios, both by mid-century (2031–2060) and later (2071–2100; Sinha et al., 2017). Moreover, stronger stratification in estuaries due to warming is expected to increase the risk of hypoxia by reducing vertical mixing (IPCC, 2019a; Hallett et al., 2018; Warwick et al., 2018; Du et al., 2018).

1.5.5 Coupled Model Intercomparison Project

Future climate change assessments like the coming IPCC sixth Assessment Report AR6 (due 2021/2022), will rely on a new generation of Earth System Models (ESMs), developed during CMIP6, which offers a wider range of scenarios than during CMIP5. In particular, scenarios aiming to limit global warming to 1.5°C and 2.0°C and overshoot scenarios including negative emissions in the second part of the century will be available.

A subset of current CMIP6 models have been shown to be more sensitive to greenhouse gases than previous generations of CMIP models. Thus, the estimated response to an instantaneous doubling of CO₂ (equilibrium climate sensitivity, ECS) is higher in CMIP6 models (1.8 – 5.6°C) than in CMIP5 models (1.5 – 4.5°C) and their predecessors (Meehl et al., 2020). Indeed, the first transient simulations with the CMIP6 EC-Earth ESM found stronger warming than with earlier versions, with about half of the increase attributed to differences between CMIP5 and CMIP6 greenhouse gas forcing (Wyser et al., 2020).

However, it turns out that models with the highest projected warmings fail to capture past warming trends well, and therefore recent studies argue that those models should not be used for climate assessments and policy decisions (Forster et al., 2020; Nijssen et al., 2019; Tokarska et al., 2020; Brunner et al., 2020). Furthermore, systematic errors in many CMIP5 and CMIP6 models prevent the simulation of the observed 1951–2014 summer warming trend in Western Europe, and that neither higher resolution nor better representation of the sea surface is likely to improve this (Boé et al., 2020).

2 Methods

2.1 Assessment of literature

33 variables representing the components of the Earth system (atmosphere, land, terrestrial biosphere, cryosphere, ocean and sediment, marine biosphere) of the Baltic Sea region were selected (Table 1) and with respect to past, present and future climate changes corresponding scientific peer-reviewed publications and reports of scientific institutes since 2013 were assessed by 48 experts (see the section about author contributions). The year 2013 was chosen as a starting point because earlier material was included in the last assessment by the BACC II Author Team (2015). Information about climate change available in the BEARs (Section 2.1) was summarized and cross-references can be found in Table 1.

For the selected 33 variables and even more general, knowledge gaps (Section 6) and key messages (Section 7) as well as overall conclusions (Section 8) were formulated. Key messages, new compared to the results of the BACC



II Author Team (2015), were marked. The identified changes of the selected variables of the Earth system and their uncertainties, following the definitions of the IPCC reports as outlined in Section 2.3, are summarized in Table 10. The attribution of a changing variable to climate change, here the deterministic response to changes in external anthropogenic forcing such as greenhouse gas and aerosol emissions, is illustrated by Figure 34. This study does not claim to be complete, neither with regard to the importance of the limited selection of variables for the Earth system nor with regard to the discussed and assessed publications.

The assessment was done without influence from any political, economic or ideological group or party. The results of the BEARs including this summary about climate change impacts in the Baltic Sea region were used by the joint HELCOM-Baltic Earth Expert Network of Climate Change (EN CLIME) for the compilation of a Climate Change Fact Sheet for the Baltic Sea region.

For further details about the assessment methods, the reader is referred to the BACC Author Team (2008) and the BACC II Author Team (2015).

2.2 Proxy data, instrumental measurements and climate model data

In addition to selected figures that are reproduced from the literature, for the assessment previously published datasets were analyzed and discussed.

2.2.1 Past climate

For the Holocene climate evolution, paleo-pollen data with a decadal resolution, reconstructing seasonal temperature and precipitation changes compared to preindustrial climate (Mauri et al., 2015), were analyzed (Fig. 3). More accurate tree-ring data, resolving annual summer mean temperatures, are available for the past millennium (Luterbacher et al., 2016) and have been discussed here (Fig. 4). For further details, the reader is referred to Section 3.1.2.

2.2.2 Present climate

Historical station data of sea level pressure and sea surface temperature (SST) were used to calculate climate indices such as the NAO (sea level pressure differences, Fig. 5) and the AMO (SST anomalies, Fig. 6), describing decadal to multidecadal variability of the large-scale atmosphere circulation. Furthermore, selected records of variables such as air temperature (Fig. 8), river runoff (Fig. 10), land nutrient inputs (Fig. 11, Table 5), glacier masses (Fig. 12, Table 6), maximum sea ice extent (Fig. 14), ice thickness data (Figs. 15 and 16), length of the ice season (Fig. 17), sea level (Fig. 24) and gridded data sets of air temperature, e.g. the land-based CRUTEM4 data (Jones et al., 2012; Fig. 7, Table 5), and of precipitation, e.g. Copernicus data (Fig. 9), were analyzed.

For the Baltic Sea, intensive environmental monitoring started more than 100 years ago. Since 1898 an agreement between various Baltic Sea countries on simultaneous investigations on a regular basis at a few selected deep stations was signed and 1902 the International Council of the Exploration of the Sea (ICES) started its work. Examples from the national monitoring programs for water temperature (Figs. 18, 19, 20) and salinity (Figs. 21 and 22) are shown, illustrating climate variability and climate change of the Baltic Sea.



In addition, some institutes such as the Swedish Meteorological and Hydrological Institute (SMHI) provide environmental/climate indices, e.g. averaged sea level station data corrected for land uplift (Fig. 23) and hypoxic and anoxic areas (Fig. 25).

Since 1979 satellite data have become available, complementing traditional Earth observing systems and having the advantage of spatially high resolution (e.g. Karlsson and Devasthale, 2018).

Atmospheric reanalysis products, i.e. the combination of model data and observations (e.g. NCEP/NCAR, ERA40, ERA-Interim, ERA5, UERRA), were important for calculating water and energy budgets of the Baltic Sea region (BACC Author Team, 2008; BACC II Author Team, 2015). More recently, also ocean reanalysis products have been developed (e.g. Liu et al., 2017; Axell et al., 2019; Liu et al., 2019) and were, for instance, used for the evaluation of models (e.g. Placke et al., 2018).

Furthermore, various gridded datasets for North Sea SSTs exist and were compared (Fig. 33).

All data sets presented here are publicly online available. For further details on various datasets, the reader is referred to Rutgersson et al. (2021).

2.2.3 Future climate

For the BEARs, regionalizations of Global Climate Models (GCMs) or ESMs from CMIP3 and CMIP5 analyzed by (IPCC, 2014b) and (IPCC, 2019b) are assessed. The scenario simulations of CMIP5 are driven by greenhouse gas concentration scenarios, the Representative Concentration Pathways, RCP2.6, 4.5 and 8.5 (see Section 1.5).

Uncoupled atmospheric regional climate simulations for the 21st century from the Euro-CORDEX framework, calculated with several Regional Climate Models (RCMs) and global ESMs were analyzed by Christensen et al. (2021) and conclusions are summarized here.

Furthermore, coupled atmosphere – sea ice – ocean simulations for the Baltic Sea and North Sea regions with one, so-called Regional Climate System Model (RCSM, Dieterich et al., 2013; Bülow et al., 2014; Dieterich et al., 2019; Wang et al., 2015; Gröger et al., 2015; Gröger et al., 2019; Gröger et al., 2021b) driven by eight ESMs and three greenhouse gas concentration scenarios, i.e. RCP2.6, 4.5 and 8.5, were compared by (Christensen et al., 2021). In this study, we present figures of these consistent results from the coupled atmosphere-ice-ocean scenario simulations, e.g. for air temperature and precipitation (Fig. 26, Tables 7 and 8), and for sea surface temperature (Fig. 29, Table 9). The state-of-the-art of coupled modeling is discussed by Gröger et al. (2021a). For further details about the comparison between coupled and uncoupled scenario simulations, the reader is referred to Christensen et al. (2021).

Novel compared to the assessment by the BACC II Author Team (2015) are high-resolution projections of glacier masses including Scandinavian glaciers (Hock et al., 2019). In Figure 27 results are reproduced.



Oceanographic regional climate model projections for the Baltic Sea until 2100 driven by the atmospheric surface fields of the above mentioned RCM by Dieterich et al. (2019) have been developed and analyzed by Saraiva et al. (2019a; 2019b) and Meier et al. (2021a; 2021b). In Meier et al. (2021b), global sea level rise was also considered, a driver of the Baltic Sea climate variability that were previously neglected (cf. Hordoir et al., 2015; Arneborg, 2016; Meier et al., 2017). Here, we compare the latest scenario simulation results by Saraiva et al. (2019b) with previous projections by Meier et al. (2011a; 2011c) for, e.g. SST, sea surface and bottom salinities, sea level (Fig. 30), bottom oxygen concentration (Fig. 31), and Secchi depth (Fig. 32).

For further details about the latest oceanographic regional climate model projections for the Baltic Sea, the reader is referred to Meier et al. (2021a).

2.3 Uncertainty estimates

Uncertainties of future projections were estimated following the IPCC (2014a) guidance note for lead authors of the Fifth Assessment Report on consistent treatment of uncertainties (Mastrandrea et al., 2010). These uncertainty estimates are based upon a matrix of consensus and evidence reported in the literature. For the high confidence of a statement, high levels of both consensus and cases of evidence are required.

In this assessment, we applied a three-level confidence scale measuring low, medium and high confidence of identified climate changes (as defined in Section 2.1) of the selected 33 Earth system variables according to current knowledge (Table 10). We assessed the sign of a change but not its magnitude. Only detected or projected changes undoubtedly attributed to climate change were considered and synthesized in Figure 34. Changes likely not caused by increasing greenhouse gas concentrations or changing aerosol emissions were not considered. Other external drivers of climate variability are internal “random” variations of the climate system, land use, eutrophication, contaminants, litter, river regulations, fishery, aquaculture, underwater noise, traffic, spatial planning, etc. (see Reckermann et al., 2021).

Key messages of this assessment that are new compared to the previous assessment by the BACC II Author Team (2015) are specially marked (Section 7).

3 Current state of knowledge

3.1 Past climate change

3.1.1 Key messages from previous assessments

Climate variations may be triggered by changes in drivers external to the climate system or may be due to internal processes that reflect the non-linear, chaotic interactions between the different components of the climate system. The analysis of past climate variations is, therefore, useful for two purposes. One is to estimate the reaction of the climate to changes in the external forcing. The second is to better understand the mechanisms of internal climate variations. Since future climate change will include a mixture of both types of climate variations, the analysis of past climate variations is also necessary for better estimations of future climate change.



634 The past climate of the Baltic Sea region can be reconstructed from paleo-pollen and dendroclimatological records,
 635 with different time resolutions and degrees of accuracy. Paleo-pollen in lake sediments give information about the
 636 dominant plant species of a certain period. Combining the environmental ranges of those species in terms of annual
 637 maximum temperatures, minimum temperatures and total annual precipitation allows an approximate
 638 reconstruction of past climate conditions over the past millennia, with time resolutions of a few decades (e.g. Kühl
 639 et al., 2002). Dendroclimatological data of tree ring widths, wood density and sometimes also carbon and oxygen
 640 isotopic composition in tree-rings can be dated as exactly as at annual scales.

641
 642 As described by the BACC II Author Team (2015), the climate history of the Baltic Sea region during the
 643 Holocene, i.e. the last 12000 years, involved very large climate changes, much larger than those during the 20th
 644 century. These climate changes were caused by strong changes in external forcing factors, in particular the Earth's
 645 orbit. These changes first brought about a warming that terminated the Last Ice Age about 13000 years BP, then
 646 caused a period of very warm temperatures (~ 3°C above preindustrial levels) centered around 6000 years BP (the
 647 Holocene Thermal Maximum), followed by a slow temperature decline towards preindustrial levels. During this
 648 long period, other shorter-lived climate events, with durations of a few centuries, caused abrupt drops of
 649 temperature. These events, e.g. the Younger Dryas (12000 years BP) or the 8.2K event (8200 years BP) were
 650 possibly related to abrupt changes in the North Atlantic circulation, when sudden melting of portions of the
 651 remnants of the North American ice-sheet disturbed the circulation of the North Atlantic Ocean and disrupted the
 652 poleward heat transport.

653
 654 In general, annual precipitation is believed to have changed with the slow multicentennial-scale changes in
 655 temperature. Warmer periods, in particular the Mid-Holocene Optimum, tended to be wetter, although the regional
 656 heterogeneity may have been larger than for temperature.

657
 658 Following the end of the last glaciation, the coastlines of the Baltic Sea underwent changes due to the interplay
 659 between the rising global sea-level and the local rebound of the Earth's crust after the disappearance of the
 660 Fennoscandian ice sheet. The weight of this ice sheet depressed Fennoscandia by about 500 meters, and its slow,
 661 viscous rebound continues today, with a rate of about 10 mm year⁻¹ at the northern Baltic Sea coast. Due to this
 662 interplay, the Baltic Sea experienced periods of open or closed connections to the North Sea that governed the
 663 transport of salinity and heat and the nature of the Baltic Sea ecosystems (Groß et al., 2018).

664
 665 The climate evolution during more recent historical times – the past 1000 years (Section 3.1.3) - can be
 666 reconstructed with better accuracy and higher time resolution due to better dendrochronological data availability.
 667 These data show the imprint of the Medieval Warm Period (approx. 900-1350 AD), the Little Ice Age
 668 (approx. 1550-1850 AD) and the Contemporary Warm Period (1850-present) on the Baltic Sea region. These
 669 periods were likely caused by changes in the external forcing (volcanic eruptions and solar radiation), and during
 670 the Contemporary Warm Period also by the increase in anthropogenic greenhouse gases (Hegerl et al., 2003).

671
 672 In the Baltic Sea, this succession of warm-cold-warm temperatures was accompanied by changes in the deep water
 673 oxygen content, with low oxygen conditions in warmer periods (next section). The reasons for these oxygen



variations are still not fully understood, but may be relevant for the future, should future warming also cause lower oxygen concentrations.

3.1.2 New paleoclimate reconstructions

Since the publication of the BACC II report (BACC II Author Team, 2015), new reconstructions of the evolution of the European climate over the Holocene and over the past millennium have become available. Like previous reconstructions, the new ones are based on paleo-pollen data and now comprise summer and winter temperatures and summer and winter precipitation. They are available for 1000-year time segments (Mauri et al., 2015). The reconstructions of the late-spring-summer temperature evolution over the past millennium are based on dendroclimatological data, as the previous reconstructions, but they are now based on wood density measurements, which reflect the slow climate variations better than tree-ring width. These reconstructions are available for Western Europe from 755 AD onwards (Luterbacher et al., 2016). In this study, only the results for a regular geographical box approximately covering the Baltic Sea region are discussed.

In addition, new regional climate simulations since the publication of the BACC II report better demonstrate the connections between the Baltic Sea and North Atlantic climates on multidecadal timescales (Börgel et al., 2018; Börgel et al., 2020; Kniebusch et al., 2019a).

3.1.3 Holocene climate evolution

The picture of the Holocene climate evolution from the BACC II report (BACC II Author Team, 2015) is essentially confirmed, but the regional details are now clearer (Mauri et al., 2015). Between 7000 and 5000 years BP, the Baltic Sea region (especially the western Baltic Sea) experienced a period with summer temperatures about 2.5–3°C warmer than in the preindustrial reference period (before the 20th century). However, according to these reconstructions, the Eastern Baltic Sea region (Finland and the Baltic Republics) did not experience a Mid-Holocene Optimum in summer, when temperatures were similar to the preindustrial period. In contrast, winter temperatures showed a clear Mid-Holocene Optimum over the whole Baltic Sea region, lasting about 8000–4000 BP, with winter temperatures roughly 3°C warmer than during the preindustrial period. In the eastern Baltic Sea, winter temperatures were even slightly higher, especially between 6000–5000 BP. As a result, annual mean temperatures during the millennia of the Mid-Holocene optimum, were generally warmer than in the preindustrial period. This warming was limited to the winter in the eastern Baltic Sea, where the amplitude of the annual temperature cycle was clearly lower than in the preindustrial period.

The warm temperatures in the Baltic Sea region during the Mid-Holocene-Optimum, are not surprising, and basically agree with the previous review (BACC II Author Team, 2015). They also agree with evidence from regions further north, indicating that the Arctic Ocean in summer may have been ice-free during this period (Jakobsson et al., 2010). These findings do not contradict the anthropogenic effect on climate observed during recent decades. During the Mid-Holocene Optimum, the orbital configuration of the Earth was different and favored warmer temperatures at northern high latitudes, especially in summer, as explained later. For the analysis of climate impacts on ecosystems it is relevant that high latitudes were exposed to warm temperatures and reduced sea-ice cover just a few millennia ago. However, at that time temperature changed at a much slower pace – around 2–3°C over several millennia – compared to present and projected rates of about 2°C in just a few decades.



713

714 For precipitation, the new reconstructions give a regionally more nuanced view of climate evolution during the
 715 Holocene. The BACC II report (BACC II Author Team, 2015) indicated that warmer climates were generally more
 716 humid. The new reconstructions (Mauri et al., 2015) modulate this vision and constrain the wetter conditions to
 717 the eastern Baltic Sea region, both in summer and winter seasons, with a stronger signal in winter. Precipitation
 718 anomalies in the eastern Baltic Sea region were of the order of + 1-2 mm month⁻¹ relative to preindustrial climate.
 719 In the western Baltic Sea region, the Mid-Holocene Optimum tended to be slightly drier than the preindustrial
 720 reference period both in summer and winter, with precipitation deficits of the order of 1-2 mm month⁻¹.

721

722 The main external forcing that drove the millennial climate evolution over the Holocene Period is the changing
 723 orbital configuration of the Earth (the so-called Milanković cycles), as explained in the BACC-II report (BACC II
 724 Author Team, 2015), and especially the variation in the time of the year of the perihelion (when Earth is nearest
 725 to the sun). The perihelion is now at the beginning of January, but ~10000 BP it was in July. This changes the
 726 seasonal distribution of solar insolation and determines the rate of melting of winter snow and its possible survival
 727 into the next winter. The solar insolation at 60°N at the top of the atmosphere during the Holocene, derived from
 728 Laskar et al. (2004) is depicted in Figure 3. The shift of the perihelion from summer to winter diminishes summer
 729 insolation - and in principle summer temperature - and increases winter insolation during the past few millennia.
 730 The long-term evolution of temperatures would, however, not be a linear response to the long-term evolution of
 731 the seasonal insolation. For instance, the presence or absence of ice-sheets may influence the timing of the response
 732 to increasing insolation during the early Holocene, delaying the Holocene temperature maximum with respect to
 733 the annual insolation maximum. In wintertime, the insolation is rather weak, so that its effect may be overwhelmed
 734 by other factors, such as changes in the large-scale atmospheric and oceanic heat transports.

735

736 For the last IPCC report, the mid-Holocene climate was simulated with 14 global Earth System models within
 737 CMIP (Schmidt et al., 2011). These models were essentially the same as those used for future climate projections,
 738 although in some cases with a few simplifications required by limitations in computer power and by the long
 739 timescales involved. These models were driven by known external forcings, including the orbital forcing. The
 740 common evaluation of these simulations with reconstructions helps to interpret the reconstructions and sheds light
 741 on model limitations. An important aspect in this comparison was that the spatial resolution of global models was
 742 relatively coarse, about 2 x 2 degrees longitude x latitude, so that smaller details within the Baltic Sea region
 743 cannot be properly represented.

744

745 The simulations showed some agreements with the reconstructions, but also clear, not yet resolved disagreements
 746 (Mauri et al., 2014). In summer, all models showed temperatures 2-3°C warmer during the Mid-Holocene
 747 Optimum than in the preindustrial climate (Fig. 3). However, no model showed the gradient with clearer warming
 748 in the western Baltic Sea, seen in the reconstructions. For wintertime, the disagreement was much clearer. Whereas
 749 reconstructions show a clear warming over the whole region, the 14 models displayed widely varying patterns of
 750 temperature change. Only three models agreed with the reconstructions. For precipitation, the models disagreed
 751 with the west (wet) – east (dry) dipole shown by the reconstructions for summer and winter precipitation (Fig. 3).
 752 Not a single simulation showed this pattern of summer precipitation change, and in general the simulated
 753 precipitation deviations were much smaller than in the reconstructions. This disagreement regarding temperature



(especially in winter) and precipitation, known as the mid-Holocene conundrum, is not unique for the Baltic Sea region, but was also found for the Mediterranean (Mauri et al., 2014; Liu et al., 2014). Errors in the applied external (orbital) forcing can be ruled out, as this forcing can be accurately calculated during this period. The reasons for the disagreement are still unknown. They may involve the influence of chaotic internal climate variations (unlikely over such long time scales), model deficiencies, or reconstruction uncertainties.

3.1.4 The past millennium

For shorter periods closer to the present, like the past one or two millennia, the data available for reconstructing past climate are denser and more accurate. Abundant dendroclimatological information is available, dated to the exact year, in contrast to the uncertain decadal-scale dating of paleo-pollen data. Recently, temperature reconstructions for Western Europe, spatially resolved and approximately covering the last 1200 years, have become available (Luterbacher et al., 2016) and are presented here in some detail for the Baltic Sea region. These data are based on analysis of wood density in tree rings. Wood density is more sensitive to growing season temperature than tree-ring width. In addition, tree-ring width variations usually contain too weak multidecadal scale variations, even when the year-to-year variations in temperature may be well captured. This makes wood density a better proxy for temperature reconstructions at these latitudes.

Figure 4 shows the reconstructed growing-season temperature (spring-early summer) for the period 755-2000 AD, averaged over the Baltic Sea region, based on the European reconstructions by Luterbacher et al. (2016). The reconstructed temperature displays warmer conditions around 950 AD, confirmed also by the previous pollen-based reconstructions (Mauri et al., 2015), colder conditions between 1200 and 1850 AD, followed by the recent warming. This temperature evolution confirms that presented by the BACC II Author Team (2015). The relative temperature difference between the Medieval Warm Period and the Contemporary Warm Period (mid 20th century) agree within their respective uncertainties. According to these reconstructions, the Little Ice Age was on average about 0.8 °C colder than the 20th century.

There is no new analysis of the causes of this temperature evolution specific for the Baltic Sea region. For Europe as a whole, for which the reconstructions display a similar temporal pattern, the main identified forcings were volcanic activity - more intense during the Little Ice Age and weaker during the Medieval Warm Period - and solar activity, with roughly the reverse temporal signal (Luterbacher et al., 2016). With industrialization, greenhouse gases have become dominant.

The CMIP5 project also included simulations of the past millennium with Earth System Models, although with fewer models than for the mid-Holocene. These simulations have been compared with the temperature reconstructions for Europe, in general yielding agreement. However, for the Baltic Sea region, the simulated temperature changes tend to be smaller, especially for the transition between the Medieval Warm Period and the Little Ice Age, with a modelled temperature difference of only ~0.2°C (compare with Figure 4 by Luterbacher et al., 2016).

Climate fluctuations are driven not only by the external forcings but also by chaotic internal dynamics of the Earth system. Regional climate simulations indicate that North Atlantic temperature variability influences Baltic Sea



temperatures (Kniebusch et al., 2019a) and precipitation (Börgel et al., 2018). North Atlantic temperatures tend to fluctuate internally at multidecadal timescales, the AMO, and influences the atmospheric circulation of the Baltic Sea region. Further, the interaction between internal modes of climate variability has recently been identified as a key driver for the state of the Baltic Sea. Internal fluctuations in the North Atlantic are likely to influence the spatial position of the NAO, affecting the regional importance of this climate mode for the Baltic Sea (Börgel et al., 2020).

Climate simulations also indicate an impact of internally driven climate variability on the frequency of wind extremes. In the present climate, the wintertime wind regime in the Baltic Sea is linked to the NAO, but at the longer time scales of the preindustrial period, variations in wind extremes appear related neither to the mean wind conditions nor to the external climate forcings (Bierstedt et al., 2015). In the recent centuries, the main driver of trends of wind extremes over land appears to be land-use changes such as de- and reforestation (Bierstedt et al., 2015; Gröger et al., 2021a, and references therein).

An important question is how North Atlantic variations can influence the state of the Baltic Sea, especially its oxygen conditions, since freshwater input and water temperature (less strongly) affect the stratification of the water column, and therefore the exchange of oxygen between the surface and deeper layers. Temperature also modulates algal blooms and thus dissolved oxygen, when bacteria use oxygen to decompose dead algae. Analysis of sediment cores indicated that the mid-Holocene Optimum, the Roman Period (2000 BP), and the Medieval Warm Period were all periods of oxygen deficiency at the bottom of the Baltic Sea. Low oxygen conditions are also observed during the Contemporary Warm Period, unique in their extent on a thousand year perspective (Norbäck Ivarsson et al., 2019, and references therein). Hence, factors other than temperature, like nutrients input into the Baltic Sea, can also affect oxygen conditions, and thus the reasons for those hypoxic phases during the past millennia are not yet completely clear (Schimanke et al., 2012). It had been suggested that agricultural nutrient input was large enough to influence oxygen conditions already during the Medieval Warm Period, perhaps also modulated by changes in river runoff due to the described climate fluctuations (Zillén and Conley, 2010). However, a detailed analysis of new sediments records find little evidence of anthropogenic eutrophication before the industrial period (Norbäck Ivarsson et al., 2019; Ning et al., 2018; van Helmond et al., 2018). In view of the large temperature increases projected for this region in the next decades, further study of the influence of climate on oxygen conditions is warranted.

3.2 Present climate change

This section assesses our knowledge of Baltic Sea region climate variability during the past ~200 years, based on instrumental records, model based reconstructions and reanalyses. We focus on changes in means, extremes, trends and decadal to multidecadal climate variability.

3.2.1 Atmosphere

3.2.1.1 Large-scale atmospheric circulation

Long-term trends in NAO could not be detected (e.g. Deser et al., 2017; Marshall et al., 2020). For the period 1960-1990, a positive trend in NAO, with more zonal circulation, mild and wet winters and increased storminess in central and Northern Europe was found (Hurrell et al., 2003; Gillett et al., 2013; Ruosteenoja et al., 2020).



833 However, after the mid-1990s, there was a tendency towards more negative NAO indices, i.e. a more meridional
 834 circulation and more cold spells in winter (Fig. 5).

835

836 There is no consensus on how strongly the interannual NAO variability is forced externally (Stephenson et al.,
 837 2000; Feldstein, 2002; Rennert and Wallace, 2009). Several external forcing mechanisms have been proposed,
 838 most prominently SST (Rodwell et al., 1999; Marshall et al., 2001) and sea ice in the Arctic (Strong and
 839 Magnusdottir, 2011; Peings and Magnusdottir, 2016; Kim et al., 2014; Nakamura et al., 2015). Other authors
 840 (Screen et al., 2013; Sun et al., 2016; Boland et al., 2017) found no dependence on sea-ice extent. Furthermore,
 841 the impact of changes in the Arctic on midlatitude dynamics are still under debate (Dethloff et al., 2006; Francis
 842 and Vavrus, 2012; Barnes, 2013; Cattiaux and Cassou, 2013; Vihma, 2017).

843

844 A weakening of the zonal wind, eddy kinetic energy and amplitude of Rossby waves in summer (Coumou et al.,
 845 2015) as well as an increased waviness of the jet stream associated with Arctic warming (Francis and Vavrus,
 846 2015) in winter have been identified, which may be linked to an increase in blocking frequencies. Blackport and
 847 Screen (2020) argued that previously observed correlations between surface temperature gradients and the
 848 amplitude of Rossby waves have broken down in recent years. Therefore, previously observed correlations may
 849 have to be reinterpreted as internal variability. On the other hand, it has been shown that observed trends in
 850 blocking are sensitive to the choice of the blocking index, and that there is a huge natural variability that
 851 complicates the detection of forced trends (Woollings et al., 2018), compromising the robustness of observed
 852 changes in blocking.

853

854 With ongoing global warming, the Arctic will warm faster than the rest of the earth. This decrease of the poleward
 855 temperature gradient will tend to weaken the westerlies and increase the likelihood of blockings. On the other
 856 hand, maximum warming (compared to other tropospheric levels) will occur just below the tropical tropopause
 857 due to the enhanced release of latent heat, which tends to increase the poleward gradient, strengthen upper-level
 858 westerlies and affect the vertical stability, thus altering the vertical shear in midlatitudes. It is not clear which of
 859 these two factors will have the largest effect on the jet streams (Stendel et al., 2021).

860

861 The atmospheric circulation over Europe naturally varies significantly on decadal time scales (Dong et al., 2017;
 862 Ravestein et al., 2018). Proposed drivers for these circulation changes include polar and tropical amplification,
 863 stratospheric dynamics and the AMOC (Haarsma et al., 2015; Shepherd et al., 2018; Zappa and Shepherd, 2017).
 864 The attribution of drivers is more straightforward for local changes, in particular for the soil-moisture feedback,
 865 for which an enhancement of heat waves due to a lack of soil moisture has been demonstrated (Seneviratne et al.,
 866 2013; Teuling, 2018; Whan et al., 2015). Räisänen (2019) found only a weak effect of circulation changes on the
 867 observed annual mean temperature trends in Finland, but circulation changes have considerably modified the
 868 trends in individual months. In particular, circulation changes explain the lack of observed warming in June, the
 869 very modest warming in October in southern Finland, and about a half of the very large warming in December.

870

871 As part of its natural variability, the North Atlantic warmed from the late 1970s to 2014 (Fig. 6). Recently, the
 872 AMO began transitioning to a negative phase again (Frajka-Williams et al., 2017). Paleoclimate reconstructions
 873 and model simulations suggest that the AMO might change its dominant frequency over time (Knudsen et al.,



2011; Wang et al., 2017). The impact of the AMO on climate is, however, independent of its frequency (Börgel et al., 2018; Börgel et al., 2020). Its influence on regional climate has been analyzed in several studies (Enfield et al., 2001; Knight et al., 2006; Sutton and Hodson, 2005; Ting et al., 2011; Casanueva et al., 2014; Ruprich-Robert et al., 2017; Peings and Magnusdottir, 2014), some dealing with the Baltic Sea (Börgel et al., 2018; Börgel et al., 2020; Kniebusch et al., 2019a). Kniebusch et al. (2019a) suggested that the influence of the AMO on temperature during 1980–2008 might have been at least as strong as that induced by humans (IPCC, 2014b).

3.2.1.2 Air temperature

A significant increase in surface air temperature in the Baltic Sea region during the last century has been shown previously (e.g. BACC Author Team, 2008; Rutgersson et al., 2014; BACC II Author Team, 2015). The temperature increase was not monotonous but accompanied by large multidecadal variations that divided the 20th century into three main phases: (1) warming from the beginning of the century until the 1930s; (2) slight cooling until 1960s; and (3) a distinct warming during the last decades of the time series that has continued also during 2014–2020 (Figs. 7 and 8 and Table 4).

Linear trends of the annual mean temperature anomalies during 1878–2020 were $0.10\text{ }^{\circ}\text{C decade}^{-1}$ north of 60°N as well as south of 60°N in the Baltic Sea region. This is larger than the global mean temperature trend and slightly larger compared to the earlier BACC reports. Over the Baltic Sea surface air temperature trends were smaller than over land. During 1856–2005, surface air temperature over the Baltic Sea increased by 0.06 and $0.08\text{ }^{\circ}\text{C decade}^{-1}$ in the central Baltic Sea and in the Bothnian Bay, respectively (Kniebusch et al., 2019a).

There is a large variability in annual and seasonal mean temperatures, in particularly during winter, but the warming is seen for all seasons (being largest during spring in the northern part of the region).

Both daily minimum and daily maximum temperatures have increased. A decrease in the daily temperature range (DTR) have been observed in many regions of the world, but there is no clear signal for the entire Baltic Sea region (see for example, Jaagus et al., 2014, for DTR analysis of the Baltic countries).

These changes have also resulted in seasonality changes: the growing season has lengthened by about 5 days decade^{-1} in the period 1965–2016 (Cornes et al., 2019). From this follows that the cold season has become shorter.

Extreme air temperatures can be high or low, but extended periods of extreme temperatures (spells or waves) are often the most influential. Averaged over land areas, warm spell duration increased during recent decades (Rutgersson et al., 2021). For some regions, the average annual days defined as warm spells increased from 6–8 to 14 during recent decades. Along with more frequent and longer warm spells came decreases in the frequency, duration and severity of cold spells, based both on observations (Easterling et al., 2016) and model results. The length of the frost season and the annual number of frost days also decreased (Sillmann et al., 2013).

3.2.1.3 Solar radiation and cloudiness

Multidecadal variations of solar radiation at the Earth's surface, called “dimming” and “brightening”, have been observed in Europe and other parts of the world, particularly in the northern hemisphere (Wild et al., 2005; Wild, 2012; Wild et al., 2017).



913

914 One of the world's longest time series of global radiation, i.e. incoming solar radiation at the Earth's surface, is
 915 from Stockholm, where measurements started in 1922. Recently, a first attempt to homogenize this time series was
 916 made by Josefsson (2019). No significant trend was found over the whole time series, but there were large
 917 variations over one to three decades. Other long time series of global radiation in Northern Europe are from
 918 Potsdam, Germany, (Wild et al., 2021), and Tõravere, Estonia, (Russak, 2009). All three time series show a
 919 minimum in global radiation around the mid-1980s. Then a clear increase or "brightening" of about 5-8% followed,
 920 until at least 2005. Before the 1980s minimum there was a period of "dimming" at all stations but with differences
 921 in the details. In Potsdam, there was rather stable dimming all the time from the late 1940s. In Tõravere, there was
 922 a maximum dimming around mid-1960s, while Stockholm recorded high values both around 1950 and 1970, with
 923 a minimum dimming in between. The data also suggest an early brightening in Stockholm time series, but this is
 924 still uncertain, especially before 1950.

925

926 Current twentieth century reanalyses models provide results for surface solar radiation. However, most of them
 927 fail to capture multidecadal surface radiation variability in central and southern Europe (Wohland et al., 2020).
 928 The model CERA20C, which shows best results for central and southern Europe still gives questionable results
 929 over Scandinavia, showing a weak increase instead of a decrease in surface solar radiation during the presumed
 930 dimming period before 1980.

931

932 Satellite data allowing analyses of cloudiness and solar radiation at the Earth's surface are available since the early
 933 1980s. For Europe, important work has been done within the EUMETSAT Satellite Application Facility on
 934 Climate Monitoring (CM SAF). Several satellite data records have been validated and used in climate studies (e.g.
 935 Urraca et al., 2017; Pfeifroth et al., 2018). At the highest latitudes of the Baltic Sea region there are however larger
 936 uncertainties (Riihelä et al., 2015) or often no data at all, due to low standing Sun and slant viewing geometry from
 937 the satellites.

938

939 The satellite data only cover the latest brightening period observed at ground-based stations in Europe. While the
 940 geographical patterns of global average cloud conditions agree well among several satellite cloud-data sets, there
 941 are clear differences in the distribution and size of trends (Karlsson and Devasthale, 2018). However, there seems
 942 to be consensus on a decreasing trend in total cloud fraction of about 1-2% per decade over the Baltic Sea region
 943 during 1984-2009.

944

945 Recent CM SAF satellite products on solar irradiance at the Earth's surface, the SARAH-2 and CLARA-A2
 946 datasets, both agree well with station data according to Pfeifroth et al. (2018). In many cases this holds both for
 947 climatological averages and for trend detection. The average trend for the period 1983-2015 is about $+3 \text{ W m}^{-2}$
 948 decade^{-1} both at the stations closest to the Baltic Sea and in the SARAH-2 dataset. The three long-term stations
 949 mentioned above are all used as reference stations for the satellite data validation. For example, the on-going
 950 monitoring at stations spread over all Sweden show an average increase of about 8% (corresponding to $+4 \text{ W m}^{-2}$
 951 decade^{-1}) from 1983 until 2005-2006 (SMHI, 2021). In later years the solar radiation leveled off, but *inter alia* the
 952 extremely sunny 2018 in Northern Europe contributed to keeping the trend increasing over time.

953



954 The multidecadal variations in the solar radiation at the Earth's surface were most probably caused by a
 955 combination of changes in cloudiness and in anthropogenic aerosols. Which of the two drivers is the largest
 956 contribution is still an open question, and might differ among regions. Aerosol concentrations over Northern
 957 Europe decreased during the brightening period from the mid-80s onwards (Ruckstuhl et al., 2008; Russak, 2009;
 958 Markowicz and Uscka-Kowalkowska, 2015; Glantz et al., 2019). Russak (2009) considered changes in cloudiness
 959 caused by variations in atmospheric circulation to be the most important factor in Estonia, but aerosol changes also
 960 played a role. In an early study of the modern radiation measurements in Sweden the strong increase in solar
 961 radiation 1983-1997 was also accompanied by a clear decrease in total cloud cover, especially during the half-year
 962 of summer (Persson, 1999). The satellite datasets SARAH-2 and CLARA-A2 where both derived using an aerosol
 963 climatology as input. This underlines the important role of changes in cloudiness for surface solar radiation. Stjern
 964 et al. (2009) also stressed the importance of the contribution of clouds and the atmospheric circulation for dimming
 965 and brightening periods in Northern Europe.

966
 967 Other studies, e.g. Ruckstuhl et al. (2008) and Wild et al. (2021), concluded that aerosol effects under clear skies
 968 is the main contributor to the multidecadal variations of solar radiation in central Europe. Aerosol-induced
 969 multidecadal variations in surface solar radiation could be expected also over oceans (Wild, 2016), but long-term
 970 measurements are lacking. The interaction between aerosols and clouds, the indirect aerosol effects, needs also be
 971 better understood and quantified.

972 **3.2.1.4 Precipitation**

973 During the twentieth century in the Baltic Sea region, changes in precipitation were spatially more variable than
 974 for temperature (BACC II Author Team, 2015). Irregularly distributed precipitation measurement stations make it
 975 difficult to determine statistically significant trends and regime shifts. Sweden shows an overall wetting trend since
 976 the 1900s, in particular since the mid 20th century (Chen et al., 2020). In Finland, the overall increase detected for
 977 1961-2010 is neither regionally consistent nor always statistically significant (Aalto et al., 2016). The same holds
 978 for the Baltic States (Jaagus et al., 2018). In the south of the Baltic Sea region, changes were small and not
 979 significant. Nevertheless, precipitation averaged over the Baltic Sea catchment area has increased since 1950 due
 980 to an increase in winter (Fig. 9).

981
 982 The number of heavy precipitation days is largest in summer. Compared to southern Europe, precipitation extremes
 983 in the Baltic Sea region are not as intense, with daily amounts typically ranging from 8 to 20 mm (Cardell et al.,
 984 2020). Extreme precipitation intensity increased during the period 1960-2018. An index for the maximum annual
 985 five consecutive days of precipitation (Rx5d) shows significant increases of up to 5 mm per decade over the eastern
 986 part of the Baltic Sea catchment (EEA, 2019b). The change is more pronounced in winter than in summer.

987 **3.2.1.5 Wind**

988 In situ observations allow direct analysis of winds, in particular over sea (e.g. Woodruff et al., 2011). However, in
 989 situ measurements, especially over land, are often locally influenced, and inhomogeneities make the
 990 straightforward use of such data difficult, even for recent decades. Therefore, many studies use reanalyses rather
 991 than direct wind observations. But analysis of storm-track activity for longer periods using reanalysis data suffers
 992 necessarily from uncertainties associated with changing data assimilation and observations before and after the



introduction of satellites, resulting in large variations of storm-track changes across assessments (Wang et al., 2016; Chang and Yau, 2016).

Owing to inherent inhomogeneities and the large climate variability in the Baltic Sea region, it is unclear whether there is a general trend in wind speed in the recent climate. Results regarding changes or trends in the wind climate are strongly dependent on period and region considered (Feser et al., 2015). Due to the strong link to large-scale atmospheric variability over the North Atlantic, conclusions about changes over the Baltic Sea region are perhaps best made in a wider spatial context, considering *inter alia* the NAO.

Recent trend estimates for the total number of cyclones over the Northern Hemisphere extratropics during 1979-2010 revealed a large spread across the reanalysis products, strong seasonal differences, as well as decadal-scale variability (Tilina et al., 2013; Wang et al., 2016; Chang et al., 2016; Matthews et al., 2016; Chang et al., 2012). Common to all reanalysis datasets is a weak upward trend in the number of moderately deep and shallow cyclones (7 to 11% per decade for both winter and summer), but a decrease in the number of deep cyclones, in particular for the period 1989-2010. Chang et al. (2016) reported a minor reduction in cyclone activity in the Northern Hemisphere summer due to a decrease in baroclinic instability as a consequence of Arctic temperatures rising faster than at low latitudes. Chang et al. (2012) also noticed that state-of-the art models (CMIP5) generally underestimate this trend. In the Northern Hemisphere winter, recent studies reported a decrease in storm track activity related to Arctic warming (Ceppi and Hartmann, 2015; Shaw et al., 2016; Wills et al., 2019; Stendel et al., 2021).

Despite large decadal variations, there is still a positive trend in the number of deep cyclones (< 980 hPa) over the last six decades, which is consistent with results based on the NCEP reanalysis since 1958 over the northern North Atlantic Ocean (Lehmann et al., 2011). Using an analogue-based field reconstruction of daily pressure fields over central to Northern Europe (Schenk and Zorita, 2012), the increase in deep lows over the region might be unprecedented since 1850 (Schenk, 2015). However, for limited areas the conclusions were rather uncertain.

The effect of differential temperature trends on storm tracks has been recently addressed, both in terms of upper tropospheric tropical warming (Zappa and Shepherd, 2017) and lower tropospheric Arctic amplification (Wang et al., 2017), including the direct role of Arctic sea-ice loss (Zappa et al., 2018), and a possible interaction of these factors (Shaw et al., 2016). The remote and local SST influence has been further examined by Ciaso et al. (2016), who confirmed the sensitivity of the storm tracks to the SST trends generated by the models and suggested that the primary greenhouse gas influence on storm track changes was indirect, acting through its influence on SSTs. The importance of the stratospheric polar vortex for storm track changes has received more attention (Zappa and Shepherd, 2017). In an aqua planet simulation, Sinclair et al. (2020) found a decrease in the number of extratropical cyclones and a poleward and downstream displacement due to an increase in diabatic heating.

3.2.1.6 Air pollution, air quality and atmospheric nutrient deposition

Air pollution continues to significantly impair the health of the European population, particularly in urban areas. Brandt et al. (2013) estimated the total number of premature deaths due to air pollution in Europe in the year 2000 to be ~680 000 year⁻¹. Although this number was predicted to decrease to approximately 450 000 by 2020, it is



1033 still a matter of grave concern. Particulate matter concentrations were reported to be the primary reason for adverse
 1034 health effects. Estimates indicated that PM_{2.5} concentrations in 2016 were responsible for ~412 000 premature
 1035 deaths in Europe, due to long-term exposure (EEA, 2019a).

1036

1037 The state of air pollution is often expressed as air quality, when human health is in focus. The ambient air quality
 1038 in the Baltic Sea region is dominated by anthropogenic emissions, and natural emissions play only a minor role.
 1039 These emissions show an overall decreasing trend in recent years (EEA, 2019a), as reflected in the ambient
 1040 concentrations reported in the EMEP status report (EMEP, 2018). To quantify air quality concentrations of certain
 1041 gases and particulate matter are used as measures. The general conclusions in the field of air quality reported by
 1042 the BACC II Author Team (2015) still hold today, i.e. that land-based emissions and concentrations of major
 1043 constituents continue to decrease due to emission control measures, with the possible exception of certain
 1044 emissions from the shipping sector. Sulphur emissions from shipping have continued to decrease strongly in the
 1045 Baltic Sea from 2015, due to much lower limit values for the sulphur content of ship fuel in the emission control
 1046 areas. A noticeable decrease in nitrogen emissions due to the newly (2021) implemented nitrogen emission control
 1047 area (NECA) is expected in the next decade.

1048

1049 In Europe, the pollutants most harmful to human health are particulate matter (PM), nitrogen dioxide (NO₂) and
 1050 ground-level ozone (O₃). About 14% of the EU-28 urban population was exposed to O₃ concentrations above the
 1051 EU target value threshold (EEA, 2019a). When compared to other European countries, air pollution was relatively
 1052 low in Scandinavia, exception for a few urban traffic hotspots, with annual mean NO₂ concentrations elevated to
 1053 near the limit value (EEA, 2019b). In this comparison, Northern Germany is located in the lower mid-field, while
 1054 northern Poland is among the more polluted countries, especially with PM. Biomonitoring samples analyzed for
 1055 toxic metals by Schröder et al. (2016) tended to show lowest concentrations in Northern Europe.

1056

1057 Contributions to air pollution and pollutant deposition in coastal areas by shipping can be substantial. Major
 1058 pollutants from shipping are SO₂, NO_x and PM (including black carbon). The BACC II assessment estimated
 1059 emissions from shipping in the Baltic Sea region. Several studies have recently been published, including Jonson
 1060 et al. (2015), Claremar et al. (2017), and Karl et al. (2019b), which use chemistry transport models to predicted
 1061 ambient concentrations from known emissions. They show, as expected, that the highest air pollution
 1062 concentrations due to ship exhaust are found near major shipping lanes and harbors, but also that considerable
 1063 concentrations of NO₂ and PM reach populated land areas. This effect is pronounced in the south-western Baltic
 1064 Sea area (Quante et al., 2021). Exact numbers from such modelling should still be interpreted with care, as shown
 1065 by Karl et al. (2019a), who compared output from three state-of-the-art chemistry transport models for the Baltic
 1066 Sea area.

1067

1068 The most important recent change in shipping emissions in the North and Baltic seas are due to the 2015
 1069 strengthening of the fuel sulphur content limit for the Sulphur Emission Control Areas (SECAs), by lowering the
 1070 maximum allowed sulphur content from 1 to 0.1%. Model calculations indicate large reductions in sulphur
 1071 deposition in countries bordering these two sea areas after the implementation of the lowered sulphur limit (Gauss
 1072 et al., 2017). Barregard et al. (2019) estimated the contribution of Baltic Sea shipping emissions to PM_{2.5} before
 1073 2014 and after 2016 the new SECA regulation of marine fuel sulphur was implemented. These authors also



1074 estimated human exposure to PM_{2.5} from shipping and its health effects in the countries around the Baltic Sea.
 1075 They concluded that PM_{2.5} emissions from Baltic Sea shipping, and resulting health impacts decreased
 1076 substantially after the 2015 SECA regulation. Population exposure studies estimating the influence of shipping
 1077 emissions for selected Baltic Sea harbor cities were published for Rostock, Riga and Gdańsk–Gdynia by Ramacher
 1078 et al. (2019) and for Gothenburg by Tang et al. (2020). Ramacher et al. (2019) found that shipping emissions
 1079 strongly influence NO₂ exposure in the port areas (50–80 %), while the average influence in home, work and other
 1080 environments is lower (3–14 %) but still with strong influence close to the ports. It should, however, be noted that
 1081 reduction of sulphur emissions to the atmosphere by the use of new cleaning techniques (e.g. open loop scrubbers)
 1082 can increase the risk of acidification and marine pollution (Turner et al., 2017; 2018).

1083

1084 Johansson et al. (2020) published a first comprehensive assessment of emissions from leisure boats in the Baltic
 1085 Sea. While the modeled NO_x and PM_{2.5} emissions from leisure boats are clearly lower than those from commercial
 1086 shipping, these first estimates suggest that carbon monoxide (CO) emissions from leisure boats equal 70 % of the
 1087 registered shipping emissions and non-methane volatile organic carbon (NMVOC) emissions equal 160 %. It
 1088 should be noted that most of the leisure boat emissions occur in summer, and often occur near areas for nature
 1089 conservation and tourism. Most of these emissions can be attributed to Swedish, Finnish and Danish leisure boats,
 1090 but the leisure boat fleet has the potential for large future increases also in Russia, Estonia, Latvia, Lithuania and
 1091 Poland.

1092

1093 Air pollution leads to environmental degradation by affecting natural ecosystems and biodiversity. Ground-level
 1094 ozone (O₃) can damage crops, forests and other vegetation, impairing growth and reducing biodiversity. According
 1095 to a recent study by Proietti et al. (2021), assessed trends of the O₃ mean concentration in Northern Europe were
 1096 not statistically significant for the time period from 2000 to 2014. The annual mean ozone concentration is reported
 1097 to be slightly below 35 ppb, as compared to 43 to 45 ppb in the Mediterranean Region, for which a significant
 1098 decreasing trend is found. The exposure index AOT40 (sum of the hourly exceedances above 40 ppb, for daylight
 1099 hours during the growing season) significantly declined in all European regions except for Northern Europe, for
 1100 which a positive but not significant trend is seen. On the nation level among the six European countries showing
 1101 a positive trend were Denmark, Germany, Sweden (Proietti et al., 2021). A clear difference in trends between rural
 1102 sites and other station typologies is found for Europe for the period 2000 to 2017. I.e. for traffic sites a substantial
 1103 increase of annual mean O₃ concentration was observed, in contrast to rural stations, for which a slight decrease
 1104 was found (Colette and Rouïl, 2020). Regarding the monitored population exposed to ozone all countries in Europe
 1105 show a decrease from 2000 to 2014 (NDGT60 > 25 days per year; Fleming et al., 2018).

1106

1107 Harmful exposure and impacts of air pollutants on ecosystems are assessed using the concept of critical loads
 1108 (CLs; Nilsson and Grennfelt, 1988). The CL is the amount of pollutants that an ecosystem can tolerate without
 1109 risking unacceptable damage. The most harmful air pollutants in terms of damage to ecosystems in addition to O₃
 1110 are ammonia (NH₃) and nitrogen oxides (NO_x). It is estimated that about 62% of the European ecosystem area is
 1111 still exposed to high levels of NO_x, leading to exceedances of CLs for eutrophication in all countries in 2016
 1112 (EEA, 2019a). Hotspots of exceedances of CLs for acidification in 2016 were the Netherlands and its borders with
 1113 Germany and Belgium, southern Germany and also Czechia. However, most of Europe including the Baltic Sea
 1114 region did not exceed the CLs for acidification (EEA, 2019a).



1115

1116 Since the 1980s, the total nitrogen deposition on the Baltic Sea has decreased substantially, due to an overall
 1117 reduction of European emissions, but emission and deposition reductions have stalled since the mid-2000s (Colette
 1118 et al., 2015; Gauss et al., 2021). Atmospheric phosphorus deposition remains highly uncertain in amount and trends
 1119 (HELCOM, 2015; Kanakidou et al., 2018; Ruoho-Airola et al., 2012).

1120

1121 Air quality and climate interact in several ways. On the one hand, air pollutants can affect climate both directly
 1122 and indirectly by changing the radiative balance of the atmosphere. On the other hand, climate change alters
 1123 meteorological conditions, which may affect concentrations of air pollutants via several pathways, since air quality
 1124 is strongly dependent on weather (Jacob and Winner, 2009). The effects of important meteorological and climate
 1125 variables on surface O₃ and PM were discussed in a comprehensive review by Doherty et al. (2017). The
 1126 connection between high temperatures and increased ground-level ozone concentrations is well established.
 1127 Increases in temperature related to climate change (i.e. during heat waves) are expected to lead to higher ozone
 1128 concentrations in certain regions with the required precursor concentrations. Other important meteorological
 1129 factors influencing air pollution concentrations are a possible change in the number of midlatitude cyclones and
 1130 in the number of occurrences and duration of stagnant weather conditions (Jacob and Winner, 2009).

1131 3.2.2 Land

1132 3.2.2.1 River discharge

1133 The total river discharge to the Baltic Sea is approximately 14,000 m³ s⁻¹ (Bergström and Carlsson, 1994). This is
 1134 substantially more than the direct net precipitation (precipitation minus evaporation) on the Baltic Sea itself, which
 1135 has been estimated at 1,000-2,000 m³ s⁻¹ (Meier and Kauker, 2003; Meier and Döschner, 2002; Meier et al., 2019d),
 1136 see also the discussion by Leppäranta and Myrberg, (2009). In other words, most of the fresh water entering the
 1137 Baltic Sea comes from the terrestrial part of the catchment. Therefore, the fresh water input to the Baltic Sea cannot
 1138 be described entirely with only climatic parameters. Non-climatic drivers of runoff include river regulation by
 1139 dams and reservoirs, land-use changes in the catchment, and water uptake for irrigation. Although dams are known
 1140 to have altered the seasonality of discharge (e.g. McClelland et al., 2004; Adam et al., 2007; Adam and
 1141 Lettenmaier, 2008), they do not seem to be responsible for annual discharge changes. In the long-term, net
 1142 precipitation over the catchment area and river runoff are strongly correlated (Meier and Kauker, 2003).

1143

1144 For the period 1850-2008, the total river discharge from the Baltic Sea catchment area, reconstructed from
 1145 observations (Bergström and Carlsson, 1994; Cyberski et al., 2000; Hansson et al., 2011; Mikulski, 1986) and
 1146 hydrological model results (Graham, 1999), showed no statistically significant trend but a pronounced
 1147 multidecadal variability, with a period of about 30 years (Meier et al., 2019d). Furthermore, summed river flow
 1148 observations in the period 1900-2018 (Lindström, 2019) and a historical reconstruction of the annual river
 1149 discharge for the past 500 years showed no statistically significant trend either (Hansson et al., 2011). However,
 1150 river runoff from northern Sweden, a part of the catchment area of the Bothnian Bay, significantly increased since
 1151 the 1980s compared to 1911-2018 (Lindström, 2019).

1152

1153 There are indeed substantial regional and decadal variations in the river flow. Stahl et al. (2010) studied near-
 1154 natural rivers of Europe over the period 1942-2004 and found a clear overall pattern of positive trends in annual



streamflow in the northern areas. (Kniebusch et al., 2019b) also identified a statistically significant positive trend in the river discharge to the Bothnian Bay for 1921-2004. In Estonian rivers, regime shifts in annual specific runoff corresponded to the alternation of wet and dry periods (Jaagus et al., 2017). A dry period started in 1963/1964, followed by a wet period from 1978, with the latest dry period commencing at the beginning of the 21st century.

For the period 1920-2005, positive trends in stream flow at stations of a pan-Nordic dataset dominate annual mean, winter and spring figures whereas summer trends are statistically not significant (Wilson et al., 2010). A clear signal of earlier snow-melt floods and a tendency towards more severe summer droughts in southern and eastern Norway were found.

The observed temperature increases have affected stream flow in the northern Baltic Sea region for 1920-2002 in a manner corresponding well to the projected consequences of a continued rise in global temperature (Hisdal et al., 2010). Regarding precipitation, however, the regional impacts of both the observed and projected changes on stream flow are still unclear.

In the northern Baltic Sea region, all the way south to the Gulf of Finland, runoff is strongly linked to the climate indices air temperature, wind and rotational circulation components. In the southern region, runoff is associated more with the strength and torque of the cyclonic or anticyclonic pressure systems (Hansson et al., 2011).

In the Baltic states (Lithuania, Latvia and Estonia), changes in streamflow over the 20th century showed a redistribution of runoff over the year, with a significant increase in winter and a tendency for decreasing spring floods (Reihan et al., 2007; Sarauskiene et al., 2015; Jaagus et al., 2017). A similar winter trend was found also for the reconstructed river discharge to the entire Baltic Sea since the 1970s (Meier and Kauker, 2003).

For the period 1911-2010, a trend of observed annual maximum daily flows in Sweden could not be detected (Arheimer and Lindström, 2015). However, in particular the annual minimum daily flows in northern Sweden considerably increased in the period 1911-2018 (Lindström, 2019). Analyzing a pan-European database, Blöschl et al. (2017) showed that river floods over the past five decades occurred earlier in spring due to (1) an earlier spring snow melt in northeastern Europe, (2) delayed winter storms associated with polar warming around the North Sea, and (3) earlier soil moisture maxima in Western Europe.

3.2.2.2 Land nutrient inputs

The Baltic Sea catchment area of 1.7 million km², which is more than four times larger than the sea surface area (cf. Fig. 1), is populated by over 84 million inhabitants. Stretching between 49° - 69°N and 10° - 38°E, the catchment exhibits significant gradients in both natural (precipitation, river discharge, temperature, etc.) and anthropogenic (population density and occupation, agricultural and industrial development, etc.) environmental factors. These factors change both in time (phenological changes, long-term trends and lags due to land cover processes) and space (north-south gradients in climate and land use, east-west gradients in socio-economic features and climate) thus determining heterogeneity and variation of land nutrient inputs that drive long-term eutrophication of the Baltic Sea (Savchuk, 2018, and references therein; Kuliński et al., 2021).



Estimates of nutrient inputs had been attempted since the 1980s (e.g. Larsson et al., 1985; Stålnacke et al., 1999) and are now being compiled within a permanent process of the HELCOM Pollution Load Compilation (PLC, e.g. HELCOM, 2019). However, these data officially reported to HELCOM by the participating riparian states have been and still are suffering from gaps and inconsistencies. Therefore, the “best available estimates” have been reconstructed in attempts to both fill in such gaps and correct possible sources of inconsistencies (Savchuk et al., 2012; the present study based on Svendsen and Gustafsson, 2020). For long-term studies of the Baltic Sea ecosystem, a historical reconstruction of nutrient inputs since 1850 is available (Gustafsson et al., 2012).

According to HELCOM (2018a; Savchuk, 2018) and updated estimates (HELCOM, 2018c), substantial reductions of land nutrient inputs, comprising riverine inputs and direct point sources at the coast, have been achieved since the 1980s (Fig. 11, Table 5). Since there are no statistically significant trends in annual river discharge (Section 3.2.2.1), these reductions are attributed to socio-economic development, including expansion of the wastewater treatment and reduction of atmospheric nitrogen deposition (Gauss et al., 2021) over the entire Baltic Sea drainage basin, and not to climate related effects (HELCOM, 2018a; Svendsen and Gustafsson, 2020). As an example, the coastal point sources of TN and TP decreased three- and ten-fold, respectively, comparing to the 1990s (Savchuk et al., 2012) and today contribute to the Baltic Sea less nutrients than they did in 1900 (Savchuk et al., 2008; Kuliński et al., 2021).

Agriculture is the main source of anthropogenic diffuse nutrient inputs, which comprise 47% of the riverine nitrogen and 36% of the riverine phosphorus inputs (HELCOM, 2018c). In turn, mineral fertilizer dominates the anthropogenic nutrient inputs to the Baltic Sea catchment, in particular in its intensely farmed southern part (Hong et al., 2017). However, during 2000-2010 only about 17% of the net anthropogenic nitrogen and only about 4.7% of the net anthropogenic phosphorus input to the catchment are currently exported with rivers to the sea (Hong et al., 2017). While denitrification might have removed part of the nitrogen applied in agriculture, the remaining phosphorus has accumulated in the drainage basin. A global budget estimated that agriculture has increased the soil storage of phosphorus in the drainage basin by 50 Mt during 1900-2010 (Bouwman et al., 2013) and a regional approach calculated an increase by 40 Mt during 1900-2013 (McCrackin et al., 2018). However, McCrackin et al. (2018) estimated that about 60% of these phosphorus inputs were retained in a stable pool and did not contribute noticeably to the riverine export. About 40% accumulated in a mobile pool with a residence time of 27 years. McCrackin et al. (2018) suggested that leakage from this mobile legacy pool is, though slowly declining, the dominant source of present riverine phosphorus inputs.

3.2.3 Terrestrial biosphere

Previous assessments

The comprehensive review of climate-related changes in terrestrial ecosystems in the first BACC report (BACC Author Team, 2008), Smith et al. (2008) concluded that climate change during the preceding 30-50 years had already caused measurable changes in terrestrial ecosystems in the Baltic Sea region, e.g. an advancement of spring phenological phases in some plants, upslope displacement of the alpine tree-line and increased land-surface greenness in response to improved growth conditions and a richer CO₂ supply. But as nearly all ecosystems in the region were managed to some extent, the climate impacts might be alleviated or intensified by human interventions, e.g. by choosing favorable tree species in forestry. The observed trends were expected to continue



for at least several decades, assuming that continued future increases in the atmospheric CO₂ concentration will cause continued warming.

In the second BACC report (BACC II Author Team, 2015), climate effects on terrestrial ecosystems were less in focus, with a section related to forests and natural vegetation in the chapter on environmental impacts on coastal ecosystems, birds and forests (Niemelä et al., 2015) and as part of the chapter on socioeconomic impacts on forestry and agriculture (Krug et al., 2015). On the other hand, the second BACC report also considered anthropogenic land-cover changes as a driver of regional climate change (Gaillard et al., 2015). Niemelä et al. (2015) concluded that the observed positive effects of climate change on forest growth would continue, in particular for boreal forest stands that benefitted more than temperate forest stands. The species composition of natural vegetation in the Baltic Sea region was expected to undergo changes, with a predominantly northward shift of the hemiboreal and temperate mixed forests. Terrestrial carbon storage was likely to increase in the region, but land-use change could play an important modifying role, affecting this storage both positively and negatively. Krug et al. (2015) concluded that there were regional differences in how the vulnerability and productivity of forestry systems were affected by climate change, with the southern and eastern parts of the Baltic Sea region likely to experience reduced production and the northern and western parts increased production. Gaillard et al. (2015) found no indication that deforestation in the Baltic Sea region since 1850 could have been a major cause of the observed climate warming.

Acknowledging the importance of the land component in the climate system, the IPCC recently published its special report entitled ‘Climate Change and Land’ on climate change, desertification land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems (IPCC, 2019b). This is because land, including its water bodies, provides the basis for human livelihoods and well-being through primary productivity, the supply of food, freshwater, and multiple other ecosystem services.

Biophysical and biogeochemical interactions

The land surface and its terrestrial ecosystems in the Baltic Sea region interact with the atmosphere and are, thus, coupled to the local and regional climate. These interactions determine the exchanges of heat, water and momentum between the land surface and the atmosphere via biophysical processes, the exchange of greenhouse gases, e.g. CO₂, CH₄ and N₂O, and emissions of black carbon, aerosol precursors, e.g. biogenic volatile compounds, or organic carbon aerosols via biogeochemical processes, altering the atmospheric composition.

The nature of the biophysical and biogeochemical interactions between the land surface and the atmosphere and their effects on climate are studied by comparing the effects of forests with that of open land, e.g. grassland, pastures or cropland (e.g. Bonan, 2016). Details on the nature of these feedbacks are given in Gröger et al. (2021b).

On average, across the globe, forests absorb atmospheric CO₂ and, thus, reduce net radiation and have a cooling effect on climate. In equilibrium, forest ecosystems, are expected to be carbon neutral, with carbon loss through phenological turnover, mortality and decomposition over large areas on average matching plant productivity. CO₂ fertilization is considered a strong driver for the terrestrial carbon sink, but demographic recovery following past land use, e.g. afforestation and replanting of harvested forest stands, likely provides an equally important explanation for net carbon uptake by forests in industrialized regions of North America, Europe and Asia (Pugh et



al., 2019). Deforestation, on the other hand, can lead to a release of carbon and to an increase in net radiation and, thus, has a warming effect on climate (Friedlingstein et al., 2020, and references therein). In contrast to the biophysical effects described above and some of the other biogeochemical interactions, the biogeochemical interactions associated with the carbon cycle have global impacts and operate at very long time scales.

Anthropogenic land-use and land cover changes

Using remote sensing data, Jin et al. (2019) investigated recent trends in springtime plant phenology in the Baltic Sea region and the sensitivities of phenological trends to temperature and precipitation, in spring, winter and summer. Considering the entire region and combining all vegetation types, the authors found an advancement of the growing season by 0.30 day year⁻¹ over the period 2000–2016. The advancement was particularly strong for evergreen needle-leaved forests (0.47 day year⁻¹) and weaker for cropland and grassland (0.14 day year⁻¹). Evergreen needle-leaved forests, together with deciduous broadleaf forests, dominate the northern part of the Baltic Sea region, while the southern part is mainly grassland and cropland. Jin et al. (2019) found that the most important driver of the advancement of the growing season is spring mean temperature, with an advancement rate of 2.47 day (°C)⁻¹ of spring warming, considering the entire area and all vegetation types. Spring drying could further increase the advancement rate by 0.18 day cm⁻¹. While the sensitivity of the start of the growing season to climate conditions in spring is comparable for the entire Baltic Sea region, the sensitivity to climate conditions in the summer and winter seasons differs between the northern and southern parts of the region. In both seasons an increase in the mean temperature was found to advance the growing season in the southern part of the Baltic Sea region but to delay the start of the growing season in the northern part, in contrast to the spring warming. These sensitivities were markedly stronger for summer than for winter. These trends in plant phenology in spring result in changes of the land cover early in the year, thus affecting climate through biophysical and biogeochemical interactions with the atmosphere.

Observations reveal a local impact of changes in forest cover on near-surface temperatures, due to biophysical effects that depend on the geographical latitude, roughly separating the boreal regions and the temperate zone of the Northern Hemisphere. When investigating the effects of small-scale clearings at sites in the Americas and Asia, Zhang et al. (2014a) found on both continents that annual mean temperatures cooled over open land north of about 35°N and warmed south of this latitude. Changes in forest cover have, however, different effects on daily minimum and daily maximum temperatures. Zhang et al. (2014a) found that the warming effect over open land south of 35°N was related to an increase in daily maximum temperatures, with little change in daily minimum temperatures, while the cooling effect to the north was due to a decrease in daily minimum temperatures. Lee et al. (2011) showed consistent results for North America, where the cooling effect of 0.85±0.44°C over non-forested areas north of 45°N was due to a decrease in daily minimum temperatures, associated with the reduced roughness length. At night, open land cools more than forests, regardless of geographical latitude. This is confirmed by Alkama and Cescatti (2016), who analyzed the impacts of recent losses in forest cover on near-surface and land-surface temperatures in the boreal zone. For both, the authors found cooling trends in daily minimum and warming trends in daily maximum temperatures in response to deforestation and opposite tendencies after afforestation. These effects were somewhat stronger for land-surface temperatures than air temperatures.



1310 Regional Climate Models (RCMs) have been used to investigate the biophysical effects of changes in forest cover
 1311 on climate in Europe. Strandberg and Kjellström (2019), for instance, used simulations with the Rossby Centre
 1312 Atmosphere (RCA) RCM to assess the climate effect of maximal afforestation or deforestation in Europe, focusing
 1313 on seasonal mean temperatures and precipitation, as well as daily temperature extremes. Maximum afforestation
 1314 and deforestation were inferred from a simulation with the LPJ-GUESS dynamical vegetation model, providing a
 1315 map for potential natural forest cover for Europe in equilibrium with present-day climate (Gröger et al., 2021a).
 1316 To effect maximum afforestation, present-day land cover classes, which represent considerable agricultural
 1317 activity in Europe (particularly in western, central and southern Europe), were replaced by the potential natural
 1318 forest cover. In the case of deforestation, on the other hand, the potential natural forest cover was converted to
 1319 grassland in the model.

1320 The simulations indicated that afforestation in Europe generally increased evapotranspiration, which, in turn, led
 1321 to colder near-surface temperatures. In western, central and southern Europe, the cooling in winter due to
 1322 afforestation was between 0.5 and 2.5°C. The cooling effect was somewhat stronger in summer, exceeding 2.5°C
 1323 in large parts of western and southeastern Europe. Deforestation had the opposite effect, warmer near-surface
 1324 temperatures due to decreased evapotranspiration, typically in the range between 0.5 and 2°C in western and
 1325 central Europe and reaching up to 3°C in southeastern Europe. In regions with low evapotranspiration, however,
 1326 changes in the surface albedo were relatively more important for temperatures. During summer, warming by
 1327 deforestation affected the entire Baltic Sea region (in the range between 0.5 and 1.5°C), while the cooling
 1328 associated with afforestation only affected its southern part. Over parts of Scandinavia, afforestation actually
 1329 resulted in a slight warming of about 0.5°C. In winter, the cooling effect of afforestation was only evident over the
 1330 southern part of the Baltic Sea region, while deforestation had no effect on winter temperatures.

1331 Strandberg and Kjellström (2019) found relatively strong biophysical effects of afforestation or deforestation in
 1332 Europe on daily maximum temperatures in summer. Deforestation markedly increased daily maximum
 1333 temperatures over the entire Baltic Sea region (typically between 2 and 6°C), while afforestation lowered daily
 1334 maximum temperatures in the southern part of the region by about 2 to 6°C and slightly increased over parts of
 1335 Scandinavia. In contrast to its cooling effect on mean winter temperatures, afforestation lead to a warming of the
 1336 daily minimum temperatures in the southern part of the Baltic Sea region in the range between 2 and 6°C.

1337 In a similar study with the Regional Model (REMO) RCM, Gálos et al. (2013) investigated the biophysical effects
 1338 of afforestation in Europe on climate, and also compared these effects to the climatic changes expected from future
 1339 global warming. Potential afforestation was implemented by specifying deciduous forest cover at all vegetated
 1340 areas that were not covered by forests at the end of the 20th century, mainly in western, Central and Eastern Europe.
 1341 Given the strong historical deforestation in Central Europe, there is potential for rather extensive afforestation in
 1342 the southern part of the Baltic Sea region, but lesser potential in the northern part, i.e. Scandinavia and Finland.
 1343 The results indicated a cooling effect of the re-established forests in boreal summer exceeding 0.3°C, mainly
 1344 related to increased evapotranspiration from the trees, in combination with intensified fluxes of latent heat due to
 1345 stronger vertical mixing (see above). The stronger latent heat fluxes also enhanced precipitation by more than 10%
 1346 in some regions. These effects of potential afforestation counteracted the projected future changes in climate, i.e.
 1347 somewhat reduced the magnitude of the pronounced future warming and markedly reduced the future drying in
 1348 the boreal summer. In some cases, such as in northern Germany, the enhanced precipitation was found to



completely offset the drying effect of future warming. More recent results (Meier et al., 2021c) have used rain-gauge data to estimate precipitation changes induced by land cover change. Meier et al. (2021c) created a statistical model to show that reforestation of agricultural land can increase precipitation locally, especially in winter, and were able to separate the effects on both local and downwind precipitation regionally and seasonally. They also found that climate change induced summer precipitation reductions could be offset by reforestation, with a particularly strong effect in southwest Europe, though their analyses also indicate small precipitation increases in the Baltic Sea region, relative to a baseline scenario with no land cover change, consistent with the results of Gálos et al. (2013).

In a more regionalized study, Gao et al. (2014) applied the REMO RCM to investigate the biophysical effects of peatland forestation in Finland before (1920) and after drainage (2000s). In Finland, as in other northern European countries, vast areas of naturally tree-less or sparsely tree-covered peatland were drained for timber production in the second half of the 20th century. The total peatland area of Finland was estimated to be 9.7 million ha in the 1950s, but at the beginning of the 21st century the area of peatland drained for forestry was estimated to 5.5 million ha. The authors found that the peatland forestation caused warming in spring, i.e. during the snow-melt season, and slight cooling in the growing season (May through October). The spring warming was mainly caused by decreased surface albedo and the cooling in the growing season by increased evapotranspiration.

3.2.4 Cryosphere

3.2.4.1 Snow

In the Baltic Sea region, snow cover is an important feature that greatly affects cold season weather conditions. It is characterized by very high interannual and spatial variability. Snow cover is a sensitive indicator of climate change, and its variations are closely related to air temperature in many regions. General climate warming is expected to reduce snow cover. Several thaw periods now interrupt snow cover, making it less stable. Total winter snowfall in Northern Europe is projected to decrease, but is still expected to increase in mid-winter in the very coldest regions (Räsänen, 2016).

Previous climate change assessments demonstrated a number of snow-cover trends in recent decades in the Baltic Sea region (BACC Author Team, 2008; BACC II Author Team, 2015). A decrease in snow cover was observed in the south, while an increase in snow storage and duration of snow cover was detected in the north-east, and in the Scandinavian mountains. The spring snow melt has become earlier in most of the region. As a result, the spring maximum river discharge has become smaller and earlier, in many regions shifting from April to March.

Recent investigations confirm these results. Snow cover in the Northern Hemisphere has decreased since mid-20th century (IPCC, 2014a), as also shown by satellite measurements (Estilow et al., 2015). The largest decline in the extent of snow cover has been observed in March-April and also in summer. Using the satellite-based NOAA-CDR data for the period 1970–2019, it was shown that the annual snow cover fraction has reduced over most areas of the Northern Hemisphere by up to 2% decade⁻¹ (Zhu et al., 2021). Thereby, the annual snow cover area has reduced by $2 \times 10^5 \text{ km}^2 \text{ decade}^{-1}$.



1387 In 1980–2008, snow-cover duration in Northern Europe decreased by about 3–7 days per decade, and the trend
 1388 was significant at many stations (Peng et al., 2013). Most of the reduction happened in spring, with the end-date
 1389 of snow cover five days earlier per decade, on average (Peng et al., 2013). Snow-cover variability over Europe is
 1390 closely related to temperature fluctuations, which, in turn, are determined by large-scale atmospheric circulation
 1391 during the cold season (Ye and Lau, 2017). A recent study of European snow-depth data in 1951–2017
 1392 demonstrated an accelerated decrease after the 1980s (Fontrodona Bach et al., 2018), with an average decline,
 1393 excluding the coldest climates, of 12.2% per decade for mean snow depth and 11.4% per decade for its maximum.
 1394 A decreasing trend in snow density was detected in the eastern Baltic Sea region in 1966–2008 (Zhong et al.,
 1395 2014).

1396

1397 In Poland, rather large changes in snow cover parameters were found for 1952–2013 (Szwed et al., 2017). The
 1398 duration of snow cover decreased in almost the whole country but this change is mostly not statistically significant.
 1399 The total reduction in snow cover duration was 1–3 weeks over the 62 years, but the mean and maximum snow
 1400 depths did not change. The start date of snow cover has not changed, but the end date moved slightly earlier (Szwed
 1401 et al., 2017). A recent study found a statistically significant decreasing trends in snow cover duration as well as of
 1402 in snow depth, based on 40 Polish stations in 1967–2020 (Tomczyk et al., 2021). The trend values for the number
 1403 of days with snow cover were from –3.5 to –4.9 days per decade.

1404

1405 The snow-cover regime at 57 stations in the eastern Baltic Sea region (Lithuania, Latvia and Estonia) in 1961–
 1406 2015 was analyzed by Rimkus et al. (2018). The mean decrease in snow-cover duration was 3.3 days per decade,
 1407 and was statistically significant at 35% of the measuring sites, mostly in the southern part of the region. There
 1408 were no trends in maximum snow depth. An earlier study for Lithuania found similar results (Rimkus et al., 2014).

1409

1410 A detailed study of snow cover data at 22 stations in Estonia during the period 1950/51–2015/16 revealed
 1411 remarkable decreasing trends (Viru and Jaagus, 2020). Snow-cover duration decreased significantly at 16 stations,
 1412 and the mean decrease was 4 days per decade. Start dates for permanent snow cover had a non-significant tendency
 1413 to occur later. Permanent snow cover had a statistically significant trend to end earlier at almost all stations. There
 1414 were no overall trends in maximum snow depth in Estonia in 1951–2016 (Viru and Jaagus, 2020).

1415

1416 Significant decreases in snow depth parameters were found in Finland in recent decades (Aalto et al., 2016;
 1417 Luomaranta et al., 2019). Regional differences were substantial. In 1961–2014, the largest decrease in snow depth
 1418 occurred in the southern, western and central parts of Finland in late winter and early spring. In northern Finland,
 1419 a decrease in snow depth was most evident in spring, with no change in the winter months, even though the amount
 1420 of solid precipitation was found to increase in December–February (Luomaranta et al., 2019). Winter mean snow
 1421 depth (Jylhä et al., 2014) as well as the annual maximum snow depth (Lehtonen, 2015) has decreased significantly
 1422 at many stations in southern Finland. At the same time, the annual maximum snow depth has not changed in
 1423 Finnish Lapland (Lépy and Pasanen, 2017; Merkouriadi et al., 2017).

1424

1425 For the century 1909–2008, a general decrease was detected in many snow-cover parameters at three stations in
 1426 different parts of Finland (Irannezhad et al., 2016). A sharp decline in annual peak snow-water equivalent was
 1427 detected since 1959. The period of permanent snow cover shortened by 21–32 days per century.



1428

1429 A decline in snow cover parameters was found also in Norway for 1961–2010 (Dyrrdal et al., 2012; Rizzi et al.,
 1430 2017).

1431 3.2.4.2 Glaciers

1432 A recent basic inventory of Scandinavian glaciers is available through the Randolph Glacier Inventory (RGI
 1433 Consortium, 2017; Pfeffer et al., 2014), a collection of digital outlines of the world's glaciers prepared to meet the
 1434 needs of the Fifth IPCC Assessment Report (IPCC, 2014b; Vaughan et al., 2014). It has since been updated in
 1435 support of the IPCC Special Report on the Ocean and Cryosphere in a Changing Climate (IPCC, 2019a; Hock et
 1436 al., 2019). Of the 3417 Scandinavian glaciers reported in the Randolph Glacier Inventory (v6.0), 365, with a
 1437 combined area of c. 360 km², lie within the Baltic Sea Drainage Basin as defined by (Vogt et al., 2007; Vogt et
 1438 al., 2008; all in the Scandinavian mountains). The combined glacier volume estimate (following Farinotti et al.,
 1439 2019) for the Scandinavian glaciers reported in Hock et al. (Hock et al., 2019) is 0.7 ± 0.2 mm global sea level
 1440 equivalent or 254 ± 72 Gt. The rate of mass loss for all Scandinavian glaciers (not only in the Baltic Sea Drainage
 1441 Basin) for the period 2006–2015 is 2 ± 1 Gt year⁻¹ corresponding to a negligible 0.01 ± 0.00 mm year⁻¹ of global
 1442 sea level rise equivalent, yet with potential importance for local streamflow (Hock et al., 2019; Zemp et al., 2019).
 1443 With a 90–100% likelihood, atmospheric warming is the primary driver of glacier mass loss (Hock et al., 2019).

1444

1445 Most of the 365 glaciers in the Baltic Sea Drainage Basin are in Sweden (c. 72% by number, c. 75% by area), with
 1446 the remaining ones in Norway. Both Sweden and Norway do nationally coordinated glacier monitoring, with the
 1447 most recent results summarized in the Global Glacier Change Bulletin (GGCB) No. 3 (World Glacier Monitoring
 1448 Service, 2020; Zemp et al., 2020) as, among others, glacier mass balance changes. A glacier mass balance year
 1449 usually covers the period from September 1st to August 31st in the subsequent year. The GGCB includes four
 1450 Swedish glaciers, two of which (Rabots glaciär and Storglaciären in the Kebnekaise Massif, which has the world's
 1451 longest continuous mass balance record, starting in 1945/46) are so-called reference glaciers, meaning that their
 1452 dynamics are not dominated by non-climatically driven dynamics such as calving or surging, and that more than
 1453 30 years of ongoing measurements are available (Fig. 12). In Table 6, mass balances of the Swedish GGCB glaciers
 1454 following World Glacier Monitoring Service (2020) are summarized. None of the Norwegian GGCB glaciers are
 1455 in the Baltic Sea drainage basin.

1456

1457 These with time increasingly negative mass balances coincide with globally increasing air temperatures, with the
 1458 latest six years, 2015–2020, the warmest since instrumental recording began (World Meteorological Organization,
 1459 2020). Regional and local deviations in mass loss from that expected from long-term global warming is, however,
 1460 expected. The slightly positive mass balances for Mårmaglaciär, Storglaciären and Riwojekna in 2016/2017 is the
 1461 result of a cold summer, explained by the glacier mass balance years starting on September 1, and that the summer
 1462 months (June, July, August) included in the 2016/2017 balance are therefore June, July, and August 2017 – a
 1463 period during which average air temperature at Tarfala Research Station was measured to 5.8°C, compared to
 1464 6.4°C and 7.4°C in the previous and subsequent mass balance periods (Swedish Infrastructure for Ecosystem
 1465 Science, 2020; World Glacier Monitoring Service, 2020).

1466



The ice summit of Kebnekaise Sydtopp (South Peak) lost its status as Sweden's highest in September 2019 and 2020 when due to melting of its ice-covered summit its elevation dropped below that of the rocky, non-ice-covered Kebnekaise Nordtopp (North Peak; Stockholm University - Department of Physical Geography, 2017, 2019, 2020).

3.2.4.3 Permafrost

The drainage basin of the Bothnian Bay is characterized by low mean annual air temperatures and includes boreal forest and mountain ecosystems as well as large peatland areas. In this region, permafrost — ground frozen for at least two consecutive years — exists in high alpine environments and in peatlands. While almost all Baltic Sea Drainage Basin permafrost is found in the Bothnian Bay catchment, some isolated occurrences of permafrost are found in the upper reaches of Alpine rivers further south (mainly Umeälven headwaters). Permafrost is a thermal state of ground, rock, soil or sediment, and occurs in regions with low mean annual air temperatures (MAAT). Temperature is the strongest control on permafrost, but thin winter snow depth also favors permafrost aggradation and stability. In alpine permafrost, insolation is an important factor. In the upper Torne River catchment, incoming shortwave summer radiation causes a difference in the altitude of alpine permafrost (permafrost $p > 0.8$) from 850 m a.s.l. on shaded slopes to 1100 m a.s.l. on south-facing slopes (Ridefelt et al., 2008). In lowland areas, permafrost is more likely to form and persist in peatlands, where the low thermal conductivity of peat insulates the ground from warm summer air (Seppälä, 1986). Mire complexes with palsas, elevated peat mounds with an ice-rich permafrost core, are the most common form of lowland permafrost in the Baltic Sea drainage basin (Luoto et al., 2004). Palsa mires in the Baltic Sea basin are predominantly found in regions with MAAT $< -3^{\circ}\text{C}$ and low mean annual precipitation (often < 450 mm), based on the 1961-1990 climate period (Fronzek et al., 2006).

Earlier maps of northern hemisphere permafrost extent showed relatively extensive Fennoscandian permafrost, especially in Alpine regions (Brown et al., 1997). Recent advances in permafrost modeling reveal a more nuanced picture, where permafrost is spatially patchy, persisting at high elevations and in lowland regions with low precipitation and large expanses of peat plateaus (Obu et al., 2019; Gislén et al., 2017). Consistent with observation of permafrost warming and thawing (Biskaborn et al., 2019), models project substantial permafrost losses in recent decades. High resolution modelling (1 km pixels) driven by remotely sensed land surface temperature showed that c. 6,200 km² of permafrost in 1997, was reduced to 4,800 km² in 2018 (Obu et al., 2020). Most of this loss is modeled alpine permafrost (here defined as > 700 m a.s.l.) decreasing from 4,700 to 3,700 km², while lowland permafrost has decreased from 1,500 km² to 1,100 km² (Fig. 13).

3.2.4.4 Sea ice

Introduction

Sea ice is an essential indicator of climate change and variability in the Baltic Sea region. Not only does existence of sea ice indicate the general severity of the winter due to its close correlation with winter air temperature, but in addition parameters such as annual maximum sea-ice extent of the Baltic Sea (MIB), duration of ice season and maximum thickness of level ice have been monitored regularly in the Baltic Sea since the late 19th century.

The BACC II Author Team (2015) concluded that all sea-ice observations demonstrated large inter-annual variations, but with a long-term, statistically significant trend to milder ice conditions that is projected to continue



(Fig. 14). In this section, we review recent ice-climate research in the Baltic Sea and provide updated figures on sea ice trends and projections that largely confirm previous conclusions.

1508

An important indicator of advancing climate change in the Baltic Sea region is that two of the latest five ice winters have been extremely mild (2015-2020). In winter 2015, the Bothnian Bay was never fully covered by ice (with a MIB of 51 000 km²), the first such extreme winter observed with certainty. The winter of 2020 was even milder, with a MIB of only 37 000 km², the lowest value in a time series that began in 1720.

1513

Sea-ice conditions in the Baltic Sea

On recent average, the northern sea areas of the Baltic Sea are ice-covered every year, from December to May. During the mildest winters, only the Bothnian Bay (in a few years even only partially) and coastal zones of other basins are ice-covered. In the past, the entire Baltic Sea was ice-covered only in the most severe winters, e.g. 1940, 1942 and 1947 (Vihma and Haapala, 2009).

1519

In the fast ice regions near the coast, sea ice grows by thermodynamic processes only. Maximum sea ice thickness in the fast ice regions typically amounts to 40 – 70 cm in the Bothnian Bay and 10 – 40 cm in the Bothnian Sea, Gulf of Finland and Gulf of Riga. Ice thickness is regularly monitored at tens of fast ice sites.

1523

Observations of sea-ice thickness in drift ice are much more limited. A recent study combined all airborne electromagnetic ice thickness measurement in the Bothnian Bay and derived the first estimate of basin-scale ice thickness distribution in the Baltic Sea (Fig. 15; Ronkainen et al., 2018). An important finding of that study was that mean ice thickness in drift ice regions is greater than the thickness of fast ice, and also greater than the ice thickness indicated on the ice charts. As expected, the data showed large inter-annual variability, but temporal and spatial coverage was not sufficient for conclusion on changes in drift ice thickness.

1530

Individual ice ridges caused by compression and shearing of ice drift can be 30 meters thick. The largest gradients in ice motion are found in the coastal zone (Leppäranta et al., 2012), where mean ice thickness over several km² can be 1-3 meters (Ronkainen et al., 2018).

1534

In some circumstances, sea ice can accumulate towards the coast and cause spectacular on-shore ridges or ride-up of several hundred meters from shore to land, causing damage to build structures (Leppäranta, 2013). Such events have been observed in exposed coastal regions where the stability of fast ice can be overcome by combinations of storms, currents and water level (Leppäranta, 2013). In the Bothnian Bay, such events can occur regardless of the severity of ice seasons. In the southern Baltic Sea, on-shore ice has been common during severe winters. During the last ten years, such events were observed in 2010, 2011, 2012 and 2019 (Girjatowicz and Łabuz, 2020).

1541

Observed changes

Long-term changes of the MIB (Seinä and Palosuo, 1996; Niskanen et al., 2009) are shown in Figure 14. The trend of the MIB during the last 100 years (1921-2020) is -6,400 km² per decade. This is almost twice the trend reported by the BACC II Author Team (2015), based on the period 1910-2011. Since 1987 no severe ice winters and since



1546 2012 only average, mild or extremely mild winters have been observed. The latter sea-ice conditions explain the
 1547 accelerated trend after 2011.

1548

1549 The recent 30-year period (1991-2020) is definitely the mildest since 1720 (Uotila et al., 2015). The probability
 1550 distribution of the MIB has shifted towards zero, with severe winters very rare (Fig. 14). The 30-year mean MIB
 1551 is now $139 \cdot 10^3 \text{ km}^2$ and the winter 2021 with a MIB of $127\,000 \text{ km}^2$ on 15 February 2021 (Jouni Vainio, FMI,
 1552 personal communication) was close to this mean. During the second mildest 30-year period (1909-1938), the
 1553 average MIB was $184 \cdot 10^3 \text{ km}^2$ and during the last 100 years it was $182 \cdot 10^3 \text{ km}^2$. The shape of the MIB probability
 1554 distribution has changed, also indicating a change in sea-ice extremes. According to the ice season classification
 1555 (Seinä and Palosuo, 1996), the recent 30-year period includes only one severe or extremely severe ice winter and
 1556 13 mild or extremely mild ice winters.

1557

1558 Present ice conditions differ from the past to the extent that Rjazin and Pärn (2020) even suggested defining this
 1559 change as a regime shift. They analyzed changes in sea ice extent and air temperature in the Baltic Sea in 1982 –
 1560 2016, using a method of splitting the time series in two and concluded that a regime shift towards milder ice
 1561 conditions occurred in 2006-2007.

1562

1563 Other studies complement these and BACC II conclusions. Kiani et al. (2018) examined the influence of
 1564 atmospheric changes on ice roads between Oulu and Hailuoto. They used air temperature data to calculate freezing
 1565 and thawing degree days and found that freezing degree days decreased and thawing degree days increased
 1566 significantly during 1974 – 2009. As a consequence, the ice road season started later and ended earlier.

1567

1568 Merkouriadi and Leppäranta (2014) analyzed ice thickness and freezing and breakup dates collected at Tvärminne
 1569 Zoological Station, at the entrance to the Gulf of Finland. They found a decrease of almost 30 days in the ice-
 1570 covered period and a reduction of 8 cm in maximum annual ice thickness in the last 40 years. Laakso et al. (2018)
 1571 used observations from the Utö Atmospheric and Marine Research station during 1914-2016 and concluded that
 1572 the length of the ice season has decreased from 10-70 days before 1988 to 0-35 days after 1988 in the northern
 1573 Baltic Sea proper. Figures 16 and 17 show level-ice thickness at Kemi and Loviisa and the length of the ice season
 1574 at Kemi, Loviisa and Utö. All graphs show statistically significant decreasing trends, except level-ice thickness at
 1575 station Kemi, which was probably be influenced by snow cover or changes in measurement location.

1576 3.2.4.5 Lake ice

1577 The recent change in ice phenology is probably the single most important climatically induced alteration in lake
 1578 environments within the Baltic Sea catchment. New literature demonstrates almost unanimously significant
 1579 changes towards earlier ice break-up, later freeze-up, and shorter duration of ice cover across the Baltic Sea
 1580 catchment, apart from the coldest climate regime in Lapland. The available centennial data indicate that the ice-
 1581 cover duration has decreased by several days per century, whereas the intensified warming in recent decades has
 1582 produced a similar change per decade (Efremova et al., 2013; Filazzola et al., 2020; Kļaviņš et al., 2016; Knoll et
 1583 al., 2019; Korhonen, 2019; Lopez et al., 2019; Nöges and Nöges, 2014; O'Reilly et al., 2015; Ptak et al., 2020;
 1584 Sharma et al., 2016; Sharma et al., 2020; Wrzesiński et al., 2015). Some lakes have, however, responded only
 1585 weakly to the warming trend, such a lake Peipsi in Estonia, probably due to increasing snowfall. In individual



1586 years, a positive wintertime NAO seems to be an important factor causing a short ice-cover duration. Among the
 1587 main properties that affect the ice cover of individual lakes are size, depth, and shoreline complexity.

1588 3.2.5 Ocean and marine sediments

1589 3.2.5.1 Water temperature

1590 The main driver of annual mean water temperature variations and long-term changes is air temperature (Meier et
 1591 al., 2019d; 2019c; Kniebusch et al., 2019a). Baltic Sea water temperature has risen fastest at the sea surface (Meier
 1592 et al., 2021a). With time the heat spreads downward through different processes, such as lateral inflows, vertical
 1593 down-welling and diffusion, and eventually the whole water column warms up, with smallest trends in the cold
 1594 intermediate layer between the thermo- and halocline (Meier et al., 2021a).

1595
 1596 Since the 1980s, marginal seas around the globe have warmed faster than the global ocean (Belkin, 2009), and the
 1597 Baltic Sea has warmed the most (Belkin, 2009). Climate change and decadal variability led to an annual mean,
 1598 area averaged increase in Baltic Sea SST of $+0.59^{\circ}\text{C decade}^{-1}$ for 1990-2018 (Siegel and Gerth, 2019) and of
 1599 $+0.5^{\circ}\text{C decade}^{-1}$ for 1982-2013 (Stramska and Białogrodzka, 2015). Both figures were derived from satellite data.
 1600 In accordance with earlier investigations (BACC Author Team, 2008; BACC II Author Team, 2015), SST
 1601 variability in winter can be linked to the NAO (Stramska and Białogrodzka, 2015). However, the spatial maps of
 1602 SST trends by Stramska and Białogrodzka (2015) differ from those by Lehmann et al. (2011), perhaps because of
 1603 the differing horizontal resolution of the satellite data products. Linear trends for the Baltic Sea during 1982-2012
 1604 of $0.41^{\circ}\text{C decade}^{-1}$ are slightly larger than $0.37^{\circ}\text{C decade}^{-1}$ for the North Sea (Høyer and Karagali, 2016).

1605
 1606 Using monitoring data, Liblik and Lips (2019) found that the upper layer has warmed by $0.3\text{--}0.6^{\circ}\text{C decade}^{-1}$ and
 1607 the sub-halocline deep layer by $0.4\text{--}0.6^{\circ}\text{C decade}^{-1}$ in most of the Baltic Sea during 1982-2016. The total warming
 1608 in the whole Baltic Sea was 1.07°C over 35 years, approximately twice that of the upper 100 m in the Atlantic
 1609 Ocean.

1610
 1611 During 1856–2005, the reconstructed Baltic Sea average, annual mean SST increased by 0.03 and $0.06^{\circ}\text{C decade}^{-1}$
 1612 ¹ in the northeastern and southwestern areas, respectively (Kniebusch et al., 2019a). The largest SST increase
 1613 trends were found in the summer season in the northern Baltic Sea (Bothnian Bay). Bottom water temperature
 1614 trends were smaller than SST trends, with the largest increase in the Bornholm Basin. Independent monitoring
 1615 data support the results of the long-term reconstruction (Figs. 18 and 19), see also Meier et al. (2019c; 2019d). The
 1616 largest SST warming occurred in summer (May to September), while trends in winter were smaller (Kniebusch et
 1617 al., 2019a; Liblik and Lips, 2019).

1618
 1619 During the more recent period of 1978–2007, the annual mean SST trend was tenfold higher, with a mean of 0.4°C
 1620 decade^{-1} (Kniebusch et al., 2019a). Trends increased more in the northeastern areas than in the southwestern, and
 1621 exceeded the contemporary trends in air temperature. See also MARNET station data at Darss Sill and Arkona
 1622 Deep in the southwestern Baltic Sea (Fig. 20).

1623
 1624 The seasonal ice cover clearly plays an important role in the Baltic Sea by decoupling the ocean and the atmosphere
 1625 in winter and spring. Hence, the large trends in air temperature in winter were not reflected by the SST trends



because the air temperature was still below the freezing point. During the melting period, the ice-albedo feedback led to larger trends in SST than during the ice-covered period, because of a prolonged warming period of sea water.

It has been suggested that the accelerated warming in 1982-2006 might partly be explained by a dominance of the positive phase of the AMO (Kniebusch et al., 2019a). Historical eutrophication re-distributed the heat in the ocean by warming the surface layer more than the underlying layers, in particular during spring and summer, because the increased water turbidity caused an enhanced absorption of sunlight at the sea surface. However, modeling studies suggest that the historical eutrophication had no impact on SST trends (Löptien and Meier, 2011).

The summer of 2018 was the warmest on instrumental record in Europe, and also the warmest summer in the past 30 years in the southern half of the Baltic Sea (Naumann et al., 2019), with surface-water temperatures 4-5°C above the 1990-2018 long-term mean. This heat wave was also observed in the bottom temperatures (Humborg et al., 2019). However, systematic studies on changes in heat waves are not available.

3.2.5.2 Salinity and saltwater inflows

During the last decade, many new insights have been gained about the salt balance of the Baltic Sea and the dynamics of inflow and mixing processes. The Major Baltic Inflow (MBI) in December 2014, in particular, triggered new investigations and is by far the most intensively observed and modeled inflow event. Pathways and timing of the inflowing water were tracked by observations (Mohrholz et al., 2015), and numerical modelling (Gräwe et al., 2015) could reproduce the salt mass and volume of the inflow, calculated from observations. The inflow was found to be barotropic (pressure) controlled in the Danish straits but dominated by baroclinic (density stratification) processes on the further pathway into the Baltic proper. At the Bornholm Gat and the Słupsk Furrow the water exchange showed the clear two-layer flow pattern of an estuarine circulation. The inflow-related studies were underpinned by theoretical work based on the famous Knudsen Relation (Knudsen, 1900) for estuarine exchange flow, and its extension to total exchange flow by Burchard et al. (2018). The contribution of the inflowing saline water to the spatial distribution of salt and the total salt budget depends essentially on mixing with ambient brackish waters. In the course of the inflow path, the character of the mixing process between the deep salty layer and the brackish water above changes with increasing depth and decreasing current velocities. In the entrance area the mixing is dominated by entrainment of brackish water into the eastward spreading saline bottom water, due to turbulence generated by shear instability. The sills between the consecutive Baltic basins are particular mixing hotspots (Neumann et al., 2017). In the deeper basins of the Baltic proper, boundary mixing driven by the interaction of currents and internal waves with the topography, and mixing processes at sill overflows contribute to the upward salinity flux (Reissmann et al., 2009). Mixing in the eastern Gotland Basin was investigated during the Baltic Sea Tracer Experiment (BATRE). Using an inert tracer gas, the basin-scale vertical diffusivities were estimated to $10^{-5} \text{ m}^2 \text{ s}^{-1}$, whereas the diffusivities in the basins interior were one order of magnitude lower (Holtermann et al., 2012). This finding holds also for the inflow of saline water in course of an MBI (Holtermann et al., 2017). The interior mixing is often controlled by double diffusive convection that leads to a typical stair-case-like vertical stratification structure (Umlauf et al., 2018). The crucial role of boundary mixing at the basin rim was confirmed by Holtermann and Umlauf (2012), and Lappe and Umlauf (2016). Both studies identified near-boundary turbulence as the key processes for basin-scale mixing. Main energy sources for boundary mixing are basin-scale topographic waves, deep rim currents, and near-inertial waves.



1666
 1667 The temporal statistics of barotropic saline inflows was reviewed by Mohrholz (2018). In contrast to earlier
 1668 investigations he found no long-term trend in inflow frequency, but a pronounced multidecadal variability of 25
 1669 to 30 years. Lehmann and Post (2015) and Lehmann et al. (2017) who studied the frequency and intensity of large
 1670 volume changes in the Baltic Sea due to inflows, likewise could not find a long-term trend. The distinction between
 1671 MBIs and smaller inflows is artificial and does not correspond to the frequency distribution of the inflows, which
 1672 shows an exponential decrease in frequency with increasing inflow intensity (Mohrholz, 2018). The classical MBIs
 1673 are only responsible for about 20% of the total salt input, while the rest is accounted for by medium and small
 1674 inflows with much less pronounced interannual variability.
 1675
 1676 Paleoclimate simulations covering nearly the recent millennium (Schimanke and Meier, 2016) have provided new
 1677 insights into the long-term behavior of the mean salinity of the Baltic Sea. In accordance with previous historical
 1678 reconstruction studies (Schimanke and Meier, 2016; Meier and Kauker, 2003), Schimanke and Meier (2016)
 1679 identified river discharge, net precipitation and zonal winds as main drivers of the decadal variability in Baltic Sea
 1680 salinity. However, their relative contributions are not constant. Extreme periods with strong salinity decrease for
 1681 about 10 years occurred once per century. Thus, the long stagnation period from 1976 to 1992 was obviously a
 1682 rare but natural event, although its extreme duration might be caused by anthropogenic effects. The Baltic Sea
 1683 salinity also has a natural centennial variability. Based on the same numerical simulations, Börgel et al. (2018)
 1684 could show a strong coherence between the AMO climate mode and river runoff on timescales between 60 and
 1685 180 years. Accordingly, the Baltic Sea salinity and the AMO are correlated, probably due to the dominating impact
 1686 of river discharge on salinity. The river runoff leads salinity changes by about 20 years during the entire modeling
 1687 period of 850 years. Schimanke and Meier (2016) reported a similar lag of 15 years between river runoff and Baltic
 1688 Sea salinity.
 1689
 1690 According to model results, multidecadal variations in runoff (Gailiusis et al., 2011; Meier et al., 2019d) explain
 1691 about half the long-term variability of volume-averaged Baltic Sea salinity (Meier and Kauker, 2003). Radtke et
 1692 al. (2020) found that the direct dilution effect was only responsible for about one fourth of the multidecadal
 1693 variability and proposed a link between river runoff and inflow activity. Furthermore, they found that the influence
 1694 of vertical turbulent mixing is small. Salt water inflows contribute to the multidecadal salinity variability, in
 1695 particular for the bottom layer salinity. The positive trend of river runoff in the northern catchment area led to a
 1696 significant increase in the North-South salinity gradient in the Baltic Sea surface water layer (Kniebusch et al.,
 1697 2019b). Additionally, their model based study revealed a multidecadal oscillation of salinity, river runoff and
 1698 saltwater inflows of about 30 years, consistent with the long term observations.
 1699
 1700 From observations during 1982–2016, Liblik and Lips (2019) detected decreasing surface (see also Vuorinen et
 1701 al., 2015) and increasing bottom salinities, but no long term trend in the total salt budget were found (cf. Fig. 21).
 1702 Both temperature and salinity contribute to strengthening of the vertical stratification. Enhanced freshwater fluxes
 1703 combined with higher deep water salinities intensify the vertical density gradient throughout the year.



1704 3.2.5.3 Stratification and overturning circulation

1705 A direct consequence of increasing stratification is that mixing weakens between well ventilated surface waters
 1706 and badly ventilated deep waters weakens, making the Baltic Sea vulnerable to deoxygenation of bottom waters
 1707 (Conley et al., 2002). An increase in seasonal thermal stratification (e.g. Gröger et al., 2019) can additionally lower
 1708 the vertical nutrient transport from deeper layers to the euphotic zone, thereby limiting nutrient supply and
 1709 potentially affecting algal and cyanobacterial blooms, at least at the species level. The latter potential effect has
 1710 not yet been thoroughly investigated. However, the hypothesis was supported by the results of Lips and Lips (2008)
 1711 who found a correlation between cyanobacteria bloom intensity in the Gulf of Finland and the frequency of
 1712 upwelling events along both coasts.

1713
 1714 Since the start of regular salinity measurements at the end of the 19th century, the haline stratification has been
 1715 dominated by sporadic inflows from the adjacent North Sea and variations in river discharge (Fig. 22). While no
 1716 long-term trends could be demonstrated in Baltic Sea salinity during 1921-2004 (Kniebusch et al., 2019b) or in
 1717 halocline depth during 1961-2007 (Väli et al., 2013), a trend towards increased horizontal salinity difference
 1718 between the northern and southern Baltic Sea was found during 1921-2004 (Kniebusch et al., 2019b). Modeling
 1719 studies by Kniebusch et al. (2019b) attributed this trend to increased river runoff from the northernmost catchment
 1720 area. Stratification increased in most of the Baltic Sea during 1982-2016, with the seasonal thermocline
 1721 strengthening by 0.33–0.39 kg m⁻³ and the perennial halocline by 0.70–0.88 kg m⁻³ (Liblik and Lips, 2019).

1722
 1723 Sensitivity studies with a numerical model suggest that the basin-wide overturning circulation will decrease if the
 1724 climate warms or when river runoff increases, but will tend to increase if global sea level rises (Placke et al., 2021).
 1725 However, historical multidecadal variations of the overturning circulation are mainly wind-driven. Multidecadal
 1726 variations in neither river runoff nor saltwater inflow had an impact, according to Placke et al. (2021).

1727 3.2.5.4 Sea level

1728 For the era of continuously operated satellite altimetry, absolute mean sea level (relative to the reference geoid)
 1729 increased in the Baltic Sea. Available estimates vary depending on the exact period considered, but are broadly
 1730 consistent with or slightly above the global average (3-4 mm year⁻¹; Oppenheimer et al., 2019; Nerem et al., 2018).
 1731 For the period 1992-2012, Stramska and Chudziak (2013) estimated an increase of 3.3 mm year⁻¹, and for the
 1732 period 1993-2015 Madsen et al. (2019a) an increase of 4 mm year⁻¹ in the Baltic Sea absolute mean sea level. For
 1733 the period 1886/1889-2018, the analysis of Swedish mareograph data suggest a sea level rise of about 1-2 mm
 1734 year⁻¹ (Figs. 23 and 24). Passaro et al. (2021) showed that the increase is not uniform across the Baltic Sea but
 1735 varies between about 2 mm year⁻¹ in the western Baltic Sea and more than 5 mm year⁻¹ in the Gulf of Bothnia for
 1736 the period 1995-2019. The acceleration of sea level rise in the Baltic Sea was studied by Hünicke and Zorita
 1737 (2016). They found that present acceleration is small and could only be detected through averaging of observations.

1738
 1739 Sea level changes relative to the coast are more complex, since land is rising in the northern Baltic Sea, by up to
 1740 about 8 mm year⁻¹, and sinking in the southern Baltic Sea, by about 1 mm year⁻¹ (Hünicke et al., 2015; Groh et al.,
 1741 2017). In addition to the global mechanisms (thermal expansion due to warming and land-ice melting), sea level
 1742 changes in the Baltic Sea are also affected by the changes in atmospheric circulation, water inflow from the North
 1743 Sea, and changes in the freshwater budget (river runoff, precipitation and evaporation). Precipitation and river



runoff are linked to westerly winds and affect salinity and the salinity gradient across the Baltic Sea (Kniebusch et al., 2019b) and thus the sea level height and its gradient. Stronger than normal westerly winds are associated with increased transports across the Danish straits which leads to an increase in Baltic mean sea level. The correlations between sea level height and westerly wind are higher in the eastern and northern parts and lower in the southern and western parts of the Baltic Sea. Westerly winds in the region became more intense until the early 1990s, but have weakened somewhat thereafter (Feser et al., 2015). Over longer periods, no significant long-term trend is detected (Feser et al., 2015).

The Baltic mean sea level shows a pronounced seasonal cycle with a minimum in spring and maxima in late summer (in 1900-1930) or winter (in 1970-1998). According to Hünicke and Zorita (2008), the amplitude of the seasonal cycle increased over the 20th century. Other authors found different periods without systematic long-term trends (Barbosa and Donner, 2016) or even regional decreases (Männikus et al., 2020).

Baltic sea level extremes are caused by strong atmospheric cyclones, or more seldom by wind-induced meteotsunamis (Pellikka et al., 2020) and seiches (Neumann, 1941; Wübbler and Krauss, 1979). Cyclones are associated with strong winds that cause coastal storm surges over one or two days, and if their pathway is aligned along the west-east direction, cyclones may also increase the total volume of the Baltic Sea over one week by pushing in water masses from the North Sea into the Baltic Sea. Coastal storm surges can then reach 20 cm above the spatially averaged level (Weisse and Weidemann, 2017), in the eastern Gulf of Finland in Neva Bay even more. Extreme sea levels over a predefined threshold become more frequent with rising mean sea level (Pindsoo and Soomere, 2020). In addition, model results and analysis of observations indicate that atmospheric forcing is responsible for the long-term increases in storm surges in some localized areas of the eastern Baltic Sea (Ribeiro et al., 2014). The presence of sea-ice impedes the development of extreme sea levels by shielding the ocean surface from forcing by the wind, and coastal ice protects the coast from erosion by extreme sea levels.

Storm surges caused by strong onshore winds represent a substantial hazard for the low-lying parts of the Baltic Sea coast, in particular, the southwestern parts (Wolski et al., 2014), the Gulf of Finland (e.g. Suursaar and Sooäär, 2016), the Gulf of Riga (e.g. Männikus et al., 2019), and the Gulf of Bothnia (Averkiew and Klevanny, 2010). Highest surges were reported for the Gulf of Finland (about 4 m in 1824) and the western Baltic Sea (more than 3 m in 1871, Wolski and Wiśniewski, 2020). For the Gulf of Riga and the western Baltic Sea values around 2 and 1-1.5 m are frequent, respectively (Wolski and Wiśniewski, 2020). Hundred-year storm surges are higher (up to 2.4 m) at the inner end of the basins, furthest away from the Baltic proper, than in center of the Baltic Sea (up to 1.2 m). No consistent long-term trend for an increase in extreme sea levels relative to the mean sea level of the Baltic Sea has been found, in agreement with earlier assessments (BACC II Author Team, 2015). This finding is supported by paleoclimate model studies that show no influence on extremes sea levels in the North Sea in warmer climate periods compared to colder periods (Lang and Mikolajewicz, 2019), and by recent studies of sea level records that suggest a pronounced decadal to multidecadal variability in storm surges relative to the mean sea level (Marcos et al., 2015; Marcos and Woodworth, 2017; Wahl and Chambers, 2016). Although (Ribeiro et al., 2014) argued for an increase in annual maximum sea level during 1916-2005, especially in the northern Baltic Sea, and attributed the trends to changes in wind, these results were likely affected by the long-term internal variability.



1784 Furthermore, extreme sea level in the Gulf of Finland, especially in Neva Bay, are very sensitive to the position of
 1785 storm tracks (Suursaar and Sooäär, 2007).

1786 3.2.5.5 Waves

1787 Instrumental wave measurements in the Baltic Sea have been made since the 1970s, first as measurement
 1788 campaigns and since the 1990s as continuous monitoring (e.g. Broman et al., 2006; Tuomi et al., 2011). As the
 1789 spatial coverage of wave measurements is still quite sparse, and there are long-term data from few locations only,
 1790 wave hindcasts have become a valuable tool for estimating the Baltic Sea wave climate (e.g. Björkqvist et al.,
 1791 2018). Lately, satellite altimeter measurements have been used to estimate the changes in the Baltic Sea wave
 1792 climate (Kudryavtseva and Soomere, 2017).

1793
 1794 Hindcast studies (e.g. Räämet and Soomere, 2010; Tuomi et al., 2011; Björkqvist et al., 2020) are in good
 1795 agreement, and estimate the annual mean significant wave height (SWH) at 0.5-1.5 m in the open sea areas of the
 1796 Baltic Sea. Wave growth in the Baltic Sea is hampered by the shape and small size of the basins. The highest mean
 1797 values are recorded in the Baltic proper, with the longest and widest fetches. The gulfs have less severe wave
 1798 climates (Björkqvist et al., 2020). In addition, wave growth in the northern Baltic Sea in winter is limited by the
 1799 seasonal ice cover, leading to considerably lower mean and maximum values of SWH in the northernmost Gulf of
 1800 Bothnia and the easternmost Gulf of Finland.

1801
 1802 Although the measurement and hindcast periods have so far been quite short, some studies have also analyzed
 1803 trends in Baltic Sea SWH. For example, Soomere and Räämet (2011) and Kudryavtseva and Soomere (2017)
 1804 suggest an increasing trend in SWH since the 1990s, but results are site-specific and so far rather inconclusive.

1805
 1806 The seasonal wave climate is driven by the wind climate. The highest mean and maximum values of SWH are
 1807 reached in autumn and winter, while summer typically has the mildest wave climate. In sub-basins with long ice
 1808 season and large ice extent, such as the Bothnian Bay, the seasonal variation in the SWH is slightly different, since
 1809 waves are damped by the ice.

1810
 1811 So far the Northern Baltic proper (NBP) holds the record measured value of Baltic SWH. In December 2004 a
 1812 SWH of 8.2 m was measured by the NBP wave buoy, with a highest individual wave of c. 14 m (Tuomi et al.,
 1813 2011; Björkqvist et al., 2018). As the spatial coverage of the wave measurements has increased and milder ice
 1814 winters have allowed late autumn and even winter measurements also in the northern parts of the Baltic Sea, 8 m
 1815 SWH has been measured also in the Bothnian Sea, in January 2019. Björkqvist et al. (2020) estimated a return
 1816 period of 104 years for this event in the present climate. Hindcast statistics have suggested that even higher
 1817 maximum values between 9.5 – 10.5 m may occur in areas and times for which wave buoy measurements are not
 1818 available (Soomere et al., 2008; Tuomi et al., 2011; Björkqvist et al., 2018).

1819 3.2.5.6 Sedimentation and coastal erosion

1820 The Glacial Isostatic Adjustment and eustatic sea level change impose a first-order control on Baltic Sea coastal
 1821 landscape change (Harff et al., 2007). In the subsiding southern Baltic Sea region, wind-driven coastal currents



1822 and waves are the major drivers for erosion and sedimentation, especially along the sandy and clayey sections of
 1823 sandy beaches, dunes and soft moraine cliffs (Zhang et al., 2015; Harff et al., 2017).

1824

1825 Owing to spatial variation in aero- and hydrodynamic conditions (winds, waves and longshore currents) and
 1826 underlying geological structure (lithology, sediment composition), a diversity of morphological patterns have
 1827 developed along the Baltic Sea coast. Because of the dominant westerly winds that blow 60% of the year (Zhang
 1828 et al., 2011) and a sheltering by land in the west, wind-waves are larger in the south-eastern Baltic Sea than in the
 1829 south-western. As a result, sediment transport and dune development are more active and dynamic along the south-
 1830 eastern coast. Thus, the largest coastal dunes are found along the Polish coast, with wave length >100 m and height
 1831 > 20 m (Ludwig, 2017), while dunes along the German coast normally have wave length less than 60 m and height
 1832 below 6 m (e.g. Lampe and Lampe, 2018). Under conditions favorable for wind-driven sand accumulation along
 1833 sandy Baltic Sea coasts, a typical cross-shore profile features one or several foredune ridges, generally with a
 1834 height of between 3 and 12 m above the mean sea level (Zhang et al., 2015; Łabuz et al., 2018). At the backshore
 1835 behind the established foredune ridges, drifting or stabilized dunes in transgressive forms, mainly parabolic or
 1836 barchanoid types, are commonly developed. The source of sediment for dune development includes fluvio-glacial
 1837 sands from eroded cliffs, river-discharged sands, and older eroded dunes (Łabuz, 2015).

1838

1839 Because the wind-wave energy increases from west to east, so does erosion along the Baltic Sea coast. The mean
 1840 annual erosional rate of the soft moraine cliffs and sandy dunes along the north side of the southwestern Baltic Sea
 1841 coast (southern Sweden and Denmark) is 1-2 m, larger than 0.4-1 m along the south side of the southwestern Baltic
 1842 Sea coast (Germany). Erosion along the southern Baltic Sea coast increases eastward, with a mean annual rate of
 1843 0.5-1.5 m in Poland, and 0.5-4 m in Latvia, Lithuania and Russia (BACC II Author Team, 2015). Severe coastal
 1844 erosion in the Baltic Sea region is often caused by storms. The maximal storm-induced erosion increases eastward
 1845 from 2-3 m year⁻¹ at the southwestern Baltic Sea coast (southern Sweden, Denmark and Germany) to 3-6 m year⁻¹
 1846 along the Polish coast, and ~10 m year⁻¹ along the coast of Lithuania and Russia (Kaliningrad). Each storm can
 1847 erode soft Latvian cliffs 3-6 m, with a maximum of up to 20-30 m locally. Many sandy beaches along the Gulf of
 1848 Finland have recently been severely damaged by frequent storm surges, despite extensive protective measures
 1849 (BACC II Author Team, 2015).

1850

1851 The prevailing wind-wave pattern controls the spatial variations of not only coastline change rates but also
 1852 submarine morphologies (Deng et al., 2019). In the southwestern Baltic Sea coast where wind-wave energy is
 1853 relatively small, nearshore submarine morphology is generally featured by smooth transition from beach-
 1854 dunes/moraine cliffs to deeper water perturbed by one or two longshore bars. Morphological perturbations (e.g.
 1855 the number and amplitude of longshore bars and rip current channels) become increasingly larger toward the east
 1856 due to increased wind-wave energy. The wave incidence angle also impacts the nearshore submarine morphology,
 1857 with in general a smaller angle leading to a larger morphological heterogeneity, i.e. a larger amplitude of
 1858 perturbations. The amplitude of nearshore morphological perturbations may significantly affect coastal erosion
 1859 because rip currents act as efficient conduit for offshore sediment transport, despite that they only occur
 1860 sporadically along the Baltic Sea coast (Schönhofer and Dudkowska, 2021).

1861



The sediments eroded from the soft moraine cliffs are composed of grain sizes from clay to pebbles. The fine-grained sediments are mostly transported outwards to the deeper seafloor (i.e. the Baltic Sea basins), either suspended in the water column or in a concentrated benthic fluffy layer (Emeis et al., 2002). Eroded fine-grained sediments from the moraine cliffs have been found to contribute to a major portion (40-70%) of the Holocene deposits in the muddy Baltic Sea basins (Porz et al., 2021). Coarser material, such as sands, stays mostly nearshore, partly in the water, partly transported onto the beach and the dunes (Deng et al., 2014; Zhang et al., 2015).

3.2.5.7 Marine carbonate system and biogeochemistry

Studies summarized in BACC II showed that nitrate and phosphate concentrations in winter surface water of the Baltic proper had increased by a factor of about three in the second half of the twentieth century, and reached a peak between 1980 and 1990. This change was consistent with the enhanced nutrients inputs to the Baltic Sea and caused eutrophication in the affected basin. Based on the available CO₂ system data, it has been estimated that the net ecosystem production in the Baltic Sea has increased since the 1930s by a factor of about 2.5. Increase in net ecosystem production and poor ventilation of the deep water layers due to the permanent stratification of the water column caused significant expansion of the anoxic and hypoxic areas in the Baltic Sea. Since the 1980s, nutrients inputs to the Baltic Sea have decreased. This led to a decrease in winter surface-water nitrate concentrations. No decrease was, however, observed for phosphate, due to the long residence time of P in the Baltic Sea and reduced P storage in oxygen-deficient sediments by binding to Fe-oxyhydroxides.

3.2.5.7.1 Oxygen and nutrients

In the Baltic Sea, hypoxia (oxygen deficiency) and even anoxia has expanded considerably since the first oxygen measurements in 1898 (Gustafsson et al., 2012; Carstensen et al., 2014a; Fig. 25). In 2016, the maximum hypoxia area was about 70,000 km², almost the combined area of Belgium and the Netherlands, whereas it was presumably very small or even absent 150 years ago (Carstensen et al., 2014b; Carstensen et al., 2014a; Meier et al., 2019c; 2019d). Hypoxia was caused mainly by increasing land nutrient inputs and atmospheric deposition that led to eutrophication of the Baltic Sea (Andersen et al., 2017; Savchuk, 2018). The impacts of other drivers like observed warming and eustatic sea level rise were smaller, but still important (Carstensen et al., 2014a; Meier et al., 2019c). On annual to decadal time scales, halocline variations also had considerable influence on the hypoxic area (Conley et al., 2002; Väli et al., 2013).

Besides its detrimental effects on biota, hypoxia is responsible for the redox alterations of nitrogen and phosphorus integral stocks reaching in the Baltic proper hundreds of thousand tonnes annually: the DIN pool is being depleted by denitrification, while the DIP pool increases due to phosphate release in the water and sediment anoxic environments (e.g. Savchuk, 2010; 2018 and references therein). Resulting changes of nitrate and phosphate concentrations at the upper boundary of the halocline affect also neighboring gulfs, exporting to the Bothnian Sea and Gulf of Finland waters with elevated phosphorus concentration (Rolff and Elfving, 2015; Lehtoranta et al., 2017; Savchuk, 2018).

Despite the decrease of land nutrient inputs after the 1980s, the extent of hypoxia in the Baltic Sea remains unaltered. This is due to the long response time of the system to reductions in N and P inputs. According to recent computations, the residence times for TN and TP in the water and sediments of the Baltic Sea combined are 9 and



1901 49 years, respectively (Gustafsson et al., 2017; Savchuk, 2018). Furthermore, recently observed oxygen
 1902 consumption rates in the Baltic Sea are higher than earlier observed, shortening oxygen relieves from natural
 1903 ventilation by oxygen-rich saltwater intrusions from the North Sea (Meier et al., 2018b). Although sediments are
 1904 still the most important sinks of oxygen in the Baltic Sea, the increased rates of oxygen consumption was largely
 1905 driven by water column processes with, for instance, bacterial nitrification as the most prominent. Also
 1906 zooplankton and higher trophic level respiration were suggested to contribute more to oxygen consumption than
 1907 30 years ago (Meier et al., 2018b). However, the importance of the latter processes is still unknown. The present
 1908 total oxygen consumption rate in the water column below 60 m depth in the Baltic proper, Gulf of Riga and Gulf
 1909 of Finland is estimated to be about five times that in the period 1850-1950 (Meier et al., 2018b).

1910

1911 Hypoxia remains an important problem also in the Baltic Sea coastal zone (Conley et al., 2011). Coastal hypoxia
 1912 most often has episodic or temporary character and is driven by the seasonal variations in organic matter supply,
 1913 advective transports and water column stratification (Carstensen and Conley, 2019). The latter is mostly caused
 1914 by seasonal temperature changes, but may in some areas be due to occasional inflows of saltier water and lower
 1915 winds during summer, changing the vertical stratification. The coastal regions most affected by hypoxia are
 1916 estuaries in the Danish straits and parts of Swedish and Finnish archipelagos located in the Baltic proper and Gulf
 1917 of Bothnia (Conley et al., 2011). In contrast, hypoxia is rare along the southern and south-eastern coastline (from
 1918 Poland to Estonia) due to enhanced water circulation as well as in the less productive coastal zone of the northern
 1919 Baltic Sea. Despite the recently reduced nutrient inputs, bottom water oxygen concentrations have improved only
 1920 in a few coastal ecosystems that have experienced the largest reductions (for instance in the Stockholm
 1921 Archipelago). In most of the 33 coastal sites, evaluated by Caballero-Alfonso et al. (2015), oxygen conditions have
 1922 deteriorated, especially along the Danish and Finnish coasts. This finding was explained as a coupled effect of
 1923 climate changes, especially warming, which reduces oxygen solubility in water and strengthens thermal
 1924 stratification as well as a delay of the system in responding to nutrient reduction.

1925

1926 N and P are removed from the Baltic Sea by burial in sediments, but much N is also lost by denitrification
 1927 (Gustafsson et al., 2017). Coastal regions constitute an efficient nutrient filter (Almroth-Rosell et al., 2016; Asmala
 1928 et al., 2017) that remove about 16% of N (by denitrification) and as much as 53% of P (by burial) delivered from
 1929 land (Asmala et al., 2017). The filter effect of the coastal zone is, however, highly diverse. Denitrification rates
 1930 are highest in lagoons that receive large inputs of nitrate and labile organic material, while P is most efficiently
 1931 buried in archipelagos (Carstensen et al., 2020). Additionally, Hoikkala et al. (2015) argued that dissolved organic
 1932 matter (DOM) plays an important role for nutrient cycling in the Baltic Sea, since more than 25% of bioavailable
 1933 nutrients in riverine inputs and surface waters can be in organic form.

1934

1935 Furthermore, the exchange of nutrients between the coastal zone and the open sea (Eilola et al., 2012) and the role
 1936 of MBIs for the phosphorus cycling (Eilola et al., 2014) were analyzed. Eilola et al. (2014) concluded that the
 1937 overall impact of MBIs on the annual uplift of nutrients from below the halocline to the surface waters is small
 1938 because vertical transports are comparably large also during periods without MBIs. Instead, phosphorus released
 1939 from the sediments between 60 and 100 m depth in the eastern Gotland Basin contributes to the eutrophication,
 1940 especially in the coastal regions of the eastern Baltic proper.

1941



1942 The cycling between nutrients and phytoplankton biomass was studied by Hieronymus et al. (2018) who found a
 1943 regime shift between nutrient-limited phytoplankton variations before 1950 and a less nutrient-limited regime after
 1944 1950, with a larger impact of other variations such as those in water temperature.

1945 **3.2.5.7.2 Marine CO₂ system**

1946 The marine CO₂ system in the Baltic Sea is greatly influenced by the production and remineralization of organic
 1947 matter, as well as inputs of organic and inorganic carbon from land (Kuliński et al., 2017). The combination of all
 1948 these factors makes Baltic Sea pH and partial pressure of CO₂ (pCO₂) highly variable in space and time (Carstensen
 1949 and Duarte, 2019). The Baltic Sea surface water is in almost permanent pCO₂ disequilibrium with the atmosphere
 1950 throughout the year (Schneider and Müller, 2018). In spring and summer, the surface seawater is undersaturated
 1951 with respect to atmospheric CO₂, as a consequence of biological production and the shallowing mixed layer depth.
 1952 Thus, seawater pCO₂ typically has two minima corresponding to the spring bloom and the mid-summer nitrogen
 1953 fixation period. In autumn and winter, pCO₂ increases due to shifting balance between autotrophy and heterotrophy
 1954 and entrainment of deeper CO₂-rich waters.

1955
 1956 Remineralization of terrestrial organic matter plays an important role in shaping pCO₂ fields in the Baltic Sea.
 1957 Kuliński et al. (2016) found that about 20% of the dissolved organic carbon (DOC) delivered from the Vistula and
 1958 Odra rivers is bioavailable, while Gustafsson et al. (2014) even estimated that 56% of allochthonous (originating
 1959 outside the Baltic Sea) DOC is remineralized in the Baltic Sea. High inputs of terrestrial organic matter that is
 1960 subsequently partially remineralized in seawater turned the basins most affected by riverine runoff (Gulf of
 1961 Bothnia, Gulf of Finland, Gulf of Riga) to net CO₂ sources to the atmosphere in the period 1980-2005. This
 1962 outgassing was more than compensated by the high CO₂ uptake in the open Baltic proper. In 1980-2005, the whole
 1963 Baltic Sea was found to be on average a minor sink for atmospheric CO₂, absorbing $4.3 \pm 3.9 \text{ g C m}^{-2} \text{ yr}^{-1}$, with
 1964 the rate of atmospheric CO₂ exchange highly sensitive to the inputs of terrestrial organic matter (Gustafsson et al.,
 1965 2014).

1966
 1967 The high seasonal variability of pCO₂, enhanced by eutrophication, causes large seasonal fluctuations in surface
 1968 water pH, amounting to about 0.5 in the central Baltic Sea (Kuliński et al., 2017) and even more, often exceeding
 1969 1, in productive coastal ecosystems (Carstensen and Duarte, 2019; Stokowski et al., 2021). Furthermore, low total
 1970 alkalinity (A_T, measures buffer capacity), prominent in the northern basins, makes the Baltic Sea potentially
 1971 vulnerable to Ocean Acidification (OA), i.e. pH decrease caused by rising pCO₂ in the atmosphere and thus also
 1972 in seawater. However, Müller et al. (2016) showed that A_T in the Baltic Sea has increased over time, which may
 1973 partly be due to increasing inputs from land (Duarte et al., 2013). The highest trend, $7.0 \mu\text{mol kg}^{-1} \text{ yr}^{-1}$, found in
 1974 the Gulf of Bothnia, almost entirely mitigates the pH drop expected from rising pCO₂ in the atmosphere alone. In
 1975 the southern Baltic Sea, the A_T increase is lower ($3.4 \mu\text{mol kg}^{-1} \text{ yr}^{-1}$) and reduces OA by about half. High seasonal
 1976 pH variability, increasing A_T and variable productivity imply that OA is not measurable in the central and northern
 1977 Baltic Sea. In the Danish Straits, where no A_T increase has been detected (Müller et al., 2016), a mean pH decrease
 1978 of 0.004 yr^{-1} was identified in coastal waters in the period of 1972-2016 (Carstensen et al., 2018), approximately
 1979 twice the ocean trend.

1980



Recent studies showed that the Baltic Sea CO₂ system functions differently from the open ocean waters. These differences include a large CO₂ input from remineralization of terrestrial organic matter (Gustafsson et al., 2014), a considerable contribution by organic alkalinity (Kuliński et al., 2014; Ulfsbo et al., 2015; Hammer et al., 2017), A_T generation under hypoxic and anoxic conditions (Gustafsson et al., 2019; Łukawska-Matuszewska and Graca, 2018), and a borate-alkalinity anomaly (Kuliński et al., 2018), making modeling a challenge. Due to the insufficient understanding of the processes involved, state-of-the-art biogeochemical models cannot yet reproduce the positive A_T trend in the Baltic Sea.

3.2.6 Marine biosphere

3.2.6.1 Pelagic habitats

3.2.6.1.1 Microbial communities

Microbial communities respond to increases in sea surface temperature and river runoff that enhance metabolism and augment the amount of substrate available for bacteria. By using long time-series from 1994 to 2006, increased input of riverine dissolved organic matter (DOM) in the Bothnian Bay and Bothnian Sea was shown to suppress phytoplankton biomass production and shift the carbon flow towards microbial heterotrophy (Wikner and Andersson, 2012; Paczkowska et al., 2020). (Berner et al., 2018) presented further evidence of changes in marine microbial communities.

3.2.6.1.2 Phytoplankton and cyanobacteria

The phytoplankton growing season has become considerably prolonged in recent decades (Kahru et al., 2016; Groetsch et al., 2016; Hjerne et al., 2019; Wasmund et al., 2019). In the Baltic proper, the duration of the growing season arbitrarily indicated by a threshold of 3 mg Chl m⁻³ of a satellite-derived chlorophyll has doubled from approximately 110 days in 1998 to 220 days in 2013 (Kahru et al., 2016). In the western Baltic Sea, it now extends from February to December (Wasmund et al., 2019). Wasmund et al. (2019) analyzed data on chlorophyll a and microscopically determined biomass from 1988-2017 and found an earlier start of the growing season, which correlated with a slight increase in sunshine duration in spring, and a later end to the growing season, which correlated with warmer water in autumn. The shifts in the spring and autumn blooms led to a prolongation of the summer biomass minimum. However, time series were rather short (30 years) and trends in irradiance might be caused by internal variability (see Section 3.2.1.3). The spring phytoplankton communities have shifted from a preponderance of early-blooming diatoms to dominance by later-blooming dinoflagellates (Wasmund, 2017; Wasmund et al., 2017) and the autotrophic ciliate *Mesodinium rubrum* (Klais et al., 2011; Hållfors et al., 2013; Hjerne et al., 2019), perhaps due to reduced ice thickness and increased winter wind-speed since the 1970s (Klais et al., 2013). Wasmund (2017) suggested that the decline in the ratio between diatom and dinoflagellate biomasses between 1984 and 1991 was caused by warmer winters. Confidence in these results is, however, low.

In summer, the amount of cyanobacteria has increased and the phytoplankton biomass maximum, which in the 1980s was in spring, is now in July-August. This shift has been explained by a complex interaction between warming, eutrophication and increased top-down pressure (Suikkanen et al., 2013). There are, however, different opinions concerning the relative effects of eutrophication and climate on changes in phytoplankton biomass and community composition. In the long-term data, results vary according to area and species group (Wasmund et al.,



2019 2011; Groetsch et al., 2016). Some studies saw evidence of eutrophication effects, modified by climate-induced
 2020 variations in temperature and salinity (Hällfors et al., 2013; Olofsson et al., 2020). Others found no explanation
 2021 for the gradual change in community composition, and concluded that the Baltic Sea phytoplankton community is
 2022 not in a steady state (Olli et al., 2011; Griffiths et al., 2020).

2023

2024 Cyanobacteria accumulations derived from satellite data for 1979–2018 show both short-term (two to three year)
 2025 oscillations and decadal-scale variations (Kahru and Elmgren, 2014; Kahru et al., 2018; Kahru et al., 2020).
 2026 Cyanobacteria accumulations in the Baltic proper were common in the 1970s and early 1980s, but rare during
 2027 1985–1990. They increased again starting in 1991 and, especially since 1998. In the 1980s, the annual chlorophyll
 2028 maximum in the Baltic proper was caused by the spring diatom bloom, but has in recent decades shifted to the
 2029 summer cyanobacteria bloom in July; the timing of this bloom has also advanced by about 20 days, from the end
 2030 to the beginning of July (Kahru et al., 2016).

2031

2032 In the Baltic proper, the hypoxia-induced decrease in N:P ratio and increase of phosphate pool left over in the
 2033 surface layer after the spring bloom led to intensification of the “vicious circle” (Vahtera et al., 2007), further
 2034 augmented by increasing water temperature, and resulted in conspicuous expansion of the surface diazotrophic
 2035 cyanobacteria accumulations, covering in the 21st century 150–200 thousand square kilometers (e.g. Kahru and
 2036 Elmgren, 2014; Savchuk, 2018, and references therein). Although mechanisms of the interannual oscillations of
 2037 two to three years remain unexplained (Kahru et al., 2018), there is a strong correlation of the accumulations with
 2038 hypoxia-related biogeochemical variables and water temperature at the decadal scale of five to twenty years (Kahru
 2039 et al., 2020). In the Bothnia Sea, the decreased nitrogen import and increased phosphorus import from the Baltic
 2040 proper has shifted the nutrient balance and made cyanobacteria accumulations a permanent feature (Kahru and
 2041 Elmgren, 2014; Kuosa et al., 2017) and probably also increased production and sedimentation (Ahlgren et al.,
 2042 2017; Kahru et al., 2018; Kahru et al., 2020; Kuosa et al., 2017; Lehtoranta et al., 2017; Rolff and Elfving, 2015;
 2043 Savchuk, 2010; Vahtera et al., 2007).

2044

2045 Experimental evidence supports the idea that climate change can and will drive changes in the pelagic primary
 2046 production (Sommer et al., 2012), and a thorough review of benthic-pelagic coupling in the Baltic Sea
 2047 demonstrates ecosystem-wide consequences of altered pelagic primary production (Griffiths et al., 2017).

2048 3.2.6.1.3 Zooplankton

2049 Several studies have confirmed that marine copepod species have declined in abundance since the 1980s, while
 2050 euryhaline or limnetic, often small species have increased (Hänninen et al., 2015; Suikkanen et al., 2013; Kortsch
 2051 et al., 2021). The observed decline of marine taxa has been linked to the reduction in surface-water salinity since
 2052 the 1980s (Vuorinen et al., 2015), whereas the increase of brackish-water taxa has been positively influenced by
 2053 the temperature increase, directly or indirectly (Mäkinen et al., 2017). Small-scale effects on individual species
 2054 may affect reproductive success, and hence influence both populations and communities (Möller et al., 2015).



2055 3.2.6.2 Benthic habitats

2056 3.2.6.2.1 Macroalgae and vascular plants

2057 Long-term changes in Baltic Sea macroalgae and charophytes have been attributed to changes in salinity, wind
 2058 exposure, nutrient availability and water transparency (Gubelit, 2015; Blindow et al., 2016; Rinne and Salovius-
 2059 Laurén, 2020), and biotic interactions may also play a role (Haavisto and Jormalainen, 2014; Korpinen et al.,
 2060 2007). The long-term decrease of water transparency from 1936 to 2017 has been estimated to have reduced sea
 2061 floor areas in the northern Baltic Sea favorable for *Fucus* spp. by 45% (Sahla et al., 2020). Overall, it is expected
 2062 that climate change and its interaction with other environmental factors (e.g. eutrophication) will cause complex
 2063 responses and influence carbon storage in both macroalgae and vascular plants in the Baltic Sea (Jonsson et al.,
 2064 2018; Takolander et al., 2017; Röhr et al., 2016; Perry et al., 2019; Salo et al., 2020; Bobsien et al., 2021).

2065 3.2.6.2.2 Zoobenthos

2066 Soft-sediment benthic communities depend on variables that are influenced by climatic variability. On the south-
 2067 western coast of Finland, amphipods have been replaced by Baltic clam *Limecola balthica* and the invasive
 2068 polychaetes *Marenzelleria* spp., a change attributed to an increase in near-bottom temperature and fluctuations in
 2069 salinity and oxygen (Rousi et al., 2013). Variations of zoobenthos in the Åland archipelago during 1983-2012 were
 2070 associated with a salinity decline (Snickars et al., 2015), and effects related to climate change have acted as drivers
 2071 for the long-term progression of zoobenthic communities (Rousi et al., 2019; Weigel et al., 2015; Ehrnsten et al.,
 2072 2020).

2073 3.2.6.3 Non-indigenous species

2074 Numerous non-indigenous species have gained a stronghold in the Baltic Sea ecosystem during the past few
 2075 decades, and in many cases these species have wider tolerance-ranges than the native ones, thus making them
 2076 highly competitive under changing climate including warmer, and possibly less saline water, further impacted by
 2077 other drivers such as eutrophication. The ecological impacts of these species may vary from filling vacant
 2078 ecological niches to potentially outcompeting native species, and thus influencing the entire food web structure
 2079 and functioning (Weigel et al., 2015; Griffiths et al., 2017; Ojaveer et al., 2017).

2080 3.2.6.4 Fish

2081 Sprat and herring in the Baltic Sea are influenced by multiple factors, including fisheries, predation, food
 2082 availability and climatic variations. Sprat has benefited from the seawater warming (Voss et al., 2011). In 1990-
 2083 2020, sprat populations were affected both by climate and top-down control, i.e. fisheries and predation by cod
 2084 (Eero et al., 2016). In the 1980s, overfishing and a partly climate change-induced decline in suitable spawning
 2085 habitat, 'reproductive volume', interacted to drastically reduce the cod population (Hinrichsen et al., 2011; Casini
 2086 et al., 2016), with cascade effects on its main prey, sprat and herring, as well as zooplankton (Casini et al., 2008).
 2087

2088 The various effects of temperature and salinity on sprat and cod also resulted in a spatial mismatch between these
 2089 species, which contributed to an increase of sprat stocks (Reusch et al., 2018). The freshening of the Baltic Sea
 2090 surface water, with the associated decline in marine copepods (Hänninen et al., 2015), contributed to a halving of
 2091 weight-at-age of 3-year old herring, from 50–70 g in the late 1970s to 25–30 g in 2000s (Dippner et al., 2019).



2092

2093 Among coastal fish, pikeperch (*Sander lucioperca*) has recently expanded its distribution northwards along the
 2094 coasts of the Bothnian Sea, apparently aided by warmer waters (Pekcan-Hekim et al., 2011). For many coastal
 2095 piscivores (perch, pike, pike-perch), as well as for cyprinids, coastal eutrophication is, however, equally or more
 2096 important than climate (Bergström et al., 2016; Snickars et al., 2015). Long-term studies illustrate that it is hard
 2097 to disentangle abiotic and biotic interactions, e.g. between fish and their food (benthos), and climate-related drivers
 2098 thus appear significant on a multidecadal time-scale across a large spatial scale (Törnroos et al., 2019).

2099 3.2.6.5 Marine mammals

2100 The breeding distributions of the ice-breeding seals in the Baltic Sea have evolved with ice coverage, with the
 2101 seals breeding where and when ice optimal for breeding occurs. Breeding ringed seals need ice throughout their
 2102 relatively long lactation period (>6 weeks), and also use ice as moulting habitat. Ringed seals prefer compact or
 2103 consolidated pack ice as it provides cavities and snowdrifts suitable for the construction of the lairs, most
 2104 importantly the breeding lair (Sundqvist et al., 2012).

2105

2106 Sea ice changes, along with implementation of specific management- and protection measures, have had a rapid
 2107 influence on the populations of several Baltic Sea mammal populations, in particular seals (Reusch et al., 2018).
 2108 Reusch et al. (2018) attributed these changes also to reduced exposure to harmful substances and increases in
 2109 overall fish stocks as a consequence of eutrophication (including reduced stocks of several commercial fish
 2110 species). Thus, specific climate change-related impacts on seals are hard to establish, although reconstructions of
 2111 distributional histories since the last glaciation have been attempted for some seal species (Ukkonen et al., 2014).

2112

2113 The availability of suitable breeding ice for ringed seals in the Bothnian Bay is decreasing (Section 3.2.4.4). The
 2114 breeding success of ringed seal was probably reduced by the exceptionally mild winter of 2007–2008 (Jüssi, 2012)
 2115 and several similar or even milder ice-winters have followed (Ilmatieteen laitos, 2020). The winters 2019–2020,
 2116 2007–2008 and 2014–2015 are the mildest in the annual ice cover statistics for the Baltic Sea (Uotila et al., 2015).
 2117 The southern breeding populations of the ringed seal in the Gulf of Finland, the Gulf of Riga and Archipelago Sea
 2118 are already facing the challenges of milder winters: the ice covered area during the breeding season has been
 2119 reduced and overlying snow for breeding lairs has been absent from the southern areas of the ringed seal breeding
 2120 range in most winters of the past decade (Ilmatieteen laitos, 2020). Thus, the only available breeding ice in the
 2121 Gulf of Finland in 2020 was found very near St. Petersburg (Halkka, 2020).

2122

2123 For grey seals, the lower availability of suitable breeding ice in its core distribution area has led to more breeding
 2124 on land in areas where drift ice used to be found (Jüssi et al., 2008). Grey seals are known to gather to breed in
 2125 certain sea areas regardless of the winter severity, so some land colonies may become overpopulated. As an
 2126 example, in 2016, 3,000 grey seal pups were born on three islets in the northern Gulf of Riga.

2127

2128 Flooding of seal haul-outs due to sea level rise will first occur in the southernmost Baltic Sea, where relative sea
 2129 level rise will be most rapid (EEA, 2019c), and haul-out sites are mainly low sand or shingle banks. In Kattegat,
 2130 relative sea level rise is estimated to be lower, and while the haul-outs in the western part are low sand and shingle
 2131 banks similar to those in the southern Baltic Sea, haul-outs in eastern Kattegat are skerries with a higher profile.



2132 In the central and northern Baltic Sea, haul-outs are mainly skerries and here relative sea level rise is estimated to
 2133 be low or even negative in the 21st century (EEA, 2019c). Yet, in recent history, winter storm surges have been
 2134 observed to flood grey seal breeding colonies and push limited ice with ringed seal pups onto shore in the Gulf of
 2135 Riga and Pärnu Bay. In the southern Baltic Sea and western Kattegat, increasing sea levels may turn parts of larger
 2136 islands or previously inhabited islands into suitable seal haul-outs, but this is hard to project and depends, among
 2137 other things, on the future management and protection of such areas.

2138

2139 Sea levels in the southern Baltic Sea have been rising at up to 3 mm year⁻¹ since the 1970s (EEA, 2019c) and the
 2140 available haul-out areas have thus seen reductions already. However, during this time, the relevant harbour and
 2141 grey seal populations have been recovering at high rates from past depletion (HELCOM, 2018b), with no
 2142 documented or suspected effects of rising sea levels published.

2143

2144 The only cetacean resident in the Baltic Sea, the harbour porpoise (*Phocoena phocoena*), is a wide-spread species
 2145 and seems to be rather tolerant of different temperatures as well as habitats. There are harbour porpoise populations
 2146 in the waters around Greenland as well as along the coast of the Iberian Peninsula, Morocco, West Sahara and in
 2147 the Black Sea. However, the Baltic proper harbour porpoise population has been shown to differ genetically (Lah
 2148 et al., 2016) and morphologically (Galatius et al., 2012) from neighbouring populations, which may imply local
 2149 adaptations that we are currently unaware of. Sea ice limits the range available to the Baltic proper harbour
 2150 porpoise population since they need to come to the surface to breathe every 1-5 minutes. Hence, a decreasing ice
 2151 cover is likely to increase the available range for the population. However, a change in the prey community
 2152 resulting from climate-change related factors could potentially have serious effects on this critically endangered
 2153 (Hammond et al., 2008) population of a small whale, which is dependent on constant access to prey (Wisniewska
 2154 et al., 2016).

2155 3.2.6.6 Waterbirds

2156 The winter distribution of many waterbirds has extended northwards in response to the increase in temperature
 2157 and the decreasing extent of sea ice cover. This can be observed as an overall increase in winter abundance of
 2158 waterbirds, because part of the population of some species (mainly diving ducks) that formerly wintered further to
 2159 the southwest now remain in the Baltic Sea (Pavón-Jordán et al., 2019). Many species show decreasing trends in
 2160 abundance in the southern parts of their wintering ranges (typically in western and southern Europe) but increases
 2161 near the northern edge of their distribution, typically the Baltic Sea region (MacLean et al., 2008; Skov et al., 2011;
 2162 Aarvak et al., 2013; Lehikoinen et al., 2013; Pavón-Jordán et al., 2015; Nilsson and Haas, 2016; Marchowski et
 2163 al., 2017; Fox et al., 2019). Similar shifts are seen in species that traditionally wintered in the Baltic Sea, but
 2164 currently show declining wintering numbers there, as part of the population now winters in the White, Barents and
 2165 Kara seas (Fox et al., 2019).

2166

2167 Although the community composition changes rapidly, the changes are not fast enough to track the thermal isocline
 2168 shifts (Devictor et al., 2012; Galet et al., 2021). How species respond to changes in winter temperature seems,
 2169 however, to be highly species- or group-specific (Pavón-Jordán et al., 2019). Many species now winter closer to
 2170 their breeding areas, shortening migration distances (Lehikoinen et al., 2006; Rainio et al., 2006; Gunnarsson et
 2171 al., 2012).



2172
 2173 Mainly owing to milder spring temperatures and related effects on vegetation and prey, many waterbirds migrate
 2174 earlier in spring (Rainio et al., 2006), and hence arrive earlier in the breeding area (Vähätalo et al., 2004), and
 2175 some also start breeding earlier (van der Jeugd et al., 2009). Delayed autumn migrations have also been noted, but
 2176 their relation to climate change is less clear (Lehikoinen and Jaatinen, 2012).

2177
 2178 Earlier loss of sea ice was found to improve pre-breeding body condition of female common eiders, leading to
 2179 increasing fledging success in offspring (Lehikoinen et al., 2006). On the other hand, algal blooms promoted by
 2180 higher seawater temperature has in some cases caused low quality in bivalve prey for common eiders, leading
 2181 more birds to skip breeding (Larsson et al., 2014). Warmer seawater in winter also increases the energy expenditure
 2182 of mussels, thus directly reducing their quality as prey for eiders (Waldeck and Larsson, 2013).

2183
 2184 Most Baltic Sea waterbird species are migratory and affected by climate change also outside the Baltic Sea region,
 2185 in the Arctic (breeding season) and in southern Europe and western Africa (wintering; Fox et al., 2015). This is
 2186 important, given that climate warming is most pronounced in the Arctic and northern Eurasia and above average
 2187 also in southern Europe and northern Africa (Allen et al., 2018).

2188 3.2.6.7 Marine food webs

2189 The entire marine food web of the Baltic Sea has been greatly impacted by climate change-related drivers that
 2190 have altered the physical environment and the physiological tolerance limits of several species, by causing micro-
 2191 evolution of Baltic Sea species, and by interactive effects of climate change with other environmental drivers, such
 2192 as eutrophication and hypoxia/anoxia (Niiranen et al., 2013; Wikner and Andersson, 2012; Schmidt et al., 2020;
 2193 Pecuchet et al., 2020).

2194
 2195 Integrated approaches encompassing all of the ecosystem-components discussed above, are needed in order to
 2196 understand and manage the linkages between large-scale and long-term changes driven by synergistic impacts of
 2197 over-arching climate change-related physical and chemical drivers in combination with other factors such as
 2198 eutrophication, which may complicate human adaptation to the changing marine ecosystem (Blenckner et al.,
 2199 2015; Stenseth et al., 2020; Bonsdorff, 2021).

2200 3.3 Future climate change

2201 3.3.1 Atmosphere

2202 3.3.1.1 Large-scale atmospheric circulation

2203 In the future, the NAO is very likely to continue to exhibit large natural variations, similar to those observed in the
 2204 past. In response to global warming, it is likely to become slightly more positive on average (Knudsen et al., 2011).
 2205 Trends in the intensity and persistence of blocking remain uncertain (IPCC, 2014b). The AMO is expected to be
 2206 very sensitive even to weak global warming, shortening the time scale of its response and weakening in amplitude
 2207 (Wu et al., 2018; Wu and Liu, 2020). This will likely reduce the decadal variability of SSTs in the Northern
 2208 Hemisphere. Recent studies indicate a degree of decadal predictability for blocking and the NAO influenced by
 2209 the AMO (Athanasiadis et al., 2020; Wills et al., 2018; Jackson et al., 2015).



2210 3.3.1.2 Air temperature

2211 Table 7 lists the air temperature changes over the Baltic Sea catchment area and the Baltic Sea calculated from an
 2212 ensemble of regional coupled atmosphere-ocean simulations (Gröger et al., 2021b). Due to the ice/snow-albedo
 2213 feedback, warming is larger in winter than in summer, and the land is warming faster than the Baltic Sea (Fig.
 2214 26a). Due to its proximity to the Arctic, the Baltic Sea region including both land and sea is warming faster than
 2215 the global mean figures (Section 1.5.1.1). The surface air temperature increase is expected to be largest in the
 2216 northern Baltic Sea region especially in winter. These statements are true for both uncoupled atmosphere
 2217 (Christensen et al., 2021) and coupled atmosphere-ocean regional climate simulations (Gröger et al., 2019; Gröger
 2218 et al., 2021b).

2219
 2220 For RCP2.6, the global annual mean surface air temperature change averaged over the simulations is 1.0°C. The
 2221 corresponding global changes for RCP4.5 and RCP8.5 are 1.9 and 3.5°C, respectively. Over land in the Baltic Sea
 2222 region, the warming is larger in each of the three scenarios, amounting to 1.5, 2.6 and 4.3°C, respectively (Table
 2223 7). Over the Baltic Sea, the increase is slightly smaller than over land (1.4, 2.4 and 3.9°C, respectively) but still
 2224 larger than the corresponding global mean increase in surface air temperature. The latter result was expected and
 2225 found in coupled atmosphere-ocean scenario simulations (Table 7), but not in all atmosphere-only runs
 2226 (Christensen et al., 2021).

2227

2228 *Extreme Air temperatures:*

2229 Changes in daily minimum and maximum temperatures have similar spatial patterns as the mean air temperature
 2230 changes, with the expected greater warming for minimum temperature (Christensen et al., 2021). According to
 2231 Christensen et al. (2021) and previous studies (BACC II Author Team, 2015), the latter result is explained by the
 2232 reduced outgoing long-wave radiation under increased greenhouse gas concentrations. The long-wave radiation
 2233 acts to cool the surface, especially when the ground is warmer than the air, e.g. during winter and during nights.
 2234 The number of hot spells are projected to increase, in particular in the southern Baltic Sea region (Gröger et al.,
 2235 2021b). In coupled atmosphere-ocean simulations, the strongest increases in the annual mean number of
 2236 consecutive days of tropical nights and the annual maximum number of tropical nights (with temperature above
 2237 20°C all night) in the Baltic Sea region were projected to occur over the open sea (Gröger et al., 2021b). In contrast,
 2238 projections of tropical nights with atmosphere-only models show no significant change (Gröger et al., 2021b;
 2239 Meier et al., 2019a). Due to the sea ice/snow albedo feedback, the largest decline in the number of frost days was
 2240 projected to occur over the northeastern Baltic Sea region, i.e. northern Scandinavia and adjacent northern Russia
 2241 (Gröger et al., 2021b).

2242 3.3.1.3 Solar radiation and cloudiness

2243 There are a few studies on projected future solar radiation over Europe. Global climate models of the CMIP5
 2244 generation indicated an increase in surface solar radiation, highest over southern Europe and decreasing towards
 2245 north, but still with a slight increase over the Baltic Sea (Bartók et al., 2017; Müller et al., 2019). However, some
 2246 regional climate models instead showed a decrease in surface solar radiation over the Baltic Sea area, in winter by
 2247 about 10% over most of the catchment (Bartók et al., 2017; Christensen et al., 2021). This change was largely
 2248 attributed to increasing future cloud-cover, due to a more zonal airflow, and was accompanied by increased winter



precipitation. Thus, there are large differences in modelled surface solar radiation between global and regional models (Bartók et al., 2017). Unknown future aerosol emissions add to the uncertainty.

2251

Global mean energy balance components have improved with every new climate model generation. For the latest CMIP6, models show good agreement for clear sky shortwave energy fluxes in today's climate, both between models and compared to reference data (Wild et al., 2021). However, there are still substantial discrepancies among the various CMIP6 models in their representation of several of the global annual mean energy balance components, and the inter-model spread increases further on regional, seasonal and diurnal scales. Thus, future changes in solar radiation and cloudiness remain highly uncertain, not least on the regional scale.

2258 3.3.1.4 Precipitation

Precipitation in winter and spring is projected to increase over the entire Baltic Sea catchment, while summer precipitation is projected to increase in the northern half of the basin only (Christensen et al., 2021). In the south, summer precipitation is projected to change very little, although with a large spread between different models including both increases and decreases. The projected increase in the north is a rather robust feature among the regional climate models but with a large spread in the amount. Ensemble mean precipitation changes from coupled atmosphere-ocean simulations are summarized in Table 8. For the Baltic Sea catchment area, projected annual mean precipitation changes for the three RCP scenarios amount to 5, 9 and 15% (Table 8) and are much larger than global averages (Section 1.5.1.2). Over the Baltic Sea, the changes are similar than over the land area (6, 8 and 16%).

2268

Expressed by the Clausius-Clapeyron equation, warming increases the potential for extreme precipitation due to intensification of the hydrological cycle associated with the growth of atmospheric moisture content. For Northern Europe, regional climate models indicate an overall increase in the frequency and intensity of heavy precipitation events in all seasons and longer wet and dry spells (Christensen and Kjellström, 2018; Rajczak and Schär, 2017; Christensen et al., 2021, and references therein). The largest increase in the number of high precipitation days is projected for autumn. The number of drought events per year are expected to decrease, while their length is expected to increase (Christensen and Kjellström, 2018). Changes in more extreme events, like 10-, 20- or 50-year events, are less certain.

2277 3.3.1.5 Wind

In general, projected changes in wind speed over the Baltic Sea region are not robust among Earth System Models (Kjellström et al., 2018; Gröger et al., 2021b). However, Ruosteenoja et al. (2019) found in CMIP5 projections a slight but significant wind speed increase in autumn and a decrease in spring over Europe and the North Atlantic. Furthermore, over sea areas where the ice cover is projected to diminish on average, such as the Bothnian Sea and the eastern Gulf of Finland, the mean wind is projected to increase systematically because of a warmer sea surface and reduced stability of the planetary boundary layer (Meier et al., 2011c; Gröger et al., 2021b; Räisänen, 2017).

2284

Projection of the future behavior of extratropical cyclones are uncertain because changes in several drivers result in opposite effects on cyclone activity. With global warming, the lower troposphere temperature gradient between low and high latitudes decreases due to polar amplification. Near the tropopause and in the lower stratosphere, the



opposite is true, thus implying changes in baroclinicity (Grise and Polvani, 2014; Shaw et al., 2016; Stendel et al., 2021). An increase in water vapour enhances diabatic heating and tends to increase the intensity of extratropical cyclones (Willison et al., 2015; Shaw et al., 2016) and contribute to their propagation further poleward (Tamarin-Brodsky and Kaspi, 2017; Tamarin and Kaspi, 2017). The opposite is true in parts of the North Atlantic region, e.g. south of Greenland. For this region the North-South gradient is increasing, as the weakest warming in the entire Northern Hemisphere is over ocean areas south of Greenland. North of this local minimum the opposite is true. The increase in the North-South gradient over the North Atlantic may be responsible for some Earth System Models showing an intensification of the low pressure activity and thereby high wind speed over a region from the British Isles and through parts of north-central Europe (Leckebusch and Ulbrich, 2004; Ulbrich et al., 2008). These projections have been confirmed by (Harvey et al., 2012). They compared the ensemble storm track response of CMIP3 and CMIP5 model simulations and found that both projections show an increase in storm activity in the midlatitudes, with a smaller spread in the CMIP5 simulations. In contrast to CMIP3, the CMIP5 ensemble showed a significant decrease in cyclone track density north of 60°N. Hence, pre-CMIP3 and CMIP3 studies showed a clear poleward shift of the North Atlantic storm track (e.g. Fischer-Bruns et al., 2005; Yin, 2005; Bengtsson et al., 2009), whereas the CMIP5 ensemble predicts only an eastward extension of the North Atlantic storm track (Zappa et al., 2013). The newest generation of models from CMIP6 resulted in significant reduction of biases in storm track representation compared to CMIP3 and CMIP5, but the response to climate change is quite similar compared to the previous assessments (Harvey et al., 2020). The eastward extension of the North Atlantic storm track seems to be a robust result as it is found in pre-CMIP3, CMIP3 and CMIP5 simulations (Feser et al., 2015).

In summary, there is no clear consensus among climate change projections in how changes in frequency and/or intensity of extratropical cyclones will affect the Baltic Sea region (Räisänen, 2017). However, in future climate the frequency of severe wind gusts in summer associated with thunderstorms may increase (Rädler et al., 2019).

3.3.1.6 Air pollution, air quality and atmospheric nutrient deposition

The main conclusions by the BACC II Author Team (2015) concerning projections of air quality in the Baltic Sea region still hold. The main factor determining future air quality in the region is regional emissions of air pollutants, not changes in meteorological factors related to climate change or in intercontinental pollution transport (see e.g. Langner et al., 2012; Hedegaard et al., 2013).

Recent post-BACC II air quality modelling studies for the Baltic Sea area are Colette et al. (2013), Varotsos et al. (2013), Colette et al. (2015), Hendriks et al. (2016), and Watson et al. (2016). They concentrate mainly on particulate matter (PM) and ground-level ozone (O₃), the pollutants most likely to be affected by changing climate parameters. They agree with current day air quality trends in that the Baltic Sea region in general is less exposed to air pollution than the rest of Europe.

Jacob and Winner (2009) showed that climate change is likely to increase ground-level ozone in central and southern Europe. In a meta-analysis, Colette et al. (2015) assessed the significance and robustness of the impact of climate change on European ground-level ozone based on 25 model projections, including some driven by SRES (Special Report on Emission Scenarios by Nakicenovic et al., 2000) and RCP scenarios. They indicate that an increase in ground-level ozone is not expected for the Baltic Sea region. A latitudinal gradient was found from



2328 increase in large parts of continental Europe (+ 5 ppbv), but a small decrease over Scandinavia (up to -1 ppbv).
 2329 Studies that explicitly compared the magnitude of projected climate and anthropogenic emission changes (Langner
 2330 et al., 2012; Colette et al., 2013; Varotsos et al., 2013) all confirmed that changes in emission of ozone precursors
 2331 (NO_x , VOCs) had the larger effect. For Northern Europe, Varotsos et al. (2013) estimated that reductions in snow
 2332 cover and solar radiation in a SRES A1B scenario lead to an ozone decrease of about 2 ppb by 2050, compared to
 2333 present conditions.
 2334
 2335 Varotsos et al. (2013) stress the importance of future biogenic isoprene emissions for ozone concentrations. In the
 2336 2050 climate, increases in ozone concentrations are associated with increased biogenic isoprene emissions due to
 2337 increased temperatures, whereas increased water vapour over the sea, as well as increased wind speeds, are
 2338 associated with decreases. Hendriks et al. (2016) emphasise that isoprene emissions may increase significantly in
 2339 coming decades if short-rotation coppice plantations are greatly expanded, to meet the increased biofuel demand
 2340 resulting from the EU decarbonisation targets. They investigate the competing effects of anticipated trends in land
 2341 use, anthropogenic emissions of ozone precursors and climate change on European ground-level ozone
 2342 concentrations and related health and environmental effects by 2050. They found that increased ozone
 2343 concentrations and associated health damage caused by a warming climate (+ 2 to 5°C across Europe in summer)
 2344 might be more than the reduction that can be achieved by cutting emissions of anthropogenic ozone precursors in
 2345 Europe.
 2346
 2347 Orru et al. (2013, 2019) and Geels et al. (2015) studied the effect of climate change on ozone-related mortality in
 2348 Europe. Orru et al. (2019) present their results on country level, including all Baltic Sea EU-countries. They
 2349 conclude that although mortality related to ground-level ozone is projected to be lower in the future (mainly due
 2350 to decrease precursor emissions), the reduction could have been larger, without climate change and an increasingly
 2351 susceptible population.
 2352
 2353 In parts of the Baltic Sea region, a considerable air pollution is due to shipping. Ship traffic in the region is
 2354 projected to increase over the coming decades, which could lead to larger emissions (i.e. NO_x and PM) than today,
 2355 unless stricter air quality regulations counter this potential trend. For the Baltic Sea, a nitrogen emission control
 2356 area (NECA) will become effective in 2021. Karl et al. (2019a) designed future scenarios to study the effect of
 2357 current and planned regulations of ship emissions and the expected fuel efficiency development on air quality in
 2358 the Baltic Sea region. They showed that in a business-as-usual scenario for 2040 (SECA-0.1% and fuel efficiency
 2359 regulation effective starting in 2015), the introduction of the NECA will reduce NO_x emissions from ship traffic
 2360 in the Baltic Sea by about 80% in 2040. The reduction in NO_x emissions from shipping translates to a ~60%
 2361 decrease in NO_2 summer mean concentrations in a wide corridor around the ship routes. The coastal population of
 2362 northern Germany, Denmark and western Sweden will be exposed to less NO_2 in 2040 due to the introduction of
 2363 the NECA. With lower atmospheric NO_x levels, less ozone will be formed, and the estimated daily maximum O_3
 2364 concentration over the Baltic Sea in summer 2040 will on average be 6% lower than without the NECA. Compared
 2365 to today, the introduction of the NECA will also reduce ship-related $\text{PM}_{2.5}$ emissions by 72% by 2040, compared
 2366 to -48% without the NECA. Simulated nitrogen deposition on the Baltic Sea decreases 40-44% on average between
 2367 2012 and 2040. A similar study by Jonson et al. (2019) estimated that the contributions of Baltic Sea shipping to



2368 NO₂ and PM_{2.5} concentrations, and to the deposition of nitrogen, will be reduced by 40-50 % from 2016 to 2030,
 2369 mainly as a result of NECA.

2370 3.3.2 Land

2371 3.3.2.1 River discharge

2372 Climate change is likely to have a clear influence on the seasonal river flow regime, as a direct response to changes
 2373 in air temperature, precipitation and evapotranspiration (BACC II Author Team, 2015; Blöschl et al., 2017).

2374
 2375 For areas in the northern Baltic Sea region presently characterized by spring floods due to snow melt, the floods
 2376 are likely to occur earlier in the year and their magnitude is likely to decrease owing to less snowfall, shorter snow
 2377 accumulation period, and repeated melting during winter. As a consequence, sediment transport and the risk of
 2378 inundation are likely to decrease.

2379
 2380 In the southern part of the Baltic Sea region, increasing winter precipitation is projected to result in increased river
 2381 discharge in winter. In addition, groundwater recharge is projected to increase in areas where infiltration capacity
 2382 is not currently exceeded, resulting in higher groundwater levels. Decreasing precipitation combined with rising
 2383 temperature and evapotranspiration during summer is projected to result in drying of the root zone, increasing
 2384 demands for irrigation in the southern Baltic Sea region.

2385
 2386 Projections with a process-oriented hydrological model suggested that, under the RCP4.5 and RCP8.5 scenarios,
 2387 the total river flow during 2069-2098 relative to 1976-2005 will increase 1-21% and 6-20%, respectively,
 2388 illustrating the large uncertainty in hydrological projections (Saraiva et al., 2019a; Meier et al., 2021b). According
 2389 to these and previous projections, the increase of river flow will mainly take place in the north, while total river
 2390 flow to the south will decrease (Stonevičius et al., 2017; Šarauskienė et al., 2017). Winter flow will increase due
 2391 to intermittent melting (Stonevičius et al., 2017). Projected discharge changes attributed to increasing air
 2392 temperature are reflected in observed trends (Section 3.2.2.1), whereas changes attributed to increasing
 2393 precipitation are necessarily not (Wilson et al., 2010).

2394
 2395 Since the publication of BACC II (BACC II Author Team, 2015), ensemble sizes of scenario simulations with
 2396 hydrological models have increased, enabling the estimate of uncertainties in projections (e.g. Roudier et al., 2016;
 2397 Donnelly et al., 2017). Donnelly et al. (2014) focused on projecting changes in discharge to the Baltic Sea, by
 2398 using a semi-distributed conceptual hydrological model for the BSDb (Balt-HYPE), combined with a small
 2399 ensemble of climate projections under the SRES A1B and A2 scenarios. Results showed an increased overall
 2400 discharge to the Baltic Sea, with a seasonal shift towards higher winter and lower summer flows and diminished
 2401 seasonal snow-melt peaks. Efforts were made to assess the uncertainty in the model chain, and change magnitudes
 2402 were shown to be within the range of the overall uncertainty estimates, highlighting the importance of such
 2403 uncertainty assessments in effect studies to frame the quantitative model results.

2404
 2405 Arheimer and Lindström (2015) studied future changes in annual maximum and minimum daily flows. Their
 2406 projections suggested that snow-driven spring floods in the northern–central part of Sweden may occur about one



month earlier than today and rain-driven floods in the southern part of Sweden may become more frequent. The boundary between the two flood regimes is projected to shift northward.

Past observations (see Section 3.2.2.1) and future projections (e.g. Graham, 2004) suggest a temporal shift in the seasonality of the river discharge, with decreasing flow in spring/summer and increasing flow in winter. Global warming and river regulation due to hydropower production cause similar changes. However, in snow-fed rivers globally the impact of climate change is projected to be minor compared to river regulation (Arheimer et al., 2017).

3.3.2.2 Land nutrient inputs

Projected changes in riverine discharge and nutrient inputs from the Baltic Sea Drainage Basin (BSDB) to Baltic Sea coastal waters have been studied using a number of modelling frameworks in recent years. Projecting the regional effects of future climate and environmental change on hydrology and nutrient turnover poses challenges in terms of (i) the complex nature of the modelled system, including human influence on riverine nutrient inputs and transport processes alike, which necessitates long projection model chains and leads to uncertainty in modelled hydrologically driven responses, and (ii) the significance of changes in human behaviour, e.g. in terms of land management, population, or nutrient emissions from point sources, which adds complexity to the formulation scenarios for future change, on top of the climate change signal. Hydrological impact studies in the BSDB (and elsewhere) therefore often explicitly use simplifying assumptions in order to reduce complexity of the modelled system and to put focus on certain aspects of impacts of projected changes.

Hesse et al. (2015) also reported increasing discharges in a similar model study of the Vistula lagoon catchment, using a hydrological model (SWIM), which also allows for nutrient load assessment, and climate change impact modelling based on a climate model ensemble. On average, results showed decreasing trends for nitrogen and phosphorus inputs, but a wide range of projections with individual ensemble members.

Hägg et al. (2014) used a split model approach to project changes in TN and TP inputs to Baltic Sea sub-basins. Changes in discharge were estimated with a hydrological model (CSIM) combined with a climate projection ensemble, which sampled a range of climate model and emission scenario combinations. Inputs were then calculated with a statistical model, based on modelled discharges and population as a proxy for human nutrient emissions, combining population change assumptions with climate projections. Results showed a general trend towards higher nutrient inputs across the region as a result of climate change, and a significant (i.e. potentially trend-changing) influence of human adaptation scenarios, particularly in the southern half of the BSDB.

Øygarden et al. (2014) used measurements in a number of small agricultural catchments to establish functional relationships between precipitation, runoff, and N losses from agricultural land, and qualitatively related their findings to projected precipitation changes across the BSDB under climate change scenarios, as well as to mitigation measures to counter the climate-driven effects. The analyses showed a positive relationship between runoff and N losses as well as between rainfall intensity and N losses, but stressed the wide range of feedback loops possible between climate change effects and adaptation measures, through management or policy changes. Such data-driven approaches avoid uncertainties related to effect-model chains at the expense of direct BSDB-wide quantitative effect projections.



2447

2448 The potential effects of socio-economic adaptation under climate change conditions were investigated by Huttunen
 2449 et al. (2015) in a study of Finnish catchments draining to the Baltic Sea. A national nutrient load model (VEMALA)
 2450 was combined with a mini-ensemble of climate effects, and then a number of agricultural adaptation scenarios
 2451 were derived, based on crop yield and policy changes, and an economic model (DREMFIA) was used to translate
 2452 scenario assumptions to changes in the nutrient load model for evaluation of effects. On average, increased
 2453 precipitation led to increased annual discharge and a shift from spring to winter peaks, with total nitrogen (TN)
 2454 and total phosphorus (TP) inputs increasing with the discharge. Here, adaptation scenarios had less effect than
 2455 climate change, with some regional variation, but significantly different load reductions were found among
 2456 assessed adaptation strategies, leading to the conclusion that adaptation measures are important for overall climate
 2457 change effect mitigation in the region.

2458

2459 The relative importance of management decisions for TN and TP load effects was studied also by Bartosova et al.
 2460 (2019), using the hydrological model E-HYPE on the full BSDB. The ensemble approach combined climate and
 2461 socioeconomic pathways based on IPCC fifth assessment data (Zandersen et al., 2019), where socioeconomic
 2462 changes were directly translated into changes of the effect model setup. The influence of socioeconomic adaptation
 2463 choices on nutrient inputs to the Baltic Sea were shown to be in the same magnitude range as climate effects, thus
 2464 indicating the importance of effective mitigation strategies for the region. In order to increase this efficiency,
 2465 Refsgaard et al. (2019) developed and explored the concept of spatially differentiated measures for TN load
 2466 reductions in the BSDB, based on the realization that measures are not uniformly efficient over large area, and
 2467 should therefore not be uniformly applied either.

2468 3.3.3 Terrestrial biosphere

2469 In the following, we focus on the European drought in 2018, to study the impact of very warm conditions on the
 2470 terrestrial ecosystem, and on projections for the terrestrial ecosystems in the Arctic, because of the particularly
 2471 strong climate warming in the Arctic and potentially strong feedbacks from the release of CO₂ and CH₄ in the
 2472 northernmost part of the Baltic Sea region. Finally, we discuss mitigation scenarios for land use and land-cover
 2473 changes associated with the Paris Agreement.

2474 *Terrestrial ecosystems in the European drought year 2018*

2475 The summer of 2018 saw extremely anomalous weather conditions over Europe, with high temperatures
 2476 everywhere, as well as low precipitation and high incoming radiation in western, central and Northern Europe
 2477 (Peters et al., 2020). These extreme weather conditions resulted in severe drought (indicated by soil moisture
 2478 anomalies) in western, central and Northern Europe, including the entire Baltic Sea region. The impacts of the
 2479 severe drought and heatwave in Europe in 2018 were investigated in a series of papers, ranging from individual
 2480 sites to the continental scale (Peters et al., 2020).

2481 Graf et al. (2020) studied the effects of the 2018 drought conditions on the annual energy balance at the land
 2482 surface, in particular the balance between sensible and latent heat fluxes, across different terrestrial ecosystems at
 2483 various sites in Europe. Graf et al. (2020) found a 9% higher incoming solar radiation compared to their reference
 2484 period across the drought-affected sites. The outgoing shortwave radiation mostly followed the incoming radiation,



with an increase of 11.5%, indicating a small increase in surface albedo. The incoming longwave radiation, on the other hand, did not change significantly, indicating that effects of higher atmospheric temperatures and reduced cloudiness cancelled out, while outgoing longwave radiation increased by 1.3% as a result of higher land surface temperatures. Overall, the net radiation increased by 6.3% due to the extreme drought conditions. As for the non-radiative surface energy fluxes, the sensible heat flux showed a strong increase by 32%, while the latent heat fluxes did not change significantly on average. Graf et al. (2020) attributed the negligible effect on latent heat fluxes to the opposing roles of increased grass reference evapotranspiration on the one hand and soil water depletion, stomatal closure and plant development on the other. Evapotranspiration increased where and when sufficient water was available and later decreased only where stored soil water was depleted. As a consequence, latent heat fluxes typically decreased at sites with a severe precipitation deficit, but often increased at sites with a comparable surplus of grass reference evapotranspiration but only a moderate precipitation deficit. Consistent with this, peatlands were identified as the only ecosystem with very strong increases in latent heat fluxes but insignificant changes in sensible heat fluxes under drought conditions. Crop sites, on the other hand, showed significant decreases in latent heat fluxes.

Lindroth et al. (2020) analysed the impact of the drought on Scandinavian forests, based on 11 forest ecosystem sites differing in species composition, i.e. spruce, pine, mixed and deciduous. Compared to their reference year, in 2018 the forest ecosystem showed a slight decrease in evaporation at two of the sites, was nearly unchanged at most sites and increased at two sites with pine forest. At the same time, the mean surface conductance during the growing season was reduced 40-60% and the evaporative demand increased 15-65% due to the warm and dry weather conditions. The annual net ecosystem productivity (NEP) decreased at most sites, but the reasons differed. At some sites, the NEP decrease was due to an increase in ecosystem respiration (RE), while at others both RE and the gross primary productivity (GPP) decreased, with the decrease in GPP exceeding that in RE. At six sites, the annual NEP decreased by over 50 g C m⁻² year⁻¹ in 2018. Across all sites considered, NEP anomalies varied from -389 to +74 g C m⁻² year⁻¹. A multi-linear regression analysis revealed that the anomalous NEP could to a very large extent (93%) be explained by anomalous heterotrophic respiration and reduced precipitation, with most of the variation (77%) due to the heterotrophic component.

Rinne et al. (2020) studied the effects of the drought on greenhouse gas exchange in five northern mire ecosystems in Sweden and Finland. Due to low precipitation and high temperatures, the water table sank in most of the mires. This led not only to a lower CO₂ uptake, but also to lower CH₄ emissions by the ecosystems. Three out of the five mires switched from sinks to sources of CO₂. Estimates of the radiative forcing expected from the drought-related changes in greenhouse gas fluxes indicated an initial cooling effect due to the reduced CH₄ emissions, lasting up to several decades, followed by a warming caused by the lower CO₂ uptake. However, it is unknown whether these results can be generalized to all wetlands of the Baltic Sea region.

Terrestrial ecosystems in the Arctic region

Climate warming has been particularly strong at high northern latitudes, and climate change projections indicate that this trend will continue, due to the anticipated increase in anthropogenic climate forcing. This strong warming is expected to have major consequences for terrestrial ecosystems in Arctic and sub-Arctic regions.



2522 Zhang et al. (2013) used the Arctic version of a dynamic global vegetation model (LPJ-GUESS, Smith et al.,
 2523 2001), forced with a regionalized climate scenario (A1B anthropogenic emission scenario), to investigate land
 2524 surface feedbacks from vegetation shifts and biogeochemical cycling in terrestrial ecosystems under future climate
 2525 warming. They found marked changes in vegetation by the second half of the 21st century (2051-2080), i.e. a
 2526 poleward advance of the boundary between forests and tundra, expansion of tundra covered with tall shrubs and a
 2527 shift from deciduous trees, e.g. birch, to evergreen boreal coniferous forest. These changes in vegetation were
 2528 associated with decreases in surface albedo, particularly in winter due to the snow-masking effect, and with
 2529 increases in evapotranspiration. The reduced surface albedo would tend to enhance the projected warming (positive
 2530 feedback), while increased evapotranspiration would dampen it (negative effect). The terrestrial ecosystems
 2531 continued to act as carbon sinks during the 21st century, but at diminished rates in the second half of the century.
 2532 The initial increase in carbon sequestration, due to a longer growing season and CO₂ fertilisation, could be reduced
 2533 and eventually reversed by increased soil respiration and greater CO₂ release from increased wildfires. Peatlands
 2534 were identified as hotspots of CH₄ release, which would further enhance the projected warming (positive
 2535 feedback).

2536 Using a regional Earth System Model (RCA-GUESS; Smith et al., 2011) over the Arctic region, Zhang et al.
 2537 (2014b) investigated the role that the biophysical effects of the projected future changes of the land surface play
 2538 for the terrestrial carbon sink in the Arctic region under a future climate scenario based on a high emission scenario
 2539 (RCP8.5). Two simulations were performed to determine the role of the biophysical interactions, one with and one
 2540 without the biophysical feedbacks resulting from the simulated climatic changes to the terrestrial ecosystems in
 2541 the model. In both simulations the Arctic terrestrial ecosystems continued to sequester carbon until the 2060-
 2542 2070s, after which they were projected to turn into weak sources of carbon, due to increased soil respiration and
 2543 biomass burning. The biophysical effects were found to markedly enhance the terrestrial ecosystem carbon sink,
 2544 particularly in the tundra areas. Two opposing feedback mechanisms, mediated by changes in surface albedo and
 2545 evapotranspiration, contributed to the additional carbon sequestration. The decreased surface albedo in winter and
 2546 spring notably amplified warming in spring (positive feedback), while the increased evapotranspiration led to a
 2547 marked cooling during summer (negative feedback). These feedbacks stimulated vegetation growth due to an
 2548 earlier start of the growing season, leading to changes in woody plant species and the distribution of vegetation.
 2549 In a later study, Zhang et al. (2018) found that these biophysical feedbacks play essential roles also in climate
 2550 scenario simulations with weaker anthropogenic climate forcing.

2551 *Mitigation*

2552 The beneficial effects of carbon sequestration by forest ecosystems on climate change may be reinforced,
 2553 counteracted or even offset by management-induced changes in surface albedo, land-surface roughness, emissions
 2554 of biogenic volatile compounds, transpiration and sensible heat flux (see above). Luyssaert et al. (2018)
 2555 investigated the trade-offs associated with using European forests to meet the climate objectives in the Paris
 2556 Agreement. A central argument of this study was that the agreement requires more than that forest management
 2557 should dampen the rise in atmospheric CO₂ and reduce the radiative imbalance at the top of the atmosphere. The
 2558 authors suggested two additional targets, that forest management should neither increase the near-surface
 2559 temperature nor decrease precipitation, because climate effects arising from the changes in the terrestrial biosphere
 2560 would make adaptation to climate change more demanding. Analysing different forest management portfolios in



Europe designed to maximize the carbon sink, maximize the forest albedo or reduce near-surface temperatures, Luyssaert et al. (2018) found that only the portfolio designed to reduce near-surface temperatures accomplished two of the objectives, i.e. to dampen the rise in atmospheric CO₂ and to reduce near-surface temperatures. This portfolio featured a decrease in the area of coniferous forest in favour of a considerable increase in the area of deciduous forest in Northern Europe, from 130,000 to 480,000 km².

3.3.4 Cryosphere

3.3.4.1 Snow

There is agreement among models that the average amount of snow accumulated in winter will decrease by over 70% in most of the Baltic Sea region. The high Scandinavian mountains, where the warming temperature will not reach the freezing point as often as in lower-lying regions, are an exception (Christensen et al., 2021). The reduction in snow amount is slightly larger than in maps presented by the BACC II Author Team (2015), which is consistent with the stronger average warming projected in the RCP8.5 scenario, compared to the SRES A1B scenario analyzed by the BACC II Author Team (2015).

For Poland, two additional downscaling experiments were made to produce reliable high-resolution climate projections of precipitation and temperature, using the RCP4.5 and RCP8.5 scenarios (Szwed et al., 2019). The results were used as input to a snow model (seNorge), to transform bias-adjusted daily temperature and precipitation into daily snow conditions. The snow model projected future snow depth to decrease in autumn, winter and spring, in both the near and far future. The maximum snow depth was projected to decrease 15-20% by 2021-2050 and at least double that decrease by 2071–2100 (Szwed et al., 2019).

3.3.4.2 Glaciers

The Special Report on the Ocean and Cryosphere in a Changing Climate (IPCC, 2019a) provides the most recent assessment of future projected glacier mass reduction under various RCPs, and treats Scandinavian glaciers separately (Hock et al., 2019). Previous projections, summarized in the Fifth Assessment Report of the IPCC (IPCC, 2014a; Vaughan et al., 2014), did not specifically focus on Scandinavian glaciers.

By 2100, likely (i.e. with a likelihood of 60-100%) mass losses for high-mountain glaciers are 22-44% (RCP2.6) to 37-57% (RCP8.5) of their mass in 2015. These losses exceed global projections for glacier mass loss of $18 \pm 7\%$ for RCP2.6, and $36 \pm 11\%$ for RCP8.5 (likely ranges, IPCC, 2019a). Glaciers in Scandinavia will lose over 80% of their current mass by 2100 under RCP8.5 (medium confidence), and many are projected to disappear, regardless of future emission scenarios (Fig. 27). Furthermore, river runoff from glaciers is projected to change regardless of emission scenario (high confidence), and to result in increased average winter runoff (high confidence) and in earlier spring peaks (high confidence; Hock et al., 2019).

Projections of future glacier mass loss depend crucially on climate projections providing surface air temperature and precipitation as forcing factors in process-based glacier models. For high mountain glaciers, such as those along the Scandinavian mountains that drain into the Baltic Sea, this is challenging, as the interplay of regional effects such as high-mountain meteorology and elevation-dependent warming (Wang et al., 2016; Qixiang et al., 2018) with global climate is poorly understood (Hock et al., 2019). Surface air temperatures in mountain regions



are projected to increase at an average rate of $0.3 \pm 0.2^{\circ}\text{C}$ per decade until 2050 (very high confidence), i.e. faster than the present global average of $0.2 \pm 0.1^{\circ}\text{C}$ (Hock et al., 2019). Beyond 2050, and depending on the emission scenario, air surface temperatures in high mountain regions are projected to either stabilize at the 2015-2050 rate or to increase further (IPCC, 2019a).

Projected changes in surface air temperature for the period 2071–2100 (compared to 1971–2000, under various emission scenarios) for the part of the Baltic Sea Drainage Basin that extends along the Scandinavian mountains (SMHI, 2020; Kjellström et al., 2016) will be of great importance for the assessment of future mass loss from glaciers draining into the Baltic Sea.

3.3.4.3 Permafrost

Due to recent warming more than 20% of the permafrost in the region was already lost in 1997-2018 (Figure 13; Obu et al., 2020). As warming increases, so will loss of permafrost (Section 1.5.2). Global projections show very limited permafrost in the region already at $+2^{\circ}\text{C}$ (Chadburn et al., 2017), but global projection (including Chadburn et al., 2017) do not account for peatland permafrost, which can persist for centuries outside of its climate equilibrium (Osterkamp and Romanovsky, 1999). Much of the permafrost in Baltic Sea region was very close to its climatic boundary even before the recent acceleration of climate warming. Much of the lowland permafrost in palsas and peat plateaus in this region is very close to the 0°C thawing point, and is likely relict permafrost, persisting from the Little Ice Age (Sannel et al., 2016). Observations also show that lowland permafrost thaw has been going on for decades (Åkerman and Johansson, 2008). Preliminary analyses of permafrost loss in 1997-2018 suggests that this was roughly equally divided between alpine and lowland permafrost (22 and 24%, respectively, Figure 13), in agreement with projections of loss of all types of Baltic permafrost in the future.

Permafrost thaw by climate warming is known to affect river runoff and its loads of carbon, nutrients and contaminants, such as mercury (Schuster et al., 2018; Vonk et al., 2015). The local effect of permafrost thaw in alpine headwaters can be significant (Lyon et al., 2009), but in the Baltic Sea Basin alpine permafrost thaw will likely have limited influence on the characteristics of river transport at their mouths on the Baltic Sea. This is because the alpine permafrost in the Baltic Sea drainage basin mainly affects solid bedrock or regolith, with almost no soil organic matter stored in permafrost (Fuchs et al., 2015). Thaw of permafrost in peatlands affects soils with very large stocks of organic material, and has been suggested to cause large losses of peat carbon and nutrients into aquatic ecosystems (Hugelius et al., 2020). However, these projections are highly uncertain and based on studies of peatland thaw chronosequences in North America that may not be applicable to Fennoscandian permafrost peatlands (though see Tang et al., 2018).

The extent of permafrost in the Baltic Sea drainage basin may decrease significantly in this century, and depending on which warming trajectory the Earth takes, may disappear altogether in the coming century. The thaw of alpine permafrost will have little effect on flows of water, carbon and nutrients to the Baltic Sea. Thawing peatlands may increase the loads of carbon, nutrients and mercury to the Baltic Sea, but these projections remain highly uncertain.



2637 **3.3.4.4 Sea ice**

2638 Two new projections for sea ice in the Baltic Sea have been produced after BACC II (BACC II Author Team,
 2639 2015). Luomaranta et al. (2014) used simplified regression and analytical models to estimate changes in sea-ice
 2640 extent (Fig. 28) and fast-ice thickness. Due to their less demanding computational approach, they could base
 2641 estimates on 28 CMIP5 models. As in the Arctic Ocean (Section 1.5.2), maximum annual ice extent and thickness
 2642 were both estimated to decline in the future, but some sea ice will still form every year, even by the end of the
 2643 century, in agreement with earlier studies (e.g. Haapala et al., 2001; Meier, 2002; Meier et al., 2004a). Under the
 2644 RCP4.5 and RCP8.5 scenarios, the modelled mean maximum ice thicknesses in Kemi were projected to be 60 cm
 2645 and nearly 40 cm, respectively, in 2081–2090. However, under the RCP8.5 scenario, two models projected Kemi
 2646 to be ice-free.

2647
 2648 Höglund et al. (2017) used a more advanced approach to examine changes in sea ice conditions with a coupled
 2649 ice-ocean model (Hordoir et al., 2019; Pemberton et al., 2017). They used downscaled atmospheric data from the
 2650 EC-Earth and the Max Planck Institute Earth System models and simulated the response of the ice for the RCP4.5
 2651 and RCP8.5 projections. Average annual maximum ice extent at the end of the century was projected to be 90 –
 2652 100 10³ km² and 30 – 40 10³ km², for the medium and high emission scenarios, respectively, and ice thickness to
 2653 decrease 3 – 6 cm decade⁻¹. Höglund et al. (2017) also projected the mobility of the ice to increase, but with little
 2654 effect on future ridged ice production.

2655 **3.3.4.5 Lake ice**

2656 The latest model experiments demonstrate that the Baltic Sea catchment will experience a substantial reduction in
 2657 lake ice cover in the future, with many lakes becoming ice covered only intermittently (Maberly et al., 2020;
 2658 Sharma et al., 2019; Sharma et al., 2021; Shatwell et al., 2019). This change will commence in the south and move
 2659 northwards gradually. Lithuanian and Latvian lakes will lose their ice cover after +2°C warming, and further
 2660 warming will gradually move winter ice loss northwards, so that at +8°C warming, only lakes in northernmost
 2661 Lapland will retain a winter ice cover (Maberly et al., 2020; Sharma et al., 2019; Sharma et al., 2021; Shatwell et
 2662 al., 2019).

2663 **3.3.5 Ocean and marine sediments**

2664 **3.3.5.1 Water temperature**

2665 Ocean temperatures are rising at accelerating rates (IPCC, 2019a; Section 1.5.3.2). For the end of this century,
 2666 scenarios for the Baltic Sea project a sea surface temperature increase of 1.1°C (0.8–1.6°C, RCP2.6) to 3.2°C (2.5–
 2667 4.1°C, RCP8.5) compared to 1976–2005 (Gröger et al., 2019; Gröger et al., 2021b), see Table 9. In brackets, the
 2668 ensemble spreads indicated by the 5th and 95th percentiles are listed. These changes are slightly larger than the
 2669 projected global sea surface temperature changes (Section 1.5.3.2). Other ensembles than the one by Gröger et al.
 2670 (2019) give similar results that vary between 1.9°C (RCP4.5) and 2.9°C (RCP8.5) for the ensemble mean
 2671 temperature increase (Meier et al., 2021b; see also Meier and Saraiva, 2020). By the end of the century, sea surface
 2672 temperature changes for the RCP8.5 scenarios significantly exceed natural variability. Largest open-sea warming
 2673 is found in summer in the northern Baltic Sea, due to earlier melting of the sea ice (Figs. 29 and 30). Even higher



2674 warming of +2–6°C (the range denotes RCP2.6 and RCP8.5 scenarios) is projected for the Curonian Lagoon by
 2675 the year 2100 (Jakimavičius et al., 2018).

2676

2677 The main driver of interannual variations of monthly mean sea surface temperature is air temperature, through the
 2678 sensible heat fluxes (Meier et al., 2021a). The second most important drivers are cloudiness over the open sea and
 2679 latent heat and meridional and zonal wind velocities over coastal areas, the latter probably because of upwelling
 2680 (Meier et al., 2021a). In the vertical, the surface layer is warming more than the winter water, which is sandwiched
 2681 between the surface layer and the halocline. Hence, the spring and summer thermoclines are getting more intense
 2682 (Gröger et al., 2019). Water temperature trends in the deep water of those sub-basins such as Bornholm Basin and
 2683 Gotland Basin that are sporadically ventilated by salt water inflows originating from surface water are projected
 2684 to be elevated as well (Meier et al., 2021a). Projected changes of the vertical water temperature distribution are
 2685 similar than those observed since 1850 (Kniebusch et al., 2019a).

2686

2687 For extreme events, projections suggest, inter alia, more tropical nights over the Baltic Sea, increasing the risk of
 2688 record-breaking water temperatures (Meier et al., 2019a).

2689 3.3.5.2 Salinity and saltwater inflows

2690 Future changes in salinity will depend on changes in the wind fields over the Baltic Sea region (Lass and Matthäus,
 2691 1996), river runoff from the Baltic Sea catchment (Schinke and Matthäus, 1998) and mean sea level rise relative
 2692 to the seabed of the sills in the entrance area (Meier et al., 2017; Meier et al., 2021b). A projected increase in river
 2693 runoff will tend to decrease salinity, but sea level rise will have the opposing effect of tending to increase salinity,
 2694 because the water level above the sills at the Baltic Sea entrance would be higher, increasing the cross-sectional
 2695 area of the Danish straits. As a result, saltwater imports from Kattegat would be larger. A 0.5 m higher sea level
 2696 relative to the sill bottom at the end of the century would increase estimated Gotland Deep surface salinity by 0.7
 2697 g kg⁻¹ and bottom salinity by 0.9 g kg⁻¹ (Meier et al., 2017; Meier et al., 2021b). Due to the large uncertainty in
 2698 projected changes in wind fields over the Baltic Sea region (Section 3.3.1.5), in changes of the freshwater supply
 2699 from the catchment (section 3.3.2.1) and in global sea level rise (Section 3.3.5.4), salinity projections show a wide
 2700 spread. No robust changes were identified because the two main drivers, river runoff and sea level rise,
 2701 approximately compensate each other (Meier et al., 2021b). According to Saraiva et al. (2019b) river runoff would
 2702 increase by about 1 to 21% at the end of the century depending on the climate model under both RCP4.5 and
 2703 RCP8.5, in the ensemble mean causing a decrease in surface and bottom salinity at Gotland Deep of about 0.6–0.7
 2704 g kg⁻¹, with a large spread among the ensemble members. Assuming a negligible global sea level rise, the intensity
 2705 and frequency of MBIs were projected to slightly increase due to changes in the wind fields (Schimanke et al.,
 2706 2014). Hence, in ensemble studies that considered all potential drivers, no significant change in salinity were
 2707 projected as the ensemble mean (Meier et al., 2021b). In case of salinity, global climate model uncertainty was
 2708 identified to be the largest of all uncertainties (Meier et al., 2021b).

2709 3.3.5.3 Stratification and overturning circulation

2710 Model based estimations of future stratification are still rare and depend critically on how well the models project
 2711 changes in the three-dimensional distributions of temperature and salinity. A first systematic attempt using a high
 2712 resolution coupled ocean - atmosphere model and five different global climate models (Gröger et al., 2019)



2713 explored future stratification under RCP8.5. They assumed a 10% increase in river runoff (approximately the
 2714 ensemble mean in Saraiva et al., 2019b) and an unchanged mean sea level in the North Sea at the end of the
 2715 century. The ensemble consistently indicated a basin-wide intensification of the pycnocline (by 9–35%) for nearly
 2716 the whole Baltic Sea, and a shallowing of the pycnocline depth in most regions, except the Gulf of Bothnia (Gröger
 2717 et al., 2019). The area with a pycnocline intensity $> 0.05 \text{ kg m}^{-3}\text{m}^{-1}$ increased 23-100%. The warm season
 2718 thermocline likewise intensified in nearly the entire Baltic Sea (Gröger et al., 2019).

2719
 2720 All ensemble members indicate a strengthening of the zonal, wind driven near-surface overturning circulation in
 2721 the southwestern Baltic Sea towards the end of the 21st century, whereas the zonal overturning at depth is reduced
 2722 by ~ 25% (Gröger et al., 2019). In the Baltic proper, the meridional overturning shows no clear climate change
 2723 signal. However, three out of five ensemble members indicate at least a northward expansion of the main
 2724 overturning cell. In the Bothnian Sea, all ensemble members show a significant weakening of the meridional
 2725 overturning.

2726
 2727 As the study by Gröger et al. (2019) and previous projections (e.g. Meier et al., 2006) do not consider global sea
 2728 level rise, these scenario simulations are no longer considered plausible (Meier et al., 2021a; 2021b). Considering
 2729 all drivers of changes in salinity in the Baltic Sea (wind, river runoff, global sea level rise), neither the haline
 2730 induced stratification nor the overturning circulation is projected to change systematically among climate models
 2731 (Meier et al., 2021a). It was found that under a RCP4.5 or RCP8.5 scenario a linearly rising mean sea level by the
 2732 figures suggested by IPCC (2019b) would approximately counteract the effects of projected river runoff increases
 2733 and wind changes on salinity.

2734 3.3.5.4 Sea level

2735 Global mean and thus Baltic Sea level will continue to rise at an increasing rate. During this century, melting ice
 2736 sheets in Antarctica and Greenland are expected to contribute more to the total sea level than in the past (e.g.
 2737 Mitrovica et al., 2018). The fingerprints from melting ice sheets in Antarctica on sea level rise will be more
 2738 pronounced in the northern hemisphere and introduce large uncertainties for Baltic sea level rise. Furthermore, the
 2739 sea level in shelf seas such as the Baltic Sea will rise more strongly than one would expect from the thermostatic
 2740 expansion of the local water column only, due to spill-over effects from the open ocean (Landerer et al., 2007;
 2741 Bingham and Hughes, 2012). In addition, the long-term rate of coastal land rise is not easy to estimate accurately,
 2742 due to the limited length of Global Positioning System (GPS) measurements, and frequently revised geological
 2743 model values.

2744
 2745 Estimates for the ensemble mean global sea level rise by 2100 ranged from 43 cm (RCP2.6) to 84 cm (RCP8.5),
 2746 with likely ranges of 29-56 cm and 61-110 cm, respectively (IPCC, 2019a), cf. Section 1.5.3.1. In particular for
 2747 RCP8.5, sea level rise projections by the fifth IPCC assessment report (IPCC, 2014a) somewhat differ from the
 2748 more recent Special Report on the Ocean and Cryosphere in a Changing Climate (IPCC, 2019a) because of the
 2749 updated contribution from Antarctica based upon new ice-sheet modeling.

2750
 2751 The projected sea level rise relative to land in the Baltic Sea entrance area was estimated to about 80% of the
 2752 global rate (Grinsted et al., 2015; Grinsted, 2015). These results were confirmed by other studies (e.g. Kopp et al.,



2014) and summarized by Pellikka et al. (2020) who found a regional ensemble mean absolute sea level rise of 87% of the global sea level rise. Altogether, considering land uplift and eustatic sea level rise, very likely ranges (5-95% probability) of relative sea level change under the most pessimistic IPCC emissions scenario (RCP8.5) were projected to, e.g. 29-162 cm in Copenhagen (median 68 cm), -13 -117 cm in Stockholm (median 25 cm), and 21-151cm in St. Petersburg (median 59 cm) (Grinsted et al., 2015). For coastal sites in the northern Baltic Sea, relative sea level changes in the Gulf of Finland in 2000–2100 were projected to be +29 cm (–22 to+92 cm), –5 cm (–66 to+65 cm) for the Bothnian Sea, and –27 cm (–72 to +28 cm) for the Bothnian Bay, where the land uplift is larger (Johansson et al., 2014). The ranges in the latter study were estimated from the 5% and 95% cumulative probabilities considering several published scenarios from the third and fourth IPCC assessment reports. In a recent study based upon IPCC (2019a), Hieronymus and Kalén (2020) also estimated a sea-level fall in the northern Baltic Sea and a 70 cm rise in the south by 2100. These upper bounds of the sea level rise projections imply a very strong future acceleration of present rates. Current observations seems to show an acceleration, but its present magnitude is still small (Hünicke and Zorita, 2016; see Section 3.2.5.4).

Recent efforts since the IPCC AR5 report (IPCC, 2014a) that focused on the contribution of Antarctic ice sheets to global mean sea level rise have shown that the interaction of warming ocean water, melting the ice sheets from below can lead to instabilities in the ice sheet dynamics. The ice sheets flowing from land into the ocean are in contact with the ocean floor out to the grounding line. From there on outward the ocean is melting the ice from below and the ice sheets become thinner and lighter. If the weight of the ice sheet becomes less than the weight of the ocean water it replaces, it floats up and away. The grounding line retreats inland where the ice sheet is thicker and the ice flow larger and reinforces the ice loss (Mercer, 1978). This and related feedback loops could lead to an extra meter of sea level rise until the end of the century (e.g. Sweet et al., 2017). The most recent estimates based on expert judgement (Bamber et al., 2019) for global mean sea level rise in 2100 relative to 2000, including these potential contributions (including land water storage) are 69 cm and 111 cm for low and high sea level scenarios, respectively. For the high sea level scenario the likely range (5 to 95%) is between 62 cm and 238 cm.

Future changes in sea level extremes in the Baltic Sea depend on future changes in mean sea level and future developments in large-scale atmospheric conditions associated with changing wind patterns. Model projections disagree regarding atmospheric circulation changes and therefore their relevance for extreme future sea levels remains unclear (Räisänen, 2017). Absolute mean sea levels will continue to rise in the entire Baltic Sea, but exact rates remain uncertain and depend on models and greenhouse gas emission scenarios (Grinsted, 2015; Hieronymus and Kalén, 2020). Relative sea level changes will strongly vary across the Baltic Sea because of the existing spatial gradient in glacial isostatic adjustment and the spatial inhomogeneity associated with the uncertain relative contributions of melting from Antarctica and Greenland (e.g. Hieronymus and Kalén, 2020). For the Baltic Sea, changing mean sea levels are expected to have larger effects on future extremes than changing atmospheric circulation (Gräwe and Burchard, 2012). Sea ice loss in the future will further directly expose the northern Baltic coastline to stronger storm surges.

Recent projections of extreme sea levels along Europe’s coasts have considered all drivers by linear superposition, i.e. absolute mean sea level rise and land uplift, tides (small in the Baltic Sea), storm surges and waves (Vousdoukas et al., 2016; Vousdoukas et al., 2017). The results suggest that extreme sea levels will increase more



than the mean sea level, due to small changes in the large-scale atmospheric circulation, such as a northward shift of the Northern Hemisphere storm tracks and westerlies, and increases in the NAO/AO (IPCC, 2014b). These changes in the large-scale atmospheric circulation of the Baltic Sea region are, however, not robust among GCMs, giving the projections of extreme sea levels by Vousdoukas et al. (2016; 2017) low confidence.

3.3.5.5 Waves

The few existing wave climate projections for the Baltic Sea indicate an increase in the mean wave conditions, either in the whole area (Groll et al., 2017) or in its northern part (Bonaduce et al., 2019). This increase in the mean conditions has been linked to two main drivers: 1) increased wind speeds and 2) reduced seasonal ice cover.

Groll et al. (2017) projected wave climate at the end of 21st century, based on two different scenarios. They found a slight increase in the median wind speeds for most of the Baltic Sea area, which led to an increase of up to 15 % in median Significant Wave Height (SWH). Using only one climate scenario, Bonaduce et al. (2019) found that decreased wind speed in the southern Baltic Sea led to a decrease in mean SWH, whereas increased wind speeds in the north, especially in winter, led to increased mean SWH. As neither study used multi-model ensembles of scenario simulations (an exception for the western Baltic Sea is the work by Dreier et al. (2021), and there is high uncertainty in the projected wind speeds and directions, which is not attributed to the decline in ice cover, the results may not be representative. The projected changes in SWH estimates are therefore inconclusive.

Ruosteenoja et al. (2019) estimated based CMIP5 simulations that in future, mean and extreme scalar wind speeds are not likely to significantly change in the Baltic Sea area. Hence, mean wave conditions would not change. They also estimated that frequency of strong westerly winds will increase while strong easterly winds will become less common. These type of changes might have more significance on the frequency of extreme SWH values and their spatial patterns.

For extreme values, these studies give even less reliable results. The results of Groll et al. (2017), Suursaar et al. (2016) and Bonaduce et al. (2019) all indicated large spatial variability in how the projected extremes changed. In addition to the wind speed, extreme values are quite sensitive to wind direction, since fetch varies with direction due to the geometry of the Baltic Sea. Mäll et al. (2020) simulated how wave conditions during three historical Baltic Sea storms would change under climate change conditions. The results showed slight, but not significant changes in extreme SWH values during the storms.

Future changes in seasonal sea-ice conditions in the northern Baltic Sea are more reliable, and their effect on the wave climate easier to estimate (Rutgersson et al., 2021). Mild ice winters have already become common, and new records of lowest annual maximum ice extent have been recorded. In the Baltic Sea, the ice season partly overlaps with the seasons of the strongest winds, namely autumn and winter. The mean and extreme values of SWH are therefore expected to increase in areas like the Bothnian Sea, which now typically has ice cover in winter, but will have loose it in the future Baltic Sea climate.



2831 3.3.5.6 Sedimentation and coastal erosion

2832 As a consequence of the probably accelerating sea level rise, coastal erosion will increase regionally, to fill the
 2833 increased underwater accommodation space. How much erosion will increase will depend not only on the rate of
 2834 sea level rise, but also on the intensity of storms (Zhang et al., 2017).

2835
 2836 Coastal erosion, accretion and alongshore sediment transport are primarily controlled by winds and wind-induced
 2837 waves in the Baltic Sea. Projecting the future rate of coastal erosion or accretion in the Baltic Sea is highly
 2838 uncertain because of a lack of consensus in the prediction of future storms. Neglecting potential change in future
 2839 storms and assuming an intermediate sea level rise scenario (RCP4.5), an increment of 0.1-0.3 m year⁻¹ in coastline
 2840 erosion has been projected for some parts of the southern Baltic Sea coast (Zhang et al., 2017; Deng et al., 2015).
 2841 Due to the prevailing westerly winds, the dominant sediment transport will continue to be eastwards along most
 2842 of the southern Baltic Sea coast, but with high variability along coastal sections with a small incidence angle of
 2843 incoming wind-waves (Dudzińska-Nowak, 2017). It has been found that even a minor climate-change-driven
 2844 rotation of the predominant wind directions over the Baltic Sea may substantially alter the structural patterns and
 2845 pathways of wave-driven transport along large sections of the coastline (Viška and Soomere, 2013).

2846
 2847 The presence of sea ice is an important factor moderating coastal erosion. Storm surges and wave run-up on the
 2848 beach are much higher in ice-free periods than when there is even partial ice cover. The hydrodynamic forces are
 2849 particularly effective in reshaping the shoreline when there is no ice and sediment is mobile (Ryabchuk et al.,
 2850 2011). Due to global warming, both the area and duration of ice cover in the Baltic Sea will be reduced in future
 2851 (Section 3.3.4.4), thus increasing coastal erosion.

2852
 2853 Foredunes will likely continue to form on prograding coasts, but at rates influenced by the accelerating sea-level
 2854 rise (Zhang et al., 2017). Foredunes may tend to become higher, but with reduced prograding rate and wavelength,
 2855 if sea level rise is accelerated or storm frequency increased. If the wind-wave climate is stable, the height of coastal
 2856 foredunes on a prograding coast remains stable or increases linearly with a low to intermediate rate (<1.5 mm year⁻¹)
 2857 of sea level rise. An accelerating rate of sea level rise and/or changing storm frequency will lead to a nonlinear
 2858 growth in height (following a quadratic or a higher power law; Zhang et al., 2017; Lampe and Lampe, 2018). The
 2859 critical threshold that separates linear and non-linear foredune growth in response to sea level rise is likely to be
 2860 reached before 2050 in the RCP8.5 scenario (Zhang et al., 2017).

2861
 2862 Anthropogenic influence imposes further uncertainty in sediment transport and coastal erosion. Sediment transport
 2863 and coastal erosion are relevant for coastal management, construction and protection strategies. In general two
 2864 main types of management strategies exist for the Baltic Sea coast: 1) coastal protection by soft or hard measures;
 2865 and 2) adaptation to coastal change, accepting that in some places the coast would be left in its natural state (BACC
 2866 II Author Team, 2015). However, administrative efforts for coastal protection differ among Baltic Sea countries,
 2867 even between neighboring states or nations. It has been found that engineering structures (e.g. piers, seawalls) may
 2868 influence coastline change at a much larger spatial scale than the dimension of the structure itself.



2869 3.3.5.7 Marine carbonate system and biogeochemistry

2870 The BACC II Author Team (2015) concluded that model simulations indicated that climate change has a potential
 2871 to intensify eutrophication in the Baltic Sea. However, they also showed that the implementation of nutrient load
 2872 reductions according to the Baltic Sea Action Plan (BSAP, HELCOM, 2013a) may not only mitigate this effect
 2873 but may slightly decrease hypoxic and anoxic areas in the Baltic Sea, because the nutrient load abatement strategy
 2874 of the BSAP did not take the effect of climate change into account. In contrast, the business as usual nutrients
 2875 input scenario may increase by about 30% hypoxic area and even more than double the area affected by anoxia by
 2876 2100.

2877
 2878 As atmospheric CO₂ rises, so will the concentration of CO₂ in the Baltic Sea surface water. This will influence the
 2879 mean future pH, while eutrophication and enhanced organic matter production/remineralization will increase the
 2880 amplitude of daily and seasonal pH fluctuations without much affecting the mean values. It was also shown that
 2881 pH in the Baltic Sea surface water will decrease, in the worst-case emission scenario (atmospheric pCO₂ of 850
 2882 ppm) the pH will drop by about 0.40 by 2100, while the decrease in a more optimistic emission scenario (550 ppm)
 2883 will be smaller, about 0.26.

2884 3.3.5.7.1 Oxygen and nutrients

2885 Projected warming and global mean sea level rise may worsen eutrophication and oxygen depletion in the Baltic
 2886 Sea by reducing air-sea fluxes and vertical transports of oxygen in the water column, intensifying internal nutrient
 2887 cycling, and increasing river-borne nutrient loads due to increased river runoff (Meier et al., 2011a; Meier et al.,
 2888 2012b; Meier et al., 2012c). However, the future response of deep-water oxygen conditions will depend mainly on
 2889 nutrient loads from land (Saraiva et al., 2019a, b; Meier et al., 2021b; cf. Fig. 31). In contrast to the global ocean
 2890 (see Section 1.5.4), future nutrient supplies will have a relatively larger effect on oxygen conditions and primary
 2891 production than warming. With high nutrient loads, the changing climate will have a considerable negative effect,
 2892 but if loads are kept low, climate effects can be small or negligible. Scenario simulations suggest that full
 2893 implementation of the nutrient load reductions required by BSAP will significantly improve the eutrophication
 2894 status of the Baltic Sea, irrespective of the driving global climate model (Saraiva et al., 2019b; Meier et al., 2021b)
 2895 and regional coupled climate-environmental model (Meier et al., 2018a). Despite large uncertainties of future
 2896 projections, modeling studies suggested that the future Baltic Sea ecosystem may unprecedentedly change
 2897 compared to the past 150 years (Meier et al., 2012a).

2898
 2899 By the end of the century (2069-2098), the ensemble mean hypoxic area is projected to change only slightly under
 2900 reference (-14% for RCP4.5 and -5% for RCP8.5) and high (-2% for RCP4.5 and +5% for RCP8.5) nutrient load
 2901 scenarios, compared to 1976-2005 (Saraiva et al., 2019b). Nutrient loads in the reference scenario are the average
 2902 loads in 2010-2012. The high, or worst, scenario assumes changes caused by a ‘fossil-fuelled development’,
 2903 coupled to increasing river runoff (Saraiva et al., 2019a). Changes in nitrogen and phosphorus loads were estimated
 2904 from assumptions on regional population growth, changes in agricultural practices, such as land and fertilizer use,
 2905 and developments in sewage treatment (Zandersen et al., 2019; Pihlainen et al., 2020). Under the BSAP scenario,
 2906 the ensemble mean hypoxic area will be reduced by 50-60% at the end of the century, in comparison with 1976-
 2907 2005 (Saraiva et al., 2019a). The relative reductions in hypoxic area may decrease with increasing sea level (Meier
 2908 et al., 2021b).



2909

2910 In the same model ensemble (Saraiva et al., 2019a), BSAP implementation is projected to reduce the water column
 2911 phosphate pool in the Baltic Sea by 59% (RCP4.5) and 56% (RCP8.5) by the end of the century, and even the
 2912 reference loads would lead to a decline by 24% (RCP4.5) and 18% (RCP8.5). Also, a larger ensemble (Meier et
 2913 al., 2018a) of 8 biogeochemical models forced by outputs of 7 ESMs downscaled by 4 different RCMs projected
 2914 that the BSAP reduced phosphate concentrations in the Baltic proper, Gulf of Finland and Bothnian Sea despite
 2915 climate change, with largest reductions in surface concentrations by approximately 3 mmol m^{-3} in the Gulf of
 2916 Finland. Present day nutrient loads led to small increases in surface phosphate concentration in the Baltic proper,
 2917 a small decline in the Gulf of Finland and little change in Bothnian Sea and Bothnian Bay. Little change was
 2918 predicted for DIN concentrations in the Baltic proper, whereas simulations showed an increase in the Gulf of
 2919 Finland and the Bothnian Sea, regardless whether nutrient loads were kept at present level or whether loads were
 2920 reduced.

2921

2922 Furthermore, future projections suggested that the sea-ice decline in the northern Baltic Sea may have considerable
 2923 consequences for the marine biogeochemistry, because of changing underwater light conditions and wave climate
 2924 (Eilola et al., 2013). Eilola et al. (2013) found that, by the end of the century, the spring bloom would start by up
 2925 to one month earlier and winds and wave-induced resuspension would increase, causing an increased transport of
 2926 nutrients from the productive coastal zone into the deeper areas.

2927

2928 For the Baltic proper, the internal nutrient cycling and exchanges between shallow and deeper waters were
 2929 projected to be intensified, and the internal removal of phosphorus may become weaker in future climate (Eilola
 2930 et al., 2012). These effects may counteract the efforts of planned nutrient input reductions.

2931

2932 Uncertainties in projections from Baltic Sea ecosystem models have recently been systematically assessed for the
 2933 first time (Meier et al., 2018a; Meier et al., 2019b; Meier and Saraiva, 2020; Meier et al., 2021b). One of the larger
 2934 sources of uncertainty is biases in global and regional climate models, in particular concerning global mean sea
 2935 level rise and regional water cycling (Meier et al., 2019b). The mechanism behind the correlation between large-
 2936 scale meteorological conditions in the different climate periods and oxygen conditions in the Baltic Sea is not well
 2937 understood and subject to ongoing research. With respect to nutrient concentrations, also uncertainties in
 2938 conditions at the North Sea boundary as well as difficulties in simulating the long-term response of the Baltic Sea
 2939 biogeochemical system to changes in nutrient inputs, play a role.

2940

2941 Under the BSAP scenario, mean nitrogen fixation would decrease (Meier et al., 2021b) and record-breaking
 2942 cyanobacteria blooms may no longer occur in the future, but record-breaking events may reappear at the end of
 2943 the century in a business-as-usual nutrient load scenario (Meier et al., 2019a).

2944 **3.3.5.7.2 Marine CO₂ system**

2945 The rising atmospheric pCO₂ due to anthropogenic emissions will increase the mean pCO₂ of surface seawater and
 2946 thus has the potential to lower the pH. However, the magnitude of pH changes will also depend on the development
 2947 of total alkalinity concentrations (A_T; Omstedt et al., 2012). Future A_T changes in the Baltic Sea will be shaped by
 2948 both external inputs (riverine runoff and inflows from the North Sea) and internal generation. The latter is due to



biogeochemical processes of organic matter production and remineralization, especially under euxinic conditions. Kuznetsov and Neumann (2013), who used A_T as a tracer in a model (no internal processes included) showed that on average A_T in surface Baltic Sea waters should decrease by about $150 \mu\text{mol kg}^{-1}$ by 2100, a change corresponding to an assumed decrease in salinity. Simulations by Gustafsson et al. (2019) that include most of the biogeochemical processes affecting A_T (except S burial and Fe-oxide availability) showed that A_T in the central Baltic Sea in the “business as usual” scenario will first increase, by about $100 \mu\text{mol kg}^{-1}$ by 2050, and then revert to present levels by 2100. If BSAP is implemented, A_T will decrease by about $150 \mu\text{mol kg}^{-1}$ in 2100 from present levels. Irrespective of the nutrient load scenario, pH is eventually expected to decrease in the central Baltic Sea due to anthropogenic CO_2 emissions. Assuming the A1B CO_2 emission scenario (pCO_2 increase to $700 \mu\text{atm}$ by 2100), pH will drop to about 7.9 and 7.8 under “business as usual” and BSAP scenarios, respectively.

3.3.6 Marine biosphere

3.3.6.1 Pelagic habitats

3.3.6.1.1 Microbial communities

The effects of climate change on microbes and the functioning of the microbial loop have been studied by experiments in which temperature, salinity, dissolved organic matter (DOM), and ocean acidification (OA) were manipulated. In general, microbial activity and biomass increased with increasing DOM and temperature (Ducklow et al., 2009), but effects can be mixed. For instance, an increase in DOM in the northern Gulf of Bothnia enhanced the abundance of bacteria, whereas a temperature increase (from 12 to 15°C) decreased their abundance, probably due to a simultaneous increase of bacterivorous flagellates (Nydahl et al., 2013).

In the southern Baltic Sea the impact of OA was limited, and the bacterial community responded primarily to temperature and phytoplankton succession (Bergen et al., 2016). In experiments where CO_2 was increased and salinity decreased (from 6 to 3), heterotrophic bacteria declined (Wulff et al., 2018). In experiments with increasing temperature (from 16 to $18\text{--}20^\circ\text{C}$) and reduced salinity (from 6.9 to 5.9 g kg^{-1}), the Baltic proper microbial community also showed mixed responses, probably due to indirect food web effects (Berner et al., 2018).

3.3.6.1.2 Phytoplankton and cyanobacteria

The projected increase in precipitation is expected to increase nutrient loads, especially into the northern Baltic Sea (Huttunen et al., 2015), and together with increased internal loading of nutrients, several modelling studies project an increased phytoplankton biomass by the end of the century (Meier et al., 2012b; Meier et al., 2012c; Skogen et al., 2014; Ryabchenko et al., 2016).

Several mesocosm studies have investigated the effects of warming on southern Baltic Sea phytoplankton communities. Warming accelerated the phytoplankton spring bloom and increased primary productivity (Sommer and Lewandowska, 2011; Lewandowska et al., 2012; Paul et al., 2015). The total phytoplankton biomass still decreased, due to negative effects of warming on nutrient flux (Lewandowska et al., 2012; Lewandowska et al., 2014).



2986 Ocean acidification (OA) may enhance phytoplankton productivity by increasing the CO₂ concentration in the
 2987 water. The biomass of southern Baltic Sea autumn phytoplankton increased in mesocosms simulating OA
 2988 (Sommer et al., 2015). In many experiments OA had, however, little effect on phytoplankton community
 2989 composition, fatty acid composition or biovolume in spring or autumn (Paul et al., 2015; Bermúdez et al., 2016;
 2990 Olofsson et al., 2019).

2991
 2992 It has been suggested that climate change may increase the blooming of toxic species, such as the dinoflagellate
 2993 *Alexandrium ostenfeldii* (Kremp et al., 2012; Kremp et al., 2016) and the cyanobacterium *Dolichospermum* sp.
 2994 (Brutemark et al., 2015; Wulff et al., 2018). There are also contradictory results, indicating that OA and warming
 2995 may decrease the biomass of *Nodularia* sp. and *Dolichospermum* sp. (Eichner et al., 2014; Berner et al., 2018).
 2996 Several modelling studies project increases in cyanobacteria in the warmer and more stratified future Baltic Sea
 2997 (Meier et al., 2011b; Andersson et al., 2015; Neumann et al., 2012; Chust et al., 2014; Hense et al., 2013), but
 2998 other modelling studies project that the environmental state of the Baltic Sea will be significantly improved, and
 2999 extreme cyanobacteria blooms will no longer occur if BSAP is fully implemented (Meier et al., 2018a; Meier et
 3000 al., 2019a; Saraiva et al., 2019a; see Figure 32).

3001 3.3.6.1.3 Zooplankton

3002 The effects of increasing temperature and ocean acidification (OA) on zooplankton have been studied
 3003 experimentally. In *Acartia* sp., a dominant copepod in the northern Baltic Sea, warming decreased egg viability,
 3004 nauplii development and adult survival, and both warming and OA had negative effects on adult female size
 3005 (Garzke et al., 2015; Vehmaa et al., 2016; Vehmaa et al., 2013).

3006
 3007 In contrast, the effects of climate change on microzooplankton (MZP) seem to be mostly beneficial. Warming
 3008 improved the growth rate of southern Baltic Sea MZP, which led to a reduced time-lag between phytoplankton
 3009 and MZP maxima, improving the food supply to microzooplankton in warm conditions (Horn et al., 2015). (Aberle
 3010 et al., 2015) showed that while protozooplankton escaped predation by slower growing copepods at low
 3011 temperatures, at warmer temperatures small ciliates in particular became more strongly controlled by copepod
 3012 predation.

3013 3.3.6.2 Benthic habitats

3014 3.3.6.2.1 Macroalgae and vascular plants

3015 The effects of climate change on bladder wrack, *Fucus vesiculosus*, have been studied in a number of experiments.
 3016 Ocean acidification (OA) appears to have a relatively small effect on macroalgae (Al-Janabi et al., 2016; Wahl et
 3017 al., 2020), while temperature effects can be significant. The effects of increasing temperature are not linear,
 3018 however. Growth or photosynthesis is not impaired under projected temperature increase (from 15 to 17.5°C) but
 3019 at extreme temperatures (27 to 29°C), photosynthesis declines, growth ceases and necrosis starts (Graiff et al.,
 3020 2015; Takolander et al., 2017). In very low salinity (2.5 g kg⁻¹), sexual reproduction of *F. vesiculosus* ceases
 3021 (Rothäusler et al., 2018).

3022
 3023 The direct and indirect effects of changes in temperature, salinity and pH may alter the geographic distribution of
 3024 many species in the Baltic Sea. Retreat of marine species has been predicted for bladder wrack, eelgrass and blue



3025 mussel, and up to 50 other species affiliated to these keystone species (Vuorinen et al., 2015). Species distribution
 3026 modelling has indicated that a decrease of bladder wrack will have large effects on the biodiversity and functioning
 3027 of the shallow-water communities of the northern Baltic Sea (Jonsson et al., 2018; Kotta et al., 2019). The
 3028 responses of eelgrass, *Zostera marina*, to climate change and eutrophication mitigation have recently been modeled
 3029 by Bobsien et al. (2021).

3030
 3031 Experiments on climate change effects have been made also with other macroalgae and vascular plants. Thus, OA
 3032 increased the growth of the opportunistic green alga *Ulva intestinalis* in the Gulf of Riga (Pajusalu et al., 2013;
 3033 Pajusalu et al., 2016). Other studies showed that charophyte photosynthesis increased under high pCO₂, whereas
 3034 eelgrass did not respond to the elevated pCO₂ alone (Pajusalu et al., 2015). Salinity decline is projected to decrease
 3035 the distribution of *Z. marina* and the red alga *Furcellaria lumbricalis*, whereas warming will probably favour
 3036 charophytes (Torn et al., 2020).

3037 3.3.6.2.2 Zoobenthos

3038 The effects of warming on invertebrates are non-linear. Respiration and growth of the isopod *Idotea balthica*
 3039 increased up to 20°C, and then decreased at 25°C (Ito et al., 2019). Many marine invertebrates, including isopods,
 3040 will also directly and indirectly suffer from decreasing salinity (Kotta et al., 2019; Rugiu et al., 2017), as well as
 3041 ocean acidification (OA). The size and time to settlement of the pelagic larvae of the Baltic clam *Limecola balthica*
 3042 (syn *Macoma balthica*) increased with OA, suggesting a developmental delay (Jansson et al., 2016), whereas OA
 3043 had no effect on the isopod *Saduria entomon* (Jakubowska et al., 2013) or larvae of the barnacle *Amphibalanus*
 3044 *improvisus* (Pansch et al., 2012).

3045
 3046 Several modelling studies have estimated the relative effects of hydrodynamics, oxygen and food availability on
 3047 Baltic Sea zoobenthos. In previously hypoxic areas, benthic biomass was projected to increase (until 2100) by up
 3048 to 200% after re-oxygenating bottom waters, whereas in permanently oxygenated areas macrofauna may decrease
 3049 by 35% due to lowered food supply to the benthic ecosystem (Timmermann et al., 2012). It has, however, been
 3050 concluded that nutrient reductions will be a stronger driver for Baltic Sea ecosystem than climate change (Friedland
 3051 et al., 2012; Niiranen et al., 2013; Ehrnsten et al., 2019). These studies suggest that benthic-pelagic coupling will
 3052 weaken in a warmer and less eutrophic Baltic Sea, resulting in gradually decreasing benthic biomass (Ehrnsten et
 3053 al., 2020).

3054 3.3.6.3 Non-indigenous species

3055 It is often suggested that climate change will favour invasions by non-indigenous species worldwide (Jones and
 3056 Cheung, 2014). It has been shown that non-native benthic species typically occur in areas with reduced salinity,
 3057 high temperatures, high proportion of soft seabed and low wave exposure, whereas most native species show an
 3058 opposite pattern (Jänes et al., 2017). Modelled temperature and salinity scenarios suggests an increase of Ponto-
 3059 Caspian cladocerans in the pelagic community, and an increase in dreissenid bivalves, amphipods and mysids in
 3060 the benthos of coastal areas in the northern Baltic Sea by 2100 (Holopainen et al., 2016). Disentangling factors
 3061 facilitating establishment of non-native species demands long-term surveys, and data from multiple environments
 3062 in order to distinguish climate-related effects from other ecosystem-level drivers (Bailey et al., 2020). In addition,
 3063 studies on changing connectivity are needed (e.g. Jonsson et al., 2020).



3064 **3.3.6.4 Fish**

3065 Climate change may affect Baltic Sea fish through effects on water temperature, salinity, oxygen and pH, as well
 3066 as nutrient loads, which indirectly affect food availability for fish. The responses of cod larvae to ocean
 3067 acidification and warming have been studied experimentally. Some studies found no effect on hatching, survival
 3068 or development rates of cod larvae (Frommel et al., 2013), while in others mortality of cod larvae doubled when
 3069 exposed to high-end OA projections (RCP8.5). Several modelling studies however project low abundances of cod
 3070 towards the end of the century, due to continued poor oxygen conditions (Niiranen et al., 2013; Wåhlström et al.,
 3071 2020).

3072
 3073 Climate change may also be positive for fish stocks. Warmer spring and summer temperatures have been projected
 3074 to increase productivity of sprat (Voss et al., 2011; MacKenzie et al., 2012; Niiranen et al., 2013). For herring,
 3075 results are more varied: both increase (Bartolino et al., 2014) and a short-term decrease (Niiranen et al., 2013)
 3076 have been projected.

3077
 3078 Multi-species modelling has also emphasized the role of climate for cod stocks. If fishing is intense but climate
 3079 remains unchanged, cod declines, but not very dramatically, while if climate change proceeds as projected, cod
 3080 disappeared in two models out of seven, even with the current low fishing effort (Gårdmark et al., 2013). Different
 3081 scenarios yield very different outcomes, however. A medium CO₂ concentration scenario (RCP4.5), low nutrients
 3082 and sustainable fisheries resulted in high numbers of cod and flounder, while high emissions (RCP8.5) and high
 3083 nutrient loads resulted in high abundance of sprat (Bauer et al., 2018; Bauer et al., 2019). All these studies assumed
 3084 a more or less pronounced decrease in salinity.

3085 **3.3.6.5 Marine mammals**

3086 *Ringed seal and grey seal – sea ice*

3087 Climate change is projected to drastically reduce the extent of seasonal sea ice in the Baltic Sea (Luomaranta et
 3088 al., 2014; Meier, 2006; Meier, 2015; Meier et al., 2004b). At the end of the 21st century, ice will probably in most
 3089 years be confined to the Bothnian Bay, the eastern Gulf of Finland, the Archipelago sea, and the Moonsund (Sound
 3090 between the Estonian mainland and the offshore western islands Saaremaa, Hiiumaa, Muhu and Vormsi) and
 3091 eastern parts of the Gulf of Riga such as Pärnu Bay (Meier et al., 2004b), with corresponding changes in the
 3092 breeding and moulting distribution of ringed seals. Aside from these projections, ice cover has been even more
 3093 limited in all the southern areas in recent years. Extirpation of one or more of the three southern breeding ringed
 3094 seal populations is possible (Sundqvist et al., 2012; Meier et al., 2004b).

3095
 3096 The ringed seal is an obligatory ice breeder that digs lairs in the snowdrifts on offshore ice for protection of the
 3097 pup (e.g. Smith and Stirling, 1975). The Baltic grey seal prefers loose floes of drift ice (Hook et al., 1972), but can
 3098 also breed on land (Jüssi et al., 2008). Overall pup survival in land breeding grey seals is probably lower than for
 3099 ice breeders (Jüssi et al., 2008). Absence or low quality of sea ice will adversely affect pup survival and quality in
 3100 ice-breeding seals. The effects can be seen by the end of the breeding season, and beyond (Jüssi et al., 2008). Grey
 3101 and ringed seals are capital breeders, i.e., their pup quality depends on effective transfer of maternal energy (fatty
 3102 milk) during a short, intensive lactation period. Timing of birth for both species is strongly adapted to the
 3103 availability of the optimal breeding platform, sea ice. The height of the pupping season is around February-early



3104 March, when the extent and strength of the sea-ice is usually greatest. The immediate breeding success can be
 3105 defined as survival and quality of the offspring at the end of the breeding season, but breeding conditions may
 3106 have population consequences by affecting the survival and fitness of the pups throughout their lives (McNamara
 3107 and Houston, 1996; Kauhala and Kurkilahti, 2020). A warming climate with higher air and water temperatures
 3108 will decrease the extent of ice-cover, the ice thickness and the overlaying snow-cover as well as the stability and
 3109 duration of the ice.

3110

3111 Loss of habitat is critical for reproductive success of the ice-associated seals, especially the ringed seal, and can
 3112 eventually lead to local population decreases and changes in breeding distribution, starting in the southernmost
 3113 parts of its range. The ringed seal populations breeding in the Gulf of Finland, Gulf of Riga and Archipelago Sea
 3114 (SW-Finland) are already small and vulnerable to any negative changes in habitat quality.

3115

3116 *Harbour seal and grey seal – flooding of haul-outs*

3117 Harbour seals and grey seals rely on undisturbed haul-out areas for key life cycle events such as breeding, moulting
 3118 and resting (Allen et al., 1984; Thompson, 1989; Watts, 1996; Reder et al., 2003). In the southern Baltic Sea,
 3119 relative sea levels have risen by 1 to 3 mm per year over the interval 1970-2016 (section 3.2.5.4), and increased
 3120 rates of sea level rise are expected in the future (EEA, 2019c; Grinsted, 2015). A low emissions scenario for the
 3121 21st century projects an additional sea-level rise of 0.29-0.59 m, a high emissions scenario an extra 0.61-1.10 m,
 3122 but substantially higher values cannot be ruled out (Grinsted, 2015; IPCC, 2019a). A high emission scenario is
 3123 thus likely to flood all current seal haul-outs in the southern Baltic Sea and many important localities in Kattegat,
 3124 while under a low emission scenario, most haul-outs in the southern Baltic Sea will be flooded, while others will
 3125 be reduced to small fractions of their current area. In the northern and central Baltic Sea and eastern Kattegat
 3126 archipelago areas, seals will have alternative islets and skerries and are not likely to be affected to the same degree
 3127 as in the south and in eastern Kattegat. In parts of the Gulf of Bothnia, relative sealevels may even fall, due to
 3128 post-glacial rebound (EEA, 2019c).

3129

3130 *Harbour porpoise*

3131 There are no direct studies of the effects of climate change on harbour porpoises in the Baltic Sea, hence the
 3132 following is based mostly on informed guesswork and on a few studies in other areas. There are a multitude of
 3133 ways that changes in one parameter can affect others and we do not currently have the knowledge to predict the
 3134 cumulative effects this might have on the Baltic Sea harbour porpoise population.

3135

3136 Harbour porpoises are present from Greenland to the African coast and the Black Sea and seem to have a rather
 3137 wide thermal tolerance. Therefore, even though it is predicted that we will see a 1.2-3.2° increase in SST in the
 3138 Baltic Sea (Section 3.3.5.1), it seems unlikely that this will directly affect harbour porpoise distribution, unless the
 3139 Baltic Sea harbour porpoise is specifically adapted to colder temperatures. If this is the case, a northwards range
 3140 shift might occur. With the expected future decrease in sea ice extent, the winter habitat available for the harbour
 3141 porpoise in the northern Baltic Sea would increase.

3142

3143 Harbour porpoises are small cetaceans with limited capacity to store energy that mostly live in cold environments.
 3144 Hence, they need to eat almost constantly (Read and Hohn, 1995; Wisniewska et al., 2016) and are therefore



3145 expected to be tightly dependent on their prey (Sveegaard et al., 2012). Their main prey species in the Baltic proper
 3146 are cod (at least before the recent cod stock collapse), sprat, herring, gobies and sand eel (where present). Climate-
 3147 induced changes in for example SST, fronts, stratification and to some degree currents will affect the distribution,
 3148 abundance and possibly the quality of prey species, and in turn the harbour porpoise population. Their distribution
 3149 may shift as they follow their prey, and potential food shortages might lead to starvation, with possible population
 3150 effects.

3151

3152 It has been hypothesized that the susceptibility of marine mammals to disease may increase as temperature
 3153 increases. Higher temperatures can increase pathogen development and survival rates, facilitate transmission
 3154 among individuals and increase individual susceptibility to disease. The negative effects of disease as well as
 3155 environmental contaminants on individual fitness will obviously worsen if the animal is also under nutritional
 3156 stress.

3157

3158 *Seals and changes in the distribution of prey species*

3159 Any large alteration of the ecosystem can affect the distribution of seals if there are climate-related changes in the
 3160 abundance and distribution of their main prey species, such as herring, sprat and cod, as is possible with climate
 3161 change. Such changes in top consumer distribution have been modeled in other sea areas, such as the UK
 3162 continental shelf, where the current distribution of harbour seals did not match well the projected future distribution
 3163 of their prey (Sadykova et al. 2020). There are large differences in salinity between Baltic Sea models (Saraiva et
 3164 al., 2019b), and other factors such as temperature, eutrophication, predation and competition also affect fish
 3165 distributions. Thus, future changes in abundance and distribution of seal prey species, such as herring and cod, are
 3166 hard to predict (Dippner et al., 2008; Lindegren et al., 2010; Vuorinen et al., 2015; Dippner et al., 2019).

3167 **3.3.6.6 Waterbirds**

3168 Climate change scenarios agree in projecting a strong temperature increase in the Arctic and sub-Arctic. This will
 3169 likely cause a northward expansion of species ranges, with colonization by new breeding and wintering species,
 3170 as well as local species declines following migration of populations to ice-free northern waters (Pavón-Jordán et
 3171 al., 2019; Fox et al., 2019).

3172

3173 If salinity in the Baltic Sea decreases, invertebrate species serving as prey for waterbirds (e.g. blue mussels for
 3174 common eiders) are likely to change in distribution, body size and quality as food, with consequences for the
 3175 distribution, reproduction and survival of the waterbirds that eat them (Fox et al., 2015). Predicting the
 3176 consequences of climate change for piscivorous seabirds is complex, because effects are not uniform among Baltic
 3177 Sea fish species. For example, expected increase of recruitment and abundance in an important prey species (sprat;
 3178 MacKenzie et al., 2012; Lindegren et al., 2012) as well as declining numbers of large piscivorous fish (cod) may
 3179 favour fish-eating birds, although management efforts to improve cod stocks may counteract the expected increase
 3180 in sprat and lead to population declines of their main bird predator, the common guillemot (Kadin et al., 2019).
 3181 Herring, another important prey species, is reported to be negatively affected by decreasing salinity (declining
 3182 energy content; Rajasilta et al., 2018).

3183



A rising sea level will reduce the area of saltmarshes available for the breeding of waders and foraging by geese (Clausen et al., 2013), and other coastal habitats would likewise be affected (Clausen and Clausen, 2014). Sea level rise in combination with storms may cause loss by erosion of current coastal breeding habitats, and flood breeding sites, thus affecting the breeding success of coastal waterbirds. Climate change can also be expected to affect waterbirds in the Baltic Sea by changing the incidence of diseases and parasites (Fox et al., 2015).

3.3.6.7 Marine food webs

Climate change and other anthropogenic environmental drivers are expected to change entire marine food webs, from coastal to off-shore, from shallow to deep, from pelagic to benthic (sedimentary), as species-distributions are impacted, and key nodes and linkages in the food webs are altered or lost (Lindegren et al., 2010; Niiranen et al., 2013; Leidenberger et al., 2015; Griffiths et al., 2017; Kotta et al., 2019; Gårdmark and Huss, 2020). These climate-driven changes will also, when combined with societal changes, affect aquatic ecosystem services, for instance future primary production (a supportive ecosystem service) and fish catches (a provisioning ecosystem service; Hyttiäinen et al., 2021).

4 Interactions of climate with other anthropogenic drivers

The term “driver” in this section is defined as something affecting or being affected by another force. In this respect, climate is a force affecting other drivers, e.g. land use or shipping. On the other hand, (regional) climate may be affected by other drivers, e.g. land use or shipping. This section summarizes plausible two-way dependencies that have been described in the literature. For a deeper analysis, see Reckermann et al. (2021).

Climate change affects air and water temperature as well as precipitation, with a clear impact on land use and land cover. Growth conditions are affected by these changes, but also by political or management decisions, which may in turn be influenced by climate change (Yli-Pelkonen, 2008). Agriculture is the most important land use in the southern part of the Baltic Sea basin. Climate change strongly influences the choice of crops, as crops differ in their requirements for water availability and soil type (Fronzek and Carter, 2007; Smith et al., 2008). Still, socio-economic considerations may be even more important than climate in determining agricultural land use (Rounsevell et al., 2005; Pihlainen et al., 2020).

Land use and cover can influence the regional climate, through geophysical (albedo) and biogeochemical (CO₂ drawdown) effects. Bright surfaces like agricultural fields reflect more energy than dark surfaces, like forests and open waters. Thus, the type of land cover may affect regional warming, but its relative contribution is disputed (Gaillard et al., 2015; Strandberg and Kjellström, 2019). Increasing droughts with lower river flow at certain times of the year may influence water management and shipping in regulated rivers, especially in the southern catchment basins. On the other hand, extreme rain events may lead to inundations (Kundzewicz et al., 2005).

Climate change will strongly affect coastal structures through sea level rise and intensified coastal erosion. Storm surges, which run up higher as sea level rises, as well as changed currents and sediment relocations will endanger levees, groynes and other coastal structures, and have to be handled by coastal management (Le Cozannet et al., 2017; Labuz, 2015).



3222

3223 We can expect a considerable increase in offshore wind energy production worldwide, in order to counteract
 3224 climate warming. Although projections of future winds are uncertain, the number of off-shore wind farms can be
 3225 expected to increase due to the politically driven shift to renewable energies, and the limited space and low
 3226 acceptance for wind mills on land. Offshore wind farms may in turn affect the regional climate by absorbing
 3227 atmospheric energy on the regional scale (Akhtar et al., 2021), but the magnitude of this effect is uncertain
 3228 (Lundquist et al., 2019).

3229

3230 Shipping is affected by climate change. Perils at sea for ships are all climate sensitive, ranging from storms, waves,
 3231 currents, ice conditions, visibility to sea level affecting navigational fairways. Winter navigation will be facilitated
 3232 as drastically decreasing winter sea-ice cover is projected, but search and rescue missions in winter may increase
 3233 because engine power may in the future be adapted to the lower expected ice cover. Further aspects are a potential
 3234 increase in leisure boating, a potentially temperature-dependent functioning of antifouling paints, and different
 3235 noise propagation through warmer water. The efficiency of SO_x scrubbing depends on the temperature, salinity
 3236 and pH of the seawater, and eventually ends up contaminating the Baltic Sea (Turner et al., 2018). Shipping itself
 3237 affects climate through combusting fossil fuels.

3238

3239 Coastal processes, e.g. erosion and the translocation of sediments through erosion, currents and accretion, are
 3240 affected by climate change though sea level rise and changes in storm frequency, severity and tracks (Defeo et al.,
 3241 2009).

3242

3243 Climate change affects the amount of nutrients entering the sea in precipitation and by land runoff, which in turn
 3244 is affected by precipitation, air temperature and runoff pattern, e.g. Arheimer et al. (2012) and Bartosova et al.
 3245 (2019). How fertilization practices, crops grown, and land use will change in response to climate change is largely
 3246 unknown. Climate-related changes in the Baltic Sea, like warmer temperatures, changed stratification and altered
 3247 ecosystems and biogeochemical pathways may change the fate of nutrients in the sea, e.g. Kuliński et al. (2021).

3248

3249 There is not much evidence of a direct climate influence on the quantity and quality of submarine groundwater
 3250 discharge, but considering the driving forces (topography-driven flow, wave set-up, precipitation, sea level rise
 3251 and convection), an effect is highly plausible, but its magnitude and relevance is unknown (e.g. Taniguchi et al.,
 3252 2019).

3253

3254 Fisheries are strongly affected by climate change through its effect on the resources, i.e. the commercially
 3255 interesting fish populations in the Baltic Sea, mostly cod, sprat and herring (Möllmann, 2019). Climate affects
 3256 salinity and temperature in the Baltic Sea, thereby influencing the productivity of several fish species (MacKenzie
 3257 and Schiedek, 2007; Köster et al., 2016), and the resources that fisheries exploit. Growth of planktivorous species
 3258 or life stages is also affected by climatic conditions that regulate zooplankton dynamics (Casini et al., 2011; Köster
 3259 et al., 2016).

3260

3261 Climatic change is a plausible driver for the migration and occurrence of non-indigenous species, although there
 3262 is little direct evidence. Shipping has been identified as a major vector for the introduction of new marine species



into the Baltic Sea ecosystem, through ballast water or attachment to hulls or elimination of physical barriers (e.g. though the construction of canals between water bodies; Ojaveer et al., 2017). A northward migration of terrestrial (Smith et al., 2008) and marine species, including fish, is documented and expected to continue (MacKenzie and Schiedek, 2007; Holopainen et al., 2016).

Climate change affects contaminants in the Baltic Sea through an array of processes, like partitioning between environmental phase-pairs such as air-water, air-aerosols, air-soil, air-vegetation, leading to a different distribution between environmental compartments (Macdonald et al., 2003). Atmospheric transport and air-water exchange can be influenced by changes in wind fields and wind speeds (Lamon et al., 2009; Kong et al., 2014). Changing precipitation patterns influence chemical transport via atmospheric deposition (rain dissolution and scavenging of particles, (Armitage et al., 2011) and runoff, transporting terrestrial organic carbon (Ripszam et al., 2015). As ice-cover of lakes and the sea decreases, more organic contaminants may volatilize to the atmosphere (Macdonald et al., 2003; Undeman et al., 2015).

Dumped military ammunition threaten the Baltic Sea in the future, as poisonous substances are expected to leak due to advanced corrosion of hulls and containers. This process may be affected by climate, as corrosion rates depend on temperature and oxygen, so that warming and good ventilation of dumping sites can be expected to enhance corrosion rates (Vanninen et al., 2020). This is an urgent problem since the location of the dumped military material is only partially known.

There is no evident direct impact of climate change on marine litter or microplastics, but there may be a connection via increased temperature- and photolysis-dependent rates of degradation and dissolution of microplastics, and on distribution by currents.

5 Comparison with the North Sea region

5.1 The North Sea region

Like the Baltic Sea basin, the North Sea region is both a precious natural environment and a place for settlement and commerce for millions of people, with a rich cultural heritage. The North Sea is one of the world's richest fishing grounds as well as one of the busiest seas with respect to shipping and infrastructure for oil and gas extraction, and of enormous economic value. In recent years the area has also become a major site for wind energy, with many large offshore wind farms.

As climate change is expected to have profound effects on North Sea ecosystems and economic development, an independent, voluntary, international team of scientists from across the region compiled the North Sea Region Climate Change Assessment (NOSCCA; Quante and Colijn, 2016). The NOSCCA approach is similar to BACC in format and intention. The assessment provides a comprehensive overview of all aspects of a changing climate, discussing a wide range of topics including past, current and future climate change, and climate-related changes in marine, terrestrial and freshwater ecosystems. It also explores the impact of climate change on some socio-economic sectors, such as fisheries, agriculture, coastal zone management, coastal protection, urban climate, recreation/tourism, offshore activities/energy, and air pollution.



3302

3303 The North Sea is a semi-enclosed marginal sea of the North Atlantic Ocean, situated on the north-west European
 3304 shelf. It opens widely into the Atlantic Ocean at its northern boundary, with a smaller connection to the Atlantic
 3305 Ocean via the Dover Strait and English Channel in the south-west. To the east it connects to the Baltic Sea. The
 3306 Kattegat, a transition zone between the North and Baltic seas, is located between the Skagerrak and the Danish
 3307 straits. Comprehensive reviews of North Sea physical oceanography are provided by Otto et al. (1990), Rodhe
 3308 (1998) and Stündermann and Pohlmann (2011). Physical-chemical-biological interaction processes within the
 3309 North Sea are reviewed by Rodhe et al. (2006) and Emeis et al. (2015), and a description of the North Sea marine
 3310 ecosystem was compiled by McGlade (2002).

3311

3312 Among the most striking differences between the North and Baltic seas is the wide, direct opening of the North
 3313 Sea to the North Eastern Atlantic, allowing free exchange of matter, heat and momentum between the two seas.
 3314 As a result the North Sea water has a much higher salinity than the Baltic Sea. The North Sea dynamics are greatly
 3315 influenced by tides, while Baltic Sea tides are much weaker than in the North Sea, where tidal amplitudes vary
 3316 spatially from a few decimeters to several meters. In addition to the wind-driven circulation, which dominates the
 3317 mean cyclonic current system, North Sea tidal currents show non-vanishing residual currents (due to nonlinear
 3318 processes), which cannot be neglected. Tidal currents cause strong mixing. Low pressure systems often travel from
 3319 the Atlantic with minimum blockage and cause strong storm surges, which are the greatest potential natural hazards
 3320 for coastal communities in the North Sea region.

3321

3322 Only selected examples from NOSCCA will be presented here. In general, the North Sea region already
 3323 experiences a changing climate and projections indicate that further, partly accelerating, changes are to be expected
 3324 (warming of air and water, changing precipitation intensities and patterns, sea level rise, seawater acidification).
 3325 Changes in ecosystems (marine, coastal, terrestrial) are observed, and are projected to strengthen, with degree
 3326 depending on scenario. Observational as well as modelling studies have revealed a large natural variability in the
 3327 North Sea region (from annual to multidecadal time scales), making it difficult to identify regional climate change
 3328 signals and impacts for some parameters. Projecting regional climate change and impacts for the North Sea region
 3329 is currently limited by the small number of regional coupled model runs available and the lack of consistent
 3330 downscaling approaches, both for marine and terrestrial impacts. The wide spread in results from multi-model
 3331 ensembles indicates the present uncertainty in the amplitude and spatial pattern of the projected changes in sea
 3332 level, temperature, salinity and primary production. For moderate climate change, anthropogenic drivers such as
 3333 changes in land use, agricultural practice, river flow management or pollutant emissions often seem more
 3334 important for impacts on ecosystems than climate change.

3335 5.2 A few selected and highly aggregated results from NOSCCA

3336 *Atmosphere:* Observations reveal that the near-surface atmospheric temperature has increased everywhere in the
 3337 North Sea region, especially in spring and in the north. The rise was faster over land than over the sea. Linear
 3338 trends in the annual mean land temperature are about +0.39°C per decade for the period 1980–2010. Generally,
 3339 more warm extremes and fewer cold ones were observed. A north-eastward shift in storm tracks was observed, in
 3340 agreement with projections from climate models forced by increased greenhouse gas concentrations. Overall,
 3341 precipitation has increased in the northern North Sea region and decreased in the south, summers have become



warmer and drier and winters have become wetter. Heavy precipitation events have become more extreme. A marked further mean warming of 1.7–3.2°C is projected for the end of the 21st century (2071–2100, with respect to 1971–2000) for different scenarios (RCP4.5 and RCP8.5, respectively), with stronger warming in winter than in summer and particularly strong warming over southern Norway.

North Sea: There is strong evidence of surface warming in the North Sea, especially since the 1980s (Fig. 33). Warming is greatest in the south-east, exceeding 1°C since the end of the 19th century. Absolute mean sea level in the North Sea rose by about 1.6 mm/year over the past 100–120 years, in agreement with the global rise. The North Sea is a sink for atmospheric carbon dioxide (CO₂); this uptake declined over the last decade, due to lower pH and warmer water. Models consistently project the surface water to warm further by the end of the century (by about 1–3°C; A1B scenario). Exact numbers are not given due to differences in spatial averaging and reference periods from published studies. Coherent findings from published climate change studies include an overall rise in sea level, an increase in ocean acidification and a decrease in primary production. Uncertainties are large for projected changes in extreme sea level and waves as well as for decreases in net primary production, which range from 1 to 36 %.

Rivers: To date, no significant trends in response to climate change are apparent for most individual rivers discharging into the North Sea. Nevertheless, climate models project increased socio-economically important risks for the region, due to more intense hydrological extremes in the North Sea region, such as flooding along rivers, droughts and water scarcity. The exposure and vulnerability of cities in the North Sea region to changes in extreme hydrometeorological and hydrological conditions are expected to increase, due to greater urban land use and rising urban populations.

Ecosystems: Long-term knowledge from exploitation of the North Sea indicates that climate affects marine biota in complex ways. Climate change influences the distribution of all taxa, but other factors (fishing, biological interactions) are also important. The distribution and abundance of many species have changed. Warm-water species have become more common and species richness has increased. Among coastal ecosystems, estuaries and most mainland marshes will survive sea-level rise, while back-barrier salt marshes with lower suspended sediment concentrations and tidal ranges are probably more vulnerable. Plant and animal communities can suffer habitat loss in dunes and salt marshes through high wave energy, and are affected by changes in temperature and precipitation and by atmospheric deposition of nitrogen. Lakes in the North Sea region have experienced a range of physical, chemical and biological changes due to climatic drivers over past decades. Lake temperatures have increased, ice-cover duration has decreased. For terrestrial ecosystems there is strong empirical evidence of changes in phenology in many plant and animal taxa and northward range expansions of mobile heat-loving animals. Climate change projections and effect studies suggest a northward shift of vegetation zones, with terrestrial net primary production likely to increase in the North Sea region, due to warmer conditions and longer growing seasons.

Socio-economic effects: The assessments of climate change effects on the different socio-economic sectors in the North Sea region find that adaptation measures are essential for all of them, e.g. for coastal protection and in agriculture. For North Sea fisheries, the rapid temperature rise is already being felt in terms of shifts in species



distribution and variability in stock recruitment. In agriculture an increased risk of summer drought and associated effects will be a challenge, particularly in the South. In general, extreme weather events are likely to more often severely disrupt crop production. Offshore and onshore activities in the North Sea energy sector (dominated by oil, gas and wind) are highly vulnerable to extreme weather events, in terms of extreme wave heights, storms and storm surges. All coastal countries around the North Sea with areas vulnerable to flooding by storm surges are preparing for the challenges expected due to climate change, but coastal protection strategies differ widely from country to country. Due to uncertainty concerning the extent and timing of climate-driven impacts, current coastal zone adaptation plans focus on no-regret measures.

5.3 Some differences in climate change effects between the North and Baltic Seas

Many of the climate change signals in the Baltic and North Seas show similar behaviours and trends. But there are also some notable differences between the two regions, which are listed below. They are based on findings reported in the appropriate chapters of the recent assessments BACC II (BACC II Author Team, 2015) and NOSCCA (Quante and Colijn, 2016).

- In recent decades, the surface air temperature in the Baltic Sea and North Sea regions rose in a similar way, on the order of 1 °C in the past century. Projections of the surface air temperature as obtained by Euro-CORDEX downscaling for a moderate scenario (RCP4.5) indicate a stronger winter and spring warming (> 1 K) at the end of the century (2071-2100) relative to present day (1971-2000) for most parts of the Baltic Sea region than for the western part of the North Sea region. In the summer and autumn months the projected warming is at the same level.
- The North Sea is vigorously ventilated by the Atlantic (overturning time ~1 to 4 years). Therefore, climate change signals from the Atlantic are rapidly transferred to the North Sea, while climate change in the North Sea can be expected to be damped by the large thermal inertia of the Atlantic Ocean. By contrast, the Baltic Sea is more prone to changes in mean meteorological conditions as its connection to the World Ocean is very narrow.
- Projected changes in mean precipitation show a distinctive difference between the two sea regions for the summer (JJA) and autumn months (SON). In the Baltic Sea region the mean precipitation for a RCP4.5 scenario is projected to increase for most land areas (5 to 25%), whereas no noticeable change (5 to -5%) is projected along the western and southern shores of the North Sea region.
- SST is currently rising and projected to rise further for both sea areas, but the spatial pattern of the SST increase is different. In the southern North Sea SST rises more than in the northern North Sea, while SST warming trends are higher in the north-eastern part of the Baltic Sea (Bothnian Sea and Gulf of Finland) than in the southern part. These spatial differences are explained by water depth (North Sea) and the ice-albedo feedback (Baltic Sea). In addition, the northern North Sea is affected by Atlantic water inflow at the western side of the Norwegian trench.
- The coastal regions of the North Sea experience increases in both mean sea level (MSL, as measured by satellites) and relative mean sea level (RMSL, as measured by tide gauges). Trends in RMSL vary significantly across the North Sea region due to the influence of vertical land movement (uplift in northern Scotland, Norway and Denmark, and subsidence elsewhere). But the trend of RMSL is still positive everywhere in the North Sea coasts. In contrast, sea levels relative to land along the northern Baltic Sea coast are sinking because land levels continue rising, due to post-glacial rebound since the last ice age.



3423 The northern Baltic Sea will experience considerable land rise also in future. As a result, the sea level
 3424 will probably continue to decrease relative to land in this region. As positive trends in RMSL are more
 3425 relevant for coastal protection, all countries around the North Sea with coastal areas vulnerable to flooding
 3426 due to storm surges face similar challenges, while in the Baltic Sea region coastal protection is of greater
 3427 concern for the countries in the south.

- 3428 • The frequency of sea ice occurrence in the North Sea has decreased since about 1961, with a similar
 3429 development in the western Baltic Sea. In contrast, ice still forms in the northern Baltic Sea, where it will
 3430 remain a prominent feature for many years, covering about 50 to $200 \times 10^3 \text{ km}^2$, with high interannual
 3431 variability, even though a linear trend of 2% decrease per decade is reported.

3432 6 Knowledge gaps and research needs

3433 Knowledge gaps and research needs have been intensively discussed within the grand challenge working groups
 3434 of Baltic Earth and are summarized by the BEARs (Lehmann et al., 2021; Kuliński et al., 2021; Rutgersson et al.,
 3435 2021; Weisse et al., 2021; Christensen et al., 2021; Gröger et al., 2021a; Meier et al., 2021a; Viitasalo, 2021;
 3436 Reckermann et al., 2021).

3437

3438 In summary, we conclude that the processes that control the variability of salinity in the Baltic Sea and its entire
 3439 water and energy cycles are still not fully understood (Lehmann et al., 2021). The time-dependence of the haline
 3440 stratification and its links to climate change are in special need of further study. Salinity dynamics is important for
 3441 its dominant role in stratification, concerning both mixing conditions and ecosystem composition and functioning.
 3442 The environmental and biological factors favoring certain biogeochemical pathways through complex interactions,
 3443 the pools of dissolved organic matter, and sediment biogeochemical processes are poorly understood (Kuliński et
 3444 al., 2021). Although initial studies on the coastal filter capacity have been made, coastal zone models for the entire
 3445 Baltic Sea and an overall estimate of bioavailable nutrients and carbon loads from land to the open sea do not exist
 3446 (Kuliński et al., 2021). Considering the large internal variability, investigations of changes in extremes are limited
 3447 because high-resolution observational time series are too short and model ensembles too small (Rutgersson et al.,
 3448 2021). Global mean sea level rise, land uplift and wind field changes control sea level of the Baltic. However, the
 3449 future evolution of these drivers, which are needed for projections, is rather uncertain (Weisse et al., 2021).
 3450 Furthermore, databases for coastline changes and erosion and basin-scale models of coastal change under sea-level
 3451 rise do not exist (Weisse et al., 2021).

3452

3453 Fully coupled regional ESMs for the Baltic Sea including the various compartments of the Earth system,
 3454 atmosphere, land, ocean, sea ice, waves, terrestrial and marine ecosystems are under development but are not yet
 3455 available for dynamical downscaling (Gröger et al., 2021a). The numerical estimation of water and energy cycles
 3456 suffers from both model deficiencies and natural variability. For climate projections, large ensembles of regional
 3457 atmosphere models are available but only one ensemble with 22 members utilized a coupled atmosphere-ice-ocean
 3458 model (Christensen et al., 2021). In the past, the ocean ensembles have had too few members to address well the
 3459 uncertainty related to the large multidecadal variability in the ocean (Meier et al., 2021a). Furthermore, the global
 3460 sea level rise needs to be considered when making salinity projections (Meier et al., 2021b). The large uncertainty
 3461 in future projections of salinity fundamentally affects the projections of the marine ecosystem (Viitasalo, 2021).



The response of food web interactions to climate change are largely unknown. The uncertainties of scenario simulations with coupled physical-biogeochemical ocean models were discussed by Meier et al. (2018a; 2019b; 2021b). They found that in addition to natural variability the largest uncertainties are caused by (i) poorly known current bioavailable nutrient loads from land and atmosphere and uncertain assumptions about future loads, (ii) uncertainties of models including global sea level rise, and (iii) poorly known long-term future greenhouse gas emissions.

Finally, the regional Earth system is driven by multiple drivers, of which climate change is just one. Multi-driver studies are just beginning to be made and only a few have yet been published (Reckermann et al., 2021).

In the following, we list a few selected knowledge gaps related to the variables addressed by this study.

6.1 Large-scale atmospheric circulation

The interactions between atmospheric modes of variability of importance for the Baltic Sea region are still not well known. For instance, while climate models are able to simulate the main features of the NAO, the frequency of blocking over the Euro-Atlantic sector is still underestimated (IPCC, 2014b). Since observational records are relatively short, our understanding of the AMO and its possible changes depends largely on models, and these cannot be reliably evaluated for time scales longer than the AMO period (Knight, 2009). However, while possible changes in these climate phenomena do contribute to the uncertainty in near-term climate projections, they are not the main driver of the projected warming over Europe by the end of the century (Cattiaux et al., 2013; IPCC, 2014b).

6.2 Air temperature

Temperature and its extremes are to a large extent determined by the large-scale circulation patterns. There is limited knowledge primarily concerning changes in large-scale atmospheric circulation patterns in a changing climate, as mirrored by climate model discrepancies. Nevertheless, the heat cycle of the Baltic Sea region is probably better understood than the water cycle.

6.3 Solar radiation and cloudiness

Multidecadal variations in surface solar radiation (SSR) are generally not well captured by current climate model simulations (Allen et al., 2013; Storelvmo et al., 2018). The extent to which the observed variations in SSR are caused by natural variation in cloudiness induced by atmospheric dynamic variability (Stanhill et al., 2014; Parding et al., 2014), or by anthropogenic aerosol emissions (Wild, 2012; Ruckstuhl et al., 2008; Philipona et al., 2009; Storelvmo et al., 2018), or perhaps additional causes, is not understood. Future cloudiness trends in global and regional models differ in their sign (Bartók et al., 2017).

6.4 Precipitation

Even if climate scenarios are becoming more frequent and there is now a growing ensemble of relatively high-resolution regional climate scenarios for Europe, they still represent only a subset of the global climate model projections assessed by the IPCC. This means that the uncertainties of future climate change in the Baltic Sea region are not fully captured at the horizontal resolution needed for detailed studies of climate change effects in



the region (Christensen et al., 2021). Very high-resolution so called “convective-permitting” climate models operating at grid spacing of 1-3 km are lacking for the Baltic Sea region. In other regions, such models have better agreed with observations of precipitation extremes and sometimes also given a larger climate change signal than the more traditional “high-resolution” models operating at c. 10 km grid spacing (Christensen et al., 2021). Land use change and cover, including changes in forests, can induce both local and downwind precipitation change (Meier et al., 2021c), and need to be included in projections.

6.5 Wind

Historical wind measurements suffer from inhomogeneity and records too short for detecting changes, considering the large internal variability in the Baltic Sea region. Projected changes are not robust among the few available downscaled ESMs.

6.6 Air pollution

The spatially and time resolved air quality status of a region is often assessed by means of model systems, typically with emission, meteorological and chemistry transport submodels. These model systems, used for the calculation of atmospheric concentrations and deposition of pollutants, need further developments and validation. Uncertainties are often connected to the emission segment of the modelling chain. Improvements of the implemented time profiles for the different emission sectors are especially necessary (Matthias et al., 2018). For projections of air quality with climate change models, more work is needed to establish a set of emission scenarios for air pollutants consistent with regional socio-economic pathways, like those developed by Zandersen et al. (2019). The shipping sector is currently a considerable source of air pollution in the Baltic Sea region. More research and development is needed on new fuel types and emission factors for air pollutants, relevant for politically and technologically driven abatement measures. To better address exposure and health impacts of shipping emissions more studies are required like those of Ramacher et al. (2019) and Barregard et al. (2019), especially at the harbour and city scale. Better knowledge and reduced uncertainties will improve quantification of air pollution as part of the environmental imprint of shipping in the Baltic Sea region, as developed by Moldanová et al. (2021).

6.7 River discharge

Precipitation from regional atmosphere models is biased and the bias correction methods applied for hydrological modeling affect the sensitivity of hydrological models to climate change (Donnelly et al., 2014). Natural variability and model uncertainties may explain the large spread in current river discharge projections (Roudier et al., 2016; Donnelly et al., 2017). The values of the parameters of a hydrological model are normally found through calibration against historical data and are always associated with uncertainty. This uncertainty will translate into uncertainty in the projected changes.

6.8 Nutrient inputs from land

The time scales for exchange of the nutrient pools in soils are not well known (McCrackin et al., 2018). Long-term observations do not exist. Future projections of river discharge and nutrient inputs in the Baltic Sea drainage basin agree on key aspects (e.g. increased annual discharge), but also highlight the uncertainty of the projections. To improve assessments, studies should be designed to allow explicit semi-quantitative comparisons of the effects of



the incorporated change factors, e.g. climate, land management, policy. In the case of nitrogen inputs, the effect of changes in anthropogenic atmospheric deposition should also be included in future projections.

6.9 Terrestrial biosphere

Terrestrial ecosystems in the Baltic Sea region are governed by human activities, both changes in climate due to anthropogenic climate forcing and anthropogenic changes in land use and land cover. In return, terrestrial ecosystems affect climate by altering the composition and the energy and water cycles of the atmosphere. Biophysical interactions between the land surface and the atmosphere have been incorporated into regional ESMs, in order to assess the impacts of changes in land use and land cover on regional climate and terrestrial ecosystems. Still, biogeochemical processes related to the carbon cycle are lacking, as are explicit forest management actions (Lindeskog et al., 2021), while explicit descriptions of some disturbances (e.g. wildfires, major storms, insect attacks) are under development in ESMs. Only when all these interactions are incorporated can the effects of national or international (e.g. in the European Union) climate policies on regional climate and terrestrial ecosystems be fully assessed for compliance with the goals of the Paris Agreement.

6.10 Snow

A general decrease in snow-cover duration in the Baltic Sea region is well documented, especially for the southern part. Changes in snow depth due to climate warming are much more unclear. Some evidence of increasing snow depth in recent decades have been reported from the northern part of the study region and from mountainous areas. Changes in sea-effect snowfall events during present climate are unknown.

6.11 Glaciers

It is presently not known how glacier-fed lakes react to competing environmental drivers, such as the general Arctic warming, and the simultaneous warming-triggered lake cooling caused by increased inflow of cold glacier meltwater, potentially carrying high sediment, nutrient, and organic matter loads. Understanding changing lake thermal regimes and vertical mixing dynamics as well as timing and duration of seasonal ice cover is important because ecological, biological, chemical processes, including carbon-cycling, will be affected (Lundin et al., 2015; Smol et al., 2005; Jansen et al., 2019). Since Scandinavian glaciers are predicted to decline 80% in volume by 2100 under RCP8.5, Scandinavian glacier-fed lakes could be used as natural observatories, where changes in processes, timescales, and effects in response to competing drivers can be studied before they occur at other glacial lake sites, where glaciers melt more slowly (Kirchner et al., 2021).

6.12 Permafrost

Thawing permafrost peatlands may potentially release large amounts of organic matter, nutrients and greenhouse gases to aquatic systems locally, but the timing and magnitude of such releases remain highly uncertain.

6.13 Sea ice

While the extent of the sea ice cover is well observed, observations of ice thickness are scarce. Ice thickness is regularly monitored only at a few coastal sites with fast ice. Long records of the various ice classes, such as ridged ice, do not exist. Sea ice models do not represent sea ice classes correctly. Since the last assessment by the BACC II Author Team (2015) only two new scenario simulation studies on sea ice were published.



3572 **6.14 Lake ice**

3573 Research is required to better understand the reasons for regional and temporal differences in the patterns of change
 3574 in lake ice phenology and its relationship to large-scale climatic forcing. There is a need to better understand how
 3575 loss of lake ice cover modifies gas exchange between lake and atmosphere, mixing of the water column,
 3576 biogeochemical cycling, and ecosystem structure and function. The socioeconomic and cultural importance of
 3577 winter ice also deserves further research.

3578 **6.15 Water temperature**

3579 The causes of the pronounced natural variability of Baltic Sea temperature and its connection to large-scale patterns
 3580 of climate variability is not well known. The occurrence of marine heatwaves is projected to increase. However,
 3581 only a few studies of their impacts on the marine ecosystem exists. Furthermore, sea surface temperature trends
 3582 also depend on coastal upwelling, which affects large areas of the Baltic Sea surface (Lehmann et al., 2012).
 3583 Projected changes in upwelling are, however, very uncertain (Meier et al., 2021a).

3584 **6.16 Salinity and saltwater inflows**

3585 Salinity change depends on wind, river discharge, net precipitation on the sea and global sea level rise. Due to
 3586 considerable uncertainty in all drivers and the different signs in the response of salinity to these drivers, the relative
 3587 uncertainty in salinity projections is large, and larger ensembles of scenario simulations are needed (Meier et al.,
 3588 2021b). This knowledge gap is compounded by the uncertainty of whether saltwater inflows from the North Sea
 3589 will change. As salinity is a very important variable for the circulation in the Baltic Sea and for the marine
 3590 ecosystem, projections for the Baltic Sea are a priority.

3591 **6.17 Stratification and overturning circulation**

3592 Stratification depends on mixing as well as on gradients in water temperature and salinity, making changes in
 3593 stratification uncertain. Mixing processes such as thermal and haline convection, entrainment, double diffusive
 3594 convection or boundary mixing are not fully understood. Initial results on the sensitivity of the vertical overturning
 3595 circulation rely on model studies only. Hence, more measurements on the fine-structure of horizontal and vertical
 3596 turbulence are needed.

3597 **6.18 Sea level**

3598 The regional variability of processes which drive sea-level changes, along with their uncertainties and relative
 3599 importance over different timescales, display long-term developments that still require an explanation and are a
 3600 challenge to planning by coastal communities (Hamlington et al., 2020). For instance, the annual cycle in Baltic
 3601 Sea mean sea level (winter maxima minus spring minima) shows a basin-wide widening in the period 1800-2000
 3602 (Hünicke and Zorita, 2008). The precise mechanisms responsible for this effect are not yet completely understood,
 3603 although it seems strongly controlled by atmospheric forcing (Barbosa and Donner, 2016). Furthermore, at the
 3604 longer time-scales relevant for anthropogenic climate change, Baltic Sea and North Atlantic sea levels are strongly
 3605 affected by the very uncertain future melting of the Antarctic ice sheet. Current estimates are mostly based on
 3606 heuristic expert knowledge, as models are still under development. This is probably the largest knowledge gap
 3607 affecting projections of future Baltic sea-level rise (Bamber et al., 2019). Finally, long-term relative sea level



trends are strongly affected by the vertical land movement due to glacial isostatic adjustment. This can be as large as, or even larger, than global sea-level rise. Currently, it is estimated from relatively short GPS measurements and from geo-elastic models. Both are uncertain, as point GPS measurements are strongly affected by other geological and anthropogenic effects on vertical land velocities and results from model geo-elastic models are often revised. In addition, the glacial isostatic adjustment may affect the flow intensity of river runoff into the northern Baltic Sea (coastal regions rising relative to inland regions), the effects of which, e.g. on salinity and water levels, have not been explored.

6.19 Waves

The lack of long-term instrumental wave measurements and gaps in the data due to the ice season complicate the analysis of extreme values. Although wave hindcasts provide a good alternative, the accuracy naturally does not match that of measured data. Furthermore, Björkqvist et al. (2020) showed that the calculation of return periods of extreme events may depend on the sampling frequency. Adding sampling variability typical for in-situ measurements to simulated hindcast data, will result in consistently shorter estimates of return periods for high significant wave heights than using the original hindcast data.

6.20 Sedimentation and coastal erosion

We lack a comprehensive understanding of alongshore sediment transport and its associated spatial and temporal variability along the Baltic Sea coast. In general, an eastward transport dominates along most of the southern Baltic Sea coast due to the prevailing westerly winds. However, the intensity of secondary transport induced by easterly and northerly winds is much less understood. Its combination with storm surges will expose sand dunes and cliffs to the greatest erosional impact, further complicating understanding (Musielak et al., 2017). Due to the orientation of the coastline, transport along some parts of the Baltic Sea coastline is very sensitive to the angle of incidence of the waves. For example, the incidence angle of westerly wind-waves at the western part of the Wolin Island in Poland (Dudzińska-Nowak, 2017) and the coast of Lithuania and Latvia (Soomere et al., 2017) is very small and even a slight change in the wind direction (e.g. by 10 degrees) could lead to a reversal of the direction of alongshore transport. Coastline changes at these sections vary greatly, and will hence be extremely sensitive to future changes in wind wave climate (Viška and Soomere, 2013). Another knowledge gap in understanding coastal erosion in response to future climate change concerns the impact of water levels and the submergence of the beach. Water level plays a key role in dune toe erosion and also limits aeolian sand transport on the beach. The relationship between the intensity of the forcing (wave energy, run-up) and the morphological response (erosion at the beach and dunes) during storms is not straightforward (Dudzińska-Nowak, 2017; Zhang et al., 2017). At some sites (e.g. Miedzyzdroje), dune erosion is well correlated with maximum storm surge level and storm frequency, but at others (e.g. Swinoujście), the beach morphology is more important in determining the effect of erosion than the storm surge level.

6.21 Oxygen and nutrients

There are significant knowledge gaps related to the identification and quantification of oxygen sinks and sources in the Baltic Sea. In particular, more understanding is required on the dynamics of seawater inflows from the North Sea, the role of mixing processes in the ventilation of the deep water, rates of oxygen consumption in water column and sediments and how they depend on climate change. Knowledge gaps also exist concerning the transport and



transformations of DOM (including terrestrial DOM) and better quantification of the processes occurring in the microbial loop is needed to understand the nutrient (but also C and O) dynamics in the Baltic Sea.

The direct effects of climate change are likely to be detectable first in the coastal zone, e.g. indicated by increasing seasonal hypoxia due to warming. However, long-term records from the coastal zone are rare. More important could be the intensification of the proposed hypoxia-related “vicious circle” in the Baltic proper, due to the warming of both surface and deep-water layers in the Baltic proper (Savchuk, 2018; Meier et al., 2018b). The consequent expansion of cyanobacteria blooms and increased nitrogen fixation in the Baltic proper and neighboring basins could further counteract nitrogen load reductions and maintain hypoxia, with all its detrimental effects. However, there are still no biogeochemical and ecosystem models capable of producing reliable long-term scenario simulations of these processes, with sufficient confidence and precision (see Meier et al., 2018a; Meier et al., 2019b).

6.22 Marine CO₂ system

Due to the high spatial and temporal variability of air-sea carbon fluxes, it is not known whether the Baltic Sea as a whole is a net sink or a net source of CO₂. The source of the alkalinity increase observed in the Baltic Sea is still unclear. Plausible hypotheses indicate increased weathering in the catchment and processes related to anoxic remineralization of organic matter. There is high uncertainty in quantifying sediment/water fluxes of C, N and P, which are important bottlenecks for understanding the dynamics of the marine CO₂ system and the C, N, P and O₂ cycling generally, especially in the deep water layers. The lack of system understanding is particularly evident in the Gulf of Bothnia. Fransner et al. (2018) suggested that non-Redfieldian stoichiometry in phytoplankton production could explain pCO₂ fields in these basins, but confirmation by observations is still lacking.

6.23 Marine biosphere

6.23.1 Lower trophic levels

The summer cyanobacteria bloom in the Baltic proper, and increasingly in recent years in the Bothnian Sea, is considered one of the main problems of Baltic Sea eutrophication, and the nitrogen fixation it carries out is an important process in Baltic ecosystem models (Munkes et al., 2021). It has long been considered limited by the availability of phosphorus (Larsson et al., 1985; Granéli et al., 1990). It is therefore remarkable that it is not possible to predict inter-annual variations in cyanobacteria blooms observed by satellites from water chemistry (Kahru et al., 2020).

There are significant knowledge gaps related to the quantification of nitrogen fixation and the fate of the fixed nitrogen in the Baltic Sea pelagic zone. Direct nitrogen fixation measurements were until recently dogged by method problems, and even if these are now hopefully largely resolved (Klawonn et al., 2015), the enormous patchiness of cyanobacteria blooms remains a huge problem. The alternative approach of directly measuring the increase in total combined nitrogen during the bloom (Larsson et al., 2001) requires very high precision, also suffers from patchiness problems (Rolf et al., 2007), and has not been much used. Finally, the amount of nitrogen fixed can be estimated by modelling, based on uptake of CO₂ or phosphorus and assuming a Redfield N:P or C:N ratio. This theoretically highly attractive approach (Eggert and Schneider, 2015) is hampered by the possibility of



3684 non-Redfieldian ratios, and has made some biologists skeptical by predicting high nitrogen fixation in spring, when
 3685 there are not sufficient known nitrogen-fixing autotrophs in the water to carry out this nitrogen fixation.

3686

3687 While total nitrogen in the water column clearly increases during the summer cyanobacterial bloom, just a couple
 3688 of months later this increase seems largely to have disappeared, even though sediment traps find little evidence
 3689 that it has settled out of the upper mixed layer. Are sediment trap measurements gross underestimates, or are there
 3690 unidentified sites of denitrification or other overlooked nitrogen sinks in the water column? Nitrogen fixation is a
 3691 central process in Baltic ecosystem models, and better observationally based estimates of processes in the nitrogen
 3692 cycle are required for assessing their credibility (Munkes et al., 2021).

3693 **6.23.2 Marine mammals**

3694 Seal and porpoise foraging distribution and the relation of seals to haul-out sites is not well known. The
 3695 requirement of sea ice for successful breeding of ringed seals has not been sufficiently assessed. Land-breeding of
 3696 grey seals is not monitored regularly in most Baltic countries. The effects of interspecific competition on
 3697 distributions are not known. Range contraction can be conceptualized as three stages (Bates et al., 2014):
 3698 performance decline, population decrease and local extinction, all of which should be studied. For example, studies
 3699 on performance decline, such as physiological conditions that reduce reproductive potential (Helle, 1980; Jüssi et
 3700 al., 2008; Kauhala et al., 2017; Kauhala et al., 2019) are important (Bates et al., 2014). Breeding success of Baltic
 3701 Sea ringed seals in normal winters is poorly known, as the lairs in pack ice snowdrifts are rarely found. Likewise,
 3702 observations of the effects of poor ice-conditions on breeding success of ringed seals in mild winters are very
 3703 limited, but the lack of protection against harsh weather and predators is assumed to be highly negative.

3704 **6.23.3 Waterbirds**

3705 The complex interaction between many primary parameters affected by climate change makes it hard to identify
 3706 which environmental changes are actually causing changes in waterbird populations. It is currently not known in
 3707 detail how shifts in distribution and timing of migration match the availability and quality of food, and thus the
 3708 importance of potential temporal mismatches between food availability and requirements is unknown. In addition,
 3709 changes in waterbird distribution are likely to alter inter- and intraspecific competition. Resolving these issues
 3710 requires investigation of effects at other levels of the food web (e.g. loss of bivalves from areas of reduced salinity,
 3711 species and size-class composition of fish communities) and their consequences for waterbirds. So far, knowledge
 3712 of climate change effects on waterbirds in the Baltic Sea are mostly restricted to ducks (including diving and
 3713 dabbling ducks), with much less known for other quantitatively important components of the waterbird
 3714 community, i.e. divers, grebes, waders, gulls and auks. Interactions between fish and piscivorous waterbirds in
 3715 particular need more attention. Responses to climate change are likely to vary between waterbird species and
 3716 groups. There is still little information on which species are most affected (negatively or positively) by changes in
 3717 climatic conditions and the uncertainty is therefore large on how species in future waterbird assemblages will
 3718 interact and the consequences for the functioning of the Baltic Sea. To gain a better understanding on how single
 3719 species (or groups with similar ecology, often closely-related) will respond to climate change is critical for
 3720 projecting effects of climate change on waterbirds around the Baltic Sea.



3721 **6.23.4 Marine food webs**

3722 Some changes observed in marine food webs have been partly by attributed to warming, brightening and sea ice
 3723 decline on long time-scales. Other drivers, such as eutrophication or fisheries, may however predominate and many
 3724 records are too short to allow attributing the observed changes to climate change. Although effects of warming,
 3725 ocean acidification and dissolved organic matter on some ecosystem functions have been identified in mesocosm
 3726 experiments, changing food web interactions are still impossible to project. It is, however, important to include
 3727 the marine biosphere in management-strategies for tackling the complex interactive aspects of climate change-
 3728 related effects on the marine ecosystem and human adaptations to them (Andersson et al., 2015; Stenseth et al.,
 3729 2020).

3730 **7 Key messages**

3731 The following lists selected key messages from this assessment that either confirm the conclusions of previous
 3732 assessments or are novel (marked with NEW). The estimated level of confidence based upon agreement and
 3733 evidence (see Section 2.3) of each key message refers to whether a systematic change in the considered variable
 3734 was detected and attributed to climate change. Climate change is here defined as the change in climate due to
 3735 human impact only (BACC II Author Team, 2015; see Section 2.1). Key messages referring to observed or
 3736 simulated changes in the Baltic Sea region caused by other drivers than climate change (e.g. afforestation,
 3737 eutrophication, fisheries, etc.) are not classified by a confidence level. A summary of all key messages related to
 3738 climate change is presented in Table 10 and Figure 34.

3739 **7.1 Past climate changes**

- 3740 • Large-scale circulation: The AMO has undergone frequency changes, but its influence on climate
 3741 variability in the Baltic Sea region remained similar, independent of the dominant frequency [NEW].
- 3742 • Air temperature: During the Holocene, The Baltic Sea region experienced periods as warm as the 20th
 3743 century, such as the mid-Holocene Optimum and the Medieval Warm Period. The implied rate of change
 3744 was, however, much slower than the present. The past warming signal was regionally markedly
 3745 heterogeneous, mostly along a west-east gradient [NEW].
- 3746 • Oxygen: The previous warm periods were accompanied by oxygen deficiency in the deeper waters of the
 3747 Baltic Sea, which cannot be attributed to eutrophication and was likely a result of climate forcing [NEW].

3748 **7.2 Present climate changes**

- 3749 • Large-scale atmospheric circulation: Systematic changes in large-scale atmospheric circulation related to
 3750 climate change could not be detected [low confidence]. The AMO is an important driver of climate
 3751 variability in the Baltic Sea region, affecting *inter alia* the correlation of regional climate variables with
 3752 the NAO.
- 3753 • Air temperature: Linear trends of the annual mean temperature anomalies during 1876–2018 were +0.10
 3754 °C decade⁻¹ north of 60°N and +0.09 °C decade⁻¹ south of 60°N in the Baltic Sea region [high confidence].
 3755 This is larger than the global mean temperature trend and slightly larger than estimated in the earlier
 3756 BACC reports [NEW]. The warm spell duration index has increased during 1950–2018 [medium
 3757 confidence]. Statistically significant decreases in winter cold spell duration index across the period 1979–



- 2013 have been widespread in Norway and Sweden, but less prevalent in eastern Finland, while changes in summer cold spell have been small in general [medium confidence].
- Solar radiation and cloudiness: Various satellite data products suggest a small but robust decline in cloudiness over the Baltic Sea region since the 1980s [low confidence, NEW]. However, whether this signal is an indicator of a changing climate or due to internal variability is unknown.
 - Precipitation: Since 1950, annual mean precipitation has generally increased in the northern part of the Baltic Sea region. There is some evidence of a long-term trend [low confidence]. However, long-term records suffer from inhomogeneity due to the increasing number of rain gauges. Frequency and intensity of heavy precipitation events have increased [medium confidence]. Drought frequency has increased across southern Europe and most of central Europe since 1950, but decreased in many parts of Northern Europe [low confidence].
 - Wind: Owing to the large internal variability, it is unclear whether there is an overall trend in mean wind speed. There has been an increase in the number of deep cyclones over central and Northern Europe since the late 1950s, but no evidence for a long-term trend [low confidence].
 - Air pollution: The influence of climate change on air pollution is small and undetectable, given the dominance of other human activities [low confidence]. Land-based emissions are declining due to emission control measures, but some emissions from the shipping sector may be increasing.
 - River discharge: For the period 1900-2008, no trend in total river discharge was found, but there was a pronounced 30-year variability. Data for some rivers in the northern Baltic Sea catchment indicate a long-term trend, but the confidence in these reconstructions is low. Since the 1970s, the total river winter discharge is increasing, perhaps due to warming or river regulations [low confidence]. Due to earlier snow-melt, driven by temperature increases in the region and a decreasing frequency of arctic air mass advection, high flow events in the Baltic Sea region shifted from late March to February. In Sweden, trends in the magnitude of high flow events over the past 100 years are not statistically significant [low confidence, NEW].
 - Riverine nutrient loads: The effect of changing climate on riverine nutrient loads is small and not detectable [low confidence].
 - Terrestrial biosphere: Combining all vegetation types in the entire Baltic Sea region, satellite observations suggest an advancement of the growing season by 0.30 day/year over the period 2000-2016. The most important driver of the advancement of the growing season is spring mean temperature, with an advancement rate of 2.47 day/°C of spring warming [medium confidence, NEW]. Observations and model results suggest cooling trends in daily minimum and warming trends in daily maximum temperatures in response to deforestation, and the opposite tendencies for afforestation. [NEW]
 - Snow: The decrease in snow cover has accelerated in recent decades, except in the mountain areas and the north-eastern part of the Baltic Sea region [high confidence, NEW]. On average, the number of days with snow cover has declined by 3–5 days per decade, [high confidence]. Mean and maximum snow depth has also decreased, most clearly in the southern and central part of the region [high confidence]. Whether sea-effect snowfall events have changed is unknown [low confidence].
 - Glaciers: Inventories of all Scandinavian glaciers, available only since 2006, show that they have lost 20 Gt of ice (~8% of their total mass) during 2006-2015. Atmospheric warming is very likely the primary driver of glacier mass loss [high confidence]. [NEW]



- 3799 • Permafrost: Recent warming has caused losses of over 20% of the original 6200 km² of permafrost in the
 3800 Baltic Sea catchment area during 1997-2018 [medium confidence, NEW].
- 3801 • Sea ice: Long-term decreases in sea ice in the Baltic Sea have exceeded the large natural climatological
 3802 variability and can only be attributed to global climate change [high confidence]. In addition,
 3803 unprecedented mild ice seasons have occurred in the last ten years, and 100-year trends in sea ice cover
 3804 showed an accelerated decline in 1921-2020 compared to 1910-2011 [high confidence, NEW].
- 3805 • Lake ice: Warming in the Baltic Sea catchment during recent decades has resulted in earlier ice break-
 3806 up, later freeze-up, and hence shorter ice cover duration on the lakes in the region. [high confidence,
 3807 NEW]
- 3808 • Water temperature: Monitoring data, satellite data and model-based historical reconstructions indicate an
 3809 increase in annual mean sea surface temperature averaged over the Baltic Sea of 0.4–0.6 °C decade⁻¹ or
 3810 ~1 – 2 °C since the 1980s [high confidence]. During 1856–2005, reconstructed SSTs increased by 0.03
 3811 and 0.06 °C decade⁻¹ in northeastern and southwestern areas, respectively. Hence, recent warming trends
 3812 have accelerated tenfold [NEW]. Long-term measurements at Tvärminne, on the north coast of the Gulf
 3813 of Finland, indicate that marine heat waves have increased since 1926 [low confidence].
- 3814 • Salinity and saltwater inflows: The record of major Baltic inflows (MBIs) has been revised and the earlier
 3815 reported decreasing trend is now seen as artifactual. On centennial time-scales, there are no statistically
 3816 significant trends in salinity averaged over the Baltic Sea (1920-2008) or in MBIs (1887-2017), but
 3817 pronounced multidecadal variability, with a period of about 30 years. Model results suggest that a decade
 3818 of decreasing salinity, like the 1983-1992 stagnation, happens about once a century due to natural
 3819 variability. Due to increased river runoff in the northern catchment, the North-South gradient in sea
 3820 surface salinity likely increased in 1900–2008 [low confidence, NEW].
- 3821 • Stratification and overturning circulation: No long-term trend in stratification was detected, but during
 3822 1982-2016 stratification increased in most of the Baltic Sea, with the seasonal thermocline and the
 3823 perennial halocline strengthening by 0.33–0.39 and 0.70–0.88 kg m⁻³, respectively [low confidence,
 3824 NEW].
- 3825 • Sea level: Since 1886 the mean sea level in the Baltic Sea relative to the geoid has increased by about 1-
 3826 2 mm per year, similar to the global mean rate [high confidence]. However, in the northern Baltic Sea
 3827 rapid land uplift causes a relative sea level decrease [high confidence]. Although an acceleration of the
 3828 mean sea level rise at individual stations could not yet be detected, the all-station-average-record showed
 3829 an almost statistically significant acceleration [low confidence, NEW]. Basin-wide, no statistically
 3830 significant, long-term changes in extreme sea levels relative to the mean sea level of the Baltic Sea could
 3831 be documented [low confidence, NEW].
- 3832 • Waves: Wave hindcasts and observations are too short for studies of climate-relevant trends [low
 3833 confidence].
- 3834 • Sedimentation and coastal erosion: Dominance of mobile sediments makes the southern and eastern
 3835 coasts more vulnerable to wind-wave induced transport than other Baltic Sea coasts [high confidence].
 3836 Prevailing westerly winds lead to mainly west-east sediment transport and an alternation of glacial till
 3837 cliffs (sources), sandy beaches and spits (sinks). No statistically significant, long-term changes were
 3838 found [low confidence, NEW].



- 3839 • Oxygen and nutrients: Reconstructions of oxygen conditions in the Baltic Sea for the period 1898-2012
 3840 suggest a tenfold increase of the hypoxic area, with current values of up to 70,000 km². This increase was
 3841 attributed mainly to increased nutrient loads, with a minor contribution from climate warming [low
 3842 confidence, NEW]. Furthermore, recently estimated oxygen consumption rates in the Baltic Sea are
 3843 higher than observed before, reducing the duration of improved oxygen conditions after natural
 3844 ventilation events by oxygen-enriched saltwater inflows.
- 3845 • Marine CO₂ system – air-sea exchange: In the period 1980-2005, sub-basins affected by high riverine
 3846 runoff and related high loads of terrestrial organic matter (e.g. Gulf of Bothnia) were found to be on
 3847 average a source of CO₂ to the atmosphere. This outgassing was more than compensated by the high CO₂
 3848 uptake by the open waters of the Baltic proper [medium confidence, NEW].
- 3849 • Marine CO₂ system – alkalinity: During 1900-2015, a long-term trend in alkalinity was observed, with
 3850 largest increases in the Gulf of Bothnia, where it almost entirely cancelled the pH decrease expected from
 3851 rising atmospheric pCO₂. The smaller alkalinity increase in the southern Baltic Sea compensated ocean
 3852 acidification by about 50%. Due to the high seasonal variability in pH, large interannual variability in
 3853 productivity and the identified alkalinity trend, no acidification was measurable in the central and northern
 3854 Baltic Sea [medium confidence, NEW].
- 3855 • Microbial communities: Long-term time series from 1994 to 2006 show that increased riverine dissolved
 3856 organic matter suppresses phytoplankton biomass production and shifts the carbon flow towards
 3857 heterotrophic microbes [low confidence, NEW].
- 3858 • Phytoplankton and cyanobacteria: The growing season for phytoplankton and cyanobacteria has
 3859 lengthened significantly in the past few decades [medium confidence] and the ratio between diatom and
 3860 dinoflagellate biomasses declined during the past century, probably due to warmer winters [low
 3861 confidence, NEW]. The annual chlorophyll maximum, in the 1980s associated with the spring diatom
 3862 bloom, has shifted to coincide with the summer cyanobacteria bloom [low confidence, NEW]. Although
 3863 inter-annually oscillating, surface cyanobacteria accumulations became a recurrent summer feature of the
 3864 southern Bothnian Sea in the 2010s [medium confidence, NEW].
- 3865 • Macroalgae: Long-term changes in Baltic Sea macroalgae and charophytes have been attributed to
 3866 changes in salinity, wind exposure, nutrient availability and water transparency as well as biotic
 3867 interactions [low confidence, NEW]. However, the role of climate change is unclear.
- 3868 • Zoobenthos: Increasing near-bottom temperature may partially explain the spreading of non-indigenous
 3869 species, such as polychaetes of the genus *Marenzelleria*. The effects on zoobenthos are primarily
 3870 synergistic, through e.g. eutrophication and hypoxia [low confidence, NEW].
- 3871 • Fish: Changes in temperature, salinity and species interactions can affect the stocks of cod, sprat and
 3872 herring. However, the dominant driver is the fishery. For coastal fish, the distribution of pikeperch
 3873 expanded northwards along the coasts of the Bothnian Sea, apparently due to the warming waters. For
 3874 many coastal fish species eutrophication is, however, equally or more important than climate change [low
 3875 confidence, NEW].
- 3876 • Marine mammals: Populations of ice-breeding seals, especially southern populations of the ringed seal,
 3877 have likely suffered from the sea ice decline [medium confidence]. However, this is based on occasional
 3878 ringed seal moult counts that indicating no population growth, while monitoring data on reproductive
 3879 success are missing.



- 3880 • Waterbirds: Many waterbird species have shifted their wintering range northwards [high confidence].
- 3881 They now migrate earlier in spring [medium confidence]. Effects of warming sea temperature are
- 3882 inconsistent, because both positive and negative effects on foraging conditions and food quality have
- 3883 been found [low confidence]. Most migrating Baltic Sea waterbirds are also affected by climate change
- 3884 outside the Baltic Sea [medium confidence].
- 3885 • Marine food webs: Significant alterations in food web structure and functioning such as the shift from
- 3886 early diatom to later dinoflagellate dominated blooms have been observed. However, the causes of these
- 3887 changes are unknown [low confidence].

3888 7.3 Future climate changes

- 3889 • Large-scale circulation: Projections suggest a more zonal flow over Northern Europe and a northward
- 3890 shift in the mean summer position of the westerlies at the end of the century [low confidence].
- 3891 • Air temperature: Coupled atmosphere-ocean regional climate models project an increase in annual mean
- 3892 air temperature by between 1.5 and 4.3°C over the Baltic Sea catchment area at the end of the century.
- 3893 The range indicates ensemble mean values for RCP2.6 and RCP8.5 scenarios. On average, air over
- 3894 surrounding land will warm about 0.1 to 0.4°C more than the air over the Baltic Sea [high confidence,
- 3895 NEW]. A bias-adjusted median estimate of increase in warm spell duration index in Scandinavia for the
- 3896 period 2071-2100, compared to 1981-2010, was about 15 days under RCP8.5, with an uncertainty range
- 3897 of about 5-20 days [medium confidence]. The cold spell duration index in Northern Europe is projected
- 3898 to decrease in the future, with a likely range of from -5 to -8 days per year by 2071-2100, compared to
- 3899 1971-2000 [medium confidence].
- 3900 • Solar radiation and cloudiness: Projections for solar radiation and cloudiness differ systematically in sign
- 3901 between global and regional climate models, indicating high uncertainty [low confidence, NEW].
- 3902 • Precipitation: Annual mean precipitation is projected to increase over the entire Baltic Sea catchment at
- 3903 the end of the century [medium confidence]. The signal is robust for winter among the various regional
- 3904 climate models but is highly uncertain for summer in the south. The intensity and frequency of heavy
- 3905 rainfall events are projected to increase. These increases are even larger for convection-resolving models
- 3906 [high confidence, NEW]. Projections show that the number of dry days in the southern and central parts
- 3907 of the Baltic Sea basin increases mainly in summer [low confidence].
- 3908 • Wind: Changes in wind over the Baltic Sea region are highly uncertain [low confidence]. Over sea areas
- 3909 where the average ice cover is projected to diminish, such as the Bothnian Sea and the eastern Gulf of
- 3910 Finland, the mean wind is projected to increase because of a warmer sea surface and reduced stability of
- 3911 the planetary boundary layer [low confidence].
- 3912 • Air pollution: The impact of climate change on air quality and atmospheric deposition is smaller than the
- 3913 assumed impact of future changes in emissions [low confidence].
- 3914 • River discharge: River runoff is projected to increase 2–22% in RCP4.5 and 7–22% in RCP8.5. River
- 3915 discharge is projected to increase to the northern and decrease to the southern sub-basins [low
- 3916 confidence]. High flows are projected to decrease in spring and increase in autumn and winter due to
- 3917 earlier snow melt and more winter rain. Over much of continental Europe, an increase in intensity of high
- 3918 flow events is projected with increasing temperature [low confidence].



- 3919 • Land nutrient inputs: The impact of climate change on land nutrient inputs is smaller than the impact of
 3920 changes in land management, populations and nutrient point-source releases. In any given river, larger
 3921 runoff would lead to larger nutrient inputs [low confidence].
- 3922 • Terrestrial growing season: Projections suggested that decreasing surface albedo in the Arctic region in
 3923 winter and spring will notably amplify the future warming in spring (positive feedback), while the
 3924 increased evapotranspiration will lead to a marked cooling during summer (negative feedback). These
 3925 feedbacks will stimulate vegetation growth, due to an earlier start of the growing season, leading to
 3926 compositional changes in woody plants and the distribution of vegetation. Arctic terrestrial ecosystems
 3927 could continue to sequester carbon until the 2060-2070s, after which the terrestrial ecosystems are
 3928 projected to turn into weak sources of carbon due to increased soil respiration and biomass burning [low
 3929 confidence, NEW].
- 3930 • Terrestrial carbon sequestration: Mitigation scenarios that decrease the fraction of coniferous forest in
 3931 favour of deciduous forest, and increase the area of deciduous forest in Northern Europe from 130,000 to
 3932 480,000 km², were projected to reduce near-surface temperatures and give maximum carbon
 3933 sequestration. [NEW]
- 3934 • Snow: Projections under RCP8.5 suggest a reduction of the average snow amount by more 70% for most
 3935 areas, with the exception of the high Scandinavian mountains, where the warming temperature does not
 3936 reach the freezing point as often as in lower-lying regions [high confidence]. Sea-effect snowfall events
 3937 in future climate have not been investigated yet.
- 3938 • Glaciers: Scandinavian glaciers will lose more than 80% of their current mass by 2100 under RCP8.5,
 3939 and many are projected to disappear, regardless of future emission scenarios [high confidence, NEW].
 3940 Furthermore, river runoff from glaciers is also projected to change regardless of the emission scenario,
 3941 and to result in increased average winter runoff and in earlier spring peaks [high confidence, NEW].
- 3942 • Permafrost: In the future climate, the on-going loss of permafrost in the Baltic Sea catchment will very
 3943 likely accelerate [high confidence].
- 3944 • Sea ice: Regional climate projections consistently project shrinking and thinning of Baltic Sea ice cover
 3945 [high confidence], but still estimate that some ice will be formed even in mildest future winters. However,
 3946 those estimates are based on a limited number of ensembles and may not represent future climate
 3947 variability correctly.
- 3948 • Lake ice: The observed trends of earlier ice break-up, later freeze-up, and shorter ice cover duration on
 3949 lakes in the region are projected to continue with future warming, and lakes with intermittent winter ice
 3950 will consequently become increasingly abundant [high confidence, NEW].
- 3951 • Water temperature: Coupled atmosphere-ocean regional climate models project an increase in annual
 3952 mean SST of between 1.2 and 3.2°C, averaged for the Baltic Sea the end of the century. The range
 3953 indicates ensemble mean values for RCP2.6 and RCP8.5 scenarios. Warming will be largest in summer
 3954 in the northern Baltic Sea [high confidence]. Under both RCP4.5 and RCP8.5, record-breaking summer
 3955 mean SSTs were projected to increase at the end of the century [medium confidence, NEW]. However,
 3956 due to the pronounced internal variability there might be decades in the near future without record-
 3957 breaking events.
- 3958 • Salinity and saltwater inflows: An increase in river runoff or westerly winds will tend to decrease salinity,
 3959 but a global sea level rise will tend to increase it, because an enlarged cross-sectional area of the Danish



- 3960 Straits will increase the saltwater imports from the Kattegat. Due to the large uncertainty in projected
 3961 river runoff, wind and global sea level rise, salinity projections show a wide spread, from increasing to
 3962 decreasing salinities, and no robust changes were identified [low confidence, NEW].
- 3963 • Stratification and overturning circulation: Considering all potential drivers of changes in salinity in the
 3964 Baltic Sea (wind, river runoff, net precipitation, global sea level rise), neither the haline-induced
 3965 stratification nor the overturning circulation is projected to change [low confidence, NEW]. Projections
 3966 consistently show that the seasonal thermocline during summer will intensify across nearly the whole
 3967 Baltic Sea [high confidence, NEW].
 - 3968 • Sea level: Future absolute sea level in the Baltic Sea will continue to rise with the global mean sea level
 3969 [high confidence]. Its regional manifestation is, however, modulated by the future melting of Antarctica,
 3970 which affects the Baltic Sea more strongly than the melting of Greenland [low confidence]. Using current
 3971 estimates, the regional mean sea level is projected to rise by about 87% of the global mean sea level. Land
 3972 uplift is roughly known but difficult to estimate accurately in practice, as many regional geological factors
 3973 blur the signature of the glacial isostatic adjustment. Trends in sea level extremes will be determined by
 3974 the changing mean sea level and possible future changes in storminess. The uncertainty in the latter driver
 3975 is very large [low confidence].
 - 3976 • Waves: The projected decrease in seasonal sea ice cover will have considerable effects on the wave
 3977 climate in the northernmost Baltic Sea [high confidence, NEW]. Otherwise, there are no conclusive
 3978 results on possible changes in the wave climate and wave extremes, because of the uncertainty about
 3979 changes in wind fields [low confidence].
 - 3980 • Sedimentation and coastal erosion: Changes in sea level, wind, waves and sea ice all affect sediment
 3981 transport and coastal erosion. Hence, available projections are highly uncertain [low confidence, NEW].
 - 3982 • Oxygen and nutrients: The future response of deep water oxygen conditions will mainly depend on future
 3983 nutrient inputs from land [medium confidence]. However, coastal hypoxia might increase due to warming
 3984 of the water in shallow areas [medium confidence]. Implementation of the BSAP will lead to declining
 3985 phosphorus concentrations [medium confidence, NEW].
 - 3986 • Marine CO₂ system: Due to anthropogenic emissions, atmospheric pCO₂ will rise, and consequently also
 3987 the mean pCO₂ of Baltic surface seawater, which has the potential to lower pH [high confidence].
 3988 However, the magnitude of the pH change also depends on alkalinity trends, which are highly uncertain
 3989 [low confidence]. Hence, projections for the Baltic Sea are different from the global ocean.
 - 3990 • Microbial communities: The impact of climate change on microbes and the functioning of the microbial
 3991 loop have been studied experimentally. In the northern Gulf of Bothnia, adding DOM increased the
 3992 abundance of bacteria, whereas a temperature increase (from 12 to 15°C) reduced their abundance [low
 3993 confidence, NEW].
 - 3994 • Phytoplankton and cyanobacteria: The effect of climate change on phytoplankton and cyanobacteria
 3995 blooms is larger under high nutrient concentrations, but nutrient loads are the dominant driver. If the
 3996 BSAP is fully implemented, the projected environmental status of the Baltic Sea will be significantly
 3997 improved, and extreme cyanobacteria blooms will be rare or absent [low confidence, NEW].
 - 3998 • Zooplankton: Experimental studies suggested improved conditions for microzooplankton due to warming
 3999 but negative effects on some larger zooplankton species [low confidence, NEW].



- 4000 • Macroalgae and vascular plants: The direct and indirect effects of changes in temperature, salinity and
 4001 pH are likely to change the geographic distribution of Baltic Sea macrophytes. However, neither
 4002 experimental studies nor past observed changes provide conclusive projections for the effects of climate
 4003 change [low confidence, NEW].
- 4004 • Zoobenthos: In a warmer and less eutrophic Baltic Sea, benthic-pelagic coupling will be weaker, resulting
 4005 in decreasing benthic biomass [low confidence, NEW].
- 4006 • Non-indigenous species: Climate change may favour invasions of non-indigenous species. However, it is
 4007 impossible to project which species may enter the Baltic Sea in future [low confidence].
- 4008 • Fish: Projected changes in temperature and salinity will affect the stocks of cod, sprat and herring.
 4009 However, nutrient loads and especially fishing mortality are also important drivers. Although multi-driver
 4010 modeling studies have been performed, the impact of climate change is uncertain [low confidence, NEW].
- 4011 • Marine mammals: Mild winters are known to negatively affect Baltic ringed seals (*Phoca hispida*
 4012 *botnica*) because without their sea ice lair, the pups are more vulnerable to weather and predators, and it
 4013 has been projected that the growth rates of ringed seal populations will decline in the next 90 years. Also
 4014 for grey seals (*Halichoerus grypus*), it has been suggested that reduced ice cover in combination with
 4015 (partly climate-driven) changes in the food web, may affect their body condition and birth rate [low
 4016 confidence].
- 4017 • Waterbirds: The northward distributional shifts of waterbirds are expected to continue [medium
 4018 confidence]. Effects on waterbird food will be manifold, but consequences are difficult to predict [low
 4019 confidence]. The rising sea level and erosion are expected to reduce the availability of breeding habitats
 4020 [low confidence].
- 4021 • Marine food webs: Significant alterations in food web structure and functioning can be expected, since
 4022 species distributions and abundances are expected to change with warming sea water. The consequences
 4023 are difficult to project, as research into the long-term dynamics of food webs is still scarce [low
 4024 confidence].

4025 8 Concluding remarks

4026 We found that

- 4027 1. The overall conclusions of the BACC I and BACC II assessments remain valid.
- 4028 2. However, new coupled models (atmosphere-ice-ocean, atmosphere-land), larger ensembles of scenario
 4029 simulations (CORDEX), new mesocosm experiments (warming, ocean acidification, and dissolved
 4030 organic matter), extended monitoring (glaciers, satellite data) and homogenized records of observations
 4031 (MBIs) have led to new insights into past and future climate variability.
- 4032 3. Improved paleoclimate simulations of the Holocene, new dendroclimatological reconstructions of the
 4033 past 1000 years and new climate regionalizations have added regional details (east-west gradients over
 4034 the Baltic Sea region) and improved our understanding of internal variability (sea level extremes,
 4035 stagnation periods) and the remote impact of low-frequency North Atlantic variability on the Baltic Sea
 4036 region (AMO, Baltic Sea salinity). New sediment cores suggest that hypoxia during the Medieval Climate
 4037 Anomaly was caused by climate variability, rather than by human influence, as claimed earlier.



4. Natural variability of many variables of the Earth system is larger than previously realized, requiring larger model ensembles for convincing future projections. Although the first, relatively large ensemble of scenario simulations utilizing a regional coupled atmosphere-ice-ocean model has become available, uncertainty estimates are still incomplete.
5. New regional ESMs including additional components of the Earth system are under development. However, the simulated water cycle is still biased.
6. The first complex multiple-driver study with focus on present and future climates addressing for instance eutrophication of the Baltic Sea, fisheries and climate change has become available and an overall assessment of the various drivers in the Baltic Sea region is part of the BEARs. However, further research on the interplay between drivers is needed.
7. More research on changing extremes was performed, acknowledging that the impact of changing extremes may be more important than that of changing means. However, most observational records are either too short or too heterogeneous for statistical studies of extremes.
8. The climate change signal is still confined to increases in observed air and water temperatures, to decreases in sea and lake ice, snow cover, permafrost and glacier mass, to the rise in mean sea level, and to variables directly related to temperature and the cryosphere, such as ringed seal habitats. Compared to the previous BACC report, changes in air temperature, sea ice, snow cover and sea level were shown to have accelerated.
9. Intensive research on the land-sea interface focussing on the coastal filter has been performed and nutrient retention in the coastal zone was estimated for the first time. Uncertainty concerning the bioavailability of nutrient loads was identified as one of the foremost challenges for marine biogeochemistry. However, a model for the entire Baltic Sea coastal zone is still missing and the effect of climate change on the coastal filter capacity is still unknown.
10. In contradiction to earlier results, observed MBIs have no declining trend. Due to the uncertainties in projections of the regional wind, regional precipitation and evaporation, river discharge and global mean sea level rise, projections of salinity in the Baltic Sea are uncertain and it remains unknown whether the Baltic Sea will become less or more salty. As salinity is a crucial variable for the marine ecosystem and for Baltic circulation, projections for the Baltic Sea as a whole are regarded as uncertain.
11. The Baltic Sea may become more acidic in the future, but the decrease in pH may partly be compensated by an alkalinity increase, as in the past. Hence, past changes in Baltic carbonate chemistry were different from the global ocean acidification, and pH changes may differ also in future.
12. Large marine food web changes were observed, which could partly be attributed to warming, brightening and sea ice decline. However other factors also play important roles, and many records are too short for attribution studies.

Author contributions

Chapter	Title	Authors
1	Introduction	
1.1	Overview	H.E.M. Meier
1.2	The BACC process	M. Reckermann, H.E.M. Meier



1.3	Summary of BACC I and BACC II key messages	H.E.M. Meier
1.4	Baltic Sea Region characteristics	K. Myrberg
1.5	Global climate change	M. Gröger
2	Methods	
2.1	Assessment of literature	H.E.M. Meier
2.2	Climate model data	H.E.M. Meier
2.3	Uncertainty estimates	H.E.M. Meier
3	Current state of knowledge	
3.1	Past climate change	E. Zorita
3.2	Present climate change	
3.2.1	Present climate change - Atmosphere	
3.2.1.1	Large-scale circulation	M. Stendel, C. Frauen, F. Börgel, H.E.M. Meier, M. Kniebusch
3.2.1.2	Air temperature	A. Rutgersson
3.2.1.3	Solar radiation	T. Carlund, A. Rutgersson
3.2.1.4	Precipitation	J. Käyhkö, E. Kjellström
3.2.1.5	Wind	M. Stendel
3.2.1.6	Air pollution, air quality and atmospheric nutrient deposition	M. Quante
3.2.2	Present climate change - Land	
3.2.2.1	River discharge	J. Käyhkö, G. Lindström
3.2.2.2	Land nutrient inputs	O.P. Savchuk, B. Müller-Karulis
3.2.3	Terrestrial biosphere	W. May, P.A. Miller
3.2.4	Present climate change - Cryosphere	
3.2.4.1	Snow	J. Jaagus
3.2.4.2	Glaciers	N. Kirchner
3.2.4.3	Permafrost	G. Hugelius
3.2.4.4	Sea ice	J.J. Haapala
3.2.4.5	Lake ice	J. Käyhkö
3.2.5	Present climate change – Ocean and marine sediments	
3.2.5.1	Water temperature	C. Dieterich, H.E.M. Meier
3.2.5.2	Salinity and saltwater inflows	V. Mohrholz, H.E.M. Meier, K. Myrberg, A. Lehmann



3.2.5.3	Stratification and overturning circulation	M. Gröger, H.E.M. Meier, K. Myrberg
3.2.5.4	Sea level	B. Hünicke, E. Zorita, C. Dieterich, R. Weisse
3.2.5.5	Waves	L. Tuomi
3.2.5.6	Sedimentation and coastal erosion	W. Zhang
3.2.5.7	Marine carbonate and biogeochemistry	K. Kulinski, J. Carstensen, B. Müller-Karulis, O. Savchuk
3.2.6	Marine biosphere	M. Viitasalo, E. Bonsdorff, R. Elmgren, A. Galatius, M. Ahola, V. Dierschke, I. Carlen, M. Frederiksen, E. Gaget, A. Halkka, M. Jüssi, D. Pavon-Jordan
3.3	Future climate change	All
3.3.1	Future climate change - Atmosphere	
3.3.1.1	Large-scale circulation	A. Rutgersson, F. Börgel, C. Frauen
3.3.1.2	Air temperature	A. Rutgersson, E. Kjellström, O.B. Christensen
3.3.1.3	Solar radiation	A. Rutgersson, T. Carlund
3.3.1.4	Precipitation	E. Kjellström, O.B. Christensen
3.3.1.5	Wind	M. Stendel, H.E.M. Meier
3.3.1.6	Air pollution, air quality and atmospheric nutrient deposition	M. Quante
3.3.2	Future climate change - Land	
3.3.2.1	River discharge	J Käyhkö
3.3.2.2	Riverine nutrient loads	R. Capell, A. Bartosova
3.3.3	Terrestrial biosphere	W. May, P.A. Miller
3.3.4	Future climate change - Cryosphere	
3.3.4.1	Snow	O.B. Christensen
3.3.4.2	Glaciers	N. Kirchner
3.3.4.3	Permafrost	G. Hugelius
3.3.4.4	Sea ice	J.J. Haapala
3.3.4.5	Lake ice	J. Käyhkö
3.3.5	Future climate change – Ocean and marine sediments	
3.3.5.1	Water temperature	C. Dieterich, H.E.M. Meier, M. Gröger
3.3.5.2	Salinity and saltwater inflows	H.E.M. Meier
3.3.5.3	Stratification and overturning circulation	M. Gröger, C. Dieterich, H.E.M. Meier



3.3.5.4	Sea level	B. Hünicke, E. Zorita, C. Dieterich, R. Weisse
3.3.5.5	Waves	L. Tuomi
3.3.5.6	Sedimentation and coastal erosion	W. Zhang
3.3.5.7	Marine carbonate and biogeochemistry	K. Kulinski, J. Carstensen, B. Müller-Karulis, O. Savchuk
3.3.6	Marine biosphere	M. Viitasalo, E. Bonsdorff, R. Elmgren, A. Galatius, M. Ahola, V. Dierschke, I. Carlen, M. Frederiksen, E. Gaget, A. Halkka, M. Jüssi, D. Pavon-Jordan
4	Interactions with other drivers	M. Reckermann
5	Comparison with the North Sea region	M. Quante
6	Knowledge gaps	H.E.M. Meier and All
7	Key messages	H.E.M. Meier and All
8	Concluding remarks	H.E.M. Meier and All
Figures and Tables	Analysis of observed time series	M. Kniebusch
Figures and Tables	Analysis of scenario simulations	M. Kniebusch, H.E.M. Meier, C. Dieterich, M. Gröger, O.P. Savchuk, G. Lindström, N. Kirchner, E. Zorita, G. Hugelius

4073 Acknowledgements

4074 The research presented in this study is part of the Baltic Earth Assessment Reports project of the Baltic
 4075 Earth program (Earth System Science for the Baltic Sea region, see <http://www.baltic.earth>). Glacier mass
 4076 loss data were obtained from the SITES Data Portal (<https://data.fieldsites.se/portal/>, see World Glacier
 4077 Monitoring Service (WGMS, 2020) and Swedish Infrastructure for Ecosystem Science (SITES, 2021a, b, c).
 4078 We thank Berit Recklebe for technical support and preparation of the reference list.
 4079



4080 **Figures**




Figure 1: The Baltic Sea and its catchment area, showing climatological mean salinity (in g kg^{-1}) and river runoff (in mm year^{-1}). (Source: Meier et al., 2014)





Figure 2: Bottom topography of the Baltic Sea and locations of the monitoring stations Arkona Deep (BY2), Bornholm Deep (BY5), Gdansk Deep (BMPL1), Gotland Deep (BY15), Northern Deep (OMTF 0286), Landsort Deep (BY15), and Åland Sea (F64). The Baltic proper comprises the Arkona Basin, Bornholm Basin and Gotland Basin.



4112



4113

4114 **Figure 3:** Orbital forcing (irradiance) at 60°N in January and July (derived from Laskar et al., 2004) and the
 4115 anomalies of reconstructed seasonal temperature and precipitation compared to preindustrial climate (Mauri et al.,
 4116 2015) in the Baltic Sea region at the Mid-Holocene Optimum (6000 before present).

4117





Figure 4: Reconstructed spring-early-summer air temperature in the Baltic Sea region (land areas in the box 0-40°E x 55-70°N, deviations from the 20th century mean) derived from Luterbacher et al. (2016). The record is smoothed by a 30-year low-pass filter. The approximate uncertainty range has been estimated here from the data provided by the original publication at interannual and grid-cell scale.




4124


4125

4126 **Figure 5:** Normalized winter (December through March; DJFM) mean NAO index during 1821/22-2018/19. Red:
 4127 positive, blue: negative, black: 10-year running mean. Normalization: $(\text{data} - \text{mean}(\text{data})) / \text{standard deviation}(\text{data})$.
 4128 (Data source: <https://crudata.uea.ac.uk/cru/data/nao/nao.dat>, compiled by Madline Kniebusch, Leibniz Institute for
 4129 Baltic Sea Research Warnemünde)


4130




4131
 4132 **Figure 6:** Normalized annual mean AMO index during 1882-2018. Red: positive, blue: negative, black: 10-year
 4133 running mean. Normalization: $(\text{data} - \text{mean}(\text{data})) / \text{standard deviation}(\text{data})$. (Data source:
 4134 https://climexp.knmi.nl/data/iamo_hadsst_ts.dat, compiled by Madline Kniebusch, Leibniz Institute for Baltic Sea
 4135 Research Warnemünde)
 4136



4137
 4138 **Figure 7:** Annual and seasonal mean near-surface air temperature anomalies for the Baltic Sea basin for
 4139 1871–2020, taken from the CRUTEM4v dataset (Jones et al., 2012), compiled by Anna Rutgersson, Uppsala
 4140 University. Blue, red: Baltic Sea basin region north and south, respectively, of 60°N. Dots: individual years.
 4141 Smoothed curves: variability on timescales longer than 10 years.
 4142



4143
 4144 **Figure 8:** Homogenized annual and seasonal mean temperature in Stockholm during 1756-2018 measured at Bolin
 4145 Centre, Stockholm. Each colored bar show the annual mean temperature, in red or blue, depending on whether the
 4146 temperature is above or below the average during the reference period 1961-1990. The black curve represents
 4147 smoothed 10-year mean temperatures. (Source: <https://bolin.su.se/data/stockholm-historical-temps-monthly>)
 4148



4149
 4150 **Figure 9:** Mean annual precipitation over land in mm year^{-1} in the Baltic Sea catchment area during 1950-2018.
 4151 Blue: whole catchment area, green: North of 59°N , Red: South of 59°N . Bars: annual sum, black: 10-year running
 4152 mean. (Data source: http://surfobs.climate.copernicus.eu/dataaccess/access_eobs.php#datafiles, compiled by
 4153 Madline Kniebusch, Leibniz Institute for Baltic Sea Research Warnemünde). Trends: $1.44 \text{ mm year}^{-1}$ (total), 1.51
 4154 mm year^{-1} (North), $1.37 \text{ mm year}^{-1}$ (South), significant on 99% using the phase-scrambling method (Kniebusch et
 4155 al., 2019b).
 4156




Figure 10: Area weighted river runoff anomalies relative to 1960-1990 (in %) from Sweden to the Baltic Sea. The black solid curve denotes Gaussian filtered data with a standard deviation of three years. (Source: Göran Lindström, Swedish Meteorological and Hydrological Institute).






Figure 11: Long-term dynamics (1970–2017) of annual nitrogen (A) and phosphorus (B) land inputs to the major Baltic Sea basins: BB - Bothnian Bay; BS – Bothnian Sea; BP - Baltic proper; GF - Gulf of Finland; GR - Gulf of Riga; DS – Danish straits; KT – Kattegat. Time (years) is on the horizontal axis. (Source: O.P. Savchuk, Stockholm University)



4168
 4169 **Figure 12:** Cumulative mass loss for four Swedish glaciers: Storglaciären (since 1946), Rabots glaciär (since 1982,
 4170 no data for 2004 and 2007 and hence interpolated), Riukojietna (since 1986, data for 2004 interpolated), och
 4171 Mårnaglaciär (since 1990, no data for 2020). Data are accessible from the SITES Data Portal,
 4172 <https://data.fieldsites.se/portal/> (Swedish Infrastructure for Ecosystem Science, 2021a, c, b). (Source: Nina
 4173 Kirchner, Stockholm University)
 4174



4175
 4176 **Figure 13:** Modeled permafrost extent of alpine (> 700 m a.s.l.) and lowland permafrost for the years 1997-2018
 4177 in the Baltic Sea drainage basin. Permafrost data from (Obu et al., 2020), extent of catchment from (Hannerz and
 4178 Destouni, 2006) and elevation data from USGS Global Multi-resolution Terrain Elevation Data 2010
 4179 (GMTED2010). Analyses performed at 1 km resolution in an equal area projection. (Source: Gustav Hugelius,
 4180 Stockholm University)
 4181




Figure 14: Left: Annual maximum sea ice extent of the Baltic Sea (MIB) in km^2 during 1720-2020. Blue bars: annual, red: 15-year running mean. Right: 30-year distribution functions of MIB during 1909-1938 and 1991-2020. (Data sources: https://www.eea.europa.eu/data-and-maps/daviz/maximum-extent-of-ice-cover-3#tab-chart_1, website Finnish Meteorological Institute: <https://en.ilmatieteenlaitos.fi/ice-season-in-the-baltic-sea>). (Source: Jari Haapala, Finnish Meteorological Institute)






Figure 15: An average sea ice thickness distribution in the Bay of Bothnia. Statistics is based on helicopter electromagnetic measurements conducted in winters 2003, 2004, 2005, 2007 and 2011 (Source: Ronkainen et al., 2018).



4194
 4195 **Figure 16:** Level ice thickness at Kemi, Finland and Loviisa, Finland during 1912-2019. Points: annual mean
 4196 values, lines: linear trend with 95% confidence intervals (Data source: Jari Haapala, Finnish Meteorological
 4197 Institute).
 4198



4199



4200

4201

Figure 17: Length of the ice season in days at Kemi, Loviisa and Utö (Finland) during 1890-2019. Points: annual mean, lines: linear trend with 95% confidence intervals (Data source: Jari Haapala, Finnish Meteorological Institute).

4202

4203

4204




Figure 18: Annual mean values of de-seasonalized daily sea surface (left) and bottom (right) temperature (red points) at seven monitoring stations during 1877-2018. For the location of the stations see Figure 2. The grey lines show the period when every station has data for every year (1954-2018). For Figures 18, 19, 21 and 22, ICES data (<https://ocean.ices.dk/HydChem/>) for temperature and salinity (bottle data, i.e. from specific depths) were used. Post processing of the data was done following Radtke et al. (2020) in order to overcome possible seasonal biases due to missing values in the observations. Therefore, gaps were statistically filled using a GAMM model (general additive mixed models) taking the seasonality into account. (Source: Madline Kniebusch, Leibniz Institute for Baltic Sea Research Warnemünde)




Figure 19: Seasonal mean sea surface and bottom temperature values during 1877-2018 at Gotland Deep (BY15). Blue: winter, green: spring, yellow: summer and orange: autumn. The grey lines show the period when every station has data for every year (1954-2018). (Source: Madline Kniebusch, Leibniz Institute for Baltic Sea Research Warnemünde)




Figure 20: Annual sea surface temperature anomalies to the reference period 2002-2018 of de-seasonalized measurements at BY2 and the MARNET stations Darss Sill and Arkona Basin during 1980-2018. (Source: Madline Kniebusch, Leibniz Institute for Baltic Sea Research Warnemünde)




Figure 21: Annual mean values of de-seasonalized daily sea surface (left) and bottom (right) salinity (blue points) at seven important stations during 1877-2018. The grey lines show the period when every station has data for every year (1954-2018). (Source: Madline Kniebusch, Leibniz Institute for Baltic Sea Research Warnemünde)





Figure 22: Difference between bottom and surface salinity as a measure for the vertical stratification (blue points) during 1877-2018. Only time steps when both values were available are considered. The grey lines show the period when every station has data for every year (1954-2018). (Source: Madline Kniebusch, Leibniz Institute for Baltic Sea Research Warnemünde)



4237
 4238 **Figure 23:** Annual mean sea level changes in centimeters for 14 Swedish mareographs since 1886. The data are
 4239 corrected for land uplift. The grey line shows a smoothed curve. (Source: Swedish Meteorological and
 4240 Hydrological Institute)
 4241






Figure 24: Annual mean sea level in Stockholm during 1774-2018. Grey bars: historic time series from Ekman (1988), red: SMHI data (RH2000, 1889-2018), blue: trend computed for 1774-1884 (estimated land uplift: 4.9 mm year⁻¹) and extrapolated until 2018. The SMHI data has been bias corrected (mean difference during overlapping time period) to make both time series comparable. Sea level rise of SMHI data corrected by estimated land uplift amounts 1.13 mm year⁻¹ during 1889-2018. (Source: Madline Kniebusch, Leibniz Institute for Baltic Sea Research Warnemünde)



4250

4251 **Figure 25:** Extent of hypoxic ($< 2 \text{ mL O}_2 \text{ L}^{-1}$) and anoxic ($< 0 \text{ mL O}_2 \text{ L}^{-1}$) bottom water (in km^2) in the Baltic
 4252 proper, Gulf of Finland and Gulf of Riga during cruise in August–October 1960–2020. (Source: Swedish
 4253 Meteorological and Hydrological Institute)



T2m (2070-2099) minus (1970-1999)




Figure 26: (a) Ensemble mean 2 m air temperature change (°C) between 2070-2099 and 1970-1999 for winter (December through January, upper panels) and summer (June through August, lower panels) under RCP2.6, RCP4.5 and RCP8.5. (b) as (a) but for precipitation change (mm day⁻¹). Eight different Earth System Models are used. (Data source: Gröger et al., 2021b)






Figure 27: Mean projected glacier mass evolution between 2015 and 2100 relative to each region's glacier mass in 2015 (in %) and ± 1 standard deviation under RCP2.6, RCP4.5 and RCP8.5. (Source: Hock et al., 2019)




4265
 4266 **Figure 28:** Median, 5th and 95th percentiles of the annual maximum ice extent of the Baltic Sea estimated from 28
 4267 CMIP5 models. The vertical axis shows upper class limits for mild and average ice winters. (a) RCP4.5, (b)
 4268 RCP8.5. (Source: Luomaranta et al., 2014)
 4269



4270

SST (2070-2099) minus (1970-1999)



4271

Standard deviation

4272 **Figure 29:** (a) Ensemble mean sea surface temperature change (°C) between 2070-2099 and 1970-1999 for winter
 4273 (December through January, upper panels) and summer (June through August, lower panels) under RCP2.6,
 4274 RCP4.5 and RCP8.5. (b) as (a) but for the standard deviation of the change, i.e. the ensemble spread (°C). Eight
 4275 different Earth System Models are used. (Source: Gröger et al., 2019)




Figure 30: From left to right changes of summer (June – August) mean sea surface temperature (SST; °C), annual mean sea surface salinity (SSS; g kg⁻¹), annual mean bottom salinity (BS; g kg⁻¹), and winter (December – February) mean sea level (SL; cm) between 1978-2007 and 2069-2098 are shown. From top to bottom results of the ensembles by Meier et al. (2011a) under the A1B/A2 greenhouse gas emission scenario (white background), and by Saraiva et al. (2019b), RCP 4.5 (grey background) and RCP 8.5 (grey background) are depicted.




Figure 31: Ensemble mean summer (June – August) bottom dissolved oxygen concentration changes (mL L^{-1}) between 1978–2007 and 2069–2098. From left to right results of the nutrient load scenarios Baltic Sea Action Plan (BSAP), Reference (REF) and Business-As-Usual (BAU) are shown. From top to bottom results of the ensembles by Meier et al. (2011a) under the A1B/A2 greenhouse gas emission scenario (white background), and by Saraiva et al. (2019b), RCP 4.5 (grey background) and RCP 8.5 (grey background) are depicted.




Figure 32: As Figure 30 but for annual mean Secchi depth changes (m). Secchi depth changes indicate changes in water transparency caused by phytoplankton and detritus concentration changes.





Figure 33: Sea surface temperature anomaly in the Greater North Sea region from 1870 to 2016 (relative to the mean 1971 to 2000), according to different data sets. (Huthnance et al., 2016), updated by Elizabeth Kent, Southampton)



4299
 4300 **Figure 34:** Synthesis of the knowledge on present and future climate changes. Shown are the anthropogenic
 4301 climate changes in 33 Earth system variables (bubbles) of the atmosphere (yellow), land surface (brown), terrestrial
 4302 biosphere (dark green), cryosphere (grey), ocean and sediment (blue), and marine biosphere (light green). The sign
 4303 of a change (plus/minus) is shown together with the level of confidence denoted by the number of signs, i.e. one
 4304 to three signs correspond to low, medium and high confidence levels, as the result of the literature assessment
 4305 reflecting consensus and evidence following the IPCC definitions (Section 2.3). Sign colours indicate the direction
 4306 of past (black) and future (red) changes following Table 10. Uncertain changes (+/-) are not displayed. Investigated
 4307 external anthropogenic drivers of the Earth system are shown as red squares, i.e. greenhouse gases, in particular
 4308 CO₂, and aerosol emissions. Climate change attribution relationships with sufficiently high confidence are shown
 4309 by arrows (black: heat cycle, orange: water cycle, pink: carbon cycle). Projections of carbon sequestration of Arctic
 4310 terrestrial ecosystems for the 21st century showed first increased uptake and later a carbon source (Section 3.3.3),
 4311 denoted by 1. + 2. -.
 4312



4313

4314 **Tables**

4315 **Table 1:** Variables of this assessment and further reference (1: Lehmann et al., 2021; 2: Kuliński et al., 2021; 3:
 4316 Rutgersson et al., 2021; 4: Weisse et al., 2021; 5: Reckermann et al., 2021; 6: Gröger et al., 2021a; 7: Christensen
 4317 et al., 2021; 8: Meier et al., 2021a; 9: Viitasalo, 2021)

Number	Variable	Past and present climates		Future climate	
Atmosphere					
1	Large-scale circulation	3.2.1.1	3	3.3.1.1	3, 7
2	Air temperature	3.1.2, 3.1.3, 3.1.4		3.3.1.2	7
	Warm spell	3.2.1.2	3		3
	Cold spell		3		3
3	Solar radiation and cloudiness	3.2.1.3		3.3.1.3	7
4	Precipitation	3.1.2, 3.1.3, 3.1.4		3.3.1.4	7
	Heavy precipitation	3.2.1.4	3		3
	Drought		3		3
5	Wind	3.2.1.5		3.3.1.5	7
	Storm		3		3
6	Air pollution, air quality and atmospheric deposition	3.2.1.6		3.3.1.6	
Land					
7	River discharge	3.2.2.1		3.3.2.1	8
	High flow		3		3
8	Land nutrient inputs	3.2.2.2		3.3.2.2	8
Terrestrial biosphere					
9	Land cover (forest, crops, grassland, peatland, mires)	3.2.3	6	3.3.3	
10	Carbon sequestration			3.3.3	
Cryosphere					
11	Snow	3.2.4.1		3.3.4.1	7
	Sea-effect snowfall		3		3
12	Glaciers	3.2.4.2		3.3.4.2	



13	Permafrost	3.2.4.3		3.3.4.3	
14	Sea ice	3.2.4.4		3.3.4.4	8
	Extreme mild winter		3		3
	Severe winter		3		3
	Ice ridging		3		3
15	Lake ice	3.2.4.5		3.3.4.5	
Ocean and marine sediments					
16	Water temperature	3.2.5.1		3.3.5.1	8
	Marine heat wave		3		3
17	Salinity and saltwater inflows	3.2.5.2	1	3.3.5.2	8
18	Stratification and overturning circulation	3.2.5.3	1	3.3.5.3	8
19	Sea level	3.2.5.4	4	3.3.5.4	8
	Sea level extreme		3		3
20	Waves	3.2.5.5	4	3.3.5.5	
	Extreme waves		3		3
21	Sedimentation and coastal erosion	3.2.5.6	4	3.3.5.6	
22	Oxygen and nutrients	3.1.4 3.2.5.7.1	2	3.3.5.7.1	8
23	Marine CO ₂ system	3.2.5.7.2	2	3.3.5.7.2	
Marine biosphere					
24	Pelagic habitats: Microbial communities	3.2.6.1.1	2, 9	3.3.6.1.1	9
25	Pelagic habitats: Phytoplankton and cyanobacteria	3.2.6.1.2	2, 3, 9	3.3.6.1.2	3, 9
26	Pelagic habitats: Zooplankton	3.2.6.1.3	9	3.3.6.1.3	9
27	Benthic habitats: Macroalgae and vascular plants	3.2.6.2.1	9	3.3.6.2.1	9
28	Benthic habitats: Zoobenthos	3.2.6.2.2	9	3.3.6.2.2	9
29	Non-indigenous species	3.2.6.3	9	3.3.6.3	9
30	Fish	3.2.6.4	9	3.3.6.4	9



31	Marine mammals	3.2.6.5	9	3.3.6.5	9
32	Waterbirds	3.2.6.6	9	3.3.6.6	9
33	Marine food web	3.2.6.7	9	3.3.6.7	9

4318



4319 **Table 2:** Comparison of the Baltic Sea with other intra-continental seas and large lakes.

Basin	Area	Mean depth	Mean salinity	Fresh water budget	Ice cover on average	Location Centre
<u>Unit</u>	10 ³ km ²	m	g kg ⁻¹			Lat Long

4320

Baltic Sea	393	54	7½	+	Half	60°N 20°E
Black Sea	436	1197	20	+	Northeast	43°N 35°E
Gulf of Ob	41	12	5	+	All	73°N 74°E
Chesapeake Bay	12	6	15	+	Shores	38°N 76°W
Hudson Bay	1232	128	30	+	All	58°N 85°W
Red Sea	438	491	40	–	None	22°N 38°E
Persian Gulf	239	25	40	–	None	27°N 52°E
Caspian Sea	374	211	12	0	North	43°N 50°E
Lake Superior	82	149	< 0.1	0	All	48°N 88°W

4321



Table 3: The main characteristics of the physical features of the European Seas (Leppäranta and Myrberg, 2009; Sündermann and Pohlmann, 2011; www.ospar.org; British Oceanographic Data Centre). Greater North Sea – being the neighbouring sea area to the Baltic Sea - is shown as a sub-region of the NE-Atlantic, but other European sub-regions are not listed (from Myrberg et al., 2019).

Basin	Area 10 ³ km ²	Mean depth m	Mean salinity g kg ⁻¹	Fresh water budget	Ice cover on average	Tides	Water residence time (years)
Baltic Sea	393	54	7.4	Pos.	37 % ¹⁾	Weak	40
Black Sea	422	1 200	18	Pos.	Northeast only	Weak	3 000
Greater North Sea	750	80	34–35	Pos.	No	Strong	Not applicable
Mediterranean Sea	2 970	1 500	38	Neg.	No	Weak/ Moderate	80-100
NE Atlantic shelf	13 500 ²⁾	1 500	34–35	Not applicable	No	Strong	Not applicable
1) Mean maximal ice cover between 2000-2017, see Fig. 3.2.4.4.1							
2) defined as the OSPAR convention area, incl. the Greater North Sea							

4326



Table 4: Linear surface air temperature trends (K decade^{-1}) for the period 1876–2020 over the northern ($>60^\circ\text{N}$) and southern ($<60^\circ\text{N}$) Baltic Sea basin (1878–2020 is selected for comparison with (Rutgersson et al., 2014), with an equally long time period). Bold: significance at $p < 0.05$. Data from the updated CRUTEM4v dataset (Jones et al., 2012).

	Annual	Winter	Spring	Summer	Autumn
North	0.10	0.11	0.14	0.08	0.08
South	0.10	0.13	0.10	0.09	0.09



Table 5: Average (2013-2017) riverine and coastal nutrient inputs (10^3 t N (P) yr^{-1}) to the major basins of the Baltic Sea (Source: Oleg P. Savchuk, Stockholm University). For abbreviated basin names see Figure 11.

	BB	BS	BP	GF	GR	DS	KT	Entire BS
TN river	48	40	251	85	73	32	46	575
TN coast	3.6	4.3	6.5	9.3	0.5	2.8	2.0	29
TP river	2.4	1.7	11.8	3.4	2.0	1.1	1.3	24.0
TP coast	0.1	0.2	0.5	0.4	0.1	0.2	0.1	1.6



Table 6: Mass balances for the Swedish glaciers Storglaciären, Rabots glaciär, Mårmaglaciär, and Riuokjietna. General references are given as footnotes in connection with balance years and long-term monitoring intervals, respectively. Selected specific references are given as footnotes in connection with the glacier names, and include also neighboring glaciers Kårsa glacier and Kebnepakteglaciär. (Source: Nina Kirchner, Stockholm University)

	Recent mass balance years. Gains and losses in mm w.e. (millimeter water equivalent). Note that the unit mm w.e. is interchangeable with the unit kg m ⁻²			Long-term mass balance, losses per year in mm w.e.	
	2015/ 2016 ³	2016/ 2017 ³	2017/ 2018 ³	1980-2010 ⁴	1985-2015 ⁵
Storglaciären ^{6,7,8}	-240	+470	-1600	-113	-153
Rabots glaciär ^{9,10}	-650	-170	-1680	-394	-465
Mårmaglaciär	-370	+260		-430	-460
Riuokjietna	-1060	+150		-592	-592

³ World Glacier Monitoring Service, 2020

⁴ Blunden and Arendt, 2015

⁵ Hartfield et al., 2018

⁶ Mercer, 2016

⁷ Holmlund and Holmlund, 2019

⁸ Kirchner et al., 2019

⁹ Brugger and Pankratz, 2015

¹⁰ Williams et al., 2016



Table 7: Air temperature (T_{2m}) changes ($^{\circ}\text{C}$) between 1976 - 2005 and 2069 - 2098 averaged over each season and annual mean over the Baltic Sea catchment area and over the Baltic Sea calculated from nine regionalized ESMs (Data source: Gröger et al., 2021b, compiled by Christian Dieterich, Swedish Meteorological and Hydrological Institute). In addition to the ensemble mean change, the 5th and 95th percentiles indicating the ensemble spread are listed (in brackets).

	Annual	Winter	Spring	Summer	Autumn
Total land					
RCP2.6	1.5 (1.2, 2.0)	2.1 (1.5, 3.3)	1.5 (1.2, 2.0)	1.3 (0.8, 2.1)	1.3 (0.9, 1.8)
RCP4.5	2.6 (1.6, 3.2)	3.2 (2.1, 4.2)	2.4 (1.5, 3.3)	2.1 (1.3, 3.1)	2.3 (1.4, 2.8)
RCP8.5	4.3 (3.5, 5.2)	5.0 (3.4, 6.3)	3.8 (3.1, 4.5)	3.7 (2.5, 5.0)	3.8 (2.6, 4.8)
Land north of 60°N					
RCP2.6	1.7 (1.4, 2.4)	2.5 (1.9, 3.1)	1.7 (1.2, 2.3)	1.4 (0.8, 2.3)	1.5 (1.1, 2.1)
RCP4.5	2.9 (2.0, 3.7)	4.0 (2.9, 5.0)	2.8 (1.8, 3.8)	2.3 (1.3, 3.4)	2.5 (1.7, 3.2)
RCP8.5	4.9 (3.9, 5.9)	6.0 (4.2, 7.5)	4.2 (3.5, 5.1)	3.9 (2.8, 5.1)	4.2 (2.9, 5.3)
Land south of 60°N					
RCP2.6	1.4 (1.0, 1.8)	1.7 (1.1, 3.4)	1.3 (0.9, 1.7)	1.3 (0.9, 1.9)	1.2 (0.7, 1.6)
RCP4.5	2.2 (1.3, 2.8)	2.6 (1.5, 4.0)	2.2 (1.3, 2.9)	2.0 (1.1, 3.0)	2.1 (1.2, 2.7)
RCP8.5	3.9 (3.2, 4.7)	4.2 (2.9, 5.7)	3.4 (2.9, 4.0)	3.5 (2.2, 4.9)	3.5 (2.3, 4.5)
Baltic Sea					
RCP2.6	1.4 (1.2, 1.9)	1.9 (1.3, 2.8)	1.5 (1.1, 1.9)	1.2 (0.6, 1.8)	1.2 (0.9, 1.7)
RCP4.5	2.4 (1.4, 2.9)	2.9 (1.8, 3.7)	2.5 (1.5, 3.1)	2.0 (1.2, 2.7)	2.1 (1.2, 2.7)
RCP8.5	3.9 (3.1, 4.8)	4.6 (3.2, 5.8)	3.9 (3.0, 4.9)	3.5 (2.4, 4.6)	3.6 (2.6, 4.6)



Table 8: Relative precipitation changes (%) between 1976 - 2005 and 2069 - 2098 averaged over each season and annual mean over the Baltic Sea catchment area and over the Baltic Sea calculated from nine regionalized ESMs (Data source: Gröger et al., 2021b, compiled by Christian Dieterich, Swedish Meteorological and Hydrological Institute). In addition to the ensemble mean change, the 5th and 95th percentiles indicating the ensemble spread are listed (in brackets).

	Annual	Winter	Spring	Summer	Autumn
Total land					
RCP2.6	5 (2, 14)	7 (1, 22)	8 (2, 12)	3 (-2, 13)	4 (-4, 12)
RCP4.5	9 (6, 14)	12 (4, 24)	13 (8, 17)	4 (1, 11)	6 (-5, 12)
RCP8.5	15 (11, 22)	22 (11, 38)	20 (7, 26)	5 (-4, 15)	13 (-1, 18)
Land north of 60°N					
RCP2.6	6 (2, 15)	7 (2, 23)	8 (0, 13)	5 (1, 17)	5 (-5, 14)
RCP4.5	11 (7, 18)	13 (6, 27)	15 (2, 21)	9 (4, 14)	8 (-3, 17)
RCP8.5	19 (12, 30)	22 (12, 41)	24 (7, 35)	13 (-1, 30)	17 (1, 26)
Land south of 60°N					
RCP2.6	5 (0, 13)	7 (-1, 22)	7 (3, 13)	2 (-5, 10)	3 (-7, 11)
RCP4.5	7 (4, 11)	12 (1, 22)	12 (6, 20)	1 (-5, 11)	4 (-8, 11)
RCP8.5	12 (8, 18)	21 (9, 35)	18 (7, 26)	-1 (-14, 9)	9 (-3, 17)
Baltic Sea					
RCP2.6	6 (0, 15)	5 (-3, 15)	4 (-1, 8)	8 (0, 22)	5 (-3, 13)
RCP4.5	8 (3, 13)	9 (-4, 20)	11 (1, 17)	6 (-1, 16)	6 (-3, 15)
RCP8.5	16 (8, 23)	18 (3, 31)	19 (-3, 32)	10 (-9, 22)	15 (4, 26)



Table 9: Sea surface temperature (SST) changes (°C) between 1976 - 2005 and 2069 - 2098 averaged over each season and annual mean over the Baltic Sea calculated from nine regionalized ESMs (Data source: Gröger et al., 2021b, compiled by Christian Dieterich, Swedish Meteorological and Hydrological Institute). In addition to the ensemble mean change, the 5th and 95th percentiles indicating the ensemble spread are listed (in brackets).

Baltic Sea	Annual	Winter	Spring	Summer	Autumn
RCP2.6	1.1 (0.8, 1.6)	1.0 (0.9, 1.4)	1.1 (0.9, 1.6)	1.2 (0.6, 1.7)	0.9 (0.7, 1.6)
RCP4.5	1.8 (1.1, 2.5)	1.7 (1.0, 2.3)	1.9 (1.2, 2.6)	2.0 (1.2, 2.6)	1.8 (1.1, 2.4)
RCP8.5	3.2 (2.5, 4.1)	3.0 (2.3, 3.8)	3.2 (2.5, 3.9)	3.4 (2.4, 4.5)	3.1 (2.4, 4.1)



Table 10: Summary of key messages about the impact of global warming on selected variables. The sign of a change (plus/minus) is listed together with the level of confidence denoted by the number of signs, i.e. one to three signs correspond to low, medium and high confidence levels. +/- means no detected or projected change due to climate change. Key messages of this assessment that are new compared to the previous assessment by BACC II Author Team (2015) are marked and a brief explanation is provided in the neighboring column. (NA = North Atlantic)

Number	Variable	Present climates		Future climate	
Atmosphere					
1	Large-scale circulation	+/-	Remote influence of the multi-decadal variability in the NA on the Baltic Sea	+/-	Impact of warming Arctic with declining sea ice might be relevant
2	Air temperature	+++	Accelerated warming	+++	Greater confidence due to increased ensemble size, coupled atmosphere-ocean models
	Warm spell	++		++	
	Cold spell	--		--	
3	Solar radiation	+	Comparison between various satellite products	+/-	GCM and RCM systematically differ
4	Precipitation	+		++	convection-resolving models became available
	Heavy precipitation	++		+++	
	Drought north (south) of 59°N	-(+)		-(+)	
5	Wind	+/-		+	Small systematic increase in winter in the northern Baltic where the sea ice will melt
	Number of deep cyclones	+		+/-	
6	Air pollution, air quality and atmospheric deposition	+/-		+/-	
Land					
7	River discharge	+/-	Dataset of observed time series for the past century from Sweden merged with high-resolution dynamic model	+	Changing seasonality (decrease of river discharge in spring, increase in winter)
	High flow ¹¹ in the north (south)	+/- (+/-)		- (+)	

¹¹ Based upon annual maximum river discharges of daily data for Sweden with 10- and 100-year repeat periods (Roudier et al., 2016) and for Finland with 100-year repeat period (Veijalainen et al., 2010)



			projections of the upcoming century		may affect the occurrence of floods ^{12,13}
8	Land nutrient inputs	+/-		+	
Terrestrial biosphere					
9	Growing season in the Baltic Sea region	++	Study based on satellite data available	+/-	No new study
10	Carbon sequestration in northern terrestrial ecosystems	+/-		+, later -	First increasing sinks. Weak sources of carbon after 2060-2070s due to increased soil respiration and biomass burning
Cryosphere					
11	Snow	---		---	
	Sea-effect snowfall	+/-		+/-	
12	Ice mass of glaciers	---	Since 2006 inventories of all Scandinavian glaciers have become available	---	High-resolution projections of Scandinavian glaciers available
13	Permafrost	--	High-resolution modeling	---	
14	Sea ice cover	---		---	
	Extreme mild winter	+++		+++	
	Extreme severe winter	---		---	
	Ice ridging	+/-		-	
15	Lake ice	---	Systematic assessment available	---	Projections for global lake ice available
Ocean and marine sediments					
16	Water temperature	+++	Accelerated warming	+++	
	Marine heat wave	+		+++	Increasing number of record-breaking summer mean SST events and number of heat waves
17	Salinity and saltwater inflows	+/-	Homogenous data of saltwater inflows, north-south salinity	+/-	Uncertainty sources of salinity due to wind, river discharge and global sea

¹² Roudier et al., 2016

¹³ Veijalainen et al., 2010



			gradient has increased		level rise changes were assessed
18	Stratification and overturning circulation	+/-	Systematic study of monitoring data since the 1980s	+/-	Intensified seasonal thermoclines during summer but no change of the halocline and overturning circulation
19	Absolute sea level Storm surge relative to the mean sea level	+++ +/-	Paleoclimate study on sea level extremes did not show systematic changes in changing climate, dissensus in the literature ^{14,15}	+++ +/-	Dissensus in the literature ^{16,17}
20	Waves Extreme waves	+/- +/-		+ +	Small increase in winter in the northern Baltic Sea
21	Sedimentation and coastal erosion	+/-		+/-	First modeling studies available
22	Hypoxic area	+	Warming contributed to the historical spread of hypoxia in the deep water and in the coastal zone, sediment cores suggest that changing climate caused hypoxia during the Medieval Climate Anomaly instead of agriculture	++	Oxygen decline in the coastal zone due to warming
23	CO ₂ uptake pH southern (northern) Baltic Sea	++ --(+/-)	New observations and modeling, positive alkalinity trends identified	+/- +/-	

¹⁴ Ribeiro et al., 2014

¹⁵ Marcos and Woodworth, 2017

¹⁶ Vousdoukas et al., 2016

¹⁷ Vousdoukas et al., 2017



Marine biosphere					
24	Microbial communities	+	In the northern Baltic Sea increased riverine dissolved organic matter suppressed phytoplankton biomass production and shifts the carbon flow towards microbial heterotrophy	+/-	Increase of dissolved organic matter and temperature will enhance and decrease the abundance of bacteria, respectively
25	growing season of phytoplankton (cyanobacteria)	++	new indicator for the environmental status developed	+/-	Warming causes prolonged and intensified cyanobacteria blooms but the nutrient control is dominating
	cyanobacteria biomass	+/-		+	
	ratio between diatom and dinoflagellate biomasses since 1901	-		+/-	
26	Zooplankton	+/-		+/-	Increasing microzooplankton biomass
27	Macroalgae and vascular plants	+/-	Systematic studies on benthic ecosystems	+/-	
28	Zoobenthos	+/-	Systematic studies on benthic ecosystems, spreading of non-indigenous such as polychaete <i>Marenzelleria</i> spp.	+/-	Weaker benthic-pelagic coupling and decreasing benthic biomass in a warmer and less eutrophic Baltic
29	Non-indigenous species	+		++	
30	Fish	+/-	Food web modeling including fisheries	+/-	Multi-driver (climate change, eutrophication,



					fisheries) food web projections were performed
31	Populations of marine mammals	+/-		-	
32	Waterbird migration	+++	Northward shift of the wintering range of waterbirds	++	Controlled by food availability
33	Marine food web	+/-		+/-	

4363



4364 References

- 4365 Aalto, J., Pirinen, P., and Jylhä, K.: New gridded daily climatology of Finland: Permutation-based uncertainty
 4366 estimates and temporal trends in climate, *J. Geophys. Res-Atmos.*, 121, 3807-3823,
 4367 <https://doi.org/10.1002/2015JD024651>, 2016.
- 4368 Aarvak, T., Jostein Øien, I., Krasnov, Y. V., Gavrilov, M. V., and Shavykin, A. A.: The European wintering
 4369 population of Steller's Eider *Polysticta stelleri* reassessed, *Bird Conserv. Int.*, 23, 337-343,
 4370 <https://doi.org/10.1017/S0959270912000251>, 2013.
- 4371 Aberle, N., Malzahn, A. M., Lewandowska, A. M., and Sommer, U.: Some like it hot: the protozooplankton-
 4372 copepod link in a warming ocean, *Mar. Ecol. Prog. Ser.*, 519, 103-113, <https://doi.org/10.3354/meps11081> 2015.
- 4373 Adam, J. C., Haddeland, I., Su, F., and Lettenmaier, D. P.: Simulation of reservoir influences on annual and
 4374 seasonal streamflow changes for the Lena, Yenisei, and Ob' rivers, *J. Geophys. Res-Atmos.*, 112,
 4375 <https://doi.org/10.1029/2007JD008525>, 2007.
- 4376 Adam, J. C., and Lettenmaier, D. P.: Application of New Precipitation and Reconstructed Streamflow Products
 4377 to Streamflow Trend Attribution in Northern Eurasia, *J. Climate*, 21, 1807-1828,
 4378 <https://doi.org/10.1175/2007jcli1535.1>, 2008.
- 4379 Ahlgren, J., Grimvall, A., Omstedt, A., Rolff, C., and Wikner, J.: Temperature, DOC level and basin interactions
 4380 explain the declining oxygen concentrations in the Bothnian Sea, *J. Mar. Sys.*, 170, 22-30,
 4381 <https://doi.org/10.1016/j.jmarsys.2016.12.010>, 2017.
- 4382 Åkerman, H. J., and Johansson, M.: Thawing permafrost and thicker active layers in sub-arctic Sweden,
 4383 *Permafrost Periglac.*, 19, 279-292, <https://doi.org/10.1002/ppp.626>, 2008.
- 4384 Akhtar, N., Geyer, B., Rockel, B., Sommer, P. S., and Schrum, C.: Accelerating deployment of offshore wind
 4385 energy alter wind climate and reduce future power generation potentials, *Sci. Rep.-UK*, 11, 11826,
 4386 <https://doi.org/10.1038/s41598-021-91283-3>, 2021.
- 4387 Al-Janabi, B., Kruse, I., Graiff, A., Karsten, U., and Wahl, M.: Genotypic variation influences tolerance to
 4388 warming and acidification of early life-stage *Fucus vesiculosus* L. (Phaeophyceae) in a seasonally fluctuating
 4389 environment, *Mar. Biol.*, 163, 14, <https://doi.org/10.1007/s00227-015-2804-8>, 2016.
- 4390 Alkama, R., and Cescatti, A.: Biophysical climate impacts of recent changes in global forest cover, *Science*, 351,
 4391 600-604, <https://doi.org/10.1126/science.aac8083>, 2016.
- 4392 Allen, M. R., Dube, O. P., Solecki, W., Aragón-Durand, F., Cramer, W., Humphreys, S., Kainuma, M., Kala, J.,
 4393 Mahowald, N., Mulugetta, Y., Perez, R., Wairiu, M., and Zickfeld, K.: Framing and Context, in: *Global*
 4394 *Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial*
 4395 *levels and related global greenhouse gas emission pathways, in the context of strengthening the global response*
 4396 *to the threat of climate change, sustainable development, and efforts to eradicate poverty*, edited by: Masson-
 4397 Delmotte, V., Zhai, P., Pörtner, H.-O., Roberts, D., Skea, J., Shukla, P. R., Pirani, A., Moufouma-Okia, W.,
 4398 Péan, C., Pidcock, R., Connors, S., Matthews, J. B. R., Chen, Y., Zhou, X., Gomis, M. I., Lonnoy, E., Maycock,
 4399 T., Tignor, M., and Waterfield, T., <https://www.ipcc.ch/sr15/chapter/chapter-1/>, 2018.
- 4400 Allen, R. J., Norris, J. R., and Wild, M.: Evaluation of multidecadal variability in CMIP5 surface solar radiation
 4401 and inferred underestimation of aerosol direct effects over Europe, China, Japan, and India, *J. Geophys. Res-*
 4402 *Atmos.*, 118, 6311-6336, <https://doi.org/10.1002/jgrd.50426>, 2013.
- 4403 Allen, S. G., Ainley, D. G., Page, G. W., and Ribic, C. A.: The effect on distribution on harbor seal haul out
 4404 patterns at Bolinas Lagoon, California, in: *Fish. Bull.*, edited by: Richards, W. J., Collette, B. B., Houde, E. D.,
 4405 Ingham, M. C., Lasker, R., Malins, D. C., Pella, J. J., Quast, J. C., Sindermann, C. J., and Fukuyama, M. S.,
 4406 National Marine Fisheries Service, NOAA, Seattle, WA, USA, 493-500, 1984.
- 4407 Almroth-Rosell, E., Edman, M., Eilola, K., Meier, H. E. M., and Sahlberg, J.: Modelling nutrient retention in the
 4408 coastal zone of an eutrophic sea, *Biogeosciences*, 13, 5753-5769, <https://doi.org/10.5194/bg-13-5753-2016>,
 4409 2016.
- 4410 Andersen, J. H., Carstensen, J., Conley, D. J., Dromph, K., Fleming-Lehtinen, V., Gustafsson, B. G., Josefson,
 4411 A. B., Norkko, A., Villnäs, A., and Murray, C.: Long-term temporal and spatial trends in eutrophication status of
 4412 the Baltic Sea, *Biol. Rev.*, 92, 135-149, <https://doi.org/10.1111/brv.12221>, 2017.



- 4413 Andersson, A., Meier, H. E. M., Ripszam, M., Rowe, O., Wikner, J., Haglund, P., Eilola, K., Legrand, C.,
 4414 Figueroa, D., Paczkowska, J., Lindehoff, E., Tysklind, M., and Elmgren, R.: Projected future climate change and
 4415 Baltic Sea ecosystem management, *Ambio* 44, 345-356, <https://doi.org/10.1007/s13280-015-0654-8>, 2015.
- 4416 Andersson, H. C.: Influence of long-term regional and large-scale atmospheric circulation on the Baltic sea level,
 4417 *Tellus A*, 54, 76-88, <https://doi.org/10.3402/tellusa.v54i1.12125>, 2002.
- 4418 Arheimer, B., Dahné, J., and Donnelly, C.: Climate change impact on riverine nutrient load and land-based
 4419 remedial measures of the Baltic Sea Action Plan, *Ambio*, 41, 600-612, [https://doi.org/10.1007/s13280-012-0323-](https://doi.org/10.1007/s13280-012-0323-0)
 4420 0, 2012.
- 4421 Arheimer, B., and Lindström, G.: Climate impact on floods: changes in high flows in Sweden in the past and the
 4422 future (1911-2100), *Hydrol. Earth Syst. Sci.*, 19, 771-784, <https://doi.org/10.5194/hess-19-771-2015>, 2015.
- 4423 Arheimer, B., Donnelly, C., and Lindström, G.: Regulation of snow-fed rivers affects flow regimes more than
 4424 climate change, *Nat. Commun.*, 8, 62, <https://doi.org/10.1038/s41467-017-00092-8>, 2017.
- 4425 Armitage, J. M., Quinn, C. L., and Wania, F.: Global climate change and contaminants—an overview of
 4426 opportunities and priorities for modelling the potential implications for long-term human exposure to organic
 4427 compounds in the Arctic, *J. Environ. Monitor.*, 13, 1532-1546, <https://doi.org/10.1039/C1EM10131E>, 2011.
- 4428 Arneborg, L.: Comment on “Influence of sea level rise on the dynamics of salt inflows in the Baltic Sea” by R.
 4429 Hordoir, L. Axell, U. Löptien, H. Dietze, and I. Kuznetsov, *J. Geophys. Res-Oceans*, 121, 2035-2040,
 4430 <https://doi.org/10.1002/2015JC011451>, 2016.
- 4431 Asmala, E., Carstensen, J., Conley, D. J., Slomp, C. P., Stadmark, J., and Voss, M.: Efficiency of the coastal
 4432 filter: Nitrogen and phosphorus removal in the Baltic Sea, *Limnol. Oceanogr.*, 62,
 4433 <https://doi.org/10.1002/lno.10644>, 2017.
- 4434 Athanasiadis, P. J., Yeager, S., Kwon, Y.-O., Bellucci, A., Smith, D. W., and Tibaldi, S.: Decadal predictability
 4435 of North Atlantic blocking and the NAO, *npj Clim. Atmos. Sci.*, 3, 20, [https://doi.org/10.1038/s41612-020-0120-](https://doi.org/10.1038/s41612-020-0120-0)
 4436 0, 2020.
- 4437 Averkiev, A. S., and Klevanny, K. A.: A case study of the impact of cyclonic trajectories on sea-level extremes
 4438 in the Gulf of Finland, *Cont. Shelf Res.*, 30, 707-714, <https://doi.org/10.1016/j.csr.2009.10.010>, 2010.
- 4439 Axell, L., Liu, Y., Jandt, S., Lorkowski, I., Lindenthal, A., Verjovkina, S., and Schwichtenberg, F.: Baltic Sea
 4440 Production Centre BALTICSEA_REANALYSIS_BIO_003_012 COPERNICUS Marine Environment
 4441 Monitoring Service, Quality Information Document, 51pp, 2019.
- 4442 BACC Author Team: Assessment of climate change for the Baltic Sea basin, *Regional Climate Studies*, Springer
 4443 Science & Business Media, Berlin, Heidelberg, 473 pp., 2008.
- 4444 BACC II Author Team: Second Assessment of Climate Change for the Baltic Sea Basin, *Regional Climate*
 4445 *Studies*, Springer International Publishing, Cham, 2015.
- 4446 Bailey, S. A., Brown, L., Campbell, M. L., Canning-Clode, J., Carlton, J. T., Castro, N., Chainho, P., Chan, F.
 4447 T., Creed, J. C., Curd, A., Darling, J., Fofonoff, P., Galil, B. S., Hewitt, C. L., Inglis, G. J., Keith, I., Mandrak, N.
 4448 E., Marchini, A., McKenzie, C. H., Occhipinti-Ambrogi, A., Ojaveer, H., Pires-Teixeira, L. M., Robinson, T. B.,
 4449 Ruiz, G. M., Seaward, K., Schwindt, E., Son, M. O., Theriault, T. W., and Zhan, A.: Trends in the detection of
 4450 aquatic non-indigenous species across global marine, estuarine and freshwater ecosystems: A 50-year
 4451 perspective, *Divers. Distrib.*, 26, 1780-1797, <https://doi.org/10.1111/ddi.13167>, 2020.
- 4452 BALTEX: Minutes of 17th Meeting of the BALTEX Science Steering Group held at Institute of Meteorology
 4453 and Water Management (IMGW), Poznan, Poland, 24 - 26 November 2004, International BALTEX Secretariat
 4454 Publication No. 32, 100pp, 2005.
- 4455 Bamber, J. L., Oppenheimer, M., Kopp, R. E., Aspinall, W. P., and Cooke, R. M.: Ice sheet contributions to
 4456 future sea-level rise from structured expert judgment, *P. Natl. Acad. Sci. USA*, 116, 11195,
 4457 <https://doi.org/10.1073/pnas.1817205116>, 2019.
- 4458 Barbosa, S. M., and Donner, R. V.: Long-term changes in the seasonality of Baltic sea level, *Tellus A*, 68,
 4459 30540, <https://doi.org/10.3402/tellusa.v68.30540>, 2016.
- 4460 Barnes, E. A.: Revisiting the evidence linking Arctic amplification to extreme weather in midlatitudes, *Geophys.*
 4461 *Res. Lett.*, 40, 4734-4739, <https://doi.org/10.1002/grl.50880>, 2013.



- 4462 Barregard, L., Molnàr, P., Jonson, J. E., and Stockfelt, L.: Impact on Population Health of Baltic Shipping
 4463 Emissions, *Int. J. Environ. Res. Public Health*, 16, 1954, <https://doi.org/10.3390/ijerph16111954>, 2019.
- 4464 Bartók, B., Wild, M., Folini, D., Lüthi, D., Kotlarski, S., Schär, C., Vautard, R., Jerez, S., and Imecs, Z.:
 4465 Projected changes in surface solar radiation in CMIP5 global climate models and in EURO-CORDEX regional
 4466 climate models for Europe, *Clim. Dyn.*, 49, 2665-2683, <https://doi.org/10.1007/s00382-016-3471-2>, 2017.
- 4467 Bartolino, V., Margonski, P., Lindegren, M., Linderholm, H. W., Cardinale, M., Rayner, D., Wennhage, H., and
 4468 Casini, M.: Forecasting fish stock dynamics under climate change: Baltic herring (*Clupea harengus*) as a case
 4469 study, *Fish. Oceanogr.*, 23, 258-269, <https://doi.org/10.1111/fog.12060>, 2014.
- 4470 Bartosova, A., Capell, R., Olesen, J. E., Jabloun, M., Refsgaard, J. C., Donnelly, C., Hyytiäinen, K., Pihlainen,
 4471 S., Zandersen, M., and Arheimer, B.: Future socioeconomic conditions may have a larger impact than climate
 4472 change on nutrient loads to the Baltic Sea, *Ambio*, 48, 1325-1336, <https://doi.org/10.1007/s13280-019-01243-5>,
 4473 2019.
- 4474 Bates, A. E., Pecl, G. T., Frusher, S., Hobday, A. J., Wernberg, T., Smale, D. A., Sunday, J. M., Hill, N. A.,
 4475 Dulvy, N. K., Colwell, R. K., Holbrook, N. J., Fulton, E. A., Slawinski, D., Feng, M., Edgar, G. J., Radford, B.
 4476 T., Thompson, P. A., and Watson, R. A.: Defining and observing stages of climate-mediated range shifts in
 4477 marine systems, *Glob. Environ. Change*, 26, 27-38, <https://doi.org/10.1016/j.gloenvcha.2014.03.009>, 2014.
- 4478 Bauer, B., Meier, H. E. M., Casini, M., Hoff, A., Margoński, P., Orio, A., Saraiva, S., Steenbeek, J., and
 4479 Tomczak, M. T.: Reducing eutrophication increases spatial extent of communities supporting commercial
 4480 fisheries: a model case study, *ICES J. Mar. Sci.*, 75, 1306-1317, <https://doi.org/10.1093/icesjms/fsy023>, 2018.
- 4481 Bauer, B., Gustafsson, B. G., Hyytiäinen, K., Meier, H. E. M., Müller-Karulis, B., Saraiva, S., and Tomczak, M.
 4482 T.: Food web and fisheries in the future Baltic Sea, *Ambio*, <https://doi.org/10.1007/s13280-019-01229-3>, 2019.
- 4483 Belkin, I. M.: Rapid warming of large marine ecosystems, *Prog. Oceanogr.*, 81, 207-213,
 4484 <https://doi.org/10.1016/j.pcean.2009.04.011>, 2009.
- 4485 Bengtsson, L., Hodges, K. I., and Keenlyside, N.: Will Extratropical Storms Intensify in a Warmer Climate?, *J.*
 4486 *Climate*, 22, 2276-2301, <https://doi.org/10.1175/2008jcli2678.1>, 2009.
- 4487 Bergen, B., Endres, S., Engel, A., Zark, M., Dittmar, T., Sommer, U., and Jürgens, K.: Acidification and
 4488 warming affect prominent bacteria in two seasonal phytoplankton bloom mesocosms, *Environ. Microbiol.*, 18,
 4489 4579-4595, <https://doi.org/10.1111/1462-2920.13549>, 2016.
- 4490 Bergström, L., Heikinheimo, O., Svirgdsen, R., Kruze, E., Ložys, L., Lappalainen, A., Saks, L., Minde, A.,
 4491 Dainys, J., Jakubavičiūtė, E., Ådjers, K., and Olsson, J.: Long term changes in the status of coastal fish in the
 4492 Baltic Sea, *Estuar. Coast. Shelf Sci.*, 169, 74-84, <https://doi.org/10.1016/j.ecss.2015.12.013>, 2016.
- 4493 Bergström, S., and Carlsson, B.: River runoff to the Baltic Sea - 1950-1990, *Ambio*, 23, 280-287, 1994.
- 4494 Bermúdez, R., Winder, M., Stühr, A., Almén, A. K., Engström-Öst, J., and Riebesell, U.: Effect of ocean
 4495 acidification on the structure and fatty acid composition of a natural plankton community in the Baltic Sea,
 4496 *Biogeosciences*, 13, 6625-6635, <https://doi.org/10.5194/bg-13-6625-2016>, 2016.
- 4497 Berner, C., Bertos-Fortis, M., Pinhassi, J., and Legrand, C.: Response of Microbial Communities to Changing
 4498 Climate Conditions During Summer Cyanobacterial Blooms in the Baltic Sea, *Front. Microbiol.*, 9,
 4499 <https://doi.org/10.3389/fmicb.2018.01562>, 2018.
- 4500 Bierstedt, S. E., Hünicke, B., and Zorita, E.: Variability of wind direction statistics of mean and extreme wind
 4501 events over the Baltic Sea region, *Tellus A*, 67, 29073, <https://doi.org/10.3402/tellusa.v67.29073>, 2015.
- 4502 Bingham, R. J., and Hughes, C. W.: Local diagnostics to estimate density-induced sea level variations over
 4503 topography and along coastlines, *J. Geophys. Res.-Oceans*, 117, <https://doi.org/10.1029/2011jc007276>, 2012.
- 4504 Biskaborn, B. K., Smith, S. L., Noetzli, J., Matthes, H., Vieira, G., Streletskiy, D. A., Schoeneich, P.,
 4505 Romanovsky, V. E., Lewkowicz, A. G., Abramov, A., Allard, M., Boike, J., Cable, W. L., Christiansen, H. H.,
 4506 Delaloye, R., Diekmann, B., Drozdov, D., Etzelmüller, B., Grosse, G., Guglielmin, M., Ingeman-Nielsen, T.,
 4507 Isaksen, K., Ishikawa, M., Johansson, M., Johansson, H., Joo, A., Kaverin, D., Kholodov, A., Konstantinov, P.,
 4508 Kröger, T., Lambiel, C., Lanckman, J.-P., Luo, D., Malkova, G., Meiklejohn, I., Moskalenko, N., Oliva, M.,
 4509 Phillips, M., Ramos, M., Sannel, A. B. K., Sergeev, D., Seybold, C., Skryabin, P., Vasiliev, A., Wu, Q.,
 4510 Yoshikawa, K., Zheleznyak, M., and Lantuit, H.: Permafrost is warming at a global scale, *Nat. Commun.*, 10,
 4511 264, <https://doi.org/10.1038/s41467-018-08240-4>, 2019.



- 4512 Björkqvist, J.-V., Lukas, I., Alari, V., van Vledder, G. P., Hulst, S., Pettersson, H., Behrens, A., and Männik, A.:
 4513 Comparing a 41-year model hindcast with decades of wave measurements from the Baltic Sea, *Ocean Eng.*, 152,
 4514 57-71, <https://doi.org/10.1016/j.oceaneng.2018.01.048>, 2018.
- 4515 Björkqvist, J. V., Rikka, S., Alari, V., Männik, A., Tuomi, L., and Pettersson, H.: Wave height return periods
 4516 from combined measurement–model data: A Baltic Sea case study, *Nat. Hazard. Earth Syst. Sci. Discuss.*, 2020,
 4517 1-25, <https://doi.org/10.5194/nhess-2020-190>, 2020.
- 4518 Blackport, R., and Screen, J. A.: Insignificant effect of Arctic amplification on the amplitude of midlatitude
 4519 atmospheric waves, *Sci. Adv.*, 6, eaay2880, <https://doi.org/10.1126/sciadv.aay2880>, 2020.
- 4520 Blenckner, T., Österblom, H., Larsson, P., Andersson, A., and Elmgren, R.: Baltic Sea ecosystem-based
 4521 management under climate change: Synthesis and future challenges, *Ambio* 44, 507-515,
 4522 <https://doi.org/10.1007/s13280-015-0661-9>, 2015.
- 4523 Blindow, I., Dahlke, S., Dewart, A., Flügge, S., Hendreschke, M., Kerkow, A., and Meyer, J.: Long-term and
 4524 interannual changes of submerged macrophytes and their associated diaspore reservoir in a shallow southern
 4525 Baltic Sea bay: influence of eutrophication and climate, *Hydrobiologia*, 778, 121-136,
 4526 <https://doi.org/10.1007/s10750-016-2655-4>, 2016.
- 4527 Blöschl, G., Hall, J., Parajka, J., Perdigão, R. A. P., Merz, B., Arheimer, B., Aronica, G. T., Bilibashi, A.,
 4528 Bonacci, O., Borga, M., Čanjevac, I., Castellarin, A., Chirico, G. B., Claps, P., Fiala, K., Frolova, N.,
 4529 Gorbachova, L., Gül, A., Hannaford, J., Harrigan, S., Kireeva, M., Kiss, A., Kjeldsen, T. R., Kohnová, S.,
 4530 Koskela, J. J., Ledvinka, O., Macdonald, N., Mavrova-Guirguinova, M., Mediero, L., Merz, R., Molnar, P.,
 4531 Montanari, A., Murphy, C., Osuch, M., Ovcharuk, V., Radevski, I., Rogger, M., Salinas, J. L., Sauquet, E., Šraj,
 4532 M., Szolgay, J., Viglione, A., Volpi, E., Wilson, D., Zaimi, K., and Živković, N.: Changing climate shifts timing
 4533 of European floods, *Science*, 357, 588, <https://doi.org/10.1126/science.aan2506>, 2017.
- 4534 Blunden, J., and Arndt, D. S.: State of the Climate in 2014, *Bull. Amer. Meteor. Soc.*, 96, ES1-ES32,
 4535 <https://doi.org/10.1175/2015BAMSStateoftheClimate.1>, 2015.
- 4536 Bobsien, I. C., Hukriede, W., Schlamkow, C., Friedland, R., Dreier, N., Schubert, P. R., Karez, R., and Reusch,
 4537 T. B. H.: Modeling eelgrass spatial response to nutrient abatement measures in a changing climate, *Ambio*, 50,
 4538 400-412, <https://doi.org/10.1007/s13280-020-01364-2>, 2021.
- 4539 Boé, J., Terray, L., Moine, M.-P., Valcke, S., Bellucci, A., Drijfhout, S., Haarsma, R., Lohmann, K., Putrasahan,
 4540 D. A., Roberts, C., Roberts, M., Scoccimarro, E., Seddon, J., Senan, R., and Wyser, K.: Past long-term summer
 4541 warming over western Europe in new generation climate models: role of large-scale atmospheric circulation,
 4542 *Environ. Res. Lett.*, 15, 084038, <https://doi.org/10.1088/1748-9326/ab8a89>, 2020.
- 4543 Boland, E. J. D., Bracegirdle, T. J., and Shuckburgh, E. F.: Assessment of sea ice-atmosphere links in CMIP5
 4544 models, *Clim. Dyn.*, 49, 683-702, <https://doi.org/10.1007/s00382-016-3367-1>, 2017.
- 4545 Bonaduce, A., Staneva, J., Behrens, A., Bidlot, J.-R., and Wilcke, R. A. I.: Wave Climate Change in the North
 4546 Sea and Baltic Sea, *J. Mar. Sci. Eng.*, 7, 166, <https://doi.org/10.3390/jmse7060166>, 2019.
- 4547 Bonan, G. B.: Forests, Climate, and Public Policy: A 500-Year Interdisciplinary Odyssey, *Annu. Rev. Ecol.*
 4548 *Evol. S.*, 47, 97-121, <https://doi.org/10.1146/annurev-ecolsys-121415-032359>, 2016.
- 4549 Bonsdorff, E.: Eutrophication: Early warning signals, ecosystem-level and societal responses, and ways forward,
 4550 *Ambio*, <https://doi.org/10.1007/s13280-020-01432-7>, 2021.
- 4551 Börgel, F., Frauen, C., Neumann, T., Schimanke, S., and Meier, H. E. M.: Impact of the Atlantic Multidecadal
 4552 Oscillation on Baltic Sea Variability, *Geophys. Res. Lett.*, 45, 9880-9888,
 4553 <https://doi.org/10.1029/2018GL078943>, 2018.
- 4554 Börgel, F., Frauen, C., Neumann, T., and Meier, H. E. M.: The Atlantic Multidecadal Oscillation controls the
 4555 impact of the North Atlantic Oscillation on North European climate, *Environ. Res. Lett.*,
 4556 <https://doi.org/10.1088/1748-9326/aba925>, 2020.
- 4557 Bouwman, L., Goldewijk, K. K., Van Der Hoek, K. W., Beusen, A. H. W., Van Vuuren, D. P., Willems, J.,
 4558 Rufino, M. C., and Stehfest, E.: Exploring global changes in nitrogen and phosphorus cycles in agriculture
 4559 induced by livestock production over the 1900–2050 period, *P. Natl. Acad. Sci. USA*, 110, 20882,
 4560 <https://doi.org/10.1073/pnas.1012878108>, 2013.



- 4561 Brandt, J., Silver, J. D., Christensen, J. H., Andersen, M. S., Bønløkke, J. H., Sigsgaard, T., Geels, C., Gross, A.,
 4562 Hansen, A. B., Hansen, K. M., Hedegaard, G. B., Kaas, E., and Frohn, L. M.: Assessment of past, present and
 4563 future health-cost externalities of air pollution in Europe and the contribution from international ship traffic
 4564 using the EVA model system, *Atmos. Chem. Phys.*, 13, 7747-7764, <https://doi.org/10.5194/acp-13-7747-2013>,
 4565 2013.
- 4566 Broman, B., Hammarklint, T., Rannat, K., Soomere, T., and Valdmann, A.: Trends and extremes of wave fields
 4567 in the north-eastern part of the Baltic Proper, *Oceanologia*, 48, 2006.
- 4568 Brown, J., Ferrians Jr, O. J., Heginbottom, J. A., and Melnikov, E. S.: Circum-Arctic map of permafrost and
 4569 ground-ice conditions, U.S. Geological Survey, U.S.A., Report 45, <https://doi.org/10.3133/cp45>, 1997.
- 4570 Brugger, K. A., and Pankratz, L.: Changes in the geometry and volume of rabots glaciär, sweden, 2003–2011:
 4571 recent accelerated volume loss linked to more negative summer balances, *Geogr. Ann. A*, 97, 265-278,
 4572 <https://doi.org/10.1111/geoa.12062>, 2015.
- 4573 Brunner, L., Pendergrass, A. G., Lehner, F., Merrifield, A. L., Lorenz, R., and Knutti, R.: Reduced global
 4574 warming from CMIP6 projections when weighting models by performance and independence, *Earth Syst.*
 4575 *Dynam. Discuss.*, 2020, 1-23, <https://doi.org/10.5194/esd-2020-23>, 2020.
- 4576 Brutemark, A., Engström-Öst, J., Vehmaa, A., and Gorokhova, E.: Growth, toxicity and oxidative stress of a
 4577 cultured cyanobacterium (*Dolichospermum* sp.) under different CO₂/pH and temperature conditions, *Phycol.*
 4578 *Res.*, 63, 56-63, <https://doi.org/10.1111/pre.12075>, 2015.
- 4579 Bülow, K., Dietrich, C., Elizalde, A., Gröger, M., Heinrich, H., Hüttel-Kabos, S., Klein, B., Mayer, B., Meier, H.
 4580 E. M., and Mikolajewicz, U.: Comparison of three regional coupled ocean atmosphere models for the North Sea
 4581 under today's and future climate conditions (KLIWAS Schriftenreihe; KLIWAS-27/2014), Bundesanstalt für
 4582 Gewässerkunde, 2014.
- 4583 Burchard, H., Bolding, K., Feistel, R., Gräwe, U., Klingbeil, K., MacCready, P., Mohrholz, V., Umlauf, L., and
 4584 van der Lee, E. M.: The Knudsen theorem and the Total Exchange Flow analysis framework applied to the Baltic
 4585 Sea, *Prog. Oceanogr.*, 165, 268-286, <https://doi.org/10.1016/j.pocean.2018.04.004>, 2018.
- 4586 Caballero-Alfonso, A. M., Carstensen, J., and Conley, D. J.: Biogeochemical and environmental drivers of
 4587 coastal hypoxia, *J. Mar. Syst.*, 141, 190-199, <https://doi.org/10.1016/j.jmarsys.2014.04.008>, 2015.
- 4588 Cardell, M. F., Amengual, A., Romero, R., and Ramis, C.: Future extremes of temperature and precipitation in
 4589 Europe derived from a combination of dynamical and statistical approaches, *Int. J. Climatol.*, 40, 4800-4827,
 4590 <https://doi.org/10.1002/joc.6490>, 2020.
- 4591 Carstensen, J., Andersen, J. H., Gustafsson, B. G., and Conley, D. J.: Deoxygenation of the Baltic Sea during the
 4592 last century, *P. Natl. Acad. Sci. USA*, 111, 5628-5633, <https://doi.org/10.1073/pnas.1323156111>, 2014a.
- 4593 Carstensen, J., Conley, D. J., Bonsdorff, E., Gustafsson, B. G., Hietanen, S., Janas, U., Jilbert, T., Maximov, A.,
 4594 Norkko, A., Norkko, J., and others: Hypoxia in the Baltic Sea: Biogeochemical cycles, benthic fauna, and
 4595 management, *Ambio*, 43, 26-36, <https://doi.org/10.1007/s13280-013-0474-7>, 2014b.
- 4596 Carstensen, J., Chierici, M., Gustafsson, B. G., and Gustafsson, E.: Long-Term and Seasonal Trends in Estuarine
 4597 and Coastal Carbonate Systems, *Global Biogeochem. Cy.*, 32, 497-513, <https://doi.org/10.1002/2017gb005781>,
 4598 2018.
- 4599 Carstensen, J., and Conley, D. J.: Baltic Sea Hypoxia Takes Many Shapes and Sizes, *Limnol. Oceanogr. Bull.*,
 4600 28, 125-129, <https://doi.org/10.1002/lob.10350>, 2019.
- 4601 Carstensen, J., and Duarte, C. M.: Drivers of pH Variability in Coastal Ecosystems, *Environ. Sci. Technol.*, 53,
 4602 4020-4029, <https://doi.org/10.1021/acs.est.8b03655>, 2019.
- 4603 Carstensen, J., Conley, D. J., Almroth-Rosell, E., Asmala, E., Bonsdorff, E., Fleming-Lehtinen, V., Gustafsson,
 4604 B. G., Gustafsson, C., Heiskanen, A.-S., Janas, U., Norkko, A., Slomp, C., Villnäs, A., Voss, M., and Zilius, M.:
 4605 Factors regulating the coastal nutrient filter in the Baltic Sea, *Ambio*, 49, 1194-1210,
 4606 <https://doi.org/10.1007/s13280-019-01282-y>, 2020.
- 4607 Casanueva, A., Rodríguez-Puebla, C., Frías, M. D., and González-Reviriego, N.: Variability of extreme
 4608 precipitation over Europe and its relationships with teleconnection patterns, *Hydrol. Earth Syst. Sci.*, 18, 709-
 4609 725, <https://doi.org/10.5194/hess-18-709-2014>, 2014.



- 4610 Casini, M., Lövgren, J., Hjelm, J., Cardinale, M., Molinero, J.-C., and Kornilovs, G.: Multi-level trophic
 4611 cascades in a heavily exploited open marine ecosystem, *P. Roy. Soc. B-Biol. Sci.*, 275, 1793-1801,
 4612 <https://doi.org/10.1098/rspb.2007.1752> 2008.
- 4613 Casini, M., Kornilovs, G., Cardinale, M., Möllmann, C., Grygiel, W., Jonsson, P., Raid, T., Flinkman, J., and
 4614 Feldman, V.: Spatial and temporal density dependence regulates the condition of central Baltic Sea clupeids:
 4615 compelling evidence using an extensive international acoustic survey, *Popul. Ecol.*, 53, 511-523,
 4616 <https://doi.org/10.1007/s10144-011-0269-2>, 2011.
- 4617 Casini, M., Käll, F., Hansson, M., Plikshs, M., Baranova, T., Karlsson, O., Lundström, K., Neuenfeldt, S.,
 4618 Gårdmark, A., and Hjelm, J.: Hypoxic areas, density-dependence and food limitation drive the body condition of
 4619 a heavily exploited marine fish predator, *Roy. Soc. Open Sci.*, 3, 160416, <https://doi.org/10.1098/rsos.160416>,
 4620 2016.
- 4621 Cattiaux, J., and Cassou, C.: Opposite CMIP3/CMIP5 trends in the wintertime Northern Annular Mode
 4622 explained by combined local sea ice and remote tropical influences, *Geophys. Res. Lett.*, 40, 3682-3687,
 4623 <https://doi.org/10.1002/grl.50643>, 2013.
- 4624 Cattiaux, J., Douville, H., and Peings, Y.: European temperatures in CMIP5: origins of present-day biases and
 4625 future uncertainties, *Clim. Dyn.*, 41, 2889-2907, <https://doi.org/10.1007/s00382-013-1731-y>, 2013.
- 4626 Ceppi, P., and Hartmann, D. L.: Connections Between Clouds, Radiation, and Midlatitude Dynamics: a Review,
 4627 *Curr. Clim. Chang. Rep.*, 1, 94-102, <https://doi.org/10.1007/s40641-015-0010-x>, 2015.
- 4628 Chadburn, S. E., Burke, E. J., Cox, P. M., Friedlingstein, P., Hugelius, G., and Westermann, S.: An observation-
 4629 based constraint on permafrost loss as a function of global warming, *Nat. Clim. Change*, 7, 340-344,
 4630 <https://doi.org/10.1038/nclimate3262>, 2017.
- 4631 Chang, E. K. M., Guo, Y., and Xia, X.: CMIP5 multimodel ensemble projection of storm track change under
 4632 global warming, *J. Geophys. Res-Atmos.*, 117, <https://doi.org/10.1029/2012JD018578>, 2012.
- 4633 Chang, E. K. M., Ma, C.-G., Zheng, C., and Yau, A. M. W.: Observed and projected decrease in Northern
 4634 Hemisphere extratropical cyclone activity in summer and its impacts on maximum temperature, *Geophys. Res.*
 4635 *Lett.*, 43, 2200-2208, <https://doi.org/10.1002/2016GL068172>, 2016.
- 4636 Chang, E. K. M., and Yau, A. M. W.: Northern Hemisphere winter storm track trends since 1959 derived from
 4637 multiple reanalysis datasets, *Clim. Dyn.*, 47, 1435-1454, <https://doi.org/10.1007/s00382-015-2911-8>, 2016.
- 4638 Chen, D., Zhang, P., Seftigen, K., Ou, T., Giese, M., and Barthel, R.: Hydroclimate changes over Sweden in the
 4639 twentieth and twenty-first centuries: a millennium perspective, *Geogr. Ann. A*, 1-29,
 4640 <https://doi.org/10.1080/04353676.2020.1841410>, 2020.
- 4641 Christensen, J. H., Larsen, M. A. D., Christensen, O. B., Drews, M., and Stendel, M.: Robustness of European
 4642 climate projections from dynamical downscaling, *Clim. Dyn.*, 53, 4857-4869, <https://doi.org/10.1007/s00382-019-04831-z>, 2019.
- 4644 Christensen, O. B., and Kjellström, E.: Projections for Temperature, Precipitation, Wind, and Snow in the Baltic
 4645 Sea Region until 2100, *Oxford Research Encyclopedia of Climate Science*, Oxford University Press,
 4646 <https://doi.org/10.1093/acrefore/9780190228620.013.695>, 2018.
- 4647 Christensen, O. B., Kjellström, E., Dieterich, C., Gröger, M., and Meier, H. E. M.: Atmospheric regional climate
 4648 projections for the Baltic Sea Region until 2100, *Earth Syst. Dynam.*, 2021.
- 4649 Chust, G., Allen, J. I., Bopp, L., Schrum, C., Holt, J., Tsiaras, K., Zavatarelli, M., Chifflet, M., Cannaby, H.,
 4650 Dadou, I., Daewel, U., Wakelin, S. L., Machu, E., Pushpadas, D., Butenschon, M., Artioli, Y., Petihakis, G.,
 4651 Smith, C., Garçon, V., Goubanova, K., Le Vu, B., Fach, B. A., Salihoglu, B., Clementi, E., and Irigoien, X.:
 4652 Biomass changes and trophic amplification of plankton in a warmer ocean, *Global Change Biol.*, 20, 2124-2139,
 4653 <https://doi.org/10.1111/gcb.12562>, 2014.
- 4654 Ciasto, L. M., Li, C., Wettstein, J. J., and Kvamstø, N. G.: North Atlantic Storm-Track Sensitivity to Projected
 4655 Sea Surface Temperature: Local versus Remote Influences, *J. Climate*, 29, 6973-6991,
 4656 <https://doi.org/10.1175/jcli-d-15-0860.1>, 2016.
- 4657 Claremar, B., Haglund, K., and Rutgersson, A.: Ship emissions and the use of current air cleaning technology:
 4658 contributions to air pollution and acidification in the Baltic Sea, *Earth Syst. Dynam.*, 8, 901-919,
 4659 <https://doi.org/10.5194/esd-8-901-2017>, 2017.



- 4660 Clausen, K. K., Stjernholm, M., and Clausen, P.: Grazing management can counteract the impacts of climate
 4661 change-induced sea level rise on salt marsh-dependent waterbirds, *J. Appl. Ecol.*, 50, 528-537,
 4662 <https://doi.org/10.1111/1365-2664.12043>, 2013.
- 4663 Clausen, K. K., and Clausen, P.: Forecasting future drowning of coastal waterbird habitats reveals a major
 4664 conservation concern, *Biol. Conserv.*, 171, 177-185, <https://doi.org/10.1016/j.biocon.2014.01.033>, 2014.
- 4665 Colette, A., Bessagnet, B., Vautard, R., Szopa, S., Rao, S., Schucht, S., Klimont, Z., Menut, L., Clain, G.,
 4666 Meleux, F., CURCI, G., and Rouil, L.: European atmosphere in 2050, a regional air quality and climate
 4667 perspective under CMIP5 scenarios, *Atmos. Chem. Phys.*, 13, 7451-7471, [https://doi.org/10.5194/acp-13-7451-](https://doi.org/10.5194/acp-13-7451-2013)
 4668 2013, 2013.
- 4669 Colette, A., Andersson, C., Baklanov, A., Bessagnet, B., Brandt, J., Christensen, J. H., Doherty, R., Engardt, M.,
 4670 Geels, C., Giannakopoulos, C., Hedegaard, G. B., Katragkou, E., Langner, J., Lei, H., Manders, A., Melas, D.,
 4671 Meleux, F., Rouil, L., Sofiev, M., Soares, J., Stevenson, D. S., Tombrou-Tzella, M., Varotsos, K. V., and Young,
 4672 P.: Is the ozone climate penalty robust in Europe?, *Environ. Res. Lett.*, 10, 084015, [https://doi.org/10.1088/1748-](https://doi.org/10.1088/1748-9326/10/8/084015)
 4673 9326/10/8/084015, 2015.
- 4674 Colette, A., and Rouil, L.: Air Quality Trends in Europe: 2000-2017, European Environment Agency, Kjeller,
 4675 Norway, Eionet Report - ETC/ATNI 2019/16, 36pp, 2020.
- 4676 Collins, M., Knutti, R., Arblaster, J., Dufresne, J.-L., Fichet, T., Friedlingstein, P., Gao, X., Gutowski, W. J.,
 4677 Johns, T., Krinner, G., and others: Long-term climate change: projections, commitments and irreversibility, In:
 4678 IPCC Chapter 12, Climate Change 2013: The Physical Science Basis. IPCC Working Group I Contribution to
 4679 IPCC AR5; Cambridge University Press, Cambridge, UK and New York, USA, 1029-1136, 2013.
- 4680 Conley, D. J., Humborg, C., Rahm, L., Savchuk, O. P., and Wulff, F.: Hypoxia in the Baltic Sea and basin-scale
 4681 changes in phosphorus biogeochemistry, *Environ. Sci. Technol.*, 36, 5315-5320,
 4682 <https://doi.org/10.1021/es025763w>, 2002.
- 4683 Conley, D. J., Carstensen, J., Aigars, J., Axe, P., Bonsdorff, E., Eremina, T., Haahti, B.-M., Humborg, C.,
 4684 Jonsson, P., Kotta, J., Lännegren, C., Larsson, U., Maximov, A., Medina, M. R., Lysiak-Pastuszak, E.,
 4685 Remeikaitė-Nikienė, N., Walve, J., Wilhelms, S., and Zillén, L.: Hypoxia Is Increasing in the Coastal Zone of
 4686 the Baltic Sea, *Environ. Sci. Technol.*, 45, 6777-6783, <https://doi.org/10.1021/es201212r>, 2011.
- 4687 Cornes, R. C., van der Schrier, G., and Squintu, A. A.: A reappraisal of the thermal growing season length across
 4688 Europe, *Int. J. Climatol.*, 39, 1787-1795, <https://doi.org/10.1002/joc.5913>, 2019.
- 4689 Coumou, D., Lehmann, J., and Beckmann, J.: The weakening summer circulation in the Northern Hemisphere
 4690 mid-latitudes, *Science*, 348, 324, <https://doi.org/10.1126/science.1261768>, 2015.
- 4691 Cyberski, J., Wróblewski, A., and Stewart, J.: Riverine water inflows and the Baltic Sea water volume 1901-
 4692 1990, *Hydrol. Earth Syst. Sci.*, 4, 1-11, <https://doi.org/10.5194/hess-4-1-2000>, 2000.
- 4693 Defeo, O., McLachlan, A., Schoeman, D. S., Schlacher, T. A., Dugan, J., Jones, A., Lastra, M., and Scapini, F.:
 4694 Threats to sandy beach ecosystems: A review, *Estuar. Coast. Shelf Sci.*, 81, 1-12,
 4695 <https://doi.org/10.1016/j.ecss.2008.09.022>, 2009.
- 4696 Deng, J., Zhang, W., Harff, J., Schneider, R., Dudzinska-Nowak, J., Terefenko, P., Giza, A., and Furmanczyk,
 4697 K.: A numerical approach for approximating the historical morphology of wave-dominated coasts—A case study
 4698 of the Pomeranian Bight, southern Baltic Sea, *Geomorphology*, 204, 425-443,
 4699 <https://doi.org/10.1016/j.geomorph.2013.08.023>, 2014.
- 4700 Deng, J., Harff, J., Schimanke, S., and Meier, H. E. M.: A method for assessing the coastline recession due to the
 4701 sea level rise by assuming stationary wind-wave climate, *Oceanol. Hydrobiol. Stud.*, Vol. 44, No. 3, 362-380,
 4702 <https://doi.org/10.1515/ohs-2015-0035>, 2015.
- 4703 Deng, J., Wu, J., Zhang, W., Dudzinska-Nowak, J., and Harff, J.: Characterising the relaxation distance of
 4704 nearshore submarine morphology: A southern Baltic Sea case study, *Geomorphology*, 327, 365-376,
 4705 <https://doi.org/10.1016/j.geomorph.2018.11.018>, 2019.
- 4706 Deser, C., Hurrell, J. W., and Phillips, A. S.: The role of the North Atlantic Oscillation in European climate
 4707 projections, *Clim. Dyn.*, 49, 3141-3157, <https://doi.org/10.1007/s00382-016-3502-z>, 2017.
- 4708 Dethloff, K., Rinke, A., Benkel, A., Körtzow, M., Sokolova, E., Kumar Saha, S., Handorf, D., Dorn, W., Rockel,
 4709 B., von Storch, H., Haugen, J. E., Røed, L. P., Roeckner, E., Christensen, J. H., and Stendel, M.: A dynamical



- 4710 link between the Arctic and the global climate system, *Geophys. Res. Lett.*, 33,
 4711 <https://doi.org/10.1029/2005GL025245>, 2006.
- 4712 Devictor, V., van Swaay, C., Brereton, T., Brotons, L., Chamberlain, D., Heliölä, J., Herrando, S., Julliard, R.,
 4713 Kuussaari, M., Lindström, Å., Reif, J., Roy, D. B., Schweiger, O., Settele, J., Stefanescu, C., Van Strien, A., Van
 4714 Turnhout, C., Vermouzek, Z., WallisDeVries, M., Wynhoff, I., and Jiguet, F.: Differences in the climatic debts
 4715 of birds and butterflies at a continental scale, *Nat. Clim. Change*, 2, 121-124,
 4716 <https://doi.org/10.1038/nclimate1347>, 2012.
- 4717 Dieterich, C., Schimanke, S., Wang, S., Väli, G., Liu, Y., Hordoir, R., Höglund, A., and Meier, H. E. M.:
 4718 Evaluation of the SMHI coupled atmosphere-ice-ocean model RCA4-NEMO, SMHI, Norrköping, Sweden, 80,
 4719 2013.
- 4720 Dieterich, C., Wang, S., Schimanke, S., Gröger, M., Klein, B., Hordoir, R., Samuelsson, P., Liu, Y., Axell, L.,
 4721 and Höglund, A.: Surface heat budget over the North Sea in climate change simulations, *Atmosphere*, 10, 272,
 4722 <https://doi.org/10.3390/atmos10050272>, 2019.
- 4723 Dippner, J. W., Vuorinen, I., Daunys, D., Flinkman, J., Halkka, A., Köster, F. W., Lehtikoinen, E., MacKenzie,
 4724 B. R., Möllmann, C., Møhlenberg, F., Olenin, S., Schiedek, D., Skov, H., and Wasmund, N.: Climate-related
 4725 Marine Ecosystem Change, in: *Assessment of Climate Change for the Baltic Sea Basin*, Springer Berlin
 4726 Heidelberg, Berlin, Heidelberg, 309-377, https://doi.org/10.1007/978-3-540-72786-6_5, 2008.
- 4727 Dippner, J. W., Fründt, B., and Hammer, C.: Lake or Sea? The Unknown Future of Central Baltic Sea Herring,
 4728 *Front. Ecol. Evol.*, 7, <https://doi.org/10.3389/fevo.2019.00143>, 2019.
- 4729 Doherty, R. M., Heal, M. R., and O'Connor, F. M.: Climate change impacts on human health over Europe
 4730 through its effect on air quality, *Environ. Health-Glob.*, 16, 118, <https://doi.org/10.1186/s12940-017-0325-2>,
 4731 2017.
- 4732 Dong, B., Sutton, R. T., and Shaffrey, L.: Understanding the rapid summer warming and changes in temperature
 4733 extremes since the mid-1990s over Western Europe, *Clim. Dyn.*, 48, 1537-1554, [https://doi.org/10.1007/s00382-](https://doi.org/10.1007/s00382-016-3158-8)
 4734 016-3158-8, 2017.
- 4735 Donnelly, C., Yang, W., and Dahné, J.: River discharge to the Baltic Sea in a future climate, *Clim. Change*, 122,
 4736 157-170, <https://doi.org/10.1007/s10584-013-0941-y>, 2014.
- 4737 Donnelly, C., Greuell, W., Andersson, J., Gerten, D., Pisacane, G., Roudier, P., and Ludwig, F.: Impacts of
 4738 climate change on European hydrology at 1.5, 2 and 3 degrees mean global warming above preindustrial level,
 4739 *Clim. Change*, 143, 13-26, <https://doi.org/10.1007/s10584-017-1971-7>, 2017.
- 4740 Dreier, N., Nehlsen, E., Fröhle, P., Rechid, D., Bouwer, L. M., and Pfeifer, S.: Future Changes in Wave
 4741 Conditions at the German Baltic Sea Coast Based on a Hybrid Approach Using an Ensemble of Regional
 4742 Climate Change Projections, *Water*, 13, <https://doi.org/10.3390/w13020167>, 2021.
- 4743 Du, J., Shen, J., Park, K., Wang, Y. P., and Yu, X.: Worsened physical condition due to climate change
 4744 contributes to the increasing hypoxia in Chesapeake Bay, *Sci. Total Environ.*, 630, 707-717,
 4745 <https://doi.org/10.1016/j.scitotenv.2018.02.265>, 2018.
- 4746 Duarte, C. M., Hendriks, I. E., Moore, T. S., Olsen, Y. S., Steckbauer, A., Ramajo, L., Carstensen, J., Trotter, J.
 4747 A., and McCulloch, M.: Is Ocean Acidification an Open-Ocean Syndrome? Understanding Anthropogenic
 4748 Impacts on Seawater pH, *Estuar. Coast.*, 36, 221-236, <https://doi.org/10.1007/s12237-013-9594-3>, 2013.
- 4749 Ducklow, H. W., Morán, X. A. G., and Murray, A. E.: Bacteria in the Greenhouse: Marine Microbes and
 4750 Climate Change, in: *Environ. Microbiol.*, edited by: Mitchell, R., and Gu, J. D., Wiley-Blackwell, Hoboken,
 4751 N.J., USA, 1-31, <https://doi.org/10.1002/9780470495117.ch1>, 2009.
- 4752 Dudzińska-Nowak, J.: Morphodynamic Processes of the Swina Gate Coastal Zone Development (Southern
 4753 Baltic Sea), in: *Coastline Changes of the Baltic Sea from South to East: Past and Future Projection*, edited by:
 4754 Harff, J., Furmańczyk, K., and von Storch, H., Springer International Publishing, Cham, 219-255,
 4755 https://doi.org/10.1007/978-3-319-49894-2_11, 2017.
- 4756 Dyrørdal, A. V., Saloranta, T., Skaugen, T., and Stranden, H. B.: Changes in snow depth in Norway during the
 4757 period 1961–2010, *Hydrol. Res.*, 44, 169-179, <https://doi.org/10.2166/nh.2012.064>, 2012.
- 4758 Easterling, D. R., Kunkel, K. E., Wehner, M. F., and Sun, L.: Detection and attribution of climate extremes in
 4759 the observed record, *Weather. Clim. Extremes*, 11, 17-27, <https://doi.org/10.1016/j.wace.2016.01.001>, 2016.



- 4760 Eero, M., Andersson, H. C., Almroth-Rosell, E., and MacKenzie, B. R.: Has eutrophication promoted forage fish
 4761 production in the Baltic Sea?, *Ambio*, 45, 649-660, <https://doi.org/10.1007/s13280-016-0788-3>, 2016.
- 4762 Efremova, T., Palshin, N., and Zdorovenov, R.: Long-term characteristics of ice phenology in Karelian lakes,
 4763 *Est. J. Earth Sci.*, 62, 33-41, <https://doi.org/10.3176/earth.2013.04>, 2013.
- 4764 Eggert, A., and Schneider, B.: A nitrogen source in spring in the surface mixed-layer of the Baltic Sea: evidence
 4765 from total nitrogen and total phosphorus data, *Journal of Marine Systems*, 148, 39-47,
 4766 <https://doi.org/10.1016/j.jmarsys.2015.01.005>, 2015.
- 4767 Ehrnsten, E., Bauer, B., and Gustafsson, B. G.: Combined Effects of Environmental Drivers on Marine Trophic
 4768 Groups – A Systematic Model Comparison, *Front. Mar. Sci.*, 6, <https://doi.org/10.3389/fmars.2019.00492>, 2019.
- 4769 Ehrnsten, E., Norkko, A., Müller-Karulis, B., Gustafsson, E., and Gustafsson, B. G.: The meagre future of
 4770 benthic fauna in a coastal sea—Benthic responses to recovery from eutrophication in a changing climate, *Global
 4771 Change Biol.*, 26, 2235-2250, <https://doi.org/10.1111/gcb.15014>, 2020.
- 4772 Eichner, M., Rost, B., and Kranz, S. A.: Diversity of ocean acidification effects on marine N₂ fixers, *J. Exp.
 4773 Mar. Biol. Ecol.*, 457, 199-207, <https://doi.org/10.1016/j.jembe.2014.04.015>, 2014.
- 4774 Eilola, K., Rosell, E. A., Dieterich, C., Fransner, F., Höglund, A., and Meier, H. E. M.: Modeling Nutrient
 4775 Transports and Exchanges of Nutrients Between Shallow Regions and the Open Baltic Sea in Present and Future
 4776 Climate, *Ambio*, 41, 586-599, <https://doi.org/10.1007/s13280-012-0322-1>, 2012.
- 4777 Eilola, K., Mårtensson, S., and Meier, H. E. M.: Modeling the impact of reduced sea ice cover in future climate
 4778 on the Baltic Sea biogeochemistry, *Geophys. Res. Lett.*, 40, 149-154, <https://doi.org/10.1029/2012GL054375>,
 4779 2013.
- 4780 Eilola, K., Almroth-Rosell, E., and Meier, H. E. M.: Impact of saltwater inflows on phosphorus cycling and
 4781 eutrophication in the Baltic Sea: a 3D model study, *Tellus A*, 66, 23985,
 4782 <https://doi.org/10.3402/tellusa.v66.23985>, 2014.
- 4783 Emeis, K.-C., van Beusekom, J., Callies, U., Ebinghaus, R., Kannen, A., Kraus, G., Kröncke, I., Lenhart, H.,
 4784 Lorkowski, L., Matthias, V., Möllmann, C., Pätsch, J., Scharfe, M., Thomas, H., Weisse, R., and Zorita, E.: The
 4785 North Sea — A shelf sea in the Anthropocene, *J. Mar. Syst.*, 141, 18-33,
 4786 <https://doi.org/10.1016/j.jmarsys.2014.03.012>, 2015.
- 4787 Emeis, K., Christiansen, C., Edelvang, K., Jähmlich, S., Kozuch, J., Laima, M., Leipe, T., Löffler, A., Lund-
 4788 Hansen, L. C., Miltner, A., Pazdro, K., Pempkowiak, J., Pollehne, F., Shimmield, T., Voss, M., and Witt, G.:
 4789 Material transport from the near shore to the basinal environment in the southern Baltic Sea: II: Synthesis of data
 4790 on origin and properties of material, *J. Mar. Syst.*, 35, 151-168, [https://doi.org/10.1016/S0924-7963\(02\)00127-6](https://doi.org/10.1016/S0924-7963(02)00127-6),
 4791 2002.
- 4792 EMEP: Transboundary particulate matter, photo-oxidants, acidifying and eutrophying components, Norwegian
 4793 Meteorological Institute, Oslo, Norway, EMEP Status Report 1/2018, 204pp, 2018.
- 4794 Enfield, D. B., Mestas-Núñez, A. M., and Trimble, P. J.: The Atlantic Multidecadal Oscillation and its relation to
 4795 rainfall and river flows in the continental U.S., *Geophys. Res. Lett.*, 28, 2077-2080,
 4796 <https://doi.org/10.1029/2000gl012745>, 2001.
- 4797 Estilow, T. W., Young, A. H., and Robinson, D. A.: A long-term Northern Hemisphere snow cover extent data
 4798 record for climate studies and monitoring, *Earth Syst. Sci. Data*, 7, 137-142, <https://doi.org/10.5194/essd-7-137-2015>,
 4799 2015.
- 4800 EEA - European Environment Agency: Air quality in Europe - 2019 report, Publications Office of the European
 4801 Union, Luxemburg, EEA Report No. 10/2019, 99pp, <https://doi.org/10.2800/822355>, 2019a.
- 4802 EEA - European Environment Agency: Heavy precipitation in Europe. Indicator Assessment CLIM 004. IND-
 4803 92-en: https://www.eea.europa.eu/ds_resolveuid/998fbc113cc84e9a978becd87079f874, access: 24-03-2021,
 4804 2019b.
- 4805 EEA - European Environment Agency: Global and European sea-level rise: <https://www.eea.europa.eu/data-and-maps/indicators/sea-level-rise-6/assessment>, access: 08-12-2020, 2019c.



- 4807 Eyering, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the
 4808 Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geosci. Model*
 4809 *Dev.*, 9, 1937-1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- 4810 Farinotti, D., Huss, M., Fürst, J. J., Landmann, J., Machguth, H., Maussion, F., and Pandit, A.: A consensus
 4811 estimate for the ice thickness distribution of all glaciers on Earth, *Nat. Geosci.*, 12, 168-173,
 4812 <https://doi.org/10.1038/s41561-019-0300-3>, 2019.
- 4813 Feistel, R., Nausch, G., and Wasmund, N.: State and evolution of the Baltic Sea, 1952-2005: a detailed 50-year
 4814 survey of meteorology and climate, physics, chemistry, biology, and marine environment, John Wiley & Sons,
 4815 Hoboken, NJ, 2008.
- 4816 Feldstein, S. B.: The Recent Trend and Variance Increase of the Annular Mode, *J. Climate*, 15, 88-94,
 4817 [https://doi.org/10.1175/1520-0442\(2002\)015<0088:Trtavi>2.0.Co;2](https://doi.org/10.1175/1520-0442(2002)015<0088:Trtavi>2.0.Co;2), 2002.
- 4818 Feser, F., Barcikowska, M., Krueger, O., Schenk, F., Weisse, R., and Xia, L.: Storminess over the North Atlantic
 4819 and northwestern Europe—A review, *Q. J. Roy. Meteor. Soc.*, 141, 350-382, <https://doi.org/10.1002/qj.2364>,
 4820 2015.
- 4821 Filazzola, A., Blagrove, K., Imrit, M. A., and Sharma, S.: Climate Change Drives Increases in Extreme Events
 4822 for Lake Ice in the Northern Hemisphere, *Geophys. Res. Lett.*, 47, e2020GL089608,
 4823 <https://doi.org/10.1029/2020GL089608>, 2020.
- 4824 Fischer-Bruns, I., Storch, H. v., González-Rouco, J. F., and Zorita, E.: Modelling the variability of midlatitude
 4825 storm activity on decadal to century time scales, *Clim. Dyn.*, 25, 461-476, [https://doi.org/10.1007/s00382-005-](https://doi.org/10.1007/s00382-005-0036-1)
 4826 0036-1, 2005.
- 4827 Fleming, Z. L., Doherty, R. M., von Schneidmesser, E., Malley, C. S., Cooper, O. R., Pinto, J. P., Colette, A.,
 4828 Xu, X., Simpson, D., Schultz, M. G., Lefohn, A. S., Hamad, S., Moolla, R., Solberg, S., and Feng, Z.:
 4829 Tropospheric Ozone Assessment Report: Present-day ozone distribution and trends relevant to human health,
 4830 *Elementa-Sci. Anthropol.*, 6, <https://doi.org/10.1525/elementa.273>, 2018.
- 4831 Fontrodona Bach, A., van der Schrier, G., Melsen, L. A., Klein Tank, A. M. G., and Teuling, A. J.: Widespread
 4832 and Accelerated Decrease of Observed Mean and Extreme Snow Depth Over Europe, *Geophys. Res. Lett.*, 45,
 4833 12,312-312,319, <https://doi.org/10.1029/2018GL079799>, 2018.
- 4834 Forster, P. M., Maycock, A. C., McKenna, C. M., and Smith, C. J.: Latest climate models confirm need for
 4835 urgent mitigation, *Nat. Clim. Change*, 10, 7-10, <https://doi.org/10.1038/s41558-019-0660-0>, 2020.
- 4836 Fox, A. D., Jónsson, J. E., Aarvak, T., Bregnballe, T., Christensen, T. K., Clausen, K. K., Clausen, P., Dalby, L.,
 4837 Holm, T. E., Pavón-Jordan, D., Laursen, K., Lehtikoinen, A., Lorentsen, S.-H., Møller, A. P., Nordström, M.,
 4838 Öst, M., Söderquist, P., and Roland Therkildsen, O.: Current and Potential Threats to Nordic Duck Populations
 4839 — A Horizon Scanning Exercise, *Annal. Zool.*, 52, 193-220, 128, 2015.
- 4840 Fox, A. D., Nielsen, R. D., and Petersen, I. K.: Climate-change not only threatens bird populations but also
 4841 challenges our ability to monitor them, *Ibis*, 161, 467-474, <https://doi.org/10.1111/ibi.12675>, 2019.
- 4842 Frajka-Williams, E., Beaulieu, C., and Duchez, A.: Emerging negative Atlantic Multidecadal Oscillation index in
 4843 spite of warm subtropics, *Sci. Rep.-UK*, 7, 11224, <https://doi.org/10.1038/s41598-017-11046-x>, 2017.
- 4844 Francis, J. A., and Vavrus, S. J.: Evidence linking Arctic amplification to extreme weather in mid-latitudes,
 4845 *Geophys. Res. Lett.*, 39, <https://doi.org/10.1029/2012GL051000>, 2012.
- 4846 Francis, J. A., and Vavrus, S. J.: Evidence for a wavier jet stream in response to rapid Arctic warming, *Environ.*
 4847 *Res. Lett.*, 10, 014005, <https://doi.org/10.1088/1748-9326/10/1/014005>, 2015.
- 4848 Fransner, F., Gustafsson, E., Tedesco, L., Vichi, M., Hordoir, R., Roquet, F., Spilling, K., Kuznetsov, I., Eilola,
 4849 K., Mörth, C.-M., Humborg, C., and Nycander, J.: Non-Redfieldian Dynamics Explain Seasonal pCO₂
 4850 Drawdown in the Gulf of Bothnia, *J. Geophys. Res.-Oceans*, 123, 166-188,
 4851 <https://doi.org/10.1002/2017JC013019>, 2018.
- 4852 Friedland, R., Neumann, T., and Schernewski, G.: Climate change and the Baltic Sea action plan: model
 4853 simulations on the future of the western Baltic Sea, *J. Mar. Syst.*, 105-108, 175-186,
 4854 <https://doi.org/10.1016/j.jmarsys.2012.08.002>, 2012.



- 4855 Friedlingstein, P., O'Sullivan, M., Jones, M. W., Andrew, R. M., Hauck, J., Olsen, A., Peters, G. P., Peters, W.,
 4856 Pongratz, J., Sitch, S., Quéré, C. L., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S., Aragão, L. E. O. C.,
 4857 Arneeth, A., Arora, V., Bates, N. R., Becker, M., Benoit-Cattin, A., Bittig, H. C., Bopp, L., Bultan, S., Chandra,
 4858 N., Chevallier, F., Chini, L. P., Evans, W., Florentie, L., Forster, P. M., Gasser, T., Gehlen, M., Gilfillan, D.,
 4859 Gkritzalis, T., Gregor, L., Gruber, N., Harris, I., Hartung, K., Haverd, V., Houghton, R. A., Ilyina, T., Jain, A.
 4860 K., Joetzjer, E., Kadono, K., Kato, E., Kitidis, V., Korsbakken, J. I., Landschützer, P., Lefèvre, N., Lenton, A.,
 4861 Lienert, S., Liu, Z., Lombardozzi, D., Marland, G., Metzl, N., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-I.,
 4862 Niwa, Y., O'Brien, K., Ono, T., Palmer, P. I., Pierrot, D., Poulter, B., Resplandy, L., Robertson, E., Rödenbeck,
 4863 C., Schwinger, J., Séférian, R., Skjelvan, I., Smith, A. J. P., Sutton, A. J., Tanhua, T., Tans, P. P., Tian, H.,
 4864 Tilbrook, B., Werf, G. v. d., Vuichard, N., Walker, A. P., Wanninkhof, R., Watson, A. J., Willis, D., Wiltshire,
 4865 A. J., Yuan, W., Yue, X., and Zaehle, S.: Global carbon budget 2020, *Earth Syst. Sci. Data*, 12, 3269–3340,
 4866 <https://doi.org/10.5194/essd-12-3269-2020>, 2020.
- 4867 Frommel, A. Y., Schubert, A., Piatkowski, U., and Clemmesen, C.: Egg and early larval stages of Baltic cod,
 4868 *Gadus morhua*, are robust to high levels of ocean acidification, *Mar. Biol.*, 160, 1825–1834,
 4869 <https://doi.org/10.1007/s00227-011-1876-3>, 2013.
- 4870 Fronzek, S., Luoto, M., and Carter, T. R.: Potential effect of climate change on the distribution of palustrine mires in
 4871 subarctic Fennoscandia, *Clim. Res.*, 32, 1–12, <https://doi.org/10.3354/cr032001>, 2006.
- 4872 Fronzek, S., and Carter, T. R.: Assessing uncertainties in climate change impacts on resource potential for
 4873 Europe based on projections from RCMs and GCMs, *Clim. Change*, 81, 357–371,
 4874 <https://doi.org/10.1007/s10584-006-9214-3>, 2007.
- 4875 Fuchs, M., Kuhry, P., and Hugelius, G.: Low below-ground organic carbon storage in a subarctic Alpine
 4876 permafrost environment, *Cryosphere*, 9, 427–438, <https://doi.org/10.5194/tc-9-427-2015>, 2015.
- 4877 Gaget, E., Pavón-Jordán, D., Johnston, A., Lehtikainen, A., Hochachka, W. M., Sandercock, B. K., Soultan, A.,
 4878 Azafzaf, H., Bendjedda, N., Bino, T., Božić, L., Clausen, P., Dakki, M., Devos, K., Domsa, C., Encarnação, V.,
 4879 Erciyas-Yavuz, K., Faragó, S., Frost, T., Gaudard, C., Gosztonyi, L., Haas, F., Hornman, M., Langendoen, T.,
 4880 Ieronymidou, C., Kostyushin, V. A., Lewis, L. J., Lorentsen, S.-H., Luijckx, L., Meissner, W., Mikuska, T.,
 4881 Molina, B., Musilová, Z., Natykanets, V., Paquet, J.-Y., Petkov, N., Portolou, D., Ridzoň, J., Sayoud, S., Šćiban,
 4882 M., Sniauksta, L., Stipnice, A., Strebel, N., Teufelbauer, N., Topić, G., Uzunova, D., Vizi, A., Wahl, J.,
 4883 Zenatello, M., and Brommer, J. E.: Benefits of protected areas for nonbreeding waterbirds adjusting their
 4884 distributions under climate warming, *Conserv. Biol.*, n/a, <https://doi.org/10.1111/cobi.13648>, 2021.
- 4885 Gailiūšis, B., Kriaučiūnienė, J., Jakimavičius, D., and Šarausienė, D.: The variability of long-term runoff series
 4886 in the Baltic Sea drainage basin, *Baltica*, 24, 45–54, 2011.
- 4887 Gaillard, M.-J., Kleinen, T., Samuelsson, P., Nielsen, A. B., Bergh, J., Kaplan, J., Poska, A., Sandström, C.,
 4888 Strandberg, G., Trondman, A.-K., and Wramneby, A.: Causes of Regional Change—Land Cover, in: *Second*
 4889 *Assessment of Climate Change for the Baltic Sea Basin*, edited by: BACC II Author Team, Springer
 4890 International Publishing, Cham, 453–477, https://doi.org/10.1007/978-3-319-16006-1_25, 2015.
- 4891 Galatius, A., Kinze, C. C., and Teilmann, J.: Population structure of harbour porpoises in the Baltic region:
 4892 evidence of separation based on geometric morphometric comparisons, *J. Mar. Biol. Assoc. UK*, 92, 1669–1676,
 4893 <https://doi.org/10.1017/S0025315412000513>, 2012.
- 4894 Gálos, B., Hagemann, S., Hänsler, A., Kindermann, G., Rechid, D., Sieck, K., Teichmann, C., and Jacob, D.:
 4895 Case study for the assessment of the biogeophysical effects of a potential afforestation in Europe, *Carbon Bal.*
 4896 *Manage.*, 8, 3, <https://doi.org/10.1186/1750-0680-8-3>, 2013.
- 4897 Gao, Y., Markkanen, T., Backman, L., Henttonen, H. M., Pietikäinen, J. P., Mäkelä, H. M., and Laaksonen, A.:
 4898 Biogeophysical impacts of peatland forestation on regional climate changes in Finland, *Biogeosciences*, 11,
 4899 7251–7267, <https://doi.org/10.5194/bg-11-7251-2014>, 2014.
- 4900 Gårdmark, A., Lindegren, M., Neuenfeldt, S., Blenckner, T., Heikinheimo, O., Müller-Karulis, B., Niiranen, S.,
 4901 Tomczak, M. T., Aro, E., Wikström, A., and Möllmann, C.: Biological ensemble modeling to evaluate potential
 4902 futures of living marine resources, *Ecol. Appl.*, 23, 742–754, <https://doi.org/10.1890/12-0267.1>, 2013.
- 4903 Gårdmark, A., and Huss, M.: Individual variation and interactions explain food web responses to global
 4904 warming, *Philos. T. Roy. Soc. B.*, 375, 20190449, <https://doi.org/10.1098/rstb.2019.0449>, 2020.



- 4905 Garzke, J., Ismar, S. M. H., and Sommer, U.: Climate change affects low trophic level marine consumers:
 4906 warming decreases copepod size and abundance, *Oecologia*, 177, 849-860, <https://doi.org/10.1007/s00442-014->
 4907 3130-4, 2015.
- 4908 Gauss, M., Jonson, J. E., and Nyíri, Á.: Emissions from international shipping, The Norwegian Meteorological
 4909 Institute Oslo, Norway, 129-133, 2017.
- 4910 Gauss, M., Bartnicki, J., Jalkanen, J.-P., Nyiri, A., Klein, H., Fagerli, H., and Klimont, Z.: Airborne nitrogen
 4911 deposition to the Baltic Sea: Past trends, source allocation and future projections, *Atmos. Environ.*, 253, 118377,
 4912 <https://doi.org/10.1016/j.atmosenv.2021.118377>, 2021.
- 4913 Geels, C., Andersson, C., Hänninen, O., Lansø, A. S., Schwarze, P. E., Skjøth, C. A., and Brandt, J.: Future
 4914 Premature Mortality Due to O₃, Secondary Inorganic Aerosols and Primary PM in Europe — Sensitivity to
 4915 Changes in Climate, Anthropogenic Emissions, Population and Building Stock, *Int. J. Env. Res. Pub. He.*, 12,
 4916 2837-2869, 2015.
- 4917 Gillett, N. P., Arora, V. K., Matthews, D., and Allen, M. R.: Constraining the Ratio of Global Warming to
 4918 Cumulative CO₂ Emissions Using CMIP5 Simulations, *J. Climate*, 26, 6844-6858, <https://doi.org/10.1175/jcli-d->
 4919 12-00476.1, 2013.
- 4920 Girjatowicz, J. P., and Łabuz, T. A.: Forms of piled ice at the southern coast of the Baltic Sea, *Estuar. Coast.*
 4921 *Shelf Sci.*, 239, 106746, <https://doi.org/10.1016/j.ecss.2020.106746>, 2020.
- 4922 Gisnås, K., Eitzelmüller, B., Lussana, C., Hjort, J., Sannel, A. B. K., Isaksen, K., Westermann, S., Kuhry, P.,
 4923 Christiansen, H. H., Frampton, A., and Åkerman, J.: Permafrost Map for Norway, Sweden and Finland,
 4924 *Permafrost Periglac.*, 28, 359-378, <https://doi.org/10.1002/ppp.1922>, 2017.
- 4925 Glantz, P., Freud, E., Johansson, C., Noone, K. J., and Tesche, M.: Trends in MODIS and AERONET derived
 4926 aerosol optical thickness over Northern Europe, *Tellus B*, 71, 1554414,
 4927 <https://doi.org/10.1080/16000889.2018.1554414>, 2019.
- 4928 Graf, A., Klosterhalfen, A., Arriga, N., Bernhofer, C., Bogen, H., Bornet, F., Brüggemann, N., Brümmer, C.,
 4929 Buchmann, N., Chi, J., Chipeaux, C., Cremonese, E., Cuntz, M., Dušek, J., El-Madany, T. S., Fares, S., Fischer,
 4930 M., Foltýnová, L., Gharun, M., Ghiasi, S., Gielen, B., Gottschalk, P., Grünwald, T., Heinemann, G., Heinesch,
 4931 B., Heliasz, M., Holst, J., Hörtnagl, L., Ibrom, A., Ingwersen, J., Jurasinski, G., Klatt, J., Knohl, A., Koebsch, F.,
 4932 Konopka, J., Korkiakoski, M., Kowalska, N., Kremer, P., Kruijt, B., Lafont, S., Léonard, J., Ligne, A. D.,
 4933 Longdoz, B., Loustau, D., Magliulo, V., Mammarella, I., Manca, G., Mauder, M., Migliavacca, M., Mölder, M.,
 4934 Neiryneck, J., Ney, P., Nilsson, M., Paul-Limoges, E., Peichl, M., Pitacco, A., Poyda, A., Rebmann, C., Roland,
 4935 M., Sachs, T., Schmidt, M., Schrader, F., Siebke, L., Šigut, L., Tuittila, E.-S., Varlagin, A., Vendrame, N.,
 4936 Vincke, C., Völksch, I., Weber, S., Wille, C., Wizenmann, H.-D., Zeeman, M., and Vereecken, H.: Altered energy
 4937 partitioning across terrestrial ecosystems in the European drought year 2018, *Philos. T. Roy. Soc. B.*, 375,
 4938 20190524, <https://doi.org/10.1098/rstb.2019.0524>, 2020.
- 4939 Graham, L. P.: Climate Change Effects on River Flow to the Baltic Sea, *Ambio* 33, 235-241, 237,
 4940 <https://doi.org/10.1579/0044-7447-33.4.235>, 2004.
- 4941 Graham, P.: Modeling runoff to the Baltic Sea, *Ambio*, 28, 328-334, 1999.
- 4942 Graiff, A., Liesner, D., Karsten, U., and Bartsch, I.: Temperature tolerance of western Baltic Sea *Fucus*
 4943 *vesiculosus* – growth, photosynthesis and survival, *J. Exp. Mar. Biol. Ecol.*, 471, 8-16,
 4944 <https://doi.org/10.1016/j.jembe.2015.05.009>, 2015.
- 4945 Granéli, E., Kerstin, W., Larsson, U., Granéli, W., and Elmgren, R.: Nutrient Limitation of Primary Production
 4946 in the Baltic Sea Area, *Ambio*, 19, 142-151, 1990.
- 4947 Gräwe, U., and Burchard, H.: Storm surges in the Western Baltic Sea: the present and a possible future, *Clim.*
 4948 *Dyn.*, 39, 165-183, <https://doi.org/10.1007/s00382-011-1185-z>, 2012.
- 4949 Gräwe, U., Naumann, M., Mohrholz, V., and Burchard, H.: Anatomizing one of the largest saltwater inflows into
 4950 the Baltic Sea in December 2014, *J. Geophys. Res-Oceans*, 120, 7676-7697,
 4951 <https://doi.org/10.1002/2015JC011269>, 2015.
- 4952 Griffiths, J. R., Kadin, M., Nascimento, F. J. A., Tamelander, T., Törnroos, A., Bonaglia, S., Bonsdorff, E.,
 4953 Brüchert, V., Gärdmark, A., Järnström, M., Kotta, J., Lindegren, M., Nordström, M. C., Norkko, A., Olsson, J.,
 4954 Weigel, B., Żydelski, R., Blenckner, T., Niiranen, S., and Winder, M.: The importance of benthic–pelagic



- 4955 coupling for marine ecosystem functioning in a changing world, *Global Change Biol.*, 23, 2179-2196,
 4956 <https://doi.org/10.1111/gcb.13642>, 2017.
- 4957 Griffiths, J. R., Lehtinen, S., Suikkanen, S., and Winder, M.: Limited evidence for common interannual trends in
 4958 Baltic Sea summer phytoplankton biomass, *PLoS ONE*, 15, e0231690,
 4959 <https://doi.org/10.1371/journal.pone.0231690>, 2020.
- 4960 Grinsted, A.: Projected Change - Sea Level, in: Second Assessment of Climate Change for the Baltic Sea Basin,
 4961 edited by: BACC II Author Team, *Regional Climate Studies*, Springer International Publishing, Cham, 253-263,
 4962 https://doi.org/10.1007/978-3-319-16006-1_14, 2015.
- 4963 Grinsted, A., Jevrejeva, S., Riva, R. E. M., and Dahl-Jensen, D.: Sea level rise projections for northern Europe
 4964 under RCP8.5, *Clim. Res.*, 64, 15-23, <https://doi.org/10.3354/cr01309> 2015.
- 4965 Grise, K. M., and Polvani, L. M.: The response of midlatitude jets to increased CO₂: Distinguishing the roles of
 4966 sea surface temperature and direct radiative forcing, *Geophys. Res. Lett.*, 41, 6863-6871,
 4967 <https://doi.org/10.1002/2014GL061638>, 2014.
- 4968 Groetsch, P. M. M., Simis, S. G. H., Eleveld, M. A., and Peters, S. W. M.: Spring blooms in the Baltic Sea have
 4969 weakened but lengthened from 2000 to 2014, *Biogeosciences*, 13, 4959-4973, [https://doi.org/10.5194/bg-13-](https://doi.org/10.5194/bg-13-4959-2016)
 4970 4959-2016, 2016.
- 4971 Gröger, M., Dieterich, C., Meier, H. E. M., and Schimanke, S.: Thermal air-sea coupling in hindcast simulations
 4972 for the North Sea and Baltic Sea on the NW European shelf, *Tellus A*, 67, 26911,
 4973 <https://doi.org/10.3402/tellusa.v67.26911>, 2015.
- 4974 Gröger, M., Arneborg, L., Dieterich, C., Höglund, A., and Meier, H. E. M.: Summer hydrographic changes in the
 4975 Baltic Sea, Kattegat and Skagerrak projected in an ensemble of climate scenarios downscaled with a coupled
 4976 regional ocean–sea ice–atmosphere model, *Clim. Dyn.*, <https://doi.org/10.1007/s00382-019-04908-9>, 2019.
- 4977 Gröger, M., Dieterich, C., Ho-Hagemann, H. T. M., Hagemann, S., Jakacki, J., May, W., Meier, H. E. M.,
 4978 Miller, P. A., Rutgersson, A., and Wu, L.: Coupled regional Earth system modelling in the Baltic Sea region,
 4979 *Earth Syst. Dynam. Discuss.*, 1-50, <https://doi.org/10.5194/esd-2021-14>, 2021a.
- 4980 Gröger, M., Dieterich, C., and Meier, H. E. M.: Is interactive air sea coupling relevant for simulating the future
 4981 climate of Europe?, *Clim. Dyn.*, 56, 491-514, <https://doi.org/10.1007/s00382-020-05489-8>, 2021b.
- 4982 Groh, A., Richter, A., and Dietrich, R.: Recent Baltic Sea Level Changes Induced by Past and Present Ice
 4983 Masses, in: *Coastline Changes of the Baltic Sea from South to East: Past and Future Projection*, edited by: Harff,
 4984 J., Furmańczyk, K., and von Storch, H., Springer International Publishing, Cham, 55-68,
 4985 https://doi.org/10.1007/978-3-319-49894-2_4, 2017.
- 4986 Groll, N., Grabemann, I., Hünicke, B., and Meese, M.: Baltic Sea wave conditions under climate change
 4987 scenarios, *Boreal Environ. Res.*, 22, 1-12, 2017.
- 4988 Groß, D., Zander, A., Boethius, A., Dreibrodt, S., Grøn, O., Hansson, A., Jessen, C., Koivisto, S., Larsson, L.,
 4989 Lübke, H., and Nilsson, B.: People, lakes and seashores: Studies from the Baltic Sea basin and adjacent areas in
 4990 the early and Mid-Holocene, *Quaternary Sci. Rev.*, 185, 27-40, <https://doi.org/10.1016/j.quascirev.2018.01.021>,
 4991 2018.
- 4992 Gubelit, Y. I.: Climatic impact on community of filamentous macroalgae in the Neva estuary (eastern Baltic
 4993 Sea), *Mar. Pollut. Bull.*, 91, 166-172, <https://doi.org/10.1016/j.marpolbul.2014.12.009>, 2015.
- 4994 Gunnarsson, G., Waldenström, J., and Fransson, T.: Direct and indirect effects of winter harshness on the
 4995 survival of Mallards *Anas platyrhynchos* in northwest Europe, *Ibis*, 154, 307-317, [https://doi.org/10.1111/j.1474-](https://doi.org/10.1111/j.1474-919X.2011.01206.x)
 4996 919X.2011.01206.x, 2012.
- 4997 Gustafsson, B. G., Schenk, F., Blenckner, T., Eilola, K., Meier, H. E. M., Müller-Karulis, B., Neumann, T.,
 4998 Ruoho-Airola, T., Savchuk, O. P., and Zorita, E.: Reconstructing the development of Baltic Sea eutrophication
 4999 1850-2006, *Ambio*, 41, 534-548, <https://doi.org/10.1007/s13280-012-0318-x>, 2012.
- 5000 Gustafsson, E., Deutsch, B., Gustafsson, B. G., Humborg, C., and Mörrth, C. M.: Carbon cycling in the Baltic Sea
 5001 — The fate of allochthonous organic carbon and its impact on air–sea CO₂ exchange, *J. Mar. Syst.*, 129, 289-
 5002 302, <https://doi.org/10.1016/j.jmarsys.2013.07.005>, 2014.



- 5003 Gustafsson, E., Savchuk, O. P., Gustafsson, B. G., and Müller-Karulis, B.: Key processes in the coupled carbon,
 5004 nitrogen, and phosphorus cycling of the Baltic Sea, *Biogeochemistry*, 134, 301-317,
 5005 <https://doi.org/10.1007/s10533-017-0361-6>, 2017.
- 5006 Gustafsson, E., Hagens, M., Sun, X., Reed, D. C., Humborg, C., Slomp, C. P., and Gustafsson, B. G.:
 5007 Sedimentary alkalinity generation and long-term alkalinity development in the Baltic Sea, *Biogeosciences*, 16,
 5008 437-456, <https://doi.org/10.5194/bg-2018-313>, 2019.
- 5009 Haapala, J., Meier, H. E. M., and Rinne, J.: Numerical investigations of future ice conditions in the Baltic Sea,
 5010 *Ambio*, 30, 237-244, <https://doi.org/10.1579/0044-7447-30.4.237>, 2001.
- 5011 Haarsma, R. J., Selten, F. M., and Drijfhout, S. S.: Decelerating Atlantic meridional overturning circulation main
 5012 cause of future west European summer atmospheric circulation changes, *Environ. Res. Lett.*, 10, 094007,
 5013 <https://doi.org/10.1088/1748-9326/10/9/094007>, 2015.
- 5014 Haavisto, F., and Jormalainen, V.: Seasonality elicits herbivores' escape from trophic control and favors induced
 5015 resistance in a temperate macroalga, *Ecology*, 95, 3035-3045, <https://doi.org/10.1890/13-2387.1>, 2014.
- 5016 Hägg, H. E., Lyon, S. W., Wällstedt, T., Möhrh, C.-M., Claremar, B., and Humborg, C.: Future Nutrient Load
 5017 Scenarios for the Baltic Sea Due to Climate and Lifestyle Changes, *Ambio* 43, 337-351,
 5018 <https://doi.org/10.1007/s13280-013-0416-4>, 2014.
- 5019 Halkka, A.: Changing climate and the Baltic region biota, Doctoral dissertation, Faculty of Biological and
 5020 Environmental Sciences, University of Helsinki, Helsinki, Finland, 52 pp., 2020.
- 5021 Hallett, C. S., Hobday, A. J., Tweedley, J. R., Thompson, P. A., McMahon, K., and Valesini, F. J.: Observed and
 5022 predicted impacts of climate change on the estuaries of south-western Australia, a Mediterranean climate region,
 5023 *Reg. Environ. Change*, 18, 1357-1373, <https://doi.org/10.1007/s10113-017-1264-8>, 2018.
- 5024 Hällfors, H., Backer, H., Leppänen, J.-M., Hällfors, S., Hällfors, G., and Kuosa, H.: The northern Baltic Sea
 5025 phytoplankton communities in 1903–1911 and 1993–2005: a comparison of historical and modern species data,
 5026 *Hydrobiologia*, 707, 109-133, <https://doi.org/10.1007/s10750-012-1414-4>, 2013.
- 5027 Hamlington, B. D., Gardner, A. S., Ivins, E., Lenaerts, J. T. M., Reager, J. T., Trossman, D. S., Zaron, E. D.,
 5028 Adhikari, S., Arendt, A., Aschwanden, A., Beckley, B. D., Bekaert, D. P. S., Blewitt, G., Caron, L., Chambers,
 5029 D. P., Chandanpurkar, H. A., Christianson, K., Csatho, B., Cullather, R. I., DeConto, R. M., Fasullo, J. T.,
 5030 Frederikse, T., Freymueller, J. T., Gilford, D. M., Giroto, M., Hammond, W. C., Hock, R., Holschuh, N., Kopp,
 5031 R. E., Landerer, F., Larour, E., Menemenlis, D., Merrifield, M., Mitrovica, J. X., Nerem, R. S., Nias, I. J.,
 5032 Nieves, V., Nowicki, S., Pangaluru, K., Piecuch, C. G., Ray, R. D., Rounce, D. R., Schlegel, N.-J., Seroussi, H.,
 5033 Shirzaei, M., Sweet, W. V., Velicogna, I., Vinogradova, N., Wahl, T., Wiese, D. N., and Willis, M. J.:
 5034 Understanding of Contemporary Regional Sea-Level Change and the Implications for the Future, *Rev. Geophys.*,
 5035 58, e2019RG000672, <https://doi.org/10.1029/2019RG000672>, 2020.
- 5036 Hammer, K., Schneider, B., Kuliński, K., and Schulz-Bull, D. E.: Acid-base properties of Baltic Sea dissolved
 5037 organic matter, *J. Mar. Syst.*, 173, 114-121, <https://doi.org/10.1016/j.jmarsys.2017.04.007>, 2017.
- 5038 Hammond, P., Bearzi, G., Bjørge, A., Forney, K., Karczmarski, L., Kasuya, T., Perrin, W., Scott, M., Wang, J.,
 5039 and Wells, R.: *Phocoena phocoena* - Baltic Sea subpopulation (errata version published in 2016)(No. e.
 5040 T17031A98831650), The IUCN Red List of Threatened Species, 2008.
- 5041 Hannerz, F., and Destouni, G.: Spatial Characterization of the Baltic Sea Drainage Basin and Its Unmonitored
 5042 Catchments, *Ambio* 35, 214-219, 216, <https://doi.org/10.1579/05-A-022R.1>, 2006.
- 5043 Hänninen, J., Vuorinen, I., Rajasilta, M., and Reid, P. C.: Response of the Baltic and North Seas to river runoff
 5044 from the Baltic watershed – Physical and biological changes, *Prog. Oceanogr.*, 138, 91-104,
 5045 <https://doi.org/10.1016/j.pocan.2015.09.001>, 2015.
- 5046 Hansson, D., Eriksson, C., Omstedt, A., and Chen, D.: Reconstruction of river runoff to the Baltic Sea, AD
 5047 1500–1995, *Int. J. Climatol.*, 31, 696-703, <https://doi.org/10.1002/joc.2097>, 2011.
- 5048 Harff, J., Deng, J., Dudzińska-Nowak, J., Fröhle, P., Groh, A., Hünicke, B., Soomere, T., and Zhang, W.: What
 5049 Determines the Change of Coastlines in the Baltic Sea?, in: *Coastline Changes of the Baltic Sea from South to*
 5050 *East: Past and Future Projection*, edited by: Harff, J., Furmańczyk, K., and von Storch, H., Springer International
 5051 Publishing, Cham, 15-35, https://doi.org/10.1007/978-3-319-49894-2_2, 2017.



- 5052 Hartfield, G., Blunden, J., and Arndt, D. S.: State of the Climate in 2017, Bull. Amer. Meteor. Soc., 99, Si-S310,
 5053 <https://doi.org/10.1175/2018BAMSStateoftheClimate.1>, 2018.
- 5054 Harvey, B. J., Shaffrey, L. C., Woollings, T. J., Zappa, G., and Hodges, K. I.: How large are projected 21st
 5055 century storm track changes?, Geophys. Res. Lett., 39, <https://doi.org/10.1029/2012GL052873>, 2012.
- 5056 Harvey, B. J., Cook, P., Shaffrey, L. C., and Schiemann, R.: The Response of the Northern Hemisphere Storm
 5057 Tracks and Jet Streams to Climate Change in the CMIP3, CMIP5, and CMIP6 Climate Models, J. Geophys. Res-
 5058 Atmos., 125, e2020JD032701, <https://doi.org/10.1029/2020JD032701>, 2020.
- 5059 Hausfather, Z., and Peters, G. P.: Emissions - the 'business as usual' story is misleading, Nature, 577, 618-620,
 5060 <https://doi.org/10.1038/d41586-020-00177-3>, 2020.
- 5061 Hedegaard, G. B., Christensen, J. H., and Brandt, J.: The relative importance of impacts from climate change vs.
 5062 emissions change on air pollution levels in the 21st century, Atmos. Chem. Phys., 13, 3569-3585,
 5063 <https://doi.org/10.5194/acp-13-3569-2013>, 2013.
- 5064 Hegerl, G. C., Crowley, T. J., Baum, S. K., Kim, K.-Y., and Hyde, W. T.: Detection of volcanic, solar and
 5065 greenhouse gas signals in paleo-reconstructions of Northern Hemispheric temperature, Geophys. Res. Lett., 30,
 5066 <https://doi.org/10.1029/2002GL016635>, 2003.
- 5067 HELCOM: Climate Change in the Baltic Sea Area - HELCOM Thematic Assessment in 2007, Helsinki, Finland,
 5068 Baltic Sea Environment Proceedings No. 111, 54pp, 2007.
- 5069 HELCOM: HELCOM Copenhagen Ministerial Declaration - Taking Further Action to Implement the Baltic Sea
 5070 Action Plan - Reaching Good Environmental Status for a healthy Baltic Sea, Copenhagen, Denmark, HELCOM
 5071 Ministerial Meeting, 2013a.
- 5072 HELCOM: Climate Change in the Baltic Sea Area - HELCOM Thematic Assessment in 2013, Helsinki, Finland,
 5073 Baltic Sea Environment Proceedings No. 137, 70pp, 2013b.
- 5074 HELCOM: Updated Fifth Baltic Sea Pollution Load Compilation (PLC-5.5), Helsinki, Finland, Baltic Sea
 5075 Environment Proceedings No. 145, 143pp, 2015.
- 5076 HELCOM: HELCOM thematic assessment of eutrophication 2011-2016., Helsinki, Finland, Baltic Sea
 5077 Environment Proceedings No. 156, 83pp, 2018a.
- 5078 HELCOM: Population trends and abundance of seals, HELCOM core indicator report, 34pp, 2018b.
- 5079 HELCOM: The Sixth Pollution Load Compilation (PLC-6), Helsinki, Finland, 15pp, 2018c.
- 5080 Helle, E.: Reproduction, size and structure of the Baltic ringed seal population of the Bothnian Bay., PhD, Acta
 5081 Universitatis Ouluensis series A Scientiae rerum naturalium No. 106 Biologica No. 11, 47 pp., 1980.
- 5082 Hendriks, C., Forsell, N., Kieseewetter, G., Schaap, M., and Schöpp, W.: Ozone concentrations and damage for
 5083 realistic future European climate and air quality scenarios, Atmos. Environ., 144, 208-219,
 5084 <https://doi.org/10.1016/j.atmosenv.2016.08.026>, 2016.
- 5085 Hense, I., Meier, H. E. M., and Sonntag, S.: Projected climate change impact on Baltic Sea cyanobacteria, Clim.
 5086 Change, 119, 391-406, <https://doi.org/10.1007/s10584-013-0702-y>, 2013.
- 5087 Hesse, C., Krysanova, V., Stefanova, A., Bielecka, M., and Domnin, D. A.: Assessment of climate change
 5088 impacts on water quantity and quality of the multi-river Vistula Lagoon catchment, Hydrol. Sci. J., 60, 890-911,
 5089 <https://doi.org/10.1080/02626667.2014.967247>, 2015.
- 5090 Hieronymus, J., Eilola, K., Hieronymus, M., Meier, H. E. M., Saraiva, S., and Karlson, B.: Causes of simulated
 5091 long-term changes in phytoplankton biomass in the Baltic proper: a wavelet analysis, Biogeosciences, 15, 5113-
 5092 5129, <https://doi.org/10.5194/bg-15-5113-2018>, 2018.
- 5093 Hieronymus, M., and Kalén, O.: Sea-level rise projections for Sweden based on the new IPCC special report:
 5094 The ocean and cryosphere in a changing climate, Ambio, <https://doi.org/10.1007/s13280-019-01313-8>, 2020.
- 5095 Hinrichsen, H.-H., Huwer, B., Makarchouk, A., Petereit, C., Schaber, M., and Voss, R.: Climate-driven long-
 5096 term trends in Baltic Sea oxygen concentrations and the potential consequences for eastern Baltic cod (*Gadus*
 5097 *morhua*), ICES J. Mar. Sci., 68, 2019-2028, <https://doi.org/10.1093/icesjms/fsr145>, 2011.



- 5098 Hisdal, H., Holmqvist, E., Jonsdottir, J. F., Jonsson, P., Kuusisto, E., Lindstroem, G., and Roald, L. A.: Has
 5099 streamflow changed in the Nordic countries?, Norwegian Water Resources and Energy Directorate (NVE),
 5100 NorwayNVE-1/2010, 28pp, 2010.
- 5101 Hjerne, O., Hajdu, S., Larsson, U., Downing, A. S., and Winder, M.: Climate Driven Changes in Timing,
 5102 Composition and Magnitude of the Baltic Sea Phytoplankton Spring Bloom, *Front. Mar. Sci.*, 6,
 5103 <https://doi.org/10.3389/fmars.2019.00482>, 2019.
- 5104 Hock, R., Rasul, G., Adler, C., Cáceres, B., Gruber, S., Hirabayashi, Y., Jackson, M., Kääb, A., Kang, S.,
 5105 Kutuzov, S., Milner, A., Molau, U., Morin, S., Orlove, B., and Steltzer, H.: High Mountain Areas, in: IPCC
 5106 Special Report on the Ocean and Cryosphere in a Changing Climate, edited by: Pörtner, H.-O., Roberts, D. C.,
 5107 Masson-Delmotte, V., Zhai, P., Tignor, M., Poloczanska, E., Mintenbeck, K., Alegría, A., Nicolai, M., Okem,
 5108 A., Petzold, J., Rama, B., and Weyer, N. M., 2019.
- 5109 Höglund, A., Pemberton, P., Hordoir, R., and Schimanke, S.: Ice conditions for maritime traffic in the Baltic Sea
 5110 in future climate, *Boreal Environ. Res.*, 22, 245-265, 2017.
- 5111 Hoikkala, L., Kortelainen, P., Soinne, H., and Kuosa, H.: Dissolved organic matter in the Baltic Sea, *J. Mar.*
 5112 *Syst.*, 142, 47-61, <https://doi.org/10.1016/j.jmarsys.2014.10.005>, 2015.
- 5113 Holmlund, E. S., and Holmlund, P.: Constraining 135 years of mass balance with historic structure-from-motion
 5114 photogrammetry on Storglaciären, Sweden, *Geogr. Ann. A*, 101, 195-210,
 5115 <https://doi.org/10.1080/04353676.2019.1588543>, 2019.
- 5116 Holopainen, R., Lehtiniemi, M., Meier, H. E. M., Albertsson, J., Gorokhova, E., Kotta, J., and Viitasalo, M.:
 5117 Impacts of changing climate on the non-indigenous invertebrates in the northern Baltic Sea by end of the twenty-
 5118 first century, *Biol. Invasions*, 18, 3015-3032, <https://doi.org/10.1007/s10530-016-1197-z>, 2016.
- 5119 Holtermann, P. L., and Umlauf, L.: The Baltic Sea Tracer Release Experiment: 2. Mixing processes, *J. Geophys.*
 5120 *Res-Oceans*, 117, <https://doi.org/10.1029/2011JC007445>, 2012.
- 5121 Holtermann, P. L., Umlauf, L., Tanhua, T., Schmale, O., Rehder, G., and Waniek, J. J.: The Baltic Sea Tracer
 5122 Release Experiment: 1. Mixing rates, *J. Geophys. Res-Oceans*, 117, <https://doi.org/10.1029/2011JC007439>,
 5123 2012.
- 5124 Holtermann, P. L., Prien, R., Naumann, M., Mohrholz, V., and Umlauf, L.: Deepwater dynamics and mixing
 5125 processes during a major inflow event in the central Baltic Sea, *J. Geophys. Res-Oceans*, 122, 6648-6667,
 5126 <https://doi.org/10.1002/2017JC013050>, 2017.
- 5127 Hong, B., Swaney, D. P., McCrackin, M., Svanbäck, A., Humborg, C., Gustafsson, B., Yershova, A., and
 5128 Pakhomau, A.: Advances in NANI and NAPI accounting for the Baltic drainage basin: spatial and temporal
 5129 trends and relationships to watershed TN and TP fluxes, *Biogeochemistry*, 133, 245-261,
 5130 <https://doi.org/10.1007/s10533-017-0330-0>, 2017.
- 5131 Hook, O., Johnels, A. G., and Matthews, L. H.: The breeding and distribution of the grey seal (*Halichoerus*
 5132 *grypus* Fab.) in the Baltic Sea, with observations on other seals of the area, *P. Roy. Soc. B-Biol. Sci.*, 182, 37-58,
 5133 <https://doi.org/10.1098/rspb.1972.0065>, 1972.
- 5134 Hordoir, R., Axell, L., Löptien, U., Dietze, H., and Kuznetsov, I.: Influence of sea level rise on the dynamics of
 5135 salt inflows in the Baltic Sea, *J. Geophys. Res-Oceans*, 120, 6653-6668, <https://doi.org/10.1002/2014JC010642>,
 5136 2015.
- 5137 Hordoir, R., Axell, L., Höglund, A., Dieterich, C., Fransner, F., Gröger, M., Liu, Y., Pemberton, P., Schimanke,
 5138 S., Andersson, H., Ljungemyr, P., Nygren, P., Falahat, S., Nord, A., Jönsson, A., Lake, I., Döös, K.,
 5139 Hieronymus, M., Dietze, H., Löptien, U., Kuznetsov, I., Westerlund, A., Tuomi, L., and Haapala, J.: Nemo-
 5140 Nordic 1.0: a NEMO-based ocean model for the Baltic and North seas – research and operational applications,
 5141 *Geosci. Model Dev.*, 12, 363-386, <https://doi.org/10.5194/gmd-12-363-2019>, 2019.
- 5142 Horn, H. G., Boersma, M., Garzke, J., Löder, M. G. J., Sommer, U., and Aberle, N.: Effects of high CO₂ and
 5143 warming on a Baltic Sea microzooplankton community, *ICES J. Mar. Sci.*, 73, 772-782,
 5144 <https://doi.org/10.1093/icesjms/fsv198>, 2015.
- 5145 Høyer, J. L., and Karagali, I.: Sea Surface Temperature Climate Data Record for the North Sea and Baltic Sea, *J.*
 5146 *Climate*, 29, 2529-2541, <https://doi.org/10.1175/jcli-d-15-0663.1>, 2016.



- 5147 Hugelius, G., Loisel, J., Chadburn, S., Jackson, R. B., Jones, M., MacDonald, G., Marushchak, M., Olefeldt, D.,
 5148 Packalen, M., Siewert, M. B., Treat, C., Turetsky, M., Voigt, C., and Yu, Z.: Large stocks of peatland carbon and
 5149 nitrogen are vulnerable to permafrost thaw, *P. Natl. Acad. Sci. USA*, 117, 20438–20446,
 5150 <https://doi.org/10.1073/pnas.1916387117>, 2020.
- 5151 Humborg, C., Geibel, M. C., Sun, X., McCrackin, M., Mörrth, C.-M., Stranne, C., Jakobsson, M., Gustafsson, B.,
 5152 Sokolov, A., Norkko, A., and Norkko, J.: High Emissions of Carbon Dioxide and Methane From the Coastal
 5153 Baltic Sea at the End of a Summer Heat Wave, *Front. Mar. Sci.*, 6, <https://doi.org/10.3389/fmars.2019.00493>,
 5154 2019.
- 5155 Hünicke, B., and Zorita, E.: Trends in the amplitude of Baltic Sea level annual cycle, *Tellus A*, 60, 154–164,
 5156 <https://doi.org/10.1111/j.1600-0870.2007.00277.x>, 2008.
- 5157 Hünicke, B., Zorita, E., Soomere, T., Madsen, K. S., Johansson, M., and Suursaar, Ü.: Recent Change—Sea
 5158 Level and Wind Waves, in: *Second Assessment of Climate Change for the Baltic Sea Basin*, edited by: BACC II
 5159 Author Team, Springer International Publishing, Cham, 155–185, 10.1007/978-3-319-16006-1_9, 2015.
- 5160 Hünicke, B., and Zorita, E.: Statistical Analysis of the Acceleration of Baltic Mean Sea-Level Rise, 1900–2012,
 5161 *Front. Mar. Sci.*, 3, <https://doi.org/10.3389/fmars.2016.00125>, 2016.
- 5162 Hurrell, J. W.: Decadal trends in the North Atlantic Oscillation: regional temperatures and precipitation, *Science-*
 5163 *AAAS-Weekly Paper Edition*, 269, 676–679, <https://doi.org/10.1126/science.269.5224.676> 1995.
- 5164 Hurrell, J. W., Kushnir, Y., Ottersen, G., and Visbeck, M.: An overview of the North Atlantic oscillation,
 5165 *Geophys. Monogr. Ser.*, 134, 1–36, 2003.
- 5166 Huthnance, J., Weisse, R., Wahl, T., Thomas, H., Pietrzak, J., Souza, A. J., van Heteren, S., Schmelzer, N., van
 5167 Beusekom, J., Colijn, F., Haigh, I., Hjøllø, S., Holfort, J., Kent, E. C., Kühn, W., Loewe, P., Lorkowski, I.,
 5168 Mork, K. A., Pätsch, J., Quante, M., Salt, L., Siddorn, J., Smyth, T., Sterl, A., and Woodworth, P.: Recent
 5169 Change—North Sea, in: *North Sea Region Climate Change Assessment*, edited by: Quante, M., and Colijn, F.,
 5170 Springer International Publishing, Cham, 85–136, https://doi.org/10.1007/978-3-319-39745-0_3, 2016.
- 5171 Huttunen, I., Lehtonen, H., Huttunen, M., Piirainen, V., Korppoo, M., Veijalainen, N., Viitasalo, M., and
 5172 Vehviläinen, B.: Effects of climate change and agricultural adaptation on nutrient loading from Finnish
 5173 catchments to the Baltic Sea, *Sci. Total Environ.*, 529, 168–181, <https://doi.org/10.1016/j.scitotenv.2015.05.055>,
 5174 2015.
- 5175 Hyytiäinen, K., Bauer, B., Bly Joyce, K., Ehrnsten, E., Eilola, K., Gustafsson, B. G., Meier, H. E. M., Norkko,
 5176 A., Saraiva, S., Tomczak, M., and Zandersen, M.: Provision of aquatic ecosystem services as a consequence of
 5177 societal changes: The case of the Baltic Sea, *Popul. Ecol.*, 63, 61–74, <https://doi.org/10.1002/1438-390X.12033>,
 5178 2021.
- 5179 Ilmatieteen laitos: Jäätalvi Itämerellä: <https://www.ilmatieteenlaitos.fi/jaatalvi-itamerella>, The ice cover of the
 5180 Baltic 1961–2018. Public record of Finnish Meteorological Institute, access: 10.12.2020, 2020.
- 5181 IPCC: Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth
 5182 Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Team, C. W., Pachauri, R. K.,
 5183 and Meyer, L. A., Geneva, 2014a.
- 5184 IPCC: Climate Change 2013 – The Physical Science Basis: Working Group I Contribution to the Fifth
 5185 Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Stocker, T. F., Qin, D.,
 5186 Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M.,
 5187 Cambridge University Press, Cambridge, 2014b.
- 5188 IPCC: IPCC Special Report on the Ocean and Cryosphere in a Changing Climate, edited by: Pörtner, H.-O.,
 5189 Roberts, D. C., Masson-Delmotte, V., Zhai, P., Tignor, M., Poloczanska, E., Mintenbeck, K., Alegría, A.,
 5190 Nicolai, M., Okem, A., Petzold, J., Rama, B., and Weyer, N. M., 2019a.
- 5191 IPCC: Climate Change and Land: an IPCC special report on climate change, desertification, land degradation,
 5192 sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems, edited by:
 5193 Shukla, P. R., Skea, J., Buendia, E. C., Masson-Delmotte, V., Pörtner, H.-O., Roberts, D. C., Zhai, P., Slade, R.,
 5194 Connors, S., Diemen, R. v., Ferrat, M., Haughey, E., Luz, S., Neogi, S., Pathak, M., Petzold, J., Pereira, J. P.,
 5195 Vyas, P., Huntley, E., Kissick, K., Belkacemi, M., and Malley, J., 2019b.
- 5196 Irannezhad, M., Ronkanen, A.-K., and Kløve, B.: Wintertime climate factors controlling snow resource decline
 5197 in Finland, *Int. J. Climatol.*, 36, 110–131, <https://doi.org/10.1002/joc.4332>, 2016.



- 5198 Ito, M., Scotti, M., Franz, M., Barboza, F. R., Buchholz, B., Zimmer, M., Guy-Haim, T., and Wahl, M.: Effects
 5199 of temperature on carbon circulation in macroalgal food webs are mediated by herbivores, *Mar. Biol.*, 166, 158,
 5200 <https://doi.org/10.1007/s00227-019-3596-z>, 2019.
- 5201 Ivanov, V. A., and Belokopytov, V. N.: Oceanography of the Black Sea, National Academy of Sciences of
 5202 Ukraine, Marine Hydrophysical Institute, Sevastopol, Ukraine, 2013.
- 5203 Jaagus, J., Briede, A., Rimkus, E., and Remm, K.: Variability and trends in daily minimum and maximum
 5204 temperatures and in the diurnal temperature range in Lithuania, Latvia and Estonia in 1951–2010, *Theor. Appl.*
 5205 *Climatol.*, 118, 57–68, <https://doi.org/10.1007/s00704-013-1041-7>, 2014.
- 5206 Jaagus, J., Sepp, M., Tamm, T., Järvet, A., and Möisja, K.: Trends and regime shifts in climatic conditions and
 5207 river runoff in Estonia during 1951–2015, *Earth Syst. Dynam.*, 8, 963–976, [https://doi.org/10.5194/esd-8-963-](https://doi.org/10.5194/esd-8-963-2017)
 5208 2017, 2017.
- 5209 Jaagus, J., Briede, A., Rimkus, E., and Sepp, M.: Changes in precipitation regime in the Baltic countries in
 5210 1966–2015, *Theor. Appl. Climatol.*, 131, 433–443, <https://doi.org/10.1007/s00704-016-1990-8>, 2018.
- 5211 Jackson, L. C., Kahana, R., Graham, T., Ringer, M. A., Woollings, T., Mecking, J. V., and Wood, R. A.: Global
 5212 and European climate impacts of a slowdown of the AMOC in a high resolution GCM, *Clim. Dyn.*, 45, 3299–
 5213 3316, <https://doi.org/10.1007/s00382-015-2540-2>, 2015.
- 5214 Jacob, D., Kotova, L., Teichmann, C., Sobolowski, S. P., Vautard, R., Donnelly, C., Koutroulis, A. G., Grillakis,
 5215 M. G., Tsanis, I. K., Damm, A., Sakalli, A., and van Vliet, M. T. H.: Climate Impacts in Europe Under +1.5°C
 5216 Global Warming, *Earths Future*, 6, 264–285, <https://doi.org/10.1002/2017ef000710>, 2018.
- 5217 Jacob, D. J., and Winner, D. A.: Effect of climate change on air quality, *Atmos. Environ.*, 43, 51–63,
 5218 <https://doi.org/10.1016/j.atmosenv.2008.09.051>, 2009.
- 5219 Jakimavičius, D., Kriaučiūnienė, J., and Šarauskienė, D.: Impact of climate change on the Curonian Lagoon
 5220 water balance components, salinity and water temperature in the 21st century, *Oceanologia*, 60, 378–389,
 5221 <https://doi.org/10.1016/j.oceano.2018.02.003>, 2018.
- 5222 Jakobsson, M., Long, A., Ingólfsson, Ó., Kjær, K. H., and Spielhagen, R. F.: New insights on Arctic Quaternary
 5223 climate variability from palaeo-records and numerical modelling, *Quaternary Sci. Rev.*, 29, 3349–3358,
 5224 <https://doi.org/10.1016/j.quascirev.2010.08.016>, 2010.
- 5225 Jakubowska, M., Jerzak, M., Normant, M., Burska, D., and Drzazgowski, J.: Effect of Carbon Dioxide-Induced
 5226 Water Acidification on the Physiological Processes of the Baltic Isopod *Saduria entomon*, *J. Shellfish Res.*, 32,
 5227 825–834, 810, <https://doi.org/10.2983/035.032.0326>, 2013.
- 5228 Jänes, H., Herkül, K., and Kotta, J.: Environmental niche separation between native and non-native benthic
 5229 invertebrate species: Case study of the northern Baltic Sea, *Mar. Environ. Res.*, 131, 123–133,
 5230 <https://doi.org/10.1016/j.marenvres.2017.08.001>, 2017.
- 5231 Jansen, J., Thornton, B. F., Jammet, M. M., Wik, M., Cortés, A., Friborg, T., MacIntyre, S., and Crill, P. M.:
 5232 Climate-Sensitive Controls on Large Spring Emissions of CH₄ and CO₂ From Northern Lakes, *J. Geophys. Res-*
 5233 *Biogeo*, 124, 2379–2399, <https://doi.org/10.1029/2019JG005094>, 2019.
- 5234 Jansson, A., Lischka, S., Boxhammer, T., Schulz, K. G., and Norkko, J.: Survival and settling of larval *Macoma*
 5235 *balthica* in a large-scale mesocosm experiment at different fCO₂ levels, *Biogeosciences*, 13, 3377–3385,
 5236 <https://doi.org/10.5194/bg-13-3377-2016>, 2016.
- 5237 Jin, H., Jönsson, A. M., Olsson, C., Lindström, J., Jönsson, P., and Eklundh, L.: New satellite-based estimates
 5238 show significant trends in spring phenology and complex sensitivities to temperature and precipitation at
 5239 northern European latitudes, *Int. J. Biometeorol.*, 63, 763–775, <https://doi.org/10.1007/s00484-019-01690-5>,
 5240 2019.
- 5241 Johansson, L., Ytreberg, E., Jalkanen, J. P., Fridell, E., Eriksson, K. M., Lagerström, M., Maljutenko, I.,
 5242 Raudsepp, U., Fischer, V., and Roth, E.: Model for leisure boat activities and emissions – implementation for the
 5243 Baltic Sea, *Ocean Sci.*, 16, 1143–1163, <https://doi.org/10.5194/os-16-1143-2020>, 2020.
- 5244 Johansson, M. M., Pellikka, H., Kahma, K. K., and Ruosteenoja, K.: Global sea level rise scenarios adapted to
 5245 the Finnish coast, *J. Mar. Syst.*, 129, 35–46, <https://doi.org/10.1016/j.jmarsys.2012.08.007>, 2014.



- 5246 Jones, M. C., and Cheung, W. W. L.: Multi-model ensemble projections of climate change effects on global
 5247 marine biodiversity, *ICES J. Mar. Sci.*, 72, 741-752, <https://doi.org/10.1093/icesjms/fsu172>, 2014.
- 5248 Jones, P. D., Lister, D. H., Osborn, T. J., Harpham, C., Salmon, M., and Morice, C. P.: Hemispheric and large-
 5249 scale land-surface air temperature variations: An extensive revision and an update to 2010, *J. Geophys. Res-*
 5250 *Atmos.*, 117, <https://doi.org/10.1029/2011jd017139>, 2012.
- 5251 Jonson, J. E., Jalkanen, J. P., Johansson, L., Gauss, M., and van der Gon, H. A. C. D.: Model calculations of the
 5252 effects of present and future emissions of air pollutants from shipping in the Baltic Sea and the North Sea,
 5253 *Atmos. Chem. Phys.*, 15, 783-783, 2015.
- 5254 Jonson, J. E., Gauss, M., Jalkanen, J.-P., and Johansson, L.: Effects of strengthening the Baltic Sea ECA
 5255 regulations, *Atmos. Chem. Phys.*, 19, 2019.
- 5256 Jonsson, P. R., Kotta, J., Andersson, H. C., Herkül, K., Virtanen, E., Sandman, A. N., and Johannesson, K.: High
 5257 climate velocity and population fragmentation may constrain climate-driven range shift of the key habitat former
 5258 *Fucus vesiculosus*, *Divers. Distrib.*, 24, 892-905, <https://doi.org/10.1111/ddi.12733>, 2018.
- 5259 Jonsson, P. R., Moksnes, P.-O., Corell, H., Bonsdorff, E., and Nilsson Jacobi, M.: Ecological coherence of
 5260 Marine Protected Areas: New tools applied to the Baltic Sea network, *Aquat. Conserv.*, 30, 743-760,
 5261 <https://doi.org/10.1002/aqc.3286>, 2020.
- 5262 Josefsson, W.: Long-term global radiation in Stockholm, 1922-2018, SMHI (Swedish Meteorological and
 5263 Hydrological Institute), Norrköping, Sweden, 92pp, 2019.
- 5264 Jüssi, M., Härkönen, T., Helle, E., and Jüssi, I.: Decreasing Ice Coverage Will Reduce the Breeding Success of
 5265 Baltic Grey Seal (*Halichoerus grypu*) Females, *Ambio* 37, 80-85, 86, 2008.
- 5266 Jüssi, M.: Living on an edge: land-locked seals in changing climate, 2012.
- 5267 Jylhä, K., Laapas, M., Ruosteenoja, K., Arvola, L., Drebs, A., Kersalo, J., Saku, S., Gregow, H., Hannula, H.-R.,
 5268 and Pirinen, P. J. B. E. R.: Climate variability and trends in the Valkea-Kotinen region, southern Finland:
 5269 comparisons between the past, current and projected climates, *Boreal Environ. Res.*, 19, 4-30, 2014.
- 5270 Kadin, M., Frederiksen, M., Niiranen, S., and Converse, S. J.: Linking demographic and food-web models to
 5271 understand management trade-offs, *Ecol. Evol.*, 9, 8587-8600, <https://doi.org/10.1002/ece3.5385>, 2019.
- 5272 Kahru, M., and Elmgren, R.: Multidecadal time series of satellite-detected accumulations of cyanobacteria in the
 5273 Baltic Sea, *Biogeosciences*, 11, 3619-3633, <https://doi.org/10.5194/bg-11-3619-2014>, 2014.
- 5274 Kahru, M., Elmgren, R., and Savchuk, O. P.: Changing seasonality of the Baltic Sea, *Biogeosciences*, 13, 1009-
 5275 1018, <https://doi.org/10.5194/bg-13-1009-2016>, 2016.
- 5276 Kahru, M., Elmgren, R., Di Lorenzo, E., and Savchuk, O.: Unexplained interannual oscillations of
 5277 cyanobacterial blooms in the Baltic Sea, *Sci. Rep.-UK*, 8, 6365, <https://doi.org/10.1038/s41598-018-24829-7>,
 5278 2018.
- 5279 Kahru, M., Elmgren, R., Kaiser, J., Wasmund, N., and Savchuk, O.: Cyanobacterial blooms in the Baltic Sea:
 5280 Correlations with environmental factors, *Harmful Algae*, 92, 101739, <https://doi.org/10.1016/j.hal.2019.101739>,
 5281 2020.
- 5282 Kanakidou, M., Myriokefalitakis, S., and Tsigaridis, K.: Aerosols in atmospheric chemistry and biogeochemical
 5283 cycles of nutrients, *Environ. Res. Lett.*, 13, 063004, <https://doi.org/10.1088/1748-9326/aabedb>, 2018.
- 5284 Karl, M., Bieser, J., Geyer, B., Matthias, V., Jalkanen, J.-P., Johansson, L., and Fridell, E.: Impact of a nitrogen
 5285 emission control area (NECA) on the future air quality and nitrogen deposition to seawater in the Baltic Sea
 5286 region, *Atmos. Chem. Phys.*, 19, 1721-1752, <https://doi.org/10.5194/acp-2018-1107>, 2019a.
- 5287 Karl, M., Jonson, J. E., Uppstu, A., Aulinger, A., Prank, M., Sofiev, M., Jalkanen, J. P., Johansson, L., Quante,
 5288 M., and Matthias, V.: Effects of ship emissions on air quality in the Baltic Sea region simulated with three
 5289 different chemistry transport models, *Atmos. Chem. Phys.*, 19, 7019-7053, <https://doi.org/10.5194/acp-19-7019-2019>, 2019b.
- 5290
- 5291 Karlsson, K.-G., and Devasthale, A.: Inter-comparison and evaluation of the four longest satellite-derived cloud
 5292 climate data records: CLARA-A2, ESA Cloud CCI V3, ISCCP-HGM, and PATMOS-x, *Remote Sens.-Basel*, 10,
 5293 1567, <https://doi.org/10.3390/rs10101567>, 2018.



- 5294 Kauhala, K., Bäcklin, B.-M., Raitaniemi, J., and Harding, K. C.: The effect of prey quality and ice conditions on
 5295 the nutritional status of Baltic gray seals of different age groups, *Mammal Res.*, 62, 351-362,
 5296 <https://doi.org/10.1007/s13364-017-0329-x>, 2017.
- 5297 Kauhala, K., Bergenius, M., Isomursu, M., and Raitaniemi, J.: Reproductive rate and nutritional status of Baltic
 5298 ringed seals, *Mammal Res.*, 64, 109-120, 2019.
- 5299 Kauhala, K., and Kurkilahti, M.: Delayed effects of prey fish quality and winter temperature during the birth
 5300 year on adult size and reproductive rate of Baltic grey seals, *Mammal Res.*, 65, 117-126,
 5301 <https://doi.org/10.1007/s13364-019-00454-1>, 2020.
- 5302 Kauker, F., and Meier, H. E. M.: Modeling decadal variability of the Baltic Sea: 1. Reconstructing atmospheric
 5303 surface data for the period 1902-1998, *J. Geophys. Res-Oceans*, 108, <https://doi.org/10.1029/2003JC001797>,
 5304 2003.
- 5305 Kiani, S., Irannezhad, M., Ronkanen, A.-K., Moradkhani, H., and Kløve, B.: Effects of recent temperature
 5306 variability and warming on the Oulu-Hailuoto ice road season in the northern Baltic Sea, *Cold Reg. Sci.*
 5307 *Technol.*, 151, 1-8, <https://doi.org/10.1016/j.coldregions.2018.02.010>, 2018.
- 5308 Kim, B.-M., Son, S.-W., Min, S.-K., Jeong, J.-H., Kim, S.-J., Zhang, X., Shim, T., and Yoon, J.-H.: Weakening
 5309 of the stratospheric polar vortex by Arctic sea-ice loss, *Nat. Commun.*, 5, 4646,
 5310 <https://doi.org/10.1038/ncomms5646>, 2014.
- 5311 Kirchner, N., Noormets, R., Kutteneuler, J., Erstorp, E. S., Holmlund, E. S., Rosqvist, G., Holmlund, P.,
 5312 Wennbom, M., and Karlin, T.: High-resolution bathymetric mapping reveals subaqueous glacial landforms in the
 5313 Arctic alpine lake Tarfala, Sweden, *J. Quat. Sci.*, 34, 452-462, <https://doi.org/10.1002/jqs.3112>, 2019.
- 5314 Kirchner, N., Kutteneuler, K., Rosqvist, G., Hancke, M., Granebeck, A., Weckström, J., Weckström, K.,
 5315 Schenk, F., Korhola, A., and Eriksson, P.: A first continuous three-year temperature record from the dimictic
 5316 arctic-alpine Lake Tarfala, northern Sweden, *Arct. Antarct. Alp. Res.*, 2021.
- 5317 Kjellström, E., Bärring, L., Nikulin, G., Nilsson, C., Persson, G., and Strandberg, G.: Production and use of
 5318 regional climate model projections – A Swedish perspective on building climate services, *Clim. Serv.*, 2-3, 15-
 5319 29, <https://doi.org/10.1016/j.cliser.2016.06.004>, 2016.
- 5320 Kjellström, E., Nikulin, G., Strandberg, G., Christensen, O. B., Jacob, D., Keuler, K., Lenderink, G., van
 5321 Meijgaard, E., Schär, C., and Somot, S.: European climate change at global mean temperature increases of 1.5
 5322 and 2 degrees C above pre-industrial conditions as simulated by the EURO-CORDEX regional climate models,
 5323 *Earth Syst. Dynam.*, 9, 459-478, <https://doi.org/10.5194/esd-9-459-2018>, 2018.
- 5324 Kjerfve, B.: Oceanography of Chesapeake Bay, in: *Hydrodynamics of Estuaries*, edited by: Kjerfve, B., Taylor
 5325 & Francis group, CRC Press, 1988.
- 5326 Klais, R., Tamminen, T., Kremp, A., Spilling, K., and Olli, K.: Decadal-scale changes of dinoflagellates and
 5327 diatoms in the anomalous Baltic Sea spring bloom, *PLOS ONE* 6, e21567,
 5328 <https://doi.org/10.1371/journal.pone.0021567>, 2011.
- 5329 Klais, R., Tamminen, T., Kremp, A., Spilling, K., An, B. W., Hajdu, S., and Olli, K.: Spring phytoplankton
 5330 communities shaped by interannual weather variability and dispersal limitation: Mechanisms of climate change
 5331 effects on key coastal primary producers, *Limnol. Oceanogr.*, 58, 753-762,
 5332 <https://doi.org/10.4319/lo.2013.58.2.0753>, 2013.
- 5333 Kļaviņš, M., Avotniece, Z., and Rodinova, V.: Dynamics and Impacting Factors of Ice Regimes in Latvia Inland
 5334 and Coastal Waters, *Proceedings of the Latvian Academy of Sciences. Section B. Natural, Exact, and Applied*
 5335 *Sciences.*, 70, 400-408, <https://doi.org/10.1515/prolas-2016-0059>, 2016.
- 5336 Klawonn, I., Lavik, G., Böning, P., Marchant, H., Dekaezemacker, J., Mohr, W., and Ploug, H.: Simple approach
 5337 for the preparation of 15-15N₂-enriched water for nitrogen fixation assessments: evaluation, application and
 5338 recommendations, *Front. Microbiol.*, 6, <https://doi.org/10.3389/fmicb.2015.00769>, 2015.
- 5339 Kniebusch, M., Meier, H. E. M., Neumann, T., and Börgel, F.: Temperature Variability of the Baltic Sea Since
 5340 1850 and Attribution to Atmospheric Forcing Variables, *J. Geophys. Res-Oceans*, 124, 4168-4187,
 5341 <https://doi.org/10.1029/2018jc013948>, 2019a.



- 5342 Kniebusch, M., Meier, H. E. M., and Radtke, H.: Changing Salinity Gradients in the Baltic Sea As a
 5343 Consequence of Altered Freshwater Budgets, *Geophys. Res. Lett.*, 46, 9739-9747,
 5344 <https://doi.org/10.1029/2019GL083902>, 2019b.
- 5345 Knight, J. R., Folland, C. K., and Scaife, A. A.: Climate impacts of the Atlantic Multidecadal Oscillation,
 5346 *Geophys. Res. Lett.*, 33, <https://doi.org/10.1029/2006gl026242>, 2006.
- 5347 Knight, J. R.: The Atlantic Multidecadal Oscillation Inferred from the Forced Climate Response in Coupled
 5348 General Circulation Models, *J. Climate*, 22, 1610-1625, <https://doi.org/10.1175/2008jcli2628.1>, 2009.
- 5349 Knoll, L. B., Sharma, S., Denfeld, B. A., Flaim, G., Hori, Y., Magnuson, J. J., Straile, D., and Weyhenmeyer, G.
 5350 A.: Consequences of lake and river ice loss on cultural ecosystem services, *Limnol. Oceanogr. Lett.*, 4, 119-131,
 5351 <https://doi.org/10.1002/lol2.10116>, 2019.
- 5352 Knudsen, M.: Ein hydrographischer lehrsatz, *Ann. Hydr. u. mar. Meteor.*, 28, 316-320, 1900.
- 5353 Knudsen, M. F., Seidenkrantz, M.-S., Jacobsen, B. H., and Kuijpers, A.: Tracking the Atlantic Multidecadal
 5354 Oscillation through the last 8,000 years, *Nat. Commun.*, 2, 178, <https://doi.org/10.1038/ncomms1186>, 2011.
- 5355 Kong, D., MacLeod, M., and Cousins, I. T.: Modelling the influence of climate change on the chemical
 5356 concentrations in the Baltic Sea region with the POPCYCLING-Baltic model, *Chemosphere*, 110, 31-40,
 5357 <https://doi.org/10.1016/j.chemosphere.2014.02.044>, 2014.
- 5358 Kopp, R. E., Horton, R. M., Little, C. M., Mitrovica, J. X., Oppenheimer, M., Rasmussen, D. J., Strauss, B. H.,
 5359 and Tebaldi, C.: Probabilistic 21st and 22nd century sea-level projections at a global network of tide-gauge sites,
 5360 *Earth's Future*, 2, 383-406, <https://doi.org/10.1002/2014EF000239>, 2014.
- 5361 Korhonen, J.: Long-term changes and variability of the winter and spring season hydrological regime in Finland,
 5362 Faculty of Science, Institute for Atmospheric and Earth System Research, University of Helsinki, Finland, 2019.
- 5363 Korpinen, S., Honkanen, T., Vesakoski, O., Hemmi, A., Koivikko, R., Lopenen, J., and Jormalainen, V.:
 5364 Macroalgal Communities Face the Challenge of Changing Biotic Interactions: Review with Focus on the Baltic
 5365 Sea, *Ambio* 36, 203-211, 209, [https://doi.org/10.1579/0044-7447\(2007\)36\[203:MCFTCO\]2.0.CO;2](https://doi.org/10.1579/0044-7447(2007)36[203:MCFTCO]2.0.CO;2), 2007.
- 5366 Kortsch, S., Frelat, R., Pecuchet, L., Olivier, P., Putnis, I., Bonsdorff, E., Ojaveer, H., Jurgensone, I., Stråke, S.,
 5367 Rubene, G., Krüze, Ē., and Nordström, M. C.: Disentangling temporal food web dynamics facilitates
 5368 understanding of ecosystem functioning, *J. Anim. Ecol.*, <https://doi.org/10.1111/1365-2656.13447>, 2021.
- 5369 Köster, F. W., Huwer, B., Hinrichsen, H.-H., Neumann, V., Makarchouk, A., Eero, M., Dewitz, B. V., Hüßy,
 5370 K., Tomkiewicz, J., Margonski, P., Temming, A., Hermann, J.-P., Oesterwind, D., Dierking, J., Kotterba, P., and
 5371 Plikshs, M.: Eastern Baltic cod recruitment revisited—dynamics and impacting factors, *ICES J. Mar. Sci.*, 74, 3-
 5372 19, <https://doi.org/10.1093/icesjms/fsw172>, 2016.
- 5373 Kotta, J., Vanhatalo, J., Jänes, H., Orav-Kotta, H., Rugiu, L., Jormalainen, V., Bobsien, I., Viitasalo, M.,
 5374 Virtanen, E., Sandman, A. N., Isaeus, M., Leidenberger, S., Jonsson, P. R., and Johannesson, K.: Integrating
 5375 experimental and distribution data to predict future species patterns, *Sci. Rep.-UK*, 9, 1821,
 5376 <https://doi.org/10.1038/s41598-018-38416-3>, 2019.
- 5377 Kremp, A., Godhe, A., Egardt, J., Dupont, S., Suikkanen, S., Casabianca, S., and Penna, A.: Intraspecific
 5378 variability in the response of bloom-forming marine microalgae to changed climate conditions, *Ecol. Evol.*, 2,
 5379 1195-1207, <https://doi.org/10.1002/ece3.245>, 2012.
- 5380 Kremp, A., Oja, J., LeTortorec, A. H., Hakanen, P., Tahvanainen, P., Tuimala, J., and Suikkanen, S.: Diverse
 5381 seed banks favour adaptation of microalgal populations to future climate conditions, *Environ. Microbiol.*, 18,
 5382 679-691, <https://doi.org/10.1111/1462-2920.13070>, 2016.
- 5383 Krug, J., Eriksson, H., Heidecke, C., Kellomäki, S., Köhl, M., Lindner, M., and Saikkonen, K.: Socio-economic
 5384 Impacts—Forestry and Agriculture, in: Second Assessment of Climate Change for the Baltic Sea Basin, edited
 5385 by: BACC II Author Team, Springer International Publishing, Cham, 399-409, https://doi.org/10.1007/978-3-319-16006-1_21, 2015.
- 5387 Kudryavtseva, N., and Soomere, T.: Satellite altimetry reveals spatial patterns of variations in the Baltic Sea
 5388 wave climate, *Earth Syst. Dynam.*, 8, 697-706, <https://doi.org/10.5194/esd-8-697-2017>, 2017.



- 5389 Kühl, N., Gebhardt, C., Litt, T., and Hense, A.: Probability Density Functions as Botanical-Climatological
 5390 Transfer Functions for Climate Reconstruction, *Quat. Res.*, 58, 381-392, <https://doi.org/10.1006/qres.2002.2380>,
 5391 2002.
- 5392 Kuliński, K., Schneider, B., Hammer, K., Machulik, U., and Schulz-Bull, D.: The influence of dissolved organic
 5393 matter on the acid-base system of the Baltic Sea, *J. Mar. Syst.*, 132, 106-115,
 5394 <https://doi.org/10.1016/j.jmarsys.2014.01.011>, 2014.
- 5395 Kuliński, K., Hammer, K., Schneider, B., and Schulz-Bull, D.: Remineralization of terrestrial dissolved organic
 5396 carbon in the Baltic Sea, *Mar. Chem.*, 181, 10-17, <https://doi.org/10.1016/j.marchem.2016.03.002>, 2016.
- 5397 Kuliński, K., Schneider, B., Szymczycha, B., and Stokowski, M.: Structure and functioning of the acid-base
 5398 system in the Baltic Sea, *Earth Syst. Dynam.*, 8, 1107-1120, <https://doi.org/10.5194/esd-8-1107-2017>, 2017.
- 5399 Kuliński, K., Szymczycha, B., Koziorowska, K., Hammer, K., and Schneider, B.: Anomaly of total boron
 5400 concentration in the brackish waters of the Baltic Sea and its consequence for the CO₂ system calculations, *Mar.*
 5401 *Chem.*, 204, 11-19, <https://doi.org/10.1016/j.marchem.2018.05.007>, 2018.
- 5402 Kuliński, K., Rehder, G., Asmala, E., Bartosova, A., Carstensen, J., Gustafsson, B., Hall, P. O. J., Humborg, C.,
 5403 Jilbert, T., Jürgens, K., Meier, H. E. M., Müller-Karulis, B., Naumann, M., Olesen, J. E., Savchuk, O., Schramm,
 5404 A., Slomp, C. P., Sofiev, M., Sobek, A., Szymczycha, B., and Undeman, E.: Baltic Earth Assessment Report on
 5405 the biogeochemistry of the Baltic Sea, *Earth Syst. Dynam. Discuss.*, 1-93, <https://doi.org/10.5194/esd-2021-33>,
 5406 2021.
- 5407 Kundzewicz, Z. W., Ulbrich, U., Brücher, T., Graczyk, D., Krüger, A., Leckebusch, G. C., Menzel, L., Pińskwar,
 5408 I., Radziejewski, M., and Szwed, M.: Summer Floods in Central Europe – Climate Change Track?, *Nat. Hazards*,
 5409 36, 165-189, <https://doi.org/10.1007/s11069-004-4547-6>, 2005.
- 5410 Kuosa, H., Fleming-Lehtinen, V., Lehtinen, S., Lehtiniemi, M., Nygård, H., Raateoja, M., Raitaniemi, J.,
 5411 Tuimala, J., Uusitalo, L., and Suikkanen, S.: A retrospective view of the development of the Gulf of Bothnia
 5412 ecosystem, *J. Mar. Syst.*, 167, 78-92, <https://doi.org/10.1016/j.jmarsys.2016.11.020>, 2017.
- 5413 Kuznetsov, I., and Neumann, T.: Simulation of carbon dynamics in the Baltic Sea with a 3D model, *J. Mar.*
 5414 *Syst.*, 111-112, 167-174, <https://doi.org/10.1016/j.jmarsys.2012.10.011>, 2013.
- 5415 Laakso, L., Mikkonen, S., Drebs, A., Karjalainen, A., Pirinen, P., and Alenius, P.: 100 years of atmospheric and
 5416 marine observations at the Finnish Utö Island in the Baltic Sea, *Ocean Sci.*, 14, 617-632,
 5417 <https://doi.org/10.5194/os-14-617-2018>, 2018.
- 5418 Łabuz, T. A.: Environmental Impacts—Coastal Erosion and Coastline Changes, in: *Second Assessment of*
 5419 *Climate Change for the Baltic Sea Basin*, edited by: BACC II Author Team, Springer International Publishing,
 5420 Cham, 381-396, https://doi.org/10.1007/978-3-319-16006-1_20, 2015.
- 5421 Łabuz, T. A., Grunewald, R., Bobykina, V., Chubarenko, B., Česnulevičius, A., Baurénas, A., Morkūnaitė, R.,
 5422 and Tönnis, H.: Coastal Dunes of the Baltic Sea Shores: A Review, *Quaest. Geogr.*, 37, 47-71,
 5423 <https://doi.org/10.2478/quageo-2018-0005>, 2018.
- 5424 Lah, L., Trense, D., Benke, H., Berggren, P., Gunnlaugsson, P., Lockyer, C., Öztürk, A., Öztürk, B., Pawliczka,
 5425 I., Roos, A., Siebert, U., Skóra, K., Víkingsson, G., and Tiedemann, R.: Spatially Explicit Analysis of Genome-
 5426 Wide SNPs Detects Subtle Population Structure in a Mobile Marine Mammal, the Harbor Porpoise, *PLoS ONE*,
 5427 11, e0162792, <https://doi.org/10.1371/journal.pone.0162792>, 2016.
- 5428 Lamon, L., von Waldow, H., MacLeod, M., Scheringer, M., Marcomini, A., and Hungerbühler, K.: Modeling the
 5429 Global Levels and Distribution of Polychlorinated Biphenyls in Air under a Climate Change Scenario, *Environ.*
 5430 *Sci. Technol.*, 43, 5818-5824, <https://doi.org/10.1021/es900438j>, 2009.
- 5431 Lampe, M., and Lampe, R.: Evolution of a large Baltic beach ridge plain (Neudarss, NE Germany): A
 5432 continuous record of sea-level and wind-field variation since the Homeric Minimum, *Earth Surf. Process. Landf.*,
 5433 43, 3042-3056, <https://doi.org/10.1002/esp.4468>, 2018.
- 5434 Landerer, F. W., Jungclaus, J. H., and Marotzke, J.: Regional dynamic and steric sea level change in response to
 5435 the IPCC-A1B scenario, *J. Phys. Oceanogr.*, 37, 296-312, <https://doi.org/10.1175/JPO3013.1>, 2007.
- 5436 Lang, A., and Mikolajewicz, U.: The long-term variability of extreme sea levels in the German Bight, *Ocean*
 5437 *Sci.*, 15, 651-668, 2019.



- 5438 Langner, J., Engardt, M., Baklanov, A., Christensen, J. H., Gauss, M., Geels, C., Hedegaard, G. B., Nuterman,
 5439 R., Simpson, D., and Soares, J.: A multi-model study of impacts of climate change on surface ozone in Europe,
 5440 *Atmos. Chem. Phys.*, 12, 10423-10440, 2012.
- 5441 Lappe, C., and Umlauf, L.: Efficient boundary mixing due to near-inertial waves in a nontidal basin:
 5442 Observations from the Baltic Sea, *J. Geophys. Res.-Oceans*, 121, 8287-8304,
 5443 <https://doi.org/10.1002/2016JC011985>, 2016.
- 5444 Larsson, K., Hajdu, S., Kilpi, M., Larsson, R., Leito, A., and Lyngs, P.: Effects of an extensive *Prymnesium*
 5445 *polylepis* bloom on breeding eiders in the Baltic Sea, *J. Sea Res.*, 88, 21-28,
 5446 <https://doi.org/10.1016/j.seares.2013.12.017>, 2014.
- 5447 Larsson, U., Elmgren, R., and Wulff, F.: Eutrophication and the Baltic Sea: causes and consequences;
 5448 *L'eutrophisation et la mer Baltique: causes et conséquences*, *Ambio*, 14, 9-14, 1985.
- 5449 Larsson, U., Hajdu, S., Walve, J., and Elmgren, R.: Baltic Sea nitrogen fixation estimated from the summer
 5450 increase in upper mixed layer total nitrogen, *Limnol. Oceanogr.*, 46, 811-820,
 5451 <https://doi.org/10.4319/lo.2001.46.4.0811>, 2001.
- 5452 Laskar, J., Robutel, P., Joutel, F., Gastineau, M., Correia, A. C. M., and Levrard, B.: A long-term numerical
 5453 solution for the insolation quantities of the Earth, *Astron. Astrophys.*, 428, 261-285,
 5454 <https://doi.org/10.1051/0004-6361:20041335>, 2004.
- 5455 Lass, H. U., and Matthäus, W.: On temporal wind variations forcing salt water inflows into the Baltic Sea, *Tellus*
 5456 *A*, 48, 663-671, <https://doi.org/10.1034/j.1600-0870.1996.t01-4-00005.x>, 1996.
- 5457 Le Cozannet, G., Nicholls, R. J., Hinkel, J., Sweet, W. V., McInnes, K. L., Van de Wal, R. S. W., Slangen, A. B.
 5458 A., Lowe, J. A., and White, K. D.: Sea Level Change and Coastal Climate Services: The Way Forward, *J. Mar.*
 5459 *Sci. Eng.*, 5, <https://doi.org/10.3390/jmse5040049>, 2017.
- 5460 Leckebusch, G. C., and Ulbrich, U.: On the relationship between cyclones and extreme windstorm events over
 5461 Europe under climate change, *Global Planet. Change*, 44, 181-193,
 5462 <https://doi.org/10.1016/j.gloplacha.2004.06.011>, 2004.
- 5463 Lee, X., Goulden, M. L., Hollinger, D. Y., Barr, A., Black, T. A., Bohrer, G., Bracho, R., Drake, B., Goldstein,
 5464 A., Gu, L., Katul, G., Kolb, T., Law, B. E., Margolis, H., Meyers, T., Monson, R., Munger, W., Oren, R., Paw U,
 5465 K. T., Richardson, A. D., Schmid, H. P., Staebler, R., Wofsy, S., and Zhao, L.: Observed increase in local
 5466 cooling effect of deforestation at higher latitudes, *Nature*, 479, 384-387, <https://doi.org/10.1038/nature10588>,
 5467 2011.
- 5468 Lehikoinen, A., Kilpi, M., and Öst, M.: Winter climate affects subsequent breeding success of common eiders,
 5469 *Global Change Biol.*, 12, 1355-1365, <https://doi.org/10.1111/j.1365-2486.2006.01162.x>, 2006.
- 5470 Lehikoinen, A., and Jaatinen, K.: Delayed autumn migration in northern European waterfowl, *J. Ornithol.*, 153,
 5471 563-570, 2012.
- 5472 Lehikoinen, A., Jaatinen, K., Vähätalo, A. V., Clausen, P., Crowe, O., Deceuninck, B., Hearn, R., Holt, C. A.,
 5473 Hornman, M., Keller, V., Nilsson, L., Langendoen, T., Tománková, I., Wahl, J., and Fox, A. D.: Rapid climate
 5474 driven shifts in wintering distributions of three common waterbird species, *Global Change Biol.*, 19, 2071-2081,
 5475 <https://doi.org/10.1111/gcb.12200>, 2013.
- 5476 Lehmann, A., Krauß, W., and Hinrichsen, H.-H.: Effects of remote and local atmospheric forcing on circulation
 5477 and upwelling in the Baltic Sea, *Tellus A*, 54, 299-316, <https://doi.org/10.3402/tellusa.v54i3.12138>, 2002.
- 5478 Lehmann, A., Getzlaff, K., and Harlaß, J.: Detailed assessment of climate variability in the Baltic Sea area for
 5479 the period 1958 to 2009, *Clim. Res.*, 46, 185-196, <https://doi.org/10.3354/cr00876>, 2011.
- 5480 Lehmann, A., Myrberg, K., and Höflich, K.: A statistical approach to coastal upwelling in the Baltic Sea based
 5481 on the analysis of satellite data for 1990-2009, *Oceanologia*, 54, 369-393, <https://doi.org/10.5697/oc.54-3.369>,
 5482 2012.
- 5483 Lehmann, A., and Post, P.: Variability of atmospheric circulation patterns associated with large volume changes
 5484 of the Baltic Sea, *Adv. Sci. Res.*, 12, 219-225, <https://doi.org/10.5194/asr-12-219-2015>, 2015.



- 5485 Lehmann, A., Höflich, K., Post, P., and Myrberg, K.: Pathways of deep cyclones associated with large volume
 5486 changes (LVCs) and major Baltic inflows (MBIs), *J. Mar. Syst.*, 167, 11-18,
 5487 <https://doi.org/10.1016/j.jmarsys.2016.10.014>, 2017.
- 5488 Lehmann, A., Myrberg, K., Post, P., Chubarenko, I., Dailidienė, I., Hinrichsen, H.-H., Hüseyin, K., Liblik, T.,
 5489 Lips, U., Meier, H. E. M., and Bukanova, T.: Salinity dynamics of the Baltic Sea, *Earth Syst. Dynam. Discuss.*,
 5490 1-36, <https://doi.org/10.5194/esd-2021-15>, 2021.
- 5491 Lehtonen, I.: Four consecutive snow-rich winters in Southern Finland: 2009/2010–2012/2013, *Weather* 70, 3-8,
 5492 <https://doi.org/10.1002/wea.2360>, 2015.
- 5493 Lehtoranta, J., Savchuk, O. P., Elken, J., Dahlbo, K., Kuosa, H., Raateoja, M., Kauppila, P., Räike, A., and
 5494 Pitkänen, H.: Atmospheric forcing controlling inter-annual nutrient dynamics in the open Gulf of Finland, *J.*
 5495 *Mar. Syst.*, 171, 4-20, <https://doi.org/10.1016/j.jmarsys.2017.02.001>, 2017.
- 5496 Leidenberger, S., De Giovanni, R., Kulawik, R., Williams, A. R., and Bourlat, S. J.: Mapping present and future
 5497 potential distribution patterns for a meso-grazer guild in the Baltic Sea, *J. Biogeogr.*, 42, 241-254,
 5498 <https://doi.org/10.1111/jbi.12395>, 2015.
- 5499 Lenggenhager, S., and Martius, O.: Atmospheric blocks modulate the odds of heavy precipitation events in
 5500 Europe, *Clim. Dyn.*, 53, 4155-4171, <https://doi.org/10.1007/s00382-019-04779-0>, 2019.
- 5501 Leppäranta, M., and Myrberg, K.: Physical oceanography of the Baltic Sea, Springer Science & Business Media,
 5502 Berlin, 2009.
- 5503 Leppäranta, M., Oikkonen, A., Shirasawa, K., and Fukumachi, Y.: A treatise on frequency spectrum of drift ice
 5504 velocity, *Cold Reg. Sci. Technol.*, 76, 83-91, <https://doi.org/10.1016/j.coldregions.2011.12.005>, 2012.
- 5505 Leppäranta, M.: Land-ice interaction in the Baltic Sea, *Est. J. Earth Sci.*, 62, 2013.
- 5506 Lépy, É., and Pasanen, L.: Observed Regional Climate Variability during the Last 50 Years in Reindeer Herding
 5507 Cooperatives of Finnish Fell Lapland, *Climate*, 5, 81, <https://doi.org/10.3390/cli5040081>, 2017.
- 5508 Lewandowska, A. M., Breithaupt, P., Hillebrand, H., Hoppe, H.-G., Jürgens, K., and Sommer, U.: Responses of
 5509 primary productivity to increased temperature and phytoplankton diversity, *J. Sea Res.*, 72, 87-93,
 5510 <https://doi.org/10.1016/j.seares.2011.10.003>, 2012.
- 5511 Lewandowska, A. M., Boyce, D. G., Hofmann, M., Matthiessen, B., Sommer, U., and Worm, B.: Effects of sea
 5512 surface warming on marine plankton, *Ecol. Lett.*, 17, 614-623, <https://doi.org/10.1111/ele.12265>, 2014.
- 5513 Liblik, T., and Lips, U.: Stratification Has Strengthened in the Baltic Sea – An Analysis of 35 Years of
 5514 Observational Data, *Front. Earth Sci.*, 7, <https://doi.org/10.3389/feart.2019.00174>, 2019.
- 5515 Lindegren, M., Möllmann, C., Nielsen, A., Brander, K., MacKenzie, B. R., and Stenseth, N. C.: Ecological
 5516 forecasting under climate change: the case of Baltic cod, *P. Roy. Soc. B-Biol. Sci.*, 277, 2121-2130,
 5517 <https://doi.org/10.1098/rspb.2010.0353>, 2010.
- 5518 Lindegren, M., Blenckner, T., and Stenseth, N. C.: Nutrient reduction and climate change cause a potential shift
 5519 from pelagic to benthic pathways in a eutrophic marine ecosystem, *Global Change Biol.*, 18, 3491-3503,
 5520 <https://doi.org/10.1111/j.1365-2486.2012.02799.x>, 2012.
- 5521 Lindeskog, M., Lagergren, F., Smith, B., and Rammig, A.: Accounting for forest management in the estimation
 5522 of forest carbon balance using the dynamic vegetation model LPJ-GUESS (v4.0, r9333): Implementation and
 5523 evaluation of simulations for Europe, *Geosci. Model Dev. Discuss.*, 2021, 1-42, <https://doi.org/10.5194/gmd-2020-440>, 2021.
- 5525 Lindroth, A., Holst, J., Linderson, M.-L., Aurela, M., Biermann, T., Heliasz, M., Chi, J., Ibrom, A., Kolari, P.,
 5526 Klemetsson, L., Krasnova, A., Laurila, T., Lehner, I., Lohila, A., Mammarella, I., Mölder, M., Löfvenius, M.,
 5527 O., Peichl, M., Pilegaard, K., Soosaar, K., Vesala, T., Vestin, P., Weslien, P., and Nilsson, M.: Effects of drought
 5528 and meteorological forcing on carbon and water fluxes in Nordic forests during the dry summer of 2018, *Philos.*
 5529 *T. Roy. Soc. B.*, 375, 20190516, <https://doi.org/10.1098/rstb.2019.0516>, 2020.
- 5530 Lindström, G.: Hydrologiska aspekter på åtgärder mot vattenbrist och torka inom avrinningsområden02837722
 5531 (ISSN), 60pp, 2019.
- 5532 Lips, I., and Lips, U.: Abiotic factors influencing cyanobacterial bloom development in the Gulf of Finland
 5533 (Baltic Sea), *Hydrobiologia*, 614, 133-140, <https://doi.org/10.1007/s10750-008-9449-2>, 2008.



- 5534 Liu, Y., Meier, H. E. M., and Eilola, K.: Nutrient transports in the Baltic Sea – results from a 30-year physical–
 5535 biogeochemical reanalysis, *Biogeosciences*, 14, 2113–2131, <https://doi.org/10.5194/bg-14-2113-2017>, 2017.
- 5536 Liu, Y., Axell, L., Jandt, S., Lorkowski, I., Lindenthal, A., Verjovkina, S., and Schwichtenberg, F.: Baltic Sea
 5537 Production Centre BALTICSEA_REANALYSIS_PHY_003_012 COPERNICUS Marine Environment
 5538 Monitoring Service, Quality Information Document, 35pp, 2019.
- 5539 Liu, Z., Zhu, J., Rosenthal, Y., Zhang, X., Otto-Bliesner, B. L., Timmermann, A., Smith, R. S., Lohmann, G.,
 5540 Zheng, W., and Elison Timm, O.: The Holocene temperature conundrum, *Proceedings of the National Academy*
 5541 *of Sciences*, 111, E3501, <https://doi.org/10.1073/pnas.1407229111>, 2014.
- 5542 Lopez, L. S., Hewitt, B. A., and Sharma, S.: Reaching a breaking point: How is climate change influencing the
 5543 timing of ice breakup in lakes across the northern hemisphere?, *Limnol. Oceanogr.*, 64, 2621–2631,
 5544 <https://doi.org/10.1002/lno.11239>, 2019.
- 5545 Löptien, U., and Meier, H. E. M.: The influence of increasing water turbidity on the sea surface temperature in
 5546 the Baltic Sea: a model sensitivity study, *J. Mar. Syst.*, 88, 323–331,
 5547 <https://doi.org/10.1016/j.jmarsys.2011.06.001>, 2011.
- 5548 Ludwig, J.: *Climate Signals in Coastal Deposits*, Staats- und Universitätsbibliothek Hamburg, Hamburg, 2017.
- 5549 Łukawska-Matuszewska, K., and Graca, B.: Pore water alkalinity below the permanent halocline in the Gdańsk
 5550 Deep (Baltic Sea) - Concentration variability and benthic fluxes, *Mar. Chem.*, 204, 49–61,
 5551 <https://doi.org/10.1016/j.marchem.2018.05.011>, 2018.
- 5552 Lundin, E. J., Klaminder, J., Bastviken, D., Olid, C., Hansson, S. V., and Karlsson, J.: Large difference in carbon
 5553 emission – burial balances between boreal and arctic lakes, *Sci. Rep.-UK*, 5, 14248,
 5554 <https://doi.org/10.1038/srep14248>, 2015.
- 5555 Lundquist, J. K., DuVivier, K. K., Kaffine, D., and Tomaszewski, J. M.: Costs and consequences of wind turbine
 5556 wake effects arising from uncoordinated wind energy development, *Nat. Energy*, 4, 26–34,
 5557 <https://doi.org/10.1038/s41560-018-0281-2>, 2019.
- 5558 Luomaranta, A., Ruosteenoja, K., Jylhä, K., Gregow, H., Haapala, J., and Laaksonen, A.: Multimodel estimates
 5559 of the changes in the Baltic Sea ice cover during the present century, *Tellus A*, 66, 22617,
 5560 <https://doi.org/10.3402/tellusa.v66.22617>, 2014.
- 5561 Luomaranta, A., Aalto, J., and Jylhä, K.: Snow cover trends in Finland over 1961–2014 based on gridded snow
 5562 depth observations, *Int. J. Climatol.*, 39, 3147–3159, <https://doi.org/10.1002/joc.6007>, 2019.
- 5563 Luoto, M., Fronzek, S., and Zuidhoff, F. S.: Spatial modelling of palsa mires in relation to climate in northern
 5564 Europe, *Earth Surface Processes and Landforms: The Journal of the British Geomorphological Research Group*,
 5565 29, 1373–1387, <https://doi.org/10.1002/esp.1099>, 2004.
- 5566 Luterbacher, J., Werner, J. P., Smerdon, J. E., Fernández-Donado, L., González-Rouco, F. J., Barriopedro, D.,
 5567 Ljungqvist, F. C., Büntgen, U., Zorita, E., Wagner, S., Esper, J., McCarroll, D., Toreti, A., Frank, D., Jungclauss,
 5568 J. H., Barriendos, M., Bertolin, C., Bothe, O., Brázdil, R., Camuffo, D., Dobrovolný, P., Gagen, M., García-
 5569 Bustamante, E., Ge, Q., Gómez-Navarro, J. J., Guiot, J., Hao, Z., Hegerl, G. C., Holmgren, K., Klimenko, V. V.,
 5570 Martín-Chivelet, J., Pfister, C., Roberts, N., Schindler, A., Schurer, A., Solomina, O., von Gunten, L., Wahl, E.,
 5571 Wanner, H., Wetter, O., Xoplaki, E., Yuan, N., Zanchettin, D., Zhang, H., and Zerefos, C.: European summer
 5572 temperatures since Roman times, *Environ. Res. Lett.*, 11, 024001, [https://doi.org/10.1088/1748-](https://doi.org/10.1088/1748-9326/11/2/024001)
 5573 9326/11/2/024001, 2016.
- 5574 Luyssaert, S., Marie, G., Valade, A., Chen, Y.-Y., Njakou Djomo, S., Ryder, J., Otto, J., Naudts, K., Lansø, A.
 5575 S., Ghattas, J., and McGrath, M. J.: Trade-offs in using European forests to meet climate objectives, *Nature*, 562,
 5576 259–262, <https://doi.org/10.1038/s41586-018-0577-1>, 2018.
- 5577 Lyon, S. W., Destouni, G., Giesler, R., Humborg, C., Mörtz, M., Seibert, J., Karlsson, J., and Troch, P. A.:
 5578 Estimation of permafrost thawing rates in a sub-arctic catchment using recession flow analysis, *Hydrol. Earth*
 5579 *Syst. Sci.*, 13, 595–604, <https://doi.org/10.5194/hess-13-595-2009>, 2009.
- 5580 Maberly, S. C., O'Donnell, R. A., Woolway, R. I., Cutler, M. E. J., Gong, M., Jones, I. D., Merchant, C. J.,
 5581 Miller, C. A., Politi, E., Scott, E. M., Thackeray, S. J., and Tyler, A. N.: Global lake thermal regions shift under
 5582 climate change, *Nat. Commun.*, 11, 1232, <https://doi.org/10.1038/s41467-020-15108-z>, 2020.



- 5583 Macdonald, R. W., Mackay, D., Li, Y. F., and Hickie, B.: How Will Global Climate Change Affect Risks from
 5584 Long-Range Transport of Persistent Organic Pollutants?, *Hum. Ecol. Risk Assess.*, 9, 643-660,
 5585 <https://doi.org/10.1080/713609959>, 2003.
- 5586 MacKenzie, B. R., and Schiedek, D.: Daily ocean monitoring since the 1860s shows record warming of northern
 5587 European seas, *Global Change Biol.*, 13, 1335-1347, <https://doi.org/10.1111/j.1365-2486.2007.01360.x>, 2007.
- 5588 MacKenzie, B. R., Meier, H. E. M., Lindegren, M., Neuenfeldt, S., Eero, M., Blenckner, T., Tomczak, M. T.,
 5589 and Niiranen, S.: Impact of Climate Change on Fish Population Dynamics in the Baltic Sea: A Dynamical
 5590 Downscaling Investigation, *Ambio* 41, 626-636, <https://doi.org/10.1007/s13280-012-0325-y>, 2012.
- 5591 MacLean, I. M. D., Austin, G. E., Rehfish, M. M., Blew, J., Crowe, O., Delany, S., Devos, K., Deceuninck, B.,
 5592 Günther, K., Laursen, K., von Roomen, M., and Wahl, J.: Climate change causes rapid changes in the
 5593 distribution and site abundance of birds in winter, *Global Change Biol.*, 14, 2489-2500,
 5594 <https://doi.org/10.1111/j.1365-2486.2008.01666.x>, 2008.
- 5595 Magaard, L., and Rheinheimer, G.: *Meereskunde der Ostsee*, Springer-Verlag, New York, N.Y., 1974.
- 5596 Mäkinen, K., Vuorinen, I., and Hänninen, J.: Climate-induced hydrography change favours small-bodied
 5597 zooplankton in a coastal ecosystem, *Hydrobiologia*, 792, 83-96, <https://doi.org/10.1007/s10750-016-3046-6>,
 5598 2017.
- 5599 Mäll, M., Nakamura, R., Suursaar, Ü., and Shibayama, T.: Pseudo-climate modelling study on projected changes
 5600 in extreme extratropical cyclones, storm waves and surges under CMIP5 multi-model ensemble: Baltic Sea
 5601 perspective, *Nat. Hazards*, 102, 67-99, <https://doi.org/10.1007/s11069-020-03911-2>, 2020.
- 5602 Mann, M. E., Steinman, B. A., Brouillette, D. J., and Miller, S. K.: Multidecadal climate oscillations during the
 5603 past millennium driven by volcanic forcing, *Science*, 371, 1014, <https://doi.org/10.1126/science.abc5810>, 2021.
- 5604 Männikus, R., Soomere, T., and Viška, M.: Variations in the mean, seasonal and extreme water level on the
 5605 Latvian coast, the eastern Baltic Sea, during 1961–2018, *Estuar. Coast. Shelf Sci.*, 245, 106827,
 5606 <https://doi.org/10.1016/j.ecss.2020.106827>, 2020.
- 5607 Marchowski, D., Jankowiak, Ł., Wysocki, D., Ławicki, Ł., and Girjatowicz, J.: Ducks change wintering patterns
 5608 due to changing climate in the important wintering waters of the Odra River Estuary, *PeerJ*, 5, e3604,
 5609 <https://doi.org/10.7717/peerj.3604>, 2017.
- 5610 Marcos, M., Calafat, F. M., Berihuete, Á., and Dangendorf, S.: Long-term variations in global sea level
 5611 extremes, *J. Geophys. Res-Oceans*, 120, 8115-8134, <https://doi.org/10.1002/2015jc011173>, 2015.
- 5612 Marcos, M., and Woodworth, P. L.: Spatiotemporal changes in extreme sea levels along the coasts of the North
 5613 Atlantic and the Gulf of Mexico, *J. Geophys. Res-Oceans*, 122, 7031-7048,
 5614 <https://doi.org/10.1002/2017JC013065>, 2017.
- 5615 Markowicz, K. M., and Uscka-Kowalkowska, J.: Long-term and seasonal variability of the aerosol optical depth
 5616 at Mount Kasprowy Wierch (Poland), *J. Geophys. Res-Atmos.*, 120, 1865-1879,
 5617 <https://doi.org/10.1002/2014JD022580>, 2015.
- 5618 Marshall, G. J., Jylhä, K., Kivinen, S., Laapas, M., and Dyrddal, A. V.: The role of atmospheric circulation
 5619 patterns in driving recent changes in indices of extreme seasonal precipitation across Arctic Fennoscandia, *Clim.*
 5620 *Change*, 162, 741-759, <https://doi.org/10.1007/s10584-020-02747-w>, 2020.
- 5621 Marshall, J., Johnson, H., and Goodman, J.: A Study of the Interaction of the North Atlantic Oscillation with
 5622 Ocean Circulation, *J. Climate*, 14, 1399-1421, [https://doi.org/10.1175/1520-0442\(2001\)014<1399:ASOTIO>2.0.CO;2](https://doi.org/10.1175/1520-0442(2001)014<1399:ASOTIO>2.0.CO;2), 2001.
- 5624 Mastrandrea, M. D., Field, C. B., Stocker, T. F., Edenhofer, O., Ebi, K. L., Frame, D. J., Held, H., Kriegler, E.,
 5625 Mach, K. J., Matschoss, P. R., Plattne, G.-K., Yohe, G. W., and Zwiers, F. W.: Guidance Note for Lead Authors
 5626 of the IPCC Fifth Assessment Report on Consistent Treatment of Uncertainties, Intergovernmental Panel on
 5627 Climate Change (IPCC), <http://www.ipcc.ch>, 2010.
- 5628 Matthäus, W., and Franck, H.: Characteristics of major Baltic inflows—a statistical analysis, *Cont. Shelf Res.*,
 5629 12, 1375-1400, [https://doi.org/10.1016/0278-4343\(92\)90060-W](https://doi.org/10.1016/0278-4343(92)90060-W), 1992.
- 5630 Matthews, T., Murphy, C., Wilby, R. L., and Harrigan, S.: A cyclone climatology of the British-Irish Isles 1871–
 5631 2012, *Int. J. Climatol.*, 36, 1299-1312, <https://doi.org/10.1002/joc.4425>, 2016.



- 5632 Matthias, V., Arndt, J. A., Aulinger, A., Bieser, J., Denier van der Gon, H., Kranenburg, R., Kuenen, J.,
 5633 Neumann, D., Pouliot, G., and Quante, M.: Modeling emissions for three-dimensional atmospheric chemistry
 5634 transport models, *J. Air Waste Manag.*, 68, 763-800, <https://doi.org/10.1080/10962247.2018.1424057>, 2018.
- 5635 Mauri, A., Davis, B. A. S., Collins, P. M., and Kaplan, J. O.: The influence of atmospheric circulation on the
 5636 mid-Holocene climate of Europe: a data-model comparison, *Clim. Past*, 10, 1925-1938,
 5637 <https://doi.org/10.5194/cp-10-1925-2014>, 2014.
- 5638 Mauri, A., Davis, B. A. S., Collins, P. M., and Kaplan, J. O.: The climate of Europe during the Holocene: a
 5639 gridded pollen-based reconstruction and its multi-proxy evaluation, *Quaternary Sci. Rev.*, 112, 109-127,
 5640 <https://doi.org/10.1016/j.quascirev.2015.01.013>, 2015.
- 5641 McClelland, J. W., Holmes, R. M., Peterson, B. J., and Stieglitz, M.: Increasing river discharge in the Eurasian
 5642 Arctic: Consideration of dams, permafrost thaw, and fires as potential agents of change, *J. Geophys. Res.-Atmos.*,
 5643 109, <https://doi.org/10.1029/2004JD004583>, 2004.
- 5644 McCrackin, M. L., Muller-Karulis, B., Gustafsson, B. G., Howarth, R. W., Humborg, C., Svanbäck, A., and
 5645 Swaney, D. P.: A Century of Legacy Phosphorus Dynamics in a Large Drainage Basin, *Global Biogeochem.*
 5646 *Cy.*, 32, 1107-1122, <https://doi.org/10.1029/2018GB005914>, 2018.
- 5647 McGlade, J. M.: 12 The North Sea Large Marine Ecosystem, in: *Large Marine Ecosystems*, edited by: Sherman,
 5648 K., and Skjoldal, H. R., Elsevier, 339-412, [https://doi.org/10.1016/S1570-0461\(02\)80064-7](https://doi.org/10.1016/S1570-0461(02)80064-7), 2002.
- 5649 McNamara, J. M., and Houston, A. I.: State-dependent life histories, *Nature*, 380, 215-221,
 5650 <https://doi.org/10.1038/380215a0>, 1996.
- 5651 Meehl, G. A., Senior, C. A., Eyring, V., Flato, G., Lamarque, J.-F., Stouffer, R. J., Taylor, K. E., and Schlund,
 5652 M.: Context for interpreting equilibrium climate sensitivity and transient climate response from the CMIP6 Earth
 5653 system models, *Sci. Adv.*, 6, eaba1981, <https://doi.org/10.1126/sciadv.aba1981>, 2020.
- 5654 Meier, H. E. M.: Regional ocean climate simulations with a 3D ice-ocean model for the Baltic Sea. Part 2:
 5655 results for sea ice, *Clim. Dyn.*, 19, 255-266, <https://doi.org/10.1007/s00382-001-0225-5>, 2002.
- 5656 Meier, H. E. M., and Döscher, R.: Simulated water and heat cycles of the Baltic Sea using a 3D coupled
 5657 atmosphere-ice-ocean model, *Boreal Environ. Res.*, 7, 327-334, 2002.
- 5658 Meier, H. E. M., and Kauker, F.: Modeling decadal variability of the Baltic Sea: 2. Role of freshwater inflow and
 5659 large-scale atmospheric circulation for salinity, *J. Geophys. Res.-Oceans*, 108,
 5660 <https://doi.org/10.1029/2003JC001799>, 2003.
- 5661 Meier, H. E. M., Broman, B., and Kjellström, E.: Simulated sea level in past and future climates of the Baltic
 5662 Sea, *Clim. Res.*, 27, 59-75, <https://doi.org/10.3354/cr027059>, 2004a.
- 5663 Meier, H. E. M., Döscher, R., and Halkka, A.: Simulated distributions of Baltic Sea-ice in warming climate and
 5664 consequences for the winter habitat of the Baltic ringed seal, *Ambio*, 33, 249-256, <https://doi.org/10.1579/0044-7447-33.4.249>, 2004b.
- 5666 Meier, H. E. M.: Baltic Sea climate in the late twenty-first century: a dynamical downscaling approach using two
 5667 global models and two emission scenarios, *Clim. Dyn.*, 27, 39-68, <https://doi.org/10.1007/s00382-006-0124-x>,
 5668 2006.
- 5669 Meier, H. E. M., Kjellström, E., and Graham, L. P.: Estimating uncertainties of projected Baltic Sea salinity in
 5670 the late 21st century, *Geophys. Res. Lett.*, 33, L15705, <https://doi.org/10.1029/2006GL026488>, 2006.
- 5671 Meier, H. E. M., Andersson, H. C., Eilola, K., Gustafsson, B. G., Kuznetsov, I., Müller-Karulis, B., Neumann,
 5672 T., and Savchuk, O. P.: Hypoxia in future climates: A model ensemble study for the Baltic Sea, *Geophys. Res.*
 5673 *Lett.*, 38, L24608, <https://doi.org/10.1029/2011GL049929>, 2011a.
- 5674 Meier, H. E. M., Eilola, K., and Almroth, E.: Climate-related changes in marine ecosystems simulated with a 3-
 5675 dimensional coupled physical-biogeochemical model of the Baltic Sea, *Clim. Res.*, 48, 31-55,
 5676 <https://doi.org/10.3354/cr00968> 2011b.
- 5677 Meier, H. E. M., Höglund, A., Döscher, R., Andersson, H., Löptien, U., and Kjellström, E.: Quality assessment
 5678 of atmospheric surface fields over the Baltic Sea from an ensemble of regional climate model simulations with
 5679 respect to ocean dynamics, *Oceanologia*, 53, 193-227, <https://doi.org/10.5697/oc.53-1-TI.193>, 2011c.



- 5680 Meier, H. E. M., Andersson, H. C., Arheimer, B., Blenckner, T., Chubarenko, B., Donnelly, C., Eilola, K.,
 5681 Gustafsson, B. G., Hansson, A., Havenhand, J., and others: Comparing reconstructed past variations and future
 5682 projections of the Baltic Sea ecosystem—first results from multi-model ensemble simulations, *Environ. Res.*
 5683 *Letts.*, 7, 34005, <https://doi.org/10.1088/1748-9326/7/3/034005>, 2012a.
- 5684 Meier, H. E. M., Hordoir, R., Andersson, H. C., Dieterich, C., Eilola, K., Gustafsson, B. G., Höglund, A., and
 5685 Schimanke, S.: Modeling the combined impact of changing climate and changing nutrient loads on the Baltic
 5686 Sea environment in an ensemble of transient simulations for 1961–2099, *Clim. Dyn.*, 39, 2421–2441,
 5687 <https://doi.org/10.1007/s00382-012-1339-7>, 2012b.
- 5688 Meier, H. E. M., Müller-Karulis, B., Andersson, H. C., Dieterich, C., Eilola, K., Gustafsson, B. G., Höglund, A.,
 5689 Hordoir, R., Kuznetsov, I., Neumann, T., and others: Impact of climate change on ecological quality indicators
 5690 and biogeochemical fluxes in the Baltic Sea: a multi-model ensemble study, *Ambio*, 41, 558–573,
 5691 <https://doi.org/10.1007/s13280-012-0320-3>, 2012c.
- 5692 Meier, H. E. M., Rutgersson, A., and Reckermann, M.: An Earth System Science Program for the Baltic Sea
 5693 Region, *Eos Trans. AGU*, 95, 109–110, <https://doi.org/10.1002/2014EO130001>, 2014.
- 5694 Meier, H. E. M.: Projected Change—Marine Physics, in: Second Assessment of Climate Change for the Baltic
 5695 Sea Basin, edited by: BACC II Author Team, Springer International Publishing, Cham, 243–252,
 5696 https://doi.org/10.1007/978-3-319-16006-1_13, 2015.
- 5697 Meier, H. E. M., Höglund, A., Eilola, K., and Almroth-Rosell, E.: Impact of accelerated future global mean sea
 5698 level rise on hypoxia in the Baltic Sea, *Clim. Dyn.*, 49, 163–172, <https://doi.org/10.1007/s00382-016-3333-y>,
 5699 2017.
- 5700 Meier, H. E. M., Edman, M. K., Eilola, K. J., Placke, M., Neumann, T., Andersson, H. C., Brunnabend, S.-E.,
 5701 Dieterich, C., Frauen, C., Friedland, R., Gröger, M., Gustafsson, B. G., Gustafsson, E., Isaev, A., Kniebusch, M.,
 5702 Kuznetsov, I., Müller-Karulis, B., Omstedt, A., Ryabchenko, V., Saraiva, S., and Savchuk, O. P.: Assessment of
 5703 Eutrophication Abatement Scenarios for the Baltic Sea by Multi-Model Ensemble Simulations, *Front. Mar. Sci.*,
 5704 5, <https://doi.org/10.3389/fmars.2018.00440>, 2018a.
- 5705 Meier, H. E. M., Väli, G., Naumann, M., Eilola, K., and Frauen, C.: Recently accelerated oxygen consumption
 5706 rates amplify deoxygenation in the Baltic Sea, *J. Geophys. Res.-Oceans*, 123, 3227–3240,
 5707 <https://doi.org/10.1029/2017JC013686>, 2018b.
- 5708 Meier, H. E. M., Dieterich, C., Eilola, K., Gröger, M., Höglund, A., Radtke, H., Saraiva, S., and Wählström, I.:
 5709 Future projections of record-breaking sea surface temperature and cyanobacteria bloom events in the Baltic Sea,
 5710 *Ambio*, 48, 1362–1376, <https://doi.org/10.1007/s13280-019-01235-5>, 2019a.
- 5711 Meier, H. E. M., Edman, M., Eilola, K., Placke, M., Neumann, T., Andersson, H. C., Brunnabend, S.-E.,
 5712 Dieterich, C., Frauen, C., Friedland, R., Gröger, M., Gustafsson, B. G., Gustafsson, E., Isaev, A., Kniebusch, M.,
 5713 Kuznetsov, I., Müller-Karulis, B., Naumann, M., Omstedt, A., Ryabchenko, V., Saraiva, S., and Savchuk, O. P.:
 5714 Assessment of Uncertainties in Scenario Simulations of Biogeochemical Cycles in the Baltic Sea, *Front. Mar.*
 5715 *Sci.*, 6, <https://doi.org/10.3389/fmars.2019.00046>, 2019b.
- 5716 Meier, H. E. M., Eilola, K., Almroth-Rosell, E., Schimanke, S., Kniebusch, M., Höglund, A., Pemberton, P., Liu,
 5717 Y., Väli, G., and Saraiva, S.: Correction to: Disentangling the impact of nutrient load and climate changes on
 5718 Baltic Sea hypoxia and eutrophication since 1850, *Clim. Dyn.*, 53, 1167–1169, <https://doi.org/10.1007/s00382-018-4483-x>, 2019c.
- 5720 Meier, H. E. M., Eilola, K., Almroth-Rosell, E., Schimanke, S., Kniebusch, M., Höglund, A., Pemberton, P., Liu,
 5721 Y., Väli, G., and Saraiva, S.: Disentangling the impact of nutrient load and climate changes on Baltic Sea
 5722 hypoxia and eutrophication since 1850, *Clim. Dyn.*, 53, 1145–1166, <https://doi.org/10.1007/s00382-018-4296-y>,
 5723 2019d.
- 5724 Meier, H. E. M., and Saraiva, S.: Projected Oceanographical Changes in the Baltic Sea until 2100, Oxford
 5725 Research Encyclopedia of Climate Science, <https://doi.org/10.1093/acrefore/9780190228620.013.699>, 2020.
- 5726 Meier, H. E. M., Börgel, F., Bøssing-Christensen, O., Dieterich, C., Dutheil, C., Gröger, M., and Kjellström, E.:
 5727 Oceanographic regional climate projections for the Baltic Sea until 2100, *Earth Syst. Dynam.*, 2021a.
- 5728 Meier, H. E. M., Dieterich, C., and Gröger, M.: Natural variability is a large source of uncertainty in future
 5729 projections of hypoxia in the Baltic Sea, *Commun. Earth Environ.*, 2, 50, <https://doi.org/10.1038/s43247-021-00115-9>, 2021b.



- 5731 Meier, R., Schwaab, J., Seneviratne, S. I., Sprenger, M., Lewis, E., and Davin, E. L.: Empirical estimate of
 5732 forestation-induced precipitation changes in Europe, *Nat. Geosci.*, 14, 473–478, [https://doi.org/10.1038/s41561-](https://doi.org/10.1038/s41561-021-00773-6)
 5733 021-00773-6, 2021c.
- 5734 Mercer, A.: A DEM of the 2010 surface topography of Storglaciären, Sweden, *J. Maps*, 12, 1112–1118,
 5735 <https://doi.org/10.1080/17445647.2015.1131754>, 2016.
- 5736 Mercer, J. H.: West Antarctic ice sheet and CO₂ greenhouse effect: a threat of disaster, *Nature*, 271, 321–325,
 5737 1978.
- 5738 Merkouriadi, I., and Leppäranta, M.: Long-term analysis of hydrography and sea-ice data in Tvärminne, Gulf of
 5739 Finland, Baltic Sea, *Clim. Change*, 124, 849–859, <https://doi.org/10.1007/s10584-014-1130-3>, 2014.
- 5740 Merkouriadi, I., Leppäranta, M., and Järvinen, O.: Interannual variability and trends in winter weather and snow
 5741 conditions in Finnish Lapland, *Est. J. Earth Sci.*, 66, 47–57, <https://doi.org/10.3176/earth.2017.03>, 2017.
- 5742 Mikulski, Z.: Inflow from drainage basin, *Water Balance of the Baltic Sea-Baltic Sea Environment Proceedings*,
 5743 16, 24–34, Baltic Marine Environment Protection Commission, Helsinki, Finland, 1986.
- 5744 Mitrovica, J. X., Hay, C. C., Kopp, R. E., Harig, C., and Latychev, K.: Quantifying the Sensitivity of Sea Level
 5745 Change in Coastal Localities to the Geometry of Polar Ice Mass Flux, *J. Climate*, 31, 3701–3709,
 5746 <https://doi.org/10.1175/jcli-d-17-0465.1>, 2018.
- 5747 Mohrholz, V., Naumann, M., Nausch, G., Krüger, S., and Gräwe, U.: Fresh oxygen for the Baltic Sea—An
 5748 exceptional saline inflow after a decade of stagnation, *J. Mar. Syst.*, 148, 152–166,
 5749 <https://doi.org/10.1016/j.jmarsys.2015.03.005>, 2015.
- 5750 Mohrholz, V.: Major Baltic Inflow Statistics – Revised, *Front. Mar. Sci.*, 5,
 5751 <https://doi.org/10.3389/fmars.2018.00384>, 2018.
- 5752 Moldanová, J., Hasselöv, I.-M., Matthias, V., Fridell, E., Jalkanen, J.-P., Ytreberg, E., Quante, M., Tröltzsch, J.,
 5753 Maljutenko, I., Raudsepp, U., and Eriksson, K. M.: Framework for the environmental impact assessment of
 5754 operational shipping, *Ambio*, <https://doi.org/10.1007/s13280-021-01597-9>, 2021.
- 5755 Möller, K. O., Schmidt, J. O., St. John, M., Temming, A., Diekmann, R., Peters, J., Floeter, J., Sell, A. F.,
 5756 Herrmann, J.-P., and Möllmann, C.: Effects of climate-induced habitat changes on a key zooplankton species, *J.*
 5757 *Plankton Res.*, 37, 530–541, <https://doi.org/10.1093/plankt/fbv033>, 2015.
- 5758 Möllmann, C.: Effects of Climate Change and Fisheries on the Marine Ecosystem of the Baltic Sea, *Oxford*
 5759 *Research Encyclopedia of Climate Science*, <https://doi.org/10.1093/acrefore/9780190228620.013.682>, 2019.
- 5760 Müller, J., Folini, D., Wild, M., and Pfenninger, S.: CMIP-5 models project photovoltaics are a no-regrets
 5761 investment in Europe irrespective of climate change, *Energy*, 171, 135–148,
 5762 <https://doi.org/10.1016/j.energy.2018.12.139>, 2019.
- 5763 Müller, J. D., Schneider, B., and Rehder, G.: Long-term alkalinity trends in the Baltic Sea and their implications
 5764 for CO₂-induced acidification, *Limnol. Oceanogr.*, 61, 1984–2002, <https://doi.org/10.1002/lno.10349>, 2016.
- 5765 Munkes, B., Löptien, U., and Dietze, H.: Cyanobacteria blooms in the Baltic Sea: a review of models and facts,
 5766 *Biogeosciences*, 18, 2347–2378, <https://doi.org/10.5194/bg-18-2347-2021>, 2021.
- 5767 Musielak, S., Furmańczyk, K., and Bugajny, N.: Factors and Processes Forming the Polish Southern Baltic Sea
 5768 Coast on Various Temporal and Spatial Scales, in: *Coastline Changes of the Baltic Sea from South to East: Past*
 5769 *and Future Projection*, edited by: Harff, J., Furmańczyk, K., and von Storch, H., Springer International
 5770 Publishing, Cham, 69–85, https://doi.org/10.1007/978-3-319-49894-2_5, 2017.
- 5771 Myrberg, K., Korpinen, S., and Uusitalo, L.: Physical oceanography sets the scene for the Marine Strategy
 5772 Framework Directive implementation in the Baltic Sea, *Mar. Policy*, 107, 103591,
 5773 <https://doi.org/10.1016/j.marpol.2019.103591>, 2019.
- 5774 Nakamura, T., Yamazaki, K., Iwamoto, K., Honda, M., Miyoshi, Y., Ogawa, Y., and Ukita, J.: A negative phase
 5775 shift of the winter AO/NAO due to the recent Arctic sea-ice reduction in late autumn, *J. Geophys. Res.-Atmos.*,
 5776 120, 3209–3227, <https://doi.org/10.1002/2014JD022848>, 2015.
- 5777 Nakicenovic, N., Alcamo, J., Grubler, A., Riahi, K., Roehrl, R. A., Rogner, H.-H., and Victor, N.: Special report
 5778 on emissions scenarios (SRES), a special report of Working Group III of the intergovernmental panel on climate
 5779 change, IPCC Special Report, 2000.



- 5780 Naumann, M., Gräwe, U., Mohrholz, V., Kuss, J., Siegel, H., Waniek, J. J., and Schulz-Bull, D. E.:
 5781 Hydrographic-hydrochemical assessment of the Baltic Sea 2018, *Meereswiss. Ber.*, Warnemünde, 110,
 5782 <https://doi.io-warnemuende.de/10.12754/msr-2019-0110>, 2019.
- 5783 Nerem, R. S., Beckley, B. D., Fasullo, J. T., Hamlington, B. D., Masters, D., and Mitchum, G. T.: Climate-
 5784 change-driven accelerated sea-level rise detected in the altimeter era, *Proceedings of the National Academy of*
 5785 *Sciences*, 115, 2022, <https://doi.org/10.1073/pnas.1717312115>, 2018.
- 5786 Neumann, G.: *Eigenschwingungen der Ostsee*, Archiv der Deutschen Seewarte und des Marineobservatoriums,
 5787 Hamburg, 1941.
- 5788 Neumann, T., Eilola, K., Gustafsson, B., Müller-Karulis, B., Kuznetsov, I., Meier, H. E. M., and Savchuk, O. P.:
 5789 Extremes of temperature, oxygen and blooms in the Baltic Sea in a changing climate, *Ambio*, 41, 574-585,
 5790 <https://doi.org/10.1007/s13280-012-0321-2>, 2012.
- 5791 Neumann, T., Radtke, H., and Seifert, T.: On the importance of Major Baltic Inflows for oxygenation of the
 5792 central Baltic Sea, *J. Geophys. Res-Oceans*, 122, 1090-1101, <https://doi.org/10.1002/2016JC012525>, 2017.
- 5793 Niemelä, P., Tolvanen, H., Rönkä, M., Kellomäki, S., Krug, J., Schurgers, G., Lehikoinen, E., and Kalliola, R.:
 5794 Environmental Impacts—Coastal Ecosystems, Birds and Forests, in: *Second Assessment of Climate Change for*
 5795 *the Baltic Sea Basin*, edited by: BACC II Author Team, Springer International Publishing, Cham, 291-306,
 5796 https://doi.org/10.1007/978-3-319-16006-1_16, 2015.
- 5797 Niiranen, S., Yletyinen, J., Tomczak, M. T., Blenckner, T., Hjerne, O., MacKenzie, B. R., Müller-Karulis, B.,
 5798 Neumann, T., and Meier, H. E. M.: Combined effects of global climate change and regional ecosystem drivers
 5799 on an exploited marine food web, *Global Change Biol.*, 19, 3327-3342, <https://doi.org/10.1111/gcb.12309>, 2013.
- 5800 Nijse, F. J. M. M., Cox, P. M., Huntingford, C., and Williamson, M. S.: Decadal global temperature variability
 5801 increases strongly with climate sensitivity, *Nat. Clim. Change*, 9, 598-601, [https://doi.org/10.1038/s41558-019-](https://doi.org/10.1038/s41558-019-0527-4)
 5802 [0527-4](https://doi.org/10.1038/s41558-019-0527-4), 2019.
- 5803 Nilsson, J., and Grennfelt, P.: *Critical loads for sulphur and nitrogen*, Report from Skokloster Workshop.
 5804 Skokloster. Sweden, 1988.
- 5805 Nilsson, L., and Haas, F.: Distribution and numbers of wintering waterbirds in Sweden in 2015 and changes
 5806 during the last fifty years, *Ornis Svecica*, 26, 3–54-53–54, 2016.
- 5807 Ning, W., Nielsen, A. B., Ivarsson, L. N., Jilbert, T., Åkesson, C. M., Slomp, C. P., Andrén, E., Broström, A.,
 5808 and Filipsson, H. L.: Anthropogenic and climatic impacts on a coastal environment in the Baltic Sea over the last
 5809 1000 years, *Anthropocene*, 21, 66-79, <https://doi.org/10.1016/j.ancene.2018.02.003>, 2018.
- 5810 Niskanen, T., Vainio, J., Eriksson, P., and Heiler, I.: Maximum extent of the Baltic sea ice recalculated for the
 5811 period 1971-2008, *Report Series in Geophysics*, 164, 2009.
- 5812 Nöges, P., and Nöges, T.: Weak trends in ice phenology of Estonian large lakes despite significant warming
 5813 trends, *Hydrobiologia*, 731, 5-18, <https://doi.org/10.1007/s10750-013-1572-z>, 2014.
- 5814 Norbäck Ivarsson, L., Andrén, T., Moros, M., Andersen, T. J., Lönn, M., and Andrén, E.: Baltic Sea Coastal
 5815 Eutrophication in a Thousand Year Perspective, *Front. Environ. Sci.*, 7,
 5816 <https://doi.org/10.3389/fenvs.2019.00088>, 2019.
- 5817 Nydahl, A., Panigrahi, S., and Wikner, J.: Increased microbial activity in a warmer and wetter climate enhances
 5818 the risk of coastal hypoxia, *FEMS Microbiol. Ecol.*, 85, 338-347, <https://doi.org/10.1111/1574-6941.12123>,
 5819 2013.
- 5820 O'Reilly, C. M., Sharma, S., Gray, D. K., Hampton, S. E., Read, J. S., Rowley, R. J., Schneider, P., Lenters, J.
 5821 D., McIntyre, P. B., Kraemer, B. M., Weyhenmeyer, G. A., Straile, D., Dong, B., Adrian, R., Allan, M. G.,
 5822 Anneville, O., Arvola, L., Austin, J., Bailey, J. L., Baron, J. S., Brookes, J. D., de Eyto, E., Dokulil, M. T.,
 5823 Hamilton, D. P., Havens, K., Hetherington, A. L., Higgins, S. N., Hook, S., Izmet'eva, L. R., Joehnk, K. D.,
 5824 Kangur, K., Kasprzak, P., Kumagai, M., Kuusisto, E., Leshkevich, G., Livingstone, D. M., MacIntyre, S., May,
 5825 L., Melack, J. M., Mueller-Navarra, D. C., Naumenko, M., Nöges, P., Nöges, T., North, R. P., Plisnier, P.-D.,
 5826 Rigosi, A., Rimmer, A., Rogora, M., Rudstam, L. G., Rusak, J. A., Salmaso, N., Samal, N. R., Schindler, D. E.,
 5827 Schladow, S. G., Schmid, M., Schmidt, S. R., Silow, E., Soylu, M. E., Teubner, K., Verburg, P., Voutilainen, A.,
 5828 Watkinson, A., Williamson, C. E., and Zhang, G.: Rapid and highly variable warming of lake surface waters
 5829 around the globe, *Geophys. Res. Lett.*, 42, 10,773-710,781, <https://doi.org/10.1002/2015GL066235>, 2015.



- Obu, J., Westermann, S., Bartsch, A., Berdnikov, N., Christiansen, H. H., Dashtseren, A., Delaloye, R., Elberling, B., Etzelmüller, B., Kholodov, A., Khomutov, A., Kääb, A., Leibman, M. O., Lewkowicz, A. G., Panda, S. K., Romanovsky, V., Way, R. G., Westergaard-Nielsen, A., Wu, T., Yamkhin, J., and Zou, D.: Northern Hemisphere permafrost map based on TTOP modelling for 2000–2016 at 1 km² scale, *Earth Sci. Rev.*, 193, 299–316, <https://doi.org/10.1016/j.earscirev.2019.04.023>, 2019.
- Obu, J., Westermann, S., Barboux, C., Bartsch, A., Delaloye, R., Grosse, G., Heim, B., Hugelius, G., Irrgang, A., Kääb, A. M., Kroisleitner, C., Matthes, H., Nitze, I., Pellet, C., Seifert, F. M., Strozzi, T., Wegmüller, U., Wieczorek, M., and Wiesmann, A.: ESA Permafrost Climate Change Initiative (Permafrost_cci): Permafrost extent for the Northern Hemisphere, v2.0., Centre for Environmental Data Analysis, <http://dx.doi.org/10.5285/28e889210f884b469d7168fde4b4e54f>, 2020.
- Ojaveer, H., Olenin, S., Naršcius, A., Florin, A.-B., Ezhova, E., Gollasch, S., Jensen, K. R., Lehtiniemi, M., Minchin, D., Normant-Saremba, M., and Sträke, S.: Dynamics of biological invasions and pathways over time: a case study of a temperate coastal sea, *Biol. Invasions*, 19, 799–813, <https://doi.org/10.1007/s10530-016-1316-x>, 2017.
- Olli, K., Klais, R., Tamminen, T., Ptacnik, R., and Andersen, T.: Long term changes in the Baltic Sea phytoplankton community, *Boreal Environ. Res.*, 16 (suppl. A), 3–14, 2011.
- Olofsson, M., Torstensson, A., Karlberg, M., Steinhoff, F. S., Dinasquet, J., Riemann, L., Chierici, M., and Wulff, A.: Limited response of a spring bloom community inoculated with filamentous cyanobacteria to elevated temperature and pCO₂, *Bot. Mar.*, 62, 3–16, <https://doi.org/10.1515/bot-2018-0005>, 2019.
- Olofsson, M., Suikkanen, S., Kobos, J., Wasmund, N., and Karlson, B.: Basin-specific changes in filamentous cyanobacteria community composition across four decades in the Baltic Sea, *Harmful Algae*, 91, 101685, <https://doi.org/10.1016/j.hal.2019.101685>, 2020.
- Omstedt, A., and Chen, D.: Influence of atmospheric circulation on the maximum ice extent in the Baltic Sea, *J. Geophys. Res.-Oceans*, 106, 4493–4500, <https://doi.org/10.1029/1999JC000173>, 2001.
- Omstedt, A., Edman, M., Claremar, B., Frodin, P., Gustafsson, E., Humborg, C., Hägg, H., Mörh, M., Rutgersson, A., Schurgers, G., and others: Future changes in the Baltic Sea acid–base (pH) and oxygen balances, *Tellus B*, 64, 19586, <https://doi.org/10.3402/tellusb.v64i0.19586>, 2012.
- Omstedt, A., Elken, J., Lehmann, A., Leppäranta, M., Meier, H. E. M., Myrberg, K., and Rutgersson, A.: Progress in physical oceanography of the Baltic Sea during the 2003–2014 period, *Prog. Oceanogr.*, 128, 139–171, <https://doi.org/10.1016/j.pocean.2014.08.010>, 2014.
- Oppenheimer, M., Glavovic, B. C., Hinkel, J., Wal, R. v. d., Magnan, A. K., Abd-Elgawad, A., Cai, R., Cifuentes-Jara, M., DeConto, R. M., Ghosh, T., Hay, J., Isla, F., Marzeion, B., Meyssignac, B., and Sebesvari, Z.: Sea Level Rise and Implications for Low-Lying Islands, Coasts and Communities, in: *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*, edited by: Pörtner, H.-O., Roberts, D. C., Masson-Delmotte, V., Zhai, P., Tignor, M., Poloczanska, E., Mintenbeck, K., Alegría, A., Nicolai, M., Okem, A., Petzold, J., Rama, B., and Weyer, N. M., 2019.
- Orru, H., Andersson, C., Ebi, K. L., Langner, J., Åström, C., and Forsberg, B.: Impact of climate change on ozone-related mortality and morbidity in Europe, *Eur. Respir. J.*, 41, 285–294, <https://doi.org/10.1183/09031936.00210411>, 2013.
- Orru, H., Åström, C., Andersson, C., Tamm, T., Ebi, K. L., and Forsberg, B.: Ozone and heat-related mortality in Europe in 2050 significantly affected by changes in climate, population and greenhouse gas emission, *Environ. Res. Lett.*, 14, 074013, 2019.
- Osterkamp, T., and Romanovsky, V.: Evidence for warming and thawing of discontinuous permafrost in Alaska, *Permafrost Periglac.*, 10, 17–37, [https://doi.org/10.1002/\(SICI\)1099-1530\(199901/03\)10:1<17::AID-PPP303>3.0.CO;2-4](https://doi.org/10.1002/(SICI)1099-1530(199901/03)10:1<17::AID-PPP303>3.0.CO;2-4), 1999.
- Otto, L., Zimmerman, J. T. F., Furnes, G. K., Mork, M., Saetre, R., and Becker, G.: Review of the physical oceanography of the North Sea, *Neth. J. Sea Res.*, 26, 161–238, [https://doi.org/10.1016/0077-7579\(90\)90091-T](https://doi.org/10.1016/0077-7579(90)90091-T), 1990.
- Øygarden, L., Deelstra, J., Lagzdins, A., Bechmann, M., Greipsland, I., Kyllmar, K., Povilaitis, A., and Iital, A.: Climate change and the potential effects on runoff and nitrogen losses in the Nordic–Baltic region, *Agric. Ecosyst. Environ.*, 198, 114–126, <https://doi.org/10.1016/j.agee.2014.06.025>, 2014.



- 5881 Paczkowska, J., Brugel, S., Rowe, O., Lefébure, R., Brutemark, A., and Andersson, A.: Response of Coastal
 5882 Phytoplankton to High Inflows of Terrestrial Matter, *Front. Mar. Sci.*, 7,
 5883 <https://doi.org/10.3389/fmars.2020.00080>, 2020.
- 5884 Pajusalu, L., Martin, G., Pöllumäe, A., and Paalme, T.: Results of laboratory and field experiments of the direct
 5885 effect of increasing CO₂ on net primary production of macroalgal species in brackish-water ecosystems, *P. Est.*
 5886 *Acad. Sci.*, 62, 148, <https://doi.org/10.3176/proc.2013.2.09>, 2013.
- 5887 Pajusalu, L., Martin, G., Pöllumäe, A., Torn, K., and Paalme, T.: Direct effects of increased CO₂ concentrations
 5888 in seawater on the net primary production of charophytes in a shallow, coastal, brackish-water ecosystem, *Boreal*
 5889 *Environ. Res.*, 20, 413–422, 2015.
- 5890 Pajusalu, L., Martin, G., Paalme, T., and Pöllumäe, A.: The effect of CO₂ enrichment on net photosynthesis of
 5891 the red alga *Furcellaria lumbricalis* in a brackish water environment, *PeerJ*, 4, e2505,
 5892 <https://doi.org/10.7717/peerj.2505>, 2016.
- 5893 Pansch, C., Nasrolahi, A., Appelhans, Y. S., and Wahl, M.: Impacts of ocean warming and acidification on the
 5894 larval development of the barnacle *Amphibalanus improvisus*, *J. Exp. Mar. Biol. Ecol.*, 420–421, 48–55,
 5895 <https://doi.org/10.1016/j.jembe.2012.03.023>, 2012.
- 5896 Parding, K., Olseth, J. A., Dagestad, K. F., and Liepert, B. G.: Decadal variability of clouds, solar radiation and
 5897 temperature at a high-latitude coastal site in Norway, *Tellus B*, 66, 25897,
 5898 <https://doi.org/10.3402/tellusb.v66.25897>, 2014.
- 5899 Paul, C., Matthiessen, B., and Sommer, U.: Warming, but not enhanced CO₂ concentration, quantitatively and
 5900 qualitatively affects phytoplankton biomass, *Mar. Ecol. Prog. Ser.*, 528, 39–51,
 5901 <https://doi.org/10.3354/meps11264> 2015.
- 5902 Pavón-Jordán, D., Fox, A. D., Clausen, P., Dagys, M., Deceuninck, B., Devos, K., Hearn, R. D., Holt, C. A.,
 5903 Hornman, M., Keller, V., Langendoen, T., Ławicki, Ł., Lorentsen, S. H., Luigujõe, L., Meissner, W., Musil, P.,
 5904 Nilsson, L., Paquet, J.-Y., Stipniece, A., Stroud, D. A., Wahl, J., Zenatello, M., and Lehtikainen, A.: Climate-
 5905 driven changes in winter abundance of a migratory waterbird in relation to EU protected areas, *Divers. Distrib.*,
 5906 21, 571–582, <https://doi.org/10.1111/ddi.12300>, 2015.
- 5907 Pavón-Jordán, D., Clausen, P., Dagys, M., Devos, K., Encarnação, V., Fox, A. D., Frost, T., Gaudard, C.,
 5908 Hornman, M., Keller, V., Langendoen, T., Ławicki, Ł., Lewis, L. J., Lorentsen, S.-H., Luigujõe, L., Meissner,
 5909 W., Molina, B., Musil, P., Musilova, Z., Nilsson, L., Paquet, J.-Y., Ridzon, J., Stipniece, A., Teufelbauer, N.,
 5910 Wahl, J., Zenatello, M., and Lehtikainen, A.: Habitat- and species-mediated short- and long-term distributional
 5911 changes in waterbird abundance linked to variation in European winter weather, *Divers. Distrib.*, 25, 225–239,
 5912 <https://doi.org/10.1111/ddi.12855>, 2019.
- 5913 Pecuchet, L., Lindegren, M., Kortsch, S., Calkiewicz, J., Jurgensone, I., Margonski, P., Otto, S. A., Putnis, I.,
 5914 Stråke, S., and Nordström, M. C.: Spatio-temporal dynamics of multi-trophic communities reveal ecosystem-
 5915 wide functional reorganization, *Ecography*, 43, 197–208, <https://doi.org/10.1111/ecog.04643>, 2020.
- 5916 Peings, Y., and Magnusdottir, G.: Forcing of the wintertime atmospheric circulation by the multidecadal
 5917 fluctuations of the North Atlantic ocean, *Environ. Res. Lett.*, 9, 034018, <https://doi.org/10.1088/1748-9326/9/3/034018>, 2014.
- 5919 Peings, Y., and Magnusdottir, G.: Wintertime atmospheric response to Atlantic multidecadal variability: effect of
 5920 stratospheric representation and ocean–atmosphere coupling, *Clim. Dyn.*, 47, 1029–1047,
 5921 <https://doi.org/10.1007/s00382-015-2887-4>, 2016.
- 5922 Pekcan-Hekim, Z., Urho, L., Auvinen, H., Heikinheimo, O., Lappalainen, J., Raitaniemi, J., and Söderkultalahti,
 5923 P.: Climate Warming and Pikeperch Year-Class Catches in the Baltic Sea, *Ambio*, 40, 447–456,
 5924 <https://doi.org/10.1007/s13280-011-0143-7>, 2011.
- 5925 Pellikka, H., Särkkä, J., Johansson, M., and Pettersson, H.: Probability distributions for mean sea level and storm
 5926 contribution up to 2100 AD at Forsmark, Swedish Nuclear Fuel and Waste Management Company (SKB), 49pp,
 5927 2020.
- 5928 Pemberton, P., Löptien, U., Hordoir, R., Höglund, A., Schimanke, S., Axell, L., and Haapala, J.: Sea-ice
 5929 evaluation of NEMO-Nordic 1.0: a NEMO–LIM3. 6-based ocean–sea-ice model setup for the North Sea and
 5930 Baltic Sea, *Geosci. Model Dev.*, 10, 3105, <https://doi.org/10.5194/gmd-10-3105-2017>, 2017.



- 5931 Peng, S., Piao, S., Ciais, P., Friedlingstein, P., Zhou, L., and Wang, T.: Change in snow phenology and its
 5932 potential feedback to temperature in the Northern Hemisphere over the last three decades, *Environ. Res. Lett.*, 8,
 5933 014008, <https://doi.org/10.1088/1748-9326/8/1/014008>, 2013.
- 5934 Perry, D., Staveley, T., Deyanova, D., Baden, S., Dupont, S., Hernroth, B., Wood, H., Björk, M., and Gullström,
 5935 M.: Global environmental changes negatively impact temperate seagrass ecosystems, *Ecosphere*, 10, e02986,
 5936 <https://doi.org/10.1002/ecs2.2986>, 2019.
- 5937 Persson, T.: Solar radiation climate in Sweden, *Physics and Chemistry of the Earth, Part B: Hydrology, Oceans*
 5938 *and Atmosphere*, 24, 275-279, [https://doi.org/10.1016/S1464-1909\(98\)00050-1](https://doi.org/10.1016/S1464-1909(98)00050-1), 1999.
- 5939 Peters, W., Bastos, A., Ciais, P., and Vermeulen, A.: A historical, geographical and ecological perspective on the
 5940 2018 European summer drought, *Philos. T. Roy. Soc. B.*, 375, 20190505, <https://doi.org/10.1098/rstb.2019.0505>,
 5941 2020.
- 5942 Pfeffer, W. T., Arendt, A. A., Bliss, A., Bolch, T., Cogley, J. G., Gardner, A. S., Hagen, J.-O., Hock, R., Kaser,
 5943 G., Kienholz, C., Miles, E. S., Moholdt, G., Mölg, N., Paul, F., Radić, V., Rastner, P., Raup, B. H., Rich, J., and
 5944 Sharp, M. J.: The Randolph Glacier Inventory: a globally complete inventory of glaciers, *J. Glaciol.*, 60, 537-
 5945 552, <https://doi.org/10.3189/2014JG13J176>, 2014.
- 5946 Pfeifroth, U., Sanchez-Lorenzo, A., Manara, V., Trentmann, J., and Hollmann, R.: Trends and Variability of
 5947 Surface Solar Radiation in Europe Based On Surface- and Satellite-Based Data Records, *J. Geophys. Res-*
 5948 *Atmos.*, 123, 1735-1754, <https://doi.org/10.1002/2017JD027418>, 2018.
- 5949 Philipona, R., Behrens, K., and Ruckstuhl, C.: How declining aerosols and rising greenhouse gases forced rapid
 5950 warming in Europe since the 1980s, *Geophys. Res. Lett.*, 36, <https://doi.org/10.1029/2008GL036350>, 2009.
- 5951 Pihlainen, S., Zandersen, M., Hyytiäinen, K., Andersen, H. E., Bartosova, A., Gustafsson, B., Jabloun, M.,
 5952 McCrackin, M., Meier, H. E. M., Olesen, J. E., Saraiva, S., Swaney, D., and Thodsen, H.: Impacts of changing
 5953 society and climate on nutrient loading to the Baltic Sea, *Sci. Total Environ.*, 138935,
 5954 <https://doi.org/10.1016/j.scitotenv.2020.138935>, 2020.
- 5955 Pindsoo, K., and Soomere, T.: Basin-wide variations in trends in water level maxima in the Baltic Sea, *Cont.*
 5956 *Shelf Res.*, 193, 104029, <https://doi.org/10.1016/j.csr.2019.104029>, 2020.
- 5957 Placke, M., Meier, H. E. M., Gräwe, U., Neumann, T., Frauen, C., and Liu, Y.: Long-Term Mean Circulation of
 5958 the Baltic Sea as Represented by Various Ocean Circulation Models, *Front. Mar. Sci.*, 5,
 5959 <https://doi.org/10.3389/fmars.2018.00287>, 2018.
- 5960 Placke, M., Meier, H. E. M., and Neumann, T.: Sensitivity of the Baltic Sea Overturning Circulation to Long-
 5961 Term Atmospheric and Hydrological Changes, *J. Geophys. Res-Oceans*, 126, e2020JC016079,
 5962 <https://doi.org/10.1029/2020JC016079>, 2021.
- 5963 Porz, L., Zhang, W., and Schrum, C.: Density-driven bottom currents control development of muddy basins in
 5964 the southwestern Baltic Sea, *Mar. Geol.*, 438, 106523, <https://doi.org/10.1016/j.margeo.2021.106523>, 2021.
- 5965 Proietti, C., Fornasier, M. F., Sicard, P., Anav, A., Paoletti, E., and De Marco, A.: Trends in tropospheric ozone
 5966 concentrations and forest impact metrics in Europe over the time period 2000–2014, *J. For. Res.*, 32, 543-551,
 5967 <https://doi.org/10.1007/s11676-020-01226-3>, 2021.
- 5968 Ptak, M., Sojka, M., and Nowak, B.: Effect of climate warming on a change in thermal and ice conditions in the
 5969 largest lake in Poland – Lake Śniardwy, *J. Hydrol. Hydromech.*, 68, 260-270, <https://doi.org/10.2478/johh-2020-0024>, 2020.
- 5971 Pugh, T. A. M., Lindeskog, M., Smith, B., Poulter, B., Arneth, A., Haverd, V., and Calle, L.: Role of forest
 5972 regrowth in global carbon sink dynamics, *P. Natl. Acad. Sci. USA*, 116, 4382,
 5973 <https://doi.org/10.1073/pnas.1810512116>, 2019.
- 5974 Qixiang, W., Wang, M., and Fan, X.: Seasonal patterns of warming amplification of high-elevation stations
 5975 across the globe, *Int. J. Climatol.*, 38, 3466-3473, <https://doi.org/10.1002/joc.5509>, 2018.
- 5976 Quante, M., and Colijn, F.: North Sea region climate change assessment, *Regional Climate Studies*,
 5977 SpringerOpen, Cham, 2016.
- 5978 Räämet, A., and Soomere, T.: The wave climate and its seasonal variability in the northeastern Baltic Sea, *Est. J.*
 5979 *Earth Sci.*, 59, <https://doi.org/10.3176/earth.2010.1.08>, 2010.



- 5980 Rädler, A. T., Groenemeijer, P. H., Faust, E., Sausen, R., and Púčik, T.: Frequency of severe thunderstorms
 5981 across Europe expected to increase in the 21st century due to rising instability, *npj Clim. Atmos. Sci.*, 2, 30,
 5982 <https://doi.org/10.1038/s41612-019-0083-7>, 2019.
- 5983 Radtke, H., Brunnabend, S. E., Gräwe, U., and Meier, H. E. M.: Investigating interdecadal salinity changes in the
 5984 Baltic Sea in a 1850–2008 hindcast simulation, *Clim. Past*, 16, 1617–1642, [https://doi.org/10.5194/cp-16-1617-](https://doi.org/10.5194/cp-16-1617-2020)
 5985 2020, 2020.
- 5986 Rainio, K., Laaksonen, T., Ahola, M., Vähätalo, A. V., and Lehikoinen, E.: Climatic responses in spring
 5987 migration of boreal and arctic birds in relation to wintering area and taxonomy, *J. Avian Biol.*, 37, 507–515,
 5988 <https://doi.org/10.1111/j.0908-8857.2006.03740.x>, 2006.
- 5989 Räisänen, J.: Twenty-first century changes in snowfall climate in Northern Europe in ENSEMBLES regional
 5990 climate models, *Clim. Dyn.*, 46, 339–353, <https://doi.org/10.1007/s00382-015-2587-0>, 2016.
- 5991 Räisänen, J.: Effect of atmospheric circulation on recent temperature changes in Finland, *Clim. Dyn.*, 53, 5675–
 5992 5687, <https://doi.org/10.1007/s00382-019-04890-2>, 2019.
- 5993 Rajasilta, M., Hänninen, J., Laaksonen, L., Laine, P., Suomela, J.-P., Vuorinen, I., and Mäkinen, K.: Influence of
 5994 environmental conditions, population density, and prey type on the lipid content in Baltic herring (*Clupea*
 5995 *harengus membras*) from the northern Baltic Sea, *Can. J. Fish. Aquat. Sci.*, 76, 576–585,
 5996 <https://doi.org/10.1139/cjfas-2017-0504>, 2018.
- 5997 Rajczak, J., and Schär, C.: Projections of Future Precipitation Extremes Over Europe: A Multimodel Assessment
 5998 of Climate Simulations, *J. Geophys. Res.-Atmos.*, 122, 10,773–710,800, <https://doi.org/10.1002/2017JD027176>,
 5999 2017.
- 6000 Ramacher, M. O. P., Karl, M., Bieser, J., Jalkanen, J.-P., and Johansson, L.: Urban population exposure to
 6001 NO₂ emissions from local shipping in three Baltic Sea harbour cities - a generic approach, *Atmos. Chem.*
 6002 *Phys.*, 19, 9153, 2019.
- 6003 Ravestein, P., van der Schrier, G., Haarsma, R., Scheele, R., and van den Broek, M.: Vulnerability of European
 6004 intermittent renewable energy supply to climate change and climate variability, *Renew. Sust. Energ. Rev.*, 97,
 6005 497–508, <https://doi.org/10.1016/j.rser.2018.08.057>, 2018.
- 6006 Read, A. J., and Hohn, A. A.: LIFE IN THE FAST LANE: THE LIFE HISTORY OF HARBOR PORPOISES
 6007 FROM THE GULF OF MAINE, *Mar. Mammal Sci.*, 11, 423–440, [https://doi.org/10.1111/j.1748-](https://doi.org/10.1111/j.1748-7692.1995.tb00667.x)
 6008 7692.1995.tb00667.x, 1995.
- 6009 Reckermann, M., Langner, J., Omstedt, A., von Storch, H., Keevallik, S., Schneider, B., Arheimer, B., Meier, H.
 6010 E. M., and Hünicke, B.: BALTTEX—an interdisciplinary research network for the Baltic Sea region, *Environ.*
 6011 *Res. Lett.*, 6, 045205, <http://doi.org/10.1088/1748-9326/6/4/045205>, 2011.
- 6012 Reckermann, M., Omstedt, A., Soomere, T., Aigars, J., Akhtar, N., Beldowski, J., Brauer, C. P.-S. d., Cronin,
 6013 T., Czub, M., Eero, M., Hyttiäinen, K., Jalkanen, J.-P., Kiessling, A., Kjellström, E., Larsén, X. G., McCrackin,
 6014 M., Meier, H. E. M., Oberbeckmann, S., Parnell, K., Poska, A., Saarinen, J., Szymczycha, B., Undeman, E.,
 6015 Viitasalo, M., Wörman, A., and Zorita, E.: Human impacts and their interactions in the Baltic Sea region, *Earth*
 6016 *Syst. Dynam. Discuss.*, 1–127, <https://doi.org/10.5194/esd-2021-54>, 2021.
- 6017 Reder, S., Lydersen, C., Arnold, W., and Kovacs, K. M.: Haulout behaviour of High Arctic harbour seals (*Phoca*
 6018 *vitulina vitulina*) in Svalbard, Norway, *Polar Biol.*, 27, 6–16, <https://doi.org/10.1007/s00300-003-0557-1>, 2003.
- 6019 Refsgaard, J. C., Hansen, A. L., Højberg, A. L., Olesen, J. E., Hashemi, F., Wachniew, P., Wörman, A.,
 6020 Bartosova, A., Steljes, N., and Chubarenko, B.: Spatially differentiated regulation: Can it save the Baltic Sea
 6021 from excessive N-loads?, *Ambio*, 48, 1278–1289, <https://doi.org/10.1007/s13280-019-01195-w>, 2019.
- 6022 Reihan, A., Koltsova, T., Kriaciuniene, J., Lizuma, L., and Meilutyte-Barauskiene, D.: Changes in water
 6023 discharges of the Baltic states rivers in the 20th century and its relation to climate change, *Hydrol. Res.*, 38, 401–
 6024 412, <https://doi.org/10.2166/nh.2007.020>, 2007.
- 6025 Reissmann, J. H., Burchard, H., Feistel, R., Hagen, E., Lass, H. U., Mohrholz, V., Nausch, G., Umlauf, L., and
 6026 Wiczorek, G.: Vertical mixing in the Baltic Sea and consequences for eutrophication—A review, *Prog.*
 6027 *Oceanogr.*, 82, 47–80, <https://doi.org/10.1016/j.pocean.2007.10.004>, 2009.
- 6028 Rennert, K. J., and Wallace, J. M.: Cross-Frequency Coupling, Skewness, and Blocking in the Northern
 6029 Hemisphere Winter Circulation, *J. Climate*, 22, 5650–5666, <https://doi.org/10.1175/2009jcli2669.1>, 2009.



- 6030 Reusch, T. B. H., Dierking, J., Andersson, H. C., Bonsdorff, E., Carstensen, J., Casini, M., Czajkowski, M.,
 6031 Hasler, B., Hinsby, K., Hyytiäinen, K., Johannesson, K., Jomaa, S., Jormalainen, V., Kuosa, H., Kurland, S.,
 6032 Laikre, L., MacKenzie, B. R., Margonski, P., Melzner, F., Oesterwind, D., Ojaveer, H., Refsgaard, J. C.,
 6033 Sandström, A., Schwarz, G., Tonderski, K., Winder, M., and Zandersen, M.: The Baltic Sea as a time machine
 6034 for the future coastal ocean, *Sci. Adv.*, 4, <https://doi.org/10.1126/sciadv.aar8195> 2018.
- 6035 RGI Consortium: Randolph Glacier Inventory – A Dataset of Global Glacier Outlines: Version 6.0: Technical
 6036 Report, Global Land Ice Measurements from Space, Colorado, USA. Digital Media, [https://doi.org/10.7265/N5-](https://doi.org/10.7265/N5-RGI-60)
 6037 RGI-60, 2017.
- 6038 Riahi, K., Rao, S., Krey, V., Cho, C., Chirkov, V., Fischer, G., Kindermann, G., Nakicenovic, N., and Rafaj, P.:
 6039 RCP 8.5—A scenario of comparatively high greenhouse gas emissions, *Clim. Change*, 109, 33,
 6040 <https://doi.org/10.1007/s10584-011-0149-y>, 2011.
- 6041 Ribeiro, A., Barbosa, S. M., Scotto, M. G., and Donner, R. V.: Changes in extreme sea-levels in the Baltic Sea,
 6042 *Tellus A*, 66, 20921, <https://doi.org/10.3402/tellusa.v66.20921>, 2014.
- 6043 Ridefelt, H., Etzelmüller, B., Boelhouwers, J., and Jonasson, C.: Statistic-empirical modelling of mountain
 6044 permafrost distribution in the Abisko region, sub-Arctic northern Sweden, *Norsk Geografisk Tidsskrift-*
 6045 *Norwegian Journal of Geography*, 62, 278-289, <https://doi.org/10.1080/00291950802517890>, 2008.
- 6046 Riihelä, A., Carlund, T., Trentmann, J., Müller, R., and Lindfors, A. V.: Validation of CM SAF Surface Solar
 6047 Radiation Datasets over Finland and Sweden, *Remote Sens.-Basel*, 7, 6663-6682,
 6048 <https://doi.org/10.3390/rs70606663>, 2015.
- 6049 Rimkus, E., Kažys, J., Valiukas, D., and Stankūnavičius, G.: The atmospheric circulation patterns during dry
 6050 periods in Lithuania, *Oceanologia*, 56, 223-239, <https://doi.org/10.5697/oc.56-2.223>, 2014.
- 6051 Rimkus, E., Briede, A., Jaagus, J., Stonevicius, E., Kilpys, J., and Viru, B.: Snow-cover regime in Lithuania,
 6052 Latvia and Estonia and its relationship to climatic and geographical factors in 1961–2015, *Boreal Environ. Res.*,
 6053 23, 193-208, 2018.
- 6054 Rinne, H., and Salovius-Laurén, S.: The status of brown macroalgae *Fucus* spp. and its relation to environmental
 6055 variation in the Finnish marine area, northern Baltic Sea, *Ambio*, 49, 118-129, [https://doi.org/10.1007/s13280-](https://doi.org/10.1007/s13280-019-01175-0)
 6056 019-01175-0, 2020.
- 6057 Rinne, J., Tuovinen, J.-P., Klemetsson, L., Aurela, M., Holst, J., Lohila, A., Weslien, P., Vestin, P., Łakomiec,
 6058 P., Peichl, M., Tuittila, E.-S., Heiskanen, L., Laurila, T., Li, X., Alekseychik, P., Mammarella, I., Ström, L.,
 6059 Crill, P., and Nilsson, M. B.: Effect of the 2018 European drought on methane and carbon dioxide exchange of
 6060 northern mire ecosystems, *Philos. T. Roy. Soc. B.*, 375, 20190517, <https://doi.org/10.1098/rstb.2019.0517>, 2020.
- 6061 Ripszám, M., Paczkowska, J., Figueira, J., Veenaas, C., and Haglund, P.: Dissolved Organic Carbon Quality and
 6062 Sorption of Organic Pollutants in the Baltic Sea in Light of Future Climate Change, *Environ. Sci. Technol.*, 49,
 6063 1445-1452, <https://doi.org/10.1021/es504437s>, 2015.
- 6064 Rizzi, J., Nilsen, I. B., Stagge, J. H., Gislås, K., and Tallaksen, L. M.: Five decades of warming: impacts on
 6065 snow cover in Norway, *Hydrol. Res.*, 49, 670-688, <https://doi.org/10.2166/nh.2017.051>, 2017.
- 6066 Rjazin, J., and Pärn, O.: Determining the Regime Shift of the Baltic Sea Ice Seasons during 1982–2016, *NAŠE*
 6067 *MORE: znanstveni časopis za more i pomorstvo*, 67, 53-59, 2020.
- 6068 Rodhe, J.: The Baltic and North Seas: A process-oriented review of the physical oceanography. Coastal segment.
 6069 , in: *The Sea, Volume 11: The Global Coastal Ocean: Regional Studies and Syntheses*, edited by: Robinson, A.
 6070 R., and Brink, K. H., John Wiley & Sons, Inc., 699-732, 1998.
- 6071 Rodhe, J., Tett, P., and Wulff, F.: The Baltic and North Seas: A Regional Review of some important Physical-
 6072 Chemical-Biological Interaction Processes, in: *The Sea, Volume 14B: The Global Coastal Ocean*, edited by:
 6073 Robinson, A. R., and Brink, K. H., Harvard University Press, Cambridge, MA, USA, 1033-1075, 2006.
- 6074 Rodwell, M. J., Rowell, D. P., and Folland, C. K.: Oceanic forcing of the wintertime North Atlantic Oscillation
 6075 and European climate, *Nature*, 398, 320-323, <https://doi.org/10.1038/18648>, 1999.
- 6076 Roff, J. C., and Legendre, L.: Chapter 14 Physico-Chemical and Biological Oceanography of Hudson Bay, in:
 6077 *Elsevier Oceanography Series*, edited by: Martini, I. P., Elsevier, 265-292, [https://doi.org/10.1016/S0422-](https://doi.org/10.1016/S0422-9894(08)70907-3)
 6078 9894(08)70907-3, 1986.



- 6079 Röhr, M. E., Boström, C., Canal-Vergés, P., and Holmer, M.: Blue carbon stocks in Baltic Sea eelgrass (*Zostera*
 6080 marina) meadows, *Biogeosciences*, 13, 6139-6153, <https://doi.org/10.5194/bg-13-6139-2016>, 2016.
- 6081 Rolff, C., Almesjö, L., and Elmgren, R.: Nitrogen fixation and abundance of the diazotrophic cyanobacterium
 6082 *Aphanizomenon* sp. in the Baltic Proper, *Mar. Ecol. Prog. Ser.*, 332, 107-118,
 6083 <https://doi.org/10.3354/meps332107>, 2007.
- 6084 Rolff, C., and Elfving, T.: Increasing nitrogen limitation in the Bothnian Sea, potentially caused by inflow of
 6085 phosphate-rich water from the Baltic Proper, *Ambio*, 44, 601-611, <https://doi.org/10.1007/s13280-015-0675-3>,
 6086 2015.
- 6087 Ronkainen, I., Lehtiranta, J., Lensu, M., Rinne, E., Haapala, J., and Haas, C.: Interannual sea ice thickness
 6088 variability in the Bay of Bothnia, *Cryosphere*, 12, 3459-3476, <https://doi.org/10.5194/tc-12-3459-2018>, 2018.
- 6089 Rothäusler, E., Rugiu, L., and Jormalainen, V.: Forecast climate change conditions sustain growth and
 6090 physiology but hamper reproduction in range-margin populations of a foundation rockweed species, *Mar.*
 6091 *Environ. Res.*, 141, 205-213, <https://doi.org/10.1016/j.marenvres.2018.09.014>, 2018.
- 6092 Roudier, P., Andersson, J. C. M., Donnelly, C., Feyen, L., Greuell, W., and Ludwig, F.: Projections of future
 6093 floods and hydrological droughts in Europe under a +2°C global warming, *Clim. Change*, 135, 341-355,
 6094 <https://doi.org/10.1007/s10584-015-1570-4>, 2016.
- 6095 Rounsevell, M. D. A., Ewert, F., Reginster, I., Leemans, R., and Carter, T. R.: Future scenarios of European
 6096 agricultural land use: II. Projecting changes in cropland and grassland, *Agric. Ecosyst. Environ.*, 107, 117-135,
 6097 <https://doi.org/10.1016/j.agee.2004.12.002>, 2005.
- 6098 Rousi, H., Korpinen, S., and Bonsdorff, E.: Brackish-Water Benthic Fauna Under Fluctuating Environmental
 6099 Conditions: The Role of Eutrophication, Hypoxia, and Global Change, *Front. Mar. Sci.*, 6,
 6100 <https://doi.org/10.3389/fmars.2019.00464>, 2019.
- 6101 Ruckstuhl, C., Philipona, R., Behrens, K., Collaud Coen, M., Dürr, B., Heimo, A., Mätzler, C., Nyeki, S.,
 6102 Ohmura, A., Vuilleumier, L., Weller, M., Wehrli, C., and Zelenka, A.: Aerosol and cloud effects on solar
 6103 brightening and the recent rapid warming, *Geophys. Res. Lett.*, 35, L12708,
 6104 <https://doi.org/10.1029/2008GL034228>, 2008.
- 6105 Rugiu, L., Manninen, I., Sjöroos, J., and Jormalainen, V.: Variations in tolerance to climate change in a key
 6106 littoral herbivore, *Mar. Biol.*, 165, 18, <https://doi.org/10.1007/s00227-017-3275-x>, 2017.
- 6107 Ruoho-Airola, T., Eilola, K., Savchuk, O. P., Parviainen, M., and Tarvainen, V.: Atmospheric Nutrient Input to
 6108 the Baltic Sea from 1850 to 2006: A Reconstruction from Modeling Results and Historical Data, *Ambio* 41, 549-
 6109 557, <https://doi.org/10.1007/s13280-012-0319-9>, 2012.
- 6110 Ruosteenoja, K., Vihma, T., and Venäläinen, A.: Projected Changes in European and North Atlantic Seasonal
 6111 Wind Climate Derived from CMIP5 Simulations, *J. Climate*, 32, 6467-6490, <https://doi.org/10.1175/JCLI-D-19-0023.1>, 2019.
- 6113 Ruosteenoja, K., Markkanen, T., and Räisänen, J.: Thermal seasons in northern Europe in projected future
 6114 climate, *Int. J. Climatol.*, 40, 4444-4462, <https://doi.org/10.1002/joc.6466>, 2020.
- 6115 Ruprich-Robert, Y., Msadek, R., Castruccio, F., Yeager, S., Delworth, T., and Danabasoglu, G.: Assessing the
 6116 Climate Impacts of the Observed Atlantic Multidecadal Variability Using the GFDL CM2.1 and NCAR CESM1
 6117 Global Coupled Models, *J. Climate*, 30, 2785-2810, <https://doi.org/10.1175/jcli-d-16-0127.1>, 2017.
- 6118 Russak, V.: Changes in solar radiation and their influence on temperature trend in Estonia (1955–2007), *J.*
 6119 *Geophys. Res.-Atmos.*, 114, <https://doi.org/10.1029/2008JD010613>, 2009.
- 6120 Rutgersson, A., Jaagus, J., Schenk, F., and Stendel, M.: Observed changes and variability of atmospheric
 6121 parameters in the Baltic Sea region during the last 200 years, *Clim. Res.*, 61, 177-190,
 6122 <https://doi.org/10.3354/cr01244>, 2014.
- 6123 Rutgersson, A., Kjellström, E., Haapala, J., Stendel, M., Danilovich, I., Drews, M., Jylhä, K., Kujala, P., Larsén,
 6124 X. G., Halsnæs, K., Lehtonen, I., Luomaranta, A., Nilsson, E., Olsson, T., Särkkä, J., Tuomi, L., and Wasmund,
 6125 N.: Natural Hazards and Extreme Events in the Baltic Sea region, *Earth Syst. Dynam. Discuss.*, 1-80,
 6126 <https://doi.org/10.5194/esd-2021-13>, 2021.



- 6127 Ryabchenko, V. A., Karlin, L. N., Isaev, A. V., Vankevich, R. E., Eremina, T. R., Molchanov, M. S., and
 6128 Savchuk, O. P.: Model estimates of the eutrophication of the Baltic Sea in the contemporary and future climate,
 6129 *Oceanology+*, 56, 36-45, <https://doi.org/10.1134/S0001437016010161>, 2016.
- 6130 Ryabchuk, D., Kolesov, A., Chubarenko, B., Spiridonov, M., Kurennoy, D., and Soomere, T.: Coastal erosion
 6131 processes in the eastern Gulf of Finland and their links with geological and hydrometeorological factors, *Boreal*
 6132 *Environ. Res.*, 16 (Suppl. A), 117-137, 2011.
- 6133 Sahla, M., Tolvanen, H., Ruuskanen, A., and Kurvinen, L.: Assessing long term change of *Fucus* spp.
 6134 communities in the northern Baltic Sea using monitoring data and spatial modeling, *Estuar. Coast. Shelf Sci.*,
 6135 245, 107023, <https://doi.org/10.1016/j.ecss.2020.107023>, 2020.
- 6136 Salo, T., Mattila, J., and Eklöf, J.: Long-term warming affects ecosystem functioning through species turnover
 6137 and intraspecific trait variation, *Oikos*, 129, 283-295, <https://doi.org/10.1111/oik.06698>, 2020.
- 6138 Sannel, A. B. K., Hugelius, G., Jansson, P., and Kuhry, P.: Permafrost Warming in a Subarctic Peatland – Which
 6139 Meteorological Controls are Most Important?, *Permafrost Periglac.*, 27, 177-188,
 6140 <https://doi.org/10.1002/ppp.1862>, 2016.
- 6141 Saraiva, S., Meier, H. E. M., Andersson, H., Höglund, A., Dieterich, C., Gröger, M., Hordoir, R., and Eilola, K.:
 6142 Baltic Sea ecosystem response to various nutrient load scenarios in present and future climates, *Clim. Dyn.*, 52,
 6143 3369-3387, <https://doi.org/10.1007/s00382-018-4330-0>, 2019a.
- 6144 Saraiva, S., Meier, H. E. M., Andersson, H., Höglund, A., Dieterich, C., Gröger, M., Hordoir, R., and Eilola, K.:
 6145 Uncertainties in Projections of the Baltic Sea Ecosystem Driven by an Ensemble of Global Climate Models,
 6146 *Front. Earth Sci.*, 6, <https://doi.org/10.3389/feart.2018.00244>, 2019b.
- 6147 Sarauskienė, D., Kriauciūnienė, J., Reihan, A., and Klavins, M.: Flood pattern changes in the rivers of the Baltic
 6148 countries, *J. Environ. Eng. Landsc. Manag.*, 23, 28-38, <https://doi.org/10.3846/16486897.2014.937438>, 2015.
- 6149 Šarauskienė, D., Akstinas, V., Kriauciūnienė, J., Jakimavičius, D., Bukantis, A., Kažys, J., Povilaitis, A., Ložys,
 6150 L., Kesminas, V., Virbickas, T., and Pliuraitė, V.: Projection of Lithuanian river runoff, temperature and their
 6151 extremes under climate change, *Hydrol. Res.*, 49, 344-362, <https://doi.org/10.2166/nh.2017.007>, 2017.
- 6152 Savchuk, O. P., Wulff, F., Hille, S., Humborg, C., and Pollehne, F.: The Baltic Sea a century ago — a
 6153 reconstruction from model simulations, verified by observations, *J. Mar. Syst.*, 74, 485-494,
 6154 <https://doi.org/10.1016/j.jmarsys.2008.03.008>, 2008.
- 6155 Savchuk, O. P., Gustafsson, B. G., Rodríguez Medina, M., Sokolov, A. V., and Wulff, F.: External nutrient loads
 6156 to the Baltic Sea, 1970-2006, *Baltic Nest Institute Techn. Rep.*, 22, 2012.
- 6157 Savchuk, O. P.: Large-Scale Nutrient Dynamics in the Baltic Sea, 1970-2016, *Front. Mar. Sci.*, 5, 95,
 6158 <https://doi.org/10.3389/fmars.2018.00095>, 2018.
- 6159 Scheff, J., and Frierson, D. M. W.: Robust future precipitation declines in CMIP5 largely reflect the poleward
 6160 expansion of model subtropical dry zones, *Geophys. Res. Lett.*, 39, <https://doi.org/10.1029/2012GL052910>,
 6161 2012.
- 6162 Schenk, F., and Zorita, E.: Reconstruction of high resolution atmospheric fields for Northern Europe using
 6163 analog-upscaling, *Clim. Past Discuss.*, 8, 819-868, <https://doi.org/10.5194/cp-8-1681-2012>, 2012.
- 6164 Schenk, F.: The analog-method as statistical upscaling tool for meteorological field reconstructions over
 6165 Northern Europe since 1850, *Doctoral Dissertation, University of Hamburg, Hamburg*, 2015.
- 6166 Schimanke, S., Meier, H. E. M., Kjellström, E., Strandberg, G., and Hordoir, R.: The climate in the Baltic Sea
 6167 region during the last millennium simulated with a regional climate model, *Clim. Past*, 8, 1419,
 6168 <https://doi.org/10.5194/cp-8-1419-2012>, 2012.
- 6169 Schimanke, S., Dieterich, C., and Meier, H. E. M.: An algorithm based on sea-level pressure fluctuations to
 6170 identify major Baltic inflow events, *Tellus A*, 66, 23452, <http://dx.doi.org/10.3402/tellusa.v66.23452>, 2014.
- 6171 Schimanke, S., and Meier, H. E. M.: Decadal-to-Centennial Variability of Salinity in the Baltic Sea, *J. Climate*,
 6172 29, 7173-7188, <https://doi.org/10.1175/JCLI-D-15-0443.1>, 2016.
- 6173 Schinke, H., and Matthäus, W.: On the causes of major Baltic inflows - an analysis of long time series, *Cont.*
 6174 *Shelf Res.*, 18, 67-97, [https://doi.org/10.1016/S0278-4343\(97\)00071-X](https://doi.org/10.1016/S0278-4343(97)00071-X), 1998.



- 6175 Schmidt, G. A., Jungelaus, J. H., Ammann, C. M., Bard, E., Braconnot, P., Crowley, T. J., Delaygue, G., Joos,
 6176 F., Krivova, N. A., Muscheler, R., Otto-Bliesner, B. L., Pongratz, J., Shindell, D. T., Solanki, S. K., Steinhilber,
 6177 F., and Vieira, L. E. A.: Climate forcing reconstructions for use in PMIP simulations of the last millennium
 6178 (v1.0), *Geosci. Model Dev.*, 4, 33-45, <https://doi.org/10.5194/gmd-4-33-2011>, 2011.
- 6179 Schmidt, K., Birchill, A. J., Atkinson, A., Brewin, R. J. W., Clark, J. R., Hickman, A. E., Johns, D. G., Lohan,
 6180 M. C., Milne, A., Pardo, S., Polimene, L., Smyth, T. J., Tarran, G. A., Widdicombe, C. E., Woodward, E. M. S.,
 6181 and Ussher, S. J.: Increasing picocyanobacteria success in shelf waters contributes to long-term food web
 6182 degradation, *Global Change Biol.*, 26, 5574-5587, <https://doi.org/10.1111/gcb.15161>, 2020.
- 6183 Schneider, B., and Müller, J. D.: Biogeochemical Transformations in the Baltic Sea. Observations Through
 6184 Carbon Dioxide Glasses. , 110, Springer International Publishing, Cham, 2018.
- 6185 Schönhofer, J., and Dudkowska, A.: Rip currents in the southern Baltic Sea multi-bar nearshore zone, *Cont.*
 6186 *Shelf Res.*, 212, 104324, <https://doi.org/10.1016/j.csr.2020.104324>, 2021.
- 6187 Schröder, W., Nickel, S., Schönrock, S., Meyer, M., Wosniok, W., Harmens, H., Frontasyeva, M. V., Alber, R.,
 6188 Aleksiyenak, J., Barandovski, L., Carballeira, A., Danielsson, H., de Temmermann, L., Godzik, B., Jeran, Z.,
 6189 Karlsson, G. P., Lazo, P., Leblond, S., Lindroos, A.-J., Liiv, S., Magnússon, S. H., Mankovska, B., Martínez-
 6190 Abaigar, J., Piispanen, J., Poikolainen, J., Popescu, I. V., Qarri, F., Santamaria, J. M., Skudnik, M., Špirić, Z.,
 6191 Stafilov, T., Steinnes, E., Stihl, C., Thöni, L., Uggerud, H. T., and Zechmeister, H. G.: Spatially valid data of
 6192 atmospheric deposition of heavy metals and nitrogen derived by moss surveys for pollution risk assessments of
 6193 ecosystems, *Environ. Sci. Pollut. Res.*, 23, 10457-10476, <https://doi.org/10.1007/s11356-016-6577-5>, 2016.
- 6194 Schubert, S. D., Wang, H., Koster, R. D., Suarez, M. J., and Groisman, P. Y.: Northern Eurasian Heat Waves
 6195 and Droughts, *J. Climate*, 27, 3169-3207, <https://doi.org/10.1175/jcli-d-13-00360.1>, 2014.
- 6196 Schuster, P. F., Schaefer, K. M., Aiken, G. R., Antweiler, R. C., Dewild, J. F., Gryziec, J. D., Gusmeroli, A.,
 6197 Hugelius, G., Jafarov, E., Krabbenhoft, D. P., Liu, L., Herman-Mercer, N., Mu, C., Roth, D. A., Schaefer, T.,
 6198 Striegl, R. G., Wickland, K. P., and Zhang, T.: Permafrost Stores a Globally Significant Amount of Mercury,
 6199 *Geophys. Res. Lett.*, 45, 1463-1471, <https://doi.org/10.1002/2017GL075571>, 2018.
- 6200 Screen, J. A., Simmonds, I., Deser, C., and Tomas, R.: The Atmospheric Response to Three Decades of
 6201 Observed Arctic Sea Ice Loss, *J. Climate*, 26, 1230-1248, <https://doi.org/10.1175/jcli-d-12-00063.1>, 2013.
- 6202 Seager, R., Naik, N., and Vecchi, G. A.: Thermodynamic and Dynamic Mechanisms for Large-Scale Changes in
 6203 the Hydrological Cycle in Response to Global Warming*, *J. Climate*, 23, 4651-4668,
 6204 <https://doi.org/10.1175/2010JCLI3655.1>, 2010.
- 6205 Seinä, A., and Palosuo, E.: The classification of the maximum annual extent of ice cover in the Baltic Sea 1720–
 6206 1995, *Meri*, 27, 79-91, 1996.
- 6207 Seneviratne, S. I., Wilhelm, M., Stanelle, T., van den Hurk, B., Hagemann, S., Berg, A., Cheruy, F., Higgins, M.
 6208 E., Meier, A., Brovkin, V., Claussen, M., Ducharne, A., Dufresne, J.-L., Findell, K. L., Ghattas, J., Lawrence, D.
 6209 M., Malyshev, S., Rummukainen, M., and Smith, B.: Impact of soil moisture-climate feedbacks on CMIP5
 6210 projections: First results from the GLACE-CMIP5 experiment, *Geophys. Res. Lett.*, 40, 5212-5217,
 6211 <https://doi.org/10.1002/grl.50956>, 2013.
- 6212 Seppälä, M.: The origin of palsas, *Geogr. Ann. A*, 68, 141-147, <https://doi.org/10.2307/521453>, 1986.
- 6213 Sharma, S., Magnuson, J. J., Batt, R. D., Winslow, L. A., Korhonen, J., and Aono, Y.: Direct observations of ice
 6214 seasonality reveal changes in climate over the past 320–570 years, *Sci. Rep.-UK*, 6, 25061,
 6215 <https://doi.org/10.1038/srep25061>, 2016.
- 6216 Sharma, S., Blagrove, K., Magnuson, J. J., O'Reilly, C. M., Oliver, S., Batt, R. D., Magee, M. R., Straile, D.,
 6217 Weyhenmeyer, G. A., Winslow, L., and Woolway, R. I.: Widespread loss of lake ice around the Northern
 6218 Hemisphere in a warming world, *Nat. Clim. Change*, 9, 227-231, <https://doi.org/10.1038/s41558-018-0393-5>,
 6219 2019.
- 6220 Sharma, S., Meyer, M. F., Culpepper, J., Yang, X., Hampton, S., Berger, S. A., Brouil, M. R., Fradkin, S. C.,
 6221 Higgins, S. N., Jankowski, K. J., Kirillin, G., Smits, A. P., Whitaker, E. C., Yousef, F., and Zhang, S.:
 6222 Integrating Perspectives to Understand Lake Ice Dynamics in a Changing World, *J. Geophys. Res.-Biogeo*, 125,
 6223 e2020JG005799, <https://doi.org/10.1029/2020JG005799>, 2020.



- 6224 Sharma, S., Blagrove, K., Filazzola, A., Imrit, M. A., and Hendricks Franssen, H.-J.: Forecasting the Permanent
 6225 Loss of Lake Ice in the Northern Hemisphere Within the 21st Century, *Geophys. Res. Lett.*, 48,
 6226 e2020GL091108, <https://doi.org/10.1029/2020GL091108>, 2021.
- 6227 Shatwell, T., Thiery, W., and Kirillin, G.: Future projections of temperature and mixing regime of European
 6228 temperate lakes, *Hydrol. Earth Syst. Sci.*, 23, 1533-1551, <https://doi.org/10.5194/hess-23-1533-2019>, 2019.
- 6229 Shaw, T. A., Baldwin, M., Barnes, E. A., Caballero, R., Garfinkel, C. I., Hwang, Y. T., Li, C., O'Gorman, P. A.,
 6230 Rivière, G., Simpson, I. R., and Voigt, A.: Storm track processes and the opposing influences of climate change,
 6231 *Nat. Geosci.*, 9, 656-664, <https://doi.org/10.1038/ngeo2783>, 2016.
- 6232 Shepherd, T. G., Boyd, E., Calel, R. A., Chapman, S. C., Dessai, S., Dima-West, I. M., Fowler, H. J., James, R.,
 6233 Maraun, D., Martius, O., Senior, C. A., Sobel, A. H., Stainforth, D. A., Tett, S. F. B., Trenberth, K. E., van den
 6234 Hurk, B. J. J. M., Watkins, N. W., Wilby, R. L., and Zenghelis, D. A.: Storylines: an alternative approach to
 6235 representing uncertainty in physical aspects of climate change, *Clim. Change*, 151, 555-571,
 6236 <https://doi.org/10.1007/s10584-018-2317-9>, 2018.
- 6237 Siegel, H., and Gerth, M.: Sea Surface Temperature in the Baltic Sea 2018, HELCOM Baltic Sea Environment
 6238 Fact Sheets 2019, 7pp, [https://helcom.fi/wp-content/uploads/2020/07/BSEFS-Sea-Surface-Temperature-in-the-](https://helcom.fi/wp-content/uploads/2020/07/BSEFS-Sea-Surface-Temperature-in-the-Baltic-Sea-2018.pdf)
 6239 [Baltic-Sea-2018.pdf](https://helcom.fi/wp-content/uploads/2020/07/BSEFS-Sea-Surface-Temperature-in-the-Baltic-Sea-2018.pdf), 2019.
- 6240 Sillmann, J., Kharin, V. V., Zwiers, F. W., Zhang, X., and Bronaugh, D.: Climate extremes indices in the CMIP5
 6241 multimodel ensemble: Part 2. Future climate projections, 118, 2473-2493, <https://doi.org/10.1002/jgrd.50188>,
 6242 2013.
- 6243 Sinclair, V. A., Rantanen, M., Haapanala, P., Räisänen, J., and Järvinen, H.: The characteristics and structure of
 6244 extra-tropical cyclones in a warmer climate, *Weather Clim. Dynam.*, 1, 1-25, [10.5194/wcd-1-1-2020](https://doi.org/10.5194/wcd-1-1-2020), 2020.
- 6245 Sinha, E., Michalak, A. M., and Balaji, V.: Eutrophication will increase during the 21st century as a result of
 6246 precipitation changes, *Science*, 357, 405, <https://doi.org/10.1126/science.aan2409>, 2017.
- 6247 Skogen, M. D., Eilola, K., Hansen, J. L. S., Meier, H. E. M., Molchanov, M. S., and Ryabchenko, V. A.:
 6248 Eutrophication status of the North Sea, Skagerrak, Kattegat and the Baltic Sea in present and future climates: A
 6249 model study, *J. Mar. Syst.*, 132, 174-184, <https://doi.org/10.1016/j.jmarsys.2014.02.004>, 2014.
- 6250 Skov, H., Heinänen, S., Žydelis R., Bellebaum J., Bzoma S., Dagys M., Durinck J., Garthe S., Grishanov G., Hario M,
 6251 Kieckbusch JJ, Kube J, Kuresoo A, Larsson K, Luigujoe L, Meissner W, Nehls HW, Nilsson L, Petersen IK,
 6252 Roos MM, Pihl S, Sonntag N, Stock A, and A, S.: Waterbird populations and pressures in the Baltic Sea, Nordic
 6253 Council of Ministers, Copenhagen, 2011.
- 6254 Smith, B., Prentice, I. C., and Sykes, M. T.: Representation of Vegetation Dynamics in the Modelling of
 6255 Terrestrial Ecosystems: Comparing Two Contrasting Approaches within European Climate Space, *Global Ecol.*
 6256 *Biogeogr.*, 10, 621-637, 2001.
- 6257 Smith, B., Aasa, A., Ahas, R., Blenckner, T., Callaghan, T. V., de Chazal, J., Humborg, C., Jönsson, A. M.,
 6258 Kellomäki, S., Kull, A., Lehtikainen, E., Mander, Ü., Nöges, P., Nöges, T., Rounsevell, M., Sofiev, M.,
 6259 Tryjanowski, P., and Wolf, A.: Climate-related Change in Terrestrial and Freshwater Ecosystems, in:
 6260 Assessment of Climate Change for the Baltic Sea Basin, edited by: BACC Author Team, Springer Berlin
 6261 Heidelberg, Berlin, Heidelberg, 221-308, https://doi.org/10.1007/978-3-540-72786-6_4, 2008.
- 6262 Smith, B., Samuelsson, P., Wramneby, A., and Rummukainen, M.: A model of the coupled dynamics of climate,
 6263 vegetation and terrestrial ecosystem biogeochemistry for regional applications, *Tellus A*, 63, 87-106,
 6264 <https://doi.org/10.1111/j.1600-0870.2010.00477.x>, 2011.
- 6265 Smith, T. G., and Stirling, I.: The breeding habitat of the ringed seal (*Phoca hispida*). The birth lair and
 6266 associated structures, *Can. J. Zool.*, 53, 1297-1305, <https://doi.org/10.1139/z75-155>, 1975.
- 6267 Smol, J. P., Wolfe, A. P., Birks, H. J. B., Douglas, M. S. V., Jones, V. J., Korhola, A., Pienitz, R., Rühland, K.,
 6268 Sorvari, S., Antoniades, D., Brooks, S. J., Fallu, M.-A., Hughes, M., Keatley, B. E., Laing, T. E., Michelutti, N.,
 6269 Nazarova, L., Nyman, M., Paterson, A. M., Perren, B., Quinlan, R., Rautio, M., Saulnier-Talbot, É., Siitonen, S.,
 6270 Solovieva, N., and Weckström, J.: Climate-driven regime shifts in the biological communities of arctic lakes, *P.*
 6271 *Natl. Acad. Sci. USA*, 102, 4397, <https://doi.org/10.1073/pnas.0500245102>, 2005.
- 6272 Snickars, M., Weigel, B., and Bonsdorff, E.: Impact of eutrophication and climate change on fish and
 6273 zoobenthos in coastal waters of the Baltic Sea, *Mar. Biol.*, 162, 141-151, <https://doi.org/10.1007/s00227-014-2579-3>, 2015.



- 6275 Sommer, U., and Lewandowska, A.: Climate change and the phytoplankton spring bloom: warming and
 6276 overwintering zooplankton have similar effects on phytoplankton, *Global Change Biol.*, 17, 154-162,
 6277 <https://doi.org/10.1111/j.1365-2486.2010.02182.x>, 2011.
- 6278 Sommer, U., Aberle, N., Lengfellner, K., and Lewandowska, A.: The Baltic Sea spring phytoplankton bloom in
 6279 a changing climate: an experimental approach, *Mar. Biol.*, 159, 2479-2490, [https://doi.org/10.1007/s00227-012-](https://doi.org/10.1007/s00227-012-1897-6)
 6280 1897-6, 2012.
- 6281 Sommer, U., Paul, C., and Moustaka-Gouni, M.: Warming and Ocean Acidification Effects on Phytoplankton—
 6282 From Species Shifts to Size Shifts within Species in a Mesocosm Experiment, *PLoS ONE*, 10, e0125239,
 6283 <https://doi.org/10.1371/journal.pone.0125239>, 2015.
- 6284 Soomere, T., Behrens, A., Tuomi, L., and Nielsen, J. W.: Wave conditions in the Baltic Proper and in the Gulf of
 6285 Finland during windstorm Gudrun, *Natural Hazards and Earth System Sciences*, 8, 37-46,
 6286 <https://doi.org/10.5194/nhess-8-37-2008>, 2008.
- 6287 Soomere, T., and Räämet, A.: Spatial patterns of the wave climate in the Baltic Proper and the Gulf of
 6288 Finland**This study was supported by the Estonian Science Foundation (grant No. 7413), targeted financing by
 6289 the Estonian Ministry of Education and Research (grants SF0140077s08 and SF0140007s11) and partially
 6290 performed within the framework of the BalticWay project, which is supported by funding from the European
 6291 Community's Seventh Framework Programme (FP/2007-2013) under grant agreement No. 217246 made with
 6292 the joint Baltic Sea research and development programme BONUS, *Oceanologia*, 53, 335-371,
 6293 <https://doi.org/10.5697/oc.53-1-TI.335>, 2011.
- 6294 Soomere, T., Viška, M., and Pindsoo, K.: Retrieving the Signal of Climate Change from Numerically Simulated
 6295 Sediment Transport Along the Eastern Baltic Sea Coast, in: *Coastline Changes of the Baltic Sea from South to*
 6296 *East: Past and Future Projection*, edited by: Harff, J., Furmańczyk, K., and von Storch, H., Springer International
 6297 Publishing, Cham, 327-361, https://doi.org/10.1007/978-3-319-49894-2_15, 2017.
- 6298 Sousa, P. M., Trigo, R. M., Barriopedro, D., Soares, P. M. M., Ramos, A. M., and Liberato, M. L. R.: Responses
 6299 of European precipitation distributions and regimes to different blocking locations, *Clim. Dyn.*, 48, 1141-1160,
 6300 <https://doi.org/10.1007/s00382-016-3132-5>, 2017.
- 6301 Stahl, K., Hisdal, H., Hannaford, J., Tallaksen, L. M., van Lanen, H. A. J., Sauquet, E., Demuth, S., Fendekova,
 6302 M., and Jódar, J.: Streamflow trends in Europe: evidence from a dataset of near-natural catchments, *Hydrol.*
 6303 *Earth Syst. Sci.*, 14, 2367-2382, <https://doi.org/10.5194/hess-14-2367-2010>, 2010.
- 6304 Stålnacke, P., Grimvall, A., Sundblad, K., and Tonderski, A.: Estimation of riverine loads of nitrogen and
 6305 phosphorus to the Baltic Sea, 1970–1993, *Environ. Monit. Assess.*, 58, 173-200,
 6306 <https://doi.org/10.1023/a:1006073015871>, 1999.
- 6307 Stanhill, G., Achiman, O., Rosa, R., and Cohen, S.: The cause of solar dimming and brightening at the Earth's
 6308 surface during the last half century: Evidence from measurements of sunshine duration, *J. Geophys. Res.-Atmos.*,
 6309 119, 902-910, 911, <https://doi.org/10.1002/2013JD021308>, 2014.
- 6310 Stendel, M., Francis, J., White, R., Williams, P. D., and Woollings, T.: Chapter 15 - The jet stream and climate
 6311 change, in: *Climate Change (Third Edition)*, edited by: Letcher, T. M., Elsevier, 327-357,
 6312 <https://doi.org/10.1016/B978-0-12-821575-3.00015-3>, 2021.
- 6313 Stenseth, N. C., Payne, M. R., Bonsdorff, E., Dankel, D. J., Durant, J. M., Anderson, L. G., Armstrong, C. W.,
 6314 Blenckner, T., Brakstad, A., Dupont, S., Eikeset, A. M., Goksøyr, A., Jónsson, S., Kuparinen, A., Våge, K.,
 6315 Österblom, H., and Paasche, Ø.: Attuning to a changing ocean, *P. Natl. Acad. Sci. USA*, 117, 20363,
 6316 <https://doi.org/10.1073/pnas.1915352117>, 2020.
- 6317 Stephenson, D. B., Pavan, V., and Bojariu, R.: Is the North Atlantic Oscillation a random walk?, *Int. J. Climatol.*,
 6318 20, 1-18, [https://doi.org/10.1002/\(SICI\)1097-0088\(200001\)20:1<1::AID-JOC456>3.0.CO;2-P](https://doi.org/10.1002/(SICI)1097-0088(200001)20:1<1::AID-JOC456>3.0.CO;2-P), 2000.
- 6319 Stjern, C. W., Kristjánsson, J. E., and Hansen, A. W.: Global dimming and global brightening—an analysis of
 6320 surface radiation and cloud cover data in northern Europe, *Int. J. Climatol.*, 29, 643-653,
 6321 <https://doi.org/10.1002/joc.1735>, 2009.
- 6322 Stockholm University - Department of Physical Geography: Kebnekaise's south peak still highest in Sweden:
 6323 <https://www.su.se/english/research/kebnekaise-s-south-peak-still-highest-in-sweden-1.343483>, access:
 6324 14.12.2020, 2017.



- 6325 Stockholm University - Department of Physical Geography: Southern peak of Kebnekaise, lowest height ever
 6326 measured: [https://www.natgeo.su.se/english/research/resarch-news/southern-peak-of-kebnekaise-lowest-height-](https://www.natgeo.su.se/english/research/resarch-news/southern-peak-of-kebnekaise-lowest-height-ever-measured-1.452263)
 6327 ever-measured-1.452263, access: 14.12.2020, 2019.
- 6328 Stockholm University - Department of Physical Geography: Tarfala Research Station:
 6329 <https://www.natgeo.su.se/english/tarfala-research-station>, access: 14.12.2020, 2020.
- 6330 Stokowski, M., Winogradow, A., Szymczycha, B., Carstensen, J., and Kuliński, K.: The CO₂ system dynamics
 6331 in the vicinity of the Vistula River mouth (the southern Baltic Sea): A baseline investigation, *Estuar. Coast. Shelf*
 6332 *Sci.*, 258, 107444, <https://doi.org/10.1016/j.ecss.2021.107444>, 2021.
- 6333 Stonevičius, E., Rimkus, E., Štaras, A., Kažys, J., and Valiuškevičius, G.: Climate change impact on the
 6334 Nemunas River basin hydrology in the 21st century, *Boreal Environ. Res.*, 22, 49-65, 2017.
- 6335 Storelvmo, T., Heede, U. K., Leirvik, T., Phillips, P. C. B., Arndt, P., and Wild, M.: Lethargic Response to
 6336 Aerosol Emissions in Current Climate Models, *Geophys. Res. Lett.*, 45, 9814-9823,
 6337 <https://doi.org/10.1029/2018GL078298>, 2018.
- 6338 Stramska, M., and Białogrodzka, J.: Spatial and temporal variability of sea surface temperature in the Baltic Sea
 6339 based on 32-years (1982–2013) of satellite data, *Oceanologia*, 57, 223-235,
 6340 <https://doi.org/10.1016/j.oceano.2015.04.004>, 2015.
- 6341 Strandberg, G., and Kjellström, E.: Climate Impacts from Afforestation and Deforestation in Europe, *Earth*
 6342 *Interact.*, 23, 1-27, <https://doi.org/10.1175/ei-d-17-0033.1>, 2019.
- 6343 Strong, C., and Magnusdottir, G.: Dependence of NAO variability on coupling with sea ice, *Clim. Dyn.*, 36,
 6344 1681-1689, <https://doi.org/10.1007/s00382-010-0752-z>, 2011.
- 6345 Suikkanen, S., Pulina, S., Engström-Öst, J., Lehtiniemi, M., Lehtinen, S., and Brutemark, A.: Climate Change
 6346 and Eutrophication Induced Shifts in Northern Summer Plankton Communities, *PLoS ONE*, 8, e66475,
 6347 <https://doi.org/10.1371/journal.pone.0066475>, 2013.
- 6348 Sun, L., Perlwitz, J., and Hoerling, M.: What caused the recent “Warm Arctic, Cold Continents” trend pattern in
 6349 winter temperatures?, *Geophys. Res. Lett.*, 43, 5345-5352, <https://doi.org/10.1002/2016GL069024>, 2016.
- 6350 Sündermann, J., and Pohlmann, T.: A brief analysis of North Sea physics, *Oceanologia*, 53, 663-689,
 6351 <https://doi.org/10.5697/oc.53-3.663>, 2011.
- 6352 Sundqvist, L., Harkonen, T., Svensson, C. J., and Harding, K. C.: Linking Climate Trends to Population
 6353 Dynamics in the Baltic Ringed Seal: Impacts of Historical and Future Winter Temperatures, *Ambio* 41, 865-872,
 6354 <https://doi.org/10.1007/s13280-012-0334-x>, 2012.
- 6355 Sutton, R. T., and Hodson, D. L. R.: Atlantic Ocean Forcing of North American and European Summer Climate,
 6356 *Science*, 309, 115-118, <https://doi.org/10.1126/science.1109496>, 2005.
- 6357 Suursaar, Ü., and Sooäär, J.: Decadal variations in mean and extreme sea level values along the Estonian coast of
 6358 the Baltic Sea, *Tellus A*, 59, 249-260, <https://doi.org/10.1111/j.1600-0870.2006.00220.x>, 2007.
- 6359 Suursaar, Ü., Tõnisson, H., Alari, V., Raudsepp, U., Rästas, H., and Anderson, A.: Projected Changes in Wave
 6360 Conditions in the Baltic Sea by the end of 21st Century and the Corresponding Shoreline Changes, *J. Coastal*
 6361 *Res.*, 1012-1016, <https://doi.org/10.2112/si75-203.1>, 2016.
- 6362 Sveegaard, S., Andreassen, H., Mouritsen, K. N., Jeppesen, J. P., Teilmann, J., and Kinze, C. C.: Correlation
 6363 between the seasonal distribution of harbour porpoises and their prey in the Sound, Baltic Sea, *Mar. Biol.*, 159,
 6364 1029-1037, <https://doi.org/10.1007/s00227-012-1883-z>, 2012.
- 6365 Svendsen, L. M., and Gustafsson, B.: Waterborne nitrogen and phosphorus inputs and water flow to the Baltic
 6366 Sea 1995-2018, HELCOM Baltic Sea Environmental Fact Sheet, 25pp., 2020.
- 6367 Swedish Infrastructure for Ecosystem Science: Meteorological data from Tarfala Research Station:
 6368 <https://data.fieldsites.se/portal/>, access: 14.12.2020, 2020.
- 6369 Swedish Infrastructure for Ecosystem Science: Glacier data - surface mass balance from Rabots Glaciär, 1982–
 6370 2020, <https://meta.fieldsites.se/objects/KI06fLDs-YUzlvq2GmVKTNg>, 2021a.
- 6371 Swedish Infrastructure for Ecosystem Science: Glacier data - surface mass balance from Mårnaglaciären, 1986–
 6372 2020, <https://meta.fieldsites.se/objects/EXzEgEOQbn9ofJJrTX2fMHH>, 2021b.



- 6373 Swedish Infrastructure for Ecosystem Science: Glacier data - surface mass balance from Riukojietna, 1986–
 6374 2020, <https://meta.fieldsites.se/objects/BZvbLykTx4cnEVaEuMSFVQeK>, 2021c.
- 6375 SMHI - Swedish Meteorological and Hydrological Institute: Climate scenarios:
 6376 <http://www.smhi.se/en/climate/future-climate/climate-scenarios/>, access: 14.12.2020, 2020.
- 6377 SMHI - Swedish Meteorological and Hydrological Institute: Klimatindikator – Globalstrålning:
 6378 <https://www.smhi.se/klimat/klimatet-da-och-nu/klimatindikatorer/stralning-1.17841>, access: 12.02.2021, 2021.
- 6379 Sweet, W., Kopp, R. E., Weaver, C. P., Obeysekera, J. T. B., Horton, R. M., Thieler, E. R., and Zervas, C. E.:
 6380 Global and regional sea level rise scenarios for the United States, <https://doi.org/10.7289/v5/tr-nos-coops-083>,
 6381 2017.
- 6382 Szwed, M., Pińskwar, I., Kundzewicz, Z. W., Graczyk, D., and Mezghani, A.: Changes of snow cover in Poland,
 6383 *Acta Geophys.*, 65, 65-76, <https://doi.org/10.1007/s11600-017-0007-z>, 2017.
- 6384 Szwed, M., Dobler, A., Mezghani, A., and Saloranta, T. M.: Change of maximum snow cover depth in Poland -
 6385 Trends and projections, *Időjárás*, 123, 487-500, <https://doi.org/10.28974/idojaras.2019.4.5>, 2019.
- 6386 Takolander, A., Cabeza, M., and Leskinen, E.: Climate change can cause complex responses in Baltic Sea
 6387 macroalgae: A systematic review, *J. Sea Res.*, 123, 16-29, <https://doi.org/10.1016/j.seares.2017.03.007>, 2017.
- 6388 Tamarin-Brodsky, T., and Kaspi, Y.: Enhanced poleward propagation of storms under climate change, *Nat.*
 6389 *Geosci.*, 10, 908-913, <https://doi.org/10.1038/s41561-017-0001-8>, 2017.
- 6390 Tamarin, T., and Kaspi, Y.: The poleward shift of storm tracks under global warming: A Lagrangian perspective,
 6391 *Geophys. Res. Lett.*, 44, 10,666-610,674, <https://doi.org/10.1002/2017GL073633>, 2017.
- 6392 Tang, J., Yurova, A. Y., Schurgers, G., Miller, P. A., Olin, S., Smith, B., Siewert, M. B., Olefeldt, D., Pilesjö, P.,
 6393 and Poska, A.: Drivers of dissolved organic carbon export in a subarctic catchment: Importance of microbial
 6394 decomposition, sorption-desorption, peatland and lateral flow, *Sci. Total Environ.*, 622-623, 260-274,
 6395 <https://doi.org/10.1016/j.scitotenv.2017.11.252>, 2018.
- 6396 Tang, L., Ramacher, M. O. P., Moldanová, J., Matthias, V., Karl, M., Johansson, L., Jalkanen, J. P., Yaramenka,
 6397 K., Aulinger, A., and Gustafsson, M.: The impact of ship emissions on air quality and human health in the
 6398 Gothenburg area – Part 1: 2012 emissions, *Atmos. Chem. Phys.*, 20, 7509-7530, <https://doi.org/10.5194/acp-20-7509-2020>, 2020.
- 6400 Taniguchi, M., Dulai, H., Burnett, K. M., Santos, I. R., Sugimoto, R., Stieglitz, T., Kim, G., Moosdorf, N., and
 6401 Burnett, W. C.: Submarine Groundwater Discharge: Updates on Its Measurement Techniques, Geophysical
 6402 Drivers, Magnitudes, and Effects, *Front. Environ. Sci.*, 7, <https://doi.org/10.3389/fenvs.2019.00141>, 2019.
- 6403 Teichmann, C., Bülow, K., Otto, J., Pfeifer, S., Rechid, D., Sieck, K., and Jacob, D.: Avoiding Extremes:
 6404 Benefits of Staying below +1.5 °C Compared to +2.0 °C and +3.0 °C Global Warming, *Atmosphere*, 9, 115,
 6405 <https://doi.org/10.3390/atmos9040115>, 2018.
- 6406 Teuling, A. J.: A hot future for European droughts, *Nat. Clim. Change*, 8, 364-365,
 6407 <https://doi.org/10.1038/s41558-018-0154-5>, 2018.
- 6408 Thompson, P. M.: Seasonal changes in the distribution and composition of common seal (*Phoca vitulina*) haul-
 6409 out groups, *J. Zool.*, 217, 281-294, <https://doi.org/10.1111/j.1469-7998.1989.tb02488.x>, 1989.
- 6410 Thomson, A. M., Calvin, K. V., Smith, S. J., Kyle, G. P., Volke, A., Patel, P., Delgado-Arias, S., Bond-
 6411 Lamberty, B., Wise, M. A., Clarke, L. E., and Edmonds, J. A.: RCP4.5: a pathway for stabilization of radiative
 6412 forcing by 2100, *Clim. Change*, 109, 77, <https://doi.org/10.1007/s10584-011-0151-4>, 2011.
- 6413 Tikkanen, M., and Oksanen, J.: Late Weichselian and Holocene shore displacement history of the Baltic Sea in
 6414 Finland, Fennia - International Journal of Geography, 180, 9,20, 2002.
- 6415 Tilinina, N., Gulev, S. K., Rudeva, I., and Koltermann, P.: Comparing Cyclone Life Cycle Characteristics and
 6416 Their Interannual Variability in Different Reanalyses, *J. Climate*, 26, 6419-6438, <https://doi.org/10.1175/jcli-d-12-00777.1>, 2013.
- 6418 Timmermann, K., Norkko, J., Janas, U., Norkko, A., Gustafsson, B. G., and Bonsdorff, E.: Modelling
 6419 macrofaunal biomass in relation to hypoxia and nutrient loading, *J. Mar. Syst.*, 105-108, 60-69,
 6420 <https://doi.org/10.1016/j.jmarsys.2012.06.001>, 2012.



- 6421 Ting, M., Kushnir, Y., Seager, R., and Li, C.: Robust features of Atlantic multi-decadal variability and its
 6422 climate impacts, *Geophys. Res. Lett.*, 38, <https://doi.org/10.1029/2011GL048712>, 2011.
- 6423 Tinz, B.: On the relation between annual maximum extent of ice cover in the Baltic Sea and sea level pressure as
 6424 well as air temperature field, *Geophysica*, 32, 319-341, 1996.
- 6425 Tokarska, K. B., Stolpe, M. B., Sippel, S., Fischer, E. M., Smith, C. J., Lehner, F., and Knutti, R.: Past warming
 6426 trend constrains future warming in CMIP6 models, *Sci. Adv.*, 6, eaaz9549,
 6427 <https://doi.org/10.1126/sciadv.aaz9549>, 2020.
- 6428 Tomczyk, A. M., Bednorz, E., and Szyga-Pluta, K.: Changes in Air Temperature and Snow Cover in Winter in
 6429 Poland, *Atmosphere*, 12, 68, <https://doi.org/10.3390/atmos12010068>, 2021.
- 6430 Torn, K., Peterson, A., and Herkül, K.: Predicting the Impact of Climate Change on the Distribution of the Key
 6431 Habitat-Forming Species in the Ne Baltic Sea, *J. Coastal Res.*, 95, 177-181, 175, <https://doi.org/10.2112/SI95-035.1>, 2020.
- 6433 Törnroos, A., Pecuchet, L., Olsson, J., Gårdmark, A., Blomqvist, M., Lindegren, M., and Bonsdorff, E.: Four
 6434 decades of functional community change reveals gradual trends and low interlinkage across trophic groups in a
 6435 large marine ecosystem, *Global Change Biol.*, 25, 1235-1246, <https://doi.org/10.1111/gcb.14552>, 2019.
- 6436 Tuomi, L., Kahma, K. K., and Pettersson, H.: Wave hindcast statistics in the seasonally ice-covered Baltic Sea,
 6437 *Boreal Environ. Res.*, 16, 451-472, 2011.
- 6438 Turner, D. R., Hassellöv, I.-M., Ytreberg, E., and Rutgersson, A.: Shipping and the environment: Smokestack
 6439 emissions, scrubbers and unregulated oceanic consequences, *Elementa-Sci. Anthropol.*, 5,
 6440 <https://doi.org/10.1525/elementa.167>, 2017.
- 6441 Turner, D. R., Edman, M., Gallego-Urrea, J. A., Claremar, B., Hassellöv, I.-M., Omstedt, A., and Rutgersson,
 6442 A.: The potential future contribution of shipping to acidification of the Baltic Sea, *Ambio*, 47, 368-378,
 6443 <https://doi.org/10.1007/s13280-017-0950-6>, 2018.
- 6444 Ukkonen, P., Aaris-Sørensen, K., Arppe, L., Daugnora, L., Halkka, A., Lõugas, L., Oinonen, M. J., Pilot, M.,
 6445 and Storå, J.: An Arctic seal in temperate waters: History of the ringed seal (*Pusa hispida*) in the Baltic Sea and
 6446 its adaptation to the changing environment, *Holocene*, 24, 1694-1706,
 6447 <https://doi.org/10.1177/0959683614551226>, 2014.
- 6448 Ulbrich, U., Pinto, J. G., Kupfer, H., Leckebusch, G. C., Spanghel, T., and Reyers, M.: Changing Northern
 6449 Hemisphere Storm Tracks in an Ensemble of IPCC Climate Change Simulations, *J. Climate*, 21, 1669-1679,
 6450 <https://doi.org/10.1175/2007jcli1992.1>, 2008.
- 6451 Ulfsbo, A., Kuliński, K., Anderson, L. G., and Turner, D. R.: Modelling organic alkalinity in the Baltic Sea
 6452 using a Humic-Pitzer approach, *Mar. Chem.*, 168, 18-26, <https://doi.org/10.1016/j.marchem.2014.10.013>, 2015.
- 6453 Umlauf, L., Holtermann, P. L., Gillner, C. A., Prien, R. D., Merkelbach, L., and Carpenter, J. R.: Diffusive
 6454 Convection under Rapidly Varying Conditions, *J. Phys. Oceanogr.*, 48, 1731-1747, <https://doi.org/10.1175/jpo-d-18-0018.1>, 2018.
- 6456 Undeman, E., Gustafsson, B. G., Humborg, C., and McLachlan, M. S.: Application of a novel modeling tool
 6457 with multistressor functionality to support management of organic contaminants in the Baltic Sea, *Ambio* 44,
 6458 498-506, <https://doi.org/10.1007/s13280-015-0668-2>, 2015.
- 6459 United Nations Climate Change: Paris Agreement, 27pp, <https://unfccc.int/documents/37107>, 2015.
- 6460 Uotila, P., Vihma, T., and Haapala, J.: Atmospheric and oceanic conditions and the extremely low Bothnian Bay
 6461 sea ice extent in 2014/2015, *Geophys. Res. Lett.*, 42, 7740-7749, <https://doi.org/10.1002/2015GL064901>, 2015.
- 6462 Urraca, R., Gracia-Amillo, A. M., Koubli, E., Huld, T., Trentmann, J., Riihelä, A., Lindfors, A. V., Palmer, D.,
 6463 Gottschalg, R., and Antonanzas-Torres, F.: Extensive validation of CM SAF surface radiation products over
 6464 Europe, *Remote Sens. Environ.*, 199, 171-186, <https://doi.org/10.1016/j.rse.2017.07.013>, 2017.
- 6465 Vähätalo, A. V., Rainio, K., Lehtikoinen, A., and Lehtikoinen, E.: Spring arrival of birds depends on the North
 6466 Atlantic Oscillation, *J. Avian Biol.*, 35, 210-216, <https://doi.org/10.1111/j.0908-8857.2004.03199.x>, 2004.
- 6467 Vahtera, E., Conley, D. J., Gustafsson, B. G., Kuosa, H., Pitkänen, H., Savchuk, O. P., Tamminen, T., Viitasalo,
 6468 M., Voss, M., Wasmund, N., and others: Internal ecosystem feedbacks enhance nitrogen-fixing cyanobacteria



- 6469 blooms and complicate management in the Baltic Sea, *Ambio*, 36, 186-194, [https://doi.org/10.1579/0044-](https://doi.org/10.1579/0044-7447(2007)36[186:IEFENC]2.0.CO;2)
 6470 7447(2007)36[186:IEFENC]2.0.CO;2, 2007.
- 6471 Väli, G., Meier, H. E. M., and Elken, J.: Simulated halocline variability in the Baltic Sea and its impact on
 6472 hypoxia during 1961-2007, *J. Geophys. Res-Oceans*, 118, 6982-7000, <https://doi.org/10.1002/2013JC009192>,
 6473 2013.
- 6474 van der Jeugd, H. P., Eichhorn, G., Litvin, K. E., Stahl, J., Larsson, K., van der Graaf, A. J., and Drent, R. H.:
 6475 Keeping up with early springs: rapid range expansion in an avian herbivore incurs a mismatch between
 6476 reproductive timing and food supply, *Global Change Biol.*, 15, 1057-1071, [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2486.2008.01804.x)
 6477 2486.2008.01804.x, 2009.
- 6478 van Helmond, N. A. G. M., Jilbert, T., and Slomp, C. P.: Hypoxia in the Holocene Baltic Sea: Comparing
 6479 modern versus past intervals using sedimentary trace metals, *Chem. Geol.*, 493, 478-490,
 6480 <https://doi.org/10.1016/j.chemgeo.2018.06.028>, 2018.
- 6481 van Vuuren, D. P., Stehfest, E., den Elzen, M. G. J., Kram, T., van Vliet, J., Deetman, S., Isaac, M., Klein
 6482 Goldewijk, K., Hof, A., Mendoza Beltran, A., Oostenrijk, R., and van Ruijven, B.: RCP2.6: exploring the
 6483 possibility to keep global mean temperature increase below 2°C, *Clim. Change*, 109, 95,
 6484 <https://doi.org/10.1007/s10584-011-0152-3>, 2011.
- 6485 Vanninen, P., Östin, A., Beldowski, J., Pedersen, E. A., Söderström, M., Szubska, M., Grabowski, M.,
 6486 Siedlewicz, G., Czub, M., Popiel, S., Nawala, J., Dziedzic, D., Jakacki, J., and Pączek, B.: Exposure status of
 6487 sea-dumped chemical warfare agents in the Baltic Sea, *Mar. Environ. Res.*, 161, 105112,
 6488 <https://doi.org/10.1016/j.marenvres.2020.105112>, 2020.
- 6489 Varotsos, K. V., Giannakopoulos, C., and Tombrou, M.: Assessment of the Impacts of Climate Change on
 6490 European Ozone Levels, Water, Air & Soil Pollution, 224, 1596, <https://doi.org/10.1007/s11270-013-1596-z>,
 6491 2013.
- 6492 Vaughan, D. G., Comiso, J. C., Allison, I., Carrasco, J., Kaser, G., Kwok, R., Mote, P., Murray, T., Paul, F., Ren,
 6493 J., Rignot, E., Solomina, O., Steffen, K., and Zhang, T.: Observations: Cryosphere, in: *Climate Change 2013 –*
 6494 *The Physical Science Basis: Working Group I Contribution to the Fifth Assessment Report of the*
 6495 *Intergovernmental Panel on Climate Change*, edited by: Intergovernmental Panel on Climate, C., Cambridge
 6496 University Press, Cambridge, 317-382, <https://doi.org/10.1017/CBO9781107415324.012>, 2014.
- 6497 Vehmaa, A., Hogfors, H., Gorokhova, E., Brutemark, A., Holmborn, T., and Engström-Öst, J.: Projected marine
 6498 climate change: effects on copepod oxidative status and reproduction, *Ecol. Evol.*, 3, 4548-4557,
 6499 <https://doi.org/10.1002/ece3.839>, 2013.
- 6500 Vehmaa, A., Almén, A. K., Brutemark, A., Paul, A., Riebesell, U., Furuhaugen, S., and Engström-Öst, J.: Ocean
 6501 acidification challenges copepod phenotypic plasticity, *Biogeosciences*, 13, 6171-6182,
 6502 <https://doi.org/10.5194/bg-13-6171-2016>, 2016.
- 6503 Veijalainen, N., Lotsari, E., Vehviläinen, B., Alho, P., and Käyhkö, J.: Changes in floods in Finland due to
 6504 climate change: General assessment on national scale, 6th Alexander von Humboldt International Conference on
 6505 Climate Change, 2010.
- 6506 Vihma, T., and Haapala, J.: Geophysics of sea ice in the Baltic Sea: A review, *Prog. Oceanogr.*, 80, 129-148,
 6507 <https://doi.org/10.1016/j.pocean.2009.02.002>, 2009.
- 6508 Vihma, T.: Weather Extremes Linked to Interaction of the Arctic and Midlatitudes, in: *Climate Extremes*, edited
 6509 by: Wang, S. Y. S., Yoon, J. H., Funk, C. C., and Gillies, R. R., 39-50,
 6510 <https://doi.org/10.1002/9781119068020.ch2>, 2017.
- 6511 Viitasalo, M.: Climate change and the Baltic Sea ecosystem: direct and indirect effects on species, communities
 6512 and ecosystem function *Earth Syst. Dynam.*, 2021.
- 6513 Viru, B., and Jaagus, J.: Spatio-temporal variability and seasonal dynamics of snow cover regime in Estonia,
 6514 *Theor. Appl. Climatol.*, 139, 759-771, <https://doi.org/10.1007/s00704-019-03013-5>, 2020.
- 6515 Viška, M., and Soomere, T.: Simulated and observed reversals of wave-driven alongshore sediment transport at
 6516 the eastern Baltic Sea coast, *Baltica*, 26, 145-156, 2013.



- 6517 Vogt, J., Soille, P., Jager, A. d., Rimaviciute, E., Mehl, W., Foisneau, S., Bódis, K., Dusart, J., Paracchini, M. L.,
 6518 Hastrup, P., and Bamps, C.: A pan-European River and Catchment Database, European Commission - Joint
 6519 Research Centre - Institute for Environment and Sustainability, Ispra, Italy, 124pp, 2007.
- 6520 Vogt, J., Rimaviciute, E., and Jager, A. d.: CCM2 River and Catchment Database for Europe Version 2.1
 6521 Release Notes, European Commission - Joint Research Centre - Institute for Environment and Sustainability,
 6522 Ispra, Italy, 5pp, 2008.
- 6523 Voipio, A.: The Baltic Sea, Elsevier Oceanographic Series, Amsterdam, 1981.
- 6524 Volkov, V. A., Johannessen, O. M., Borodachev, V. E., Voinov, G. N., Pettersson, L. H., Bobylev, L. P., and
 6525 Kouraev, A. V.: Polar seas oceanography: an integrated case study of the Kara Sea, Springer Science & Business
 6526 Media, Berlin, Heidelberg, 2002.
- 6527 Vonk, J. E., Tank, S. E., Bowden, W. B., Laurion, I., Vincent, W. F., Alekseychik, P., Amyot, M., Billet, M.,
 6528 Canário, J., and Cory, R. M.: Reviews and syntheses: Effects of permafrost thaw on Arctic aquatic ecosystems,
 6529 Biogeosciences, 12, 7129-7167, <https://doi.org/10.5194/bg-12-7129-2015>, 2015.
- 6530 Voss, R., Hinrichsen, H.-H., Quaas, M. F., Schmidt, J. O., and Tahvonen, O.: Temperature change and Baltic
 6531 sprat: from observations to ecological-economic modelling, ICES J. Mar. Sci., 68, 1244-1256,
 6532 <https://doi.org/10.1093/icesjms/fsr063>, 2011.
- 6533 Vousdoukas, M. I., Voulkouvalas, E., Annunziato, A., Giardino, A., and Feyen, L.: Projections of extreme storm
 6534 surge levels along Europe, Clim. Dyn., 47, 3171-3190, <https://doi.org/10.1007/s00382-016-3019-5>, 2016.
- 6535 Vousdoukas, M. I., Mentaschi, L., Voulkouvalas, E., Verlaan, M., and Feyen, L.: Extreme sea levels on the rise
 6536 along Europe's coasts, Earths Future, 5, 304-323, <https://doi.org/10.1002/2016EF000505>, 2017.
- 6537 Vuorinen, I., Hänninen, J., Rajasilta, M., Laine, P., Eklund, J., Montesino-Pouzols, F., Corona, F., Junker, K.,
 6538 Meier, H. E. M., and Dippner, J. W.: Scenario simulations of future salinity and ecological consequences in the
 6539 Baltic Sea and adjacent North Sea areas—implications for environmental monitoring, Ecol. Indic., 50, 196-205,
 6540 <https://doi.org/10.1016/j.ecolind.2014.10.019>, 2015.
- 6541 Wahl, M., Werner, F. J., Buchholz, B., Raddatz, S., Graiff, A., Matthiessen, B., Karsten, U., Hiebenthal, C.,
 6542 Hamer, J., Ito, M., Gültow, E., Rilov, G., and Guy-Haim, T.: Season affects strength and direction of the
 6543 interactive impacts of ocean warming and biotic stress in a coastal seaweed ecosystem, Limnol. Oceanogr., 65,
 6544 807-827, <https://doi.org/10.1002/lno.11350>, 2020.
- 6545 Wahl, T., and Chambers, D. P.: Climate controls multidecadal variability in U. S. extreme sea level records, J.
 6546 Geophys. Res.-Oceans, 121, 1274-1290, <https://doi.org/10.1002/2015JC011057>, 2016.
- 6547 Wählström, I., Höglund, A., Almroth-Rosell, E., MacKenzie, B. R., Gröger, M., Eilola, K., Plikshs, M., and
 6548 Andersson, H. C.: Combined climate change and nutrient load impacts on future habitats and eutrophication
 6549 indicators in a eutrophic coastal sea, Limnol. Oceanogr., 65, 2170-2187, <https://doi.org/10.1002/lno.11446>,
 6550 2020.
- 6551 Waldeck, P., and Larsson, K.: Effects of winter water temperature on mass loss in Baltic blue mussels:
 6552 Implications for foraging sea ducks, J. Exp. Mar. Biol. Ecol., 444, 24-30,
 6553 <https://doi.org/10.1016/j.jembe.2013.03.007>, 2013.
- 6554 Wang, J., Yang, B., Ljungqvist, F. C., Luterbacher, J., Osborn, Timothy J., Briffa, K. R., and Zorita, E.: Internal
 6555 and external forcing of multidecadal Atlantic climate variability over the past 1,200 years, Nat. Geosci., 10, 512-
 6556 517, <https://doi.org/10.1038/ngeo2962>, 2017.
- 6557 Wang, Q., Fan, X., and Wang, M.: Evidence of high-elevation amplification versus Arctic amplification, Sci.
 6558 Rep.-UK, 6, 19219, <https://doi.org/10.1038/srep19219>, 2016.
- 6559 Wang, S., Dieterich, C., Döscher, R., Höglund, A., Hordoir, R., Meier, H. E. M., Samuelsson, P., and
 6560 Schimanke, S.: Development and evaluation of a new regional coupled atmosphere-ocean model in the North
 6561 Sea and Baltic Sea, Tellus A, 67, 24284, <https://doi.org/10.3402/tellusa.v67.24284>, 2015.
- 6562 Warwick, R. M., Tweedley, J. R., and Potter, I. C.: Microtidal estuaries warrant special management measures
 6563 that recognise their critical vulnerability to pollution and climate change, Mar. Pollut. Bull., 135, 41-46,
 6564 <https://doi.org/10.1016/j.marpolbul.2018.06.062>, 2018.



- 6565 Wasmund, N., Tuimala, J., Suikkanen, S., Vandepitte, L., and Kraberg, A.: Long-term trends in phytoplankton
 6566 composition in the western and central Baltic Sea, *J. Mar. Syst.*, 87, 145-159,
 6567 <https://doi.org/10.1016/j.jmarsys.2011.03.010>, 2011.
- 6568 Wasmund, N.: The Diatom/Dinoflagellate Index as an Indicator of Ecosystem Changes in the Baltic Sea. 2.
 6569 Historical Data for Use in Determination of Good Environmental Status, *Front. Mar. Sci.*, 4,
 6570 <https://doi.org/10.3389/fmars.2017.00153>, 2017.
- 6571 Wasmund, N., Kownacka, J., Göbel, J., Jaanus, A., Johansen, M., Jurgensone, I., Lehtinen, S., and Powilleit, M.:
 6572 The Diatom/Dinoflagellate Index as an Indicator of Ecosystem Changes in the Baltic Sea 1. Principle and
 6573 Handling Instruction, *Front. Mar. Sci.*, 4, <https://doi.org/10.3389/fmars.2017.00022>, 2017.
- 6574 Wasmund, N., Nausch, G., Gerth, M., Busch, S., Burmeister, C., Hansen, R., and Sadkowiak, B.: Extension of
 6575 the growing season of phytoplankton in the western Baltic Sea in response to climate change, *Mar. Ecol. Prog.
 6576 Ser.*, 622, 1-16, <https://doi.org/10.3354/meps12994>, 2019.
- 6577 Watson, L., Lacrosonnière, G., Gauss, M., Engardt, M., Andersson, C., Josse, B., Marécal, V., Nyiri, A.,
 6578 Sobolowski, S., Siour, G., Szopa, S., and Vautard, R.: Impact of emissions and +2 °C climate change upon future
 6579 ozone and nitrogen dioxide over Europe, *Atmos. Environ.*, 142, 271-285,
 6580 <https://doi.org/10.1016/j.atmosenv.2016.07.051>, 2016.
- 6581 Watts, P.: The diel hauling-out cycle of harbour seals in an open marine environment: correlates and constraints,
 6582 *J. Zool.*, 240, 175-200, <https://doi.org/10.1111/j.1469-7998.1996.tb05494.x>, 1996.
- 6583 Weigel, B., Andersson, H. C., Meier, H. E. M., Blenckner, T., Snickars, M., and Bonsdorff, E.: Long-term
 6584 progression and drivers of coastal zoobenthos in a changing system, *Mar. Ecol. Prog. Ser.*, 528, 141-159,
 6585 <https://doi.org/10.3354/meps11279>, 2015.
- 6586 Weisse, R., and Weidemann, H.: Baltic Sea extreme sea levels 1948-2011: Contributions from atmospheric
 6587 forcing, *Proc. IUTAM*, 25, 65-69, <https://doi.org/10.1016/j.piutam.2017.09.010>, 2017.
- 6588 Weisse, R., Dailidienė, I., Hünicke, B., Kahma, K., Madsen, K., Omstedt, A., Parnell, K., Schöne, T., Soomere,
 6589 T., Zhang, W., and Zorita, E.: Sea Level Dynamics and Coastal Erosion in the Baltic Sea Region, *Earth Syst.
 6590 Dynam. Discuss.*, 1-40, <https://doi.org/10.5194/esd-2021-6>, 2021.
- 6591 Whan, K., Zscheischler, J., Orth, R., Shongwe, M., Rahimi, M., Asare, E. O., and Seneviratne, S. I.: Impact of
 6592 soil moisture on extreme maximum temperatures in Europe, *Weather. Clim. Extremes*, 9, 57-67,
 6593 <https://doi.org/10.1016/j.wace.2015.05.001>, 2015.
- 6594 Wikner, J., and Andersson, A.: Increased freshwater discharge shifts the trophic balance in the coastal zone of
 6595 the northern Baltic Sea, *Global Change Biol.*, 18, 2509-2519, <https://doi.org/10.1111/j.1365-2486.2012.02718.x>,
 6596 2012.
- 6597 Wild, M., Gilgen, H., Roesch, A., Ohmura, A., Long, C. N., Dutton, E. G., Forgan, B., Kallis, A., Russak, V.,
 6598 and Tsvetkov, A.: From Dimming to Brightening: Decadal Changes in Solar Radiation at Earth's Surface,
 6599 *Science*, 308, 847, <https://doi.org/10.1126/science.1103215>, 2005.
- 6600 Wild, M.: Enlightening Global Dimming and Brightening, *Bull. Amer. Meteor. Soc.*, 93, 27-37,
 6601 <https://doi.org/10.1175/bams-d-11-00074.1>, 2012.
- 6602 Wild, M.: Decadal changes in radiative fluxes at land and ocean surfaces and their relevance for global warming,
 6603 *Wires. Clim. Change*, 7, 91-107, <https://doi.org/10.1002/wcc.372>, 2016.
- 6604 Wild, M., Ohmura, A., Schär, C., Müller, G., Folini, D., Schwarz, M., Hakuba, M. Z., and Sanchez-Lorenzo, A.:
 6605 The Global Energy Balance Archive (GEBA) version 2017: a database for worldwide measured surface energy
 6606 fluxes, *Earth Syst. Sci. Data*, 9, 601-613, <https://doi.org/10.5194/essd-9-601-2017>, 2017.
- 6607 Wild, M., Wacker, S., Yang, S., and Sanchez-Lorenzo, A.: Evidence for Clear-sky Dimming and Brightening in
 6608 Central Europe, *Geophys. Res. Lett.*, n/a, e2020GL092216, <https://doi.org/10.1029/2020GL092216>, 2021.
- 6609 Williams, C. N., Carrivick, J. L., Evans, A. J., and Rippin, D. M.: Quantifying uncertainty in using multiple
 6610 datasets to determine spatiotemporal ice mass loss over 101 years at kårsaglaciären, sub-arctic sweden, *Geogr.
 6611 Ann. A*, 98, 61-79, <https://doi.org/10.1111/geoa.12123>, 2016.
- 6612 Willison, J., Robinson, W. A., and Lackmann, G. M.: North Atlantic Storm-Track Sensitivity to Warming
 6613 Increases with Model Resolution, *J. Climate*, 28, 4513-4524, <https://doi.org/10.1175/jcli-d-14-00715.1>, 2015.



- 6614 Wills, R. C. J., Armour, K. C., Battisti, D. S., and Hartmann, D. L.: Ocean–Atmosphere Dynamical Coupling
 6615 Fundamental to the Atlantic Multidecadal Oscillation, *J. Climate*, 32, 251–272, [https://doi.org/10.1175/jcli-d-18-](https://doi.org/10.1175/jcli-d-18-0269.1)
 6616 0269.1, 2018.
- 6617 Wills, R. C. J., White, R. H., and Levine, X. J.: Northern Hemisphere Stationary Waves in a Changing Climate,
 6618 *Curr. Clim. Chang. Rep.*, 5, 372–389, <https://doi.org/10.1007/s40641-019-00147-6>, 2019.
- 6619 Wilson, D., Hisdal, H., and Lawrence, D.: Has streamflow changed in the Nordic countries? – Recent trends and
 6620 comparisons to hydrological projections, *J. Hydrol.*, 394, 334–346, <https://doi.org/10.1016/j.jhydrol.2010.09.010>,
 6621 2010.
- 6622 Wisniewska, Danuta M., Johnson, M., Teilmann, J., Rojano-Doñate, L., Shearer, J., Sveegaard, S., Miller,
 6623 Lee A., Siebert, U., and Madsen, Peter T.: Ultra-High Foraging Rates of Harbor Porpoises Make Them
 6624 Vulnerable to Anthropogenic Disturbance, *Curr. Biol.*, 26, 1441–1446,
 6625 <https://doi.org/10.1016/j.cub.2016.03.069>, 2016.
- 6626 Wohland, J., Brayshaw, D., Bloomfield, H., and Wild, M.: European multidecadal solar variability badly
 6627 captured in all centennial reanalyses except CERA20C, *Environ. Res. Lett.*, 15, 104021,
 6628 <https://doi.org/10.1088/1748-9326/aba7e6>, 2020.
- 6629 Woodruff, S. D., Worley, S. J., Lubker, S. J., Ji, Z., Eric Freeman, J., Berry, D. I., Brohan, P., Kent, E. C.,
 6630 Reynolds, R. W., Smith, S. R., and Wilkinson, C.: ICOADS Release 2.5: extensions and enhancements to the
 6631 surface marine meteorological archive, *Int. J. Climatol.*, 31, 951–967, <https://doi.org/10.1002/joc.2103>, 2011.
- 6632 Woollings, T., Barriopedro, D., Methven, J., Son, S.-W., Martius, O., Harvey, B., Sillmann, J., Lupo, A. R., and
 6633 Seneviratne, S.: Blocking and its Response to Climate Change, *Curr. Clim. Chang. Rep.*, 4, 287–300,
 6634 <https://doi.org/10.1007/s40641-018-0108-z>, 2018.
- 6635 World Glacier Monitoring Service: Fluctuations of Glaciers (FoG) Database:
 6636 https://wgms.ch/data_databaseversions/, access: 14.12.2020, 2020.
- 6637 World Meteorological Organization: State of the Global Climate 2020, 38pp, 2020.
- 6638 Wrzesiński, D., Chosiński, A., Ptak, M., and Skowron, R.: Effect of the North Atlantic Oscillation on the Pattern
 6639 of Lake Ice Phenology in Poland, *Acta Geophys.*, 63, 1664–1684, <https://doi.org/10.1515/acgeo-2015-0055>,
 6640 2015.
- 6641 Wu, S., Liu, Z.-Y., Cheng, J., and Li, C.: Response of North Pacific and North Atlantic decadal variability to
 6642 weak global warming, *Adv. Clim. Chang. Res.*, 9, 95–101, <https://doi.org/10.1016/j.accre.2018.03.001>, 2018.
- 6643 Wu, S., and Liu, Z.-Y.: Decadal Variability in the North Pacific and North Atlantic under Global Warming: The
 6644 Weakening Response and Its Mechanism, *J. Climate*, 33, 9181–9193, <https://doi.org/10.1175/jcli-d-19-1012.1>,
 6645 2020.
- 6646 Wübbler, C., and Krauss, W.: The two dimensional seiches of the Baltic Sea, *Oceanol. Acta*, 2, 435–446, 1979.
- 6647 Wulff, A., Karlberg, M., Olofsson, M., Torstensson, A., Riemann, L., Steinhoff, F. S., Mohlin, M., Ekstrand, N.,
 6648 and Chierici, M.: Ocean acidification and desalination: climate-driven change in a Baltic Sea summer
 6649 microplanktonic community, *Mar. Biol.*, 165, 63, <https://doi.org/10.1007/s00227-018-3321-3>, 2018.
- 6650 Wyser, K., van Noije, T., Yang, S., von Hardenberg, J., O'Donnell, D., and Döschner, R.: On the increased
 6651 climate sensitivity in the EC-Earth model from CMIP5 to CMIP6, *Geosci. Model Dev.*, 13, 3465–3474,
 6652 <https://doi.org/10.5194/gmd-13-3465-2020>, 2020.
- 6653 Ye, K., and Lau, N.-C.: Influences of surface air temperature and atmospheric circulation on winter snow cover
 6654 variability over Europe, *Int. J. Climatol.*, 37, 2606–2619, <https://doi.org/10.1002/joc.4868>, 2017.
- 6655 Yin, J. H.: A consistent poleward shift of the storm tracks in simulations of 21st century climate, *Geophys. Res.*
 6656 *Lett.*, 32, <https://doi.org/10.1029/2005GL023684>, 2005.
- 6657 Yli-Pelkonen, V.: Ecological information in the political decision making of urban land-use planning, *J. Environ.*
 6658 *Plan. Manag.*, 51, 345–362, <https://doi.org/10.1080/09640560801977224>, 2008.
- 6659 Zandersen, M., Hyytiäinen, K., Meier, H. E. M., Tomczak, M. T., Bauer, B., Haapasaari, P. E., Olesen, J. E.,
 6660 Gustafsson, B. G., Refsgaard, J. C., Fridell, E., Pihlainen, S., Le Tissier, M. D. A., Kosenius, A.-K., and Van
 6661 Vuuren, D. P.: Shared socio-economic pathways extended for the Baltic Sea: exploring long-term environmental
 6662 problems, *Reg. Environ. Change*, 19, 1073–1086, <https://doi.org/10.1007/s10113-018-1453-0>, 2019.



- 6663 Zappa, G., Shaffrey, L. C., Hodges, K. I., Sansom, P. G., and Stephenson, D. B.: A Multimodel Assessment of
 6664 Future Projections of North Atlantic and European Extratropical Cyclones in the CMIP5 Climate Models, *J.*
 6665 *Climate*, 26, 5846–5862, <https://doi.org/10.1175/jcli-d-12-00573.1>, 2013.
- 6666 Zappa, G., and Shepherd, T. G.: Storylines of Atmospheric Circulation Change for European Regional Climate
 6667 Impact Assessment, *J. Climate*, 30, 6561–6577, <https://doi.org/10.1175/jcli-d-16-0807.1>, 2017.
- 6668 Zappa, G., Pithan, F., and Shepherd, T. G.: Multimodel Evidence for an Atmospheric Circulation Response to
 6669 Arctic Sea Ice Loss in the CMIP5 Future Projections, *Geophys. Res. Lett.*, 45, 1011–1019,
 6670 <https://doi.org/10.1002/2017GL076096>, 2018.
- 6671 Zemp, M., Huss, M., Thibert, E., Eckert, N., McNabb, R., Huber, J., Barandun, M., Machguth, H., Nussbaumer,
 6672 S. U., Gärtner-Roer, I., Thomson, L., Paul, F., Maussion, F., Kutuzov, S., and Cogley, J. G.: Global glacier mass
 6673 changes and their contributions to sea-level rise from 1961 to 2016, *Nature*, 568, 382–386,
 6674 <https://doi.org/10.1038/s41586-019-1071-0>, 2019.
- 6675 Zemp, M., Gärtner-Roer, I., Nussbaumer, S. U., Bannwart, J., Rastner, P., Paul, F., Hoelzle, M., and [eds.]:
 6676 Global Glacier Change Bulletin No. 3 (2016–2017) World Glacier Monitoring Service, Zürich,
 6677 <https://doi.org/10.5167/uzh-191797>, 2020.
- 6678 Zhang, M., Lee, X., Yu, G., Han, S., Wang, H., Yan, J., Zhang, Y., Li, Y., Ohta, T., Hirano, T., Kim, J.,
 6679 Yoshifuji, N., and Wang, W.: Response of surface air temperature to small-scale land clearing across latitudes,
 6680 *Environ. Res. Lett.*, 9, 034002, <https://doi.org/10.1088/1748-9326/9/3/034002>, 2014a.
- 6681 Zhang, W., Harff, J., and Schneider, R.: Analysis of 50-year wind data of the southern Baltic Sea for modelling
 6682 coastal morphological evolution – a case study from the Darss-Zingst Peninsula, *Oceanologia*, 53, 489–518,
 6683 <https://doi.org/10.5697/oc.53-1-TI.489>, 2011.
- 6684 Zhang, W., Miller, P. A., Smith, B., Wania, R., Koenig, T., and Döscher, R.: Tundra shrubification and tree-
 6685 line advance amplify arctic climate warming: results from an individual-based dynamic vegetation model,
 6686 *Environ. Res. Lett.*, 8, 034023, <https://doi.org/10.1088/1748-9326/8/3/034023>, 2013.
- 6687 Zhang, W., Jansson, C., Miller, P. A., Smith, B., and Samuelsson, P.: Biogeophysical feedbacks enhance the
 6688 Arctic terrestrial carbon sink in regional Earth system dynamics, *Biogeosciences*, 11, 5503–5519,
 6689 <https://doi.org/10.5194/bg-11-5503-2014>, 2014b.
- 6690 Zhang, W., Schneider, R., Kolb, J., Teichmann, T., Dudzinska-Nowak, J., Harff, J., and Hanebuth, T. J. J.:
 6691 Land–sea interaction and morphogenesis of coastal foredunes — A modeling case study from the southern Baltic
 6692 Sea coast, *Coast. Eng.*, 99, 148–166, <https://doi.org/10.1016/j.coastaleng.2015.03.005>, 2015.
- 6693 Zhang, W., Schneider, R., Harff, J., Hünicke, B., and Fröhle, P.: Modelling of Medium-Term (Decadal) Coastal
 6694 Foredune Morphodynamics- Historical Hindcast and Future Scenarios of the Świna Gate Barrier Coast
 6695 (Southern Baltic Sea), in: *Coastline Changes of the Baltic Sea from South to East: Past and Future Projection*,
 6696 edited by: Harff, J., Furmańczyk, K., and von Storch, H., Springer International Publishing, Cham, 107–135,
 6697 https://doi.org/10.1007/978-3-319-49894-2_7, 2017.
- 6698 Zhang, W., Miller, P. A., Jansson, C., Samuelsson, P., Mao, J., and Smith, B.: Self-Amplifying Feedbacks
 6699 Accelerate Greening and Warming of the Arctic, *Geophys. Res. Lett.*, 45, 7102–7111,
 6700 <https://doi.org/10.1029/2018GL077830>, 2018.
- 6701 Zhong, X., Zhang, T., and Wang, K.: Snow density climatology across the former USSR, *Cryosphere*, 8, 785–
 6702 799, <https://doi.org/10.5194/tc-8-785-2014>, 2014.
- 6703 Zhu, X., Lee, S.-Y., Wen, X., Wei, Z., Ji, Z., Zheng, Z., and Dong, W.: Historical evolution and future trend of
 6704 Northern Hemisphere snow cover in CMIP5 and CMIP6 models, *Environ. Res. Lett.*, 16, 065013,
 6705 <https://doi.org/10.1088/1748-9326/ac0662>, 2021.
- 6706 Zillén, L., and Conley, D. J.: Hypoxia and cyanobacteria blooms-are they really natural features of the late
 6707 Holocene history of the Baltic Sea?, *Biogeosciences*, 7, 2567, <https://doi.org/10.5194/bg-7-2567-2010>, 2010.
- 6708 Zorita, E., and Laine, A.: Dependence of salinity and oxygen concentrations in the Baltic Sea on large-scale
 6709 atmospheric circulation, *Clim. Res.*, 14, 25–41, <https://doi.org/10.3354/cr014025>, 2000.