

The Deployment Length of Solar Radiation Modification: An Interplay of Mitigation, Net-negative Emissions and Climate Uncertainty

Susanne Baur^{1,2}, Alexander Nauels^{2,3}, Zebedee Nicholls³, Benjamin M. Sanderson⁵, Carl-Friedrich
5 Schleussner^{2,4}

¹CECI, Université de Toulouse, CERFACS, CNRS, Toulouse, 31100, France

²Climate Analytics, Berlin, 10969, Germany

³Australian-German Climate & Energy College, The University of Melbourne, Parkville, VIC 3010, Australia

⁴Geography Department and IRI THESys, Humboldt-Universität zu Berlin, Berlin, Germany

10 ⁵Centre for International Climate and Environmental Research (CICERO), Oslo, Norway

Correspondence to: Susanne Baur (susanne.baur@cerfacs.fr)

Abstract. A growing body of literature investigates the effects of Solar Radiation Modification (SRM) on global and regional
15 climates. Previous studies have focused on potentials and side-effects of SRM with little attention given to possible deployment
timescales and the levels of Carbon Dioxide Removal required for a phase out. Here, we investigate the deployment timescales
of SRM and how they are affected by different levels of mitigation, net-negative emissions (NNEs) and climate uncertainty.
We generate a large dataset of 355 emission scenarios in which SRM is deployed to keep warming levels at 1.5°C global mean
temperature. Probabilistic climate projections from this ensemble result in a large range of plausible future warming and
20 cooling rates that lead to various SRM deployment timescales. In all pathways consistent with extrapolated current ambition,
SRM deployment would exceed 100 years even under the most optimistic assumptions on climate response. As soon as the
temperature threshold is exceeded, neither mitigation nor NNEs or climate sensitivity alone can guarantee short deployment
timescales. Any deployment of SRM therefore comes with the risk of multi-century legacies of deployment, implying pan-
generational commitments of costs, risks and negative side effects of SRM and NNEs combined. Our study points towards an
25 additional risk of SRM that so far has received limited attention: Initialization and commitment to SRM would happen under
the uncertainty of climate reversibility and effects of SRM on mitigation ambition, as well as an implicit assumption that
Carbon Dioxide Removal can be scaled up sufficiently to allow SRM to be phased out again. All these parameters influence
SRM deployment length but will be mostly unknown at its initialisation time, rendering it impossible to predict how
'temporary' any SRM deployment would be.

30

1 Introduction

Emission pathways that reflect the level of climate ambition of current Nationally Determined Contributions (NDCs) until the end of the century are estimated to lead to an average global warming of around 2.4°C (CAT, 2022). This is still a lot higher than the warming limit agreed in the Paris Agreement of 2015 that entails holding warming to well below 2°C and pursuing efforts of limiting warming to 1.5°C (UNFCCC, 2016). The growing concern around overshooting the Paris Agreement’s long-term temperature target has led to a discussion of Solar Radiation Modification (SRM) which could in theory halt global temperature increase very rapidly, but only as long as it is actively supported (Irvine et al., 2016; Keith, 2000). SRM techniques intend to artificially lower global mean surface air temperature (GSAT) by modifying the radiative energy budget of the Earth system. Proposed methods include Stratospheric Aerosol Injection (SAI), Cirrus Cloud Thinning (CCT) and Marine Cloud Brightening (MCB) (Lawrence et al., 2018). SRM methods generally operate on one of the key impacts of climate change, temperature increase, without addressing its cause, anthropogenic greenhouse gas (GHG) emissions, or other impacts e.g. ocean acidification. Without explicit emission reductions as well as the removal of some of these climate forcers from the atmosphere in the long term, i.e. through Carbon Dioxide Removal (CDR) (Fuss et al., 2018), GHG emissions commit us to millennia of elevated temperature levels. Therefore, SRM deployment would only be temporary if combined with emission reductions and CDR.

Achieving the Paris Agreement’s target of 1.5°C relies on stringent mitigation with large near-term emission reductions as shown in the 1.5°C-compatible pathways assessed by the Intergovernmental Panel on Climate Change (IPCC) (Rogelj et al., 2018; IPCC, 2021; IPCC, 2022). It has been discussed that, in the absence of this strong near-term mitigation, SRM could be a tool to avoid the impacts associated with overshooting 1.5°C until emission reductions and CDR are sufficiently scaled up that SRM is no longer needed to artificially lower GSAT (Belaia et al., 2021; Buck et al., 2020; MacMartin et al., 2018; Neuber & Ott, 2020; Allen et al., 2018). This ‘buying-time’-approach, although criticised for relying on uncertain promises of SRM and CDR and increasing the risk of ‘climate debt’ (Asayama & Hulme, 2019), currently remains the dominant framing for any SRM deployment (Neuber & Ott, 2020). Surprisingly little analysis, however, has been done on the timescales this type of SRM deployment could entail. Tilmes et al. (2016) analysed climate impacts of pathways whose temperature would peak at 3°C by the end of the 21st century and use CDR and SRM to limit temperature increase to 2.5°C and 2°C. Similarly, MacMartin et al. (2018) chose an experimental setup where mitigation, CDR and SRM are used to meet the 1.5°C-goal from a ‘business-as-usual’ starting point. Both Tilmes et al. (2016) and MacMartin et al. (2018), did not discuss the length of SRM deployment and looked at selected illustrative pathways that cannot capture the many possible futures where a ‘buying-time’ approach of SRM could be embedded.

In this study, we generate a large dataset of scenarios that use SRM to avoid overshooting the 1.5°C warming target. The underlying emission scenarios reflect current NDCs until 2030 and subsequently diverge to spread a wide range of conditions by the end of the century. We employ this large scenario set to explore a large variety of futures of SRM interlinkages with

mitigation ambition and different magnitudes to which CDR could be scaled up. There is large uncertainty regarding the evolution of emissions under SRM with some studies arguing that SRM could be deployed for ‘peak-shaving’ under already
65 ambitious mitigation scenarios (Coninck et al., 2018) or does not negatively affect the public's willingness to engage in mitigation behaviours (Andrews et al., 2022; Austin & Converse, 2021; Fairbrother, 2016; Kahan et al., 2015; Merk et al., 2016), while others fear it could undermine mitigation ambition even further (Baatz, 2016; Corner & Pidgeon, 2014; Pierrehumbert, 2019; Raimi et al., 2019) and present a ‘moral hazard’ risk (Belaia et al., 2021; Bellamy et al., 2016; Burns et al., 2016; Keith, 2000; McLaren, 2016; Merk et al., 2016; Moreno-Cruz, 2015; Wibeck et al., 2015). Here, we do not investigate
70 to what extent SRM will or will not change mitigation ambition and instead highlight what various emission reduction assumptions could mean in terms of SRM deployment length. Thanks to the large dataset underlying this study, we can analyse several other factors that influence the length of SRM deployment such as the amount of annual net-negative emissions (NNEs) realised through large-scale CDR and climate system uncertainty.

There is a large body of literature dedicated to CDR and its application in mitigation pathways (Fuss et al., 2014, 2018;
75 Johansson et al., 2020; Rogelj et al., 2019; Schleussner et al., 2016). Large uncertainties regarding overshoot pathways and CDR remain, especially since the technology and the resulting temperature declines are unproven at scale (Pathak et al., 2022). Uncertainties are related to the response of the climate system to negative emissions, the state of the climate system post overshoot in general, environmental and economic side-effects of large-scale deployment, and the level to which CDR can be scaled up (Fuss et al., 2018; Matthews et al., 2020; Rogelj et al., 2019; Schleussner et al., 2016; Zickfeld et al., 2016). In this
80 study, we do not account for feasibility constraints or potential side-effects of CDR deployment, but rather explicitly assess the sensitivity of our results to any such constraints being in place. As part of our sensitivity assessment, we consider a wide range for maximum annual NNEs reaching up to 40 GtCO₂/yr (the highest potential the IPCC 6th Assessment Report (AR6) assigns to any carbon removal technology that does not interfere with ocean chemistry (IPCC, 2022)) as well as the uncertainty surrounding the temperature decline in response to net-negative emissions.

85 It is important to examine SRM deployment length in the context of climate uncertainty. In contrast to mitigation and CDR, climate uncertainty is beyond human control and, as this study demonstrates, would result in greatly differing SRM deployment outcomes for the same levels of emissions. We address climate uncertainty in two ways. First, by considering a large range of plausible climate simulators. And secondly, by calculating two climate metrics that relate emissions to temperature change and affect the duration of the overshoot and therefore SRM deployment length.

90 While many different methods for CDR (Fuss et al., 2018) and SRM (Boucher et al., 2013; Lawrence et al., 2018) exist that come with different specificities, this study does not differentiate between these specific technological approaches as our results are independent of the boundary conditions for individual SRM and CDR techniques. We acknowledge that forecasting technology this far into the future is highly speculative and this analysis is by no means intended to be a realistic representation of SRM and CDR pathways. Therefore, this paper does not address issues of feasibility or environmental side effects of SRM

95 or CDR, of which there are many (Lee et al., 2021; Canadell et al., 2021; Douville et al., 2021). Neither does it propose potential implementation strategies and designs or poses questions relating to economic, political or ethical concerns. With this contribution, we aim to provide a conceptual framework for exploring SRM deployment length in the context of scenarios that use the technology as a temporary (albeit potentially multi-century) measure.

2 Methods

100 2.1 Emissions data and pathway extension

As underlying data, we use all scenarios from the IPCCs' 6th Assessment Reports' database that are in line with the 2030 NDCs (Riahi et al., 2022) and have decreasing or stagnating emissions in the last five years of the 21st century. The policy categories we consider when identifying the scenarios that pass through NDC emissions are P3b, P1c and P0_3b (Riahi et al., 2022). This amounts to a total of 355 scenarios covering the years 2015 to 2100 that originate in a similar climatic state in 105 2030 but diverge afterwards to cover a large variety of emission levels in 2100. We employ this broad range of pathways to analyse SRM under many different developments of mitigation ambition under SRM, such as large increases and decreases, as well as under the scrutiny of climate change uncertainty by running a probabilistic ensemble of 600 members (see 2.2). To be able to estimate potential SRM timescales, we extend all pathways until 2500. In order to explore a wide range of possible future development, we randomly sample three parameters for the extension of each scenario: the change in rate of 110 decarbonisation after 2100 (0-3% increase), the maximum net-negative fossil-CO₂ emissions (log-normal distribution from 0-40 GtCO₂/yr) realised through large-scale CDR (Figure 1c) and the floor for CO₂ emissions from agriculture, forestry and other land use (AFOLU-CO₂) (normal distribution -1+1 GtCO₂/yr) (Figure 1f). We choose a log-normal distribution of 0 to 40 GtCO₂/yr with positive skew for net-fossil-CDR. The range depicts the maximum mitigation potential that the IPCC AR6 assigns to industrial carbon removal technologies (Direct Air Carbon Capture and Storage (DACCS)) while the distribution 115 with positive skew represents the tendency of limits to be at the lower rather than higher end of the chosen spectrum. The mean of the distribution is set to be 15 GtCO₂/yr, representing the rounded up median potential experts assign to DACCS in 2100 (Grant et al., 2021). The storage capacity of technologically captured CO₂ is uncertain but likely in the range of 8000 to 55000 GtCO₂ (Dooley, 2013; Kearns et al., 2017). This capacity potential does not impact our results because even the lower end is sufficient for the majority of our pathways (Figure 4b). We therefore choose no constraint on maximum negative cumulative 120 fossil-CO₂ emissions. In contrast, nature-based CDR solutions represented by AFOLU negative emissions are seriously limited in the total amount they can remove. Current literature suggests that AFOLU-CDR is a solution of the 21st century and that saturation could be hit soon after (Fuss et al., 2018). Therefore, we remove the need to make decisions about the role of AFOLU-CDR (which would be a relatively small part of the picture over the timescales we are considering) by setting the AFOLU long-term emissions level to be close but not exactly equal to 0 (-1 to 1 GtCO₂/yr in line with the Global Carbon Budget 2022 (Friedlingstein et al., 2022)). In the analysis of our results, we sum AFOLU and fossil net-negative emissions to 125 one variable which we call Net-Negative Emissions (NNEs).

The modified decarbonisation rate of the last ten years of the 21st century is linearly extrapolated until meeting the maximum net-negative fossil-CO₂ emissions for the respective pathway (Figure 1b,e). In the rare cases where scenarios have increasing AFOLU- or fossil-CO₂ emissions in the last 10 years (we only allow scenarios with stagnating or decreasing *combined* CO₂ emissions in the last few years, meaning fossil-CDR or AFOLU-CDR individually can still have increasing emissions or decreasing negative emissions in the last decade) are assigned decreasing emissions after 2100 with the rate of increasing emissions they had before plus the change in rate that is randomly prescribed (0-3%) (see uppermost lines in Figure 1b,e). We assume that saturation for nature-based carbon removal is hit in 2150 and all pathways linearly move towards their randomly chosen floor level after 2100 with a rate that allows reaching the assigned floor by 2150 (Figure 1e). For simplicity, we hold non-CO₂ emissions constant at 2100 emission levels until 2500, which gives some residual warming signal but is relatively small compared to the CO₂ contribution.




Figure 1: Extension of emission pathways consistent with NDCs from the AR6 WG3 database to 2500. A distribution of extension options is used to sample the range of possible outcomes consistent with current literature. For simplicity, in all cases, non-CO₂ emissions are kept constant after 2100. a) Fossil CO₂ emissions from selected AR6 WG3 database scenarios. b) Illustration of extension algorithm used for fossil CO₂ emissions - each pathway is extended at a fixed rate of decline until it hits some prescribed value, after which emissions are held constant (see 2.1 for full description). c) Distribution of minimum fossil CO₂ emissions used in this study. d) Land-use CO₂ emissions from selected AR6 WG3 database scenarios. e) Illustration


145 *of extension algorithm used for land-use CO₂ emissions – each pathway is extended such that it reaches a specific value by*
2150, after which emissions are held constant (see 2.1 for full description). f) Distribution of steady state land-use CO₂
emissions used in this study. g) resulting temperature trajectories, light gray covering the 5th-95th percentiles over all
scenarios with all 600 ensemble members, dark gray the 5th-95th percentile of an exemplary scenario (MESSAGEix-
GLOBIOM_1.1 EN_INDCi2030_1800f_NDCp) with the black line being the median of it. We span a wide range of trajectories,
150 *from always remaining below 1.5°C to never coming back below 1.5°C after crossing 1.5°C in the early 21st century.*

2.2 MAGICC setup and SRM-pathway construction

All global climate model simulations are conducted with the climate emulator MAGICC7.5.3 (Meinshausen et al, 2020). This
155 emulator is commonly used in several leading Integrated Assessment Models and consecutively in the IPCC assessments,
including the most recent 6th Assessment Cycle. The model includes a simplified terrestrial and ocean carbon cycle
(Meinshausen et al., 2009, 2011, 2020). We apply a probabilistic setup with an ensemble of 600 runs derived by a Markov-
Chain Monte-Carlo approach and display all ensemble members except if indicated otherwise. The range of the ensemble
members depicts the Equilibrium Climate Sensitivity uncertainty range of the IPCC 6th Assessment Report (Forster et al.,
160 2021) and the C4MIP carbon cycle ranges (Forster et al., 2021) and as a consequence offers good coverage of climate system
and model uncertainty. The ensemble members span a Transient Climate Response to Cumulative Carbon Emissions (TCRE)
range of 0.87 to 3.47 [K / 1000 PgC] (17-83% range 1.37-2.19).

The purpose of SRM in our scenario setup is to cool the temperature overshoot pathways down to global average warming of
165 1.5°C starting in 2030. Because SRM is implemented in the model by modifying the Effective Radiative Forcing (ERF), it is
necessary to determine what forcing pathway is equivalent to following a 1.5°C-compliant trajectory for each member of the
ensemble (Figure 2). This 1.5°C-trajectory represents the ‘SRM-pathway’. Due to computational efficiency and close
resemblance to WG3 NDC-pathways, we choose an SSP2-4.5 pathway as a starting point for this SRM-pathway construction
(Figure 2a). This reduces the required computing time by a factor of 10 while still retaining sufficient accuracy to make useful
170 quantifications of required SRM deployment times. Using 2030 NDC emission levels as the starting point for our analysis, we
also assume that radiative forcing can be modified earliest in 2030. Depending on the ensemble member this leads to either a
smooth approach to 1.5°C where an overshoot is avoided in the SRM-pathway (Figure 2b) or to an overshoot that is
subsequently brought down to 1.5°C (Figure 2c). Whether the SRM-pathway for a given ensemble member overshoots is
determined by its SSP2-4.5 2035 warming level had it followed its 10-year gradient from 2030 to 2040 for five more years
175 after 2030. If this 2035 warming level is higher than 1.5°C, the pathway will overshoot to the respective 2035 level and
subsequently descend to 1.5°C in a sigmoidal pattern to reach 1.5°C in 2050.

Since each of the 355 scenarios is prescribed different assumptions (see 2.1) and each ensemble member in MAGICC's probabilistic distribution has different physics, we must calculate the SRM on a scenario-ensemble member basis where the SRM required is the difference between the 1.5°C-pathway and the extended NDC-scenario without SRM (NDC-extension described in 2.1). The start date of SRM is determined by the date where the extended NDC-scenario exceeds the designed SRM-pathway for the respective ensemble member. SRM termination is assumed to happen once 1.5°C global mean warming is reattained in the NDC-scenario.



185 **Figure 2: Calculating required SRM.** a) Calculating warming to 2035 using an NDC-like pathway (in this case SSP2-4.5). b) Determining a 1.5°C temperature trajectory for ensemble members that have not already overshoot 1.5°C by 2035. c) Determining a 1.5°C temperature trajectory for ensemble members that have already overshoot 1.5°C by 2035. d), e) Calculating required solar radiation manipulation (SRM) for each scenario-ensemble member combination, whether it overshoots 1.5°C (e) or remains below 1.5°C at all times (d).

190

To address climate uncertainty and depict the whole range of possible outcomes, we use all members of our ensemble instead of only focusing on medians. Additionally, we calculate the uncertainty surrounding the rise in temperature for a specific amount of CO₂ emissions and non-CO₂ forcing, called effective Transient Response to Cumulative CO₂ Emissions (eTCRE,

195 subsequently called eTCRE-up) (Gregory et al., 2009; Matthews et al., 2009, 2020; Vakilifard et al., 2022). We also consider uncertainty in the temperature change as a result of reducing the concentrations of CO₂ and associated non-CO₂ gases in the atmosphere (subsequently called eTCRE-down). Due to uncertain Earth system feedbacks such as a lagged ocean response, it is possible that eTCRE-up and eTCRE-down are not identical (Matthews et al., 2020; Zickfeld et al., 2016) and this asymmetry is reflected in the MAGICC ensemble. eTCRE-up informs whether temperature targets are exceeded for a specific amount of cumulative emissions and therefore whether SRM is deployed in our scenarios or not, as well as which peak warming is hit when. eTCRE-down informs us on how effective NNEs are at cooling, meaning how many cumulative NNEs we need and how long it takes to return to our target temperature increase of 1.5°C.

We define eTCRE-up and eTCRE-down as:

$$(1) \ eTCRE_{up} = \frac{\Delta T_{2030 \rightarrow peak_warming}}{\Sigma_{2030}^{peak_warming} CO_2 \ emissions} [K \ 1000PgC^{-1}]$$

205

$$(2) \ eTCRE_{down} = \frac{\Delta T_{1.5^\circ C \rightarrow peak_warming}}{\Sigma_{peak_warming}^{year_of_return_to_1.5^\circ C} CO_2 \ emissions} [K \ 1000PgC^{-1}]$$

While the change in temperature is related to both CO₂ and non-CO₂ gases, the cumulative emissions are only related to CO₂. Due to a few extreme outliers both metrics are constrained to their 1-99th percentile.

210

3 Results

The 355 NDC-scenarios including extensions and probabilistic MAGICC7 simulations lead to 213 000 different realisations with a large range of warming outcomes throughout the centuries (Figure 1g). While few realisations peak at or below 1.5°C, the vast majority of the baseline simulations without SRM overshoot the temperature target temporarily. Others do not return back to 1.5°C before 2500 at all. The 5-95th percentile range of peak-warming in our dataset ranges from 1.60°C to 3.91°C with the median at 2.14°C. Therefore, even though all realisations originate in 2030-NDC levels, due to the range of possible developments of mitigation and NNEs under SRM, the resulting SRM deployment length ranges from 0 to >470 years (Figure 3), with 5% of realisations not requiring SRM to limit warming to 1.5°C and 15% having deployment times that exceed 470 years. In the following analysis we only include realisations that fall into the 1-470 year range.

220




Figure 3: SRM deployment length for all scenarios and all ensemble members. 1 bar spans a range of 50 years. Marked in black are pathways consistent with current 2100-warming projections for NDCs (2.4°C) (CAT, 2022).

225

Figure 4 breaks down the three indicators mitigation, negative emissions and climate uncertainty and their interlinkages with SRM deployment length. The relationship between the emission pathway, i.e. the cumulative emissions from 2030 until net-zero, and SRM deployment length is weak (Figure 4a). The triangular shape of the data points towards a tendency for very high cumulative emissions (>3000 GtCO₂ at time of net-zero) to lead to longer SRM timescales, whereas scenarios with lower cumulative emissions are spread across the whole range of SRM deployment times with a skew towards the shorter end. Realisations above 1500 GtCO₂ (apart from a few exceptions) require at least a few decades of deployment to keep warming at 1.5°C, no matter the climate uncertainty. The plot looks stratified because the cumulative emissions value is member-independent and therefore stays equal across all ensemble-members of each scenario.

230

235

The way our scenarios are set up, the choice of maximum potential NNEs is random and therefore independent of the amount of cumulative emissions. However, *average* annual NNEs are not entirely arbitrarily spread: shorter deployment timescales are constrained to lower annual NNEs because of the limited time these scenarios have to scale up their net-negative emissions. Similarly, high emission scenarios take longer to reach high net-negative emissions. This is demonstrated by scenarios with cumulative emissions above 3500 GtCO₂ that almost all reach annual NNEs above 15 GtCO₂/yr by 2500 (see Figure S01 in S.I.) but most never reach this level in average terms because of the time required to scale up.

240


Focussing on the interlinkages with negative emissions, Figure 4b shows cumulative emissions between net-zero and reattaining 1.5°C against SRM deployment. The relationship is much clearer and the triangle shape of Figure 4b considerably more defined; low cumulative negative emissions are represented across the whole range of SRM deployment length but high cumulative negative emissions are constrained to long SRM timeframes. For example, cumulative negative emission requirements above -2000 GtCO₂ imply more than 100 years of SRM deployment. Similarly, -6000 GtCO₂ lead to more than 200 years of SRM deployment and -10000 GtCO₂ to more than 300 years. Part of this effect is the simple fact that you only get very large amounts of cumulative negative emissions if you're deploying SRM for long time periods.

245

Where in Figure 4a the annual average NNEs are partly random, a clear patterning becomes visible in 4b. The higher the annual average NNEs the shorter the SRM timescale for the same amount of cumulative emissions. For higher total negative emission

250 requirements low NNEs are not sufficient in limiting the deployment to 470 years and are thus not shown. Very small amounts of NNEs ($< 1 \text{ GtCO}_2/\text{yr}$) are constrained to pathways with no or very small amounts of required negative emissions to get down to 1.5°C of warming.

Regarding climate uncertainty, the calculated 5-95th percentile range of eTCRE-up is $1.0 - 6.0 \text{ [K/1000PgC]}$, eTCRE-down is
255 $-6.9 - 12.4 \text{ [K/1000PgC]}$. Figure 4c sets the eTCRE-ratio (defined as eTCRE-up / eTCRE-down) against the SRM timeframes. The spread in uncertainty increases with SRM deployment length. Realisations with an absolute eTCRE-ratio of >1 have a lower sensitivity to negative emissions than positive ones, causing them to deploy SRM for longer periods even though net-zero is hit earlier than for other scenarios (see Figure S02 S.I.). Negative eTCRE-ratios (dark purple data points) are the result of negative eTCRE-down values, all of which arise in realisations that have positive cumulative emissions in the cooling phase.
260 This could be due to a high negative Zero Emissions Commitment (ZEC) (Jenkins et al., 2022; C. D. Jones et al., 2019; MacDougall et al., 2020) or non-CO₂ effects that allow temperatures to drop despite cumulative positive CO₂ emissions.



265 *Figure 4: Interdependencies of mitigation, negative emissions and climate uncertainty with SRM deployment length a) Relationship between cumulative CO₂ emissions from 2030 until net-zero CO₂ and SRM deployment length. Colour-coding according to annual average NNEs in GtCO₂/yr. b) Relationship between cumulative CO₂ emissions from net-zero CO₂ until reattaining 1.5°C and SRM deployment length. Colour-coding according to annual average NNEs in GtCO₂/yr. c) Relationship between eTCRE-ratio and SRM deployment length. Colour-coding according to cumulative CO₂ emissions from time of peak-warming until reattaining 1.5°C. Plot shows data points that fall in the 1-99th percentile range.*

270 4 Discussion

In current literature, SRM is often framed in the context of a stopgap measure (Asayama & Hulme, 2019; Buck et al., 2020; Neuber & Ott, 2020). Our study focuses on the question of what range of timeframes would be consistent with an intended temporary SRM deployment due to uncertainty in mitigation ambition, negative emissions and climate uncertainty.


We show that the range of possible deployment timescales is vast even for pathways that have similar conditions at the start
275 of SRM deployment, in our case in 2030 (Figure 3). This is due to the uncertain evolution of mitigation ambition and annual NNEs under SRM and the effects of climate uncertainty. We find that neither of these three indicators (mitigation, net-negative emissions or climate uncertainty) alone can determine SRM deployment length. Mitigation and negative emissions represent a bounding condition on SRM deployment length rather than a linear correlation, to some degree due to the climate sensitivity of increasing and decreasing emissions (Figure 4c). This means that positive emissions after SRM initialisation on the medium
280 to lower end could imply both short and long term SRM deployment (Figure 4a). Similarly, low cumulative negative emission requirements are no guarantee for short SRM timeframes (Figure 4b). However, the faster and higher annual NNEs are scaled up the shorter the time-wise commitment to SRM (Figure 4b, 5a). Our data suggest that for the range of emission pathways, NNEs and in particular the limited analysis length of 470 years in our experiments, it might be easier to determine a lower limit of deployment length than an upper limit.

285 The relationship between the eTCRE-ratio and SRM deployment length versus cumulative emissions that results from our data is complex and requires further study (Figure 4c, 5b). Several aspects play into the variables used to calculate the two climate uncertainty metrics. For example, the temperature change in the eTCRE calculation is related to CO₂ as well as non-CO₂ gases while the cumulative emissions are only a function of CO₂. Additionally, the experimental setup of this study is such that CO₂ and non-CO₂ emissions are not linearly related: Non-CO₂ emissions vary between 2030 and 2100 and then stay constant at the
290 2100 level while CO₂ emissions fall consistently after 2100.

Our calculated eTCRE-up is higher than the pure range of TCRE that the MAGICC7 ensemble members have, i.e. the TCRE range in the AR6 of the IPCC (0.87-3.47). This is expected and has been observed before because of the impact of non-CO₂ climate forcings (Matthews et al., 2020, 2021). Assumptions regarding the constant ratio of cumulative emissions of CO₂ to temperature (TCRE) are suggested to go up to at least 3000 PgC of positive cumulative CO₂ emissions (Leduc et al., 2015;
295 Tokarska et al., 2016). Our values exceed this limit. However, this might not be directly transferable to our eTCRE-up metric especially since these values are expectedly higher than TCRE (Matthews et al., 2020). Studies with intermediate complexity climate models have found a higher TCRE-up than TCRE-down, suggesting a greater impact of positive CO₂ emissions on temperatures than negative emissions (Zickfeld et al., 2016). This can be explained by a lagged ocean response that leads to continued warming after the start of carbon removal (Zickfeld et al., 2016). However, ocean effects might differ in cases where

300 SRM is deployed and complex earth system models show a range of responses after emissions are halted (Jenkins et al., 2022; MacDougall et al., 2020), let alone negative for a sustained period.

The inconclusive relationship of the eTCRE-ratio and negative emissions against SRM deployment length (Figure 5) emphasises the fact that while ending SRM hinges fundamentally on the decline of temperatures (when SRM is used for the purposes described in this paper), climate reversibility is currently marked by large uncertainties.



305

Figure 5: Relationship between cumulative CO₂ emissions from net-zero CO₂ until reattaining 1.5°C and SRM deployment length. a) Colour-coded by the maximum deployed annual NNEs. b) Colour-coded by the eTCRE-ratio. Clean figure without descriptions in S.I (Figure S03).

310 The shape of the triangle in Figure 4a,b and Figure 5a,b is partly due to the effects of climate uncertainty and partly due to the experimental setup with a limit of yearly NNEs at 40 GtCO₂/yr. However, the area of the plot with short SRM deployment under around a century would not be covered even for higher amounts of annual NNEs because the scaling up requires time. To contextualise, a yearly removal of 40 GtCO₂ equates to today's yearly fossil CO₂ emissions (Friedlingstein et al., 2022) and would entail a massive industrial effort by itself, which would take decades to build up. So even though very high NNEs
 315 could shorten SRM timescales, it likely would not be available for short SRM deployment. In those cases, the best bet is to

require a small amount of total net-negative emissions to return back down to 1.5°C. This would imply low cumulative positive emissions and a high eTCRE-down.

The upper limit of NNEs goes to the heart of concerns around CDR: the magnitude of yearly removal to which it can be scaled up. Because it is impossible to predict technological development this far into the future we reflect all plausible options by assuming NNEs that range from barely being able to compensate residual positive emissions, to being consistent with current sustainable levels, to much larger amounts that future technological development could enable but far exceed many estimates in current literature (Fuss et al., 2018; Grant et al., 2021; Coninck et al., 2018; Pathak et al., 2022). It needs to be highlighted that this study looks at net-negative emissions and not CDR. This means that our estimates of net-negative emissions are a lower bound on the amount of CDR needed. There is a broad discussion in current literature on negative side effects and sustainability concerns with CDR in mitigation scenarios (Brack & King, 2020; Fuss et al., 2018; Smith et al., 2015) as well as the question of equity in the allocation of CDR burden (Fyson et al., 2020). Major concerns are the considerable land, water, energy and financial requirements and constraints in long-term storage of removed CO₂ that increase for higher yearly removal rates. These recognized environmental and social concerns of CDR are at least as applicable for pathways where CDR exceeds current sustainable ranges and needs to be sustained for decades up to centuries. This paper quantifies the importance of decades-long large-scale CDR for ‘peak-shaving’ SRM scenarios, which has previously been pointed out by Asayama & Hulme, 2019 and others. In addition, we find that initialisation and commitment to SRM happen under the promise that CDR can be scaled up high enough to end SRM again. Belaia et al., 2021 even demonstrate that a cost-optimal portfolio to meet 2°C GSAT increase would consist of SRM and mitigation first and later CDR deployment. With CDR being unproven at scale and peak-and-decline scenarios marked by large uncertainties, this is a risky promise. Our scenarios show what happens if the scalability of CDR is not as high as assumed or TCRE-down is low: SRM deployment will have to be extended for a much longer period of time (Figure 4b).

In scenarios where the level of warming in 2100 follows the level of mitigation ambition reflected in 2030 NDC commitments until the end of the century (CAT, 2022), no pathway deploys SRM for a shorter period than 100 years, while most require 150-300 years in order to return to 1.5°C GSAT increase (black bars in Figure 3; 25-75% percentile range 159-294 years, excluding all pathways that exceed 470 years in the statistic). Even for scenarios with a somewhat smaller overshoot above 1.5°C potential deployment length remains considerable. Half of all scenario realisations that would peak around 1.8 °C require SRM deployment for 96-195 years to stay at 1.5°C. Therefore, ‘short’ peak-shaving SRM deployment perhaps implies a length of the order of 100-200 years rather than 10-50. These timeframes are much longer than the hotly debated CDR deployment lengths in GHG emission reduction pathways for limiting warming to 1.5°C in 2100 after a high or no/limited overshoot assessed in the IPCC SR1.5 (Rogelj et al., 2018) or IPCC AR6 WG3 (Riahi et al., 2022).

MacMartin et al. (2018) created an exemplary scenario where emissions would lead to a peak warming of 2.7°C without SRM and deploy 15 Gt CDR per year. Their scenario requires around 235 years of SRM deployment to limit warming to 1.5°C. Constraining our dataset to their benchmark data results in an SRM deployment of 99-404 years with a mean at 221 - a range

that encompasses their result with a mean that is not far off from their estimate. Tilmes et al. (2016) constructed a scenario that peaks at 3°C and uses SRM to cool down to 2.0°C with 18.5 GtCO₂ CDR per year. Simulated with the CESM Earth System Model they indicate a deployment time of 160 years to compensate the overshoot over 2°C. While these results are not fully comparable, compensating 1°C of overshoot with NNEs of 18.5 GtCO₂ per year in our scenarios results in an average deployment length of 204 years and a full range of 98-393 years. Our results thus compare well with previous explorations of this question documented in the literature, yet provide a much more holistic perspective on the uncertainties involved. As we demonstrate, these are just few of the many possible future outcomes of SRM deployment and including climate uncertainty shows where the limits of our control lie.

The challenge in defining SRM deployment length at its initialisation poses a clear risk as there are several issues that might arise with a multi-century SRM deployment: One of the arguably biggest difficulties of SRM deployment is international cooperation and coordination (Möller, 2020; Reynolds, 2019; Shepherd et al., 2009). Maintaining international cooperation will be even more difficult if deployment needs to be sustained over many decades as priorities and interests of countries and their leaders might change. Furthermore, long-lasting SRM deployment would require substantial financial resources; due to the difficulty in predicting SRM deployment length as pointed out in this study, costs could end up being much higher than originally anticipated if deployment ends up being longer than planned. This needs to be put into context with the overall cost of responding to climate change and avoided costs of climate change damages through SRM (Arino et al., 2016; Belaia et al., 2021). Moreover, a key risk of SRM is the so-called ‘termination shock’, a rapid warming response to a sudden stop of SRM deployment (Parker & Irvine, 2018). The longer the deployment, the longer the risk for such an occurrence. And lastly, long-term deployment would bind future generations to SRM which raises substantial ethical and moral questions (Flegal et al., 2019; Goeschl et al., 2013; Svoboda et al., 2011). The idea of imposing a technology on individuals who have not yet been born and do not have a say in the matter can be considered to be a violation of their rights and autonomy.

In addition to all the risks, costs and potential side-effects that come with SRM, due to the dependency, future generations would be burdened by large-scale deployment of CDR which might compete with efforts to secure their own requirements by increasing risks of biodiversity loss and food and water scarcity (Dooley & Kartha, 2018; Shue, 2017; Asayama & Hulme, 2019; Svoboda et al., 2011). With this study we add quantitative data to the literature calling for precautionary and ethical approaches to technology development. The identified risks need to be weighed against the risk of not deploying SRM. Our research highlights the substantial dependencies of SRM and CDR deployment which imply side-effects, risks, costs and uncertainties of both, SRM and CDR.

In this analysis, we use SRM for the prevention of overshooting the 1.5°C-target and the deployment lengths we indicate only hold under the stated conditions of the analysis. Possible avenues for prior phase-outs have been discussed in Keith & MacMartin (2015), MacMartin et al. (2018) and Parker & Irvine (2018). However, these avenues would violate the 1.5°C-

target and, as suggested by Parker & Irvine (2018), depending on the amount of Wm^{-2} compensated could take several decades. Therefore, even if SRM was, due to technical and / or environmental considerations, phased out earlier, the phase-out itself could become a multi-decadal undertaking in itself.

385 Being a reduced-complexity model, MAGICC7 has its caveats and constraints related to the physical and spatial resolution of relevant climate and carbon cycle processes. Nevertheless, MAGICC has been used successfully in many instances to robustly analyse long-term perspectives (Meinshausen et al., 2020; Meinshausen, Smith, et al., 2011; Nauels et al., 2017). Therefore, we consider it also appropriate for this first quantification of hypothetical SRM deployment length which could motivate further study in more complex models.

It is important to highlight that even if global temperature was stabilised with the help of SRM, this will not provide a solution with respect to regional impacts (A. C. Jones et al., 2018) and other impacts of high GHG concentration levels such as ocean acidification (Tjiputra et al., 2016). In our study, however, we do not aim to provide a comprehensive analysis of impacts or side-effects of such a climate intervention nor do we provide a likely or desirable implementation strategy but rather explore a concept to determine hypothetical SRM deployment lengths.

395 5 Conclusions

In this paper, we generate a large dataset of pathways to analyse SRM deployment length in scenarios that are consistent with 2030 NDC levels and use SRM to cool down to 1.5°C of warming. We find a large spread of SRM deployment lengths, ranging from no NNE and SRM requirements at all to massive NNE and SRM requirements past 2500 to limit global warming to 1.5°C . Most of our simulations require around 150-300 years of SRM. Deployment timeframes are considerably dependent on mitigation, negative emissions and climate uncertainty yet none of these factors alone can predict its length: Large cumulative positive emissions lead to long SRM deployments. However, small cumulative emissions are no guarantee for short deployment timescales. Similarly, realisations that require large cumulative negative emissions lead to long SRM deployments. But, small cumulative negative emission requirements do not necessarily imply short SRM timeframes. A large part of this uncertainty can be attributed to the uncertainty surrounding eTCRE-up and eTCRE-down.

405 For all realisations that follow current NDC median 2100-warming projections (2.4°C), none deploy SRM for a shorter period than 100 years even under the most optimistic assumptions on the eTCRE-ratio. For SRM deployment to truly be ‘temporary’, carbon removal technologies need to be massively scaled up within a relatively short timeframe, except in cases of very low emission requirements and extremely high negative ZEC. Larger average annual NNEs do generally imply shorter SRM timescales.

410 Our study shows that the range of possible deployment timescales is vast even if pathways start at a similar point at the beginning of SRM deployment because the evolution of mitigation under SRM, the availability of carbon removal technologies

and the effects of climate reversibility are not precisely known. Since these effects will be mostly uncertain at the time of SRM initialisation, a precedent prediction of deployment length seems unlikely with possibilities ranging from decades to multiple centuries. This is a knowledge gap that must be considered before any SRM proposal is seriously considered.

415

Code and data availability statement

420 The code is available at <https://gitlab.com/susannebaur/deployment-length-srm>. Upon final publication the code will be made publicly available with a DOI under zenodo.

Author Contribution

425 All authors designed the experiments and SB and ZN carried them out. SB prepared the manuscript with contributions from all authors.

Ethics Declaration

The authors declare that they have no conflict of interest.

Acknowledgments

430 Funding was provided by the European Union's Horizon 2020 Research and Innovation Programme under grant nos. 820829 (CONSTRAIN) and 101003687 (PROVIDE).

References

- Allen, M. R., Dube, O. P., Solecki, W., Aragón-Durand, F., Cramer, W., Humphreys, S., Kainuma, M., Kala, J., Mahowald, N., Mulugetta, Y., Perez, R., Wairiu, M., & Zickfeld, K. (2018). Framing and Context. In *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change*, (Vol. 2, Issue October).
- Andrews, T. M., Delton, A. W., & Kline, R. (2022). Anticipating moral hazard undermines climate mitigation in an experimental geoengineering game. *Ecological Economics*, 196, 107421. <https://doi.org/10.1016/j.ecolecon.2022.107421>
- Arino, Y., Akimoto, K., Sano, F., Homma, T., Oda, J., & Tomoda, T. (2016). Estimating option values of solar radiation management assuming that climate sensitivity is uncertain. *Proceedings of the National Academy of Sciences*, 113(21), 5886–5891. <https://doi.org/10.1073/pnas.1520795113>
- Asayama, S., & Hulme, M. (2019). Engineering climate debt: Temperature overshoot and peak-shaving as risky subprime mortgage lending. *Climate Policy*, 19(8), 937–946. <https://doi.org/10.1080/14693062.2019.1623165>
- Austin, M. M. K., & Converse, B. A. (2021). In search of weakened resolve: Does climate-engineering awareness decrease individuals' commitment to mitigation? *Journal of Environmental Psychology*, 78, 101690. <https://doi.org/10.1016/j.jenvp.2021.101690>
- Baatz, C. (2016). Can we have it both ways? On potential trade-offs between mitigation and solar radiation management. *Environmental Values*, 25(1), 29–49. <https://doi.org/10.3197/096327115X14497392134847>
- Belaia, M., Moreno-Cruz, J. B., & Keith, D. W. (2021). OPTIMAL CLIMATE POLICY IN 3D: MITIGATION, CARBON REMOVAL, AND SOLAR GEOENGINEERING. *Climate Change Economics*, 12(03), 2150008. <https://doi.org/10.1142/S2010007821500081>
- Bellamy, R., Chilvers, J., & Vaughan, N. E. (2016). Deliberative Mapping of options for tackling climate change: Citizens and specialists 'open up' appraisal of geoengineering. *Public Understanding of Science*, 25(3), 269–286. <https://doi.org/10.1177/0963662514548628>
- Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V. M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S. K., Sherwood, S., Stevens, B., & Zhang, X. Y. (2013). Clouds and aerosols. *Climate Change 2013 the Physical Science Basis: Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, 9781107057, 571–658. <https://doi.org/10.1017/CBO9781107415324.016>
- Brack, D., & King, R. (2020). Managing Land-based CDR: BECCS, Forests and Carbon Sequestration: Managing Land-based CDR. *Global Policy*, 12. <https://doi.org/10.1111/1758-5899.12827>
- Buck, H. J., Martin, L. J., Geden, O., Kareiva, P., Koslov, L., Krantz, W., Kravitz, B., Noël, J., Parson, E. A., Preston, C. J., Sanchez, D. L., Scarlett, L., & Talati, S. (2020). Evaluating the efficacy and equity of environmental stopgap measures. *Nature Sustainability*, 3(7), 499–504. <https://doi.org/10.1038/s41893-020-0497-6>
- Burns, E. T., Flegal, J. A., Keith, D. W., Mahajan, A., Tingley, D., & Wagner, G. (2016). What do people think when they think about solar geoengineering? A review of empirical social science literature, and prospects for future research. *Earth's Future*, 4(11), 536–542. <https://doi.org/10.1002/2016EF000461>
- Canadell, J.G., P.M.S. Monteiro, M.H. Costa, L. Cotrim da Cunha, P.M. Cox, A.V. Eliseev, S. Henson, M. Ishii, S. Jaccard, C. Koven, A. Lohila, P.K. Patra, S. Piao, J. Rogelj, S. Syampungani, S. Zaehle, and K. Zickfeld, 2021: Global Carbon and other Biogeochemical Cycles and Feedbacks. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M.

Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 673–816, doi: 10.1017/9781009157896.007.

- 480 CAT (Climate Action Tracker) (2022). 2100 Warming Projections: Emissions and expected warming based on pledges and current policies. November 2022. Available at: <https://climateactiontracker.org/global/temperatures/>. Copyright ©2022 by Climate Analytics and NewClimate Institute. All rights reserved.
- Coninck, H., Revi, A., Babiker, M., Bertoldi, P., Buckeridge, M., Cartwright, A., Dong, W., Ford, J., Fuss, S., Hourcade, J.-C., Ley, D., Mechler, R., Newman, P., Revokatova, A., Schultz, S., Steg, L., & Sugiyama, T. (2018). Chapter 4 - Strengthening and implementing the global response. In: Global warming of 1.5°C. 313–443. https://www.ipcc.ch/site/assets/uploads/sites/2/2018/11/SR15_Chapter4_Low_Res.pdf
- 485 Corner, A., & Pidgeon, N. (2014). Geoengineering, climate change scepticism and the ‘moral hazard’ argument: An experimental study of UK public perceptions. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 372(2031), 20140063. <https://doi.org/10.1098/rsta.2014.0063>
- 490 Dooley, J. J. (2013). Estimating the supply and demand for deep geologic CO₂ storage capacity over the course of the 21st century: A meta-analysis of the literature. *Energy Procedia*, 37, 5141–5150. <https://doi.org/10.1016/j.egypro.2013.06.429>
- Dooley, K., & Kartha, S. (2018). Land-based negative emissions: Risks for climate mitigation and impacts on sustainable development. *International Environmental Agreements: Politics, Law and Economics*, 18(1), 79–98. doi:10.1007/s10784-017-9382-9
- 495 Douville, H., K. Raghavan, J. Renwick, R.P. Allan, P.A. Arias, M. Barlow, R. Cerezo-Mota, A. Cherchi, T.Y. Gan, J. Gergis, D. Jiang, A. Khan, W. Pokam Mba, D. Rosenfeld, J. Tierney, and O. Zolina, 2021: Water Cycle Changes. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 1055–1210, doi: 10.1017/9781009157896.010.
- 500 Fairbrother, M. (2016). Geoengineering, moral hazard, and trust in climate science: Evidence from a survey experiment in Britain. *Climatic Change*, 139(3–4), 477–489. <https://doi.org/10.1007/s10584-016-1818-7>
- Flegal, J. A., Hubert, A. M., Morrow, D. R., & Moreno-Cruz, J. B. (2019). Solar Geoengineering: Social Science, Legal, Ethical, and Economic Frameworks. *Annual Review of Environment and Resources*, 44, 399–423. <https://doi.org/10.1146/annurev-environ-102017-030032>
- 505 Forster, P., T. Storelvmo, K. Armour, W. Collins, J.-L. Dufresne, D. Frame, D.J. Lunt, T. Mauritsen, M.D. Palmer, M. Watanabe, M. Wild, and H. Zhang, 2021: The Earth’s Energy Budget, Climate Feedbacks, and Climate Sensitivity. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 923–1054, doi:10.1017/9781009157896.009.
- 510 Friedlingstein, P., O’Sullivan, M., Jones, M. W., Andrew, R. M., Gregor, L., Hauck, J., Le Quéré, C., Lujikx, I. T., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Alkama, R., ... Zheng, B. (2022). Global Carbon Budget 2022. *Earth System Science Data*, 14(11), 4811–4900. <https://doi.org/10.5194/essd-14-4811-2022>
- 515

- 520 Fuss, S., Canadell, J. G., Peters, G. P., Tavoni, M., Andrew, R. M., Ciais, P., Jackson, R. B., Jones, C. D., Kraxner, F., Nakicenovic, N., Le Quéré, C., Raupach, M. R., Sharifi, A., Smith, P., & Yamagata, Y. (2014). Betting on negative emissions. *Nature Climate Change*, 4(10), Article 10. <https://doi.org/10.1038/nclimate2392>
- Fuss, S., Lamb, W. F., Callaghan, M. W., Hilaire, J., Creutzig, F., Amann, T., Beringer, T., De Oliveira Garcia, W., Hartmann, J., Khanna, T., Luderer, G., Nemet, G. F., Rogelj, J., Smith, P., Vicente, J. V., Wilcox, J., Del Mar Zamora Dominguez, M., & Minx, J. C. (2018). Negative emissions—Part 2: Costs, potentials and side effects. *Environmental Research Letters*, 13(6). <https://doi.org/10.1088/1748-9326/aabf9f>
- 525 Fyson, C. L., Baur, S., Gidden, M., & Schleussner, C. F. (2020). Fair-share carbon dioxide removal increases major emitter responsibility. *Nature Climate Change*, 10(9), 836–841. <https://doi.org/10.1038/s41558-020-0857-2>
- Goeschl, T., Heyen, D., & Moreno-Cruz, J. (2013). The Intergenerational Transfer of Solar Radiation Management Capabilities and Atmospheric Carbon Stocks. *Environmental and Resource Economics*, 56(1), 85–104. <https://doi.org/10.1007/s10640-013-9647-x>
- 530 Grant, N., Hawkes, A., Mittal, S., & Gambhir, A. (2021). The policy implications of an uncertain carbon dioxide removal potential. *Joule*, 5(10), 2593–2605. <https://doi.org/10.1016/j.joule.2021.09.004>
- Gregory, J. M., Jones, C. D., Cadule, P., & Friedlingstein, P. (2009). Quantifying Carbon Cycle Feedbacks. *Journal of Climate*, 22(19), 5232–5250. <https://doi.org/10.1175/2009JCLI2949.1>
- 535 IPCC, 2021: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2391 pp. doi:10.1017/9781009157896.
- 540 IPCC, 2022: Climate Change 2022: Mitigation of Climate Change. Contribution of Working Group III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [P.R. Shukla, J. Skea, R. Slade, A. Al Khourdajie, R. van Diemen, D. McCollum, M. Pathak, S. Some, P. Vyas, R. Fradera, M. Belkacemi, A. Hasija, G. Lisboa, S. Luz, J. Malley, (eds.)]. Cambridge University Press, Cambridge, UK and New York, NY, USA. doi: 10.1017/9781009157926
- Irvine, P. J., Kravitz, B., Lawrence, M. G., & Muri, H. (2016). An overview of the Earth system science of solar geoengineering. *Wiley Interdisciplinary Reviews: Climate Change*, 7(6), 815–833. <https://doi.org/10.1002/wcc.423>
- 545 Jenkins, S., Sanderson, B., Peters, G., Frölicher, T. L., Friedlingstein, P., & Allen, M. (2022). The Multi-Decadal Response to Net Zero CO₂ Emissions and Implications for Emissions Policy. *Geophysical Research Letters*, 49(23). <https://doi.org/10.1029/2022GL101047>
- 550 Johansson, D. J. A., Azar, C., Lehtveer, M., & Peters, G. P. (2020). The role of negative carbon emissions in reaching the Paris climate targets: The impact of target formulation in integrated assessment models. *Environmental Research Letters*, 15(12), 124024. <https://doi.org/10.1088/1748-9326/abc3f0>
- Jones, A. C., Hawcroft, M. K., Haywood, J. M., Jones, A., Guo, X., & Moore, J. C. (2018). Regional Climate Impacts of Stabilizing Global Warming at 1.5 K Using Solar Geoengineering. *Earth's Future*, 6(2), 230–251. <https://doi.org/10.1002/2017EF000720>
- 555 Jones, C. D., Frölicher, T. L., Koven, C., MacDougall, A. H., Matthews, H. D., Zickfeld, K., Rogelj, J., Tokarska, K. B., Gillett, N. P., Ilyina, T., Meinshausen, M., Mengis, N., Séférian, R., Eby, M., & Burger, F. A. (2019). The Zero Emissions Commitment Model Intercomparison Project (ZECMIP) contribution to C4MIP: Quantifying committed climate changes following zero carbon emissions. *Geoscientific Model Development*, 12(10), 4375–4385. <https://doi.org/10.5194/gmd-12-4375-2019>

- 560 Kahan, D. M., Jenkins-Smith, H., Tarantola, T., Silva, C. L., & Braman, D. (2015). Geoengineering and Climate Change Polarization: Testing a Two-Channel Model of Science Communication. *The ANNALS of the American Academy of Political and Social Science*, 658(1), 192–222. <https://doi.org/10.1177/0002716214559002>
- Kearns, J., Teletzke, G., Palmer, J., Thomann, H., Kheshgi, H., Chen, Y.-H. H., Paltsev, S., & Herzog, H. (2017). Developing a Consistent Database for Regional Geologic CO₂ Storage Capacity Worldwide. *Energy Procedia*, 114, 4697–4709. <https://doi.org/10.1016/j.egypro.2017.03.1603>
- 565 Keith, D. W. (2000). Geoengineering the climate: History and Prospect. *Annu. Rev. Energy Environ.*, 25:245–84.
- Keith, D. W., & MacMartin, D. G. (2015). A temporary, moderate and responsive scenario for solar geoengineering. *Nature Climate Change*, 5(3), 201–206. <https://doi.org/10.1038/nclimate2493>
- Lawrence, M. G., Schäfer, S., Muri, H., Scott, V., Oeschies, A., Vaughan, N. E., Boucher, O., Schmidt, H., Haywood, J., & Scheffran, J. (2018). Evaluating climate geoengineering proposals in the context of the Paris Agreement temperature goals. *Nature Communications*, 9(1). <https://doi.org/10.1038/s41467-018-05938-3>
- 570 Leduc, M., Matthews, H. D., & Elía, R. de. (2015). Quantifying the Limits of a Linear Temperature Response to Cumulative CO₂ Emissions. *Journal of Climate*, 28(24), 9955–9968. <https://doi.org/10.1175/JCLI-D-14-00500.1>
- Lee, J.-Y., J. Marotzke, G. Bala, L. Cao, S. Corti, J.P. Dunne, F. Engelbrecht, E. Fischer, J.C. Fyfe, C. Jones, A. Maycock, J. Mutemi, O. Ndiaye, S. Panickal, and T. Zhou, 2021: Future Global Climate: Scenario-Based Projections and Near-Term Information. In Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 553–672, doi:10.1017/9781009157896.006.
- 575 MacDougall, A. H., Frölicher, T. L., Jones, C. D., Rogelj, J., Matthews, H. D., Zickfeld, K., Arora, V. K., Barrett, N. J., Brovkin, V., Burger, F. A., Eby, M., Eliseev, A. V., Hajima, T., Holden, P. B., Jeltsch-Thömmes, A., Koven, C., Mengis, N., Menviel, L., Michou, M., ... Ziehn, T. (2020). Is there warming in the pipeline? A multi-model analysis of the Zero Emissions Commitment from CO₂. *Biogeosciences*, 17(11), 2987–3016. <https://doi.org/10.5194/bg-17-2987-2020>
- 585 MacMartin, D. G., Ricke, K. L., & Keith, D. W. (2018). Solar geoengineering as part of an overall strategy for meeting the 1.5°C Paris target. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 376(2119). <https://doi.org/10.1098/rsta.2016.0454>
- 590 Matthews, H. D., Gillett, N. P., Stott, P. A., & Zickfeld, K. (2009). The proportionality of global warming to cumulative carbon emissions. *Nature*, 459(7248), Article 7248. <https://doi.org/10.1038/nature08047>
- Matthews, H. D., Tokarska, K. B., Nicholls, Z. R. J., Rogelj, J., Canadell, J. G., Friedlingstein, P., Frölicher, T. L., Forster, P. M., Gillett, N. P., Ilyina, T., Jackson, R. B., Jones, C. D., Koven, C., Knutti, R., MacDougall, A. H., Meinshausen, M., Mengis, N., Séférian, R., & Zickfeld, K. (2020). Opportunities and challenges in using remaining carbon budgets to guide climate policy. *Nature Geoscience*, 13(12), 769–779. <https://doi.org/10.1038/s41561-020-00663-3>
- 595 Matthews, H. D., Tokarska, K. B., Rogelj, J., Smith, C. J., MacDougall, A. H., Haustein, K., Mengis, N., Sippel, S., Forster, P. M., & Knutti, R. (2021). An integrated approach to quantifying uncertainties in the remaining carbon budget. *Communications Earth & Environment*, 2(1), 7. <https://doi.org/10.1038/s43247-020-00064-9>
- 600 McLaren, D. (2016). Mitigation deterrence and the “moral hazard” of solar radiation management. *Earth’s Future*, 4(12), 596–602. <https://doi.org/10.1002/2016EF000445>
- Meinshausen, M., Meinshausen, N., Hare, W., Raper, S. C. B., Frieler, K., Knutti, R., Frame, D. J., & Allen, M. R. (2009). Greenhouse-gas emission targets for limiting global warming to 2°C. *Nature*, 458(7242), 1158–1162. <https://doi.org/10.1038/nature08017>

- 605 Meinshausen, M., Nicholls, Z. R. J., Lewis, J., Gidden, M. J., Vogel, E., Freund, M., Beyerle, U., Gessner, C., Nauels, A.,
Bauer, N., Canadell, J. G., Daniel, J. S., John, A., Krummel, P. B., Luderer, G., Meinshausen, N., Montzka, S. A.,
Rayner, P. J., Reimann, S., ... Wang, R. H. J. (2020). The shared socio-economic pathway (SSP) greenhouse gas
concentrations and their extensions to 2500. *Geoscientific Model Development*, 13(8), 3571–3605.
<https://doi.org/10.5194/gmd-13-3571-2020>
- 610 Meinshausen, M., Raper, S. C. B., & Wigley, T. M. L. (2011). Emulating coupled atmosphere-ocean and carbon cycle models
with a simpler model, MAGICC6—Part 1: Model description and calibration. *Atmospheric Chemistry and Physics*,
11(4), 1417–1456. <https://doi.org/10.5194/acp-11-1417-2011>
- Meinshausen, M., Smith, S. J., Calvin, K., Daniel, J. S., Kainuma, M. L. T., Lamarque, J., Matsumoto, K., Montzka, S. A.,
Raper, S. C. B., Riahi, K., Thomson, A., Velders, G. J. M., & van Vuuren, D. P. P. (2011). The RCP greenhouse gas
concentrations and their extensions from 1765 to 2300. *Climatic Change*, 109(1), 213–241.
615 <https://doi.org/10.1007/s10584-011-0156-z>
- Merk, C., Pönitzsch, G., & Rehdanz, K. (2016). Knowledge about aerosol injection does not reduce individual mitigation
efforts. *Environmental Research Letters*, 11(5). <https://doi.org/10.1088/1748-9326/11/5/054009>
- Moreno-Cruz, J. (2015). Mitigation and the geoengineering threat. *Resource and Energy Economics*, 41, 248–263.
<https://doi.org/10.1016/j.reseneeco.2015.06.001>
- 620 Möller, I. (2020). Political Perspectives on Geoengineering: Navigating Problem Definition and Institutional Fit. *Global
Environmental Politics*, 20(2), 57–82. https://doi.org/10.1162/glep_a_00547
- Nauels, A., Meinshausen, M., Mengel, M., Lorbacher, K., & Wigley, T. M. L. (2017). Synthesizing long-Term sea level rise
projections-The MAGICC sea level model v2.0. *Geoscientific Model Development*, 10(6), 2495–2524.
<https://doi.org/10.5194/gmd-10-2495-2017>
- 625 Neuber, F., & Ott, K. (2020). The buying time argument within the solar radiation management discourse. *Applied Sciences
(Switzerland)*, 10(13). <https://doi.org/10.3390/app10134637>
- Parker, A., & Irvine, P. J. (2018). The Risk of Termination Shock From Solar Geoengineering. *Earth's Future*, 6(3), 456–467.
<https://doi.org/10.1002/2017EF000735>
- 630 Pathak, R. Slade, P.R. Shukla, J. Skea, R. Pichs-Madruga, D. Üрге-Vorsatz,2022: Technical Summary. In: Climate Change
2022: Mitigation of Climate Change. Contribution of Working Group III to the Sixth Assessment Report of the
Intergovernmental Panel on Climate Change [P.R. Shukla, J. Skea, R. Slade, A. Al Khourdajie, R. van Diemen, D.
McCollum, M. Pathak, S. Some, P. Vyas, R. Fradera, M. Belkacemi, A. Hasija, G. Lisboa, S. Luz, J. Malley, (eds.)].
Cambridge University Press, Cambridge, UK and New York, NY, USA. doi: 10.1017/9781009157926.002.
- Pierrehumbert, R. (2019). There is no Plan B for dealing with the climate crisis. *Bulletin of the Atomic Scientists*, 75(5), 215–
635 221. <https://doi.org/10.1080/00963402.2019.1654255>
- Raimi, K. T., Maki, A., Dana, D., & Vandenbergh, M. P. (2019). Framing of Geoengineering Affects Support for Climate
Change Mitigation. *Environmental Communication*, 13(3), 300–319.
<https://doi.org/10.1080/17524032.2019.1575258>
- Reynolds, J. L. (2019). Solar geoengineering to reduce climate change: A review of governance proposals. *Proceedings of the
640 Royal Society A: Mathematical, Physical and Engineering Sciences*, 475(2229).
<https://doi.org/10.1098/rspa.2019.0255>
- Riahi, K., R. Schaeffer, J. Arango, K. Calvin, C. Guivarch, T. Hasegawa, K. Jiang, E. Kriegler, R. Matthews, G.P. Peters, A.
Rao, S. Robertson, A.M. Sebbit, J. Steinberger, M. Tavoni, D.P. van Vuuren, 2022: Mitigation pathways compatible
with long-term goals. In IPCC, 2022: Climate Change 2022: Mitigation of Climate Change. Contribution of Working
645 Group III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [P.R. Shukla, J. Skea,
R. Slade, A. Al Khourdajie, R. van Diemen, D. McCollum, M. Pathak, S. Some, P. Vyas, R. Fradera, M. Belkacemi,

A. Hasija, G. Lisboa, S. Luz, J. Malley, (eds.)). Cambridge University Press, Cambridge, UK and New York, NY, USA. doi: 10.1017/9781009157926.005

- 650 Rogelj, J., Huppmann, D., Krey, V., Riahi, K., Clarke, L., Gidden, M., Nicholls, Z., & Meinshausen, M. (2019). A new scenario logic for the Paris Agreement long-term temperature goal. *Nature*, *573*(7774), 357–363. <https://doi.org/10.1038/s41586-019-1541-4>
- Rogelj, J., Shindell, D., Jiang, K., Fifita, S., Forster, P., Ginzburg, V., Handa, C., Kheshgi, H., Kobayashi, S., Kriegler, E., Mundaca, L., Séférian, R., & Vilariño, M. V. (2018). Mitigation Pathways Compatible with 1.5°C in the Context of Sustainable Development. In: *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathw.* IPCC Special Report Global Warming of 1.5 oC, 82pp. https://www.ipcc.ch/site/assets/uploads/sites/2/2019/02/SR15_Chapter2_Low_Res.pdf
- 655 Schleussner, C.-F., Rogelj, J., Schaeffer, M., Lissner, T., Licker, R., Fischer, E. M., Knutti, R., Levermann, A., Frieler, K., & Hare, W. (2016). Science and policy characteristics of the Paris Agreement temperature goal. *Nature Climate Change*, *6*(9), Article 9. <https://doi.org/10.1038/nclimate3096>
- 660 Shepherd J, Caldeira K, Haigh J, Keith D, Launder B, Mace G, MacKerron G, Pyle J, Rayner S, Redgwell C, Watson A (2009) *Geoengineering the climate: science, governance and uncertainty.* The Royal Academy
- Shue, H. (2017). Climate dreaming: Negative emissions, risk transfer, and irreversibility. *Journal of Human Rights and the Environment*, *8* (2), 203–216. doi:10.4337/jhre.2017.02.02
- Smith, P., Davis, S. J., Creutzig, F., Fuss, S., Minx, J., Smith, P. (2011). Agricultural greenhouse gas mitigation potential globally, in Europe and in the UK: What have we learnt in the last 20 years? *Global Change Biology*, *18*(1), 35–43. <https://doi.org/10.1111/j.1365-2486.2011.02517.x>
- 665 Smith, P., Davis, S. J., Creutzig, F., Fuss, S., Minx, J., Gabrielle, B., Kato, E., Jackson, R. B., Cowie, A., Kriegler, E., Van Vuuren, D. P., Rogelj, J., Ciais, P., Milne, J., Canadell, J. G., McCollum, D., Peters, G., Andrew, R., Krey, V., ... Yongsung, C. (2015). Biophysical and economic limits to negative CO₂ emissions. *Nature Climate Change*, *6*(1), 42–50. <https://doi.org/10.1038/nclimate2870>
- 670 Svoboda, T., Keller, K., Goes, M., Tuana, N. 2011: Sulfate aerosol geoengineering: the question of justice. *Public Aff. Q.* *25*:157-79.
- Tilmes, S., Sanderson, B. M., & O'Neill, B. C. (2016). Climate impacts of geoengineering in a delayed mitigation scenario. *Geophysical Research Letters*, *43*(15), 8222–8229. <https://doi.org/10.1002/2016GL070122>
- Tjiputra, J. F., Grini, A., & Lee, H. (2016). Impact of idealized future stratospheric aerosol injection on the large-scale ocean and land carbon cycles. *Journal of Geophysical Research: Biogeosciences*, *121*(1), 2–27. <https://doi.org/10.1002/2015JG003045>
- 675 Tokarska, K. B., Gillett, N. P., Weaver, A. J., Arora, V. K., & Eby, M. (2016). The climate response to five trillion tonnes of carbon. *Nature Climate Change*, *6*(9), Article 9. <https://doi.org/10.1038/nclimate3036>
- UNFCCC. (2016). Second conference of the parties. <https://unfccc.int/resource/%0Adocs/2015/cop21/eng/10a01.pdf>
- 680 Wibeck, V., Hansson, A., Anshelm, J. (2015). Questioning the technological fix to climate change – Lay sense-making of geoengineering in Sweden. *Energy Research & Social Science*, *7*, 23-30. <https://doi.org/10.1016/j.erss.2015.03.001>.
- Vakilifard, N., Williams, R. G., Holden, P. B., Turner, K., Edwards, N. R., & Beerling, D. J. (2022). Impact of negative and positive CO₂ emissions on global warming metrics using an ensemble of Earth system model simulations. *Biogeosciences*, *19*(17), 4249–4265. <https://doi.org/10.5194/bg-19-4249-2022>
- 685 Wibeck, V., Hansson, A., Anshelm, J. (2015). Questioning the technological fix to climate change – Lay sense-making of geoengineering in Sweden. *Energy Research & Social Science*, *7*, 23-30. <https://doi.org/10.1016/j.erss.2015.03.001>.
- Zickfeld, K., MacDougall, A. H., & Matthews, H. D. (2016). On the proportionality between global temperature change and cumulative CO₂ emissions during periods of net negative CO₂ emissions. *Environmental Research Letters*, *11*(5), 055006. <https://doi.org/10.1088/1748-9326/11/5/055006>
- 690

