

1 Developing the Svalbard Integrated Arctic Earth Observing System - SIOS

2 Hanne [H. Christiansen](#)¹, [Ilkka Seppo Olavi Matero](#)², Lisa Baddeley^{1,8,3}, [Kim Holmén](#)⁴, Clara Julie Marie
3 [Hoppe](#)⁵, Maarten J.J.E. [Loonen](#)⁶, Rune [Storvold](#)⁷, Vito [Vitale](#)⁸, Agata [Zaborska](#)⁹, Heikki [Lihavainen](#)²

4 ¹ Arctic Geophysics Department, The University Centre in Svalbard, UNIS, 9170 Longyearbyen, Norway

5 ² Svalbard Integrated Arctic Earth Observing System (SIOS), SIOS Knowledge Centre, 9170 Longyearbyen, Norway

6 ^{8,3} Birkeland Centre for Space Science, University of Bergen, Bergen, Norway

7 ⁴ [Norwegian Polar Institute, Framsenteret, 9296 Tromsø, Norway](#)

8 ⁵ Alfred Wegener Institute - Helmholtzcenter for Polar and Marine Research, 27570 Bremerhaven, Germany

9 ⁶ University of Groningen, Arctic Centre, Aweg 30, 9718 CW Groningen, Netherlands

10 ⁷ Norwegian Research Centre, 9019 Tromsø, Norway

11 ⁸ National Research Council, Institute of Polar Sciences, 40129 Bologna, Italy

12 ⁹ Institute of Oceanology Polish Academy of Sciences, 81-712 Sopot, Poland

13 *Correspondence to:* Hanne H. Christiansen (HanneC@unis.no)

14 **Abstract.** ~~We~~Based on the ongoing large climatic and environmental changes and the history of science coordination in
15 Svalbard leading to the development of Svalbard Integrated Arctic Earth Observing System (SIOS), we present an overview
16 of the current gaps in knowledge and infrastructure based on ~~an overall~~a synthesis of ~~all the~~ recommendations ~~developed~~
17 ~~as presented in~~ the ~~main outcome of the~~ annual State of Environmental Science in Svalbard (SESS) reporting of ~~the Svalbard~~
18 ~~Integrated Arctic Earth Observing System (SIOS)-SIOS~~. Recommendations from the first four years of SESS reporting
19 represent the point of view of the wide scientific community operating the large observing system implemented in Svalbard
20 (SIOS) since 2018, and aim to identify the scientific potential to further develop the observing system. The recommendations
21 are bottom-up inputs for a continuous process that aims to accomplish the vision and mission of SIOS: optimising, integrating
22 and further developing the observing system in an Earth System Science (ESS) perspective. The primary outcome of the
23 synthesis work is the evidence that ESS in SIOS has, during the first 4 years of operation, naturally developed from individual
24 scientists or smaller groups of scientists to larger disciplinary international groups of scientists working together within the
25 different environments (atmosphere, cryosphere, marine and terrestrial environments). It is clear that strategic efforts towards
26 interdisciplinarity are necessary for operating fully at ESS scale in Svalbard. As Svalbard is experiencing the largest ongoing
27 warming in the Arctic and worldwide, SIOS is in a unique position to perform a full-scale study of all processes impacting
28 ESS dynamics and controlling the water cycle, using all parts of the SIOS observation network, with a large potential for
29 increasing the understanding of key mechanisms in the Earth System. We also identify the potential to upscale Svalbard-based
30 observations collected in SIOS to pan-Arctic scale, and to global scale, contributing to full scale ESS.

31 1. Introduction to ongoing climatic and environmental changes in Svalbard

32 The Arctic Earth System is experiencing rapid transformations driven by climate change. The archipelago of Svalbard (74°-
33 81°N), located halfway between Norway and the North Pole, is experiencing some of the largest climatic changes during the
34 last decades, making it an Arctic hotspot. Svalbard has the longest high Arctic meteorological record from the Longyearbyen
35 area extending back to 1898. It clearly shows how surface warming has been ongoing during the last half century (Fig 1).
36 Isaksen et al. (2022) identify a statistically significant and exceptional to the Arctic and globally record-high annual surface
37 air temperature warming of up to 2.7°C per decade, with a maximum in autumn of up to 4.0°C per decade for north easternmost
38 Svalbard, on the Karl XII-øya islands (1991-2020). In central Svalbard the surface air temperature at Longyearbyen airport
39 has had a linear increase of 3.8°C in the 1899–2018 period. This is about 3.5 times more than for the global mean temperature
40 during the same period (Nordli et al, 2020). Also, the annual warming rates have accelerated since the 1980s up to the latest last
41 decade ~~since 1981~~ (Isaksen et al., 2022). This warming is closely linked to substantial reduction in sea ice and an increase in
42 sea surface temperatures in the Northern Barents Sea. For this region, summers (JJA) have had the lowest decadal surface air
43 temperature warming ranging from 0 to 0.7°C, but with Karl XXII-øya experiencing a decadal warming rate of 1.3°C, while
44 winters (DJF) have seen up to 3.8°C per decade, and springs (MAM) 2.1°C per decade, all numbers based on the 2001-2020
45 period (Isaksen et al., 2022). The sea surface temperature has increased by 0.8°C per decade for the last two decades along
46 western and southern Svalbard, and in the southeastern Barents Sea, representing some of the largest sea surface warming rates
47 observed in the Northern Hemisphere, and reflecting the larger and warmer inflow of Atlantic Water by the West Spitsbergen
48 Current to this region (Isaksen et al., 2022). This inflow may also influence the late freeze-up of the Northern Barents Sea and
49 Franz Josef Land areas. Clearly, the ongoing environmental changes show how different parts of the Earth System are affected
50 and interacting in their responses to the ongoing surface warming both on land and in the sea.

51 In the mesosphere from 76-90 km, the hydroxyl (OH*) airglow temperature series from Longyearbyen is one of the longest
52 continuous measurement records of winter temperatures in the world extending from 1983 to the present (Wurst et al., 2023).
53 The overall daily average mesospheric temperature for the 2005 – 2012 seasons was -67.15°C, 3°C colder than studies from
54 earlier time periods (Holmen et al. 2014). Temperatures at 90 km altitude above Svalbard have also been determined using a
55 meteor wind radar and subsequently calibrated by satellite measurements for the period autumn 2002 to 2019. The cooling
56 rate during summer months is $9.9 \pm 2.9^\circ\text{C decade}^{-1}$ between 2002 and 2012, and $4.3 \pm 1.2^\circ\text{C decade}^{-1}$ between 2002 and 2019
57 (Hall et al. 2020). Carbon dioxide is a major driver of long-term trends in the upper atmosphere. This, along with other
58 greenhouse gas emissions, causes a general cooling effect of about -1 to -3 °C/yr at 250 km altitude (Zhang et al. 2016). Radar
59 observations have also shown that the situation is different between the low and high latitude regions, with the high latitudes
60 showing less cooling, even some warming above 300 km altitude (Lastovicka 2017). Similarities between the observations of
61 atmospheric vertical temperature trends and climate model projections are pointed out by the IPCC (IPCC, 2021), which state
62 that new techniques permit more robust quantification of temperature values and trends, allowing an improved confidence in
63 the vertical structure of temperature changes.

64 The Svalbard region is furthermore experiencing Atlantification of the surrounding seas, as increased ocean heat is transported
65 to the region from the west in the form of Atlantic water masses (Efstathiou et al., 2022). Concurrent changes in oceanographic
66 conditions (e.g., water temperature, nutrient loading, stratification), together with transport of organisms from further south,
67 may significantly alter species composition and productivity of Svalbard's coastal marine ecosystems (Bischof et al. 2019,
68 Assmy et al. 2023). Arctic terrestrial flora and fauna are generally sensitive to warming, but trends over a period of time are
69 heterogeneous and complex (Bjorkman et al., 2020; O'Connell et al. 2006, Pedersen et al. 2022). Most terrestrial Arctic
70 endemic species in Svalbard are experiencing negative consequences induced by the warming environment (Descamps, 2017).
71 In the terrestrial ecosystem, increased winter air temperatures are often accompanied by increases in the frequency of 'rain-
72 on-snow' events, one of the most important facets of climate change with respect to impacts on flora, fauna and society (Hansen
73 et al., 2014). Also, given that all the cryosphere components are inherently sensitive to temperature change especially around
74 0°C, Svalbard has experienced a continued net loss of ice (Meredith et al., 2019). The latter is a natural integrator of climate
75 variability and provides some of the most visible signatures of climate change, with retreating glaciers, shorter snow-covered
76 seasons and thawing permafrost (Constable et al., 2022).

77 **4.12 Science coordination in Svalbard leading to Svalbard Integrated Arctic Earth Observing System – SIOS**

78 ~~The potential to develop Svalbard as a full ESS regional distributed Research Infrastructure (RI) to study Arctic environmental~~
79 ~~change was identified already in 2007. To reach this goal, cooperation between the numerous national research stations,~~
80 ~~institutes, initiatives, and data collections in Svalbard had to be greatly improved. To achieve this, the concept of Svalbard~~
81 ~~Integrated Arctic Earth Observing System (SIOS)~~Research in Svalbard has a long tradition, going back at least to the “La
82 Recherche” expeditions in 1838 and 1839. Early work was in expedition form, but some permanent observatories emerged
83 from the International Geophysical Year (IGY) in 1957 (Norges Forskningsråd, 1997). A French station (“Corbel”) was
84 established outside Ny-Ålesund, as well as the Polish station in Hornsund. The first more permanent research establishment
85 came in 1967 when the Kongsfjord Telemetry Station was set up in Ny-Ålesund for communication with the European Space
86 Research Organization satellites. This was in close collaboration with The Norwegian Polar Institute, which established a
87 research station in Ny-Ålesund in 1968. The telemetry station was discontinued in 1974. A diversification of research in Ny-
88 Ålesund began with the Norwegian Institute for Air Research and University of Tromsø conducting activities as of the middle
89 of the 1970-ies. Norwegian universities and other institutions began sending summer expeditions to Ny-Ålesund, followed by
90 expansion of activities after the 1990s with institutions from numerous nations establishing a presence. It soon became apparent
91 that many projects worked on similar topics, and were partially duplicating logistical and scientific efforts. This was one of
92 the motivations for establishing the Ny-Ålesund Science Managers Committee (NySMAC) in 1994 as an organisation to
93 coordinate research and logistics. As NySMAC matured there was an increased desire to coordinate also the scientific questions
94 addressed. This led to the formulation of the NySMAC flagship programs, which were essentially established around the four
95 themes that *de facto* were most studied in Ny-Ålesund (Atmospheric science, Glaciology, Marine Science and Terrestrial
96 Science).

97 The research landscape in Svalbard then further developed. In Barentsburg there have been scientific activities under the
98 auspices of the Russian Arctic Antarctic Research Institute for decades. In Longyearbyen there were several research projects
99 present from around the time of the opening of the airport in 1975. The University Centre in Svalbard (UNIS) was established
100 in Longyearbyen in 1993, and has since grown to its present scale of ~50 scientific personnel and an annual throughflow of
101 about 750 students (200 student year equivalents per year). In addition to research activities in these permanent sites, a large
102 number of studies and projects have been carried out in the surrounding parts of the archipelago, with the scope to study the
103 glaciers, permafrost, vegetation and ecosystems in a changing climate.

104 With the diversity of research spreading across Svalbard, a need for better sharing of information about ongoing research was
105 seen and Svalbard Science Forum (SSF) was established in May 1998 under the auspices of the Norwegian Research Council
106 (St.meld.nr.22, 2009). SSF has expanded its activities with a new mandate in 2005 and rapidly evolved to its present
107 organisation with a permanent office in the Svalbard Science Centre in Longyearbyen. The initial task of SSF was to establish
108 the Research in Svalbard (RiS) database which then was a paper-based annual report. Today RiS is a fully digital direct
109 information platform with past, ongoing, and planned research projects in Svalbard. Everyone doing research in Svalbard is
110 expected to contribute to the RiS database. SSF has two meetings per year for Svalbard-wide research, providing information
111 exchange between the major research nodes in Svalbard (Barentsburg not participating since February 2022). SSF does not
112 make scientific priorities or recommendations *per se*, but facilitates the flow of information. SSF also provides some funding
113 opportunities to further scientific cooperation in Svalbard and enable the recruitment of young scientists to Arctic research, as
114 well as organising a conference focusing on Svalbard Science every other year.

115 The next challenge for the scientific community was to dare formulate and tackle larger Svalbard-wide endeavours where
116 questions larger than any single of the research entities would be able to pursue on their own, e.g. an Earth System Science
117 perspective. Another challenge was harmonising data and making it openly available using machine readable formats, so that
118 results from different contributors can be utilised together with minimal data revision. These were visions and roles that none
119 of the existing structures were designed to undertake. The creation of Svalbard Integrated Arctic Earth Observing System
120 (SIOS), an Earth System Science (ESS) -guided independent international consortium was therefore a next step (St.meld.nr.22,
121 2009) to bring together the full scientific benefits of the previously established practical research cooperation bodies. Towards
122 this backdrop the potential to develop a full ESS regional distributed Research Infrastructure (RI) to study Arctic environmental
123 change in Svalbard was identified already in 2007. To achieve this, the concept of SIOS was developed through an EU funded
124 preparatory phase project 2010-2013, and the operation continued with an interim phase 2014-2017. Thanks to great efforts
125 during the previous 10 years, SIOS entered its operational phase in 2018.

126 The mission of SIOS is to study the environment and climate in and around Svalbard to develop an efficient observing system,
127 share technology, experience and data, close knowledge gaps and decrease the environmental footprint of science. The aim of
128 SIOS is to perform an integrated assessment of how Arctic Earth System changes are developing and interacting with a clear

129 aim to connect the different scientific subdisciplines for improved ESS understanding. SIOS presents opportunities for research
130 and the acquisition of key knowledge on global environmental change, with a focus on processes and their interactions between
131 different ~~spheres~~environments, i.e., biosphere, geosphere, atmosphere, cryosphere, and hydrosphere. The vision of SIOS is to
132 be the leading long-term observing system in the Arctic to serve Earth System Science ~~and~~for society- ([https://sios-](https://sios-svalbard.org)
133 [svalbard.org](https://sios-svalbard.org)).

134 SIOS is currently a consortium of ~~29~~28 international research institutions from 10 different countries (<https://sios-svalbard.org>)
135 that own or operate research facilities in the Svalbard region, or that provide research data relevant for the consortium. Together
136 the consortium develops and maintains a regional observational system (Fig.2) for long-term measurements in and around
137 Svalbard, addressing Earth System Science questions related to Global Change. The members own and give access to their
138 research infrastructure. ~~Data produced by SIOS members follow~~The SIOS data policy states that new data contributions
139 from the consortium are to be made available through the SIOS data management system (SDMS, [https://sios-](https://sios-svalbard.org/Data)
140 [svalbard.org/Data](https://sios-svalbard.org/Data)), and that they need to follow the FAIR (Findable, Accessible, Interoperable, Reusable) guiding principles
141 for scientific data management and stewardship (Wilkinson et al. 2016), and they are accessible through SIOS data management
142 system-2016). SDMS includes a federated database system, in which the individual datasets are hosted at contributing data
143 centres across the globe and metadata records about these data are stored in a SIOS metadata catalogue. The datasets can be
144 accessed through a portal hosted on the SIOS website (<https://sios-svalbard.org/metsis/search>). At the time of writing the
145 database provides access to over 550000 datasets, with the longest time series spanning over 70 years back to 1945. The
146 currently available datasets follow the FAIR guiding principles to a varying degree, and a large fraction of them are from the
147 Norwegian National Ground Segment for satellite data (<https://www.satellitdata.no/en/metsis/search>). SIOS also hosts a
148 separate portal (<https://www.sios-svalbard.org/sios-ri-catalogue>), in which the current, historic and planned observation
149 facilities collecting SIOS data are documented.

150 The operations of the SIOS consortium are coordinated by the SIOS Knowledge Centre (SIOS KC), the central hub of SIOS.
151 SIOS has five active working groups with different tasks: Science Optimisation Advisory Group, Research Infrastructure
152 Coordination Committee, ~~SIOS~~-Data Management System working group, Remote Sensing Working Group and Information
153 Advisory Group. The mandate of the Science Optimisation Advisory Group (SOAG) is to prioritise ideas and initiatives for
154 ~~the~~ observing system development, considering scientific and societal relevance, feasibility and realism.

155 The previously-mentioned flagship programs of Ny-Ålesund are developing and taking an increasing role in formulating
156 upcoming research projects and programs which have direct similarities with the priority processes in SIOS. An intimate
157 relationship between SIOS and the NySMAC flagship programs is a logical and emerging development. SIOS and SSF have
158 some overlapping activities in supporting projects, but whereas SSF does not prioritise science, SIOS seeks to actively coax
159 member institutions towards embracing new and bigger scientific endeavours.

2.1.2 State of Environmental Science in Svalbard

As SIOS entered operation in 2018, an important overview tool for the observing system was initiated, the annual report series: State of Environmental Science in Svalbard (SESS). The overall aim of the reports is to summarise the state of current knowledge of key ESS parameters and analyse how these interact. The SESS reports contain peer-reviewed scientific chapters and associated outreach summaries. During the first 4 years of operation (2018-2021), a total of 40 individual chapters presented 169 recommendations (Orr et al., 2019; Van den Heuvel et al., 2020; Moreno-Ibáñez et al, 2021 & Feldner et al, 2022). Most SESS contributions have been reviews, but data summaries and updates to earlier SESS contributions have also been included. The SESS contributions have been authored by international and sometimes multidisciplinary groups.

~~The SESS recommendations are a bottom-up process to develop the SIOS RI, as topics are proposed by international research groups and selected through a review process. Proposals have been concerning new topics or simply updates to earlier SESS reporting. SESS reports are the integral part of the SIOS work program aimed to develop a coherent Svalbard observing system for many uses and users (Fig. 3). SIOS, i.e. the SIOS member institutions, the SIOS working groups, and the SIOS Knowledge Centre are the internal forces that try to move the Svalbard scientific community forward to provide better Earth system science data in Svalbard and thus, serve societal needs related to climate change, pollution and biodiversity loss. The different aspects of the Earth System and the means to improve the observing system are the preoccupation of the many SIOS working units, including task forces that are formed from different SIOS working groups, individuals from SIOS member institutions working in secondments on a specific topic within SIOS, and expert residents such as the SIOS Chair targeting to utilise SIOS data in scientific research. With the principles of cooperation, innovation and data harmonisation this work ensures that the SIOS community produces relevant long-term data series, the SIOS core data (<https://sios-svalbard.org/CoreData>), and improved research infrastructure (<https://www.sios-svalbard.org/sios-ri-catalogue>) as basis for new research projects and capacity building. The SESS report allows research groups from member institutions to identify observational gaps and provide recommendations on how to close those in a bottom-up process. The SESS reports are released annually in the last week of January, during an annual SIOS conference called the Polar Night Week. This event is also the main venue for stakeholders, researchers, and SIOS working groups to meet for discussing and finalising consortium activity plans for the following years. The plans are then further aligned with strategic aspects by the General Assembly of SIOS in a top-down process, allowing the work program to develop with input from all levels of SIOS.~~

~~The SESS reports help develop the SIOS research infrastructure, as topics are proposed by international research groups and selected through a review process.~~ The calls for input to the SESS reports have been open to all interested researchers from the SIOS member institutions and from other institutions with research activity in Svalbard. Each SESS contribution has been urged to connect interdisciplinarily with the different ESS spheres to allow for improved ESS understanding. SESS recommendations within the reports form an integral part of the contributions and help identify the scientific knowledge gaps, recommend improvements in terms of ~~research infrastructure~~RI and rising societal needs in Svalbard. The topics of SESS

192 chapters have ranged across the many sub-disciplines of ESS from atmosphere, cryosphere to the marine and terrestrial
193 environments. The objective of this paper is to 1) present the synthesis of the SESS recommendations and 2) discuss how to
194 further develop SIOS to serve the pan-Arctic ESS community.

195 2.2 The State of interdisciplinary Earth System Science in Svalbard

196 Understanding the value and accessing the work performed in SIOS and published in the SESS reports, the state of the
197 interdisciplinary scientific background is obviously essential. There are numerous publications on the state of science within
198 the separate ESS disciplines in Svalbard. Only a few of these address interdisciplinary research questions cutting across
199 disciplines. However, at few exist such as Peeters et al. (2019), who identified a climate-cryosphere regime shift stressing the
200 linkages between the ongoing meteorological changes towards more winter rain precipitation and thicker ground ice covers
201 affecting the cryosphere, but also affecting the biological environment with animal foraging being moved to ice-free grounds.
202 This changing form of precipitation is thus identified as important for changes in the high-arctic terrestrial environment (Peeters
203 et al., 2019). It has also been identified at Arctic scale (Bintanja & Selten (2014) that increased winter precipitation is primarily
204 due to intensified local evaporation typically from open sea, thus clearly linking the ongoing increases in Arctic winter
205 precipitation to the marine environment as well. The most recent review of snow research in Svalbard (Zdanowicz et al., 2023),
206 largely based on SIOS work and collaboration, identifies knowledge gaps and research needs in snow science. In doing so the
207 review relates these gaps and needs to other cryospheric fields such as glacier mass balance and active layer thickness above
208 permafrost, but also to atmospheric conditions in relation to snowpack pollution and to terrestrial ecology and hydrology.
209 Clearly some scientific fields do recognise their interdisciplinary scientific connections. With respect to extreme precipitation
210 events over Svalbard, it is well-known that these are caused by moisture transport extending thousands of kilometres south
211 into the subtropical Atlantic and often characterised as ‘atmospheric rivers’ (Serreze et al, 2015; Müller et al., 2022). The
212 increase in extreme precipitation events over the last four decades in Svalbard have been directly linked to the sea ice extent
213 east of Greenland, as this sea ice shields the west coast of Svalbard from incoming southerly moist air (Müller et al., 2022).
214 Thus results show clear linking of meteorological changes with changes in the marine environment and with well-known
215 consequences for ecosystems, cryosphere and society at Arctic scale.

216 2.1. 3. Methods

217 SOAG performed the first synthesis of the output of the SESS reporting to identify how to improve and develop SIOS. A task
218 force was established from SOAG and SIOS_KC which included experts from all ESS environments. This approach was
219 considered most natural and sustainable when reviewing and condensing the recommendations, focusing on the atmosphere,
220 the cryosphere and the marine and terrestrial environments. The main aim of the synthesis was to identify, condense and
221 prioritise the recommendations into key recommendations for the four environments and use these to identify joint
222 interdisciplinary or overall recommendations that expand across the entire Svalbard Earth System.

223

224 An open process took place with as much participation from all SESS authors as possible, inviting lead authors from all SESS
225 chapters within the four environments to thematic workshops, with the focus on condensing the range of recommendations
226 within each of their environments. The task force has worked fully online, and due to COVID-19, the four thematic
227 environment workshops had, unfortunately, to be held online in early winter 2022. The cryosphere workshop brought together
228 6 SESS authors covering widely different parts of the cryosphere SESS contributions. For the atmosphere, 5 authors and
229 experts largely discussed the list of recommendations extracted for this domain.

230 In each environment workshop, the recommendations were grouped and condensed to obtain an overview of the main
231 recommendations within each environment. Based on this thematic condensation of the key recommendations from the four
232 environments, the task force then discussed and further condensed the recommendations across for the entire ESS focusing on
233 interdisciplinarity. Finally, the entire SOAG provided feedback and the Board of Directors of SIOS approved the report as a
234 guiding document for developing SIOS RI.

235 **3.2. 4. Results**

236 **34.1 Key recommendations from the atmosphere, cryosphere, marine and terrestrial environments**

237 Here we briefly present the main recommendations from the four different environments of the SIOS ESS.

238 **34.1.1 Atmosphere**

239 —In the mesosphere / lower thermosphere -ionosphere (M/LTI) system (40 - 400 km), research focuses on understanding the
240 interaction and coupling between the neutral and ionised atmospheric components, energetic particle fluxes, ionospheric
241 instabilities, and the cusp auroral region. In the troposphere, stratosphere (T/S) system (0 - 40 km) research focuses on such
242 topics as aerosol-cloud interactions, atmospheric chemistry in the Arctic boundary layer, stratospheric ozone and surface
243 spectral UV fluxes. Several recommendations focus particularly on identifying methodological, technological or observational
244 gaps along with the need to continue existing long time series measurements- [\(Table 1\)](#). For the M/LTI region this is centred
245 around maintaining well established, larger facilities, such as radars, which can provide multiple parameters such as
246 temperatures, densities, atmospheric motion **en**ergy dissipation rates, across multiple scales, with an established, extensive
247 international use base. The facilities have built up extensive long-term databases and provide the opportunity for additional
248 instrumentation to be deployed in their vicinity.

249 Suggested improvements for the lower atmosphere include filling gaps in observations of spectral UV, black carbon, local
250 sources of biological aerosol precursors as well as further meteorological and hydrological measurements. Recent
251 developments in unmanned aerial vehicles (UAV) and miniaturisation of electronics and instruments should be exploited to
252 develop more mobile observational platforms. There should also be a push to increase the number of parameters observed by

253 AWS (Automatic Weather Station), especially in Eastern and Northern Svalbard. There should also be a stronger connection
254 and integration with measurements performed in East Greenland.

255 **34.1.2 Cryosphere**

256 The cryosphere synthesis is based on contributions within hydrology, glaciology, seismology, permafrost, snow, and on UVA
257 of the cryosphere. SIOS has fostered international groups collaborating in Svalbard within the different cryospheric
258 subdisciplines. The group on snow research has been leading the way with collaborating extensively and is now running a
259 SIOS pilot snow project. The group has also submitted a joint manuscript for peer-review based on this collaboration. The four
260 main recommendations from the cryosphere group are listed in Table 1. The changing water cycle during climate change was
261 identified as an overall scientific knowledge gap that needs all parts of the cryosphere to be involved, and in addition needs
262 extended links to the atmosphere. Integration across SIOS was found to be increased by establishing supersites building on
263 existing SIOS observations. Longyearbyen-Adventdalen area was identified as an obvious candidate where repetitive central
264 observations in combination with remote sensing can serve society and students and even with possibilities for citizen science
265 to be developed. There is also a wish for methodological coordination across the cryospheric research fields with, for instance,
266 time-lapse cameras, remote sensing (InSAR), fibre-optic cables for high spatial observations, ground penetrating radar,
267 seismics/cryoseismics passive/active seismic and cryoseismic measurements, and potentially networks of these types of
268 measurements across different SIOS supersites. Supersites are also instrumental for developing and harmonising methods to
269 ensure that all collected data are comparable and immediately usable.

270 Finally, it would be optimal to equip SIOS for observing sudden cryospheric events affecting the cryosphere such as glacial
271 lake drainage (GLOFs) or meteorological extreme events such as autumn or winter rainstorms causing landslides and increased
272 runoff due to extensive snow melting. These topics are also important for interdisciplinary studies.

273 **34.1.3 Marine**

274 The marine synthesis is based on contributions covering the following topics: oceanic circulation, ocean-atmosphere
275 interactions, plankton monitoring, sea ice thickness, microplastic pollution and the status of Svalbard coastal waters. Major
276 recommendations identified a need for a marine observing infrastructure development that provides data needed to fill critical
277 gaps in the observing system. Specifically, these included autonomous observational sites for co-located long-term monitoring
278 of ocean physics, biogeochemistry and ecology (with the latter two currently being under sampled with continuous methods),
279 enhanced sampling capabilities during the winter and early spring season, as well as spatially increased data collection by a
280 wider involvement of ships of opportunity. This recommendation could be addressed by specific funding via SIOS access and
281 optimisation calls that provide access to existing infrastructure and aid the development of new infrastructure. Homogeneous
282 data collection in disparate areas and by various teams could be ensured by a ~~dedicated~~ programme for harmonisation of
283 marine measurements around Svalbard. To fulfil this need, marine infrastructure network workshops have been organised by

284 SIOS, encouraging further cooperation on joint scientific programs. This network would be well suited to develop a handbook
285 of best practices for ocean observing approaches and data management. The four most essential recommendations for marine
286 ESS Observing System development were identified: (i) to extend the geographical and temporal coverage of research
287 activities, (ii) to enhance year-round observations to resolve seasonal variability in the ocean, (iii) to support research activities
288 exploring linkages between fjord, shelf and open ocean systems, and (iv) to establish long-term year-round monitoring of
289 marine biota.

290 **34.1.4 Terrestrial**

291 Several recommendations relate to infrastructure, wider data collection and remote sensing. Measurements on varied spatial
292 scales and development of models are mentioned frequently, but still need further implementation. Notably, there is no
293 contribution on lake systems and just one recommendation for experimental design. Overall, there is a need for co-location of
294 measurements to integrate various disciplines, temporal and spatial scales, experimental manipulations and to facilitate model-
295 based quantitative analysis. Current examples include the terrestrial flagship in Ny-Ålesund (Pedersen et al. 2022), the COAT
296 program led by the Norwegian Polar Institute (Ims et al., 2013), and the Bjørndalen Integrated Gradients (BIG) studies
297 initiated by the University Centre in Svalbard. COAT is a national ecosystem-based monitoring system of low and high Arctic
298 terrestrial environments, developed as a joint effort by the Fram centre institutions already in 2013 before SIOS was initiated.
299 COAT is now fully integrated into SIOS, and the two projects have received joint Norwegian infrastructure funding. These
300 studies should further develop, with new methods and technologies facilitating more spatially and temporally extensive and
301 high-resolution automated measurements of biological parameters. There needs to be a focus on long-term ecosystem-based
302 monitoring to establish how various anthropogenic pressures affect the Arctic environment. Climate change is likely to
303 transform Arctic terrestrial ecosystems beyond scientists' current abilities to make predictions. Therefore it is important to
304 have scientifically robust ecosystem-based, integrated systems to detect the fast changes from climate change, which is the
305 overall overriding threat, on terrestrial ecosystems.

306 **34.2 Knowledge gaps and associated improvement potential in Svalbard Earth System Science**

307 In addition to these environment-specific recommendations, interdisciplinary and overall recommendations that all four SESS
308 environments have brought up have been identified, and further synthesised for increasing and improving the interdisciplinarity
309 of SIOS.

310 With Svalbard experiencing the largest ongoing warming in the Arctic and worldwide and with the closely associated further
311 environmental changes, this puts SIOS in a unique position to provide coordinated datasets to investigate ongoing processes
312 in the Earth-system in a coupled manner. SIOS can quantify and understand in detail all physical processes that ultimately
313 warm the atmosphere, turning ice into water with all the resulting consequences this has on the entire Earth System. This
314 warming is controlled by the atmosphere and changes in ocean currents and impacts the cryosphere, as well as the terrestrial

315 and marine environments. By using the entire observation network of SIOS to perform a full-scale study of all the processes
316 which impact Earth System dynamics controlling the water cycle, a major potential for increasing the understanding of key
317 mechanisms in the Earth System has been identified. Such a study has potential to provide large knowledge gain, and thus
318 further the understanding of effects on the Svalbard Earth System.

319 Many cross-cutting individual SESS recommendations indicate the relevance and need of atmospheric parameters and
320 observations for studying the other environments in SIOS. Also, cross-cutting actions between the atmosphere, hydrosphere
321 and cryosphere have been indicated as areas of clear interest. Influences on the upper atmosphere from within the Sun-Earth
322 system, unique to the polar regions, have been shown to effect ozone depletion down to stratospheric altitudes (30km) and that
323 this may become more prominent with climate change (Maliniemi et al. 2020). In turn, lower atmospheric processes, such as
324 waves and tides, can influence the dynamics and behaviours of the upper atmosphere (Stober et al. 2021). The degree of
325 influence is a topic of debate, with global models failing to reproduce some of the observed effects (e.g., Oliver et al. 2013,
326 Cnossen 2020). Precipitation and wind action in the lower atmosphere result in long- and short-range transport of dust and
327 aerosols and accumulation of black carbon on snow and glaciers. Arctic terrestrial ecosystems, such as vegetation and its
328 components, and microbial communities, are impacted by air temperature, wind and precipitation, all of which are directly
329 controlled by the atmosphere. For instance, spectral characteristics of downwelling flux influence vegetation and microbial
330 community. Melting of snow and glaciers, coupled with thawing of permafrost (and an increase in active layer depth), supply
331 water, sediment, microbes and aqueous chemical species through the terrestrial environment into the marine environment, via
332 the hydrological system. The pathway from cryosphere through terrestrial to marine thus links all spheres in SIOS. Thus, all
333 these identified cross-cutting topics clearly have large potential if coordinated to fill the identified knowledge gap of the
334 water cycle and its consequences.

335 With the bottom-up approach of the SESS contributions, the development of a perspective from within the four environments
336 was clearly a very natural starting point for synthesising. To start increasing the interdisciplinarity and cross-cutting nature of
337 SIOS, a next step is to increase the dialogue between the different environments. Potentially, dedicated workshops on specific
338 topics such as radiative forcing between two or more SIOS environments could be a starting point. Such workshops should
339 also be tasked with pointing out optimal observations needed for datasets to fill and address the overall identified cross-cutting
340 knowledge gap.

341 Additionally, all environments of SIOS depend on obtaining better spatial and temporal coverage of observations in and around
342 Svalbard. This can be achieved through increased and coordinated instrument deployments that better cover the
343 physiographically different regions of Svalbard including various landscape/seascape/icescape types and boundaries between
344 them (e.g., the coastal zone). As warming is greatest in the eastern part of the Svalbard area (Isaksen et al., 2022), it is the
345 ambition to expand the SIOS observation system to also cover the eastern and northern part of Svalbard. Given the high
346 ambition for preserving these relatively untouched parts of the archipelago of Svalbard, such expansion might have to be
347 mainly based on remote sensing observations and/or dedicated supersites based mainly on the use of new low-maintenance

348 technology. For example, the findings regarding Arctic vegetation and its climatic control by Bjorkman et al. (2020) highlight
349 the need for more geographically widespread, integrated, and comprehensive monitoring efforts that can better resolve the
350 interacting effects of warming and other local and regional ecological factors. In this case, studies across the entire vegetation
351 zonation of Svalbard are important to fully understand the effects of climatic changes to the entire flora of the different Arctic
352 vegetation zones. The spatially distributed network approach by COAT with large spatial and temporal coverage and co-
353 location of measurements covering the terrestrial above-ground food web is another example. COAT measures vegetation
354 from 57 module stations covering large parts of Nordenskiöldsland, Isfjorden and Brøggerhalvøya in central Svalbard. There
355 is a clear need across all the environments for special focus on the autumn and spring periods.

356 While the current focus on the major science hubs represents a disadvantage in terms of covering more different areas and
357 habitat types as detailed above, there is clear potential ~~identified~~ for co-location of all the different types of observations from
358 all parts of the ESS, this way developing real supersites. This should complement existing infrastructure, with a high level of
359 coordination, long-term planning, and funding. It would have the advantage of fewer sites to maintain, allowing for a more
360 sustainable observation system. It would also allow for focusing on much closer collaboration on new methods and
361 technologies across SIOS, with a potential for more efficient sharing of knowledge. Knowledge gained from the supersites
362 may be used to guide the selection of a subset of measurement types for observations with higher spatial and temporal coverage.
363 An ambitious tool identified is dedicated method-based action forces, that would make SIOS able to respond quickly to extreme
364 events such as solar storms, sudden stratospheric warmings (SSWs), glacial lake outburst floods (GLOFs) or various
365 meteorological extremes, such as rainstorms and avalanches, the latter of which are also highly relevant to be able to perform
366 a full-scale study of the water cycle changes and its consequences.
367 Likewise, the need for real-time access to SIOS observations are important, particularly for extreme events, for increasing the
368 return of data from desirable but high-risk deployments (such as moorings or instruments near or on surging glaciers, or rocket
369 launches into specific atmospheric phenomena), but also for more widespread use of the SIOS data.

370 **4.5. Discussion and conclusion**

371 This synthesis of knowledge gaps based on the first four years of operation of the Svalbard Integrated Arctic Earth Observing
372 System (SIOS) builds on a unique and strong international collaboration. No other observing system exists which is this wide-
373 ranging, covering all part of ESS within a specific regional area, yet at the same time supported so widely internationally and
374 so well-coordinated. When developing a full-scale Earth science observation system as SIOS, with the comprehensive full
375 scale ESS ambition that SIOS has, it is very natural for researchers within the individual disciplines and even subdisciplines
376 to first start working closer together before being able to focus more on the interdisciplinarity necessary for performing full-
377 scale ESS analysing how the observed parameters influence each other. This is clearly the case for SIOS after the first four
378 years of operation.

379 Ways to accommodate this important next step include the implementation of above-mentioned workshops on interdisciplinary
380 topics as well as the development of supersites. Also, a workshop across all of SIOS working groups is planned to increase the
381 collaboration and interdisciplinarity of SIOS. Hopefully also scientific collaboration between two environments might start
382 being more developed in SIOS, such as already developed in the fully ecosystem-based observation system of COAT
383 combining climate and ecology, designed to observe, understand and predict outcomes from climate change (drivers) on tundra
384 ecosystems (Ims & Yoccoz, 2017). In addition SIOS is operating training activities in the form of small courses on different
385 key SIOS observation techniques, SIOS data availability and SIOS remote sensing opportunities for all interested. Increased
386 interdisciplinarity can also be obtained by further developing the major science hubs in Svalbard into real supersites
387 complementing existing infrastructure allowing even closer collaboration on particular new technologies and methods
388 including real-time access to observations across SIOS. Just as obtaining better spatial and temporal observation coverage in
389 and around Svalbard, including the physiographically different regions of Svalbard, both the various land-, sea- and icescapes
390 and boundaries between them in the coastal zone, would lead to increased potential for interdisciplinarity.

391 Other ways forward will be to increase the use of the SIOS observations through the SIOS data catalogue for Earth System
392 modelling, to obtain improved process understanding that bridges science disciplines. This will demand working to expand
393 the SIOS community with modellers from different scientific fields, just as increased collaboration between modellers and the
394 observational groups in SIOS needs to be established to enable observing at the relevant scales and resolutions, and of missing
395 parameters. The Earth system modelling community could make little use of high resolution regional data in the early stages
396 of SIOS but presently as model resolution approaches sub-regional scales it becomes increasingly relevant to utilise models
397 for observation design as well as accrued data to calibrate and verify model output. The SIOS Observation Facility Catalogue
398 and the Research in Svalbard database are important tools for developing interdisciplinarity in Svalbard ESS. Just as the SESS
399 reports are ~~also~~ an integral part of the strategic development of the observing system, ~~called the SIOS science wheel~~. This
400 bottom-up approach, which is complemented with rising societal needs and member priorities, leads to the outcomes from the
401 SIOS ~~science~~ Science wheel: (Fig. 3). In addition, the harmonised SIOS core data, new research infrastructure and research
402 projects are all innovations leading to increased interdisciplinarity, highly encouraged through the SIOS optimization and
403 access calls. Finally, SIOS offers ~~innovation awards~~ Innovation Awards to encourage further developing, also high risk – high
404 gain, technological and/or other ESS key activities that address knowledge gaps identified by the SIOS community.

405 Further, as Svalbard is experiencing the largest ongoing warming in the Arctic and worldwide, this puts SIOS in a unique
406 position to perform a full-scale study of all processes impacting ESS dynamics controlling the water cycle using all parts of
407 the SIOS observation network, with large potential for increasing the understanding of key mechanisms in the Earth System.
408 The clear identification of a joint scientific focus, studying the water cycle changes and its consequences using all parts of the
409 SIOS ESS observation network, will function well also as a focal point for cross-disciplinary work. Despite a disciplinary
410 focus, snow, precipitation and hydrology have been identified as overlapping topics in many SESS contributions, but not
411 prominent in the marine environment. These topics are therefore connected to all environments primarily on land, but observed

412 and treated in different ways within the different environments. This is important for the identified scientific focus on the
413 changing water cycle. Hence, SIOS has potential to develop a multidisciplinary study with data products and modelling
414 approaches based on the variety of observations that is being collected and supplement this with additional necessary new
415 observations leading to increased Earth System understanding of current and future dynamics.

416 There is also a need for increased regional collaboration using SIOS in scales broader than just Svalbard, when aiming for full
417 Earth System understanding within the Arctic. This means collaborating primarily with the rest of the Arctic. The international
418 ‘Sustaining Arctic Observing Networks’ (SAON) is the regional facilitator aiming at coordinating, improving, integrating and
419 sustaining pan-Arctic observations during rapid environmental and social change, with a main focus on specifically addressing
420 key gaps in coordination (Starkweather et al., 2021). With SIOS in the warmest part of the Arctic, and the largest climatic
421 gradient in the high Arctic existing between Svalbard and cold North Greenland, closer collaboration with Greenland could be
422 a first natural step. In Greenland, the main aim of the ‘Greenland Integrated Observing System’ (GIOS), <https://gios.org/> is
423 to resolve and understand the mechanisms behind climate and environmental change in Greenland and beyond. GIOS is
424 developing a network of sustainable long-term research infrastructures in and around Greenland observing the changing air,
425 ice, land, and ocean conditions. GIOS is, as opposed to SIOS, a national research infrastructure linking all institutions and
426 universities currently carrying out Arctic research in the Danish Realm. SIOS and GIOS have started discussing ~~to increase~~
427 ~~the increasing~~ collaboration within research, higher education and logistics. Interhemispheric co-operation is also another
428 obvious step with networks in the Antarctic Region such as SCAR (Scientific Committee on Antarctic Research), as an
429 obvious collaboration partner. Initiatives in specific disciplines, such as GRAPE (Global Navigation Satellite System Research
430 and Application for Polar Environment) are already in place. The research outlined through SIOS would benefit from being
431 part of global observing systems such as the SuperDARN (Super Dual Auroral Radar, Chisham et al. 2007) atmospheric
432 network. Such global networks, with commensurable data products, allow regional effects to be placed in a global context.

433 SIOS is making use of the Earth Observing System (EOS) program of NASA with its different satellite missions and scientific
434 instruments comprised of a series of coordinated polar-orbiting satellites designed to monitor and understand key components
435 of the climate system and their interactions through long-term global observations of the land surface, the biosphere,
436 atmosphere and oceans (Platnik, 2022). This also enables direct comparison between Svalbard and particularly the rest of the
437 Arctic, of the different ESS components which are being remotely observed. To make best use of this data, links between
438 existing EOS data ~~needs~~need to be established, and data need to be available in formats that global Earth system modelling
439 can make use of.

440 Finally, many of the observations collected by regional observation systems like SIOS have large potential to add to improved
441 understanding of how people living in the Arctic are affected by climatic changes in many different ways (e.g., by landslides,
442 avalanches, coastal erosion). ~~Such understanding~~COAT has a clear goal of providing management relevance. Clearly, also
443 SIOS can assist humans and societies to develop management of climate change impacts. Such assistance is needed for

444 developing resilience and adaptation to enable Arctic communities to deal with the consequences of the ongoing climatic
445 changes. Presently, with the move towards involving local communities directly in climate change observations and research,
446 this might place SIOS in a new role as also contributing more directly to societal needs.

447 **Data Availability**

448 Data for Figure 1 is available through the Norwegian Meteorological Institute online service. Data for Figure 2 is
449 available in the SIOS Observation facility catalogue.

450 **Author contributions**

451 All authors have contributed to the SIOS synthesis process, and all have contributed to writing this manuscript.

452 **Competing interests**

453 The authors declare that they have no conflict of interest.

454 **Disclaimer**

455 **Acknowledgements**

456 The authors would like to acknowledge the support from the Research Council of Norway through projects No. 291644,
457 No.322387 & No. 269927 and Norwegian Space Agency through contract No. NIT 12.20.5. We thank our SIOS colleagues
458 who have contributed with SESS reporting, and those who assisted us during the thematic workshops to start synthesising the
459 SESS recommendations. Thanks to Åshild Ø. Pedersen, Norwegian Polar Institute for comments improving the revised
460 manuscript on terrestrial biology. Finally, we very much appreciate the comments by two anonymous reviewers, which have
461 largely improved the revised manuscript.

462 **References**

- 463 [Assmy, P., Kvernvik, A.C., Hop, H., Hoppe, C.J.M., Chierici, M., David, D.T., Duarte, P., Fransson, A., Martin Garcia, L., Kwasniewski, S.,](#)
464 [Maturilli, M., Pavlova, O., Tatarek, A., Wiktor, J.M., Wold, A., Wolf, K.K.E., Bailey, A., 2023, Plankton dynamics in Kongsfjorden during](#)
465 [two years of contrasting environmental conditions. *Progress in Oceanography*, <https://doi.org/10.1016/j.pocean.2023.102996>.](#)
- 466 [Bintanja, R., Selten, F., \(2014\). Future increases in Arctic precipitation linked to local evaporation and sea-ice retreat. *Nature* 509, 479–482.](#)
467 <https://doi.org/10.1038/nature13259>.
- 468 [Bischof, K., Convey, P., Duarte, P., Gattuso, J.-P., Granberg, M., Hop, H., Hoppe, C., Jiménez, C., Lisitsyn, L., Martinez, B., Roleda, M.Y.,](#)
469 [Thor, P., Wiktor, J.M., Gabrielsen, G.W., 2019. Kongsfjorden as Harbinger of the Future Arctic: Knowns, Unknowns and Research](#)
470 [Priorities, in: Hop, H., Wiencke, C. \(Eds.\), *The Ecosystem of Kongsfjorden, Svalbard. Springer International Publishing, Cham*, pp. 537–](#)
471 [562. \[https://doi.org/10.1007/978-3-319-46425-1_14\]\(https://doi.org/10.1007/978-3-319-46425-1_14\)](https://doi.org/10.1007/978-3-319-46425-1_14)
- 472 [Bjorkman, A.D., García Criado, M., Myers-Smith, I.H. et al., 2020. Status and trends in Arctic vegetation: Evidence from experimental](#)
473 [warming and long-term monitoring. *Ambio* 49, 678–692.](#)

- 474 [Chisham G, Lester M, Milan SE, Freeman MP, Bristow WA, Grocott A, McWilliams KA, Ruohoniemi JM, Yeoman TK, Dyson PL,](#)
475 [Greenwald RA, 2007. A decade of the Super Dual Auroral Radar Network \(SuperDARN\): Scientific achievements, new techniques and](#)
476 [future directions. *Surveys in Geophysics* 28\(1\):33–109 437](#)
- 477 [Cnossen, I., 2020. Analysis and attribution of climate change in the upper atmosphere from 1950 to 2015 simulated by WACCM-X.](#)
478 [Journal of Geophysical Research: *Space Physics*, 125, e2020JA028623. <https://doi.org/10.1029/2020JA028623>](#)
- 479 [Constable, A.J., S. Harper, J. Dawson, K. Holsman, T. Mustonen, D. Piepenburg, and B. Rost, 2022. Cross-Chapter Paper 6: Polar](#)
480 [Regions. In: *Climate Change 2022: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Sixth Assessment*](#)
481 [Report of the Intergovernmental Panel on Climate Change \[H.-O. Pörtner, D.C. Roberts, M. Tignor, E.S. Poloczanska, K. Mintenbeck, A.](#)
482 [Alegría, M. Craig, S. Langsdorf, S. Löschke, V. Möller, A. Okem, B. Rama \(eds.\)\]. Cambridge University Press, Cambridge, UK and New](#)
483 [York, NY, USA, pp. 2319–2368, <https://doi:10.1017/9781009325844.023>](#)
- 484 [Descamps, S., Aars, J., Fuglei, E., Kovacs, K.M., Lydersen, C., Pavlova, O., Pedersen, Å.Ø., Ravolainen, V. and Strøm, H., 2017. Climate](#)
485 [change impacts on wildlife in a High Arctic archipelago – Svalbard, Norway. *Glob Change Biol*, 23: 490-502.](#)
486 [https://doi.org/10.1111/gcb.13381](#)
- 487 [Efstathiou, E., Eldevik, T., Arthun, M. & Lind, S., 2022. Spatial patterns, mechanisms and predictability of Barents Sea ice change. *J.*](#)
488 [Clim. 35\(10\), 2961–2973. <https://doi.org/10.1175/JCLI-D-21-0044.1>](#)
- 489 [Feldner J., Hübner C., Lihavainen H., Neuber R., Zaborska A. \(eds\), 2022. *SESS report 2021*, Svalbard Integrated Arctic Earth Observing](#)
490 [System, Longyearbyen. \[https://sios-svalbard.org/SESS_Issue4\]\(https://sios-svalbard.org/SESS_Issue4\).](#)
- 491 [Hall, C.M., Tsutsumi, M., 2020. Neutral temperatures at 90 km altitude over Svalbard \(78°N 16°E\), 2002–2019, derived from meteor radar](#)
492 [observations, *Polar Science*, Volume 24, 100530, ISSN 1873-9652, <https://doi.org/10.1016/j.polar.2020.100530>.](#)
- 493 [Hansen B.B., Isaksen K., Benestad R.E. et al., 2014. Warmer and wetter winters: characteristics and implications of an extreme weather](#)
494 [event in the High Arctic. *Environmental Research Letter*, 9, 114021](#)
- 495 [Holmen, S.E., Dyrland, M.E. & Sigernes, F., 2014. Mesospheric temperatures derived from three decades of hydroxyl airglow](#)
496 [measurements from Longyearbyen, Svalbard \(78°N\). *Acta Geophys*. 62, 302–315. <https://doi.org/10.2478/s11600-013-0159-4>](#)
- 497 [Ims, R.A., Jepsen, J.U., Stien, A. & Yoccoz, N.G., 2013. Science plan for COAT: Climate-ecological Observatory for Arctic Tundra.](#)
498 [Fram Centre Report Series 1, Fram Centre, Norway, 177 pages.](#)
- 499 [Ims, R.A. & Yoccoz, N.G., 2017. Ecosystem-based monitoring in the age of rapid climate change and new technologies, *Current Opinion*](#)
500 [in *Environmental Sustainability*, 29, 170-176, <https://doi.org/10.1016/j.cosust.2018.01.003>.](#)
- 501 [IPCC Climate Change 2021: The Physical Science Basis. Working Group I Contribution to the IPCC Sixth Assessment Report.](#)
- 502 [Isaksen, K., Nordli, Ø., Ivanov, B., Køltzow, M.A.Ø., Aaboe, S., Gjelten, H.M., Mezghani, A., Eastwood, S., Fjørland, E., Benestad, R.E.,](#)
503 [Hanssen-Bauer, I., Brækkan, R., Sviashchennikov, P., Demin, V., Revina, A., Karandasheva, T., 2022. Exceptional warming over the](#)
504 [Barents area. *Scientific Reports* 12:9371. <https://doi.org/10.1038/s41598-022-13568-5>](#)
- 505 [Lastovicka, J., Urbar, J., Kozubek, M., 2017. Long-term trends in the total electron content. *Geophys. Res. Lett.*, 44, 8168–8172,](#)
506 [https://doi:10.1002/2017GL075063](#)
- 507 [Maliniemi, V., Marsh, R Daniel, Tyssøy, H.N, Smith-Johnsen, C, 2020. Will climate change impact polar nox produced by energetic](#)
508 [particle precipitation?. *Geophysical Research Letters* 47\(9\), 2020–087041](#)
- 509 [Meredith, M., M. Sommerkorn, S. Cassotta, C. Derksen, A. Ekaykin, A. Hollowed, G. Kofinas, A. Mackintosh, J. Melbourne-Thomas,](#)
510 [M.M.C. Muelbert, G. Ottersen, H. Pritchard, and E.A.G. Schuur, 2019: Polar Regions. In: *IPCC Special Report on the Ocean and*](#)
511 [Cryosphere in a Changing Climate \[H.-O. Pörtner, D.C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K.](#)
512 [Mintenbeck, A. Alegría, M. Nicolai, A. Okem, J. . Petzold, B. Rama, N.M. Weyer \(eds.\)\].](#)

513 [Moreno-Ibáñez M., Hagen J.O., Hübner C., Lihavainen H., Zaborska A. \(eds\), 2021. *SESS report 2020*, Svalbard Integrated Arctic Earth](https://sios-svalbard.org/SESS_Issue3)
514 [Observing System, Longyearbyen. https://sios-svalbard.org/SESS_Issue3.](https://sios-svalbard.org/SESS_Issue3)

515 [Müller, M. et al., 2022. Decline of sea-ice in the Greenland Sea intensifies extreme precipitation over Svalbard. *Weather and Climate*](https://doi.org/10.1016/j.wace.2022.100437)
516 [Extremes, 36, 100437, https://doi.org/10.1016/j.wace.2022.100437.](https://doi.org/10.1016/j.wace.2022.100437)

517 [Nordli, O. et al., 2020. Revisiting the extended Svalbard Airport monthly temperature series, and the compiled corresponding daily series](https://doi.org/10.33265/polar.v39.3614)
518 [1898–2018. *Polar Res.* https://doi.org/10.33265/polar.v39.3614](https://doi.org/10.33265/polar.v39.3614)

519 [Norges forskningsråd, 1997. Organisering av forskningen på Svalbard, Norwegian Research Council report, ISBN 82-12-00864-9.](https://doi.org/10.33265/polar.v39.3614)

520 [Oliver, W. L., S.-R. Zhang, and L. P. Goncharenko, 2013. Is thermospheric global cooling caused by gravity waves?, *J. Geophys. Res.*](https://doi.org/10.1002/jgra.50370)
521 [*Space Physics*, 118, 3898–3908, https://doi.org/10.1002/jgra.50370](https://doi.org/10.1002/jgra.50370)

522 [O'Connell, M.J., Huiskes A.H.L., Loonen M.J.J.E., Madsen J., Klaassen M. & M. Rounsevell, 2006. Developing an integrated approach to](https://doi.org/10.33265/polar.v41.6310)
523 [understanding the effects of climate change and other environmental alterations at a flyway level. *Waterbirds around the world*. Eds. G.C.](https://doi.org/10.33265/polar.v41.6310)
524 [Boere, C.A. Galbraith & D.A. Stroud. pp. 385-397.](https://doi.org/10.33265/polar.v41.6310)

525 [Orr, E., Hansen, G., Lappalainen H., Hübner C., Lihavainen, H. \(eds\) 2019: *SESS report 2018*, Svalbard Integrated Arctic Earth Observing](https://sios-svalbard.org/SESS_Issue1)
526 [System, Longyearbyen. https://sios-svalbard.org/SESS_Issue1](https://sios-svalbard.org/SESS_Issue1)

527 [Pedersen, Å. Ø., Stien, J., Albon, S., Fuglei, E., Isaksen, K., Liston, G. & Ims, R. A., 2020. Climate-ecological observatory for Arctic](https://doi.org/10.1088/1748-9326/aaefb3)
528 [Tundra \(COAT\). *State of Environmental Science in Svalbard \(SESS\) Report 2019*, 58-83.](https://doi.org/10.1088/1748-9326/aaefb3)

529 [Pedersen, Å. Ø., Convey, P., Newsham, K. K., Mosbacher, J. B., Fuglei, E., Ravolainen, V., Hansen, B. B., Jensen, T. C., Augusti, A.,](https://doi.org/10.33265/polar.v41.6310)
530 [Biersma, E. M., Cooper, E. J., Coulson, S. J., Gabrielsen, G. W., Gallet, J. C., Karsten, U., Kristiansen, S. M., Svenning, M. M., Tveit, A.](https://doi.org/10.33265/polar.v41.6310)
531 [T., Uchida, M., ... Loonen, M. J. J. E., 2022. Five decades of terrestrial and freshwater research at Ny-Ålesund, Svalbard. *Polar Research*,](https://doi.org/10.33265/polar.v41.6310)
532 [41, \[6310\]. https://doi.org/10.33265/polar.v41.6310](https://doi.org/10.33265/polar.v41.6310)

533 [Peeters, B. et al., 2019. Spatiotemporal patterns of rain-on-snow and basal ice in high Arctic Svalbard: detection of a climate-cryosphere](https://doi.org/10.1088/1748-9326/aaefb3)
534 [regime shift. *Environmental Research Letters*, 14, 015002. https://doi.org/10.1088/1748-9326/aaefb3](https://doi.org/10.1088/1748-9326/aaefb3)

535 [Platnik, S., 2022. "Earth Observing System". *NASA's Earth Observing System*.](https://doi.org/10.1088/1748-9326/aaefb3)

536 [Serreze, et al., 2015. Extreme daily precipitation events at Spitsbergen, an Arctic Island. *International Journal of Climatology*, 35, 4574-](https://doi.org/10.1002/joc.4308)
537 [4588. https://doi.org/10.1002/joc.4308](https://doi.org/10.1002/joc.4308)


538 [St.meld.nr.22, 2009, Stortingsmelding nr. 22 \(2008-2009\) Svalbard, Det Kongelige Justis- og Politidepartement, Norwegian Parliament](https://doi.org/10.1088/1748-9326/aaefb3)
539 [White paper, in Norwegian.](https://doi.org/10.1088/1748-9326/aaefb3)

540 [Starkweather, S., Larsen, J.R., Kruemmel, E., Eicken, H., Arthurs, D., Bradley, A.C., Carlo, N., Christensen, T., Daniel, R., Danielsen, F.,](https://doi.org/10.14430/arctic74330)
541 [Kalhok, S., Karcher, M., Johansson, M., Jóhannsson, H., Kodama, Y., Lund, S., Murray, M.S., Petäjä, T., Pulsifer, P.L., Sandven, S.,](https://doi.org/10.14430/arctic74330)
542 [Sankar, R.D., Strahlendorff, M., & Wilkinson, J., 2021. Sustaining Arctic Observing Networks' \(SAON\) Roadmap for Arctic Observing](https://doi.org/10.14430/arctic74330)
543 [and Data Systems \(ROADS\). *Arctic*, 74, SUPPL. 1 \(2021\) P. 56 – 68, https://doi.org/10.14430/arctic74330](https://doi.org/10.14430/arctic74330)

544 [Stober, G., Kuchar, A., Pokhotelov, D., Liu, H., Liu, H.-L., Schmidt, H., Jacobi, C., Baumgarten, K., Brown, P., Janches, D., Murphy, D.,](https://doi.org/10.5194/acp-21-13855-2021)
545 [Kozlovsky, A., Lester, M., Belova, E., Kero, J., and Mitchell, N., 2021. Interhemispheric differences of mesosphere–lower thermosphere](https://doi.org/10.5194/acp-21-13855-2021)
546 [winds and tides investigated from three whole-atmosphere models and meteor radar observations, *Atmos. Chem. Phys.*, 21, 13855–13902.](https://doi.org/10.5194/acp-21-13855-2021)
547 <https://doi.org/10.5194/acp-21-13855-2021>


548 [Van den Heuvel F., Hübner C., Błaszczyk M., Heimann M., Lihavainen H. \(eds\) 2020: *SESS report 2019*, Svalbard Integrated Arctic Earth](https://sios-svalbard.org/SESS_Issue2)
549 [Observing System, Longyearbyen. https://sios-svalbard.org/SESS_Issue2](https://sios-svalbard.org/SESS_Issue2)

- 550 [Wurst, Sabine, M. Bittner, P. J. Espy, W. John R. French, and F. J. Mulligan, \(2023\) Hydroxyl airglow observations for investigating](https://doi.org/10.5194/acp-23-1599-2023)
551 [atmospheric dynamics: results and challenges, *Atmos. Chem. Phys.*, 23, pp. 1599–1618, https://doi.org/10.5194/acp-23-1599-2023](https://doi.org/10.5194/acp-23-1599-2023)
- 552 [Zdanowicz C., Gallet J.-C., Salvatori R., Malnes E., Isaksen K., Hübner C., Jones E., & Lihavainen H., 2023. An agenda for the future of](https://doi.org/10.33265/polar.v42.8827)
553 [Arctic snow research: the view from Svalbard. *Polar Research*, 42. https://doi.org/10.33265/polar.v42.8827](https://doi.org/10.33265/polar.v42.8827)
- 554 [Zhang, Shun-Rong, et al., 2016. "Ionospheric ion temperature climate and upper atmospheric long-term cooling." *Journal of Geophysical*](https://doi.org/10.1002/2016JA022971)
555 [*Research: Space Physics* 121.9 \(2016\): 8951-8968. https://doi.org/10.1002/2016JA022971](https://doi.org/10.1002/2016JA022971)



557 **Figure 1. The meteorological record from the Longyearbyen area is composed of the homogenised data series from the**
558 **Norwegian Meteorological Institute (1898-2022).**

559



561 **Figure 2. SIOS observation infrastructure in Svalbard, with the three main science hubs Longyearbyen, Hornsund and**
 562 **Ny-Ålesund inset to present all the various types of observations in more detail.**

563

AN OBSERVING SYSTEM FOR MANY



564

565 **Figure 3:** SIOS Science wheel that provides an overview of developing the observing system for many uses and
566 users.

567

Atmosphere	Cryosphere	Marine	Terrestrial
Support continued funding of large scale, multi-disciplinary infrastructure, such as the Svalbard SuperDARN radar	Science based coordination of the cryospheric observations	Development of the marine environmental observing infrastructure - autonomous observational sites	Focus on co-location of basic cryospheric observations and subsequent long-term studies on biota at similar spatial and temporal scales
Provision of data management and online open access portal to facilitate easier sharing of datasets	Integrated SIOS observations supersites	Provision of new datasets by e.g., enhancing measurement capabilities	Focus on new methods and technologies using automated sensors, animal trackers and cameras
Better harmonisation of the geographical and temporal coverage of atmospheric observations and applying standardized observation methods	Common cryospheric methodological infrastructure priorities	Harmonised methodologies for in situ observations and data harmonisation	Develop model-based quantitative analyses of ecosystem processes using SIOS datasets
Enhance the spatial scale of the atmospheric observational network	Establish SIOS action force for extreme events	Support for research exploring linkages between fjord, shelf and open ocean systems	Develop interphases between monitoring-based ecosystem science and end-users

Table 1. Key synthesised recommendations identified with the four environments of SIOS.

570 **5. Supplementary information**

571 **S1. Table of names, acronyms, countries and homepages of the 28 SIOS institutions**

<u>Institution full name</u>	<u>Acronym</u>	<u>Country</u>	<u>Website</u>
<u>Alfred Wegener Institute, Helmholtz centre for polar and marine research</u>	<u>AWI</u>	<u>Germany</u>	<u>https://www.awi.de/</u>
<u>Andoya</u>	<u>ASP</u>	<u>Norway</u>	<u>https://www.andoyaspace.no/</u>
<u>Geological Survey of Norway</u>	<u>NGU</u>	<u>Norway</u>	<u>http://www.ngu.no/en/</u>
<u>Institute for Atmospheric and Earth System Research of the University of Helsinki</u>	<u>INAR</u>	<u>Finland</u>	<u>https://www.helsinki.fi/en/inar-institute-for-atmospheric-and-earth-system-research</u>
<u>Institute of Geophysics, Polish Academy of Sciences</u>	<u>IG PAS</u>	<u>Poland</u>	<u>http://www.igf.edu.pl/home.php</u>
<u>Institute of Marine Research</u>	<u>IMR</u>	<u>Norway</u>	<u>http://www.imr.no/en</u>
<u>Korea Polar Research Institute</u>	<u>KOPRI</u>	<u>Rep. of Korea</u>	<u>https://eng.kopri.re.kr/</u>
<u>NILU</u>	<u>NILU</u>	<u>Norway</u>	<u>https://www.nilu.no/</u>
<u>Norwegian Research Centre AS</u>	<u>NORCE</u>	<u>Norway</u>	<u>https://www.norceresearch.no/en/</u>
<u>NORSAR</u>	<u>NORSAR</u>	<u>Norway</u>	<u>https://www.norsar.no/home/</u>
<u>Norwegian University of Science and Technology</u>	<u>NTNU</u>	<u>Norway</u>	<u>https://www.ntnu.edu/</u>
<u>Nansen Environmental and Remote Sensing Center</u>	<u>NERSC</u>	<u>Norway</u>	<u>https://www.nersc.no/</u>
<u>National Centre for Polar and Ocean Research</u>	<u>NCPOR</u>	<u>India</u>	<u>http://www.ncaor.gov.in/</u>
<u>National Institute for Polar Research</u>	<u>NIPR</u>	<u>Japan</u>	<u>http://www.nipr.ac.jp/english/</u>
<u>National Research Council of Italy</u>	<u>CNR</u>	<u>Italy</u>	<u>https://www.cnr.it/en</u>
<u>Norwegian Institute for Nature Research</u>	<u>NINA</u>	<u>Norway</u>	<u>https://www.nina.no/</u>
<u>Norwegian Institute for Water Research</u>	<u>NIVA</u>	<u>Norway</u>	<u>https://www.niva.no/en</u>
<u>Norwegian Meteorological Institute</u>	<u>MET Norway</u>	<u>Norway</u>	<u>https://www.met.no/en</u>
<u>Norwegian Polar Institute</u>	<u>NPI</u>	<u>Norway</u>	<u>http://www.npolar.no/en/</u>
<u>Norwegian Water Resources and Energy Directorate</u>	<u>NVE</u>	<u>Norway</u>	<u>https://www.nve.no/english/</u>
<u>Swedish Polar Research Secretariat</u>	<u>SPRS</u>	<u>Sweden</u>	<u>https://polar.se/en/</u>
<u>The Norwegian Geotechnical Institute</u>	<u>NGI</u>	<u>Norway</u>	<u>https://www.ngi.no/</u>
<u>The University Centre in Svalbard</u>	<u>UNIS</u>	<u>Norway</u>	<u>https://www.unis.no/</u>
<u>The Arctic University of Norway</u>	<u>UiT</u>	<u>Norway</u>	<u>https://en.uit.no/</u>
<u>University of Bergen</u>	<u>UiB</u>	<u>Norway</u>	<u>http://www.uib.no/en</u>
<u>University of Groningen</u>	<u>UG</u>	<u>Netherlands</u>	<u>https://www.rug.nl/</u>
<u>University of Oslo</u>	<u>UiO</u>	<u>Norway</u>	<u>http://www.uio.no/english/</u>
<u>University of Silesia, Poland</u>	<u>US</u>	<u>Poland</u>	<u>http://english.us.edu.pl/</u>