

1 Developing the Svalbard Integrated Arctic Earth Observing System - SIOS

2 Hanne H. Christiansen¹, Ilkka Seppo Olavi Matero², Lisa Baddeley^{1,3}, Kim Holmén⁴, Clara Julie Marie
3 Hoppe⁵, Maarten J.J.E. Loonen⁶, Rune Storvold⁷, Vito Vitale⁸, Agata Zaborska⁹, Heikki Lihavainen²

4 ¹ Arctic Geophysics Department, The University Centre in Svalbard, UNIS, 9170 Longyearbyen, Norway

5 ² Svalbard Integrated Arctic Earth Observing System (SIOS), SIOS Knowledge Centre, 9170 Longyearbyen, Norway

6 ³ Birkeland Centre for Space Science, University of Bergen, Bergen, Norway

7 ⁴ Norwegian Polar Institute, Framsenteret, 9296 Tromsø, Norway

8 ⁵ Alfred Wegener Institute - Helmholtzcenter for Polar and Marine Research, 27570 Bremerhaven, Germany

9 ⁶ University of Groningen, Arctic Centre, Aweg 30, 9718 CW Groningen, Netherlands

10 ⁷ Norwegian Research Centre, 9019 Tromsø, Norway

11 ⁸ National Research Council, Institute of Polar Sciences, 40129 Bologna, Italy

12 ⁹ Institute of Oceanology Polish Academy of Sciences, 81-712 Sopot, Poland

13 *Correspondence to:* Hanne H. Christiansen (HanneC@unis.no)

14 **Abstract.** Based on the ongoing large climatic and environmental changes and the history of science coordination in Svalbard
15 leading to the development of Svalbard Integrated Arctic Earth Observing System (SIOS), we present an overview of the
16 current gaps in knowledge and infrastructure based on a synthesis of the recommendations presented in the annual State of
17 Environmental Science in Svalbard (SESS) reporting of SIOS. Recommendations from the first four years of SESS reporting
18 represent the point of view of the wide scientific community operating the large observing system implemented in Svalbard
19 (SIOS) since 2018, and aim to identify the scientific potential to further develop the observing system. The recommendations
20 are bottom-up inputs for a continuous process that aims to accomplish the vision and mission of SIOS: optimising, integrating
21 and further developing the observing system in an Earth System Science (ESS) perspective. The primary outcome of the
22 synthesis work is the evidence that ESS in SIOS has, during the first 4 years of operation, naturally developed from individual
23 scientists or smaller groups of scientists to larger disciplinary international groups of scientists working together within the
24 different environments (atmosphere, cryosphere, marine and terrestrial environments). It is clear that strategic efforts towards
25 interdisciplinarity are necessary for operating fully at ESS scale in Svalbard. As Svalbard is experiencing the largest ongoing
26 warming in the Arctic and worldwide, SIOS is in a unique position to perform a full-scale study of all processes impacting
27 ESS dynamics and controlling the water cycle, using all parts of the SIOS observation network, with a large potential for
28 increasing the understanding of key mechanisms in the Earth System. We also identify the potential to upscale Svalbard-based
29 observations collected in SIOS to pan-Arctic scale, and to global scale, contributing to full scale ESS.

1. Introduction to ongoing climatic and environmental changes in Svalbard

The Arctic Earth System is experiencing rapid transformations driven by climate change. The archipelago of Svalbard (74°-81°N), located halfway between Norway and the North Pole, is experiencing some of the largest climatic changes during the last decades, making it an Arctic hotspot. Svalbard has the longest high Arctic meteorological record from the Longyearbyen area extending back to 1898. It clearly shows how surface warming has been ongoing during the last half century (Fig 1). Isaksen et al. (2022) identify a statistically significant and exceptional to the Arctic and globally record-high annual surface air temperature warming of up to 2.7°C per decade, with a maximum in autumn of up to 4.0°C per decade for north easternmost Svalbard, on the Karl XII-øya islands (1991-2020). In central Svalbard the surface air temperature at Longyearbyen airport has had a linear increase of 3.8°C in the 1899–2018 period. This is about 3.5 times more than for the global mean temperature during the same period (Nordli et al, 2020). Also, the annual warming rates have accelerated since the 1980s up to the last decade (Isaksen et al., 2022). This warming is closely linked to substantial reduction in sea ice and an increase in sea surface temperatures in the Northern Barents Sea. For this region, summers (JJA) have had the lowest decadal surface air temperature warming ranging from 0 to 0.7°C, but with Karl XXII-øya experiencing a decadal warming rate of 1.3°C, while winters (DJF) have seen up to 3.8°C per decade, and springs (MAM) 2.1°C per decade, all numbers based on the 2001-2020 period (Isaksen et al., 2022). The sea surface temperature has increased by 0.8°C per decade for the last two decades along western and southern Svalbard, and in the southeastern Barents Sea, representing some of the largest sea surface warming rates observed in the Northern Hemisphere, and reflecting the larger and warmer inflow of Atlantic Water by the West Spitsbergen Current to this region (Isaksen et al., 2022). This inflow may also influence the late freeze-up of the Northern Barents Sea and Franz Josef Land areas. Clearly, the ongoing environmental changes show how different parts of the Earth System are affected and interacting in their responses to the ongoing surface warming both on land and in the sea.

In the mesosphere from 76-90 km, the hydroxyl (OH*) airglow temperature series from Longyearbyen is one of the longest continuous measurement records of winter temperatures in the world extending from 1983 to the present (Wurst et al., 2023). The overall daily average mesospheric temperature for the 2005 – 2012 seasons was -67.15°C, 3°C colder than studies from earlier time periods (Holmen et al. 2014). Temperatures at 90 km altitude above Svalbard have also been determined using a meteor wind radar and subsequently calibrated by satellite measurements for the period autumn 2002 to 2019. The cooling rate during summer months is $9.9 \pm 2.9^\circ\text{C decade}^{-1}$ between 2002 and 2012, and $4.3 \pm 1.2^\circ\text{C decade}^{-1}$ between 2002 and 2019 (Hall et al. 2020). Carbon dioxide is a major driver of long-term trends in the upper atmosphere. This, along with other greenhouse gas emissions, causes a general cooling effect of about -1 to -3 °C/yr at 250 km altitude (Zhang et al. 2016). Radar observations have also shown that the situation is different between the low and high latitude regions, with the high latitudes showing less cooling, even some warming above 300 km altitude (Lastovicka 2017). Similarities between the observations of atmospheric vertical temperature trends and climate model projections are pointed out by the IPCC (IPCC, 2021), which state that new techniques permit more robust quantification of temperature values and trends, allowing an improved confidence in the vertical structure of temperature changes.

63 The Svalbard region is furthermore experiencing Atlantification of the surrounding seas, as increased ocean heat is transported
64 to the region from the west in the form of Atlantic water masses (Efstathiou et al., 2022). Concurrent changes in oceanographic
65 conditions (e.g., water temperature, nutrient loading, stratification), together with transport of organisms from further south,
66 may significantly alter species composition and productivity of Svalbard's coastal marine ecosystems (Bischof et al. 2019,
67 Assmy et al. 2023). Arctic terrestrial flora and fauna are generally sensitive to warming, but trends over a period of time are
68 heterogeneous and complex (Bjorkman et al., 2020; O'Connell et al. 2006, Pedersen et al. 2020). Most terrestrial Arctic
69 endemic species in Svalbard are experiencing negative consequences induced by the warming environment (Descamps, 2017).
70 In the terrestrial ecosystem, increased winter air temperatures are often accompanied by increases in the frequency of 'rain-
71 on-snow' events, one of the most important facets of climate change with respect to impacts on flora, fauna and society (Hansen
72 et al., 2014). Also, given that all the cryosphere components are inherently sensitive to temperature change especially around
73 0°C, Svalbard has experienced a continued net loss of ice (Meredith et al., 2019). The latter is a natural integrator of climate
74 variability and provides some of the most visible signatures of climate change, with retreating glaciers, shorter snow-covered
75 seasons and thawing permafrost (Constable et al., 2022).

76 **2. Science coordination in Svalbard leading to Svalbard Integrated Arctic Earth Observing System – SIOS**

77 Research in Svalbard has a long tradition, going back at least to the “La Recherche” expeditions in 1838 and 1839. Early work
78 was in expedition form, but some permanent observatories emerged from the International Geophysical Year (IGY) in 1957
79 (Norges Forskningsråd, 1997). A French station (“Corbel”) was established outside Ny-Ålesund, as well as the Polish station
80 in Hornsund. The first more permanent research establishment came in 1967 when the Kongsfjord Telemetry Station was set
81 up in Ny-Ålesund for communication with the European Space Research Organization satellites. This was in close
82 collaboration with The Norwegian Polar Institute, which established a research station in Ny-Ålesund in 1968. The telemetry
83 station was discontinued in 1974. A diversification of research in Ny-Ålesund began with the Norwegian Institute for Air
84 Research and University of Tromsø conducting activities as of the middle of the 1970-ies. Norwegian universities and other
85 institutions began sending summer expeditions to Ny-Ålesund, followed by expansion of activities after the 1990s with
86 institutions from numerous nations establishing a presence. It soon became apparent that many projects worked on similar
87 topics, and were partially duplicating logistical and scientific efforts. This was one of the motivations for establishing the Ny-
88 Ålesund Science Managers Committee (NySMAC) in 1994 as an organisation to coordinate research and logistics. As
89 NySMAC matured there was an increased desire to coordinate also the scientific questions addressed. This led to the
90 formulation of the NySMAC flagship programs, which were essentially established around the four themes that *de facto* were
91 most studied in Ny-Ålesund (Atmospheric science, Glaciology, Marine Science and Terrestrial Science).

92 The research landscape in Svalbard then further developed. In Barentsburg there have been scientific activities under the
93 auspices of the Russian Arctic Antarctic Research Institute for decades. In Longyearbyen there were several research projects
94 present from around the time of the opening of the airport in 1975. The University Centre in Svalbard was established in

95 Longyearbyen in 1993, and has since grown to its present scale of ~50 scientific personnel and an annual throughflow of about
96 750 students (200 student year equivalents per year). In addition to research activities in these permanent sites, a large number
97 of studies and projects have been carried out in the surrounding parts of the archipelago, with the scope to study the glaciers,
98 permafrost, vegetation and ecosystems in a changing climate.

99 With the diversity of research spreading across Svalbard, a need for better sharing of information about ongoing research was
100 seen and Svalbard Science Forum (SSF) was established in May 1998 under the auspices of the Norwegian Research Council
101 (St.meld.nr.22, 2009). SSF has expanded its activities with a new mandate in 2005 and rapidly evolved to its present
102 organisation with a permanent office in the Svalbard Science Centre in Longyearbyen. The initial task of SSF was to establish
103 the Research in Svalbard (RiS) database which then was a paper-based annual report. Today RiS is a fully digital direct
104 information platform with past, ongoing, and planned research projects in Svalbard. Everyone doing research in Svalbard is
105 expected to contribute to the RiS database. SSF has two meetings per year for Svalbard-wide research, providing information
106 exchange between the major research nodes in Svalbard (Barentsburg not participating since February 2022). SSF does not
107 make scientific priorities or recommendations *per se*, but facilitates the flow of information. SSF also provides some funding
108 opportunities to further scientific cooperation in Svalbard and enable the recruitment of young scientists to Arctic research, as
109 well as organising a conference focusing on Svalbard Science every other year.

110 The next challenge for the scientific community was to dare formulate and tackle larger Svalbard-wide endeavours where
111 questions larger than any single of the research entities would be able to pursue on their own, e.g. an Earth System Science
112 perspective. Another challenge was harmonising data and making it openly available using machine readable formats, so that
113 results from different contributors can be utilised together with minimal data revision. These were visions and roles that none
114 of the existing structures were designed to undertake. The creation of Svalbard Integrated Arctic Earth Observing System
115 (SIOS), an Earth System Science (ESS) -guided independent international consortium was therefore a next step (St.meld.nr.22,
116 2009) to bring together the full scientific benefits of the previously established practical research cooperation bodies. Towards
117 this backdrop the potential to develop a full ESS regional distributed Research Infrastructure (RI) to study Arctic environmental
118 change in Svalbard was identified already in 2007. To achieve this, the concept of SIOS was developed through an EU funded
119 preparatory phase project 2010-2013, and the operation continued with an interim phase 2014-2017. Thanks to great efforts
120 during the previous 10 years, SIOS entered its operational phase in 2018.

121 The mission of SIOS is to study the environment and climate in and around Svalbard to develop an efficient observing system,
122 share technology, experience and data, close knowledge gaps and decrease the environmental footprint of science. The aim of
123 SIOS is to perform an integrated assessment of how Arctic Earth System changes are developing and interacting with a clear
124 aim to connect the different scientific subdisciplines for improved ESS understanding. SIOS presents opportunities for research
125 and the acquisition of key knowledge on global environmental change, with a focus on processes and their interactions between

126 different environments, i.e., biosphere, geosphere, atmosphere, cryosphere, and hydrosphere. The vision of SIOS is to be the
127 leading long-term observing system in the Arctic to serve Earth System Science for society (<https://sios-svalbard.org>).

128 SIOS is currently a consortium of 28 international research institutions from 10 different countries (<https://sios-svalbard.org>)
129 that own or operate research facilities in the Svalbard region, or that provide research data relevant for the consortium. Together
130 the consortium develops and maintains a regional observational system (Fig.2) for long-term measurements in and around
131 Svalbard, addressing Earth System Science questions related to Global Change. The members own and give access to their
132 research infrastructure. The SIOS data policy states that new data contributions from the consortium are to be made available
133 through the SIOS data management system (SDMS, <https://sios-svalbard.org/Data>), and that they need to follow the FAIR
134 (Findable, Accessible, Interoperable, Reusable) guiding principles for scientific data management and stewardship (Wilkinson
135 et al. 2016). SDMS includes a federated database system, in which the individual datasets are hosted at contributing data
136 centres across the globe and metadata records about these data are stored in a SIOS metadata catalogue. The datasets can be
137 accessed through a portal hosted on the SIOS website (<https://sios-svalbard.org/metsis/search>). At the time of writing the
138 database provides access to over 550000 datasets, with the longest time series spanning over 70 years back to 1945. The
139 currently available datasets follow the FAIR guiding principles to a varying degree, and a large fraction of them are from the
140 Norwegian National Ground Segment for satellite data (<https://www.satellittdata.no/en/metsis/search>). SIOS also hosts a
141 separate portal (<https://www.sios-svalbard.org/sios-ri-catalogue>), in which the current, historic and planned observation
142 facilities collecting SIOS data are documented.

143 The operations of the SIOS consortium are coordinated by the SIOS Knowledge Centre, the central hub of SIOS. SIOS has
144 five active working groups with different tasks: Science Optimisation Advisory Group, Research Infrastructure Coordination
145 Committee, Data Management System working group, Remote Sensing Working Group and Information Advisory Group.
146 The mandate of the Science Optimisation Advisory Group (SOAG) is to prioritise ideas and initiatives for observing system
147 development, considering scientific and societal relevance, feasibility and realism.

148 The previously-mentioned flagship programs of Ny-Ålesund are developing and taking an increasing role in formulating
149 upcoming research projects and programs which have direct similarities with the priority processes in SIOS. An intimate
150 relationship between SIOS and the NySMAC flagship programs is a logical and emerging development. SIOS and SSF have
151 some overlapping activities in supporting projects, but whereas SSF does not prioritise science, SIOS seeks to actively coax
152 member institutions towards embracing new and bigger scientific endeavours.

153 **2.1 State of Environmental Science in Svalbard**

154 As SIOS entered operation in 2018, an important overview tool for the observing system was initiated, the annual report series:
155 State of Environmental Science in Svalbard (SESS). The overall aim of the reports is to summarise the state of current
156 knowledge of key ESS parameters and analyse how these interact. The SESS reports contain peer-reviewed scientific chapters

157 and associated outreach summaries. During the first 4 years of operation (2018-2021), a total of 40 individual chapters
158 presented 169 recommendations (Orr et al., 2019; Van den Heuvel et al., 2020; Moreno-Ibáñez et al, 2021 & Feldner et al,
159 2022). Most SESS contributions have been reviews, but data summaries and updates to earlier SESS contributions have also
160 been included. The SESS contributions have been authored by international and sometimes multidisciplinary groups.

161 SESS reports are the integral part of the SIOS work program aimed to develop a coherent Svalbard observing system for many
162 uses and users (Fig. 3). SIOS, i.e. the SIOS member institutions, the SIOS working groups, and the SIOS Knowledge Centre
163 are the internal forces that try to move the Svalbard scientific community forward to provide better Earth system science data
164 in Svalbard and thus, serve societal needs related to climate change, pollution and biodiversity loss. The different aspects of
165 the Earth System and the means to improve the observing system are the preoccupation of the many SIOS working units,
166 including task forces that are formed from different SIOS working groups, individuals from SIOS member institutions working
167 in secondments on a specific topic within SIOS, and expert residents such as the SIOS Chair targeting to utilise SIOS data in
168 scientific research. With the principles of cooperation, innovation and data harmonisation this work ensures that the SIOS
169 community produces relevant long-term data series, the SIOS core data (<https://sios-svalbard.org/CoreData>), and improved
170 research infrastructure (<https://www.sios-svalbard.org/sios-ri-catalogue>) as basis for new research projects and capacity
171 building. The SESS report allows research groups from member institutions to identify observational gaps and provide
172 recommendations on how to close those in a bottom-up process. The SESS reports are released annually in the last week of
173 January, during an annual SIOS conference called the Polar Night Week. This event is also the main venue for stakeholders,
174 researchers, and SIOS working groups to meet for discussing and finalising consortium activity plans for the following years.
175 The plans are then further aligned with strategic aspects by the General Assembly of SIOS in a top-down process, allowing
176 the work program to develop with input from all levels of SIOS.

177 The SESS reports help develop the SIOS research infrastructure, as topics are proposed by international research groups and
178 selected through a review process. The calls for input to the SESS reports have been open to all interested researchers from
179 the SIOS member institutions and from other institutions with research activity in Svalbard. Each SESS contribution has been
180 urged to connect interdisciplinarily with the different ESS spheres to allow for improved ESS understanding. SESS
181 recommendations within the reports form an integral part of the contributions and help identify the scientific knowledge gaps,
182 recommend improvements in terms of RI and rising societal needs in Svalbard. The topics of SESS chapters have ranged
183 across the many sub-disciplines of ESS from atmosphere, cryosphere to the marine and terrestrial environments. The objective
184 of this paper is to 1) present the synthesis of the SESS recommendations and 2) discuss how to further develop SIOS to serve
185 the pan-Arctic ESS community.

186 **2.2 The State of interdisciplinary Earth System Science in Svalbard**

187 Understanding the value and accessing the work performed in SIOS and published in the SESS reports, the state of the
188 interdisciplinary scientific background is obviously essential. There are numerous publications on the state of science within
189 the separate ESS disciplines in Svalbard. Only a few of these address interdisciplinary research questions cutting across
190 disciplines. However, at few exist such as Peeters et al. (2019), who identified a climate-cryosphere regime shift stressing the
191 linkages between the ongoing meteorological changes towards more winter rain precipitation and thicker ground ice covers
192 affecting the cryosphere, but also affecting the biological environment with animal foraging being moved to ice-free grounds.
193 This changing form of precipitation is thus identified as important for changes in the high-arctic terrestrial environment (Peeters
194 et al., 2019). It has also been identified at Arctic scale (Bintanja & Selten, 2014) that increased winter precipitation is primarily
195 due to intensified local evaporation typically from open sea, thus clearly linking the ongoing increases in Arctic winter
196 precipitation to the marine environment as well. The most recent review of snow research in Svalbard (Zdanowicz et al., 2023),
197 largely based on SIOS work and collaboration, identifies knowledge gaps and research needs in snow science. In doing so the
198 review relates these gaps and needs to other cryospheric fields such as glacier mass balance and active layer thickness above
199 permafrost, but also to atmospheric conditions in relation to snowpack pollution and to terrestrial ecology and hydrology.
200 Clearly some scientific fields do recognise their interdisciplinary scientific connections. With respect to extreme precipitation
201 events over Svalbard, it is well-known that these are caused by moisture transport extending thousands of kilometres south
202 into the subtropical Atlantic and often characterised as ‘atmospheric rivers’ (Serreze et al, 2015; Müller et al., 2022). The
203 increase in extreme precipitation events over the last four decades in Svalbard have been directly linked to the sea ice extent
204 east of Greenland, as this sea ice shields the west coast of Svalbard from incoming southerly moist air (Müller et al., 2022).
205 Thus results show clear linking of meteorological changes with changes in the marine environment and with well-known
206 consequences for ecosystems, cryosphere and society at Arctic scale.

207 **3. Methods**

208 SOAG performed the first synthesis of the output of the SESS reporting to identify how to improve and develop SIOS. A task
209 force was established from SOAG and SIOS Knowledge Centre which included experts from all ESS environments. This
210 approach was considered most natural and sustainable when reviewing and condensing the recommendations, focusing on the
211 atmosphere, the cryosphere and the marine and terrestrial environments. The main aim of the synthesis was to identify,
212 condense and prioritise the recommendations into key recommendations for the four environments and use these to identify
213 joint interdisciplinary or overall recommendations that expand across the entire Svalbard Earth System.

214
215 An open process took place with as much participation from all SESS authors as possible, inviting lead authors from all SESS
216 chapters within the four environments to thematic workshops, with the focus on condensing the range of recommendations
217 within each of their environments. The task force has worked fully online, and due to COVID-19, the four thematic
218 environment workshops had, unfortunately, to be held online in early winter 2022. The cryosphere workshop brought together

219 6 SESS authors covering widely different parts of the cryosphere SESS contributions. For the atmosphere, 5 authors and
220 experts largely discussed the list of recommendations extracted for this domain.

221 In each environment workshop, the recommendations were grouped and condensed to obtain an overview of the main
222 recommendations within each environment. Based on this thematic condensation of the key recommendations from the four
223 environments, the task force then discussed and further condensed the recommendations across for the entire ESS focusing on
224 interdisciplinarity. Finally, the entire SOAG provided feedback and the Board of Directors of SIOS approved the report as a
225 guiding document for developing SIOS RI.

226 **4. Results**

227 **4.1 Key recommendations from the atmosphere, cryosphere, marine and terrestrial environments**

228 Here we briefly present the main recommendations from the four different environments of the SIOS ESS.

229 **4.1.1 Atmosphere**

230 In the mesosphere / lower thermosphere -ionosphere (M/LTI) system (40 - 400 km), research focuses on understanding the
231 interaction and coupling between the neutral and ionised atmospheric components, energetic particle fluxes, ionospheric
232 instabilities, and the cusp auroral region. In the troposphere, stratosphere (T/S) system (0 - 40 km) research focuses on such
233 topics as aerosol-cloud interactions, atmospheric chemistry in the Arctic boundary layer, stratospheric ozone and surface
234 spectral UV fluxes. Several recommendations focus particularly on identifying methodological, technological or observational
235 gaps along with the need to continue existing long time series measurements (Table 1). For the M/LTI region this is centred
236 around maintaining well established, larger facilities, such as radars, which can provide multiple parameters such as
237 temperatures, densities, atmospheric motion, energy dissipation rates, across multiple scales, with an established, extensive
238 international use base. The facilities have built up extensive long-term databases and provide the opportunity for additional
239 instrumentation to be deployed in their vicinity.

240 Suggested improvements for the lower atmosphere include filling gaps in observations of spectral UV, black carbon, local
241 sources of biological aerosol precursors as well as further meteorological and hydrological measurements. Recent
242 developments in unmanned aerial vehicles (UAV) and miniaturisation of electronics and instruments should be exploited to
243 develop more mobile observational platforms. There should also be a push to increase the number of parameters observed by
244 AWS (Automatic Weather Station), especially in Eastern and Northern Svalbard. There should also be a stronger connection
245 and integration with measurements performed in East Greenland.

246 **4.1.2 Cryosphere**

247 The cryosphere synthesis is based on contributions within hydrology, glaciology, seismology, permafrost, snow, and on UVA
248 of the cryosphere. SIOS has fostered international groups collaborating in Svalbard within the different cryospheric
249 subdisciplines. The group on snow research has been leading the way with collaborating extensively and is now running a
250 SIOS pilot snow project. The group has also submitted a joint manuscript for peer-review based on this collaboration. The four
251 main recommendations from the cryosphere group are listed in Table 1. The changing water cycle during climate change was
252 identified as an overall scientific knowledge gap that needs all parts of the cryosphere to be involved, and in addition needs
253 extended links to the atmosphere. Integration across SIOS was found to be increased by establishing supersites building on
254 existing SIOS observations. Longyearbyen-Adventdalen area was identified as an obvious candidate where repetitive central
255 observations in combination with remote sensing can serve society and students and even with possibilities for citizen science
256 to be developed. There is also a wish for methodological coordination across the cryospheric research fields with, for instance,
257 time-lapse cameras, remote sensing (InSAR), fibre-optic cables for high spatial observations, ground penetrating radar,
258 passive/active seismic and cryoseismic measurements, and potentially networks of these types of measurements across
259 different SIOS supersites. Supersites are also instrumental for developing and harmonising methods to ensure that all collected
260 data are comparable and immediately usable.

261 Finally, it would be optimal to equip SIOS for observing sudden cryospheric events affecting the cryosphere such as glacial
262 lake drainage (GLOFs) or meteorological extreme events such as autumn or winter rainstorms causing landslides and increased
263 runoff due to extensive snow melting. These topics are also important for interdisciplinary studies.

264 **4.1.3 Marine**

265 The marine synthesis is based on contributions covering the following topics: oceanic circulation, ocean-atmosphere
266 interactions, plankton monitoring, sea ice thickness, microplastic pollution and the status of Svalbard coastal waters. Major
267 recommendations identified a need for a marine observing infrastructure development that provides data needed to fill critical
268 gaps in the observing system. Specifically, these included autonomous observational sites for co-located long-term monitoring
269 of ocean physics, biogeochemistry and ecology (with the latter two currently being under sampled with continuous methods),
270 enhanced sampling capabilities during the winter and early spring season, as well as spatially increased data collection by a
271 wider involvement of ships of opportunity. This recommendation could be addressed by specific funding via SIOS access and
272 optimisation calls that provide access to existing infrastructure and aid the development of new infrastructure. Homogeneous
273 data collection in disparate areas and by various teams could be ensured by a dedicated programme for harmonisation of marine
274 measurements around Svalbard. To fulfil this need, marine infrastructure network workshops have been organised by SIOS,
275 encouraging further cooperation on joint scientific programs. This network would be well suited to develop a handbook of best
276 practices for ocean observing approaches and data management. The four most essential recommendations for marine ESS
277 Observing System development were identified: (i) to extend the geographical and temporal coverage of research activities,

278 (ii) to enhance year-round observations to resolve seasonal variability in the ocean, (iii) to support research activities exploring
279 linkages between fjord, shelf and open ocean systems, and (iv) to establish long-term year-round monitoring of marine biota.

280 **4.1.4 Terrestrial**

281 Several recommendations relate to infrastructure, wider data collection and remote sensing. Measurements on varied spatial
282 scales and development of models are mentioned frequently, but still need further implementation. Notably, there is no
283 contribution on lake systems and just one recommendation for experimental design. Overall, there is a need for co-location of
284 measurements to integrate various disciplines, temporal and spatial scales, experimental manipulations and to facilitate model-
285 based quantitative analysis. Current examples include the terrestrial flagship in Ny-Ålesund (Pedersen et al. 2022), the COAT
286 program led by the Norwegian Polar Institute (Ims et al., 2013), and the Bjørndalen Integrated Gradients studies initiated by
287 the University Centre in Svalbard. COAT is a national ecosystem-based monitoring system of low and high Arctic terrestrial
288 environments, developed as a joint effort by the Fram centre institutions already in 2013 before SIOS was initiated. COAT is
289 now fully integrated into SIOS, and the two projects have received joint Norwegian infrastructure funding. These studies
290 should further develop, with new methods and technologies facilitating more spatially and temporally extensive and high-
291 resolution automated measurements of biological parameters. There needs to be a focus on long-term ecosystem-based
292 monitoring to establish how various anthropogenic pressures affect the Arctic environment. Climate change is likely to
293 transform Arctic terrestrial ecosystems beyond scientists' current abilities to make predictions. Therefore it is important to
294 have scientifically robust ecosystem-based, integrated systems to detect the fast changes from climate change, which is the
295 overall overriding threat, on terrestrial ecosystems.

296 **4.2 Knowledge gaps and associated improvement potential in Svalbard Earth System Science**

297 In addition to these environment-specific recommendations, interdisciplinary and overall recommendations that all four SESS
298 environments have brought up have been identified, and further synthesised for increasing and improving the interdisciplinarity
299 of SIOS.

300 With Svalbard experiencing the largest ongoing warming in the Arctic and worldwide and with the closely associated further
301 environmental changes, this puts SIOS in a unique position to provide coordinated datasets to investigate ongoing processes
302 in the Earth-system in a coupled manner. SIOS can quantify and understand in detail all physical processes that ultimately
303 warm the atmosphere, turning ice into water with all the resulting consequences this has on the entire Earth System. This
304 warming is controlled by the atmosphere and changes in ocean currents and impacts the cryosphere, as well as the terrestrial
305 and marine environments. By using the entire observation network of SIOS to perform a full-scale study of all the processes
306 which impact Earth System dynamics controlling the water cycle, a major potential for increasing the understanding of key
307 mechanisms in the Earth System has been identified. Such a study has potential to provide large knowledge gain, and thus
308 further the understanding of effects on the Svalbard Earth System.

309 Many cross-cutting individual SESS recommendations indicate the relevance and need of atmospheric parameters and
310 observations for studying the other environments in SIOS. Also, cross-cutting actions between the atmosphere, hydrosphere
311 and cryosphere have been indicated as areas of clear interest. Influences on the upper atmosphere from within the Sun-Earth
312 system, unique to the polar regions, have been shown to effect ozone depletion down to stratospheric altitudes (30km) and that
313 this may become more prominent with climate change (Maliniemi et al. 2020). In turn, lower atmospheric processes, such as
314 waves and tides, can influence the dynamics and behaviours of the upper atmosphere (Stober et al. 2021). The degree of
315 influence is a topic of debate, with global models failing to reproduce some of the observed effects (e.g., Oliver et al. 2013,
316 Cnossen 2020). Precipitation and wind action in the lower atmosphere result in long- and short-range transport of dust and
317 aerosols and accumulation of black carbon on snow and glaciers. Arctic terrestrial ecosystems, such as vegetation and its
318 components, and microbial communities, are impacted by air temperature, wind and precipitation, all of which are directly
319 controlled by the atmosphere. For instance, spectral characteristics of downwelling flux influence vegetation and microbial
320 community. Melting of snow and glaciers, coupled with thawing of permafrost and an increase in active layer depth, supply
321 water, sediment, microbes and aqueous chemical species through the terrestrial environment into the marine environment, via
322 the hydrological system. The pathway from cryosphere through terrestrial to marine thus links all spheres in SIOS. Thus, all
323 these identified cross-cutting topics clearly have large potential if coordinated to fill the identified knowledge gap of the water
324 cycle and its consequences.

325 With the bottom-up approach of the SESS contributions, the development of a perspective from within the four environments
326 was clearly a very natural starting point for synthesising. To start increasing the interdisciplinarity and cross-cutting nature of
327 SIOS, a next step is to increase the dialogue between the different environments. Potentially, dedicated workshops on specific
328 topics such as radiative forcing between two or more SIOS environments could be a starting point. Such workshops should
329 also be tasked with pointing out optimal observations needed for datasets to fill and address the overall identified cross-cutting
330 knowledge gap.

331 Additionally, all environments of SIOS depend on obtaining better spatial and temporal coverage of observations in and around
332 Svalbard. This can be achieved through increased and coordinated instrument deployments that better cover the
333 physiographically different regions of Svalbard including various landscape/seascape/icescape types and boundaries between
334 them (e.g., the coastal zone). As warming is greatest in the eastern part of the Svalbard area (Isaksen et al., 2022), it is the
335 ambition to expand the SIOS observation system to also cover the eastern and northern part of Svalbard. Given the high
336 ambition for preserving these relatively untouched parts of the archipelago of Svalbard, such expansion might have to be
337 mainly based on remote sensing observations and/or dedicated supersites based mainly on the use of new low-maintenance
338 technology. For example, the findings regarding Arctic vegetation and its climatic control by Bjorkman et al. (2020) highlight
339 the need for more geographically widespread, integrated, and comprehensive monitoring efforts that can better resolve the
340 interacting effects of warming and other local and regional ecological factors. In this case, studies across the entire vegetation
341 zonation of Svalbard are important to fully understand the effects of climatic changes to the entire flora of the different Arctic

342 vegetation zones. The spatially distributed network approach by COAT with large spatial and temporal coverage and co-
343 location of measurements covering the terrestrial above-ground food web is another example. COAT measures vegetation
344 from 57 module stations covering large parts of Nordenskiöldland, Isfjorden and Brøggerhalvøya in central Svalbard. There
345 is a clear need across all the environments for special focus on the autumn and spring periods.

346 While the current focus on the major science hubs represents a disadvantage in terms of covering more different areas and
347 habitat types as detailed above, there is clear potential for co-location of all the different types of observations from all parts
348 of the ESS, this way developing real supersites. This should complement existing infrastructure, with a high level of
349 coordination, long-term planning, and funding. It would have the advantage of fewer sites to maintain, allowing for a more
350 sustainable observation system. It would also allow for focusing on much closer collaboration on new methods and
351 technologies across SIOS, with a potential for more efficient sharing of knowledge. Knowledge gained from the supersites
352 may be used to guide the selection of a subset of measurement types for observations with higher spatial and temporal coverage.
353 An ambitious tool identified is dedicated method-based action forces, that would make SIOS able to respond quickly to extreme
354 events such as solar storms, sudden stratospheric warmings (SSWs), glacial lake outburst floods (GLOFs) or various
355 meteorological extremes, such as rainstorms and avalanches, the latter of which are also highly relevant to be able to perform
356 a full-scale study of the water cycle changes and its consequences.

357 Likewise, the need for real-time access to SIOS observations are important, particularly for extreme events, for increasing the
358 return of data from desirable but high-risk deployments (such as moorings or instruments near or on surging glaciers, or rocket
359 launches into specific atmospheric phenomena), but also for more widespread use of the SIOS data.

360 **5. Discussion and conclusion**

361 This synthesis of knowledge gaps based on the first four years of operation of the Svalbard Integrated Arctic Earth Observing
362 System (SIOS) builds on a unique and strong international collaboration. No other observing system exists which is this wide-
363 ranging, covering all part of ESS within a specific regional area, yet at the same time supported so widely internationally and
364 so well-coordinated. When developing a full-scale Earth science observation system as SIOS, with the comprehensive full
365 scale ESS ambition that SIOS has, it is very natural for researchers within the individual disciplines and even subdisciplines
366 to first start working closer together before being able to focus more on the interdisciplinarity necessary for performing full-
367 scale ESS analysing how the observed parameters influence each other. This is clearly the case for SIOS after the first four
368 years of operation.

369 Ways to accommodate this important next step include the implementation of above-mentioned workshops on interdisciplinary
370 topics as well as the development of supersites. Also, a workshop across all of SIOS working groups is planned to increase the
371 collaboration and interdisciplinarity of SIOS. Hopefully also scientific collaboration between two environments might start
372 being more developed in SIOS, such as already developed in the fully ecosystem-based observation system of COAT
373 combining climate and ecology, designed to observe, understand and predict outcomes from climate change (drivers) on tundra

374 ecosystems (Ims & Yoccoz, 2017). In addition SIOS is operating training activities in the form of small courses on different
375 key SIOS observation techniques, SIOS data availability and SIOS remote sensing opportunities for all interested. Increased
376 interdisciplinarity can also be obtained by further developing the major science hubs in Svalbard into real supersites
377 complementing existing infrastructure allowing even closer collaboration on particular new technologies and methods
378 including real-time access to observations across SIOS. Just as obtaining better spatial and temporal observation coverage in
379 and around Svalbard, including the physiographically different regions of Svalbard, both the various land-, sea- and icescapes
380 and boundaries between them in the coastal zone, would lead to increased potential for interdisciplinarity.

381 Other ways forward will be to increase the use of the SIOS observations through the SIOS data catalogue for Earth System
382 modelling, to obtain improved process understanding that bridges science disciplines. This will demand working to expand
383 the SIOS community with modellers from different scientific fields, just as increased collaboration between modellers and the
384 observational groups in SIOS needs to be established to enable observing at the relevant scales and resolutions, and of missing
385 parameters. The Earth system modelling community could make little use of high resolution regional data in the early stages
386 of SIOS but presently as model resolution approaches sub-regional scales it becomes increasingly relevant to utilise models
387 for observation design as well as accrued data to calibrate and verify model output. The SIOS Observation Facility Catalogue
388 and the Research in Svalbard database are important tools for developing interdisciplinarity in Svalbard ESS. Just as the SESS
389 reports are an integral part of the strategic development of the observing system. This bottom-up approach, which is
390 complemented with rising societal needs and member priorities, leads to the outcomes from the SIOS Science wheel (Fig. 3).
391 In addition, the harmonised SIOS core data, new research infrastructure and research projects are all innovations leading to
392 increased interdisciplinarity, highly encouraged through the SIOS optimization and access calls. Finally, SIOS offers
393 Innovation Awards to encourage further developing, also high risk – high gain, technological and/or other ESS key activities
394 that address knowledge gaps identified by the SIOS community.

395 Further, as Svalbard is experiencing the largest ongoing warming in the Arctic and worldwide, this puts SIOS in a unique
396 position to perform a full-scale study of all processes impacting ESS dynamics controlling the water cycle using all parts of
397 the SIOS observation network, with large potential for increasing the understanding of key mechanisms in the Earth System.
398 The clear identification of a joint scientific focus, studying the water cycle changes and its consequences using all parts of the
399 SIOS ESS observation network, will function well also as a focal point for cross-disciplinary work. Despite a disciplinary
400 focus, snow, precipitation and hydrology have been identified as overlapping topics in many SESS contributions, but not
401 prominent in the marine environment. These topics are therefore connected to all environments primarily on land, but observed
402 and treated in different ways within the different environments. This is important for the identified scientific focus on the
403 changing water cycle. Hence, SIOS has potential to develop a multidisciplinary study with data products and modelling
404 approaches based on the variety of observations that is being collected and supplement this with additional necessary new
405 observations leading to increased Earth System understanding of current and future dynamics.

406 There is also a need for increased regional collaboration using SIOS in scales broader than just Svalbard, when aiming for full
407 Earth System understanding within the Arctic. This means collaborating primarily with the rest of the Arctic. The international
408 ‘Sustaining Arctic Observing Networks’ (SAON) is the regional facilitator aiming at coordinating, improving, integrating and
409 sustaining pan-Arctic observations during rapid environmental and social change, with a main focus on specifically addressing
410 key gaps in coordination (Starkweather et al., 2021). With SIOS in the warmest part of the Arctic, and the largest climatic
411 gradient in the high Arctic existing between Svalbard and cold North Greenland, closer collaboration with Greenland could be
412 a first natural step. In Greenland, the main aim of the ‘Greenland Integrated Observing System’ (GIOS, <https://gios.org/>) is to
413 resolve and understand the mechanisms behind climate and environmental change in Greenland and beyond. GIOS is
414 developing a network of sustainable long-term research infrastructures in and around Greenland observing the changing air,
415 ice, land, and ocean conditions. GIOS is, as opposed to SIOS, a national research infrastructure linking all institutions and
416 universities currently carrying out Arctic research in the Danish Realm. SIOS and GIOS have started discussing increasing
417 collaboration within research, higher education and logistics. Interhemispheric co-operation is also another obvious step with
418 networks in the Antarctic Region such as SCAR (Scientific Committee on Antarctic Research) as an obvious collaboration
419 partner. Initiatives in specific disciplines, such as GRAPE (Global Navigation Satellite System Research and Application for
420 Polar Environment) are already in place. The research outlined through SIOS would benefit from being part of global observing
421 systems such as the SuperDARN (Super Dual Auroral Radar, Chisham et al. 2007) atmospheric network. Such global
422 networks, with commensurable data products, allow regional effects to be placed in a global context.

423 SIOS is making use of the Earth Observing System (EOS) program of NASA with its different satellite missions and scientific
424 instruments comprised of a series of coordinated polar-orbiting satellites designed to monitor and understand key components
425 of the climate system and their interactions through long-term global observations of the land surface, the biosphere,
426 atmosphere and oceans (Platnik, 2022). This also enables direct comparison between Svalbard and particularly the rest of the
427 Arctic, of the different ESS components which are being remotely observed. To make best use of this data, links between
428 existing EOS data need to be established, and data need to be available in formats that global Earth system modelling can make
429 use of.

430 Finally, many of the observations collected by regional observation systems like SIOS have large potential to add to improved
431 understanding of how people living in the Arctic are affected by climatic changes in many different ways (e.g., by landslides,
432 avalanches, coastal erosion). COAT has a clear goal of providing management relevance. Clearly, also SIOS can assist humans
433 and societies to develop management of climate change impacts. Such assistance is needed for developing resilience and
434 adaptation to enable Arctic communities to deal with the consequences of the ongoing climatic changes. Presently, with the
435 move towards involving local communities directly in climate change observations and research, this might place SIOS in a
436 new role as also contributing more directly to societal needs.

437 **Data Availability**

438 Data for Figure 1 is available through the Norwegian Meteorological Institute online service. Data for Figure 2 is
439 available in the SIOS Observation facility catalogue.

440 **Author contributions**

441 All authors have contributed to the SIOS synthesis process, and all have contributed to writing this manuscript.

442 **Competing interests**

443 The authors declare that they have no conflict of interest.

444 **Disclaimer**

445 **Acknowledgements**

446 The authors would like to acknowledge the support from the Research Council of Norway through projects No. 291644,
447 No.322387 & No. 269927 and Norwegian Space Agency through contract No. NIT 12.20.5. We thank our SIOS colleagues
448 who have contributed with SESS reporting, and those who assisted us during the thematic workshops to start synthesising the
449 SESS recommendations. Thanks to Åshild Ø. Pedersen, Norwegian Polar Institute for comments improving the revised
450 manuscript on terrestrial biology. Finally, we very much appreciate the comments by two anonymous reviewers, which have
451 largely improved the revised manuscript.

452 **References**

453 Assmy, P, Kvernvik, A.C., Hop, H, Hoppe, C.J.M., Chierici, M., David, D.T., Duarte, P., Fransson, A., Martin Garcia, L., Kwasniewski, S.,
454 Maturilli, M., Pavlova, O., Tatarek, A., Wiktor, J.M., Wold, A., Wolf, K.K.E., Bailey, A., 2023, Plankton dynamics in Kongsfjorden during
455 two years of contrasting environmental conditions. *Progress in Oceanography*, <https://doi.org/10.1016/j.pocean.2023.102996>.

456 Bintanja, R., Selten, F., (2014). Future increases in Arctic precipitation linked to local evaporation and sea-ice retreat. *Nature* 509, 479–482.
457 <https://doi.org/10.1038/nature13259>

458 Bischof, K., Convey, P., Duarte, P., Gattuso, J.-P., Granberg, M., Hop, H., Hoppe, C., Jiménez, C., Lisitsyn, L., Martinez, B., Roleda, M.Y.,
459 Thor, P., Wiktor, J.M., Gabrielsen, G.W., 2019. Kongsfjorden as Harbinger of the Future Arctic: Knowns, Unknowns and Research
460 Priorities, in: Hop, H., Wiencke, C. (Eds.), *The Ecosystem of Kongsfjorden, Svalbard*. Springer International Publishing, Cham, pp. 537–
461 562. https://doi.org/10.1007/978-3-319-46425-1_14


462 Bjorkman, A.D., García Criado, M., Myers-Smith, I.H. et al., 2020. Status and trends in Arctic vegetation: Evidence from experimental
463 warming and long-term monitoring. *Ambio* 49, 678–692.

464 Chisham G, Lester M, Milan SE, Freeman MP, Bristow WA, Grocott A, McWilliams KA, Ruohoniemi JM, Yeoman TK, Dyson PL,
465 Greenwald RA, 2007. A decade of the Super Dual Auroral Radar Network (SuperDARN): Scientific achievements, new techniques and
466 future directions. *Surveys in Geophysics* 28(1):33–109 437

- 467 Cnossen, I., 2020. Analysis and attribution of climate change in the upper atmosphere from 1950 to 2015 simulated by WACCM-X.
468 Journal of Geophysical Research: *Space Physics*, 125, e2020JA028623. <https://doi.org/10.1029/2020JA028623>
- 469 Constable, A.J., S. Harper, J. Dawson, K. Holsman, T. Mustonen, D. Piepenburg, and B. Rost, 2022. Cross-Chapter Paper 6: Polar
470 Regions. In: Climate Change 2022: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Sixth Assessment
471 Report of the Intergovernmental Panel on Climate Change [H.-O. Pörtner, D.C. Roberts, M. Tignor, E.S. Poloczanska, K. Mintenbeck, A.
472 Alegría, M. Craig, S. Langsdorf, S. Löschke, V. Möller, A. Okem, B. Rama (eds.)]. Cambridge University Press, Cambridge, UK and New
473 York, NY, USA, pp. 2319–2368, <https://doi:10.1017/9781009325844.023>
- 474 Descamps, S., Aars, J., Fuglei, E., Kovacs, K.M., Lydersen, C., Pavlova, O., Pedersen, Å.Ø., Ravolainen, V. and Strøm, H., 2017. Climate
475 change impacts on wildlife in a High Arctic archipelago – Svalbard, Norway. *Glob Change Biol*, 23: 490-502.
476 <https://doi.org/10.1111/gcb.13381>
- 477 Efstathiou, E., Eldevik, T., Arthun, M. & Lind, S., 2022. Spatial patterns, mechanisms and predictability of Barents Sea ice change. *J.*
478 *Clim.* 35(10), 2961–2973. <https://doi.org/10.1175/JCLI-D-21-0044.1>
- 479 Feldner J., Hübner C., Lihavainen H., Neuber R., Zaborska A. (eds), 2022. *SESS report 2021*, Svalbard Integrated Arctic Earth Observing
480 System, Longyearbyen. https://sios-svalbard.org/SESS_Issue4.
- 481 Hall, C.M., Tsutsumi, M., 2020. Neutral temperatures at 90 km altitude over Svalbard (78°N 16°E), 2002–2019, derived from meteor radar
482 observations, *Polar Science*, Volume 24, 100530, ISSN 1873-9652, <https://doi.org/10.1016/j.polar.2020.100530>.
- 483 Hansen B.B., Isaksen K., Benestad R.E. et al., 2014. Warmer and wetter winters: characteristics and implications of an extreme weather
484 event in the High Arctic. *Environmental Research Letter*, 9, 114021
- 485 Holmen, S.E., Dyrland, M.E. & Sigernes, F., 2014. Mesospheric temperatures derived from three decades of hydroxyl airglow
486 measurements from Longyearbyen, Svalbard (78°N). *Acta Geophys.* 62, 302–315. <https://doi.org/10.2478/s11600-013-0159-4>
- 487 Ims, R.A., Jepsen, J.U., Stien, A. & Yoccoz, N.G., 2013. Science plan for COAT: Climate-ecological Observatory for Arctic Tundra.
488 *Fram Centre Report Series 1*, Fram Centre, Norway, 177 pages.
- 489 Ims, R.A. & Yoccoz, N.G., 2017. Ecosystem-based monitoring in the age of rapid climate change and new technologies, Current Opinion
490 in *Environmental Sustainability*, 29, 170-176, <https://doi.org/10.1016/j.cosust.2018.01.003>.
- 491 IPCC Climate Change 2021: The Physical Science Basis. Working Group I Contribution to the IPCC Sixth Assessment Report.
- 492 Isaksen, K., Nordli, Ø., Ivanov, B., Køltzow, M.A.Ø., Aaboe, S., Gjeltén, H.M., Mezghani, A., Eastwood, S., Førland, E., Benestad, R.E.,
493 Hanssen-Bauer, I., Brækkan, R., Sviashchennikov, P., Demin, V., Revina, A., Karandasheva, T., 2022. Exceptional warming over the
494 Barents area. *Scientific Reports* 12:9371. <https://doi.org/10.1038/s41598-022-13568-5>
- 495 Lastovicka, J., Urbar, J., Kozubek, M., 2017. Long-term trends in the total electron content. *Geophys. Res. Lett.*, 44, 8168–8172,
496 <https://doi:10.1002/2017GL075063>
- 497 Maliniemi, V, Marsh, R Daniel, Tyssøy, H.N, Smith-Johnsen, C, 2020. Will climate change impact polar nox produced by energetic
498 particle precipitation?, *Geophysical Research Letters* 47(9), 2020–087041
- 499 Meredith, M., M. Sommerkorn, S. Cassotta, C. Derksen, A. Ekaykin, A. Hollowed, G. Kofinas, A. Mackintosh, J. Melbourne-Thomas,
500 M.M.C. Muelbert, G. Ottersen, H. Pritchard, and E.A.G. Schuur, 2019: Polar Regions. In: IPCC Special Report on the Ocean and
501 Cryosphere in a Changing Climate [H.-O. Pörtner, D.C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K.
502 Mintenbeck, A. Alegría, M. Nicolai, A. Okem, J. . Petzold, B. Rama, N.M. Weyer (eds.)].
- 503 Moreno-Ibáñez M., Hagen J.O., Hübner C., Lihavainen H., Zaborska A. (eds), 2021. *SESS report 2020*, Svalbard Integrated Arctic Earth
504 Observing System, Longyearbyen. https://sios-svalbard.org/SESS_Issue3.

- 505 Müller, M. et al., 2022. Decline of sea-ice in the Greenland Sea intensifies extreme precipitation over Svalbard. *Weather and Climate*
506 *Extremes*, 36, 100437, <https://doi.org/10.1016/j.wace.2022.100437>.
- 507 Nordli, O. et al., 2020. Revisiting the extended Svalbard Airport monthly temperature series, and the compiled corresponding daily series
508 1898–2018. *Polar Res.* <https://doi.org/10.33265/polar.v39.3614>
- 509 Norges forskningsråd, 1997. Organisering av forskningen på Svalbard, Norwegian Research Council report, ISBN 82-12-00864-9.
- 510 Oliver, W. L., S.-R. Zhang, and L. P. Goncharenko, 2013. Is thermospheric global cooling caused by gravity waves?, *J. Geophys. Res.*
511 *Space Physics*, 118, 3898–3908, <https://doi.org/10.1002/jgra.50370>
- 512 O'Connell, M.J., Huiskes A.H.L., Loonen M.J.J.E., Madsen J., Klaassen M. & M. Rounsevell, 2006. Developing an integrated approach to
513 understanding the effects of climate change and other environmental alterations at a flyway level. Waterbirds around the world. Eds. G.C.
514 Boere, C.A. Galbraith & D.A. Stroud. pp. 385-397.
- 515 Orr, E., Hansen, G., Lappalainen H., Hübner C., Lihavainen, H. (eds) 2019: *SESS report 2018*, Svalbard Integrated Arctic Earth Observing
516 System, Longyearbyen. https://sios-svalbard.org/SESS_Issue1
- 517 Pedersen, Å. Ø., Stien, J., Albon, S., Fuglei, E., Isaksen, K., Liston, G. & Ims, R. A., 2020. Climate-ecological observatory for Arctic
518 Tundra (COAT). *State of Environmental Science in Svalbard (SESS) Report 2019*, 58-83.
- 519 Pedersen, Å. Ø., Convey, P., Newsham, K. K., Mosbacher, J. B., Fuglei, E., Ravolainen, V., Hansen, B. B., Jensen, T. C., Augusti, A.,
520 Biersma, E. M., Cooper, E. J., Coulson, S. J., Gabrielsen, G. W., Gallet, J. C., Karsten, U., Kristiansen, S. M., Svenning, M. M., Tveit, A.
521 T., Uchida, M., ... Loonen, M. J. J. E., 2022. Five decades of terrestrial and freshwater research at Ny-Ålesund, Svalbard. *Polar Research*,
522 41, [6310]. <https://doi.org/10.33265/polar.v41.6310>
- 523 Peeters, B. et al., 2019. Spatiotemporal patterns of rain-on-snow and basal ice in high Arctic Svalbard: detection of a climate-cryosphere
524 regime shift. *Environmental Research Letters*, 14, 015002. <https://doi.org/10.1088/1748-9326/aaefb3>
- 525 Platnik, S., 2022. "Earth Observing System". *NASA's Earth Observing System*.
- 526 Serreze, et al., 2015. Extreme daily precipitation events at Spitsbergen, an Arctic Island. *International Journal of Climatology*, 35, 4574-
527 4588. <https://doi.org/10.1002/joc.4308>
- 528 St.meld.nr.22, 2009, Stortingsmelding nr. 22 (2008-2009) Svalbard, Det Kongelige Justis- og Politidepartement, Norwegian Parliament
529 White paper, in Norwegian.
- 530 Starkweather,S., Larsen,J.R., Kruemmel,E., Eicken,H., Arthurs,D., Bradley, A.C., Carlo,N., Christensen, T., Daniel,R., Danielsen,F.,
531 Kalhok, S., Karcher, M., Johansson, M., Jóhannsson, H., Kodama, Y., Lund, S., Murray, M.S., Petäjä, T., Pulsifer, P.L., Sandven, S.,
532 Sankar, R.D., Strahlendorff, M., & Wilkinson, J., 2021. Sustaining Arctic Observing Networks' (SAON) Roadmap for Arctic Observing
533 and Data Systems (ROADS). *Arctic*, 74, SUPPL. 1 (2021) P. 56 – 68, <https://doi.org/10.14430/arctic74330>
- 534 Stober, G., Kuchar, A., Pokhotelov, D., Liu, H., Liu, H.-L., Schmidt, H., Jacobi, C., Baumgarten, K., Brown, P., Janches, D., Murphy, D.,
535 Kozlovsky, A., Lester, M., Belova, E., Kero, J., and Mitchell, N., 2021. Interhemispheric differences of mesosphere–lower thermosphere
536 winds and tides investigated from three whole-atmosphere models and meteor radar observations, *Atmos. Chem. Phys.*, 21, 13855–13902,
537 <https://doi.org/10.5194/acp-21-13855-2021>
- 538 Van den Heuvel F., Hübner C., Błaszczyk M., Heimann M., Lihavainen H. (eds) 2020: *SESS report 2019*, Svalbard Integrated Arctic Earth
539 Observing System, Longyearbyen. https://sios-svalbard.org/SESS_Issue2

- 540 Wurst, Sabine, M. Bittner, P. J. Espy, W. John R. French, and F. J. Mulligan, (2023) Hydroxyl airglow observations for investigating
541 atmospheric dynamics: results and challenges, *Atmos. Chem. Phys.*, 23, pp. 1599–1618, <https://doi.org/10.5194/acp-23-1599-2023>
- 542 Zdanowicz C., Gallet J.-C., Salvatori R., Malnes E., Isaksen K., Hübner C., Jones E., & Lihavainen H., 2023. An agenda for the future of
543 Arctic snow research: the view from Svalbard. *Polar Research*, 42. <https://doi.org/10.33265/polar.v42.8827>
- 544 Zhang, Shun-Rong, et al., 2016. "Ionospheric ion temperature climate and upper atmospheric long-term cooling." *Journal of Geophysical*
545 *Research: Space Physics* 121.9 (2016): 8951-8968. <https://doi.org/10.1002/2016JA022971>




547

Figure 1. The meteorological record from the Longyearbyen area is composed of the homogenised data series from the

548

Norwegian Meteorological Institute (1898-2022).



549 **Figure 2. SIOS observation infrastructure in Svalbard, with the three main science hubs Longyearbyen, Hornsund and**
 550 **Ny-Ålesund inset to present all the various types of observations in more detail.**
 551

AN OBSERVING SYSTEM FOR MANY



552

553 **Figure 3:** SIOS Science wheel that provides an overview of developing the observing system for many uses and
554 users.

555

Atmosphere	Cryosphere	Marine	Terrestrial
Support continued funding of large scale, multi-disciplinary infrastructure, such as the Svalbard SuperDARN radar	Science based coordination of the cryospheric observations	Development of the marine environmental observing infrastructure - autonomous observational sites	Focus on co-location of basic cryospheric observations and subsequent long-term studies on biota at similar spatial and temporal scales
Provision of data management and online open access portal to facilitate easier sharing of datasets	Integrated SIOS observations supersites	Provision of new datasets by e.g., enhancing measurement capabilities	Focus on new methods and technologies using automated sensors, animal trackers and cameras
Better harmonisation of the geographical and temporal coverage of atmospheric observations and applying standardized observation methods	Common cryospheric methodological infrastructure priorities	Harmonised methodologies for in situ observations and data harmonisation	Develop model-based quantitative analyses of ecosystem processes using SIOS datasets
Enhance the spatial scale of the atmospheric observational network	Establish SIOS action force for extreme events	Support for research exploring linkages between fjord, shelf and open ocean systems	Develop interphases between monitoring-based ecosystem science and end-users

557 **Table 1. Key synthesised recommendations identified with the four environments of SIOS.**