# 1. General comments

This well-written and informative editorial summarises the main findings from the recently published Baltic Earth Assessment Reports (BEARs) and the most important knowledge gaps identified in them. It also nicely puts the BEARs in context, explaining their position in the history of Earth system research and climate change assessments for the Baltic Sea basin, as well as their connection to the stakeholder-oriented fact sheet that the Baltic Earth Expert Network on Climate Change has recently produced.

Overall, this is work well done. I only have a few minor comments on its substance and a few more suggestions for improving the wording and presentation.

Thank you very much for your positive and helpful comments. We have implemented all requested changes. Please find our detailed response in red below.

# 2. Comments on substance and presentation

1. What does "today successfully completed" refer to? Do you mean that all the work of the working groups has been completed or that scenario simulations have been completed?

The working group on scenario simulations has been completed. However, this is an unimportant detail. Hence, we have deleted the phrase in the brackets.

2. has been warming. "is warming" implicitly predicts that the observed faster warming can be extrapolated to the future, which may or may not be true.

Corrected.

3. L103-104. This sentence could be interpreted to mean that this is the third BEAR assessment. Suggested reformulation, beginning from L102: The two BACC books adopted a format inspired by the IPCC assessment reports. The present special issue in Earth System Dynamics is the third assessment. It takes a new format of Baltic Earth Assessment Reports (BEARs), encompassing 10 peer-reviewed scientific journal articles.

Thank you. We have adopted your suggestion.

4. I only found 30 years (but not 25) from Lehmann et al. (2022).

Corrected.

5. L147-148. As an advance over the previous assessments?

Very good suggestion, which we have implemented.

6. would still result (doubling of pCO<sub>2</sub> is something that international climate policy very much strives to avoid).

Corrected.

7. L181-183. I think this sentence simplifies a bit too much. Even without GIA, the mean sea level rise in the Baltic Sea would differ from the global mean, due to (e.g.) the gravitational effect of the melting of Greenland ice. Furthermore, the changes in extremes might differ from those in the mean sea level, although (as noted on L185-186) there is no consensus that there would be a large difference.

We revised the sentence "Changes in relative sea level extremes will depend on the competition between the rising global mean sea level, the gravitational effect of the melting of the Greenland and Antarctic ice sheets, changes in wind fields, etc. and the regionally differing Glacial Isostatic Adjustment (GIA) resulting in land uplift or subsidence." The relative importance of the various drivers of changes in sea level extremes is not discussed here.

8. L186, If the ensemble size (rather than model weaknesses) is indeed the limiting factor, then there is no reason to expect that statistically significant changes would occur in the real world, where only one realization of climate change will be observed. Therefore, I would replace "probably because available climate model ensembles are too small" by something like "suggesting that these changes will likely be small compared with natural variability".

We agree and modified the sentence according to the reviewer's suggestion.

9. "Antarctic" should be "Greenland". Due to the gravitational effects, melting of Greenland will increase sea level much less in the Baltic Sea than in the global mean.

In the Baltic Sea, the fingerprint from the melting Antarctic ice sheet in the Baltic Sea is larger than from the Greenland ice sheet (Grinstedt, 2015, BACC II; his Fig. 14.2) because the zero line of the Greenland fingerprint pattern is located in the Baltic Sea. To avoid confusion, we deleted the sub-clause.

10. I guess this increase mainly affects the southern parts, and less if at all the north, where the absolute sea level rise and the GIA largely compensate each other. "increase in these coastal areas" would implicitly convey this.

Correct. We added "these"

11. Perhaps "recent brightening"? While brightening (i.e., increased solar radiation) has been observed in the last few decades, it was preceded by a general dimming between the Second World War and the 1980s, presumably due to increased aerosols.

We agree and added "recent"

12. L244-245. Adding surface waves to coupled atmosphere-ocean system models?

Corrected.

13. has not been considered?

Corrected.

14. L266-269. The lack of increasing solar radiation in summer is also most likely affected by the lack of time-varying aerosols in many of the RCMs. By contrast, the projected decrease in anthropogenic aerosol emissions in the GCM simulations leads to a future increase in solar irradiation (Boé et al. 2020, Climate Dynamics,

https://link.springer.com/article/10.1007/s00382-020-05153-1). If the projected decreases in aerosol emissions are realized, the GCM simulations should probably be judged more plausible regarding the changes in solar irradiation and perhaps also summer temperature.

We agree and deleted the misleading sentence. In Meier et al. (2022, Climate change in the Baltic Sea region: a summary) an explanation for the discrepancy is provided, following your argumentation.

15. cannot exhaustively take into account?

Corrected.

- 3. Technical comments
  - 1. Delete "more than". 19 is the exact number according to Table 1.

Corrected.

2. Biogeochemical

Corrected.

3. mean surface salinity?

Yes. Corrected.

- Figure 2. The y axis label should be "Temperature anomaly" rather than "Temperature".
   Corrected.
- 5. 2022a. Furthermore, the order of this and the previous reference (Meier et al., 2022b) should be switched.

Corrected.

Please find the revised manuscript below.

Thank you very much!

Markus on behalf of the co-authors

# **The Baltic Earth Assessment Reports**

H. E. Markus Meier<sup>1</sup>, Marcus Reckermann<sup>2</sup>, Joakim Langner<sup>3</sup>, Ben Smith<sup>4,5</sup> and Ira

Didenkulova<sup>6</sup>

<sup>1</sup>Department of Physical Oceanography and Instrumentation, Leibniz Institute for Baltic Sea Research Warnemünde, Rostock, Germany

<sup>2</sup>International Baltic Earth Secretariat, Helmholtz-Zentrum Hereon, Geesthacht, Germany

<sup>3</sup>Swedish Meteorological and Hydrological Institute, Norrköping, Sweden

<sup>4</sup>Hawkesbury Institute for the Environment, Western Sydney University, Australia

<sup>5</sup>Department of Physical Geography and Ecosystem Science, Lund University, Sweden

<sup>6</sup>Department of Mathematics, University of Oslo, Norway

Correspondence to: H. E. Markus Meier (markus.meier@io-warnemuende.de)

Abstract. Baltic Earth is an independent research network of scientists from all Baltic Sea countries that promotes regional Earth system research. Within the framework of this network, the Baltic Earth Assessment Reports (BEARs) were produced in the period 2019-2022. These are a collection of 10 review articles summarising current knowledge on the environmental and climatic state of the Earth system in the Baltic Sea region and its changes in the past (palaeoclimate), present (historical period with instrumental observations) and prospective future (until 2100) caused by natural variability, climate change and other human activities. The division of topics between articles follows the grand challenges and selected themes of the Baltic Earth Science Plan, such as the regional water, biogeochemical and carbon cycles, extremes and natural hazards, sea level dynamics and coastal erosion, marine ecosystems, coupled Earth system models, scenario simulations for the regional atmosphere and the Baltic Sea, and climate change and impacts of human use. Each review article contains an introduction, the current state of knowledge, knowledge gaps, conclusions and key statements, based on which recommendations are made for future research. In parallel, Baltic Earth's ongoing outreach work has led to the publication of an information leaflet on climate change in the Baltic Sea, which has been published in two languages so far, and the organisation of stakeholder conferences and workshops.

#### **1** Introduction

#### 1.1 BALTEX/Baltic Earth history

Baltic Earth is an international research network dealing with Earth system science of the Baltic Sea region (https://baltic.earth, last access: 4 February 2023). The catchment area of the Baltic Sea is about four times larger than the Baltic Sea surface and is part of mainly the countries Belarus, Denmark, Estonia, Finland, Germany, Latvia, Lithuania, Poland, Russia, and Sweden (Fig. 1). Baltic Earth is politically independent and focusses on research on the water and energy cycles, climate variability and climate change, water management and extreme events, and related impacts on marine and terrestrial biogeochemical cycles. Human impact on the Earth system in more general terms, i.e. the anthroposphere, defined as the part of the environment that is made or modified by humans for use in human activities, was added to the 2017 Baltic Earth Science Plan (https://baltic.earth/grandchallenges, last access: 4 February 2023).

Baltic Earth is the successor of the Baltic Sea Experiment (BALTEX) programme, which was founded in 1993 as a GEWEX continental-scale experiment (Global Energy and Water Exchanges, a core project of the World Climate Research Programme) (Reckermann et al., 2011). During Phase I (1993–2002), BALTEX was primarily devoted to hydrological, meteorological and oceanographic processes in the Baltic Sea drainage basin, hence focussed on physical aspects of the Earth system. During the second phase (Phase II: 2003–2012), the programme was expanded to encompass regional climate research, carbon and biogeochemical cycles, engagement with stakeholders and decision makers via assessment reports, as well as communication and education, i.e. organizing summer and winter schools and international master courses.

In 2013, Baltic Earth was launched with a novel science plan reinforcing efforts to address Grand Challenges on (1) salinity dynamics in the Baltic Sea, (2) land-sea biogeochemical linkages, (3) natural hazards and extreme events, (4) sea level and coastal dynamics, (5) regional variability of water and energy exchanges, and (6) multiple

drivers of regional Earth system changes (Meier et al., 2014). Working groups were initiated on coupled Earth system models, the Baltic Sea Model Intercomparison Project (BMIP), education, outreach and communication, and scenario simulations for the Baltic Sea.

Baltic Earth and its predecessor BALTEX have produced three extensive regional assessment reports since 2008. The first two (The BACC Author Team, 2008, and The BACC II Author Team, 2015) emphasised climate change and its impacts in the Baltic Sea region and were published as text books, while the third, the Baltic Earth Assessment Reports (BEARs), was published in the format of a special issue in *Earth System Dynamics*, in 2022. The BEARs and BACC assessment reports fill a gap relative to the assessment reports of the Intergovernmental Panel on Climate Change (IPCC), since the latter focus on global scales, and do not provide detailed local to regional information about the current state of knowledge on climate change and its impacts in the Baltic Sea region. The BEARs provide a comprehensive and up-to-date overview of the state-of-the-art research on the compartments of the Earth system in the Baltic Sea region encompassing processes in the atmosphere, on land and in the sea, including the marine and terrestrial ecosystems as well as processes and impacts related to the anthroposphere.

The BEARs wrap together the currently available published scientific knowledge, updating the second assessment report of climate change in the Baltic Sea basin (The BACC II Author Team, 2015) based on the scientific literature. The present BEAR Special Issue comprises 10 articles on the Baltic Earth Grand Challenges and working group topics including a summary of the current knowledge about past, present, and future climate changes for the Baltic Sea region. The articles encompass contributions by 109 authors from 14 countries and reference 2822 scientific articles and institutional reports in their synthesis effort.

#### 1.2 Baltic Sea region characteristics

The Baltic Sea is a semi-enclosed, shallow sea with limited water exchange with the world ocean and small tidal amplitudes. Situated in Northern Europe, the climate of the region is highly variable because it is located in the transition zone between maritime and continental climates, influenced by the North Atlantic and Arctic regions. The river discharge from the large catchment area causes a pronounced gradient in sea surface salinity from about 20 g kg<sup>-1</sup> in the Danish Straits region to about 2 g kg<sup>-1</sup> or even less in the northern and eastern reaches of the Baltic Sea. Hence, the Baltic Sea is brackish, with habitats of maritime species in the south-west and freshwater species in the north-east. The Baltic Sea catchment area covers an area of almost 20% of the European continent. 85 million people in 14 countries live in the catchment area, which stretches from the temperate, densely populated south to the subarctic wilderness in the north.

Episodically, large amounts of saltwater from the North Sea cross the sills, located in the Danish Straits, into the Baltic Sea and ventilate the Baltic Sea deepwater. These events require a period of about 20 days of easterly winds that lower the Baltic sea level, followed by a period of about equal duration of strong westerly winds that push saltwater into the Baltic Sea. These events are called Major Baltic Inflows (MBIs) and are important for the water exchange between the North Sea and the Baltic Sea. Mixing is low compared to other seas, with an origin at the lateral boundaries, because tidal amplitudes are very small and energetically unimportant.

In recent decades, environmental conditions in the Baltic Sea have changed considerably. For instance, since 1980 the Baltic Sea has been warming more than any other coastal sea (Fig. 2), causing shorter sea ice and snow covers over land during winter. Furthermore, rising nutrient inputs from land in the 1950s/60s, caused by population growth accompanied by sewage water release to the Baltic Sea and intensified usage of fertilisers from agriculture, led to eutrophication and spreading of hypoxic and anoxic areas. Since the 1980s, nutrient input into the Baltic Sea has been steadily decreasing, but this has not yet led to a significant improvement in oxygen conditions. Recent trends in acidification are smaller than in the world ocean, in particular in the northern Baltic Sea, because positive trends in alkalinity supply counteract the acidification.

## 2 Methods

Following the BACC<sup>1</sup> Author Team (2008) and the BACC II Author Team (2015), the BEAR project is an effort to summarise scientific knowledge about climate change and other drivers of Earth system changes and their impacts on the Baltic Sea region. The two BACC books adopted a format inspired by the IPCC assessment reports. The present special issue in Earth System Dynamics is the third assessment. It takes a new format of Baltic Earth Assessment Reports (BEARs), encompassing 10 peer-reviewed scientific journal articles. The assessed knowledge was extracted from the scientific literature such as peer-reviewed articles, reports from research institutions, and published datasets. Importantly, literature from non-governmental organisations with a political or economic interest, political parties and other stakeholder organisations was excluded from the assessment, ensuring that only scientific knowledge informed the assessment. The BEARs focus on publications after 2012/2013, the year of the editorial deadline of the second assessment report. Whenever possible, uncertainty levels of the BEAR results are classified based on a matrix of consensus within the scientific literature and the documented evidence of detected changes and their attributed drivers such as climate change and human use. For a high confidence of a certain statement, high levels of both scientific consensus and evidence cases are required. Instances of disagreement and knowledge gaps are documented and discussed, informing priorities for future research.

Together with the intergovernmental Baltic Marine Environment Protection Commission (HELCOM), Baltic Earth formed an Expert Network on Climate Change (EN CLIME). The aim of the expert network is to regularly produce a climate change fact sheet (CCFS, 2021<sup>2</sup>) from the BEAR material. In 2021, it was published for the first time (http://helcom.fi/ccfs, last access: 4 February 2023). The CCFS contains some background information, a map showing regional future climate changes for selected parameters under the greenhouse gas concentration scenario RCP4.5 and information about 34 variables, directly and indirectly affected by climate change. For each parameter, a general description, past and prospective future changes, other drivers than climate change (only for the indirect parameters), knowledge gaps, policy relevance and references are presented. More than 100 scientists contributed to the compilation of the fact sheet which was coordinated by the HELCOM secretariat. Updated versions are planned for intervals of seven years. Like the BEARs, the fact sheet was peer-reviewed and quality assured. It has

<sup>&</sup>lt;sup>1</sup> Assessment of Climate Change in the Baltic Sea Basin (BACC); <u>https://baltic.earth/bacc</u>, last access: 4 February 2023

<sup>&</sup>lt;sup>2</sup> <u>https://helcom.fi/wp-content/uploads/2021/09/Baltic-Sea-Climate-Change-Fact-Sheet-2021.pdf</u>, last access: 4 February 2023

so far been translated to German (Ostsee Klimawandel Faktenblatt, 2022<sup>3</sup>) and translations to other languages are planned, enhancing accessibility to stakeholders.

In this editorial, we highlight the main findings and knowledge gaps as detailed by the BEARs and future work is proposed.

#### **3 Results**

A number of the main findings of the 10 BEARs are selected and highlighted below. The BEARs are either based on the Baltic Earth Grand Challenges or Baltic Earth Special Topics (Baltic Earth Science Plan, 2017).

- 1. Salinity dynamics of the Baltic Sea, Grand Challenge 1 (Lehmann et al., 2022): Salinity is an important parameter for the circulation and the marine ecosystem in the Baltic Sea. Any changes in salinity are caused by changes either in the freshwater inflow from rivers and net precipitation over sea or in the water exchange between the Baltic Sea and the adjacent North Sea. Although long-term records of salinity and its drivers suffer from data gaps, these records starting in the 19<sup>th</sup> century are globally unique. Major research efforts focused on the MBI event in 2014 and its consequences for water masses, oxygen concentration and biogeochemical cycling. During the event, an unexpectedly large contribution of oxic intrusions at intermediate depth and essentially nonturbulent conditions in the deep interior were found, emphasizing the importance of boundary mixing. A revised reconstruction of the long-term record of MBIs showed no trend but a pronounced multi-decadal variability with a period of about 30 years. Despite intense research activities, observed variations in the intensity and frequency of MBIs and related Large Volume Changes (LVCs) could not be attributed to atmospheric circulation variability. Hence, on time scales larger than the synoptical time scale, MBIs are not predictable. As an advance over the previous assessments, salinity dynamics of the various sub-basins and lagoons mainly based on observations have been discussed, documenting large regional differences.
- 2. **Biogeochemical functioning of the Baltic Sea**, Grand Challenge 2 (Kuliński et al., 2022): The review addresses the following topics: (1) terrestrial biogeochemical processes and nutrient inputs to the Baltic Sea, (2) the transformation of C, N and P in the coastal zone, (3) the production and remineralisation of organic matter, (4) oxygen availability, (5) the burial and turnover of C, N and P in sediments, (6) the Baltic Sea CO<sub>2</sub> system and seawater acidification, (7) the role of certain microorganisms in the biogeochemistry of the Baltic Sea and (8) the interactions between biogeochemical processes and chemical pollutants. It was found that oxygen depletion and the area of anoxic bottoms have still increased despite the reductions in nutrient inputs from land since the 1980s. Hence, the nitrogen pool has declined due to denitrification whereas the phosphorus loads, respectively, have accumulated in the Baltic Sea, while the remainder are either exported to the North Sea or lost via biogeochemical processes such as denitrification and burial. Furthermore, it was discovered that in the central and northern sub-basins the

<sup>&</sup>lt;sup>3</sup> <u>https://baltic.earth/ccfs</u>, last access: 4 February 2023

uptake of C, N and P during primary production does not correspond to the Redfield ratio, which strongly affects the relationship between primary production, export of organic matter and oxygen demand of the deep sea. While it is clear that the Baltic Sea is a  $CO_2$  sink in summer and a  $CO_2$  source in winter, the annual net balance remains unknown. The past increase in total alkalinity of unknown origin has entirely mitigated ocean acidification in the northern Baltic Sea and significantly reduced it in the central Baltic Sea. In the future, a doubling of atmospheric p $CO_2$  would still result in lower pH in the entire Baltic Sea, even if alkalinity should further increase.

- 3. Natural hazards and extreme events in the Baltic Sea region, Grand Challenge 3 (Anna Rutgersson et al., 2022): Existing knowledge is summarised about extreme events in the Baltic Sea region with a focus on the past 200 years with instrumental data as well as on future projections. Considered events are wind-storms, extreme waves, high and low sea levels, hot and cold spells in the atmosphere, marine heat waves, droughts, sea-effect snowfall, sea-ice ridging, extremely mild and extremely severe sea ice winters, heavy precipitation events, river floods, and extreme phytoplankton blooms. Furthermore, the knowledge about implications of these extreme events for society such as forest fires, coastal flooding, offshore infrastructure and shipping was assessed. With respect to the impacts of climate change, terrestrial and marine heat waves, extremely mild sea ice winters, heavy precipitation and high-flow events are expected to increase, while cold spells, severe sea ice winters and ice ridging are expected to decrease due to the increase in mean atmospheric temperature. Changes in relative sea level extremes will depend on the competition between the rising global mean sea level, the gravitational effect of the melting of the Greenland and Antarctic ice sheets, changes in wind fields, etc. and the regionally differing Glacial Isostatic Adjustment (GIA) resulting in land uplift or subsidence. Furthermore, projections suggest an increase of droughts in the southern and central parts of the Baltic Sea region mainly in summer. Significant future changes in wind-storms, extreme waves and sea level extremes relative to the mean sea level have not been found, suggesting that these changes will likely be small compared with natural variability.
- 4. Sea level dynamics and coastal erosion in the Baltic Sea region, Grand Challenge 4 (Weisse et al., 2021): In this study, the current knowledge about the diverse processes affecting mean and extreme sea level changes, coastal erosion and sedimentation with impact on coastline changes and coastal management is assessed. Such processes are GIA, contributions from global sea level changes, wind-storms, wind-waves, seiches or meteotsunamis. During 1886-2020, the mean absolute sea level in the Baltic Sea corrected for GIA increased by about 25 cm or ~2 mm year<sup>-1</sup> on average. Land uplift in the north is still faster than the absolute sea level rise while in the south the opposite is true with potential impacts on changes in coastal erosion and inundation. The current acceleration of sea level rise is small and could only be determined by spatially averaging observations at different tide gauge locations. Future sea level rise in the Baltic Sea is expected to further accelerate, probably somewhat less than the global mean sea level rise. The Baltic sea level is substantially more sensitive to melting from the Antarctic than from the Greenland ice sheet. Concerning sediment transports, the dominance of mobile sediments makes the southern and eastern Baltic Sea coasts susceptible to wind-wave induced transports, in particular during storms. Due to the global sea level rise, future sediment transports can be expected to increase in

these coastal areas, with a large spatial variability depending on the angles of incidence of incoming windwaves.

- 5. Human impacts and their interactions in the Baltic Sea region, Grand Challenge 6 (Reckermann et al., 2022): An inventory and discussion of the various man-made factors and processes affecting the environment of the Baltic Sea region and their interrelationships are presented. In total, 19 factors are addressed (Table 1). Some of the factors are natural and are only modified by human activities (e.g. climate change, coastal processes, hypoxia, acidification, submarine groundwater discharges, marine ecosystems, non-indigenous species, land use and land cover), others are entirely man-made (e.g. agriculture, aquaculture, fisheries, river regulation, offshore wind farms, shipping, chemical contamination, dumped ammunition, marine litter and microplastics, tourism and coastal management). All factors are interconnected to varying degrees. The knowledge of these linkages was assessed and analysed in depth. The main finding was that climate change has an overarching, integrating effect on all other factors and can be interpreted as a background effect that affects the other factors differently. After climate change, shipping and land use/agriculture have the strongest impacts, while fisheries, marine ecosystems and agriculture are the most affected. The results of the assessment depend on the region and may be different for other coastal seas and their catchments in the world, where different human activities prevail.
- 6. Global climate change and the Baltic Sea ecosystem: direct and indirect effects on species, communities and ecosystem functioning, Baltic Earth Special Topic (Viitasalo and Bonsdorff, 2022): Climate change has multiple impacts on species, communities and ecosystem functioning in the Baltic Sea through changes in physical and biogeochemical parameters such as temperature, salinity, oxygen, pH and nutrient levels. The associated secondary effects on species interactions, trophic dynamics and ecosystem function are also likely to be important. Climate change (warming, recent brightening, decrease in sea ice) has led to shifts in the seasonality of primary production, with a prolonged growing season of phytoplankton, an earlier onset of the spring bloom and a delayed autumn bloom. However, the development of cyanobacteria varies from species to species, and a clear causal relationship between temperature or salinity and the abundance of cyanobacteria has not been demonstrated. An increase in water temperature and river dissolved organic matter (DOM) could reduce primary production while favouring bacterial growth. If nutrient reduction proceeds, the improvement in oxygen conditions could initially increase zoobenthos biomass, but the subsequent decrease in sedimenting organic matter would likely disrupt the pelagic-benthic coupling and result in lower zoobenthos biomass. Sprat and some coastal fish species could be favoured by a rise in temperature. Regime shifts and cascading effects have already been observed in both pelagic and benthic systems as a result of climate change.
- 7. Coupled regional Earth system modeling in the Baltic Sea region, Baltic Earth Special Topic, with relevance to Baltic Earth Grand Challenge 5 (Gröger et al., 2021): Recent progress on the development of coupled climate models for the Baltic Sea region is assessed. Feedback mechanisms are important to simulate the response of the Earth system to external forcing such as greenhouse gas and aerosol emissions. In this review article, the coupling between (1) atmosphere, sea ice and ocean, (2) atmosphere

and land surface including dynamic vegetation, (3) atmosphere, ocean and waves and (4) atmosphere and hydrological components to close the water cycle is discussed. Adding surface waves to coupled atmosphere-ocean system models is becoming more important with increasing resolution, in particular when detailed information is required, for instance, for offshore wind energy applications in the coastal zone. Furthermore, the wave information is essential for the calculation of ocean mixing and resuspension. While long-term climate simulations using coupled atmosphere, sea ice and ocean models or coupled atmosphere and dynamic vegetation models have successfully been performed and their added value demonstrated, the impact of aerosols on the climate of the Baltic Sea region has not been considered. Coupling hydrology models to close the hydrological cycle is also still problematic, as the precipitation accuracy provided by the atmospheric models is, in most cases, insufficient to realistically simulate river discharge into the Baltic Sea without bias adjustments.

- 8. Atmospheric regional climate projections for the Baltic Sea region until 2100, Baltic Earth Special Topic (Christensen et al., 2022): Current climate projections based on regional climate atmosphere-only models of the EURO-CORDEX project with a horizontal resolution of 12.5 km under the scenarios RCP2.6, 4.5 and 8.5 are presented. As the number of simulations (124) is relatively large compared to previous assessments, the uncertainties can be better estimated than before. These projections indicate strong warming, especially in the north in winter, where warming approaches twice the average global warming. Precipitation is projected to increase throughout the Baltic Sea region, except in the southern half in summer, where the results are inconclusive. Extreme precipitation, here the 10-year return value, is projected to increase systematically throughout the study area, especially in summer. The large ensemble of simulations does not indicate a significant change in wind speed. Surface solar radiation is projected to remain unchanged in summer, but to decrease slightly in winter, due to increased cloud cover and possibly less snow in the future. Snow cover is projected to decrease dramatically, especially in the south of the Baltic Sea catchment. The comparison between the uncoupled model simulations of the EURO-CORDEX project and a small ensemble of scenario simulations performed with a coupled atmosphere-sea-ice-ocean model driven by a subset of global climate models indicates stronger warming in the coupled model during winter, mainly in areas that are seasonally affected by sea ice today. In summer, the coupled model shows weaker warming compared to the uncoupled models.
- 9. Oceanographic regional climate projections for the Baltic Sea until 2100, Baltic Earth Special Topic (Meier et al., 2022a): New projections of the future Baltic Sea climate with a coupled physical-biogeochemical ocean model were compared with previous projections. The differences are mainly due to different scenario assumptions and model setups. For example, the impact of future global sea level rise on salinity was previously neglected, but taken into account in the latest projections. Although the number of projections for the Baltic Sea is still small compared to regional atmospheric projections such as the EURO-CORDEX model ensemble, a relatively large ensemble of 48 scenario simulations allowed the assessment of uncertainties related to greenhouse gas emissions, global climate model differences, global sea level rise, nutrient inputs and natural variability. In the future climate, higher water temperatures, a shallower mixed layer with a sharper thermocline in summer, lower sea ice cover and stronger mixing in the northern Baltic Sea in winter compared to the current climate could be expected.

The assessment of marine heat wave changes is new. Both the frequency and duration of marine heat waves are projected to increase significantly, especially in the coastal zone of the southern Baltic Sea. Due to uncertainties in the projections regarding regional winds, precipitation and global sea level rise, no robust and statistically significant changes in salinity could be identified. The impacts of a changing climate on the biogeochemical cycle are projected to be significant, but still less than the plausible changes in nutrient inputs. Implementation of the proposed Baltic Sea Action Plan, a basin-wide nutrient input reduction plan, would lead to a significant improvement in the ecological status of the Baltic Sea, including a reduction in the size of the hypoxic area also in a future climate.

10. Climate change in the Baltic Sea region: A summary, Baltic Earth Special Topic (Meier et al., 2022b): In this comprehensive study, the recent knowledge on past (paleo-), present (historical) and projected future (< 2100) climate change in the Baltic Sea region, based upon all BEARs and >800 scientific articles, is summarised. It focuses on the atmosphere, the land surface, the cryosphere, the ocean and its sediments, and the terrestrial and marine biospheres. 33 parameters characterizing the state of these components of the Earth system were analysed (Fig. 3, Table 2). The anthroposphere is not part of this assessment but instead is discussed in detail by Reckermann et al. (2022). The main findings concerning changes of the 33 selected state parameters attributed to climate change are summarised in Figure 3. The prevailing causal relationships of climate change with sufficiently high confidence suggest a clear impact of global greenhouse gas emissions on regional heat cycles including all parameters of the cryosphere. However, changes caused by global warming of the water, momentum and carbon cycles are less clear because of either the large natural variability at regional scales or the impact of other drivers than global warming. Overall, it was concluded that the results from the previous BACC assessments mainly are still valid. However, new long-term, homogenous observational records, such as those for Scandinavian glacier inventories, sea-level-driven saltwater inflows (MBIs), or phytoplankton species distributions, and new scenario simulations with improved models, such as those for glaciers, lake ice, or marine food webs, have become available, resulting in a revised understanding of observed changes. Compared to previous assessments, observed changes in air temperature, sea ice, snow cover, and sea level were shown to have accelerated. However, natural variability is large, challenging our ability to detect observed and projected changes in climate of the Baltic Sea region. As the ensembles of scenario simulations both for the atmosphere and the ocean became larger, uncertainties can now be better estimated, although coordinated scenario simulations for the Baltic Sea based on ensembles of different regional ocean models are still missing. Furthermore, with the help of coupled models, feedbacks between several components of the Earth system have been studied, and multiple driver studies were performed, e.g., projections of the marine food web that include fisheries, eutrophication and climate change. Intensive research on the landsea interface, focusing on the coastal filter, has been performed, and nutrient retention in the coastal zone was estimated for the first time. However, a model for the entire Baltic Sea coastal zone is still missing, and the effect of climate change on the coastal filter capacity is still unknown. More research on changing extremes was performed, acknowledging that the impact of changing extremes may be more important than that of changing means (see also Rutgersson et al., 2022). However, many observational records are either too short or too heterogeneous (data gaps) for statistical studies of extremes.

## **4** Discussion

The identification of knowledge gaps in Earth system science of the Baltic Sea region, that should be further addressed in future research, was one of the main aims of the BEAR project. For specific knowledge gaps that have been identified during the project, the reader is referred to the individual assessment reports. However, as an overarching result, three new research topics are identified:

- 1) Small-scale processes and their impacts on large-scale climate dynamics and biogeochemical cycles. The number of observations in the sea is smaller than those over land. This is also true for the Baltic Sea, although the international long-term monitoring programme in the Baltic Sea started over a century ago, with measurements of temperature, salinity and oxygen concentration from the central parts of the various sub-basins. Nowadays, monitoring data are available from all sub-basins with a resolution of up to one month. Recently, many new observational systems for temporally and spatially high-resolution data have been developed or are under development, including remotely operated vehicles (ROVs) and autonomous underwater vehicles (AOVs) as well as remote sensing data. Examples of such systems operating in the Baltic Sea are continuously profiling moorings, ARGO floats, Gliders, ScanFish, and echo sounders. In addition to traditional physical parameters, measurements of turbulence, biogeochemical and biodiversity (e.g. environmental DNA) parameters are now available. Another area of research that is developing rapidly is numerical modelling of the Earth system, also on a regional scale, e.g. eddy- and submesoscale resolving multi-year simulations for the Baltic Sea. Corresponding arguments apply to the atmosphere, e.g. cloud resolving simulations to address heavy precipitation events. A novel research topic for Baltic Earth would therefore be to better understand the dynamics of small-scale atmospheric and oceanic processes that are not yet resolved in state-of-the-art numerical models or in conventional monitoring observations, and their role in the large-scale circulation on short and long time scales. Such research activities would help to answer some of the gaps in knowledge that have been raised by Lehmann et al. (2022), Kuliński et al. (2022), Rutgersson et al. (2022), Weisse et al. (2021), Viitasalo and Bonsdorff (2022), and Gröger et al. (2021). Furthermore, a realistic consideration of small-scale processes would improve the projections for the atmosphere (Christensen et al., 2022) and the ocean (Meier et al., 2022a).
- 2) Attribution of regional climate variability and change to anthropogenic radiative forcing versus other drivers. In order to unambigiously disentangle the impacts of anthropogenic climate change and other human influences from the natural climate variability of the regional Earth system, more knowledge about internal variations and feedback mechanisms is needed. For example, climate models have recently shown that multi-decadal variability originating from the North Atlantic and the Arctic significantly controls the climate of the Baltic Sea region by means of teleconnection patterns (Lehmann et al., 2022; Meier et al., 2022a; 2022b). For example, observations of precipitation and wind in the Baltic Sea region, total river discharge from the catchment, individual river flows, water temperature, sea level, MBIs and salinity in the Baltic Sea show a pronounced multidecadal variability with a quasi-periodicity of about 30 years (Meier et al., 2022b). It is assumed that the Atlantic Multidecadal Variability (AMV) and, as part of it, the variations of the North Atlantic overturning circulation are the source of these variations, although the exact mechanisms, cause and effect chains and feedback processes are still unknown. For

the development of climate prediction modeling systems, knowledge about the teleconnectivity of the Baltic Sea region with the North Atlantic and the Arctic is essential.

3) Development of integrated regional Earth system models accounting for multiple drivers of Earth system changes in the Baltic Sea region. The BEAR study by Reckermann et al. (2022) on human influences and their interactions in the Baltic Sea region is part of the relatively new Baltic Earth Grand Challenge 6 on the multiple drivers of Earth system change in the Baltic Sea region and represents an important step towards an integrated understanding of the Earth system, encompassing all traditionally considered climate compartments such as atmosphere, cryosphere, hydrosphere, lithosphere (including the pedosphere), biosphere (marine and terrestrial) and the anthroposphere. Such a holistic view is urgently needed, since in very many cases several causes are responsible for the observed changes in the Earth system and attributing them to only one factor, e.g. climate change, would be an inadmissible simplification. One example is the oxygen depletion and the large hypoxic area in the Baltic Sea caused by anthropogenic nutrient inputs from land and exacerbated by rising water temperatures (Kuliński et al., 2022). Of course, the factors discussed by Reckermann et al. (2022) cannot exhaustively take into account the whole Earth system and all interactions, and of course there is a bias towards marine-related parameters and activities. Moreover, the analysis is based on an extensive literature review by experts who reflect their subjective interpretations of the results. Nevertheless, this is the first time such an assessment has been conducted, which is a major step forward. To continue and deepen this research, the factors discussed by Reckermann et al. (2022) could be classified either by human activities (e.g. food production, energy production, transport, tourism, healthcare) or by environmental and climate state variables of the Earth system (e.g. hypoxia, acidification) (Table 3). Such a breakdown of parameters would allow the development of an integrated Earth system model that includes the anthroposphere at the regional scale.

The fact sheet on climate change in the Baltic Sea (CCFS, 2021) was positively received by various stakeholders and decision-makers. Although uncertainties regarding observed and projected future climate change and the other drivers remain high, our experience engaging with stakeholders confirms that scientific uncertainties are taken into account in different ways in management and decision-making. This is an important reason for investing in the above key issues and their potential to reduce uncertainties that currently hamper decision-making in the region.

## **5** Concluding remarks

We conclude that 1) the BEARs have been useful to identify research progress and knowledge gaps and to initiate new research foci as, for examples, suggested in the discussions; 2) regional assessments, such as the BEARs, complement the IPCC climate change assessments by adding a greater depth and scope of regional information about the specific situation of the Baltic Sea region; and 3) the BEARs provided useful information for the Expert Network on Climate Change, that produced the Baltic Earth – HELCOM climate change fact sheet for stakeholders.

#### Author contributions

H.E.M.M. wrote the first draft of the editorial. All co-authors, which acted as guest editors of the special issue in Earth System Dynamics, contributed substantially with important comments and editing of the manuscript, read and approved the submitted manuscript version.

## Acknowledgements

During 2019-2022, the Baltic Earth Assessment Reports were produced under the umbrella of the Baltic Earth programme (Earth System Science for the Baltic Sea region, see <a href="http://baltic.earth">http://baltic.earth</a>, last access: 2 February 2023). 109 co-authors from 14 countries contributed to 10 articles in the international scientific journal Earth System Dynamics and 2822 different references have been assessed. We thank the reviewers of all 10 articles of the special issue for their constructive comments that helped to improve the review articles. In particular, we thank Dr. Jouni Räisänen and Dr. Donald Boesch for their advice and many excellent comments on the individual articles and this overview article.

# Figures



**Figure 1:** The Baltic Sea and its catchment area with climatological mean sea surface salinity (in g kg<sup>-1</sup>) and river discharge (in mm year<sup>-1</sup>). (Source: Meier et al., 2014; their Fig. 1 distributed under the terms of the Creative Commons CC-BY 4.0 License, <u>http://creativecommons.org/licenses/by/4.0/</u>, last access: 4 February 2023)



**Figure 2:** Annual mean sea surface temperature anomalies to the reference period 2002-2018 from de-seasonalised measurements at the Arkona Deep monitoring station and the MARNET stations Darss Sill and Arkona Basin in the period 1980-2018. (Source: Meier et al., 2022b; their Fig. 20 distributed under the terms of the Creative Commons CC-BY 4.0 License, <u>http://creativecommons.org/licenses/by/4.0/</u>, last access: 4 February 2023)



Figure 3: Synthesis of knowledge on present and future climate change. Shown are anthropogenic climate changes in 33 Earth system variables (bubbles) of the atmosphere (yellow), land surface (brown), terrestrial biosphere (dark green), cryosphere (grey), ocean and sediments (blue) and marine biosphere (light green). The abbreviation NIS stands for non-indigenous species. The sign of a change (plus/minus) is shown together with the confidence level indicated by the number of signs, i.e. one to three signs correspond to a low, medium and high confidence level as a result of the literature assessment reflecting consensus and evidence according to IPCC definitions. The colours of the signs indicate the direction of past (black) and future (red) changes according to Meier et al. (2022b). Uncertain changes are not shown. The external anthropogenic drivers of the Earth system studied are shown as red squares, i.e. greenhouse gases, especially CO<sub>2</sub>, and aerosol emissions. The predominant climate change linkages with sufficiently high confidence are shown by arrows (black: thermal cycle, blue: hydrological cycle, orange: momentum cycle including sea level change, pink: carbon cycle). Projections of carbon sequestration of Arctic terrestrial ecosystems for the 21st century show first an increased uptake and later a carbon source, marked by "1. + 2. -". Future changes in mean sea level are dominated by the thermal expansion of the global ocean and the melting of ice sheets outside the Baltic Sea region. (Source: Meier et al., 2022b; their Fig. 35 distributed under the terms of the Creative Commons CC-BY 4.0 License, http://creativecommons.org/licenses/by/4.0/, last access: 4 February 2023)

## Tables

**Table 1:** The matrix of factors studied by Reckermann et al. (2022). + = evidence for a connection; - = no evidence for a connection; ? = no evidence, but connection plausible (according to the author's assessment). The table is read from left to right, i.e. if you go to the right in the first row "climate change", you see the factors on which climate change has an effect (or not), etc. (Source: Reckermann et al., 2022; their Table 2a distributed under the terms of the Creative Commons CC-BY 4.0 License, <u>http://creativecommons.org/licenses/by/4.0/</u>, last access: 4 February 2023)

impact by↓/on→	Climate change	Coastal processes	Hypoxia	Acidification	Subm. Groundw. Disch.	Marine ecosystems	Non-inig. species	Land cover and use	Agriculture Nutr. loads	Aquaculture	Fisheries	River regulations	Offshore wind farms	Shipping	Chem. Contamin.	Dumped milititary	Marine litter	Tourism	Coastal management
Climate change		+	+	+	?	+	?	+	+	+	+	+	+	+	+	?	?	+	+
Coastal processes	-		?	?	+	?	-	+	+	?	?	+	+	+	?	?	+	-	+
Нурохіа	-	-		+	-	+	÷.	-	+	?	+	-	-	-	+	+	-	-	-
Acidification	-	-	-			?	·			?	?	-		-	?		-	0.75	2.75
Subm. Groundw. Disch.			?	?		?	-	1.77	+				27	-	+	a		107.1	17.0
Marine ecosystems	-	-	+	+	1.00		+	1.00	-	-	+			-	-	-	-	+	-
Non-inigenous species		÷	1	8. <del>4</del> 0	1.75	+		2.40		100	+			+	+			( <b>7</b> )	?
Land cover and use	+		+	+	+	?	-		+	-	+	+	?	-	+		-	+	+
Agriculture/Nutrient loads	+	1.4	+	+	+	+	1.000	+		+	+	+	-	-	+	- <u>-</u>	-	243	?
Aquaculture	-	2	+	-	-	+	+	+	+		?	-	+	-	?		?	?	+
Fisheries			?			+	?	1.00	?	?			+	?	-		+	?	+
River regulations	-	+	?	+	?	+	-	?	?	?	+		-	-	?	-	?	-	+
Offshore wind farms	+	+	- 2	- 2	1.12	+	2	?	?	+	+	-		+	?	?	?	+	+
Shipping	+	+	-	+	-	+	+	-	+	?	+		?		+	-	+	+	+
Chemical contaminants			-	-	-	+			+	+	+		-	-		-	-	-	-
Dumped milititary material	-		5		-	?				-	+		+	-	+		-	1.72	?
Marine litter	-	-				?	-		-	?	+		1.0	-	?	-		+	?
Tourism	+	-		-	1 <del></del> 2	?	-	+	+		-	-	+	+	-	-	+		+
Coastal management	-	+-	-	-	?	?	?	?	?	?	+	?	+	+	-	+	?	+	

**Table 2:** Variables of the Meier et al. (2022b) assessment and further references to the BEARs (1: Lehmann et al., 2022; 2: Kuliński et al., 2022; 3: Rutgersson et al, 2022; 4: Weisse et al., 2021; 5: Reckermann et al, 2022; 6: Gröger et al, 2021; 7: Christensen et al, 2022; 8: Meier et al, 2022a; 9: Viitasalo and Bonsdorff, 2022). The third column lists the subsection in the study by Meier et al. (2022b) that contains further information. (Source: Meier et al., 2022b; their Table 2 distributed under the terms of the Creative Commons CC-BY 4.0 License, http://creativecommons.org/licenses/by/4.0/, last access: 4 February 2023 )

Number	Variable	Past and present c	limates	Future climate				
Atmosphere								
1	Large-scale	3.2.1.1	3	3.3.1.1	3, 7			
	atmospheric							
	circulation							
2	Air temperature	3.1.2, 3.1.3,						
		3.1.4		3.3.1.2	7			
	Warm spell	3.2.1.2	3		3			
	Cold spell		3		3			
3	Solar radiation and	3.2.1.3		3.3.1.3	7			
	cloudiness							
4	Precipitation	3.1.2, 3.1.3,						
		3.1.4		3.3.1.4	7			
	Heavy precipitation	3.2.1.4	3		3			
	Drought		3		3			
5	Wind	3.2.1.5		3.3.1.5	7			
	Storm		3		3			
6	Air pollution, air	3.2.1.6		3.3.1.6				
	quality and							
	atmospheric							
	deposition							
Land	Land							
7	River discharge	3.2.2.1		3.3.2.1	8			
	High flow		3		3			
8	Land nutrient inputs	3.2.2.2		3.3.2.2	8			
Terrestrial biosphe	Terrestrial biosphere							
9	Land cover (forest,	3.2.3	6	3.3.3				
	crops, grassland,							
	peatland, mires)							
10	Carbon			3.3.3				
	sequestration							
Cryosphere								
11	Snow	3.2.4.1		3.3.4.1	7			
	Sea-effect snowfall		3		3			

12	Glaciers	3.2.4.2		3.3.4.2	
13	Permafrost	3.2.4.3		3.3.4.3	
14	Sea ice	3.2.4.4		3.3.4.4	8
	Extreme mild winter		3		3
	Severe winter		3		3
	Ice ridging		3		3
15	Lake ice	3.2.4.5		3.3.4.5	
Ocean and marine	esediments				
16	Water temperature	3.2.5.1		3.3.5.1	8
	Marine heat wave		3		3
17	Salinity and	3.2.5.2	1	3.3.5.2	8
	saltwater inflows				
18	Stratification and	3.2.5.3	1	3.3.5.3	8
	overturning				
	circulation				
19	Sea level	3.2.5.4	4	3.3.5.4	8
	Sea level extreme		3		3
20	Waves	3.2.5.5	4	3.3.5.5	
	Extreme waves		3		3
21	Sedimentation and	3.2.5.6	4	3.3.5.6	
	coastal erosion				
22	Oxygen and	3.1.4			
	nutrients	3.2.5.7.1	2	3.3.5.7.1	8
23	Marine CO <sub>2</sub> system	3.2.5.7.2	2	3.3.5.7.2	
Marine biosphere					
24	Pelagic habitats:	3.2.6.1.1	2,9	3.3.6.1.1	9
	Microbial				
	communities				
25	Pelagic habitats:	3.2.6.1.2	2, 3, 9	3.3.6.1.2	3, 9
	Phytoplankton and				
	cyanobacteria				
26	Pelagic habitats:	3.2.6.1.3	9	3.3.6.1.3	9
	Zooplankton				
27	Benthic habitats:	3.2.6.2.1	9	3.3.6.2.1	9
	Macroalgae and				
	vascular plants				
28	Benthic habitats	3.2.6.2.2	9	3.3.6.2.2	9
	Zoobenthos	2.2.0.2.2	-	2.2.0.2.2	-
29	Non-indigenous	3263	9	3363	9
27	species	5.2.0.5		5.5.0.5	
	species				

30	Fish	3.2.6.4	9	3.3.6.4	9
31	Marine mammals	3.2.6.5	9	3.3.6.5	9
32	Waterbirds	3.2.6.6	9	3.3.6.6	9
33	Marine food web	3.2.6.7	9	3.3.6.7	9

Human activities		
Economic sectors	Factors	Comments
Primary (natural ) sector (e.g. food	Fisheries	
production)	Agriculture	
	Marine and coastal ecosystem	Factor belongs to several sectors
	services	
	Blue carbon storage capacity	Mitigation of greenhouse gases
Secondary (industrial) sector (e.g.	River regulation	
energy production)	Offshore wind farms	
	Greenhouse gas and aerosol	Emission are largest from
	emissions	industries
	Dumped warfare agents	Factor is an industrial product
Tertiary (service) sector (e.g.	Shipping	
transportation, tourism, healthcare)	Chemical contamination	Contamination is a result of several
		sectors
	Marine noise	Marine noise is a result of several
		sectors
	Marine litter and microplastics	Emission mainly by offshore
		platforms, shipping, lost
		containers, fisheries, aquaculture,
		agriculture, municipal waste and
		tourism
	Tourism	
	Coastal protection and	Also relevant for the other sectors
	management	
Quaternary (information) sector	-	
(e.g. information technology;		
media; research and development)		
Earth system	1	1
Environmental state variables	Coastal processes	
	Нурохіа	
	Submarine groundwater discharge	
	Marine ecosystems	
	Land use and land cover	
	Non-indigenious species	
	Indirect parameters such as carbon	
	and nutrient cycles, biota and	
	ecosystems	

**Table 3**: Factors discussed by Reckermann et al. (2022) sorted by related economic sectors or state variables of the Earth system.

Climate state variables	Climate change, acidification,	Superordinated concept (large-
	direct parameters of the climate	scale)
	system	
	Direct parameters of the climate	
	system	
	Acidification	

## References

Baltic Earth Science Plan, 2017. https://baltic.earth/grandchallenges, last access: 4 February 2023

CCFS, 2021. Climate Change in the Baltic Sea. 2021 Fact Sheet. Baltic Sea Environment Proceedings n°180. HELCOM/Baltic Earth, http://doi.io-warnemuende.de/10.12754/misc-2022-0001, 2021.

Christensen, O. B., Kjellström, E., Dieterich, C., Gröger, M., and Meier, H. E. M.: Atmospheric regional climate projections for the Baltic Sea Region until 2100, Earth Syst. Dynam., 13, 133–157, <u>https://doi.org/10.5194/esd-13-133-2022</u>, 2022.

Gröger, M., Dieterich, C., Haapala, J., Ho-Hagemann, H. T. M., Hagemann, S., Jakacki, J., May, W. Meier, H. E. M., Miller, P. A. Rutgersson, A. and Wu, L.: Coupled regional Earth system modeling in the Baltic Sea region, Earth Syst. Dynam., 12, 939–973, <u>https://doi.org/10.5194/esd-12-939-2021</u>, 2021.

Kuliński, K., Rehder, G., Asmala, E., Bartosova, A., Carstensen, J., Gustafsson, B., Hall, P. O. J., Humborg, C., Jilbert, T., Jürgens, K., Meier, H. E. M., Müller-Karulis, B., Naumann, M., Olesen, J. E., Savchuk, O., Schramm, A., Slomp, C. P., Sofiev, M., Sobek, A., Szymczycha, B., and Undeman, E.: Biogeochemical functioning of the Baltic Sea. Earth Syst. Dynam., 13, 633–685, <u>https://doi.org/10.5194/esd-13-633-2022</u>, 2022.

Lehmann, A., Myrberg, K., Post, P., Chubarenko, I., Dailidiene, I., Hinrichsen, H.-H., Hüssy, K., Liblik, T., Meier, H. E. M., Lips, U., and Bukanova, T.: Salinity dynamics of the Baltic Sea, Earth Syst. Dynam., 13, 373–392, https://doi.org/10.5194/esd-13-373-2022, 2022.

Meier, H.E.M., A. Rutgersson, and M. Reckermann: An Earth System Science Program for the Baltic Sea Region. Eos, 95, 109–110. https://doi.org/10.1002/2014EO130001, 2014.

Meier, H. E. M., Dieterich, C., Gröger, M., Dutheil, C., Börgel, F., Safonova, K., Christensen, O. B., and Kjellström, E.: Oceanographic regional climate projections for the Baltic Sea until 2100, Earth Syst. Dynam., 13, 159–199, <u>https://doi.org/10.5194/esd-13-159-2022</u>, 2022a.

Meier, H. E., Kniebusch, M., Dieterich, C., Gröger, M., Zorita, E., Elmgren, R., ... and Zhang, W.: Climate change in the Baltic Sea region: a summary. Earth Syst. Dynam., 13, 457-593, <u>https://doi.org/10.5194/esd-13-457-2022</u>, 2022b.

Ostsee Klimawandel Faktenblatt: Baltic Sea Climate Change Fact Sheet (in German). <u>http://doi.io-</u> warnemuende.de/10.12754/misc-2022-0003, 2022.

Reckermann, M., Omstedt, A., Soomere, T., Aigars, J., Akhtar, N., Bełdowski, J., Brauwer, C. P.-S. d., Cronin, T., Czub, M., Eero, M., Hyttiäinen, K., Jalkanen, J.-P., Kiessling, A., Kjellström, E., Larsén, X. G., McCrackin, M., Meier, H. E. M., Oberbeckmann, S., Parnell, K., Poska, A., Saarinen, J., Szymczycha, B., Undeman, E.,

Viitasalo, M., Wörman, A., and Zorita, E.: Human impacts and their interactions in the Baltic Sea region. Earth Syst. Dynam., 13, 1–80, <u>https://doi.org/10.5194/esd-13-1-2022</u>, 2022.

Rutgersson, A., Kjellström, E., Haapala, J., Stendel, M., Danilovich, I., Drews, M., Jylhä, K., Kujala, P., Larsén, X. G., Halsnæs, K., Lehtonen, I., Luomaranta, A., Nilsson, E., Olsson, T., Särkkä, J., Tuomi, L., and Wasmund, N.: Natural Hazards and Extreme Events in the Baltic Sea region, Earth Syst. Dynam., 13, 251–301, https://doi.org/10.5194/esd-13-251-2022, 2022.

The BACC Author Team, 2008. Assessment of Climate Change for the Baltic Sea Basin. Regional Climate Studies series. Heidelberg, Springer-Verlag Berlin. ISBN 978-3-540-72785-9. 473 pp. https://doi.org/10.1007/978-3-540-72786-6.

The BACC II Author Team, 2015. Second Assessment of Climate Change for the Baltic Sea Basin. Regional Climate Studies series. Cham, Springer Cham. ISBN 978-3-319-16005-4. 501 pp. https://doi.org/10.1007/978-3-319-16006-1.

Viitasalo, M., and Bonsdorff, B.: Global climate change and the Baltic Sea ecosystem: direct and indirect effects on species, communities and ecosystem functioning. Earth Syst. Dynam., 13, 711–747, https://doi.org/10.5194/esd-13-711-2022, 2022.

Weisse, R., Dailidiene, I., Hünicke, B., Kahma, K., Madsen, K., Omstedt, A., Parnell, K., Schöne, T., Soomere, T., Zhang, W., and Zorita, E.: Sea Level Dynamics and Coastal Erosion in the Baltic Sea Region. Earth Syst. Dynam., 12, 871–898, <u>https://doi.org/10.5194/esd-12-871-2021</u>, 2021.