

1 **The Baltic Earth Assessment Reports**

2 H. E. Markus Meier<sup>1</sup>, Marcus Reckermann<sup>2</sup>, Joakim Langner<sup>3</sup>, Ben Smith<sup>4,5</sup> and Ira  
3 Didenkulova<sup>6</sup>

4 <sup>1</sup>Department of Physical Oceanography and Instrumentation, Leibniz Institute for Baltic Sea Research  
5 Warnemünde, Rostock, Germany

6 <sup>2</sup>International Baltic Earth Secretariat, Helmholtz-Zentrum Hereon, Geesthacht, Germany

7 <sup>3</sup>Swedish Meteorological and Hydrological Institute, Norrköping, Sweden

8 <sup>4</sup>Hawkesbury Institute for the Environment, Western Sydney University, Australia

9 <sup>5</sup>Department of Physical Geography and Ecosystem Science, Lund University, Sweden

10 <sup>6</sup>Department of Mathematics, University of Oslo, Norway

11

12 *Correspondence to:* H. E. Markus Meier ([markus.meier@io-warnemuende.de](mailto:markus.meier@io-warnemuende.de))

13

14 **Abstract.** Baltic Earth is an independent research network of scientists from all Baltic Sea countries that promotes  
15 regional Earth system research. Within the framework of this network, the Baltic Earth Assessment Reports  
16 (BEARs) were produced in the period 2019-2022. These are a collection of 10 review articles summarising current  
17 knowledge on the environmental and climatic state of the Earth system in the Baltic Sea region and its changes in  
18 the past (palaeoclimate), present (historical period with instrumental observations) and prospective future (until  
19 2100) caused by natural variability, climate change and other human activities. The division of topics  
20 ~~amongbetween~~ articles follows the grand challenges and selected themes of the Baltic Earth Science Plan, such as  
21 the regional water, biogeochemical and carbon cycles, extremes and natural hazards, sea level dynamics and  
22 coastal erosion, marine ecosystems, coupled Earth system models, scenario simulations for the regional  
23 atmosphere and the Baltic Sea, and climate change and impacts of human use. Each review article contains an  
24 introduction, the current state of knowledge, knowledge gaps, conclusions and key ~~messagesstatements~~, based on  
25 which recommendations ~~are made~~ for future research ~~are made~~. ~~In parallel~~Based on the BEARs, Baltic Earth's  
26 ~~ongoing-has published~~outreach work has led to the publication of an information leaflet on climate change in the  
27 Baltic Sea ~~as part of its outreach work~~, which has been published in two languages so far, and ~~organisedthe~~  
28 ~~organisation-of stakeholder~~ conferences and workshops ~~for stakeholders, in collaboration with the Baltic Marine~~  
29 ~~Environment Protection Commission (HELCOM)~~.  
30

## 31 1 Introduction

### 32 1.1 BALTEX/Baltic Earth history

33 Baltic Earth<sup>1</sup> is an international research network dealing with Earth system sciences ~~inof~~ the Baltic Sea region  
34 (Fig. 1) (<https://baltic.earth>, last access: 4 February 2023). The catchment area of the Baltic Sea is about four times  
35 larger than the Baltic Sea surface and is part of mainly the countries Belarus, Denmark, Estonia, Finland, Latvia,  
36 Lithuania, Poland, Russia, and Sweden (Fig. 1). Baltic Earth ~~It~~ is politically independent and focusses on research  
37 on the water and energy cycles, climate variability and climate change, water management and extreme events,  
38 and ~~associatedrelated~~ impacts on marine and terrestrial biogeochemical cycles. Research on ~~h~~Human impact on  
39 the Earth system in more general terms, i.e. the anthroposphere, ~~defined as the part of the environment created or~~  
40 ~~modified by humans for use by human activities~~, was ~~also included in~~added to the 2017 Baltic Earth Science Plan  
41 (2017)<sup>2</sup> (<https://baltic.earth/grandchallenges>, last access: 4 February 2023).  
42

43 Baltic Earth is the successor of the Baltic Sea Experiment (BALTEX) programme, which was founded in 1993 as  
44 a GEWEX continental-scale experiment (Global Energy and Water Exchanges, a core project of the World Climate  
45 Research Programme) (Reckermann et al., 2011). ~~In~~During the first phase (Phase I) (1993–2002), BALTEX was  
46 primarily devoted to hydrological, meteorological and oceanographic processes in the Baltic Sea drainage basin

---

<sup>1</sup> <https://baltic.earth>, last access: 4 February 2023

<sup>2</sup> <https://baltic.earth/grandchallenges>, last access: 4 February 2023

47 ~~and thus, hence~~ focussed on physical aspects of the Earth system. ~~In~~During the second phase (~~Phase II~~: 2003–  
48 2012), the programme was expanded to ~~include~~~~encompass~~ regional climate research, ~~carbon and~~ biogeochemical  
49 cycles ~~including carbon~~, engagement with stakeholders and decision–makers via assessment reports, as well as  
50 communication and education, i.e. ~~the organisation of~~~~organizing~~ summer and winter schools and international  
51 master courses.

52  
53 In 2013, Baltic Earth was launched with a ~~new~~~~novel~~ science plan ~~to strengthen~~~~reinforcing~~ efforts to address Grand  
54 Challenges on (1) salinity dynamics in the Baltic Sea, (2) ~~land-sea~~ biogeochemical linkages ~~between land and sea~~,  
55 (3) natural hazards and extreme events, (4) sea level and coastal dynamics, (5) regional variability ~~in~~~~of~~ water and  
56 energy exchanges, and (6) multiple drivers of regional Earth system changes (Meier et al., 2014). Working groups  
57 ~~were initiated~~ on coupled Earth system models, the Baltic Sea Model Intercomparison Project (BMIP), ~~uncertainty~~  
58 ~~of scenario simulations for the Baltic Sea, and~~ education, outreach and communication ~~have been established, and~~  
59 ~~scenario simulations for the Baltic Sea (today successfully completed)~~.

60  
61 Baltic Earth and its predecessor BALTEX have produced three ~~comprehensive~~~~extensive~~ regional assessment  
62 reports since 2008. The first two (The BACC Author Team, 2008, and The BACC II Author Team, 2015) ~~focused~~  
63 ~~on~~~~emphasised~~ climate change and its impacts in the Baltic Sea region and were published as text books, while the  
64 third, the Baltic Earth Assessment Reports (BEARs), was published in the format of a special issue ~~of the journal~~~~in~~  
65 ~~Earth System Dynamics~~, in 2022. The ~~Assessment of Climate Change in the Baltic Sea Basin (BACC) reports~~<sup>3</sup>  
66 ~~and BEARs and BACC assessment reports~~ fill a gap ~~compared~~~~relative~~ to the assessment reports of the  
67 Intergovernmental Panel on Climate Change (IPCC), ~~as~~~~since~~ the latter focus on global scales, and do not provide  
68 detailed local to regional information ~~on~~~~about~~ the current state of knowledge on climate change and its impacts in  
69 the Baltic Sea region. The BEARs provide a comprehensive and up-to-date overview of the state-of-the-art  
70 research on the compartments of the Earth system in the Baltic Sea region encompassing processes in the  
71 atmosphere, on land and in the sea, including the marine and terrestrial ecosystems as well as processes and impacts  
72 related to the anthroposphere.

73  
74 The BEARs ~~summarise the published scientific knowledge currently available and update the second BACC report~~  
75 ~~wrap together the currently available published scientific knowledge, updating the second assessment report of~~  
76 ~~climate change in the Baltic Sea basin~~ (The BACC II Author Team, 2015) based on the ~~latest~~ scientific literature.  
77 ~~This present BEAR s~~Special iIssue ~~includes~~~~comprises~~ 10 articles on the Baltic Earth Grand Challenges and ~~Baltic~~  
78 ~~Earth Special Topics (Baltic Earth Science Plan, 2017), working group topics~~ including a summary of ~~the~~ current  
79 knowledge ~~on~~~~about~~ past, present, and future climate changes ~~in~~~~for~~ the Baltic Sea region. The articles encompass  
80 contributions ~~from~~~~by~~ 109 authors from 14 countries and reference 2822 scientific articles and institutional reports.  
81 ~~in their synthesis effort.~~

---

<sup>3</sup> ~~Assessment of Climate Change in the Baltic Sea Basin (BACC); <https://baltic.earth/bacc>. last access:~~  
~~4 February 2023~~

## 82 1.2 Baltic Sea region characteristics

83 The Baltic Sea is a semi-enclosed, shallow sea with limited water exchange with the ~~world ocean~~World Ocean and  
84 small tidal amplitudes. ~~Located~~Situated in Northern Europe, the climate of the region is highly variable ~~as~~because  
85 it is ~~located~~in the transition zone between maritime and continental climates ~~and is~~ influenced by the North  
86 Atlantic and Arctic ~~regions~~. ~~The R~~iver discharges from the large catchment area causes a pronounced gradient in  
87 ~~sea surface~~ salinity from about 20 g kg<sup>-1</sup> in the Danish ~~s~~Straits' region to about 2 g kg<sup>-1</sup> or even less in the northern  
88 and eastern reaches of the Baltic Sea. Hence, the Baltic Sea is brackish, with habitats of ~~marinemaritime~~ species  
89 in the south-west and freshwater species in the north-east. The Baltic Sea catchment area ~~is about four times the~~  
90 ~~surface area of the Baltic Sea and~~ covers an area of almost 20% of the European continent (Fig. 1). ~~85 million~~  
91 ~~people in 14 countries live in the catchment area, which stretches from the temperate, densely populated south to~~  
92 ~~the subarctic rural north.~~It stretches from the temperate, densely populated south to the subarctic wilderness in the  
93 north and is home to approximately 85 million people in 14 countries, namely Belarus, the Czech Republic,  
94 Denmark, Estonia, Finland, Germany, Latvia, Lithuania, Norway, Poland, Russia, Slovakia, Sweden and Ukraine.

95  
96 Episodically, large amounts of saline ~~t~~water flow from the North Sea ~~overeross~~ the sills, ~~located~~ in the Danish  
97 ~~s~~Straits, into the Baltic Sea and ventilate the ~~deep waters of the~~ Baltic Sea ~~deepwater~~. These events require a period  
98 of about 20 days ~~with~~of easterly winds that lower the ~~Baltic~~ sea level ~~in the Baltic Sea~~, followed by a period of  
99 about ~~the same~~equal ~~length~~duration ~~with~~of strong westerly winds that push saline ~~t~~water into the Baltic Sea. These  
100 events are called Major Baltic Inflows (MBIs) and are important for the water exchange between the North Sea  
101 and the Baltic Sea. Mixing is low compared to other seas, with an origin at the lateral boundaries, because tidal  
102 amplitudes are very small and energetically ~~insignificant~~unimportant.

103  
104 In recent decades, environmental conditions in the Baltic Sea have changed considerably. For instance, ~~since 1980~~  
105 the Baltic Sea ~~has been~~is warming more than any other coastal sea ~~since 1980~~ (Fig. 2), ~~which has led to a reduction~~  
106 ~~in sea ice and snow cover over the land in winter~~causing shorter sea ice and snow covers over land during winter.  
107 Furthermore, ~~increasing~~rising nutrient inputs from ~~the~~ land in the 1950s/60s, caused by population growth ~~and the~~  
108 ~~discharge of~~accompanied by sewage ~~water release~~ into the Baltic Sea, ~~as well as the increased~~and intensified  
109 ~~usage~~ of fertilisers ~~in~~from agriculture, led to eutrophication and ~~the spreading~~ of hypoxic and anoxic areas. Since  
110 the 1980s, nutrient inputs into the Baltic Sea ~~has~~yes been steadily decreasing, but this has not yet led to a significant  
111 improvement in oxygen conditions. Recent trends in acidification are ~~lower~~smaller than in the ~~W~~world ~~O~~ocean,  
112 ~~especially~~in particular in the northern Baltic Sea, ~~as~~because positive trends in alkalinity ~~input~~supply counteract the  
113 acidification.

## 114 2 Methods

115 ~~Succeeding~~Following ~~T~~the BACC<sup>4</sup> Author Team (2008) and ~~T~~the BACC II Author Team (2015) ~~assessments~~, the  
116 BEAR project is an ~~attempt~~~~effort~~ to summarise ~~the~~ scientific knowledge ~~on~~~~about~~ climate change and other drivers  
117 of Earth system changes and their impacts on the Baltic Sea region. ~~The two BACC books have a format inspired~~  
118 ~~by the IPCC assessment reports. This special issue in *Earth System Dynamics* is the third assessment. It has a new~~  
119 ~~format of BEARs, encompassing 10 peer-reviewed scientific journal articles.~~~~The BACC books adopted a format~~  
120 ~~inspired by the Intergovernmental Panel on Climate Change (IPCC) assessment reports. The present special issue~~  
121 ~~is the third assessment of a new format of Baltic Earth Assessment Reports (BEARs), encompassing 10 peer-~~  
122 ~~reviewed scientific articles in the scientific journal *Earth System Dynamics*.~~ The ~~assessed~~ knowledge ~~assessed~~ was  
123 extracted from the scientific literature such as peer-reviewed articles, reports from research institutions, and  
124 published datasets. Importantly, literature from non-governmental organisations with ~~a~~-political or economic  
125 interests, political parties and other stakeholder organisations was excluded from the assessment, ~~to ensure~~~~ing~~ that  
126 only scientific knowledge ~~was included in~~~~informed~~ the assessment. The BEARs focus on publications after  
127 2012/2013, the year of the editorial deadline of the second assessment report. Whenever possible, ~~the~~ uncertainty  
128 levels of the BEAR results are ~~ranked~~~~classified~~ based on a matrix of consensus within the scientific literature and  
129 ~~the~~ documented evidence of detected changes and their attributed drivers such as climate change and human use.  
130 ~~For a high confidence of a certain statement, A~~ high levels of ~~both~~ scientific consensus and evidence ~~is~~~~ases~~ are  
131 required ~~for high confidence in a particular statement. Instances of disagreement and knowledge gaps are~~  
132 ~~documented and discussed, informing priorities for future research.~~~~Disagreements and gaps in knowledge are~~  
133 ~~documented and discussed to prioritise future research.~~

134  
135 Together with the intergovernmental Baltic Marine Environment Protection Commission (HELCOM), Baltic  
136 Earth ~~has established~~~~formed~~ an Expert Network on Climate Change (EN CLIME). The aim of the expert network  
137 is to regularly produce a climate change fact sheet (CCFS, 2021<sup>5</sup>) ~~based on~~~~from~~ the BEAR ~~and BACC~~ material.  
138 In 2021, it was published for the first time<sup>6</sup> (<http://helcom.fi/ccfs>, last access: 4 February 2023). The CCFS contains  
139 some background information, a map showing regional future climate changes for selected parameters under the  
140 greenhouse gas concentration scenario RCP4.5 and information ~~on~~~~about~~ 34 variables, directly and indirectly  
141 affected by climate change. For each parameter, a general description, past and prospective future changes, other  
142 drivers than climate change (only for the indirect parameters), knowledge gaps, policy relevance and references  
143 are presented. More than 100 scientists contributed to the compilation of the ~~first~~ fact sheet, which was coordinated  
144 by the HELCOM secretariat. Updated versions are planned ~~at seven-year intervals~~~~for intervals of seven years~~. Like  
145 the BEARs, the fact sheet was peer-reviewed and quality assured. It has so far ~~only~~ been translated to German

Formatiert: Schriftart: Kursiv

<sup>4</sup> Assessment of Climate Change in the Baltic Sea Basin (BACC); <https://baltic.earth/bacc>, last access: 4 February 2023

<sup>5</sup> <https://helcom.fi/wp-content/uploads/2021/09/Baltic-Sea-Climate-Change-Fact-Sheet-2021.pdf>, last access: 4 February 2023

<sup>6</sup> <http://helcom.fi/ccfs>, last access: 4 February 2023

146 (*Ostsee-Klimawandel in der Ostsee, 2021* Faktenblatt, 2022<sup>7</sup>), ~~but~~ translations ~~in~~ into other languages are planned  
147 ~~to improve, enhancing~~ accessibility to stakeholders.

150 In this editorial, we highlight the ~~key~~main findings and knowledge gaps as ~~described~~detailed by the BEARs and  
151 ~~propose~~ future work ~~is proposed~~.

### 152 3 Results

153 ~~Some~~A number of the ~~key~~main findings of the 10 BEARs are selected and highlighted below. ~~The BEARs are~~  
154 ~~either based on the Baltic Earth Grand Challenges or Baltic Earth Special Topics (Baltic Earth Science Plan, 2017).~~

155  
156 1. **Salinity dynamics of the Baltic Sea**, Grand Challenge 1 (Lehmann et al., 2022): Salinity is an important  
157 parameter for the circulation and the marine ecosystem in the Baltic Sea. Any changes in salinity are  
158 caused by changes either in the freshwater inflow from rivers and net precipitation over sea or in the water  
159 exchange between the Baltic Sea and the adjacent North Sea. Although long-term records of salinity and  
160 its drivers suffer from data gaps, these records starting in the 19<sup>th</sup> century are globally unique. Major  
161 research efforts focused on the MBI event in 2014 and its consequences for water masses, oxygen  
162 concentration and biogeochemical cycling. During the event, an unexpectedly large contribution of oxic  
163 intrusions at intermediate depth and essentially nonturbulent conditions in the deep interior were found,  
164 ~~emphasizing~~ the importance of boundary mixing. A revised reconstruction of the long-term record of  
165 MBIs showed no trend but a pronounced multi-decadal variability with a period of about ~~25~~-30 years.  
166 Despite intense research activities, observed variations in the intensity and frequency of MBIs and related  
167 Large Volume Changes (LVCs) could not be attributed to atmospheric circulation variability. Hence, on  
168 time scales larger than the synoptical time scale, MBIs are not predictable. ~~As an advance over~~ ~~In contrast~~  
169 ~~to~~ the previous assessments, salinity dynamics of the various sub-basins and lagoons mainly based on  
170 observations have been discussed, documenting large regional differences.

171  
172 2. **Biogeochemical functioning of the Baltic Sea**, Grand Challenge 2 (Kuliński et al., 2022): The review  
173 addresses the following topics: (1) terrestrial biogeochemical processes and nutrient inputs to the Baltic  
174 Sea, (2) the transformation of C, N and P in the coastal zone, (3) the production and remineralisation of  
175 organic matter, (4) oxygen availability, (5) the burial and turnover of C, N and P in sediments, (6) the  
176 Baltic Sea CO<sub>2</sub> system and seawater acidification, (7) the role of certain microorganisms in the  
177 biogeochemistry of the Baltic Sea, and (8) the interactions between biogeochemical processes and  
178 chemical pollutants. It was found that oxygen depletion and the ~~are~~increase of anoxic ~~sea~~-bottoms have  
179 still increased despite the reductions in nutrient inputs from land since the 1980s. Hence, the nitrogen  
180 pool has declined due to denitrification whereas the phosphorus inventory has increased. Estimates

---

<sup>7</sup> <https://baltic.earth/ccfs>, last access: 4 February 2023

181 suggest that about 1% and 4% of the annual nitrogen and phosphorus loads, respectively, have  
182 accumulated in the Baltic Sea, while the remainder are either exported to the North Sea or lost via  
183 biogeochemical processes such as denitrification and burial. Furthermore, it was discovered that in the  
184 central and northern sub-basins the uptake of C, N and P during primary production does not correspond  
185 to the Redfield ratio, which strongly affects the relationship between primary production, export of  
186 organic matter and oxygen demand of the deep sea. While it is clear that the Baltic Sea is a CO<sub>2</sub> sink in  
187 summer and a CO<sub>2</sub> source in winter, the annual net balance remains unknown. The past increase in total  
188 alkalinity of unknown origin has entirely mitigated ocean acidification in the northern Baltic Sea and  
189 significantly reduced it in the central Baltic Sea. In the future, a doubling of atmospheric pCO<sub>2</sub> ~~would~~  
190 still result in lower pH in the entire Baltic Sea, even if alkalinity should further increase.

191  
192 3. **Natural hazards and extreme events in the Baltic Sea region**, Grand Challenge 3 (~~Anna~~ Rutgersson  
193 et al., 2022): Existing knowledge is summarised about extreme events in the Baltic Sea region with a  
194 focus on the past 200 years with instrumental data as well as on future projections. Considered events are  
195 wind-storms, extreme waves, high and low sea levels, hot and cold spells in the atmosphere, marine heat  
196 waves, droughts, sea-effect snowfall, sea-ice ridging, extremely mild and extremely severe sea ice  
197 winters, heavy precipitation events, river floods, and extreme phytoplankton blooms. Furthermore, the  
198 knowledge about implications of these extreme events for society such as forest fires, coastal flooding,  
199 offshore infrastructure and shipping was assessed. With respect to the impacts of climate change,  
200 terrestrial and marine heat waves, extremely mild sea ice winters, heavy precipitation and high-flow  
201 events are expected to increase, while cold spells, severe sea ice winters and sea-ice ridging are expected  
202 to decrease due to the increase in mean atmospheric temperature. Changes in relative sea level extremes  
203 will depend on the competing impacts of competition between the rising global mean sea level, the  
204 gravitational effect of the melting of the Greenland and Antarctic ice sheets, changes in wind fields,  
205 and the regionally differing Glacial Isostatic Adjustment (GIA) resulting in land uplift or subsidence.  
206 Furthermore, projections suggest an increase of droughts in the southern and central parts of the Baltic  
207 Sea region mainly in summer. Significant future changes in wind-storms, extreme waves and sea level  
208 extremes relative to the mean sea level have not been found, suggesting that these changes will likely be  
209 small compared with natural variability probably because available climate model ensembles are too  
210 small.

211  
212 4. **Sea level dynamics and coastal erosion in the Baltic Sea region**, Grand Challenge 4 (Weisse et al.,  
213 2021): In this study, the current knowledge about the diverse processes affecting mean and extreme sea  
214 level changes, coastal erosion and sedimentation with impact on coastline changes and coastal  
215 management is assessed. Such processes are GIA, contributions from global sea level changes, wind-  
216 storms, wind-waves, seiches or meteotsunamis. During 1886-2020, the mean absolute sea level in the  
217 Baltic Sea corrected for GIA increased by about 25 cm or ~2 mm year<sup>-1</sup> on average. Land uplift in the  
218 north is still faster than the absolute sea level rise while in the south the opposite is true with potential  
219 impacts on changes in coastal erosion and inundation. The current acceleration of sea level rise is small

220 and could only be determined by spatially averaging observations at different tide gauge locations. Future  
221 sea level rise in the Baltic Sea is expected to further accelerate, probably somewhat less than the global  
222 mean sea level rise ~~due to the gravitational contributions from the melting of the Antarctic ice sheet~~. The  
223 Baltic sea level is substantially more sensitive to melting from the Antarctic than from the Greenland ice  
224 sheet. Concerning sediment transports, the dominance of mobile sediments makes the southern and  
225 eastern Baltic Sea coasts susceptible to wind-wave induced transports, in particular during storms. Due  
226 to the global sea level rise, future sediment transports can be expected to increase in these coastal areas,  
227 with a large spatial variability depending on the angles of incidence of incoming wind-waves.

228  
229 5. **Human impacts and their interactions in the Baltic Sea region**, Grand Challenge 6 (Reckermann et  
230 al., 2022): An inventory and discussion of the various man-made factors and processes affecting the  
231 environment of the Baltic Sea region and their interrelationships are presented. In total, ~~more than~~ 19  
232 factors are addressed (Table 1). Some of the factors are natural and are only modified by human activities  
233 (e.g. climate change, coastal processes, hypoxia, acidification, submarine groundwater discharges, marine  
234 ecosystems, non-indigenous species, land use and land cover), others are entirely man-made (e.g.  
235 agriculture, aquaculture, fisheries, river regulation, offshore wind farms, shipping, chemical  
236 contamination, dumped ammunition, marine litter and microplastics, tourism and coastal management).  
237 All factors are interconnected to varying degrees. The knowledge of these linkages was assessed and  
238 analysed in depth. The main finding was that climate change has an overarching, integrating effect on all  
239 other factors and can be interpreted as a background effect that affects the other factors differently. After  
240 climate change, shipping and land use/agriculture ~~are the factors affecting most other factors~~ have the  
241 strongest impacts, while fisheries, marine ecosystems and agriculture in turn are the most affected. The  
242 results of the assessment depend on the region and may be different for other coastal seas and their  
243 catchments in the world, where different human activities prevail.

244  
245 6. **Global climate change and the Baltic Sea ecosystem: direct and indirect effects on species,**  
246 **communities and ecosystem functioning**, Baltic Earth Special Topic (Viitasalo and Bonsdorff, 2022):  
247 Climate change has multiple impacts on species, communities and ecosystem functioning in the Baltic  
248 Sea through changes in physical and biogeochemical parameters such as temperature, salinity, oxygen,  
249 pH and nutrient levels. The associated secondary effects on species interactions, trophic dynamics and  
250 ecosystem function are also likely to be important. Climate change (warming, recent brightening,  
251 decrease in sea ice) has led to shifts in the seasonality of primary production, with a prolonged growing  
252 season of phytoplankton, an earlier onset of the spring bloom and a delayed autumn bloom. However, the  
253 development of cyanobacteria varies from species to species, and a clear causal relationship between  
254 temperature or salinity and the abundance of cyanobacteria has not been demonstrated. An increase in  
255 water temperature and river input of dissolved organic matter (DOM) could reduce primary production  
256 while favouring bacterial growth. If nutrient reduction ~~continues~~ proceeds, the improvement in oxygen  
257 conditions could initially increase zoobenthos biomass, but the subsequent decrease in sedimenting  
258 organic matter would likely disrupt the pelagic-benthic coupling and result in lower zoobenthos biomass.



259 Sprat and some coastal fish species could be favoured by a rise in temperature. Regime shifts and  
260 cascading effects have already been observed in both pelagic and benthic systems as a result of climate  
261 change.

262

263 7. **Coupled regional Earth system modeling in the Baltic Sea region**, Baltic Earth Special Topic, with  
264 relevance to Baltic Earth Grand Challenge 5 (Gröger et al., 2021): Recent progress ~~in~~ the development  
265 of coupled climate models for the Baltic Sea region is assessed. Feedback mechanisms are important to  
266 simulate the response of the Earth system to external forcing such as greenhouse gas and aerosol  
267 emissions. In this review article, the ~~couplings~~ between (1) atmosphere, sea ice and ocean, (2) atmosphere  
268 and land surface including dynamic vegetation, (3) atmosphere, ocean and waves and (4) atmosphere and  
269 hydrological components to close the water cycle ~~are~~ discussed. Adding surface waves to a-coupled  
270 atmosphere-ocean system ~~models~~ is becoming more important with increasing resolution, in particular  
271 when detailed information is required, for instance, for offshore wind energy applications in the coastal  
272 zone. Furthermore, the wave information is essential for the calculation of ocean mixing and  
273 resuspension. While long-term climate simulations using coupled atmosphere, sea ice and ocean models  
274 or coupled atmosphere and dynamic vegetation models have successfully been performed and their added  
275 value demonstrated, the impact of aerosols on the climate ~~of the Baltic Sea region~~ ~~has not been~~ ~~is not~~  
276 considered. Coupling hydrology models to close the hydrological cycle is also still problematic, as the  
277 precipitation accuracy provided by the atmospheric models is, in most cases, insufficient to realistically  
278 simulate river discharge into the Baltic Sea without bias adjustments.

279

280 8. **Atmospheric regional climate projections for the Baltic Sea region until 2100**, Baltic Earth Special  
281 Topic (Christensen et al., 2022): Current climate projections based on regional climate atmosphere-only  
282 models of the EURO-CORDEX project with a horizontal resolution of 12.5 km under the scenarios  
283 RCP2.6, 4.5 and 8.5 are presented. As the number of simulations (124) is relatively large compared to  
284 previous assessments, the uncertainties can be better estimated than before. These projections indicate  
285 strong warming, especially in the north in winter, where warming approaches twice the average global  
286 warming. Precipitation is projected to increase throughout the Baltic Sea region, except in the southern  
287 half in summer, where the results are inconclusive. Extreme precipitation, here the 10-year return value,  
288 is projected to increase systematically throughout the study area, especially in summer. The large  
289 ensemble of simulations does not indicate a significant change in wind speed. Surface solar radiation is  
290 projected to remain unchanged in summer, but to decrease slightly in winter, due to increased cloud cover  
291 and possibly less snow in the future. ~~These results from regional climate models contrast with the~~  
292 ~~projections of many global climate models, which show an increasing trend in solar radiation, and~~  
293 ~~illustrate how important more precisely resolved spatial features such as topography and coastlines are~~  
294 ~~for climate and weather.~~ Snow cover is projected to decrease dramatically, especially in the south of the  
295 Baltic Sea catchment. The comparison between the uncoupled model simulations of the EURO-CORDEX  
296 project and a small ensemble of scenario simulations performed with a coupled atmosphere-sea-ice-ocean  
297 model driven by a subset of global climate models indicates stronger warming in the coupled model

298 during winter, mainly in areas that are seasonally affected by sea ice today. In summer, the coupled model  
299 shows weaker warming compared to the uncoupled models.  
300

301 9. **Oceanographic regional climate projections for the Baltic Sea until 2100**, Baltic Earth Special Topic  
302 (Meier et al., 2022<sup>ab</sup>): New projections of the future Baltic Sea climate with a coupled physical-  
303 biogeochemical ocean model were compared with previous projections. The differences are mainly due  
304 to different scenario assumptions and model setups. For example, the impact of future global sea level  
305 rise on salinity was previously neglected, but taken into account in the latest projections. Although the  
306 number of projections for the Baltic Sea is still small compared to regional atmospheric projections such  
307 as the EURO-CORDEX model ensemble, a relatively large ensemble of 48 scenario simulations allowed  
308 the assessment of uncertainties related to greenhouse gas emissions, global climate model differences,  
309 global sea level rise, nutrient inputs and natural variability. In the future climate, higher water  
310 temperatures, a shallower mixed layer with a sharper thermocline in summer, lower sea ice cover and  
311 stronger mixing in the northern Baltic Sea in winter compared to the current climate could be expected.  
312 The assessment of marine heat wave changes is new. Both the frequency and duration of marine heat  
313 waves are projected to increase significantly, especially in the coastal zone of the southern Baltic Sea.  
314 Due to uncertainties in the projections regarding regional winds, precipitation and global sea level rise,  
315 no robust and statistically significant changes in salinity could be identified. The impacts of a changing  
316 climate on the biogeochemical cycle are projected to be significant, but still less than the plausible  
317 changes in nutrient inputs. Implementation of the proposed Baltic Sea Action Plan, a basin-wide nutrient  
318 input reduction plan, would lead to a significant improvement in the ecological status of the Baltic Sea,  
319 including a reduction in the size of the hypoxic area also in a future climate.  
320

321 10. **Climate change in the Baltic Sea region: A summary**, Baltic Earth Special Topic (Meier et al., 2022<sup>ba</sup>):  
322 In this comprehensive study, the recent knowledge on past (paleo-), present (historical) and projected  
323 future (< 2100) climate change in the Baltic Sea region, based upon all BEARs and >800 scientific  
324 articles, is summarised. It focuses on the atmosphere, the land surface, the cryosphere, the ocean and its  
325 sediments, and the terrestrial and marine biospheres. 33 parameters characterising the state of these  
326 components of the Earth system were analysed (Fig. 3, Table 2). The anthroposphere is not part of this  
327 assessment by Meier et al. (2022b) but instead is discussed in detail by Reckermann et al. (2022). The  
328 main findings concerning changes of the 33 selected state parameters attributed to climate change are  
329 summarised in Figure 3. The prevailing causal relationships of climate change with sufficiently high  
330 confidence suggest a clear impact of global greenhouse gas emissions on regional heat cycles including  
331 all parameters of the cryosphere. However, changes caused by global warming of the water, momentum  
332 and carbon cycles are less clear because of either the large natural variability at regional scales or the  
333 impact of other drivers than global warming. For further details, the reader is referred to Meier et al.  
334 (2022b). Overall, it was concluded that the results from the previous BACC assessments mainly are still  
335 valid. However, new long-term, homogenous observational records, such as those for Scandinavian  
336 glacier inventories, sea-level-driven saltwater inflows (MBIs), or phytoplankton species distributions, and

337 new scenario simulations with improved models, such as those for glaciers, lake ice, or marine food webs,  
338 have become available, resulting in a revised understanding of observed changes. Compared to previous  
339 assessments, observed changes in air temperature, sea ice, snow cover, and sea level were shown to have  
340 accelerated. However, natural variability is large, challenging our ability to detect observed and projected  
341 changes in climate of the Baltic Sea region. As the ensembles of scenario simulations both for the  
342 atmosphere and the ocean became larger, uncertainties can now be better estimated, although coordinated  
343 scenario simulations for the Baltic Sea based on ensembles of different regional ocean models are still  
344 missing. Furthermore, with the help of coupled models, feedbacks between several components of the  
345 Earth system have been studied, and multiple driver studies were performed, e.g., projections of the  
346 marine food web that include fisheries, eutrophication and climate change. Intensive research on the land–  
347 sea interface, focusing on the coastal filter, has been performed, and nutrient retention in the coastal zone  
348 was estimated for the first time. However, a model for the entire Baltic Sea coastal zone is still missing,  
349 and the effect of climate change on the coastal filter capacity is still unknown. More research on changing  
350 extremes was performed, acknowledging that the impact of changing extremes may be more important  
351 than that of changing means (see also Rutgersson et al., 2022). However, many observational records are  
352 either too short or too heterogeneous (~~data gaps~~) for statistical studies of extremes due to data gaps.

#### 353 4 Discussion

354 ~~The identification of knowledge gaps in Earth system science of the Baltic Sea region, that should be further~~  
355 ~~addressed in future research, was one of the main aims of the BEAR project. One of the main objectives of the~~  
356 ~~BEAR project was to identify knowledge gaps in the Earth system science of the Baltic Sea region so that these~~  
357 ~~can be further addressed in future research.~~ For specific knowledge gaps that have been identified during the  
358 project, the reader is referred to the individual assessment reports. However, as an overarching result, three new  
359 research topics are identified:

360

- 361 1) **Small-scale processes and their impacts on large-scale climate dynamics and biogeochemical cycles.**  
362 The number of observations in the sea is smaller than those ~~over~~ land. This is also true for the Baltic  
363 Sea, although the international long-term monitoring programme in the Baltic Sea started ~~more than~~  
364 a century ago, with measurements of temperature, salinity and oxygen concentration ~~in from~~ the central  
365 parts of the ~~different~~ various sub-basins. Nowadays, monitoring data are available from all sub-basins with  
366 a resolution of up to one month. Recently, many new observational systems for temporally and spatially  
367 high-resolution data have been developed or are under development, including remotely operated vehicles  
368 (ROVs) and autonomous underwater vehicles (AOVs) as well as remote sensing data. Examples of such  
369 systems operating in the Baltic Sea are continuously profiling moorings, ARGO floats, Gliders, ScanFish,  
370 and echo sounders. In addition to traditional physical parameters, measurements of turbulence,  
371 biogeochemical and biodiversity (e.g. environmental DNA) parameters are now available. Another area  
372 of research that is developing rapidly is numerical modelling of the Earth system, also on a regional scale,  
373 e.g. eddy- and submesoscale resolving multi-year simulations for the Baltic Sea. ~~Similar~~ Corresponding

arguments apply to the atmosphere, e.g. cloud-resolving simulations to cope with heavy precipitation events. A novel research topic for Baltic Earth would therefore be a better understanding of the dynamics of small-scale atmospheric and oceanic processes that are not yet resolved in state-of-the-art numerical models or in-conventional monitoring-observations, and their role in the large-scale circulation on short and long time scales. Such research activities would help to fill some of the gaps in knowledge that have been raised by Lehmann et al. (2022), Kuliński et al. (2022), Rutgersson et al. (2022), Weisse et al. (2021), Viitasalo and Bonsdorff (2022), and Gröger et al. (2021). Furthermore, a realistic consideration of small-scale processes would improve the projections for the atmosphere (Christensen et al., 2022) and the ocean (Meier et al., 2022<sup>ab</sup>).

- 2) **Attribution of regional climate variability and change to anthropogenic radiative forcing and versus other drivers.** In order to unambiguously disentangle the impacts of anthropogenic climate change and other human influences from the natural climate variability of the regional Earth system, more knowledge about internal variations and feedback mechanisms is needed. For example, climate models have recently shown that multi-decadal variability emanating/originating from the North Atlantic and the Arctic significantly controls the climate of the Baltic Sea region by means of teleconnection patterns (Lehmann et al., 2022; Meier et al., 2022a; 2022b). For example, observations of precipitation and wind in the Baltic Sea region, total river discharge from the catchment, individual river flows, water temperature, sea level, MBIs and salinity in the Baltic Sea show a pronounced multidecadal variability with a quasi-periodicity of about 30 years (Meier et al., 2022<sup>ba</sup>). It is assumed that the Atlantic Multidecadal Variability (AMV) and, as part of it, the variations of the North Atlantic overturning circulation are the source of these variations, although the exact mechanisms, cause and effect chains and feedback processes are still unknown. ~~For the development of climate prediction modeling systems, knowledge about the teleconnectivity of the Baltic Sea region with the North Atlantic and the Arctic is essential.~~ Knowledge about the teleconnectivity of the Baltic Sea region with the North Atlantic and the Arctic is essential for the development of climate prediction models.
- 3) **Development of integrated regional-Earth system models accounting for anthropogenic changes/multiple drivers of Earth system changes in the Baltic Sea region.** The BEAR study by Reckermann et al. (2022) on human influences and their interactions in the Baltic Sea region is part of the relatively new Baltic Earth Grand Challenge 6 on the multiple drivers of Earth system change in the Baltic Sea region and represents an important step towards an integrated understanding of the Earth system ~~that~~, encompassing all traditionally considered climate compartments such as atmosphere, cryosphere, hydrosphere, lithosphere (including the pedosphere), biosphere (marine and terrestrial) and the anthroposphere. Such a holistic view is urgently needed, ~~as~~ in very-many cases, several ~~reasons/causes~~ are responsible for the observed changes in the Earth system and attributing them to only one factor, e.g. climate change, would be an inadmissible simplification. One example is the oxygen depletion and the large hypoxic area in the Baltic Sea caused by anthropogenic nutrient inputs from land and exacerbated by rising water temperatures (Kuliński et al., 2022). Of course, the factors discussed by

413 Reckermann et al. (2022) cannot ~~be exhaustively considered to take into account~~ the ~~entire~~whole Earth  
414 system and all interactions, and ~~of course there is a bias the selection of factors is biased~~ towards  
415 ~~ocean~~marine-related parameters and activities. Moreover, the analysis is based on an extensive literature  
416 review by experts who reflect their subjective interpretations of the results. Nevertheless, this is the first  
417 time such an assessment has been conducted, which is a major step forward. To continue and deepen this  
418 research, the factors discussed by Reckermann et al. (2022) could be ~~subdivided~~classified either by human  
419 activities (e.g. food production, energy production, transport, tourism, healthcare) or by environmental  
420 and climate state variables of the Earth system (e.g. hypoxia, acidification) (Table 3). Such a breakdown  
421 of parameters would allow the development of an integrated Earth system model that includes the  
422 anthroposphere at the regional scale. This type of research is timely, and such efforts are already  
423 underway (e.g. Korpinen et al., 2019; references in Reckermann et al., 2022).

424  
425 The fact sheet on climate change in the Baltic Sea (CCFS, 2021) was positively received by various stakeholders  
426 and decision-makers. Although uncertainties regarding observed and projected future climate change and the other  
427 drivers remain high, our experience engaging with stakeholders confirms that scientific uncertainties are taken into  
428 account in different ways in management and decision-making. This is an important reason for investing in the  
429 above key issues. They have the and their potential to reduce uncertainties that currently hamper decision-making  
430 in the region.

## 431 **5 Concluding remarks**

432 We conclude that 1) the BEARs have been useful to identify research progress and knowledge gaps and to initiate  
433 new research foci as, for examples, suggested in the discussions; 2) regional assessments, such as the BEARs,  
434 complement the IPCC climate change assessments by adding a greater depth and scope of regional information  
435 about the specific situation of the Baltic Sea region; and 3) the BEARs provided useful information for the Expert  
436 Network on Climate Change, that produced the Baltic Earth – HELCOM climate change fact sheet for  
437 stakeholders. Since the information summarised by the BEARs are used extensively in science and management,  
438 it is recommended that a new update of the reports will be conducted in about seven years.

## 439 **Author contributions**

440 H.E.M.M. wrote the first draft of the editorial. All co-authors, which acted as guest editors of the special issue in  
441 *Earth System Dynamics*, contributed ~~substantially~~ with important comments and editing of the manuscript, read  
442 and approved the submitted manuscript version.

## 443 **Acknowledgements**

444 During 2019-2022, the Baltic Earth Assessment Reports were ~~produced~~developed under the umbrella of  
445 the Baltic Earth programme (Earth System Science for the Baltic Sea region, see <http://baltic.earth>, last  
446 access: 2 February 2023). 109 co-authors from 14 countries contributed to 10 articles in the international


Formatiert: Schriftart: Kursiv

447 scientific journal *Earth System Dynamics* and 2822 different references have been assessed. We thank the  
448 reviewers of all 10 articles of the special issue for their constructive comments that helped to improve the  
449 review articles. In particular, we thank Dr. Jouni Räisänen, Dr. Donald Boesch and Dr. Andris Andrusaitis for  
450 their advice and many excellent comments on individual articles and this overview article.  
451

**Formatiert:** Schriftart: Kursiv

**Formatiert:** Englisch (USA)


452 **Figures**




453  
454 **Figure 1:** The Baltic Sea and its catchment area with climatological mean sea surface salinity (in g kg<sup>-1</sup>) and river  
455 discharge (in mm year<sup>-1</sup>). (Source: Meier et al., 2014; their Fig. 1 distributed under the terms of the Creative  
456 Commons CC-BY 4.0 License, <http://creativecommons.org/licenses/by/4.0/>, last access: 4 February 2023)

457

458




459



460

461 **Figure 2:** Annual mean sea surface temperature anomalies relative to the reference period 2002-2018 from de-  
462 seasonalised measurements at the Arkona Deep monitoring station and the MARNET stations Darss Sill and  
463 Arkona Basin in the period 1980-2018. (Source: Meier et al., 2022<sup>ba</sup>; their Fig. 20 distributed under the terms of  
464 the Creative Commons CC-BY 4.0 License, <http://creativecommons.org/licenses/by/4.0/>, last access: 4 February  
465 2023)  
466





468  
 469 **Figure 3:** Synthesis of knowledge on present and future climate change. Shown are anthropogenic climate changes  
 470 in 33 Earth system variables (bubbles) of the atmosphere (yellow), land surface (brown), terrestrial biosphere (dark  
 471 green), cryosphere (grey), ocean and sediments (blue) and marine biosphere (light green). The abbreviation NIS  
 472 stands for non-indigenous species. The sign of a change (plus/minus) is shown together with the confidence level  
 473 indicated by the number of signs, i.e. one to three signs correspond to a low, medium and high confidence level as  
 474 a result of the literature assessment reflecting consensus and evidence according to IPCC definitions. The colours  
 475 of the signs indicate the direction of past (black) and future (red) changes according to Meier et al. (2022<sup>ba</sup>).  
 476 Uncertain changes are not shown. The external anthropogenic drivers of the Earth system studied are shown as red  
 477 squares, i.e. greenhouse gases, especially CO<sub>2</sub>, and aerosol emissions. The predominant climate change linkages  
 478 with sufficiently high confidence are shown by arrows (black: thermal cycle, blue: hydrological cycle, orange:  
 479 momentum cycle including sea level change, pink: carbon cycle). Projections of carbon sequestration of Arctic  
 480 terrestrial ecosystems for the 21<sup>st</sup> century show first an increased uptake and later a carbon source, marked by “1.  
 481 + 2. -”. Future changes in mean sea level are dominated by the thermal expansion of the global ocean and the  
 482 melting of ice sheets outside the Baltic Sea region. (Source: Meier et al., 2022<sup>ba</sup>; their Fig. 35 distributed under  
 483 the terms of the Creative Commons CC-BY 4.0 License, <http://creativecommons.org/licenses/by/4.0/>, last access:  
 484 4 February 2023)

486 **Tables**

487 **Table 1:** The matrix of factors studied by Reckermann et al. (2022). += evidence for a connection; -= no evidence  
 488 for a connection; ? = no evidence, but connection plausible (according to the author's assessment). The table is  
 489 read from left to right, i.e. if you go to the right in the first row "climate change", you see the factors on which  
 490 climate change has an effect (or not), etc. (Source: Reckermann et al., 2022; their Table 2a distributed under the  
 491 terms of the Creative Commons CC-BY 4.0 License, <http://creativecommons.org/licenses/by/4.0/>, last access: 4  
 492 February 2023)

impact by ↓ / on →	Climate change	Coastal processes	Hypoxia	Acidification	Subm. Groundw. Disch.	Marine ecosystems	Non-inlg. species	Land cover and use	Agriculture Nutr. loads	Aquaculture	Fisheries	River regulations	Offshore wind farms	Shipping	Chem. Contamin.	Dumped military	Marine litter	Tourism	Coastal management	
Climate change		+	+	+	?	+	?	+	+	+	+	+	+	+	+	+	?	?	+	+
Coastal processes	-		?	?	+	?	-	+	+	?	?	+	+	+	?	?	+	-	+	
Hypoxia	-	-		+	-	+	-	-	+	?	+	-	-	-	-	+	+	-	-	
Acidification	-	-	-		?	?	-	-	-	?	?	-	-	-	-	?	-	-	-	
Subm. Groundw. Disch.	-	-	?	?		?	-	-	+	-	-	-	-	-	+	-	-	-	-	
Marine ecosystems	-	-	+	+	-		+	-	-	-	+	-	-	-	-	-	-	-	+	
Non-inigenous species	-	-	-	-	-	+		-	-	-	+	-	-	+	+	-	-	-	?	
Land cover and use	+	-	+	+	+	?	-		+	+	+	+	?	-	+	-	-	-	+	
Agriculture/Nutrient loads	+	-	+	+	+	+	-	+		+	+	+	-	-	+	-	-	-	?	
Aquaculture	-	-	+	-	-	+	+	+	+	?	-	+	-	?	-	?	-	?	?	
Fisheries	-	-	?	-	-	+	?	-	?	?		+	+	?	-	-	+	?	+	
River regulations	-	+	?	+	?	?	?	?	?	?	+		-	-	?	-	?	-	+	
Offshore wind farms	+	+	-	-	-	+	-	?	?	?	+	+	-	-	+	?	?	?	+	
Shipping	+	+	-	+	-	+	+	-	+	?	+	-	?		+	-	+	+	+	
Chemical contaminants	-	-	-	-	-	+	-	-	+	+	+	-	-	-	-	-	-	-	-	
Dumped military material	-	-	-	-	-	?	-	-	-	-	+	-	+	-	+				?	
Marine litter	-	-	-	-	-	?	-	-	-	?	-	-	-	-	?	-	-	+	?	
Tourism	+	-	-	-	-	?	-	+	+	-	-	-	+	+	-	-	+		+	
Coastal management	-	+	-	-	?	?	?	?	?	?	?	?	+	+	-	+	?	+	+	

493  
 494

495 **Table 2:** Variables of the Meier et al. (2022<sup>ba</sup>) assessment and further references to the BEARs (1: Lehmann et  
496 al., 2022; 2: Kuliński et al., 2022; 3: Rutgersson et al., 2022; 4: Weisse et al., 2021; 5: Reckermann et al., 2022; 6:  
497 Gröger et al., 2021; 7: Christensen et al., 2022; 8: Meier et al., 2022<sup>ab</sup>; 9: Viitasalo and Bonsdorff, 2022). The third  
498 column lists the subsection in the study by Meier et al. (2022<sup>ba</sup>) that contains further information. (Source: Meier  
499 et al., 2022<sup>ba</sup>; their Table 2 distributed under the terms of the Creative Commons CC-BY 4.0 License,  
500 <http://creativecommons.org/licenses/by/4.0/>, last access: 4 February 2023 )

Number	Variable	Past and present climates		Future climate	
Atmosphere					
1	Large-scale atmospheric circulation	3.2.1.1	3	3.3.1.1	3, 7
2	Air temperature	3.1.2, 3.1.3, 3.1.4		3.3.1.2	7
	Warm spell	3.2.1.2	3		3
	Cold spell		3		3
3	Solar radiation and cloudiness	3.2.1.3		3.3.1.3	7
4	Precipitation	3.1.2, 3.1.3, 3.1.4		3.3.1.4	7
	Heavy precipitation	3.2.1.4	3		3
	Drought		3		3
5	Wind	3.2.1.5		3.3.1.5	7
	Storm		3		3
6	Air pollution, air quality and atmospheric deposition	3.2.1.6		3.3.1.6	
Land					
7	River discharge	3.2.2.1		3.3.2.1	8
	High flow		3		3
8	Land nutrient inputs	3.2.2.2		3.3.2.2	8
Terrestrial biosphere					
9	Land cover (forest, crops, grassland, peatland, mires)	3.2.3	6	3.3.3	
10	Carbon sequestration			3.3.3	
Cryosphere					

11	Snow Sea-effect snowfall	3.2.4.1	3	3.3.4.1	7 3
12	Glaciers	3.2.4.2		3.3.4.2	
13	Permafrost	3.2.4.3		3.3.4.3	
14	Sea ice Extreme mild winter Severe winter Ice ridging	3.2.4.4	3 3 3	3.3.4.4	8 3 3 3
15	Lake ice	3.2.4.5		3.3.4.5	
Ocean and marine sediments					
16	Water temperature Marine heat wave	3.2.5.1	3	3.3.5.1	8 3
17	Salinity and saltwater inflows	3.2.5.2	1	3.3.5.2	8
18	Stratification and overturning circulation	3.2.5.3	1	3.3.5.3	8
19	Sea level Sea level extreme	3.2.5.4	4 3	3.3.5.4	8 3
20	Waves Extreme waves	3.2.5.5	4 3	3.3.5.5	3
21	Sedimentation and coastal erosion	3.2.5.6	4	3.3.5.6	
22	Oxygen and nutrients	3.1.4 3.2.5.7.1	2	3.3.5.7.1	8
23	Marine CO <sub>2</sub> system	3.2.5.7.2	2	3.3.5.7.2	
Marine biosphere					
24	Pelagic habitats: Microbial communities	3.2.6.1.1	2, 9	3.3.6.1.1	9
25	Pelagic habitats: Phytoplankton and cyanobacteria	3.2.6.1.2	2, 3, 9	3.3.6.1.2	3, 9
26	Pelagic habitats: Zooplankton	3.2.6.1.3	9	3.3.6.1.3	9
27	Benthic habitats: Macroalgae and vascular plants	3.2.6.2.1	9	3.3.6.2.1	9

28	Benthic habitats: Zoobenthos	3.2.6.2.2	9	3.3.6.2.2	9
29	Non-indigenous species	3.2.6.3	9	3.3.6.3	9
30	Fish	3.2.6.4	9	3.3.6.4	9
31	Marine mammals	3.2.6.5	9	3.3.6.5	9
32	Waterbirds	3.2.6.6	9	3.3.6.6	9
33	Marine food web	3.2.6.7	9	3.3.6.7	9

501

502 **Table 3:** Factors discussed by Reckermann et al. (2022) sorted by related economic sectors or state variables of  
 503 the Earth system.

<b>Human activities</b>		
Economic sectors	Factors	Comments
Primary (natural ) sector (e.g. food production)	Fisheries	
	Agriculture	
	Marine and coastal ecosystem services	Factor belongs to several sectors
	Blue carbon storage capacity	Mitigation of greenhouse gases
Secondary (industrial) sector (e.g. energy production)	River regulation	
	Offshore wind farms	
	Greenhouse gas and aerosol emissions	Emission are largest from industries
	Dumped warfare agents	Factor is an industrial product
Tertiary (service) sector (e.g. transportation, tourism, healthcare)	Shipping	
	Chemical contamination	Contamination is a result of several sectors
	Marine noise	Marine noise is a result of several sectors
	Marine litter and microplastics	Emission mainly by offshore platforms, shipping, lost containers, fisheries, aquaculture, agriculture, municipal waste and tourism
	Tourism	
	Coastal protection and management	Also relevant for the other sectors
Quaternary (information) sector (e.g. information technology; media; research and development)	-	
<b>Earth system</b>		
Environmental state variables	Coastal processes	
	Hypoxia	
	Submarine groundwater discharge	
	Marine ecosystems	
	Land use and land cover	
	Non-indigenous species	

	Indirect parameters such as carbon and nutrient cycles, biota and ecosystems	
Climate state variables	Climate change, acidification, direct parameters of the climate system	Superordinated concept (large-scale)
	Direct parameters of the climate system	
	Acidification	

504

505 **References**

506 Baltic Earth Science Plan, 2017. <https://baltic.earth/grandchallenges>, last access: 4 February 2023

507

508 CCFS, 2021. Climate Change in the Baltic Sea. 2021 Fact Sheet. Baltic Sea Environment Proceedings n° 180.

509 HELCOM/Baltic Earth, <http://doi.io-warnemuende.de/10.12754/misc-2022-0001>, 2021.

510

511 Christensen, O. B., Kjellström, E., Dieterich, C., Gröger, M., and Meier, H. E. M.: Atmospheric regional climate

512 projections for the Baltic Sea Region until 2100, *Earth Syst. Dynam.*, 13, 133–157, [https://doi.org/10.5194/esd-](https://doi.org/10.5194/esd-13-133-2022)

513 [13-133-2022](https://doi.org/10.5194/esd-13-133-2022), 2022.

514

515 Gröger, M., Dieterich, C., Haapala, J., Ho-Hagemann, H. T. M., Hagemann, S., Jakacki, J., May, W. Meier, H. E.

516 M., Müller, P. A. Rutgersson, A. and Wu, L.: Coupled regional Earth system modeling in the Baltic Sea region,

517 *Earth Syst. Dynam.*, 12, 939–973, <https://doi.org/10.5194/esd-12-939-2021>, 2021.

518

519 [Klimawandel in der Ostsee. 2021 Faktenblatt - Baltic Sea Climate Change in the Baltic Sea, 2021 Fact Sheet \(in](http://doi.io-warnemuende.de/10.12754/misc-2022-0003)

520 [German\). http://doi.io-warnemuende.de/10.12754/misc-2022-0003](http://doi.io-warnemuende.de/10.12754/misc-2022-0003), 2022.

521

522 [Korpinen, S., Klančnik, K., Peterlin, M., Nurmi, M., Laamanen, L., Zupančič, G., Popit, A., Murray, C., Harvey,](https://doi.org/10.5194/esd-12-939-2021)

523 [T., Andersen, J.H., Zenetos, A., Stein, U., Tunesi, L., Abhold, K., Piet, G., Kallenbach, E., Agnesi, S., Bolman,](https://doi.org/10.5194/esd-12-939-2021)

524 [B., Vaughan, D., Reker, J. & Royo Gelabert, E.: Multiple pressures and their combined effects in Europe's seas,](https://doi.org/10.5194/esd-12-939-2021)

525 [ETC/ICM Technical Report 4/2019: European Topic Centre on Inland, Coastal and Marine waters, 164 pp., 2019.](https://doi.org/10.5194/esd-12-939-2021)

526

527 Kuliński, K., Rehder, G., Asmala, E., Bartosova, A., Carstensen, J., Gustafsson, B., Hall, P. O. J., Humborg, C.,

528 Jilbert, T., Jürgens, K., Meier, H. E. M., Müller-Karulis, B., Naumann, M., Olesen, J. E., Savchuk, O., Schramm,

529 A., Slomp, C. P., Sofiev, M., Sobek, A., Szymczycha, B., and Undeman, E.: Biogeochemical functioning of the

530 Baltic Sea. *Earth Syst. Dynam.*, 13, 633–685, <https://doi.org/10.5194/esd-13-633-2022>, 2022.

531

532 Lehmann, A., Myrberg, K., Post, P., Chubarenko, I., Dailidienė, I., Hinrichsen, H.-H., Hüseyin, K., Liblik, T., Meier,

533 H. E. M., Lips, U., and Bukanova, T.: Salinity dynamics of the Baltic Sea, *Earth Syst. Dynam.*, 13, 373–392,

534 <https://doi.org/10.5194/esd-13-373-2022>, 2022.

535

536 Meier, H.E.M., A. Rutgersson, and M. Reckermann: An Earth System Science Program for the Baltic Sea Region.

537 *Eos*, 95, 109–110. <https://doi.org/10.1002/2014EO130001>, 2014.

538

539 Meier, H. E. M., Dieterich, C., Gröger, M., Dutheil, C., Börgel, F., Safonova, K., Christensen, O. B., and

540 Kjellström, E.: Oceanographic regional climate projections for the Baltic Sea until 2100, *Earth Syst. Dynam.*, 13,

541 159–199, <https://doi.org/10.5194/esd-13-159-2022>, 2022<sup>ab</sup>.

542

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)



543 Meier, H. E., Kniebusch, M., Dieterich, C., Gröger, M., Zorita, E., Elmgren, R., ... and Zhang, W.: Climate change  
544 in the Baltic Sea region: a summary. *Earth Syst. Dynam.*, 13, 457-593, <https://doi.org/10.5194/esd-13-457-2022>,  
545 2022b.

546

547 ~~Ostsee Klimawandel Faktenblatt: Baltic Sea Climate Change Fact Sheet (in German).~~ [http://doi.io-](http://doi.io-warnemuende.de/10.12754/misc-2022-0003)  
548 [warnemuende.de/10.12754/misc-2022-0003](http://doi.io-warnemuende.de/10.12754/misc-2022-0003), 2022.

549

550 Reckermann, M., Omstedt, A., Soomere, T., Aigars, J., Akhtar, N., Beldowski, J., Brauwer, C. P.-S. d., Cronin,  
551 T., Czub, M., Eero, M., Hyttiäinen, K., Jalkanen, J.-P., Kiessling, A., Kjellström, E., Larsén, X. G., McCrackin,  
552 M., Meier, H. E. M., Oberbeckmann, S., Parnell, K., Poska, A., Saarinen, J., Szymczycha, B., Undeman, E.,  
553 Viitasalo, M., Wörman, A., and Zorita, E.: Human impacts and their interactions in the Baltic Sea region. *Earth*  
554 *Syst. Dynam.*, 13, 1–80, <https://doi.org/10.5194/esd-13-1-2022>, 2022.

555

556 Rutgersson, A., Kjellström, E., Haapala, J., Stendel, M., Danilovich, I., Drews, M., Jylhä, K., Kujala, P., Larsén,  
557 X. G., Halsnæs, K., Lehtonen, I., Luomaranta, A., Nilsson, E., Olsson, T., Särkkä, J., Tuomi, L., and Wasmund,  
558 N.: Natural Hazards and Extreme Events in the Baltic Sea region, *Earth Syst. Dynam.*, 13, 251–301,  
559 <https://doi.org/10.5194/esd-13-251-2022>, 2022.

560

561 The BACC Author Team, 2008. Assessment of Climate Change for the Baltic Sea Basin. Regional Climate Studies  
562 series. Heidelberg, Springer-Verlag Berlin. ISBN 978-3-540-72785-9. 473 pp. [https://doi.org/10.1007/978-3-540-](https://doi.org/10.1007/978-3-540-72786-6)  
563 [72786-6](https://doi.org/10.1007/978-3-540-72786-6).

564

565 The BACC II Author Team, 2015. Second Assessment of Climate Change for the Baltic Sea Basin. Regional  
566 Climate Studies series. Cham, Springer Cham. ISBN 978-3-319-16005-4. 501 pp. [https://doi.org/10.1007/978-3-](https://doi.org/10.1007/978-3-319-16006-1)  
567 [319-16006-1](https://doi.org/10.1007/978-3-319-16006-1).

568

569 Viitasalo, M., and Bonsdorff, B.: Global climate change and the Baltic Sea ecosystem: direct and indirect effects  
570 on species, communities and ecosystem functioning. *Earth Syst. Dynam.*, 13, 711–747,  
571 <https://doi.org/10.5194/esd-13-711-2022>, 2022.

572

573 Weisse, R., Dailidienė, I., Hünicke, B., Kahma, K., Madsen, K., Omstedt, A., Parnell, K., Schöne, T., Soomere,  
574 T., Zhang, W., and Zorita, E.: Sea Level Dynamics and Coastal Erosion in the Baltic Sea Region. *Earth Syst.*  
575 *Dynam.*, 12, 871–898, <https://doi.org/10.5194/esd-12-871-2021>, 2021.