

1 **The Baltic Earth Assessment Reports**

2 H. E. Markus Meier¹, Marcus Reckermann², Joakim Langner³, Ben Smith^{4,5} and Ira
3 Didenkulova⁶

4 ¹Department of Physical Oceanography and Instrumentation, Leibniz Institute for Baltic Sea Research
5 Warnemünde, Rostock, Germany

6 ²International Baltic Earth Secretariat, Helmholtz-Zentrum Hereon, Geesthacht, Germany

7 ³Swedish Meteorological and Hydrological Institute, Norrköping, Sweden

8 ⁴Hawkesbury Institute for the Environment, Western Sydney University, Australia

9 ⁵Department of Physical Geography and Ecosystem Science, Lund University, Sweden

10 ⁶Department of Mathematics, University of Oslo, Norway

11

12 *Correspondence to:* H. E. Markus Meier (markus.meier@io-warnemuende.de)

13

14 **Abstract.** Baltic Earth is an independent research network of scientists from all Baltic Sea countries that promotes
15 regional Earth system research. Within the framework of this network, the Baltic Earth Assessment Reports
16 (BEARs) were produced in the period 2019-2022. These are a collection of 10 review articles summarising current
17 knowledge on the environmental and climatic state of the Earth system in the Baltic Sea region and its changes in
18 the past (palaeoclimate), present (historical period with instrumental observations) and prospective future (until
19 2100) caused by natural variability, climate change and other human activities. The division of topics among
20 articles follows the grand challenges and selected themes of the Baltic Earth Science Plan, such as the regional
21 water, biogeochemical and carbon cycles, extremes and natural hazards, sea level dynamics and coastal erosion,
22 marine ecosystems, coupled Earth system models, scenario simulations for the regional atmosphere and the Baltic
23 Sea, and climate change and impacts of human use. Each review article contains an introduction, the current state
24 of knowledge, knowledge gaps, conclusions and key messages, based on which recommendations for future
25 research are made. Based on the BEARs, Baltic Earth has published an information leaflet on climate change in
26 the Baltic Sea as part of its outreach work, which has been published in two languages so far, and organised
27 conferences and workshops for stakeholders, in collaboration with the Baltic Marine Environment Protection
28 Commission (HELCOM).

29 **1 Introduction**

30 **1.1 BALTEX/Baltic Earth history**

31 Baltic Earth¹ is an international research network dealing with Earth system sciences in the Baltic Sea region (Fig.
32 1). It is politically independent and focuses on research on the water and energy cycles, climate variability and
33 climate change, water management and extreme events, and associated impacts on marine and terrestrial
34 biogeochemical cycles. Research on human impact on the Earth system in more general terms, i.e. the
35 anthroposphere, defined as the part of the environment created or modified by humans for use by human activities,
36 was also included in the Baltic Earth Science Plan (2017)².

37

38 Baltic Earth is the successor of the Baltic Sea Experiment (BALTEX) programme, which was founded in 1993 as
39 a GEWEX continental-scale experiment (Global Energy and Water Exchanges, a core project of the World Climate
40 Research Programme) (Reckermann et al., 2011). In the first phase (1993–2002), BALTEX was primarily devoted
41 to hydrological, meteorological and oceanographic processes in the Baltic Sea drainage basin and thus focused on
42 physical aspects of the Earth system. In the second phase (2003–2012), the programme was expanded to include
43 regional climate research, biogeochemical cycles including carbon, engagement with stakeholders and decision-
44 makers via assessment reports, as well as communication and education, i.e. the organisation of summer and winter
45 schools and international master courses.

46

¹ <https://baltic.earth>, last access: 4 February 2023

² <https://baltic.earth/grandchallenges>, last access: 4 February 2023

47 In 2013, Baltic Earth was launched with a new science plan to strengthen efforts to address Grand Challenges on
48 (1) salinity dynamics in the Baltic Sea, (2) biogeochemical linkages between land and sea, (3) natural hazards and
49 extreme events, (4) sea level and coastal dynamics, (5) regional variability in water and energy exchanges, and (6)
50 multiple drivers of regional Earth system changes (Meier et al., 2014). Working groups on coupled Earth system
51 models, the Baltic Sea Model Intercomparison Project (BMIP), uncertainty of scenario simulations for the Baltic
52 Sea, and education, outreach and communication have been established.

53
54 Baltic Earth and its predecessor BALTEX have produced three comprehensive regional assessment reports since
55 2008. The first two (The BACC Author Team, 2008, and The BACC II Author Team, 2015) focused on climate
56 change and its impacts in the Baltic Sea region and were published as text books, while the third, the Baltic Earth
57 Assessment Reports (BEARs), was published in the format of a special issue of the journal *Earth System Dynamics*
58 in 2022. The Assessment of Climate Change in the Baltic Sea Basin (BACC) reports³ and BEARs fill a gap
59 compared to the assessment reports of the Intergovernmental Panel on Climate Change (IPCC), as the latter focus
60 on global scales, and do not provide detailed local to regional information on the current state of knowledge on
61 climate change and its impacts in the Baltic Sea region. The BEARs provide a comprehensive and up-to-date
62 overview of the state-of-the-art research on the compartments of the Earth system in the Baltic Sea region
63 encompassing processes in the atmosphere, on land and in the sea, including the marine and terrestrial ecosystems
64 as well as processes and impacts related to the anthroposphere.

65
66 The BEARs summarise the published scientific knowledge currently available and update the second BACC report
67 (The BACC II Author Team, 2015) based on the latest scientific literature. This BEAR special issue includes 10
68 articles on the Baltic Earth Grand Challenges and Baltic Earth Special Topics (Baltic Earth Science Plan, 2017),
69 including a summary of current knowledge on past, present, and future climate change in the Baltic Sea region.
70 The articles encompass contributions from 109 authors from 14 countries and reference 2822 scientific articles
71 and institutional reports.

72 **1.2 Baltic Sea region characteristics**

73 The Baltic Sea is a semi-enclosed, shallow sea with limited water exchange with the World Ocean and small tidal
74 amplitudes. Located in Northern Europe, the climate of the region is highly variable as it is in the transition zone
75 between maritime and continental climates and is influenced by the North Atlantic and Arctic. River discharges
76 from the large catchment area cause a pronounced gradient in sea surface salinity from about 20 g kg⁻¹ in the
77 Danish straits' region to about 2 g kg⁻¹ or even less in the northern and eastern reaches of the Baltic Sea. Hence,
78 the Baltic Sea is brackish, with habitats of marine species in the south-west and freshwater species in the north-
79 east. The Baltic Sea catchment area is about four times the surface area of the Baltic Sea and covers an area of
80 almost 20% of the European continent (Fig. 1). It stretches from the temperate, densely populated south to the

³ Assessment of Climate Change in the Baltic Sea Basin (BACC); <https://baltic.earth/bacc>, last access: 4 February 2023

81 subarctic wilderness in the north and is home to approximately 85 million people in 14 countries, namely Belarus,
82 the Czech Republic, Denmark, Estonia, Finland, Germany, Latvia, Lithuania, Norway, Poland, Russia, Slovakia,
83 Sweden and Ukraine.

84
85 Episodically, large amounts of saline water flow from the North Sea over the sills in the Danish straits into the
86 Baltic Sea and ventilate the deep waters of the Baltic Sea. These events require a period of about 20 days with
87 easterly winds that lower the sea level in the Baltic Sea, followed by a period of about the same length with strong
88 westerly winds that push saline water into the Baltic Sea. These events are called Major Baltic Inflows (MBIs) and
89 are important for the water exchange between the North Sea and the Baltic Sea. Mixing is low compared to other
90 seas, with an origin at the lateral boundaries, because tidal amplitudes are very small and energetically
91 insignificant.

92
93 In recent decades, environmental conditions in the Baltic Sea have changed considerably. For instance, the Baltic
94 Sea has been warming more than any other coastal sea since 1980 (Fig. 2), which has led to a reduction in sea ice
95 and snow cover over the land in winter. Furthermore, increasing nutrient input from the land in the 1950s/60s,
96 caused by population growth and the discharge of sewage into the Baltic Sea, as well as the increased use of
97 fertilisers in agriculture, led to eutrophication and the spread of hypoxic and anoxic areas. Since the 1980s, nutrient
98 inputs into the Baltic Sea have been steadily decreasing, but this has not yet led to a significant improvement in
99 oxygen conditions. Recent trends in acidification are lower than in the World Ocean, especially in the northern
100 Baltic Sea, as positive trends in alkalinity input counteract acidification.

101 **2 Methods**

102 Succeeding The BACC Author Team (2008) and The BACC II Author Team (2015) assessments, the BEAR
103 project is an attempt to summarise the scientific knowledge on climate change and other drivers of Earth system
104 changes and their impacts on the Baltic Sea region. The two BACC books have a format inspired by the IPCC
105 assessment reports. This special issue in *Earth System Dynamics* is the third assessment. It has a new format of
106 BEARs, encompassing 10 peer-reviewed scientific journal articles. The knowledge assessed was extracted from
107 the scientific literature such as peer-reviewed articles, reports from research institutions, and published datasets.
108 Importantly, literature from non-governmental organisations with political or economic interests, political parties
109 and other stakeholder organisations was excluded from the assessment to ensure that only scientific knowledge
110 was included in the assessment. The BEARs focus on publications after 2012/2013, the year of the editorial
111 deadline of the second assessment report. Whenever possible, the uncertainty levels of the BEAR results are ranked
112 based on a matrix of consensus within the scientific literature and documented evidence of detected changes and
113 their attributed drivers such as climate change and human use. A high level of scientific consensus and evidence
114 is required for high confidence in a particular statement. Disagreements and gaps in knowledge are documented
115 and discussed to prioritise future research.

116

117 Together with the intergovernmental Baltic Marine Environment Protection Commission (HELCOM), Baltic
118 Earth has established an Expert Network on Climate Change (EN CLIME). The aim of the expert network is to
119 regularly produce a climate change fact sheet (CCFS, 2021⁴) based on the BEAR and BACC material. In 2021, it
120 was published for the first time⁵. The CCFS contains some background information, a map showing regional future
121 climate changes for selected parameters under the greenhouse gas concentration scenario RCP4.5 and information
122 on 34 variables, directly and indirectly affected by climate change. For each parameter, a general description, past
123 and prospective future changes, other drivers than climate change (only for the indirect parameters), knowledge
124 gaps, policy relevance and references are presented. More than 100 scientists contributed to the compilation of the
125 first fact sheet, which was coordinated by the HELCOM secretariat. Updated versions are planned at seven-year
126 intervals. Like the BEARs, the fact sheet was peer-reviewed and quality assured. It has so far only been translated
127 to German (Klimawandel in der Ostsee, 2021 Faktenblatt, 2022⁶), but translations into other languages are planned
128 to improve accessibility to stakeholders.

129

130 In this editorial, we highlight the key findings and knowledge gaps as described by the BEARs and propose future
131 work.

132 **3 Results**

133 Some of the key findings of the 10 BEARs are selected and highlighted below.

134

- 135 1. **Salinity dynamics of the Baltic Sea**, Grand Challenge 1 (Lehmann et al., 2022): Salinity is an important
136 parameter for the circulation and the marine ecosystem in the Baltic Sea. Any changes in salinity are
137 caused by changes either in the freshwater inflow from rivers and net precipitation over sea or in the water
138 exchange between the Baltic Sea and the adjacent North Sea. Although long-term records of salinity and
139 its drivers suffer from data gaps, these records starting in the 19th century are globally unique. Major
140 research efforts focused on the MBI event in 2014 and its consequences for water masses, oxygen
141 concentration and biogeochemical cycling. During the event, an unexpectedly large contribution of oxic
142 intrusions at intermediate depth and essentially nonturbulent conditions in the deep interior were found,
143 emphasising the importance of boundary mixing. A revised reconstruction of the long-term record of
144 MBIs showed no trend but a pronounced multi-decadal variability with a period of about 30 years. Despite
145 intense research activities, observed variations in the intensity and frequency of MBIs and related Large
146 Volume Changes (LVCs) could not be attributed to atmospheric circulation variability. Hence, on time
147 scales larger than the synoptical time scale, MBIs are not predictable. As an advance over the previous

⁴ <https://helcom.fi/wp-content/uploads/2021/09/Baltic-Sea-Climate-Change-Fact-Sheet-2021.pdf>, last access: 4 February 2023

⁵ <http://helcom.fi/ccfs>, last access: 4 February 2023

⁶ <https://baltic.earth/ccfs>, last access: 4 February 2023

148 assessments, salinity dynamics of the various sub-basins and lagoons mainly based on observations have
149 been discussed, documenting large regional differences.

150

151 2. **Biogeochemical functioning of the Baltic Sea**, Grand Challenge 2 (Kuliński et al., 2022): The review
152 addresses the following topics: (1) terrestrial biogeochemical processes and nutrient inputs to the Baltic
153 Sea, (2) the transformation of C, N and P in the coastal zone, (3) the production and remineralisation of
154 organic matter, (4) oxygen availability, (5) the burial and turnover of C, N and P in sediments, (6) the
155 Baltic Sea CO₂ system and seawater acidification, (7) the role of certain microorganisms in the
156 biogeochemistry of the Baltic Sea, and (8) the interactions between biogeochemical processes and
157 chemical pollutants. It was found that oxygen depletion and the area of anoxic bottoms have still increased
158 despite the reductions in nutrient inputs from land since the 1980s. Hence, the nitrogen pool has declined
159 due to denitrification whereas the phosphorus inventory has increased. Estimates suggest that about 1%
160 and 4% of the annual nitrogen and phosphorus loads, respectively, have accumulated in the Baltic Sea,
161 while the remainder are either exported to the North Sea or lost via biogeochemical processes such as
162 denitrification and burial. Furthermore, it was discovered that in the central and northern sub-basins the
163 uptake of C, N and P during primary production does not correspond to the Redfield ratio, which strongly
164 affects the relationship between primary production, export of organic matter and oxygen demand of the
165 deep sea. While it is clear that the Baltic Sea is a CO₂ sink in summer and a CO₂ source in winter, the
166 annual net balance remains unknown. The past increase in total alkalinity of unknown origin has entirely
167 mitigated ocean acidification in the northern Baltic Sea and significantly reduced it in the central Baltic
168 Sea. In the future, a doubling of atmospheric pCO₂ would still result in lower pH in the entire Baltic Sea,
169 even if alkalinity should further increase.

170

171 3. **Natural hazards and extreme events in the Baltic Sea region**, Grand Challenge 3 (Rutgersson et al.,
172 2022): Existing knowledge is summarised about extreme events in the Baltic Sea region with a focus on
173 the past 200 years with instrumental data as well as on future projections. Considered events are wind-
174 storms, extreme waves, high and low sea levels, hot and cold spells in the atmosphere, marine heat waves,
175 droughts, sea-effect snowfall, sea-ice ridging, extremely mild and extremely severe sea ice winters, heavy
176 precipitation events, river floods, and extreme phytoplankton blooms. Furthermore, the knowledge about
177 implications of these extreme events for society such as forest fires, coastal flooding, offshore
178 infrastructure and shipping was assessed. With respect to the impacts of climate change, terrestrial and
179 marine heat waves, extremely mild sea ice winters, heavy precipitation and high-flow events are expected
180 to increase, while cold spells, severe sea ice winters and sea-ice ridging are expected to decrease due to
181 the increase in mean atmospheric temperature. Changes in relative sea level extremes will depend on the
182 competing impacts of the rising global mean sea level, the gravitational effect of the melting of the
183 Greenland and Antarctic ice sheets, changes in wind fields, and the regionally differing Glacial Isostatic
184 Adjustment (GIA) resulting in land uplift or subsidence. Furthermore, projections suggest an increase of
185 droughts in the southern and central parts of the Baltic Sea region mainly in summer. Significant future

186 changes in wind-storms, extreme waves and sea level extremes relative to the mean sea level have not
187 been found, suggesting that these changes will likely be small compared with natural variability.

188
189 4. **Sea level dynamics and coastal erosion in the Baltic Sea region**, Grand Challenge 4 (Weisse et al.,
190 2021): In this study, the current knowledge about the diverse processes affecting mean and extreme sea
191 level changes, coastal erosion and sedimentation with impact on coastline changes and coastal
192 management is assessed. Such processes are GIA, contributions from global sea level changes, wind-
193 storms, wind-waves, seiches or meteotsunamis. During 1886-2020, the mean absolute sea level in the
194 Baltic Sea corrected for GIA increased by about 25 cm or ~ 2 mm year⁻¹ on average. Land uplift in the
195 north is still faster than the absolute sea level rise while in the south the opposite is true with potential
196 impacts on changes in coastal erosion and inundation. The current acceleration of sea level rise is small
197 and could only be determined by spatially averaging observations at different tide gauge locations. Future
198 sea level rise in the Baltic Sea is expected to further accelerate, probably somewhat less than the global
199 mean sea level rise. The Baltic sea level is substantially more sensitive to melting from the Antarctic than
200 from the Greenland ice sheet. Concerning sediment transports, the dominance of mobile sediments makes
201 the southern and eastern Baltic Sea coasts susceptible to wind-wave induced transports, in particular
202 during storms. Due to the global sea level rise, future sediment transports can be expected to increase in
203 these coastal areas, with a large spatial variability depending on the angles of incidence of incoming wind-
204 waves.

205
206 5. **Human impacts and their interactions in the Baltic Sea region**, Grand Challenge 6 (Reckermann et
207 al., 2022): An inventory and discussion of the various man-made factors and processes affecting the
208 environment of the Baltic Sea region and their interrelationships are presented. In total, 19 factors are
209 addressed (Table 1). Some of the factors are natural and are only modified by human activities (e.g.
210 climate change, coastal processes, hypoxia, acidification, submarine groundwater discharges, marine
211 ecosystems, non-indigenous species, land use and land cover), others are entirely man-made (e.g.
212 agriculture, aquaculture, fisheries, river regulation, offshore wind farms, shipping, chemical
213 contamination, dumped ammunition, marine litter and microplastics, tourism and coastal management).
214 All factors are interconnected to varying degrees. The knowledge of these linkages was assessed and
215 analysed in depth. The main finding was that climate change has an overarching, integrating effect on all
216 other factors and can be interpreted as a background effect that affects the other factors differently. After
217 climate change, shipping and land use/agriculture are the factors affecting most other factors, while
218 fisheries, marine ecosystems and agriculture in turn are the most affected. The results of the assessment
219 depend on the region and may be different for other coastal seas and their catchments in the world, where
220 different human activities prevail.

221
222 6. **Global climate change and the Baltic Sea ecosystem: direct and indirect effects on species,**
223 **communities and ecosystem functioning**, Baltic Earth Special Topic (Viitasalo and Bonsdorff, 2022):
224 Climate change has multiple impacts on species, communities and ecosystem functioning in the Baltic

225 Sea through changes in physical and biogeochemical parameters such as temperature, salinity, oxygen,
226 pH and nutrient levels. The associated secondary effects on species interactions, trophic dynamics and
227 ecosystem function are also likely to be important. Climate change (warming, recent brightening,
228 decrease in sea ice) has led to shifts in the seasonality of primary production, with a prolonged growing
229 season of phytoplankton, an earlier onset of the spring bloom and a delayed autumn bloom. However, the
230 development of cyanobacteria varies from species to species, and a clear causal relationship between
231 temperature or salinity and the abundance of cyanobacteria has not been demonstrated. An increase in
232 water temperature and river input of dissolved organic matter (DOM) could reduce primary production
233 while favouring bacterial growth. If nutrient reduction continues, the improvement in oxygen conditions
234 could initially increase zoobenthos biomass, but the subsequent decrease in sedimenting organic matter
235 would likely disrupt the pelagic-benthic coupling and result in lower zoobenthos biomass. Sprat and some
236 coastal fish species could be favoured by a rise in temperature. Regime shifts and cascading effects have
237 already been observed in both pelagic and benthic systems as a result of climate change.
238

239 7. **Coupled regional Earth system modeling in the Baltic Sea region**, Baltic Earth Special Topic, with
240 relevance to Baltic Earth Grand Challenge 5 (Gröger et al., 2021): Recent progress in the development of
241 coupled climate models for the Baltic Sea region is assessed. Feedback mechanisms are important to
242 simulate the response of the Earth system to external forcing such as greenhouse gas and aerosol
243 emissions. In this review article, the couplings between (1) atmosphere, sea ice and ocean, (2) atmosphere
244 and land surface including dynamic vegetation, (3) atmosphere, ocean and waves and (4) atmosphere and
245 hydrological components to close the water cycle are discussed. Adding surface waves to coupled
246 atmosphere-ocean system models is becoming more important with increasing resolution, in particular
247 when detailed information is required, for instance, for offshore wind energy applications in the coastal
248 zone. Furthermore, the wave information is essential for the calculation of ocean mixing and
249 resuspension. While long-term climate simulations using coupled atmosphere, sea ice and ocean models
250 or coupled atmosphere and dynamic vegetation models have successfully been performed and their added
251 value demonstrated, the impact of aerosols on the climate of the Baltic Sea region has not been considered.
252 Coupling hydrology models to close the hydrological cycle is also still problematic, as the precipitation
253 accuracy provided by the atmospheric models is, in most cases, insufficient to realistically simulate river
254 discharge into the Baltic Sea without bias adjustments.
255

256 8. **Atmospheric regional climate projections for the Baltic Sea region until 2100**, Baltic Earth Special
257 Topic (Christensen et al., 2022): Current climate projections based on regional climate atmosphere-only
258 models of the EURO-CORDEX project with a horizontal resolution of 12.5 km under the scenarios
259 RCP2.6, 4.5 and 8.5 are presented. As the number of simulations (124) is relatively large compared to
260 previous assessments, the uncertainties can be better estimated than before. These projections indicate
261 strong warming, especially in the north in winter, where warming approaches twice the average global
262 warming. Precipitation is projected to increase throughout the Baltic Sea region, except in the southern
263 half in summer, where the results are inconclusive. Extreme precipitation, here the 10-year return value,

264 is projected to increase systematically throughout the study area, especially in summer. The large
265 ensemble of simulations does not indicate a significant change in wind speed. Surface solar radiation is
266 projected to remain unchanged in summer, but to decrease slightly in winter, due to increased cloud cover
267 and possibly less snow in the future. Snow cover is projected to decrease dramatically, especially in the
268 south of the Baltic Sea catchment. The comparison between the uncoupled model simulations of the
269 EURO-CORDEX project and a small ensemble of scenario simulations performed with a coupled
270 atmosphere-sea-ice-ocean model driven by a subset of global climate models indicates stronger warming
271 in the coupled model during winter, mainly in areas that are seasonally affected by sea ice today. In
272 summer, the coupled model shows weaker warming compared to the uncoupled models.

273
274 9. **Oceanographic regional climate projections for the Baltic Sea until 2100**, Baltic Earth Special Topic
275 (Meier et al., 2022a): New projections of the future Baltic Sea climate with a coupled physical-
276 biogeochemical ocean model were compared with previous projections. The differences are mainly due
277 to different scenario assumptions and model setups. For example, the impact of future global sea level
278 rise on salinity was previously neglected, but taken into account in the latest projections. Although the
279 number of projections for the Baltic Sea is still small compared to regional atmospheric projections such
280 as the EURO-CORDEX model ensemble, a relatively large ensemble of 48 scenario simulations allowed
281 the assessment of uncertainties related to greenhouse gas emissions, global climate model differences,
282 global sea level rise, nutrient inputs and natural variability. In the future climate, higher water
283 temperatures, a shallower mixed layer with a sharper thermocline in summer, lower sea ice cover and
284 stronger mixing in the northern Baltic Sea in winter compared to the current climate could be expected.
285 The assessment of marine heat wave changes is new. Both the frequency and duration of marine heat
286 waves are projected to increase significantly, especially in the coastal zone of the southern Baltic Sea.
287 Due to uncertainties in the projections regarding regional winds, precipitation and global sea level rise,
288 no robust and statistically significant changes in salinity could be identified. The impacts of a changing
289 climate on the biogeochemical cycle are projected to be significant, but still less than the plausible
290 changes in nutrient inputs. Implementation of the proposed Baltic Sea Action Plan, a basin-wide nutrient
291 input reduction plan, would lead to a significant improvement in the ecological status of the Baltic Sea,
292 including a reduction in the size of the hypoxic area also in a future climate.

293
294 10. **Climate change in the Baltic Sea region: A summary**, Baltic Earth Special Topic (Meier et al., 2022b):
295 In this comprehensive study, the recent knowledge on past (paleo-), present (historical) and projected
296 future (< 2100) climate change in the Baltic Sea region, based upon all BEARs and >800 scientific
297 articles, is summarised. It focuses on the atmosphere, the land surface, the cryosphere, the ocean and its
298 sediments, and the terrestrial and marine biospheres. 33 parameters characterising the state of these
299 components of the Earth system were analysed (Fig. 3, Table 2). The anthroposphere is not part of this
300 assessment by Meier et al. (2022b) but instead is discussed in detail by Reckermann et al. (2022). The
301 main findings concerning changes of the 33 selected state parameters attributed to climate change are
302 summarised in Figure 3. The prevailing causal relationships of climate change with sufficiently high

303 confidence suggest a clear impact of global greenhouse gas emissions on regional heat cycles including
304 all parameters of the cryosphere. However, changes caused by global warming of the water, momentum
305 and carbon cycles are less clear because of either the large natural variability at regional scales or the
306 impact of other drivers than global warming. For further details, the reader is referred to Meier et al.
307 (2022b). Overall, it was concluded that the results from the previous BACC assessments mainly are still
308 valid. However, new long-term, homogenous observational records, such as those for Scandinavian
309 glacier inventories, sea-level-driven saltwater inflows (MBIs), or phytoplankton species distributions, and
310 new scenario simulations with improved models, such as those for glaciers, lake ice, or marine food webs,
311 have become available, resulting in a revised understanding of observed changes. Compared to previous
312 assessments, observed changes in air temperature, sea ice, snow cover, and sea level were shown to have
313 accelerated. However, natural variability is large, challenging our ability to detect observed and projected
314 changes in climate of the Baltic Sea region. As the ensembles of scenario simulations both for the
315 atmosphere and the ocean became larger, uncertainties can now be better estimated, although coordinated
316 scenario simulations for the Baltic Sea based on ensembles of different regional ocean models are still
317 missing. Furthermore, with the help of coupled models, feedbacks between several components of the
318 Earth system have been studied, and multiple driver studies were performed, e.g., projections of the
319 marine food web that include fisheries, eutrophication and climate change. Intensive research on the land–
320 sea interface, focusing on the coastal filter, has been performed, and nutrient retention in the coastal zone
321 was estimated for the first time. However, a model for the entire Baltic Sea coastal zone is still missing,
322 and the effect of climate change on the coastal filter capacity is still unknown. More research on changing
323 extremes was performed, acknowledging that the impact of changing extremes may be more important
324 than that of changing means (see also Rutgersson et al., 2022). However, many observational records are
325 either too short or too heterogeneous for statistical studies of extremes due to data gaps.

326 **4 Discussion**

327 One of the main objectives of the BEAR project was to identify knowledge gaps in the Earth system science of
328 the Baltic Sea region so that these can be further addressed in future research. For specific knowledge gaps that
329 have been identified during the project, the reader is referred to the individual assessment reports. However, as an
330 overarching result, three new research topics are identified:

- 331
- 332 1) **Small-scale processes and their impact on large-scale climate dynamics and biogeochemical cycles.**
333 The number of observations in the sea is smaller than those on land. This is also true for the Baltic Sea,
334 although the international long-term monitoring programme in the Baltic Sea started more than a century
335 ago, with measurements of temperature, salinity and oxygen concentration in the central parts of the
336 different sub-basins. Nowadays, monitoring data are available from all sub-basins with a resolution of up
337 to one month. Recently, many new observational systems for temporally and spatially high-resolution
338 data have been developed or are under development, including remotely operated vehicles (ROVs) and
339 autonomous underwater vehicles (AOVs) as well as remote sensing data. Examples of such systems

340 operating in the Baltic Sea are continuously profiling moorings, ARGO floats, Gliders, ScanFish, and
341 echo sounders. In addition to traditional physical parameters, measurements of turbulence,
342 biogeochemical and biodiversity (e.g. environmental DNA) parameters are now available. Another area
343 of research that is developing rapidly is numerical modelling of the Earth system, also on a regional scale,
344 e.g. eddy- and submesoscale resolving multi-year simulations for the Baltic Sea. Similar arguments apply
345 to the atmosphere, e.g. cloud-resolving simulations to cope with heavy precipitation events. A novel
346 research topic for Baltic Earth would therefore be a better understanding of the dynamics of small-scale
347 atmospheric and oceanic processes that are not yet resolved in state-of-the-art numerical models or
348 conventional observations, and their role in the large-scale circulation on short and long time scales. Such
349 research activities would help to fill some of the gaps in knowledge that have been raised by Lehmann et
350 al. (2022), Kuliński et al. (2022), Rutgersson et al. (2022), Weisse et al. (2021), Viitasalo and Bonsdorff
351 (2022), and Gröger et al. (2021). Furthermore, a realistic consideration of small-scale processes would
352 improve the projections for the atmosphere (Christensen et al., 2022) and the ocean (Meier et al., 2022a).

353
354 2) **Attribution of regional climate variability and change to anthropogenic radiative forcing and other**
355 **drivers.** In order to unambiguously disentangle the impacts of anthropogenic climate change and other
356 human influences from the natural climate variability of the regional Earth system, more knowledge about
357 internal variations and feedback mechanisms is needed. For example, climate models have recently
358 shown that multi-decadal variability emanating from the North Atlantic and the Arctic significantly
359 controls the climate of the Baltic Sea region by means of teleconnection patterns (Lehmann et al., 2022;
360 Meier et al., 2022a; 2022b). For example, observations of precipitation and wind in the Baltic Sea region,
361 total river discharge from the catchment, individual river flows, water temperature, sea level, MBIs and
362 salinity in the Baltic Sea show a pronounced multidecadal variability with a quasi-periodicity of about 30
363 years (Meier et al., 2022b). It is assumed that the Atlantic Multidecadal Variability (AMV) and, as part
364 of it, the variations of the North Atlantic overturning circulation are the source of these variations,
365 although the exact mechanisms, cause and effect chains and feedback processes are still unknown.
366 Knowledge about the teleconnectivity of the Baltic Sea region with the North Atlantic and the Arctic is
367 essential for the development of climate prediction models.

368
369 3) **Development of integrated Earth system models accounting for anthropogenic changes in the Baltic**
370 **Sea region.** The BEAR study by Reckermann et al. (2022) on human influences and their interactions in
371 the Baltic Sea region is part of the relatively new Baltic Earth Grand Challenge 6 on the multiple drivers
372 of Earth system change in the Baltic Sea region and represents an important step towards an integrated
373 understanding of the Earth system that encompasses all traditionally considered climate compartments
374 such as atmosphere, cryosphere, hydrosphere, lithosphere (including the pedosphere), biosphere (marine
375 and terrestrial) and the anthroposphere. Such a holistic view is urgently needed, as in many cases, several
376 reasons are responsible for the observed changes in the Earth system and attributing them to only one
377 factor, e.g. climate change would be an inadmissible simplification. One example is the oxygen depletion
378 and the large hypoxic area in the Baltic Sea caused by anthropogenic nutrient inputs from land and

379 exacerbated by rising water temperatures (Kuliński et al., 2022). Of course, the factors discussed by
380 Reckermann et al. (2022) cannot exhaustively consider the entire Earth system and all interactions, and
381 the selection of factors is biased towards ocean-related parameters and activities. Moreover, the analysis
382 is based on an extensive literature review by experts who reflect their subjective interpretations of the
383 results. Nevertheless, this is the first time such an assessment has been conducted, which is a major step
384 forward. To continue and deepen this research, the factors discussed by Reckermann et al. (2022) could
385 be subdivided either by human activities (e.g. food production, energy production, transport, tourism,
386 healthcare) or by environmental and climate state variables of the Earth system (e.g. hypoxia,
387 acidification) (Table 3). Such a breakdown of parameters would allow the development of an integrated
388 Earth system model that includes the anthroposphere at the regional scale. This type of research is timely,
389 and such efforts are already underway (e.g. Korpinen et al., 2019; references in Reckermann et al., 2022).

390

391 The fact sheet on climate change in the Baltic Sea (CCFS, 2021) was positively received by various stakeholders
392 and decision-makers. Although uncertainties regarding observed and projected future climate change and the other
393 drivers remain high, our experience engaging with stakeholders confirms that scientific uncertainties are taken into
394 account in different ways in management and decision-making. This is an important reason for investing in the
395 above key issues. They have the potential to reduce uncertainties that currently hamper decision-making in the
396 region.

397 **5 Concluding remarks**

398 We conclude that 1) the BEARs have been useful to identify research progress and knowledge gaps and to initiate
399 new research foci as, for examples, suggested in the discussions; 2) regional assessments, such as the BEARs,
400 complement the IPCC climate change assessments by adding a greater depth and scope of regional information
401 about the specific situation of the Baltic Sea region; and 3) the BEARs provided useful information for the Expert
402 Network on Climate Change, that produced the Baltic Earth – HELCOM climate change fact sheet for
403 stakeholders. Since the information summarised by the BEARs are used extensively in science and management,
404 it is recommended that a new update of the reports will be conducted in about seven years.


405 **Author contributions**

406 H.E.M.M. wrote the first draft of the editorial. All co-authors, which acted as guest editors of the special issue in
407 *Earth System Dynamics*, contributed with important comments and editing of the manuscript, read and approved
408 the submitted manuscript version.

409 **Acknowledgements**

410 During 2019-2022, the Baltic Earth Assessment Reports were produced under the umbrella of the Baltic
411 Earth programme (Earth System Science for the Baltic Sea region, see <http://baltic.earth>, last access: 2
412 February 2023). 109 co-authors from 14 countries contributed to 10 articles in the international scientific


413 journal *Earth System Dynamics* and 2822 different references have been assessed. We thank the reviewers
414 of all 10 articles of the special issue for their constructive comments that helped to improve the review
415 articles. In particular, we thank Dr. Jouni Räisänen, Dr. Donald Boesch and Dr. Andris Andrusaitis for their
416 advice and many excellent comments on individual articles and this overview article.
417



419

420 **Figure 1:** The Baltic Sea and its catchment area with climatological mean sea surface salinity (in g kg^{-1}) and river
 421 discharge (in mm year^{-1}). (Source: Meier et al., 2014; their Fig. 1 distributed under the terms of the Creative
 422 Commons CC-BY 4.0 License, <http://creativecommons.org/licenses/by/4.0/>, last access: 4 February 2023)


423



425

426 **Figure 2:** Annual mean sea surface temperature anomalies relative to the reference period 2002-2018 from de-
 427 seasonalised measurements at the Arkona Deep monitoring station and the MARNET stations Darss Sill and
 428 Arkona Basin in the period 1980-2018. (Source: Meier et al., 2022b; their Fig. 20 distributed under the terms of
 429 the Creative Commons CC-BY 4.0 License, <http://creativecommons.org/licenses/by/4.0/>, last access: 4 February
 430 2023)

431



433

434 **Figure 3:** Synthesis of knowledge on present and future climate change. Shown are anthropogenic climate changes
 435 in 33 Earth system variables (bubbles) of the atmosphere (yellow), land surface (brown), terrestrial biosphere (dark
 436 green), cryosphere (grey), ocean and sediments (blue) and marine biosphere (light green). The abbreviation NIS
 437 stands for non-indigenous species. The sign of a change (plus/minus) is shown together with the confidence level
 438 indicated by the number of signs, i.e. one to three signs correspond to a low, medium and high confidence level as
 439 a result of the literature assessment reflecting consensus and evidence according to IPCC definitions. The colours
 440 of the signs indicate the direction of past (black) and future (red) changes according to Meier et al. (2022b).
 441 Uncertain changes are not shown. The external anthropogenic drivers of the Earth system studied are shown as red
 442 squares, i.e. greenhouse gases, especially CO₂, and aerosol emissions. The predominant climate change linkages
 443 with sufficiently high confidence are shown by arrows (black: thermal cycle, blue: hydrological cycle, orange:
 444 momentum cycle including sea level change, pink: carbon cycle). Projections of carbon sequestration of Arctic
 445 terrestrial ecosystems for the 21st century show first an increased uptake and later a carbon source, marked by “1.
 446 + 2. –“. Future changes in mean sea level are dominated by the thermal expansion of the global ocean and the
 447 melting of ice sheets outside the Baltic Sea region. (Source: Meier et al., 2022b; their Fig. 35 distributed under the
 448 terms of the Creative Commons CC-BY 4.0 License, <http://creativecommons.org/licenses/by/4.0/>, last access: 4
 449 February 2023)

450

451 **Tables**

452 **Table 1:** The matrix of factors studied by Reckermann et al. (2022). + = evidence for a connection; - = no evidence
 453 for a connection; ? = no evidence, but connection plausible (according to the author's assessment). The table is
 454 read from left to right, i.e. if you go to the right in the first row "climate change", you see the factors on which
 455 climate change has an effect (or not), etc. (Source: Reckermann et al., 2022; their Table 2a distributed under the
 456 terms of the Creative Commons CC-BY 4.0 License, <http://creativecommons.org/licenses/by/4.0/>, last access: 4
 457 February 2023)

	Climate change	Coastal processes	Hypoxia	Acidification	Subm. Groundw. Disch.	Marine ecosystems	Non-inig. species	Land cover and use	Agriculture Nutr. loads	Aquaculture	Fisheries	River regulations	Offshore wind farms	Shipping	Chem. Contamin.	Dumped military	Marine litter	Tourism	Coastal management
impact by ↓/on→																			
Climate change		+	+	+	?	+	?	+	+	+	+	+	+	+	+	?	?	+	+
Coastal processes	-		?	?	+	?	-	+	+	?	?	+	+	+	?	?	+	-	+
Hypoxia	-	-		+	-	+	-	-	+	?	+	-	-	-	+	+	-	-	-
Acidification	-	-	-			?	-	-	-	?	?	-	-	-	?	-	-	-	-
Subm. Groundw. Disch.	-	-	?	?		?	-	-	+	-	-	-	-	-	+	-	-	-	-
Marine ecosystems	-	-	+	+	-		+	-	-	-	+	-	-	-	-	-	-	+	-
Non-inigenous species	-	-	-	-	-	+				-	+	-	-	+	+	-	-	-	?
Land cover and use	+	-	+	+	+	?	-		+		+	+	?	-	+	-	-	+	+
Agriculture/Nutrient loads	+	-	+	+	+	+	-	+		+	+	+	-	-	+	-	-	-	?
Aquaculture	-	-	+	-	-	+	+	+	+		?	-	+	-	?	-	?	?	+
Fisheries	-	-	?	-	-	+	?	-	?	?		-	+	?	-	-	+	?	+
River regulations	-	+	?	+	?	+	-	?	?	?	+		-	-	?	-	?	-	+
Offshore wind farms	+	+	-	-	-	+	-	?	?	?	+	+		+	?	?	?	+	+
Shipping	+	+	-	+	-	+	+	-	+	?	+	-	?		+	-	+	+	+
Chemical contaminants	-	-	-	-	-	+	-	-	+	+	+	-	-	-		-	-	-	-
Dumped military material	-	-	-	-	-	?	-	-	-	-	+	+	+	-	+		-	-	?
Marine litter	-	-	-	-	-	?	-	-	-	?	+	-	-	-	?	-		+	?
Tourism	+	-	-	-	-	?	-	+	+	-	-	-	+	+	-	-	+		+
Coastal management	-	+	-	-	?	?	?	?	?	?	+	?	+	+	-	+	?	+	

458
 459

460 **Table 2:** Variables of the Meier et al. (2022b) assessment and further references to the BEARs (1: Lehmann et al.,
 461 2022; 2: Kuliński et al., 2022; 3: Rutgersson et al, 2022; 4: Weisse et al., 2021; 5: Reckermann et al, 2022; 6:
 462 Gröger et al, 2021; 7: Christensen et al, 2022; 8: Meier et al, 2022a; 9: Viitasalo and Bonsdorff, 2022). The third
 463 column lists the subsection in the study by Meier et al. (2022b) that contains further information. (Source: Meier
 464 et al., 2022b; their Table 2 distributed under the terms of the Creative Commons CC-BY 4.0 License,
 465 <http://creativecommons.org/licenses/by/4.0/>, last access: 4 February 2023)

Number	Variable	Past and present climates		Future climate	
Atmosphere					
1	Large-scale atmospheric circulation	3.2.1.1	3	3.3.1.1	3, 7
2	Air temperature	3.1.2, 3.1.3, 3.1.4		3.3.1.2	7
	Warm spell	3.2.1.2	3		3
	Cold spell		3		3
3	Solar radiation and cloudiness	3.2.1.3		3.3.1.3	7
4	Precipitation	3.1.2, 3.1.3, 3.1.4		3.3.1.4	7
	Heavy precipitation	3.2.1.4	3		3
	Drought		3		3
5	Wind	3.2.1.5		3.3.1.5	7
	Storm		3		3
6	Air pollution, air quality and atmospheric deposition	3.2.1.6		3.3.1.6	
Land					
7	River discharge	3.2.2.1		3.3.2.1	8
	High flow		3		3
8	Land nutrient inputs	3.2.2.2		3.3.2.2	8
Terrestrial biosphere					
9	Land cover (forest, crops, grassland, peatland, mires)	3.2.3	6	3.3.3	
10	Carbon sequestration			3.3.3	
Cryosphere					

11	Snow Sea-effect snowfall	3.2.4.1	3	3.3.4.1	7 3
12	Glaciers	3.2.4.2		3.3.4.2	
13	Permafrost	3.2.4.3		3.3.4.3	
14	Sea ice Extreme mild winter Severe winter Ice ridging	3.2.4.4	3 3 3	3.3.4.4	8 3 3 3
15	Lake ice	3.2.4.5		3.3.4.5	
Ocean and marine sediments					
16	Water temperature Marine heat wave	3.2.5.1	3	3.3.5.1	8 3
17	Salinity and saltwater inflows	3.2.5.2	1	3.3.5.2	8
18	Stratification and overturning circulation	3.2.5.3	1	3.3.5.3	8
19	Sea level Sea level extreme	3.2.5.4	4 3	3.3.5.4	8 3
20	Waves Extreme waves	3.2.5.5	4 3	3.3.5.5	3
21	Sedimentation and coastal erosion	3.2.5.6	4	3.3.5.6	
22	Oxygen and nutrients	3.1.4 3.2.5.7.1	2	3.3.5.7.1	8
23	Marine CO ₂ system	3.2.5.7.2	2	3.3.5.7.2	
Marine biosphere					
24	Pelagic habitats: Microbial communities	3.2.6.1.1	2, 9	3.3.6.1.1	9
25	Pelagic habitats: Phytoplankton and cyanobacteria	3.2.6.1.2	2, 3, 9	3.3.6.1.2	3, 9
26	Pelagic habitats: Zooplankton	3.2.6.1.3	9	3.3.6.1.3	9
27	Benthic habitats: Macroalgae and vascular plants	3.2.6.2.1	9	3.3.6.2.1	9

28	Benthic habitats: Zoobenthos	3.2.6.2.2	9	3.3.6.2.2	9
29	Non-indigenous species	3.2.6.3	9	3.3.6.3	9
30	Fish	3.2.6.4	9	3.3.6.4	9
31	Marine mammals	3.2.6.5	9	3.3.6.5	9
32	Waterbirds	3.2.6.6	9	3.3.6.6	9
33	Marine food web	3.2.6.7	9	3.3.6.7	9

466

467 **Table 3:** Factors discussed by Reckermann et al. (2022) sorted by related economic sectors or state variables of
 468 the Earth system.

Human activities		
Economic sectors	Factors	Comments
Primary (natural) sector (e.g. food production)	Fisheries	
	Agriculture	
	Marine and coastal ecosystem services	Factor belongs to several sectors
	Blue carbon storage capacity	Mitigation of greenhouse gases
Secondary (industrial) sector (e.g. energy production)	River regulation	
	Offshore wind farms	
	Greenhouse gas and aerosol emissions	Emission are largest from industries
	Dumped warfare agents	Factor is an industrial product
Tertiary (service) sector (e.g. transportation, tourism, healthcare)	Shipping	
	Chemical contamination	Contamination is a result of several sectors
	Marine noise	Marine noise is a result of several sectors
	Marine litter and microplastics	Emission mainly by offshore platforms, shipping, lost containers, fisheries, aquaculture, agriculture, municipal waste and tourism
	Tourism	
	Coastal protection and management	Also relevant for the other sectors
Quaternary (information) sector (e.g. information technology; media; research and development)	-	
Earth system		
Environmental state variables	Coastal processes	
	Hypoxia	
	Submarine groundwater discharge	
	Marine ecosystems	
	Land use and land cover	
	Non-indigenous species	

	Indirect parameters such as carbon and nutrient cycles, biota and ecosystems	
Climate state variables	Climate change, acidification, direct parameters of the climate system	Superordinated concept (large-scale)
	Direct parameters of the climate system	
	Acidification	

469

470 **References**

- 471 Baltic Earth Science Plan, 2017. <https://baltic.earth/grandchallenges>, last access: 4 February 2023
- 472
- 473 CCFS, 2021. Climate Change in the Baltic Sea. 2021 Fact Sheet. Baltic Sea Environment Proceedings n°180.
- 474 HELCOM/Baltic Earth, <http://doi.io-warnemuende.de/10.12754/misc-2022-0001>, 2021.
- 475
- 476 Christensen, O. B., Kjellström, E., Dieterich, C., Gröger, M., and Meier, H. E. M.: Atmospheric regional climate
- 477 projections for the Baltic Sea Region until 2100, *Earth Syst. Dynam.*, 13, 133–157, [https://doi.org/10.5194/esd-](https://doi.org/10.5194/esd-13-133-2022)
- 478 [13-133-2022](https://doi.org/10.5194/esd-13-133-2022), 2022.
- 479
- 480 Gröger, M., Dieterich, C., Haapala, J., Ho-Hagemann, H. T. M., Hagemann, S., Jakacki, J., May, W. Meier, H. E.
- 481 M., Miller, P. A. Rutgersson, A. and Wu, L.: Coupled regional Earth system modeling in the Baltic Sea region,
- 482 *Earth Syst. Dynam.*, 12, 939–973, <https://doi.org/10.5194/esd-12-939-2021>, 2021.
- 483
- 484 Klimawandel in der Ostsee, 2021 Faktenblatt - Baltic Sea Climate Change in the Baltic Sea, 2021 Fact Sheet (in
- 485 German). <http://doi.io-warnemuende.de/10.12754/misc-2022-0003>, 2022.
- 486
- 487 Korpinen, S., Klančnik, K., Peterlin, M., Nurmi, M., Laamanen, L., Zupančič, G., Popit, A., Murray, C., Harvey,
- 488 T., Andersen, J.H., Zenetos, A., Stein, U., Tunesi, L., Abhold, K., Piet, G., Kallenbach, E., Agnesi, S., Bolman,
- 489 B., Vaughan, D., Reker, J. & Royo Gelabert, E.: Multiple pressures and their combined effects in Europe’s seas,
- 490 ETC/ICM Technical Report 4/2019: European Topic Centre on Inland, Coastal and Marine waters, 164 pp., 2019.
- 491
- 492 Kuliński, K., Rehder, G., Asmala, E., Bartosova, A., Carstensen, J., Gustafsson, B., Hall, P. O. J., Humborg, C.,
- 493 Jilbert, T., Jürgens, K., Meier, H. E. M., Müller-Karulis, B., Naumann, M., Olesen, J. E., Savchuk, O., Schramm,
- 494 A., Slomp, C. P., Sofiev, M., Sobek, A., Szymczycha, B., and Undeman, E.: Biogeochemical functioning of the
- 495 Baltic Sea. *Earth Syst. Dynam.*, 13, 633–685, <https://doi.org/10.5194/esd-13-633-2022>, 2022.
- 496
- 497 Lehmann, A., Myrberg, K., Post, P., Chubarenko, I., Dailidienė, I., Hinrichsen, H.-H., Hüseyin, K., Liblik, T., Meier,
- 498 H. E. M., Lips, U., and Bukanova, T.: Salinity dynamics of the Baltic Sea, *Earth Syst. Dynam.*, 13, 373–392,
- 499 <https://doi.org/10.5194/esd-13-373-2022>, 2022.
- 500
- 501 Meier, H.E.M., A. Rutgersson, and M. Reckermann: An Earth System Science Program for the Baltic Sea Region.
- 502 *Eos*, 95, 109–110. <https://doi.org/10.1002/2014EO130001>, 2014.
- 503
- 504 Meier, H. E. M., Dieterich, C., Gröger, M., Dutheil, C., Börgel, F., Safonova, K., Christensen, O. B., and
- 505 Kjellström, E.: Oceanographic regional climate projections for the Baltic Sea until 2100, *Earth Syst. Dynam.*, 13,
- 506 159–199, <https://doi.org/10.5194/esd-13-159-2022>, 2022a.
- 507

508 Meier, H. E., Kniebusch, M., Dieterich, C., Gröger, M., Zorita, E., Elmgren, R., ... and Zhang, W.: Climate change
509 in the Baltic Sea region: a summary. *Earth Syst. Dynam.*, 13, 457-593, <https://doi.org/10.5194/esd-13-457-2022>,
510 2022b.

511

512 Reckermann, M., Omstedt, A., Soomere, T., Aigars, J., Akhtar, N., Beldowski, J., Brauwer, C. P.-S. d., Cronin,
513 T., Czub, M., Eero, M., Hyttiäinen, K., Jalkanen, J.-P., Kiessling, A., Kjellström, E., Larsén, X. G., McCrackin,
514 M., Meier, H. E. M., Oberbeckmann, S., Parnell, K., Poska, A., Saarinen, J., Szymczycha, B., Undeman, E.,
515 Viitasalo, M., Wörman, A., and Zorita, E.: Human impacts and their interactions in the Baltic Sea region. *Earth*
516 *Syst. Dynam.*, 13, 1–80, <https://doi.org/10.5194/esd-13-1-2022>, 2022.

517

518 Rutgersson, A., Kjellström, E., Haapala, J., Stendel, M., Danilovich, I., Drews, M., Jylhä, K., Kujala, P., Larsén,
519 X. G., Halsnæs, K., Lehtonen, I., Luomaranta, A., Nilsson, E., Olsson, T., Särkkä, J., Tuomi, L., and Wasmund,
520 N.: Natural Hazards and Extreme Events in the Baltic Sea region, *Earth Syst. Dynam.*, 13, 251–301,
521 <https://doi.org/10.5194/esd-13-251-2022>, 2022.

522

523 The BACC Author Team, 2008. Assessment of Climate Change for the Baltic Sea Basin. Regional Climate Studies
524 series. Heidelberg, Springer-Verlag Berlin. ISBN 978-3-540-72785-9. 473 pp. [https://doi.org/10.1007/978-3-540-](https://doi.org/10.1007/978-3-540-72786-6)
525 [72786-6](https://doi.org/10.1007/978-3-540-72786-6).

526

527 The BACC II Author Team, 2015. Second Assessment of Climate Change for the Baltic Sea Basin. Regional
528 Climate Studies series. Cham, Springer Cham. ISBN 978-3-319-16005-4. 501 pp. [https://doi.org/10.1007/978-3-](https://doi.org/10.1007/978-3-319-16006-1)
529 [319-16006-1](https://doi.org/10.1007/978-3-319-16006-1).

530

531 Viitasalo, M., and Bonsdorff, B.: Global climate change and the Baltic Sea ecosystem: direct and indirect effects
532 on species, communities and ecosystem functioning. *Earth Syst. Dynam.*, 13, 711–747,
533 <https://doi.org/10.5194/esd-13-711-2022>, 2022.

534

535 Weisse, R., Dailidienė, I., Hünicke, B., Kahma, K., Madsen, K., Omstedt, A., Parnell, K., Schöne, T., Soomere,
536 T., Zhang, W., and Zorita, E.: Sea Level Dynamics and Coastal Erosion in the Baltic Sea Region. *Earth Syst.*
537 *Dynam.*, 12, 871–898, <https://doi.org/10.5194/esd-12-871-2021>, 2021.