

Effects of idealised land cover and land management changes on the atmospheric water cycle

Steven J. De Hertog¹, Carmen E. Lopez-Fabara¹, Ruud van der Ent², Jessica Keune³, Diego G. Miralles³, Raphael Portmann⁴, Sebastian Schemm⁵, Felix Havermann⁶, Suqi Guo⁶, Fei Luo^{7,8,9}, Iris Manola⁷, Quentin Lejeune¹⁰, Julia Pongratz^{6,11}, Carl-Friedrich Schleussner¹⁰, Sonia I. Seneviratne⁵, and Wim Thiery¹

¹Vrije Universiteit Brussel, Department of Water and Climate, Brussels, Belgium

²Delft University of Technology, Department of water management, Delft, Netherlands

³Ghent University, Hydro-Climate Extremes Lab, Ghent, Belgium

⁴Agroscope Reckenholz, Division of Agroecology and Environment, Zurich, Switzerland

⁵ETH Zurich, Institute for Atmospheric and Climate Science, Zurich, Switzerland.

⁶Ludwig-Maximilians-University Munich, Department of Geography, Munich, Germany

⁷Vrije Universiteit Amsterdam, Institute for Environmental studies, Amsterdam, Netherlands

⁸Royal Netherlands Meteorological Institute (KNMI), De Bilt, Netherlands.

⁹Centre for Climate Research Singapore (CCRS), Singapore.

¹⁰Climate Analytics, Berlin, Germany

¹¹Max Planck Institute for Meteorology, Hamburg, Germany

Correspondence: Steven De Hertog (steven.de.hertog@vub.be)

Abstract. Land cover and land management changes (~~LCLMCs~~LCLMC) play an important role in achieving low-end warming scenarios through land-based mitigation. However, their effects on moisture fluxes and recycling remain uncertain although they have important implications for the future viability of such strategies. Here, we analyse the impact of idealised LCLMC scenarios on atmospheric moisture transport in three different ESMs: the Community Earth System Model (CESM), the Max
5 Planck Institute Earth System Model (MPI-ESM) and the European Consortium Earth System Model (EC-EARTH). The LCLMC scenarios comprise of a full cropland world, a fully afforested world, and a cropland world with unlimited irrigation expansion. The effects of these ~~LCLMCs~~LCLMC in the different ESMs are analysed for precipitation, evaporation and vertically integrated moisture flux convergence to understand the LCLMC-induced changes in the atmospheric moisture cycle. Then, a moisture tracking algorithm is applied to assess the effects of ~~LCLMCs~~LCLMC on moisture recycling at the
10 local (grid cell level) and the global scale (continental moisture recycling). ~~Our results indicate that LCLMCs are generally inducing consistent effects on moisture fluxes over land in all ESMs. Cropland expansion causes~~ By applying a moisture tracking algorithm on fully coupled ESM simulations we are able to quantify the complete effects of LCLMC on moisture recycling (including circulation changes) which are generally not considered in moisture recycling studies. Our results indicate that cropland expansion is generally causing a drying and reduced local moisture recycling ~~in all ESMs~~, while afforestation and
15 irrigation expansion generally cause wetting and increased local moisture recycling. However, the strength of this ~~influence varies in time and space and across the~~ effect varies across ESMs and shows a ~~strong~~large dependency on the dominant driver: Some ESMs show a dominance of large scale atmospheric circulation changes while other ESMs show a dominance of local

to regional changes in the atmospheric water cycle only within the vicinity of the LCLMC. Overall, these results corroborate that ~~LCLMCs can induce large~~ LCLMC can induce substantial effects on the atmospheric water cycle and moisture recycling, ~~but both through local effects and changes in atmospheric circulation. However,~~ more research is needed to constrain the uncertainty of these effects within ESMs ~~and better evaluate~~ to better inform future land-based mitigation strategies.

1 Introduction

Currently, about three quarters ($\sim 100 * 10^6$ km²) of the ice-free land surface has undergone some kind of anthropogenic land cover or land management change (LCLMC) (Luyssaert et al., 2014; Mbow et al., 2017). All these modifications are important drivers of climate change as they alter the carbon cycle (biogeochemical effects) and affect surface properties, which impact the energy and water balance (biogeophysical effects) (Pongratz et al., 2010; Bonan, 2008; Pongratz et al., 2021) and feed back on the local to global climate (Winckler et al., 2019; Boysen et al., 2020; Portmann et al., 2022; De Hertog et al., 2023). Therefore, future ~~LCLMCs~~ LCLMC are increasingly seen as a viable tool for land based mitigation and play a crucial role within low warming emission scenarios (Rogelj et al., 2018; Seneviratne et al., 2018). Hence, exploring and understanding the extent to which ~~LCLMCs~~ LCLMC influence climate has become key to develop effective mitigation and adaptation strategies (Lawrence et al., 2016).

From a biogeophysical perspective, ~~LCLMCs~~ LCLMC lead to changes in the albedo, aerodynamic conductance and the partitioning between the sensible and latent heat flux which has an impact on atmospheric temperature and moisture content (~~Bowen, 1926; Wang et al., 2009~~) (Bowen, 1926; Wang et al., 2009; Cui et al., 2022). For example, tropical deforestation is expected to further dry and warm the regional climate (~~Bonan, 2008; Akkermans et al., 2014; Spracklen et al., 2018~~) (Bonan, 2008; Akkermans et al., 2014; Spracklen et al., 2012). In contrast, irrigation expansion can cause a local to regional cooling and moistening of the atmosphere (Mahmood et al., 2014; Thiery et al., 2017, 2020; Hauser et al., 2019; Tuinenburg et al., 2014). Evaporation, being the link between the surface energy and the water balance (Shukla and Mintz, 1982), modulates the influence of ~~LCLMCs~~ LCLMC on atmospheric conditions (van der Ent et al., 2010; Spracklen et al., 2012). Tracking the origins of precipitation back to evaporation and determining the fraction of terrestrial precipitation that originates from land — here referred to as continental precipitation recycling (van der Ent et al., 2010) — can increase our understanding of the effects of future ~~LCLMCs~~ LCLMC on the climate. On the other hand, the fate of land evaporation can be determined and illustrates the reach of local ~~LCLMCs~~ LCLMC; the fraction of terrestrial evaporation precipitating over land is often referred to as continental evaporation recycling (van der Ent et al., 2010). Even though, it is well established that LCLMC can affect these moisture recycling strengths (Wang-Erlandsson et al., 2018; Benedict et al., 2020) — i.e., the degree to which terrestrial precipitation depends on land evaporation — this is rarely quantified within dedicated ESM studies. Most studies that quantify the effects of ~~LCLMCs~~ LCLMC on the atmospheric moisture cycle focus on the changes in moisture fluxes, but often cannot unravel the role of local and continental moisture recycling in these differences (Tuinenburg et al., 2020; Hoek van Dijke et al., 2022; Baudena et al., 2021; Wunderling et al., 2022; Staal et al., 2018). Those studies that do account for moisture recycling in assessing the effects of ~~LCLMC~~ generally apply reanalysis-based recycling ratios

(Tuinenburg et al., 2020; Hoek van Dijke et al., 2022; Baudena et al., 2021; Wunderling et al., 2022; Staal et al., 2018) future LCLMC (Hoek van Dijke et al., 2022; Baudena et al., 2021; Wunderling et al., 2022; Staal et al., 2018) generally apply reanalysis-based recycling ratios (such as those presented in Tuinenburg et al. (2020)) which do not include the two-way interactions of circulation changes and the water cycle second-order effect on moisture recycling caused by LCLMC-induced circulation changes (te Wierik et al., 202

55 . By analysing dedicated ESM simulations for LCLMC we are able to address these shortcomings and include the potential effects of atmospheric circulation changes on moisture recycling.

Idealized or extensive implementations of LCLMCs-LCLMC within ESM simulations are used to cope with weather-induced noise that dampens climatic responses (Winckler et al., 2017a; Boysen et al., 2020). Within such simulations, large-scale at-
60 mospheric circulation changes have been shown to occur as a consequence of LCLMC (Goessling and Reick, 2011; Boysen et al., 2020; Portmann et al., 2022; Devaraju et al., 2018; Laguë et al., 2019). However, most studies have only focused on one LCLMC type (e.g. Boysen et al., 2020; Laguë et al., 2019; Devaraju et al., 2018)) and only used a single ESM (e.g. Portmann et al., 2022; de Vrese et al., 2016). Further, these studies generally cannot distinguish explicitly between the influence of local processes (directly induced by the LCLMC) and non-local or remote processes (induced by LCLMC elsewhere, including
65 circulation and advection changes). The study of De Hertog et al. (2023) presented a first multi-model intercomparison using three different ESMs and four different LCLMC types in which a clear distinction between local and non-local biogeophysical effects was established through the checkerboard LCLMC implementation as developed by Winckler et al. (2017a). These simulations facilitated the comparison of the climate changes induced by different LCLMC types and to grasp the multi-model uncertainty.

70

Here, we assess the atmospheric water cycle responses to idealised LCLMC scenarios using global simulations of three different ESMs (De Hertog et al., 2023). The simulations comprise different idealised LCLMC scenarios — from afforestation, over cropland expansion to irrigation expansion — and have been implemented in a checkerboard pattern. The simulation setup and the moisture tracking algorithm and its derived metrics are described below (Section 2). We first analyse the ESM output
75 for changes in the atmospheric water cycle including evaporation, precipitation, and atmospheric moisture flux convergence (Section 3.1). Second, we analyse the moisture tracking algorithm output to assess the direct effects of LCLMCs-LCLMC on moisture recycling and unravel local and remote drivers of the analysed moisture flux changes. This is done on a local scale using the concept of 'length scales' of moisture recycling (Section 3.2), and on a continental scale using continental recycling ratios (Section 3.3). Finally we highlight the most important findings and implications of this research (Section 4. and 5.).

80

2 Methods

2.1 ESM simulations

The ESM simulations analysed here were conducted within the LAnd Management for CLimate Mitigation and Adaptation (LAMA CLIMA) project and are presented in detail in De Hertog et al. (2023). In this project, different sensitivity experiments were performed for three ESMs, i.e., the Community Earth System Model version 2.1.3 (CESM), the European Community Earth-System Model 3-Veg v3.3.3.1 (EC-EARTH), and the Max-Planck Institute Earth System Model 1.2 low resolution (MPI-ESM). We refer to De Hertog et al. (2023) for detailed model descriptions. The experiments comprise four simulations of LCLMC scenarios. One scenario is the control case (hereafter referred to as CTL), which is conducted with a constant land cover corresponding to the year 2015. The three remaining scenarios represent an extreme case of single specific LCLMC, namely afforestation (FRST), cropland expansion (CROP), and irrigation expansion (IRR). Here, irrigation is applied on top of the cropland expansion. Thus, while the impact of afforestation and cropland expansion is always evaluated with respect to the control simulation (i.e., differences are calculated as FRST-CTL and CROP-CTL), the impact of irrigation is evaluated with respect to the cropland expansion simulation (IRR-CROP). All simulations cover a period of 160 years under a present-day climate forcing (corresponding to the year 2015).

95

The LCLMC scenarios are generated from the CTL scenario land cover by inducing the LCLMC in a checkerboard-like pattern as presented in Winckler et al. (2017a). The resulting LCLMC is shown in Figure 1. This implies that the different ~~LCLMCs~~ LCLMC are implemented in every other pixel (i.e. only 50% of hospitable land grid cells have undergone LCLMC), while all other forcings (i.e. greenhouse gas, stratospheric aerosols,...) remain identical to the initial CTL scenario configurations. Even though, a structured approach was taken to implement the ~~LCLMCs~~ LCLMC in the different ESMs, the geographical extent of irrigation and afforestation differs strongly among different ESMs due to each model's native schemes on irrigation and the transition to forest. This is especially the case for the EC-EARTH afforestation simulation, in which the afforestation simulated was extremely low compared to the other ESMs (Figure 1d,e,f). Therefore, the afforestation scenario from EC-EARTH is not considered in this study. Likewise, large discrepancies regarding the simulated irrigation expansion are related to different irrigation parameterisations being implemented in the different ESMs (see appendix B in De Hertog et al., 2023). Within the model version of EC-EARTH used in this study, irrigation does not cause any effect on atmospheric moisture content (i.e. the water cycle of the land and atmosphere model are not coupled). Hence, the irrigation expansion scenario from EC-EARTH is also not considered in this study.

100

105




Figure 1. Land cover and land management changes as implemented in the three different ESMs. Cropland expansion (CROP-CTL ; a, b, c), afforestation (FRST-CTL ; d, e, f), and irrigation expansion (IRR-CROP ; g, h, i) implemented in CESM, MPI-ESM, and EC-EARTH, respectively. Both, cropland expansion and afforestation, are shown as a change in area fraction (%) while irrigation expansion is shown through the irrigation flux in mm/year.

This implementation of the ~~LCLMCs~~ LCLMC in the ESM land cover maps following a checkerboard pattern enables a signal separation of the ESM response into local and non-local components (Winckler et al., 2017a; De Hertog et al., 2023). The local effects refer to climate effects directly induced by the LCLMC within the grid cell, while the non-local effects refer to climate effects induced by LCLMC elsewhere through changes in atmospheric circulation or advection. These non-local effects can be directly derived from the simulation output as the climate changes occurring over grid cells where no land cover change occurred can only be caused by non-local mechanisms. By interpolating the values of non-local effects within the 'no change' grid cells to the 'change' grid cells we create a global map of non-local effects due to a certain LCLMC. This global map is then used to subtract the values found in the 'change' grid cells, which contain the total signal (i.e. local and non-local effects). After subtraction we obtain values for the local effects over all 'change' grid cells. These are again interpolated to create a global map representing the local climate effects. This approach has been explained in detail in previous studies (Winckler et al., 2017a; De Hertog et al., 2023) and is also explained in more detail here in Appendix A.

120

This separation is only applicable to (near-)surface variables and not to variables representing processes that extend higher into the atmosphere, as there is lateral mixing between different adjacent atmospheric grid cells above the surface. Therefore, the signal separation approach is not applied to the atmospheric variables (i.e. variables that have a vertical dimension into the atmosphere). ~~As the~~ The analysis presented here focuses on atmospheric processes, specifically moisture recycling, which is computed through a moisture tracking algorithm requiring atmospheric variables, ~~we are not able to separate local and~~

125

~~non-local effects for all results in this study. Instead.~~ Therefore, we analyse the raw ESM output directly, which represents an extreme case of LCLMC applied in a checkerboard pattern. For the variables where signal separation can be applied, we provide those results to support interpretations of these signals. All calculations are applied over the last 30 years of the simulations and at each ESM’s native spatial resolution (latitude x longitude) (i.e., MPI-ESM: 1.88° x 1.88°, CESM: 0.90° x 1.25°, EC-EARTH: 0.7°x 0.7°).

2.2 LCLMC-induced impact on the net water fluxes

To understand the net change in the atmospheric water cycle induced by the different ~~LCLMCs~~LCLMC, we first analyse their effects on evaporation and precipitation and compare them to the reference simulation of each LCLMC scenario. In addition, the vertically integrated moisture flux convergence (MFC) is computed using the basic principles of conservation of water vapor (Banacos and Schultz, 2005; Cook, 2009; Thiery et al., 2016; Van de Walle et al., 2020), as shown in Eq. 1 below.

$$P - E \approx -\frac{1}{g\rho_w} \int_{p_s}^{p^{TOA}} (\nabla \cdot q\mathbf{v}) dp \quad (1)$$

Where g is the gravitational acceleration [m/s^2], ρ_w is the density of water (1000 kg/m^3), p_s and p^{TOA} is the pressure at surface level and top of the atmosphere respectively, q represents the specific humidity of an air parcel [kg/kg], \mathbf{v} its horizontal wind vector [m/s], P is the precipitation flux per unit area [m/s], E is the surface evaporation flux per unit area [m/s]. $\nabla \cdot (q\mathbf{v})$ is the atmospheric moisture convergence from the surface to the top of the atmosphere (TOA). The MFC is computed based on 6-hourly data along the available pressure levels of each ESM. For EC-EARTH, only eight atmospheric levels were available, which is insufficient to compute MFC. Hence, $P - E$ is used as a proxy for the MFC in EC-EARTH. Over land, MFC or $P - E$ are often used a proxy for water availability (Van de Walle et al., 2020; Thiery et al., 2016), and changes of these measures can help understand the impacts of ~~LCLMCs~~LCLMC on the redistribution of water over land. The maps for MFC and $P - E$ changes are included in Appendix D.

The comparison of P , E , and MFC changes with respect to the corresponding reference simulation is performed for the three different LCLMC scenarios, i.e., cropland expansion, afforestation, and irrigation. We focus on annual mean values for the analysis. However, seasonal means (DJF: December, January, February; JJA: June, July, August) are shown in Appendix C. As explained in Sect. 2.1 we only apply the signal separation to P and E and not MFC as this metric is computed based on variables covering the entire atmospheric column.

2.3 Moisture tracking analysis

To further quantify the direct influence of ~~LCLMCs~~LCLMC on precipitation and unravel the reach of locally-induced LCLMC on precipitation and water availability, we perform a moisture tracking analysis. Here, we apply the Eulerian moisture tracking

model WAM-2layers (van der Ent et al., 2014; Benedict et al., 2020) to identify the origin of precipitation and the fate of evaporation in the ESM simulations and to evaluate the impact of LCLMC-induced evaporation changes on precipitation and water availability. The output of WAM-2layers is then used to compute several metrics relevant to moisture recycling, which can help uncover LCLMC-induced effects within the different ESMs. In this study, we focus on two spatial scales of moisture recycling: (i) local recycling and (ii) continental recycling. Note that local recycling does not imply that only the local effects (following the signal separation approach explained above) are used for the computation of this metric. The moisture tracking algorithm is applied on the raw ESM data as it requires atmospheric variables which cannot be signal separated. Local moisture recycling is defined on the grid-cell area of each ESM, which differs by definition (see Sect. 2.1) and makes them hard to compare across the ESMs. Therefore, additional area-independent metrics are used. Evaporation and precipitation length scales (van der Ent et al., 2010) illustrate the distance that moisture travels on average to or from a given grid cell. In the following, all recycling metrics are presented at annual time scales. Details on the setup of WAM-2layers and the definition of moisture recycling metrics are presented in more detail in the following sections.

2.3.1 WAM-2layers

A moisture tracking algorithm, the Water Accounting Model - 2 layers (WAM-2layers, van der Ent et al. (2014)), is applied to analyse the effects of the different ~~LCLMCs~~LCLMC on moisture recycling. We use a recent version of this algorithm, which was modified to ingest climate model data with limited vertical levels (Benedict et al., 2020). This moisture tracking algorithm uses a Eulerian approach to solve the atmospheric moisture balance over each grid cell and a specified time step (van der Ent et al., 2014). Model outputs comprise the origins of precipitation or evaporation at the local scale or continental scale depending on which tracking is performed, and facilitate the quantification of local and continental moisture recycling measures (van der Ent et al., 2010; van der Ent et al., 2014, see below). The algorithm has been applied numerous times in recent years for ESM output (Benedict et al., 2020; Guo et al., 2020; Findell et al., 2019; Bosmans et al., 2020).

Here, the surface and atmospheric data from all ESM simulations at the original spatial resolution (see Sect. 2.1) and the finest temporal resolution (CESM: 6 hours, MPI-ESM: 3 hours for surface variables and 6hr for atmospheric variables, EC-EARTH: 6 hours for surface variables and daily for atmospheric variables) are used as (offline) inputs for WAM-2layers. To avoid stability problems related to the numerical discretisation in WAM-2layers (van der Ent et al., 2014), all ESM forcings are linearly interpolated to 15-min time steps. The moisture tracking is applied to the last 30 years within the 160-year simulation period.

2.3.2 Local moisture recycling

Here, two definitions of local moisture recycling are used. First, the local precipitation recycling ratio ρ_r , i.e. the fraction of precipitation over a region r that originates from evaporation over the same region (see Eq. 2), is calculated. The remaining fraction of precipitation over that region (i.e., $1 - \rho_r$) originates from evaporation upwind that is advected into the region r , and can be of either land or oceanic origin. Second, the local evaporation recycling ratio ϵ_r , i.e. the fraction of evaporation from a

region r that falls as precipitation over the same region (see Eq. 3), is used. The remaining fraction of local evaporation ($1 - \epsilon_r$) is transported away from the region and falls downwind of that region as precipitation.

Using the output from WAM-2layers, precipitation over the region r with area A_r , here referred to as P , can be separated into the precipitation originating from the same region (P_r) and the remaining precipitation that originates from upwind regions (P_a), so that $P = P_r + P_a$. Using these outputs, the local precipitation recycling ratio can be calculated as

$$\rho_r = \frac{P_r}{P}. \quad (2)$$

Similarly, the local evaporation recycling ratio can be calculated using evaporation from the region r , here referred to as E , and the evaporation that falls as precipitation over the same region (E_r), i.e.

$$\epsilon_r = \frac{E_r}{E}. \quad (3)$$

Both local recycling definitions are subject to the area of the region considered (A_r). Here, local recycling is defined on the area of a grid cell. It is noted that this area varies with latitude per definition and, in addition, varies for the ESM simulations employed here (see Sect. 2.1). The differences between local recycling ratios thus need to be interpreted with caution and are not comparable across different data sources or ESMs. To overcome these shortcomings we compute length ratios and continental recycling ratios.

2.3.3 Precipitation and evaporation length scales

To assess local moisture recycling independently of the ESM, we compute the length scale of the moisture recycling process as introduced by van der Ent and Savenije (2011). Here we summarise the derivation of extracting the length scale metric from local moisture recycling ratios, for the complete derivation of how length scales are defined we refer to van der Ent and Savenije (2011). Following (Dominguez et al., 2006) [Dominguez et al. \(2006\)](#), the moisture recycling ratio is a function of the distance travelled along a streamline (x), the horizontal wind speed (u) and the moisture flux (M) normalized by the atmospheric storage capacity (S_a):

$$\gamma_r(x) = 1 - \left[\exp\left(-\frac{M}{S_a u} x\right) \right] \quad (4)$$

Here, $\gamma_r(x)$ is the moisture recycling ratio (which can be evaporation or precipitation recycling as defined in Eqs.(2) and (3)), and M is evaporation (E) for the evaporation recycling ratio and precipitation (P) for the precipitation recycling ratio. This equation is valid for a streamline starting at distance $x=0$. By definition, the recycling ratio equals one at the origin ($x=0$) and also in case of zero moisture uptake ($M=0$). With increasing moisture uptake and distance to the source region, the ratio tends exponentially towards zero, which indicates that the original moisture taken up at the source location is removed from the air. van der Ent and Savenije (2011) subsequently rewrote this equation of the local moisture recycling ratio as a function of the length scale (λ_γ):

$$\lambda_\gamma = \frac{S_a u}{M} \quad (5)$$

By then filling in Eq. 5 in Eq. 4 and integrating it for the distance x an equation is obtained for the regional average moisture recycling γ_r :

$$\gamma_r = \frac{\Delta x + \lambda_\gamma \exp(\frac{\Delta x}{\lambda_\gamma}) - \lambda_\gamma}{\Delta x} \quad (6)$$

Here, γ_r is the moisture recycling ratio over a given region r , with r representing a single grid cell in order to compute the length scale at the local, grid cell level. Δx equals the representative length of a grid cell and is computed from the zonal and meridional lengths of a grid cell weighted by the moisture fluxes in the corresponding directions (van der Ent and Savenije, 2011). From this equation the length scale can be isolated and solved based on the local moisture recycling to get the following expression using W as the Lambert W-Function (e.g. Corless et al., 1996):

$$\lambda_\gamma = \frac{\Delta x}{W\left[\frac{\exp(\frac{1}{\gamma_r - 1})}{\gamma_r - 1}\right] + \frac{1}{1 - \gamma_r}} \quad (7)$$

Length scales overcome one of the major shortcomings of local recycling ratios, which are dependent on the shape and size of the source region they are computed over (van der Ent and Savenije, 2011; Theeuwens et al., 2023), ~~which is normally problematic with~~. This occurs when using regular latitude-longitude grids in which grid cells vary both in shape and size within a model as well as between models that applied apply different resolutions. Length scales of local moisture recycling, in contrast, are designed to be area- and shape-independent at least for the region over which one can assume the same climatological conditions (i.e., the variables in Eq. 7 do not change much across nearby grid cells within a given grid cell). The length scales give an indication of the distance over which moisture would travel on average to or from a given grid cell under the given local hydrological and climatological conditions (van der Ent and Savenije, 2011). Hence, they should not be interpreted as an actual travel distance, but rather as which depends strongly on environmental conditions over the thousands of kilometres moisture typically travels and thus do not represent recycling strength in the local scope which is aimed for here. The local scope (i.e. ESM grid cell size) is preferred as it allows us to understand changes to moisture recycling as much disconnected from remote changes in LCLMC as possible, thus giving an indication of the local sensitivity of moisture recycling to LCLMC independent on downwind/upwind environmental conditions (see also Appendix E). Length scales should thus be interpreted as a local process-based metric of moisture recycling strength (Eq. 5) expressed in distance units (km). A short length scale indicates that local recycling is strong and a long length scale indicates that local recycling is weak.

Like local recycling ratios, the length scales can be calculated from a precipitation- or an evaporation-centric perspective (i.e., precipitation recycling ratios λ_ρ or evaporation recycling ratios λ_ϵ). Here the precipitation length scale represents the

length scale of precipitation raining down in a given grid cell, and the evaporation length scale represents the length scale of evaporation that travels from a given grid cell before precipitating. Both length scales can be derived from the local recycling ratios presented above (see section 2.3.2). By computing these different length scales for the different simulations representing the different LCLMC, we can assess how the LCLMC affects precipitation and evaporation recycling strength.

250

2.3.4 Continental moisture recycling

To study the continental contribution to moisture recycling, we compute the continental recycling ratios. Analogous to previous studies (e.g. Brubaker et al., 1993; van der Ent et al., 2010; Gimeno et al., 2012; Findell et al., 2019; Gimeno et al., 2020), we define continental precipitation recycling ratio ρ_c as the fraction of precipitation over land that originates from land evaporation.

255 The precipitation recycling ratio answers the question 'how much of the moisture precipitating over land originates from land?'. The remaining fraction $(1 - \rho_c)$ of the precipitation over land originates from evaporation over oceans. Similarly, continental

evaporation recycling ratio ϵ_c is defined as the fraction of land evaporation that falls as precipitation over land. These ratios can be computed through equations (2) and (3) by defining the region r as all land areas. Hence, in contrast to the local recycling ratios, continental recycling ratios refer to the same area, i.e. the area of all continental land regions A_c , which facilitates a direct

260 comparison of recycling ratios between the ESMs with different spatial resolution employed here. Continental evaporation and precipitation are computed by tracking all continental moisture fluxes at the same time.

3 Results

3.1 Changes in atmospheric moisture fluxes due to LCLMC

All ESMs generally show a decrease in evaporation over land due to cropland expansion and an increase in evaporation due to afforestation and irrigation expansion (Figure 2). However, some of the ESMs' regional signals deviate from this general

265 pattern. For cropland expansion (Figure 2a,d,g), CESM and MPI-ESM simulate a quasi-global decrease in evaporation over all land areas. However, in contrast, some regions also show an increase such as central U.S. in CESM, as well as East Africa and western Australia in MPI-ESM. The effects over the mid-latitudes exhibit a strong-large seasonality, with an increase in evaporation in JJA and a decrease in DJF. This impact is clearly visible in CESM and also slightly visible in MPI-ESM (Figure

270 C1 and Figure C2). In EC-EARTH, the annual patterns are less clear, with a strong-large decrease in evaporation following cropland expansion over tropical forests and a slight decrease over the mid-latitudes, but a clear increase over sub-tropical and tropical regions, such as the Sahel, East Africa, India and Australia. Moreover, all models clearly distinguish between effects over deforested grid cells and those that have remained unaltered, following the checkerboard implementation of LCLMC (see Sect. 2.1). In EC-EARTH, there are deforested patches that show a distinct decrease in evaporation while the nearby unaltered

275 grid cells instead show a strong-large increase.




Figure 2. The mean annual impacts of land cover and land management changes on evaporation (mm/day), for cropland expansion (CROP-CTL; a,d), afforestation (FRST-CTL; b,e) and irrigation expansion (IRR-CROP; c,f) for CESM and MPI-ESM respectively. Cropland expansion for EC-EARTH is shown in (g).

Regarding afforestation, MPI-ESM shows opposite patterns compared to cropland expansion, with mostly an increase in evaporation (Figure 2b,e). However, in CESM this increase is only apparent over the Southern Hemisphere tropics while the Northern-Hemisphere extratropics and the Sahel show a clear ~~increase~~ decrease in annual evaporation due to afforestation. Whereby it should be noted that the decrease in the extratropics is clearly linked to the boreal summer season (Figure C1 and Figure C2). Over the North Atlantic, CESM simulates a wide-spread and ~~strong~~ large decrease of evaporation, which may be linked to the widespread cooling of the North Atlantic in this ESM (De Hertog et al., 2023). Regarding irrigation expansion, both models, for which this simulation is analysed (CESM and MPI-ESM), strongly agree on the sign of evaporation change over land and simulate a global increase (Figure 2c,f). Differences between both ESMs are mostly related to the extent to which irrigation is applied within the different ESMs (see Figure 1).

The effects of LCLMC on precipitation are less similar across the models, ~~but~~ despite some regionally consistent patterns ~~emerge~~ (Figure 3). ~~Globally, cropland expansion causes a decrease in precipitation while afforestation and irrigation expansion cause an increase.~~ Under the cropland expansion scenario, there is a decrease of precipitation over land in MPI-ESM (Figure 3b) and CESM, except for Central ~~America~~ U.S., the Congo basin and Eastern Africa (Figure 3a). The patterns of precipitation changes around the tropics in CESM are similar to those found in Portmann et al. (2022) ~~-In their study,~~




Figure 3. The mean annual impacts of land cover and land management changes on precipitation (mm/day), for cropland expansion (CROP-CTL; a,d), afforestation (FRST-CTL; b,e) and irrigation expansion (IRR-CROP; c,f) for CESM and MPI-ESM respectively. Cropland expansion for EC-EARTH is shown in (g).

Portmann et al. (2022) which showed that deforestation in CESM is cooling-induced cooling over the Northern extratropics, which leads to changes in causes large scale circulation changes such as the intensity of the Hadley cell and the position of the intertropical convergence zone, which could also explain the simulated pattern here. In line with CESM, the cropland expansion simulation with EC-EARTH simulate simulates the largest changes in precipitation over the ocean and forest areas within the tropics. However, there is tropics which seems to indicate a shift of these changes that is causing less precipitation north of Australia and more precipitation over the tropical Pacific tropical rainfall bands. The effects of LCLMCs LCLMC on MFC also show substantial regional difference differences between CESM and MPI-ESM (Figure D1). Overall, the patterns in MFC are highly similar to those seen for precipitation (Figure 3) which indicates that these precipitation changes are likely driven by the changes in MFC.

Afforestation causes widespread increases in precipitation over land (Figure 3b,e). This increase is quasi-global in MPI-ESM, while in CESM some areas show a precipitation decrease, such as the Indian subcontinent, the Sahel, and Europe. In CESM, the afforestation-induced precipitation differences over the intertropical convergence zone are again similar to those found by Portmann et al. (2022) which indicates that shifts of the large-scale circulation determine the precipitation patterns in

this ESM.

For the irrigation expansion scenario, all models simulate a global increase in precipitation. In CESM, it is apparent that Southeast Asia is an exception to this pattern and shows a clear reduction of precipitation despite being an area of large-scale irrigation. This ~~finding might be linked to~~ is in line with the hypothesis that ~~the~~ regional temperature decreases as a consequence of irrigation expansion (De Hertog et al., 2023) ~~, further causing~~ cause a weakening of the Indian summer monsoon and a decrease of precipitation over ~~large parts of the continent~~ South East Asia, a feedback mechanism that has ~~also~~ been documented in several previous studies (de Vrese et al., 2016; Guimberteau et al., 2012; Thiery et al., 2017). This decrease in precipitation over India is to some extent also present in MPI-ESM, although it is not as ~~strong~~ large as in CESM. ~~However, for~~ For both ESMs it is clear that the response over this region occurs mainly during JJA (Figure C3 and Figure C4). This regional decrease is also apparent for MFC (Figure D1c,f), which further supports the weakened Indian Summer Monsoon hypothesis.

3.2 Local and non-local effects on precipitation

The checkerboard implementation allows for a consistent separation between local induced effects, which are directly caused by the LCLMC, and non-local induced effects due to LCLMC elsewhere (See Section 2.1 and Appendix A). In this section we focus on the signal separated results of precipitation for CROP-CTL and IRR-CROP to illustrate this separation in our simulations. The signal separated maps for FRST-CTL and for evaporation are provided in Appendix B.




Figure 4. Annual mean precipitation response in mm/day to cropland expansion (CROP-CTL) of CESM, MPI-ESM and EC-EARTH. The local effect in CESM (a), the non-local effect (b) and the total effect (c). The latitudinal average of the local (blue), non-local (yellow) and total (green) signals of CESM (d). (e-h): same as (a-d), but for MPI-ESM. (i-l): same as (a-d), but for EC-EARTH. The stippling on the maps shows grid cells where all the sign of change is consistent throughout the simulation.

Under the cropland expansion scenario, there is a decrease of local precipitation over land in MPI-ESM and EC-EARTH which is especially clear over tropical latitudes (Figure 4e,i). In CESM, in contrast, the majority of the effects are non-local, largely a decrease in precipitation, with only a relatively small local increase in some parts of the tropics. In MPI-ESM the non-local effects shows a regional decrease in precipitation mostly around areas of intense local effects. In EC-EARTH the non-local effects consist of a strong increase of precipitation, counteracting the local decreases. This is also confirmed by Figure B1 which shows the signal-separated effects of evaporation for the cropland expansion simulations. Using the checkerboard implementation to separate local and non-local effects, EC-EARTH simulates a clear local decrease in evaporation due to cropland expansion, while the non-local effect ~~cause~~ causes an increase in evaporation over the tropics, resulting in attenuated total effects (Figure 2). This pattern of strong decreases over areas of LCLMC and decreases in neighbouring grid cells is also visible for $P - E$, used here as proxy for MFC, and indicates a mesoscale circulation effects induced by the LCLMC implementation in a checkerboard pattern (see Figure D2g). In both MPI-ESM and EC-EARTH the local effects dominate the total signals while for CESM the total ~~signal~~ effect is mostly determined by the non-local effects. This further indicates that the results in CESM appear to be dominated by circulation changes which is also apparent in the signal separated results for afforestation (Figure B4).




Figure 5. Annual mean precipitation response in mm/day to irrigation expansion (IRR-CROP) of CESM and MPI-ESM. The local effect in CESM (a), the non-local effect (b) and the total effect (c). The latitudinal average of the local (blue), non-local (yellow) and total (green) signals of CESM (d). (e-h): same as (a-d), but for MPI-ESM. The stippling on the maps shows grid cells where all the sign of change is consistent throughout the simulation.

For irrigation expansion, both MPI-ESM and CESM indicate a dominant non-local increase in precipitation (Figure 5) which is mostly apparent in regions of intense irrigation (Figure 1). In MPI-ESM there are some smaller local increases in precipitation as well while in CESM the local effect appears unimportant. This general pattern does not hold over India where both ESMS show a non-local decrease in precipitation over this region despite a strong local increase in evaporation (Figure B3) which indicates that [changes in](#) circulation patterns are dominantly driving this effect. These signal separated results help in characterising the effects on the atmospheric water cycle within the different ESMS. However, they do not provide any information on the effects on moisture recycling which is analysed using the WAM-2layer moisture tracking algorithm in the next section.

3.3 Changes in local precipitation and evaporation length scales due to LCLMC

To unravel the direct impact of LCLMC-induced evaporation changes on ~~the~~ precipitation over land (and the other way around), we evaluate the outputs from WAM-2layers and ~~illustrate~~ [show](#) local evaporation and precipitation length scales for each model and LCLMC scenario. ~~An overview of consistent patterns for change in moisture fluxes and length scales is shown in Table ??.~~ Even in their control simulations, the different ESMS show very different magnitudes of length scales of moisture recycling, both for the precipitation length scale (Figure 6) and the evaporation length scale (Figure 7). CESM shows the largest length scales indicating that the importance of local recycling is relatively small. EC-EARTH, in contrast, generally shows very low values of the length scale indicating that local recycling is more important within this ESMS. The length scale values captured by the three ESMS in our study lie within the broad range of values found in literature (van der

Ent and Savenije, 2011; Theeuwen et al., 2023), even if ~~the-our~~ results are not directly comparable to literature as they are based on reanalysis data. To our knowledge, no studies exist presenting length scales for ESMs. However, it is known that ESMs simulate substantial differences in the global water cycle (Li et al., 2022), which logically also translates to substantial differences in length scales. Both reanalysis-based studies (~~van der Ent and Savenije, 2011; Theeuwen et al., 2023, e.g.-~~ e.g. van der Ent and Savenije, 2011; Theeuwen et al., 2023) present values for the length scales ranging roughly within 1000-7000 km for the majority of the land surface. In general, CESM overestimates the length scales compared to literature while EC-EARTH shows values at the lower end. Yet, the spatial patterns are quite similar among the ESMs and comparable to those in literature (van der Ent and Savenije, 2011; Theeuwen et al., 2023). The precipitation length scale is smallest over tropical rainforests and mountain ranges (see for example the Tibetan Plateau) indicating that these locations mostly get precipitation from nearby evaporation. ~~These patterns also occur~~ Similar patterns are apparent for the evaporation length scale, although ~~here~~ locations with a dry and continental climate such as Siberia and Western North America also show very low values. This implies that evaporation occurring within these locations generally precipitates nearby.




Figure 6. The annual mean precipitation length scale (km) for the control (CTL) simulation in CESM (a), MPI-ESM (b) and EC-EARTH (c). The effect of cropland expansion (CROP-CTL) on the annual mean precipitation length scale is shown for CESM (d), MPI-ESM (e) and EC-EARTH (f). The effect of afforestation (FRST-CTL) is shown for CESM (g) and MPI-ESM (h) and finally the effect of irrigation expansion (IRR-CROP) is shown for CESM (i) and MPI-ESM (j). Note that in the difference plots in (d)–(j), the areas with a reference evaporation length scale higher than 10000 km are cropped out.




Figure 7. The annual mean evaporation length scale (km) for the CTL simulation in CESM (a), MPI-ESM (b) and EC-EARTH (c). The effect of cropland expansion (CROP-CTL) on the annual mean evaporation length scale is shown for CESM (d), MPI-ESM (e) and EC-EARTH (f). The effect of afforestation (FRST-CTL) is shown for CESM (g) and MPI-ESM (h) and finally the effect of irrigation expansion (IRR-CROP) is shown for CESM (i) and MPI-ESM (j). Note that in the difference plots in (d)–(j), the areas with a reference precipitation length scale higher than 10000 km are cropped out.

370 In general, both the precipitation and evaporation length scale increase as a consequence of cropland expansion (see Figures 6d,e,f and 7d,e,f). In MPI-ESM, it seems that the Congo Basin is an exception with no clear changes in local recycling occurring there. In EC-EARTH, the patterns are more blurred than in the other ESMs with a decrease in length scales in some regions, such as **South Africa and spots over the Central U.S. and South Africa**. However, over regions where the largest cropland expansion occurred (such as Amazon basin and China, see also Figure 1), the patterns are consistent with the other ESMs.

375 Over Latin America, a dipole pattern of the change in the evaporation length scale in both EC-EARTH and CESM appears, showing an increase in the West and a decrease in the East. The general increase in length scale due to cropland expansion implies that the LCLMC induces a decrease in local recycling.

Afforestation (Figures 6g,h and 7g,h), induces a pattern that is opposite to the cropland expansion case, with a decrease in length scale for both precipitation and evaporation. However, in some areas, the patterns diverge from the general trend, e.g. afforestation causes an increase of the evaporation length scale over the Amazon in CESM, and an increase of the precipitation

length scale over the tropics in MPI-ESM. In general, the changes in both length scales are ~~stronger~~ larger for the extratropics, which is particularly visible for the evaporation length scale. This ~~effect on the evaporation length scale discrepancy~~ may be explained by the fact that the tropics are already densely forested in the CTL scenario ~~and thus the additional trees do not alter~~
385 ~~the local recycling favouring conditions~~, in contrast to the sparsely forested extratropics (Figure 1).

Regarding irrigation expansion (Figures 6i,j and 7i,j), the effects on the evaporation length scale are less clear and generally of small magnitude. Irrigation-induced differences show a tendency towards a decreased evaporation length scale, which is rather consistent in MPI-ESM but in CESM this pattern is less clear. The effects on the precipitation length scale in both ESMs
390 are larger and ~~more~~ consistently decreasing due to irrigation expansion. The change in precipitation length scale is small over the tropics due to the small amount of irrigation applied in this region (Figure 1). ~~We even observe a slight increase in length scale over the tropical forest in CESM and over some areas in the U.S. in MPI-ESM, which might imply circulation changes in those regions.~~

~~Summary of the effects of moisture fluxes over land due to the different LCLMC. Each cell indicates where the changes in moisture response are consistent in sign. LCLMC E P λ_c λ_p CROP-CTL decrease over tropics regional increase/decrease mostly decrease mostly increase FRST-CTL increase over tropics increase over tropics global increase mostly decrease IRR-CROP global increase global increase global decrease global decrease~~

395

3.4 Changes in continental moisture recycling due to LCLMC

While there are substantial differences in the local effects of LCLMC on the water cycle, their net impact on water availability
400 over land might not be the same. Here, we evaluate how LCLMC impacts E, P and P-E over land, and we identify the direct impact of LCLMC-induced effects on these fluxes via continental moisture recycling (see Section 2.3.4). The values of total annual precipitation (P), continental precipitation (P_c), continental precipitation recycling (ρ_c), evaporation (E), continental evaporation (E_c) and continental evaporation recycling (ϵ_c) are included in Appendix EF. Cropland expansion causes a net decrease of evaporation from land in CESM and MPI-ESM, while EC-EARTH simulates a small net increase of continental
405 evaporation (Figure 8a). Through this decrease of evaporation in CESM and MPI-ESM, less moisture is available for continental moisture recycling (dark bars in Figure 8a) and for precipitation over oceans (light bars in Figure 8a). Analogously, cropland expansion is causing a net decrease of precipitation over land in CESM and MPI-ESM, but a net increase in EC-EARTH, which is due to contrasting signs of change in different parts of the globe (Figure F2). In the former two models, the simulated decrease of precipitation mainly results from decreased moisture imports from the ocean (light bars in Figure 8b),
410 and only 42% and 26% respectively of the precipitation deficit is estimated to be of continental origin (dark bars in Figure 8b).

The effects on evaporation from land due to afforestation are consistent in sign and generally cause an increase for both ESMs. A large part of this increase is available for continental moisture recycling (dark bars in Figure 8a) in CESM (48%), but is negligible for MPI-ESM (2%). In MPI-ESM the increase in land evaporation mainly rains out over the oceans (light bars in Figure 8a). Evaluated over all land regions, afforestation increases precipitation over land in both MPI-ESM and CESM
415 Figure 8b). The magnitude is much smaller in CESM due to the large spatial heterogeneity in precipitation effects (Figure F1),

which cancel each other out causing only a small net increase of precipitation over land. For MPI-ESM, there is a large heterogeneity within the signal of change (Figure F1), causing diverging contributions of moisture for continental precipitation from ocean and land. Atmospheric circulation changes in this model cause an increase in precipitation of oceanic origin on land, while less precipitation is estimated to be of continental originsorigin.

420

Regarding irrigation expansion, there is a stronglarge increase of evaporation from land in CESM, of which most rains out over the oceans (light bar in Figure 8a), and only a small fraction (17%) is available for continental moisture recycling (dark bar in Figure 8a). In MPI-ESM, the effect is smaller due to some areas of decreased recycling (Figure F2) but the results still show an overall increase in evaporation of which most rains out over the ocean. For precipitation over land, the effect is not
 425 consistent across the ESMs. In CESM there is an increase in precipitation over land of which 34% is available for continental recycling. In contrast, for MPI-ESM there is a global decrease of continental precipitation, of which 12% stems from the precipitation deficit over land.




Figure 8. The global change in annual mean evaporation (a) and precipitation (b) on land is shown for CESM in blue, MPI-ESM in orange and EC-EARTH in green for cropland expansion (CROP-CTL), afforestation (FRST-CTL) and irrigation expansion (IRR-CROP). The contribution of continental moisture recycling changes is shown in a darker shade of the ESMs respective colours.




Figure 9. The total annual mean P-E on land (a) and change in P-E on land (b) is shown for CESM in blue, MPI-ESM in orange and EC-EARTH in green for cropland expansion (CROP-CTL), afforestation (FRST-CTL) and irrigation expansion (IRR-CROP). The contribution of continental moisture recycling changes is shown in a darker tone of the ESMs respective colours.

Finally, we can quantify the global effects of LCLMC on global water exports from land towards the ocean by analysing the effects on P-E. In general, $P - E$ is positive for all three ESMs, being substantially larger in CESM than in MPI-ESM and EC-EARTH (Figure 9a), indicating that the land receives more water from the atmosphere, and of oceanic origin, than it provides through evaporation. Both evaporation and precipitation decreases for MPI-ESM and CESM due to cropland expansion (Figure 8), but the decrease in evaporation is larger causing a net surplus of water at the land surface (i.e., $\Delta(P - E) > 0$). There is also a net surplus of water for EC-EARTH although this is mostly due to increased continental precipitation (Figure 8). In EC-EARTH and CESM, the contribution due to continental recycling is 16% and 20% respectively (light bars in Figure 8 and Figure 9b), but the majority comes from changes in oceanic moisture. This differs from MPI-ESM, where less moisture is recycled (dark bar in Figure 9b). Afforestation, in turn, is causing a net loss of water at the land surface for both ESMs. CESM simulates an increase in water export from land through an increase of land evaporation that rains out over the ocean (Figure 9b). In contrast MPI-ESM shows a slight increase in water availability over land due to oceanic moisture, with the effect of continental moisture changes dominating the global decrease in water availability. Following irrigation expansion, both CESM and MPI-ESM simulate a decrease in water available on land. In both simulations, enhanced evaporation through irrigation from land (see Figure 8a) mostly rains out over the ocean (light bars in Figure 9b), thus a loss of water on land to the ocean with the contribution of continental recycling being 8% and 5% for CESM and MPI-ESM, respectively.

4 Discussion

LCLMC can have substantial effects on atmospheric moisture fluxes and the local and continental recycling of moisture that determine water availability on land. ~~Common patterns emerge from our~~ The different ESMs show large differences in their hydrological responses to the different LCLMC scenarios. However, some common patterns do emerge from this multi-model analysis, despite strong differences in the implementation of LCLMC and the simulation of the hydrological response in the

~~different ESMs analysis.~~ For cropland expansion, all three ESMs agree that there is a general decrease ~~over land in evaporation~~
450 ~~and precipitation over most regions as well as a decreased in land evaporation, for precipitation the patterns are less clear with~~
~~large regional differences across ESMs. These changes in moisture fluxes affect the~~ local recycling strength ~~with a general~~
~~decrease.~~ In contrast, afforestation and irrigation expansion show ~~an opposite pattern of both increased a general increase in~~
precipitation and evaporation over most regions ~~with enhanced.~~ ~~Similarly,~~ local recycling strength ~~mostly increases despite~~
455 ~~large regional differences.~~ Here we will discuss some of the discrepancies between the different ESMs and their implications
on moisture fluxes and moisture recycling.

4.1 Different hydroclimatic responses of ESMs to LCLMC

The effects of LCLMC within the different ESMs have strong regional variations (e.g., Figure 2 and Figure 3). The differing
length scales between ESMs (Figure 7-6 and Figure 67) illustrate that different processes dominate within the different ESMs:
EC-EARTH shows a stronger importance of local processes in contrast to CESM, where atmospheric circulation seems to
460 dominate the effects. This is also illustrated through the signal separated results (Figure 4 and 5) where CESM mainly shows
non-local effects, while MPI-ESM and EC-EARTH show a stronger importance of local effects. This difference is also clear
from the effects on moisture fluxes as EC-EARTH simulates strong mesoscale effects (10 to 100 km), while in CESM global
circulation changes appear to dominate. CESM is known to be an ESM with a strong natural variability, as was shown in several
other studies (Deser et al., 2012, 2020). It has also been shown to simulate large-scale circulation shifts as a consequence to
465 land cover change (Portmann et al., 2022; Devaraju et al., 2018).

Discrepancies in the CTL length scales estimated for each ESM could also stem from the different spatial resolutions em-
ployed here (CESM:0.90°x1.25°, MPI-ESM:1.88°x1.88° and EC-EARTH:0.7°x0.7°). Although the concept of length scales is
independent of the spatial resolution (van der Ent and Savenije, 2011), the capability of ESMs to resolve processes explicitly is
470 resolution-dependent. This implies that certain processes, such as mesoscale convection, are potentially better resolved within
EC-EARTH than in CESM and MPI-ESM. These strong differences in length scales, despite showing effects from induced
LCLMC on local length scales which are consistent in sign, are important as they clearly illustrate some of the challenges
involved in modelling the atmospheric water cycle using ESMs. The LCLMC effects on climate simulated by CESM are dom-
inated by circulation effects with less importance of moisture recycling than a world as simulated by EC-EARTH. A next step
475 to better understand these strong differences could be to evaluate the outputs of ESMs to reanalysis products in order to better
understand potential biases in evaporation and precipitation fluxes that could help in explaining these differences.

The way LCLMC is implemented in the different ESMs also causes some discrepancies. Some of the ESMs only represent
crops by few generic crop types (such as MPI-ESM) while others have different crop types representing different biophysical
480 properties. CESM has eight different crop types representing common crops around the world (Lombardozzi et al., 2020). In
CESM, maize has high evaporation rates which might explain why afforestation over the Northern Hemisphere extratropics is
causing a decrease in evaporation, with particularly strong effects during summer (Figure C1). The discrepancy between the

effects due to afforestation and cropland expansion can be partially explained by a saturation effect, as the effects of adding trees are likely non-linear (Winckler et al., 2017b). For example, in the tropics, extreme deforestation will have larger impacts on the hydrological cycle than adding trees in an already densely forested region. This effect could explain some differences between these simulations such as the smaller precipitation length scale changes in afforestation over the tropics.

The implementation of irrigation also causes substantial differences in climatic responses among the ESMs, as the maps of irrigation extent and amounts differ strongly (Figure 1). Both MPI-ESM and CESM show an increase in precipitation, except for the Indian subcontinent where both ESMs show a decrease in precipitation. As there is a cooling over all irrigated areas (De Hertog et al., 2023), there is a lower land–ocean temperature contrast, which reduces convection over land and therefore precipitation (Figure 3 and Figure D1). This occurs despite the increases in evaporation (Figure 2) and enhanced local precipitation recycling (Figure 6). Considering all the above, it is likely that the reduced precipitation shown here is caused by a weakened Indian Summer Monsoon as was highlighted by previous studies (Puma and Cook, 2010; de Vrese et al., 2016; Thiery et al., 2017, 2020).

4.2 Implications of changes in moisture recycling due to LCLMC

LCLMC strongly affects the redistribution of moisture over land in the ESMs. While the absolute length scales of moisture recycling differ among the ESMs, LCLMC-induced changes ~~are consistent in local recycling~~ are typically consistent in sign across the ESMs, with cropland expansion mostly causing decreased recycling and afforestation and irrigation expansion mostly causing enhanced local recycling (Figure 7-6 and Figure 67). The effects of LCLMC on continental recycling and the continental contribution to ~~precipitation over land and evaporation from land~~ from precipitation/evaporation are less consistent across ESMs (Figure 8 and Figure 9), but also geographically more heterogeneous within the ESMs (Figure F1 and Figure F2). This is due to the complex interactions of local effects with non-local effects, such as advection and circulation changes, which all affect the redistribution of water globally.

Although the effects of LCLMC on the precipitation and evaporation changes are substantial, they are not as large as could be expected based on literature (Tuinenburg et al., 2020; Hoek van Dijke et al., 2022; Baudena et al., 2021; Wunderling et al., 2022; Staal et al., 2018). This could partially be due to the less extensive LCLMC scenarios considered here (only 50% change due to checkerboard approach). However, differences are expected because most previous studies are based on reanalyses and can only estimate the impact of upwind LCLMC changes on downwind precipitation using constant recycling ratios, neglecting any other interactions such as through changes in atmospheric circulation. Therefore, to fully capture the impact of LCLMC on moisture recycling, LCLMC model simulations should be compared to a control simulation, as done here. In fact, the resulting (substantial) differences in recycling ratios show that the interactions that are not considered when basing the studies on reanalysis are not negligible (at least within the context of extensive LCLMC scenarios as considered here). As LCLMC becomes increasingly relevant as a climate mitigation strategy it is important to consider a more holistic view of the influence

of these strategies on the water cycle. Consequently, more research is needed to better constrain the effects of LCLMC on moisture recycling, aiming to support science that can guide future land cover planning.

520 4.3 Circulation effects induced by checkerboard LCLMC implementation

The specific setup of these simulations, with a checkerboard pattern LCLMC, also has limitations and causes some artefacts within the results. This is, for example, illustrated in the clear checkerboard pattern visible for evaporation (Figure 2) and precipitation changes (Figure 3) from EC-EARTH, especially over the tropics. Due to the area dependence of the effects of land cover changes on moisture fluxes (Spracklen et al., 2018), mesoscale circulation effects occur in EC-EARTH but do not appear in the other (coarser) ESMs (see for example Figure 4). This checkerboard-like effect would likely not occur if a full land cover change was simulated instead of the checkerboard implementation of the LCLMC. This implementation could have some important implications, as the non-local effects for EC-EARTH do not represent the effects one would get in a full land cover change simulation, implying that the assumptions behind the checkerboard approach are not met (Winckler et al., 2017a; De Hertog et al., 2023). Moreover, the LCLMC-induced effects on atmospheric circulation and moisture fluxes also affect other climate variables, such as temperature. These checkerboard-induced circulation changes could also explain the differences between the temperature effects found in De Hertog et al. (2023): here, the checkerboard-implementation of cropland expansion in EC-EARTH caused tropical warming, and the simulations from Boysen et al. (2020) with EC-EARTH that simulated full deforestation changes (forest to grass conversion), showed tropical cooling. Further research is required to completely understand the implications of checkerboard-induced climate effects. For example, the LCLMC could be implemented in different densities (1/8, 1/4, 1/2) next to a full deforestation experiment to assess whether these effects are true artefacts of the LCLMC patterns. However, this might imply that the checkerboard approach for signal separation requires a rough spatial implementation to avoid mesoscale circulation effects as seen here for EC-EARTH.

5 Conclusions

In this study, we analysed the effects of land cover and land management changes (LCLMC) on the atmospheric water cycle in a slate of idealised simulations (cropland expansion, afforestation and irrigation expansion) performed by three different Earth System Models (ESMs). We showed that the effects on moisture fluxes are substantial ~~with, generally, decreased evaporation and precipitation over land due to cropland expansion and the opposite effects for~~ but differ strongly across the ESMs. Cropland expansion typically causes a decrease in evaporation, while for precipitation the sign of change depends on the region and ESM. For afforestation and irrigation expansion ~~. However, substantial discrepancies there is generally an~~ increase in both precipitation and evaporation although regional differences are important. The results presented here clearly illustrate that substantial discrepancies exist between the different ESMs ~~exist~~, with EC-EARTH displaying important local recycling and mesoscale circulation effects, while CESM shows a dominance of large-scale atmospheric circulation shifts. These differences can have various causes, such as model parameterisations of crucial processes (e.g., convection) or the extent to

550 which different land cover types are implemented within the ESMs on a global scale. Because some of these effects might have been indirectly influenced by the checkerboard LCLMC pattern used in this study, we advocate for more research to assess the implications of possible checkerboard-induced climate effects and the applicability of this approach for signal separation into local and non-local effects. Despite the strong differences between ESMs, the effects on local recycling are generally consistent in sign (with notable regional exceptions), with cropland expansion causing a decreased recycling strength, and afforestation and irrigation expansion generally causing an increased recycling strength. Overall we find that cropland expansion causes a net increase in water availability on land while afforestation and irrigation expansion cause a net decrease. However, these effects on water availability are caused by different changes in continental recycling due to LCLMC across the different ESMs. Our simulations show that changes due to atmospheric circulation patterns play an important role in explaining these patterns and should be taken into account when assessing the effects of future LCLMC on moisture recycling.

560 This is the first study ~~to~~ to our knowledge ~~to~~ to explicitly consider moisture recycling when assessing the LCLMC induced effects on moisture fluxes using multiple ESMs. Our results show that the effects of LCLMC on moisture recycling are substantial both on the local and global scale, with clear implications for water availability on land. ~~The~~ Our results highlight that large differences between the ESMs remain, which require more research. However, despite these inter-ESM differences it is clear that LCLMC will substantially affect the atmospheric water cycle. Therefore, the inclusion of potential effects of LCLMC on the atmospheric water cycle should ~~therefore~~ be considered in future land cover planning.

Code and data availability. The scripts used for the analysis of the moisture fluxes and the adapted version of WAM-2layers can be found on the GitHub page of the Department of Hydrology and Hydraulic Engineering of VUB (https://github.com/VUB-HYDR/2023_DeHertog_etal_ESD). The simulation data and postprocessed data used in this paper are available through dkrz (https://www.wdc-climate.de/ui/entry?acronym=DKRZ_LTA_1147_ds00003).

570 **Appendix A: Summary of checkerboard approach for signal separation**

The checkerboard approach was developed by Winckler et al. (2017a) for the signal separation into local and non-local effects within dedicated simulations with the MPI-ESM. Subsequently it has been applied to the LCLMC simulations performed within the LAMACLIMA project in a multi-model framework (De Hertog et al., 2023). 'Local effects' are defined as the direct climate effect due to the changes in land surface properties within a given grid cell. While 'non-local effects' are climate effects caused by changes in atmospheric circulation or advection as a consequence of LCLMC elsewhere (Winckler et al., 2017a). Here we summarise the approach and highlight technical processing steps taken to extract the local and non-local signal from the raw ESM output.

The checkerboard approach allows a robust signal separation of local and non-local effects by implementing the LCLMC in a checkerboard pattern, i.e. alternating 'change' grid cells, where LCLMC occurs, with 'no change' grid cells, which retain the

580 CTL land cover (see Figure 1). It can then be assumed that any effects in climate variables over the ‘no change’ grid cells can be attributed as a non-local effect while in the ‘change’ grid cells of the raw model output both local and non-local effects occur (Winckler et al., 2017a). The 150 year-simulation with constant external forcings is split into 5 slices of 30 years each. To account for natural variability, we treat each slice as a member of a perturbed initial condition ensemble. In order to apply the signal separation, a multi-year monthly mean is computed over each of these ensemble members (De Hertog et al., 2023).

585 (1) We subtract a reference member from the LCLMC member of interest, i.e., for CROP and FRST this reference member is CTL while for the IRR simulation it is CROP. (2) We mask the LCLMC grid cells of this difference map, hence, retaining only those grid cells where only non-local effects occur. (3) The grid cells with only non-local effects are spatially interpolated to fill the masked LCLMC grid cells in order to create a global map of the non-local effect. Consequently, to create a global map of the local effects, we mask all ‘no change’ grid cells from the difference map created in (1), thus only retaining the grid cells

590 where both local and non-local effects occur. The values of these grid cells are then corrected by the interpolated non-local values in order to only retain the local effects. Subsequently, the values of the resulting grid cells are again spatially interpolated in order to attain a full global map of local effects. Finally, the sum of both, the local and non-local effects, then represents the total effect. This is the effect we would expect if an actual idealised LCLMC simulation had been performed without an LCLMC checkerboard pattern, although it should be noted that this total effect can be considered a lower limit as the non-local

595 effects represent only a 50% change simulation (Winckler et al., 2017a). The checkerboard approach is implemented to each model grid at its native resolution. Hence, grid cell sizes vary across the different ESMs. As we have five ensemble members of 30 years for each simulation, we can extract local and non-local signals for each ensemble member, which are then used as a measure of uncertainty coming from natural variability (De Hertog et al., 2023). Within this study we focus on interpreting the raw ESM output (as this is used in the moisture tracking analysis) and only apply the signal separation for completeness

600 and to support the interpretation of the presented results.

Appendix B: Signal separated plots of evaporation and precipitation




Figure B1. Annual mean evaporation response in mm/day to cropland expansion (CROP-CTL) of CESM, MPI-ESM and EC-EARTH. The local effect in CESM (a), the non-local effect (b) and the total effect (c). The latitudinal average of the local (blue), non-local (yellow) and total (green) signals of CESM (d). (e-h): same as (a-d), but for MPI-ESM. (i-l): same as (a-d), but for EC-EARTH. The stippling on the maps shows grid cells where all the sign of change is consistent throughout the simulation.




Figure B2. Annual mean evaporation response in mm/day to afforestation (FRST-CTL) of CESM and MPI-ESM. The local effect in CESM (a), the non-local effect (b) and the total effect (c). The latitudinal average of the local (blue), non-local (yellow) and total (green) signals of CESM (d). (e-h): same as (a-d), but for MPI-ESM. The stippling on the maps shows grid cells where all the sign of change is consistent throughout the simulation.




Figure B3. Annual mean evaporation response in mm/day to irrigation expansion (IRR-CROP) of CESM and MPI-ESM. The local effect in CESM (a), the non-local effect (b) and the total effect (c). The latitudinal average of the local (blue), non-local (yellow) and total (green) signals of CESM (d). (e-h): same as (a-d), but for MPI-ESM. The stippling on the maps shows grid cells where all the sign of change is consistent throughout the simulation.




Figure B4. Annual mean precipitation response in mm/day to afforestation (FRST-CTL) of CESM and MPI-ESM. The local effect in CESM (a), the non-local effect (b) and the total effect (c). The latitudinal average of the local (blue), non-local (yellow) and total (green) signals of CESM (d). (e-h): same as (a-d), but for MPI-ESM. The stippling on the maps shows grid cells where all the sign of change is consistent throughout the simulation.

Appendix C: Seasonal effects on evaporation and precipitation




Figure C1. The seasonal mean (JJA) effects on evaporation in mm/day as a consequence of cropland expansion (CROP-CTL) in CESM (a), MPI-ESM (d) and EC-EARTH (g), for afforestation (FRST-CTL) in CESM (b) and MPI-ESM (e) and irrigation expansion (IRR-CROP) for CESM (c) and MPI-ESM (f).




Figure C2. The seasonal mean (DJF) effects on evaporation in mm/day as a consequence of cropland expansion (CROP-CTL) in CESM (a), MPI-ESM (d) and EC-EARTH (g), for afforestation (FRST-CTL) in CESM (b) and MPI-ESM (e) and irrigation expansion (IRR-CROP) for CESM (c) and MPI-ESM (f).




Figure C3. The seasonal mean (JJA) effects on precipitation in mm/day as a consequence of cropland expansion (CROP-CTL) in CESM (a), MPI-ESM (d) and EC-EARTH (g), for afforestation (FRST-CTL) in CESM (b) and MPI-ESM (e) and irrigation expansion (IRR-CROP) for CESM (c) and MPI-ESM (f).




Figure C4. The seasonal mean (DJF) effects on precipitation in mm/day as a consequence of cropland expansion (CROP-CTL) in CESM (a), MPI-ESM (d) and EC-EARTH (g), for afforestation (FRST-CTL) in CESM (b) and MPI-ESM (e) and irrigation expansion (IRR-CROP) for CESM (c) and MPI-ESM (f).

Appendix D: Moisture Flux Convergence and P-E




Figure D1. The mean annual impacts of land cover and land management changes on Moisture Flux Convergence (MFC) in mm/day, for cropland expansion (CROP-CTL; a,d), afforestation (FRST-CTL; b,e) and irrigation expansion (IRR-CROP; c,f) for CESM and MPI-ESM respectively. For EC-EARTH the P-E plots are available in Appendix D.




Figure D2. The annual mean effects on P-E in mm/day as a consequence of cropland expansion (CROP-CTL) in CESM (a), MPI-ESM (d) and EC-EARTH (g) for afforestation (FRST-CTL) in CESM (b) and MPI-ESM (e) and irrigation expansion (IRR-CROP) for CESM (c) and MPI-ESM (f).

605 **Appendix E: Moisture fluxes of continental origin** Difference between length scale and actual travel distance

The length scales are a shape- and size- independent metric defined to represent the local moisture recycling strength (van der Ent and Savenije, 2010). They are represented in distance units (typically km) but should not be confused with actual travel distances, an actual travel distance of moisture is not representative of local recycling strength. The actual travel distances as inferred from an air parcel trajectory accumulates the influences of downstream regions on the moisture content of the air parcel along its trajectory, which is typically within the order of thousands of kilometers. The length scale, in contrast, retains solely the local impact and is further independent to other effects, such as the grid-cell size.

610

To illustrate the difference between actual travel distance and local moisture recycling strength we refer to Figure 2 from van der Ent and Savenije (2011), shown below. This figure shows the relationship between precipitation recycling strength and distance travelled by the moisture as derived by different authors (Figure E1). These relationship strongly differs among the different studies, which is to be expected, as each of these studies refers to a different region over which the recycling ratio is defined. However, they all illustrate a certain basic pattern of moisture recycling: the recycling ratio is low at small distances and becomes larger with distance travelled.

615

620 Here, we add three additional lines to the original figure of van der Ent and Savenije (2011) to illustrate the influence of the
environment on the actual travel distance. These lines illustrate two hypothetical extreme cases where the moisture travels over
a hypothetical infinite desert (purple full line) and a hypothetical mountain range (orange full line). For both these cases the
local moisture recycling (i.e. moisture recycling ratio near the source region illustrated by the blue box) is the same as the
black full line. However, in the case of the mountain range (orange full line) the atmospheric flow is blocked by the mountain.
625 Consequently, the actual travel distance is low and due to the formation of precipitation the moisture recycling goes towards
one (i.e. all water has left the air parcel) which is illustrated on Figure E1 at the distance of the orange dot striped line. For
the case of the desert, the water particle reaches the hypothetical infinite desert at which no interaction with the ground occurs,
hence the atmospheric flow is not blocked. All moisture is retained in the air parcel and no additional moisture exchanges are
observed. Hence, the moisture recycling ratio remains constant while the actual travel distance will reach near infinite (purple
630 dot striped line in Figure E1).

This example clearly illustrates the local nature of the length scale, as this metric is not affected by the different environments
across the trajectory which do strongly affect the actual travel distance. The length scale will only be influenced by the local
changes (i.e. within the small distance of an ESM grid cell as illustrated by the blue box) and represents the distance moisture
635 would travel if the conditions over this local area would continue along the entire trajectory. This is illustrated by the thin black
line which results in a length scale of 2500 km here (as indicated by the dotted line). This local scope is preferred in this study
as it provides the local sensitivity of moisture recycling to local LCLMC, without considering any other upstream effects.




Figure E1. The relationship between the precipitation recycling ratios and distance using different formulas as was shown as Figure 2 in van der Ent et al. (2011). The orange and purple full lines are added here as they illustrate an extreme cases where moisture travels over a desert and mountain range respectively. They have the same local recycling as the black full line but the distance travelled is very different. The orange dot striped line indicates the distance at which the atmospheric flow is blocked by the mountain range and the purple dotted line indicates the moisture recycling ratio while the atmospheric flow travels the desert. Both illustrate very different moisture recycling patterns at a large distance but have the same local recycling (blue box). The length scale then represents the distance if those local conditions would be extrapolated (thin black line) which results in a distance of 2500 km for this case (dotted black line).

Appendix F: Moisture fluxes of continental origin




Figure F1. The annual mean continental evaporation is shown in mm/day for the CTL simulation in CESM (a), MPI-ESM (b) and EC-EARTH (c). The effect of cropland expansion (CROP-CTL) on the annual mean continental evaporation is shown for CESM (d), MPI-ESM (e) and EC-EARTH (f). The effect of afforestation (FRST-CTL) is shown for CESM (g) and MPI-ESM (h) and finally the effect of irrigation expansion (IRR-CROP) is shown for CESM (i) and MPI-ESM (j).




Figure F2. The annual mean continental precipitation is shown in mm/day for the CTL simulation in CESM (a), MPI-ESM (b) and EC-EARTH (c). The effect of cropland expansion (CROP-CTL) on the annual mean continental precipitation is shown for CESM (d), MPI-ESM (e) and EC-EARTH (f). The effect of afforestation (FRST-CTL) is shown for CESM (g) and MPI-ESM (h) and finally the effect of irrigation expansion (IRR-CROP) is shown for CESM (i) and MPI-ESM (j).

Table F1. Summary of annual total values of P , P_c , ρ_c , E , E_c , ϵ_c for the different simulations and ESMs. The absolute values are given in $km^3/year$.

CESM	P	P_c	ρ_c	E	E_c	ϵ_c
CTL	4359380	677025	0.155	2519998	659515	0.262
CROP	4244858	628946	0.148	2329935	599010	0.257
FRST	4356876	688149	0.158	2544179	671143	0.264
IRR	4300741	647669	0.151	2492105	626196	0.251
MPI-ESM	P	P_c	ρ_c	E	E_c	ϵ_c
CTL	3304814	468189	0.141	2065653	434761	0.210
CROP	3193252	438837	0.137	1942738	413053	0.213
FRST	3411430	43719	0.133	2168825	437156	0.201
IRR	3130711	431297	0.138	1988772	410919	0.207
EC-EARTH	P	P_c	ρ_c	E	E_c	ϵ_c
CTL	3597150	749179	0.208	2437720	72247	0.296
CROP	3625020	754114	0.208	2433458	720834	0.296

Author contributions. W.T., S.D.H., Q.L., J.P., F.L., S.G., I.M., F.L., C.E.L.F., J.K., D.M. and R.v.d.E. set up the idea for the analysis and
640 helped design the study. S.D.H. and C.E.L.F. performed the analysis and adapted the WAM-2layers model to ingest the LAMACLIMA sim-
ulation data. S.D.H. performed the calculations and wrote the paper. All authors commented on the paper and provided feedback throughout
the data analysis.

Competing interests. Sonia Seneviratne is member of the editorial board of the journal ESD. The authors have no other competing interests
to declare.

645 *Acknowledgements.* This work was funded by the LAMACLIMA (“LAnd MAnagement for CLimate Mitigation and Adaptation”) project
which is part of AXIS, an ERA-NET initiated by JPI Climate, and funded by DLR/BMBF (DE, Grant No. 01LS1905A), NWO (NL, Grant
No. 438.19.904), RCN (NO, Grant No. 300478) and BELSPO (BE, Grant No. B2/181/P1) with co-funding by the European Union (Grant No.
776608). The computational resources and services used in this work for the simulations and storage of CESM data and postprocessing with
650 and the Flemish Government – department EWI. For the storage of signal separated results and the simulations of MPI-ESM, this work used
resources of the Deutsches Klimarechenzentrum (DKRZ) granted by its Scientific Steering Committee (WLA) under project ID bm1147.
F.L. acknowledge VIDI-award from Netherlands Organization for Scientific Research (NWO) (Persistent Summer Extremes “PERSIST”
project: 016.Vidi.171.011). J.K. is grateful for the support from the Research Foundation–Flanders (FWO) under grant 1244122N. All the
simulations from EC-EARTH were carried out on European Center for Medium-Range Weather Forecast (ECMWF) platforms.

655 **References**

- Akkermans, T., Thiery, W., and Van Lipzig, N. P.: The regional climate impact of a realistic future deforestation scenario in the Congo Basin, *Journal of Climate*, 27, 2714–2734, 2014.
- Banacos, P. and Schultz, D.: The Use of Moisture Flux Convergence in Forecasting Convective Initiation: Historical and Operational Perspectives, *Weather and Forecasting*, 20, 351–366, <https://doi.org/10.1175/WAF858.1>, 2005.
- 660 Baudena, M., Tuinenburg, O. A., Ferdinand, P. A., and Staal, A.: Effects of land-use change in the Amazon on precipitation are likely underestimated, *Global Change Biology*, 27, 5580–5587, 2021.
- Benedict, I., van Heerwaarden, C., van der Ent, R., Weerts, A., and Hazeleger, W.: Decline in Terrestrial Moisture Sources of the Mississippi River Basin in a Future Climate, *Journal of Hydrometeorology*, 21, <https://doi.org/10.1175/JHM-D-19-0094.1>, 2020.
- Bonan, G. B.: Forests and climate change: forcings, feedbacks, and the climate benefits of forests, *science*, 320, 1444–1449, 2008.
- 665 Bosmans, J., van der Ent, R., Haarsma, R., Drijfhout, S., and Hilgen, F.: Precession-and obliquity-induced changes in moisture sources for enhanced precipitation over the Mediterranean Sea, *Paleoceanography and Paleoclimatology*, 35, e2019PA003 655, 2020.
- Bowen, I. S.: The ratio of heat losses by conduction and by evaporation from any water surface, *Physical review*, 27, 779, 1926.
- Boysen, L. R., Brovkin, V., Pongratz, J., Lawrence, D. M., Lawrence, P., Vuichard, N., Peylin, P., Liddicoat, S., Hajima, T., Zhang, Y., Rocher, M., Delire, C., Séférian, R., Arora, V. K., Nieradzic, L., Anthoni, P., Thiery, W., Laguë, M. M., Lawrence, D., and Lo, M. H.: Global climate
670 response to idealized deforestation in CMIP6 models, *Biogeosciences*, 17, 5615–5638, <https://doi.org/10.5194/bg-17-5615-2020>, 2020.
- Brubaker, K. L., Entekhabi, D., and Eagleson, P.: Estimation of continental precipitation recycling, *Journal of Climate*, 6, 1077–1089, 1993.
- Cook, K.: South American Climate Variability and Change: Remote and Regional Forcing Processes, pp. 193–212, https://doi.org/10.1007/978-90-481-2672-9_8, 2009.
- Corless, R. M., Gonnet, G. H., Hare, D. E., Jeffrey, D. J., and Knuth, D. E.: On the LambertW function, *Advances in Computational
675 mathematics*, 5, 329–359, 1996.
- Cui, J., Lian, X., Huntingford, C., Gimeno, L., Wang, T., Ding, J., He, M., Xu, H., Chen, A., Gentine, P., et al.: Global water availability boosted by vegetation-driven changes in atmospheric moisture transport, *Nature Geoscience*, 15, 982–988, 2022.
- De Hertog, S. J., Havermann, F., Vanderkelen, I., Guo, S., Luo, F., Manola, I., Coumou, D., Davin, E. L., Duveiller, G., Lejeune, Q., Pongratz, J., Schleussner, C. F., Seneviratne, S. I., and Thiery, W.: The biogeophysical effects of idealized land cover and land management changes
680 in Earth system models, *Earth System Dynamics*, 14, 629–667, <https://doi.org/10.5194/egusphere-2023-253>, 2023.
- de Vrese, P., Hagemann, S., and Claussen, M.: Asian irrigation, African rain: Remote impacts of irrigation, *Geophysical Research Letters*, 43, 3737–3745, <https://doi.org/10.1002/2016GL068146>, 2016.
- Deser, C., Knutti, R., Solomon, S., and Phillips, A. S.: Communication of the role of natural variability in future North American climate, *Nature Climate Change*, 2, 775–779, 2012.
- 685 Deser, C., Lehner, F., Rodgers, K. B., Ault, T., Delworth, T. L., DiNezio, P. N., Fiore, A., Frankignoul, C., Fyfe, J. C., Horton, D. E., et al.: Insights from Earth system model initial-condition large ensembles and future prospects, *Nature Climate Change*, 10, 277–286, 2020.
- Devaraju, N., de Noblet-Ducoudré, N., Quesada, B., and Bala, G.: Quantifying the relative importance of direct and indirect biophysical effects of deforestation on surface temperature and teleconnections, *Journal of Climate*, 31, 3811–3829, <https://doi.org/10.1175/JCLI-D-17-0563.1>, 2018.
- 690 Dominguez, F., Kumar, P., Liang, X.-Z., and Ting, M.: Impact of atmospheric moisture storage on precipitation recycling, *Journal of climate*, 19, 1513–1530, 2006.

- Findell, K. L., Keys, P. W., van der Ent, R. J., Lintner, B. R., Berg, A., and Krasting, J. P.: Rising temperatures increase importance of oceanic evaporation as a source for continental precipitation, *Journal of Climate*, 32, 7713–7726, 2019.
- 695 Gimeno, L., Stohl, A., Trigo, R. M., Dominguez, F., Yoshimura, K., Yu, L., Drumond, A., Durán-Quesada, A. M., and Nieto, R.: Oceanic and terrestrial sources of continental precipitation, *Reviews of Geophysics*, 50, 2012.
- Gimeno, L., Vázquez, M., Eiras-Barca, J., Sorí, R., Stojanovic, M., Algarra, I., Nieto, R., Ramos, A. M., Durán-Quesada, A. M., and Dominguez, F.: Recent progress on the sources of continental precipitation as revealed by moisture transport analysis, *Earth-Science Reviews*, 201, 103 070, 2020.
- 700 Goessling, H. and Reick, C.: What do moisture recycling estimates tell us? Exploring the extreme case of non-evaporating continents, *Hydrology and Earth System Sciences*, 15, 3217–3235, 2011.
- Guimberteau, M., Laval, K., Perrier, A., and Polcher, J.: Global effect of irrigation and its impact on the onset of the Indian summer monsoon, *Climate Dynamics*, 39, 1329–1348, 2012.
- Guo, L., Van Der Ent, R. J., Klingaman, N. P., Demory, M.-E., Vidale, P. L., Turner, A. G., Stephan, C. C., and Chevuturi, A.: Effects of horizontal resolution and air–sea coupling on simulated moisture source for East Asian precipitation in MetUM GA6/GC2, *Geoscientific Model Development*, 13, 6011–6028, 2020.
- 705 Hauser, M., Thiery, W., and Seneviratne, S. I.: Potential of global land water recycling to mitigate local temperature extremes, *Earth System Dynamics*, 10, 157–169, 2019.
- Hoek van Dijke, A. J., Herold, M., Mallick, K., Benedict, I., Machwitz, M., Schlerf, M., Pranindita, A., Theeuwens, J. J., Bastin, J.-F., and Teuling, A. J.: Shifts in regional water availability due to global tree restoration, *Nature Geoscience*, 15, 363–368, 2022.
- 710 Laguë, M. M., Bonan, G. B., and Swann, A. L.: Separating the impact of individual land surface properties on the terrestrial surface energy budget in both the coupled and uncoupled land–atmosphere system, *Journal of Climate*, 32, 5725–5744, <https://doi.org/10.1175/JCLI-D-18-0812.1>, 2019.
- Lawrence, D. M., Hurtt, G. C., Arneeth, A., Brovkin, V., Calvin, K. V., Jones, A. D., Jones, C. D., Lawrence, P. J., de Noblet-Ducoudré, N., Pongratz, J., et al.: The Land Use Model Intercomparison Project (LUMIP) contribution to CMIP6: rationale and experimental design, *Geoscientific Model Development*, 9, 2973–2998, 2016.
- 715 Li, Z., Liu, T., Huang, Y., Peng, J., and Ling, Y.: Evaluation of the CMIP6 precipitation simulations over global land, *Earth’s Future*, 10, e2021EF002 500, 2022.
- Lombardozi, D. L., Lu, Y., Lawrence, P. J., Lawrence, D. M., Swenson, S., Oleson, K. W., Wieder, W. R., and Ainsworth, E. A.: Simulating agriculture in the Community Land Model version 5, *Journal of Geophysical Research: Biogeosciences*, 125, e2019JG005 529, 2020.
- 720 Luysaert, S., Jammot, M., Stoy, P. C., Estel, S., Pongratz, J., Ceschia, E., Churkina, G., Don, A., Erb, K., Ferlicoq, M., et al.: Land management and land-cover change have impacts of similar magnitude on surface temperature, *Nature Climate Change*, 4, 389–393, 2014.
- Mahmood, R., Pielke Sr, R. A., Hubbard, K. G., Niyogi, D., Dirmeyer, P. A., McAlpine, C., Carleton, A. M., Hale, R., Gameda, S., Beltrán-Przekurat, A., et al.: Land cover changes and their biogeophysical effects on climate, *International journal of climatology*, 34, 929–953, 2014.
- 725 Mbow, H.-O. P., Reisinger, A., Canadell, J., and O’Brien, P.: Special Report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems (SR2), Ginevra, IPCC, 2017.
- Pongratz, J., Reick, C., Raddatz, T., and Claussen, M.: Biogeophysical versus biogeochemical climate response to historical anthropogenic land cover change, *Geophysical Research Letters*, 37, 2010.

- Pongratz, J., Schwingshackl, C., Bultan, S., Obermeier, W., Havermann, F., and Guo, S.: Land use effects on climate: current state, recent progress, and emerging topics, *Current Climate Change Reports*, pp. 1–22, 2021.
- 730 Portmann, R., Beyerle, U., Davin, E., Fischer, E. M., De Hertog, S., and Schemm, S.: Global forestation and deforestation affect remote climate via adjusted atmosphere and ocean circulation, *Nature Communications*, 13, 1–11, 2022.
- Puma, M. and Cook, B.: Effects of irrigation on global climate during the 20th century, *Journal of Geophysical Research: Atmospheres*, 115, 2010.
- 735 Rogelj, J., Shindell, D., Jiang, K., Ffifita, S., Forster, P., Ginzburg, V., Handa, C., Kheshgi, H., Kobayashi, S., Kriegler, E., et al.: Mitigation pathways compatible with 1.5 C in the context of sustainable development, pp. 93–174, 2018.
- Seneviratne, S. I., Wartenburger, R., Guillod, B. P., Hirsch, A. L., Vogel, M. M., Brovkin, V., van Vuuren, D. P., Schaller, N., Boysen, L., Calvin, K. V., et al.: Climate extremes, land–climate feedbacks and land-use forcing at 1.5 C, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 376, 20160450, 2018.
- 740 Shukla, J. and Mintz, Y.: Influence of land-surface evapotranspiration on the Earth’s climate, *Science*, 215, 1498–1501, 1982.
- Smith, C. and Spracklen, D. V.: reductions in rainfall due to tropical deforestation, *Nature*, 5690, 1, 2023.
- Spracklen, D., Baker, J., Garcia-Carreras, L., and Marsham, J.: The effects of tropical vegetation on rainfall, *Annual Review of Environment and Resources*, 2018.
- Spracklen, D. V., Arnold, S. R., and Taylor, C.: Observations of increased tropical rainfall preceded by air passage over forests, *Nature*, 489, 282–285, 2012.
- 745 Staal, A., Tuinenburg, O. A., Bosmans, J. H., Holmgren, M., van Nes, E. H., Scheffer, M., Zemp, D. C., and Dekker, S. C.: Forest-rainfall cascades buffer against drought across the Amazon, *Nature Climate Change*, 8, 539–543, 2018.
- te Wierik, S. A., Cammeraat, E. L., Gupta, J., and Artzy-Randrup, Y. A.: Reviewing the impact of land use and land-use change on moisture recycling and precipitation patterns, *Water Resources Research*, 57, e2020WR029234, 2021.
- 750 Theeuwens, J. J., Staal, A., Tuinenburg, O. A., Hamelers, B. V., and Dekker, S. C.: Local moisture recycling across the globe, *Hydrology and Earth System Sciences*, 27, 1457–1476, 2023.
- Thiery, W., Davin, E. L., Seneviratne, S. I., Bedka, K., Lhermitte, S., and van Lipzig, N. P.: Hazardous thunderstorm intensification over Lake Victoria, *Nature communications*, 7, 1–7, 2016.
- Thiery, W., Davin, E. L., Lawrence, D. M., Hirsch, A. L., Hauser, M., and Seneviratne, S. I.: Present-day irrigation mitigates heat extremes, *Journal of Geophysical Research*, 122, 1403–1422, <https://doi.org/10.1002/2016JD025740>, 2017.
- 755 Thiery, W., Visser, A. J., Fischer, E. M., Hauser, M., Hirsch, A. L., Lawrence, D. M., Lejeune, Q., Davin, E. L., and Seneviratne, S. I.: Warming of hot extremes alleviated by expanding irrigation, *Nature communications*, 11, 1–7, 2020.
- Tuinenburg, O., Hutjes, R., Stacke, T., Wiltshire, A., and Lucas-Picher, P.: Effects of irrigation in India on the atmospheric water budget, *Journal of Hydrometeorology*, 15, 1028–1050, 2014.
- 760 Tuinenburg, O. A., Theeuwens, J. J., and Staal, A.: High-resolution global atmospheric moisture connections from evaporation to precipitation, *Earth System Science Data*, 12, 3177–3188, 2020.
- Van de Walle, J., Thiery, W., Brousse, O., Souverijns, N., Demuzere, M., and Lipzig, N.: A convection-permitting model for the Lake Victoria Basin: evaluation and insight into the mesoscale versus synoptic atmospheric dynamics, *Climate Dynamics*, 54, 1779–1799, <https://doi.org/10.1007/s00382-019-05088-2>, 2020.
- 765 van der Ent, R. and Savenije, H.: Length and time scales of atmospheric moisture recycling, *Atmospheric Chemistry and Physics*, 11, 1853–1863, 2011.

- van der Ent, R., Wang-Erlandsson, L., Keys, P. W., and Savenije, H.: Contrasting roles of interception and transpiration in the hydrological cycle—Part 2: Moisture recycling, *Earth System Dynamics*, 5, 471–489, 2014.
- van der Ent, R. J., Savenije, H. H., Schaeffli, B., and Steele-Dunne, S. C.: Origin and fate of atmospheric moisture over continents, *Water Resources Research*, 46, 1–12, <https://doi.org/10.1029/2010WR009127>, 2010.
- 770 Wang, J., Chagnon, F., Williams, E., Betts, A., Rennó, N., Machado, L., Bisht, G., Knox, R., and Bras, R.: Impact of deforestation in the Amazon Basin on cloud climatology, *Proceedings of the National Academy of Sciences of the United States of America*, 106, 3670–4, <https://doi.org/10.1073/pnas.0810156106>, 2009.
- Wang-Erlandsson, L., Fetzer, I., Keys, P., Ent, R., Savenije, H., and Gordon, L.: Remote land use impacts on river flows through atmospheric teleconnections, *Hydrology and Earth System Sciences*, 22, 4311–4328, <https://doi.org/10.5194/hess-22-4311-2018>, 2018.
- 775 Winckler, J., Reick, C. H., and Pongratz, J.: Robust identification of local biogeophysical effects of land-cover change in a global climate model, *Journal of Climate*, 30, 1159–1176, 2017a.
- Winckler, J., Reick, C. H., and Pongratz, J.: Why does the locally induced temperature response to land cover change differ across scenarios?, *Geophysical Research Letters*, 44, 3833–3840, 2017b.
- 780 Winckler, J., Lejeune, Q., Reick, C. H., and Pongratz, J.: Nonlocal effects dominate the global mean surface temperature response to the biogeophysical effects of deforestation, *Geophysical Research Letters*, 46, 745–755, 2019.
- Wunderling, N., Staal, A., Sakschewski, B., Hirota, M., Tuinenburg, O. A., Donges, J. F., Barbosa, H. M., and Winkelmann, R.: Recurrent droughts increase risk of cascading tipping events by outpacing adaptive capacities in the Amazon rainforest, *Proceedings of the National Academy of Sciences*, 119, e2120777 119, 2022.