



# First comprehensive assessment of industrial era land heat uptake from multiple sources

Félix García-Pereira<sup>1,2</sup>, Jesús Fidel González-Rouco<sup>1,2</sup>, Camilo Melo-Aguilar<sup>3</sup>, Norman Julius Steinert<sup>4</sup>, Elena García-Bustamante<sup>5</sup>, Philip de Vrese<sup>6</sup>, Johann Jungclaus<sup>6</sup>, Stephan Lorenz<sup>6</sup>, Stefan Hagemann<sup>7</sup>, Francisco José Cuesta-Valero<sup>8</sup>, Almudena García-García<sup>8</sup>, and Hugo Beltrami<sup>9</sup>

<sup>1</sup>Complutense University of Madrid, Faculty of Physical Sciences, Madrid, Spain

<sup>2</sup>Geosciences Institute, IGEO (UCM-CSIC), Madrid, Spain

<sup>3</sup>Balearic Ocean Centre, Spanish Institute of Oceanography (IEO-CSIC), Palma de Mallorca, Spain

<sup>4</sup>NORCE, Norwegian Research Centre, Climate and Environment, Bergen, Norway

<sup>5</sup>Research Center for Energy, Environment, and Technology (CIEMAT), Madrid, Spain

<sup>6</sup>Max Planck Institute for Meteorology, Hamburg, Germany

<sup>7</sup>Helmholtz-Zentrum Hereon, Geesthacht, Germany

<sup>8</sup>Department Remote Sensing, Helmholtz Centre for Environmental Research - UFZ, Leipzig, Germany

<sup>9</sup>Climate and Atmospheric Sciences Institute, St. Francis Xavier University, Antigonish, Canada

**Correspondence:** Félix García-Pereira (felgar03@ucm.es)

**Abstract.** The anthropogenically-intensified greenhouse effect has caused a radiative imbalance at the top of the atmosphere during the industrial period. This, in turn, has led to an energy surplus in various components of the Earth system, with the ocean storing the largest part. The land contribution ranks second with the latest observational estimates based on borehole temperature profiles, which quantify the terrestrial energy surplus to be 6 % in the last five decades, whereas studies based on state-of-the-art climate models scale it down to 2 %. This underestimation stems from land surface models (LSMs) having a too shallow subsurface, which severely constrains the land heat uptake simulated by Earth System Models (ESMs). A forced simulation of the last 2000 years with the Max Planck Institute ESM (MPI-ESM) using a deep LSM captures 4 times more heat than the standard shallow MPI-ESM simulations in the historical period, well above the estimates provided by other ESMs. However, deepening the LSM does not remarkably affect the simulated surface temperature. It is shown that the heat stored during the historical period by an ESM using a deep LSM component can be accurately estimated by considering the surface temperatures simulated by the ESM using a shallow LSM and propagating them with a standalone forward model. This result is used to derive estimates of land heat uptake using all available observational datasets, reanalysis products, and state-of-the-art ESM experiments. This approach yields values of 10.5-16.0 ZJ for 1971-2018, slightly smaller than the latest borehole-based estimates (18.2 ZJ).

## 15 1 Introduction

The climate system has been experiencing a net heat gain in all its components (von Schuckmann et al., 2020) during the industrial period due to a radiative imbalance at the top of the atmosphere (Wild, 2020; Forster et al., 2021). The radiative imbalance is a consequence of the anthropogenically-intensified greenhouse effect (Gulev et al., 2021). Accurately estimating



the energy surplus partitioning among the different components helps in understanding their interactions and climate change  
20 influences on near-surface climate (Forster et al., 2021).

As a result of the energy imbalance at the surface, the land stored around 6 % of the terrestrial energy surplus in the last five  
decades, as derived from observational estimates (von Schuckmann et al., 2020; Cuesta-Valero et al., 2021a, 2023), being the  
second largest contributor after the ocean (ca. 90 %; Levitus et al., 2012; Abraham et al., 2013; von Schuckmann et al., 2020) to  
the total Earth System energy gain. Regardless of whether the amount associated with land heat uptake storage is comparatively  
25 small, climate models should realistically represent the partitioning of energy. State-of-the-art climate models estimate land  
contribution to be 2 %, misrepresenting observational results. A misrepresentation of land heat uptake in climate models  
can alter the surface energy balance (Mottaghy and Rath, 2006; García-García et al., 2023), with impacts on soil hydrology  
(Krakauer et al., 2013), particularly in permafrost regions (Andresen et al., 2020). The amount of energy absorbed by the  
ground is also relevant for other soil processes, such as respiration or productivity (Pries et al., 2017), and its biogeochemical  
30 activity (Soong et al., 2021).

Observational land heat uptake estimates have been derived from borehole temperature profile (BTP) log collections (Mareschal  
and Beltrami, 1992; Huang et al., 2000; Beltrami et al., 2002; Cuesta-Valero et al., 2021b, 2023). BTPs record transient tem-  
perature perturbations relative to the steady geothermal gradient. Assuming that the subsurface acts as a half-infinite fully  
conductive medium (Carslaw and Jaeger, 1959), BTPs provide information about the land heat uptake up to their logging date  
35 (Cuesta-Valero et al., 2021b). BTP logs are scarce and show uneven distributions of sampling dates and locations, as well as  
variable vertical resolution and depths. These factors may potentially have some impact on resulting estimates of land heat  
uptake (Beltrami et al., 2015; Melo-Aguilar et al., 2018, 2020). Nevertheless, estimates derived from BTPs are the only ob-  
servational source of information to quantify the land contribution to terrestrial energy budget so far (Forster et al., 2021; von  
Schuckmann et al., 2022).

State-of-the-art Earth System Models (ESMs; Eyring et al., 2016) have been incapable of properly reproducing observa-  
tional land heat uptake estimates. Their limited volume to store energy stems from imposing a bottom boundary condition  
placement (BBCP; Smerdon and Stieglitz, 2006; Stevens et al., 2007) in their land surface models (LSMs), ca. from 3 to 10 m  
in the majority of Coupled Model Intercomparison Project Phase 5 (CMIP5; Taylor et al., 2012) and Phase 6 (CMIP6; Eyring  
et al., 2016) models. These depths are insufficient to account for the heat penetrated into the subsurface in the industrial period,  
45 rendering them inadequate for estimating long-term heat storage (Alexeev et al., 2007; MacDougall et al., 2008; González-  
Rouco et al., 2009; Cuesta-Valero et al., 2016; Steinert et al., 2021a). Only a small number of CMIP6 ESMs impose a deeper  
BBCP (ca. 40 m), most of them using the Community Land Model (CLM; Lawrence et al., 2019) as the LSM (see Table 2).  
Further, longer-term simulations spanning multi-centennial or millennial time scales, like the ones developed within the Pale-  
oclimate Modelling Intercomparison Project (PMIP; Kageyama et al., 2018), should include deeper BBCPs to accommodate  
50 longer lasting and deeper subsurface energy transfer (Smerdon and Stieglitz, 2006; Alexeev et al., 2007; Steinert et al., 2021a).

Some simulation-based efforts have been devoted to deriving unbiased estimates of land heat uptake using deepened BBCP  
LSMs. Standalone one-dimensional half-infinite space heat conduction forward models (FMs; González-Rouco et al., 2006, 2009)  
were used to yield artificial global mean surface temperature histories and heat uptake estimates, forced by either observational-



based SAT (Huang, 2006) or ESM simulated ground surface temperature (GST) data (Stevens et al., 2007; MacDougall et al.,  
55 2008). More recently, modifications of the BBCP (Hermoso de Mendoza et al., 2020; González-Rouco et al., 2021) and soil  
hydrology (Steinert et al., 2021b) have been introduced into LSMs in standalone mode, and their impact on heat uptake and  
subsurface temperature variability extensively discussed. However, the impact of these LSM changes has not yet been explored  
within a fully-coupled ESM.

This work analyzes for the first time a long-term ESM simulation including a deep LSM. A version of the MPI-ESM  
60 (Mauritsen et al., 2019), modified to include a deep BBCP, is used to produce a 2000-year-long simulation including a PMIP4  
representation of natural and external forcings (Jungclaus et al., 2017). This simulation is compared with another experiment  
that was made with the same setup, albeit with the standard shallow LSM version and different initial conditions. The land  
energy storage is computed in the industrial period for the deep and shallow simulations considering the progressive subsurface  
warming down to their BBCP. In addition, land energy estimates were derived using an offline FM driven by the GSTs of the  
65 shallow and deep simulations as boundary conditions. This constitutes a first assessment of the impacts of deepening the BBCP  
on land energy storage and is extended to all available instrumental and modeling datasets. CMIP6 simulations and hybrid  
model-observational products (i.e. reanalysis) allow, for the first time, for having a variety of observational and model-based  
land heat uptake estimates that enable an exhaustive comparison with BTP-based estimates.

## 2 Data

### 70 2.1 MPI-ESM simulations

The first part of this work focuses on two simulations performed with version 1.2 of the Max Planck Institute for Meteorology  
ESM (MPI-ESM; Mauritsen et al., 2019), spanning the time interval 0-1850 CE and extended over the historical period and  
the SSP585 scenario. The interval encompassing the years 0-2100 CE is noted hereafter as P2k+. One of the experiments  
includes a standard shallow BBCP while the other one uses a deep BBCP (P2k+s and P2k+d, respectively). The shallow  
75 experiment used hereafter is the MPI-ESM contribution to PMIP4 (van Dijk et al., 2022). MPI-ESM comprises the ocean  
model MPIOM1.6 and the atmosphere model ECHAM6.3 (Stevens et al., 2013). The latter is directly coupled to the LSM  
JSBACH3.2 (JSBACH; Reick et al., 2021) through the surface exchange of mass, momentum, and heat. The low-resolution  
configuration (MPI-ESM1.2-LR) has an atmospheric horizontal resolution truncated to T63, corresponding approximately to a  
200-km grid cell size. This resolution is shared by the LSM. The CMIP6 version of JSBACH uses a 5-soil-layer discretization,  
80 with a BBCP imposed at 10 m (P2k+s), which is too shallow to properly account for the conductive propagation of decadal  
and centennial temperature perturbations (Steinert et al., 2021a). The deeper version of JSBACH (González-Rouco et al.,  
2021; Steinert et al., 2021b) has a 12-soil-layer scheme down to 1417 m and is used here in the P2k+d experiment. The P2k+  
simulations allow for evaluating the effect of having a deep LSM on surface temperatures at long time scales. Also, having  
a long millennial P2k+d experiment provides realistic, i.e. consistent with external forcing, preindustrial initial conditions for  
85 subsurface temperatures down to the BBCP in 1850.



**Table 1.** Characteristics of the observational-type global surface temperature data sets used in this work and their trends for various periods. The first seven rows account for reanalysis data, while the last five are for instrumental data. GST data was used, except in the cases marked with (1), for which surface air temperature (SAT) data were used. The soil temperature at the first layer was taken as GST in every case. Glacier areas (Antarctica and Greenland) were excluded from the calculations. Temperature trends in the three different periods assessed in Figure 3 (1950-2000, 1960-2020, and 1971-2018) for every data set and the means for reanalyses and observational subsets are also provided. (2) indicates cases in which the interval of trend evaluation is shorter due to limited data availability.

Name	Time span	Resolution (N. gridpoints)	Institution	1950-2000 [Kdec <sup>-1</sup> ]	1960-2020 [Kdec <sup>-1</sup> ]	1971-2018 [Kdec <sup>-1</sup> ]	Reference
20CRv3	1836-2015	1° (16779)	NOAA/ESRL/PSD	0.11	0.20 <sup>2</sup>	0.25 <sup>2</sup>	Slivinski et al. (2019)
CERA20c	1901-2010	1.125° (13184)	ECMWF	0.06	0.17 <sup>2</sup>	0.27 <sup>2</sup>	Laloyaux et al. (2018)
ERA20c	1900-2010	1.125° (13184)	ECMWF	0.09	0.22 <sup>2</sup>	0.35 <sup>2</sup>	Poli et al. (2016)
ERA5L	1950-2021	0.1° (1668600)	ECMWF	0.09	0.20	0.25	Muñoz Sabater et al. (2021)
NCEP1	1948-2021	2.5° (2707)	NCEP/NCAR	0.03	0.15	0.19	Kalnay et al. (1996)
JRA55	1958-2021	1.25° (10679)	JMA	0.10 <sup>2</sup>	0.20	0.26	Ebita et al. (2011)
LMRv2.1 <sup>1</sup>	1-2000	2° (4174)	NOAA	0.13	0.20 <sup>2</sup>	0.27 <sup>2</sup>	Tardif et al. (2019)
<b>MEAN TREND</b>				<b>0.09</b>	<b>0.19</b>	<b>0.24</b>	
BEST <sup>1</sup>	1750-2022	1° (16779)	Berkeley Earth	0.16	0.25	0.29	Rohde and Hausfather (2020)
CRUTEM5 <sup>1</sup>	1857-2021	5° (667)	MOHC	0.13	0.24	0.28	Osborn et al. (2021)
GISTEMPv4 <sup>1</sup>	1880-2021	2° (4174)	NASA-GISS	0.16	0.27	0.31	Lenssen et al. (2019)
NOAGlobalTemp <sup>1</sup>	1880-2021	5° (667)	NOAA	0.14	0.24	0.27	Zhang et al. (2019)
UDEL <sup>1</sup>	1900-2017	0.5° (66744)	Univ. of Delaware	0.13	0.23	0.28	Willmott and Matsuura (2018)
<b>MEAN TREND</b>				<b>0.14</b>	<b>0.25</b>	<b>0.29</b>	

<sup>1</sup> SAT used instead of GST. <sup>2</sup> Temperature trend value is given for a shorter period due to limited data availability.

In addition to the P2k+ experiments, an ensemble of 30 historical and SSP585 simulations (hereafter denoted as 30ENS) run with the CMIP6 version of the MPI-ESM, MPI-ESM1.2-LR, are considered. The experiments use the same model version as P2k+s and serve the purpose of tackling the uncertainties in historical warming due to internal variability.

## 2.2 Reanalysis, observational and CMIP6 model products

90 Global mean temperatures at the ground surface over land coming from 7 global reanalyses and 5 gridded observational  
 databases have also been used (Table 1) to derive land heat uptake estimates. In all cases, glacier areas were excluded. GST  
 data was used from 6 of the sources (see Table 1), whilst SAT was taken as a surrogate of the GST evolution when GST was  
 not available (6 data sources, see Table 1). This substitution leans on the assumption that SAT and GST are strongly coupled in  
 long-term scales (Melo-Aguilar et al., 2018). All sources cover the whole 20th century, except for ERA5-Land, NCEP1, and  
 95 JRA55. Further, all of them except LMRv2.1 cover the first decade of the 21st century. These datasets will allow for obtaining  
 land heat uptake estimates complementary to BTP-based values.

Additionally, GST and subsurface temperature simulations of the historical period and SSP585 scenario from the available  
 CMIP6 ESMs (Eyring et al., 2016) were used (see Table 2). This added up to 36 simulations stemming from 22 different  
 ESMs and some model variants, 38 if accounting for P2k+ simulations since 1850. Both for reanalyses and CMIP6 ESMs, the



**Table 2.** CMIP6 experiments used in this work (Col. 1) and their corresponding horizontal resolution (Col. 2) and LSM component (Col. 3). The number of layers (Col. 4) in the LSM and the BBCP depth (Col. 5), as well as temperature trends in the three different periods assessed in Fig. 3 (1950-2000, 1960-2020, and 1971-2018; Cols. 6-8) for every CMIP6 model and the multi-model mean are also provided. Information for the deep LSM configuration of the MPI-ESM1.2-LR is also included and shown in bold.

	ESM	Resolution N. gridpoints	LSM	N. of layers	BBCP [m]	1950-2000 [Kdec <sup>-1</sup> ]	1960-2020 [Kdec <sup>-1</sup> ]	1971-2018 [Kdec <sup>-1</sup> ]	Reference
1	ACCESS-CM2	1.88°x1.25° (6240)	CABLE2.4	6	4.0	0.08	0.28	0.32	Bi et al. (2020)
	ACCESS-ESM1-5	1.88°x1.25° (6240)	CABLE2.4	6	4.0	0.14	0.33	0.40	Ziehn et al. (2020)
2	BCC-CSM2-MR	1.13° (11583)	BCC-AVIM2	10	3.4	0.06	0.20	0.24	Wu et al. (2019)
3	CAMS-CSM1-0	1.13° (11583)	CLM1.0	10	3.4	0.00	0.12	0.14	Chen et al. (2019)
4	CanESM5	2.81° (1849)	CLASS3.6/CTEM1.2	3	4.0	0.20	0.38	0.40	Swart et al. (2019)
5	CAS-ESM2-0	1.41° (7322)	CLM4.0	15	42.1	0.03	0.20	0.25	Zhang et al. (2020)
6	CESM2	1.25°x0.94° (12376)	CLM5.0	25	49.6	0.11	0.27	0.29	Danabasoglu et al. (2020)
	CESM2-WACCM	1.25°x0.94° (12376)	CLM5.0	25	49.6	0.08	0.30	0.36	Gottelman et al. (2019)
7	CMCC-CM2-SR5	1.25°x0.94° (12376)	CLM4.5	15	42.1	0.09	0.23	0.28	Cherchi et al. (2019)
	CMCC-ESM2	1.25°x0.94° (12376)	CLM4.5	15	42.1	0.13	0.21	0.22	Lovato et al. (2022)
8	CNRM-CM6-1	1.41° (7437)	Surfex 8.0c	14	12.0	0.19	0.25	0.26	Volodire et al. (2019)
	CNRM-CM6-1-HR	0.5° (58356)	Surfex 8.0c	14	12.0	0.12	0.24	0.26	Volodire et al. (2019)
	CNRM-ESM2-1	1.41° (7437)	Surfex 8.0c	14	12.0	0.15	0.24	0.30	Séférian et al. (2019)
9	E3SM-1-1	1° (14545)	ELM1.0	15	42.1	0.03	0.30	0.37	Golaz et al. (2019)
	E3SM-1-1-ECA	1° (14545)	ELM1.0	15	42.1	0.06	0.28	0.35	Golaz et al. (2019)
10	EC-Earth3	0.7° (29572)	HTESSEL	4	1.9	0.16	0.39	0.46	Döscher et al. (2021)
	EC-Earth3-Veg-LR	1.13° (11583)	HTESSEL	4	1.9	0.12	0.26	0.29	Döscher et al. (2021)
11	FGOALS-f3-L	1.25°x0.94° (12376)	CLM4.0	15	42.1	0.15	0.25	0.28	HE et al. (2020)
	FGOALS-g3	2° (3503)	CLM4.0	15	42.1	0.14	0.24	0.23	Li et al. (2020)
12	GFDL-CM4.0	1.25°x1° (11642)	GFDL-LM4.0.1	20	10.0	0.07	0.29	0.35	Held et al. (2019)
	GFDL-ESM4.0	1.25°x1° (11642)	GFDL-LM4.0.1	20	10.0	0.02	0.21	0.26	Dunne et al. (2020)
13	GISS-E2-1-G	2.5°x2° (2908)	GISS LSM	6	3.5	0.01	0.23	0.24	Kelley et al. (2020)
	GISS-E2-1-H	2.5°x2° (2908)	GISS LSM	6	3.5	0.08	0.20	0.23	Kelley et al. (2020)
	GISS-E2-2-G	2.5°x2° (2908)	GISS LSM	6	3.5	-0.05	0.11	0.12	Rind et al. (2020)
14	HadGEM3-GC31-LL	1.88°x1.25° (6215)	JULES-GL7.0	4	3.0	0.04	0.32	0.40	Williams et al. (2018)
	HadGEM3-GC31-MM	0.83°x0.55° (31481)	JULES-GL7.0	4	3.0	0.05	0.28	0.32	Williams et al. (2018)
15	IPSL-CM6A-LR	2.5x1.26° (4617)	ORCHIDE2.0	18	90.0	0.13	0.26	0.32	Boucher et al. (2020)
16	KACE-1-0-G	1.88°x1.25° (6215)	JULES-GL7.0	4	3.0	0.10	0.28	0.33	Lee et al. (2019)
	MIROC6	1.41° (1849)	MATSIRO6.0	6	14.0	0.06	0.20	0.23	Tatebe et al. (2019)
17	MIROC-ES2L	2.81° (1849)	MATSIRO6.0	6	14.0	0.07	0.23	0.26	Hajima et al. (2020)
	MPI-ESM1.2-LR	1.88° (4172)	JSBACH <sub>s</sub>	5	9.8	0.13	0.24	0.30	Mauritsen et al. (2019)
18	<b>MPI-ESM1.2-LR</b>	<b>1.88° (4172)</b>	<b>JSBACH<sub>d</sub></b>	<b>12</b>	<b>1416.0</b>	<b>0.07</b>	<b>0.24</b>	<b>0.27</b>	<b>González-Rouco et al. (2021)</b>
19	MRI-ESM2-0	1.13° (11583)	HAL 1.0	14	10.0	0.05	0.23	0.27	Yukimoto et al. (2019)
20	NorESM2-LM	2.5x1.9° (3057)	CLM5.0	25	49.6	0.05	0.27	0.33	Seland et al. (2020)
	NorESM2-MM	1.25°x0.94° (12376)	CLM5.0	25	49.6	0.06	0.26	0.33	Seland et al. (2020)
21	TaiESM1	1.25°x0.94° (12376)	CLM4.0	15	42.1	0.03	0.27	0.34	Wang et al. (2021)
22	UKESM1-0-LL	1.88°x1.25° (6215)	JULES-GL7.0	4	3.0	0.06	0.32	0.41	Sellar et al. (2019)
	<b>MEAN TREND</b>					<b>0.09</b>	<b>0.26</b>	<b>0.30</b>	



100 temperature at the first soil layer was used as GST. Only the first ensemble member was selected for every specific ESM. In the  
case of the CMIP6 historical and the SSP585 simulation ensemble with the MPI-ESM1.2-LR (30ENS), only the first member  
was considered for the analyses hereafter and is referred to as MPI-ESM1.2-LR. Some features of the LSM of every CMIP6  
ESM, such as the number of soil layers, BBCP, and horizontal resolution can be found in Table 2. The ensemble of CMIP6  
simulations complements land heat uptake estimates derived from the reanalysis and observational data with ESM experiments  
105 that considered changes in external forcing since 1850 CE, allowing for tackling the system energy input regardless of the  
different realization of the internal variability. It is noteworthy that most CMIP6 ESMs include LSM depths lower than 14 m,  
which is not sufficient to properly represent the propagation of the temperature annual cycle into the soil column (Smerdon  
and Stieglitz, 2006). Only 14 out of 38 members impose a BBCP below 40 m, still shallower than the 200 m LSM depth  
recommended to correctly simulate the propagation of the centennial climate change signal (González-Rouco et al., 2021;  
110 Steinert et al., 2021a). Thus, land heat uptake estimates for CMIP6 in this work will be derived from the integration of the  
LSM subsurface temperatures from every simulation, and also corrected estimates will be provided by using a standalone FM  
approach (see Section 3).

### 3 Methods

#### 3.1 Subsurface temperatures generation

115 A half-infinite one-dimensional heat conduction FM (Mareschal and Beltrami, 1992; Cuesta-Valero et al., 2022b, 2023) is used  
to propagate yearly surface temperature data from the sources in Tables 1 and 2, and the P2k+ simulations (both d and s) and  
produce forward temperature anomaly profiles (FTP hereafter):

$$FTP(t_N, z) = \sum_{i=1}^N T(t_i) \left[ \operatorname{erfc} \left( \frac{z}{2\sqrt{\kappa t_i}} \right) - \operatorname{erfc} \left( \frac{z}{2\sqrt{\kappa t_{i-1}}} \right) \right] \quad (1)$$

where  $T(t_i)$  is annual mean surface temperature anomaly at time step  $t_i$ ,  $\kappa$  is thermal diffusivity, and  $z$  is depth. To compute  
120 FTPs at a certain time step, surface temperature data are trimmed to the interval spanning from the initial year ( $t_1$ ; e.g.,  
1850, 1950, 1960, and 1971) to  $t_i$ . Temperature anomalies are calculated by subtracting the temperature value at  $t_1$  from the  
temperature values of the trimmed series to depart from equilibrium initial conditions. These FTPs are equivalent to the BTPs  
used to yield observational land heat uptake estimates, which are derived by subtracting the geothermal gradient (mean state)  
from the absolute log temperature values (Mareschal and Beltrami, 1992; Huang et al., 2000). An ensemble of 3 different FTPs  
125 per time step was obtained using different thermal diffusivity values resulting from a Monte Carlo random sampling method  
of local properties (Davison and Hinkley, 1997), which accounts for the uncertainty in temporal and spatial soil heterogeneity  
(Cuesta-Valero et al., 2022b). The Monte Carlo method consisted of averaging 100 different global maps of thermal diffusivity  
with values randomly sampled from a realistic range of 0.5 to 1.5  $10^{-6} \text{ m}^2\text{s}^{-1}$ . The sample size was the original number of  
gridpoints over land for every data source in Tables 1 and 2. Then, percentiles 2.5, 50, and 97.5 of the 100 values were taken  
130 as the three thermal diffusivity values for computing the global FTPs. This is the approach for every data source except for the





P2k+ simulations, for which  $0.79 \cdot 10^{-6} \text{ m}^2\text{s}^{-1}$  is taken as the thermal diffusivity to produce the FTP, the actual value used by JSBACH to resolve the temperature vertical scheme (Reick et al., 2021). All the FTPs were resolved down to a common depth of 1417 m so that results could be comparable to the estimates derived from the P2k+d simulation, limited by a BBCP at 1417 m. This comparison is possible since this BBCP depth ensures the thermal decoupling between the ground surface and the  
135 bottom layer, which is equivalent to the subsurface acting as a half-infinite conductive medium (González-Rouco et al., 2021; Steinert et al., 2021a). As the uncertainties associated with local thermal properties and subsurface depth are tackled when calculating the FTPs, global differences between mean FTPs and land heat uptake estimates from different sources cannot be due to these factors.

In addition, both subsurface temperature data coming from P2k+ and CMIP6 simulations are used. Temperatures at different  
140 subsurface levels for every time step are aggregated to yield subsurface temperature profiles. Subsurface temperature anomaly profiles (STPs hereafter) are subsequently computed by subtracting the initial temperature profile from this  $T(t_i, z)$ , as follows:

$$STP(t_i, z) = T(t_i, z) - T(t_1, z) \quad (2)$$


where  $T(t_i, z)$  is the subsurface temperature at depth  $z$  and time step  $t_i$ , and, i.e. the subsurface temperature profile at time step  $t_i$ , and  $T(t_1, z)$  is the initial subsurface temperature profile.

### 145 3.2 Land heat uptake estimation

The annual time evolution of the land heat uptake,  $Q_s(t)$ , can be estimated by integrating the FTP anomalies along the whole soil column at every yearly time step as follows:


$$Q_s(t) = A \sum_{j=1}^n \Delta z(j) C_v(j) \frac{FTP(t, j) + FTP(t, j + 1)}{2} \quad (3)$$

where  $\Delta z(j)$ ,  $C_v(j)$ , and  $FTP(t, j)$  are thickness, volumetric heat capacity, and global mean FTP of layer  $j$ , respectively, and  
150  $A$  is the Earth's land surface area excluding glaciers ( $1.34 \cdot 10^{14} \text{ m}^2$ ; Cuesta-Valero et al., 2023). FTPs were calculated using an evenly spaced layering of 1 m thickness. Land heat uptake from both P2k+ and CMIP6 (Table 2) was also calculated from the simulated model subsurface temperatures down to the BBCP, i.e. STPs, by making use of Equation 2, albeit temperature layers are unevenly spaced in these cases. Soil layering thickness for both shallow and deep configurations of JSBACH is given by González-Rouco et al. (2021), whilst it can be found from the references in Table 2 for CMIP6 models. Analogous to what it  
155 is done to compute thermal diffusivity (see Section 3.1), three values of volumetric heat capacity were derived using a Monte Carlo approach (Davison and Hinkley, 1997; Cuesta-Valero et al., 2022b) for every source in Tables 1 and 2 using randomly sampled values between  $2.5$  and  $3.5 \cdot 10^6 \text{ Jm}^{-3}\text{K}^{-1}$ . The combination of three values for volumetric heat capacity and three for thermal diffusivity yield nine land heat uptake estimates, which are represented hereby in terms of mean and variability (two standard deviations).



**Figure 1. Temperature evolution in MPI-ESM.** Past 2k, historical, and SSP585 (P2k+) global means for a) SAT, b) GST, and c) ST5 anomalies over land areas (excluding Greenland and Antarctica) in 0-2000 CE relative to 1850-1900 for a simulation run with the MPI-ESM1.2-LR with a shallow (subscript s, red lines) and a deep (subscript d, blue lines) version of its LSM, JSBACH. Annual time series are shown in light, while 31-year moving averages are portrayed in dark colors. (d) 31-year moving averages in a-c (see legend for colors) are represented here together with the ensemble of 30 MPI-ESM1.2-LR CMIP6 standard shallow historical and SSP585 simulations (30ENS, grey lines) in 1850-2100 CE for comparison. (e) SAT (black), GST (purple), and ST5 (pink) differences between P2k+s and P2k+d, filtered with a 31-year moving average. For a,b,c,d, and e, vertical solid (dashed) lines indicate the 6 (19) strongest (weakest) volcanic events of 0-1850 CE. (f,g,h) Range of temperature variability of annual (A) and 31-yr moving average (MA31) time series in the preindustrial period (0-1850 CE) for SAT (f), GST (g), and ST5 (h). Whiskers represent percentiles 10 and 90, and boxes 25 and 75, respectively. (i) Temperature anomaly at 2100 with respect to 0-1850 CE (K) in P2k+ (see legend for colors). In this case, whiskers show the range of variability (percentiles 10, and 90) of the total warming (K) in 30ENS.






**Figure 2. Land heat uptake in the MPI-ESM.** (a) FTPs (continuous lines) and MPI-ESM STPs (symbols, see legend) for P2k+d (blue) and P2k+s (red) and for the 30-member ensemble mean of MPI-ESM historical and SSP585 simulations (30ENS, black) in years 2014 and 2100. Y-axis is logarithmic. (b) Land heat uptake derived from the yearly time step-wise vertical integration of FTPs and STPs for P2k+d (blue), P2k+s (red), and 30ENS (black). The time x-axis is unevenly spaced to enhance land heat gain since 1850. For the 30ENS, the confidence interval is also shown in grey ( $p < 0.05$ ).

## 160 4 Results

### 4.1 Influence of a realistically deep LSM

The comparison of the MPI-ESM P2k+s and P2k+d simulated SAT, GST, and soil temperature at layer 5 (ST5) allows for assessing the thermal impact of deepening the BBCP (Fig. 1). ST5 is selected because it is the deepest level for P2k+s, reaching 9.83 m, and can be compared with the corresponding layer in P2k+d. Figure 1a to d shows the time evolution of SAT, GST, and ST5 for the two simulations both in annual averages and with 31-year running means to emphasize the low-frequency response. The response to the common external forcing during the last two millennia (Jungclaus et al., 2017) generates similar interannual (e.g. volcanic events), multidecadal, and centennial variability for P2k+s and P2k+d for SAT and GST (Fig. 1a, b and d), indicating no systematic changes in preindustrial mean state (Fig. 1f) and variability (Fig. 1g and h) from using the deep or shallow versions of the model. The resulting SAT and GST warming of about 5.5 °C at the end of the 21st century agrees well with the range of warming in the ensemble of MPI-ESM runs (Fig. 1e) and with a climate sensitivity of 3.2 °C (Mauritsen and Roeckner, 2020). Note that for each simulation SAT and GST virtually overlap, showing long-term SAT-GST coupling (Melo-Aguilar et al., 2018). However, the impacts of deepening the BBCP are noticeable in ST5, consistent with González-Rouco et al. (2021), with P2k+d showing lower high-frequency volcanic cooling (e.g., 1257, 1457 and 1815, Fig.



**Figure 3. Decadal multi-source surface temperature and heat uptake variability.** (a) Global land mean temperature anomaly in 1850–2020 with respect to 1950 for reanalyses, observational, and a 38-member CMIP6 ensemble. SAT is plotted both for observational and CMIP6 sources and LMRv2.1 (tags in bold), whereas GST is given for the remaining reanalyses and CMIP6 simulations (see Tables 1,2). GST and SAT spread ranges for the CMIP6 ensemble are also portrayed ( $p < 0.05$ ). (b) Decadal global land heat uptake in the industrial period (1850–2020) from the different data sources in (a). Red (blue) tones in the heatmap represent decadal energy gains (losses). The black horizontal line separates CMIP6 models (38 members), including MPI-ESM P2k+ simulations, from reanalysis and observational data sources (12 members). CMIP6 sources are sorted downwards alphabetically, while the rest are ordered by length of decreasing periods. The last row represents the multi-source decadal mean (denoted as MEAN).



175 1c), preindustrial variability (Fig. 1f, g, and h) and warming ( $0.8\text{ }^{\circ}\text{C}$  by 2100, Fig. 1e and i) than P2k+s. The reduced warming is significant if compared to the range of 21st-century warming values produced by the 30ENS of MPI-ESM simulations.

Therefore, increasing the depth of the BBCP produces significant changes in the subsurface temperatures, yet it does not change surface temperatures. The GST of the P2k+s and P2k+d can be considered equal in terms of their range of variability and long-term trends, indicating that the surface temperatures simulated by a LSM are not affected by its BBCP depth. This result offers the potential to use GSTs from different sources, e.g. reanalysis or CMIP6 simulations, as a boundary condition for forcing an infinite half-space standalone FM and deriving realistic land energy storage estimates. This is first verified in Fig. 2 using the P2k+ runs. Figure 2a shows both STPs from P2k+s, P2k+d, and the ensemble mean STP from 30ENS, and their respective standalone FM-derived FTPs by the end of the historical period (2014) and SSP585 (2100). Both STPs and FTPs can capture the different degrees of warming between 2014 and 2100 and agree to portray its penetration down to a depth of about 150 m (Fig. 2a). However, there is a remarkable difference when comparing the FTPs and STPs for P2k+s, P2k+d, and the 30ENS. Whilst P2k+d FTP is capable of reproducing its analogous STP, FTPs, and STPs for P2k+s diverge in depth. This is because the FM is run with no BBCP imposed, which can be considered analogous to the deep configuration of JSBACH (1417 m). Consequently, FTP warm anomalies are smaller in depth than the ones corresponding to the STP, which complies with a BBCP at 10 m. Imposing a shallow BBCP generates a warm bias with depth since the ground heat flux is halted from penetrating downwards due to the existence of a zero-flux condition (Smerdon and Stieglitz, 2006; Alexeev et al., 2007; Steinert et al., 2021a). Thus, FTPs derived from P2k+s and 30ENS reproduce P2k+d STP and FTP, realistically representing the half-infinite space thermal conduction that governs heat propagation in observational BTPs (Mareschal and Beltrami, 1992; Cuesta-Valero et al., 2021c).

Figure 2b shows the land heat uptake since 0 CE resulting from the stepwise integration of FTPs and STPs for the P2k+ simulations, and FTPs for 30ENS. Both STP and FTP-derived heat energy uptake estimates for P2k+d and the FTP-derived for P2k+s depict a preindustrial land heat loss, which gets more acute in the Little Ice Age (from 1400 to 1850; Miller et al., 2012), and a noticeable heat uptake since 1900. The STP-based heat uptake evolution for P2k+s misses to simulate the heat loss and largely underestimates the magnitude of the subsequent heat uptake shown by P2k+d, yielding a heat uptake of around 18 ZJ by the end of the 21st century, which is 4 times smaller than the values derived from deep FTPs and P2k+d STP (about 75 ZJ). The agreement of P2k+s and P2k+d on the land heat uptake when the FM approach is used evidences the viability of using surface temperatures as boundary conditions to derive FTPs and estimate heat uptake when subsurface thermal schemes are too shallow.

## 4.2 Multi-source land heat uptake estimates


The previous analysis opens the possibility of estimating land heat uptake from a variety of sources. Global mean land surface temperature anomalies with respect to 1950 for the seven reanalyses, five observational databases, and the CMIP6 model ensemble are shown in Fig. 3a. By averaging 38 CMIP6 members, the effect of internal variability is minimized, thus emphasizing the response to forced variability. It can be seen that both SAT and GST CMIP6 multi-model means (thick lines) are fully coupled up to 2000, depicting sudden cooling events in the 1880 and 1990 decades, associated with the volcanic



activity of Krakatoa or later El Chichón and Pinatubo (Self and Rampino, 1981; McCormick et al., 1995). They also capture the cooling during the 1950s and 1960s and the subsequent warming. However, the warming trend is slightly higher for SAT than GST, which produces a small temperature offset of about 0.1 K by 2020. This offset is negligible compared to the range of temperature variability of the 38 members (SAT, shaded grey; GST, scratched gold), so SAT and GST can be considered as coupled, and therefore SAT can be used as alternative boundary information to feed the FM when GST data is not available. Regarding the observational and reanalysis datasets, surface temperatures lie within the range of CMIP6 multi-model variability. Nevertheless, observational data render higher temperature anomaly values than all the reanalyses, showing higher temperature trends in 1950-2000 and 1960-2020 (Table 1). Moreover, NCEP1 and JRA55 depict the coldest GST evolutions of all observational-based sources, even lying beneath the multi-model spread of CMIP6 models at some instants in the mid-70s.

The decadal evolution of the different surface temperature datasets impacts their land heat uptake, as it is shown in Fig. 3b, where decadal heat gains and losses are assembled for the time span of each of the 50 datasets collected in this work. All the sources agree on depicting a progressive land warming, only interrupted by transient coolings in the 1880s, 1900s, and 1960s, that coincide with the volcanic eruptions of Krakatoa (Self and Rampino, 1981), Péléé (Gueugneau et al., 2020) and Agung (Hansen et al., 1978). Moreover, heat gain is enhanced in 1980-2020, rendering a land uptake of around 14 ZJ. Even though land heat uptake intensifies in these decades, its increase is not linear, with 1990-2000 and 2010-2020 showing a deceleration in the heat uptake trend (0.4 and 0.8 ZJ, respectively) with respect to the 1970-2020 values (increasing land heat uptake of 1.2 ZJ per decade on average). The uncertainty associated with variability in soil properties is negligible and accounts for less than 0.1 ZJ decadally. Differences between sources in the intensity or the time occurrence of decadal heat gains (losses) can result from differences in the representation of internal variability, as suggested by the higher consistency among estimates from observations compared to those from CMIP6 models. Products showing the greatest decadal temperature trends in Tables 1 and 2, such as ACCESS-ESM1-5 or GISTEMPv4, generate the largest storage of energy, with 16.4 and 17.2 ZJ during 1980-2020, respectively.

The estimates of land heat uptake derived from reanalyses, observational data, and CMIP6 models (Fig. 3) are compared to values previously reported in the literature derived from logged BTPs (Beltrami et al., 2002; Beltrami, 2002; von Schuckmann et al., 2020; Cuesta-Valero et al., 2023) and CRUTEM2 SAT (Huang, 2006) for the periods 1950-2000, 1960-2020, and 1971-2018 (Fig. 4). The selection of these time intervals enables the comparison with previous works. In the case of CMIP6 models, both the heat estimates obtained through direct integration of the STPs and FTPs are depicted. The MPI-ESM P2k+ estimates (Fig. 2) are also included. All the estimates from this work and the previous literature show an increasing trend of land heat uptake towards the present. The 1950-2000 interval (Fig. 4a) shows less warming in the reanalyses than in the observational derived estimates, which is particularly noteworthy for LMRv2.1 and JRA55. These two reanalyses show no heat uptake in 1950-2000, which stems from a long-lasting negative temperature anomaly in 1950-1980 that is not shown by the rest of the sources (Fig. 3a). Furthermore, CMIP6 depth corrected estimates using FTPs are in range with the reanalysis and slightly below the observational estimates. Hence, all estimates based on FTPs agree overall with the previous estimates, with von Schuckmann et al. (2020) giving the highest land heat uptake values across all data sets. Figure 4b and c portray similar results for the periods 1960-2020 and 1971-2018, respectively. In both cases, the estimates derived from the different FTPs are



**Figure 4. Land heat uptake estimates from different sources and periods.** Land heat uptake during 1950-2000 (a), 1960-2020 (b), and 1971-2018 (c) derived from FTPs (hollow points) of reanalysis and observational databases (Table 1), CMIP6 FTPs (right hollow) and direct integration of CMIP6 STPs (left solid points, Table 2), and previous estimates: B02 (Beltrami, 2002), B02b (Beltrami et al., 2002), H06 (Huang, 2006), vS20 (von Schuckmann et al., 2020), and CV23 (Cuesta-Valero et al., 2023), orange crosses. The vS20 and CV23 estimates for 1950-2000 refer in fact to 1960-2000 and are marked with a star. BBCP depth for every CMIP6 model used in this work is given by a grey shading (see legend), while the multi-model mean both for the STP and FTP-based values is given by gold crosses. MPI-ESM P2k+d (P2k+s) estimates are plotted in dark blue (red).



consistent, with observational estimates at the upper end and some reanalyses showing the smallest land heat uptake values. For instance, land heat uptake estimates for the period 1971-2018 (Fig. 4c) range from 10.5 to 16.0 ZJ, yielding an ensemble mean of 13.9 ZJ for the CMIP6 FTPs, 12.2 ZJ for the reanalyses, and 15.0 ZJ for the observational-driven estimates. These differences are due to the smaller temperature trends shown by reanalyses and CMIP6 models when compared to observational data sets from the 1970s to the present, as shown by Fig. 3a (see Tables 1, 2 for the exact trend values).

Furthermore, Fig. 4 shows the dependency of land heat uptake on BBCP depth for CMIP6 STP-derived estimates (note gray shading on the left side), evidencing a significant increase when the FM is applied to correct estimates stemming from shallow BBCP ESMs (from 6.9 to 12.6 ZJ in 1960-2020, Fig. 4b, and from 7.0 to 13.9 ZJ in 1971-2018, Fig. 4c). Since 1950-2000 warming is very small for most of the CMIP6 models and very sensitive to the internal variability of each one (see Table 2), correcting the estimates in this case does not make a remarkable difference (Fig. 4a). Even after correcting the BBCP bias on land heat uptake, the range of CMIP6 multi-model variability is still very large, with a standard deviation of 5.4 (5.1) ZJ in 1960-2020 (1971-2018). This large variability is explained by some outlier estimates (e.g. CAMS-CSM1-0 and GISS-E2-2-G) which yield very small or even negative land heat uptakes, linked to the small climate sensitivity of these models in the periods of study, as shown by Table 2. By contrast, the CMIP6 simulations showing the greatest temperature trends also show the largest corrected FTP values, some of them yielding greater estimates than the state-of-the-art estimates from von Schuckmann et al. (2020); Cuesta-Valero et al. (2023). In all periods shown in Fig. 4, the multi-model CMIP6 mean is consistent with estimates yielded by reanalyses and observational data sets. Despite the consistency among the estimates obtained from the different data sets in this work, all of them yield results in the range of past literature values for the 1950-2000 period (Beltrami et al., 2002; Beltrami, 2002; Huang, 2006), but are lower than those in recent literature for 1960-2020 and 1971-2018, which quantified land heat uptake to be between 18.6 and 25 ZJ in 1971-2018 (von Schuckmann et al., 2020; Forster et al., 2021), and subsequently narrowed and lowered this range to be 18.0-18.4 ZJ (Cuesta-Valero et al., 2023).

## 5 Conclusions and discussion

We compare land surface and soil temperatures as well as land heat uptake from two simulations of the period 0-2100 CE made with two versions of the MPI-ESM, one including a standard shallow and the other one a more realistic deep BBCP. The deep LSM included in MPI-ESM generates about four times more heat uptake than the shallow version by the end of the historical period. This underestimation by the shallow LSM version only grows up to the end of the 21st century, reaching 18 (75) ZJ for the shallow (deep) BBCP versions. Also, while it produces relatively colder near-surface soil temperatures after 1850 CE, GST low-frequency variability and trends remain unchanged. This finding suggests the potential utility of employing GSTs to drive a FM and obtaining deep FTPs and subsequently unbiased estimates of land heat uptake. The concurrence of estimates from both STPs of P2k+d and FTPs of P2k+s and P2k+d supports the application of the FM method to correct insufficient land heat uptake values stemming from shallow LSMs. The results portray a land heat gain in the industrial period, which has intensified in the last decades of the twentieth and the first two decades of the twenty-first century. The FTP strategy is also applied to estimate land heat uptake from both observation-based products and CMIP6 simulations. Results are compared with previous estimates





based on observational BTPs, thus providing additional information to assess consistency between model and observational-based estimates. Land heat uptake in 1971-2018 is between 10 and 16 ZJ for all sources, being around 14 ZJ for CMIP6 models. This figure doubles the 7 ZJ CMIP6 models capture when their heat uptake capability is constrained by imposing a shallow BBCP, which shows the relevance of FM in correcting model-based estimates. Reanalysis-based estimates are slightly smaller than those obtained from applying the FM driven by observational gridded datasets or corrected CMIP6 simulations, in consistency with the generally smaller reanalyses warming since 1950 relative to other data sets. SAT observation-based estimates render the highest values, in consistency with SAT trends being slightly larger than GST's. SAT-related values can be thus regarded as an upper-limit land heat uptake estimate.

Nevertheless, all observational-based and corrected CMIP6 estimates underestimate the most recent values of land heat uptake based on BTPs. This discrepancy may stem from different sources of uncertainty. One possibility is that since borehole-derived surface temperature and ground heat flux stories have a low time resolution (e.g. 25-year time step in Cuesta-Valero et al., 2021b), they may not consider the decadal cold anomalies occurring in 1960-1980 (Fig. 3). This discrepancy may play a major role since decadal surface temperature differences might alter inter-decadal land heat imbalance due to subsurface thermal inertia (Beltrami et al., 2015). Moreover, BTP logs are scarce in the last decades and non-existent after the 2000s, so borehole-based estimates for the last two decades stemming from von Schuckmann et al. (2020) and Cuesta-Valero et al. (2023) for 1960-2020 and 1971-2018 are yielded extrapolating the ground heat flux temperature reconstructed in the last decades of the 20th century to the 21st century (2000-2018, and 2000-2020, respectively). This could be overlooking the global warming hiatus (1998-2012; Karl et al., 2015) effect on a relative slowdown in land heat uptake. Furthermore, the different methodological approaches used to yield observational and simulation-based (STP, FTP) heat uptake estimates might play a role. While STP and FTP-based heat uptake values in a certain time period are rendered by integrating the differences in the temperature profiles between the initial and the final time steps of that period, previous observational land heat uptake values are yielded by estimating ground heat flux stories based on single-time BTPs at different locations to then reconstruct the global heat uptake evolution (Beltrami et al., 2002; Cuesta-Valero et al., 2023). That entails BTP-based estimates do not represent net land heat variations in a certain time span, but a continuous variation of land heat uptake by aggregating logs taken in different dates and locations. This subsequently implies that this methodology preserves land energy changes due to past climate signals (Beltrami et al., 2017), which may contribute to comparatively greater heat uptake estimates in recent decades. With these differences reconciled, this work contributes to bringing observational and model-based land heat uptake estimates to the same frame of comparison, by removing the bias associated with imposing the BBCP. The comparison between CMIP6-based and observational data-driven land heat uptake estimates can also serve as an additional metric to assess the realism of ESMs in reproducing current global warming.

*Code and data availability.* MPI-ESM P2k+d SAT, GST, and subsurface temperature data in yearly resolution were stored in the following Zenodo repository (<https://doi.org/10.5281/zenodo.10364950>). This repository also includes the land mask, heat capacity, and volumetric heat capacity maps used by MPI-ESM to resolve the vertical subsurface thermal scheme. Both MPI-ESM P2k+s (MPI-ESM1.2-LR, Experiment



310 ID: past2k) and CMIP6 surface and subsurface temperature data are publicly available on the different data servers provided by ESGF  
(<https://esgf-node.llnl.gov/projects/cmip6/>). Reanalysis and observational gridded data sets stem from different open sources, whose links  
can be found following the referenced papers in Table 1. Previous observational estimates for the land heat uptake were taken directly  
from the published papers for Beltrami (2002), Beltrami et al. (2002), and Huang (2006), and from GCOS EHI database ([https://www.wdc-climate.de/ui/entry?acronym=GCOS\\_EHI\\_1960-2020](https://www.wdc-climate.de/ui/entry?acronym=GCOS_EHI_1960-2020)) for von Schuckmann et al. (2020) and Cuesta-Valero et al. (2023). The half-infinite  
FM used in this work is provided by CIBOR (Cuesta-Valero, 2022; Cuesta-Valero et al., 2022b). A numerical version of the FM, ConForM  
315 (v1.0), which allows for imposing a finite-depth (non-)zero flux bottom boundary condition and a geothermal gradient, is also available in  
the following Zenodo repository (<https://doi.org/10.5281/zenodo.10371439>).

*Author contributions.* FGP and JFGR conceptualized the study. FGP, CMA, and NJS assembled and processed the data, and FGP performed  
the calculations. All the co-authors contributed to the analysis and discussion of the results. FGP and JFGR prepared the manuscript with  
contributions from all the co-authors.

320 *Competing interests.* The authors declare no conflicts of interest.

*Acknowledgements.* This work has been developed within the frame of the GreatModelS (RTI2018-102305-B-C21) and SMILEME (PID2021-  
126696OB-C21) projects from the Spanish Ministry of Science and Innovation (MICINN) and CSIC Interdisciplinary Thematic Platform  
(PTI) Polar zone observatory (PTI-POLARCSIC). FGP was funded by contract PRE2019-090694 of the MICINN. FJCV was funded by  
the Alexander von Humboldt Foundation. HB is supported by the Natural Sciences and Engineering Research Council of Canada Discovery  
325 Grant (NSERC DG 140576948), the Canada Research Chairs Program (CRC 230687) and AceNet. We would also thank the Deutsches Kli-  
marechenzentrum (DKRZ) for the resources granted by its Scientific Steering Committee (WLA) to run MPI-ESM1.2-LR P2k+d simulation  
under project ID bm1026. We wanted also to appreciate the effort of the World Climate Research Programme's Working Group on Coupled  
Modeling, responsible for CMIP, and the climate modeling groups responsible for the model simulations used herein (Table 2) in making  
their simulation outputs accessible.



## 330 References

- Abraham, J. P., Baringer, M., Bindoff, N. L., Boyer, T., Cheng, L. J., Church, J. A., Conroy, J. L., Domingues, C. M., Fasullo, J. T., Gilson, J., Goni, G., Good, S. A., Gorman, J. M., Gouretski, V., Ishii, M., Johnson, G. C., Kizu, S., Lyman, J. M., Macdonald, A. M., Minkowycz, W. J., Moffitt, S. E., Palmer, M. D., Piola, A. R., Reseghetti, F., Schuckmann, K., Trenberth, K. E., Velicogna, I., and Willis, J. K.: A review of global ocean temperature observations: Implications for ocean heat content estimates and climate change, *Reviews of Geophysics*, 51, 450–483, <https://doi.org/10.1002/rog.20022>, 2013.
- Alexeev, V. A., Nicolsky, D. J., Romanovsky, V. E., and Lawrence, D. M.: An evaluation of deep soil configurations in the CLM3 for improved representation of permafrost, *Geophysical Research Letters*, 34, <https://doi.org/10.1029/2007GL029536>, 2007.
- Andresen, C. G., Lawrence, D. M., Wilson, C. J., McGuire, A. D., Koven, C., Schaefer, K., Jafarov, E., Peng, S., Chen, X., Gouttevin, I., Burke, E., Chadburn, S., Ji, D., Chen, G., Hayes, D., and Zhang, W.: Soil moisture and hydrology projections of the permafrost region – a model intercomparison, *The Cryosphere*, 14, 445–459, <https://doi.org/10.5194/tc-14-445-2020>, 2020.
- Beltrami, H.: Climate from borehole data: Energy fluxes and temperatures since 1500, *Geophysical Research Letters*, 29, 26–1–26–4, <https://doi.org/10.1029/2002GL015702>, 2002.
- Beltrami, H., Smerdon, J. E., Pollack, H. N., and Huang, S.: Continental heat gain in the global climate system, *Geophysical Research Letters*, 29, 8–1–8–3, <https://doi.org/10.1029/2001GL014310>, 2002.
- Beltrami, H., Matharoo, G. S., and Smerdon, J. E.: Impact of borehole depths on reconstructed estimates of ground surface temperature histories and energy storage, *Journal of Geophysical Research: Earth Surface*, 120, 763–778, <https://doi.org/10.1002/2014JF003382>, 2015.
- Beltrami, H., Matharoo, G. S., Smerdon, J. E., Illanes, L., and Tarasov, L.: Impacts of the Last Glacial Cycle on ground surface temperature reconstructions over the last millennium, *Geophysical Research Letters*, 44, 355–364, <https://doi.org/https://doi.org/10.1002/2016GL071317>, 2017.
- Bi, D., Dix, M., Marsland, S., O’Farrell, S., Sullivan, A., Bodman, R., Law, R., Harman, I., Jhan, S., Rashid, H., Dobrohotoff, P., Chloe, M., Hailin, Y., Tony, H., Savita, A., Dias, F., Fiedler, R., and Heerdegen, A.: Configuration and spin-up of ACCESS-CM2, the new generation Australian Community Climate and Earth System Simulator Coupled Model, *Journal of Southern Hemisphere Earth System Science*, 70, <https://doi.org/10.1071/ES19040>, 2020.
- Boucher, O., Servonnat, J., Albright, A. L., Aumont, O., Balkanski, Y., Bastrikov, V., Bekki, S., Bonnet, R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Caubel, A., Cheruy, F., Codron, F., Cozic, A., Cugnet, D., D’Andrea, F., Davini, P., de Lavergne, C., Denvil, S., Deshayes, J., Devilliers, M., Ducharne, A., Dufresne, J.-L., Dupont, E., Éthé, C., Fairhead, L., Falletti, L., Flavoni, S., Foujols, M.-A., Gardoll, S., Gastineau, G., Ghattas, J., Grandpeix, J.-Y., Guenet, B., Guez, Lionel, E., Guilyardi, E., Guimberteau, M., Hauglustaine, D., Hourdin, F., Idelkadi, A., Joussaume, S., Kageyama, M., Khodri, M., Krinner, G., Lebas, N., Levvasseur, G., Lévy, C., Li, L., Lott, F., Lurton, T., Luyssaert, S., Madec, G., Madeleine, J.-B., Maignan, F., Marchand, M., Marti, O., Mellul, L., Meurdesoif, Y., Mignot, J., Musat, I., Otlé, C., Peylin, P., Planton, Y., Polcher, J., Rio, C., Rochetin, N., Rousset, C., Sepulchre, P., Sima, A., Swingedouw, D., Thiéblemont, R., Traore, A. K., Vancoppenolle, M., Vial, J., Vialard, J., Viovy, N., and Vuichard, N.: Presentation and Evaluation of the IPSL-CM6A-LR Climate Model, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS002 010, <https://doi.org/10.1029/2019MS002010>, 2020.
- Carslaw, H. S. and Jaeger, J. C.: *Conduction of heat in solids*, vol. 2nd ed., Oxford University Press, New York, 1959.
- Chen, H.-M., Li, J., Su, J.-Z., Hua, L.-J., Rong, X.-Y., Xin, Y.-F., and Zhang, Z.-Q.: Introduction of CAMS-CSM model and its participation in CMIP6, *Advances in Climate Change Research*, 15, 540, <https://doi.org/10.12006/j.issn.1673-1719.2019.186>, 2019.



- Cherchi, A., Fogli, P. G., Lovato, T., Peano, D., Iovino, D., Gualdi, S., Masina, S., Scoccimarro, E., Materia, S., Bellucci, A., and Navarra, A.: Global Mean Climate and Main Patterns of Variability in the CMCC-CM2 Coupled Model, *Journal of Advances in Modeling Earth Systems*, 11, 185–209, <https://doi.org/10.1029/2018MS001369>, 2019.
- Cuesta-Valero, F. J.: CIBOR: Codes for Inverting BOREholes (Version 1.0.0), <https://github.com/cuestavalero/CIBOR>, 2022.
- 370 Cuesta-Valero, F. J., García-García, A., Beltrami, H., and Smerdon, J. E.: First assessment of continental energy storage in CMIP5 simulations, *Geophysical Research Letters*, 43, 5326–5335, <https://doi.org/10.1002/2016GL068496>, 2016.
- Cuesta-Valero, F. J., García-García, A., Beltrami, H., and Finnis, J.: First assessment of the earth heat inventory within CMIP5 historical simulations, *Earth System Dynamics*, 12, 581–600, <https://doi.org/10.5194/esd-12-581-2021>, 2021a.
- Cuesta-Valero, F. J., García-García, A., Beltrami, H., González-Rouco, J. F., and García-Bustamante, E.: Long-term global ground heat flux  
375 and continental heat storage from geothermal data, *Climate of the Past*, 17, 451–468, <https://doi.org/10.5194/cp-17-451-2021>, 2021b.
- Cuesta-Valero, F. J., García-García, A., Beltrami, H., González-Rouco, J. F., and García-Bustamante, E.: Xibalbá: Underground Temperature Database, <https://doi.org/10.6084/m9.figshare.13516487.v4>, 2021c.
- Cuesta-Valero, F. J., Beltrami, H., Gruber, S., García-García, A., and González-Rouco, J. F.: A new bootstrap technique to quantify uncertainty in estimates of ground surface temperature and ground heat flux histories from geothermal data, *Geoscientific Model Development*,  
380 15, 7913–7932, <https://doi.org/10.5194/gmd-15-7913-2022>, 2022b.
- Cuesta-Valero, F. J., Beltrami, H., García-García, A., Krinner, G., Langer, M., MacDougall, A. H., Nitzbon, J., Peng, J., von Schuckmann, K., Seneviratne, S. I., Thiery, W., Vanderkelen, I., and Wu, T.: Continental heat storage: contributions from the ground, inland waters, and permafrost thawing, *Earth System Dynamics*, 14, 609–627, <https://doi.org/10.5194/esd-14-609-2023>, 2023.
- Danabasoglu, G., Lamarque, J.-F., Bacmeister, J., Bailey, D. A., DuVivier, A. K., Edwards, J., Emmons, L. K., Fasullo, J., Garcia, R.,  
385 Gettelman, A., Hannay, C., Holland, M. M., Large, W. G., Lauritzen, P. H., Lawrence, D. M., Lenaerts, J. T. M., Lindsay, K., Lipscomb, W. H., Mills, M. J., Neale, R., Oleson, K. W., Otto-Bliesner, B., Phillips, A. S., Sacks, W., Tilmes, S., van Kampenhout, L., Versteijn, M., Bertini, A., Dennis, J., Deser, C., Fischer, C., Fox-Kemper, B., Kay, J. E., Kinnison, D., Kushner, P. J., Larson, V. E., Long, M. C., Mickelson, S., Moore, J. K., Nienhouse, E., Polvani, L., Rasch, P. J., and Strand, W. G.: The Community Earth System Model Version 2 (CESM2), *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001916, <https://doi.org/10.1029/2019MS001916>, 2020.
- 390 Davison, A. C. and Hinkley, D. V.: *Bootstrap Methods and their Application*, Cambridge Series in Statistical and Probabilistic Mathematics, Cambridge University Press, <https://doi.org/10.1017/CBO9780511802843>, 1997.
- Döscher, R., Acosta, M., Alessandri, A., Anthoni, P., Arneth, A., Arsouze, T., Bergmann, T., Bernadello, R., Bousetta, S., Caron, L.-P., Carver, G., Castrillo, M., Catalano, F., Cvijanovic, I., Davini, P., Dekker, E., Doblus-Reyes, F. J., Docquier, D., Echevarria, P., Fladrich, U., Fuentes-Franco, R., Gröger, M., v. Hardenberg, J., Hieronymus, J., Karami, M. P., Keskinen, J.-P., Koenig, T., Makkonen, R., Massonnet,  
395 F., Ménégos, M., Miller, P. A., Moreno-Chamarro, E., Nieradzic, L., van Noije, T., Nolan, P., O'Donnell, D., Ollinaho, P., van den Oord, G., Ortega, P., Prims, O. T., Ramos, A., Reerink, T., Rousset, C., Ruprich-Robert, Y., Le Sager, P., Schmith, T., Schrödner, R., Serva, F., Sicardi, V., Sloth Madsen, M., Smith, B., Tian, T., Tourigny, E., Uotila, P., Vancoppenolle, M., Wang, S., Wärlind, D., Willén, U., Wyser, K., Yang, S., Yepes-Arbós, X., and Zhang, Q.: The EC-Earth3 Earth System Model for the Climate Model Intercomparison Project 6, *Geoscientific Model Development Discussions*, 2021, 1–90, <https://doi.org/10.5194/gmd-2020-446>, 2021.
- 400 Dunne, J. P., Horowitz, L. W., Adcroft, A. J., Ginoux, P., Held, I. M., John, J. G., Krasting, J. P., Malyshev, S., Naik, V., Paulot, F., Shevliakova, E., Stock, C. A., Zadeh, N., Balaji, V., Blanton, C., Dunne, K. A., Dupuis, C., Durachta, J., Dussin, R., Gauthier, P. P. G., Griffies, S. M., Guo, H., Hallberg, R. W., Harrison, M., He, J., Hurlin, W., McHugh, C., Menzel, R., Milly, P. C. D., Nikonov, S., Paynter, D. J., Ploshay, J., Radhakrishnan, A., Rand, K., Reichl, B. G., Robinson, T., Schwarzkopf, D. M., Sentman, L. T., Underwood, S., Vahlenkamp, H., Winton,



- M., Wittenberg, A. T., Wyman, B., Zeng, Y., and Zhao, M.: The GFDL Earth System Model Version 4.1 (GFDL-ESM 4.1): Overall  
405 Coupled Model Description and Simulation Characteristics, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS002015,  
<https://doi.org/10.1029/2019MS002015>, 2020.
- Ebita, A., Kobayashi, S., Ota, Y., Moriya, M., Kumabe, R., Onogi, K., Harada, Y., Yasui, S., Miyaoka, K., Takahashi, K., Kamahori, H.,  
Kobayashi, C., Endo, H., Soma, M., Oikawa, Y., and Ishimizu, T.: The Japanese 55-year Reanalysis JRA-55: An Interim Report, *SOLA*,  
7, 149–152, <https://doi.org/10.2151/sola.2011-038>, 2011.
- 410 Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model  
Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*, 9, 1937–1958,  
<https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- Forster, P., Storelvmo, T., Armour, K., Collins, W., Dufresne, J.-L., Frame, D., Lunt, D., Mauritsen, T., Palmer, M., Watanabe, M., Wild, M.,  
and Zhang, H.: The Earth’s Energy Budget, Climate Feedbacks, and Climate Sensitivity, in: *Climate Change 2021: The Physical Science*  
415 *Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-  
Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E.  
Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]*, edited by van Dijk, W. R., pp. 923–1054,  
Cambridge University Press, Cambridge, <https://doi.org/10.1017/9781009157896.009>, 2021.
- García-García, A., Cuesta-Valero, F. J., Miralles, D., Mahecha, M., Quaas, J., Reichstein, M., Zscheischler, J., and Peng, J.: Soil heat extremes  
420 can outpace air temperature extremes, *Nature Climate Change*, pp. 1–5, <https://doi.org/10.1038/s41558-023-01812-3>, 2023.
- Gottelman, A., Mills, M. J., Kinnison, D. E., Garcia, R. R., Smith, A. K., Marsh, D. R., Tilmes, S., Vitt, F., Bardeen, C. G., McInerney, J., Liu,  
H.-L., Solomon, S. C., Polvani, L. M., Emmons, L. K., Lamarque, J.-F., Richter, J. H., Glanville, A. S., Bacmeister, J. T., Phillips, A. S.,  
Neale, R. B., Simpson, I. R., DuVivier, A. K., Hodzic, A., and Randel, W. J.: The Whole Atmosphere Community Climate Model Version  
6 (WACCM6), *Journal of Geophysical Research: Atmospheres*, 124, 12 380–12 403, <https://doi.org/10.1029/2019JD030943>, 2019.
- 425 Golaz, J.-C., Caldwell, P. M., Van Roekel, L. P., Petersen, M. R., Tang, Q., Wolfe, J. D., Abeshu, G., Anantharaj, V., Asay-Davis, X. S.,  
Bader, D. C., Baldwin, S. A., Bisht, G., Bogenschutz, P. A., Branstetter, M., Brunke, M. A., Brus, S. R., Burrows, S. M., Cameron-Smith,  
P. J., Donahue, A. S., Deakin, M., Easter, R. C., Evans, K. J., Feng, Y., Flanner, M., Foucar, J. G., Fyke, J. G., Griffin, B. M., Hannay, C.,  
Harrop, B. E., Hoffman, M. J., Hunke, E. C., Jacob, R. L., Jacobsen, D. W., Jeffery, N., Jones, P. W., Keen, N. D., Klein, S. A., Larson,  
V. E., Leung, L. R., Li, H.-Y., Lin, W., Lipscomb, W. H., Ma, P.-L., Mahajan, S., Maltrud, M. E., Mamatjanov, A., McClean, J. L., McCoy,  
430 R. B., Neale, R. B., Price, S. F., Qian, Y., Rasch, P. J., Reeves Eyre, J. E. J., Riley, W. J., Ringler, T. D., Roberts, A. F., Roesler, E. L.,  
Salinger, A. G., Shaheen, Z., Shi, X., Singh, B., Tang, J., Taylor, M. A., Thornton, P. E., Turner, A. K., Veneziani, M., Wan, H., Wang, H.,  
Wang, S., Williams, D. N., Wolfram, P. J., Worley, P. H., Xie, S., Yang, Y., Yoon, J.-H., Zelinka, M. D., Zender, C. S., Zeng, X., Zhang, C.,  
Zhang, K., Zhang, Y., Zheng, X., Zhou, T., and Zhu, Q.: The DOE E3SM Coupled Model Version 1: Overview and Evaluation at Standard  
Resolution, *Journal of Advances in Modeling Earth Systems*, 11, 2089–2129, <https://doi.org/10.1029/2018MS001603>, 2019.
- 435 González-Rouco, J. F., Beltrami, H., Zorita, E., and von Storch, H.: Simulation and inversion of borehole temperature profiles in surrogate  
climates: Spatial distribution and surface coupling, *Geophysical Research Letters*, 33, <https://doi.org/10.1029/2005GL024693>, 2006.
- González-Rouco, J. F., Beltrami, H., Zorita, E., and Stevens, M. B.: Borehole climatology: a discussion based on contributions from climate  
modeling, *Climate of the Past*, 5, 97–127, <https://doi.org/10.5194/cp-5-97-2009>, 2009.
- González-Rouco, J. F., Steinert, N. J., García-Bustamante, E., Hagemann, S., de Vrese, P., Jungclaus, J. H., Lorenz, S. J., Melo-Aguilar, C.,  
440 García-Pereira, F., and Navarro, J.: Increasing the Depth of a Land Surface Model. Part I: Impacts on the Subsurface Thermal Regime and  
Energy Storage, *Journal of Hydrometeorology*, 22, 3211 – 3230, <https://doi.org/10.1175/JHM-D-21-0024.1>, 2021.



- Gueugneau, V., Kelfoun, K., Charbonnier, S., Germa, A., and Carazzo, G.: Dynamics and Impacts of the May 8th, 1902 Pyroclastic Current at Mount Pelée (Martinique): New Insights From Numerical Modeling, *Frontiers in Earth Science*, 8, <https://doi.org/10.3389/feart.2020.00279>, 2020.
- 445 Gulev, S., Thorne, P., Ahn, J., Dentener, F., Domingues, C., Gerland, S., Gong, D., Kaufman, D., Nnamchi, H., Quaas, J., Rivera, J., Sathyendranath, S., Smith, S., Trewin, B., von Schuckmann, K., and Vose, R.: Changing State of the Climate System, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)], edited by van Dijk, 450 W. R., pp. 287–422, Cambridge University Press, Cambridge, <https://doi.org/10.1017/9781009157896.004>, 2021.
- Hajima, T., Watanabe, M., Yamamoto, A., Tatebe, H., Noguchi, M. A., Abe, M., Ohgaito, R., Ito, A., Yamazaki, D., Okajima, H., Ito, A., Takata, K., Ogochi, K., Watanabe, S., and Kawamiya, M.: Development of the MIROC-ES2L Earth system model and the evaluation of biogeochemical processes and feedbacks, *Geoscientific Model Development*, 13, 2197–2244, <https://doi.org/10.5194/gmd-13-2197-2020>, 2020.
- 455 Hansen, J. E., Wang, W.-C., and Lacis, A. A.: Mount Agung Eruption Provides Test of a Global Climatic Perturbation, *Science*, 199, 1065–1068, <https://doi.org/10.1126/science.199.4333.1065>, 1978.
- HE, B., YU, Y., BAO, Q., LIN, P., LIU, H., LI, J., WANG, L., LIU, Y., WU, G., CHEN, K., GUO, Y., ZHAO, S., ZHANG, X., SONG, M., and XIE, J.: CAS FGOALS-f3-L model dataset descriptions for CMIP6 DECK experiments, *Atmospheric and Oceanic Science Letters*, 13, 582–588, <https://doi.org/10.1080/16742834.2020.1778419>, 2020.
- 460 Held, I. M., Guo, H., Adcroft, A., Dunne, J. P., Horowitz, L. W., Krasting, J., Shevliakova, E., Winton, M., Zhao, M., Bushuk, M., Wittenberg, A. T., Wyman, B., Xiang, B., Zhang, R., Anderson, W., Balaji, V., Donner, L., Dunne, K., Durachta, J., Gauthier, P. P. G., Ginoux, P., Golaz, J.-C., Griffies, S. M., Hallberg, R., Harris, L., Harrison, M., Hurlin, W., John, J., Lin, P., Lin, S.-J., Malyshev, S., Menzel, R., Milly, P. C. D., Ming, Y., Naik, V., Paynter, D., Paulot, F., Rammasswamy, V., Reichl, B., Robinson, T., Rosati, A., Seman, C., Silvers, L. G., Underwood, S., and Zadeh, N.: Structure and Performance of GFDL's CM4.0 Climate Model, *Journal of Advances in Modeling Earth* 465 *Systems*, 11, 3691–3727, <https://doi.org/10.1029/2019MS001829>, 2019.
- Hermoso de Mendoza, I., Beltrami, H., MacDougall, A. H., and Mareschal, J.-C.: Lower boundary conditions in land surface models – effects on the permafrost and the carbon pools: a case study with CLM4.5, *Geoscientific Model Development*, 13, 1663–1683, <https://doi.org/10.5194/gmd-13-1663-2020>, 2020.
- Huang, S.: 1851–2004 annual heat budget of the continental landmasses, *Geophysical Research Letters*, 33, 470 <https://doi.org/10.1029/2005GL025300>, 2006.
- Huang, S., Pollack, H., and Shen, P.: Temperature trends over the five past centuries reconstructed from borehole temperature, *Nature*, 403, 756–8, <https://doi.org/10.1038/35001556>, 2000.
- Jungclaus, J. H., Bard, E., Baroni, M., Braconnot, P., Cao, J., Chini, L. P., Egorova, T., Evans, M., González-Rouco, J. F., Goosse, H., Hurrell, G. C., Joos, F., Kaplan, J. O., Khodri, M., Klein Goldewijk, K., Krivova, N., LeGrande, A. N., Lorenz, S. J., Luterbacher, J., Man, 475 W., Maycock, A. C., Meinshausen, M., Moberg, A., Muscheler, R., Nehrbass-Ahles, C., Otto-Bliesner, B. I., Phipps, S. J., Pongratz, J., Rozanov, E., Schmidt, G. A., Schmidt, H., Schmutz, W., Schurer, A., Shapiro, A. I., Sigl, M., Smerdon, J. E., Solanki, S. K., Timmreck, C., Toohey, M., Usoskin, I. G., Wagner, S., Wu, C.-J., Yeo, K. L., Zanchettin, D., Zhang, Q., and Zorita, E.: The PMIP4 contribution to CMIP6 – Part 3: The last millennium, scientific objective, and experimental design for the PMIP4 *past1000* simulations, *Geoscientific Model Development*, 10, 4005–4033, <https://doi.org/10.5194/gmd-10-4005-2017>, 2017.





- 480 Kageyama, M., Braconnot, P., Harrison, S. P., Haywood, A. M., Jungclaus, J. H., Otto-Bliesner, B. L., Peterschmitt, J.-Y., Abe-Ouchi, A., Albani, S., Bartlein, P. J., Brierley, C., Crucifix, M., Dolan, A., Fernandez-Donado, L., Fischer, H., Hopcroft, P. O., Ivanovic, R. F., Lambert, F., Lunt, D. J., Mahowald, N. M., Peltier, W. R., Phipps, S. J., Roche, D. M., Schmidt, G. A., Tarasov, L., Valdes, P. J., Zhang, Q., and Zhou, T.: The PMIP4 contribution to CMIP6 – Part 1: Overview and over-arching analysis plan, *Geoscientific Model Development*, 11, 1033–1057, <https://doi.org/10.5194/gmd-11-1033-2018>, 2018.
- 485 Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Leetmaa, A., Reynolds, R., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-Year Reanalysis Project, *Bulletin of the American Meteorological Society*, 77, 437 – 472, [https://doi.org/10.1175/1520-0477\(1996\)077<0437:TNYRP>2.0.CO;2](https://doi.org/10.1175/1520-0477(1996)077<0437:TNYRP>2.0.CO;2), 1996.
- Karl, T. R., Arguez, A., Huang, B., Lawrimore, J. H., McMahon, J. R., Menne, M. J., Peterson, T. C., Vose, R. S., and Zhang, H.-M.: Possible  
490 artifacts of data biases in the recent global surface warming hiatus, *Science*, 348, 1469–1472, <https://doi.org/10.1126/science.aaa5632>, 2015.
- Kelley, M., Schmidt, G. A., Nazarenko, L. S., Bauer, S. E., Ruedy, R., Russell, G. L., Ackerman, A. S., Aleinov, I., Bauer, M., Bleck, R., Canuto, V., Cesana, G., Cheng, Y., Clune, T. L., Cook, B. I., Cruz, C. A., Del Genio, A. D., Elsaesser, G. S., Faluvegi, G., Kiang, N. Y., Kim, D., Lacis, A. A., Leboissetier, A., LeGrande, A. N., Lo, K. K., Marshall, J., Matthews, E. E., McDermid, S., Mezzuman, K., Miller,  
495 R. L., Murray, L. T., Oinas, V., Orbe, C., García-Pando, C. P., Perlwitz, J. P., Puma, M. J., Rind, D., Romanou, A., Shindell, D. T., Sun, S., Tausnev, N., Tsigaridis, K., Tselioudis, G., Weng, E., Wu, J., and Yao, M.-S.: GISS-E2.1: Configurations and Climatology, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS002 025, <https://doi.org/10.1029/2019MS002025>, 2020.
- Krakauer, N. Y., Puma, M. J., and Cook, B. I.: Impacts of soil–aquifer heat and water fluxes on simulated global climate, *Hydrology and Earth System Sciences*, 17, 1963–1974, <https://doi.org/10.5194/hess-17-1963-2013>, 2013.
- 500 Laloyaux, P., de Boissesson, E., Balmaseda, M., Bidlot, J.-R., Broennimann, S., Buizza, R., Dalhgren, P., Dee, D., Haimberger, L., Hersbach, H., Kosaka, Y., Martin, M., Poli, P., Rayner, N., Rustemeier, E., and Schepers, D.: CERA-20C: A Coupled Reanalysis of the Twentieth Century, *Journal of Advances in Modeling Earth Systems*, 10, 1172–1195, <https://doi.org/10.1029/2018MS001273>, 2018.
- Lawrence, D. M., Fisher, R. A., Koven, C. D., Oleson, K. W., Swenson, S. C., Bonan, G., Collier, N., Ghimire, B., van Kampenhou, L., Kennedy, D., Kluzek, E., Lawrence, P. J., Li, F., Li, H., Lombardozzi, D., Riley, W. J., Sacks, W. J., Shi, M., Vertenstein, M., Wieder,  
505 W. R., Xu, C., Ali, A. A., Badger, A. M., Bisht, G., van den Broeke, M., Brunke, M. A., Burns, S. P., Buzan, J., Clark, M., Craig, A., Dahlin, K., Drewniak, B., Fisher, J. B., Flanner, M., Fox, A. M., Gentine, P., Hoffman, F., Keppel-Aleks, G., Knox, R., Kumar, S., Lenaerts, J., Leung, L. R., Lipscomb, W. H., Lu, Y., Pandey, A., Pelletier, J. D., Perket, J., Randerson, J. T., Ricciuto, D. M., Sanderson, B. M., Slater, A., Subin, Z. M., Tang, J., Thomas, R. Q., Val Martin, M., and Zeng, X.: The Community Land Model Version 5: Description of New Features, Benchmarking, and Impact of Forcing Uncertainty, *Journal of Advances in Modeling Earth Systems*, 11, 4245–4287,  
510 <https://doi.org/https://doi.org/10.1029/2018MS001583>, 2019.
- Lee, J., Kim, J., Sun, M.-A., Kim, B.-H., Moon, H., Sung, H. M., Kim, J., and Byun, Y.-H.: Evaluation of the Korea Meteorological Administration Advanced Community Earth-System model (K-ACE), *Asia-Pacific Journal of Atmospheric Sciences*, 56, <https://doi.org/10.1007/s13143-019-00144-7>, 2019.
- Lenssen, N. J. L., Schmidt, G. A., Hansen, J. E., Menne, M. J., Persin, A., Ruedy, R., and Zyss, D.: Improvements in the GISTEMP  
515 Uncertainty Model, *Journal of Geophysical Research: Atmospheres*, 124, 6307–6326, <https://doi.org/10.1029/2018JD029522>, 2019.



- Levitus, S., Antonov, J. I., Boyer, T. P., Baranova, O. K., Garcia, H. E., Locarnini, R. A., Mishonov, A. V., Reagan, J. R., Seidov, D., Yarosh, E. S., and Zweng, M. M.: World ocean heat content and thermocline sea level change (0–2000 m), 1955–2010, *Geophysical Research Letters*, 39, <https://doi.org/10.1029/2012GL051106>, 2012.
- Li, L., Yu, Y., Tang, Y., Lin, P., Xie, J., Song, M., Dong, L., Zhou, T., Liu, L., Wang, L., Pu, Y., Chen, X., Chen, L., Xie, Z., Liu, H., Zhang, L., Huang, X., Feng, T., Zheng, W., Xia, K., Liu, H., Liu, J., Wang, Y., Wang, L., Jia, B., Xie, F., Wang, B., Zhao, S., Yu, Z., Zhao, B., and Wei, J.: The Flexible Global Ocean-Atmosphere-Land System Model Grid-Point Version 3 (FGOALS-g3): Description and Evaluation, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS002012, <https://doi.org/10.1029/2019MS002012>, 2020.
- Lovato, T., Peano, D., Butenschön, M., Materia, S., Iovino, D., Scoccimarro, E., Fogli, P. G., Cherchi, A., Bellucci, A., Gualdi, S., Masina, S., and Navarra, A.: CMIP6 Simulations With the CMCC Earth System Model (CMCC-ESM2), *Journal of Advances in Modeling Earth Systems*, 14, e2021MS002814, <https://doi.org/10.1029/2021MS002814>, 2022.
- MacDougall, A. H., González-Rouco, J. F., Stevens, M. B., and Beltrami, H.: Quantification of subsurface heat storage in a GCM simulation, *Geophysical Research Letters*, 35, <https://doi.org/10.1029/2008GL034639>, 2008.
- Mareschal, J. and Beltrami, H.: Evidence for recent warming from perturbed geothermal gradients: examples from eastern Canada, *Climate Dynamics*, pp. 135–143, <https://doi.org/10.1007/BF00193525>, 1992.
- Mauritsen, T. and Roeckner, E.: Tuning the MPI-ESM1.2 Global Climate Model to Improve the Match With Instrumental Record Warming by Lowering Its Climate Sensitivity, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS002037, <https://doi.org/10.1029/2019MS002037>, 2020.
- Mauritsen, T., Bader, J., Becker, T., Behrens, J., Bittner, M., Brokopf, R., Brovkin, V., Claussen, M., Crueger, T., Esch, M., Fast, I., Fiedler, S., Fläschner, D., Gayler, V., Giorgetta, M., Goll, D. S., Haak, H., Hagemann, S., Hedemann, C., Hohenegger, C., Ilyina, T., Jahns, T., Jimenez-de-la Cuesta, D., Jungclaus, J., Kleinen, T., Kloster, S., Kracher, D., Kinne, S., Kleberg, D., Lasslop, G., Kornbluh, L., Marotzke, J., Matei, D., Meraner, K., Mikolajewicz, U., Modali, K., Möbis, B., Müller, W. A., Nabel, J. E. M. S., Nam, C. C. W., Notz, D., Nyawira, S.-S., Paulsen, H., Peters, K., Pincus, R., Pohlmann, H., Pongratz, J., Popp, M., Raddatz, T. J., Rast, S., Redler, R., Reick, C. H., Rohrschneider, T., Schemann, V., Schmidt, H., Schnur, R., Schulzweida, U., Six, K. D., Stein, L., Stemmler, I., Stevens, B., von Storch, J.-S., Tian, F., Voigt, A., Vrese, P., Wieners, K.-H., Wilkenskjaeld, S., Winkler, A., and Roeckner, E.: Developments in the MPI-M Earth System Model version 1.2 (MPI-ESM1.2) and Its Response to Increasing CO<sub>2</sub>, *Journal of Advances in Modeling Earth Systems*, 11, 998–1038, <https://doi.org/10.1029/2018MS001400>, 2019.
- Mccormick, M., Thomason, L., and Trepte, C.: Atmospheric effects of the Mt Pinatubo eruption, *Nature*, 373, 399–404, <https://doi.org/10.1038/373399a0>, 1995.
- Melo-Aguilar, C., González-Rouco, J. F., García-Bustamante, E., Navarro-Montesinos, J., and Steinert, N.: Influence of radiative forcing factors on ground–air temperature coupling during the last millennium: implications for borehole climatology, *Climate of the Past*, 14, 1583–1606, <https://doi.org/10.5194/cp-14-1583-2018>, 2018.
- Melo-Aguilar, C., González-Rouco, J. F., García-Bustamante, E., Steinert, N., Jungclaus, J. H., Navarro, J., and Roldán-Gómez, P. J.: Methodological and physical biases in global to subcontinental borehole temperature reconstructions: an assessment from a pseudo-proxy perspective, *Climate of the Past*, 16, 453–474, <https://doi.org/10.5194/cp-16-453-2020>, 2020.
- Miller, G. H., Geirsdóttir, A., Zhong, Y., Larsen, D. J., Otto-Bliesner, B. L., Holland, M. M., Bailey, D. A., Refsnider, K. A., Lehman, S. J., Southon, J. R., Anderson, C., Björnsson, H., and Thordarson, T.: Abrupt onset of the Little Ice Age triggered by volcanism and sustained by sea-ice/ocean feedbacks, *Geophysical Research Letters*, 39, <https://doi.org/10.1029/2011GL050168>, 2012.



- Mottaghy, D. and Rath, V.: Latent heat effects in subsurface heat transport modelling and their impact on palaeotemperature reconstructions, *Geophysical Journal International*, 164, 236–245, <https://doi.org/10.1111/j.1365-246X.2005.02843.x>, 2006.
- 555 Muñoz Sabater, J., Dutra, E., Agustí-Panareda, A., Albergel, C., Arduini, G., Balsamo, G., Boussetta, S., Choulga, M., Harrigan, S., Hersbach, H., Martens, B., Miralles, D. G., Piles, M., Rodríguez-Fernández, N. J., Zsoter, E., Buontempo, C., and Thépaut, J.-N.: ERA5-Land: a state-of-the-art global reanalysis dataset for land applications, *Earth System Science Data*, 13, 4349–4383, <https://doi.org/10.5194/essd-13-4349-2021>, 2021.
- Osborn, T. J., Jones, P. D., Lister, D. H., Morice, C. P., Simpson, I. R., Winn, J. P., Hogan, E., and Harris, I. C.: Land Surface Air Temperature Variations Across the Globe Updated to 2019: The CRUTEM5 Data Set, *Journal of Geophysical Research: Atmospheres*, 126, e2019JD032352, <https://doi.org/10.1029/2019JD032352>, 2021.
- 560 Poli, P., Hersbach, H., Dee, D. P., Berrisford, P., Simmons, A. J., Vitart, F., Laloyaux, P., Tan, D. G. H., Peubey, C., Thépaut, J.-N., Trémolet, Y., Hólm, E. V., Bonavita, M., Isaksen, L., and Fisher, M.: ERA-20C: An Atmospheric Reanalysis of the Twentieth Century, *Journal of Climate*, 29, 4083 – 4097, <https://doi.org/10.1175/JCLI-D-15-0556.1>, 2016.
- 565 Pries, C. E. H., Castanha, C., Porras, R. C., and Torn, M. S.: The whole-soil carbon flux in response to warming, *Science*, 355, 1420–1423, <https://doi.org/10.1126/science.aal1319>, 2017.
- Reick, C. H., Gayler, V., Goll, D., Hagemann, S., Heidkamp, M., Nabel, J. E. M. S., Raddatz, T., Roeckner, E., Schnur, R., and Wilkenskield, S.: JSBACH 3 - The land component of the MPI Earth System Model: documentation of version 3.2, Hamburg: MPI für Meteorologie, <https://doi.org/10.17617/2.3279802>, 2021.
- 570 Rind, D., Orbe, C., Jonas, J., Nazarenko, L., Zhou, T., Kelley, M., Lacis, A., Shindell, D., Faluvegi, G., Romanou, A., Russell, G., Tausnev, N., Bauer, M., and Schmidt, G.: GISS Model E2.2: A Climate Model Optimized for the Middle Atmosphere—Model Structure, Climatology, Variability, and Climate Sensitivity, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD032204, <https://doi.org/10.1029/2019JD032204>, 2020.
- Rohde, R. A. and Hausfather, Z.: The Berkeley Earth Land/Ocean Temperature Record, *Earth System Science Data*, 12, 3469–3479, <https://doi.org/10.5194/essd-12-3469-2020>, 2020.
- 575 Seland, Ø., Bentsen, M., Oliví, D., Toniazzo, T., Gjermundsen, A., Graff, L. S., Debernard, J. B., Gupta, A. K., He, Y.-C., Kirkevåg, A., Schwinger, J., Tjiputra, J., Aas, K. S., Bethke, I., Fan, Y., Griesfeller, J., Grini, A., Guo, C., Ilicak, M., Karset, I. H. H., Landgren, O., Liakka, J., Moseid, K. O., Nummelin, A., Spensberger, C., Tang, H., Zhang, Z., Heinze, C., Iversen, T., and Schulz, M.: Overview of the Norwegian Earth System Model (NorESM2) and key climate response of CMIP6 DECK, historical, and scenario simulations, *Geoscientific Model Development*, 13, 6165–6200, <https://doi.org/10.5194/gmd-13-6165-2020>, 2020.
- 580 Self, S. and Rampino, M. R.: The 1883 eruption of Krakatau, *Nature*, 294, 699–704, <https://doi.org/10.1038/294699a0>, 1981.
- Sellar, A. A., Jones, C. G., Mulcahy, J. P., Tang, Y., Yool, A., Wiltshire, A., O'Connor, F. M., Stringer, M., Hill, R., Palmieri, J., Woodward, S., de Mora, L., Kuhlbrodt, T., Rumbold, S. T., Kelley, D. I., Ellis, R., Johnson, C. E., Walton, J., Abraham, N. L., Andrews, M. B., Andrews, T., Archibald, A. T., Berthou, S., Burke, E., Blockley, E., Carslaw, K., Dalvi, M., Edwards, J., Folberth, G. A., Gedney, N., Griffiths, P. T., Harper, A. B., Hendry, M. A., Hewitt, A. J., Johnson, B., Jones, A., Jones, C. D., Keeble, J., Liddicoat, S., Morgenstern, O., Parker, R. J., Predoi, V., Robertson, E., Siahann, A., Smith, R. S., Swaminathan, R., Woodhouse, M. T., Zeng, G., and Zerroukat, M.: UKESM1: Description and Evaluation of the U.K. Earth System Model, *Journal of Advances in Modeling Earth Systems*, 11, 4513–4558, <https://doi.org/10.1029/2019MS001739>, 2019.
- 590 Slivinski, L. C., Compo, G. P., Whitaker, J. S., Sardeshmukh, P. D., Giese, B. S., McColl, C., Allan, R., Yin, X., Vose, R., Titchner, H., Kennedy, J., Spencer, L. J., Ashcroft, L., Brönnimann, S., Brunet, M., Camuffo, D., Cornes, R., Cram, T. A., Crouthamel, R., Domínguez-



- 595 Castro, F., Freeman, J. E., Gergis, J., Hawkins, E., Jones, P. D., Jourdain, S., Kaplan, A., Kubota, H., Blancq, F. L., Lee, T.-C., Lorrey, A., Luterbacher, J., Maugeri, M., Mock, C. J., Moore, G. K., Przybylak, R., Pudmenzky, C., Reason, C., Slonosky, V. C., Smith, C. A., Tinz, B., Trewin, B., Valente, M. A., Wang, X. L., Wilkinson, C., Wood, K., and Wyszyński, P.: Towards a more reliable historical reanalysis: Improvements for version 3 of the Twentieth Century Reanalysis system, *Quarterly Journal of the Royal Meteorological Society*, 145, 2876–2908, <https://doi.org/10.1002/qj.3598>, 2019.
- Smerdon, J. E. and Stieglitz, M.: Simulating heat transport of harmonic temperature signals in the Earth’s shallow subsurface: Lower-boundary sensitivities, *Geophysical Research Letters*, 33, <https://doi.org/10.1029/2006GL026816>, 2006.
- 600 Soong, J. L., Castanha, C., Pries, C. E. H., Ofiti, N., Porras, R. C., Riley, W. J., Schmidt, M. W., and Torn, M. S.: Five years of whole-soil warming led to loss of subsoil carbon stocks and increased CO<sub>2</sub> efflux, *Science Advances*, 7, eabd1343, <https://doi.org/10.1126/sciadv.abd1343>, 2021.
- Steinert, N. J., González-Rouco, J. F., Melo-Aguilar, C. A., García-Pereira, F., García-Bustamante, E., de Vrese, P., Alexeev, V., Jungclaus, J. H., Lorenz, S. J., and Hagemann, S.: Agreement of Analytical and Simulation-Based Estimates of the Required Land Depth in Climate Models, *Geophysical Research Letters*, 48, e2021GL094273, <https://doi.org/10.1029/2021GL094273>, 2021a.
- 605 Steinert, N. J., González-Rouco, J. F., de Vrese, P., García-Bustamante, E., Hagemann, S., Melo-Aguilar, C. A., Jungclaus, J. H., and Lorenz, S. J.: Increasing the depth of a Land Surface Model. Part II: Sensitivity to improved coupling between soil hydrology and thermodynamics and associated permafrost response., *Journal of Hydrometeorology*, 22, 3231 – 3254, <https://doi.org/10.1175/JHM-D-21-0023.1>, 2021b.
- Stevens, B., Smerdon, J. E., González-Rouco, J. F., Stieglitz, M., and Beltrami, H.: Effects of bottom boundary placement on subsurface heat storage: Implications for climate model simulations, *Geophysical Research Letters*, 34, <https://doi.org/10.1029/2006GL028546>, 2007.
- 610 Stevens, B., Giorgetta, M., Esch, M., Mauritsen, T., Crueger, T., Rast, S., Salzmann, M., Schmidt, H., Bader, J., Block, K., Brokopf, R., Fast, I., Kinne, S., Kornblueh, L., Lohmann, U., Pincus, R., Reichler, T., and Roeckner, E.: Atmospheric component of the MPI-M Earth System Model: ECHAM6, *Journal of Advances in Modeling Earth Systems*, 5, 146–172, <https://doi.org/10.1002/jame.20015>, 2013.
- Swart, N. C., Cole, J. N. S., Kharin, V. V., Lazare, M., Scinocca, J. F., Gillett, N. P., Anstey, J., Arora, V., Christian, J. R., Hanna, S., Jiao, Y., Lee, W. G., Majaess, F., Saenko, O. A., Seiler, C., Seinen, C., Shao, A., Sigmond, M., Solheim, L., von Salzen, K., Yang, D., and Winter, B.: The Canadian Earth System Model version 5 (CanESM5.0.3), *Geoscientific Model Development*, 12, 4823–4873, <https://doi.org/10.5194/gmd-12-4823-2019>, 2019.
- 615 Sférian, R., Nabat, P., Michou, M., Saint-Martin, D., Voldoire, A., Colin, J., Decharme, B., Delire, C., Berthet, S., Chevallier, M., Sénési, S., Franchisteguy, L., Vial, J., Mallet, M., Joetzjer, E., Geoffroy, O., Guérémy, J.-F., Moine, M.-P., Msadek, R., Ribes, A., Rocher, M., Roehrig, R., Salas-y Mélia, D., Sanchez, E., Terray, L., Valcke, S., Waldman, R., Aumont, O., Bopp, L., Deshayes, J., Éthé, C., and Madec, G.: Evaluation of CNRM Earth System Model, CNRM-ESM2-1: Role of Earth System Processes in Present-Day and Future Climate, *Journal of Advances in Modeling Earth Systems*, 11, 4182–4227, <https://doi.org/10.1029/2019MS001791>, 2019.
- 620 Tardif, R., Hakim, G. J., Perkins, W. A., Horlick, K. A., Erb, M. P., Emile-Geay, J., Anderson, D. M., Steig, E. J., and Noone, D.: Last Millennium Reanalysis with an expanded proxy database and seasonal proxy modeling, *Climate of the Past*, 15, 1251–1273, <https://doi.org/10.5194/cp-15-1251-2019>, 2019.
- 625 Tatebe, H., Ogura, T., Nitta, T., Komuro, Y., Ogochi, K., Takemura, T., Sudo, K., Sekiguchi, M., Abe, M., Saito, F., Chikira, M., Watanabe, S., Mori, M., Hirota, N., Kawatani, Y., Mochizuki, T., Yoshimura, K., Takata, K., O’ishi, R., Yamazaki, D., Suzuki, T., Kurogi, M., Kataoka, T., Watanabe, M., and Kimoto, M.: Description and basic evaluation of simulated mean state, internal variability, and climate sensitivity in MIROC6, *Geoscientific Model Development*, 12, 2727–2765, <https://doi.org/10.5194/gmd-12-2727-2019>, 2019.



- Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An Overview of CMIP5 and the Experiment Design, *Bulletin of the American Meteorological Society*, 93, 485 – 498, <https://doi.org/10.1175/BAMS-D-11-00094.1>, 2012.
- 630 van Dijk, E., Jungclaus, J., Lorenz, S., Timmreck, C., and Krüger, K.: Was there a volcanic-induced long-lasting cooling over the Northern Hemisphere in the mid-6th–7th century?, *Climate of the Past*, 18, 1601–1623, <https://doi.org/10.5194/cp-18-1601-2022>, 2022.
- Voldoire, A., Saint-Martin, D., Sénési, S., Decharme, B., Alias, A., Chevallier, M., Colin, J., Guérémy, J.-F., Michou, M., Moine, M.-P., Nabat, P., Roehrig, R., Salas y Mélia, D., Sférian, R., Valcke, S., Beau, I., Belamari, S., Berthet, S., Cassou, C., Cattiaux, J., Deshayes, J., Douville, H., Ethé, C., Franchistéguy, L., Geoffroy, O., Lévy, C., Madec, G., Meurdesoif, Y., Msadek, R., Ribes, A., Sanchez-Gomez, E., Terray, L., and Waldman, R.: Evaluation of CMIP6 DECK Experiments With CNRM-CM6-1, *Journal of Advances in Modeling Earth Systems*, 11, 2177–2213, <https://doi.org/10.1029/2019MS001683>, 2019.
- 635 von Schuckmann, K., Cheng, L., Palmer, M. D., Hansen, J., Tassone, C., Aich, V., Adusumilli, S., Beltrami, H., Boyer, T., Cuesta-Valero, F. J., Desbruyères, D., Domingues, C., García-García, A., Gentine, P., Gilson, J., Gorfer, M., Haimberger, L., Ishii, M., Johnson, G. C., Killick, R., King, B. A., Kirchengast, G., Kolodziejczyk, N., Lyman, J., Marzeion, B., Mayer, M., Monier, M., Monselesan, D. P., Purkey, S., Roemmich, D., Schweiger, A., Seneviratne, S. I., Shepherd, A., Slater, D. A., Steiner, A. K., Straneo, F., Timmermans, M.-L., and Wjiffels, S. E.: Heat stored in the Earth system: where does the energy go?, *Earth System Science Data*, 12, 2013–2041, <https://doi.org/10.5194/essd-12-2013-2020>, 2020.
- 640 von Schuckmann, K., Minère, A., Gues, F., Cuesta-Valero, F. J., Kirchengast, G., Adusumilli, S., Straneo, F., Allan, R., Barker, P. M., Beltrami, H., Boyer, T., Cheng, L., Church, J., Desbruyeres, D., Dolman, H., Domingues, C. M., García-García, A., Giglio, D., Gilson, J. E., Gorfer, M., Haimberger, L., Hendricks, S., Hosoda, S., Johnson, G. C., Killick, R., King, B., Kolodziejczyk, N., Korosov, A., Krinner, G., Kuusela, M., Langer, M., Lavergne, T., Lawrence, I., Li, Y., Lyman, J., Marzeion, B., Mayer, M., MacDougall, A. H., McDougall, T., Monselesan, D. P., Nitzbon, J., Ootosaka, I., Peng, J., Purkey, S., Roemmich, D., Sato, K., Sato, K., Savita, A., Schweiger, A., Shepherd, A., Seneviratne, S. I., Simons, L., Slater, D. A., Slater, T., Smith, N., Steiner, A., Suga, T., Szekely, T., Thiery, W., Timmermans, M.-L., Vanderkelen, I., Wjiffels, S. E., Wu, T., and Zemp, M.: Heat stored in the Earth system 1960–2020: Where does the energy go?, *Earth System Science Data Discussions*, 2022, 1–55, <https://doi.org/10.5194/essd-2022-239>, 2022.
- 645 Wang, Y.-C., Hsu, H.-H., Chen, C.-A., Tseng, W.-L., Hsu, P.-C., Lin, C.-W., Chen, Y.-L., Jiang, L.-C., Lee, Y.-C., Liang, H.-C., Chang, W.-M., Lee, W.-L., and Shiu, C.-J.: Performance of the Taiwan Earth System Model in Simulating Climate Variability Compared With Observations and CMIP6 Model Simulations, *Journal of Advances in Modeling Earth Systems*, 13, e2020MS002353, <https://doi.org/10.1029/2020MS002353>, 2021.
- 650 Wild, M.: The global energy balance as represented in CMIP6 climate models, *Climate Dynamics*, 55, 553–577, <https://doi.org/10.1007/s00382-020-05282-7>, 2020.
- Williams, K. D., Copsey, D., Blockley, E. W., Bodas-Salcedo, A., Calvert, D., Comer, R., Davis, P., Graham, T., Hewitt, H. T., Hill, R., Hyder, P., Ineson, S., Johns, T. C., Keen, A. B., Lee, R. W., Megann, A., Milton, S. F., Rae, J. G. L., Roberts, M. J., Scaife, A. A., Schiemann, R., Storkey, D., Thorpe, L., Watterson, I. G., Walters, D. N., West, A., Wood, R. A., Woollings, T., and Xavier, P. K.: The Met Office Global Coupled Model 3.0 and 3.1 (GC3.0 and GC3.1) Configurations, *Journal of Advances in Modeling Earth Systems*, 10, 357–380, <https://doi.org/10.1002/2017MS001115>, 2018.
- 660 Willmott, C. J. and Matsuura, K.: Terrestrial Air Temperature and Precipitation: Monthly and Annual Time Series (1900 - 2017), [https://doi.org/http://climate.geog.udel.edu/climate/html\\_pages/README.ghcn\\_ts2.html](https://doi.org/http://climate.geog.udel.edu/climate/html_pages/README.ghcn_ts2.html), 2018.



- 665 Wu, T., Lu, Y., Fang, Y., Xin, X., Li, L., Li, W., Jie, W., Zhang, J., Liu, Y., Zhang, L., Zhang, F., Zhang, Y., Wu, F., Li, J., Chu, M., Wang, Z.,  
Shi, X., Liu, X., Wei, M., Huang, A., Zhang, Y., and Liu, X.: The Beijing Climate Center Climate System Model (BCC-CSM): the main  
progress from CMIP5 to CMIP6, *Geoscientific Model Development*, 12, 1573–1600, <https://doi.org/10.5194/gmd-12-1573-2019>, 2019.
- 670 Yukimoto, S., Kawai, H., Koshiro, T., Oshima, N., Yoshida, K., Urakawa, S., Tsujino, H., Deushi, M., Tanaka, T., Hosaka, M., Yabu, S.,  
Yoshimura, H., Shindo, E., Mizuta, R., Obata, A., Adachi, Y., and Ishii, M.: The Meteorological Research Institute Earth System Model  
Version 2.0, MRI-ESM2.0: Description and Basic Evaluation of the Physical Component, *Journal of the Meteorological Society of Japan*.  
Ser. II, 97, 931–965, <https://doi.org/10.2151/jmsj.2019-051>, 2019.
- Zhang, H., Zhang, M., Jin, J., Fei, K., Ji, D., Wu, C., Zhu, J., He, J., Chai, Z., Xie, J., Dong, X., Zhang, D., Bi, X., Cao, H., Chen, H., Chen,  
K., Chen, X., Gao, X., Hao, H., Jiang, J., Kong, X., Li, S., Li, Y., Lin, P., Lin, Z., Liu, H., Liu, X., Shi, Y., Song, M., Wang, H., Wang, T.,  
Wang, X., Wang, Z., Wei, Y., Wu, B., Xie, Z., Xu, Y., Yu, Y., Yuan, L., Zeng, Q., Zeng, X., Zhao, S., Zhou, G., and Zhu, J.: Description  
and Climate Simulation Performance of CAS-ESM Version 2, *Journal of Advances in Modeling Earth Systems*, 12, e2020MS002210,  
675 <https://doi.org/10.1029/2020MS002210>, 2020.
- Zhang, H.-M., Huang, B., Lawrimore, J. H., Menne, M. J., and Smith, T. M.: NOAA Global Surface Temperature Dataset (NOAAGlobal-  
Temp), Version 5.0, NOAA National Centers for Environmental Information, <https://doi.org/10.25921/9qth-2p70>, 2019.
- Ziehn, T., Chamberlain, M., Law, R., Lenton, A., Bodman, R., Dix, M., Stevens, L., Wang, Y., and Srbinovsky, J.: The Australian Earth  
System Model: ACCESS-ESM1.5, *Journal of Southern Hemisphere Earth Systems Science*, 70, <https://doi.org/10.1071/ES19035>, 2020.