

# A quantitative assessment of air-sea heat flux trends from ERA5 since 1950 in the North Atlantic basin

Johannes Mayer<sup>1</sup>, Leopold Haimberger<sup>1</sup>, and Michael Mayer<sup>1,2</sup>

<sup>1</sup>Department of Meteorology and Geophysics, University of Vienna, Vienna, Austria

<sup>2</sup>European Centre for Medium-Range Weather Forecasts, Bonn, Germany

**Correspondence:** Johannes Mayer (johannes.mayer@univie.ac.at)

**Abstract.** This work aims to investigate the temporal stability and reliability of trends in air-sea heat fluxes from ERA5 forecasts over the North Atlantic basin for the period 1950–2019. Driving forces of the trends are investigated using analyzed state quantities from ERA5. Estimating trends from reanalysis data can be challenging as changes in the observing system may introduce temporal inconsistencies. To this end, the impact of analysis increments is discussed. For individual sub-regions in the North Atlantic basin, parametrization formulas for latent and sensible heat fluxes are linearized to quantitatively attribute trends to long-term changes in wind speed, moisture, and temperature. Our results suggest good temporal stability and reliability of air-sea heat fluxes from ERA5 forecasts on sub-basin ~~scale~~ scales and below. Regional averages show that trends are largely driven by changes in the skin temperature and atmospheric advection (e.g., of warmer or drier air masses). The influence of ~~climate variability modes~~ modes of climate variability, such as the North Atlantic Oscillation (NAO) and Atlantic Multidecadal Oscillation ~~on the found patterns~~, on the patterns found is discussed as well. Results indicate a significant impact on trends in the Irminger and Labrador ~~Sea-Sea~~ Sea-Sea associated with more positive NAO phases during the past 4 decades. Finally, we use basin-wide trends of air-sea heat fluxes in combination with an observational ocean heat content estimate to provide an energy-budget-based trend estimate of the Atlantic meridional overturning circulation (AMOC). A decrease of area-averaged air-sea heat fluxes in the North Atlantic basin suggests a decline of the AMOC over the study period. However, basin-wide flux trends are deemed partially artificial, as indicated by temporally varying moisture increments. Thus, the exact magnitude of change is uncertain, but its sign appears robust and adds complementary evidence that the AMOC has weakened over the past 70 years.

## 1 Introduction

The North Atlantic Ocean plays a central role for weather and climate in Europe and eastern North America. For instance, the formation of tropical cyclones and severe weather systems along the Gulf Stream and its extension have a significant impact on our economy, agriculture, and society. The North Atlantic Oscillation [NAO; Hurrell (1995); Visbeck et al. (2001)], a periodic change in strength of the Azores High and Icelandic Low, impacts the moisture transport in the northern hemisphere on seasonal timescales and thus influences temperature and precipitation in wide areas of Europe and North America. In addition, the Gulf Stream current in the western North Atlantic is responsible for the poleward transport of oceanic energy that is taken up in tropical latitudes and released to the atmosphere further north via air-sea fluxes. The associated cooling of the waters is

25 required to trigger deep water formation, which is the main driver of the Atlantic meridional overturning circulation [AMOC; Rahmstorf et al. (2015)] and consequently also of the global thermohaline circulation. Previous research used climate models to demonstrate that the AMOC has declined over past decades as a result of anthropogenic global warming, which could have further effects on storms and weather patterns (Fox-Kemper et al., 2021). However, direct observations of the AMOC are limited in time and do not show clear evidence of an externally forced slowdown (Baehr et al., 2007; Roberts et al., 2014; 30 Worthington et al., 2021).

In all these processes, the exchange of energy (and momentum) between [the](#) atmosphere and underlying ocean is of vital importance. Long-term changes in air-sea heat fluxes over the North Atlantic Ocean can thus have a wide range of implications. Consequently, it is of high relevance to accurately estimate air-sea heat flux trends, which also helps to understand numerous aspects of climate variability in the North Atlantic basin. Nonetheless, observation-based estimates are spatially limited and 35 are available only for the past two to three decades (see, e.g., oceansites flux data provided by NOAA, <https://www.pmel.noaa.gov/gtmba/>) making the distinction between anthropogenic changes and natural variability on decadal to multi-decadal timescales demanding. For instance, the Atlantic multidecadal oscillation [AMO; Kerr (2000)] describes a natural variability mode of basin-wide SST anomalies on timescales of 70–80 years so that its long-term effect on air-sea heat fluxes can not be determined adequately from observations.

40 An alternative to observations is given by recent reanalysis data such as the fifth generation global reanalysis data produced by ECMWF [ERA5; Hersbach et al. (2020)] providing global gridded data for more than seven decades thanks to its recent [preliminary](#) back-extension (Bell et al., 2021). Reanalysis data are constructed from past model forecasts constrained by observational data (such as in-situ, satellite, airplanes, or radiosondes) through data assimilation, which warrants optimal combination and reduction of biases. However, changes in the observing system can result in temporal discontinuities and 45 introduce increments between forecasts and analyses that may alter the atmospheric state [moisture, temperature, and wind; see Chiodo and Haimberger (2010) and Mayer et al. (2021)] and consequently also air-sea heat fluxes making climate trend studies with reanalysis data challenging.

Reanalysis data can also be used to indirectly estimate [net](#) air-sea heat fluxes ( $F_s$ ) by evaluating the atmospheric energy budget (Trenberth, 1991; Mayer et al., 2016; Trenberth and Fasullo, 2017; Liu et al., 2017; Mayer et al., 2019, 2021, 2022; 50 Liu et al., 2022). This method does not inherently reduce temporal discontinuities, but allows the application of a global wind field correction (Trenberth, 1991; Fasullo and Trenberth, 2008; Mayer and Haimberger, 2012), which diminishes both artificial noise over high topography and temporal discontinuities introduced by changes in the observing system (Mayer et al., 2021). Although air-sea heat fluxes derived from ERA5 in this way are available only for the period from 1985 onward [[due to the limited availability of observationally constrained TOA fluxes, see Liu et al. \(2020\) and Mayer et al. \(2022\)](#)], they are proven 55 to be temporally relatively stable over the global ocean (around  $1.7 \text{ W m}^{-2}$  mean based on 1985–2018) and thus can serve as reference to test the reliability of other commonly used air-sea heat flux products (Mayer et al., 2021, 2022).

This work aims to investigate the reliability and temporal stability of long-term trends of winter months (December–February) net [surface-air-sea](#) heat fluxes based on ERA5 data over the North Atlantic ocean during 1950–2019. Main drivers of trends in latent and sensible heat fluxes are identified based on analyzed state quantities, and the impact of the assimilation

60 process and climate variability modes, such as [the](#) NAO and AMO, are discussed. Whenever possible, net air-sea heat fluxes from ERA5 are compared with indirect estimates from Mayer et al. (2022). In four individual sub-regions, turbulent air-sea heat flux trends are quantitatively attributed to long-term changes in wind speed, moisture, and temperature using linearized flux parameterization formulae. Finally, we use basin-wide [net](#) air-sea heat fluxes from ERA5 forecasts and an observation-based ocean product to indirectly estimate trends of the AMOC over the past 70 years, and discuss sources of uncertainties and  
65 reliability of the trend estimate.

The data we use in this study are introduced in section 2. Section 3 describes the methodology. Results are presented in section 4 and summarized and discussed in section 5.

## 2 Data

The data we primarily use in this study are from ECMWF's most recent reanalysis ERA5 (Hersbach et al., 2020). ERA5  
70 provides [hourly estimates of](#) a variety of meteorological variables as ~~12-hourly twice-daily forecasts as well as~~ analyzed state quantities [as well as 12-hourly twice-daily forecasts](#) on a Gaussian grid equivalent to 0.25 degree spatial resolution [see Hersbach et al. (2020) for details; [Hersbach et al. \(2023b\)](#)]. Individual components of the net air-sea heat flux (i.e., short-wave and long-wave radiation, and sensible and latent heat flux) are taken as monthly means and are available only as forecasts (~~denoted as model-based~~. [These net air-sea heat fluxes from ERA5 forecasts](#) ) [are denoted as model-based  \$F\_S\$  hereafter](#). We also ~~use~~  
75 ~~single-level atmospheric moisture, temperature, pressure, and~~ [employ temperature and moisture in the lowest model level as well as surface pressure](#), 10 metre wind, [near-surface atmospheric density, and skin temperature](#) fields as monthly means from both analyses and forecasts to compute ~~surface turbulent air-sea~~ heat fluxes using parameterizations as implemented in the Integrated Forecast System [IFS; see ECMWF (2021)], which allows ~~to estimate an estimation of~~ the role of changes in single input ~~parameters variables (moisture, temperature, and wind speed; trends in air density are in general not discussed as they~~  
80 ~~are negligibly small)~~. The 3D horizontal wind fields used to compute the meridional mass stream function are also taken from ERA5 but on pressure levels and [a](#) regular  $0.25 \times 0.25$  grid.

Whenever possible, we compare model-based [heat](#) fluxes with indirectly estimated net ~~surface-air-sea~~ heat fluxes (denoted as inferred ~~surface fluxes~~  $F_S$ ) from Mayer et al. (2022), which are derived from the DEEP-C TOA flux product [see Liu et al. (2020)] and atmospheric energy transports from Mayer et al. (2022b) [see also Mayer et al. (2021) and section 2 in Mayer et al.  
85 (2022) for details of the computation and assessment]. Inferred ~~surface-heat~~ fluxes are provided as monthly averages on a one degree regular grid covering 1985–2020.

Observationally constrained ocean heat content (OHC) data are provided by the Institute of Atmospheric Physics [IAP; Cheng et al. (2017); available at <http://www.ocean.iap.ac.cn/pages/dataService/dataService.html>] on a regular  $1 \times 1^\circ$  grid covering the whole study period 1950–2019. We use the 0–300m OHC monthly data for the correlation and comparison with the  
90 skin temperature from ERA5, and the 0–2000m OHC to indirectly estimate the AMOC trend from the oceanic heat budget. The IAP dataset is constructed based on a modified version of the ensemble optimal interpolation method proposed by Cheng and Zhu (2016). Sampling errors are minimized using observational data and a prior guess from Coupled Model Intercomparison

Project Phase 5 (CMIP5) multimodel simulations. The small sampling error indicates a robust reconstruction of the temperature signal in all ocean basins. Furthermore, the dataset is bias-corrected using [in-situ observations \(CBT and MBT data in-situ observations \(from expendable and mechanical bathythermographs\)\)](#) and thus appears well suited for our evaluations.

In addition, we use mooring-derived and volume-conserving monthly ocean heat transport (OHT) estimates from Tsubouchi et al. (2018) and Tsubouchi et al. (2020) as [the](#) northern choke point of the oceanic heat budget. Tsubouchi et al. (2018) offer transport estimates from the Davis Strait (DS; see <http://metadata.nmdc.no/metadata-api/landingpage/0a2ae0e42ef7af767a920811e83784b>) covering the period 1993–2016. Tsubouchi et al. (2020) provide observations for 2005–2010 from the Fram Strait and Barents Sea Opening (FS and BSO; see <https://doi.pangaea.de/10.1594/PANGAEA.909966>).

### 3 Methods

#### 3.1 Bulk formulas

In this study, sensible and latent heat fluxes are taken from 12-hourly ERA5 forecasts. However, to i) estimate the [relationship between analysis increments and impact of analysis increments on](#) long-term flux trends, and ii) for regression on input variables, we compute fluxes from scratch as described in the following.

Net [surface energy fluxes air-sea heat fluxes \( \$F\_S\$ \)](#) are the sum of radiative and turbulent heat fluxes. Radiative fluxes contain short- and long-wave radiation and are not discussed in detail here. Turbulent heat fluxes are the sum of latent and sensible heat fluxes and can be approximated by the commonly used bulk formulas [see Fairall et al. (2003), Cronin et al. (2019), and ECMWF (2021)], which are written as follows

110

$$F_{LH} = C_Q \rho |U_{ml}| (L_v q_{ml} - L_v q_{sfc}) \quad \text{with} \quad F_{LH} = F_{LH}(\rho, U_{10m}, q_{ml}, q_{sfc}, t), \quad q_{sfc} = q_{sfc}(p_{sfc}, T_{skin}) \quad (1)$$

$$F_{SH} = C_H \rho |U_{ml}| (c_p T_{ml} - c_p T_{skin} + g z_{ml}) \quad \text{with} \quad F_{SH} = F_{SH}(\rho, U_{10m}, T_{ml}, T_{skin}, t) \quad (2)$$

115 where  $C_Q$  and  $C_H$  are non-constant transfer coefficients [see also ECMWF (2021)],  $\rho$  is the air density above ocean,  $L_v$  ( $2.5008 \times 10^6 \text{ J kg}^{-1}$ ) is the latent heat of vaporization,  $c_p$  ( $1004.709 \text{ J kg}^{-1} \text{ K}^{-1}$ ) is the specific heat capacity of dry air,  $g$  ( $9.80665 \text{ m s}^{-2}$ ) is the gravitational acceleration, and  $z_{ml}$  is the height of the lowest model level.  $|U_{ml}|$ ,  $q_{ml}$ , and  $T_{ml}$  are the wind speed, specific humidity, and air temperature at [the](#) lowest model level.  $q_{sfc}$  is the surface saturation humidity, and  $T_{skin}$  is the skin temperature (as used in the IFS instead of the sea surface temperature).  $q_{sfc}$  depends on surface pressure  $p_{sfc}$  and  $T_{skin}$  and can be derived from the Clausius-Clapeyron relation for 100 % relative humidity. [Parameters at Input variables at the](#) lowest model level and above ocean are with good approximation 10 metres above the surface. According to Eq. (1) and 120 (2), fluxes from the atmosphere into the ocean are positive.



Whenever fluxes are computed with the above formulae, transfer coefficients are indirectly approximated for each grid point by dividing latent and sensible heat fluxes from ERA5 forecasts with the other terms of the right hand side of Eq. (1) and (2) (also from forecasts), which has to be done before flux anomalies are computed. This procedure works remarkably well for the ice-free ocean. Over sea ice, however, differences between model-level and surface quantities can be very small (i.e., mean climatology of absolute temperature and moisture differences can be  $\leq 0.01$  K and  $\leq 0.01$  g kg<sup>-1</sup>) so that the division by small numbers introduces artificial noise.

Seasonal trends are computed from monthly anomalies by subtracting the climatology of each grid point and subsequently averaging over December–February. The statistical significance of seasonal trends is computed with the 95 % confidence level and consideration of lag-1 autocorrelation. Analysis increments of model-level ~~parameters-variables~~ (temperature and humidity) are calculated at 06 and 18 UTC from the difference between analyzed state quantities and ~~12-hourly-12-hour~~ forecasts valid at ~~the analysis time. The impact of increments on flux trends is then those times.~~ During data assimilation, changes in the observing system can introduce analysis increments that may add or remove temperature or humidity from the model resulting in unrealistic trends (Bengtsson et al., 2004; Trenberth et al., 2011; Hersbach et al., 2020; Mayer et al., 2021). Uncertainties in model-based  $F_S$  trends caused by analysis increments are thus estimated by taking the difference between fluxes computed with analyzed state quantities and with short-term forecasts of those quantities.

In this study, we closely follow the mathematical interpretation of air-sea heat fluxes [i.e., Eq. (1) and (2)] and assume that the derived flux trends are solely caused by changes in their input variables (air density, wind, moisture, and temperature). We are aware that this is not the physically correct interpretation as changes in fluxes also influence the input variables due to their mutual dependency (e.g., an increase of latent heat flux increases the moisture in the lowest model-level, which in turn reduces latent heat fluxes). However, we argue that in equilibrium (i.e., considering long-term changes over multiple decades) this mutual influence is irrelevant for such discussions so that long-term flux trends can be considered as direct result of trends in input variables.

### 145 3.2 Linearization of turbulent heat fluxes and partial trends

To attribute trends in latent and sensible heat fluxes to trends in their input variables in a more quantitative way, we linearize the bulk formulas by decomposing each input variable  $x$  ( $\rho, U, q$ , and  $T$ ) into their mean state  $\bar{x}$  and deviation from its mean  $x'$  [known as Reynolds decomposition; see also Tanimoto et al. (2003) and Yang et al. (2016)]; that is, each input variable on the right side of Eq. (1) and (2) is described by  $\bar{x} + x'$ . After some calculus, computed turbulent heat fluxes can further be separated into a non-linear and linear part (see appendix A). The latter contains all products with at most one deviation term on which input variables are regressed to obtain a linear relation between trend in surface air-sea heat flux  $F$  (sensible or latent heat) and trend of each input variable, which reads as follows

$$\frac{\partial F}{\partial t} = \frac{\partial F}{\partial \rho} \frac{\partial \rho}{\partial t} + \frac{\partial F}{\partial |U_{10m}|} \frac{\partial |U_{10m}|}{\partial t} + \frac{\partial F}{\partial \xi_{ml}} \frac{\partial \xi_{ml}}{\partial t} + \frac{\partial F}{\partial \xi_{sfc}} \frac{\partial \xi_{sfc}}{\partial t} \quad \text{with} \quad F = F(\rho, U_{10m}, \xi_{ml}, \xi_{sfc}, t), \quad (3)$$

where  $\xi$  is a placeholder for either q or T. We term the expressions on the right *partial trends*, which are the product of the mean sensitivity  $\partial F/\partial x$  (i.e., regression of F on x using the whole period of time) and linear trend  $\partial x/\partial t$  of input variable x. The mean sensitivity describes how F changes when x is changed. Consequently, partial trends on the right side tell us how much of the flux trend ( $\partial F/\partial t$ ) is explained by the trend in one of the input variables ( $\partial x/\partial t$ ). Note that this procedure neglects trends in transfer coefficients and the non-linear part, which is a sufficient assumption for the purpose of this study as their contribution to the total turbulent heat flux trend is rather small (the linear part explains  $\geq 93\%$  of the total turbulent flux trend). Furthermore, we do not show partial trends of  $\rho$  in our evaluations as they are negligibly small.

### 3.3 Study area and box averages

Our general study area includes the ice-free ocean between 0–90° N and 90° W–30° E, but particular focus is laid on four  $8 \times 8^\circ$  boxes for which partial trends of latent and sensible heat fluxes are computed. The four focus regions are located in the Norwegian Sea (NWS; 68–76° N 2–10° E), over the northern flank of the North Atlantic Warming Hole (NAWH; 55–63° N 37–29° W), along the Gulf Stream extension (GS, 35–43° N 66–58° W), and in the tropical North Atlantic (TNA; 15–23° N 40–32° W) as trends in these regions are associated with distinct atmospheric and oceanic thermodynamics (see Fig. 1). Spatially averaging over these areas reduces the variance of fluxes and computational cost while trends are still captured reasonably well.

### 3.4 Meridional mass stream function

The seasonal mean meridional mass stream function  $\Psi_m$  is obtained by

$$\Psi_m = \frac{r_E}{g} \int_{\lambda=0}^{\lambda=2\pi} \int_0^p \bar{v}^* dp d\lambda \quad (4)$$

where  $r_E$  is the Earth's radius and  $\bar{v}^*$  is the seasonal mean of the deviation of the meridional wind component (on pressure levels) from its meridional mean, which is vertically integrated from the TOA to the pressure level of interest (p) and over all longitudes  $\lambda$ .

### 3.5 Indirect estimation of oceanic heat transports

We indirectly estimate the vertically integrated oceanic heat transport at a specific latitude of interest  $\varphi$  in the North Atlantic basin using the oceanic heat budget equation in the following form

$$OHT_\varphi = OHT_{\varphi_C} - \left[ F_S - \rho_0 c_p \frac{\partial}{\partial t} \int_0^Z (T_o - T_{ref}) dz \right]_{\varphi}^{\varphi_C} - R \Big|_{global} \quad (5)$$

where  $OHT_{\varphi_C}$  is the heat transport through the choke point  $\varphi_C$  (from DS+FS+BSO mooring-derived estimates; see red lines in Fig. 1), and the second term on the right side ~~describes the~~ is the difference between net air-sea heat flux ( $F_S$ ; from

ERA5 forecasts) and temporal ocean heat content tendency (OHCT; from 0–2000m IAP data) ~~subtracted from the net surface heat flux (from ERA5 forecasts) and averaged~~ integrated over the ocean area between  $\varphi_C$  and  $\varphi$  (the Mediterranean Sea is excluded; units are Watts). This is the same approach as used by Trenberth and Fasullo (2017), Mayer et al. (2022), and Baker et al. (2022) to indirectly estimate heat transports through the RAPID [Johns et al. (2011); McCarthy et al. (2015); Bryden et al. (2020); see also Fig. 1 in Mayer et al. (2022)] and SAMBA ~~arrayarrays~~, respectively.  $OHT_\varphi$  is calculated for every fifth latitude between 0–60° N ( $\varphi_C$  is situated between 67–80° N). Additionally, we adjust ~~heat fluxes in the North Atlantic basin~~ net heat fluxes by subtracting the monthly difference between global ocean mean vertically integrated 0–2000m OHCT and  $F_S$  (denoted as  $R$ ) uniformly from each grid point ~~of the global ocean, as done, e.g., by Trenberth et al. (2019)~~. This removes inconsistencies between ~~surface-air-sea~~ heat fluxes and OHCT and guarantees temporal consistency in a global manner [see also Trenberth et al. (2019); Liu et al. (2020); Mayer et al. (2022)].

To indirectly estimate the AMOC trend over the whole study period, we extend the 2005–10 OHT climatology of the choke point DS+FS+BSO to 1950–2019, under the assumption that the upward OHT trend at high latitudes is relatively weak (Mulwijk et al., 2018; Wang et al., 2019, 2020; Docquier and Koenigk, 2021) compared to trends at low latitudes and basin-wide heat flux trends. Trends of indirectly estimated OHT are computed in two ways, with monthly data and 5-year means, using the procedure described by Loeb et al. (2022) (see section 3.2 therein) to estimate trend uncertainties; that is, the effective sample size takes into account all significant autocorrelation functions  $\rho_\gamma$  up to lag  $m$  where  $\rho_{m+1} < 0$  and  $\rho_{m+1} + \rho_{m+2} < 0$   ~~$\gamma_{m+1} < 0$  and  $\gamma_{m+1} + \gamma_{m+2} < 0$~~  is satisfied. If ~~this~~ these conditions are not satisfied for the autocorrelation at any lag, the true instead of the effective sample size is used to estimate uncertainties.

### 3.6 Computation of climate indices

The AMO index is calculated similarly to the approach suggested by Trenberth and Shea (2006), where global-mean SST anomalies are subtracted from the spatially averaged SST time series of the North Atlantic basin (0–60° N and 0–80° W). The NAO index is derived from an EOF analysis applied to monthly surface pressure fields from ERA5 between 20–80° N and 90° W–40° E. The normalized principle component of the first EOF then describes the NAO index (Hurrell, 1995; Hurrell and Deser, 2009).

## 4 Results

In the following, we split the study period 1950–2019 equally and consider the periods 1950–84 and 1985–2019 separately as ~~inferred air-sea heat fluxes from Mayer et al. (2022) are~~ the inferred  $F_S$  from Mayer et al. (2022) is available only for the period from 1985 onward. Moreover, it has been shown that the global warming trend has accelerated in the past few decades (Cheng et al., 2017; Fox-Kemper et al., 2021) making the separation into two periods reasonable.

Air-sea heat fluxes in the North Atlantic ocean exhibit a distinct annual cycle, with the largest ocean heat loss to the atmosphere in boreal winter and strongest heat gain in summer. In boreal winter, fluxes are widely negative (heat loss from the ocean to atmosphere) over the ocean basin, in particular at high latitudes and along the Gulf Stream where an immense amount of oceanic energy is transported northward and long-term averages of net surface-air-sea heat fluxes can be as large as  $-400$   $\text{W m}^{-2}$  (Fig. 1). During summer, heat fluxes are positive (the ocean gains energy from the atmosphere) across the ocean basin, with values ranging from zero (along the Gulf Stream and tropical North Atlantic) to  $200 \text{ W m}^{-2}$  in coastal areas of North America and Africa (not shown). Here, we focus on winter-month (December–February) heat fluxes as they feature the most pronounced heat flux trends among all calendar months.

Over the past 35 years (1985–2019), several prominent regions with significant positive or negative trends have emerged (Fig. 2a). Negative surface-air-sea heat flux trends (stronger loss of energy from the ocean to atmosphere) can be found along the Gulf Stream, and in regions of strong sea ice retreat that is driven by recent global warming (Fox-Kemper et al., 2021), e.g., along the East Greenland current, in the Baffin Bay and the western part of the Labrador Sea, and in the northern part of the Barents Sea. The retreat allows the ocean to cool in areas that were otherwise covered by sea ice resulting in strong negative heat flux trends. Surface heat loss in the tropical North Atlantic also strengthens significantly, but to a lesser extent than in the Gulf Stream or region-regions of strong sea ice retreat.

Positive trends (weakening of negative net surface-air-sea heat fluxes during winter months) are prominent in the Norwegian and-Nordic Seas and in the eastern part of the Labrador Sea, but also in the region where Gulf Stream water masses bifurcate and form the North Atlantic Drift Current further north and the equatorward propagating Azores current in the south (between  $40^{\circ}$ – $50^{\circ}$  N and  $45^{\circ}$ – $25^{\circ}$  W). This region of strongly positive trends appears spatially more extended for inferred surface-heat fluxes- $F_S$  from Mayer et al. (2022) (see Fig. 2b), but with similar peak value-values of about  $29 \text{ W m}^{-2} \text{ dec}^{-1}$  (dec = 10 years). At other locations of the North Atlantic Ocean, both flux products exhibit qualitatively similar trends indicating that ERA5 flux trends seem reliable in terms of spatial structure, at least for the chosen study area and period of time. However, note that the trends in many areas are statistically insignificant (e.g., the positive trends in the Labrador and Irminger Sea, or at the southern flank of the Gulf Stream) and thus should be treated with caution when interpreting them.

Main-contributor-The main contributors to the  $F_S$  trend-trends are turbulent heat fluxes (THF; Fig. 2c), whereas trends in radiative fluxes (RHF; Fig. 2d) are usually an order of magnitude weaker (except for the Arctic ocean, which is not further discussed here). Spatial means over the whole study area are  $-1.6 \text{ W m}^{-2} \text{ dec}^{-1}$  for THF and  $\sim 0.1 \text{ W m}^{-2} \text{ dec}^{-1}$  for RHF resulting in negative  $F_S$  trends of about  $-1.4 \text{ W m}^{-2} \text{ dec}^{-1}$  during 1985–2019 (see Table 1). For comparison, the inferred  $F_S$  exhibits a weak positive trend of  $0.3 \text{ W m}^{-2} \text{ dec}^{-1}$  owing to the spatially more extended positive trends.

We also computed mean trends of globally adjusted  $F_S$  (see section 3) and net surface-air-sea heat fluxes from Liu et al. (2020) (also known as DEEP-C dataset; publicly available at <https://doi.org/10.17864/1947.000347> for the period 1985–2017) over the whole study area. The DEEP-C product is based on our inferred  $F_S$  but unrealistic surface fluxes over land are subsequently redistributed to the ocean, which removes spurious trends in the late 1990s and 2000s (Liu et al., 2017, 2020;

Mayer et al., 2022). As a consequence,  $F_S$  from DEEP-C exhibits a realistic global ocean mean that matches the observed  
245 mean ocean heat uptake and is thus well suited as reference for long-term trend studies (Mayer et al., 2022). For 1985–2017,  
we find a mean trend of  $1.4 \text{ W m}^{-2} \text{ dec}^{-1}$  for globally adjusted  $F_S$  and  $1.1 \text{ W m}^{-2} \text{ dec}^{-1}$  for the DEEP-C product (both are  
statistically insignificant), indicating good agreement of the two estimates. For comparison, the unadjusted model-based  $F_S$   
exhibits a 1985–2017 mean trend of  $-1.6 \text{ W m}^{-2} \text{ dec}^{-1}$  indicating that the global adjustment of  $F_S$  yields more realistic and  
reliable trend estimates. This adds confidence to our globally adjusted  $F_S$  data and its use for the full period starting in 1950.

250 In addition, we show each component of THF separately. In general, THF trends (and consequently also model-based  $F_S$   
trends) are governed by changes in latent heat flux (Fig. 2e) at low latitudes and sensible heat flux trends (Fig. 2f) at mid- and  
high-latitudes (north of  $\sim 40^\circ \text{ N}$ ). Of the  $-1.6 \text{ W m}^{-2} \text{ dec}^{-1}$  mean THF trend, about  $-1.9 \text{ W m}^{-2} \text{ dec}^{-1}$  stem from latent heat  
and  $0.3 \text{ W m}^{-2} \text{ dec}^{-1}$  from sensible heat flux trends (Table 1). Along the sea ice edge, both components contribute equally to  
the negative THF trend as both were substantially lower when ocean areas were covered by sea ice before.

255 While flux trends shown in Fig. 2 are from forecasts, the following evaluations are based on analyzed quantities as they are  
better constrained by observations than their forecast counterpart. Most differences between forecast and analyzed flux trends  
can be related to moisture analysis increments (discussed below).

Trends in latent and sensible heat fluxes can further be formally attributed to changes in 10 metre horizontal wind speed  
and temperature or humidity differences between [the](#) lowest model level and ocean surface [see bulk formulae Eq. (1) and  
260 (2); [trends in air density are not discussed as they are negligibly small](#)]. Model-level humidity (Fig. 3a) uniformly increases at  
almost all locations as expected from a warming atmosphere [a warmer atmosphere can hold more moisture; Douville et al.  
(2021)]. The statistically insignificant decline in the eastern North Atlantic and Mediterranean Sea can be attributed to stronger  
northerly winds (see Fig. 4a) and declining moisture transport into that area as related to a strengthened NAO.

Changes in model-level temperature (Fig. 3b) are qualitatively similar to those in atmospheric moisture, but are statistically  
265 significant in almost all parts of the tropical North Atlantic. As the ocean warms due to climate change (Fox-Kemper et al.,  
2021), surface saturation humidity and skin temperature (Fig. 3c and d) increase almost everywhere (note that the surface  
saturation humidity is derived from skin temperature and surface pressure according to the Clausius-Clapeyron equation). The  
moderate and statistically insignificant decreasing trend [in the Irminger Sea, to the south of the Irminger Sea \(around 20–30°  
W, 55° N\)](#), which also appears weaker in model-level [parameters variables](#), is a result of the anomalously cool ocean in the  
270 North Atlantic Warming Hole (Rahmstorf et al., 2015).

The [surface air-sea heat](#) fluxes are not so much governed by individual [parameters variables](#) at surface and model level, but  
by their differences ([Fig. 3e and f](#)) from which several observations can be made:

1. The trend pattern of  $(q_{\text{ml}} - q_{\text{sfc}})$  and  $(T_{\text{ml}} - T_{\text{skin}})$  are almost identical to that of latent and sensible heat fluxes (cf. Fig.  
2e and f; pattern correlations are  $> 0.8$ ) indicating that the horizontal wind speed (Fig. 4a) has a comparatively small  
275 impact on the spatial distribution of LHF and SHF trends.
2. Long-term changes in surface saturation humidity (governed by skin temperature trends) are in most areas of the North  
Atlantic stronger than changes in model-level humidity, and vice versa for the temperature. This results in almost uni-

280 formly negative ( $q_{ml} - q_{sfc}$ ) but positive ( $T_{ml} - T_{skin}$ ) trends. While surface saturation humidity increases with increasing skin temperature according to the Clausius-Clapeyron relation (i.e., relative humidity remains 100 %), the increase in model-level humidity is much weaker so that relative humidity decreases (Fig. 4b), especially south of 40° N. This means, near-surface air masses in the tropical North Atlantic become drier relative to the temperature increase (the Clausius-Clapeyron relation would postulate stronger humidity trends for constant relative humidity), which can be caused by several factors (discussed below).

285 3. Among the regions of strong sea ice retreat [in the Labrador and Nordic Seas](#), peak positive trends in surface humidity and temperature can be found along the [Gulf Stream region \(main current of the Gulf Stream \(35–45° N, 80–45° W\)](#); with values up to 0.5 g kg<sup>-1</sup> dec<sup>-1</sup> and 0.8 K dec<sup>-1</sup>). This leads to remarkably strong negative trends in ( $q_{ml} - q_{sfc}$ ) and ( $T_{ml} - T_{skin}$ ) highlighting the importance of the ocean in this area. It should be noted that the Gulf Stream signal is barely visible in relative humidity trends (Fig. 4b) due to a well-mixed boundary layer and strong coupling between atmosphere and the underlying warm Gulf Stream.

290 4. Over the North Atlantic Warming Hole, changes in the model-level temperature and humidity closely follow the Clausius-Clapeyron relation so that the warming hole signal is barely visible in RH trends (Fig. 4b). Trends in model-level and surface [parameters-variables](#) are of similar strength resulting in weak and statistically insignificant heat flux trends (note that the North Atlantic Warming Hole is further north to the bifurcation area of the Gulf Stream and does not coincide with peak positive trends).

295 5. The strong positive ( $T_{ml} - T_{skin}$ ) trend in the Norwegian Sea originates from positive trends in the atmosphere and somewhat less positive trends (or even negative trends east of Iceland) of the skin temperature. This can be attributed to trends towards more south-easterly winds (see Fig. 4a) advecting warmer air masses from lower latitudes to the Norwegian Sea, which is related to a strengthened Icelandic low.

300 6. In the Labrador and Nordic Seas, ( $T_{ml} - T_{skin}$ ) trends downwind [to-of](#) areas of strong sea ice retreat become widely positive (mean wintertime climatology is a northerly wind in both basins; not shown). One explanation could be that air masses that are advected from further north get heated by the enhanced fluxes where sea ice retreated. The anomalously warm air-masses damp air-sea fluxes further south resulting in largely compensating sensible heat fluxes along the wind direction (spatial average is  $\sim 2 \text{ W m}^{-2} \text{ dec}^{-1}$  over the Nordic Seas; see Table 1), with negative trends in areas of strong sea ice retreat and positive trends downwind. [This effect cannot be observed for latent heat fluxes and requires further](#)  
305 [investigation that is beyond the scope of this study.](#)

One possibility to throttle the growth in near-surface humidity in the tropical North Atlantic is a stronger advection of dry air masses through intensification of the Hadley Cell. To manifest this, we present DJF trends of the zonally averaged meridional mass stream function derived from ERA5 wind fields (Fig. 5). The dipole structure between 0–30° N indicates that the north hemispheric Hadley cell has shifted poleward and strengthened in intensity, which enhances the subsidence of dry air along  
310 the northern flank of the Hadley Cell. This also agrees with positive trends in 10 metre wind speed between  $\sim 20\text{--}30^\circ \text{ N}$  (Fig.

4a). Note that the mass stream function is obtained by integrating over all longitudes, and the intensification may take place over other ocean basins. However, a statistically significant increase of low-level cloud cover and outgoing longwave radiation (~~not shown~~) [~~not shown~~; see, e.g., [Chen et al. \(2002\)](#); [Mathew and Kumar \(2019\)](#)] in the almost entire tropical North Atlantic Ocean indicate that the Hadley cell intensification also appears over the tropical Atlantic Ocean.

315 On the other hand, Trenberth et al. (2011) and Mayer et al. (2021) noted that analysis increments introduced by the data assimilation system due to changes in the observing system can artificially remove or add atmospheric moisture, which in turn could influence near-surface humidity trends. To investigate the impact of atmospheric moisture and temperature increments in the lowest model level on air-sea heat fluxes from ERA5, we compute latent and sensible heat fluxes according to the bulk formulas (Eqs.1 and 2) using both forecast and analyzed state quantities (not shown). Differences between trends derived  
320 from analyses and forecasts can then be used as [a rough estimate for trend uncertainties caused by analysis increments](#) (note that only temporally varying analysis increments introduce an artificial trend, and not a constant offset between analyses and forecasts). [It should be highlighted that analysis increments computed in this way cannot be used to fully remove the temporal inconsistencies in model-based  \$F\_S\$  as their exact impact on both forecasts and analyzed state quantities is unknown. That is, the difference between adjusted and unadjusted model-based  \$F\_S\$  \(magnitude of the global adjustment\) is not equivalent to the](#)  
325 [difference between fluxes computed with forecasts and analyzed state quantities.](#)

We find strongest variation in humidity increments (approximated by  $q_{an}-q_{fc}$ ) in the tropics, with ~~1985–2019 trends up to  $\sim 0.05$  and~~ values of  $-0.2 \text{ g kg}^{-1}$  in the late 2000s and early 2010s (see appendix ~~B) B)~~ and ~~1985–2019 trends of about  $-0.05 \text{ g kg}^{-1} \text{ dec}^{-1}$ .~~ Humidity increments in earlier times and at higher latitudes are in general ~~less negative. Temperature increments are temporally more stable and almost independent of latitude, with values ranging between  $\pm 0.03$ , and are thus~~  
330 ~~less impactful on turbulent heat fluxes~~ [less negative](#). The latent heat flux trends derived from analyses ~~are in (not shown), for~~ the zonal mean of the North Atlantic, [are](#)  $1\text{--}2 \text{ W m}^{-2} \text{ dec}^{-1}$  ~~stronger more negative~~ than those based on forecasts (root mean square error between trends derived from analysis and forecasts is  $1.1 \text{ W m}^{-2} \text{ dec}^{-1}$  over the ice-free ocean). The negative humidity increments in the lowest model level ~~artificially~~ remove moisture from the model, which results in larger ( $q_{ml}-q_{sfc}$ ) differences and thus stronger [\(more negative\)](#) analyzed latent heat flux trends. For sensible heat fluxes, analysis increments  
335 are less important. [Temperature increments are temporally more stable and almost independent of latitude, with trends varying between  \$\pm 0.03 \text{ K dec}^{-1}\$  \(see appendix B\).](#) Zonally averaged SHF trends derived from analyses are ~~<at low (high) latitudes less than 0.2 (0.4)  $\text{W m}^{-2} \text{ dec}^{-1}$  larger more positive (negative)~~ than those derived from forecasts (RMSE over the whole study area is  $0.5 \text{ W m}^{-2} \text{ dec}^{-1}$ ). ~~Therefore, we argue that the regional impact of all relevant analysis increments introduced by the ERA5 data assimilation on air-sea heat flux trends is rather small during the 1985–2019 period. While trends from~~  
340 ~~analyses~~ [Consequently, temperature increments are less impactful on turbulent heat fluxes. Trends of turbulent heat fluxes derived from analysis \(not shown\)](#) are at all latitudes ~~stronger up to  $1 \text{ W m}^{-2} \text{ dec}^{-1}$  more negative~~ than those derived from forecasts. [However](#), the spatial pattern of the trends remains almost unaffected (~~i.e., because~~ the difference between analyses and forecasts is smaller for weaker trends, and vice versa); ~~that is, the negative LHF trend.~~ [Therefore, we argue that the regional impact of all relevant analysis increments introduced by the ERA5 data assimilation on air-sea heat flux trends is rather small](#)



345 during the 1985–2019 period. The negative LHF trends in the tropical North Atlantic is-are most likely a result of Hadley cell intensification and can not be explained by temporally varying moisture increments.

In summary, long-term changes in net surface-air-sea heat fluxes over the North Atlantic Ocean are primarily driven by latent heat flux trends (Table 1), which are associated with changes in the surface (related to changes in skin temperature) and model-level humidity (e.g., advection of drier air masses), while changes in wind speed are negligibly small. Furthermore, we  
350 conclude that temporally varying analysis increments influence the magnitude of air-sea-turbulent heat flux trends by about 1–2  $\text{W m}^{-2} \text{dec}^{-1}$ , whereas their spatial pattern remains widely unchanged.

## 4.2 1950–1984 trends

Long-term latent and sensible heat flux changes before 1985 (Fig. 6) differ in most areas substantially from those during the more recent period. Most notable are the strong positive and significant LHF trends in the Caribbean Sea, along the Gulf Stream  
355 north of  $\sim 40^\circ \text{N}$ , and in the Labrador Sea.

Trends in sensible heat flux are strongest along the sea ice edge and are absent in the Norwegian Sea, the region with the strongest-weakening-most prominent negative trends after the 1980s (cf. Fig. 2f). The widely positive trends before 1985 can be attributed to a stronger temperature decrease at the surface relative to the atmosphere (not shown). We find remarkably strong correlations (0.5–0.8) between trends of skin temperature and 0–300m OHC from IAP for all four box averages (see Fig. 1)  
360 suggesting that the The skin temperature of the ice-free ocean shows a statistically insignificant increase in wide areas of the tropical North Atlantic and a decrease almost everywhere north of  $25^\circ \text{N}$ , which coincides with the basin-wide trend pattern of the 0–300m OHC trend from IAP (not shown). Although their pattern correlation is weak ( $\sim 0.2$ ) for the ice-free ocean area due to small scale differences, the generally good match of the large-scale trend pattern in terms of sign suggests that the weakening of sensible heat fluxes is related to the ocean cooling that occurred during that time [see Hodson et al. (2014)].  
365 Consequently, For comparison, the pattern correlation for the more recent period 1985–2019, where both skin temperature and OHC increases uniformly (except for the North Atlantic Warming Hole) due to global warming, is only  $\sim 0.15$  for the ice-free ocean. This indicates a generally weak spatial correlation between OHC from IAP and skin temperature from ERA5, although their large-scale trend pattern agrees well in terms of sign. As a consequence of the ocean cooling, there is no warming hole signal in the skin and model-level temperature before 1985, which is consistent with results from Chemke et al. (2020) based  
370 on satellite-based HadISST data [see supplementary information therein; in fact, ERA5 employs the second version of this dataset as SST forcing, see Hersbach et al. (2020)]. Temperature analysis increments before 1985 do not play an important role due to their negligibly weak trends of less than  $\pm 0.02 \text{ K dec}^{-1}$  in ice-free regions. This results in differences between trends derived from analysis and forecast data of less than  $0.5 \text{ W m}^{-2} \text{dec}^{-1}$  in the zonal mean, with an RMSE of about  $0.5 \text{ W m}^{-2} \text{dec}^{-1}$  for the ice-free ocean.

375 In accordance with the ocean cooling during the 1960s and early 1970s, surface saturation humidity decreases almost everywhere before 1985 (not shown), with the strongest negative trends-decrease along the Gulf Stream. The only larger patch of positive (but statistically insignificant) trends appear in the subtropics and along the sea ice edge. Near-surface humidity also decreases where temperature decreases (not shown), but weaker than the CC-related Clausius–Clapeyron-related decrease such

that the relative humidity increases significantly in most areas (whereas negative trends in all ice-free areas are insignificant; not shown). This leads to mostly increasing humidity differences ( $q_{ml}-q_{sfc}$ ) and thus also in positive latent heat fluxes trends before 1985 (Fig. 6a).

As for the SHF, we exclude analysis increments as a possible source of uncertainties during that time because of their weak trend of  $\pm 0.02 \text{ g kg}^{-1} \text{ dec}^{-1}$  over most locations of the ice-free ocean. Therefore, differences between LHF trends derived from analyses and forecasts are less than  $0.6 \text{ W m}^{-2} \text{ dec}^{-1}$  in zonal mean before 1985 (RMSE over the ice-free ocean is  $0.5 \text{ W m}^{-2} \text{ dec}^{-1}$ ). Changes in 10-metre wind speed are mostly insignificant and of similar strength as after 1985, and thus affect turbulent heat fluxes only marginally.

### 4.3 Flux trends in focus regions

To understand long-term changes in the four thermodynamically interesting areas of the North Atlantic (see boxes in Fig. 1) in more detail, we show spatial averages of model-based and inferred  $F_S$ , and partial trends of analyzed input variables as regressed onto LHF and SHF from ERA5 forecasts, for the period 1950–2019 (see Fig. 7 and Table 1). In the Norwegian Sea (NWS), air-sea heat fluxes weaken particularly in the late 2000s and 2010s (Table 1), which is in good agreement with the enhanced oceanic heating during that time (Mork et al., 2019; von Schuckmann et al., 2021). However, the advection of warmer more humid air associated with changes in 10 metre wind direction appears to overcompensate (see Fig. 4a) appears to dominate oceanic trends (right panel in Fig. 7; partial trends of surface quantities are negative as they contribute with opposite sign to turbulent heat flux trends) so that LHF and SHF trends are relatively weak compared to those in model-level or surface quantities alone. Note that Skagseth et al. (2020) found similar changes in wind direction for the adjacent Barents Sea.

Long-term changes trends of model-based  $F_S$  in the North Atlantic Warming Hole (NAWH) are the weakest among the four areas of interest weak and statistically insignificant in both sub-periods 1950–84 and 1985–2019 (see also Table 1). Derived trends should thus be treated carefully as they also depend strongly on the chosen reference period. To test the robustness of model-based  $F_S$  trends in the four study areas, we considered various reference periods. For instance, we removed the last year from the time series and computed DJF trends based on 1950–2018. While net air-sea heat fluxes steadily increase in the NWS and decrease in the tropical North Atlantic (TNA) almost independently of the chosen reference period, trends in the NAWH and GS region are less than  $1 \text{ W m}^{-2} \text{ dec}^{-1}$  (or less than  $2 \text{ W m}^{-2} \text{ dec}^{-1}$  when considering 1950–2019, see Table 1) and statistically insignificant. From this, we cautiously argue that heat fluxes in the NAWH, and also in the GS box where model-based  $F_S$  trends from the early and late period compensate each other, do not exhibit a prominent long-term trend over the past 70 years as related to global warming, while changes in the TNA and NWS are most likely a result of global warming.

We also explore the winter-month  $F_S$  climatology along the Gulf Stream extension on decadal timescales in order to reveal any signal in air-sea heat fluxes associated with a poleward displacement due to global warming. Besides an oscillatory behaviour similar to temporal changes in the more regional GS box shown in Fig. 7, we could not find a distinct sign of a poleward shift in air-sea heat fluxes, which is consistent with findings from Yang et al. (2016).

The partial trends in Fig. 7 show that trends in model-level and surface quantities almost always act in opposite directions and thus compensate each other to some degree (except for moisture in the TNA during the first period, where both

are negative). In addition, ~~it demonstrates the partial trends demonstrate~~ qualitatively that the impact of 10 metre wind speed on heat flux trends is rather small compared to changes in moisture or temperature, especially in cases where latent or sensible  
415 heat fluxes exhibit trends of several Watts per square metre.

We also find that in the three northernmost boxes, trends of inferred and model-based ~~fluxes- $F_S$~~  have the same sign but differ by about  $2\text{--}4 \text{ W m}^{-2} \text{ dec}^{-1}$ . ~~Inferred fluxes exhibit~~ The inferred  $F_S$  exhibits stronger upward trends in the NWS and NAWH, and a weaker downward trend in the GS area (Table 1). Trends in the TNA coincide remarkably well underlining the reliability of model-based ~~fluxes from ERA5 forecasts- $F_S$~~  in that particular region. In summary, this suggests that model-based trends are  
420 largely reliable in terms of sign and spatial pattern (see also Fig. 2).

#### 4.4 Long-term impact of natural variability modes

The North Atlantic Oscillation (NAO) is a periodic oscillation in sea level pressure and wind (Visbeck et al., 2001) and can temporally and regionally influence air-sea interactions. During the last 30–40 years, the NAO tends to more positive phases (strengthened Icelandic low and Azores high) than before, which has been attributed to global warming (Gillett et al., 2003).  
425 Here, we want to explore its long-term impact on trends of air-sea heat fluxes (Fig. 8).

Long-term  $F_S$  changes over the entire study period appear to be weaker and spatially more uniform compared to those over the two sub-periods discussed before. Trends are widely positive in the western North Atlantic, in the region of the North Atlantic Warming Hole, and in the Norwegian Sea. Persistent negative flux trends occur in the tropical North Atlantic, along the  
430 Gulf Stream, and in regions of strong sea ice retreat which are largely consistent with negative changes during both sub-periods.

The December–February NAO regressed onto  $F_S$  features a basin-wide tripolar pattern (see appendix C), with strong negative values in the Irminger and Labrador Sea, negligibly weak trends in the tropical and subtropical latitudes, and positive values in between (with peak values along the Gulf Stream). The more frequent occurrence of positive NAO phases over the past 30 years, relative to 1950–90, seem to favour anomalous ocean cooling at higher latitudes and heating in the western North  
435 Atlantic ( $25\text{--}45^\circ \text{ N}$ ,  $80\text{--}40^\circ \text{ W}$ ; see Fig. 8a, and positive trends in Fig. C1a). Removing the NAO signal from  $F_S$  trends (Fig. 8b) thus weakens ocean cooling (more positive trends) in the Irminger and Labrador Sea over time and allows stronger cooling in the western North Atlantic (less positive trends). In addition, we find weak correlations of less than 0.4 between NAO index and  $F_S$  box averages in the TNA, GS, and NWS, but -0.75 for the NAWH. This indicates that the  $F_S$  trend over the North Atlantic Warming Hole box is strongly influenced by trends of the NAO and its tendency toward more positive phases, while  
440 other areas are less effected.

Despite the remarkably strong regional impact of the NAO on air-sea heat fluxes at high latitudes, its spatial mean averaged over the whole study area is less than  $0.03 \text{ W m}^{-2} \text{ dec}^{-1}$  (1950–2019). For comparison, the 1950–2019  $F_S$  trend averaged over the whole study area (as shown in Fig. 8a) is  $0.14 \text{ W m}^{-2} \text{ dec}^{-1}$ . This suggests that the trend toward more positive NAO phases only leads to a relocation of areas where oceanic heat is lost or taken up through surface-air-sea heat fluxes, rather  
445 than a steady increase in anomalous ocean heat uptake as related to global warming. This somewhat agrees with the finding of Cohen and Barlow (2005), that the global DJF warming trend during 1972–2004 may be unrelated to regional warming trends

driven by the NAO. To check the potential role of decadal NAO variability, we tested the impact of applying LOWESS and Butterworth low-pass filters on the NAO series before trend computation (not shown), but the impact on the trend results was small and the results are thus deemed robust.

450 We also regressed the AMO forcing (Kerr, 2000) onto  $F_S$  to estimate its long-term impact on flux trends (see appendix C). The AMO partial trend varies between  $\pm 2 \text{ W m}^{-2} \text{ dec}^{-1}$  over the ice-free ocean, with negative values in the Irminger and Labrador Sea and around the North Atlantic Warming Hole ( $40\text{--}60^\circ$  and  $50\text{--}20^\circ$  W), and positive values elsewhere. Although the AMO impact on flux trends in the Irminger and Labrador Sea has the same sign and similar spatial structure as the NAO forcing, its strength over the 70-year period is weaker. Additionally, we find a spatial mean of the AMO signal of  $0.22$   
455  $\text{W m}^{-2} \text{ dec}^{-1}$ , which points to a basin-wide weakening of air-sea heat fluxes, but this is likely an effect of non-zero AMO trend due to the relative shortness of the time series (the AMO does not complete a full period during the whole study period), and is likely not related to global warming.

#### 4.5 Changes in the Atlantic Meridional Overturning Circulation

In the previous sections we have diagnosed a reduction of the net ~~surface heat flux from ocean to atmosphere~~ air-sea heat flux  
460 during 1950–2019 when averaging over the North Atlantic. This reduction could be related to a cooling trend of the underlying ocean and/or a reduction of oceanic heat transports associated with the AMOC. In this section we explore both possibilities to verify the AMOC trends with observation-based data (reanalysis is a combination of observations and forecasts). The ocean heat transport at different latitudes of the North Atlantic basin is indirectly estimated from the ocean heat budget using globally adjusted  $F_S$  from ERA5 forecasts and OHC data from IAP (see section 3). Here we focus on full-year OHT estimates because  
465 it increases the signal-to-noise ratio (sub-annual OHC changes are often related to seasonally compensating trends in wind patterns) and observational uncertainties of OHCT are considered larger on sub-annual time scales. Furthermore, annual mean  $F_S$  trends are similar to seasonal DJF trends in terms of spatial pattern (pattern correlation is  $\sim 0.8$ ), but are generally weaker across the North Atlantic basin (root mean square of trends is  $1.8 \text{ W m}^{-2} \text{ dec}^{-1}$  as compared to  $3.7 \text{ W m}^{-2} \text{ dec}^{-1}$ ).

Results for  $0\text{--}60^\circ$  N are shown in Fig. 9 using two types of trend estimates (see section 3). Both estimates show more negative  
470 trends (weakened AMOC) and larger uncertainties at lower latitudes, with a maximum at the equator. While the method based on five-year means gives significant trends for all latitudes except  $35\text{--}50^\circ$  N (averaging over 5 years reduces the variance), linear regression on monthly data is statistically significant only between  $45\text{--}60^\circ$  N.

~~Main contributor~~ The main contributors to the weakened OHT in the North Atlantic basin are statistically significant long-term changes of globally adjusted air-sea heat fluxes, whereas the trend of meridionally integrated OHCT is comparably small  
475 and insignificant throughout all latitudes between the equator and  $60^\circ$  N (see Table 2 for trends integrated over the area between choke point DS+FS+BSO and  $26^\circ$  N). Removing the OHCT term from the budget equation [integral term in Eq. (5)] thus reduces trend uncertainties while the strength and meridional structure of the estimated OHT trend remain roughly the same (compare left and middle panel of Fig. 9). In other words, the AMOC weakening is primarily associated with a positive trend of globally-adjusted  $F_S$  and thus a decline of ocean-to-atmosphere heat fluxes (1950–2019 mean is  $-13.7 \text{ W m}^{-2}$ ). We

480 also computed the indirectly estimated OHT trend based on the sub-periods 1950–84 and 1985–2019 but could not find a significant AMOC weakening in either period (not shown).

## 5 Summary and discussion

In this work, we investigated the reliability and temporal stability of winter-months (December–February) trends of model-based net surface-air-sea heat fluxes from ERA5 forecasts (denoted as model-based  $F_S$ ) over the North Atlantic Ocean during 485 1950–2019. Main drivers of these trends are identified using analyzed state quantities from ERA5, and the influence of natural variability modes and analysis increments as introduced by the ERA5 data assimilation system are considered. Whenever possible, ERA5 forecast fluxes are compared with indirect estimates from Mayer et al. (2022), which are proven to be temporally stable and exhibit a small mean bias over the global ocean. Furthermore we performed a linear perturbation analysis on turbulent heat fluxes in four distinct  $8 \times 8^\circ$  boxes, which allowed us to quantitatively attribute flux trends to changes in wind speed, 490 moisture, and temperature, assuming a linear regime. In a final step, we used basin-wide annual mean air-sea heat fluxes to indirectly estimate the AMOC trends over the past 70 years, and discussed its-their reliability and sources of uncertainties.

We find that air-sea heat flux trends at low (high) latitudes are largely driven by long-term changes of differences between model level and surface humidity (temperature). We further traced surface trends back to local changes in the ocean heat content, whereas model level trends strongly depend on altered conditions of advected air masses through changes in wind di- 495 rection, and not so much in wind speed. This process likely plays a major role in the tropical North Atlantic where increasingly drier air masses are advected (likely linked with a strengthening of the Hadley cell), as well as in the Norwegian Sea, where increasingly warmer air is advected.

A more quantitative assessment of turbulent heat fluxes in four individual sub-regions reveals that the relative contribution of wind speed to turbulent heat flux trends is indeed negligible, and that surface and model level trends largely compensate 500 each other. Furthermore, it is shown that trends in the later period (1985–2019) are substantially stronger compared to the early period, which is consistent with accelerated warming in the past few decades (Cheng et al., 2017; Fox-Kemper et al., 2021). It should be noted that the strength of trends clearly depends on the chosen averaging area, especially in the Gulf Stream where north-south gradients of surface-air-sea heat flux trends are steep.

The long-term changes in air-sea heat fluxes could have some further implications on weather and climate. For instance, 505 the increased intensity of tropical cyclones during the past 40 years (Kossin et al., 2020) could possibly be linked to stronger latent heat fluxes in the tropical North Atlantic (Fig. 8a2e; similar trends can be found for the Hurricane season September–November). Similarly, the negative heat flux trends over the Gulf Stream are most likely a response to an increased storm frequency, which in further consequence favours more cyclogenesis (Shaman et al., 2010).

We also examined the impact of NAO and AMO on long-term  $F_S$  trends. The more frequent positive NAO phases during 510 the last 30–40 years significantly alter trends at high latitudes. It favours stronger ocean heat loss to the atmosphere via air-sea heat fluxes in the Irminger and Labrador sea and anomalously weak loss in the western North Atlantic, albeit the basin-wide mean heat exchange between atmosphere and ocean remains unaffected. The AMO forcing, on the other hand, is weaker than

the NAO forcing but exhibits a non-zero mean in the North Atlantic basin, but robust statements about the impact of AMO are difficult given the relative shortness of the considered time series.

515 Finally, we linked the basin-wide air-sea heat flux trend to the AMOC weakening found in other studies by evaluating the oceanic heat budget using surface air-sea heat fluxes from ERA5 forecasts, OHCT data from IAP, and ocean heat transport data from Arctic Gateways in the north (i.e., mooring-derived estimates from the Davis Strait, Fram Strait, and Barents Sea Opening; see Fig. 1). As 90 % of the observed total meridional heat transport at 26° N is carried out by the overturning circulation (Johns et al., 2011; Zhang et al., 2019), the found decrease in inferred OHT can directly be linked to a weakening  
520 of the AMOC. Trend estimates based on monthly data exhibit large uncertainties and are insignificant south of 45° N, whereas computations based on 5-year means yield significant trends at almost all latitudes (taking into account all significant autocorrelation coefficients, see section 3). Removing the OHCT term from the calculations reduces uncertainties while trends remain approximately the same (the long-term OHCT trend is small compared to that in  $F_S$  but introduces noise). Based on these results, we provide new and independent evidence for a weakening of the AMOC over the past 70 years [see also Rahmstorf  
525 et al. (2015); Caesar et al. (2018); Fox-Kemper et al. (2021); Boers (2021)], which is associated with positive heat flux trends (weakened negative fluxes) in the North Atlantic basin. We argue that the mean ocean heat transport through the choke point in the Nordic Seas is small ( $\sim 0.15$  PW) so that even relatively large changes would not have a strong impact on indirectly estimated OHT trends further south [see Muilwijk et al. (2018) for long-term simulations of ocean heat transports through Arctic gateways].

530 Analysis increments of moisture and temperature at the lowest model level (i.e., the difference between analysis and forecast) likely influence the strength of trends but not so much the basin-wide spatial pattern. At most locations, moisture is removed from the model by the assimilation process resulting in stronger (more negative) LHF trends from analysed data by about ~~1-2~~  
1  $\text{W m}^{-2} \text{dec}^{-1}$  as compared to forecast data. Strongest moisture analysis increments can be found in the tropics in the late 2000s. At higher latitudes and before 2000, moisture increments are temporally stable and have negligible impact on surface  
535 turbulent air-sea heat flux trends. Temperature increments are relatively small and stable throughout the study period and thus play only a secondary role.

In the early period, observations are temporally and spatially sparse resulting in analyzed states that are closer to the model climate (to which forecasts are drifting) than to observations. Over time, more and more observational data are assimilated pulling the analysis away from the model climate. This increases analysis increments, which can have several implications  
540 on air-sea heat flux trend estimates. When trends are weak or compensate each other such that signal-to-noise ratio becomes low (e.g., when averaging over large areas), analysis increments can have a relatively large impact on the trend estimate. For example, the heat flux trend in the tropical North Atlantic box (see Fig. 1) is only  $-2.7 \text{ W m}^{-2} \text{dec}^{-1}$  during 1985–2019 (Table 1). Analysis increments increasingly remove moisture from the atmosphere in that region (see appendix B) so that the trend based on analyzed state quantities is  $-4 \text{ W m}^{-2} \text{dec}^{-1}$  (not shown). This is a 50 % stronger trend compared to the forecast-  
545 based estimate. Nonetheless, it is important to note that this is still a factor of  $\sim 3$  smaller than the trend uncertainty listed in Table 1. A similar effect can be found for global ocean and basin-wide averages as used to estimate the AMOC weakening. Both suffer from temporal inconsistencies in the late 1990s and early 2000s, which are likely caused by changes in the atmospheric

observing system and hence analysis increments. Nevertheless, given that trends in analysis increments are spatially relatively uniform, we find that the applied global correction removes much of the effect of spurious air-sea flux trend on our inferred estimate of OHT; that is, temporal inconsistencies in basin-wide averages of model-based  $F_S$  (and thus also in the inferred OHT) are almost completely eliminated by the global correction.

From our results, we find that analysis increments are a useful tool for interpreting the trend estimates based on reanalysis data. Air-sea heat flux trends from ERA5 forecasts in the North Atlantic basin seem reliable in terms of sign (on sub-basin scale, see Fig. 7) and spatial structure, but we speculate that temporal inconsistencies in the late 1990s and 2000s [as shown by Mayer et al. (2022) for global ocean averages] and temporally varying analysis increments have a common cause which is the increasing number of observations that indicate a drier atmosphere than in the model climate. Further research is needed to fully understand their impact on both forecast and analysis-based trends.

### Appendix A: Linearized turbulent heat fluxes

Turbulent heat fluxes are linearized by decomposing each variable on the right side of Eq. (1) and (2) into a mean state (with overbar) and deviation from the mean (with prime); that is, we substitute  $\rho = \bar{\rho} + \rho'$ ,  $|U_{ml}| = \overline{|U_{ml}|} + |U_{ml}|'$ ,  $\Delta q = \overline{\Delta q} + \Delta q'$ , and  $\Delta T = \overline{\Delta T} + \Delta T'$ , where  $\Delta q = q_{ml} - q_{sfc}$  and  $\Delta T = T_{ml} - T_{skin}$ . After some calculus, turbulent heat fluxes can be separated into a non-linear and linear part, where the former contains all products with more than one deviation term (e.g., the non-linear term  $\overline{|U_{ml}|} \rho' \Delta q'$ ; not shown). The linear latent heat flux can be written as

$$F_{LH,linear} = C_Q L_v \left( \overline{|U_{ml}|} \bar{\rho} \overline{\Delta q} + \overline{|U_{ml}|} \bar{\rho} \Delta q' + \overline{|U_{ml}|} \rho' \overline{\Delta q} + |U_{ml}|' \bar{\rho} \overline{\Delta q} \right), \quad (A1)$$

and the linear sensible heat flux as

$$F_{SH,linear} = C_H c_p \left( \overline{|U_{ml}|} \bar{\rho} \overline{\Delta T} + \overline{|U_{ml}|} \bar{\rho} \Delta T' + \overline{|U_{ml}|} \rho' \overline{\Delta T} + |U_{ml}|' \bar{\rho} \overline{\Delta T} \right) + C_{SH} g z \left( \overline{|U_{ml}|} \bar{\rho} + \overline{|U_{ml}|} \rho' + |U_{ml}|' \bar{\rho} \right). \quad (A2)$$

### Appendix B: Moisture and temperature increments

Figure B1 shows 1985–2019 trends of moisture and temperature increments and corresponding TNA box averages for the whole study period. Note that moisture increments in the tropical North Atlantic before 2000 are remarkably stable around zero but rapidly decrease afterward, with minima values of about  $-0.2 \text{ g kg}^{-1}$  in 2010–15 (negative analysis increments mean that moisture is removed from the model by the data assimilation). Temperature increments show a weak increase in the early 1990s but are temporally stable between 0–0.1 K otherwise.



## 575 **Appendix C: NAO and AMO regression onto air-sea heat fluxes**

Figure C1 shows winter-months partial trends of NAO and AMO as regressed onto air-sea heat fluxes from ERA5 forecasts for the period 1950–2019.

*Author contributions.* All authors participated in the discussion and conceptual design of the paper. JM prepared the figures and wrote the manuscript under the supervision of LH and MM.

580 *Competing interests.* The contact author has declared that none of the authors has any competing interests.

*Acknowledgements.* JM and MM were financially supported by the Austrian Science Funds (FWF) project P33177. LH received support from the Austrian HRSM project GEOCLIM.

## References


- Baehr, J., Haak, H., Alderson, S., Cunningham, S. A., Jungclaus, J. H., and Marotzke, J.: Timely Detection of Changes in the Meridional  
585 Overturning Circulation at 26°N in the Atlantic, *Journal of Climate*, 20, 5827 – 5841, <https://doi.org/10.1175/2007JCLI1686.1>, 2007.
- Baker, J. A., Renshaw, R., Jackson, L. C., Dubois, C., Iovino, D., Zuo, H., Perez, R. C., Dong, S., Kersalé, M., Mayer, M., Mayer, J., Speich, S., and Lamont, T.: Overturning and heat transport variations in the South Atlantic in an ocean reanalysis ensemble, *State of the Planet Discussions*, 2022, 1–17, <https://doi.org/10.5194/sp-2022-8>, 2022.
- Bell, B., Hersbach, H., Simmons, A., Berrisford, P., Dahlgren, P., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Radu, R., Schepers, D., Soci,  
590 C., Villaume, S., Bidlot, J.-R., Haimberger, L., Woollen, J., Buontempo, C., and Thépaut, J.-N.: The ERA5 global reanalysis: Preliminary extension to 1950, *Quarterly Journal of the Royal Meteorological Society*, 147, 4186–4227, <https://doi.org/https://doi.org/10.1002/qj.4174>, 2021.
- Bengtsson, L., Hagemann, S., and Hodges, K. I.: Can climate trends be calculated from reanalysis data?, *Journal of Geophysical Research: Atmospheres*, 109, <https://doi.org/https://doi.org/10.1029/2004JD004536>, 2004.
- 595 Boers, N.: Observation-based early-warning signals for a collapse of the Atlantic Meridional Overturning Circulation, *Nature Climate Change*, 11, 680–688, <https://doi.org/10.1038/s41558-021-01097-4>, 2021.
- Bryden, H. L., Johns, W. E., King, B. A., McCarthy, G., McDonagh, E. L., Moat, B. I., and Smeed, D. A.: Reduction in Ocean Heat Transport at 26°N since 2008 Cools the Eastern Subpolar Gyre of the North Atlantic Ocean, *Journal of Climate*, 33, 1677 – 1689, <https://doi.org/10.1175/JCLI-D-19-0323.1>, 2020.
- 600 Caesar, L., Rahmstorf, S., Robinson, A., Feulner, G., and Saba, V.: Observed fingerprint of a weakening Atlantic Ocean overturning circulation, *Nature*, 556, 191–196, <https://doi.org/10.1038/s41586-018-0006-5>, 2018.
- Chemke, R., Zanna, L., and Polvani, L. M.: Identifying a human signal in the North Atlantic warming hole, *Nature Communications*, 11, 1–7, <https://doi.org/10.1038/s41467-020-15285->, 2020.
- Chen, J., Carlson, B. E., and Genio, A. D. D.: Evidence for Strengthening of the Tropical General Circulation in the 1990s, *Science*, 295,  
605 838–841, <https://doi.org/10.1126/science.1065835>, 2002.
- Cheng, L. and Zhu, J.: Benefits of CMIP5 Multimodel Ensemble in Reconstructing Historical Ocean Subsurface Temperature Variations, *Journal of Climate*, 29, 5393 – 5416, <https://doi.org/10.1175/JCLI-D-15-0730.1>, 2016.
- Cheng, L., Trenberth, K. E., Fasullo, J., Boyer, T., Abraham, J., and Zhu, J.: Improved estimates of ocean heat content from 1960 to 2015, *Science Advances*, 3, e1601 545, <https://doi.org/10.1126/sciadv.1601545>, 2017.
- 610 Chiodo, G. and Haimberger, L.: Interannual changes in mass consistent energy budgets from ERA-Interim and satellite data, *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/10.1029/2009JD012049>, 2010.
- Cohen, J. and Barlow, M.: The NAO, the AO, and Global Warming: How Closely Related?, *Journal of Climate*, 18, 4498 – 4513, <https://doi.org/10.1175/JCLI3530.1>, 2005.
- Cronin, M. F., Gentemann, C. L., Edson, J., Ueki, I., Bourassa, M., Brown, S., Clayson, C. A., Fairall, C. W., Farrar, J. T., Gille, S. T., Gulev, S., Josey, S. A., Kato, S., Katsumata, M., Kent, E., Krug, M., Minnett, P. J., Parfitt, R., Pinker, R. T., Stackhouse, P. W., Swart, S., Tomita, H., Vandemark, D., Weller, A. R., Yoneyama, K., Yu, L., and Zhang, D.: Air-Sea Fluxes With a Focus on Heat and Momentum, *Frontiers in Marine Science*, 6, 430, <https://doi.org/10.3389/fmars.2019.00430>, 2019.
- Docquier, D. and Koenigk, T.: A review of interactions between ocean heat transport and Arctic sea ice, *Environmental Research Letters*, 16, 123 002, <https://doi.org/10.1088/1748-9326/ac30be>, 2021.

- 620 Douville, H., Raghavan, K., Renwick, J., Allan, R., Arias, P., Barlow, M., Cerezo-Mota, R., Cherchi, A., Gan, T., Gergis, J., Jiang, D., Khan, A., Mba, W. P., Rosenfeld, D., Tierney, J., , and Zolina, O.: Water Cycle Changes, *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, p. 1055–1210, <https://doi.org/10.1017/9781009157896.010>, 2021.
- ECMWF: IFS Documentation CY47R3 - Part IV Physical processes, chap. IV, pp. 1–232, 4, ECMWF, <https://doi.org/10.21957/eyrpir4vj>,  
625 2021.
- Fairall, C. W., Bradley, E. F., Hare, J. E., Grachev, A. A., and Edson, J. B.: Bulk Parameterization of Air–Sea Fluxes: Updates and Verification for the COARE Algorithm, *Journal of Climate*, 16, 571 – 591, [https://doi.org/10.1175/1520-0442\(2003\)016<0571:BPOASF>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<0571:BPOASF>2.0.CO;2), 2003.
- Fasullo, J. T. and Trenberth, K. E.: The Annual Cycle of the Energy Budget. Part I: Global Mean and Land–Ocean Exchanges, *Journal of*  
630 *Climate*, 21, 2297–2312, <https://doi.org/10.1175/2007JCLI1935.1>, 2008.
- Fox-Kemper, B., Hewitt, H., Xiao, C., Aðalgeirsdóttir, G., Drijfhout, S., Edwards, T., Golledge, N., Hemer, M., Kopp, R., Krinner, G., Mix, A., Notz, D., Nowicki, S., Nurhati, I., Ruiz, L., Sallée, J.-B., Slangen, A., and Yu, Y.: Ocean, Cryosphere and Sea Level Change, *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, pp. 1211 – 1362, <https://doi.org/10.1017/9781009157896.011>, 2021.
- 635 Gillett, N. P., Graf, H. F., and Osborn, T. J.: Climate Change and the North Atlantic Oscillation, pp. 193–209, *American Geophysical Union (AGU)*, <https://doi.org/https://doi.org/10.1029/134GM09>, 2003.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J.,  
640 Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 Global Reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R., Hólm, E.,  
645 Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: Complete ERA5 global atmospheric reanalysis: Fifth generation of ECMWF atmospheric reanalyses of the global climate, <https://doi.org/10.24381/CDS.143582CF>, accessed on 04-07-2023, 2023b.
- Hodson, D. L. R., Robson, J. I., and Sutton, R. T.: An Anatomy of the Cooling of the North Atlantic Ocean in the 1960s and 1970s, *Journal*  
650 *of Climate*, 27, 8229 – 8243, <https://doi.org/10.1175/JCLI-D-14-00301.1>, 2014.
- Hurrell, J. W.: Decadal Trends in the North Atlantic Oscillation: Regional Temperatures and Precipitation, *Science*, 269, 676–679, <https://doi.org/10.1126/science.269.5224.676>, 1995.
- Hurrell, J. W. and Deser, C.: North Atlantic climate variability: The role of the North Atlantic Oscillation, *Journal of Marine Systems*, 78, 28–41, <https://doi.org/https://doi.org/10.1016/j.jmarsys.2008.11.026>, 2009.
- 655 Johns, W. E., Baringer, M. O., Beal, L. M., Cunningham, S. A., Kanzow, T., Bryden, H. L., Hirschi, J. J. M., Marotzke, J., Meinen, C. S., Shaw, B., and Curry, R.: Continuous, Array-Based Estimates of Atlantic Ocean Heat Transport at 26.5°N, *Journal of Climate*, 24, 2429 – 2449, <https://doi.org/10.1175/2010JCLI3997.1>, 2011.

- Kerr, R. A.: A North Atlantic Climate Pacemaker for the Centuries, *Science*, 288, 1984–1985, <https://doi.org/10.1126/science.288.5473.1984>, 2000.
- 660 Kossin, J. P., Knapp, K. R., Olander, T. L., and Velden, C. S.: Global increase in major tropical cyclone exceedance probability over the past four decades, *Proceedings of the National Academy of Science*, 117, 11 975–11 980, <https://doi.org/10.1073/pnas.1920849117>, 2020.
- Liu, C., Allan, R. P., Mayer, M., Hyder, P., Loeb, N. G., Roberts, C. D., Valdivieso, M., Edwards, J. M., and Vidale, P.-L.: Evaluation of satellite and reanalysis-based global net surface energy flux and uncertainty estimates, *Journal of Geophysical Research: Atmospheres*, 122, 6250–6272, <https://doi.org/10.1002/2017JD026616>, 2017.
- 665 Liu, C., Allan, R. P., Mayer, M., Hyder, P., Desbruyères, D., Cheng, L., Xu, J., Xu, F., and Zhang, Y.: Variability in the global energy budget and transports 1985–2017, *Climate Dynamics*, <https://doi.org/10.1007/s00382-020-05451-8>, 2020.
- Liu, C., Yang, Y., Liao, X., Cao, N., Liu, J., Ou, N., Allan, R. P., Jin, L., Chen, N., and Zheng, R.: Discrepancies in Simulated Ocean Net Surface Heat Fluxes over the North Atlantic, *Advances in Atmospheric Sciences*, 39, 1941–1955, <https://doi.org/10.1007/s00376-022-1360-7>, 2022.
- 670 Loeb, N. G., Mayer, M., Kato, S., Fasullo, J. T., Zuo, H., Senan, R., Lyman, J. M., Johnson, G. C., and Balmaseda, M.: Evaluating Twenty-Year Trends in Earth’s Energy Flows From Observations and Reanalyses, *Journal of Geophysical Research: Atmospheres*, 127, e2022JD036 686, <https://doi.org/https://doi.org/10.1029/2022JD036686>, 2022.
- Mathew, S. S. and Kumar, K. K.: Characterization of the long-term changes in moisture, clouds and precipitation in the ascending and descending branches of the Hadley Circulation, *Journal of Hydrology*, 570, 366–377, <https://doi.org/https://doi.org/10.1016/j.jhydrol.2018.12.047>, 2019.
- 675 Mayer, J., Mayer, M., and Haimberger, L.: Consistency and Homogeneity of Atmospheric Energy, Moisture, and Mass Budgets in ERA5, *Journal of Climate*, 34, 3955 – 3974, <https://doi.org/10.1175/JCLI-D-20-0676.1>, 2021.
- Mayer, J., Mayer, M., Haimberger, L., and Liu, C.: Comparison of Surface Energy Fluxes from Global to Local Scale, *Journal of Climate*, 35, 4551 – 4569, <https://doi.org/10.1175/JCLI-D-21-0598.1>, 2022.
- 680 Mayer, J., Mayer, M., and Haimberger, L.: Mass-consistent atmospheric energy and moisture budget monthly data from 1979 to present derived from ERA5 reanalysis, v1.0, <https://doi.org/10.24381/cds.c2451f6b>, accessed on 04-07-2023, 2022b.
- Mayer, M. and Haimberger, L.: Poleward Atmospheric Energy Transports and Their Variability as Evaluated from ECMWF Reanalysis Data, *Journal of Climate*, 25, 734–752, <https://doi.org/10.1175/JCLI-D-11-00202.1>, 2012.
- Mayer, M., Haimberger, L., Pietschnig, M., and Storto, A.: Facets of Arctic energy accumulation based on observations and reanalyses 2000–2015, *Geophysical Research Letters*, 43, 10,420–10,429, <https://doi.org/https://doi.org/10.1002/2016GL070557>, 2016.
- 685 Mayer, M., , Tietsche, S., Haimberger, L., Tsubouchi, T., Mayer, J., and Zuo, H.: An improved estimate of the coupled Arctic energy budget, *Journal of Climate*, 32, 7915–7934, <https://doi.org/10.1175/JCLI-D-19-0233.1>, 2019.
- McCarthy, G., Smeed, D., Johns, W., Frajka-Williams, E., Moat, B., Rayner, D., Baringer, M., Meinen, C., Collins, J., and Bryden, H.: Measuring the Atlantic Meridional Overturning Circulation at 26°N, *Progress in Oceanography*, 130, 91–111, <https://doi.org/https://doi.org/10.1016/j.pocean.2014.10.006>, 2015.
- 690 Mork, K. A., Øystein Skagseth, and Sjøiland, H.: Recent Warming and Freshening of the Norwegian Sea Observed by Argo Data, *Journal of Climate*, 32, 3695 – 3705, <https://doi.org/10.1175/JCLI-D-18-0591.1>, 2019.
- Muilwijk, M., Smedsrud, L. H., Ilicak, M., and Drange, H.: Atlantic Water Heat Transport Variability in the 20th Century Arctic Ocean From a Global Ocean Model and Observations, *Journal of Geophysical Research: Oceans*, 123, 8159–8179, <https://doi.org/https://doi.org/10.1029/2018JC014327>, 2018.
- 695


- Rahmstorf, S., Box, J. E., Feulner, G., Mann, M. E., Robinson, A., Rutherford, S., and Schaffernicht, E. J.: Exceptional twentieth-century slowdown in Atlantic Ocean overturning circulation, *Nature Climate Change*, 5, 475–480, <https://doi.org/10.1038/nclimate2554>, 2015.
- Roberts, C. D., Jackson, L., and McNeall, D.: Is the 2004-2012 reduction of the Atlantic meridional overturning circulation significant?, *Geophysical Research Letters*, 41, 3204–3210, <https://doi.org/https://doi.org/10.1002/2014GL059473>, 2014.
- 700 Shaman, J., Samelson, R. M., and Skillingstad, E.: Air–Sea Fluxes over the Gulf Stream Region: Atmospheric Controls and Trends, *Journal of Climate*, 23, 2651–2670, <https://doi.org/10.1175/2010JCLI3269.1>, 2010.
- Skagseth, Ø., Eldevik, T., Årthun, M., Asbjørnsen, H., Lien, V. S., and Smedsrud, L. H.: Reduced efficiency of the Barents Sea cooling machine, *Nature Climate Change*, 10, 661–666, <https://doi.org/10.1038/s41558-020-0772-6>, 2020.
- Tanimoto, Y., Nakamura, H., Kagimoto, T., and Yamane, S.: An active role of extratropical sea surface temperature anomalies in determining anomalous turbulent heat flux, *Journal of Geophysical Research: Oceans*, 108, <https://doi.org/https://doi.org/10.1029/2002JC001750>, 2003.
- 705 Trenberth, K. E.: Climate Diagnostics from Global Analyses: Conservation of Mass in ECMWF Analyses, *Journal of Climate*, 4, 707–722, [https://doi.org/10.1175/1520-0442\(1991\)004<0707:CDFGAC>2.0.CO;2](https://doi.org/10.1175/1520-0442(1991)004<0707:CDFGAC>2.0.CO;2), 1991.
- Trenberth, K. E. and Fasullo, J. T.: Atlantic meridional heat transports computed from balancing Earth’s energy locally, *Geophysical Research Letters*, 44, 1919–1927, <https://doi.org/10.1002/2016GL072475>, 2017.
- 710 Trenberth, K. E. and Shea, D. J.: Atlantic hurricanes and natural variability in 2005, *Geophysical Research Letters*, 33, <https://doi.org/https://doi.org/10.1029/2006GL026894>, 2006.
- Trenberth, K. E., Fasullo, J. T., and Mackaro, J.: Atmospheric Moisture Transports from Ocean to Land and Global Energy Flows in Reanalyses, *Journal of Climate*, 24, 4907–4924, <https://doi.org/10.1175/2011JCLI4171.1>, 2011.
- 715 Trenberth, K. E., Zhang, Y., Fasullo, J. T., and Cheng, L.: Observation-Based Estimates of Global and Basin Ocean Meridional Heat Transport Time Series, *Journal of Climate*, 32, 4567–4583, <https://doi.org/10.1175/JCLI-D-18-0872.1>, 2019.
- Tsubouchi, T., Bacon, S., Aksenov, Y., Garabato, A. C. N., Beszczynska-Möller, A., Hansen, E., de Steur, L., Curry, B., and Lee, C. M.: The Arctic Ocean Seasonal Cycles of Heat and Freshwater Fluxes: Observation-Based Inverse Estimates, *Journal of Physical Oceanography*, 48, 2029 – 2055, <https://doi.org/10.1175/JPO-D-17-0239.1>, 2018.
- 720 Tsubouchi, T., kjetil Vage, Hansen, B., Larsen, K., Osterhus, S., Johnson, C., Jonsson, S., and Valdimarsson, H.: Increased ocean heat transport into the Nordic Seas and Arctic Ocean over the period 1993-2016, *Nature Climate Change*, 2020, <https://doi.org/10.1038/s41558-020-00941-3>, 2020.
- Visbeck, M. H., Hurrell, J. W., Polvani, L., and Cullen, H. M.: The North Atlantic Oscillation: Past, present, and future, *Proceedings of the National Academy of Sciences*, 98, 12 876–12 877, <https://doi.org/10.1073/pnas.231391598>, 2001.
- 725 von Schuckmann, K., Traon, P.-Y. L., (Chair), N. S., Pascual, A., Djavidnia, S., Gattuso, J.-P., Grégoire, M., Aaboe, S., Alari, V., Alexander, B. E., Alonso-Martirena, A., Aydogdu, A., Azzopardi, J., Bajo, M., Barbariol, F., Batistić, M., Behrens, A., Ismail, S. B., Benetazzo, A., Bitetto, I., Borghini, M., Bray, L., Capet, A., Carlucci, R., Chatterjee, S., Chiggiato, J., Ciliberti, S., Cipriano, G., Clementi, E., Cochrane, P., Cossarini, G., D’Andrea, L., Davison, S., Down, E., Drago, A., Druon, J.-N., Engelhard, G., Federico, I., Garić, R., Gauci, A., Gerin, R., Geyer, G., Giesen, R., Good, S., Graham, R., Grégoire, M., Greiner, E., Gundersen, K., Hélaouët, P., Hendricks, S., Heymans, J. J., Holt, J., Hure, M., Juza, M., Kassis, D., Kellett, P., Knol-Kauffman, M., Kountouris, P., Kōuts, M., Lagemaat, P., Lavergne, T., Legeais, J.-F., Traon, P.-Y. L., Libralato, S., Lien, V. S., Lima, L., Lind, S., Liu, Y., Macías, D., Maljutenko, I., Mangin, A., Männik, A., Marinova, V., Martellucci, R., Masnadi, F., Mauri, E., Mayer, M., Menna, M., Meulders, C., Møgster, J. S., Monier, M., Mork, K. A., Müller, M., Øie Nilsen, J. E., Notarstefano, G., Oviedo, J. L., Palerme, C., Palialexis, A., Panzeri, D., Pardo, S., Peneva, E., Pezzutto, P., Pirro, A., Platt, T.,

- 735 Poulain, P.-M., Prieto, L., Querin, S., Rabenstein, L., Raj, R. P., Raudsepp, U., Reale, M., Renshaw, R., Ricchi, A., Ricker, R., Rikka, S., Ruiz, J., Russo, T., Sanchez, J., Santoleri, R., Sathyendranath, S., Scarcella, G., Schroeder, K., Sparnocchia, S., Spedicato, M. T., Stanev, E., Staneva, J., Stocker, A., Stoffelen, A., Teruzzi, A., Townhill, B., Uiboupin, R., Valcheva, N., Vandenbulcke, L., Vindenes, H., von Schuckmann, K., Vrgoč, N., Wakelin, S., and Zupa, W.: Copernicus Marine Service Ocean State Report, Issue 5, *Journal of Operational Oceanography*, 14, 1–185, <https://doi.org/10.1080/1755876X.2021.1946240>, 2021.
- 740 Wang, Q., Wang, X., Wekerle, C., Danilov, S., Jung, T., Koldunov, N., Lind, S., Sein, D., Shu, Q., and Sidorenko, D.: Ocean Heat Transport Into the Barents Sea: Distinct Controls on the Upward Trend and Interannual Variability, *Geophysical Research Letters*, 46, 13 180–13 190, <https://doi.org/https://doi.org/10.1029/2019GL083837>, 2019.
- Wang, Q., Wekerle, C., Wang, X., Danilov, S., Koldunov, N., Sein, D., Sidorenko, D., von Appen, W.-J., and Jung, T.: Intensification of the Atlantic Water Supply to the Arctic Ocean Through Fram Strait Induced by Arctic Sea Ice Decline, *Geophysical Research Letters*, 47, e2019GL086 682, <https://doi.org/https://doi.org/10.1029/2019GL086682>, 2020.
- 745 Worthington, E. L., Moat, B. I., Smeed, D. A., Mecking, J. V., Marsh, R., and McCarthy, G. D.: A 30-year reconstruction of the Atlantic meridional overturning circulation shows no decline, *Ocean Science*, 17, 285–299, <https://doi.org/10.5194/os-17-285-2021>, 2021.
- Yang, H., Lohmann, G., Wei, W., Dima, M., Ionita, M., and Liu, J.: Intensification and poleward shift of subtropical western boundary currents in a warming climate, *Journal of Geophysical Research: Oceans*, 121, 4928–4945, <https://doi.org/10.1002/2015JC011513>, 2016.
- 750 Zhang, R., Sutton, R., Danabasoglu, G., Kwon, Y.-O., Marsh, R., Yeager, S. G., Amrhein, D. E., and Little, C. M.: A Review of the Role of the Atlantic Meridional Overturning Circulation in Atlantic Multidecadal Variability and Associated Climate Impacts, *Reviews of Geophysics*, 57, 316–375, <https://doi.org/https://doi.org/10.1029/2019RG000644>, 2019.




**Figure 1.** Mean 1985–2019 December–February climatology of model-based **net air-sea heat fluxes from ERA5 forecasts**  $F_{g}$ . Black boxes indicate the four areas of interest located in the Norwegian Sea (NWS), North Atlantic Warming Hole (NAWH), Gulf Stream (GS), and tropical North Atlantic (TNA). The red lines mark the mooring locations in the Davis Strait (DS), Fram Strait (FS), and Barents Sea Opening (BSO), which are used to indirectly estimate the ocean heat transport in the North Atlantic basin. Contour lines are shown for 0,  $\pm 200$ , and  $\pm 400 \text{ W m}^{-2}$ .







**Figure 2.** Linear trends of a) model-based net surface heat fluxes from ERA5 forecasts  $F_S$  and b) inferred heat fluxes derived from atmospheric energy transports and TOA radiation  $F_S$  for the period 1985–2019. Panel Panels c) and d) show the turbulent and radiative energy flux component components of model-based  $F_S$  trends, and panel panels e) and f) illustrate latent and sensible heat fluxes separately. All trends are computed based from DJF means of anomalies. Units are  $\text{W m}^{-2} \text{dec}^{-1}$ . The shading represents areas of statistically significant trends (95 % confidence level).




**Figure 3.** Linear DJF trends of analyzed a) model-level humidity, b) model-level temperature, c) surface saturation humidity, and d) skin temperature anomalies for 1985–2019. In addition, the difference between model-level and surface e) humidity and f) temperature is shown. Units are  $\text{g kg}^{-1} \text{dec}^{-1}$  for humidity trends and  $\text{K dec}^{-1}$  for temperature trends. The shading represents areas of statistically significant trends (95 % confidence level).




**Figure 4.** Linear trend of analyzed a) 10 metre horizontal wind speed and direction anomalies ( $\text{m s}^{-1} \text{dec}^{-1}$ ) and b) model-level relative humidity ( $\% \text{dec}^{-1}$ ) for DJF 1985–2019. Anomalous wind direction trends are illustrated by black arrows (with a maximum of  $\sim 0.8 \text{ m s}^{-1} \text{dec}^{-1}$ ). The shading represents areas of statistically significant trends (95 % confidence level).



**Figure 5.** Linear trend of the meridional mass stream function from analyzed ERA5 winds over the period 1985 to 2019. Positive trends are shown as solid contour lines, negative trends as dotted lines. The stream function is integrated over 360 degrees in longitude. Units are  $10^9 \text{ kg s}^{-1} \text{ dec}^{-1}$ .




**Figure 6.** As in Fig. 2e and f, but for 1950–84.




**Figure 7.** (left) DJF anomalies and (middle) full-year climatology of model-based  $F_S$  from ERA5 forecasts (solid lines) and inferred  $F_S$  (dotted lines) for box averages (see Fig. 1) in the Norwegian Sea (NWS), North Atlantic Warming Hole (NAWH), Gulf Stream (GS), and Tropical North Atlantic (TNA). (right) Partial trends [see Eq. (3)] of 10 metre wind speed (ws), model-level humidity ( $q_{m1}$ ) and temperature ( $T_{m1}$ ), surface saturation humidity ( $q_{sfc}$ ), and skin temperature ( $T_{sfc}$ ) as regressed onto latent (LHF) and sensible heat fluxes (SHF), respectively. Anomalies Climatologies and anomalies are computed w.r.t. 1985–2019. Trends for 1950–84 (1985–2019) are shown in blue (red). The grey background in the middle panel highlights the boreal winter months December–February. Units are  $W m^{-2}$  for anomalies and annual cycles, and  $W m^{-2} dec^{-1}$  for trends.






**Figure 8.** Linear trend of a) model-based **surface heat fluxes from ERA5 forecasts**  $F_S$  for 1950–2019, and b) the same but with the partial NAO trend subtracted (see appendix C for regression pattern). The shading represents areas of statistically significant trends (95 % confidence level).






**Figure 9.** (left) 1950–2019 full-year trend of indirectly estimated Atlantic ocean heat transport as derived from the oceanic heat budget and (middle) the same but without OHCT data employed. Blue (grey) lines are trend estimates based on 5-year (monthly) means. The shading illustrates the 95 % confidence interval of the trend estimate. (right) 1950–2019 mean total indirectly estimated heat transport at each latitude. The red dot shows the 2004–2018 mean observed ocean heat transport through the RAPID array at 26.5° N.



**Figure B1.** Analysis increments (analysis minus forecast field) of (left) moisture and (right) temperature at [the](#) lowest model level. The upper panel shows DJF trend maps of analysis increments for the period 1985–2019. The lower panel shows analysis increments of the TNA box average for the whole study period. Time series are smoothed by a 12-month moving average.



**Figure C1.** Partial trends of (left) NAO and (right) AMO as regressed onto air-sea heat fluxes from ERA5 forecasts for the period 1950–2019. Partial trends are computed for each grid point by multiplying the sensitivity between climate index and  $F_S$  with the linear trend of the climate index [see explanation to Eq. (3)].

**Table 1.** Flux trends for various areas and periods of time. The study area refers to the ocean area between 0–90° N and 90° W–30° E. Nordic seas include the ocean area between 60–82° N and 45° W–30° E. Units are  $\text{W m}^{-2} \text{dec}^{-1}$ . Bold values are statistically significant trends at the 95 % confidence level. Note that the difference between globally adjusted  $F_S$  and model-based  $F_S$  in each period is the magnitude of the global adjustment (see section 3) and can also be added to other model-based  $F_S$  trend estimates of that particular period.

Averaging area	Term	DJF trend		
		1950–84	1985–2019	1950–2019
Study area	Latent heat flux	<b>1.3 ± 0.7</b>	<b>-1.9 ± 1.4</b>	-0.4 ± 0.6
	Sensible heat flux	0.2 ± 0.3	0.3 ± 0.3	<b>0.3 ± 0.1</b>
	Radiative fluxes	<b>0.4 ± 0.3</b>	0.1 ± 0.4	<b>0.3 ± 0.1</b>
	Model-based $F_S$	<b>1.9 ± 1.2</b>	-1.4 ± 1.9	0.1 ± 0.7
	Globally adjusted $F_S$	<b>3.2 ± 1.7</b>	1.2 ± 2.2	<b>1.6 ± 0.7</b>
	Inferred $F_S$	–	0.3 ± 1.4	–
NWS box	Model-based $F_S$	0.8 ± 9.3	<b>9.7 ± 8.0</b>	<b>4.8 ± 3.1</b>
	Inferred $F_S$	–	<b>12.6 ± 9.8</b>	–
NAWH box	Model-based $F_S$	2.5 ± 12.4	5.2 ± 12.5	1.8 ± 4.1
	Inferred $F_S$	–	7.0 ± 15.0	–
GS box	Model-based $F_S$	5.9 ± 12.4	-9.4 ± 13.2	-0.7 ± 4.5
	Inferred $F_S$	–	-4.8 ± 12.6	–
TNA box	Model-based $F_S$	-2.9 ± 4.5	-2.7 ± 4.2	<b>-4.7 ± 1.5</b>
	Inferred $F_S$	–	-2.3 ± 4.3	–
Nordic Seas	Sensible heat flux	0.0 ± 2.4	1.9 ± 2.1	<b>0.9 ± 0.8</b>

**Table 2.** Full-year trends of globally adjusted air-sea heat fluxes from ERA5 forecasts, ocean heat content tendency (OHCT), and indirectly estimated ocean heat transport (OHT) at 26° N for the period 1950–2019.  $F_S$  and OHCT are spatially integrated over the ocean area between the choke point in the north (see red lines in Fig. 1) and 26° N. Trends are estimated based on full-year monthly means and 5-year means. Bold values are statistically significant trends at the 95 % confidence level. The left column contains area-averaged values (relative to the area between the northern choke point and 26° N) given in  $\text{W m}^{-2} \text{dec}^{-1}$ . The values in the right column represent the area-integrated contribution to the ocean heat budget given in  $\text{PW dec}^{-1}$ .

Method	Term	Full-year trend	
		$[\text{W m}^{-2} \text{dec}^{-1}]$	$[\times 10^{-2} \text{PW dec}^{-1}]$
Monthly mean	Globally adjusted $F_S$	<b><math>0.9 \pm 0.4</math></b>	<b><math>2.3 \pm 1.1</math></b>
	OHCT	$0.2 \pm 0.9$	$0.5 \pm 2.2$
	OHT at 26° N	$-0.7 \pm 0.9$	$-1.8 \pm 2.1$
5-year mean	Globally adjusted $F_S$	<b><math>0.9 \pm 0.5</math></b>	<b><math>2.3 \pm 1.3</math></b>
	OHCT	$0.2 \pm 0.4$	$0.5 \pm 1.0$
	OHT at 26° N	<b><math>-0.7 \pm 0.6</math></b>	<b><math>-1.8 \pm 1.4</math></b>